Langlands correspondence and Bezrukavnikov's equivalence

Anna Romanov and Geordie Williamson

October 23, 2021

This document contains notes (taken by the first author) from a course (taught by the second author) at the University of Sydney. The course took place over two extended semesters, and consisted of two hours of lectures per week between 2019-2020. The course is divided in two parts, reflecting the two terms.

The first part (concluding with Lecture 11) is an attempt to give an informal introduction to what the Langlands program is about, from an arithmetical point of view. We assume the audience (like the lecturer) is a beginner in this subject, but had a first course in complex analysis, Galois theory, topology and representation theory. At times we also assume background in algebraic geometry. Not much is proved, but we try to give enough detail to convince the reader that there is a lot of marvellous mathematics here.

The second longer part (Lectures 12 through 34) tries to give enough background in geometric representation theory to understand Bezrukavnikov's equivalence. Taking the tamely ramified local Langlands correspondence as motivation, we pass through work of Deligne-Lusztig and Ginzburg giving a coherent construction of the affine Hecke algebra. Here we are roughly following the book of Chriss-Ginzburg, but our route is different at times. Bezrukavnikov's equivalence is a categorification of this isomorphism, and is our aim for the rest of the notes. In order to even understand the statement of Bezrukavnikov's equivalence, we need much of the toolbox of modern geometric representation theory: perverse sheaves; highest weight categories; Koszul duality; the geometric Satake equivalence, etc. A major role is also played by monoidal categories and their actions ("higher representation theory"). We try to spend enough time on each of these topics to make students feel somewhat comfortable with the ideas. It goes without saying that all of this took much longer to cover than anyone expected, and at the very end we arrive at a statement of Bezrukavnikov's equivalence. Its proof will have to wait for the next lecture series!

The spirit of these notes is very informal, and this is intentional. We hope that they are nonetheless useful for a casual reader trying to orient themselves in a fascinating but potentially intimidating landscape. We assume that the reader is willing to take some things on faith, and have tried to be honest. Audience members were encouraged to do exercises throughout, and this wouldn't be bad advice for any potential reader either.

Contents

Lectur	e 1: Reciprocity Laws	
1.1	Reciprocity Laws	7
1.2	0	9
1.3	What is going on here? What does this have to do with representation theory?	9
1.4	Schematic picture of the Langlands correspondence	0
1.5	Chebotarev density theorem	1
Lectur	e 2: Review of some algebraic number theory	
2.1	Number fields	3
2.2	An analogy	4
2.3	The fundamental exact sequence	5
2.4	Ramification	6
2.5	The case of Galois extensions	8
Lectur	e 3: L-functions	
3.1	The Riemann ζ -function	0
3.2	Why do we care? $\ldots \ldots 2$	1
3.3	Dirichlet L -functions $\ldots \ldots \ldots$	2
3.4	Dedekind ζ -functions	3
3.5	Walking across the bridge	4
3.6	Artin L-functions	5
Lectur	e 4: The Sato-Tate conjecture	
4.1	Equidistribution in representation theory	7
4.2	Elliptic curves and the Sato-Tate conjecture	0
4.3		4
Lectur	e 5: Infinite Galois theory and global class field theory	
5.1		7
5.2		9
5.3	Global class field theory á la Artin	0
Lectur	e 6: Local Galois groups and local class field theory	
6.1	A potted history of class field theory	3
6.2	A trip across the bridge	3
6.3	- 0	6
6.4	•	9
Lectur	e 7: Representation theory of <i>p</i> -adic groups	
7.1	Tying up some loose ends	1
7.2	The no small subgroups argument	
7.3		3
7.4	-	5
	e 8: Precise statement of local Langlands for $GL_2, p \neq 2$	
8.1	Basic representation theory of <i>p</i> -adic groups, continued	8
8.2	Canonical norms	
8.3	Smooth admissible representations of $\operatorname{GL}_1(K)$	
	i i ()	

8.4	Smooth admissible representations of $GL_2(K)$	•					61
8.5	Weil-Deligne representations						64
8.6	The local Langlands correspondence for GL_2	•		•			65
Lecture	e 9: The case of $p = 2$ and the Satake isomorphism						
9.1	Ramification filtration revisited						66
9.2	More details on Weil–Deligne representations	•					68
9.3	Why is LLC for $p = 2$ special?						69
9.4	Unramified representations						70
9.5	Hecke algebras	•					70
Lecture	e 10: The big picture						
	The very big picture						74
	10.1.1 Dimension 0						74
	10.1.2 Dimension ≥ 1						75
10.2	Local Langlands correspondence for split groups						78
	The Deligne–Langlands conjecture						78
	Geometric Satake equivalence						80
	Bezrukavnikov's equivalence						80
	e 11: Review of the first semester						
	The local Langlands correspondence						81
	The unramified story						82
	The tamely ramified with unipotent monodromy (TRUM) story .						83
	Affine Weyl groups and affine Hecke algebras						84
	e 12: The Deligne-Langlands conjecture		•	•	•	•	01
	The Iwahori–Matsumoto Hecke algebra						88
	The Deligne–Langlands conjecture						90
							90 92
	Geometric setting						92 94
	The Springer resolution	• •	• •	•	•	·	94
	e 13: Springer fibres and the Steinberg variety						0.0
	The Springer resolution, continued						96
	Many examples of Springer fibres						97
	The conormal space						101
13.4	The Steinberg variety	• •		•	•	•	103
	e 14: Springer correspondence and Borel-Moore homology						
	The Springer correspondence						105
	Borel-Moore homology						107
14.3	Geometric convolution algebras	• •		•			108
Lecture	e 15: The Kazhdan–Lusztig isomorphism						
15.1	More convolution algebras						111
15.2	Equivariant K -theory \ldots \ldots \ldots \ldots \ldots \ldots \ldots	•		· •			112
15.3	The Kazhdan–Lusztig isomorphism	•					114
	Bezrukavnikov's equivalence: rough outline						116
Lecture	e 16: The constructible side of Bezrukavnikov's equivalence						
16.1	Weil conjectures	•					118

0	121 123
Lecture 17: Constructible and perverse sheaves on curves	123
	125
	$125 \\ 126$
	131
Lecture 18: Beilinson gluing on curves	100
1 0	133
1 0 5	134
0	140
Lecture 19: The derived category of perverse sheaves	
19.1 Overview of Beilinson gluing	143
	144
19.3 $K(\pi, 1)$ spaces	145
19.4 Yoneda extensions	146
19.5 A key ingredient	147
19.6 Proof of Beilinson's theorem for curves	149
Lecture 20: Perverse sheaves for affine stratifications	
20.1 Highest weight categories	154
0 0 0	157
	160
Lecture 21: Equivariant K-theory of the Steinberg variety	
	163
	166
	168
Lecture 22: Proof of the Kazhdan–Lusztig isomorphism	
	170
*	171
	174
	114
Lecture 23: Gaitsgory's central sheaves	170
6	178
	178
	181
	186
Lecture 24: A hitchhiker's guide to the Hecke category	
	189
Č Č	190
	191
	192
24.5 Fourth (and final) try \ldots	195
Lecture 25: The categorical anti-spherical module and its symmetries	
25.1 The affine Hecke category	197
25.2 The categorical (anti-)spherical module	198

25.3 Symmetries of categories	201
25.4 A notational interlude:	202
25.5 The basic approach of Arkhipov-Bezrukavnikov	202
Lecture 26: Monoidal categories and their actions	
6	203
26.2 Modules for monoidal categories	204
26.3 Examples of modules over monoidal categories	205
	207
26.5 Example: Representations of the Verlinde category	208
Lecture 27: Abelian categories over stacks	
27.1 Modules over algebras in monoidal categories	211
27.2 Linear algebra over a stack	212
27.3 A brief review of descent	214
27.4 Back to sheaves over stacks	215
Lecture 28: (De)equivariantisation	
28.1 A simple example of this phenomenon	218
	219
28.3 Deequivariantisation	219
28.4 $\operatorname{Rep} G$ enrichments	220
28.5 Deequivariantisation principle	221
28.6 Tannakian formalism	223
Lecture 29: Coherent sheaves on base affine space	
29.1 Serre's description of coherent sheaves on \mathbb{P}^n	226
	227
	228
29.4 Arrows between coherent sheaves on $\overline{G/U}$	229
Lecture 30: Constructible sheaves on finite flag varieties and braid groups	
30.1 A fun calculation \ldots	231
30.2 The finite Hecke category	232
30.3 The Braid group	234
30.4 Further properties of the braid group action	237
Lecture 31: Affine flags, affine braids, and Wakimoto sheaves	
31.1 Wakimoto sheaves	241
31.2 Motivational interlude \ldots	243
31.3 Beilinson–Gaitsgory–Kottwitz (BGK) central sheaves	244
Lecture 32: Whittaker sheaves and Arkhipov-Bezrukavnikov theorem	
32.1 Averaging functors	248
32.2 Another fun calculation \ldots	248
	251
32.4 Bird's eye view of the rest of the proof \ldots	253
Lecture 33: Soergel bimodules, Soergel calculus, and BGK central sheaves	
	257
33.2 BGK central sheaves for $G = SL_2$	260

33.3 Braid group categorification	262
Lecture 34: Bezrukavnikov's equivalence	
34.1 Potted history of Koszul duality	265
34.2 A toy example \ldots	266
34.3 Torus monodromic sheaves á la Bezrukavnikov-Yun	267
34.4 Bezrukavnikov's equivalence	269

Lecture 1: Reciprocity Laws

If you do nothing else with this course this semester beyond attending the first lecture, you should at least try to read [Lan90].

1.1 Reciprocity Laws

We start at the natural starting place: an equation. Consider the equation

$$x^2 + 1 = 0.$$

If p is a prime, one might wonder: how many solutions does this equation have, modulo p? Some calculations will reveal the following table.

p	2	3	5	7	11	13	17	19	23]
# of sol's mod p	1	0	2	0	0	2	2	0	0]
$p \mod 4$	2	3	1	3	3	1	1	3	3	

We see a pattern. The prime 2 is weird, so we ignore it. But for the rest, it seems that

of solutions mod
$$p \neq 2 = \begin{cases} 2 & \text{if } p = 1 \mod 4, \\ 0 & \text{if } p = 3 \mod 4. \end{cases}$$

This pattern is surprising. It appears to be saying that there is a global rule governing the number of solutions mod p; that is, that the different primes somehow "talk to one another".

Here we can give a simple proof of why our claim above must be true. Assume $p \neq 2$. We have a short exact sequence

$$1 \to (\mathbb{F}_p^{\times})^2 \to \mathbb{F}_p^{\times} \to \{\pm 1\} \to 1,$$

where the third arrow is given by $x \mapsto x^{\frac{p-1}{2}}$. Therefore,

$$-1$$
 is a square mod $p \iff (-1)^{\frac{p-1}{2}} = 1 \iff \frac{p-1}{2}$ is even $\iff p = 1 \mod 4$.

Let's do another example. Consider the equation

$$x^2 - 3 = 0$$

We ask the same question and compute:

p	2	3	5	7	11	13	17	19	23	29	31	
# of sol's mod p	1	1	0	0	2	2	0	0	2	0	0]
$p \mod 12$	2	3	5	7	11	1	5	7	11	5	7	

Again, we have some small weird primes (2 and 3), so we throw them out. For the rest, we make a guess:

of solutions mod
$$p \neq 2, 3 = \begin{cases} 2 & \text{if } p = 1, 11 \mod 12, \\ 0 & \text{if } p = 5, 7 \mod 12. \end{cases}$$

To prove that this is indeed the case, we introduce a little more technology. Let $p \neq 2$ be a prime. Define

$$\epsilon(p) = \begin{cases} 0 & \text{if } p = 1 \mod 4, \\ 1 & \text{if } p = 3 \mod 4 \end{cases}$$

and the Legendre symbol

$$\left(\frac{x}{p}\right) = x^{\frac{p-1}{2}} \mod p = \begin{cases} 1 & \text{if } x \text{ is a square mod } p, \\ -1 & \text{if } x \text{ is not a square mod } p. \end{cases}$$

(For example, we saw above that $\left(\frac{-1}{p}\right) = (-1)^{\epsilon(p)}$.)

Theorem 1.1. (Gauss's Law) Let p, q be distinct primes $\neq 2$. Then

$$\left(\frac{p}{q}\right)\left(\frac{q}{p}\right) = (-1)^{\epsilon(p)\epsilon(q)}.$$

With this we can prove that our guess was correct. Assume $p \neq 2, 3$. Then

$$x^{2} - 3 \text{ has } 2 \text{ solutions mod } p \iff \left(\frac{3}{p}\right) = 1$$
$$\iff \left(\frac{p}{3}\right)(-1)^{\epsilon(p)\epsilon(3)} = 1$$
$$\iff \left(\frac{p}{3}\right)(-1)^{\epsilon(p)} = 1$$
$$\iff \begin{cases} p = 1 \mod 3 \text{ and } p = 1 \mod 4\\ p = 2 \mod 3 \text{ and } p = 3 \mod 4\\ \iff p = 1 \text{ or } -1 \mod 12. \end{cases}$$

These are examples of **reciprocity laws**. All polynomials of degree 2 can be worked out analogously to the ones above using quadratic reciprocity (Gauss's law). There was much activity on this problem starting with Gauss's work, which finally led to Artin's reciprocity law. This implied all known reciprocity laws at the time, and in particular treats polynomials of degree 3 and 4. However, we get stuck at 5. For example, consider

$$x^5 + 20x + 16 = 0.$$

We can construct a table

p	2	3	5	7	11	13	17	19	23	29	31	37	41	43	47	53
# of sol's mod p	1	0	1	2	2	0	2	0	1	2	0	1	2	2	0	0

but no obvious pattern emerges. (For a table that goes much further than this one, see the sheet on the course website.) It turns out that there is a pattern, but it is very well-hidden, and to find it, we need analysis.

1.2 Higher dimensional varieties

We could ask similar questions for polynomials in two variables. Consider the equation

$$y^2 = x^3 + 1.$$

How many solutions does this equation have modulo p? Let's try to answer this for one specific prime. Let p = 5, and we can compile our results in the following table:

$y \setminus x$	0	1	2	3	4
0	\times	×	\times	\times	\checkmark
1	\checkmark	×	×	×	×
2	×	×	\checkmark	×	×
3	×	×	\checkmark	×	×
4	\checkmark	×	×	×	×

So here we found that there are five solutions modulo 5. In general, how many solutions do we expect? Well, the map $x \mapsto x^3 + 1$ in \mathbb{F}_p is "roughly random," about half the elements of \mathbb{F}_p are squares, and for every square we get two solutions, so we expect *approximately* p *solutions*. But how often is this actually the case? We can measure the accuracy of this estimation by studying the **Sato-Tate error term**:

$$ST(p) = p - \#$$
(solutions modulo p).

Here is another table.

p	2	3	5	7	11	13	17	19	23	29	31	37	41	43	47	53]
# ST(p)	0	0	0	-4	0	2	0	8	0	0	-4	-10	0	8	0	0	

Notice how frequently the Sato-Tate error term is zero! We can now study this table and see if any patterns emerge. This is the content of the **Sato-Tate conjecture**, which is basically known thanks to recent work of Harris, Taylor, Clozel, and many others.

1.3 What is going on here? What does this have to do with representation theory?

Let $f(x) \in \mathbb{Z}[x]$ be an irreducible polynomial with integral coefficients. We can consider the **splitting field** of f

$$K = \mathbb{Q}(\alpha_1, \ldots, \alpha_n),$$

and the associated Galois group

$$\Gamma = \operatorname{Gal}(K/\mathbb{Q}),$$

which acts on the set of roots $\{\alpha_1, \ldots, \alpha_n\}$. As representation theorists, our natural instinct when we see a group action is to linearize. Doing this here results in the **permutation** representation

$$\Gamma \circlearrowright H = \bigoplus_{i=1}^n \mathbb{C}\alpha_i.$$

We can also consider the reduction of f modulo $p, \bar{f}(x) \in \mathbb{F}_p[x]$, as we did in the previous section. In general, \bar{f} will be reducible. If $p \nmid \Delta(f)$ (that is, p is not one of the "weird" primes we encountered earlier), then $\bar{f}(x)$ has n roots, $\bar{\alpha}_1, \ldots, \bar{\alpha}_n \in \mathbb{F}_{p^n}$. Recall that the Galois group $\operatorname{Gal}(\mathbb{F}_{p^n}/\mathbb{F}_p) = \mathbb{Z}/n\mathbb{Z}$ is generated by the Frobenius map

$$\overline{\mathrm{Frob}}_{\mathrm{p}}: x \mapsto x^p.$$

Then $\mathbb{F}_p = (\mathbb{F}_{p^n})^{\overline{\operatorname{Frob}_p}}$, and the number of solutions of \overline{f} is the number of fixed points of $\overline{\operatorname{Frob}_p}$ on $\{\overline{\alpha}_1, \ldots, \overline{\alpha}_n\}$. For such a p ("unramified") and after a choice ("prime in \mathbb{O} above p"), we get a bijection

$$\{\alpha_1,\ldots,\alpha_n\}\longrightarrow \{\overline{\alpha}_1,\ldots\overline{\alpha}_n\},\$$

and an element $\operatorname{Frob}_{p} \in \Gamma$ such that the action of Frob_{p} on $\{\alpha_{1}, \ldots, \alpha_{n}\}$ aligns with the action of $\overline{\operatorname{Frob}}_{p}$ on $\{\overline{\alpha}_{1}, \ldots, \overline{\alpha}_{n}\}$ under the bijection above.

Remark 1.2. Different choices of "prime in \mathbb{O} over p" result in conjugate Frob_p's. Hence, it is best to think of Frob_p as a conjugacy class instead of an individual element.

The upshot of the discussion above is that

solutions modulo
$$p = \#$$
 fixed points of Frob_p on $\{\overline{\alpha}_1, \ldots, \overline{\alpha}_n\}$
= # of fixed points of Frob_p on $\{\alpha_1, \ldots, \alpha_n\}$
= $\operatorname{Tr}(\operatorname{Frob}_p, H),$

where H is the permutation representation introduced at the beginning of this section. The number $Tr(Frob_p, H)$ is completely canonical - it doesn't depend on any of our choices! So we've reduced our question of finding solutions of polynomials modulo p to computing something that looks very much like the character of a representation.

The Punchline: If $p \nmid \Delta(f)$,

solutions of $f \mod p = \operatorname{Tr}(\operatorname{Frob}_{p}, H)$.

1.4 Schematic picture of the Langlands correspondence

(*Do not worry if this makes no sense!*) A caricature of the Langlands correspondence is captured in the diagram below.

"geometric" reps
$$V$$
 of $\operatorname{Gal}(\overline{\mathbb{Q}}/\mathbb{Q}) \longrightarrow$ "character" $\operatorname{Tr}(\operatorname{Frob}_{p}, V)$ automorphic forms
 \downarrow
L-functions (analytic)

From any "geometric" representation of $\operatorname{Gal}(\overline{\mathbb{Q}}/\mathbb{Q})$, we can take the trace of Frobenius, as we did in the previous section for the permutation representation H. We should think of this procedure as taking the character of the representation. To $\operatorname{Tr}(\operatorname{Frob}_p, V)$, we can attach the associated "*L*-function," which is an analytic object. (For example, when we start with the trivial representation of $\operatorname{Gal}(\overline{\mathbb{Q}}/\mathbb{Q})$, the resulting *L*-function is the Riemann ζ -function.) On the other hand, there is also a procedure for constructing *L*-functions from automorphic forms. The Langlands correspondence is an attempt to align these two sources of *L*-functions.

This is very deep. For example, two-dimensional representations of $\operatorname{Gal}(\mathbb{Q}/\mathbb{Q})$ result in Hecke *L*-functions, and the corresponding automorphic forms are modular forms. It turns out that working out the correspondence for 2-dimensional representations is enough to prove Fermat's last theorem.

1.5 Chebotarev density theorem

If we talk of $Tr(Frob_p, H)$ as a "character," we would like to know at least that the set $\{Frob_p\}$ for all p unramified cover the set of all conjugacy classes of Γ . This is a deep theorem.

Theorem 1.3. (Chebotarev density theorem) Fix a conjugacy class $C \subset \Gamma$. Then

$$\{p \text{ unramified } | \operatorname{Frob}_{p} = C\}$$

has density $|C|/|\Gamma|$.

Here density refers to either the natural density or the analytic density of the set of primes.

Example 1.4. Let $f(x) = x^2 + 1 \in \mathbb{Z}[x]$. The set of roots of f(x) is $\{i, -i\}$. The splitting field is $K = \mathbb{Q}(i)$ and $\operatorname{Gal}(\mathbb{Q}(i)/\mathbb{Q}) = \mathbb{Z}/2\mathbb{Z} = \{id, s\}$. In this example, all $p \neq 2$ are unramified. Then for such an unramified p,

$$\operatorname{Frob}_{p}: i \mapsto i^{p}$$
.

Hence,

$$\operatorname{Frob}_{p} = \begin{cases} id & \text{if } p = 1 \mod 4, \\ s & \text{if } p = 3 \mod 4. \end{cases}$$

Exercise 1.5. (Mandatory) Check that $Frob_p$ is indeed given as above!

Exercise 1.6. (Harder) By considering cyclotomic extensions (i.e. $\mathbb{Q}(e^{2\pi i/m})$), show that Chebotarev's density theorem implies Dirichlet's theorem on primes in arithmetic progression.

At the beginning of today's lecture, we discussed patterns in the number of solutions of a given polynomial modulo p. There is a sheet on the course webpage which shows tables of these patterns for the polynomials $x^2 + 1$, $x^2 - 3$, $x^2 + x + 1$, $x^2 + 2x + 3$, and $x^2 - x - 1$. A somewhat mysterious feature of these tables was the modulus appearing in the patterns. (For example, we showed that $x^2 - 3$ has two solutions modulo $p \neq 2, 3$ if and only if p = 1or 11 mod 12. Where did 12 come from?) We'll complete today's lecture with an example to demonstrate where this modulus comes from. **Example 1.7.** Consider the polynomial $f(x) = x^2 - x - 1$. We can see from the patterns on the handout that f(x) has 2 solutions mod p if p = 1 or 9 mod 10 and f(x) has 0 solutions mod p if p = 3 or 7 mod 10 (for p unramified). In this example, the splitting field is $K = \mathbb{Q}(\phi)$, where $\phi = \frac{1+\sqrt{5}}{2} = 2\cos(\pi/5)$ is the golden ratio, and $\operatorname{Gal}(\mathbb{Q}(\phi)/\mathbb{Q}) = \mathbb{Z}/2\mathbb{Z} = \langle s \rangle$. Note that $\phi = \zeta + \overline{\zeta}$, where $\zeta = e^{\pi i/5}$ is a fifth root of unity. Hence we can embed

$$K \hookrightarrow \mathbb{Q}(e^{2\pi i/10}) = \mathbb{Q}(\zeta)$$

via $\phi \mapsto \zeta + \overline{\zeta}$. As in the last example, for unramified p,

$$\operatorname{Frob}_{p}: \zeta \mapsto \zeta^{p}$$

Hence,

$$\operatorname{Frob}_{p}: \begin{cases} \phi \mapsto \phi & \text{if } p = 1 \text{ or } 9 \mod 10, \\ \phi \mapsto s(\phi) & \text{if } p = 3 \text{ or } 7 \mod 10. \end{cases}$$

In general, the modulus for quadratic fields are determined by embeddings of the splitting field into cyclotomic fields:

$$K \hookrightarrow \mathbb{Q}(e^{2\pi i/\text{modulus}})$$

Remark 1.8. Consider the degree 5 polynomial $f(x) = x^5 + 20x + 16$ we discussed in the first section. In all of the primes that occurred in our table, f(x) had either 0 or 2 solutions. However, we would expect that for certain primes, f(x) should have 5 solutions by Chebotarev's theorem. Should we be worried that we haven't seen any 5's in our chart? Geordie reassures us that we shouldn't be worried. In this example, $\Gamma = A_5$ has order 60. Then, by our discussion earlier,

solutions modulo
$$p = 5 \iff$$
 all solutions in \mathbb{F}_{p^m} are defined over \mathbb{F}_p
 \iff all solutions are fixed by $\overline{\text{Frob}}_p$
 \iff $\overline{\text{Frob}}_p = id.$

Since *id* is in its own conjugacy class, we expect f(x) to have 5 solutions modulo p about 1/60th of the time by Chebotarev's density theorem. We included fewer than 60 primes in our table, so we shouldn't be surprised that we haven't seen this happen yet.

It may seem like considering the number of solutions of a polynomial over a finite field is a cute, but not particularly important problem. However, it is actually of fundamental importance in number theory. A **number field** is a finite extension of \mathbb{Q} . All number fields (which are Galois extensions) are splitting fields of polynomials $f(x) \in \mathbb{Z}[x]$. One of the the most basic open questions in number theory is the following:

Question 1.9. How many number fields are there?

We can determine the field extension K corresponding to the polynomial f(x) by reducing mod p:

Theorem 1.10. The set $\{p \mid p \text{ unramified and } f(x) \text{ splits completely mod } p\}$ completely determines K.

So our motivational problem may have been cute, but it certainly isn't unimportant.

Lecture 2: Review of some algebraic number theory

Last time we discussed how by Chebotarev's density theorem, the equation $f(x) = x^5 + 20x + 16$ should have five solutions modulo p about $1/60^{th}$ of the time. Joel (+ a computer) computed that in the set of all primes below 500,000, there are 16,613 where f(x) has no solutions, 10,367 where f(x) has one solution, 13,885 with two solutions, and 673 with five solutions. In this case, we know that the Galois group is A_5 , so it is order 60, but if we didn't know the Galois group, we could use this data to predict its order.

Exercise 2.1. Check the consistency of the numbers above with Chebotarev's density theorem.

The goal of today's lecture is to give the necessary background in algebraic number theory to continue. It is roughly based on a lecture by Dick Gross [Gro11].

2.1 Number fields

A number field is a finite extension of \mathbb{Q} . Given a number field K/\mathbb{Q} of degree n (in this lecture, our field extensions will always be degree n), there is an associated ring of integers $\mathbb{O} \subset K$ consisting of all elements of K which satisfy a monic polynomial with coefficients in \mathbb{Z} . The ring of integers \mathbb{O} is a free \mathbb{Z} -module of rank n, as well as a Dedekind domain (i.e. Noetherian, normal, Krull dimension 1).

Exercise 2.2. Show that the following field extensions have the following rings of integers:

1.
$$K = \mathbb{Q}(i), \mathbb{O} = \mathbb{Z}[i].$$

2.
$$K = \mathbb{Z}(\sqrt{2}), \mathbb{O} = \mathbb{Z}[\sqrt{2}].$$

3. $K = \mathbb{Q}(\sqrt{5}), \mathbb{O} = \mathbb{Z}[\phi], \text{ where } \phi = \frac{1+\sqrt{5}}{2}.$

Generally, for a complicated extension, it is not easy to find \mathbb{O} .

We can measure how complicated a number field is using something called the **discrim**inant. It is defined as follows. Let K/\mathbb{Q} be a number field. Given $x \in K$, we get a \mathbb{Q} -linear map $x \colon K \to K$. Using this we define

$$\operatorname{Tr}: K \to \mathbb{Q}$$
$$\operatorname{Nm}: K^{\times} \to \mathbb{Q}^{\times}$$

by $Tr(x) := Tr(x \cdot)$, $Nm(x) := det(x \cdot)$. This gives us a bilinear form called the **trace form**:

$$K \times K \to \mathbb{Q}$$
$$(x, y) := \operatorname{Tr}(xy).$$

The trace form is nondegenerate over \mathbb{Q} because $\operatorname{Tr}(1) = n$, hence $\operatorname{Tr}(xx^{-1}) = n \neq 0$. Since any element of \mathbb{O} satisfies a monic polynomial with integer coefficients, the trace form restricts to a map $\mathbb{O} \times \mathbb{O} \to \mathbb{Z}$. Choose a \mathbb{Z} -basis $\{\alpha_1, \ldots, \alpha_n\}$ for \mathbb{O} . Then the **discriminant** of the field K is

$$\operatorname{Disc}(K) := \det((\alpha_i, \alpha_j)).$$

This is a close relative of the discriminant of a polynomial.

Remark 2.3. We have no idea how many number fields there are, so it is useful to have a measurement of how complicated a number field is. This is one of the reasons the discriminant is so useful.

Example 2.4. Let $K = \mathbb{Q}(i)$. Then $\mathbb{O} = \mathbb{Z}[i]$ has basis $\{\alpha_1, \alpha_2\} = \{1, i\}$. We can compute

$$((\alpha_i, \alpha_j)) = \begin{pmatrix} 2 & 0 \\ 0 & -2 \end{pmatrix},$$

so $\operatorname{Disc}(K) = \det((\alpha_i, \alpha_j)) = -4.$

Exercise 2.5. 1. Let $\alpha \in \mathbb{Z}$ be square-free. Let $K = \mathbb{Q}(\sqrt{\alpha})$. Then

$$\mathbb{O} = \begin{cases} \mathbb{Z}[\sqrt{\alpha}] & \text{if } \alpha \neq 1 \mod 4, \\ \mathbb{Z}\left[\frac{1+\sqrt{\alpha}}{2}\right] & \text{if } \alpha = 1 \mod 4. \end{cases}$$

Hence, calculate

$$\operatorname{Disc}(K) = \begin{cases} 4\alpha & \text{if } \alpha \neq 1 \mod 4, \\ \alpha & \text{if } \alpha = 1 \mod 4. \end{cases}$$

2. Calculate the discriminant of $\mathbb{Q}(e^{2\pi i/3})$.

Let K/\mathbb{Q} be an étale \mathbb{Q} -algebra (i.e. a finite separable extension of \mathbb{Q}). Then $K \otimes_{\mathbb{Q}} \mathbb{R}$ is an étale \mathbb{R} -algebra, so $K \otimes_{\mathbb{Q}} \mathbb{R} \simeq \mathbb{R}^{r_1} \times \mathbb{C}^{r_2}$, where $n = r_1 + 2r_2$. The field K is **totally real** if $r_2 = 0$ (or, equivalently, if every embedding $K \hookrightarrow \mathbb{C}$ lands in \mathbb{R}). For example, this happens if K is the splitting field of a polynomial with real roots.

Exercise 2.6. Show that the signature of the trace form on K totally real is $(r_1 + r_2, r_2)$. In particular,


K is totally real $\iff r_2 = 0 \iff (\cdot, \cdot)$ is positive definite.

2.2 An analogy

Next we will explore a useful analogy which will be a theme of this course.

$$\begin{cases} \text{finite extensions} \\ K \text{ of } \mathbb{C}(x) = \text{Frac}\mathbb{C}[x] \\ (\text{called function fields}) \end{cases} \leftarrow \left\{ \begin{array}{c} \text{smooth complex} \\ \text{projective curves} \\ C \text{ over } \mathbb{P}^1\mathbb{C} \end{array} \right\} \leftrightarrow \left\{ \begin{array}{c} \text{compact Riemann} \\ \text{surfaces with a} \\ \text{map to } \mathbb{P}^1\mathbb{C} \end{array} \right\}$$

The first arrow going left is given by taking the function field of the curve. We can also go in the other direction. Given



we obtain a map $\operatorname{Spec} \mathbb{O} \to \operatorname{Spec} \mathbb{C}[x] = \mathbb{A}^1$, so we have a unique compactification and an equivalence between the first two sets.

Similarly, there is a bijection

$$\left\{\begin{array}{c} \text{finite extensions} \\ K \text{ of } \mathbb{F}_p(x) \end{array}\right\} \leftrightarrow \left\{\begin{array}{c} \text{smooth projective} \\ \text{curves } C \text{ over } \mathbb{P}^1_{\mathbb{F}_p} \end{array}\right\}$$

Classically, people worked on problems in number theory in the algebraic world. Artin moved to the geometric world and proved deep results there. Many people in modern number theory work on the geometric side and hope to prove something about number fields. Here is a (very rough) schematic of difficulty:

function fields over $\mathbb{C} \ll$ function fields over $\mathbb{F}_p \ll$ number fields

For a very inspiring reference for all of this, see André Weil's letter to his sister Simone Weil on the role of analogy in mathematics [Kri05].

Exercise 2.7. (Use of Google allowed.) Show that Fermat's last theorem is true in function fields; i.e. if $f, g, h \in k[x]$ are relatively prime and $f^n + g^n = h^n$, then n = 2.

2.3 The fundamental exact sequence

Let K/\mathbb{Q} be a number field and \mathbb{O} the ring of integers of K. A **fractional ideal** is a finitely generated \mathbb{O} -submodule of K. Given two fractional ideals I, J, we can construct their product:

$$IJ := \left\{ \sum \alpha_i \beta_j | \alpha_i \in I, \beta_j \in J \right\}.$$

(This is the "union" in the sense of algebraic geometry.) Since \mathbb{O} is a Dedekind domain,

- every prime ideal $\mathfrak{p} \neq 0$ is maximal, and
- every fractional ideal has a unique factorization $I = \prod \mathfrak{p}_i^{e_i}$, where \mathfrak{p}_i are prime ideals.

Denote by $\mathcal{J} = \bigoplus_{\mathfrak{p}\neq 0 \text{ prime}} \mathbb{Z}\mathfrak{p}$ the group of nonzero fractional ideals under this product. We have the following fundamental exact sequence:

$$\{1\} \to \mathbb{O}^{\times} \hookrightarrow K^{\times} \to \mathcal{J} \to \mathcal{C}\ell(K) \to 0.$$

Here $\mathcal{C}\ell(K)$ is the **ideal class group** of K, which measures the failure of \mathbb{O} to be a PID. The ideal class group is difficult to calculate, and we know very little about it in general. The image of the second map in this exact sequence is \mathcal{P} , the set of all principal ideals (that is, ideals of the form $x\mathbb{O}$ for some $x \in K^{\times}$) of K.

Theorem 2.8. (Fundamental finiteness theorems)

- 1. The ideal class group $\mathcal{C}\ell(K)$ is finite.
- 2. The group \mathbb{O}^{\times} is finitely generated of rank $r_1 + r_2 1$.

Exercise 2.9. Compute \mathbb{O}^{\times} for $\mathbb{Q}(\sqrt{2})$ and $\mathbb{Q}(\sqrt{3})$. (*Hint*: Pell's equation)

We can study an analogue of this exact sequence for a smooth projective curve. Let C be a compact Riemann surface. Then under the analogy,

$$0 \neq \mathfrak{p}$$
 prime ideals = maximal ideals \leftrightarrow points of C ,

and we have the following exact sequence:

$$\{1\} \to \mathbb{C}^{\times} \to K^{\times} \to \mathcal{P} \hookrightarrow \bigoplus_{x \in C} \mathbb{Z}x \twoheadrightarrow \operatorname{Pic}(C) \to 0.$$

Here K is the function field of C, \mathcal{P} is the set of divisors of meromorphic functions ("principal divisors"), and $\operatorname{Pic}(C)$ is the Picard group of C (isomorphism classes of line bundles on C).


Remark 2.10. The group $Pic(C) = Jac(C) \times \mathbb{Z}$ is very far from finite. Also, \mathbb{C}^{\times} is not finitely generated. So in this setting, neither of the fundamental finiteness theorems hold.

Exercise 2.11. (For those who know some algebraic geometry) Show that the analogues of $\mathcal{C}\ell(K)$ and \mathbb{O}^{\times} are finite if C is an affine curve defined over a finite field.

Exercise 2.12. (If you know what K_0 is) Show that $K_0(\mathbb{O}) = \mathbb{Z} \oplus \mathcal{C}\ell(K)$.

2.4 Ramification

First consider a smooth projective curve $C \xrightarrow{f} \mathbb{P}^1 \mathbb{C}$ (e.g. $y^2 = f(x)$ for some polynomial f(x) without repeated roots.) After deleting a finite set of points from $\mathbb{P}^1 \mathbb{C}$ (the "discriminant" of f), then f is étale, and hence gives us a finite covering of open sets $U \subset \mathbb{P}^1 \mathbb{C}$. So all fibres are "the same" away from finitely many points where f is "ramified". A picture:




Over finite fields, almost the same thing happens. Again, if we have a smooth projective curve $C \xrightarrow{f} \mathbb{P}^1_{\mathbb{F}_n}$, f is étale after deleting finitely many points. For example, consider C =


Spec $\mathbb{F}_p[x, y]/(y^2 = x^3 + 1) \to \text{Spec } \mathbb{F}_p[x]$. Fibres of this map over a point $x = \lambda$ are singular if λ is a third root of unity, or else

$$\mathbb{F}_p[x,y]/(y^2 = x^3 + 1) \otimes_{\mathbb{F}_p[x]} \mathbb{F}_p = \mathbb{F}_p[y]/(y^2 = \lambda^3 + 1) = \begin{cases} \mathbb{F}_p \times \mathbb{F}_p & \text{if } \lambda^3 + 1 \text{ is a square} \\ \mathbb{F}_{p^2} & \text{if } \lambda^3 + 1 \text{ is not a square} \end{cases}$$

A picture:



A similar picture holds for number fields:



We can make this picture rigorous. Let K/\mathbb{Q} be a number field and \mathbb{O} its ring of integers. For each prime $p \in \mathbb{Z}$, the corresponding ideal in \mathbb{O} decomposes into the product of prime ideals in \mathbb{O} :

$$(p) = \prod_{i=1}^{g_p} \mathfrak{p}_i^{e_i}.$$

We say the primes \mathfrak{p}_i appearing in this decomposition are the "primes above (p)". The number e_i is the **ramification index** of \mathfrak{p}_i . For each \mathfrak{p}_i , the field $\mathbb{O}/\mathfrak{p}_i$ is a finite extension of \mathbb{F}_p (so $\mathbb{O}/\mathfrak{p}_i = \mathbb{F}_{p^{f_i}}$ for some f_i), and the degree f_i of this extension is the **inertia degree**.

Exercise 2.13. (Important!) Show that
$$n = \sum_{i=1}^{g_p} e_i f_i$$
.

Definition 2.14. (a) The ideal (p) is **unramified** if all $e_i = 1$. Otherwise it is **ramified**.

- (b) The ideal (p) splits completely if $f_i = e_i = 1$ for all i.
- (c) The ideal (p) is **inert** if $g_p = 1$ and $e_1 = 1$.

Theorem 2.15. The ideal (p) is unramified in \mathbb{O} if and only if $p \nmid \text{Disc}(K)$.

Exercise 2.16. Prove Theorem 2.15. (*Hint*: Show that a finite-dimensional commutative \mathbb{F}_{p} -algebra is étale if and only if its trace form is nondegenerate.)

Example 2.17. Let $K = \mathbb{Q}(i)$, so $\mathbb{O} = \mathbb{Z}[i]$ and Disc(K) = -4. (See Exercise 2.4.) Since $(1+i)^2 = 1+2i-1=2i$, we see that the ideal $(2) \subset \mathbb{O}$ decomposes as $(2) = (1+i)^2$, and is therefore ramified. By Theorem 2.15, all other primes are unramified. Let $p \neq 2$ be prime. Then to determine if (p) splits, we notice that

$$\mathbb{Z}[i]/(p) = \mathbb{Z}[x]/(x^2+1,p) = \mathbb{F}_p[x]/(x^2+1) = \begin{cases} \mathbb{F}_p \times \mathbb{F}_p & \text{if } \left(\frac{-1}{p}\right) = 1, \\ \mathbb{F}_{p^2} & \text{if } \left(\frac{-1}{p}\right) = -1 \end{cases}$$

Therefore (p) splits completely if and only if $p = 1 \mod 4$. (For example, (5) = (2+i)(2-i)), and (p) is inert if and only if $p = 3 \mod 4$.

We can adapt the strategy of Example 2.17 to determine splitting behavior in general. Let K/\mathbb{Q} be a number field, and choose a primitive element (i.e. a generating element) θ of K. We can assume without loss of generality that $\theta \in \mathbb{O}$. Then θ satisfies a monic polynomial $f(x) \in \mathbb{Z}[x]$ of degree n, so $\mathbb{Z}[\theta] \subset \mathbb{O}$ is of finite index m. Then for p such the $p \nmid m$ and $p \nmid \text{Disc}(K)$,

$$\mathbb{O}/(p) = \mathbb{Z}[\theta]/(p) = \mathbb{Z}[x]/(p, f(x)) = \mathbb{F}_p[x]/(f(x)) = \mathbb{F}_{p^{f_1}} \times \dots \times \mathbb{F}_{p^{f_k}}$$

where f_i 's are the degrees of irreducible factors of f(x) modulo p. So in particular,

(p) splits completely
$$\iff f$$
 is reducible modulo p ,
(p) is inert $\iff f$ is irreducible modulo p .

Hence we could have (and probably should have) phrased last week's lecture in terms of splitting of primes in a ring of integers.

2.5 The case of Galois extensions

Assume K/\mathbb{Q} is a Galois extension, with Galois group $\operatorname{Gal}(K/\mathbb{Q}) = G$. Then G acts on \mathbb{O} , and Frobenius proved the following result.

Theorem 2.18. If $(p) = \prod \mathfrak{p}_i^{e_i}$, then G acts transitively on the primes \mathfrak{p}_i .

Therefore, all e_i (resp. f_i) are equal. Denote their common value e (resp. f). Exercise 2.13 $(n = \sum e_i f_i)$ implies that $n = efg_p$. Fix a prime $\mathfrak{p}_i =: \mathfrak{p}$ lying over (p). Denote by $G_{\mathfrak{p}}$ the stabilizer of \mathfrak{p} in G (the "decomposition group"). Then

$$|G/G_{\mathfrak{p}}| = \#$$
 primes over $\mathbf{p} = g_p$,

so $|G_{\mathfrak{p}}| = ef$. Let $I_{\mathfrak{p}}$ be the subgroup of $G_{\mathfrak{p}}$ which acts trivially on $\mathbb{O}_{\mathfrak{p}}$ (the "inertia group"). This group has order e. We have the following exact sequence

$$1 \to I_{\mathfrak{p}} \to G_{\mathfrak{p}} \twoheadrightarrow \operatorname{Aut}(\mathbb{O}/\mathfrak{p}) \simeq \mathbb{Z}/f\mathbb{Z}.$$

Since $\mathbb{O}/\mathfrak{p} \simeq \mathbb{F}_{p^f}$, the group $\operatorname{Aut}(\mathbb{O}/\mathfrak{p})$ is generated by $\overline{\operatorname{Frob}}_p : x \mapsto x^p$. If p is unramified (i.e. $p \nmid \operatorname{Disc}(K)$), then e = 1, so $I_\mathfrak{p}$ is trivial and $G_\mathfrak{p} \simeq \mathbb{Z}/f\mathbb{Z}$. In this case, $\operatorname{Frob}_p \in G_\mathfrak{p}$ is defined to be the element that maps to $\overline{\operatorname{Frob}}_p \in \operatorname{Aut}(\mathbb{O}/\mathfrak{p})$. A different choice of \mathfrak{p} lying over p results in conjugate $G_\mathfrak{p}$ and conjugate Frob_p . This explains rigorously the Frob_p from the last lecture.

Lecture 3: L-functions

Last class we reviewed some algebraic number theory. This class we will review some analytic number theory. The motivation for this lecture is the following. We start with an arithmetic problem (for example, counting the number of $x \in \mathbb{F}_p$ such that $x^5 + 20x + 16 = 0$), and assign to it an *L*-function (something like a "character" for the representation theorists in the audience), which can be studied analytically.

3.1 The Riemann ζ -function

Define

$$\zeta(s) = \sum_{n \ge 1} n^{-s},$$

where $s \in \mathbb{C}$ is a complex variable. We can compare this sum to the integral

$$\int_{1}^{\infty} x^{-s} dx,$$

which converges to $\frac{1}{s-1}$ for real s > 1. When viewed as a holomorphic function, the integral converges absolutely for $s \in \mathbb{C}$ such that $\Re(s) > 1$. Hence the sum $\zeta(s)$ converges for all $s \in \mathbb{C}$ such that $\Re(s) > 1$.

This function has a rich history. Euler computed special values (e.g. $\zeta(2) = \frac{\pi^2}{2}$), and noticed that the ζ -function may also be given as the Euler product:

$$\sum_{n\geq 1} n^{-s} = (1+2^{-s}+(2^2)^{-s}+\cdots)(1+3^{-s}+(3^2)^{-s}+\cdots)\cdots = \prod_{p \text{ prime}} \left(\frac{1}{1-p^s}\right).$$

This product relates an analytic object, $\zeta(s)$, to the prime numbers. This relationship lets us study properties of primes using analysis! For example, the Euler product immediately gives us two proofs of the infinitude of primes: (1) the divergence of $\sum_{n\geq 1} \frac{1}{n}$ implies the product on the right must be infinite, and (2) since $\zeta(2) = \frac{\pi^2}{2}$, the irrationality of π^2 also implies that the right must be infinite.

Riemann showed that ζ admits a meromorphic continuation to all of \mathbb{C} . This is the **Riemann zeta function**. He also showed that $\zeta(s)$ has a simple pole at s = 1 (Exercise: Show that $\zeta(s) - \frac{1}{1-s}$ converges for $\Re(s) > 0$), and "trivial zeros" at $-2, -4, \ldots$ Furthermore, he established the **functional equation**: $\Lambda(s) := \pi^{-s/2} \Gamma(\frac{s}{2}) \zeta(s)$ satisfies

$$\Lambda(s) = \Lambda(1-s)$$

Here $\Gamma(s) = \int_{o}^{\infty} x^{s-1} e^{-x} dx$ is the gamma function. And finally, he proposed the following conjecture, which eventually became a millenium question.

Riemann hypothesis: If z is a non-trivial zero of $\zeta(z)$, then $\Re(z) = \frac{1}{2}$.

The Riemann hypothesis is known to be true for the first 10^{12} zeros of $\zeta(s)$.

3.2 Why do we care?

Here's the slogan of this story: "The zeros of the Riemann zeta function are the Fourier modes of the primes". We will spend the rest of the lecture trying to make this precise.

One of Riemann's motivations was the following theorem, which was a conjecture during his lifetime.

Theorem 3.1. (Prime Number Theorem) Let $\pi(x)$ be the number of primes less than or equal to $x \in \mathbb{R}$. Then

$$\pi(x) \sim \frac{x}{\log x}.$$

Here ~ means that $\lim_{x\to\infty} \frac{\pi(x)}{x/\log x} = 1$. Riemann was interested in two questions about the Prime Number Theorem:

- Why does it hold?
- What is the error term?

Instead of considering $\pi(x)$ directly, we can examine the **von Mangoldt function**, which "makes noise at prime powers":

$$\Lambda_{vm}(n) = \begin{cases} \log p & \text{if } n = p^m, \\ 0 & \text{otherwise.} \end{cases}$$

Define

$$\psi(n) = \sum_{m \le n} \Lambda_{vm}(m).$$

Exercise 3.2. Show that the Prime Number Theorem is equivalent to $\psi(n) \sim n$.

Riemann discovered an explicit formula for $\psi(x)$ at non-integers.

Theorem 3.3. (Riemann) For $x \in \mathbb{R} - \mathbb{Z}$, there is equality

$$\psi(x) = x - \sum_{\rho} \frac{x^{\rho}}{\rho} - \log(2\pi),$$

where the sum is taken over all zeros ρ of the Riemann ζ -function.

In other words,

$$x - \psi(x) = \log(2\pi) + \sum_{\rho} \frac{x^{\rho}}{\rho},$$

so the zeros of the Riemann ζ -function measure the error term in the prime number theorem. We can examine what effect the different types of zeros have on the right hand side of the equality above.

• Trivial zeros: The function $\frac{x^{-2n}}{-2n} = -\frac{1}{2nx^n}$ decays very quickly, so for large x, trivial zeros have almost no effect on the formula.

• A pair ρ , $\bar{\rho}$ of non-trivial conjugate zeros: Each such pair contribues

$$\lambda \cdot x^{\Re(\rho)} \cdot \cos(\gamma + \log x),$$

where λ and γ depend in a simple way on ρ and $\bar{\rho}$. So $\Re(\rho)$ is crucial in the contribution of ρ and $\bar{\rho}$ to the error term, and if the Riemann hypothesis is true, the growth of this contribution looks roughly like the product of $\cos(x)$ and $x^{1/2}$. Also, as $\Im(\rho)$ gets bigger, λ gets smaller. Thus, if the Riemann hypothesis is true, small zeros will contribute larger variations. A counterexample to the Riemann hypothesis would cause unexpectedly large fluctuations in $x - \psi(x)$.

- **Exercise 3.4.** (a) Show that the Prime Number Theorem is equivalent to $\zeta(s)$ having no zeros z with $\Re(z) = 1$.
 - (b) Show that the Riemann hypothesis is equivalent to $x \psi(x) \in O(x^{1/2})$.
 - (c) Find the error term in $\pi(x) \frac{x}{\log x}$ assuming the Riemann hypothesis.

Remark 3.5. It is unknown whether there exist non-trivial zeros ζ of $\zeta(s)$ with $\Re(z) = 1 - \epsilon$ for any $\epsilon > 0$.

3.3 Dirichlet *L*-functions

It is natural to ask questions about primes satisfying certain properties. For example, fix $m \in \mathbb{Z}_{\geq 0}$, and $a \in (\mathbb{Z}/m\mathbb{Z})^{\times}$. Consider the set

$$\{p \text{ prime } | p = a \mod m\}.$$

Is this set infinite? Is there an analogue to the Prime Number Theorem in this setting? A naive attempt to show that this set is infinite would be to consider the product

$$\prod_{p} \frac{1}{1 - p^{-s}}$$

taken over all primes p such that $p = a \mod m$ and recreate one of the arguments for the infinitude of primes given in the previous section. However, there is no Euler product in this setting, so this approach fails. Another approach is representation theory.

What do we learn from representation theory?

Consider the set of all functions from

$$(\mathbb{Z}/m\mathbb{Z})^{\times} \to \mathbb{C}$$

This set has two natural bases:

- 1. Indicator functions: $x \mapsto \delta_{x,a}$ for $a \in (\mathbb{Z}/m\mathbb{Z})^{\times}$.
- 2. Irreducible characters: $x \mapsto e^{2\pi i j/\phi(m)}$ for $j = 0, \dots, \phi(m) 1$, where $\phi(m) = |(\mathbb{Z}/m\mathbb{Z})|$.

In many ways, the basis of characters is more natural. Dirichlet borrowed this idea of characters to adapt our naive attempt above into something that works.

Definition 3.6. A Dirichlet character modulo **m** is a character $\chi : (\mathbb{Z}/m\mathbb{Z})^{\times} \to \mathbb{C}^{\times}$ extended by zero to all of \mathbb{Z} . In other words, it is a function $\chi : \mathbb{Z} \to \mathbb{C}^{\times}$ such that $\chi(n) = 0$ if $(n,m) > 1, \chi(n)$ depends only on $n \mod m$, and χ induces a character $\chi : (\mathbb{Z}/m\mathbb{Z})^{\times} \to \mathbb{C}^{\times}$.

Such a function χ has the property that $\chi(nn') = \chi(n)\chi(n')$. Using these characters, Dirichlet defined a **Dirichlet** *L*-function:

$$L(\chi, s) = \sum_{n \ge 1} \chi(n) n^{-s} = \prod_{p \text{ prime}} \frac{1}{1 - \chi(p) p^{-s}}.$$

With this adjustment by a character, the sum *does* admit an Euler product, as well as a meromorphic extension to all of \mathbb{C} , and a (rather complicated) functional equation. If χ is trivial, we recover the Riemann ζ -function. If χ is not trivial, then $L(\chi, s)$ is entire.

Dirichlet's theorem (that $\{p | p = a \mod m\}$ is infinite, with distribution $\frac{1}{\phi(m)} \frac{\pi}{\log x}$) is an easy consequence of the fact that $L(\chi, 1) \neq 0$ if χ is not trivial. Furthermore, there is an analogue of the Riemann hypothesis in this setting, usually called the "generalized Riemann hypothesis," and all non-trivial zeros of $L(\chi, s)$ lie on the critical line (i.e. satisfy $\Re(z) = \frac{1}{2}$).

3.4 Dedekind ζ -functions

Another natural question of this flavor is how primes behave in $\mathbb{Z}[i]$, or other rings of integers. To answer this question, Dedekind introduced his version of a ζ -function.

Return to the setting of last week: Let K/\mathbb{Q} be a number field with ring of integers \mathbb{O} . For a nontrivial fractional ideal $I \subset \mathbb{O}$, let $\mathbb{N}I = \#\mathbb{O}/I$. (For example, if $K = \mathbb{Q}$, $\mathbb{N}(p) = p$.) The **Dedekind** ζ -function is

$$\zeta_K(s) = \sum_{0 \neq I \subset \mathbb{O}} (\mathbb{N}I)^{-s} = \prod_{\mathfrak{p} \subset \mathbb{O} \text{ prime}} \frac{1}{1 - \mathbb{N}\mathfrak{p}^{-s}}.$$

This sum admits an Euler product because of the uniqueness of factorization of ideals in \mathbb{O} . Again, we have a meromorphic continuation and functional equation in this setting. (Historical note: The functional equation first appeared in Hecke's thesis in the 1920's, but the proof was very complicated in Hecke's work. A much simpler proof was given by Tate in his thesis in 1950.) The key example is the following.

Example 3.7. Let $K = \mathbb{Q}(i) \supset \mathbb{O} = \mathbb{Z}[i]$. Then

$$\begin{split} \zeta_{K}(s) &= \prod_{\mathfrak{p}\subset\mathbb{O} \text{ prime}} \frac{1}{1-\mathbb{N}\mathfrak{p}^{-s}} \\ &= \left(\frac{1}{1-2^{-s}}\right) \prod_{\mathfrak{p} \text{ s.t. } (p) \text{ splits}} \left(\frac{1}{1-\mathbb{N}\mathfrak{p}^{-s}}\right) \left(\frac{1}{1-\mathbb{N}\mathfrak{p}^{-s}}\right) \prod_{\mathfrak{p} \text{ s.t. } (p) \text{ is inert}} \left(\frac{1}{1-\mathbb{N}\mathfrak{p}^{-2s}}\right) \\ &= \left(\frac{1}{1-2^{-s}}\right) \prod_{p=1 \mod 4} \left(\frac{1}{1-p^{-s}}\right)^{2} \prod_{p=3 \mod 4} \left(\frac{1}{1-p^{-s}}\right) \left(\frac{1}{1+p^{-s}}\right) \\ &= \prod_{p} \left(\frac{1}{1-p^{-s}}\right) \prod_{p\neq 2} \left(\frac{1}{1-\chi(p)p^{-s}}\right) \\ &= \zeta(s)L(\chi,s), \end{split}$$

where

$$\chi(p) = \begin{cases} 1 & \text{if } p = 1 \mod 4, \\ -1 & \text{if } p = 3 \mod 4 \end{cases}$$

is a Dirichlet character on $\mathbb{Z}/4\mathbb{Z}$. The moral of this example is that we can understand $\mathbb{Z}(i)$ in terms of \mathbb{Z} ! We are witnessing the beginnings of class field theory.

3.5 Walking across the bridge

Next we will cross our bridge of analogy and see what happens in the geometric world. Recall that the objects analogous to a ring of integers \mathbb{O} contained in a number field K are smooth projective curves over \mathbb{F}_p and their function fields over \mathbb{F}_p .

Example 3.8. The simplest example of such a smooth projective curve is $\mathbb{P}^1_{\mathbb{F}_p}$. Instead we will work with $\mathbb{A}^1\mathbb{F}_p$, where the analogue of a ring of integers is $\mathcal{O}(\mathbb{A}^1\mathbb{F}_p) = \mathbb{F}_p[x]$. Here,

$$\begin{aligned} \zeta_{\mathbb{A}^{1}\mathbb{F}_{p}}(s) &= \sum_{\substack{o \neq I \subset \mathbb{F}_{p}[x]}} (\mathbb{N}I)^{-s} \\ &= \sum_{\substack{f \text{ monic}}} (\mathbb{N}(f))^{-s} \\ &= \sum_{\substack{d \geq 1}} \left(\sum_{\substack{f \text{ monic degree } d}} (p^{d})^{-s} \right) \\ &= \sum_{\substack{d \geq 0}} p^{d} (p^{d})^{-s} \\ &= \sum_{\substack{d \geq 0}} (p^{-s+1})^{d} \\ &= \frac{1}{1-p^{-s+1}}. \end{aligned}$$

So $\zeta_{\mathbb{A}^1_{\mathbb{F}_p}}(s)$ is a rational function with a unique pole at s = 1 and *no zeros*. Since $\zeta_{\mathbb{A}^1_{\mathbb{F}_p}}(s)$ measures the error term of the prime number theorem, this means that we can count primes exactly in this setting! In fact, we have (Gauss)

irred. polys of degree
$$d = \frac{1}{d} \sum_{m|d} \mu\left(\frac{d}{m}\right) p^m$$
.

Here μ is the Mobius function (i.e. $\mu(n)$ is the sum of the primitive n^{th} roots of unity). So in this setting, we know exactly how many primes there are in a given interval.

Our example was a little too simple, so we should bump it up a notch. In Artin's thesis (1923), he instead considered $X = \mathbb{F}_p[x, y]/(y^2 - f(x))$ for f(x) square free. (Under our analogy, this is the analogue of a quadratic field $\mathbb{Q}(\sqrt{a})$ for a square free.) Artin showed that

$$\zeta_X(s) = \frac{(1 - \alpha p^s)(1 - \bar{\alpha} p^s)}{1 - p^{-s+1}}$$

is again a rational function, with $\alpha \in \mathbb{C}$ of norm $p^{1/2}$. Hence all zeros of $\zeta_X(s)$ have $\Re z = \frac{1}{2}$, and the Riemann hypothesis is true in this setting. However, unlike the zeros of $\zeta(s)$, the zeros of $\zeta_X(s)$ are distributed evenly along the critical line.

The analogue to this statement for all curves was proven by Weil, and the case of arbitrary varieties was completed by Deligne (~ 1970). These accomplishments were some of the crowning glories of 20^{th} century mathematics.

3.6 Artin *L*-functions

Now we enter the non-abelian world. The year is 1927, about 31 years after Frobenius started developing the theory of group characters. Let K/\mathbb{Q} be a Galois extension, with $G = \text{Gal}(K/\mathbb{Q})$, and $d_K = \text{Disc}(K)$. Recall from previous lectures that:

- p is unramified if and only if $p \nmid d_K$, for unramified p,
- a choice of prime \mathfrak{p} over p results in an element $\operatorname{Frob}_{\mathfrak{p}} \in G$, and
- different choices of **p** lead to conjugate Frob_p.

Fix a finite dimensional complex representation

$$\rho: G \to \mathrm{GL}(V)$$

of the Galois group. Notice that the conjugacy class of $\rho(\text{Frob}_p)$ is completely determined by its characteristic polynomial det $(1 - t \operatorname{Frob}_p|_V)$. The (first approximation) of an **Artin** *L*-function is

$$L_{ur}(V,s) = \prod_{p \nmid d_K} \frac{1}{\det(1 - p^{-s} \operatorname{Frob}_p|_V)}.$$

Example 3.9. (a) V is the trivial representation. Then

$$L(V,s) = \prod_{p \nmid d_K} \frac{1}{1 - p^{-s}} = \zeta(s)$$
 up to finitely many factors.

So the Artin L-function recovers the Riemann ζ -function as a special case.

(b) Let $K = \mathbb{Z}(\sqrt{\alpha})/\mathbb{Q}$ be a quadratic field, $G = \{\pm 1\}$, and $\rho : G \to \mathrm{GL}_1(C)$ be the identity map $\{\pm 1\} \mapsto \{\pm 1\}$. Then for $p \nmid d_K$,

$$\rho(\mathrm{Frob}_{\mathrm{p}}) = \begin{cases} 1 & \text{if } p \text{ splits } \left(\iff \left(\frac{p}{\alpha} \right) = 1 \right), \\ -1 & \text{if } p \text{ is inert } \left(\iff \left(\frac{p}{\alpha} \right) = -1 \right). \end{cases}$$

Therefore, by quadratic reciprocity,

$$L_{ur}(\rho, s) = \prod_{p} \left(1 - \left(\frac{p}{\alpha}\right) p^{-s} \right)^{-1}$$
$$= \prod_{p} (1 - \chi(p)p^{-s})^{-1}$$
$$= L(\chi, s) \text{ (up to finitely many factors)}$$

for some Dirichlet character $\chi : \mathbb{Z}/4\alpha\mathbb{Z} \to \mathbb{C}^{\times}$. So Artin L-functions also recover Dirichlet *L*-functions.

Exercise 3.10. (a) Show that $L_{ur}(V_1 \oplus V_2, s) = L_{ur}(V_1, s)L_{ur}(V_2, s)$.

- (b) Show that L(V, s) converges absolutely for $\Re s > 1$. (*Hint*: Compare it to a product of dim V copies of the ζ -function.)
- (c) (Beautiful! Do it!) For a Galois extension K/\mathbb{Q} , let V_{reg} be the regular representation of G. Show that $L_{ur}(V_{reg}, s) = \zeta_K(s)$ up to finitely many factors.

A consequence of the exercise is the **Artin decomposition**:

$$\zeta_K(s) = \prod_{V \text{ irrep of } G} L_{ur}(V, s)^{\dim V} \text{ (up to finitely many factors)}.$$

Hence $\zeta_{\mathbb{Q}}(s)$ always divides $\zeta_K(s)$! This is a generalization of the factorization $\zeta_{\mathbb{Q}(i)}(s) = \zeta(s)L(\chi, s)$ that we saw in Example 3.7.

The Moral: All of the difficulty in K is contained in the difficulty of \mathbb{Q} if we allow for "non-abelian" difficulty (i.e. general representations of the Galois group).

Remark 3.11. We can get rid of the "up to finitely many factors" caveat in all of these statements by modifying $L_{ur}(V, s)$ with "general local factors" at ramified primes. See Geordie's hand-written lecture notes or Gus's lecture for a description of this procedure.

Lecture 4: The Sato-Tate conjecture

This will be our final lecture of global motivation before we zoom in on the local story. The goal of today's lecture is to describe the **Sato-Tate conjecture** and its relationship to the global Langlands picture. We will see that seemingly innocuous statements in the Langlands correspondence can have very powerful repercussions.

4.1 Equidistribution in representation theory

We say that the real numbers $\alpha_1, \alpha_2, \ldots \in [0, 1]$ are **equidistributed** if

$$\lim_{n \to \infty} \frac{1}{n} \# \{ \alpha_i \mid \alpha_i \in (a, b) \text{ for } i = 1, \dots, n \} = b - a = \int_0^1 \mathbb{1}_{(a, b)} dx$$

for any interval $(a, b) \subset [0, 1]$. Here $1_{(a,b)}$ is the indicator function on (a, b). Because indicator functions are dense in complex-valued Riemann integrable functions on [0, 1], this condition is equivalent to saying that the discrete average of any function on this set and continuous average of the same function agree; that is,

$$\lim_{n \to \infty} \frac{1}{n} \sum_{i=1}^{n} f(\alpha_i) = \int_0^1 f(x) dx$$

for all Riemann integrable $f : [0, 1] \to \mathbb{C}$. Now we can approximate any Riemann integrable $f : [0, 1] \to \mathbb{C}$ with a Fourier series, so this is in turn equivalent to

$$\lim_{n \to \infty} \frac{1}{n} \sum_{i=1}^{n} e(m\alpha_i) = \int_0^1 e(mx) dx = \begin{cases} 1 & \text{if } m = 0, \\ 0 & \text{otherwise} \end{cases}$$

for all $m \in \mathbb{Z}$. Here $e(x) = \exp(2\pi i x)$. The condition above always holds for m = 0, so to check if $\alpha_1, \alpha_2, \ldots$ are equidistributed, it suffices to check that for $m \neq 0$,

$$\lim_{n \to \infty} \sum_{i=1}^{n} e(m\alpha_i) = 0.$$

Here is an application of this observation. Let $\xi \in \mathbb{R}$ be irrational. Consider $(\xi), (2\xi), (3\xi), \ldots$ where $(m\xi) := m\xi \mod 1$. Are these numbers equidistributed? Weyl used the observation above to show that they are. Choose $m \neq 0$, and let $\eta = m\xi$. Then

$$\left|\frac{1}{n}\sum_{j=1}^{n}e(mj\xi)\right| = \left|\frac{1}{n}(e(\eta) + e(2\eta) + \dots + e(n\eta))\right| = \left|\frac{1}{n}\frac{e((n+1)\eta) - e(\eta)}{e(\eta) - 1}\right| \le \frac{1}{n}\left|\frac{2}{e(\eta) - 1}\right| \to 0$$

as $n \to \infty$.

Remark 4.1. This leads to many questions of a similar flavor (e.g. what about $(\xi), (4\xi), (9\xi), \ldots$?) Weyl's paper [Wey16] gives an affirmative answer for polynomials $f(x) \in n\mathbb{R}[n]$ whose coefficients are not all rational! We can reinterpret equidistribution via representation theory. The above reasoning shows (after identifying integers mod 1 with S^1) that a sequence $z_1, z_2, \ldots \in S^1 \subset \mathbb{C}$ equidistributes if for every nontrivial rational character χ of S^1 (e.g. $\chi : z \mapsto z^m$ for $m \neq 0$),

$$\lim_{n \to \infty} \frac{1}{n} \sum_{i=1}^{n} \chi(z_i) = 0.$$

If this condition holds, we say that the sum is "little o of n," and write

$$\sum_{i=1}^{n} \chi(z_i) = o(n).$$

Now suppose G is a compact group with Haar measure μ , and let X denote the space of conjugacy classes of G. (Recall that the **Haar measure** is the unique left (and then necessarily also right) G-invariant measure on G with $\mu(G) = 1$.) Let C(X) be the Banach space of continuous complex-valued functions on X. Two properties of irreducible characters on compact groups are the following:

Theorem 4.2. (The Peter-Weyl Theorem) The irreducible characters span a dense subspace of C(X).

Theorem 4.3. (Orthogonality of characters) If χ, χ' are irreducible characters, then

$$\int_{G} \chi(g) \overline{\chi'(g)} d\mu = \begin{cases} 1 & \text{if } \chi = \chi', \\ 0 & \text{otherwise.} \end{cases}$$

Hence, we have the following theorem about equidistribution of sequences in G.

Theorem 4.4. A sequence $\alpha_1, \alpha_2, \ldots \in X$ is equidistributed with respect to the (push forward of the) Haar measure if and only if

$$\sum_{i=1}^{n} \chi(\alpha_i) = o(n)$$

for all irreducible nontrivial characters χ .

Example 4.5. Let $G = S_3$. The conjugacy classes are determined by cycle type: $C_1 = \{id\}, C_2 = \{(12), (23), (13)\}, C_3 = \{(123), (132)\}$. The character table of S_3 is the following.

Haar measure	$\#C_i$	conj. class	triv	nat	sgn
1/6	1	C_1	1	2	1
1/2	3	C_2	1	0	-1
1/3	2	C_3	1	-1	1

We can see from this table that for a sequence to be equidistributed, it should spend twice the time in C_3 as it does in C_1 (as dictated by the column corresponding to the natural representation), and should spend as much time in C_2 as in $C_1 \cup C_3$ (as dictated by the column corresponding to the sign representation). **Example 4.6.** Let G = SU(2). Since all matrices in SU(2) are diagonalizable, conjugacy classes are determined by eigenvalues, which come in conjugate pairs. So $X = \{(\gamma, \bar{\gamma}) | \gamma \in S_1\}$ can be identified with $S^1_+ = \{z \in S^1 | \Re(z) \ge 0\} \simeq [0, \pi]$. By the Weyl character formula for SU(2), irreducible characters are of the form

$$\chi_m(z) = z^m + z^{m-2} + \dots + z^{-m},$$

where $z = e^{i\theta} \in S^1$. (Here we are using the identification $X \simeq S^1_+$ when defining these characters.) The Haar measure on the space of conjugacy classes (after identifying S^1 with $[0, 2\pi]$ via the exponential function) is then

$$\frac{2}{\pi}\sin^2\theta d\theta.$$

Exercise 4.7. Check that this is correct by showing that

$$\frac{2}{\pi} \int_0^\pi \sin^2 \theta d\theta = 1$$


and

$$\int_0^\pi \chi_m(e^{i\theta}) \sin^2\theta d\theta = 0$$

for $m \neq 0$.

The figure below is a plot of $\frac{2}{\pi}\sin^2\theta$. From this we see the distribution of eigenvalues of matrices in SU(2). A first observation is that there are many matrices in SU(2) with eigenvalues $\{i, -i\}$ (corresponding to $\theta = \frac{\pi}{2}$), and very few matrices with eigenvalues $\{1, 1\}$ and $\{-1, -1\}$ (corresponding to $\theta = 0$ and $\theta = \pi$, respectively). In fact, there is exactly one matrix in each case: I and -I.

Remark 4.8. This demonstrates that it is more likely for matrices in SU(2) to have eigenvalues which are "far away" (meaning that the angle between them in the complex plane is large), an important fact in random matrix theory.



Hence if you had a sequence of pairs of complex numbers which you suspect are eigenvalues of random matrices in SU(2), you could tell pretty quickly whether or not it was plausible.

4.2 Elliptic curves and the Sato-Tate conjecture

Next we discuss elliptic curves. (This seems to be completely unrelated, but we should have faith that it will come full circle.) Let k be a field, and E an elliptic curve. (That is, a smooth projective curve over k of genus 1 with a fixed rational point $0 \in E(k)$.) Assume the characteristic of k is not 2 or 3. Then E can be made to be of the form (the projective closure of)


 $y^{2} = x^{3} + ax + b$, with 0 = (0:0:1) its point at infinity.

Here smoothness translates into the fact that $x^3 + ax + b$ has no repeated roots, i.e. that $4a^3 - 27b^2 \neq 0$.

First assume we are working over \mathbb{C} . Then E is a compact Riemann surface of genus g = 1, so $E = \mathbb{C}/\Lambda$ for some lattice $\Lambda \simeq \mathbb{Z}^2 \hookrightarrow \mathbb{C}$. Hence

$$\left\{\begin{array}{c} \text{elliptic curves} \\ \text{over } \mathbb{C} \end{array}\right\}_{/ \text{ iso}} \simeq \left\{\begin{array}{c} \text{lattices} \\ \Lambda \subset \mathbb{C} \end{array}\right\}_{/ \text{ iso}} \xrightarrow{\text{"period ratio"}} {}_{\mathrm{SL}_2(\mathbb{Z})} \backslash \mathbb{H},$$

where \mathbb{H} is the upper half-plane. The quotient $_{SL_2(\mathbb{Z})} \setminus \mathbb{H}$ can be identified (up to some ambiguity on the boundary) with its fundamental domain



so we can consider all complex elliptic curves as points in the region above.


Certain elliptic curves have extra structure called complex multiplication. Let E be a complex elliptic curve. As a real Lie group, E is isomorphic to $S^1 \times S^1$, hence

$$\operatorname{End}_{\operatorname{Lie\,gp}}(E) = \mathbb{Z}^2$$
$$(z, w) \mapsto (z^m, w^n) \leftrightarrow (m, n)$$

But *E* has additional structure - it is an elliptic curve $(E \simeq \mathbb{C}/\Lambda)$, and a complex algebraic group. By a miracle of abelian varieties, the elliptic curve endomorphisms of *E* fixing 0 are the same as the complex algebraic group endomorphisms of *E*. Hence,

$$\operatorname{End}_{0\mapsto 0}(E) = \operatorname{End}_{\operatorname{alg gp}}(E) = \{z \mid z\Lambda \subset \Lambda\} = \begin{cases} \mathbb{Z} & \text{if } \Lambda \cdot \Lambda \not\subset \Lambda, \\ \Lambda & \text{if } \Lambda \cdot \Lambda \subset \Lambda. \end{cases}$$

In the second case, we say E has **complex multiplication**. In the fundamental domain, the elliptic curves with complex multiplication are of the form $i\sqrt{d}$ (the "corner point" in the picture above):



Thus one can think of curves with complex multiplication as being very special.

Exercise 4.9. Show that if $\Lambda \cdot \Lambda \subset \Lambda$ (i.e. *E* has complex multiplication), $\Lambda \otimes \mathbb{Q}$ is an imaginary quadratic field. In particular, if *E* has complex multiplication then Λ is what is called an *order* in an imaginary quadratic field.

Next we work over \mathbb{Q} , and consider the elliptic curve E given by $y^2 = x^3 + ax + b$ for $a, b \in \mathbb{Z}$. To understand E, we reduce modulo p and study the number of points of $E(\mathbb{F}_p)$. If $E_{\mathbb{F}_p} = E \times \operatorname{Spec} \mathbb{F}_p$ is nonsingular, then p is a prime of **good reduction**. (This is the analogue for algebraic varieties of a prime being unramified.) We can bound $\#E(\mathbb{F}_p)$ by the **Hasse-Weil bound**:

Theorem 4.10. (Hasse-Weil)

$$#E(\mathbb{F}_p) = 1 + p - \alpha_p,$$

where $|\alpha_p| \leq 2\sqrt{p}$.

The Hasse-Weil bound tells us that 1 + p is a "square root accurate" approximation for $\#E(\mathbb{F}_p)$. By our discussions last week, hopefully you are convinced that this is an analogue of the Riemann hypothesis for elliptic curves.

Big question: How do the α_p 's behave?

We can start to answer this question using the **Grothendieck–Lefshetz trace formula.** Let $H^*(E)$ be the étale cohomology of E. For all p outside a finite set, we have an action of Frob_p on (étale, but don't worry if you don't know what this is) cohomology:

dim : 1 2 1

$$H^{0}(E)$$
 $H^{1}(E)$ $H^{2}(E)$
 \circlearrowright \circlearrowright \circlearrowright
Frob_p = 1 Frob_p $\sim \begin{pmatrix} \gamma_{p} & 0\\ 0 & \overline{\gamma_{p}} \end{pmatrix}$ Frob_p = p

The Grothendieck-Lefshetz trace formula is


$$#E(\mathbb{F}_p) = \sum (-1)^i \operatorname{Tr}(\operatorname{Frob}_p : H^i) = 1 + p - (\gamma_p + \overline{\gamma_p}),$$

where $\gamma_p, \overline{\gamma_p}$ are the eigenvalues of Frobenius on $H^1(E)$. Hence to determine the number of solutions of $E(\mathbb{F}_p)$, it is enough to examine γ_p , and it is true (but not so easy to see) that the Riemann hypothesis for E is equivalent to $|\gamma_p| = \frac{1}{2}$.

We can renormalize so that

$$\theta_p := \frac{1}{\sqrt{p}} \gamma_p \in S^1_+.$$

This leaves us with a sequence $\theta_2, \theta_3, \theta_5, \ldots$ of points on a semicircle S^1_+ controlling the number of points of E modulo p.




Sato–Tate conjecture (1960): Suppose *E* does not have complex multiplication. If we identify $S^1_+ \xrightarrow{\simeq} [0, \pi]$, then


$$\lim_{n \to \infty} \frac{1}{\pi(n)} \sum_{p \le n} \mu_{\theta_p} = \frac{2}{\pi} \sin^2 \theta d\theta$$

Here μ_{θ_p} is the Dirac distribution.

In other words, the Sato-Tate conjecture is that the θ_p 's look like eigenvalues of random matrices in SU(2)! Below is a very beautiful illustration of this phenomenon. For the elliptic curve $y^2 + y = x^3 + x + 3x + 5$ (which has no complex multiplication), the following is the plot of θ_p for the first 5,000 primes. (The further out the dot, the bigger the prime.)



In contrast to this, consider the elliptic curve $y^2 = x^3 + 1$. Here complex multiplication is given by the Eisenstein integers. The plot below shows θ_p for the first 5,000 primes. Norm is linear in the prime (so, again, the further the dot, the bigger the prime).



The difference between these two curves is striking!

Remark 4.11. The case of complex multiplication is well understood. The idea (Geordie thinks) is that the extra endomorphisms force the Frob_p to lie in a subgroup of SU(2). (Geordie points out that he is "the exact opposite of an expert on this topic...")

Remark 4.12. One can think of Sato-Tate (roughly) as a higher-dimensional analogue of Chebotarev density: $\operatorname{Gal}(K/\mathbb{Q}) \leftrightarrow SU(2)$.

4.3 Equidistribution and *L*-functions

In the final part of this lecture, we will describe how the Sato–Tate conjecture follows from a simple part of the Langlands correspondence (which is still conjectural). The Sato–Tate conjecture has been proven using other methods, but the proof is very involved and heavily influenced by ideas from the Langlands program. This example is meant to showcase the power of the Langlands correspondence.

Let G be a compact group, X its space of conjugacy classes, and $x_p \in X$ a family of elements parameterized by primes p (perhaps outside some finite set of "bad" primes). For an irreducible character χ , we can define an L-function analogously to how we defined Artin L-functions for Galois groups:

$$L(\chi, s) = \prod_{p} \frac{1}{\det(1 - \rho(x_p)p^{-s})},$$

where ρ is the representation afforded by χ . This converges for $\Re(s) > 1$. Assume additionally that $L(\chi, s)$ extends to a meromorphic function on $\Re(s) \ge 1$ having neither zeros nor poles along $\Re(s) = 1$ except possibly at s = 1. Let $-c_{\chi}$ be the order of $L(\chi, s)$ at s = 1 (so $c_{\chi} > 0$ pole, $c_{\chi} < 0$ zero). With these assumptions, we have the following theorem.

Theorem 4.13.

$$\sum_{p \le n} \chi(x_p) = c_{\chi} \cdot \frac{n}{\log(n)} + o\left(\frac{n}{\log(n)}\right).$$

The proof of this theorem involves some complex analysis and tricks with sums, but is not difficult. This theorem has an important corollary.

Corollary 4.14. If for all nontrivial χ , $L(\chi, s)$ is holomorphic and nonzero at s = 1, then the x_p are equidistributed in X.

So we can use *L*-functions to test whether a sequence in a compact group is equidistributed!

Exercise 4.15. (a) Show that $L(\chi, 1) \neq 0$ implies Dirichlet's theorem.

(b) Let K be a number field. It's known that $\zeta_K(s)/\zeta(s)$ is holomorphic and nonvanishing at s = 1. Using this, deduce Chebotarev's density theorem.


An important consequence of Corollary 4.14 is that it lets us reframe the Sato-Tate conjecture in terms of *L*-functions.

Example 4.16. (Serre) Assume that for all $m \ge 1$, the symmetric *L*-power function

$$L(S_{\chi}^{m},s) := \prod_{p} \frac{1}{(1 - \theta_{p}^{m}p^{-s})(1 - \theta_{p}^{m-2}p^{-s})\cdots(1 - \theta_{p}^{-m}p^{-s})}$$

satisfies the extra assumption above. (Here S_{χ}^{m} is the m^{th} symmetric power of the representation with character χ sending $\operatorname{Frob}_{p} \mapsto \begin{pmatrix} \theta_{p} & 0 \\ 0 & \theta_{p}^{-1} \end{pmatrix}$.) Then the Sato-Tate conjecture holds!

Recall our cartoon of the Langlands correspondence:



An extremely important part of this picture is **functoriality**. That is, the diagram should be compatible with

- 1. composition of representations of the Galois group with algebraic representations of GL_n (so $\rho : \operatorname{Gal}(K) \to \operatorname{GL}_n \xrightarrow{\operatorname{alg rep'n}} \operatorname{GL}_m$ should correspond to some operation on automorphic representations), and
- 2. changing the field K.

A key piece of the Langlands correspondence is that L-functions coming from automorphic representations have many desirable properties, which are extremely difficult to establish for L-functions coming from geometric representations of Gal(K). For example, once we know that an L-function comes from an automorphic representation, we immediately know that it admits a meromorphic continuation and has a functional equation.

The Punchline: In the simple example of the algebraic representation $GL_2 \rightarrow GL_m$ via symmetric powers, the prediction of functoriality implies the Sato-Tate conjecture!


Lecture 5: Infinite Galois theory and global class field theory

The topic of today's lecture is **class field theory**. But before diving in, we will start with a motivating question and a review of infinite Galois theory.

Basic Question: What are all finite extensions of a number field K (e.g. \mathbb{Q})?

This is certainly a question of fundamental importance in number theory. One could also ask more specific questions, such as "How many number fields have a given Galois group and discriminant?" For almost every question of this sort, we have no idea what the answer is. Class field theory develops techniques for answering these questions in the abelian setting. We will say more precisely what this means later in the lecture, but for now, let's take a look at an example.

Example 5.1. Let $K = \mathbb{F}_p$, and \overline{K} its algebraic closure. For all $n \ge 1$, there is a unique subfield $K_n \subset \overline{K}$ with p^n elements, and $\overline{K} = \bigcup_{n \ge 1} K_n$. We have the following picture of field extensions and corresponding Galois groups:



Let's calculate $\operatorname{Gal}(\overline{K}/K)$. Because $\overline{K} = \bigcup_{n>1} K_n$, we have an injection

$$\operatorname{Gal}(\overline{K}/K) \hookrightarrow \prod_{n \ge 1} \operatorname{Gal}(K_n/K) = \prod_{n \ge 1} \mathbb{Z}/n\mathbb{Z}$$

For $\varphi \in \operatorname{Gal}(\overline{K}/K)$, let the sequence (φ_n) be its image in $\prod_{n\geq 1} \operatorname{Gal}(K_n/K)$. A sequence $(\gamma_n) \in \prod_{n\geq 1} \operatorname{Gal}(K_n/K)$ is in the image if and only if $\gamma_n = \gamma_m \mod m$ whenever m|n. Hence,

$$\operatorname{Gal}(\overline{K}/K) = \lim_{\longleftarrow} \mathbb{Z}/n\mathbb{Z} = \widehat{\mathbb{Z}}$$

is the "profinite completion of \mathbb{Z} ".

Exercise 5.2. (a) Show that $\widehat{Z} = \prod_{p \text{ prime}} \mathbb{Z}_p$.

(b) Show that $\widehat{\mathbb{Z}}$ has uncountably many subgroups. Hence a naive Galois correspondence cannot hold.

5.1 Infinite Galois theory

Let L/K be a Galois extension (algebraic, normal, separable, not necessarily finite). Then we have an injection

$$\operatorname{Gal}(L/K) \hookrightarrow \prod_{K \subset L' \subset L} \operatorname{Gal}(L'/K),$$

where the product is taken over all towers of field extensions $K \subset L' \subset L$, where the extension L'/K is finite Galois. For all towers of extensions $K \subset L' \subset L'' \subset L$, where the extensions L'/K and L''/K are finite Galois, there is a corresponding map $\operatorname{Gal}(L''/K) \to \operatorname{Gal}(L'/K)$. This determines the image of this injection; that is,

$$\operatorname{Gal}(L/K) = \lim_{K \subset L' \subset L} \operatorname{Gal}(L'/K).$$

This is a topological group. Indeed, if we give $\prod_{K \subset L' \subset L} \operatorname{Gal}(L'/K)$ the product topology (which is compact, by Tychonov), then $\operatorname{Gal}(L/K)$ inherits the subspace topology.

Exercise 5.3. The group $\operatorname{Gal}(L/K)$ is closed (therefore compact) in $\prod_{K \subset L' \subset L} \operatorname{Gal}(L'/K)$.

Example 5.4. We see from Exercise 5.3 and Example 5.1 that the group $\widehat{\mathbb{Z}}$ is compact, which might look strange.

Exercise 5.5. (Important, can be used as a definition) A basis of open neighborhoods of $1 \in \text{Gal}(L/K)$ is given by kernels of the maps

$$\operatorname{Gal}(L/K) \to \operatorname{Gal}(L'/L)$$


for L'/K finite Galois.

For a general group G, define

$$\widehat{G} = \lim_{\substack{H \subseteq G \\ \text{normal} \\ \text{finite index}}} G/H.$$

The group G is **profinite** if $G \xrightarrow{\simeq} \widehat{G}$, otherwise, say \widehat{G} is the **profinite completion** of G. So Galois groups are profinite groups! The key example to keep in mind is the following.

Example 5.6. Consider the group $\mathbb{Z}_p = \lim_{\longleftarrow} \mathbb{Z}/p^n \mathbb{Z}$. What does this group look like? Here's a picture for p = 3:



Exercise 5.7. Here are some fun thought experiments.

- (a) Where is -1 in this picture?
- (b) Think about \mathbb{Q}_3 .

The motto: Galois groups are fractal-like objects!

Theorem 5.8. (Fundamental theorem of infinite Galois theory) Let L/K be a Galois extension. Then there exists a canonical bijection

$$\{ K \subset L' \subset L \} \leftrightarrow \left\{ \begin{array}{l} \text{closed subgroups} \\ \text{of } \operatorname{Gal}(L/K) \end{array} \right\}$$
$$L^H \leftrightarrow H \\ L' \mapsto \operatorname{Gal}(L/L')$$

Moreover, under this bijection,

finite extensions \leftrightarrow closed and open subgroups Galois extensions \leftrightarrow normal subgroups

Exercise 5.9. Show that the only closed subgroups of $\widehat{\mathbb{Z}}$ are $n\widehat{\mathbb{Z}}$ for $n \in \mathbb{Z}$. If $n \geq 1$, then the subgroup $n\widehat{\mathbb{Z}}$ corresponds to the extension K_n under the bijection above, and n = 0 corresponds to \overline{K} . (So $0\widehat{\mathbb{Z}}$ is the only closed subgroup which isn't open.)

Now we return to the problem posed at the start of this lecture:

describe all number fields over \mathbb{Q}

or, equivalently,

describe all closed subgroups of $\operatorname{Gal}(\overline{\mathbb{Q}}/\mathbb{Q})$.

However, after some thought, one sees that this isn't really a well-defined question (from a philosophical point of view), because $\overline{\mathbb{Q}}$ involves a *choice* (or many choices), so there is no concrete canonical realisation of $\overline{\mathbb{Q}}$. Hence $\operatorname{Gal}(\overline{\mathbb{Q}}/\mathbb{Q})$ is only really a "group up to conjugacy," in the sense that any meaningful statements one can make about this group must be invariant under conjugation. (One cannot talk about individual elements.)

The Punchline:

- 1. Isomorphism classes of representations of "a group up to conjugacy" are canonical, so it makes sense to talk about representations of $\operatorname{Gal}(\overline{\mathbb{Q}}/\mathbb{Q})$. This is one reason why representations are so important in the Langlands program!
- 2. The maximal abelian extension \mathbb{Q}^{ab} of \mathbb{Q} (which is the extension corresponding to $\overline{[\operatorname{Gal}(\overline{\mathbb{Q}}/\mathbb{Q}), \operatorname{Gal}(\overline{\mathbb{Q}}/\mathbb{Q})]}$) is canonical, and we can hope to describe it by studying the maximal abelian quotient $G^{ab} := G/\overline{[G,G]}$ of $G = \operatorname{Gal}(\overline{\mathbb{Q}}/\mathbb{Q})$. This is class field theory!

5.2 Global class field theory, first version

The key example to keep in mind is the maximal abelian extension of \mathbb{Q} .

Example 5.10. Let $K_m := \mathbb{Q}(\zeta_m)$ for $\zeta_m = e^{2\pi i/m}$. Define $\mathbb{Q}(\mu_\infty) := \bigcup_{m \ge 1} K_m$. The Galois group of K_m/\mathbb{Q} is $(\mathbb{Z}/m\mathbb{Z})^{\times}$, hence

$$\operatorname{Gal}(\mathbb{Q}(\mu_{\infty})/\mathbb{Q}) = \lim_{\longleftarrow} (\mathbb{Z}/m\mathbb{Z})^{\times} = \prod_{p} \mathbb{Z}_{p}^{\times}$$

Fact: ("Kronecker Jugendtraum") $\mathbb{Q}(\mu_{\infty})$ is the maximal abelian extension of \mathbb{Q} .

This fact is not easy! (It will follow from global class field theory.) The hope of Kronecker was to predict this starting just from \mathbb{Q} , without calculating extensions.

Exercise 5.11. Use Kronecker Jugentraum to show that any continuous character χ : $\operatorname{Gal}(\overline{\mathbb{Q}}/\mathbb{Q}) \to \mathbb{C}^{\times}$ "is" a Dirichlet character. (Part of the exercise is to work out what "is" means in this context.)

Given a number field K, it is useful to consider all norms at once. Let $\mathbb{O} \subset K$ be the ring of integers. A **place** v is an equivalence class of nontrivial multiplicative norms

$$|\cdot|_v: K \to \mathbb{R}_{\geq 0}$$

on K.

Theorem 5.12. All places of a number field K are of the following form.

- Finite places: $|x|_v := (\#\mathbb{O}/\mathfrak{p})^{-\operatorname{val}_\mathfrak{p}(x)}$ for $\mathfrak{p} \subset \mathbb{O}$ prime.
- Real places: $|x|_v := |i(x)|$ for some real embedding $i: K \hookrightarrow \mathbb{R}$,
- Complex places: $|x|_v := |i(x)|^2$ for some pair of conjugate $i : K \hookrightarrow \mathbb{C}$ not landing in \mathbb{R} .

These are all possible notions of distance on a number field.

Exercise 5.13. Show that there are no nontrivial norms on a finite field.

Note that we could have chosen any scalar $\lambda > 1$ in place of $\#\mathbb{O}/\mathfrak{p}$ above. The reason for the beautiful product formula: For $x \in K^{\times}$,

$$\prod_{\text{places } v} |x|_v = 1.$$

Note that this product makes sense because all but finitely many places are 1. The function field case of this formula is the statement that the number of poles and number of zeros (with multiplicity) agree.

5.3 Global class field theory á la Artin

Fix a finite abelian Galois extension L/K with abelian Galois group $\operatorname{Gal}(L/K)$. Let $\mathbb{O}_L \subset L$ and $\mathbb{O}_K \subset K$ be the rings of integers, and let $S_f \subset \mathbb{O}_K$ be the set of ramified primes. We have seen that for a prime $\mathfrak{p} \subset \mathbb{O}_K$ such that $\mathfrak{p} \notin S_f$, there is a corresponding conjugacy class $\operatorname{Frob}_p \in \operatorname{Gal}(L/K)$. In general Frob_p is only defined up to conjugacy, but since we are assuming that $\operatorname{Gal}(L/K)$ is abelian, Frob_p is a single element. Hence we obtain a map (the **Artin map**):

$$\mathcal{J}^{S_f} = \bigoplus_{\mathfrak{p} \notin S_f} \mathbb{Z}\mathfrak{p} \to \operatorname{Gal}(L/K)$$
$$\sum m_{\mathfrak{p}}\mathfrak{p} \mapsto \prod \operatorname{Frob}_{p}^{m_{\mathfrak{p}}}$$

Here $\mathcal{J}^{S_f} \subset \mathcal{J}$ is a subgroup of the group of nonzero fractional ideals $\mathcal{J} = \bigoplus_{\text{primes } \mathfrak{p}} \mathbb{Z}\mathfrak{p}$ discussed in Lecture 2. By Chebotarev's density theorem, the Artin map is surjective.


Question: What is the kernel of the Artin map?

The answer to this question is related to an observation we made in the very first lecture. Recall our motivating problem for the course of determining the number of solutions of a polynomial modulo p for various primes p. For quadratic polynomials, we used quadratic reciprocity to find some modulus $m \in \mathbb{Z}$ such that the number of solutions of the polynomial modulo p was given by the residue of p modulo m. At first, the modulus m seemed to be somewhat mysterious, but eventually we observed that it was obtained from the *ramified primes* (that is, the "weird primes" which we ignored). For example, for the polynomial $x^2 + 1$, which has 2 solutions modulo p if $p = 1 \mod 4$ and 0 solutions if $p = 3 \mod 4$, the modulus 4 is the square of 2, our only ramified prime.

Returning to the setting of the Artin map, we define a **modulus** m supported in a set of places S to be a formal \mathbb{Z} -linear combination of places $m = \sum m_i v_i$ such that $m_i \in \{0, 1\}$ for real places and $m_i = 0$ for all complex places and places $v_i \notin S$. Given a modulus msupported in a set of places S, we can define an associated group (the Ray class group) as follows. Consider the following two subsets of K^{\times} :

$$K^S = \{\lambda \in K^{\times} \mid \operatorname{val}_{\mathfrak{p}}(\lambda) = 0 \text{ for all } \mathfrak{p} \in S\}, \text{ and}$$

 $K^{m,1} = \{\lambda \in K^s \mid \operatorname{val}_{\mathfrak{p}}(\lambda - 1) \geq m_i \text{ for finite places, and } i(\lambda) \in \mathbb{R}_{>0}^{\times} \text{ for real places } m_i = 1\}.$ The set $K^{m,1}$ is the set of λ which are "*m* close to 1". We have the following maps:



The **Ray class group** associated to the modulus m is the quotient

$$\mathcal{C}\ell_K^m := \mathcal{J}^S/\mathrm{val}(K^{m,1})$$

Example 5.14. If m = 0, then $\mathcal{C}\ell_K^0 = \mathcal{C}\ell_K$ is the class group.

Example 5.15. Let $K = \mathbb{Q}$, and m = n(p) for a prime $p \in \mathbb{Z}$. Then *m* is a modulus supported on $S = \{(p)\}$. We have

$$K^{S} = \left\{ \frac{a}{b} \mid a, b \text{ are coprime to } p \right\} = \mathbb{Z}_{(p)}^{\times}, \text{ and}$$


$$K^{m,1} = \left\{\frac{a}{b} \mid a, b \text{ coprime to } p \text{ and } \operatorname{val}_p(\frac{a}{b}-1) \ge n\right\} = \left\{\frac{a}{b} \mid \frac{a}{b} = 1 \mod p^n\right\}.$$

Hence

$$K^s/K^{m,1} = \mathbb{Z}^{\times}_{(p)}/K^{m,1} = (\mathbb{Z}/p^n\mathbb{Z})^{\times} = \mathcal{C}\ell^m_{\mathbb{Q}}$$

Theorem 5.16. For any modulus m, the Ray class group is finite and surjects onto the class group of K.

Theorem 5.17. (Artin) For any L/K as above, there exists a modulus m with $\operatorname{supp}(m) \cap \{ \text{finite places} \} = S_f$ such that $\operatorname{val}(K^{m,1})$ is contained in the kernel of the Artin map. Moreover, for any modulus m and any quotient $q : \mathcal{C}\ell_K^m \to Q$, there exists an abelian extension L/K ramified only at primes in $\operatorname{supp}(m)$ such that



commutes.

A weaker form of this theorem provides a more direct answer to our question from the beginning of this section.

Theorem 5.18. For any abelian Galois extension L/K, there exists an $\epsilon > 0$ such that if $\lambda \in K^{S_f}$ is ϵ close to 1 for all places $v \in S$, then $val(\lambda)$ is in the kernel of the Artin map.

Example 5.19. Consider the extension $\mathbb{Q}(i)/Q$, which is the splitting field of the polynomial $x^2 + 1$. The ramified primes are 2 and ∞ (but we haven't discussed what it means for ∞ to be ramified, so we are sweeping this under the rug), so $S^f = \{(2)\}$. For any $p \neq 2$, $\operatorname{Frob}_p(i) = i^p$, hence in the Galois group, $\operatorname{Frob}_p \leftrightarrow p \mod 4 \in (\mathbb{Z}/4\mathbb{Z})^{\times} = \operatorname{Gal}(\mathbb{Q}(i)/\mathbb{Q})$. We have

$$\mathcal{J}^{S_f} = \left\{ \frac{a}{b} \mid a, b > 0 \text{ coprime to } 2 \right\} = \mathbb{Z}_{(2), > 0}^{\times}$$

Hence the Artin map is the natural map

$$\mathbb{Z}^{\times}_{(2),>0} \to (\mathbb{Z}/4\mathbb{Z})^{\times}$$

Its kernel is the set of all elements satisfying a "congruence condition at 2":

$$\left\{\lambda \in \mathbb{Z}_{(2),>0}^{\times} \mid \operatorname{val}_2(\lambda-1) \ge 2\right\}.$$

We see from this example that the finitely many "weird" (ramified) primes from the first lecture are the primes determining our congruences.

Example 5.20. If m = 0, then $\mathcal{C}\ell_K^0 = \mathcal{C}\ell_K$. Hence the existence statement in Theorem 5.17 implies that there exists an unramified everywhere extension L/K with $\operatorname{Gal}(L/K) = \mathcal{C}\ell_K$. This extension is the **Hilbert class field**.

Lecture 6: Local Galois groups and local class field theory

6.1 A potted history of class field theory

Sometimes if we find something difficult, it can be comforting to know that other people also found it difficult. Accordingly, we'll start today's lecture with a potted history of class field theory, following [Con01, Miy11].


- Kronecker, Weber (1850-1880): Kronecker's Jugentraum (that $\mathbb{Q}(\mu_{\infty})$ is the maximal abelian extension of \mathbb{Q}), explicit class field theory (describing K^{ab} explicitly, not just its Galois group) for \mathbb{Q} and $\mathbb{Q}(i)$, relevance of complex multiplication.
- Dedekind, Frobenius (1880): defined Frob_p (then everyone promptly forgot for 40 years).
- Hilbert (189-1900): first correct proof of Jugentraum for \mathbb{Q} , emphasis on "places at ∞ ," introduction of Hilbert class field, 12^{th} problem on Hilbert's list was explicit class field theory for any K. (Still open! Even for $\mathbb{Q}(\sqrt{d}), d \ge 0$!)
- Hensel (1897): introduction of *p*-adic numbers, took a while to catch on.
- Takagi (1900): PhD student of Hilbert in Göttingen, thesis on $\mathbb{Q}(i)^{ab}$, proof of existence theorem during WWI (when there was no contact between Germany and Japan), result announced at the ICM in 1920.
- Hasse (1922): local global principle, first time *p*-adics were taken seriously by the broader mathematics community.
- Chebotarev (1927): density theorem.
- E. Artin (1927): introduction of Artin map (the return of Frob_p!), reciprocity theorem.
- Schmidt (1930): deduced local class field theory from global class field theory (proofs still analytic).
- Noether (1930s): local theory should be simpler and come first.
- Chevalley (1940): algebraic proof of local class field theory.

6.2 A trip across the bridge


Recall that the "bridge" in this course is the motivating analogy between number fields and function fields.

Remark 6.1. This bridge was what motivated Hensel's advocation for the introduction of the *p*-adic numbers.


Let's look at local class field theory through this analogy. Let L/K be a finite Galois extension. Across the bridge, this should correspond to a surjective map (i.e. ramified cover) $f: X \to Y$ of algebraic curves/Riemann surfaces over \mathbb{C} . Recall that our \mathbb{R} -picture and \mathbb{C} -picture of such a map are the following:



For all $y \in Y$ outside of a finite set, f is étale; that is, a smooth n : 1 cover in a neighborhood of y:




For a finite set of points $y \in Y$, f is not smooth, and locally sends $z \mapsto z^{n_i}$, such that $\sum n_i = n$:



If f is Galois, then all n_i are equal.

The Moral: The ramified cover f is determined by simple data ("local monodromies") at finitely many points ("ramified primes of Y").

Remark 6.2. We have



The algebraic closure of $\mathbb{C}((t))$ is $\mathbb{C}((t^{\mathbb{Q}})) := \bigcup_{n \ge 1} \mathbb{C}((t^{1/n}))$. Hence,

$$\operatorname{Gal}\left(\overline{\mathbb{C}((t))}/\mathbb{C}((t))\right) = \lim_{\leftarrow} \mathbb{Z}/n\mathbb{Z} = \widehat{\mathbb{Z}}.$$

This is "why" the local field information is so simple in function field/ \mathbb{C} case¹. In language to come, every extension of $\mathbb{C}((t))$ is "tamely ramified".

The upshot is that on the function field side of the bridge, local information is easy, and can be patched together to form the global picture. On the number field side of the bridge, local information is much harder, but the philosophy we learn from our analogy is that it should still be easier than global information, so we should focus on it first. Because of this, essentially the rest of this course will be local.

Now we return to number fields. Let L/K be a finite Galois extension. Fix a place v of K. If v is finite (corresponding to some $\mathfrak{p} \subset \mathbb{O}_k$), then we know what it means for a place v' (corresponding to $\mathfrak{q} \subset \mathbb{O}_L$) of L to "lie over v": v' lies over v precisely when \mathfrak{q} is a prime above \mathfrak{p} in the sense of lecture 2 (i.e. $\mathfrak{p}\mathbb{O}_L \subset \mathfrak{q}$).

If v is a real or complex place corresponding to $i: K \hookrightarrow \mathbb{K} \in \{\mathbb{R}, \mathbb{C}\}$, then a place v' of L lies over v if the corresponding injection i' fits into a commutative diagram.



¹Recall that we saw last lecture that $\operatorname{Gal}(\overline{\mathbb{F}_p}/\mathbb{F}_p) = \widehat{\mathbb{Z}}$ as well. This turns out to be a useful coincidence, but we won't comment on it further here.

Hence if v is real, then v' is either real or complex. A real place v is **ramified** if there exists a v' lying above v that is complex in L, and **unramified** if all v' above v are real. If v is complex, then all v' above v are complex, and we say the place v is **unramified**.

Fix a place v of K and a place v' of L over v. Let $L_{v'}$ (resp. K_v) be the completion of L (resp. K) with respect to the place v' (resp. v). Then we have the diagram



Set

$$G_{v} = \{ \sigma \in \operatorname{Gal}(L/K) \mid \sigma \text{ acts continuously on } L_{v} \}$$
$$= \{ \sigma \mid \sigma \text{ preserves } v' \} = \begin{cases} G_{\mathfrak{q}} & \text{if } v \text{ is finite,} \\ \{1\} & \text{if } v \text{ is unramified infinite,} \\ \mathbb{Z}/2\mathbb{Z} & \text{if } v \text{ is ramified infinite.} \end{cases}$$

Here $G_{\mathfrak{q}} \subset \operatorname{Gal}(L/K)$ is the decomposition group corresponding to the prime $\mathfrak{q} \subset \mathbb{O}_L$ determining v'.

Remark 6.3. In the last case (when v is ramified infinite), we get a canonical element $c \in \text{Gal}(L/K)$ corresponding to complex conjugation.

The point: $L_{v'}/K_v$ is a finite Galois extension of local fields with Galois group G_v . We will first try to understand such extensions for all places v, then piece together this information to understand L/K.

6.3 Local class field theory

Between the "easy" world of finite fields and the complicated world of global fields lies the world of local fields. Let K be a field equipped with discrete valuation val : $K \to \mathbb{Z} \cup \{\infty\}$. In K lies its ring of integers \mathbb{O}_K , with maximal idea \mathfrak{m} generated by a "uniformizer" $\pi \in \mathfrak{m}$:

$$K \supset \mathbb{O}_K = \operatorname{val}^{-1}(\mathbb{Z}_{\geq 0} \cup \{\infty\}) \supset \mathfrak{m} = \operatorname{val}^{-1}(\mathbb{Z}_{>0} \cup \{\infty\}) = (\pi).$$

The field $k_K = \mathbb{O}_K/\mathfrak{m}$ is the residue field. Note that in this setup, K, \mathbb{O}_K and \mathfrak{m} are all canonical, but the uniformizer $\pi \in \mathfrak{m}$ is not. For us, a **local field** will be a field K equipped with a discrete valuation as above such that

- 1. K is complete with respect to val (i.e. K has the topology coming from $\mathbb{O}_K = \lim_{\leftarrow} \mathbb{O}_K / \mathfrak{m}_K^n$), and
- 2. k_K is finite.

Exercise 6.4. Show that 1. and 2. are equivalent to K being locally compact.

Example 6.5. The field \mathbb{Q}_p is locally compact because it is covered by dilates of \mathbb{Z}_p :

$$\mathbb{Q}_p = \bigcup_{n \ge 1} p^{-n} \mathbb{Z}_p$$

Recall that \mathbb{Z}_p are compact open sets in \mathbb{Q}_p .

Remark 6.6. In some terminology, \mathbb{R} and \mathbb{C} are also referred to as local fields.

Let L/K be a finite Galois extension of local fields. Then any element $\sigma \in \text{Gal}(L/K)$ preserves $\mathbb{O}_L/\mathbb{O}_K$ and $\mathfrak{m}_L/\mathfrak{m}_K$, and thus acts on k_L/k_K . Hence we get maps

$$1 \to I_{L/k} \hookrightarrow \operatorname{Gal}(L/K) \twoheadrightarrow \operatorname{Gal}(k_L/k_K) \to 1,$$

where $I_{L/K}$ is the inertia subgroup of lecture 2. The Galois group $\operatorname{Gal}(k_L/k_K) \simeq \mathbb{Z}/d\mathbb{Z}$ is generated by Frob_q . For $L = \overline{K}$, this short exact sequence becomes

$$1 \to I_{\overline{K}/K} \hookrightarrow \operatorname{Gal}(\overline{K}/K) \twoheadrightarrow \operatorname{Gal}(\overline{\mathbb{F}_q}/\mathbb{F}_q) \simeq \widehat{\mathbb{Z}} \to 1.$$

Local class field theory: There exists a canonical map² $r_K : K^{\times} \to \operatorname{Gal}(\overline{K}/K)^{ab}$ with dense image such that r_K induces an isomorphism $\widehat{K^{\times}} \xrightarrow{\sim} \operatorname{Gal}(\overline{K}/K)^{ab}$. Here $\widehat{K^{\times}}$ is the profinite completion of K^{\times} (that is, the completion with respect to subgroups of finite index). Moreover,

1. the diagram

commutes, and

2. if L/K is finite Galois, then

$$\begin{array}{ccc} L^{\times} & \stackrel{r_L}{\longrightarrow} & \operatorname{Gal}(\overline{L}/L)^{ab} \\ & & & \downarrow^{\operatorname{res}} \\ K^{\times} & \stackrel{r_K}{\longrightarrow} & \operatorname{Gal}(\overline{L}/K)^{ab} \end{array}$$

commutes.

Remark 6.7. The analogous statements for the fields \mathbb{R} and $\mathbb{C}((t))$ are the following:

²The canonical map r_K is sometimes called the "reciprocity map".

1. $K = \mathbb{R}$: In the diagram

$$\begin{array}{ccc} \mathbb{C}^{\times} & \xrightarrow{r_{\mathbb{C}}} & \operatorname{Gal}(\mathbb{C}/\mathbb{C}) = \{1\} \\ & & & \downarrow \\ & & & \downarrow \\ \mathbb{R}^{\times} & \xrightarrow{r_{\mathbb{R}}} & \operatorname{Gal}(\mathbb{C}/\mathbb{R}) \end{array}$$


the map $r_{\mathbb{R}} : -1 \mapsto$ complex conjugation is continuous and surjective. The kernel of $r_{\mathbb{R}}$ is $\mathbb{R}^{\times}_{>0}$, the set of norms coming from \mathbb{C}^{\times} (i.e. the image of Norm_{\mathbb{C}/\mathbb{R}}).

2. $K = \mathbb{C}((t))$:³ The map

$$r_{\mathbb{C}((t))}: \mathbb{C}((t))^{\times} \to \operatorname{Gal}\left(\overline{\mathbb{C}((t))}/\mathbb{C}((t))\right) = \widehat{\mathbb{Z}}$$

has dense image, so a reasonable choice is valuation val : $\mathbb{C}((t))^{\times} \to \mathbb{Z}$.

It is useful to modify the Galois group $\operatorname{Gal}(\overline{K}/K)$ slightly. Define the **Weil group** of K to be the subgroup $W_K \subset \operatorname{Gal}(\overline{K}/K)$ of elements whose projection onto $\widehat{\mathbb{Z}}$ is an integral power of Frob_q ; that is, W_K fits into the short exact sequence



The purpose for this modification of the following fact: the reciprocity map r_K provides an isomorphism between K^{\times} and the abelianization of the Weil group:

$$r_K: K^{\times} \xrightarrow{\simeq} W_K^{ab}.$$

With this, we can state the local Langlands correspondence for $GL_n(K)$.

Theorem 6.8. (Local Langlands correspondence for $GL_n(K)$) (Harris-Taylor) There is a bijection

$$\operatorname{Hom}_{cts}(W_K, \operatorname{GL}_n(\mathbb{C}))/_{\operatorname{conj}} \stackrel{1:1}{\longleftrightarrow} \left\{ \begin{array}{c} \operatorname{irreps} \text{ of } \operatorname{GL}_n(K) \\ \operatorname{in} \mathbb{C} \operatorname{-vector spaces} \end{array} \right\}$$

The continuous group homomorphisms on the left hand side of this bijection are referred to as the **Langlands parameters** of the corresponding $GL_n(K)$ -representations on the right.

Remark 6.9. Actually, this is not quite correct. Instead we should consider Weil-Deligne reps on the left, and smooth admissible reps on the right. These issues will be addressed in coming lectures.

³Note that K does not quite fit our assumptions so LCFT does not apply, but morally it fits into the same picture.

Example 6.10. The n = 1 case of this theorem is true by local class field theory:

$$\operatorname{Hom}_{cts}(W_K, \operatorname{GL}_1(\mathbb{C}))/_{\operatorname{conj}} = \operatorname{Hom}_{cts}(W_K, \mathbb{C}^{\times})$$
$$= \operatorname{Hom}_{cts}(W_K^{ab}, \mathbb{C}^{\times})$$
$$= \operatorname{Hom}_{cts}(K^{\times}, \mathbb{C}^{\times})$$
$$= \{\operatorname{irreps} \text{ of } \operatorname{GL}_1(K)\}.$$

Example 6.11. We can see explicitly that local class field theory is true for \mathbb{Q}_p . By a local version of the Jugentraum,

$$\mathbb{Q}_p^{ab} = \mathbb{Q}_p(\mu_\infty) = \bigcup \mathbb{Q}_p(\zeta_n),$$

where ζ_n is an n^{th} root of unity. Hence

$$\mathbb{Q}_p^{ab} = \mathbb{Q}_p(\mu_{p'}) \cdot \mathbb{Q}_p(\mu_{p^{\infty}}),$$

where $\mathbb{Q}_p(\mu_{p'}) := \bigcup_{p \nmid n} \mathbb{Q}_p(\zeta_n)$ and $\mathbb{Q}_p(\mu_{p^{\infty}}) := \bigcup_{n \geq 1} \mathbb{Q}_p(\zeta_{p^n})$. As before, $\operatorname{Gal}(\mathbb{Q}_p(\zeta_{p^n})/\mathbb{Q}_p) = (\mathbb{Z}/p^n\mathbb{Z})^{\times}$, so

 $\operatorname{Gal}(\mathbb{Q}_p(\mu_{p^{\infty}})/\mathbb{Q}_p) = \mathbb{Z}_p^{\times}.$

If $p \nmid n$, note that $\mathbb{Q}_p(\zeta_n)/\mathbb{Q}_p$ is unramified, so $\mathbb{Q}(\mu_{p'}) =: \mathbb{Q}_p^{ur}$ is the maximal unramified extension of \mathbb{Q}_p . Because $\overline{\mathbb{F}_p} = \bigcup_{p \nmid n} \mathbb{F}_p(\zeta_n)$, we have that

$$\operatorname{Gal}(\mathbb{Q}_p(\mu_{p'})/\mathbb{Q}_p) = \operatorname{Gal}(\overline{\mathbb{F}_p}/\mathbb{F}_p) = \widehat{\mathbb{Z}}.$$

We conclude that

$$\operatorname{Gal}(\mathbb{Q}_p^{ab}/\mathbb{Q}_p) = \widehat{\mathbb{Z}} \times \mathbb{Z}_p^{\times} \simeq \widehat{\mathbb{Q}}_p^{\times}$$

which is exactly what is predicted by local class field theory.

6.4 Structure of Galois groups of local fields

We will finish this lecture with some remarks on the structure of local Galois groups. This section is hands-on and explicit. Morally, it should come before local class field theory.

Let L/K be a finite Galois extension of local fields, and $L \supset \mathbb{O}_L \supset \mathfrak{m}_L = (\pi_L), K \supset \mathbb{O}_K \supset \mathfrak{m}_K = (\pi_K)$ the respective rings of integers, maximal ideals, and uniformizers. As before, there is a corresponding extension of residue fields k_L/k_K . We have a short exact sequence

$$1 \to I_{L/K} \to \operatorname{Gal}(L/K) \twoheadrightarrow \operatorname{Gal}(k_L/k_K) \simeq \mathbb{Z}/n\mathbb{Z} \to 1,$$

where $I_{L/K} = \{\sigma \mid \sigma \text{ acts trivially on } k_L\}$, and $\operatorname{Gal}(k_L/k_K)$ is generated by the canonical generator Frob. Note that $\operatorname{Gal}(L/K)$ preserves \mathbb{O}_K , hence \mathfrak{m}_K , hence the valuation v_K , hence acts on $\mathbb{O}_K/\mathfrak{m}_K^j$, hence acts continuously on L.

Lemma 6.12. (Key lemma) Any $\sigma \in I_{K/L}$ is determined by its action on π_L .

Exercise 6.13. Prove Lemma 6.12. (*Hint*: use the fact that any $\sigma \in \text{Gal}(L/K)$ is automatically continuous.)

Set $I := I_{L/K}$, $I_0 := I$, and $I_j := \{ \sigma \in I \mid \sigma(\pi)\pi^{-1} \in 1 + \mathfrak{m}_L^j \}$ for $j \ge 1$.

Proposition 6.14. This defines a filtration

$$I = I_0 \supset I_1 \supset I_2 \supset \cdots$$

of I by normal subgroups. Moreover,

- 1. This is a finite filtration; i.e. $I_m = \{1\}$ for large enough m.
- 2. We have natural injections

$$\begin{split} &I_0/I_1 \xrightarrow{\sigma(\pi)\pi^{-1}} k_L^{\times}, \\ &I_j/I_{j+1} \xrightarrow{} (1+\mathfrak{m}_L^j)/(1+\mathfrak{m}_L^{j+1}) \simeq k_L \end{split}$$

for $j \ge 1$. In particular, I is solvable, I_0/I_1 is of order prime to p, and I_1 is the Sylow p subgroup of I.

This filtration is called the **ramification filtration of** *I*. The proof is an easy exercise.

Definition 6.15. We say that L/K is tamely ramified if $I_1 = \{1\}$, and L/K is unramified⁴ if $I_0 = \{1\}$.


Remark 6.16. This agrees with our earlier notion of unramified from Lecture 2.

 $^{^{4}}$ Note that a quirk of this terminology is that unramified is tamely ramified. It is strange, but we will just have to get used to it.

Lecture 7: Representation theory of *p*-adic groups

7.1 Tying up some loose ends

We start today's lecture by tying up some loose ends from previous weeks. Recall that two lectures ago, we discussed how looking for a description of $\operatorname{Gal}(\overline{\mathbb{Q}}/\mathbb{Q})$ is misguided because $\operatorname{Gal}(\overline{\mathbb{Q}}/\mathbb{Q})$ is only a "group up to conjugacy," since its definition requires a choice of $\overline{\mathbb{Q}}$. There is an analogy for this idea that Geordie learned from Kevin Buzzard which might be more familiar to us. Let X be a path-connected space. A choice of base point $x \in X$ yields the fundamental group $\pi_1(X, x)$. Another choice of base point $y \in X$ yields the isomorphic group $\pi_1(X, y)$.



An isomorphism $\pi_1(X, x) \simeq \pi(X, y)$ requires a choice of path from x to y in X. Such a choice of path is *not canonical*. Grothendieck taught us an analogue of this for extensions of fields.

 $\mathbb{Q} \leftrightarrow \quad \text{``étale site''} \operatorname{Spec} \mathbb{Q} \text{ (something like a space)}$ choice of $\overline{\mathbb{Q}} \leftrightarrow \text{choice of ``base point'' of Spec} \mathbb{Q}$

Then the étale fundamental group $\pi_1^{\text{ét}}(\operatorname{Spec} \mathbb{Q}, \mathbb{Q}) = \operatorname{Gal}(\overline{\mathbb{Q}}/\mathbb{Q}).$

The punchline: The fundamental group $\pi_1(X, x)$ depends on base point x, but $\operatorname{Rep} \pi_1(X, x) \simeq \{ \text{local systems on } X \}$ is canonical! Transporting this statement via Grothendieck's analogy we see that, although the absolute Galois group is not defined canonically, its category of (continuous) representations is. It is this category that the Langlands correspondence tries to understand.

Last lecture we stated the local Langlands correspondence for GL_n : Fix a local field K (i.e. K is a finite extension of \mathbb{Q}_p or $K \simeq \mathbb{F}_q((t))$). There is a canonical bijection

$$\left\{\begin{array}{c} \operatorname{cts reps of } W_K \\ \operatorname{in } \operatorname{GL}_n(\mathbb{C}) \end{array}\right\}_{/\mathrm{iso}} \stackrel{1:1}{\longleftrightarrow} \left\{\begin{array}{c} \operatorname{irred \ blah} \\ \operatorname{reps \ of \ } \operatorname{GL}_n(K) \end{array}\right\}_{/\mathrm{iso}}$$

Here W_K is the **Weil group** of the field K. Last week we showed why this follows from local class field theory for n = 1. However, this statement is not quite precise. On the left hand side we need to consider **Weil-Deligne representations** of W_K , and on the right hand side we need to establish exactly what conditions are captured by "blah". We'll keep stating versions of this theorem every lecture until we converge on something correct.

7.2 The no small subgroups argument

Our final piece of housekeeping is the **no small subgroups argument**. This is a very useful fact that has not fit in naturally to our story so far, so we will slot it in here.

Definition 7.1. A topological group G has **no small subgroups** if there exists a neighborhood U of the identity in G such that any subgroup contained in U is trivial.

Example 7.2. Here are some examples of groups with no small subgroups.

- 1. The circle group S^1 .
- 2. Discrete groups (e.g. finite groups). We can take $U = \{id\}$.
- 3. The real numbers \mathbb{R} and the complex numbers \mathbb{C} . (Powers of any non-identity element move far away from the identity.)
- 4. Any Lie group G. (Use that exp : Lie $G \to G$ is a local diffeomorphism and $\exp(mg) = \exp(g)^m$.)
- 5. Any topological subgroup of a group with no small subgroups has no small subgroups (e.g. $\mathbb{Q}/\mathbb{Z} \hookrightarrow S^1$ has no small subgroups).

Remark 7.3. In contrast, profinite groups have "many small subgroups," because a basis of neighbourhoods of the identity consists of subgroups of finite index.

Lemma 7.4. Let Γ be a profinite group and G a topological group with no small subgroups. Then any continuous group homomorphism $\varphi : \Gamma \to G$ has finite image.

Proof. Let $U \subset G$ be an open neighborhood of the identity containing no nontrivial subgroups, and let $\varphi : \Gamma \to G$ be a continuous group homomorphism. Then $\varphi^{-1}(U) \subset \Gamma$ is open. Since Γ is profinite, there exists a normal subgroup $N \subset \varphi^{-1}(U)$ such that G/N is finite. The image $\varphi(N) \subset U$ is a subgroup, so $\varphi(N) = \{1\}$ since G has no small subgroups. Hence Γ factors through a finite group, $\varphi : \Gamma \to \Gamma/N \to G$.

The Moral:

$$\left\{ \begin{array}{c} \text{Fractal-like objects} \\ \text{(p-adic groups, Galois groups)} \end{array} \right\} \cap \left\{ \begin{array}{c} \text{Euclidean-type objects} \\ \text{(Lie groups)} \end{array} \right\} = \{\text{finite groups}\}$$

A consequence of this is that we can not draw any good pictures of \mathbb{Z}_p in \mathbb{C} (or for that matter in any Lie group) which respect the addition or multiplication structure.

This moral gives us a new perspective of the local Langlands correspondence. The "no small subgroups" lemma implies that the left hand side of the LLC consists (roughly) of a collection of finite subgroups of $\operatorname{GL}_n(\mathbb{C})$, along with surjections from a Galois-group-type object to the subgroups. So very roughly, the LLC provides a classification of irreducible admissible representations of a Lie group over a local field by certain finite subgroups of $\operatorname{GL}_n(\mathbb{C})$.

7.3 You could have guessed the LLC for GL_2 !

The goal is this section is to give a heuristic explanation for the LLC. *Warning*: this is not precise! Everything we say here will have to be tweaked later. Geordie learned this perspecive from a series of lectures by Dipendra Prasad in Russia.

Starting place: Say we wanted to guess the representation theory of $GL_n(\mathbb{Q}_p)$. What would we do?

Step 1: We might start by figuring out the representation theory of finite reductive groups. For example, let $G = SL_2(\mathbb{F}_q)$. There are two maximal tori in G, up to conjugacy:

$$T_s = \left\{ \begin{pmatrix} a & 0 \\ 0 & a^{-1} \end{pmatrix} \mid a \in \mathbb{F}_q^{\times} \right\} \simeq \mathbb{Z}/(q-1)\mathbb{Z}, \text{ the "split torus," and}$$

$$T_a = \{\lambda \in \mathbb{F}_{q^2}^{\times} \subset \operatorname{GL}_2(\mathbb{F}_q) \mid \operatorname{Norm}(\lambda) = 1\} \simeq \mathbb{Z}/(q+1)\mathbb{Z}, \text{ the "anisotropic torus"}.$$

In the definition of T_a above, we are using the fact that $\operatorname{GL}_2(\mathbb{F}_q)$ is the group of invertible linear transformations of the \mathbb{F}_q -vector space \mathbb{F}_q^2 , so

$$\mathbb{F}_{q^2}^{\times} \subset \mathrm{GL}_{\mathbb{F}_q}(\mathbb{F}_q^2) = \mathrm{GL}_2(\mathbb{F}_q).$$

Roughly,

$$\left\{\begin{array}{c} \text{irred reps of} \\ \mathrm{SL}_2(\mathbb{F}_q) \text{ over } \mathbb{C}\end{array}\right\} \stackrel{1:1}{\longleftrightarrow} \left\{\chi: T_s \to \mathbb{C}^{\times}\right\}_{/\chi \sim \chi^{-1}} \bigsqcup_{q \to \infty} \{\theta: T_a \to \mathbb{C}^{\times}\}_{/\theta \sim \theta^{-1}} \\ \text{``principal series''} \quad \text{``discrete series''} \right\}$$

Let us check that we are not too far off by doing a count. Irreducible representations of a finite group are in bijection with conjugacy classes, and conjucagy classes are roughly in bijection with characteristic polynomials, so the sizes of the sets above are roughly

For more details and a careful construction of the irreducible representation of $SL_2(\mathbb{F}_q)$, see the notes from Joe Baine's talks on the Informal Friday Seminar webpage. The upshot is that we obtain almost all irreducible representations of $SL_2(\mathbb{F}_q)$ through some "induction" from characters of the two conjugacy classes of tori. (Note that the details are much more complicated as there is no actual induction functor. In the setting of finite reductive groups we use Deligne-Lusztig induction.)

Step 2: Once we have a good idea of the representation theory of finite reductive groups, a next natural step might be to understand the representation theory of real reductive groups. For example, let $G = SL_2(\mathbb{R})$. Again, there are two conjugacy classes of maximal tori: the "split torus"

$$T_s = \left\{ \begin{pmatrix} a & 0\\ 0 & a^{-1} \end{pmatrix} \mid a \in \mathbb{R}^{\times} \right\} \simeq \mathbb{R}^{\times},$$

and the "anisotropic torus"

$$T_a = \left\{ \begin{pmatrix} \cos\theta & \sin\theta \\ -\sin\theta & \cos\theta \end{pmatrix} \right\} \simeq SO_2.$$

Something similar happens in this setting to what we saw with the finite reductive groups. Roughly,

$$\left\{ \begin{array}{c} \text{irred admissible} \\ \text{reps of } SL_2(\mathbb{R}) \end{array} \right\} \stackrel{1:1}{\longleftrightarrow} \left\{ \begin{array}{c} \text{cts characters} \\ \text{of } T_s \simeq \mathbb{R} \times \mathbb{Z}/2\mathbb{Z} \\ \text{"principal series"} \end{array} \right\} \bigsqcup \left\{ \begin{array}{c} \text{cts characters} \\ \text{of } T_a \simeq SO_2 \simeq S^1 \\ \text{"discrete series"} \end{array} \right\}$$

So again we see that, roughly, irreducible representations are all obtained by "inducing" characters of conjugacy classes of tori.

Step 3: We dream that something similar might be true for representations of *p*-adic groups. Let $G = GL_2(K)$ for a local field K. By descent, we have the following relationship:

$$\begin{cases} \text{ conjugacy classes of} \\ \max'l \text{ tori in } \operatorname{GL}_2(K) \end{cases} \leftrightarrow \begin{cases} \text{ semisimple } K\text{-algebras} \\ L \text{ s.t. } \dim_K L = 2 \end{cases} \end{cases}$$
$$L^{\times} \leftrightarrow L$$

There are two cases:

- 1. Split torus: $L \simeq K \times K$, so maximal torus is of the form $L^{\times} \simeq K^{\times} \times K^{\times}$.
- 2. Anisotropic torus: L/K degree 2 extension, so maximal torus is of the form L^{\times} .

Applying our analogy from earlier, we might expect

$$\left\{\begin{array}{c} \text{irred blah} \\ \text{reps of } \operatorname{GL}_2(K) \end{array}\right\} \xleftarrow{\operatorname{roughly 1:1}} \left\{\begin{array}{c} \text{pairs of characters} \\ \chi_1, \chi_2 : K^{\times} \to \mathbb{C}^{\times} \end{array}\right\} \bigsqcup \left\{\begin{array}{c} \text{characters } \theta : L^{\times} \to \mathbb{C}^{\times} \\ \text{where } L/K \text{ is degree } 2 \end{array}\right\}.$$

Now, local class field theory tells us that $W_K^{ab} \simeq K^{\times}$ and $W_L^{ab} \simeq L^{\times}$. Moreover, $W_L \subset W_K$ is an index 2 subgroup, so

$$\left\{\begin{array}{c} \text{irred blah}\\ \text{reps of } \operatorname{GL}_2(K) \end{array}\right\} \xleftarrow{\operatorname{roughly 1:1}} \left\{\chi_1 \otimes \chi_2 : W_K \to \operatorname{GL}_2(\mathbb{C})\right\} \sqcup \left\{\operatorname{Ind}_{W_L}^{W_K}(\theta) : W_L \to \operatorname{GL}_2(\mathbb{C})\right\}.$$

It turns out that our dream is a reality:

Fact: If $p \neq 2$, all continuous representations of W_K are either of the form $\chi_1 \otimes \chi_2$ or $\operatorname{Ind}_{W_L}^{W_K}(\theta)$ as above.

So we guessed LLC for $GL_2(K)$! Though again, let us emphasize that this is not actually the correct version of the correspondence (it is for example not compatible with taking duals). However it will not take too much effort to make this into a correct statement next lecture.

Remark 7.5. For p = 2, the matching still works, but there are more objects on both sides.

7.4 Basic representation theory of *p*-adic groups

Let K be a local field. Then $\operatorname{GL}_n(K)$ is a topological group, with a basis of open neighborhoods of *id* given by

$$K_j = \left\{ g \in \operatorname{GL}_n(\mathbb{O}_K) \mid g = id \mod \mathfrak{m}_K^j \right\}.$$

Note that $\operatorname{GL}_n(\mathbb{O}_K)/K_j \simeq \operatorname{GL}_n(\mathbb{O}_K/\mathfrak{m}_K^j)$ is a finite group.

For example, if $K = \mathbb{Q}_p$, we have a natural surjective map

$$\operatorname{GL}_n(\mathbb{Q}_p) \supset \operatorname{GL}_n(\mathbb{Z}_p) \xrightarrow{\varphi_j} \operatorname{GL}_n(\mathbb{Z}/p^j\mathbb{Z})$$

for all $j \in \mathbb{Z}_{:} \geq 0$, and $K_j = \varphi_j^{-1}(id)$.

Remark 7.6. 1. $K_j \subset K_0$ is normal.

2. K_0 is a maximal compact subgroup.

Exercise 7.7. Let $\pi \in \mathbb{O}_K \subset K$ be a uniformizer.

1. Establish the Bruhat decomposition:


$$\operatorname{GL}_{n}(K) = \bigsqcup_{\substack{\lambda_{1} \geq \lambda_{2} \geq \ldots \geq \lambda_{n} \\ \lambda_{i} \in \mathbb{Z}}} \operatorname{GL}_{n}(\mathbb{O}_{K}) \begin{pmatrix} \pi^{\lambda_{1}} & & & \\ & \pi^{\lambda_{2}} & & \\ & & \pi^{\lambda_{3}} & & \\ & & & & \pi^{\lambda_{n}} \end{pmatrix} \operatorname{GL}_{n}(\mathbb{O}_{K})$$

$$(*)$$

(*Hint*: Gaussian elimination.)

- 2. Use (*) to classify the subgroups $\operatorname{GL}_n(\mathbb{O}_K) \subset H \subset \operatorname{GL}_n(K)$.
- 3. Hence or otherwise, show that $\operatorname{GL}_n(\mathbb{O}_K)$ is a maximal compact subgroup of $\operatorname{GL}_n(K)$.

Example 7.8. Consider $GL_1(\mathbb{Q}_3) = \mathbb{Q}_3^{\times} = \mathbb{Z} \times \mathbb{Z}_3^{\times}$. Recall our picture of the 3-adics from Example 5.6. A picture of the maximal compact subgroup K_0 of this group is:



A space is **totally disconnected** if every point admits a family of compact open neighborhoods; e.g. $\operatorname{GL}_n(K)$ is totally disconnected because each K_j is compact open. Let G be a totally disconnected topological group and V a vector space over a field \mathbb{K} of characteristic 0. We give V the discrete topology.


Definition 7.9. A representation $\rho: G \to GL(V)$ is

- 1. smooth if for all $v \in V$, stab_G v is open, and
- 2. **admissible** if for all open $K \subset G$, V^K is finite-dimensional.
- **Example 7.10.** 1. The trivial representation \mathbb{K} is smooth and admissible, \mathbb{K}^{∞} is smooth but not admissible.
 - 2. The standard representation of $\operatorname{GL}_n(K)$ on K^n is not smooth, as $\operatorname{stab}_{\operatorname{GL}_n(K)} v$ is not open for $v \neq 0$.
 - 3. The group $G = (\mathbb{Z}_p, +)$ acts on the vector space $\mathcal{F} = \{\varphi : \mathbb{Z}_p \to \mathbb{C} \mid \varphi \text{ is locally constant}\}$ in the natural way, forming the "smooth regular representation". (This is the *p*-adic analogue of $L^2(G)$.) We claim that this representation is smooth and admissible.
 - Smooth: Let $\varphi \in \mathcal{F}$. Then for all $x \in \mathbb{Z}_p$, there is a neighborhood U_x such that $\varphi|_{U_x}$ is constant. This forms a covering of \mathbb{Z}_p by open neighborhoods of the form $U_x = x + p^{n_x} \mathbb{Z}_p$. Since \mathbb{Z}_p is compact, there exists a finite subcovering U_{x_1}, \ldots, U_{x_m} . Then φ is fixed by $p^n \mathbb{Z}_p$, where $n = \text{Max}\{n_i\}$, so the the stablizer of φ is open, hence the representation is smooth.
 - Admissible: A basis of open neighborhoods of 0 is given by $p^m \mathbb{Z}_p, m \ge 0$. Then


$$\mathcal{F}^{p^m \mathbb{Z}_p} = \{ \varphi \mid \varphi \text{ is constant on } p^m \mathbb{Z}_p \text{-orbits} \}$$
$$= \{ \varphi : \mathbb{Z}/p^m \mathbb{Z} \to \mathbb{C} \}$$

is finite dimensional, so the representation is admissible.

4. (The most important example!) Recall that $\mathbb{P}^1\mathbb{C}$ is covered by the compact sets $D_{\leq 1} = \{z \mid |z| \leq 1\}$ and $D_{\leq 1}^{-1}$, so $\mathbb{P}^1\mathbb{C}$ is compact:



Similarly, $\mathbb{P}^1 K = K \cup \{\infty\}$ is covered by the compact sets $D_{\leq 1} = \{z \in K \mid |z|_p \leq 1\} = \mathbb{O}_K$ and $D_{\leq 1}^{-1}$, so $\mathbb{P}^1 K$ is compact:



The vector space

 $I = \{ f : \mathbb{P}^1 K \to \mathbb{C} \mid f \text{ is locally constant} \}$

admits a natural $\operatorname{GL}_2(K)$ -action, and the same argument as in the previous example (using compactness) shows that I is a smooth, admissible representation of $\operatorname{GL}_2(K)$. In fact,

constant functions $\hookrightarrow I \twoheadrightarrow St$

Where $St := I/\{\text{constant functions}\}$ is the **Steinberg module**. The module St is irreducible (exercise, might be hard with current technology!).

Exercise 7.11. Show that any smooth finite dimensional representation of $\operatorname{GL}_n(K)$ factors over det : $\operatorname{GL}_n(K) \to K^{\times}$. (*Hint*: the kernel of a smooth finite dimensional representation is a finite intersection of stabilizers of a basis, so it must be open and normal, hence contain $\operatorname{SL}_n(K)$).

Exercise 7.12. The representation $\mathcal{F}' = \{\varphi : \mathbb{Q}_p \to \mathbb{C} \mid \varphi \text{ is locally constant}\}$ of \mathbb{Z}_p is *not* admissible or smooth.

Lecture 8: Precise statement of local Langlands for GL_2 , $p \neq 2$

8.1 Basic representation theory of *p*-adic groups, continued

We pick up where we left off in the previous lecture. Let

$$\mathfrak{m}_K \subset \mathbb{O}_K \subset K$$

be the maximal ideal in the ring of integers of a local field. We are interested in the representation theory of the group $\operatorname{GL}_n(K)$. (Or, more generally, the representation theory of any totally disconnected group G, but for concreteness we will work with GL_n .)

Recall that the sets

$$K_j = \{ g \in \operatorname{GL}_n(\mathbb{O}_K) \mid g = id \mod \mathfrak{m}_K^j \}$$

form a basis of open neighborhoods of $id \in \operatorname{GL}_n(K)$. In addition to being open neighborhoods of the identity, the K_i are *subgroups* of $\operatorname{GL}_n(K)$. (Note the existence of such subgroups which form a basis for open neighborhoods of the identity is only possible because $\operatorname{GL}_n(K)$ is a totally disconnected group; a Lie group could not have such a family of subgroups because Lie groups have no small subgroups.)

Last week we saw that K_0 is a maximal compact subgroup of $\operatorname{GL}_n(K)$. Let V be a representation of $\operatorname{GL}_n(K)$. Because $K_0 \supset K_1 \supset K_2 \supset \cdots$, we have a chain

$$V^{K_0} \subset V^{K_1} \subset V^{K_2} \cdots$$

If V is smooth, each vector lies in V^U for some open $U \subset G$, hence lies in some V^{K_i} . So the filtration is exhaustive. If V is admissible, each V^{K_i} is finite dimensional.

Because $K_i \subset K_0$ is normal, the subspace V^{K_i} is stable under action by K_0 . The subgroup $K_i \subset K_0$ acts trivially on V^{K_i} , so the K_0 -action factors through the finite group $K_0/K_i \simeq \operatorname{GL}_n(\mathbb{O}_K/\mathfrak{m}_K^i)$; e.g. for $K = \mathbb{Q}_p$, the K_0 action on V^{K_i} factors through $\operatorname{GL}_n(\mathbb{Z}/p^i\mathbb{Z})$. (The key point here is that $K_i \subset K_0$ is normal, so the quotient K_0/K_i is a group.) Since representations of finite groups are completely reducible, we have a decomposition

$$V^{K_i} = \bigoplus_{\rho \in \widehat{K_0/K_i}} V^{K_i}(\rho),$$

where $V^{K_i}(\rho)$ is the ρ -isotypic component of V^{K_i} ; that is, $V^{K_i}(\rho)$ is the direct sum of all irreducible subrepresentations of V^{K_i} which are isomorphic to ρ . Passing to the limit, we obtain a decomposition

$$V = \bigoplus_{\rho \in \widehat{K_0}} V(\rho).$$

Here $\widehat{K_0}$ denotes all representations of K_0 which factor over some quotient K_0/K_i .

Lemma 8.1. The representation V is admissible if and only if each isotypic component $V(\rho)$ in the decomposition above is finite-dimensional.

Proof. Assume that $V(\rho)$ is infinite dimensional for some $\rho \in \widehat{K}_0$. By definition, ρ factors through K_0/K_i for some *i*. Hence, $V(\rho) \subset V^{K_i}$ is an infinite dimensional subspace and *V* is not admissible.

To prove the opposite implication, assume that each $V(\rho)$ is finite-dimensional. For each i, we have a decomposition

$$V^{K_i} = \bigoplus_{\rho \in \widehat{K_0}} V(\rho)^{K_i}.$$

But since $V(\rho)$ is the direct sum of irreducible representations which are isomorphic to ρ , we have

$$V(\rho)^{K_i} = \begin{cases} 0 & \text{if } \rho|_{K_i} \neq \text{triv,} \\ V(\rho) & \text{otherwise.} \end{cases}$$

Hence

$$V^{K_i} = \bigoplus_{\substack{\rho \in \widehat{K_0} \\ \rho|_{K_i} = \text{triv}}} V(\rho)$$

Since K_0/K_i is a finite group, there are only finitely many representations $\rho \in \widehat{K}_0$ which factor through K_0/K_i for any fixed *i*, so decomposition above is a finite direct sum of finite-dimensional representations, hence *V* is admissible.

Remark 8.2. This is like the theory of K-finite vectors in representation theory of real Lie groups. A big difference is that the representation theory of, for example $\operatorname{GL}_m(\mathbb{Z}/p^n\mathbb{Z})$ for large m and n is extremely complicated, whereas we know the representation theory of compact Lie groups rather well (highest weights, etc.).

Example 8.3. 1. Consider the \mathbb{Z}_p -representation $\mathcal{F} = \{\varphi : \mathbb{Z}_p \to \mathbb{C} \mid \varphi \text{ is locally constant}\}$ from Example 7.10.3. For an open neighborhood $p^m \mathbb{Z}_p$ of the identity, the invariants are

$$\mathcal{F}^{p^m \mathbb{Z}_p} = \{ \varphi \mid \varphi \text{ constant on } p^m \mathbb{Z}_p \text{ orbits} \}$$

= regular representation of $\mathbb{Z}_p / p^m \mathbb{Z}_p.$

Hence,

$$\mathcal{F} = igoplus_{\substack{ ext{continuous}\ \chi: \mathbb{Z}_p o \mathbb{C}^ imes}} \mathbb{C}_\chi.$$

2. Consider the $GL_2(K)$ -representation

 $I = \{ f : \mathbb{P}^1 K \to \mathbb{C} \mid f \text{ is locally constant} \}$

from example 7.10.4. Here

$$I^{K_n} = \{ \varphi : \mathbb{P}^1(\mathcal{O}_K/\mathfrak{m}_K^n) \to \mathbb{C} \},\$$

 \mathbf{SO}

$$I = \lim_{\to} \mathbb{C}[\mathbb{P}^1(\mathcal{O}_K/\mathfrak{m}_K^n)].$$

Let V be a representation of $\operatorname{GL}_n(K)$. A map $\xi : V \to \mathbb{C}$ is **smooth** if $\operatorname{stab}_{\operatorname{GL}_n(K)} \xi$ is open. Define the **smooth dual**

$$\widehat{V} = \{ \text{smooth vectors } \xi : V \to \mathbb{C} \}$$

Lemma 8.4. Assume V is a smooth representation of $\operatorname{GL}_n(K)$. If $V = \bigoplus_{\rho \in \widehat{K_0}} V(\rho)$, then $\widehat{V} = \bigoplus_{\rho \in \widehat{K_0}} V(\rho)^*$.

In particular, if V is smooth and admissible, then so is \widehat{V} , and $V \xrightarrow{\sim} \widehat{\widehat{V}}$.

Proof. The map $\xi: V \to \mathbb{C}$ is smooth if and only if ξ vanishes on all but finitely many $V(\rho)$. The lemma follows.

The goal for the remainder of this lecture will be to give a bird's eye view on the smooth admissible representations of $GL_1(K)$ and $GL_2(K)$. But first, we need a digression on norms.

8.2 Canonical norms

Recall that to make the product formula of Section 5.2 hold, we define three types of equivalence classes of multiplicative norms ("places," denoted by v) on a local field K:

- finite places: $|x|_v := (\# \mathbb{O}_K / \mathfrak{p})^{-\operatorname{val}_p(x)}$ for some prime $\mathfrak{p} \subset \mathbb{O}_K$,
- real places: $|x|_v := |i(x)|$ for some real embedding $i: K \hookrightarrow \mathbb{R}$, and
- complex places: $|x|_v := |i(x)|^2$ for some pair of conjugate embeddings $i : K \hookrightarrow \mathbb{C}$ not landing in \mathbb{R} .

Different normalizations would also yield multiplicative norms, but we chose the ones above to make the product formula

$$\prod_{\text{places } v} |x|_v = 1$$

for $x \in K^{\times}$ holds. For example, if $K = \mathbb{Q}_p$, $|p| = \epsilon$ gives a norm for any $0 < \epsilon < 1$, so why do we choose |p| = 1/p? In some sense, this choice is justified by the product formula, but it is still a little mysterious.

Tate made the following observation which further justifies this choice. For a place v, the completion K_v is is locally compact. Let μ be the additive Haar measure on K_v . The measure μ is unique up to a scalar. Define

 $|x|_v =$ factor by which $x \cdot$ scales the Haar measure;

i.e., $|x|_v = \frac{\mu(x \cdot A)}{\mu(A)}$ for $A \subset K_v$ measurable and $0 < \mu(A) < \infty$.

Example 8.5. 1. $K_v = \mathbb{R}$: For $x \in \mathbb{R}$, $|x|_v = \frac{\mu(x[0,1])}{\mu([0,1])} = \mu([0,x]) = |x|$.

2. $K_v = \mathbb{C}$: For $z \in \mathbb{C}$, and

$$\beta = \int_{0}^{1+i} \frac{\mathcal{M}(z\beta)}{\mathcal{M}(\beta)} = \mathcal{M}\left(\frac{z}{2}\right) = |z|^{2}$$

3.
$$K = \mathbb{Q}_p$$
: Recall that $\mathbb{Z}_p = \bigsqcup_{0 \le m < p} m + p\mathbb{Z}_p$, so $p\mu(p\mathbb{Z}_p) = \mu(\mathbb{Z}_p)$. Hence,
 $|x|_v = \frac{\mu(p\mathbb{Z}_p)}{\mu(\mathbb{Z}_p)} = \frac{1}{p}.$

From now on, whenever we consider a norm on a locally compact field, we will always consider this canonical norm, denoted $|\cdot|$.

8.3 Smooth admissible representations of $GL_1(K)$

Let V be a smooth admissible representation of $\operatorname{GL}_1(K) = K^{\times}$. Since V is smooth admissible,

$$V = \bigcup V^{K_i}$$

and each V^{K_i} is finite-dimensional. Furthermore, since K^{\times} is abelian, each of the subgroups $K_j := 1 + \mathfrak{m}_K^j \subset \mathbb{O}_K^{\times}$ is normal in K^{\times} , and the group

$$K^{\times}/K^{j} \simeq \mathbb{Z} \times (\mathbb{O}_{K}/\mathfrak{m}_{K}^{j})^{\times}$$

acts on V^{K_j} . Hence if V is irreducible, V is one-dimensional and determined by a character of the form $|\cdot|^c \chi: K^{\times} \to \mathbb{C}$, where $c \in \mathbb{C}$ and $\chi: \mathbb{O}_K^{\times} \to \mathbb{C}$ is a continuous character.

Remark 8.6. The category of smooth admissible representations of $GL_1(K)$ is not semisimple. For example, the representation

$$x \mapsto \begin{pmatrix} 1 & \log|x| \\ 0 & 1 \end{pmatrix}$$

is a smooth, two-dimensional admissible representation which is not semisimple.

8.4 Smooth admissible representations of $GL_2(K)$

Recall from our heuristic description of last lecture that we expect roughly two types of representations of $GL_2(K)$: "principal series" representations coming from a split torus, and "cuspidal" representations coming from an anisotropic torus.

Let $B \subset \operatorname{GL}_2(K)$ be the subgroup of upper triangular matrices. Given continuous characters $\chi_1, \chi_2 : K^{\times} \to \mathbb{C}$, define

$$I(\chi_1, \chi_2) := \{ \varphi : \operatorname{GL}_2(K) \to \mathbb{C} \mid \varphi \text{ loc. const., and } \varphi \left(\begin{pmatrix} a & b \\ 0 & d \end{pmatrix} \cdot g \right) = \chi_1(a)\chi_2(d) |\frac{a}{d}|^{1/2}\varphi(g)$$

for all $\begin{pmatrix} a & b \\ 0 & d \end{pmatrix} \in B \}.$

Example 8.7.

$$I(|\cdot|^{-1/2}, |\cdot|^{1/2}) = \{\varphi : \operatorname{GL}_2(K) \to \mathbb{C} \mid \varphi \text{ loc. const. and } \varphi \left(\begin{pmatrix} a & b \\ 0 & d \end{pmatrix} \cdot g \right) = \varphi(g) \}$$
$$= \{\varphi : \mathbb{P}^1 K \simeq G/B \to \mathbb{C} \mid \varphi \text{ locally constant} \}$$

We saw last time that $I(|\cdot|^{-1/2}, |\cdot|^{1/2})$ is smooth and admissible.

The representations $I(\chi_1, \chi_2)$ formed in this way are called **principal series representations**.

Theorem 8.8. 1. For all $\chi_1, \chi_2, I(\chi_1, \chi_2)$ is smooth and admissible.

- 2. $I(\chi_1, \chi_2) \simeq I(\chi_1^{-1}, \chi_2^{-1}).$
- 3. If $\chi_1/\chi_2 = |\cdot|^{-1}$, then we have an exact sequence of representations

$$0 \to C(\chi_1, \chi_2) \to I(\chi_1, \chi_2) \to S(\chi_1, \chi_2) \to 0$$

with dim $C(\chi_1, \chi_2) = 1$ and $S(\chi_1, \chi_2)$ irreducible.

4. If $\chi_1/\chi_2 = |\cdot|$, then we have an exact sequence of representations

$$0 \to S(\chi_1, \chi_2) \to I(\chi_2, \chi_2) \to C(\chi_1, \chi_2) \to 0$$

with dim $C(\chi_1, \chi_2) = 1$ and $S(\chi_1, \chi_2)$ irreducible.

- 5. Otherwise, $I(\chi_1, \chi_2)$ is irreducible.
- 6. If $\chi_1/\chi_2 \simeq |\cdot|^{-1}$, then $S(\chi_1,\chi_2) \simeq S(\chi_2,\chi_1)$ and $C(\chi_1,\chi_2) \simeq C(\chi_2,\chi_1)$, and if $\chi_1/\chi_2 \not\simeq |\cdot|^{\pm 1}$, then $I(\chi_1,\chi_2) \simeq I(\chi_2,\chi_1)$.

Remark 8.9. Some remarks on the theorem:

- (a) The representation $I(\chi_1, \chi_2)$ is an example of an induced representation for a totally disconnected group.
- (b) Why the strange $|\frac{a}{b}|^{1/2}$ factor? It is necessary to make 2. hold! So why does 2. hold? Consider

$$I(|\cdot|^{1/2},|\cdot|^{-1/2}) = \{\varphi \mid \varphi\left(\begin{pmatrix}a & b\\ 0 & d\end{pmatrix} \cdot g\right) = |\frac{a}{d}|\varphi(g)\}.$$

We can define a function

$$\Phi: I(|\cdot|^{1/2}, |\cdot|^{-1/2}) \to \mathbb{C}$$
$$\varphi \mapsto \int_{K_0} \varphi d\mu.$$

(One can think of $I(|\cdot|^{1/2}, |\cdot|^{-1/2})$ as being some "functions" on $\mathbb{P}^1 K$ and we are integrating over $\mathbb{P}^1 K$ to get a number. More precisely, these are "densities", but to

explain why would take us too far afield.) Here is a minor miracle: the function Φ is $\operatorname{GL}_2(K)$ -invariant, hence

$$C(|\cdot|^{1/2}, |\cdot|^{-1/2}) = \mathbb{C}$$

is the trivial representation. Now, given $\varphi \in I(\chi_1, \chi_2)$ and $\varphi' \in I(\chi_1^{-1}, \chi_2^{-1}), \varphi \varphi' \in I(|\cdot|^{1/2}, |\cdot|^{-1/2})$. By composing with Φ , we get a $\operatorname{GL}_2(K)$ -invariant pairing:

$$I(\chi_1,\chi_2) \times I(\chi_1^{-1},\chi_2^{-1}) \to \mathbb{C},$$

which turns out to be non-degenerate, establishing 2.

- (c) Part 6. is the most complicated to prove. It uses an intertwiner $I(\chi_1, \chi_2) \to I(\chi_2, \chi_1)$ via analytic continuation (there are connections to the Jantzen filtration).
- (d) Finally, note that $\operatorname{Rep} \operatorname{GL}_2(K)$ is not semisimple, so we cannot just compute homs as in the finite group case.

The other class of representations of $GL_2(K)$ are cuspidal representations.

Theorem 8.10. For every degree 2 extension L/K and continuous character $\theta : L^{\times} \to \mathbb{C}$ which does not factor through the norm, there exists an irreducible representation $BC_{L/K}(\theta)$. We have that $BC_{L/K}(\theta) \simeq BC_{L/K}(\theta')$ if and only if $\theta^{\sigma} \simeq \theta'$ for $\sigma \in \text{Gal}(L/K)$.

The construction of $BC_{L/K}(\theta)$ is complicated, via the Weil representation. What is going on metaphorically?

- Consider $\operatorname{SL}_2(\mathbb{F}_q)$. We have seen in Joe's Informal Friday Seminar talks that a character $\theta: T_a \to \mathbb{C}^{\times}$ gives rise to a local system L_{θ} on $\mathbb{P}^1_{\mathbb{F}_q} \setminus \mathbb{P}^1\mathbb{F}_q$. Taking the first cohomology yields a cuspidal representation $R^G_{T_a}(\theta)$.
- Consider $SL_2(\mathbb{R})$. A character $\theta : SO(2) \to \mathbb{C}^{\times}$ (such characters are classified by \mathbb{Z}) gives rise to a local system $\mathcal{O}(n)$ on the upper half plane, taking global sections yields a discrete series representation $\Gamma(\mathbb{H}, \mathcal{O}(n))$.
- Now take $\operatorname{GL}_2(K)$. A character $\theta: L^{\times} \to \mathbb{C}$ gives rise to a local system \mathcal{L}_{θ} on "Drinfeld space" $\mathbb{P}^1\overline{K}/\mathbb{P}^1K$. Taking first cohomology yields the representation $BC_{L/K}(\theta)$. Note that this is very technical, and is an active area of research.

This can also be viewed through the lens of Langlands functoriality. Let L/K be a degree n extension, so $W_L \subset W_K$ is an index n subgroup. We have the following diagram:

Thus an innocuous induction functor on the left hand side predicts a highly non-trivial correspondence between irreducible representation on the right hand side!

8.5 Weil-Deligne representations

We are almost ready to make a precise statement of the local Langlands correspondence for $GL_2(K)$! Recall local class field theory: there is a map

$$r_K: W_K \to K^{\times}.$$

Composing with $|\cdot|$ gives us the **norm character**

$$|\cdot|: W_K \to \mathbb{Q}^{\times}$$

An *n*-dimensional Weil-Deligne representation is a triple (ρ, V, N) , where

- V is an *n*-dimensional complex vector space,
- $\rho: W_K \to \operatorname{GL}(V)$ is a continuous representation, and
- $N \in \text{End}(V)$ is nilpotent such that

$$\rho(x)N\rho(x)^{-1} = |x|N \tag{(*)}$$

for all $x \in W_K$. (In fact, (*) forces N to be nilpotent.)

- **Example 8.11.** 1. Any *n*-dimensional continuous representation of W_K with N = 0 is a Weil-Deligne representation.
 - 2. The representation $\rho = \begin{pmatrix} |\cdot|\chi & 0\\ 0 & \chi \end{pmatrix}$ for any character $\chi : W_K \to \mathbb{C}^{\times}$ and $N = \begin{pmatrix} 0 & 1\\ 0 & 0 \end{pmatrix}$ form a Weil-Deligne representation. Indeed, for $x \in W_K$,

$$\begin{split} \rho(x)N\rho(x)^{-1} &= \begin{pmatrix} |x|\chi(x) & 0\\ 0 & \chi(x) \end{pmatrix} \begin{pmatrix} 0 & 1\\ 0 & 0 \end{pmatrix} \begin{pmatrix} |x|^{-1}\chi(x)^{-1} & 0\\ 0 & \chi(x)^{-1} \end{pmatrix} \\ &= \begin{pmatrix} 0 & |x|\\ 0 & 0 \end{pmatrix} \\ &= |x| \begin{pmatrix} 0 & 1\\ 0 & 0 \end{pmatrix}. \end{split}$$

We denote this Weil-Deligne representation $St(\chi, |\cdot|\chi)$.

A Weil-Deligne representation is F-semisimple if V is semisimple as a representation of W_K .

Exercise 8.12. Let $\widetilde{\text{Frob}} \in W_K$ be any lift of Frobenius. Show that a Weil-Deligne representation is *F*-semisimple if and only if Frob is semisimple.

8.6 The local Langlands correspondence for GL_2

Theorem 8.13. (*local Langlands correspondence for* GL_2 , $p \neq 2$) Fix a local field K of residue characteristic $p \neq 2$. There is a canonical bijection.

$$\left\{ \begin{array}{c} F\text{-semisimple} \\ 2\text{-dimensional} \\ \text{Weil-Deligne reps} \end{array} \right\}_{\simeq} \leftrightarrow \left\{ \begin{array}{c} \text{irred. smooth admiss.} \\ \text{reps of } \operatorname{GL}_2(K) \end{array} \right\}_{\simeq}$$

Moreover, this bijection is given as follows.

$$\chi_1/\chi_2 = |\cdot|^{\pm 1} : \left(\begin{pmatrix} \chi_1 & 0\\ 0 & \chi_2 \end{pmatrix}, N = 0 \right) \leftrightarrow C(\chi_1, \chi_2)$$
$$\chi_1/\chi_2 = |\cdot|^{\pm 1} : \left(\begin{pmatrix} \chi|\cdot| & 0\\ 0 & \chi \end{pmatrix}, N = \begin{pmatrix} 0 & 1\\ 0 & 0 \end{pmatrix} \right) \leftrightarrow S(\chi, |\cdot|\chi)$$
$$\chi_1/\chi_2 \neq |\cdot|^{\pm 1} : \left(\begin{pmatrix} \chi_1 & 0\\ 0 & \chi_2 \end{pmatrix}, N = 0 \right) \leftrightarrow I(\chi_1, \chi_2)$$
$$L/K, \theta : L^{\times} \to \mathbb{C}^{\times} : \left(\operatorname{Ind}_{W_K}^{W_L}(\theta), N = 0 \right) \leftrightarrow BC_{L/K}(\theta)$$


Where the character $\theta: L^{\times} \to \mathbb{C}^{\times}$ does not factor through the norm.

Lecture 9: The case of p = 2 and the Satake isomorphism

Today's lecture has two objectives: to explore what the LLC looks like when p = 2, and to examine how a simple special case of the LLC for GL_n leads to the Satake isomorphism.

9.1 Ramification filtration revisited

To begin, we revisit the ramification filtration of Section 6.4. Let L/K be a finite Galois extension where K and L are both local fields. We have the following inclusions:



Here the uniformizers π_K, π_L are the only non-canonical objects in the diagram above. We denote by $k_K = \mathbb{O}_L/\mathfrak{m}_L$ (resp. $k_L = \mathbb{O}_L/\mathfrak{m}_L$) the residue fields. There is a short exact sequence

$$I_{L/K} \hookrightarrow \operatorname{Gal}(L/K) \twoheadrightarrow \operatorname{Gal}(k_L/k_K) = \langle \operatorname{Frob} \rangle \simeq \mathbb{Z}/f\mathbb{Z}.$$

Lemma 9.1. An element $\sigma \in I_{L/K}$ in the inertia subgroup is determined by $\sigma(\pi_L)$.

This leads to the **ramification filtration** of the Galois group. Define

$$I_0 := I_{L/K}, \quad I_j := \left\{ \sigma \in I_{L/K} \mid \frac{\sigma(\pi_L)}{\pi_L} = 1 \mod \mathfrak{m}_L^j \right\}.$$

Then

$$\operatorname{Gal}(L/K) \supset I_0 \supset I_1 \supset I_2 \supset \cdots \supset \{1\}$$

is a filtration of $\operatorname{Gal}(L/K)$.

Key Facts:

- 1. The ramification filtration is a finite exhaustive filtration.
- 2. There is an injection $I_0/I_1 \hookrightarrow \mathbb{O}_L^{\times}/(1 + \mathbb{O}_L^{\times}) \simeq k_L^{\times} : \sigma \mapsto \frac{\sigma(\pi_L)}{\pi_L}$. Hence I_0/I_1 is cyclic of order prime to p.
- 3. For $j \ge 1$, there is an injection $I_j/I_{j+1} \hookrightarrow (1 + \mathfrak{m}_L^j)/(1 + \mathfrak{m}_L^{j+1}) \simeq (k_L, +) : \sigma \mapsto \frac{\sigma(\pi_L)}{\pi_L}$. Hence I_j/I_{j+1} is an abelian *p*-group, and I_1 is a Sylow *p*-subgroup of $I_{L/K}$.

This filtration leads to some nomenclature: The first subquotient $\operatorname{Gal}(L/K)/I_{L/K}$ is canonically isomorphic to $\mathbb{Z}/f\mathbb{Z}$, and is referred to as the **unramified** part of the Galois group. The second subquotient $I_{L/K}/I_1$ is cyclic of order prime to p, and is referred to as the **tamely ramified** part of the Galois group. The remaining subquotients of the ramification filtration are abelian p-groups (and hence I_1 is a solvable p-group), and are referred to as the **wildly ramified** part of the Galois group. The upshot is that there are significant constraints on which groups can appear as Galois groups of extensions of local fields. (For example, they must be solvable.) This is in sharp contrast to the number field setting, where many types of groups can appear as Galois groups of extensions of \mathbb{Q} . (Though exactly which groups appear as Galois groups of number fields is still very much an open problem, the *inverse Galois problem*.)

Example 9.2. Consider the extension $L = \mathbb{Q}_p(\sqrt[p-1]{p})$ of $K = \mathbb{Q}_p$. Here $\pi_L = \sqrt[p-1]{p}$ and $\pi_K = p$ are uniformizers. (Exercise: Show that the set μ_{p-1} of all $(p-1)^{st}$ roots of unity is contained in \mathbb{Q}_p .) The Galois group of this extension is

$$\operatorname{Gal}(L/K) = \{ \sigma_{\zeta} : \sqrt[p-1]{p} \mapsto \zeta \sqrt[p-1]{p} \text{ for } \zeta \in \mu_{p-1} \} \simeq k_K^{\times} \hookrightarrow k_L^{\times}.$$

One can check that $\operatorname{Gal}(L/K)$ is totally ramified; that is, $I_{L/K} = \operatorname{Gal}(L/K)$. (This follows from the observation that we are adjoining roots of p, whose image is zero in the residue field.) Furthermore, if $\sigma_{\zeta} \in I_{L/K} = \operatorname{Gal}(L/K)$, then $\frac{\sigma_{\zeta}(\pi_L)}{\pi_L} = \zeta \in k_L^{\times}$, hence $I_1 = \{1\}$ and $\operatorname{Gal}(L/K)$ is tamely ramified.

Remark 9.3. Examples of wild ramification are almost always hard! We really should spend a lecture on such examples, but we are quickly running out of time, so sadly we will not.

Next we will examine the structure of the absolute Galois group of a local field. For a local field K, we have an exact sequence

$$I_{\overline{K}/K} \hookrightarrow \operatorname{Gal}(\overline{K}/K) \twoheadrightarrow \widehat{\mathbb{Z}}.$$

We would like to pass to the limit to obtain a ramification filtration of the inertia subgroup $I_{\overline{K}/K}$ from the ramification filtrations of the inertia subgroups of finite extensions. However, there is a problem: if L'/L/K is a tower of finite extensions, then the ramification filtration of $I_{L'/K}$ is related to multiples of the ramification filtration of $I_{L/K}$.

This can be fixed through an "upper numbering" procedure which replaces I_j with $I_{L/K}^{\lambda}$ for $\lambda \in \mathbb{Q}_{\geq 0}$ in a way that is compatible with extensions. (Exactly how one does this appears pretty crazy at first sight. It is explained in Serre's Local Fields [Ser79].) This leads to a ramification filtration $I_{K/K}^{\lambda}$ of the inertia subgroup of the absolute Galois group indexed by rational numbers:

$$\operatorname{Gal}(\overline{K}/K) \supset I_{\overline{K}/K} \supset I_{\overline{K}/K}^{\geq 0} \supset \cdots \supset I_{\overline{K}/K}^{463/5} \supset \cdots$$

This filtration has the property that $\operatorname{Gal}(\overline{K}/K)/I_{\overline{K}/K} = \widehat{\mathbb{Z}}$ canonically, $I_{\overline{K}/K}/I_{\overline{K}/K}^{\geq 0} \simeq \prod_{\substack{l \neq p \\ \text{prime}}} \mathbb{Z}_l$

non-canonically, and other subquotients are pro-p groups.

Important points:

- 1. The first two steps depend only on the residue characteristic of the field.
- 2. Via class field theory, the image of this filtration in $W_K^{ab} \simeq K^{\times}$ corresponds to the filtration by $1 + \mathfrak{m}_K^j \subset \mathbb{O}_K$. The fact that the only jumps in this filtration are at integers (as opposed to other elements of \mathbb{Q}) is the **Hasse-Arf Theorem**.

9.2 More details on Weil–Deligne representations

Next we would like to show two things: (1) why any Weil–Deligne representation is "close" to a continuous representation of $\operatorname{Gal}(\overline{K}/K)$ and (2) why p = 2 is special in the local Langlands correspondence.

Proposition 9.4. Any indecomposible *F*-semisimple Weil–Deligne representation is isomorphic to $St_n \otimes \rho$, where ρ is an irreducible representation of W_K .

Here St_n is the Steinberg representation from the previous lecture; e.g.

$$St_4 = \left(\begin{pmatrix} |\cdot|^3 & 0 & 0 & 0\\ 0 & |\cdot|^2 & 0 & 0\\ 0 & 0 & |\cdot| & 0\\ 0 & 0 & 0 & 1 \end{pmatrix}, N = \begin{pmatrix} 0 & 1 & 0 & 0\\ 0 & 0 & 1 & 0\\ 0 & 0 & 0 & 1\\ 0 & 0 & 0 & 0 \end{pmatrix} \right)$$

The proof of this proposition is left as a somewhat tricky exercise. It becomes easier if you know what the weight filtration associated to a nilpotent operator is.

- **Proposition 9.5.** 1. Let $\rho: W_K \to \operatorname{GL}(v)$ be an irreducible representation. Then there exists a continuous character $\chi: W_K \to \mathbb{C}^{\times}$ such that $\rho \otimes \chi$ has finite image and hence defines a representation $\rho \otimes \chi: \operatorname{Gal}(\overline{K}/K) \to \operatorname{GL}(V)$.
 - 2. Suppose that $\rho: W_K \to \operatorname{GL}(V)$ is irreducible and not induced from any proper subgroup of W_K . Then the restriction to wild intertia is irreducible. In particular, dim Vis a power of p (since any irreducible module over a p-group has dimension divisible by p).

Remark 9.6. Proofs of these two statements can be found in [Tat79].

Propositions 9.4 and 9.5.1 show that every Weil–Deligne representation is "close" to a representation of the absolute Galois group, in the sense that every Weil-Deligne representation can be obtained from the Steinberg representation and an irreducible representation of W_K , and every irreducible representation of W_K can be upgraded to a representation of Gal (\overline{K}/K) by tensoring with a character.

The two statements of Proposition 9.5 are reasonably easy consequences of the following lemma.

Lemma 9.7. Suppose a group G has the form

$$\Gamma \hookrightarrow G \twoheadrightarrow \mathbb{Z}$$

for some finite group Γ . Then any irreducible *G*-module is either irreducible over Γ or induced from a subgroup of the form $\Gamma \rtimes m\mathbb{Z}$.

The proof of this lemma is a worthwhile exercise!

9.3 Why is LLC for p = 2 special?

Proposition 9.5 shows that for $p \neq 2$, all irreducible 2-dimensional representations of W_K are induced from a finite index subgroup. However, for p = 2, it's possible that there are irreducible representations of W_K which are not induced. So do such representations exist? Yes!

Consider a continuous two-dimensional representation $\rho : \operatorname{Gal}(\overline{K}/K) \to \operatorname{GL}_2(\mathbb{C})$. By the no small subgroups lemma, the image of ρ must lie in a finite subgroup of $\operatorname{GL}_2(\mathbb{C})$, so in the composition of ρ with the projection

$$\operatorname{GL}_2(\mathbb{C}) \to \operatorname{PGL}_2(\mathbb{C}),$$

the image must be conjugate to a subgroup of the maximal compact subgroup $SO_3 \subset PGL_2(\mathbb{C})$. The finite subgroups of SO_3 were classified⁵! They are of the following types:

- cyclic (orientation preserving symmetries of the product of an *m*-gon and an interval, fixing one end)
- dihedral (orientation preserving symmetries of the product of an *m*-gon and an interval)
- A_4 (orientation preserving symmetries of the tetrahedron)
- symmetries of the cube
- A_5 (orientation preserving symmetries of the icosahedron)

Reducible representations have images in cyclic subgroups of SO_3 , and induced representations have images which are dihedral groups. What about the other three? Are there any representations of $\operatorname{Gal}(\overline{K}/K)$ whose image lies in any of the final three finite subgroups? Since $\operatorname{Gal}(\overline{K}/K)$ is solvable, we can eliminate the non-solvable group A_5 from our list. Let's consider the composition series of A_4 :

$$K_4 = \mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/2\mathbb{Z} \hookrightarrow A_4 \twoheadrightarrow \mathbb{Z}/3\mathbb{Z}$$

By the structure of the ramification filtration, this subgroup structure is only possible for a local Galois group if p = 2. It turns out that it does indeed occur for some local fields!

The upshot is that for p = 2, there are more representations on each side of the LLC, and the extra representations on the Weil group side are this special class of irreducible non-induced representations whose image lies in A_4 . (Geordie isn't sure if representations corresponding to the symmetries of the cube exist. He is told that they do...)

A mystery to ponder: Let G be a compact Lie group (e.g. a finite group), and let $R(G)_{\mathbb{C}}$ be its representation ring. What is a character

$$\theta: R(G)_{\mathbb{C}} \to \mathbb{C}?$$

⁵by Klein in 1884, [Kle93]

9.4 Unramified representations

One way to convince yourself that the LLC is amazing is to see that simple special cases already have deep consequences. The first example of this that we have seen is local class field theory. The second example we will see now!

The local Langlands correspondence for $GL_n(K)$ says that there is a canonical bijection:

$$\left\{\begin{array}{c} F\text{-semisimple}\\ n\text{-dimensional}\\ \text{Weil-Deligne reps}\end{array}\right\}_{\simeq} \xleftarrow{}^{1:1} \left\{\begin{array}{c} \text{irred smooth}\\ \text{admissible}\\ \text{reps of } \mathrm{GL}_n(K)\end{array}\right\}_{\simeq}$$

On the left hand side of this bijection, we can consider a special class of **unramified Weil-Deligne representations** consisting of those representations of W_K which are trivial on the inertia subgroup. The corresponding representations on the right hand side are the **spherical representations** of $GL_n(K)$:

$$\left\{ \begin{array}{l} \text{Weil-Deligne reps} \\ \text{s.t. } N = 0 \text{ and } \rho \\ \text{factors through } \mathbb{Z}: \\ W_k \twoheadrightarrow \mathbb{Z} \hookrightarrow \text{GL}_n(\mathbb{C}) \end{array} \right\}_{/\simeq} \xleftarrow{1:1} \left\{ \begin{array}{l} \text{reps of } \text{GL}_n(K) \\ \text{admitting a} \\ \text{GL}_n(\mathbb{O}_K)\text{-fixed vector} \end{array} \right\}_{/\simeq} \right\}_{/\simeq}$$

Semisimple representations of W_K which factor through \mathbb{Z} are in bijection with semisimple elements of $\operatorname{GL}_n(\mathbb{C})$, and irreducible representations of $\operatorname{GL}_n(K)$ admitting a $\operatorname{GL}_n(\mathbb{O}_K)$ -fixed vector are in bijection with irreducible representations of the "spherical Hecke algebra" (which you are not expected to be familiar with and we will soon define). Thus, the restriction of the local Langlands correspondence to this special case results in a bijection

$$\left\{\begin{array}{c} \text{semisimple elements} \\ \text{in } \operatorname{GL}_n(\mathbb{C}) \end{array}\right\}_{/\operatorname{conj}} \stackrel{1:1}{\longleftrightarrow} \left\{\begin{array}{c} \text{irreducible reps of} \\ \mathcal{H}_{sph} := \mathcal{H}(\operatorname{GL}_n(\mathbb{O}_K), \operatorname{GL}_n(K)) \end{array}\right\}_{/\simeq}$$

This is the **Satake isomorphism**! We will spend the rest of the lecture explaining this bijection (particularly the right hand side) in more detail.

Remark 9.8. The left-hand-side of the bijection above is independent of K, and even of the residue characteristic p!

9.5 Hecke algebras

Suppose G is a finite group.

Case 1: Consider $N \subset G$ a normal subgroup. If V is a G-representation, then G acts on V^N (because for $n \in N, g \in G$, and $v \in V^N$, $n \cdot gv = g \cdot g^{-1}ng \cdot v = gv$), and the action factors over G/N. Moreover, one can check that $\operatorname{End}(\operatorname{Ind}_N^G \mathbb{C}) \simeq \mathbb{C}[G/N]$. Hence we have a bijection

$$\left\{\begin{array}{c} \text{irred } G\text{-modules} \\ \text{with an } N\text{-fixed vector} \end{array}\right\}_{/\simeq} \leftrightarrow \{\text{ irred } G/N\text{-modules}\}_{/\simeq}.$$

Case 2: Consider $H \subset G$ not necessarily normal. Given a *G*-representation *V*, what acts on V^H ? The Hecke algebra! The operator

$$\pi_H : V \to V^H$$
$$v \mapsto \frac{1}{|H|} \sum_{h \in H} h \cdot v$$

projects onto *H*-invariants. This can be used to define a "Hecke operator" [HgH] for every $g \in G$ which makes the following diagram commute:

$$V^{H} \xrightarrow{[HgH]} V^{H}$$
$$\int_{V} & \uparrow_{\pi_{H}}$$
$$V \xrightarrow{\cdot g} V$$

Note that all g in the same double coset yield the same Hecke operator. Alternatively, this operator is the sum

$$[HgH] = \frac{1}{|H|} \sum_{g' \in HgH} g'.$$

The **Hecke algebra** $\mathcal{H}(H,G)$ of the pair (H,G) is the vector space ${}^{H}\mathbb{C}[G]^{H}$ with multiplication

$$(f * f')(g) := \frac{1}{|H|} \sum_{g=hh'} f(h)f'(h).$$

This is an associative unital algebra with unit

$$1_H = \frac{1}{|H|} \sum_{h \in H} h.$$

Example 9.9. 1. If N is normal, $\mathcal{H}(N, G) = \mathbb{C}[G/N]$.

2. If
$$G = \operatorname{GL}_n(\mathbb{F}_q)$$
 and $B = \left\{ \begin{pmatrix} * & \cdots & * \\ 0 & \ddots & \vdots \\ 0 & 0 & * \end{pmatrix} \right\}$, then $\mathcal{H}(B, G)$ is the "Hecke algebra of S_n

at $q = |\mathbb{F}_q|^n$. This algebra is almost independent of q.

Exercise 9.10. (Do it!) Show that

$$\operatorname{End}(\operatorname{Ind}_{H}^{G}\mathbb{C})\simeq\mathcal{H}(H,G).$$

Hence

$$\langle \operatorname{Ind}_{H}^{G} \mathbb{C} \rangle \xrightarrow{\sim} \mathcal{H}(H, G) \operatorname{-mod}$$

(Here the angle brackets mean the smallest abelian category generated by kernels, cokernels, extensions, and direct sums.) Deduce that

$$\left\{\begin{array}{c} \text{irred. } G\text{-modules} \\ \text{with } H\text{-fixed vector} \end{array}\right\} \stackrel{1:1}{\longleftrightarrow} \left\{\text{irred } \mathcal{H}(H,G)\text{-modules}\right\}.$$

Remark 9.11. There is a tendency in the literature to consider one subgroup H at a time, but one can also consider all subgroups (or a particularly nice family of subgroups) at the same time, resulting in a "Hecke algebroid".

We can also define Hecke algebras of *p*-adic groups. Let $G = \operatorname{GL}_n(K)$ for a local field K, and $K_0 = \operatorname{GL}_n(\mathbb{O}_K)$ the maximal compact subgroup. Then the "big" Hecke algebra of G is

$$\mathcal{H}^{big} = \left\{ \varphi : G \to \mathbb{C} \middle| \begin{array}{c} \varphi \text{ locally constant} \\ \text{compact support} \end{array} \right\}.$$

An alternate description is

$$\mathcal{H}^{big} = \bigcup_{i} \left\{ \varphi : G \to \mathbb{C} \, \middle| \begin{array}{c} \varphi \text{ locally constant on } K_i \text{-double} \\ \text{cosets, non-zero on finitely many} \end{array} \right\}.$$

Exercise 9.12. Prove that the two formulations of \mathcal{H}^{big} are equivalent.

The algebra structure on \mathcal{H}^{big} is given by

$$(f * f')(g) = \int_{h \in G} f(h) f'(h^{-1}g) d\mu,$$

where μ is the Haar measure.

Example 9.13. Let 1_{K_i} be the indicator function on K_i . Then

$$1_{K_i} * 1_{K_i}(g) = \int_{h \in G} 1_{K_i}(h) 1_{K_i}(h^{-1}g) d\mu = \begin{cases} 0 & \text{if } g \notin K_i \\ \int_{K_i} 1 d\mu & \text{if } g \in K_i \end{cases}$$

In other words,

$$1_{K_i} * 1_{K_i} = \mu(K_i) 1_{K_i},$$

so 1_{K_i} is a quasi-idempotent.

Remark 9.14. Because any irreducible *G*-module has $V^{K_i} \neq 0$ for some *i*, \mathcal{H}^{big} can be used to understand all smooth admissible representations of *G*. However, it is very complicated.

Assume $\mu(K_0) = 1$ so 1_{K_0} is idempotent. The **spherical Hecke algebra** is

$$\mathcal{H}^{sph} = \mathcal{H}(K_0, G) := 1_{K_0} \mathcal{H}^{big} 1_{K_0}.$$

Exercise 9.15. (Do it!) Prove the Cartan decomposition of G:

$$G = \bigsqcup_{\substack{\lambda \\ \lambda_1 \ge \lambda_2 \ge \ldots \ge \lambda_n \\ \lambda_i \in \mathbb{Z}}} K_0 \begin{pmatrix} \pi^{\lambda_1} & & & \\ & \pi^{\lambda_2} & & \\ & & \pi^{\lambda_3} & & \\ & & & \ddots & \\ & & & & \pi^{\lambda_n} \end{pmatrix} K_0$$

Hence

$$\mathcal{H}^{sph} = \bigoplus_{\underline{\lambda}} \mathbb{C} 1_{\underline{\lambda}}.$$

There are two miracles.

Theorem 9.16. 1. The spherical Hecke algebra \mathcal{H}^{sph} is commutative.

2. (The Satake isomorphism) There exists a canonical bijection

$$\mathcal{H}^{sph} \stackrel{\sim}{\leftrightarrow} R({}^{L}\mathrm{GL}_{n}(\mathbb{C}))$$

Remark 9.17. The Langlands dual group ${}^{L}\operatorname{GL}_{n}(\mathbb{C}) \simeq \operatorname{GL}_{n}(\mathbb{C})$ so we could have replaced the right hand side of the Satake isomorphism with the representation ring of $\operatorname{GL}_{n}(\mathbb{C})$; however, the theorem also holds for general reductive groups and there the dual group is important.

Recall our mystery from earlier in the lecture: For a compact Lie group G, what is a character $\theta : R(G)_{\mathbb{C}} \to \mathbb{C}$ of its representation ring? By the Chevalley restriction theorem, $R(G)_{\mathbb{C}} \simeq R(T)_{\mathbb{C}}^W$, where $T \subset G$ is a maximal torus and W is the Weyl group of G. So a character of $R(G)_{\mathbb{C}}$ is just a choice of a semisimple conjugacy class in G!

Theorem 9.16 can be used to establish unramified LLC:

$$\begin{array}{c} \text{"spherical representations";} \\ \text{i.e. smooth admissible} \\ \text{irred reps of } G \text{ with} \\ \text{a } K_0\text{-fixed vector} \end{array} \right\}_{/\simeq} \\ \end{array} \\ \begin{array}{c} \underbrace{1:1}_{\text{Hecke algebra magic}} \left\{ \begin{array}{c} \text{irreducible} \\ \mathcal{H}(K_0, G) = \mathcal{H}^{sph} - \\ \text{modules} \end{array} \right\}_{/\simeq} \\ \end{array} \\ \begin{array}{c} \underbrace{1:1}_{\text{commutativity of } \mathcal{H}^{sph}} \left\{ \begin{array}{c} \text{characters} \\ \chi : \mathcal{H}^{sph} \to \mathbb{C} \end{array} \right\} \\ \underbrace{\frac{1:1}{\text{Satake isomorphism}}} \left\{ \begin{array}{c} \text{characters} \\ \theta : R(^L \text{GL}_n(\mathbb{C})) \to \mathbb{C} \end{array} \right\} \\ \underbrace{\frac{1:1}{\text{the mystery from earlier}}} \left\{ \begin{array}{c} \text{conjugacy classes} \\ \text{of semisimple} \\ \text{elts in } ^L \text{GL}_n(\mathbb{C}) \end{array} \right\} \\ \underbrace{\frac{1:1}{\text{definition}}} \left\{ \begin{array}{c} \text{unramified} \\ n\text{-dimensional} \\ \text{Weil-Deligne} \\ \text{representations} \end{array} \right\}_{\simeq} \end{array}$$

Lecture 10: The big picture

Today is about the big picture. We start with the very big picture, and finish with the moderately big picture. This is also the final lecture of the first term of this course!

10.1 The very big picture

10.1.1 Dimension 0

Let us go back to the beginning. Let $f(x) \in \mathbb{Z}[x]$ be a polynomial; e.g. $f(x) = x^2 + 1$. Back in March, we wondered: How many solutions does f(x) have modulo a prime p? We constructed tables:

p	2	3	5	7	11	13	17	19	23	
# of sol's mod p	1	0	2	0	0	2	2	0	0	
$p \mod 4$	2	3	1	3	3	1	1	3	3	

Then we studied this via representation theory. The Galois group $\operatorname{Gal}(\overline{\mathbb{Q}}/\mathbb{Q})$ acts on the roots $\{\sigma_1, \ldots, \sigma_n\} \subset \overline{\mathbb{Q}}$ of f(x), so we have a permutation representation

$$\operatorname{Gal}(\overline{\mathbb{Q}}/\mathbb{Q}) \to \operatorname{GL}(H),$$

where $H := \bigoplus_{i=1}^{n} = \mathbb{C}\sigma_i$. Then for unramified primes,

solutions of $f(x) \mod p = \operatorname{Tr}(\operatorname{Frob}_p, H)$.

Even in this innocent ("dimension 0") case, H is enormously complicated. To simplify things, we instead considered a collection of *local* representations $H_{\mathbb{Q}_p}$, defined as follows. For each p, consider roots $\sigma'_1, \ldots, \sigma'_n$ of f(x) in $\overline{\mathbb{Q}}_p$. Then for each p we have "local Galois representations"

$$\operatorname{Gal}(\overline{\mathbb{Q}}_p/\mathbb{Q}_p) \to \operatorname{GL}(H_{\mathbb{Q}_p})$$

where $H_{\mathbb{Q}_p} = \bigoplus \mathbb{C}\sigma'_i$. This gives us a "categorification" of the table above:

p	2	3	5	7	11	13	17	19	23]
	$H_{\mathbb{Q}_2}$	$H_{\mathbb{Q}_3}$	$H_{\mathbb{Q}_5}$	$H_{\mathbb{Q}_7}$	$H_{\mathbb{Q}_{11}}$	$H_{\mathbb{Q}_{13}}$	$H_{\mathbb{Q}_{17}}$	$H_{\mathbb{Q}_{19}}$	$H_{\mathbb{Q}_{23}}$]

If p is unramified, then the inertia subgroup $I \subset \operatorname{Gal}(\overline{\mathbb{Q}_p}/\mathbb{Q}_p)$ acts trivially on $H_{\mathbb{Q}_p}$, so the local Galois representation is unramified. By the Satake isomorphism (Theorem 9.16), this implies that $H_{\mathbb{Q}_p}$ is determined by a semisimple conjugacy class $[x] \in \operatorname{GL}_n(\mathbb{C})$, and

solutions of
$$f(x) \mod p = \operatorname{Tr}([x])$$
.

For unramified primes, the representation $H_{\mathbb{Q}_p}$ is rather simple. However, for ramified primes, the representation $H_{\mathbb{Q}_p}$ can be quite complicated:

- 1. The study of $H_{\mathbb{Q}_p}$ lets us define local factors in the Artin *L*-function.
- 2. We can hope to understand $H_{\mathbb{Q}_p}$ through the local Langlands correspondence.

Remember our slogan: There is a lot of substance at ramified primes/points!


10.1.2 Dimension ≥ 1

The classic example is that of an elliptic curve E; e.g. the projective completion of the curve $y^2 + y = x^3 + x^2 + 3x + 5$ that we studied in the lecture on the Sato-Tate conjecture, Lecture 4. What is the analogue of the Galois representation H in this setting?

Recall that E is a group, and the complex points of E are

 $E(\mathbb{C}) =$ solutions over $\mathbb{C} = \mathbb{C}/\Lambda$,


where $\Lambda \subset \mathbb{C}$ is a lattice That is, we obtain $E(\mathbb{C})$ by identifying opposite edges in this picture, where the dots represent elements of Λ :



The Galois group $\operatorname{Gal}(\overline{\mathbb{Q}}/\mathbb{Q})$ does not act in any meaningful way on $E(\mathbb{C})$. It does act on $E(\overline{\mathbb{Q}})$, but this is an enormously complicated set, a little too complicated for us! However, for any prime ℓ , we can consider the " ℓ^m -torsion points":

$$E[\ell^m] := \{ x \in E(\overline{\mathbb{Q}}) \mid \ell^m \cdot x = 0 \} \simeq (\mathbb{Z}/\ell^m \mathbb{Z})^2;$$

e.g., for $\ell = 3, m = 1$:



There is a natural action of $\operatorname{Gal}(\overline{\mathbb{Q}}/\mathbb{Q})$ on $E[\ell^m]$. The **Tate module** is formed by taking the direct limit of the $E[\ell^m]$:

$$T_{\ell}(E) := \lim_{\leftarrow} E[\ell^m] \simeq \mathbb{Z}_{\ell}^2$$

The Tate module has a continuous action of $\operatorname{Gal}(\overline{\mathbb{Q}}/\mathbb{Q})$. Moreover, if $E_{\mathbb{F}_p}$ is smooth, then

$$#E(\mathbb{F}_p) = 1 + p - \operatorname{Tr}(\operatorname{Frob}_p, T_{\ell}(E)).$$

So in this classic example of an elliptic curve, the Tate modules play the role of the representation H which appeared in the dimension 0 setting. Notice that in the previous section we constructed a single representation H, but there is one Tate module for each prime ℓ . This is an embarrassment of riches!

The Tate module $T_{\ell}(E)$ is an example of " ℓ -adic cohomology":

$$T_{\ell}(E) = H^1_{\acute{e}t}(E, \mathbb{Z}_{\ell})^*.$$

In general, given a variety X over a field k and a prime ℓ such that multiplication by ℓ is non-zero in k, there is a continuous action of $\operatorname{Gal}(\overline{k}/k)$ on the ℓ -adic cohomology groups $H^*_{\acute{e}t}(X_{\overline{k}}, \mathbb{Q}_{\ell})$. Again, it is useful to study these representations via their restriction to local Galois groups $\operatorname{Gal}(\overline{\mathbb{Q}}_p/\mathbb{Q}_p)$. We can always calculate these after base change to $\operatorname{Spec} \mathbb{Q}_p$ and at primes of good reduction after base change to $\operatorname{Spec} \mathbb{F}_p$. (Exercise: Think about what this statement means for $f(x) \in \mathbb{Z}[x]$.)


In addition to étale cohomology, one has several other methods for associating cohomology groups to the variety X:

- 1. singular cohomology: $H^i(X(\mathbb{C}), \mathbb{Z}), H^i(X(\mathbb{C}), \mathbb{F}_p)$ (related via universal coefficient theorem)
- 2. deRham cohomology: $H^i_{dR}(X)$, $H^i_{dR}(X_{\mathbb{F}_p})$ (cohomology of differential forms)
- 3. crystalline cohomology: $H^i_{crys}(X_{\mathbb{F}_p}/\mathbb{Z}_p)$ (a fancy theory that produces \mathbb{Z}_p -vector spaces for \mathbb{F}_p -schemes)

Grothendieck's philosophy: All of these cohomology groups should be shadows of a unique object, the "motive" of X.

Scholze: Perhaps the "motive" is more like a sheaf/local system on Spec $\mathbb{Z} \times \text{Spec } \mathbb{Z}$.

Scholze's ICM picture:



Scholze also predicts an archimidean theory for varieties in characteristic p which has been missing since the beginning of this subject!

Recommendation/Exercise: Read section 10 of Scholze's ICM paper.

How does one compare columns in Scholze's picture? In other words, if $G_{\mathbb{Q}_p} = \operatorname{Gal}(\mathbb{Q}_p/\mathbb{Q}_p)$, how can we compare

$$\rho: G_{\mathbb{Q}_p} \to \mathrm{GL}_n(\mathbb{Q}_\ell) \text{ and } \rho': G_{\mathbb{Q}_p} \to \mathrm{GL}_n(\mathbb{Q}_{\ell'})?$$

The problem is the topology. The solution is given by Weil–Deligne representations.

A topological group Γ is **pro-**p if it is profinite and for all open normal subgroups $N \subset \Gamma$, Γ/N is a p-group.

Example 10.1. Two examples of pro-*p*-groups are:

- 1. wild inertia $\subset G_{\mathbb{Q}_p}$, and
- 2. $1 + p \operatorname{Mat}_n \mathbb{Z}_p = K_1 \subset \operatorname{GL}_n(\mathbb{Q}_p).$

Lemma 10.2. Any continuous group homomorphism

$$\rho: \Gamma \to G$$

from a pro-p group Γ to a pro- ℓ group G is trivial.

Corollary 10.3. For a pro-p group Γ , any continuous group homomorphism

$$\rho: \Gamma \to \mathrm{GL}_n(\mathbb{Q}_p)$$

has finite image.

We have seen (via the ramification filtration) the $G_{\mathbb{Q}_p}$ has the following structure:

WILD)
$$pro-p$$

TAME) $TT Z_{q'} = Z_{l} \times TT Z_{l'}$
 Q_{p}
UNRAMIFIED) \widehat{Z}

Grothendieck showed us that the pro- ℓ' group $\prod_{\ell' \neq p,\ell} \mathbb{Z}_{\ell'} \subset I$ must have finite image. Moreover, $\rho(1_{\ell}) \in \operatorname{GL}_n(\mathbb{Q}_{\ell})$ is almost unipotent. So what Grothendieck has shown us is that we can "take logs to get Weil–Deligne representations". **Theorem 10.4.** (Grothendieck) After identifying $\overline{\mathbb{Q}}_{\ell}$ with \mathbb{C} , one has a canonical injection

$$\left\{\begin{array}{c} \text{cts. reps.}\\ \rho: G_{\mathbb{Q}_p} \to \mathrm{GL}_n(\mathbb{Q}_\ell) \end{array}\right\} \hookrightarrow \left\{\begin{array}{c} \text{Weil-Deligne reps}\\ \text{of } G_{\mathbb{Q}_p} \text{ over } \mathbb{C} \end{array}\right\}.$$

Notice that the set on the right is independent of $\ell!$

Remark 10.5. We are being a bit lazy, but one can identify the image of this injection.

10.2 Local Langlands correspondence for split groups

Let G be a split reductive algebraic group over \mathbb{Z} determined by the root datum $(X^* \supset R, X_* \supset R^{\vee})$, and \widehat{G} its dual group, determined by the opposite root datum $(X_* \supset R^{\vee}, X^* \supset R)$. Fix a local field K and set $q = |\mathbb{O}_K/\mathfrak{m}_K|$.

A Weil–Deligne representation in \widehat{G} is a pair (ρ, e) , where

- $\rho: W_K \to \widehat{G}(\mathbb{C})$ is a continuous group homomorphism, and
- $e \in \operatorname{Lie} \widehat{G}(\mathbb{C})$ is a nilpotent element

such that $\rho(g)e\rho(g)^{-1} = |g|e$ for all $g \in W_K$. A Weil–Deligne representation in \widehat{G} is *F*-semisimple if ρ is semisimple.

Example 10.6. A Weil–Deligne representation in GL_n is just an *n*-dimensional Weil–Deligne representation, as in Section 8.5.

Given a Weil–Deligne representation (ρ, e) , consider

$$Z_{\widehat{G}}(\rho, e) = \{ g \in \widehat{G} \mid g \cdot (\rho, e) = (\rho, e) \}.$$

Theorem 10.7. (Local Langlands correspondence) There is a canonical correspondence

$$\left\{\begin{array}{c}F\text{-semisimple}\\\text{WD reps in }\widehat{G}\end{array}\right\}_{/\widehat{G}\text{-conj}} \xleftarrow{\text{1:finite}} \left\{\begin{array}{c}\text{irred smooth}\\\text{admissible reps}\\\text{of }G(K)\end{array}\right\}_{/\simeq}$$

Fibres of this map should be indexed by irreducible representations of $Z_{\widehat{G}}(\rho, e)/Z_{\widehat{G}}(\rho, e)^{\circ}$ and are called "L-packets".

10.3 The Deligne–Langlands conjecture

Last week we examined (for $G = GL_n$) a simple special case of the local Langlands correspondence, the case of unramified WD representations, and found that it followed from the Satake isomorphism. Another slightly less simple special case of the LLC is given by **tamely ramified WD representations with unipotent monodromy** (TRUM). By restricting the correspondence in Theorem 10.7 to TRUM, we hope to obtain a correspondence:

$$\left\{\begin{array}{c} \text{TRUM; i.e. } (\rho, e) \\ \text{s.t. } \rho \text{ factors} \\ W_K \twoheadrightarrow \mathbb{Z}, e \text{ arbitrary} \end{array}\right\} \xleftarrow{\text{1:finite}} \left\{\begin{array}{c} \text{reps with an} \\ \text{Iwahori fixed} \\ \text{vector} \end{array}\right\}$$

By analogous arguments to the ones we made last week for GL_n , the set of ρ which factor through $W_K \to \mathbb{Z}$ is in bijection with the set of conjugacy classes of semisimple elements in \widehat{G} . Hence the left hand side of the correspondence above is in bijection with the set

$$\{(s,e) \mid s \in \widehat{G} \text{ semisimple }, e \in \operatorname{Lie} \widehat{G} \text{ nilpotent s.t. } ses^{-1} = qe \}_{/\widehat{G} \operatorname{-conj}}$$

The right hand side of the corrspondence above is in bijection with the set

{irred reps of the "Iwahori-Hecke algebra"
$$\mathcal{H}_{aff} := \mathcal{H}(I, G(K))$$
}.

This motivates the following conjecture of Deligne–Langlands.

The Deligne–Langlands conjecture: As in the set-up above, let $q = |\mathbb{O}_K/\mathfrak{m}_K|$ be the residue characteristic of the local field K. There is a bijection:

$$\begin{cases} (s, e, \chi) & s \in \widehat{G}(\mathbb{C}) \text{ semisimple,} \\ e \in \operatorname{Lie} \widehat{G} \text{ nilpotent, and} \\ \chi \text{ irred rep of } \pi_0(Z_{\widehat{G}}(\rho, e)) \\ \text{ such that } ses^{-1} = qe \end{cases} \end{cases}_{\widehat{G}\text{-conj}} \xrightarrow{\simeq} \{ \text{irred } \mathcal{H}_{\operatorname{aff}}\text{-modules} \}_{/\simeq}$$

Remark 10.8. The affine Hecke algebra $\mathcal{H}(I, G(K))$ has a presentation in which q becomes a variable. The above conjecture can either be understood with fixed $q = \#|\mathbb{O}_K/\mathfrak{m}_K|$ or with q as a variable, in which case q is also a variable on the left hand side.

Recall that the unramified LLC followed from the Satake isomorphism:

$$\mathcal{H}_{\rm sph} = \mathcal{H}(G(\mathbb{O}_K), G(K)) \xleftarrow{\simeq} R(\widehat{G}) = \mathcal{O}\left(\stackrel{\text{semisimple conj.}}{\operatorname{classes in } \widehat{G}} \right) \xleftarrow{\simeq} K^0(\mathrm{pt}/\widehat{G}) = K^{\widehat{G}}(\mathrm{pt})$$

"constructible" "coherent"

Similarly, the TRUM case of the LLC (which reduces to the Deligne–Langlands conjecture) follows from the Kazhdan–Lusztig isomorphism:

$$\mathcal{H}_{\mathrm{aff}} \xleftarrow{\simeq} K^{\widehat{G} \times \mathbb{C}^{\times}}(St)$$

"constructible" "coherent"

Indeed, if $\pi : \widetilde{\mathcal{N}} = T^*\mathcal{B} \to \mathcal{N}$ is the Springer resolution, $\mathcal{B}_e = \pi^{-1}(e)$ is the Springer fibre of a nilpotent element $e \in \mathcal{N}$, and St is the Steinberg variety, the Kazhdan–Lusztig isomorphism (which is not easy to establish!) implies that there is an action of the affine Hecke algebra \mathcal{H}_{aff} on $K^{Z_{\widehat{G} \times \mathbb{C}^{\times}}(e)}(\mathcal{B}_e)$. Here we can see that

$$Z_{\widehat{G} \times \mathbb{C}^{\times}}(e) = \{ (g, c) \mid c \cdot geg^{-1} = e \} = \{ (g, c) \mid geg^{-1} = c^{-1} \cdot e \}$$

looks very close to the parameters in the Deligne–Langlands conjecture. This action shows us that the K-theory of Springer fibres provides all simple \mathcal{H}_{aff} -modules, thus proving the Deligne–Langlands conjecture.

10.4 Geometric Satake equivalence

There is a geometric upgrade of the Satake isomorphism which has proven to be a major tool in geometric representation theory. Set K = k((t)), so $\mathbb{O}_K = k[[t]]$, where $k = \mathbb{C}$ or \mathbb{F}_q . Then

$$\mathcal{H}(G(\mathbb{O}_K), G(K)) = \frac{G(\mathbb{O}_K) \text{-invariant functions on the}}{\text{"affine Grassmanian"}} \mathcal{G}r_G := G(K)/G(\mathbb{O}_K).$$

The geometric Satake equivalence is the equivalence of categories:

$$(\operatorname{Perv}_{G(\mathbb{O}_K)}(\mathcal{G}r,\mathbb{C}),*) \xrightarrow{\simeq} (\operatorname{Rep} \widehat{G}_{\mathbb{C}},\otimes)$$

"constructible" "coherent"

This equivalence was key in recent work by V. Laffourgés giving an "automorphic to Galois" correspondence for global function fields.

10.5 Bezrukavnikov's equivalence

There is also a geometric upgrade of the Kazhdan–Lusztig isomorphism. With K = k((t)) as above, the affine Hecke algebra is

 $\mathcal{H}_{\text{aff}} =$ Iwahori-invariant functions on G(K)/I.

Here G(K)/I is the set of k-points of the "affine flag variety" $\mathcal{F}l_G$. Roughly, Bezrukavnikov's equivalence is an equivalence of categories

$$(D^b_{I \times I}(\mathcal{F}l_G), *) \xrightarrow{\simeq} (D^b \operatorname{Coh}^{G \times \mathbb{C}^{\times}}(St), *)$$

"constructible" "coherent"

Remark 10.9. This is a bit of a lie! It would take several more lecture to precisely describe the categories on each side of this equivalence.

This equivalence has many applications in geometric representation theory. For example, a mod p version of this equivalence would imply everything that we know about modular representations of algebraic groups!

Lecture 11: Review of the first semester

11.1 The local Langlands correspondence

Recall our setup from last semester. Let K be a local field (i.e. K is a finite extension of \mathbb{Q}_p or $K = \mathbb{F}_q((t))$) with ring of integers \mathbb{O} and residue field k:

$$K \supset \mathbb{O} \twoheadrightarrow k$$

Nothing is lost by just thinking in terms of the example

$$\mathbb{Q}_p \supset \mathbb{Z}_p \twoheadrightarrow \mathbb{F}_p$$

Last semester, we worked up to stating the local Langlands correspondence.

Local Langlands correspondence for GL_n : There exists a canonical bijection

$$\left\{\begin{array}{l} \text{irred. smooth admiss.} \\ \text{reps of } \operatorname{GL}_n(K) \text{ on} \\ \mathbb{C}\text{-vector spaces} \end{array}\right\}_{\simeq} \xrightarrow{1:1} \left\{\begin{array}{l} F\text{-semisimple} \\ n\text{-dimensional} \\ \text{Weil-Deligne reps} \end{array}\right\}_{\simeq}$$

It has been a while sine the last lecture, so let us remind ourselves what all of these words mean. The representations on the left hand side of this correspondence are *irreducible* representations which are usually infinite-dimensional. The adjective **smooth** means that every vector has an open stabilizer, and the adjective **admissible** means that for any open subgroup $U \subset \operatorname{GL}_n(K)$, V^U is finite-dimensional. We consider the set of such representations up to equivalence.

The objects on the right hand side are *n*-dimensional representations (where *n* is the same *n* appearing in GL_n on the left) of the Weil group attached to the field *K*, along with some extra data. Recall that the **Weil group** is a subgroup of the absolute Galois group $Gal(\overline{K}/K)$ defined by the fact that it fits into the following diagram.

Here $I_{\overline{K}/K}$ is the inertia subgroup. A Weil-Deligne representation is a triple (V, ρ, N) , where

- 1. V is an *n*-dimensional \mathbb{C} -vector space,
- 2. $\rho: W_K \to \operatorname{GL}(V)$ is a continuous representation, and
- 3. N is a nilpotent endomorphism of V such that

$$\rho(x)N\rho(x)^{-1} = |x|N$$

for all $x \in W_K$.

Here $|\cdot|$ is the canonical norm on W_k , which is uniquely determined by the property that |Frob| = |k|. (See Section 8.2 for a refresher on why this exists.) A Weil-Deligne representation is *F*-semisimple if any lift of Frob acts semisimply.

Remark 11.1. For the experts: Weil-Deligne representations are distilled out of continuous representations $W_K \to \operatorname{GL}_n(\overline{\mathbb{Q}}_\ell)$ via "log of monodromy". Thus one can think of the right hand side as secretly being genuine representations of a group. It is phrased in this way to remove the auxilary choice of a prime number ℓ .

In general, if we replace $GL_n(K)$ with any split reductive group G(K) (e.g. SL_n , Sp_{2n} , etc.), then the LLC changes as follows.

Local Langlands correspondence for split reductive groups: There exists a canonical finite-to-one map

$$\left\{\begin{array}{c} \text{irred. smooth admiss.} \\ \text{reps of } G(K) \text{ on} \\ \mathbb{C}\text{-vector spaces} \end{array}\right\}_{\simeq} \xrightarrow{\text{finite:1}} \left\{\begin{array}{c} F\text{-semisimple} \\ \text{Weil-Deligne reps} \\ \text{of } W_K \text{ in } G^{\vee}(\mathbb{C}) \end{array}\right\}_{\simeq}$$

In this setting, a Weil-Deligne representation of W_K in $G^{\vee}(\mathbb{C})$ is a pair (ρ, N) , where

1. $\rho: W_K \to G^{\vee}(\mathbb{C})$ is a continuous group homomorphism of the Weil group into the complex Langlands dual group of G, and

2.
$$N \in \operatorname{Lie} G^{\vee}(\mathbb{C})$$
 such that

$$\rho(x)N\rho(x)^{-1} = |x|N$$

for all $x \in W_K$.

Note that condition 2 forces $N \in G^{\vee}(\mathbb{C})$ to be a nilpotent element.

11.2 The unramified story

Last semester we ended the course by unpacking the simplest piece of the LLC, the case of *unramified representations*. Let us remind ourselves how this story went. By restricting each side of the correspondence above, we obtain a bijection

$$\left\{ \begin{array}{c} \text{irred. smooth admissible} \\ \text{unramified reps of } G(K) \\ (\text{i.e. reps of } G(K) \text{ that admit} \\ \text{a non-zero } G(\mathbb{O})\text{-fixed vector} \end{array} \right\}_{\simeq} \longleftrightarrow \left\{ \begin{array}{c} \text{unramified} \\ \text{Weil-Deligne reps;} \\ (\text{i.e. reps which factor} \\ W_K \to \mathbb{Z} \to G^{\vee}(\mathbb{C}) \\ \text{with } N = 0 \end{array} \right\}_{\simeq}$$

By Hecke algebra theory, representations of G(K) with a $G(\mathbb{O})$ -fixed vector are in bijection with irreducible modules for the *spherical Hecke algebra*,

$$\mathcal{H}_{\rm sph} = \mathcal{H}(G(K), G(\mathbb{O})) := (\operatorname{Fun}_{G(\mathbb{O}) \times G(\mathbb{O})}(G(K), \mathbb{C}), *)$$

The functions in this definition are continuous with compact support. On the other hand, unramified Weil-Deligne representations are in bijection with the set of conjugacy classes of semisimple elements in $G^{\vee}(\mathbb{C})$. Recall the Satake isomorphism.

Theorem 11.2. (Satake isomorphism) There exists a canonical isomorphism

$$\mathcal{H}_{\mathrm{sph}} \xrightarrow{\sim} [\operatorname{Rep} G^{\vee}] \otimes_{\mathbb{Z}} \mathbb{C}.$$

This theorem will be a big feature of the course this semester. Let $T^{\vee} \subset G^{\vee}$ be a maximal torus, and W the Weyl group of G^{\vee} . By highest weight theory we have

$$[\operatorname{Rep} G^{\vee}] \longleftrightarrow [\operatorname{Rep} T^{\vee}] = \mathbb{Z} [X^*(T^{\vee})]$$

$$\uparrow$$

$$\mathbb{Z} [X^*(T^{\vee})]^W$$

Hence

$$\operatorname{Rep} G^{\vee}] \otimes_{\mathbb{Z}} \mathbb{C} = \mathbb{C}[X^*(T^{\vee})]^W = \mathcal{O}(T^{\vee}/W) = \mathcal{O}(G_{ss}^{\vee}/\operatorname{conj}).$$

The spherical Hecke algebra is commutative. Therefore,

{irred. \mathcal{H}_{sph} -modules} $\leftrightarrow \{\chi : [\operatorname{Rep} G^{\vee}] \otimes_{\mathbb{Z}} \mathbb{C} \to \mathbb{C}\} = \{\text{semisimple conjugacy classes in } G^{\vee}(\mathbb{C})\}$

The first bijection follows from the Satake isomorphism and the commutativity of \mathcal{H}_{sph} , and the second equality is the nullstellensatz. So the Satake isomorphism implies unramified LLC!

Remark 11.3. The goal of the next two Informal Friday Seminars (September 6 and 13) will be to explain the **geometric Satake equivalence**, which is a categorification of Theorem 11.2:


 $(\operatorname{Perv}_{G(\mathbb{O})\times G(\mathbb{O})}(G(K),\mathbb{C}),*) \stackrel{\sim}{\leftrightarrow} (\operatorname{Rep} G_{\mathbb{C}}^{\vee},\otimes)$

11.3 The tamely ramified with unipotent monodromy (TRUM) story

This semester we will focus on the next simplest piece of the LLC, the case of *tamely* ramified representations with unipotent monodromy (*TRUM*). In this setting, the LLC gives us a finite-to-one map:

$$\left\{\begin{array}{c} \text{TRUM reps of } G(K) \\ \text{(i.e. those which admit} \\ \text{a non-zero } Iw\text{-fixed vector} \end{array}\right\}_{\simeq} \xrightarrow{\text{finite:1}} \left\{\begin{array}{c} \text{TRUM Weil-Deligne reps} \\ \text{(i.e. reps which factor} \\ W_K \twoheadrightarrow \mathbb{Z} \to G^{\vee}(\mathbb{C}) \\ \text{with } N \text{ arbitrary} \end{array}\right\}_{\simeq}$$

Here $Iw \subset G(K)$ is the *Iwahori subgroup*, which sits in the group G(K) in the following way:



By Hecke algebra theory, TRUM representations of G(K) are in bijection with irreducible representations of the *Iwahori-Matsumoto Hecke algebra* $\mathcal{H}_{aff} = \mathcal{H}(G(K), I_w)$. On the other hand, TRUM Weil-Deligne representations are parameterized by the set

$$\{(s,N)\in G^{\vee}(\mathbb{C})\times\mathcal{N}\mid sNs^{-1}=qN\}/_{\operatorname{conj}}.$$

Here $s \in G^{\vee}(\mathbb{C})$ is the semisimple image of Frobenius, $\mathcal{N} \subset \text{Lie} G^{\vee}$ is the nilpotent cone, and q = |k|. So the LLC predicts a parameterisation of irreducible \mathcal{H}_{aff} -modules. This prediction is the *Deligne-Langland conjecture*, and it served as an important early test case of the Langland's philosophy.

Goals for the next few weeks:

- 1. Discuss the Iwahori-Matsumoto Hecke algebra is some detail.
- 2. Discuss Kazhdan–Lusztig's realisation of the affine Hecke algebra H via equivariant K-theory:

$$H \xrightarrow{\sim} K^{G^{\vee} \times \mathbb{C}^{\times}}$$
 (Steinberg)

3. Deduce the Deligne–Langlands conjecture⁶.

Then we will pass to categorifications!

11.4 Affine Weyl groups and affine Hecke algebras

Let $(X \supset R, X^{\vee} \supset R^{\vee})$ be a root datum.

Example 11.4. 1. SL₂:
$$X = \mathbb{Z} \supset R = \{\pm 2\}, X^{\vee} = \mathbb{Z} \supset R^{\vee} = \{\pm 1\}$$

2. PGL₂: $X = \mathbb{Z} \supset R = \{\pm 1\}, X^{\vee} = \mathbb{Z} \supset R^{\vee} = \{\pm 2\}$

We see from this example that SL_2 and PGL_2 are interchanged by swapping roots and coroots, so they are Langlands dual groups.

Let W_f be the finite Weyl group associated to this root datum. Then W_f acts on both X and X^{\vee} . Assume that our root datum $(X \supset R, X^{\vee} \supset R^{\vee})$ is *adjoint*; i.e., $X = \mathbb{Z}R$. (This is the "most complicated case".)

Definition 11.5. The extended affine Weyl group⁷ is

$$W_{\text{ext}} = \mathbb{Z}X^{\vee} \rtimes W_f.$$

 $^{^{6}}$ Actually, we will not end up deducing the Deligne-Langlands conjecture. We will simply promise the reader that the Deligne-Langlands conjecture is not a long walk from where we get to.

⁷This definition is the definition according to Iwahori-Matsumoto and Bourbaki, but we warn the reader that it is *not* consistent across all sources! For example, this is not the convention employed in Chriss-Ginzburg.

The group W_{ext} acts on $X_{\mathbb{R}}^{\vee} := X^{\vee} \otimes_{\mathbb{Z}} \mathbb{R}$ by "affine transformations;" that is, $w \in W_f$ acts as usual, $w(\lambda) = w(\lambda)$, and for $\gamma \in X^{\vee}$, $t_{\gamma} \in \mathbb{Z}X^{\vee}$ acts by

$$t_{\gamma}(\lambda) = \lambda + \gamma.$$

To understand W_{ext} , we first consider the affine Weyl group $W = \mathbb{Z}R^{\vee} \rtimes W_f \subset W_{\text{ext}}$. For $\alpha \in \mathbb{R}, m \in \mathbb{Z}, \lambda \in X_{\mathbb{R}}^{\vee}$, define

$$s_{\alpha,m}(\lambda) := \lambda - \langle \lambda, \alpha \rangle \alpha^{\vee} + m \alpha^{\vee}.$$

Clearly,

$$s_{\alpha,m} = t_{m\alpha^{\vee}} \circ s_{\alpha}$$

so $s_{\alpha,m} \in W$, and $t_{m\alpha^{\vee}} \in \langle s_{\alpha,m} \mid \alpha \in R, m \in \mathbb{Z} \rangle$. We conclude that

$$W = \langle s_{\alpha,m} \mid \alpha \in R, m \in \mathbb{Z} \rangle$$

is an affine reflection group generated by reflections $s_{\alpha,m}$ through the hyperplanes

$$H_{\alpha,m} = \{\lambda \in X_{\mathbb{R}}^{\vee} \mid \langle \lambda, \alpha \rangle = m\}.$$

We call the set $\{H_{\alpha,m}\}$ the set of *reflecting hyperplanes*. Denote by \mathcal{A} the corresponding set of *alcoves*; that is, the closures of connected components of

$$X_{\mathbb{R}}^{\vee} \setminus \bigcup_{\substack{\alpha \in R \\ m \in \mathbb{Z}}} H_{\alpha,m}.$$

Fix a set of positive roots $R_+ \subset R$, and let

$$A_0 = \{ \lambda \in X_{\mathbb{R}}^{\vee} \mid 0 \le \langle \lambda, \alpha \rangle \le 1 \text{ for all } \alpha \in R_+ \} \subset \mathcal{A}$$

be the *fundamental alcove*. The general (very beautiful) theory of reflection groups gives:

- 1. W is a Coxeter group with Coxeter generators $S := \{\text{reflections in the walls of } A_0\}$.
- 2. The length function may be described by


$$\ell(w) = \#\{\text{reflecting hyperplanes separating } A_0^{int} \text{ and } w A_0^{int}\}.$$

3. A_0 is a fundamental domain for the W-action on $X_{\mathbb{R}}^{\vee}$.

So we have an identification

$$W \to \mathcal{A}$$
$$w \mapsto wA_0$$

Example 11.6. $C_2 = B_2$



Now we move on to W_{ext} . There is an action of W_{ext} on \mathcal{A} because

$$t_{\gamma} \cdot H_{\alpha,m} = H_{\alpha,m+\langle \gamma,\alpha \rangle}.$$

Define

$$\ell: W_{\text{ext}} \to \mathbb{Z}_{\geq 0}$$

$$w \mapsto \ell(w) := \#\{\text{hyperplanes separating } A_0^{int} \text{ and } w A_0^{int}\}$$

Define the *length zero* elements of W_{ext} to be

$$\Omega := \ell^{-1}(0).$$

Lemma 11.7. $W_{\text{ext}} = \Omega \ltimes W$.

Proof. Step 1: $W \subset W_{\text{ext}}$ is normal.

Let $\gamma \in X^{\vee}, \lambda \in X_{\mathbb{R}}^{\vee}$. Then

$$t_{\gamma}s_{\alpha,m}t_{\gamma}^{-1}(\lambda) = t_{\gamma}(s_{\alpha,m}(\lambda - \gamma))$$

= $\lambda - \gamma - \langle \lambda - \gamma, \alpha \rangle \alpha^{\vee} + m\alpha^{\vee} + \gamma$
= $\lambda - \langle \lambda, \alpha \rangle \alpha^{\vee} + (\langle \gamma, \alpha \rangle + m)\alpha^{\vee}$
= $s_{\alpha,\langle \gamma,\alpha \rangle + m}(\lambda).$

Step 2: $W_{\text{ext}} = W \cdot \Omega$.

Let $w \in W_{\text{ext}}$. Then $wA_0 \in \mathcal{A}$, so by 3. above, there exists $y \in W$ such that $ywA_0 = A_0$. Hence $yw = \omega$ for $\omega \in \Omega$, and $w = y^{-1}\omega$.

Step 3: $W \cap \Omega = \{id\}.$

Any $w \in W \cap \Omega$ is length zero, so w = id by 2. above and the fact that W is a Coxeter group.

Lemma 11.8. (Iwahori-Matsumoto) $w \in W_f, \lambda \in X^{\vee}$,

$$\ell(t_{\lambda}w) = \sum_{\substack{\alpha \in R^+ \\ w^{-1}(\alpha) > 0}} |\langle \lambda, \alpha \rangle| + \sum_{\substack{\alpha \in R^+ \\ w^{-1}(\alpha) < 0}} |\langle \lambda, \alpha \rangle - 1|.$$

Proof. Let $x = t_{\lambda} w$. Then

$$\ell(x) = \#\{\text{hyperplanes separating } A_0^{int} \text{ and } x A_0^{int}\} \\ = \sum_{\alpha \in R_+} \#\{m \mid H_{\alpha,m} \text{ separates } A_0^{int} \text{ and } x A_0^{int}\}.$$

Now chose a point $p \in A_0$, very close to zero. Then $\langle p, \alpha \rangle$ is small and positive, and

$$\ell(x) = \sum_{\alpha \in R_{+}} \#\{\text{integers between } \langle p, \alpha \rangle \text{ and } \langle xp = \lambda + wp, \alpha \rangle\}$$
$$= \sum_{\alpha \in R_{+}} \begin{cases} |\langle \lambda, \alpha \rangle| & \text{if } \langle wp, \alpha \rangle > 0\\ |\langle \lambda, \alpha \rangle - 1| & \text{if } \langle wp, \alpha \rangle < 0 \end{cases}$$
$$= \sum_{\substack{\alpha \in R_{+} \\ w^{-1}\alpha > 0}} |\langle \lambda, \alpha \rangle| + \sum_{\substack{\alpha \in R_{+} \\ w^{-1}\alpha < 0}} |\langle \lambda, \alpha \rangle - 1|.$$

Example 11.9. For PGL₂, $X^{\vee} = \mathbb{Z}\varpi_1$. Then

$$\ell(m\varpi_1) = |m|$$
, and $\ell(m\varpi_1 s) = |m-1|$.

Hence $\varpi_1 s$ is length zero, and $\Omega = \{id, t_{\varpi_1}s\}.$

Lecture 12: The Deligne-Langlands conjecture

12.1 The Iwahori–Matsumoto Hecke algebra

Recall our setup from last week. From an adjoint root datum $(X \supset R, X^{\vee} \supset R^{\vee})$, (i.e.; meaning $X = \mathbb{Z}R$) we construct

- the finite Weyl group W_f ,
- the affine Weyl group $W = \mathbb{Z}R^{\vee} \rtimes W_f$, and
- the extended affine Weyl group $W_{\text{ext}} = \mathbb{Z}X^{\vee} \rtimes W_f$.

We fix a set of positive roots $R^{\vee}_+ \subset R^{\vee}$, then obtain the fundamental alcove A_0 , and the corresponding set $S \subset W$ of Coxeter generators. We define a length function ℓ by

$$\ell: W_{\text{ext}} \to \mathbb{Z}$$
$$x \mapsto \# \left\{ \begin{array}{c} \text{reflecting hyperplanes} \\ \text{between } A_0^{int} \text{ and } x A_0^{int} \end{array} \right\}.$$

We denoted the length zero elements by $\Omega = \ell^{-1}(0)$, and showed that W_{ext} is a "quasi Coxeter group," meaning that

$$W_{\text{ext}} = \Omega \ltimes W,$$

W is a Coxeter group, and Ω acts on W via automorphisms of the Coxeter system.

Example 12.1. For PGL_2 ,



The affine Weyl group and extended affine Weyl group are

$$W = \langle s, t \mid s^2 = t^2 = id \rangle \subset W_{\text{ext}} = \langle s, t, \tau = \varpi s \mid s^2 = t^2 = \tau^2 = id, \tau s = t\tau \rangle.$$

We can use the extended affine Weyl group to define a Hecke algebra.

Definition 12.2. The *Iwahori–Matsumoto Hecke algebra* \mathcal{H}_{ext} is the $\mathbb{Z}[v^{\pm 1}]$ -algebra with basis $\{H_x \mid x \in W_{ext}\}$ and multiplication

$$H_x H_y = H_{xy} \text{ if } \ell(xy) = \ell(x) + \ell(y) H_s^2 = H_{id} + (v^{-1} - v)H_s \text{ if } s \in S.$$

Define $\mathcal{H} = \langle H_x \mid x \in W \rangle \subset \mathcal{H}_{\text{ext}}.$

Remark 12.3. 1. $\mathcal{H} \subset \mathcal{H}_{ext}$ is a subalgebra.

2. For all $x \in W_{\text{ext}}$, H_x is invertible.

3. If $\tau \in \Omega$, then $\ell(x\tau) = \ell(x) = \ell(\tau x)$. Hence

$$\mathcal{H}_{ext} = \Omega \ltimes \mathcal{H}$$

Question: Where did the loop presentation $W_{\text{ext}} = \mathbb{Z}X^{\vee} \rtimes W_f$ go?

For $\lambda \in X^{\vee}$, write $\lambda = \gamma - \gamma'$ with $\gamma, \gamma' \in X_{+}^{\vee}$. Define

$$H_{\lambda} := H_{t_{\gamma}} H_{t_{\gamma'}}^{-1}.$$

Note that $H_{t_{\lambda}} \neq H_{\lambda}$ in general! For example, if $\lambda \in X_{+}^{\vee}$, then $H_{\lambda} = H_{t_{\lambda}}$, but if $\lambda \in -X_{+}^{\vee}$, then $H_{\lambda} = H_{t_{-\lambda}}^{-1}$.

Why is this well-defined? Assume that $\lambda = \gamma - \gamma' = \mu - \mu'$ for $\gamma, \gamma', \mu, \mu' \in X_+^{\vee}$. To show that H_{λ} is well-defined, we need to show that

$$H_{t_{\gamma}}H_{t_{\gamma'}}^{-1} = H_{t_{\mu}}H_{t_{\mu'}}^{-1}.$$

This is equivalent to showing that for $\zeta \in X^{\vee}$,

$$H_{t_{\gamma}}H_{t_{\gamma'}}^{-1}H_{t_{\zeta}} = H_{t_{\mu}}H_{t_{\mu'}}^{-1}H_{t_{\zeta}}.$$

If we choose $\zeta \in X_+^{\vee}$ very dominant, then $\zeta - \mu'$ is dominant and

$$H_{t_{\zeta-\mu'}}H_{t_{\mu'}} = H_{t_{\zeta-\mu'}t_{\mu'}} = H_{t_{\zeta}} = H_{t_{\mu'}t_{\zeta-\mu'}} = H_{t_{\mu'}}H_{t_{\zeta-\mu'}}$$

The first and fourth equalities follow from the fact that the lengths of $t_{\zeta-\mu'}$ and $t_{\mu'}$ add by the Iwahori–Matsumoto lemma (Lemma 11.8). Hence,

$$H_{t_{\gamma}}H_{t_{\gamma'}}^{-1}H_{t_{\zeta}} = H_{t_{\gamma}}H_{t_{\zeta-\gamma'}} = H_{t_{\gamma+\zeta-\gamma'}} = H_{t_{\mu+\zeta-\mu'}} = H_{t_{\mu}}H_{t_{\zeta-\mu'}} = H_{t_{\mu}}H_{t_{\mu'}}^{-1}H_{t_{\zeta}}$$

so H_{λ} is well-defined.

The upshot: When studying representations of an algebra, it is useful to have a large commutative subalgebra. This is what we have just accomplished for the Iwahori–Matsumoto Hecke algebra: the map $\lambda \mapsto H_{\lambda}$ determines an embedding

$$\mathbb{Z}[v^{\pm 1}][X^{\vee}] \hookrightarrow \mathcal{H}_{\mathrm{ext}}.$$

We can use this commutative subalgebra to describe the center of \mathcal{H}_{ext} .

Theorem 12.4. (Bernstein) For any $\lambda \in X_+^{\vee}$, define

$$z_{\lambda} := \sum_{\mu \in W_f \lambda} H_{\mu}$$

Then the center $Z(\mathcal{H}_{ext})$ of \mathcal{H}_{ext} is a free $\mathbb{Z}[v^{\pm 1}]$ -module with basis $\{z_{\lambda} \mid \lambda \in X_{+}^{\vee}\}$, and

$$Z(\mathcal{H}_{\text{ext}}) = \left(\mathbb{Z}[v^{\pm 1}][X^{\vee}]\right)^{W_f}.$$

Now we can state the Bernstein presentation of \mathcal{H}_{ext} .

Theorem 12.5. (Bernstein presentation) The Iwahori–Matsumoto Hecke algebra admits the following presentation.

- 1. $\langle H_s \mid s \in S_f \rangle$ generate a finite Hecke algebra ("finite part").
- 2. $H_{\lambda}H_{\gamma} = H_{\lambda+\gamma}$ for all $\lambda, \gamma \in X^{\vee}$ ("lattice part").
- 3. For $\lambda \in X^{\vee}, s_{\alpha} \in S_f$,

$$H_{s_{\alpha}}H_{s_{\alpha}(\lambda)} - H_{\lambda}H_{s_{\alpha}} = (v - v^{-1})\left(\frac{H_{\lambda} - H_{s_{\alpha}(\lambda)}}{1 - H_{-\alpha}}\right)$$
$$= (v - v^{-1})(H_{\lambda} + H_{\lambda - \alpha} + \dots + H_{s_{\alpha}(\lambda) + \alpha})$$

In other words, we have

$$\mathcal{H}_{\text{ext}} \simeq \mathbb{Z}[v^{\pm 1}][X^{\vee}] \otimes \mathcal{H}_f$$

We will check the relations for PGL₂. First note that if 3. holds for λ and γ , then it holds for $\lambda + \gamma$:

$$\begin{split} H_{s_{\alpha}}H_{s_{\alpha}(\lambda)+s_{\alpha}(\gamma)} &= (v-v^{-1})\left(\frac{H_{\lambda}-H_{s_{\alpha}(\lambda)}}{1-H_{-\alpha}}\right)H_{s_{\alpha}(\gamma)} + H_{\lambda}H_{s_{\alpha}}H_{s_{\alpha}(\gamma)} \\ &= (v-v^{-1})\left(\frac{H_{\lambda+s_{\alpha}(\gamma)}-H_{s_{\alpha}(\lambda)+s_{\alpha}(\gamma)}+H_{\lambda+\gamma}-H_{\lambda-s_{\alpha}(\gamma)}}{1-H_{-\alpha}}\right) + H_{\lambda+\gamma}H_{s_{\alpha}} \\ &= (v-v^{-1})\left(\frac{H_{\lambda+\gamma}-H_{s_{\alpha}(\lambda+\gamma)}}{1-H_{-\alpha}}\right) + H_{\lambda+\gamma}H_{s_{\alpha}}. \end{split}$$

In particular, if 3. is true for λ , then it is true for $-\lambda$.

For PGL₂: We will check 3. for $\lambda = \varpi$. We have that $\langle \varpi, \alpha \rangle = 1$, $\tau = t_{\varpi} s_{\alpha}$, so $t_{\varpi} = \tau s_{\alpha}$, and $H_{\varpi} = H_{\tau} H_{s_{\alpha}}$. We compute

$$H_{s_{\alpha}}H_{-\varpi} - H_{\varpi}H_{s_{\alpha}}^{-1} = H_{s_{\alpha}}H_{s_{\alpha}}^{-1}H_{\tau} - H_{\tau}H_{s_{\alpha}}H_{s_{\alpha}}$$

= $H_{\tau} - H_{\tau}(1 + (v^{-1} - v)H_{s_{\alpha}})$
= $(v - v^{-1})H_{\tau}H_{s_{\alpha}}$
= $(v - v^{-1})H_{\varpi}.$

12.2 The Deligne–Langlands conjecture

Now we return to the LLC. Let K be a local field with ring of integers \mathbb{O} and residue field k. Define q := |k|. Let G/K be a split reductive group, and $(X \supset R, X^{\vee} \supset R^{\vee})$ the corresponding root datum.

Recall that the Deligne–Langlands conjecture tells us that we should expect the following relationships:

$$\left\{ \begin{array}{c} \text{TRUM reps} \\ \text{of } G(K) \end{array} \right\}_{\simeq} \xrightarrow{finite:1} \left\{ \begin{array}{c} \text{TRUM Weil-} \\ \text{Deligne reps} \end{array} \right\}_{\simeq} \\ & & & \uparrow^{\simeq} \\ & & & \uparrow^{\simeq} \\ \left\{ \begin{array}{c} \text{irred } \mathcal{H}(G(K), Iw)\text{-} \\ \text{modules} \end{array} \right\}_{\simeq} & \left\{ \begin{array}{c} (s, x) \in G_{\mathbb{C}}^{\vee} \times \mathfrak{g}_{\mathbb{C}}^{\vee} \\ \text{s.t. } s \text{ is ss, } x \text{ nilp} \\ \text{and } sxs^{-1} = qx \end{array} \right\}_{/conj}$$

The following theorem relates the extended affine Hecke algebra of Iwahori–Matsumoto to this story.

Theorem 12.6. (Iwahori–Matasumoto)

- 1. $G(K) = \bigsqcup_{w \in W_{\text{ext}}} Iw \cdot w \cdot Iw$ ("Bruhat decomposition")
- 2. There is an isomorphism of algebras

$$\mathcal{H}_{\mathrm{ext}} \otimes_{\mathbb{Z}[v^{\pm 1}]} \mathbb{C} \xrightarrow{\sim} \mathcal{H}(G(K), Iw)$$

where \mathbb{C} is a $\mathbb{Z}[v^{\pm 1}]$ -algebra via $v \mapsto (\sqrt{q})^{-1} \in \mathbb{R}_+ \subset \mathbb{C}$.

Moreover, under 2., H_x is mapped to the indicator function on $Iw \cdot x \cdot Iw$, up to a scalar.

Example 12.7. Let $G = GL_n$, and fix a uniformizer $\pi \in \mathbb{O}$. Then

$$W_{\text{ext}} = \left\langle \begin{array}{c} \text{permutation} \\ \text{matrices} \\ \text{"finite part"} \end{array} \right\rangle \ltimes \left\langle \left(\begin{array}{c} \pi^{\lambda_1} & \cdots & 0 \\ 0 & \ddots & 0 \\ 0 & \cdots & \pi^{\lambda_n} \end{array} \right) \right\rangle$$

By Theorem 12.6, we can understand TRUM representations of G(K) by studying irreducible \mathcal{H}_{ext} -modules. Denote by $Z := Z(\mathcal{H}_{\text{ext}}) = (\mathbb{Z}[v^{\pm 1}][X^{\vee}])^{W_f}$. By Quillen's Lemma (which is an infinite-dimensional version of Schur's Lemma), Z acts by scalars on any irreducible \mathcal{H}_{ext} -module. The Bernstein presentation tells us that \mathcal{H}_{ext} is finite over $R := \mathbb{Z}[v^{\pm 1}][X^{\vee}]$. Since R is also finite over R^{W_f} , we conclude that any irreducible \mathcal{H}_{ext} -module is finite-dimensional, and, in fact, is of dimension $\leq |W_f|^2$.

Hence, the Deligne–Langlands conjecture predicts the following relationships.

$$\begin{cases} \operatorname{irreps}/\mathbb{C} \\ \operatorname{of} \mathcal{H}_{\operatorname{ext}} \end{cases} \xrightarrow{finite:1} \begin{cases} (s,x) \in G_{\mathbb{C}}^{\vee} \times \mathfrak{g}_{\mathbb{C}}^{\vee} \\ \operatorname{s.t.} s \text{ is ss, } x \text{ nilp} \\ \operatorname{and} sxs^{-1} = qx \end{cases} _{/\simeq}$$

$$\downarrow_{\operatorname{central character}}$$

$$\{ \operatorname{irreps of} Z \}$$

$$\downarrow_{=}$$

$$\{ \operatorname{pairs} (s',v) \in G_{\mathbb{C}}^{\vee} \times \mathbb{C}^{\times} \}_{/G_{\mathbb{C}}^{\vee} \operatorname{conj}} \xrightarrow{-\cdots \rightarrow} \{ (s,x) \in G_{\mathbb{C}}^{\vee} \times \mathbb{C}^{\times} \}_{/G_{\mathbb{C}}^{\vee} \operatorname{conj}} \end{cases}$$

The dashed arrow should match $v^{-1} \leftrightarrow \sqrt{q}$.

Remark 12.8. 1. We are no longer forced to take q = |k|.

2. In the diagram above, we are repeatedly using the fact from last semester that

{characters $\chi : \mathbb{Z}[X^{\vee}]^{W_f} \to \mathbb{C}$ } \leftrightarrow {semisimple elts of G up to conjugacy}.

An easy and interesting case: Consider the case when s = id and q = 1. In this case, the right side of the dashed arrow is

 ${x \in \mathfrak{g}_{\mathbb{C}}^{\vee} \text{ nilpotent}}_{/G_{\mathbb{C}}^{\vee} \text{ conjugacy}} =$ "nilpotent orbits".

This provides a hint that we should not expect to have a good algebraic grip on the problem of understanding the irreducible representations of \mathcal{H}_{ext} , because nilpotent orbits are complicated and not combinatorial in general. For the next lecture and a half, we will dive into this geometry.

12.3 Geometric setting

For notational convience, we will temporarily swap $G \leftrightarrow G^{\vee}$ in this section. Let G/\mathbb{C} be a complex reductive group, and $\mathcal{N} \subset \text{Lie } G =: \mathfrak{g}$ the nilpotent cone.

Remark 12.9. For GL_n , it is tempting to define \mathcal{N} as the variety

$$\{x \in \mathfrak{gl}_n(\mathbb{C}) \mid x^n = 0\}.$$

However, this results in a non-reduced scheme, because the ideal corresponding to the equation $x^n = 0$ is not radical. A better definition is to consider

 $\{x \in \mathfrak{gl}_n(\mathbb{C}) \mid \text{ coefficients of the characteristic polynomial vanish}\}.$

This still captures what we know as a nilpotent matrix, but results in a better geometric object. (In particular, it is a reduced scheme.)

In general, we define the nilpotent cone as follows. Consider the adjoint quotient map

$$\mathfrak{g} \xrightarrow{q} \mathfrak{g}/G = \mathfrak{h}/W.$$

The equality $\mathfrak{g}/G = \mathfrak{h}/W$ follows from Chevalley's theorem. For \mathfrak{gl}_n , the map q is "take coefficients of the characteristic polynomial," so this captures what we wanted in Remark 12.9. We define

 $\mathcal{N} = q^{-1}(0).$

Fundamental facts about the nilpotent cone:

- 1. \mathcal{N} is irreducible, reduced, and normal.
- 2. G has finitely many orbits on \mathcal{N} , and all are even-dimensional (over \mathbb{C}).

Example 12.10. If $G = GL_n$, then

$$\left\{\begin{array}{c} \text{nilpotent} \\ \text{matrices} \end{array}\right\}_{/conj} \leftrightarrow \left\{\begin{array}{c} \text{Jordan} \\ \text{normal} \\ \text{form} \end{array}\right\} \leftrightarrow \left\{\begin{array}{c} \text{partitions} \\ \lambda \vdash n \end{array}\right\}$$

Moreover, if we denote by \mathcal{O}_{λ} the orbit corresponding to the partition λ , then $\mathcal{O}_{\mu} \subset \overline{\mathcal{O}_{\lambda}} \iff \mu \leq \lambda$ in dominance order.


We can examine these orbit relations explicitly for small n.

$$n = 2:$$


$$\mathcal{N} = \left\{ x = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \mid \operatorname{Tr} x = \det x = 0 \right\} = \left\{ x = \begin{pmatrix} a & b \\ c & -a \end{pmatrix} \mid \det x = -a^2 - bc = 0 \right\} \subset \mathbb{C}^2.$$

$$A \ \mathbb{P} \text{ picture of this is:}$$

A \mathbb{R} -picture of this is:



n = 3: For n = 3, the picture is more of a caricature.



Exercise 12.11. Try to do this for large n (perhaps 7 or 8) and see that this poset is rather ugly; in particular, it is not graded.

12.4 The Springer resolution

We can study these nilpotent orbits by using a resolution of singularities of the nilpotent cone. Let \mathcal{B} be the variety of Borel subalgebras in \mathfrak{g} . For \mathfrak{gl}_n , this is the variety of complete flags in \mathbb{C}^n .

Definition 12.12. The Springer resolution is the map

$$\begin{split} \tilde{\mathcal{N}} &= \{(x, \mathfrak{b}) \in \mathcal{N} \times \mathcal{B} \mid x \in \mathfrak{b}\} \simeq T^* \mathcal{B} \\ & \downarrow \\ & \mathcal{N} \end{split}$$


which sends $(x, \mathfrak{b}) \mapsto x$.

Next week we will study the Springer resolution more carefully, and show that it is proper, smooth, and a resolution of singularities. For GL_n ,


$$\widetilde{\mathcal{N}} = \{ (x, \{0\} = V_0 \subset V_1 \subset \cdots \vee V_n = \mathbb{C}^n) \mid x \text{ preserves flag} \iff xV^i \subset V^{i-1} \}.$$

Example 12.13. We examine GL_n for small n again.

n = 2: The variety of Borel subalgebras is $\mathcal{B} = \mathbb{P}^1 \mathbb{C} = \{\text{lines in } \mathbb{C}^2\}$, and the Springer resolution looks like:



n = 3: The variety of Borel subalgebras is $\mathcal{B} = \{(\ell, P) \mid \ell \subset P \subset \mathbb{C}^2\}$ and a caricature of the Springer resolution looks like:



We will explain in more detail how we arrived at this picture next week.

Lecture 13: Springer fibres and the Steinberg variety

13.1 The Springer resolution, continued

Today we are going to pick up where we left off last week and continue to discuss the geometry of nilpotent orbits. To begin, we will discuss in more detail our claim from last week that $\widetilde{N} \simeq T^* \mathcal{B}$.

Let G be an algebraic group and \mathfrak{g} its Lie algebra. Canonically, we can express the tangent bundle of G as

$$TG = G \times \mathfrak{g},$$

since the Lie algebra can be identified with $T_e(G)$. Hence if X is a homogeneous space for G (i.e. G acts on X transitively), then we have a surjection

$$TG = G \times \mathfrak{g} \to TX$$

and for $x \in X$, $T_x X = \mathfrak{g}/\operatorname{Lie}(\operatorname{stab}_G x)$. Similarly, there is a canonical identification of the cotangent bundle of G

$$T^*G = G \times \mathfrak{g}^*$$

and for a homogeneous space X,

$$T^*X \hookrightarrow T^*G.$$

Moreover, for $x \in X$, $T_x X = (\text{Lie}(\operatorname{stab}_G x))^{\perp} \subset \mathfrak{g}^*$.

Now assume that G is semisimple and let \mathcal{B} be the variety of Borel subalgeras of \mathfrak{g} . Once we choose a Borel subalgebra $B \subset G$, we can identify $\mathcal{B} \simeq G/B$ since $\operatorname{stab}_G \mathfrak{b} = B$. Last lecture we introduced the following space

$$\mathcal{N} = \{ (\mathfrak{b}, x) \in \mathcal{B} \times \mathcal{N} \mid x \in \mathfrak{b} \}.$$

Claim 13.1. There is a canonical isomorphism $\widetilde{\mathcal{N}} \simeq T^* \mathcal{B}$.

Proof. For a point $\mathfrak{b} \in \mathcal{B}$, the tangent space to \mathcal{B} at \mathfrak{b} is

$$T_{\mathfrak{b}}\mathcal{B} = \mathfrak{g}/(\operatorname{Lie}(\operatorname{stab}_{G}\mathfrak{b}) = \mathfrak{b}).$$

Hence,

$$T^*\mathcal{B} = \{(\mathfrak{b}, v) \in \mathcal{B} \times \mathfrak{g}^* \mid v \in \mathfrak{b}^\perp\}.$$

Recall the Killing form $\kappa : \mathfrak{g} \times \mathfrak{g} \to \mathbb{C}$, $\kappa(x, y) = tr(\operatorname{ad} x \operatorname{ad} y)$. This is a symmetric, nondegenerate bilinear form. Fix a Cartan subalgebra $\mathfrak{h} \subset \mathfrak{b}$, and we obtain a triangular decomposition $\mathfrak{g} = \mathfrak{n}_- \oplus \mathfrak{h} \oplus \mathfrak{n}_+$, with $\mathfrak{b} = \mathfrak{h} \oplus \mathfrak{n}_+$. The restriction $\kappa|_{\mathfrak{h} \times \mathfrak{h}}$ is nondegenerate, and κ gives a nondegenerate pairing

$$\kappa: \mathfrak{n}_{-} \times \mathfrak{n}_{+} \to \mathbb{C}.$$

Hence under the identification $\mathfrak{g} \simeq \mathfrak{g}^*$ via the Killing form, $\mathfrak{b}^{\perp} \subset \mathfrak{g}^*$ corresponds to $\mathfrak{n}_+ \subset \mathfrak{g}$. We conclude that

$$T^*\mathcal{B} = \{(\mathfrak{b}, x) \in \mathcal{B} \times \mathfrak{g} \mid x \in \mathfrak{b} \text{ is nilpotent} \}$$
$$= \{(\mathfrak{b}, x) \in \mathcal{B} \times \mathcal{N} \mid x \in \mathfrak{b} \}$$
$$= \widetilde{\mathcal{N}}.$$

13.2 Many examples of Springer fibres

"The richness of Springer fibres cannot be underestimated."

Recall the Springer resolution

$$\widetilde{\mathcal{N}} = \{(x, \mathfrak{b}) \in \mathcal{N} imes \mathcal{B} \mid x \in \mathfrak{b}\} \simeq T^* \mathcal{B}$$

$$\downarrow^{\pi_s}_{\mathcal{N}}$$

Definition 13.2. Let $x \in \mathcal{O}_{\lambda} \subset \mathcal{N}$ be a point in a *G*-orbit. The associated Springer fibre F_{λ} is

$$F_{\lambda} := \pi_s^{-1}(x).$$

By equivariance, this is independent of the choice of $x \in \mathcal{O}_{\lambda}$, up to isomorphism.


Let $G = SL_n$. This section is devoted to studying Springer fibres for n = 2, 3, 4. Recall that for $G = SL_n$,

$$\widetilde{\mathcal{N}} = \{ (F^{\cdot}, x) \in \frac{\text{complete}}{\text{flags in } \mathbb{C}^n} \times \mathcal{N} \mid x \text{ preserves } F^{\cdot} \} = \{ (F^{\cdot}, x) \mid xF^i \subset F^{i-1} \}.$$

Example n = 2**:** The nilpotent cone is the quadric cone

$$\mathcal{N} = \left\{ \begin{pmatrix} a & b \\ c & -a \end{pmatrix} \in \mathfrak{sl}_2 \mid -a^2 - bc = 0 \right\} \subset \mathbb{C}^3.$$


A \mathbb{R} -picture:



A table of orbits:

λ		
$\dim \mathcal{O}_{\lambda}$	2	0
$\operatorname{codim} \mathcal{O}_{\lambda}$	0	2
Springer fibre	pt	$\mathbb{P}^1\mathbb{C}$
dim fibre	0	1

Example n = 3: A caricature:



We'd like to construct a table as we did in the previous example, but to do so, we need to determine the dimension⁸ of the nilpotent orbits. This is easy for the regular nilpotent orbit and the zero orbit. But what about the other orbit \mathcal{O}_{λ} ? Well we know that the matrix

$$e = \begin{pmatrix} 0 & 1 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}$$

is in the orbit, and we observe that

$$\text{Lie}(\text{stab}_{\text{SL}_3} e) = \text{stab}_{\text{Lie} \text{SL}_3} e = \ker \operatorname{ad} e.$$

A computation shows that

$$\ker \operatorname{ad} e = \left\{ \begin{pmatrix} a & b & c \\ 0 & a & 0 \\ 0 & d & -2a \end{pmatrix} \mid a, b, c, d \in \mathbb{C} \right\}$$

⁸If you'd like to understand some variety, it is a good idea to know its dimension!

is 4-dimensional, hence the orbit \mathcal{O}_{λ} is (8-4=4)-dimensional. We also need to determine the Springer fibre corresponding to this orbit. We have that

$$F_{\lambda} = \{ (L \subset P \subset \mathbb{C}^3) \mid e\mathbb{C}^3 \subset P, eP \subset L, \text{ and } eL = 0 \},\$$

but what does this look like? Well we know that

$$0 \subset \operatorname{im} e \subset \ker e \subset \mathbb{C}^3,$$

and the condition that a flag in F_{λ} must satisfy eL = 0 implies that $L \subset \ker e$. We see that there are two possibilities for flags in F_{λ} :

- $\{L = \operatorname{im} e, \text{ free choice of } P, \text{ as long as } L \subset P\} \simeq \mathbb{P}^1 \mathbb{C}$
- $\{P = \ker e, \text{ free choice of } L, \text{ as long as } L \subset P\} \simeq \mathbb{P}^1 \mathbb{C}$

These two cases intersect when L = ime and P = ker e, which is a single point, so our Springer fibre looks like two $\mathbb{P}^1\mathbb{C}$'s joined at a point. With this, we can complete our table of orbits:

λ			
$\dim \mathcal{O}_{\lambda}$	6	4	0
$\operatorname{codim} \mathcal{O}_{\lambda}$	0	2	6
Springer fibre	pt	(^(m)	B
dimfibre	0	1	3

Example n = 4: Drawing pictures is no longer so reasonable, but we can still count dimensions and make our table. The dimension of \mathcal{B} is the number of positive roots (which for SL₄ is 6), so we know that dim $T^*\mathcal{B} = 12$.

To find the dimensions of the Springer fibres, we can play a similar game to what we did in the previous example. Consider the orbit \mathcal{O}_{λ} corresponding to

$$\lambda =$$

The matrix

$$e = \begin{pmatrix} 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 \end{pmatrix}$$

is in \mathcal{O}_{λ} , and we have that

$$0 \subset \operatorname{im} e = \ker e \subset \mathbb{C}^4.$$

The Springer fibre is

$$F_{\lambda} = \{ (L \subset P \subset H \subset \mathbb{C}^4) \mid e\mathbb{C}^4 \subset H, eH \subset P, eP \subset L, eL = 0 \}.$$

Again, we have two possibilities:

- ("easy component") { $P = ime = \ker e$, free choice of L, H as long as $L \subset P \subset H$ } $\simeq \mathbb{P}^1 \mathbb{C} \times \mathbb{P}^1 \mathbb{C} =: \Sigma_1$
- ("hard component") $\{H = e^{-1}L\} =: \Sigma_2$. There is a natural map $\Sigma_2 \to \mathbb{P}^1\mathbb{C}$ sending $(L \subset P \subset H) \mapsto L$, and the fibre over L' is $\{(L' \subset P \subset e^{-1}L' = H)\} \simeq \mathbb{P}^1\mathbb{C}$, so Σ_2 is a $\mathbb{P}^1\mathbb{C}$ -bundle over $\mathbb{P}^1\mathbb{C}$.

The diagonal $\Delta \subset \Sigma_1$ embeds into Σ_2 as the zero section, so these two components are glued together along a $\mathbb{P}^1\mathbb{C}$.

We've established that the second component Σ_2 is a $\mathbb{P}^1\mathbb{C}$ -bundle over $\mathbb{P}^1\mathbb{C}$, but which one?

Claim 13.3. $\Sigma_2 = \mathbb{P}(\mathcal{O}(1) \oplus \mathcal{O}(-1))$ is the "second Hirzebruch surface".

For justification of this claim, see Geordie's hand-written notes. We finish this example by constructing our table.

λ					
$\dim \mathcal{O}_{\lambda}$	12	10	8	6	0
$\operatorname{codim} \mathcal{O}_{\lambda}$	0	2	4	6	12
Springer fibre	pt		$\Sigma_1 \cup_{\mathbb{P}^1 \mathbb{C}} \Sigma_2$	$\mathcal{B}_{\mathrm{SL}_3} \cup 2$ other comp.	$\mathcal{B}_{\mathrm{SL}_4}$
dimfibre	0	1	2	3	6

Now that we've constructed four tables, we can make some observations about patterns we see.

Observations:

- $2\dim F_{\lambda} = \operatorname{codim} \mathcal{O}_{\lambda}$
- For any λ , one ("easy") component of F_{λ} is isomorphic to a flag variety of a smaller group.
- Fibres are equidimensional⁹ and components appear to be smooth¹⁰.

Remark 13.4. An audience member "Mr. Wiggins" also observed that in our examples, dim $T^*\mathcal{B}$ is divisible by all of the codim \mathcal{O}_{λ} , a property we coined "Wiggins divisibility". (Though we are not sure if this is a general phenomenon...)

Many of our observations hold in general. Here are some fundamental properties of Springer fibers.

⁹This is really remarkable!

¹⁰Sadly this fails in bigger examples.

Theorem 13.5. Let G be an a semisimple algebraic group.

- 1. The Springer resolution $\pi_s : \widetilde{\mathcal{N}} \to \mathcal{N}$ is a *G*-equivariant, projective resolution of singularities, and is an isomorphism over $\mathcal{N}_{\text{reg}} = \{x \in \mathcal{N} \mid \dim Z_G X = \text{rank}\mathfrak{g}\}.$
- 2. For a nilpotent orbit $\mathcal{O}_{\lambda} \subset \mathcal{N}$, the corresponding Springer fibre $F_{\lambda} := \pi_s^{-1}(x), x \in \mathcal{O}_{\lambda}$ is equidimensional, and dim $F_{\lambda} = \frac{1}{2} \operatorname{codim}(\mathcal{O}_{\lambda} \subset \mathcal{N})$.
- 3. $H_{\text{odd}}(F_{\lambda}, \mathbb{Z}) = 0.$

Remark 13.6. 1. $H_{\text{even}}(F_{\lambda}, \mathbb{Z})$ is well-studied (for example, its Betti numbers are known).

- 2. Part 2. of the theorem implies that π_s is semismall and all strata are relevent.
- 3. In type A, F_{λ} have cell decompositions, $\mathbb{C}^0 \sqcup (\mathbb{C}^1)^? \sqcup \cdots$. This cell decomposition implies Part 3. of the theorem for type A. With much more work cell decompositions have been constructed in other classical types, but existence of cell decompositions is unknown in exceptional types.
- 4. In type A,

{components of F_{λ} } = dim(irrep of S_n indexed by λ).

In fact, there exists a canonical bijection

$$\left\{\begin{array}{c} \text{standard Young} \\ \text{tableaux of shape } \lambda \end{array}\right\} \leftrightarrow \left\{\begin{array}{c} \text{components} \\ \text{of } F_{\lambda} \end{array}\right\}.$$

13.3 The conormal space

Let

$$X = \bigsqcup_{\lambda \in \Lambda} X_{\lambda}$$


be a stratefied variety. For example, $\mathbb{C} = \mathbb{C}^{\times} \sqcup \{0\}$. Consider the space

$$\bigcup_{\lambda \in \Lambda} TX_{\lambda} \subset TX.$$

This is a horrible space! In our example, $T\mathbb{C} = \mathbb{C} \times \mathbb{C}$, and

$$\bigcup TX_{\lambda} = \{ (x, y) \in \mathbb{C}^2 \mid x \neq 0 \text{ or } x = 0, y = 0 \}.$$


Here's a picture:



In contrast to this, the *conormal space*

$$T^*_{\Lambda}X := \bigcup_{\lambda \in \Lambda} T^*_{\lambda}X \subset T^*X,$$

where $T_{\lambda}^* X = \{\xi \in T_x^* X \text{ for } x \in X_{\lambda} \mid \xi \text{ vanishes on } TX_{\lambda}\}$ is very nice. In our example of $\mathbb{C} = \mathbb{C}^{\times} \sqcup \{0\}, T_{\Lambda}^* \mathbb{C} = \{(x, y) \in \mathbb{C}^2 \mid xy = 0\}$:



Properties of $T^*_{\Lambda}X$:

- 1. $T^*_{\Lambda}X$ is a closed subvariety of T^*X .
- 2. dim $T_{\lambda}^* X = \dim X_{\lambda} + \operatorname{codim}(X_{\lambda} \subset X) = \dim X$ is independent of λ ! (So we have a "democracy of strata".)
- 3. The components of the conormal space are in bijection with the strata.

The conormal space is a fundamental object in microlocal geometry.

Warning: the intersection pattern of $\overline{T_{\lambda}^*X}$ may be very complicated.

An important object in our story arises as a conormal variety, the Steinberg variety.

13.4 The Steinberg variety

Let H be a group that acts on a variety X on the right, and a variety Y on the left. We can form the *balanced product*

$$X \times_H Y := X \times Y/(xh, y) \sim (x, hy).$$

This space may not exist as a variety in general, but in all examples we will encounter, it does.

Choose a Borel subgroup $B \subset G$, so $\mathcal{B} \simeq G/B$. Then consider the variety

$$G \times_B G/B \xrightarrow{\sim} G/B \times G/B$$
$$(g, g'B) \mapsto (gg'B, g'B)$$

The set of G-orbits on $G \times_B G/B$ is equal to the set of B-orbits on G/B, which is parameterized by W by the Bruhat decomposition. So we have a stratification

$$\mathcal{B} \times \mathcal{B} = \bigsqcup_{x \in W} \mathcal{O}_x$$

In type A, this stratification is given by "pairs of flags in relative position x".

Example 13.7. There is a stratification of $\mathbb{P}^1\mathbb{C} \times \mathbb{P}^1\mathbb{C}$ given by

$$\mathbb{P}^{1}\mathbb{C} \times \mathbb{P}^{1}\mathbb{C} = \Delta \sqcup (\mathbb{P}^{1}\mathbb{C} \times \mathbb{P}^{1}\mathbb{C}) \setminus \Delta.$$


Pairs of flags in Δ are in relative position *id* (i.e. they are equal), and flags in $(\mathbb{P}^1\mathbb{C}\times\mathbb{P}^1\mathbb{C})\setminus\Delta$ are in relative position *s*.

Remark 13.8. The variety of Borel subalgebras \mathcal{B} does not depend on any choices, so the product $\mathcal{B} \times \mathcal{B}$ does not depend on any choices. Since we can define the Weyl group as the set of *G*-orbits on $\mathcal{B} \times \mathcal{B}$, this gives us a canonical definition of the Weyl group that does not depend on a choice!

Definition 13.9. The Steinberg variety

$$St = \{(\mathfrak{b}, \mathfrak{b}', x) \in \mathcal{B} \times \mathcal{B} \times \mathcal{N} \mid x \in \mathfrak{b}, x \in \mathfrak{b}'\}$$

is given by the fibre product



where the maps are

Exercise 13.10. (Solution can be found in Geordie's hand-written notes.) Show that the Steinberg variety St is the conormal variety of

$$\mathcal{B} imes \mathcal{B} = \bigsqcup_{x \in W} \mathcal{O}_x.$$

Corollary 13.11. There is a canonical bijection

$$\left\{\begin{array}{c} \text{irred. comp.} \\ \text{of } St \end{array}\right\} \leftrightarrow W.$$

To finish this lecture, let us recall a remarkable geometric construction of the Robinson-Schensted correspondence, due to Steinberg. Geordie is given to understand that this is the origin of the name "Steinberg variety". Fix λ , and two components C, C' of F_{λ} . We have the following diagram

$$\begin{array}{c} \pi_s^{-1}(\mathcal{O}_\lambda) \longleftrightarrow \mathcal{O}_\lambda \times C \\ \downarrow \qquad \qquad \downarrow \\ \mathcal{O}_\lambda = \mathcal{O}_\lambda \end{array}$$

The set $\{(x, c, c') \mid x \in \mathcal{O}_{\lambda}, c \in C, c' \in C'\}$ is a subvariety in St of dimension

 $\dim \mathcal{O}_{\lambda} + \dim C + \dim C' = \dim \mathcal{O}_{\lambda} + \frac{1}{2} \operatorname{codim}(\mathcal{O}_{\lambda} \subset \mathcal{N}) + \frac{1}{2} \operatorname{codim}(\mathcal{O}_{\lambda} \subset \mathcal{N}) = \dim \mathcal{N} = \dim \mathcal{B} \times \mathcal{B}.$

The components of St all have dimension $\dim \mathcal{B} \times \mathcal{B}$, so we have a bijection

{components of St} \leftrightarrow {($\mathcal{O}_{\lambda}, C, C'$) as above}.

In type A, this is a bijection between

 $W \leftrightarrow \{(\lambda, T, T') \mid T, T' \text{ are standard Young tableaux of shape } \lambda\}.$

This is the Robinson-Schensted correspondence!

Lecture 14: Springer correspondence and Borel-Moore homology

14.1 The Springer correspondence

We pick up in the setting of the last lecture, the **Springer resolution**:

$$\begin{array}{ccc} \widetilde{\mathcal{N}} \simeq T^* \mathcal{B} & & F_x := f^{-1}(x) \\ & & & & \downarrow \\ & & & & \downarrow \\ \mathcal{N} \subset \mathfrak{g} & & & x \end{array}$$

Roughly, the **Springer correspondence** states that W acts on $H^*(F_x; \mathbb{Q})$, and one obtains all irreducible representations of W in top cohomology. Note that this does *not* come from an action of W on F_x ! This is in contrast to Deligne-Lusztig theory and other settings where we obtain representations of a group in the cohomology of varieties on which the group acts. In the first part of today's lecture, we'll work towards a precise statement of the Springer correspondence.

Remark 14.1. In type A, the Springer correspondence explains why irreducible representations of the symmetric group and nilpotent orbits are both classified by the same combinatorial data (Young diagrams).

Grothendieck-Springer alteration: Let

$$\widetilde{\mathfrak{g}} = \{(x, \mathfrak{b}) \in \mathfrak{g} \times \mathcal{B} \mid x \in \mathfrak{b}\}.$$

In type A, this is equal to

$$\{(x,F) \in \mathfrak{g} \times \mathcal{F} \ell ags \mid xF^i \subset F^i\}.$$

Remark 14.2. Note that $\tilde{\mathfrak{g}} \subset \mathcal{B} \times \mathfrak{g}$, and its dual is $\tilde{\mathcal{N}} = T^*\mathcal{B}$. (This is because the dual of $x \in \mathfrak{b}$ is $x \in \mathfrak{b}^{\perp} = \mathfrak{n}$.)

The key diagram is the following:

where $\mathfrak{g}_{r.s.} = \{x \in \mathfrak{g} \mid x \text{ regular semisimple}\}$. Note that $\mathfrak{g} \neq \mathcal{N} \cup \mathfrak{g}_{r.s.}$.

Theorem 14.3. 1. The map π_s is semismall.

2. The map π_G is small (i.e., $\pi_{G*}k_{\tilde{\mathfrak{g}}}[\dim \mathfrak{g}] = IC(\mathfrak{g}_{r.s.}, \mathcal{L})).$

3. Over $\mathfrak{g}_{r.s.}$, π_G is a W-torsor. (Here W is the Weyl group of G.)

For example, in type A,

 $\mathfrak{g}_{r.s.} = \{x \in \mathfrak{g} \mid x \text{ is semisimple with distinct eigenvalues}\}.$

To give a flag F preserved by x is equivalent to an ordering of the eigenvalues of x. This is a S_n -torsor.

Now we are ready to state the Springer correspondence precisely. Given $x \in \mathcal{N}$, let $A_G(x)$ be the component group of the centralizer: $A_G(x) := C_G(x)/C_G(x)^\circ$.

Theorem 14.4. (Springer correspondence)

1.
$$\pi_{s*}\mathbb{Q}_{\widetilde{\mathcal{N}}}[\dim\widetilde{\mathcal{N}}] = \bigoplus_{\substack{x\in\mathcal{N}/G,\\\rho\in\operatorname{Irr}A_G(x)}} H_{top}(F_x)_{\rho}\otimes IC(\overline{G\cdot x},\mathcal{L}_{\rho}).$$

2. (**Most important**) $\operatorname{End}(\pi_{s*}\mathbb{Q}_{\widetilde{\mathcal{N}}}[\dim \widetilde{\mathcal{N}}]) = \operatorname{End}(\pi_{G*}\mathbb{Q}_{\widetilde{\mathfrak{g}}}[\dim \widetilde{\mathfrak{g}}]) = \mathbb{Q}W.$

3. $\{H_{top}(F_x)_{\rho}\}$ are all irreducible representations of W.

Remark 14.5. It is not difficult to see that in fact 1. and 3. are consequences of 2.

Remark 14.6. Part 2. of the theorem is true over \mathbb{Z} , and indeed over any ring. But part 1. fails over arbitrary rings because the decomposition theorem fails.

How might we approach part 2.? Three possible approaches:

- (a) **Borho MacPherson**:
 - First note that because $\widetilde{\mathfrak{g}} \to \mathfrak{g}$ is small, the fact that $\operatorname{End}(\pi_{G*}\mathbb{Q}_{\widetilde{\mathfrak{g}}}[\dim \mathfrak{g}]) = \mathbb{Q}W$ is obvious. Indeed,

 $\operatorname{End}(\pi_{G*}\mathbb{Q}_{\widetilde{\mathfrak{g}}}) = \operatorname{End}(\pi_{G*}\mathbb{Q}_{\widetilde{\mathfrak{g}}}|_{req.ss.}) = \operatorname{End}_{\mathbb{Q}W}(\mathbb{Q}W) = \mathbb{Q}W,$

with the first equality following from smallness.

• Then BM point out that we have a homomorphism

 $\operatorname{End}(\pi_{G*}\mathbb{Q}_{\widetilde{\mathfrak{g}}}) \xrightarrow{r} \operatorname{End}(i^*\pi_{G*}\mathbb{Q}_{\widetilde{\mathfrak{g}}}) = \operatorname{End}(\pi_{s*}\mathbb{Q}_{\widetilde{\mathcal{N}}})$

where $i : \mathcal{N} \hookrightarrow \mathfrak{g}$; and, miraculously, r is an isomorphism. (Proof sketch: The map r is injective because the action of W on $H^*(\mathcal{B})$ is faithful, then compare dimensions.)

(b) Fourier transform: (Springer's original approach) Because $\widetilde{\mathcal{N}}$ and $\widetilde{\mathfrak{g}}$ are dual in $\mathcal{B} \times \mathfrak{g}$, $k_{\widetilde{\mathcal{N}}}$ and $k_{\widetilde{\mathfrak{g}}}$ are Fourier transforms of one another. This implies that the endomorphism rings are equal:

$$\operatorname{End}(\pi_{s*}k_{\widetilde{\mathcal{N}}}) = \operatorname{End}(\pi_{G*}k_{\widetilde{\mathfrak{q}}}) = kW.$$

(c) **Convolution algebras**: (in Chris-Ginzburg)

The next part of the lecture will explain this approach. For impatient readers, we'll give away the ending:

The punchline: There is a notion of homology (Borel-Moore homology, H_{BM}) such that $\operatorname{End}(\pi_{s*}k_{\widetilde{N}})$ is canonically equal to H_{BM} (Steinberg variety). This gives a concrete realisation of this endomorphism algebra.

14.2 Borel-Moore homology

Let X be an algebraic variety, k a field, $p: X \to pt$, and k_X the constant sheaf on X. Here are four notions of (co)homology:

- $H^*(X;k) = H^*(p_*k_X)$ cohomology (cochains)
- $H_c^*(X;k) = H^*(p_!k_X)$ cohomology with compact support
- $H_*(X;k) = H^{-*}(p_!\omega_X)$ homology (chains)
- $H^{BM}_*(X;k) = H^{-*}(p_*\omega_X)$ Borel-Moore homology (locally finite chains)

Here ω_X is the dualising sheaf, $\mathbb{D}k_X$. The canonical example is the following.

Example 14.7. Let $X = \{z \in \mathbb{C} \mid 0 < |z| < 1\}$. Then

$$H_0(X;\mathbb{Z}) = \mathbb{Z}\left\{\left\{ \begin{array}{c} \mathbf{x} \\ \mathbf{x} \end{array}\right\}, H_1(X;\mathbb{Z}) = \mathbb{Z}\left\{\left\{ \begin{array}{c} \mathbf{x} \\ \mathbf{x} \end{array}\right\}, H_2(X,\mathbb{Z}) = 0 \right\}\right\}$$

If we compute Borel-Moore homology, we get

$$H_0^{BM}(X;\mathbb{Z}) = 0, \ H_1^{BM}(X,\mathbb{Z}) = \mathbb{Z}$$
 ($H_2^{BM}(X;\mathbb{Z}) = \mathbb{Z}$ "fundamental class".

In the first computation (H_0^{BM}) , the formal generator of H_0 is now a boundary, so we get 0. In the second computation (H_1^{BM}) , we can now have cycles from the edge to the center, and these are not boundaries. In the third computation (H_2^{BM}) , the "fundamental class" is a triangulation of X.

From now on in this course, $H_* = H_*^{BM}$.

Key properties of Borel-Moore homology:

1. If $X \stackrel{\iota}{\hookrightarrow} M$ is a closed embedding of X into a smooth, \mathbb{C} -dimension d variety M, then we have local Poincaré duality: $\omega_M \simeq k_M[2d]$. Hence

$$H_*(X;k) = H^{-*}(X,\omega_X) = H^{-*}(X,i^!\omega_M) = H^{-*}(M,i_!i^!k_M[2d]) = H^{2d-*}(M,i_!i^!k_M) = H^{2d-*}(M,M\backslash X;k).$$

2. $H_*(-)$ is not functorial for arbitrary maps, but for proper maps p, we have p_* .

3. For an open inclusion $U \hookrightarrow X$, we have restriction

$$H_*(X) \to H_*(U).$$

(Thought exercise: Why? In terms of chains?)

4. If X is equidimensional of dimension d with components X_1, X_2, \ldots, X_m , then

$$H_{2d}(X) = \bigoplus \mathbb{Z}[X_i],$$

where $[X_i]$ are the fundamental classes of the components.

14.3 Geometric convolution algebras


Let X_1, X_2, X_3 be smooth varieties of dimensions d_1, d_2, d_3 , respectively, and

$$Z_{12} \subset X_1 \times X_2, \quad Z_{23} \subset X_2 \times X_3$$

closed subvarieties ("correspondences"). (For example, we could take $Z_{12} = \operatorname{graph}(f)$ for some $f: X_1 \to X_2$.) Define

$$Z_{12} \circ Z_{23} := \{ (x_1, x_3) \in X_1 \times X_3 \mid \text{ there exists } x_2 \in X_2 \text{ s.t. } (x_1, x_2) \in Z_{12}, (x_2, x_3) \in Z_{23} \}.$$

(So in our example, $graph(f) \circ graph(g) = graph(g \circ f)$.) We have projections



We make the following **properness assumption**: From now on, assume that the map

$$p_{13}: p_{12}^{-1}(Z_{12}) \cap p_{23}^{-1}(Z_{23}) \to X_1 \times X_3$$

is proper.

Definition 14.8. We define a **convolution product** on homology:

$$H_i(Z_{12}) \times H_j(Z_{23}) \to H_{i+j-2d_2}(Z_{12} \circ Z_{23})$$
$$(c_{12}, c_{23}) \mapsto p_{13*}(p_{12}^*c_{12} \cap p_{23}^*c_{23})$$

Here \cap is the intersection product in Borel-Moore homology.

Most important cases: Let $\widetilde{X} \xrightarrow{f} X$ be a proper map of a smooth variety \widetilde{X} to X, and let $X_i = \widetilde{X}$ for i = 1, 2, 3. Here are two cases of the construction above:


1. Let $Z_{12} = Z_{23} = Z_{12} \circ Z_{23} = \widetilde{X} \times_X \widetilde{X} \subset \widetilde{X} \times \widetilde{X}$. Then the convolution product gives an associative algebra structure on $H_*(\widetilde{X} \times_X \widetilde{X})$ (with messy gradings). 2. Let $Z_{12} = \widetilde{X} \times_X \widetilde{X}$, $Z_{23} = f^{-1}(X)$, $Z_{12} \circ Z_{23} = Z_{23}$. Then the convolution product gives us a map

$$H_*(X \times_X X) \times H_*(f^{-1}(X)) \to H_*(f^{-1}(X)).$$

This gives $H_*(f^{-1}(X))$ the structure of a $H_*(\widetilde{X} \times \widetilde{X})$ -module.

So whenever we have a smooth proper map into our variety, we obtain from this formalism an associative algebra and a collection of modules (one for each fibre) over that algebra. Seems promising!

Conceptual explanation of convolution algebras: Let f be as above, with the fibre product diagram



Claim: End $(f_*k_{\widetilde{X}}) = H_*(\widetilde{X} \times_X \widetilde{X})$

Proof. We will only check the statement on the level of vector spaces. Let $d_X := \dim_{\mathbb{C}} X$. We have

$$\operatorname{Hom}^{\cdot}(f_{*}k_{\widetilde{X}}, f_{*}k_{\widetilde{X}}) = \operatorname{Hom}^{\cdot}(f^{*}f_{*}k_{\widetilde{X}}, k_{\widetilde{X}})$$
$$= \operatorname{Hom}^{\cdot}(f^{*}f_{!}k_{\widetilde{X}}, k_{\widetilde{X}})$$
$$= \operatorname{Hom}^{\cdot}(g_{!}g^{*}k_{\widetilde{X}}, k_{\widetilde{X}})$$
$$= \operatorname{Hom}^{\cdot}(k_{\widetilde{X} \times_{X} \widetilde{X}}, g^{!}k_{\widetilde{X}})$$
$$= \operatorname{Hom}^{\cdot}(k_{\widetilde{X} \times_{X} \widetilde{X}}, g^{!}\omega_{\widetilde{X}}[-2d_{X}])$$
$$= H^{*-2d_{X}}(\widetilde{X} \times_{X} \widetilde{X}, \omega_{\widetilde{X} \times \widetilde{X}})$$
$$= H_{2d_{X}-*}(\widetilde{X} \times_{X} \widetilde{X}).$$

Here we are using properness of f (second equality), adjunctions (first equality, fourth equality), proper base change (third equality), and local Poincaré duality (fifth equality).

The Upshot: Up to gradings,

$$H_*(\widetilde{X} \times_X \widetilde{X}) = \operatorname{End}^{\cdot}(f_*k_{\widetilde{X}}),$$

and with some work we can show that multiplication matches on both sides. Similarly,

$$(f_*k_{\widetilde{X}})_x = H_*(f^{-1}(X)),$$

and the module structure comes from the action $\operatorname{End}^{\cdot}(f_*k_{\widetilde{X}}) \circlearrowright (f_*k_{\widetilde{X}})_x$.

Connection to the Spring correspondence:

Let $\widetilde{\mathcal{N}} \xrightarrow{\pi_s} \mathcal{N}$ be the Springer resolution, and

$$St := \widetilde{\mathcal{N}} \times_{\mathcal{N}} \widetilde{\mathcal{N}} = \begin{array}{c} \text{conormal space to} \\ \text{the } G \text{-space } \mathcal{B} \times \mathcal{B} \end{array} = \bigcup_{x \in W} T_x^*,$$

the Steinberg variety. (See lecture 13.) Here T_x^* is the conormal bundle to the *G*-orbit $\mathcal{O}_x \subset \mathcal{B} \times \mathcal{B}$. Recall that *St* is equidimensional and all components have dimension equal to $N := \dim \mathcal{B} \times \mathcal{B} = \dim T^* \mathcal{B}$.

The convolution product in Borel-Moore homology gives us a map

$$H_{2N}(St) \times H_{2N}(St) \xrightarrow{*} H_{4N-2\dim T^*\mathcal{B}}(St) = H_{2N}(St).$$

This gives $H_{2N}(St)$ the structure of an algebra!

Theorem 14.9. As algebras,

$$H_{2N}(St) = kW.$$


By the theory earlier, the action of kW on $H_i(F_x)$ yields the Springer action. In this way we end up with W-modules everywhere!

Lecture 15: The Kazhdan–Lusztig isomorphism

15.1 More convolution algebras

Last week we ended with a discussion on convolution algebras. We'll start today by continuing this discussion. Consider the following two settings:

- 1. Let G be a group and $H \subset G$ a subgroup. With the operation * of convolution of functions, the vector space $\operatorname{Fun}(G, \mathbb{C})$ of complex-valued functions on G has the structure of an algebra. This is just the group algebra, $\operatorname{Fun}(G, \mathbb{C}) \simeq \mathbb{C}[G]$. The subspace $\operatorname{Fun}_{H \times H}(G, \mathbb{C})$ of H-biinvariant functions forms a subalgebra. This is a *Hecke algebra*. As many of us are aware, its representation theory is very complicated in general!
- 2. Now let X be a finite set. The vector space $\operatorname{Fun}(X \times X, \mathbb{C})$ of complex-valued functions on $X \times X$ can also be given the structure of an algebra. In this setting, the convolution product is given as follows. Let



be the natural projections. Then for $f, g \in \operatorname{Fun}(X \times X, \mathbb{C})$, define $f * g \in \operatorname{Fun}(X \times X, \mathbb{C})$ by

$$(f*g)(x,z) := \sum_{y \in X} f(x,y)g(y,z)$$

for $(x, z) \in X \times X$. In other words,

$$f * g = p_{13!}(p_{12}^* f \boxtimes p_{23}^* g).$$

With *, (Fun($X \times X, \mathbb{C}$), *) is an algebra. In fact, it's a familiar¹¹ algebra. Let $e_{x,y}$ be the indicator function on $(x, y) \in X \times X$. Then $e_{x,y} * e_{y',z} = \delta_{y,y'} e_{x,z}$. Hence,

$$(\operatorname{Fun}(X \times X, \mathbb{C}), *) \simeq \operatorname{Mat}_{X \times X}(\mathbb{C}).$$

Some Variants:

(a) Let $X \to Y$ be a map of sets, then we can construct the convolution algebra

$$\operatorname{Fun}(X \times_Y X, \mathbb{C}) \simeq \prod_{y \in Y} \operatorname{Mat}_{f^{-1}(y) \times f^{-1}(y)}(\mathbb{C}).$$

We can formulate Mashke's theorem in this language. For a finite group G, let $Y = \{\text{irreps of } G\}, X = \bigsqcup_{\rho \in Y} \{\text{basis of } \rho\}, \text{ and } X \to Y \text{ the map which assigns to a basis element the corresponding representation. Then Mashke's Theorem that the group algebra is semisimple is the following statement:$

¹¹This is why we don't meet this algebra as often as we meet the Hecke algebra - it's "too easy," in the sense that it is just a matrix ring. However, when we categorify, it becomes more interesting, so we are more likely to encounter it (or regognize it!) in categorified settings.

Theorem 15.1. (Mashke's Theorem)

 $(\operatorname{Fun}(G,\mathbb{C}),*) \simeq (\operatorname{Fun}(X \times_Y X,\mathbb{C}),*).$

This realizes a complicated algebra (the group algebra) in terms of much simpler pieces (matrix rings).

(b) If our sets X, Y come with the additional structure of an action by a group Γ , and $X \to Y$ is a Γ -equivariant map of sets, then we can construct the convolution algebra

$$\operatorname{Fun}^{\Gamma}(X \times_Y X, \mathbb{C}).$$

The Upshot: There are two types of convolution algebras, one is hard (Hecke algebras), and the other is easy (Fun $(X \times X, \mathbb{C})$). The hard one is of great significance in representation theory. A strategy that we use in representation theory is to try to realise the hard type as the easy type.

15.2 Equivariant *K*-theory

Let G be an algebraic group acting on a variety X. Then we can formulate the notion of an equivariant coherent sheaf,

$$\mathcal{F} \in \operatorname{Coh}_G(X),$$

as a coherent sheaf \mathcal{F} on X, coupled with some extra data¹² Roughly speaking, one can think of an equivariant coherent sheaf on X as being an equivariant sheaf together with an algebraic action on its sections.

Remark 15.2. If \mathcal{F} is locally free, then \mathcal{F} corresponds to a vector bundle $V \to X$ on X. In this setting, *G*-equivariance of \mathcal{F} corresponds to an algebraic *G*-action on V which is compatible with projection. In particular,

$$\operatorname{Coh}_G(pt) \simeq \operatorname{Rep} G,$$

where $\operatorname{Rep} G$ is the category of algebraic representations of G.

Define two types of equivariant K-groups:

 $K^G(X) :=$ Grothendieck group of *G*-equivariant coherent sheaves on *X*; $K_G(X) :=$ Grothendieck group of *G*-equivariant vector bundles on *X*.

Remark 15.3. (a) Very loosely,

 $K^G(X) \leftrightarrow$ "Borel–Moore homology," $K_G(X) \leftrightarrow$ "cohomology".

¹²For an excellent description of this construction, see notes from Emily's Sept 20, 2019 talk in the Informal Friday Seminar. Notes from Emily's (and all other) IFS talks can be found at https://sites.google.com/view/ifssydney/home.

- (b) There are higher K-groups, $K_i^G(X), K_G^i(X)$, which we will ignore here. (Already $K_0^G(X) = K^G(X)$ and $K_G^0(X) = K_G(X)$ are rich enough.)
- (c) There is a natural map

$$K_G(X) \to K^G(X).$$

If X is smooth, then we can use resolutions by coherent sheaves to show that this map is an isomorphism.

(d) We have to be careful with functors between K-groups. For example, if $X \to pt$ is projection and X is affine, then $f_*\mathcal{O}_X = \Gamma(X, \mathcal{O}_X)$ is usually infinite dimensional. So in general, we need to work to justify that functors we wish to use preserve Coh, or at least $D^b(\text{Coh})$.

As we did for H_* , we have a convolution product in K-theory. We imitate the set-up of the previous lecture: Let X_1, X_2, X_3 be smooth varieties of dimensions d_1, d_2, d_3 , respectively, and

$$Z_{12} \subset X_1 \times X_2, \quad Z_{23} \subset X_2 \times X_3$$

closed subvarieties, with projections p_{ij} , i, j = 1, 2, 3 as before. Given $\mathcal{F} \in \operatorname{Coh}(Z_{12}), \mathcal{G} \in \operatorname{Coh}(Z_{23})$, define

$$\mathcal{F} * \mathcal{G} := p_{13*}(p_{12}^* \mathcal{F} \overset{L}{\otimes} p_{23}^* \mathcal{G}) \in D^b(\operatorname{Coh}(Z_{12} \circ Z_{23})).$$

As we emphasized in Remark 15.3, we need to justify that this product is well-defined. But sure enough, the push-forward p_{13*} preserves coherence because p_{13} is proper, and the pull-backs p_{12}^*, p_{23}^* are okay because p_{12} and p_{23} are flat. Hence * induces a product

$$K_0(Z_{12}) \times K_0(Z_{23}) \to K_0(Z_{12} \circ Z_{23})$$

which descends to a product

$$K^G(Z_{12}) \times K^G(Z_{23}) \to K^G(Z_{12} \circ Z_{23}).$$

Example 15.4. Let X be a finite set, and $X \to pt$ projection to a point. In this case, elements of $Coh(X \times X)$ are "matrices of vector spaces," and

$$*: \operatorname{Coh}(X \times X) \times \operatorname{Coh}(X \times X) \to \operatorname{Coh}(X \times X)$$

is "multiplication of matrices". That is, for vector spaces V_{ij} ,

$$\begin{pmatrix} V_{11} & V_{12} \\ V_{21} & V_{22} \end{pmatrix} * \begin{pmatrix} V'_{11} & V'_{12} \\ V'_{21} & V'_{22} \end{pmatrix} = \begin{pmatrix} V_{11} \otimes V'_{11} \oplus V_{12} \otimes V'_{21} & V_{11} \otimes V'_{12} \oplus V_{12} \otimes V'_{22} \\ V_{21} \otimes V'_{11} \oplus V_{22} \otimes V'_{21} & V_{21} \otimes V'_{12} \oplus V_{22} \otimes V'_{22} \end{pmatrix}.$$

The K-group is $K_0(X \times X) = \operatorname{Fun}(X \times X, \mathbb{Z}).$

Remark 15.5. Lusztig noticed that certain categories of the form $\operatorname{Coh}^{\Gamma}(X \times X)$, where Γ is a group acting on a finite set X, are central to the classification of unipotent character sheaves.

15.3 The Kazhdan–Lusztig isomorphism

With this we have enough machinery to state the Kazhdan–Lusztig isomorphism. Let $(X \supset R, X^{\vee} \supset R^{\vee})$ be a root datum, $W_{ext} = W_f \ltimes \mathbb{Z} X^{\vee}$ the extended affine Weyl group, and \mathcal{H}_{ext} the affine Hecke algebra. (See lecture 11 for a refresher on these objects.) Let G^{\vee} be the corresponding algebraic group, $\mathcal{N}^{\vee} \subset \mathfrak{g}^{\vee}$ the nilpotent cone, and $St^{\vee} := T^* \mathcal{B}^{\vee} \times_{\mathcal{N}^{\vee}} T^* \mathcal{B}^{\vee}$ the Steinberg variety. (See Lecture 13.)

Theorem 15.6. (Kazhdan–Lusztig) We have canonical isomorphisms.

$$\begin{array}{ccc} \mathcal{H}_{ext} & \stackrel{\sim}{\longrightarrow} & K^{G^{\vee} \times \mathbb{C}^{\times}}(St^{\vee}) \\ & & \downarrow^{\text{forget}} \\ \mathbb{Z}W_{ext} & \stackrel{\sim}{\longrightarrow} & K^{G^{\vee}}(St^{\vee}) \end{array}$$

Here \mathbb{C}^{\times} acts on $T^*\mathcal{B}^{\vee}$ by dilation along the fibres.

Here are two examples of how Theorem 15.6 gives us insight into the representation theory of affine Hecke algebras.

1. (Bernstein) As we have discussed, there is an inclusion of algebras:

$$\mathbb{Z}[v^{\pm 1}][X^{\vee}] \subset \mathcal{H}_{ext}$$
$$\lambda \mapsto H_{\lambda}$$

Bernstein noticed that the center of \mathcal{H}_{ext} can be realized as W_f -invariants of this subalgebra:

$$\mathbb{Z}[v^{\pm 1}][X^{\vee}]^{W_f} = Z(\mathcal{H}_{ext}).$$

We can see this in terms of the Kazhdan–Lusztig isomorphism. Recall that for a map $X \to Y$ between finite sets, we have

$$\begin{array}{cccc} X \xrightarrow{\text{diagonal}} X \times X \longrightarrow X \\ & & \downarrow \\ & & \downarrow \\ & & \chi \longrightarrow Y \end{array}$$

and Fun(diag, \mathbb{C}) \subset Fun($X \times_Y X, \mathbb{C}$) are "diagonal matrices". Applying this to the diagonal $T^*\mathcal{B}^{\vee} \subset St^{\vee}$, we can think of

$$K^{G\times\mathbb{C}^{\times}}(T^{*}\mathcal{B}^{\vee})\subset K^{G\times\mathbb{C}^{\times}}(St^{\vee})$$

as "diagonal matrices". Now by homotopy,

$$K^{G^{\vee}\times\mathbb{C}^{\times}}(T^*\mathcal{B}^{\vee})=K^{G^{\vee}\times\mathbb{C}^{\times}}(\mathcal{B}^{\vee}),$$

and since $K^{B^{\vee}}(pt) = [\operatorname{Rep} B^{\vee}] = \mathbb{Z}[X^{\vee}]$ and $K^{\mathbb{C}^{\times}}(pt) = [\operatorname{Rep} \mathbb{C}^{\times}] = \mathbb{Z}[v^{\pm 1}],$

$$K^{G^{\vee} \times \mathbb{C}^{\times}}(\mathcal{B}^{\vee}) = K^{G^{\vee} \times \mathbb{C}^{\times}}(G^{\vee}/B^{\vee}) = K^{B^{\vee} \times \mathbb{C}^{\times}}(pt) = \mathbb{Z}[v^{\pm 1}][X^{\vee}].$$

So the subalgebra $\mathbb{Z}[v^{\pm 1}][X^{\vee}]$ sits inside \mathcal{H}_{ext} as "diagonal matrices" in the $G^{\vee} \times \mathbb{C}^{\times}$ -equivariant K-group of St^{\vee} . Moreover, $K^{G^{\vee} \times \mathbb{C}^{\times}}(pt)$ is clearly central in $K^{G^{\vee} \times \mathbb{C}^{\times}}(St^{\vee})$, and this gives the Bernstein center:

$$K^{G^{\vee} \times \mathbb{C}^{\times}}(pt) = (K^{B^{\vee} \times \mathbb{C}^{\times}}(pt))^{W_f} = (\mathbb{Z}[v^{\pm 1}][X^{\vee}])^{W_f}.$$

2. The Kazhdan–Lusztig basis of \mathcal{H}_{ext} leads to the notion of (left, right, two-sided) cells¹³. Lusztig noticed that the sets

$$\left\{\begin{array}{c} \text{poset of} \\ 2\text{-sided} \\ \text{cells} \end{array}\right\} \leftrightarrow \left\{\begin{array}{c} \text{nilpotent} \\ \text{orbits} \\ \text{in } \mathcal{N}^{\vee} \end{array}\right\}$$


appear to match.

He also noticed other remarkable parallels, such as

$$\begin{array}{ccc} \text{2-sided cell} & & \text{reductive part of} \\ \text{is finite} & \longleftrightarrow & \text{the centralizer} \\ & & \text{is finite} \end{array}$$

These parallels convince one rather quickly that this correspondence is deep.

We can use Theorem 15.6 to understand this observation of Lustig. Consider the projections



Convolution makes it clear that any closed G^{\vee} -invariant subvariety $Z \subset \mathcal{N}^{\vee}$ gives rise to a two-sided ideal in $K^{G^{\vee} \times \mathbb{C}^{\times}}(St^{\vee})$:

$$\{[M] \mid \operatorname{supp} M \subset p^{-1}(Z)\}.$$

¹³Pictures of 2-sided cells in rank 2 can be found on Lusztig's webpage: http://www-math.mit.edu/ ~gyuri/picture.html.

Hence we get a filtration of \mathcal{H}_{ext} by nilpotent orbits in \mathcal{N}^{\vee} ! Lusztig and Bezrukavnikov showed that the filtration of \mathcal{H}_{ext} coming from 2-sided cells agrees with this filtration coming from geometry, explaining Lusztig's observation that the 2-sided cell order matches the order on nilpotent orbits.

3. We won't explain this in detail due to time constraints, but one can show relatively easily that Theorem 15.6 implies the Deligne-Langlands conjecture (Section 12.2) on representations of \mathcal{H}_{ext} , as long as q is not a root of unity.

15.4 Bezrukavnikov's equivalence: rough outline

Now we are finally in the position to approach the second half of the title of this course. Here is a rough outline of the flow of ideas that lead to Bezrukavnikov's equivalence:

- Langlands correspondence: A correspondence between sets satisfying a whole host of properties, and with many extraordinary consequences in number theory and beyond.
- Weil: In the function field case, an automorphic form¹⁴ can be regarded as a (very special) function

 $f: \operatorname{Bun}_{G}(\mathbb{F}_{q}) := \left\{ \begin{array}{c} \text{iso classes of} \\ G\text{-bundles on} \\ \text{smooth curves}/\mathbb{F}_{q} \end{array} \right\} \to \mathbb{C}.$

- **Grothendieck**: Functions on a variety should be understood as shadows of sheaves (function-sheaf correspondence). Here are some examples.
 - E.g. 1: Characters of $\operatorname{GL}_n(\mathbb{F}_q)$ are shadows (trace of Frob) of certain ℓ -adic sheaves on G, "character sheaves".
 - E.g. 2: Interesting analytic functions should satisfy many differential equations (i.e. should be solutions of a holonomic *D*-module).
- **Drinfeld**: The Langlands correspondence should be approached using Grothendieck's dictionary. At its most basic level, this philosophy asserts that automorphic forms in the function field case should arise as traces of Frobenius of certain sheaves on the moduli space of *G*-bundles. At a more sophisticated level, we should expect an equivalence of categories

$$D^b(\operatorname{Sh}(\operatorname{Bun}_G)) \xleftarrow{\gamma\gamma\gamma} D^b(\operatorname{Coh}(\operatorname{Loc}_G)),$$

i.e. a "geometric Langlands correspondence". This allows us to work over \mathbb{C} , and provides fertile connections to physics, higher category theory, etc.

¹⁴We haven't gone into what an automorphic form is yet in this course. For a global field K it is a function of a special form on a certain quotient of the adelic points of a reductive group. In the function field case, Weil realised that this quotient parametrises G-bundles on the corresponding curve. In other words, in the function field case an automorphic form can be regarded as a function of the form given above.

Remark 15.7. Geometric Langlands is for *function fields*. The classical Langlands correspondence isn't. So they seem to be living in different worlds, but the hope is that they are actually related. For many years this appeared to many as a pretty wild idea. However, recently the geometric Langlands program has been shown to have consequences that number theorists really care about. For example Fargues showed [Far18] that geometric Langlands for the Fargues-Fontaine curve implies (part of) the LLC for *any p*-adic group!

• **Ginzburg**: A small piece of geometric Langlands should be controlled by a categorification of the Kazhdan–Lusztig isomorphism

$$\mathcal{H}_{ext} \simeq K^{G^{\vee} \times \mathbb{C}^{\times}}(St^{\vee}).$$

What is sought is a fundamental monoidal category that arises in two Langlands dual ways. Recall that the geometric Satake equivalence categorifies $\mathcal{H}^{sph} = K^G(pt)$ (Theorem 11.2). Thus this equivalence can be seen as one layer of difficulty beyond the geometric Satake equivalence.

• Bezrukavnikov: realization of (several) such equivalences:

$$K^{G^{\vee} \times \mathbb{C}^{\times}}(St^{\vee}) \rightsquigarrow \operatorname{Coh}^{G^{\vee} \times \mathbb{C}^{\times}}(St^{\vee})$$
 "coherent side"
 $\mathcal{H}_{ext} \rightsquigarrow D^{b}_{Iw}(G((t))/Iw)$ "constructible side"

Theorem 15.8. (Bezrukavnikov's equivalence, most basic version) Let $Iw \subset G((t))$ be an Iwahori subgroup, and $Iw_0 \subset Iw$ the pro-unipotent radical. There is an equivalence of monoidal categories

$$\left(\widehat{D^b_{Iw_0}(G((t))/Iw}),*\right) \simeq \left(D^b\operatorname{Coh}^{G^{\vee}\times\mathbb{C}^{\times}}(\widetilde{St}^{\vee}),*\right).$$

where $\widetilde{St}^{\vee} = \widetilde{\mathfrak{g}}^{\vee} \times_{\mathfrak{g}^{\vee}} \widetilde{\mathfrak{g}}^{\vee}$ and the hat indicates the pro-unipotent completion.

Remark 15.9. There are several aspects of the above that need explanation, hopefully this will occur over the coming weeks and months!

Lecture 16: The constructible side of Bezrukavnikov's equivalence

Last lecture we ended by stating (modulo several undefined pieces) **Bezrukavnikov's equivalence**:

"constructible side" "coherent side"
$$\left(\widehat{D^b_{Iw_0}(G((t))/Iw)},*\right) \simeq \left(D^b\operatorname{Coh}^{G^\vee \times \mathbb{C}^\times}(\widetilde{St}^\vee),*\right).$$

Our goal for today is to motivate the constructible side of this equivalence. To do so, we will explain Grothendieck's function-sheaf correspondence in slightly more detail than is usually done. The starting place is the **Weil conjectures**.

16.1 Weil conjectures

The Weil conjectures were a collection of statements (made by Weil while in jail during the second world war) about counting the number of points on an algebraic variety over a finite field. Let X be a smooth projective algebraic variety over \mathbb{F}_q . Then we can ask about the number of points $\#X(\mathbb{F}_{q^n})$ for all n. Weil conjectured that a generating series built out of $\#X(\mathbb{F}_{q^n})$ has remarkable properties. He also pointed out that his conjecture would follow from an interpretation of $\#X(\mathbb{F}_{q^n})$ as the trace of an operator on a vector space, by taking traces of its powers. The Grothendieck–Lefschetz trace formula, which we briefly visited in the context of elliptic curves in Lecture 4, provides such an interpretation.

Grothendieck–Lefschetz trace formula: Let ℓ be a prime not dividing q, and X as above (but not necessarily projective). Then

$$#X(\mathbb{F}_{q^n}) = \operatorname{sTr}(\operatorname{Frob}_q^n \circlearrowright H_c^i(X_{\overline{\mathbb{F}}_q}, \overline{\mathbb{Q}}_\ell)) := \sum_i (-1)^i \operatorname{Tr}\left(\operatorname{Frob}_q^n \circlearrowright H_c^i(X_{\overline{\mathbb{F}}_q}, \overline{\mathbb{Q}}_\ell)\right).$$
(16.1)

Here $H_c^i(X_{\overline{\mathbb{F}}_q}, \overline{\mathbb{Q}}_\ell)$ denotes the compactly supported étale cohomology of $X_{\overline{\mathbb{F}}_q}$, the base change of X to Spec $\overline{\mathbb{F}}_q$. If X/k, then $H_c^*(X_{\overline{k}}, \overline{\mathbb{Q}}_\ell)$ has a continuous action of $\operatorname{Gal}(\overline{k}/k)$. The symbol sTr in (16.1) stands for "supertrace". Let's see what this formula means in examples.

Example 16.1. Let $X = \mathbb{P}^{1}_{\mathbb{F}_{q}}$, then the cohomology and action of Frobenius are given by the following table:

i	$H^i_c(X_{\overline{\mathbb{F}}_q}, \overline{\mathbb{Q}}_\ell)$	Frob action
2	$\overline{\mathbb{Q}}_{\ell}$	$\bigcirc q$
1	0	
0	$\overline{\mathbb{Q}}_{\ell}$	() 1

Hence,

$$\operatorname{sTr}\left(\operatorname{Frob}_{q}^{n} \circlearrowright H^{*}\right) = 1 + q^{n} = \#X(\mathbb{F}_{q^{n}}).$$

i	$H^i_c(X_{\overline{\mathbb{F}}_q},\overline{\mathbb{Q}}_\ell)$	Frob action
2	$\overline{\mathbb{Q}}_{\ell}$	$\bigcirc q$
1	$\overline{\mathbb{Q}}_{\ell}$	() 1
0	0	

Example 16.2. Let $X = \mathbb{G}_m$. Then the action of Frobenius on cohomology is:

Hence,

$$\operatorname{sTr}\left(\operatorname{Frob}_{q}^{n} \circlearrowright H^{*}\right) = q^{n} - 1 = \#\mathbb{G}_{m}(\mathbb{F}_{q^{n}}) = \#\mathbb{F}_{q^{n}}^{\times}.$$

Example 16.3. Let $X = \operatorname{Spec} \mathbb{F}_{q^2} / \operatorname{Spec} \mathbb{F}_q$. Then

$$X(\mathbb{F}_{q^n}) = \operatorname{Hom}(\operatorname{Spec} \mathbb{F}_{q^n}, \operatorname{Spec} \mathbb{F}_{q^2})$$
$$= \operatorname{Hom}_{\mathbb{F}_{q^{-}\operatorname{alg}}}(\mathbb{F}_{q^2}, \mathbb{F}_{q^n})$$
$$= \begin{cases} 0 \quad n \text{ odd,} \\ 2 \quad n \text{ even.} \end{cases}$$

On the geometric side,

$$\begin{aligned} X_{\overline{\mathbb{F}}_q} &= \operatorname{Spec} \overline{\mathbb{F}}_q \times_{\operatorname{Spec} \mathbb{F}_q} \operatorname{Spec} \mathbb{F}_{q^2} \\ &= \operatorname{Spec} (\overline{\mathbb{F}}_q \otimes_{\mathbb{F}_q} \mathbb{F}_{q^2}) \\ &= \operatorname{Spec} \overline{\mathbb{F}}_q \sqcup \operatorname{Spec} \overline{\mathbb{F}}_q \\ &\simeq \overline{\mathbb{F}}_q \times \overline{\mathbb{F}}_q. \end{aligned}$$

Remark 16.4. For analogy, an easier version of the computation above is the following. We can realize

$$\mathbb{C} \simeq \mathbb{R}[x]/(x^2+1).$$

Then

$$\mathbb{C} \otimes_{\mathbb{R}} \mathbb{C} = \mathbb{C}[x]/(x^2 + 1)$$
$$= \mathbb{C}[x]/(x - i)(x + i)$$
$$\simeq \mathbb{C} \times \mathbb{C}.$$

The Upshot: The variety $X_{\overline{\mathbb{F}}_q}$ consists of 2 points which are interchanged by $\operatorname{Gal}(\overline{\mathbb{F}}_q/\mathbb{F}_q)$. So our table of cohomology is

i	$H^i_c(X_{\overline{\mathbb{F}}_q},\overline{\mathbb{Q}}_\ell)$	Frob action
> 0	0	
0	$\overline{\mathbb{Q}}_\ell\oplus\overline{\mathbb{Q}}_\ell$	$\circlearrowleft \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$

and

$$\operatorname{sTr}\left(\operatorname{Frob}_{q}^{n} \circlearrowright H^{*}\right) = \begin{cases} 0 & n \text{ is odd,} \\ 2 & n \text{ is even.} \end{cases}$$

The Grothendieck–Lefschetz formula and the examples above can be realized as shadows of something happening on the level of sheaves. Let k be a field. Then, roughly,

$$\operatorname{Spec} k = \operatorname{pt}/\operatorname{Gal}(k/k).$$

In other words, one can think of Spec k as something like the classifying space of its absolute Galois group. There is a bijection

étale cohomology		finitely generated
sheaves on $\operatorname{Spec} k$	$\stackrel{\sim}{\longleftrightarrow}$	$\mathbb{Z}/\ell^m\mathbb{Z}$ -modules with
with $\mathbb{Z}/\ell^m\mathbb{Z}$ -coefficients		an action of $\operatorname{Gal}(\overline{k}/k)$
$\overline{\mathbf{A}}$ or \mathbf{C}		finite-dimensional continuous
\mathbb{Q}_{ℓ} -coefficients	\hookrightarrow	representations of $\operatorname{Gal}(\overline{k}/k)$
		on \mathbb{Q}_{ℓ} -vector spaces

Let $\mathcal{F} \in D^b_c(X, \overline{\mathbb{Q}}_{\ell})$ be a object in the bounded derived category of constructible $\overline{\mathbb{Q}}_{\ell}$ -sheaves on X. Then for any $x \in X(\mathbb{F}_{q^n})$, we get an inclusion

$$\operatorname{Spec} \mathbb{F}_{q^n} \stackrel{i}{\hookrightarrow} X,$$

and hence an object $i^* \mathcal{F} \in D^b_c(\operatorname{Spec} \mathbb{F}_q, \overline{\mathbb{Q}}_\ell)$. Applying the supertrace of Frobenius to this object results in an element of $\overline{\mathbb{Q}}_\ell$. In this way, we get a map

$$D^b_c(X, \overline{\mathbb{Q}}_\ell) \xrightarrow{f} \prod_{n \ge 1} \operatorname{Fun}(X(\mathbb{F}_{q^n}) \to \overline{\mathbb{Q}}_\ell).$$

If $\mathcal{F} \to \mathcal{G} \to \mathcal{F}' \xrightarrow{+1}$ is a distinguished triangle in $D_c^b(X, \overline{\mathbb{Q}}_\ell)$, then $f(\mathcal{F}) + f(\mathcal{F}') = f(\mathcal{G})$. In other words, f factors through the Grothendieck group:

$$\left[D_c^b(X,\overline{\mathbb{Q}}_\ell)\right] \xrightarrow{f} \prod_{n\geq 0} \operatorname{Fun}(X(\mathbb{F}_{q^n}),\overline{\mathbb{Q}}_\ell).$$

A consequence of the Chebotarev density theorem (Theorem 1.3) is that this map is *injective*; i.e. for a given q^n , the collection of all of the functions $f(\mathcal{F})$ completely determine the class of \mathcal{F} in the Grothendieck group. Now, Grothendieck tells us how to view this relationship.

Grothendieck's philosophy: Interesting functions $X(\mathbb{F}_{q^n}) \to \mathbb{C}, \overline{\mathbb{Q}}_{\ell}$, etc. should be shadows of interesting sheaves.

Lusztig provided us with an extraordinary example of a case where Grothendieck's philosophy is exactly true.

Example 16.5. (Lusztig's theory of character sheaves) There exists a set

$$\left\{\mathcal{F}_{\chi} \in D_c^b(\mathrm{GL}_n, \overline{\mathbb{Q}}_{\ell})\right\}$$

such that \mathcal{F}_{χ} yield all irreducible characters $\chi : \operatorname{GL}_n(\mathbb{F}_{q^n}) \to \mathbb{C}$ of all $\operatorname{GL}_n(\mathbb{F}_{q^n})$ "at once". (More precisely, for a given \mathbb{F}_{q^n} , one should only consider those character sheaves which are "defined over \mathbb{F}_q ", however we will not go into the details.)

16.2 The Hecke algebra, revisited

In the second half of this lecture, we will describe another example of Grothendieck's philosophy: the Hecke category. To start, we will recall the origin of the Hecke algebra.

Let G be a split reductive group over a finite field \mathbb{F}_q and $B \subset G$ a Borel subgroup. We can define a convolution algebra

$$H_{\mathbb{F}_q} = \left(\operatorname{Fun}_{B(\mathbb{F}_q) \times B(\mathbb{F}_q)}(G(\mathbb{F}_q), \mathbb{C}), * \right)$$

of complex-valued $B(\mathbb{F}_q)$ -biinvariant functions on $G(\mathbb{F}_q)$. A priori, the structure of this algebra seems to depend on q, but Iwahori found a presentation of $H_{\mathbb{F}_q}$ which depends almost only on the Weyl group. Let (W, S) be the Coxeter system associated to $B \subset G$. Iwahori's presentation has generators $\{T_s \mid s \in S\}$, and relations

$$T_s^2 = (q-1)T_s + q$$

$$T_sT_t \dots = T_sT_t \dots,$$

where the products on the second line are of $m_{st} = \operatorname{order}(st)$ generators. In Iwahori's presentation, the element T_s corresponds to the indicator function $\mathbb{1}_{BsB} \in \operatorname{Fun}_{B(\mathbb{F}_q) \times B(\mathbb{F}_q)}(G(\mathbb{F}_q), \mathbb{C})$. Iwahori's presentation demonstrated that the Hecke algebra $H_{\mathbb{F}_q}$ is "defined over $\mathbb{Z}[q]$ ".

The quadratic relation may look a little mysterious at first, but it has a natural geometric origin.

Origin of the quadratic relation: Take $G = \operatorname{SL}_2(\mathbb{F}_q)$ and $B = \left\{ \begin{pmatrix} * & * \\ 0 & * \end{pmatrix} \right\}$. Then on one hand,

 $\mathbb{1}_G * \mathbb{1}_G = \begin{array}{c} \text{pushforward of the constant function on } G \times_B G \\ \text{under the multiplication map } G \times_B G \xrightarrow{m} G \end{array}$

But on the other hand, there is a natural isomorphism

$$G \times_B G \xrightarrow{\sim} G/B \times G$$
$$(g,h) \mapsto (gB,gh)$$

and the multiplication map factors through this isomorphism:

Hence,

$$\mathbb{1}_G * \mathbb{1}_G = \begin{array}{l} \text{pushforward of constant function on} \\ G/B \times G \text{ under the projection to } G \\ = |G/B| \cdot \mathbb{1}_G \\ = (1+q)\mathbb{1}_G. \end{array}$$

From this, we deduce the quadratic relation: Since $\mathbb{1}_G = \mathbb{1}_{BsB} + \mathbb{1}_B = T_s + 1$, we have

$$T_s^2 + 2T_s + 1 = (T_s + 1)^2$$

= $(q + 1)(T_s + 1)$
= $(q + 1)T_s + q + 1$
= $(q - 1)T_s + q + 2T_s + 1$


Hence $T_s^2 = (q - 1)T_s + q$.

The geometric origin of the Hecke algebra suggests a categorification via Grothendieck's philosophy. Let G be a split reductive group over \mathbb{F}_q , and $B \subset G$ a Borel subgroup. Define the (first incarnation of) the **Hecke category** to be

$$\mathcal{H} := D^b_{B \times B}(G, \overline{\mathbb{Q}}_\ell),$$

the $B \times B$ -equivariant derived category of étale $\overline{\mathbb{Q}}_{\ell}$ -sheaves, in the sense of Bernstein-Lunts. We won't describe the precise construction of this category, but in its first approximation¹⁵, we can consider objects in $D^b_{B \times B}(G, \overline{\mathbb{Q}}_{\ell})$ to be étale sheaves which are constructible for $B \times B$ -orbits.

The convolution of functions in $H_{\mathbb{F}_q}$ can be upgraded to a **convolution product** on sheaves. Let $\mathcal{F}, \mathcal{G} \in D^b_{B \times B}(G)$. We have maps



Then if $\mathcal{F}, \mathcal{G} \in D^b_{B \times B}(G, \overline{\mathbb{Q}}_\ell)$, we define

$$\mathcal{F} * \mathcal{G} := \operatorname{mult}_*(\widetilde{\mathcal{FG}}) \in D^b_{B \times B}(G, \overline{\mathbb{Q}}_\ell),$$

where $\widetilde{\mathcal{FG}} \in D^b_{B \times B}(G \times_B G, \overline{\mathbb{Q}}_\ell)$ corresponds to $\operatorname{res}_{B \times B \times B \times B}^{B \times B \times B}(p_1^* \mathcal{F} \otimes p_2^* \mathcal{G})$ under the equivalence

$$D^b_{B \times B \times B \times B}(G \times G, \overline{\mathbb{Q}}_{\ell}) \xrightarrow{\sim} D^b_{B \times B \times B}(G \times_B G, \overline{\mathbb{Q}}_{\ell}).$$

Example 16.6. Let $k = \overline{\mathbb{Q}}_{\ell}$, and $G = SL_2$. Then

$$\underline{k}_{\mathrm{SL}_2} * \underline{k}_{\mathrm{SL}_2} = \mathrm{mult}_*(\underline{k}_{\mathrm{SL}_2 \times_B \mathrm{SL}_2}) = H^*(\mathbb{P}^1) \otimes \underline{k}_{\mathrm{SL}_2}.$$

This is the categorified version of the Hecke algebra equality $\mathbb{1}_G * \mathbb{1}_G = (1+q)\mathbb{1}_G$ that we discussed earlier!

¹⁵Be careful! This approximation is just an approximation and can get you in trouble if you take it too literally (see Lecture 24).

16.3 Perverse sheaves on $\mathbb{P}^1\mathbb{C}$

The time has come for an interlude. For the rest of this lecture we will shift gears and consider the variety $\mathbb{P}^1\mathbb{C}$ with the stratification Λ given by


$$\mathbb{P}^1\mathbb{C} = \{0\} \sqcup \mathbb{C}_\infty.$$

Denote by $\{0\} \xrightarrow{i} \mathbb{P}^1 \mathbb{C} \xleftarrow{j} \mathbb{C}_{\infty}$ the natural inclusions. It is very important in what is coming to understand perverse sheaves on $\mathbb{P}^1 \mathbb{C}$ with the stratification Λ , so we will discuss this (and generalizations) in the upcoming lectures. So what are the perverse sheaves on $\mathbb{P}^1 \mathbb{C}$?

Here's an algebraic answer (c.f. Emily's talk on nearby cycles in the Informal Friday Seminar¹⁶):

$$\operatorname{Perv}_{\Lambda}(\mathbb{P}^{1}\mathbb{C},k) \simeq \left\{ V_{1} \xleftarrow{c}{v} V_{0} \mid v \circ c = 0 \right\},\$$

where V_i are finite-dimensional k vector spaces representing the nearby cycles at 0 (the vector space V_1) and the vanishing cycles at 0 (the vector space V_0). From this perspective, we can see that there are five indecomposible objects:



¹⁶All IFS talk notes can be found on the IFS website https://sites.google.com/view/ifssydney/home



In the next lecture we will go into more detail and describe perverse sheaves on general curves.

Lecture 17: Constructible and perverse sheaves on curves

Today we continue our interlude of the previous lecture and discuss perverse sheaves on curves. We are working toward understanding Beilinson glueing on curves, which will be the main topic of next week's lecture.

17.1 Constructible sheaves

Let k be a field, fixed throughout, and let X/\mathbb{C} be a variety, viewed with the classical topology. Let V be a k-vector space. We have the notion of a **constant sheaf** V_X with values in V:


 $V_X(U) := \{ f : U \to V \text{ continuous} \},\$

where V is viewed with the discrete topology. This leads to the notion of a **local system**, which is a locally constant sheaf with finite-dimensional stalks.

Theorem 17.1. If X is connected, there is a bijection

$$\left\{\begin{array}{c} \text{local systems} \\ \text{on } X \end{array}\right\} \stackrel{\sim}{\leftrightarrow} \operatorname{Rep}(\pi_1(X, x)).$$

Remark 17.2. The difference between local systems and vector bundles is captured with the following picture.




Here the pink lines are intended to denote the sections of our local systems/vector bundles. Vector bundles with flat connections are equivalent to local systems. However, local systems form an abelian category, whereas vector bundles do not.

The notion of a local system leads us to the notion to a **constructible sheaf**: A sheaf \mathcal{F} on X is constructible if there exists a stratification $X = \bigsqcup_{\lambda \in \Lambda} X_{\lambda}$ of X by a finite number of subvarieties X_{λ} such that $\mathcal{F}|_{X_{\lambda}}$ is a local system for all $\lambda \in \Lambda$. Finally, the notion of a constructible sheaf leads to the notion of the **bounded derived category of constructible sheaves**:

$$D_c^b(X,k) := \left\{ \mathcal{F} \middle| \begin{array}{c} \mathcal{H}^i(\mathcal{F}) \text{ is constructible for all } i, \text{ and} \\ \mathcal{H}^i(\mathcal{F}) = 0 \text{ for } |i| >> 0 \end{array} \right\} \subset D^b \left(\begin{array}{c} \text{sheaves of } k \text{-vector} \\ \text{spaces on } X \end{array} \right)$$

From now on, assume that X is a connected, smooth curve.




Choose a set of points $\{x_1, \ldots, x_m\}$, and let U denote their complement in X. Fix the stratification Λ :

$$X = U \sqcup \{x_1\} \sqcup \cdots \sqcup \{x_m\}.$$

Theorem 17.3. There is an equivalence (a fun exercise if the reader is tempted!)

$$\left\{\begin{array}{l} \Lambda\text{-constructible}\\ \text{sheaves on }X\end{array}\right\} \xrightarrow{\sim} \left\{\begin{array}{l} \mathcal{L} \text{ local system on }U,\\ V_1,\ldots,V_m \text{ finite-dimensional vector spaces}\\ \max\phi_i:V_i\to (\mathcal{L}_{n_i})^{\mu_i}\end{array}\right\}.$$

Here $n_i \in X$ is a "nearby point" to x_i , and \mathcal{L}_{n_i} denotes the stalk, which carries a monodromy operator μ_i given by a small loop around x_i (see the diagram below). Note that the stalk depends on the choice of nearby point n_i , but the invariants in the stalk do not!



17.2 Perverse sheaves

One way to visualize an object $\mathcal{F} \in D^b_c(X)$ is via its table of stalks:

	i-1	i	i+1
U	• • •	$\mathcal{H}^i(\mathcal{F} _U)$	•••
x_1	•••	$\mathcal{H}^i(\mathcal{F}_{x_i})$	•••
x_2	•••	$\mathcal{H}^i(\mathcal{F}_{x_2})$	•••
:		•	

We will use these tables frequently in the remainder of this lecture.

The category of **perverse sheaves**

$$\operatorname{Perv}_{\Lambda}(X) \subset D^b_c(X)$$

is defined as the heart of the *t*-structure

$$D_{\Lambda}^{\leq 0} := \left\{ \mathcal{F} \left| \begin{array}{c} \mathcal{H}^{i}(\mathcal{F}|_{U}) = 0 \text{ for } i \geq 0\\ \mathcal{H}^{i}(\mathcal{F}|_{x_{j}}) = 0 \text{ for } i > 0 \end{array} \right\}, \quad D_{\Lambda}^{\geq 0} = \mathbb{D}(D_{\Lambda}^{\leq 0}).$$

Here \mathbb{D} denotes **Verdier duality**. We have two possibilities for what the table of stalks can look like for a perverse sheaf in $\text{Perv}_{\Lambda}(X)$:

A general strategy when trying to understand an abelian category is to try to produce and study exact functors from that category to a well-understood category (like Vect or representations of a group). For perverse sheaves, this strategy proves to be somewhat complicated, and brings nearby and vanishing cycles into our world.

Theorem 17.4. (to be explained)

$$\operatorname{Perv}_{\Lambda}(X) \stackrel{\sim}{\longleftrightarrow} \left\{ \begin{array}{c} \mathcal{L} \text{ local system on } U, \\ V_1, \dots, V_m \text{ finite-dimensional vector spaces ("vanishing cycles"),} \\ \max V_i \stackrel{v}{\xleftarrow{u}} \mathcal{L}_{n_i} \quad \text{s.t. } v \circ u = id - \mu_i \end{array} \right\}$$

Remark 17.5. The category $\operatorname{Perv}_{\Lambda}(X)$ is Verdier self-dual. The category of Λ -constructible sheaves is not:

$$\mathbb{D}(\Lambda\text{-constructible sheaves}) \simeq \left\{ \phi_i : \mathcal{L}_{n_i}^{\mu_i} \to V_i \right\}$$

Example 17.6. Here are some examples of perverse sheaves.

- The constant sheaf k_{x_i} on $\{x_j\}$ is perverse.
- For a local system \mathcal{L} on X, $\mathcal{L}[1]$ is perverse. (This follows because $\mathbb{D}\mathcal{L}[1] \cong \mathcal{L}^{\vee}[1]$, where \mathcal{L}^{\vee} denotes the dual local system.)
- Let $j: U \hookrightarrow X$ and \mathcal{L} a local system on U. Then we claim that $j_!\mathcal{L}[1]$ and $j_*\mathcal{L}[1]$ are both perverse. Because $j_!$ is extension by zero, computing the stalks of $j_!\mathcal{L}[1]$ is easy:

Computing the stalks of the direct image $j_* := Rj_*$ is trickier. Le $x \in X$ be a point. Then

$$H^{i}((j_{*}\mathcal{L})_{x}) = \lim_{\stackrel{\leftarrow}{B(x,\epsilon)}} H^{i}(B(x,\epsilon) \cap U, \mathcal{L})$$
$$= \begin{cases} \mathcal{L}_{x} & \text{if } x \in U, \\ H^{i}(B(x,\epsilon) \setminus \{x\}, \mathcal{L}) & \text{if } x \notin U. \end{cases}$$

How can we compute the cohomology $H^i(B(x,\epsilon) \setminus \{x\}, \mathcal{L})$? The space $B(x,\epsilon) \setminus \{x\}$ is homotopic to S^1 . We can compute the cohomology of S^1 with coefficients in \mathcal{L} using the two-term complex

$$V \xrightarrow{id-\mu} V.$$

Hence $H^0(S^1, \mathcal{L}) = V^{\mu}$, the invariants of $\mu \circlearrowright V$, and $H^1(S^1, \mathcal{L}) = V_{\mu}$, the coinvariants. (Alternatively, one can see that $H^0(S^1, \mathcal{L}) = V^{\mu}$ directly, and conclude that $H^1(S^1, \mathcal{L}) = V_{\mu}$ by Poincaré duality.) Hence, the table of stalks of $j_*\mathcal{L}[1]$ is

Because $\mathbb{D}(j_*\mathcal{L}) = j_!\mathbb{D}(\mathcal{L})$, we can conclude from these computations that $j_*\mathcal{L}[1]$ and $j_!\mathcal{L}[1]$ are both perverse. In other words, j_* and $j_!$ are exact for the perverse *t*-structure.

Next we turn our attention to classifying the simple objects in $\operatorname{Perv}_{\Lambda}(X)$. Recall that there is a natural map

$$j_! \rightarrow j_*$$
.

Given \mathcal{L} a local system on \mathcal{U} , define

$$IC(X, \mathcal{L}) := j_{!*}(\mathcal{L}) := \operatorname{Im}(j_!\mathcal{L}[1] \to j_*\mathcal{L}[1]).$$

This is not obvious, but by a construction of Deligne, we have

$$IC(X, \mathcal{L}) = \tau_{\leq -1}(j_*\mathcal{L}[1])$$

where $\tau_{\leq -1}$ is the truncation functor in the standard (not perverse) *t*-structure. Hence the table of stalks for $IC(X, \mathcal{L})$ is

	-2	-1	0	1	
U	0	\mathcal{L}	0	0.	•
x_i	0	V^{μ}	0	0	

Remark 17.7. We see from the remarks above that when X is a curve, all IC sheaves are shifts of actual constructible sheaves. This is not the case in general!

Example 17.8. Consider a local system \mathcal{L} on U such that the monodromy μ does not have 1 as an eigenvalue. Then $id - \mu$ is invertible. Hence $V_{\mu} = V^{\mu} = 0$. Recall that the general form of the tables of stalks of $j_{!}\mathcal{L}[1]$ and $j_{*}\mathcal{L}[1]$ are

So the fact that $V^{\mu} = V_{\mu} = 0$ implies that the natural map $j_{!}\mathcal{L}[1] \to j_{*}\mathcal{L}[1]$ is an isomorphism. We conclude that in this case

$$IC(X,\mathcal{L}) = j_!\mathcal{L}[1] = j_*\mathcal{L}[1].$$

Theorem 17.9. The category $\operatorname{Perv}_{\Lambda}(X)$ is a finite-length abelian category with simple objects

$$\{IC(X,\mathcal{L}) \mid \mathcal{L}|_U \text{ is irreducible}\} \cup \{i_*k \mid i : \{x_i\} \hookrightarrow X\}.$$

For any irreducible local system \mathcal{L} on U, we have the distinguished triangle

$$\tau_{\leq -1}(j_*\mathcal{L}[1]) \to j_*\mathcal{L}[1] \to \tau_{\geq 0}(j_*\mathcal{L}[1]) \xrightarrow{+1}$$
$$=$$
$$IC(X, \mathcal{L}) \to j_*\mathcal{L}[1] \to (V_{\mu})_0 \xrightarrow{+1}$$

in $D_c^b(X, k)$. There is something distinctive about this triangle: all objects are perverse sheaves! Hence, this is an exact sequence and we've found our first composition series. We can draw this composition series with an "egg diagram:"

$$j_{*}\mathcal{L}[1] = \underbrace{\left(\bigvee_{\mathcal{M}} \right)_{o}}_{\mathbb{I}((\mathbf{x}, \mathbf{j}))}$$
(17.1)

Example 17.10. Here is a special case of the construction above. Let D be a disc and

$$D^{\times} \stackrel{j}{\hookrightarrow} D \stackrel{i}{\longleftarrow} \{0\}$$

the natural inclusions. Then the composition series of $j_*k_{D^{\times}}[1]$ is

$$j_{*}k_{D^{\times}}[1] = \underbrace{\operatorname{I}(\mathbf{b}^{\mathsf{c}}\mathbf{k}_{\mathbf{b}})}_{\operatorname{I}(\mathbf{b})}$$

Example 17.11. Let

$$U \stackrel{j}{\hookrightarrow} X \stackrel{i}{\hookleftarrow} Z := X \backslash U$$

be the natural inclusions. Then in $D^b_c(X,k)$ we have the distinguished triangle

$$j_! j^! k_X \to k_X \to i_* i^* k_X \xrightarrow{+1} . \tag{17.2}$$

By turning triangles, we obtain

$$i_*k_Z \to j_!k_U[1] \to k_X[1] = IC(X) \xrightarrow{+1}$$

All objects in this triangle are perverse sheaves, so this is a composition series! Hence we have the following egg:

$$j_{*}k_{D^{\times}}[1] = \underbrace{\mathbf{I}(\mathbf{x}) \mathbf{x}_{\mathbf{x}}\mathbf{x}_{\mathbf{y}}}_{\mathbf{k}_{\mathbf{z}}}.$$

We could also have obtained this by dualizing (17.1).

Remark 17.12. Constructible sheaves on X are not necessarily finite length (see (17.2)). However, perverse sheaves are. This is one of the reasons why we are so fond of them.

Before moving on to glueing, we will do one more fun calculation. We return to the local case:


$$D^{\times} \stackrel{\jmath}{\hookrightarrow} D \stackrel{i}{\longleftrightarrow} \{0\}$$

Let \mathcal{L}_n be a local system on D^{\times} with monodromy given by a single Jordan block of size n:

$$J_n = \begin{pmatrix} 1 & 1 & & \\ & 1 & 1 & & \\ & & 1 & \ddots & \\ & & & \ddots & 1 \\ & & & & & 1 \end{pmatrix}$$

Our goal is to describe $j_{!*}\mathcal{L}_n[1] = \operatorname{Im}(j_!\mathcal{L}_n[1] \to j_*\mathcal{L}_n[1])$ in this setting. Recall that the functors $j_!$ and j_* are exact because j is affine, and we have the building blocks

In local systems on D^{\times} , we have



Remark 17.13. In this example, we see that the functor $j_{!*}$ preserves injections and surjections, but is NOT exact.

17.3 Nearby and vanishing cycles

Let D be the disc, D^{\times} the punctured disc, and $\widetilde{D^{\times}}$ its universal cover. Let

$$\{0\} \stackrel{i}{\hookrightarrow} D \stackrel{j}{\longleftrightarrow} D^{\times} \stackrel{p}{\leftarrow} \widetilde{D^{\times}}$$

be the natural maps.

Motivation: We want to define the "stalk" of a perverse sheaf at singular points.

Recall the **nearby cycles** functor ψ_f (c.f. Emily's Oct 18, 2019 IFS talk or Laurentiu's Feb 21/28, 2020 IFS series): For a local system \mathcal{F} on D^{\times} ,

$$\begin{split} \psi_f(\mathcal{F}) &= i^* j_* p_* p^* \mathcal{F} \\ &= i^* j_* p_* (p^* \mathcal{F} \otimes k_{\widetilde{D^{\times}}}) \\ &= i^* j_* (\mathcal{F} \otimes p_* k_{\widetilde{D^{\times}}}). \end{split}$$

Here the third equality follows from the projection formula. The sheaf $p_*k_{\widetilde{D^{\times}}}$ is the "universal local system on $D^{\times "}$. It is an infinite-dimensional local system with stalk $k[x^{\pm 1}]$ and monodromy x.

Recall that the monodromy $\mu \circlearrowright \psi_f(\mathcal{F})$, so we get a decomposition into generalized eigenspaces:

$$\psi_f(\mathcal{F}) = \bigoplus \psi_f^\lambda(\mathcal{F})$$

By far the most important of these eigenspaces corresponds to $\lambda = 1$, the "unipotent nearby cycles". Indeed, from this, we can recover all others:

$$\psi_f^{\lambda}(\mathcal{F}) = \psi_f^1(\mathcal{F} \otimes k_{\lambda^{-1}}).$$

From now on, set

$$\psi(\mathcal{F}) := \psi_f^1(\mathcal{F}).$$

Let \mathcal{L}_{unip} be the **universal unipotent local system**. This is the local system with stalk k[[u]] at 1 and monodromy given by multiplication by 1 + u. In other words,


$$\mathcal{L}_{unip} = \lim_{n \to \infty} \mathcal{L}_n,$$

where \mathcal{L}_n is our local system from earlier with stalk $k[u]/(u^n)$ and monodromy multiplication by 1 + u. (Think: "completion of the augmentation ideal" should give the same answer.)

Idea:

1. We should think that

$$\psi(\mathcal{F})$$
 "=" $\lim_{\longleftarrow} (i^*(j_*(\mathcal{F} \otimes \mathcal{L}_n)))$



We will see in the next lecture that after tensoring with the local system given by a big Jordan block, we can recover V and its monodromy via taking global sections. This appears to be a basic idea behind Beilinson's construction.

Hence it makes sense to define

$$\psi(\mathcal{F}) = H^1(D_{\epsilon}^{\times}, \mathcal{F} \otimes \mathcal{L}_n)$$

for n large.

Lecture 18: Beilinson gluing on curves

Throughout this lecture, we will work in the following setting. Let D be the disc, D^{\times} the punctured disc, and

$$D^{\times} \stackrel{\jmath}{\hookrightarrow} D \stackrel{\imath}{\hookleftarrow} \{0\}$$

the natural inclusions.

Our goal for this lecture is to explain the proof of Beilinson gluing¹⁷. We'll pick up where we left off last week.

18.1 A fact about unipotent monodromy

Let

$$e = \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}, \quad h = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}, \quad f = \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix}$$

be the standard basis for $\mathfrak{sl}(2,\mathbb{C}) = \mathbb{C}f \oplus \mathbb{C}h \oplus \mathbb{C}e$. Clebsch–Gordan tells us about how the tensor product of two irreducible finite-dimensional $\mathfrak{sl}(2,\mathbb{C})$ -modules decomposes:

Theorem 18.1. (Clebsch–Gordan) Let L_n, L_m be irreducible finite-dimensional $\mathfrak{sl}(2, \mathbb{C})$ modules of highest weight n, m, respectively. Then

$$L_n \otimes L_m = L_{|n-m|} \oplus \cdots \oplus L_{n+m-2} \oplus L_{n+m}.$$

A consequence of this theorem is that for n large, there are m + 1 summands, so

$$(L_n \otimes L_m)^e = \frac{\text{highest weight vectors}}{\text{in above sum}} = \mathbb{C}^{m+1} \simeq L_m$$

(Here the superscript denotes Lie algebra invariants; i.e. the kernel of the action

$$e \cdot v \otimes w = ev \otimes w + v \otimes ew$$

of e on $L_n \otimes L_m$.) Moreover, this is more than just an isomorphism of vector spaces: the action of $-e \otimes 1$ on $(L_n \otimes L_m)^e$ aligns with the action of e on L_m . Hence (be exponentiating) we have the following corollary to Clebsch–Gordan's theorem.

Corollary 18.2. Let V be a finite-dimensional vector space and ϕ a unipotent endomorphism of V. Let J_n be a (n + 1)-dimensional vector space and

$$\phi' = \begin{pmatrix} 1 & 1 & & \\ & 1 & \ddots & \\ & & \ddots & 1 \\ & & & & 1 \end{pmatrix}$$

¹⁷In addition to Beilinson's original paper [Bei87], the notes [Mor18] of Sophie Morel are an excellent reference for this material.

a Jordan block of size n + 1. For n large,

$$(V \otimes J_n)^{\phi \otimes \phi'} \simeq V,$$

and this isomorphism is equivariant with respect to the action of $1 \otimes (\phi')^{-1}$ on the LHS and ϕ on the RHS.

Hence by tensoring with a Jordan block and taking invariants, we can recover the whole vector space and its monodromy action.

Remark 18.3. (Aside for the Lie theorists) Let \mathfrak{g}/\mathbb{C} be a semisimple Lie algebra, and V a finite-dimensional \mathfrak{g} -module. For λ sufficiently dominant,

$$\Delta_{\lambda} \otimes V \simeq \bigoplus_{\gamma_i \in \Gamma} \Delta_{\lambda + \gamma_i},$$

where Δ_{μ} is the Verma module of highest weight μ and Γ is the multiset of weights of V. Moreover, there is an isomorphism of \mathfrak{n}_+ -modules:

$$(\Delta_{\lambda} \otimes V)^{\mathfrak{b}} \simeq V.$$

18.2 The unipotent vanishing cycles functor

Now we return to the world of perverse sheaves. A perverse sheaf $\mathcal{M} \in \operatorname{Perv}(D^{\times})$ is the same thing as a vector space V and monodromy endomorphism $\mu \circlearrowright V$. Let \mathcal{L}_n be the local system with stalk $k[x]/(x^n)$ and monodromy $\phi = 1 + x$, as in Lecture 17. In light of Corollary 18.2 and the third bullet point in Example 17.6, for n large enough, we could define a functor ψ by

$$\psi(\mathcal{M}) = H^0(j_*(\mathcal{M} \otimes \mathcal{L}_n)) = (V \otimes J_{n-1})^{\mu \otimes \phi} = V.$$

This will end up being our definition which allows us to see certain features in the arguments below. The key lemma of today's lecture tells us that this definition stabilizes for large n.

Lemma 18.4. Let $\mathcal{M} \in \text{Perv}(D^{\times})$. For a fixed m, the kernel and cokernel of

$$j_!(\mathcal{M}\otimes\mathcal{L}_n)\xrightarrow{x^m} j_*(\mathcal{M}\otimes\mathcal{L}_n)$$

stabilize for large n, and are canonically isomorphic.


Proof. We will prove the lemma when \mathcal{M} has rank 1 (i.e. \mathcal{M} corresponds to the vector space k with monodromy given by multiplication by $\lambda \in k^{\times}$.)

Case 1: monodromy $\lambda \neq 1$. Then

$$j_!(\mathcal{M}\otimes\mathcal{L}_n)=j_*(\mathcal{M}\otimes\mathcal{L}_n).$$

(See Example 17.8.) The theorem follows in this case because the kernel and cokernel of $\mathcal{L}_n \xrightarrow{x^m} \mathcal{L}_n$ stabilize for m fixed, n large and $j_{!}, j_{*}$ are exact.

Case 2: monodromy $\lambda = 1$; i.e. $\mathcal{M} = k_{D^{\times}}[1]$ is the trivial local system. We will illustrate what is happening in this case through an example. Let m = 2 and n = 3. We can see what the maps $\mathcal{L}_3 \xrightarrow{x^2} \mathcal{L}_3$, $j_! \mathcal{L}_3 \xrightarrow{x^2} j_! \mathcal{L}_3$ and $j_! \mathcal{L}_3 \xrightarrow{x^2} j_* \mathcal{L}_3$ are doing on egg diagrams:



As n gets larger, the eggs in the illustration above get longer and the image of $j_{!*}$ gets bigger, but the kernel and cokernel stay the same size.

Remark 18.5. What Beilinson actually proves is that

$$\lim_{\leftrightarrow} j_!(\mathcal{M} \otimes \mathcal{L}_n) \simeq \lim_{\leftrightarrow} j_*(\mathcal{M} \otimes \mathcal{L}_n),$$


where ' \lim_{\leftrightarrow} ' is appropriately defined. We'll sum this up with the slogan "middles agree". We already saw this happening above, and will see it happening again in our next example.

Example 18.6. Here is a 2-dimensional example. Let

$$\mathbb{C}^2 \xrightarrow[xy]{f} \mathbb{C},$$

and consider $\operatorname{Perv}_{\Lambda}(\mathbb{C}^2)$, where Λ is the stratification via coordinate hyperplanes and their


complement U.



Let $\mathcal{M} = k_U[2]$, and denote by $IC_1 = IC_{\{y=0\}}$, $IC_2 = IC_{\{x=0\}}$, and $IC_0 = IC_{\{x=y=0\}}$. The composition series of $j_!k_U[2]$ is given by the following egg:

$$j_!k_U[2] = \underbrace{\begin{array}{c} \mathbf{I}(\mathbf{x}) \\ \mathbf{I}(\mathbf{x})$$

Then the key Lemma can be seen on egg diagrams:




Definition 18.7. (Beilinson) Let $X \xrightarrow{f} D$, $U = f^{-1}(D^{\times})$, and $Z = f^{-1}(0)$. For a perverse sheaf $\mathcal{M} \in \operatorname{Perv}(U)$ and *n* large, define functors $\psi_f, \Xi_f : \operatorname{Perv}(U) \to \operatorname{Perv}(X)$ by


• $\psi_f(\mathcal{M}) = \ker(x^0) \simeq \operatorname{coker}(x^0)$ "unipotent nearby cycles"

• $\Xi_f(\mathcal{M}) = \ker(x^1) \simeq \operatorname{coker}(x^1)$ "maximal extension"

The perverse sheaf $\psi_f(\mathcal{M})$ is supported on Z and $\Xi_f(\mathcal{M})$ is supported everywhere. Example 18.8. Let $\mathcal{M} = k_U[1]$, then we can see the images of these functors in eggs:



Example 18.9. In our 2-dimensional example, Example 18.6 we have:



Lemma 18.10. The functors ψ_f and Ξ_f are exact, and we have functorial short exact sequences

- 1. $j_! \hookrightarrow \Xi_f \twoheadrightarrow \psi_f$
- 2. $\psi_f \hookrightarrow \Xi_f \twoheadrightarrow j_*$


Moreover, the canonical map

$$\psi_f \to \Xi_f \to \psi_f$$

agrees with monodromy -1.

Proof. See Geordie's handwritten notes on the course website.


Definition 18.11. Let $\mathcal{M} \in \operatorname{Perv}(D)$. The unipotent vanishing cycles functor is defined by



All maps in the diagram above are the canonical ones.


- **Remark 18.12.** 1. The map $j_! \to \Xi_f$ is injective, and the map $\Xi_f \to j_*$ is surjective, so the cocomplex above only has cohomology in degree 0, and hence ϕ_f is an exact functor on Perv(D).
 - 2. x induces a monodromy endomorphism of Ξ_f, ψ_f, ϕ_f .

Example 18.13. Let X = D and $\mathcal{M} = j_! k_{D^{\times}}[1]$. To compute $\phi_f(\mathcal{M})$, we need to compute the zeroeth cohomology




The image of the first map is


The kernel of the second map is



Notice that because of the sign in the second map, the image will always be contained in the kernel. Hence $\phi_f(\mathcal{M})$ is equal to




Example 18.14. If $\mathcal{M} = k_D[1]$, then using the same method as above, we compute



Here, the kernel and image are equal:

$$\operatorname{im} = \left(\begin{array}{c} \mathbb{I}(\mathbf{0}) \\ \mathbb{I}(\mathbf{0}) \\ \mathbb{I}(\mathbf{0}) \end{array} \right) = \operatorname{ker} \mathcal{Y}$$

Notice that the extra piece of the kernel in the previous example is now a subsheaf of the image:




Hence $\phi_f(\mathcal{M}) = 0.$

Exercise 18.15. Show that

$$\phi_f(\Xi_f(k_{D^{\times}}[1])) = IC_0^{\oplus 2},$$


and show that the monodromy endomorphism is not trivial.

Example 18.16. Let $\mathcal{M} = IC_X$, for $X = \mathbb{C}^2$, and $\mathbb{C}^2 \xrightarrow{f}_{xy} \mathbb{C}$ as earlier. Then



18.3 Gluing

With this machinery, Beilinson shows us how we can glue perverse sheaves. To state the theorem we return to a slightly more general setting.



Theorem 18.17. (Beilinson)

$$\operatorname{Perv}(X) \simeq \left\{ \begin{array}{cc} \mathcal{M}_U \in \operatorname{Perv}(U), \\ \mathcal{M}_Z \in \operatorname{Perv}(Z) \end{array} + \begin{array}{c} \psi_f(\mathcal{M}_U) \xrightarrow{\operatorname{monodromy} -1} \psi_f(\mathcal{M}_U) \\ \mathcal{M}_Z \in \operatorname{Perv}(Z) \end{array} + \begin{array}{c} \psi_f(\mathcal{M}_U) \xrightarrow{\operatorname{monodromy} -1} \psi_f(\mathcal{M}_U) \\ \mathcal{M}_Z & \end{array} \right\}$$


"pieces" "glue"

Remark 18.18. Beilinson's theorem glues perverse sheaves on an open set to perverse sheaves on a closed set. Gluing perverse sheaves on two open sets is much easier because perverse sheaves form a stack.

Example 18.19. Returning to our setting of $X = D = D^{\times} \sqcup \{0\}$, a perverse sheaf on D^{\times} (resp. $\{0\}$) is the same thing as a vector space with automorphism $\mu \circlearrowright V$ (resp. a vector space W). Hence

$$\begin{array}{l} \text{Perv} \\ \text{constructible} \\ \text{w.r.t.} \ D^{\times}, \{0\} \end{array} \left(D \right) = \left\{ \begin{array}{c} V \circlearrowleft \mu \text{ invertible,} \\ W \text{ vector space} \end{array} + \begin{array}{c} V \xrightarrow{\mu - id} \\ & & \\ & & \\ \end{array} \right\} \cdot \\ W \end{array} \right\}.$$

Let us explain why Beilinson gluing holds. We start with a general definition. Let \mathcal{A} be an abelian category. A **diad** in \mathcal{A} is




These form a category $Diads(\mathcal{A})$ in an obvious way.

Given a diad, we can associate a complex

$$Q^{\cdot} = C_{-} \xrightarrow{(\alpha_{-},\beta_{-})} A \oplus B \xrightarrow{(\alpha_{+},-\beta_{+})} C_{+},$$


and another diad



Lemma 18.20. $r^2 = id$

Proof. We will take this lemma as a black box.

Given this,



The second equality is obtained via applying our equivalence r on diads.

Lecture 19: The derived category of perverse sheaves

19.1 Overview of Beilinson gluing

Recall the setting of the last lecture: Let X/\mathbb{C} be a variety with the metric topology and $P_X := \text{Perv}(X, k)$. Beilinson described for us how to glue perverse sheaves on X. More specifically, we have the following set-up:

$$Z \xrightarrow{i} X \xleftarrow{j} U$$

$$\downarrow \qquad \qquad \downarrow^{f} \qquad \qquad \downarrow$$

$$\{0\} \longleftrightarrow \mathbb{A}^{1} \longleftrightarrow \mathbb{A}^{1} \setminus \{0\}$$

We can move between the bounded derived categories of constructible sheaves on X, Z, and U by pushing and pulling:

The functors $i_* = i_!, j^* = j^!$ preserve perverse sheaves, and if j is affine, then $j_*, j_!$ are exact and also preserve perverse sheaves. However, $i^*, i^!$ do not!

Last lecture, we constructed two other exact functors

 $x \circlearrowright \psi_f : P_U \to P_Z$ "unipotent nearby cycles" $x \circlearrowright \Xi_f : P_U \to P_X$ "maximal extension"


which each have a "monodromy" x. These functors fit into short exact sequences

$$0 \to j_! \mathcal{M} \to \Xi_f \mathcal{M} \to \psi_f \mathcal{M} \to 0$$
$$0 \to \psi_f \mathcal{M} \to \Xi_f \mathcal{M} \to j_* \mathcal{M} \to 0$$

for any $\mathcal{M} \in P_U$. We then used ψ_f and Ξ_f to construct another exact functor

 $x \circlearrowright \phi_f : P_X \to P_Z$ "unipotent vanishing cycles"

which we should think about as the "stalk of a perverse sheaf along Z". It was defined as




for $M \in P_X$.

Easy (and useful) fact: For $\mathcal{M} \in P_Z$,

$$\phi_f(i_*\mathcal{M})\simeq \mathcal{M}.$$

The main result of last lecture was **Beilinson gluing**, which gave an equivalence of categories:



19.2 A theorem of Beilinson

The goal of today's lecture is to use the tools above to prove the following theorem.

Theorem 19.1. $D^{b}(P_{X}) \simeq D^{b}_{c}(X)$.

This result is a beautiful example of a bigger philosophy. Often we have a triangulated category we wish to understand, and within it a collection of well-behaved objects which we understand better than the rest. Then we might hope to reconstruct the entire triangulated category from our special collection of objects. In many situations we cannot, but this is an example where we can. In the poetry of Beilinson, "the niche D where P_X dwells may be recovered from P_X ".

Recall: Let Λ be a stratification of X. Associated to this stratification is the category $\operatorname{Perv}_{\Lambda}(X)$ of Λ -constructible perverse sheaves on X. Our category P_X is the limit of such categories:

$$P_X = \operatorname{Perv}(X) = \lim_{\stackrel{\rightarrow}{\Lambda}} \operatorname{Perv}_{\Lambda}(X).$$

Remark 19.2. If Λ is a fixed stratification, usually

$$D^{b}(\operatorname{Perv}_{\Lambda}(X)) \not\simeq D^{b}_{\Lambda}(X).$$

Example 19.3. Let $X = \mathbb{P}^1 \mathbb{C}$ with the trivial stratification Λ . Then

$$\operatorname{Perv}_{\Lambda}(X) = \operatorname{local systems on } X \simeq \operatorname{Vect}_{k}^{f.d.}$$

However,

$$D^b(\operatorname{Vect}_k^{f.d.}) \not\simeq D^b_\Lambda(X)$$

because $\operatorname{Ext}^{i}(k_{X}, k_{X}) = H^{i}(X) \neq 0$ for i = 2, but the category $\operatorname{Vect}_{k}^{f.d.}$ is semisimple, so there are no higher exts.

19.3 $K(\pi, 1)$ spaces

We see in this example that differences in Ext groups prevented us from obtaining our desired derived equivalence. This illustrates a more general phenomenon which we will examine now.

Definition 19.4. A nice, path connected space X with base point $x \in X$ is $K(\pi, 1)$ if

1.
$$\pi_i(X) = 1$$
 for $i > 1$, and

2.
$$\pi_1(X) = \pi$$
.

Equivalently, X is $K(\pi, 1)$ if its universal cover is contractible.

Exercise 19.5. Show (easier) that

$$\operatorname{Rep}^{f.d.} k\pi \xrightarrow{\sim} k\text{-local systems on } X$$
$$V \mapsto \mathcal{L}_V$$

and (harder) that

$$\operatorname{Ext}^{i}(V, V') \simeq \operatorname{Ext}^{i}(\mathcal{L}_{V}, \mathcal{L}_{V'}).$$

Lemma 19.6. Let $F : \mathcal{D}_1 \to \mathcal{D}_2$ be a triangulated functor of triangulated categories. Assume that $\mathcal{D}_1, \mathcal{D}_2$ have *t*-structures with hearts $\mathcal{C}_1, \mathcal{C}_2$, respectively, such that $F : \mathcal{C}_1 \xrightarrow{\sim} \mathcal{C}_2$ is an equivalence, and $\mathcal{D}_i = \langle \mathcal{C}_i \rangle_{\Delta}$ for i = 1, 2 (i.e. the *t*-structures giving \mathcal{C}_i are nondegenerate). Then the following are equivalent:

- (a) F is an equivalence.
- (b) For any objects $M, N \in \mathcal{C}_1, F : \operatorname{Hom}^i_{\mathcal{D}_1}(M, N) \xrightarrow{\sim} \operatorname{Hom}^i_{\mathcal{D}_2}(F(M), F(N))$. In other words, the "Exts agree".
- (c) (Won't be used) Assume that $\mathcal{D}_1 = D^b(\mathcal{C}_1)$. For any $x \in \operatorname{Hom}^i_{\mathcal{D}_2}(F(M), F(N))$, there exists an injection $N \hookrightarrow N'$ such that x is zero in $\operatorname{Hom}^i_{\mathcal{D}_2}(F(M), F(N'))$; i.e. under the natural map

$$\operatorname{Hom}_{\mathcal{D}_2}^i(F(M), F(N)) \to \operatorname{Hom}_{\mathcal{D}_2}^i(F(M), F(N'))$$
$$x \mapsto 0.$$

This condition is called "effaceability".

Proof. (a) \implies (b) is immediate. (b) \implies (a) can be shown by induction and the long exact sequence.

Example 19.7. Let X be $K(\pi, 1)$. Then

$$D^b(\operatorname{Rep}^{f.d.} k\pi) \xrightarrow{\sim} D^b_{\Lambda}(X).$$

Example 19.3 illustrated that this is not necessarily the case when X is not $K(\pi, 1)$.


19.4 Yoneda extensions

To understand part (c) of Lemma 19.6, we need some facts about Yoneda extensions. Let \mathcal{A} be an abelian category. For objects M, N in \mathcal{A} , define a category $E^i(M, N)$ with

• objects: acyclic complexes

$$N \to C^1 \to C^2 \to \dots \to C^i \to M$$

• morphisms: chain complex maps of the form



Lemma 19.8. (/Definition)


$$\operatorname{Ext}^{i}(M, N) = \text{ connected components of } E^{i}(M, N)$$

 $0 \leftrightarrow \text{ split complexes in } E^{i}(M, N)$

Remark 19.9. In Lemma 19.8, two extensions are in the same "connected component" if one can pass from one to the other going along arrows in our categories *in either direction*. More formally, we can build a space with 0 (resp. 1) simplices given by objects (resp. arrows) in $E^i(M, N)$; then connected component takes on its topological meaning. One can show two objects

$$N \to C^1 \to C^2 \to \dots \to C^i \to M$$
 and $N \to \widetilde{C}^1 \to \widetilde{C}^2 \to \dots \to \widetilde{C}^i \to M$

in $E^{i}(M, N)$ are in the same connected component if there exists a commutative diagram



with

$$N \to D^1 \to D^2 \to \dots \to D^i \to M$$

also in $E^i(M, N)$.

Exercise 19.10. Show directly that with Ext^i defined as in Lemma 19.8,

$$\operatorname{Ext}^{i}(M, N) = \operatorname{Hom}_{D(\mathcal{A})}^{i}(M, N).$$

Make no assumptions on \mathcal{A} .

What about functoriality? In our usual definition of Ext^i in the derived category, it is obvious that Ext^i is a functor. With this new definition, it is not so obvious. In other words, given $N \to N'$, how to we obtain a morphism $\operatorname{Ext}^i(M, N) \to \operatorname{Ext}^i(M, N')$? Here is how:

• Given with $N \to N'$ and an element of $\operatorname{Ext}^i(M, N)$ represented by the complex

$$N \to C^1 \to C^2 \to \dots \to C^i \to M$$

we obtain a complex representing an element of $\operatorname{Ext}^{i}(M, N')$ by forming the push-out

Here $P = (C^1 \oplus N')/N$. This gives a morphism $\operatorname{Ext}^i(M, N) \to \operatorname{Ext}^i(M, N')$.

• Similarly, for $M \to M'$, we can use the pull-back to form a morphism $\operatorname{Ext}^i(M, N) \to \operatorname{Ext}^i(M', N)$.

An important consequence of this is the following. If $x \in \operatorname{Ext}^{i}(M, N)$ is represented by the complex

$$N \xrightarrow{f} C^1 \to C^2 \to \dots \to C^i \to M,$$

then applying the construction above to the morphism $f: N \to C^1$, we obtain

But since $(C^1 \oplus C^1)/N = C^1 \oplus (C^1/N)$, the lower sequence splits, and hence represents zero in $\operatorname{Ext}^i(M, C^1)$. In other words,

$$\operatorname{Ext}^{i}(M, N) \xrightarrow{\operatorname{Ext}^{i}(M, f)} \operatorname{Ext}^{i}(M, C^{1})$$
$$x \longmapsto 0.$$

This is the origin of effaceability.

Exercise 19.11. Prove (ii) \implies (iii) and (iii) \implies (ii) in Lemma 19.6. (Hint: See Geordie's hand-written notes if you are stuck.)


19.5 A key ingredient

A key ingredient in Beilinson's theorem is:

Theorem 19.12. Let X/\mathbb{C} be a smooth variety. Then there exists a Zariski open set $U \subset X$ which is $K(\pi, 1)$.

This theorem explains philosophically why Beilinson's theorem is true: for perverse sheaves, we can always refine our stratification to include this $K(\pi, 1)$ open set. We will see this more precisely when we discuss the proof of Beilinson's theorem.


Example 19.13. Any curve becomes $K(\pi, 1)$ after deleting a point!



(Here the arrows indicate homotopy equivalences and the x's are missing points.)

We will roughly explain the proof of Theorem 19.12. There are two main ingredients:

1. "Noether normalization," (**NN**): If $Z \subset \mathbb{A}^n$ is of dimension d, then a generic projection $\mathbb{A}^n \to \mathbb{A}^d$ is finite when restricted to Z. Here's a caricature:



2. "Extensions of $K(\pi, 1)$ are $K(\pi, 1)$," $(\boldsymbol{EK}(\pi, 1))$: If

$$\begin{array}{c} F \longleftrightarrow E \\ & \downarrow \\ & B \end{array}$$

is a fibre bundle, then if B is $K(\pi, 1)$, E is connected, and either F is discrete or $K(\pi, 1)$, then E is $K(\pi, 1)$. (Follows from the long exact sequence of homotopy groups.)

With these tools in our toolbox, the proof proceeds as follows.

Step 1: Reduction to $X \subset \mathbb{A}^1$, Zariski standard open¹⁸. Without loss of generality, we may assume that X is affine. By **(NN)**, we can find a finite map $f : X \to \mathbb{A}^d$. Over a standard open $U' \subset \mathbb{A}^d$, this map is étale. Hence we have a fibration

finite
$$\longleftrightarrow U$$

 \downarrow
 U'

so by $(\boldsymbol{EK}(\boldsymbol{\pi}, \mathbf{1}))$, if U' is $K(\boldsymbol{\pi}, 1)$, then U is $K(\boldsymbol{\pi}, 1)$.

Step 2: Induction on dimension. By **(NN)**, there exists a projection $f : \mathbb{A}^n \to \mathbb{A}^{n-1}$ such that on the hypersurface $Z = X \setminus U$, f is finite with fibres consisting of m points:

$$U \longleftrightarrow \mathbb{A}^n \longleftrightarrow Z \longleftrightarrow \{m \text{ points}\}$$
$$\downarrow^f \qquad \qquad \downarrow^f$$
$$\mathbb{A}^{n-1} = \mathbb{A}^{n-1}$$

Over an open $V \subset \mathbb{A}^{n-1}$, this map will be étale, hence

$$\begin{array}{ccc} U' & \longleftrightarrow & U \\ \downarrow & & \downarrow \\ V' & \longleftrightarrow & V \end{array}$$

is fibred in $\mathbb{C} \setminus \{m \text{ points}\}$. By induction, we can shrink V to V' such that V' is $K(\pi, 1)$. Hence U' is $K(\pi, 1)$ too by $(\mathbf{EK}(\pi, 1))$.

19.6 Proof of Beilinson's theorem for curves

The goal for the rest of the lecture is to prove Beilinson's theorem for curves; that is, for a curve X, we wish to show that

$$D^b(P_X) \simeq D^b_c(X).$$

To start, we have the following fact [DBB83, Bei87]. There exists a triangulated functor

real :
$$D^b(P_X) \to D^b_c(X)$$

called the "realization functor" which is the identity on P_X . (This holds for general X, not just curves.) We will show that real is an equivalence of categories.

By Lemma 19.6, it is enough to show that for $M, N \in P_X$,

$$\operatorname{Hom}_{D^b(P_X)}^i(M,N) \simeq \operatorname{Hom}_{D^b_c(X)}^i(M,N).$$

Moreover, because P_X is finite length, we can use the long exact sequence in Ext to reduce to the case where M, N are irreducible objects.

¹⁸ "Zariski standard open" means that \overline{X} is of the form $D(f) = \{z \in X \mid f(z) \neq 0\}$ for some regular function f.

Recall that in lecture 17 we classified irreducible perverse sheaves on curves. There were two types: (1) skyscrapers, and (2) IC sheaves supported everywhere.

Step 1: Let M, N be skyscrapers: $M = i_*k_x$, $N = i'_*k_y$. Then in $D^b_c(X)$,

$$\operatorname{Hom}^{j}(i_{*}k_{x}, i_{*}'k_{y}) = \operatorname{Hom}^{j}(k_{x}, i^{!}i_{*}'k_{y})$$
$$= \begin{cases} k & \text{if } x = y, j = 0, \\ 0 & \text{otherwise.} \end{cases}$$

Showing that

$$\operatorname{Hom}_{D^b(P_X)}(k_x, k_y) = 0$$

if $x \neq y$ is easier, and left as an exercise. The tricker case is show that $\operatorname{Ext}_{P_X}^i(k_x, k_x) = 0$ for i > 0. To do this, we will use the quiver description of P_X and Yoneda Exts. Recall that

$$P_X \simeq \left\{ V_1 \xleftarrow{c}{v} V_0 \mid c \circ v + id \text{ invertible} \right\}.$$

Then an element of $Ext^{1}(k_{x}, k_{y})$ has a representative of the form

$$k_k \to F \to k_x.$$

In quiver language, such an extension is a complex

$$k \to V_0 \to k.$$

But any such complex fits into a diagram

$$\begin{array}{ccc} 0 & \longrightarrow & V_1 & \longrightarrow & 0 \\ \uparrow \downarrow & & v \uparrow \downarrow c & & \uparrow \downarrow \\ k & \longrightarrow & V_0 & \longrightarrow & k \end{array}$$

The top sequence is exact, so $V_1 = 0$. But the bottom sequence is also exact, so $V_1 = 0$ implies that the bottom sequence must split. Hence our original element is zero in $\text{Ext}^1(k_x, k_y)$, so $\text{Ext}^1(k_x, k_y) = 0$.

Showing that $\operatorname{Ext}^2(k_x, k_y) = 0$ is more challenging. An element of $\operatorname{Ext}^2(k_x, k_y)$ has a representative of the form

$$k \to V_0^1 \to V_0^2 \to k.$$

This fits into a diagram

$$\begin{array}{cccc} 0 & \longrightarrow & V_1^1 & \longrightarrow & V_1^2 & \longrightarrow & 0 \\ \uparrow & & & v^1 \uparrow \downarrow c^1 & & v^2 \uparrow \downarrow c^2 & & \uparrow \downarrow \\ k & \longrightarrow & V_0^1 & \longrightarrow & V_0^2 & \longrightarrow & k \end{array}$$

We want to show that we can reduce this to an Ext supported at x. But the basic issue is that there is nothing telling us that the middle terms must be supported on a point. In sheaf language, this is equivalent to the fact that the existence of a sequence

$$i_*k_x \to \mathcal{F} \to \mathcal{G} \to i_*k_x$$

whose first and last terms are supported on a point does not imply that the middle terms \mathcal{F}, \mathcal{G} are.

There are two ways we can get around this. First, the "stupid way". We can assume that $c \circ v$ is nilpotent (exercise!), then we have a natural map

Continuing this process, we have a map

Continuing in this way, we eventually obtain a top sequence of zeros, which lets us conclude that our lower sequence splits. This procedure works for all higher Exts.

Alternatively, we could use Beilinson's approach using vanishing cycles, which works in greater generality (and in fact provides the skeleton of his argument in the general case). Choose a map $f: X \to \mathbb{C}$ with zero set $x \in X$. Let C be the complex

$$i_*k_x \to \mathcal{F}^1 \to \cdots \to \mathcal{F}^j \to i_*k_x$$

representing an element in $\operatorname{Ext}^{j}(i_{*}k_{x}, i_{*}k_{x})$. Then Beilinson showed that

$$C^{\cdot} \simeq \phi_f(C^{\cdot}), \tag{19.1}$$

where $\phi_f : P_X \to P_{\{x\}}$ is the vanishing cycles functor defined in Lecture 18. The complex $\phi_f(C^{\cdot})$ is supported on x, so this proves that $C^{\cdot} = 0 \in \operatorname{Ext}^j(i_*k_x, i_*k_x)$. To prove (19.1), we can use the exact sequences from Lecture 18:

$$C^{\cdot} \longrightarrow C^{\cdot} \oplus \Xi_{f}(C^{\cdot}) \longrightarrow (C^{\cdot} \oplus_{f} (C^{\cdot}))/j_{!}(C^{\cdot})$$

$$\uparrow$$

$$\phi_{f}(C^{\cdot})$$

Step 2: Let M be irreducible with full support and N a skyscraper supported at $\{x\}$. Let

$$j: U = X \setminus \{x\} \hookrightarrow X.$$

Recall that because the immersion j is affine, $j_{!}$ and j_{*} are exact, preserve perverse sheaves, and fit into adjoint pairs $(j^{*}, j_{*}), (j_{!}, j^{!})$. The adjunction maps give an exact sequence

$$K \hookrightarrow j_! M_U \twoheadrightarrow M,$$

where K is a perverse sheaf supported on $\{x\}$. The corresponding long exact sequence gives the diagram

Using the adjunctions, we have

$$\operatorname{Hom}_{P_X}^i(j_!M_U, N) = \operatorname{Hom}_{P_U}^i(M_U, N_U) = 0$$

because $N_U = 0$. Similarly, $\operatorname{Hom}_{D_c^b(X)}^i(j_!M_U, N) = 0$, so the middle vertical arrow is an isomorphism $0 \simeq 0$. The right vertical arrow is an isomorphism by Step 1. Hence we conclude by induction that all other vertical arrows must also be isomorphisms.

Step 3: Let M and N be irreducible objects in P_X with full support. Then by Lemma 19.12, we can choose $U \subset X$ such that M_U, N_U are (shifts of) local systems and U is $K(\pi, 1)$. Note that the inclusion $j: U \hookrightarrow X$ is still affine. Because N is irreducible, we have a short exact sequence

$$N \hookrightarrow j_* j^* N = j_* N_U \twoheadrightarrow k.$$

Then by the long exact sequence in Hom, we have the diagram

By adjunction, we have

$$\operatorname{Hom}_{P_{X}}^{i}(M, j_{*}N_{U}) = \operatorname{Hom}^{i}(M_{U}, N_{U}).$$

Hence the middle vertical arrow is an isomorphism because U is $K(\pi, 1)$. The right vertical arrow is an isomorphism by Step 2, and hence the left vertical arrow must also be an isomorphism.

The Moral: If we fix a stratification, there can be complicated structure going on in $D^b(P_X)$ which doesn't only come from fundamental groups. However, if we're allowed to refine our stratification as much as we want, then we can choose open strata which are $K(\pi, 1)$, and all information in the category comes from local systems.

Lecture 20: Perverse sheaves for affine stratifications

Recall the setting of the last few lectures: Fix k to be our field of coefficients, and let X be an algebraic variety over \mathbb{C} , with stratification

$$X = \bigsqcup_{\lambda \in \Lambda} X_{\lambda}.$$

The abelian category of Λ -constructible perverse sheaves, $\operatorname{Perv}_{\Lambda}(X)$, is a subcategory of the triangulated category $D^b_{\Lambda}(X)$, the derived category of Λ -constructible sheaves. Moreover, there is a realization functor

real :
$$D^b(\operatorname{Perv}_{\Lambda}(X)) \to D^b_{\Lambda}(X).$$

Last week we saw that for a fixed stratification, real is rarely an equivalence; however, if we are allowed to refine stratifications, we have Beilinson's theorem:

Theorem 20.1. (Beilinson) $D^b(\operatorname{Perv}(X)) \xrightarrow{\sim} D^b_c(X)$.

Today, we will show that if each strata X_{λ} is an affine space, then we have such an equivalence for a fixed stratification.

Theorem 20.2. (Beilinson–Ginzburg–Soergel) If Λ is a stratification of X such that X_{λ} is affine, then $D^b(\operatorname{Perv}_{\Lambda}(X)) \xrightarrow{\sim} D^b_{\Lambda}(X)$.

Remark 20.3. Last week we saw that the failure of the strata to be $K(\pi, 1)$ was the obstruction to real being an equivalence. Affine spaces are contractible, so this obstruction disappears¹⁹.

Remark 20.4. Serious homological algebra goes into Beilinson's construction of real. However, if we allow ourselves to use the Riemann–Hilbert correspondence, we have

$$\mathcal{D} - \operatorname{mod}_{\Lambda\operatorname{-const}}^{hol,r.s.} \longleftrightarrow \operatorname{Perv}_{\Lambda}(X)$$

$$\downarrow$$

$$D^{b} \left(\mathcal{D} - \operatorname{mod}_{\Lambda\operatorname{-const}}^{hol,r.s} \right) \xrightarrow[\mathrm{R.H.}]{} D^{b}_{\Lambda}(X)$$

Under the equivalence between perverse sheaves and \mathcal{D} -modules given by the top arrow, the bottom arrow in this diagram is real. This gives a high concept (rather than high homological algebra!) construction of the realization functor.

The BGS theorem and the techniques involved in proving it will be invaluable as we approach Bezrukavnikov's equivalence. The proof of the BGS theorem is algebraic, using what are today called *highest weight categories*.

¹⁹Caution: take this explanation with a grain of salt. The proof of Beilinson's theorem is geometric (via vanishing cycles), and the proof of the BGS theorem is algebraic (via highest weight categories), so a direct comparison doesn't exactly work.

20.1 Highest weight categories

Definition 20.5. Let \mathcal{A} be a k-linear category. We say that \mathcal{A} is highest weight if the following six conditions hold.

- 1. \mathcal{A} is finite length.
- 2. The set {simple objects in \mathcal{A} }/ \simeq is finite.
- 3. For any simple object $L \in \mathcal{A}$, $\operatorname{End}(L) = k$.

Let Λ be an indexing set for isomorphism classes of simple objects and denote by $L_{\lambda} \in \mathcal{A}$ the simple object corresponding to $\lambda \in \Lambda$. Assume that Λ is a poset (this is part of the data determining a highest weight category). For any closed subset $T \subset \Lambda$ (that is, if $\lambda' \leq \lambda$ and $\lambda \in T$, then $\lambda' \in T$), denote by


$$\mathcal{A}_T = \langle L_\lambda \mid \lambda \in T \rangle_{\text{Serre}}$$

the Serre subcategory generated by the simple objects L_{λ} with $\lambda \in T$. Assume moreover that for each $\lambda \in \Lambda$, there are objects Δ_{λ} ("standard" object), ∇_{λ} ("costandard" object) in \mathcal{A} and maps $\Delta_{\lambda} \to L_{\lambda}$, $L_{\lambda} \to \nabla_{\lambda}$.

- 4. If $T \subseteq \Lambda$ is closed, then $\Delta_{\lambda} \to L_{\lambda}$ (resp. $L_{\lambda} \to \nabla_{\lambda}$) is a projective cover²⁰ (resp. injective hull) in \mathcal{A}_T^{21} .
- 5. For $\lambda \in \Lambda$,

$$\ker(\Delta_{\lambda} \to L_{\lambda}) \in \mathcal{A}_{<\lambda}, \text{ and}$$
$$\operatorname{coker}(L_{\lambda} \to \nabla_{\lambda}) \in \mathcal{A}_{<\lambda}.$$

This implies that the composition series eggs of Δ_{λ} and ∇_{λ} must have the following form



6. $\operatorname{Ext}^2(\Delta_{\lambda}, \nabla_{\mu}) = 0$ for all $\lambda, \mu \in \Lambda$.²²

²⁰A projective cover of an object M in a category C is a morphism $P \xrightarrow{f} M$ out of a projective object $P \in C$ which is a superfluous epimorphism, meaning that every morphism $N \xrightarrow{g} P$ with the property that $f \circ g$ is an epimorphism is itself an epimorphism.

²¹Note that we are not requiring Δ_{λ} (resp. ∇_{λ}) to be projective (resp. injective) objects in \mathcal{A} , just in the Serre subcategory \mathcal{A}_T .

 $^{^{22}}$ This condition is the most mysterious, and often the hardest to show.

Theorem 20.6. (Beilinson–Ginzburg–Soergel) Let \mathcal{A} be highest weight. Then \mathcal{A} has enough projective and injective objects. Moreover, the projective cover P_{λ} of L_{λ} has a standard filtration of the form

$$P_{\lambda} = \begin{pmatrix} \Delta_{\lambda} \\ \Delta_{\lambda}'s \\ \lambda > \lambda \end{pmatrix}$$


Similarly, the injective hull I_{λ} has a cost and ard filtration of the form

$$I_{\lambda^{\pm}} \begin{pmatrix} \nabla_{\mu}'s \\ \mu > \lambda \\ \nabla_{\lambda} \end{pmatrix}$$

Proof. For full details, see [BGS96, Theorem 3.2.1].

Sketch: Instead of proving the theorem as stated, we prove that for any closed subset $T \subseteq \Lambda$, \mathcal{A}_T has enough projectives (resp. injectives), and they admit standard (resp. costandard) filtrations. We construct the projective cover P_{λ}^T of L_{λ} inductively on T:

• Let $\mu \in T$ be maximal, and $\lambda \neq \mu \in T$. Assume that the projective cover $P_{\lambda}^{T \setminus \{\mu\}}$ of L_{λ} in $\mathcal{A}_{T \setminus \{\mu\}}$ is already constructed.



- Let $E = \operatorname{Ext}^1(P_{\lambda}^{T \setminus \{\mu\}}, \nabla_{\mu})$ be the ext group.
- Every element of E gives rise to an extension

$$\Delta_{\mu} \hookrightarrow P' \twoheadrightarrow P_{\lambda}.$$

It turns out that there exists a "universal" extension

$$E^* \otimes \Delta_{\mu} \hookrightarrow \widetilde{P} \twoheadrightarrow P_{\lambda}^{T \setminus \{\mu\}}$$

from which all others are constructed via push-out. (Challenge: construct it!)

• Claim: \widetilde{P} is the projective cover we are seeking.

Proof. Use long exact sequence in Hom:


$$\operatorname{Hom}(P_{\lambda}^{T \setminus \{\mu\}}, \Delta_{\mu}) \longrightarrow \operatorname{Hom}(\widetilde{P}, \Delta_{\mu}) \longrightarrow E \otimes \operatorname{Hom}(\Delta_{\mu}, \Delta_{\lambda}) \longrightarrow \operatorname{Ext}^{1}(P_{\lambda}^{T \setminus \{\mu\}}, \Delta_{\mu}) \longrightarrow \operatorname{Ext}^{1}(\widetilde{P}, \Delta_{\mu}) \longrightarrow E \otimes \operatorname{Ext}^{1}(\Delta_{\mu}, \Delta_{\mu})$$

The last term is zero by axoim (4) of a highest weight category. Hence $\operatorname{Ext}^{1}(\widetilde{P}, \Delta_{\mu}) = 0$. A bit more work using axiom (6) shows that $\widetilde{P} = P_{\lambda}^{T}$.

Definition 20.7. In a highest weight category \mathcal{A} , an object T is *tilting* if it has a standard and costandard filtration. The indecomposable tilting objects in \mathcal{A} are also indexed by Λ .

Similar arguments give injective and tilting objects.

Important objects in a highest weight category:



If our category also has a notion of duality compatible with the highest weight structure, then the costandard/standard and projective/injective objects are dual, as indicated by the pink arrows above.

Exercise 20.8. Let $\mathcal{B} \subset \mathcal{A}$ be a Serre subcategory of an abelian category \mathcal{A} . By definition, for objects $M, N \in \mathcal{B}$,

 $\operatorname{Ext}^{1}_{\mathcal{B}}(M,N) \xrightarrow{\sim} \operatorname{Ext}^{1}_{\mathcal{A}}(M,N).$

Show that

$$\operatorname{Ext}^2_{\mathcal{B}}(M, N) \hookrightarrow \operatorname{Ext}^2_{\mathcal{A}}(M, N).$$

Hint: Use effaceability and the long exact sequence of Ext.

20.2 Perverse sheaves for an affine stratification are a highest weight category

Theorem 20.9. If $X = \bigsqcup_{\lambda \in \Lambda} X_{\lambda}$ is stratified by affine spaces, then $\operatorname{Perv}_{\Lambda}(X)$ is a highest weight category.

Proof. First, note that $j_{\lambda} : X_{\lambda} \hookrightarrow X$ is affine and hence

$$\Delta_{\lambda} := j_{\lambda !} k_{X_{\lambda}} [\dim X_{\lambda}] \text{ and}$$
$$\nabla_{\lambda} := j_{\lambda *} k_{X_{\lambda}} [\dim X_{\lambda}]$$

are perverse sheaves. Moreover, we have canonical maps

$$\Delta_{\lambda} \twoheadrightarrow IC_{\lambda} \hookrightarrow \nabla_{\lambda}, \tag{20.1}$$

where $IC_{\lambda} := j_{\lambda!*}k_{X_{\lambda}}$ is the IC sheaf corresponding to the trivial local system on X_{λ} .

- 1. $\operatorname{Perv}_{\Lambda}(X)$ is finite-length. \checkmark
- 2. Simple objects in $\operatorname{Perv}_{\Lambda}(X)$ are parameterized by pairs (\mathcal{L}, λ) , where \mathcal{L} is a local system on X_{λ} . Because X_{λ} is affine, it is contractible, and thus admits a single local system. Hence there are finitely many simple objects in $\operatorname{Perv}_{\Lambda}(X)$. \checkmark
- 3. By Schur's Lemma, $\operatorname{End}(IC_{\lambda})$ is a division algebra over k.

Claim 20.10. If D is a division algebra over a field k, any non-zero algebra homomorphism $D \rightarrow k$ is an isomorphism.

Proof. Let $\varphi : D \to k$ be a nonzero algebra homomorphism. The kernel of φ is an ideal in D, but the division algebras have no non-trivial ideals, so ker $\varphi = \{0\}$. Because φ is a k-algebra homomorphism, $\varphi(1) = 1$. Hence for all $\ell \in k$, $\varphi(\ell \cdot 1) = \ell \varphi(1) = \ell$ and $\operatorname{im} \varphi = k$.

Assume $X_{\lambda} \subset X$ is open. Then $IC_{\lambda}|_{X_{\lambda}} = k_{X_{\lambda}}[\dim X_{\lambda}]$. Hence the restriction map

$$\operatorname{End}(IC_{\lambda}) \xrightarrow{\operatorname{restriction}} \operatorname{End}(IC_{\lambda}|_{X_{\lambda}}) \simeq k$$

is a nonzero morphism from a division algebra to k. The claim lets us conclude that $\operatorname{End}(IC_{\lambda}) \simeq k$.

For any λ , the inclusion $X_{\lambda} \hookrightarrow \overline{X}_{\lambda}$ is open because X_{λ} is locally closed. There is an equivalence of categories

perverse sheaves
supported on a
closed subvariety
$$Z \subset X$$
 $\stackrel{\sim}{\leftrightarrow} \operatorname{Perv}(Z),$

 \mathbf{SO}

$$\operatorname{End}_{\operatorname{Perv}_{\Lambda}(X)}(IC_{\lambda}) \simeq \operatorname{End}_{\operatorname{Perv}_{\Lambda}(\overline{X}_{\lambda})}(IC_{\lambda}).$$

Hence by applying the argument above to the open inclusion $X_{\lambda} \hookrightarrow \overline{X}_{\lambda}$, we conclude that $\operatorname{End}(IC_{\lambda}) = k$ for all λ .

4. The maps in (20.1) give a surjection $\Delta_{\lambda} \to IC_{\lambda}$ and an injection $IC_{\lambda} \to \nabla_{\lambda}$. Because the partial order on Λ is given by closure of strata, for a closed subset $\{\leq \lambda\} \subset \Lambda$, we have

$$\operatorname{Perv}_{\Lambda}(X)_{\{\leq\lambda\}} = \operatorname{Perv}_{\Lambda}(\overline{X}_{\lambda}).$$

To establish axiom (4), we must show two things: (a) that Δ_{λ} is a projective object in $\operatorname{Perv}_{\Lambda}(\overline{X}_{\lambda})$, and (b) that $\Delta_{\lambda} \twoheadrightarrow IC_{\lambda}$ is a projective cover.

To show that Δ_{λ} is projective, we will show that $\operatorname{Hom}_{\operatorname{Perv}_{\Lambda}(\overline{X}_{\lambda})}(\Delta_{\lambda}, \cdot)$ is exact. Because $X_{\lambda} \stackrel{j}{\hookrightarrow} \overline{X}_{\lambda}$ is open, for any $\mathcal{F} \in \operatorname{Perv}_{\Lambda}(\overline{X}_{\lambda})$,

$$\operatorname{Hom}(j_{!}k_{X_{\lambda}}[\dim X_{\lambda}], \mathcal{F}) = \operatorname{Hom}(k_{X_{\lambda}}[\dim X_{\lambda}], j^{!}\mathcal{F})$$

= $\operatorname{Hom}(k_{X_{\lambda}}[\dim X_{\lambda}], j^{*}\mathcal{F})$
= $\operatorname{Hom}(k_{X_{\lambda}}[\dim X_{\lambda}], \mathcal{F}|_{X_{\lambda}})$
= $\mathcal{F}|_{X_{\lambda}}.$

Restriction to an open subvariety is an exact functor, so we conclude that $\operatorname{Hom}(\Delta_{\lambda}, \cdot)$ is exact.

Because $\operatorname{Perv}_{\Lambda}(\overline{X}_{\lambda})$ is a finite-length abelian category (in particular, it is Krull-Schmidt), to show that $\Delta \twoheadrightarrow IC_{\lambda}$ is a projective cover, it suffices to show that Δ_{λ} is indecomposable. Now, the endomorphism ring


$$\operatorname{End}(\Delta_{\lambda}) = \operatorname{Hom}(j_! k_{X_{\lambda}}, j_! k_{X_{\lambda}}) = \operatorname{Hom}(k_{X_{\lambda}}, j^! j_! k_{X_{\lambda}}) = \operatorname{End}(k_{X_{\lambda}}) = k$$

is local, so Δ_{λ} is indecomposable.

Showing that the injection $IC_\lambda \hookrightarrow \nabla_\lambda$ is an injective hull follows from a similar argument. \checkmark

5. The fact that $\ker(\Delta_{\lambda} \twoheadrightarrow IC_{\lambda})$, $\operatorname{coker}(IC_{\lambda} \hookrightarrow \nabla_{\lambda}) \in \operatorname{Perv}_{<\lambda}(X)$ is clear from the com-

position series eggs and the definition of $IC_{\lambda} = \text{Im}(J_{\lambda!} \to j_{\lambda*})$:



6. To see why condition (6) holds, first observe that by Exercise 20.8,

$$\operatorname{Ext}^2_{\operatorname{Perv}_{\Lambda}(X)}(\Delta_{\lambda}, \nabla_{\mu}) \hookrightarrow \operatorname{Ext}^2_{\operatorname{Perv}(X)}(\Delta_{\lambda}, \nabla_{\mu}).$$

Then, by Beilinson's theorem,

$$\operatorname{Ext}^{2}_{\operatorname{Perv}(X)}(\Delta_{\lambda}, \nabla_{\mu}) = \operatorname{Hom}^{2}_{D^{b}_{c}(X)}(\Delta_{\lambda}, \nabla_{\mu}).$$

Once we are in the derived category, we have full access²³ to the adjoint pairs $(j_!, j^!), (j^*, j_*)$. Hence we can use adjunctions to compute

$$\operatorname{Hom}_{D_{c}^{b}(X)}^{2}(\Delta_{\lambda}, \nabla_{\mu}) = \operatorname{Hom}_{D_{c}^{b}(X)}(j_{\lambda !}k_{X_{\lambda}}[\dim X_{\lambda}], j_{\mu *}k_{X_{\mu}}[\dim X_{\mu} + 2])$$
$$= \operatorname{Hom}_{D_{c}^{b}(X)}(j_{\mu}^{*}j_{\lambda !}k_{X_{\lambda}}[\dim X_{\lambda}], k_{X_{\mu}}[\dim X_{\mu} + 2]).$$

Because the functor $j_{\lambda !}$ is extension by zero, $j_{\mu}^* j_{\lambda !} k_{X_{\lambda}} = 0$ for $\mu \neq \lambda$. If $\mu = \lambda$, then $j_{\mu}^* j_{\lambda !} k_{X_{\lambda}}$ is a 1-dimensional vector space. There are no higher Exts in the category of vector spaces, so we conclude that

$$\operatorname{Ext}^{2}_{\operatorname{Perv}_{\lambda}(X)}(\Delta_{\lambda}, \nabla_{\mu}) = 0.\checkmark$$

Remark 20.11. It's unusual for a category of perverse sheaves to have enough projectives. By proving that $\operatorname{Perv}_{\Lambda}(X)$ is a highest weight category, we have just algebraically produced a collection of projective perverse sheaves $\{P_{\lambda}\}_{\lambda \in \Lambda}$. As far as we are aware, it is not known how to construct these perverse sheaves geometrically.

Corollary 20.12. If Λ is a stratification of X by affine spaces,

real :
$$D^b(\operatorname{Perv}_{\Lambda}(X)) \xrightarrow{\sim} D^b_{\Lambda}(X)$$

is an equivalence of categories.

²³What we mean by this is that the functor j_{λ}^* is not exact if X_{λ} is not open, so it does not preserve perverse sheaves (see beginning of lecture 19).

Proof. Both of the sets $\{\Delta_{\lambda}\}_{\lambda \in \Lambda}$ and $\{\nabla_{\lambda}\}_{\lambda \in \Lambda}$ generate $D^{b}(\operatorname{Perv}_{\Lambda}(X))$ and $D^{b}_{\Lambda}(X)$. By the upper triangularity of P_{λ} , the set $\{P_{\lambda}\}_{\lambda \in \Lambda}$ also generates each category. Hence it is enough to show

real :
$$\operatorname{Hom}_{D^b(\operatorname{Perv}_{\Lambda}(X))}^i(P_{\lambda}, \nabla_{\mu}) \xrightarrow{\sim} \operatorname{Hom}_{D_{\lambda}^b(X)}^i(P_{\lambda}, \nabla_{\mu}).$$

In $D^b(\operatorname{Perv}_{\Lambda}(X))$,

$$\operatorname{Hom}^{i}(P_{\lambda}, \nabla_{\mu}) = \begin{cases} 0 & \text{if } i > 0, \\ \operatorname{Hom}_{\operatorname{Perv}_{\Lambda}(X)}(P_{\lambda}, \nabla_{\mu}) & \text{if } i = 0, \end{cases}$$

by the projectivity of P_{λ} .

In $D^b_{\Lambda}(X)$,

$$\operatorname{Hom}^{i}(P_{\lambda}, \nabla_{\mu}) = \begin{cases} 0 & \text{if } i > 0, \\ \operatorname{Hom}_{D_{\Lambda}^{b}(X)}(P_{\lambda}, \nabla_{\mu}) & \text{if } i = 0 \end{cases}$$

as well. This is because $\operatorname{Hom}_{D_{\Lambda}^{b}(X)}^{i}(\Delta_{\lambda}, \nabla_{\mu}) = 0$ for i > 0 by adjunction, so the long exact sequence in Ext and the standard filtration of P_{λ} imply that $\operatorname{Hom}^{i}(P_{\lambda}, \nabla_{\mu}) = 0$ for i > 0. \Box

Remark 20.13. In the proof above, the vanishing of Hom^i for i > 0 happens for quite different reasons in each of the two categories. In $D^b(\operatorname{Perv}_{\Lambda}(X))$, the reason is algebraic (projectivity of an object), whereas in $D^b(X)$, the reason is topological.

20.3 Where are we going for the next few weeks?

For the rest of this lecture, we will reconnect with the big picture of this course and describe our plan for the upcoming weeks.

Let $(X \supset R, X^{\vee} \supset R^{\vee})$ be a root datum, and let G, G^{\vee} be the corresponding dual groups over \mathbb{C} . Associated to this datum, we have a finite Weyl group W_f , and an affine Weyl group $W = W_f \ltimes \mathbb{Z}X^{\vee}$. Let H be the affine Hecke algebra, and Z its center. In October and November of last year, we constructed the following diagram.

The isomorphism on the bottom line is the Kazhdan–Lusztig isomorphism, which was merely stated (not yet even sufficiently explained, let alone proved!), and the injection on the left is Bernstein's description of the center of the affine Hecke algebra. Our goal for the rest of the course is to categorify this picture. This is done via Bezrukavnikov's equivalence, which very roughly is an equivalence of the form

$$\begin{pmatrix} \text{constructible} \\ \text{affine Hecke} \\ \text{category} \end{pmatrix} \xrightarrow{\sim}_{B} \begin{pmatrix} \text{coherent} \\ \text{affine Hecke} \\ \text{category} \end{pmatrix}.$$

The constructible affine Hecke category on the LHS should be (a variant of) a category of construcible sheaves on the affine flag variety of G, and the coherent affine Hecke category on the RHS should be (a variant of) a category of $G \times \mathbb{C}^{\times}$ -equivariant coherent sheaves on the Steinberg variety. When we take Grothendieck groups, we should recover the first diagram.

Philosophy for now: Before we can understand this story on the level of categories, we need to better understand the Kazhdan–Lusztig isomorphism.

General strategy for understanding KL:

1. On the Bernstein generators $T_i, \mathcal{O}_{\lambda}$, define

 $T_i \mapsto \mathcal{Q}_i$ (to be explained next week) $\mathcal{O}_{\lambda} \mapsto \mathcal{O}_{\lambda}$ (pull-back of $\mathcal{O}(\lambda)$ on G^{\vee}/B^{\vee} to $T^*G^{\vee}/B^{\vee} \xrightarrow{\text{diagonal}} St$)

Remark 20.14. It is very challenging to verify the relations directly! (See Leonardo Maltoni's May 24, 2019 talk in the Informal Friday Seminar.)

2. The affine Hecke algebra H has two important modules, defined as follows. Let

triv:
$$H_f \to \mathbb{Z}[q^{\pm 1}], T_i \mapsto q$$

sgn: $H_f \to \mathbb{Z}[q^{\pm 1}], T_i \mapsto -1$

be the (quantized versions of) the trivial and sign representation of the finite Hecke algebra H_f . Define two *H*-modules

$$M := H \otimes_{H_f} \operatorname{triv}, \quad N := H \otimes_{H_f} \operatorname{sgn}$$

by left multiplication on the first tensor factor. These are, respectively, the **spheri**cal module and antispherical module of the Hecke algebra corresponding to the parabolic subgroup $W_f \subset W$.

Remark 20.15. By the Bernstein presentation,

$$H = \mathbb{Z}[q^{\pm 1}][X^{\vee}] \otimes H_f.$$

Hence, as $\mathbb{Z}[q^{\pm 1}]$ -modules,

$$M \cong N \cong \mathbb{Z}[q^{\pm 1}][X^{\vee}].$$

Thus, as modules over $\mathbb{Z}[q^{\pm 1}][X^{\vee}]$, both the spherical and anti-spherical modules are free of rank 1.

3. By convolution formalism, both $K^{G \times \mathbb{C}^{\times}}(G^{\vee}/B^{\vee})$ and $K^{G \times \mathbb{C}^{\times}}(T^*G^{\vee}/B^{\vee})$ are $K^{G^{\vee} \times \mathbb{C}^{\times}}(St)$ -modules.

Remark 20.16. Note that

$$\mathbb{Z}[q^{\pm 1}][X^{\vee}] = K^{B^{\vee} \times \mathbb{C}^{\times}}(\mathrm{pt}) \cong K^{G^{\vee} \times \mathbb{C}^{\times}}(G^{\vee}/B^{\vee}) \xrightarrow[\mathrm{pull-back}]{\sim} K^{G^{\vee} \times \mathbb{C}^{\times}}(T^*G^{\vee}/B^{\vee}).$$

4. Compute actions of generators in M (resp. N) and match them with actions in $K^{G^{\vee} \times \mathbb{C}^{\times}}(G^{\vee}/B^{\vee})$ (resp. $K^{G^{\vee} \times \mathbb{C}^{\times}}(T^*G^{\vee}/B^{\vee})$ under (a choice of) the above isomorphisms. Because all of these modules are faithful, this implies the Kazhdan–Lusztig isomorphism.

Remark 20.17. We really only need to do this computation for either M or N to use this argument to deduce the Kazhdan–Lusztig isomorphism.

Next week: We implement 1-4.

Following week: We take the first step toward Bezrukavnikov's equivalence by explaining the Arkhipov–Bezrukavnikov's theorem that

$$\mathcal{AS} \xrightarrow{\sim} D^b \left(\operatorname{Coh}^{G^{\vee} \times \mathbb{C}^{\times}} (T^* G^{\vee} / B^{\vee}) \right),$$

where \mathcal{AS} is the categorical antispherical module.

Lecture 21: Equivariant *K*-theory of the Steinberg variety

Our goal for the next few lectures is to prove the Kazhdan–Lusztig isomorphism:

 $H_{\text{affine}} \simeq K^{G^{\vee} \times \mathbb{C}^{\times}} (\text{Steinberg}).$

This is an isomorphism of the affine Hecke algebra with the $G^{\vee} \times \mathbb{C}^{\times}$ -equivariant K-theory of the Steinberg variety. Recall from the end of last lecture that our general strategy for proving this isomorphism is to find a vector space on which each algebra acts faithfully by the same operators.

Remark 21.1. The reader might have the (correct) impression that this is a rather indirect way of seeing that two algebras are isomorphic. However, there are several instances of very indirect techniques to obtain isomorphisms, equivalences, or correspondences in the Langlands program. Another example of this is Soergel's functor, which proves an equivalence of two categories (one geometric and one representation theoretic) by matching their images in a third world of algebra (so-called Soergel bimodules).

References for this lecture are:

- Kazhdan-Lusztig, Proof of the Deligne-Langlands conjecture for Hecke algebras, [KL87],
- Chriss-Ginzburg, Representation theory and complex geometry, Chapters 6 & 7, [CG09],
- Henderson, Notes on affine Hecke algebras and K-theory.

21.1 Equivariant *K*-theory

Let X be a scheme with an action by a group G. Associated to X are the G-equivariant K-groups for $i \in \mathbb{Z}_{\geq 0}$:

$$G \circlearrowright X \rightsquigarrow K_i^G(X).$$

When i = 0, this is

 $\begin{array}{ll} Grothendieck \mbox{ group of the exact category} \\ K_0^G(X) = & \mbox{of G-equivariant perfect complexes on X} \\ (i.e. \mbox{ bounded complexes of vector bundles}) \end{array} .$

If X is smooth, then every G-equivariant coherent sheaf has a resolution by G-equivariant vector bundles and thus

$$K_0^G(X) = {
m Grothendieck group of the category} {
m of } G$$
-equivariant coherent sheaves on X

Remark 21.2. At some future point we should expand on higher K-groups, but that time is not today. For those who are interested, Quillen's work and the groundbreaking paper

[TT90] of Thomason-Trobaugh are highly recommended. In [TT90], they show that if $U \subset X$ is open, we have a sequence²⁴

$$\cdots \to K_i(X \text{ on } Y) \to K_i(X) \to K_i(U) \to K_{i-1}(X \text{ on } Y) \to \cdots$$

which is almost exact, except that $K_0(X) \to K_0(U)$ might not be onto for singular X. (This result was already proved by Quillen for smooth X.)

In the arguments to come, we only care about K_0^G , and at certain points we will simply assert that certain boundary maps vanish; e.g., we have a short exact sequence

$$K_0^G(X \text{ on } Y) \hookrightarrow K_0^G(X) \twoheadrightarrow K_0^G(U).$$

We set $K^{G}(X) := K_{0}^{G}(X)$.

Examples

1.
$$K^G(\text{pt}) = R_G =$$
 Grothendieck group of finite-dimensional algebraic representations of G

2. We claim that $K(\mathbb{P}^1) = \mathbb{Z}[x^{\pm 1}]/(x-1)^2$. How can we see that this is true? To start, every vector bundle on \mathbb{P}^1 is a sum of line bundles. Hence there is a surjection

$$\mathbb{Z}[x^{\pm 1}] \twoheadrightarrow K(\mathbb{P}^1)$$
$$x^m \mapsto \mathcal{O}(m).$$

The tautological exact sequence of vector bundles

$$\mathcal{O}(-1) \hookrightarrow \mathcal{O}_{\mathbb{P}^1}^{\oplus 2} \twoheadrightarrow \mathcal{O}(1)$$

shows that under the surjection above,

$$x^{-1} - 2 + x \mapsto 0.$$

Hence $(x-1)^2 \mapsto 0$. This shows us that we have a map $\phi : \mathbb{Z}[x^{\pm 1}]/(x-1)^2 \to K(\mathbb{P}^1)$.

Claim: $K(\mathbb{P}^1) = \mathbb{Z}[\mathcal{O}] \oplus \mathbb{Z}[\mathcal{O}(1)].$

The arguments above imply that $[\mathcal{O}]$ and $[\mathcal{O}(1)]$ span $K(\mathbb{P}^1)$. It remains to show that they are linearly independent.

Proof 1: For X proper, K(X) carries an intersection form:

$$\langle \mathcal{F}, \mathcal{G} \rangle := \chi(\mathcal{F} \otimes \mathcal{G}),$$

 $^{^{24}}$ We can think of this like the long exact sequence in cohomology, but unlike cohomology, where the long exact sequence came almost simultaneously with its definition, this took twenty years after the initial definition of K-theory to prove.

where χ is the Euler characteristic. Computing the pairings with respect to this form, we get the following table.

$$\begin{array}{c|cccc}
 & \mathcal{O} & \mathcal{O}(1) \\
\hline
 & \mathcal{O} & 1 & 2 \\
\mathcal{O}(1) & 2 & 3 \\
\end{array}$$

We can see that the determinant is -1, so \mathcal{O} and $\mathcal{O}(1)$ are linearly independent. \Box

Proof 2: We have seen in Tom's course that

$$D^{b}(\operatorname{Coh} \mathbb{P}^{1}) \simeq D^{b} \left(\operatorname{Rep}(\bullet \Longrightarrow \bullet)\right)$$
$$\mathcal{O} \mapsto L_{0} := k \Longrightarrow 0$$
$$\mathcal{O}(1) \mapsto L_{1} := 0 \Longrightarrow k$$

The K-group of any finite length abelian category has a basis given by the classes of simple objects. In the case of Rep($\bullet \implies \bullet$), the two simple objects are L_0 and L_0 . Hence we have

$$K(\mathbb{P}^1) = K(D^b(\operatorname{Coh}(\mathbb{P}^1))) = K\left(D^b\left(\operatorname{Rep}(\bullet \Longrightarrow \bullet)\right)\right) = \mathbb{Z}[L_0] \oplus \mathbb{Z}[L_1]. \quad \Box$$

Remark 21.3. The second proof illustrates a common phenomenon: derived equivalences can have interesting consequences for K-theory.

Fact: Proper maps $f: X \to Y$ induce maps in K-theory:

$$p_*: K^G(X) \to K^G(Y)$$

The map p_* is given by $p_*(F) = \sum (-1)^i R^i f_*(F)$.

Example 21.4. 1. Projection to a point, $p : \mathbb{P}^1 \to \text{pt}$, induces the following map on (non-equivariant) *K*-theory:

$$p_*: K(\mathbb{P}^1) \cong \mathbb{Z}[x^{\pm 1}]/(x-1)^2 \to K(\mathrm{pt}) \cong \mathbb{Z}$$

 $x^m \mapsto \chi(\mathcal{O}(m)) = m+1$

2. Let $B = \left\{ \begin{pmatrix} * & * \\ 0 & * \end{pmatrix} \right\} \subset SL_2$ act on \mathbb{P}^1 in the standard way. Then the map that p induces on B-equivariant K-theory is given by the Weyl character formula. Indeed, we can identify

$$K^{G}(\mathbb{P}^{1}) = K^{B}(\mathrm{pt}) = \mathbb{Z}[X(B)] = \mathbb{Z}[x^{\pm 1}].$$

Then under the composition

$$K^{G}(\mathbb{P}^{1}) \to K^{G}(\mathrm{pt}) \to K^{B}(\mathrm{pt}) = \mathbb{Z}[x^{\pm 1}].$$

the element $x^m \in K^B(\mathbb{P}^1) \cong \mathbb{Z}[x^{\pm 1}]$ maps to $\frac{x^m - x^{-m-2}}{1 - x^{-2}} \in K^B(\text{pt}) \cong \mathbb{Z}[x^{\pm 1}].$

21.2 Equivariant *K*-theory of the Steinberg variety

Let

$$G \supset B \supset T$$

be a complex reductive group containing a Borel subgroup containing a maximal torus. Let X = X(T) be the character lattice, W_f the finite Weyl group, and $W = W_f \ltimes \mathbb{Z}X$ the affine Weyl group of the dual²⁵ root system. Let

$$\mathcal{N} \subset \mathfrak{g}$$

be the nilpotent cone in the Lie algebra of G, and

$$\widetilde{\mathcal{N}} = \{(x, \mathfrak{b}) \in \mathcal{N} \times \mathcal{B} \mid x \in \mathfrak{b}\} = T^* \mathcal{B} \to \mathcal{N}$$

be the Springer resolution. Recall that the *Steinberg variety* is defined to be


$$St := \mathcal{N} \times_{\mathcal{N}} \mathcal{N} = \{ (x, \mathfrak{b}, \mathfrak{b}') \mid x \in \mathfrak{b}, x \in \mathfrak{b}' \}.$$

We also explained in Lecture 13 how this could be realized as the conormal space to the space $\mathcal{B} \times \mathcal{B}$ with the stratification by *G*-orbits (which are parameterized by *W*).

Example 21.5. Let $G = SL_2$. Then $\mathcal{B} = \mathbb{P}^1$, and the *G*-orbit stratification on $\mathcal{B} \times \mathcal{B}$ is given by

$$\mathcal{B} \times \mathcal{B} = \Delta \sqcup Y,$$

where $Y := (\mathbb{P}^1 \times \mathbb{P}^1) \setminus \Delta$. An illustration:




Then the Steinberg variety is

$$T^*_{\Delta}(\mathbb{P}^1 \times \mathbb{P}^1) \sqcup T^*_Y(\mathbb{P}^1 \times \mathbb{P}^1) = T^*\mathbb{P}^1 \sqcup Y.$$

 $^{^{25}}$ This convention is introduced to avoid including many checks (to indicate Langlands dual groups) in the rest of this lecture and the next.

An illustration:



In the "easy part," projection onto each component of $\mathcal{B} \times \mathcal{B}$ is an isomorphism. In the "tricky part," projection onto each component hits the zero section in $T^*\mathbb{P}^1$.

The group $G \times \mathbb{C}^{\times}$ acts on St via


$$(g,z) \cdot (z,\mathfrak{b},\mathfrak{b}') = (z^2gx, g\mathfrak{b}, g\mathfrak{b}').$$
(21.1)

Algebra structure on $K^{G \times \mathbb{C}^{\times}}(St)$:

Recall that $K^{G \times \mathbb{C}^{\times}}(St)$ has an algebra structure given by convolution in $D^{b}(Coh)$ (see Lectures 14 and 15 for a refresher):

$$\mathcal{F} * \mathcal{G} := p_{13*}(p_{12}^* \mathcal{F} \otimes p_{23}^* \mathcal{G}),$$

where p_{ij} are the canonical projections




Module structure on $K^{G \times \mathbb{C}^{\times}}(\widetilde{\mathcal{N}})$:

Similarly, convolution also gives $K^{G \times \mathbb{C}^{\times}}(\widetilde{\mathcal{N}})$ the structure of a module for $K^{G \times \mathbb{C}^{\times}}(St)$:

$$\mathcal{F} * \mathcal{G} := p_{1*}(\mathcal{F} \times p_2^* \mathcal{G}),$$

where



Recall that our goal is to prove that

$$K^{G \times \mathbb{C}^{\times}}(St) \cong H.$$

We will accomplish that through a subgoal, which is to show that

 $K^{G \times \mathbb{C}^{\times}}(\widetilde{\mathcal{N}}) \cong$ anti-spherical module.

We can work toward the subgoal by examining the vector space structure of $K^{G \times \mathbb{C}^{\times}}(\widetilde{\mathcal{N}})$ more closely.

First, note that we have a map

$$K^{G \times \mathbb{C}^{\times}}(\widetilde{\mathcal{N}}) \xleftarrow{pull}{back} K^{G \times \mathbb{C}^{\times}}(\mathcal{B}) = K^{B \times \mathbb{C}^{\times}}(\mathrm{pt}) = \mathbb{Z}[v^{\pm 1}][X].$$

Because $\widetilde{\mathcal{N}}$ is a vector bundle over \mathcal{B} , pull-back is an isomorphism. Hence

$$K^{G \times \mathbb{C}^{\times}}(\widetilde{\mathcal{N}}) = \mathbb{Z}[v^{\pm 1}][X].$$

Now we turn to the vector space structure on $K^{G \times \mathbb{C}^{\times}}(St)$. The space $\mathcal{B} \times \mathcal{B}$ has a filtration via closures of G-orbits:

$$\emptyset = Z_{-1} \subset Z_0 \subset \cdots \subset Z_m = \mathcal{B} \times \mathcal{B},$$

with $Z_i \setminus Z_{i+1} \simeq G \cdot (x_i B, B) \simeq G \times_B X_i$, where $X_i = B x_i B / B$ is a Bruhat cell. This induces a filtration of the Steinberg variety:

$$\emptyset = \widetilde{Z}_{-1} \subset \widetilde{Z}_0 \subset \cdots \subset \widetilde{Z}_m = St,$$

with $\widetilde{Z}_i \setminus \widetilde{Z}_{i+1} \simeq T^*_{X_i}(\mathcal{B} \times \mathcal{B}) =: T^*_{x_i}$. We have

$$K^{G \times \mathbb{C}^{\times}}(T_{x_i}^*) = K^{G \times \mathbb{C}^{\times}}(X_i) = K^{B \times \mathbb{C}^{\times}}(X_i) = K^{B \times \mathbb{C}^{\times}}(\mathrm{pt}).$$

All boundary maps vanish when we apply $K^{G \times \mathbb{C}^{\times}}$ to the above filtration, and a little work (see [CG09]) yields:

$$K^{G \times \mathbb{C}^{\times}}(St) = \bigoplus_{x \in W_f} \mathbb{Z}[v^{\pm 1}][X][\mathcal{O}_{\overline{T_{x_i}^*}}].$$

Remark 21.6. Compare this to the decomposition

$$H = \bigoplus_{x \in W_f} \mathbb{Z}[v^{\pm 1}][X]T_x.$$

21.3 The spherical and anti-spherical modules

Recall Bernstein's presentation of the affine Hecke algebra (Theorem 12.5): H is an algebra over $\mathbb{Z}[v^{\pm 1}]$ with generators

$$\begin{cases} H_s \mid s \in S_f \\ \text{(finite part)} \end{cases} \quad \text{and} \quad \begin{cases} \theta_\lambda \mid \lambda \in X \\ \text{(lattice part)} \end{cases}.$$

The generators $\{H_s \mid s \in S_f\}$ generate a copy of the finite Hecke algebra H_f and the generators $\{\theta_\lambda \mid \lambda \in X\}$ generate $\mathbb{Z}[v^{\pm 1}][X]$. The relations are:

- 1. $H_s^2 = (v^{-1} v)H_1 + 1$ for all $s \in S$ + braid relations (finite part),
- 2. $\theta_{\lambda}\theta_{\gamma} = \theta_{\lambda+\gamma}$ for all $\lambda, \gamma \in X$ (lattice part), and
- 3. the most important relation: For a simple reflection $s = s_{\alpha} \in S$ and $\lambda \in X$,

$$H_s \theta_{s\lambda} - \theta_{\lambda} H_s = (v - v^{-1}) \left(\frac{\theta_{\lambda} - \theta_{s\lambda}}{1 - \theta_{-\alpha}} \right)$$

The first relation can be rewritten as

$$(H_s + v)(H_s - v^{-1}) = 0,$$

which implies that H_f has two natural rank 1 modules:

$$\operatorname{triv}: H_s \mapsto v^{-1},$$
$$\operatorname{sgn}: H_s \mapsto -v.$$

From these we construct two induced *H*-modules:

 $M = H \otimes_{H_f} \text{triv},$ "spherical module"

 $N = H \otimes_{H_f} \operatorname{sgn}$, "anti-spherical module"

As modules over the lattice part, each is isomorphic to $\mathbb{Z}[v^{\pm 1}][X]$.

Remark 21.7. In what follows, we will abuse notation and write

$$\theta_{\lambda} := \theta_{\lambda} \otimes 1 \in N.$$

Using the third relation in the affine Hecke algebra, we can compute the action of the generator $\theta_s \in H$ on $\theta_{s\lambda} \in N$ in the anti-spherical module:

$$H_{s} \cdot \theta_{s\lambda} = (v - v^{-1}) \left(\frac{\theta_{\lambda} - \theta_{s\lambda}}{1 - \theta_{-\alpha}} \right) + \theta_{\lambda} H_{s}$$
$$= v \left(\frac{\theta_{\lambda} - \theta_{s\lambda} - \theta_{\lambda} + \theta_{\lambda - \alpha}}{1 - \theta_{-\alpha}} \right) - v^{-1} \left(\frac{\theta_{\lambda} - \theta_{s\lambda}}{1 - \theta_{-\alpha}} \right)$$
$$= v \left(\frac{\theta_{\lambda - \alpha} - \theta_{s\lambda}}{1 - \theta_{-\alpha}} \right) - v^{-1} \left(\frac{\theta_{\lambda} - \theta_{s\lambda}}{1 - \theta_{-\alpha}} \right).$$

This formula looks nicer if we instead compute the action in terms of the Kazhdan–Lusztig generator $b_s := H_s + v$:

$$b_{s} \cdot \theta_{s\lambda} = v \left(\frac{\theta_{\lambda-\alpha} - \theta_{s\lambda}}{1 - \theta_{-\alpha}} \right) - v^{-1} \left(\frac{\theta_{\lambda} - \theta_{s\lambda}}{1 - \theta_{-\alpha}} \right) + v \theta_{s\lambda}$$
$$= v \left(\frac{\theta_{\lambda-\alpha} - \theta_{s\lambda-\alpha}}{1 - \theta_{-\alpha}} \right) - v^{-1} \left(\frac{\theta_{\lambda} - \theta_{s\lambda}}{1 - \theta_{-\alpha}} \right)$$
$$= (v \theta_{-\alpha} - v^{-1}) \left(\frac{\theta_{\lambda} - \theta_{s\lambda}}{1 - \theta_{-\alpha}} \right).$$

In other words,

$$b_s \cdot \theta_{\lambda} = (v^{-1} - v\theta_{-\alpha}) \left(\frac{\theta_{\lambda} - \theta_{s\lambda}}{1 - \theta_{-\alpha}}\right).$$
(21.2)

We'll pick up here next week with a very similar looking computation in K-theory.

Lecture 22: Proof of the Kazhdan–Lusztig isomorphism

In today's lecture we will complete the proof of the Kazhdan–Lusztig isomorphism. First, we establish and streamline notation.

22.1 Notation and set-up

Let

 $G\supset B\supset T$

be a complex reductive group, Borel subgroup, and maximal torus. Let X be the character lattice, and $W_f \subset W$ the finite and affine Weyl groups with simple reflections $S_f \subset W_f$ and $S \subset W$. We denote by \mathcal{B} the flag variety, which we realize as the variety of Borel subalgebras of $\mathfrak{g} = \text{Lie } G$. The group G acts on the product $\mathcal{B} \times \mathcal{B}$, and the orbits give a stratification with strata parameterized by W_f :

$$G \circlearrowright \mathcal{B} \times \mathcal{B} = \bigsqcup_{x \in W_f} \mathbb{O}_x.$$

The orbit \mathbb{O}_x consists of pairs of flags/Borel subalgebras in relative position x. For example, $\mathbb{O}_{id} = \Delta$ (the diagonal in $\mathcal{B} \times \mathcal{B}$), and \mathbb{O}_{w_0} (where $w_0 \in W_f$ is the longest element) is open in $\mathcal{B} \times \mathcal{B}$.

Let

$$\widetilde{\mathcal{N}} = T^* \mathcal{B} \to \mathcal{N} \subset \mathfrak{g}$$


be the Springer resolution. The Steinberg variety is

$$St = \widetilde{\mathcal{N}} \times_{\mathcal{N}} \widetilde{\mathcal{N}} = \bigsqcup_{x \in W_f} T^*_{\mathbb{O}_x}(\mathcal{B} \times \mathcal{B}) = \bigcup_{w \in W_f} \Lambda_x,$$

where $T^*_{\mathbb{O}_x}(\mathcal{B} \times \mathcal{B})$ is the conormal bundle of the *G*-orbit \mathbb{O}_x , whose closure $\Lambda_x := \overline{T^*_{\mathbb{O}_x}(\mathcal{B} \times \mathcal{B})}$ is an irreducible component of *St*.

Last week we discussed the following example.


Example 22.1. Let $G = SL_2$, so $\mathcal{B} = \mathbb{P}^1$. The *G*-orbit stratification of $\mathcal{B} \times \mathcal{B}$ is:



The corresponding stratification of the Steinberg variety is

$$St = T^*_{\mathbb{O}_{id}}(\mathcal{B} \times \mathcal{B}) \sqcup T^*_{\mathbb{O}_s}(\mathcal{B} \times \mathcal{B}).$$

The closures of the strata give the irreducible components Λ_x , $x \in W_f$ of St, which are glued together as follows:



Remark 22.2. In general, the components range from $\Lambda_{id} \simeq \widetilde{\mathcal{N}}$ to $\Lambda_{w_0} = \mathcal{B} \times \mathcal{B}$. They are not smooth in general, and their intersection pattern is extremely complicated. For example, it is not known in general when two Λ_x and Λ_y intersect in codimension 1.

Last week we introduced the anti-spherical module

$$N = H \otimes_{H_f} \operatorname{sgn} \simeq \mathbb{Z}[v^{\pm 1}][X]$$
$$\theta_{\lambda} := \theta_{\lambda} \otimes 1 \longleftrightarrow e^{\lambda}$$

for the affine Hecke algebra H and established the **Demazure–Lusztig formula**

$$b_s \cdot \theta_{\lambda} = (v^{-1} - v\theta_{-\alpha}) \left(\frac{\theta_{\lambda} - \theta_{s\lambda}}{1 - \theta_{-\alpha}}\right).$$
(22.1)

Here $H_f = \langle H_s \rangle_{s \in S_f}$ is the finite Hecke algebra and $b_s := H_s + v$ is the Kazhdan–Lusztig generator corresponding to the simple reflection $s = s_\alpha \in S_f$.

Remark 22.3. It is not immediately obvious that the formula $\begin{pmatrix} \theta_{\lambda} - \theta_{s\lambda} \\ 1 - \theta_{-\alpha} \end{pmatrix}$ gives an element in the lattice part of H. However, by rewriting

$$\left(\frac{\theta_{\lambda}-\theta_{s\lambda}}{1-\theta_{-\alpha}}\right)=(\theta_{\lambda}+\theta_{\lambda-\alpha}+\cdots+\theta_{s\lambda+\alpha}),$$

we see that $b_s \cdot \theta_\lambda \in \mathbb{Z}[v^{\pm 1}][X]$. (Here we assume $\langle \lambda, \alpha^{\vee} \rangle \geq 0$.)


We want to show that a similar formula holds for the $K^{G \times \mathbb{C}^{\times}}(St)$ -action on $K^{G \times \mathbb{C}^{\times}}(\widetilde{\mathcal{N}})$. We will first do so in the special case of $G = SL_2$, then move on to the general formulation.

22.2 The case of SL_2

We start with a baby version of the calculation. We identify $K^G(\mathbb{P}^1)$ with $\mathbb{Z}[x^{\pm 1}]$ via the chain of isomorphisms

$$K^G(\mathbb{P}^1) \simeq K^G(G/B) \simeq K^B(\mathrm{pt}) \simeq \mathbb{Z}[x^{\pm 1}]$$

Under this identification, $\mathcal{O}(m) \mapsto x^m$. We have a Cartesian square



Claim 1: The map $p_{1*}p_2^*: K^G(\mathbb{P}^1) \to K^G(\mathbb{P}^1)$ is given by $x^m \mapsto \frac{x^m - x^{-m-2}}{1 - x^{-2}}$.

Proof. By smooth base change, $p_{1*}p_2^* = p_2'^*p_{1*}'$. Then the claim follows from Weyl's character formula.

Claim 2:
$$p_{1*}(\mathcal{O}(-2,0) \otimes p_2^*(-)) : K^G(\mathbb{P}^1) \to K^G(\mathbb{P}^1)$$
 is given by $x^m \mapsto x^{-2}\left(\frac{x^m - x^{-m-2}}{1 - x^{-2}}\right)$.


Proof. The notation $\mathcal{O}(m, n)$ refers to $p_1^*\mathcal{O}(m) \otimes p_2^*\mathcal{O}(n)$. By the projection formula, $p_{1*}(p_1^*(-2) \otimes p_2^*(-)) = \mathcal{O}(-2) \otimes p_{1*}p_2^*(-)$. The claim then follows from Claim 1.

Now we move up to $\widetilde{\mathcal{N}}$. Recall that the pull-back q^* of the projection $\widetilde{\mathcal{N}} = T^* \mathbb{P}^1 \xrightarrow{q} \mathbb{P}^1$ gives an isomorphism

$$K^{G \times \mathbb{C}^{\times}}(\widetilde{\mathcal{N}}) \xleftarrow{\sim}_{q^*} K^{G \times \mathbb{C}^{\times}}(\mathbb{P}^1) = \mathbb{Z}[v^{\pm 1}][X].$$

Recall that \mathbb{C}^{\times} acts on $\widetilde{\mathcal{N}}$ via scaling by z^2 (equation (21.1)). For the rest of this section, set $\mathcal{O} := \mathcal{O}_{\widetilde{\mathcal{N}}}$.

We have projections



 Set

$$Q_s := \mathcal{O}_{\mathbb{P}^1 \times \mathbb{P}^1}(-2, 0)$$

on $\Lambda_s \simeq \mathbb{P}^1 \times \mathbb{P}^1 \subset St$. We will see that (up to some simple scalar factors), Q_s acts via convolution on $K^{G \times \mathbb{C}^{\times}}(St)$ in the same way that b_s acts on N. To do so, we want to calculate

$$p_{1*}(Q_s \otimes p_2^* \mathcal{O}(m)) \in K^{G \times \mathbb{C}^{\times}}(\widetilde{\mathcal{N}}).$$

Claim 3: Let $\mathbb{P}^1 \xrightarrow{i} \widetilde{\mathcal{N}}$. Then $p_{1*}(Q_s \otimes p_2^* \mathcal{O}(m)) = \left(\frac{x^m - x^{-m-2}}{1 - x^{-2}}\right) [i_* \mathcal{O}_{\mathbb{P}^1}(-2)].$

Exercise 22.4. 1. Suppose $\pi : E \to Y$ is a vector bundle, \mathcal{F} a quasi-coherent sheaf on Y, \mathcal{G} a locally free sheaf on Y, and $i: Y \hookrightarrow E$ the zero section. Show that

$$i_*\mathcal{F}\otimes\pi^*\mathcal{G}=i_*(\mathcal{F}\otimes\mathcal{G}).$$

2. Use 1. applied to $\widetilde{\mathcal{N}} \times \widetilde{\mathcal{N}} \to \mathbb{P}^1 \times \mathbb{P}^1$ to deduce Claim 3.

It remains to express $[i_*\mathcal{O}_{\mathbb{P}^1}(-2)]$ in terms of our basis of $K^{G\times\mathbb{C}^{\times}}(\widetilde{\mathcal{N}})$; that is, to understand $i_*\mathcal{O}_{\mathbb{P}^1}(-2)$ in terms of vector bundles.

Useful basic technique: Given a vector bundle $q: V \to Y$, there is a bijection

$$\left\{\begin{array}{c} \text{quasi-coherent sheaves}\\ \text{on the total space of } V\end{array}\right\} \simeq \left\{\begin{array}{c} \text{quasicoherent sheaves of}\\ \text{modules over } q_*\mathcal{O}_V \simeq \text{Sym}^{\bullet}(V^*)\end{array}\right\}.$$

(More generally, this holds for any affine morphism, see Hartshorne.)

In our case, the projection

$$\widetilde{\mathcal{N}} = T^* \mathbb{P}^1 \to \mathbb{P}^1$$

is the line bundle associated to $\mathcal{O}(-2)$, and

$$q_*\mathcal{O}\simeq \operatorname{Sym}^{\bullet}(\mathcal{O}_{\mathbb{P}^1}(2))=\mathcal{O}_{\mathbb{P}^1}\oplus \mathcal{O}_{\mathbb{P}^1}(2)\oplus \mathcal{O}_{\mathbb{P}^1}(4)\oplus\cdots$$

Additionally, \mathbb{C}^{\times} acts on $T^*\mathbb{P}^1 \to \mathbb{P}^1$ as multiplication by z^2 along the fibres, which corresponds to multiplication by z^{-2} in the second factor in the direct sum decomposition above. Hence we have an exact sequence

$$z^{-2}\mathcal{O}(2) \hookrightarrow \mathcal{O} \twoheadrightarrow i_*\mathcal{O}_{\mathbb{P}^1}.$$

Remark 22.5. Locally, this is the short exact sequence

$$k[x] \stackrel{\cdot x}{\hookrightarrow} k[x] \twoheadrightarrow k_{x}$$

but globally, we have some twisting. This is a special example of the Koszul resolution of the zero section of a vector bundle.

Hence in $K^{G \times \mathbb{C}^{\times}}(\widetilde{\mathcal{N}})$, we have

$$[i_*\mathcal{O}_{\mathbb{P}^1}] = [\mathcal{O}] - v^{-2}[\mathcal{O}(2)].$$

By tensoring with $\mathcal{O}(-2)$ we obtain

$$[i_*\mathcal{O}_{\mathbb{P}^1}(-2)] = [\mathcal{O}(-2)] - v^{-2}[\mathcal{O}].$$

Putting it all together, we see that

$$x^m \xrightarrow{[Q_s]^*} (x^{-2} - v^{-2}) \left(\frac{x^m - x^{-m-2}}{1 - x^{-2}}\right).$$

Hence

$$x^m \xrightarrow{-[vQ_s]^*} (v^{-1} - vx^{-2}) \left(\frac{x^m - x^{-m-2}}{1 - x^{-2}}\right)$$

We only need two last pieces of information to establish the Kazhdan–Lusztig isomorphism for SL₂, which we leave as exercises.

Exercise 22.6. The map $\theta_m \mapsto x^{m-1}$ intertwines the actions of b_s and $-[vQ_s]*$.

Exercise 22.7. Both the representation N of H and the representation $K^{G \times \mathbb{C}^{\times}}(\widetilde{\mathcal{N}})$ of $K^{G \times \mathbb{C}^{\times}}(St)$ are faithful.

22.3 The general case

Roughly speaking, in the SL_2 example, we have seen the meaning in K-theory of the Demazure–Lusztig formula


$$b_s \cdot \theta_\lambda = (v^{-1} - v\theta_{-\alpha}) \left(\frac{\theta_\lambda - \theta_{s\lambda}}{1 - \theta_{-\alpha}}\right).$$

The first factor $(v^{-1} - v\theta_{-\alpha})$ comes from the "Koszul resolution," and the second factor $\left(\frac{\theta_{\lambda}-\theta_{s\lambda}}{1-\theta_{-\alpha}}\right)$ from the "push-pull for \mathbb{P}^1 ". We will now explain that the same philosophy works in general.

Relative cotangent bundle: Let

 $Z \xrightarrow{f} Y$

be a smooth map. (Think submersion = fibration.)



The relative tangent bundle is

$$T_{Z/Y} := \ker(df : TZ \to TY).$$


We have a short exact sequence

$$T_{Z/Y} \hookrightarrow T_Z \twoheadrightarrow f^*T_Y.$$

Dually,

$$f^*T^*_Y \hookrightarrow T^*_Z \twoheadrightarrow T^*_{Z/Y}.$$


A caricature:



In our setting, we have a \mathbb{P}^1 fibration

 $\mathcal{B} \xrightarrow{\pi_s} \mathcal{B}_s,$

where \mathcal{B}_s is the variety of parabolic subalgebras of type s.



Fundamental Cartesian diagram:


$$\begin{array}{c} \overline{\mathbb{O}}_s \xrightarrow{p_1} \mathcal{B} \\ p_2 \downarrow \qquad \qquad \downarrow \\ \mathcal{B} \xrightarrow{\pi_s} \mathcal{B}_s \end{array}$$

Here $\mathbb{O}_s \subset \mathcal{B} \times \mathcal{B}$ is the *G*-orbit consisting of pairs of flags in relative position *s*. All maps in this diagram are \mathbb{P}^1 -fibrations. We identify $K^G(\mathcal{B}) \simeq \mathbb{Z}[X]$ via $\mathcal{O}_{\mathcal{B}}(\lambda) \mapsto e^{\lambda}$.

Claim 4: $\pi_{s*}\pi_s^*: K^G(\mathcal{B}) \to K^G(\mathcal{B})$ maps $e^{\lambda} \mapsto \frac{e^{\lambda} - e^{s(\lambda) - \alpha}}{1 - e^{-\alpha}}.$

Proof. Again, this is simply an instance of the Weyl character formula for \mathbb{P}^1 .

We claim that the corresponding Cartesian diagram "upstairs" is



To see that this is the correct lift of the first Cartesian square, we start with the lower right corner

$$T^*\mathcal{B}_s = \{(\mathfrak{p}, x) \mid x \in \operatorname{nilrad}(\pi_s(\mathfrak{b}))\},\$$


which implies that the upper right and lower left corners are

$$\widetilde{\mathcal{N}}_s = \pi_s^* T^*_{\mathcal{B}/\mathcal{B}_s} = \{(\mathfrak{b}, x) \mid x \in \operatorname{nilrad}(\pi_s(\mathfrak{b}))\} \subset \widetilde{\mathcal{N}}.$$

Hence the fibre product completing the diagram in the upper left corner is given by

$$\left\{ (\mathfrak{b}, \mathfrak{b}', x) \mid \begin{array}{c} \mathfrak{b}, \mathfrak{b}' \text{ in relative position } s, \text{ and} \\ x \in \operatorname{nilrad}(\pi_s(\mathfrak{b})) = \operatorname{nilrad}(\pi_s(\mathfrak{b}')) \end{array} \right\} = \Lambda_s.$$

Together, these two Cartesian squares form a "Cartesian cube:"



Define

$$Q_s := q^* \Omega_{\overline{\mathbb{O}}_s/\mathcal{B}}$$

to be the relative 1-forms with respect to the second projection. As earlier, Claim 4 gives

$$p_{1*}(Q_s \otimes p_2^* \mathcal{O}(\lambda)) = \left(\frac{e^{\lambda} - e^{s(\lambda) - \alpha}}{1 - e^{-\alpha}}\right) \left[i_* \mathcal{O}_{\widetilde{\mathcal{N}}_s}(-\alpha)\right],$$

where $\widetilde{\mathcal{N}}_s \hookrightarrow \widetilde{\mathcal{N}}$ is the inclusion. What remains is to express $[i_*\mathcal{O}_{\widetilde{\mathcal{N}}_s}(-\alpha)] \in K^{G \times \mathbb{C}^{\times}}(\widetilde{\mathcal{N}})$ in the basis of line bundles. Again, we can do so using a Koszul-type resolution. There is a short exact sequence of B-modules

$$\mathfrak{p}_s/\mathfrak{b} \hookrightarrow \mathfrak{g}/\mathfrak{b} \twoheadrightarrow \mathfrak{g}/\mathfrak{p}_s.$$

The corresponding exact sequence of vector bundles on $\mathcal{B} = G/B$ is

$$L_{\alpha} \hookrightarrow T_{\mathcal{B}} \twoheadrightarrow \pi_s^* T_{\mathcal{B}_s}$$

where L_{α} is the line bundle on \mathcal{B} associated to $\alpha \in X$. Passing to symmetric algebras, we obtain the Koszul resolution

$$z^{-2}\operatorname{Sym}^{\bullet}(T_{\mathcal{B}})(\alpha) \hookrightarrow \operatorname{Sym}^{\bullet}(T_{\mathcal{B}}) \twoheadrightarrow \operatorname{Sym}^{\bullet}(\pi_s^*T_{\mathcal{B}}).$$

Remark 22.8. This should be thought of as the vector bundle version of the short exact sequence

$$k[x_1,\ldots,x_n] \xrightarrow{\cdot x_n} k[x_1,\ldots,x_n] \twoheadrightarrow k[x_1,\ldots,x_{n-1}].$$

The z^{-2} comes from the \mathbb{C}^{\times} -action.

In other words,

$$z^{-2}\mathcal{O}_{\widetilde{\mathcal{N}}}(\alpha) \hookrightarrow \mathcal{O}_{\widetilde{\mathcal{N}}} \twoheadrightarrow i_*\mathcal{O}_{\widetilde{\mathcal{N}}_s}.$$

The result:

$$[i_*\mathcal{O}_{\widetilde{\mathcal{N}}_s}(-\alpha)] = [\mathcal{O}_{\widetilde{\mathcal{N}}}(-\alpha)] - v^{-2}[\mathcal{O}_{\widetilde{\mathcal{N}}}].$$

From here, the proof follows from two exercises analogous to those in the previous section.

Exercise 22.9. Check that $\theta_{\lambda} \mapsto e^{\lambda \pm \rho}$ intertwines $b_s \cdot$ and $-[vQ_s]*$.

Exercise 22.10. Complete the proof by showing that N (resp. $K^{G \times \mathbb{C}^{\times}}(\widetilde{\mathcal{N}})$) are faithful modules over H (resp. $K^{G \times \mathbb{C}^{\times}}(\widetilde{\mathcal{N}})$).

Remark 22.11. Geordie isn't quite sure about whether we should have α or $-\alpha$ above, so the reader should take the final arguments with a grain of salt.

Lecture 23: Gaitsgory's central sheaves

Last week we finished the proof of the Kazhdan–Lusztig isomorphism:

$$K^{G^{\vee} \times \mathbb{C}^{\times}}(St) \simeq H.$$

This is an isomorphism of algebras. We established this isomorphism by proving that two faithful modules for these algebras are isomorphic:

$$K^{G^{\vee} \times \mathbb{C}^{\times}}(\widetilde{\mathcal{N}}) \simeq N = H \otimes_{H_f} \text{sgn.}$$
 (23.1)

For the next several weeks (months?), we will work toward categorifying the Kazhdan– Lusztig isomorphism via Bezrukavnikov's equivalence. The first step towards a categorification is the Arkhipov-Bezrukavnikov theorem, which categorifies (23.1).

Theorem 23.1. [AB09]

Here D_{IW}^{mix} and D_{IW} are the mixed and unmixed version of the antispherical category.

Remark 23.2. If one wants to convince oneself that this is a deep equivalence, one only needs to note that the "easy" \mathbb{C}^{\times} -action on the LHS corresponds to the "hard" Frobenius action on the RHS!

For the remainder of the lecture, we will build the machinery of Gaitsgory's central sheaves, then focus on the example of the natural representation of GL_2 . We will return to the Arkhipov-Bezrukavnikov theorem next week.

23.1 A change of notation in the Hecke algebra

We will make a slight change of notation from previous lectures. In the affine Hecke algebra, replace

$$H_x \rightsquigarrow \delta_x$$

for all $x \in W_f$, so our standard basis of the finite Hecke algebra is now called $\{\delta_x\}_{x \in W_f}$ and the Kazhdan–Lusztig basis is $\{b_x\}_{x \in W_f}$. We make this change so that from now on, small letters indicate objects in the Hecke algebra H and big letters indicate objects in the Hecke category \mathcal{H} .

23.2 Gaitsgory's central sheaves

For more details on this construction, see notes from Emily's IFS talks on the Beilinson-Drinfeld Grassmannian²⁶.

 $^{^{26}}$ Talk notes available at https://sites.google.com/view/ifssydney/home

Recall that within the affine Hecke algebra H, we have a large commutative subalgebra

$$\mathcal{L} = \bigoplus_{\lambda \in X^{\vee}} \mathbb{Z}[v^{\pm 1}] \theta_{\lambda} \subset H.$$

In Lecture 12 we discussed Bernstein's description of the center of H in terms of this subalgebra.


Theorem 23.3.

$$Z(H) \simeq \mathcal{L}^{W_f} \simeq \mathbb{Z}[v^{\pm 1}] \otimes_{\mathbb{Z}} [\operatorname{Rep} G^{\vee}].$$


Another perspective from which we can view this description of the center of the affine Hecke algebra was discussed in Emily's IFS talk. Temporarily, let $\mathcal{K} = \mathbb{F}_q((t)) \supset \mathcal{O} = \mathbb{F}_q[[t]]$, and build Hecke algebras

$$\mathcal{H}^{\mathrm{aff}} = \mathrm{Fun}_{I}^{\mathrm{c.s.}}(\mathcal{F}\ell, \mathbb{C}), \quad \mathcal{H}^{\mathrm{sph}} = \mathrm{Fun}_{G(\mathcal{O})}^{\mathrm{c.s.}}(\mathcal{G}r, \mathbb{C}).$$

Here "c.s." stands for "compactly supported", $\mathcal{F}\ell = G(\mathcal{K})/I$ is the affine flag variety and $\mathcal{G}r = G(\mathcal{K})/G(\mathcal{O})$ is the affine Grassmannian. These function spaces gain the structure of algebras via convolution. The spherical Hecke algebra $\mathcal{H}^{\mathrm{sph}}$ is a commutative convolution algebra, $\mathcal{H}^{\mathrm{aff}}$ is not. There are natural maps



Under these maps, the image of the center $Z(\mathcal{H}^{aff})$ and the image of \mathcal{H}^{sph} agree:



This provides an isomorphism

$$Z(\mathcal{H}^{\mathrm{aff}}) \simeq \mathcal{H}^{\mathrm{sph}}.$$
 (23.2)

The analogous statement for the affine Hecke algebra H is as follows. Let $w_f \in W_f$ be the longest element, and

$$b_{w_f} = \sum_{x \in W_f} v^{\ell(w_f) - \ell(x)} \delta_x \in H$$


the corresponding Kazhdan–Lusztig basis element. Define

$$H^{\mathrm{sph}} := (b_{w_f} \cdot H) \cap (H \cdot b_{w_f}).$$

By the Satake isomorphism,

$$H^{\mathrm{sph}} \simeq \mathcal{L}^{W_f}$$

Then we have maps



This gives us the isomorphism in Theorem 23.3:

$$Z \xrightarrow{\sim} H^{\mathrm{sph}}$$
$$h \mapsto h \cdot b_w$$

Gaitsgory lifted this statement to the level of categories. For the remainder of the lecture set $\mathcal{O} = \mathbb{C}[[t]], \mathcal{K} = \mathbb{C}((t)), \ \mathcal{G}r = G(\mathcal{K})/G(\mathcal{O}), \ \mathcal{F}\ell = G(\mathcal{K})/I.$ Recall that

 $\pi: \mathcal{F}\ell \to \mathcal{G}r$

is a G/B-fibration. We replace


$$\mathcal{H}^{\mathrm{sph}} \rightsquigarrow \mathrm{Perv}_{G(\mathcal{O})}(\mathcal{G}r),$$
 "Satake category," and $\mathcal{H}^{\mathrm{aff}} \rightsquigarrow \mathrm{Perv}_{I}(\mathcal{F}\ell),$ "Hecke category".

Both of these categories obtain a monoidal structure, see Emily's IFS talk. Gaitsgory upgraded the isomorphism 23.2 to a central functor.

Theorem 23.4. There exists a central functor²⁷

$$Z : \operatorname{Perv}_{G(\mathcal{O})}(\mathcal{G}r) \to \operatorname{Perv}_I(\mathcal{F}\ell).$$

Moreover, the diagram



commutes.

²⁷This means that for all $\mathcal{F} \in \operatorname{Perv}_{G(\mathcal{O})}(\mathcal{G}r), \mathcal{G} \in \operatorname{Perv}_{I}(\mathcal{F}\ell), Z(\mathcal{F}) * \mathcal{G}$ is perverse, so is $\mathcal{G} * Z(\mathcal{F})$, and we have a canonical isomorphism $Z(\mathcal{F}) * \mathcal{G} \simeq \mathcal{G} * Z(\mathcal{F})$.

- **Remark 23.5.** 1. It is best to think of a "central functor to a monoidal category (M, *)" as a "functor to the Drinfeld centre of (M, *)". Geordie is not 100% sure, but he thinks that Z actually lands in the *symmetric center* (i.e. set of objecs with symmetric braiding) of the Drinfeld center.
 - 2. Everything that Bezrukavnikov does is built on Z, so we should get used to thinking about it!
 - 3. If $\mathcal{G}', \mathcal{G} \in \operatorname{Perv}_I(\mathcal{F}\ell)$, then usually $\mathcal{G}' * \mathcal{G}$ is not perverse. It's a miracle that $Z(\mathcal{F}) * \mathcal{G}$ always is!

23.3 Extended example: the natural representation of GL_2

For the remainder of this lecture, we will work out the details of this construction for the simplest non-trivial example: the natural representation of GL_2 . This example is very beautiful and instructive, so we will discuss it in detail.

Set $G = \operatorname{GL}_2$, $G^{\vee} = \operatorname{GL}_2$, and nat = the natural representation of GL_2 . Let $\mathcal{F}_{\operatorname{nat}} \in \operatorname{Perv}_{G(\mathcal{O})}(\mathcal{G}r)$ be the corresponding perverse sheaf under geometric Satake. Our goal is to describe $Z(\mathcal{F}_{\operatorname{nat}})$. Here is an outline of what we'll do:

Algebra: (See exercises at end of section.) To start, we have

$$H_{\mathrm{GL}_2} = H_{\mathrm{SL}_2} \ltimes \langle \delta_{\varpi} \rangle = \langle \delta_s, \delta_{s_0}, \delta_{\varpi} \rangle = H_f \otimes \mathbb{Z}[v^{\pm 1}][\theta_1, \theta_2].$$

Here δ_{ϖ} is the generator of length zero elements, $s \in S_f$ is the finite simple reflection, and s_0 is the affine simple reflection. The elements $\theta_1 = \delta_{\varpi} \delta_s$ and $\theta_2 = \delta_{\varpi} \delta_{s_0}^{-1}$ are the Bernstein generators. Hence


$$z_{\text{nat}} = \theta_1 + \theta_2 = \delta_{\varpi}(\delta_s + \delta_{s_0}^{-1}) = \delta_{\varpi}(\delta_s^{-1} + \delta_{s_0}).$$

A natural question arises: How can we produce z_{nat} geometrically?


Geometry: Let $k \in \{\mathbb{Q}, \mathbb{R}, \mathbb{C}\}$. Both $\mathcal{F}\ell$ and $\mathcal{G}r$ have components which are indexed by \mathbb{Z} (measuring the valuation of the determinant of the lattice/the polynomial degree of the representation under geometric Satake), and the projection

$$\mathcal{F}\ell_{\mathrm{GL}_2} = \mathcal{F}\ell_{\mathrm{SL}_2} \times \mathbb{Z} \to \mathcal{G}r_{\mathrm{GL}_2} = \mathcal{G}r_{\mathrm{SL}_2} \times \mathbb{Z}$$

is a \mathbb{P}^1 fibration. The natural representation corresponds to a sheaf $k_{\mathbb{P}^1}[1]$ on $\mathcal{G}r_{\mathrm{SL}_2} \times \{1\}$. Hence all of our arguments will take place on the components $\mathcal{G}r_{\mathrm{SL}_2} \times \{1\}$ and $\mathcal{F}\ell_{\mathrm{SL}_2} \times \{1\}$, so we can ignore the rest. Gaitsgory's family: Gaitsgory constructs a family of varieties over the disc:



The $\mathbb{P}^1 \times \{B/B\}$ on which the sheaf $k_{\mathbb{P}^1}[1]$ is supported in the fibre $\mathcal{G}r \times G/B$ degenerates to $\mathbb{P}^1 \times \mathbb{P}^1$ (the intersection of two Schubert curves) in $\mathcal{F}\ell$:



We will see that $z_{\rm nat}$ is categorified by nearby cycles. Moreover, we will see that the "Wakimoto filtration"

$$Z(\mathcal{J}_{nat}) = \left(\begin{array}{c} \Delta_{e_1} \\ \Delta_{e_2} \end{array} \right) = \left(\begin{array}{c} \nabla_{e_1} \\ \nabla_{e_2} \end{array} \right)$$

categorifies the relationships

$$z_{\rm nat} = \delta_{\varpi} \delta_s + \delta_{\varpi} \delta_{s_0}^{-1} = \delta_{\varpi} \delta_s^{-1} + \delta_{\varpi} \delta_{s_0}$$

in H. Here it goes!

Beilinson gluing: We want to understand perverse sheaves on $\{xy = 0\} \subset \mathbb{P}^2$; i.e. two \mathbb{P}^1 's meeting transversally at a point. To do so, we can start by examining perverse sheaves on $\{xy = 0\} \subset \mathbb{C}^2$, with stratification

$$\Lambda = \{0\} \sqcup \mathbb{C}_{x-\mathrm{axis}}^{\times} \sqcup \mathbb{C}_{y-\mathrm{axis}}^{\times}.$$

Beilinson gluing (Lectures 17 and 18) tells us how! Take f = x + y. The zero locus of f restricted to $\{xy = 0\}$ is $\{0\}$. By Beilinson gluing,

$$\operatorname{Perv}_{\Lambda}(\{xy=0\}) = \left\{ \begin{array}{ccc} \mathcal{F} \text{ perverse on } \mathbb{C}_{x}^{\times} \sqcup \mathbb{C}_{y}^{\times}, \\ V_{0} \text{ perverse on } \{0\} \end{array} + \begin{array}{c} \psi_{f}(\mathcal{F}) \xrightarrow{\mu-1} \psi_{f}(V) \\ V_{0} \end{array} \right\}$$
$$= \left\{ \begin{array}{ccc} \mu_{y} \bigcirc V_{y} & V_{x} \bigcirc \mu_{x} \\ V_{y} & V_{y} & V_{x} \bigcirc \mu_{x} \end{array} \middle| \begin{array}{c} \text{for all } z \in \{x, y\}, \\ v_{z} \circ c_{z} = \mu_{z} - 1, \text{ and} \\ c_{y} \circ v_{x} = 0 = c_{x} \circ v_{y} \end{array} \right\}.$$

Hence if $\overline{X} = \mathbb{P}^1_x \cup \mathbb{P}^1_y = \{xy = 0\} \subset \mathbb{P}^2$ (these will be our intersection of Schubert varieties later), and

$$\Lambda = \{0\} \sqcup \mathbb{C}_x \sqcup \mathbb{C}_y$$


then we obtain the same description as above, except that the point at infinity forces $\mu_x = \mu_y = 1$. In other words,

$$\operatorname{Perv}_{\Lambda}(\overline{X}) = \left\{ \begin{array}{c|c} V_y & V_x \\ & \swarrow & V_y \\ & & \swarrow & V_y \\ & & & V_0 \end{array} \middle| \begin{array}{c} v_z \circ c_z = 0 \\ \text{for } z \in \{x, y\}, \text{ and} \\ & c_y \circ v_x = 0 = c_x \circ v_y \end{array} \right\}.$$


Exercise 23.6. Let

$$\mathcal{Y} = \{((x:y:z),\lambda) \in \mathbb{P}^2 \times \mathbb{A}^1 \mid xy = \lambda z^2\} \subset \mathbb{P}^2 \times \mathbb{A}^1,$$

and $\mathcal{Y}_0 = \mathcal{Y} \setminus f^{-1}(0)$, where $f : \mathbb{P}^2 \times \mathbb{A}^1 \to \mathbb{A}^1$ is projection:




1. Show that $\psi_f k_{\mathcal{Y}_0}[2]$ is described under the equivalence above by the diagram



2. Deduce that $\psi_f k_{\mathcal{Y}_0}[2]$ has composition series

IL.
I(*@I(?
II.

3. Check that the monodromy μ is given by



$$\mathbb{C}_x \stackrel{j}{\hookrightarrow} \overline{X} \stackrel{j'}{\longleftrightarrow} \mathbb{C}_y$$

be the natural inclusions. Then

$$\mathbb{Z}(\mathfrak{F}_{nat}) = \underbrace{\left(\begin{array}{c} \mathfrak{i}_{1} \mathfrak{k}_{c_{3}} \\ \mathfrak{i}_{p} \mathfrak{k}_{c_{x}} \end{array} \right)}_{\mathfrak{i}_{p} \mathfrak{k}_{c_{x}}} = \underbrace{\left(\begin{array}{c} \mathfrak{i}_{1} \mathfrak{k}_{c_{x}} \\ \mathfrak{i}_{2} \mathfrak{k}_{c_{3}} \end{array} \right)}_{\mathfrak{i}_{p} \mathfrak{k}_{c_{3}}}$$

This is the Wakimoto filtration, which categorifies the relation

$$z_{\rm nat} = \delta_{\varpi} \delta_s + \delta_{\varpi} \delta_{s_0}^{-1} = \delta_{\varpi} \delta_s^{-1} + \delta_{\varpi} \delta_{s_0}.$$

in H.

Connect this all to Gaitsgory's picture: Let \mathscr{E}_{triv} be the trivial rank 2 vector bundle on \mathbb{A}^1 , and $\mathcal{L}_{triv} = \mathbb{C}[t]e_1 \oplus \mathbb{C}[t]e_2$ the trivial lattice in $\mathbb{C}((t))^2$. In Emily's IFS talks, she introduced the Beilison–Drinfeld Grassmannian, \mathcal{BD} , which provides a $\mathcal{G}r$ -fibration over \mathbb{A}^1 :

$$\mathcal{BD} := \left\{ (x, \mathscr{E}, \beta) \middle| \begin{array}{c} \mathscr{E} \text{ rank 2 vector bundle} \\ \beta : \mathscr{E}|_{X - \{x\}} \xrightarrow{\sim} \mathscr{E}_{\text{triv}}|_{X - \{x\}} \end{array} \right\} \xrightarrow{\mathcal{G}r} \mathbb{A}^1.$$

She also introduced Gaitsgory's souped-up version of the the Beilinson-Drinfeld Grassmannian, which in the example of GL_2 is a $\mathcal{F}\ell$ -fibration over $\{0\}$ and a $\mathcal{G}r \times \mathbb{P}^1$ -fibration over $\mathbb{A}^1 \setminus \{0\}$:

Note that a flag in \mathscr{E}_0 is simply the choice of a line in the two-dimensional vector space \mathscr{E}_0 . Within \mathcal{BD} we have a finite-dimensional closed subvariety $\mathcal{G} \subset \mathcal{BD}$. Under the lattice description of $\mathcal{G}r_{\mathrm{GL}_2}$,

$$\mathcal{G} := \{ \mathcal{L} \subset \mathcal{L}_{\mathrm{triv}} \mid \dim \mathcal{L}_{\mathrm{triv}} / \mathcal{L} = 1 \}.$$

We can cover \mathcal{G} with two charts: if λ is the coordinate on \mathbb{A}^1 ,

$$U_0 = \left\langle \begin{pmatrix} 1 \\ a \end{pmatrix}, \begin{pmatrix} 0 \\ t - \lambda \end{pmatrix} \right\rangle, \ U_\infty = \left\langle \begin{pmatrix} b \\ 1 \end{pmatrix}, \begin{pmatrix} t - \lambda \\ 0 \end{pmatrix} \right\rangle.$$

Each chart is isomorphic to \mathbb{A}^2 , and we see that \mathcal{G} is a trivial \mathbb{P}^1 -bundle over \mathbb{A}^1 .

The analogous subvariety in Gaitsgory's version is

$$\mathcal{Y} = \left\{ (\mathcal{L}, \ell) \middle| \begin{array}{c} \mathcal{L} \subset \mathcal{L}_{\text{triv}} \text{ a lattice s.t.} \\ \dim \mathcal{L}_{\text{triv}} / \mathcal{L} = 1, \text{ and} \\ \ell \subset \mathcal{L} / t \mathcal{L} \end{array} \right\} \xrightarrow{\mathbb{P}^{1}\text{-bundle}} \mathcal{G}.$$

Consider the following closed subvariety of \mathcal{Y} :

$$\mathcal{Y}^{\mathrm{Sp}} = \{ (\mathcal{L}, \ell) \mid \ell \subset \ker(\mathcal{L} \hookrightarrow \mathcal{L}_{\mathrm{triv}} \twoheadrightarrow \mathcal{L}_{\mathrm{triv}}/t\mathcal{L}_{\mathrm{triv}} = \mathbb{C} \oplus \mathbb{C} \twoheadrightarrow \mathbb{C} \oplus \mathbb{C}/(\mathbb{C} \oplus 0)) \}.$$

What are the fibres of \mathcal{Y}^{Sp} over \mathbb{A}^1 ? Well, for any $(\mathcal{L}, \ell) \in \mathcal{Y}^{Sp}$, \mathcal{L} fits into an exact sequence

$$\mathcal{L} \hookrightarrow \mathcal{L}_{\mathrm{triv}} \twoheadrightarrow \mathcal{O}_x$$

where \mathcal{O}_x is a skyscraper. There are two cases:

1. If $x \neq 0$, then

$$\mathcal{L}_0 \simeq (\mathcal{L}_{\mathrm{triv}})_0.$$


Hence ℓ is uniquely determined, and the fibre is \mathbb{P}^1 .

2. If x = 0, then

$$\mathcal{L}_0 \xrightarrow{\phi} (\mathcal{L}_{\mathrm{triv}})_0$$

has rank 1. We have two possibilities: either (1) Im $\phi = \mathbb{C} \oplus 0$, in which case ℓ is free, but \mathcal{L} is fixed, so the fibre is \mathbb{P}^1 , or (2) $\ell = \ker \phi$, in which case \mathcal{L} is free and ℓ is fixed, so the fibre is again \mathbb{P}^1 . We conclude that the fibre over 0 is $\mathbb{P}^1 \bigcup_{\{0\}} \mathbb{P}^1$, as we had hoped.

In Geordie's hand-written notes, there are charts which show that locally, this degeneration is given by the following picture:



Furthermore, one can check that this degeneration is isomorphic to the degenerating quadric in \mathbb{P}^2 discussed earlier.

23.4 Exercises

These exercises will examine the structure of the affine Hecke algebra for GL_2 . We have


$$X = \mathbb{Z}e_1 \oplus \mathbb{Z}e_2, \ X^{\vee} = \mathbb{Z}e_1^* \oplus \mathbb{Z}e_2^*, \ \Phi = \{\pm(e_1 - e_2)\}, \ \Phi^{\vee} = \{\pm(e_1^* - e_2^*)\}.$$

Remark 23.8. The group GL_2 is Langlands self-dual, so we don't need to worry too much about which side of Langlands duality we are on.

The affine Weyl group is

$$W = W_f \ltimes (\mathbb{Z}e_2 \oplus \mathbb{Z}e_2).$$

A picture:



Given $\gamma \in \mathbb{Z}e_1 \oplus \mathbb{Z}e_2$, write t_{γ} for the corresponding translation.

1. Show that the set of *length zero elements*


$$\Omega = \{ x \in W \mid x \text{ preserves strip } (0,0) \}$$

is a free abelian group generated by $t_{e_1}s \in W$, where $s = s_{\alpha}$ is the finite simple reflection.

- 2. Set $\varpi = t_{e_1}s, s_0 = t_{\alpha}s$. Check that $\varpi s = s_0 \varpi, \, \varpi s_0 = a \varpi$, and ϖ^2 is central.
- 3. Show that the Bernstein generators are given by

$$\theta_{e_1} = \delta_{\varpi} \delta_s, \ \theta_{e_2}^{-1} = \delta_{\varpi^{-1}} \delta_s \theta_{e_2} = \delta_s^{-1} \delta_{\varpi} = \delta_{\varpi} \delta_{s_0}^{-1}.$$

Hint:



4. Verify that

$$z_{\text{nat}} = \theta_{e_1} + \theta_{e_2} = \delta_{\varpi}(\delta_s + \delta_{s_0}^{-1}) = \delta_{\varpi}(\delta_s^{-1} + \delta_{s_0})$$

is central.

- 5. Check that $z_{\text{nat}} \cdot b_s = \delta_{\varpi} b_{s_0 s}$.
- 6. (Challenge) Prove that Z(H) is generated by δ_{ϖ}^2 and z_{nat} and show that

$$Z(H) \xrightarrow{\sim} [\operatorname{Rep} \operatorname{GL}_2]$$
$$z_{\operatorname{nat}} \mapsto \mathbb{C}^2$$
$$\varpi^2 \mapsto \det$$

is an isomorphism of rings.

Lecture 24: A hitchhiker's guide to the Hecke category

In Lecture 16, we motivated the constructible side of Bezrukavnikov's equivalence via Grothendieck's function-sheaf dictionary. The material from that lecture is useful background for today's lecture, and the reader may benefit by reviewing Lecture 16 before continuing.

Let X be a quasi-projective variety over \mathbb{F}_q . There is a generating series built out of $\#X(\mathbb{F}_{q^m})$ for $m \geq 1$, the **zeta function**:

$$\zeta(X,s) = \exp\left(\sum_{m \ge 0} \frac{\#X(\mathbb{F}_{q^m})}{m} q^{-ms}\right).$$

Weil noticed that when X is a smooth projective *n*-dimensional variety, its ζ function has some remarkable properties, which he packaged into the **Weil conjectures**:

- 1. Rationality: $\zeta(X, s)$ can be written as a rational function in q^{-s} .
- 2. Functional equation: $\zeta(X, s)$ and $\zeta(X, 1-s)$ agree (up to a simple and explicit scalar).
- 3. Riemann hypothesis: roots of ζ have a specific form.

Example 24.1. If $X = \mathbb{P}^1_{\mathbb{F}_q}$, then

$$\begin{aligned} \zeta(X,s) &= \exp\left(\sum_{m\geq 0} \frac{1+q^m}{m} q^{-ms}\right) \\ &= \exp\left(\sum_{m\geq 0} \frac{(q^{-s})^m}{m} + \sum_{m\geq 0} \frac{(q^{1-s})^m}{m}\right) \\ &= \frac{1}{1-q^{-s}} \cdot \frac{1}{1-q^{1-s}}. \end{aligned}$$

We see from this that: (1) $\zeta(X,s)$ is rational, (2) $\zeta(X,s) = -q^{1-s}\zeta(X,1-s)$, and (3) we can write

$$\zeta(X,s) = \frac{1}{P_0(q^{-1})P_2(q^{-s})}$$

with $P_0 = (1 - T)$ and $P_2 = 1 - qT$. Here the Riemann hypothesis is the (elementary) statement that all roots of P_0 (resp. P_2) have roots of norm 1 (resp. q).

Example 24.2. If X is a smooth elliptic curve, then

$$\zeta(X,s) = \frac{P_1(q^{-s})}{P_0(q^{-s})P_2(q^{-s})} = \frac{P_1(q^{-s})}{(1-q^{-s})(1-q^{1-s})},$$

where P_1 has two roots, which are conjugate and have norm $q^{1/2}$ ("Weil numbers of weight 1").

Remark 24.3. The case of an elliptic curve (due to Hasse) predates the Weil conjectures and was an important ingredient in their formulation. Another important ingredient was Artin's computation of the zeta function of a hyperelliptic curve, discussed in Lecture 4. **Remark 24.4.** We saw the significance of the roots of P_1 (which determine how many points E has) in Lecture 4 on the Sato-Tate conjecture.

In the Weil conjectures, **rationality** follows from the Grothendieck-Lefschetz trace formula:

$$\#X(\mathbb{F}_q) = \text{supertrace}\left(\text{Frob} \circlearrowright H^*_{\text{\acute{e}t}}(X, \overline{\mathbb{Q}}_\ell)\right).$$

The **functional equation** follows from Poincaré duality in étale cohomology, and the Riemann hypothesis (the most difficult part) follows from purity. This is the statement that the eigenvalues of Frobenius in degree i are all "Weil numbers of weight i".

Remark 24.5. Notice the strange appearance of the auxiliary prime $\ell \neq p$ in the Grothendieck-Lefschetz trace formula. Why do we need to make this choice? Such questions are usually labelled questions of "independence of ℓ " in the literature. We will also see such questions arise in the Hecke category below.

The proof of the Weil conjectures needed the full artillery of $D_c^b(X, \overline{\mathbb{Q}}_{\ell})$ and its six functor formalism. It used a relative version of the Grothendieck-Lefschetz trace formula, which associated to a sheaf \mathcal{F} a collection of functions given by "trace of Frobenius":

$$\mathcal{F} \in D^b_c(X, \overline{\mathbb{Q}}_\ell) \rightsquigarrow \left\{ f^m_{\mathcal{F}} : X(\mathbb{F}_{q^m}) \to \overline{\mathbb{Q}}_\ell \right\}.$$

Then, as we saw in Lecture 16,

$$\left[D_c^b(X,\overline{\mathbb{Q}}_\ell)\right] \hookrightarrow \prod_{m\geq 1} \operatorname{Fun}(X(\mathbb{F}_{q^m}) \to \overline{\mathbb{Q}}_\ell),$$

so the collection of functions associated to \mathcal{F} completely determines its class in the Grothendieck group. This leads to Grothendieck's philosophy of "dictionaire functions faisceaux":


functions \leftrightarrow sheaves.

This philosophy can be summed up with the slogan

"interesting functions should arise from interesting sheaves".

24.1 The Hecke category: setting the scene

Let G be split reductive over \mathbb{F}_q , and $B \subset G$ a Borel subgroup. Let $\mathcal{K} = \mathbb{F}_q((t))$, and denote by $G_{\mathcal{K}}$ the loop group and $I \subset G_{\mathcal{K}}$ the corresponding Iwahori subgroup. Recall that the Hecke algebra (either finite or affine) has its origins as a convolution algebra of bi-invariant functions on a group (see Lecture 16 for more on this perspective). Using Grothendieck's function-sheaf dictionary, a natural categorification of these functions is the following:



Remark 24.6. The loop group $G_{\mathcal{K}}$ is very infinite-dimensional, so we usually work with

$$\left(D^b_I(\mathcal{F}\ell,\overline{\mathbb{Q}}_\ell),*\right)$$

instead. (Here $\mathcal{F}\ell = G_k/I$ is the affine flag variety.)

Remark 24.7. One can see from the above that H_f is "independent of q". We have one abstract algebra which specialises to all Hecke algebras at once. This is one desirable feature that we are hoping to categorify.

Our goal for this lecture is to find a good categorification of the Hecke algebra. It might appear that we've already accomplished this. However, the categories $(D^b_{B\times B}(G, \overline{\mathbb{Q}}_{\ell}), *)$ and $(D^b_{I\times I}(G_{\mathcal{K}}, \overline{\mathbb{Q}}_{\ell}), *)$ are not quite right for several reasons. In the remainder of the lecture, we will explain why these categories are wrong, then slowly fix them. We will concentrate on the finite case (i.e. the Hecke algebra for the finite Weyl group). The affine case is similar.

What we want: A triangulated monoidal category \mathcal{H} such that:

- 1. $([\mathcal{H}], *) \simeq H$ (i.e. \mathcal{H} categorifies the Hecke algebra),
- 2. \mathcal{H} is "independent of q and ℓ ," and
- 3. \mathcal{H} admits a triangulated monoidal functor to $D^b_{B\times B}(G, \overline{\mathbb{Q}}_\ell)$ such that the diagram

$$\begin{array}{ccc} \mathcal{H} & & \longrightarrow D^b_{B \times B}(G, \overline{\mathbb{Q}}_{\ell}) \\ \downarrow & & \downarrow \\ \mathcal{H} & \stackrel{v \mapsto \frac{1}{\sqrt{q}}}{\longrightarrow} \left(\operatorname{Fun}_{B(\mathbb{F}_q) \times B(\mathbb{F}_q)}(G(\mathbb{F}_q), \mathbb{C}), * \right) \end{array}$$

commutes for all ℓ and q.

Remark 24.8. The existence of such an object is tacitly implied by Bezrukavnikov's equivalence, as ℓ and q are nowhere to be seen on the coherent side.

Now we try and fail and try and fail and try and fail, and then, finally, succeed.

24.2 First try

We start with the most obvious choice:

$$\mathcal{H} = D^{\flat}_{B \times B}(G, \mathbb{Q}_{\ell}).$$

Objections:

- 1. Depends on q and ℓ .
- 2. If G is the trivial group, then

$$\mathcal{H} = D_c^b(\operatorname{Spec} \mathbb{F}_q, \overline{\mathbb{Q}}_\ell) \hookrightarrow D_c^b \left(\begin{array}{c} \operatorname{continuous representations} \\ \operatorname{of} \pi_1^{\operatorname{\acute{e}t}}(\operatorname{Spec} \mathbb{F}_q) = \widehat{\mathbb{Z}} \text{ on} \\ \overline{\mathbb{Q}}_\ell \operatorname{-vector spaces} \end{array} \right).$$

This is almost an equivalence. In fact, one has

$$[\mathcal{H}] = \mathbb{Z}[\overline{\mathbb{Z}}_{\ell}^{\times}], \qquad (24.1)$$

which is way too big. (See [DBB83, Proposition 5.1.2] and the remark following it.) Note that the fact that this Grothendieck group is way too big persists for any group: rather than being an algebra over $\mathbb{Z}[v^{\pm 1}]$, our putative definition is an algebra over (24.1).

24.3 Second try

In the first try, we failed to obtain requirement 1 of our desired categorification because our category had a Grothendieck group which was too big. We can attempt to fix this by passing to the algebraic closure. Try

$$\mathcal{H} = D^b_{B \times B}(G, \overline{\mathbb{Q}}_\ell) \text{ for } G/\overline{\mathbb{F}}_q, \text{ or}$$
$$\mathcal{H} = D^b_{B \times B}(G, \mathbb{Q}) \text{ for } G/\mathbb{C}.$$

Objections:

1. If G is the trivial group, then

$$D^b_{B \times B}(G, \mathbb{Q}) = D^b \left(\begin{array}{c} \text{finite dimensional} \\ \text{vector spaces} \end{array} \right).$$

Hence


$$[\mathcal{H}] = \mathbb{Z},$$

which is too small. (We expect $\mathbb{Z}[v^{\pm 1}]$ for the Grothendieck group in this case.) But perhaps this is just a trivial-case phenomenon, we can test with a slightly bigger example.

2. Recall (c.f. Lecture 16) that in SL_2 , the quadratic relation in $Fun_{B\times B}(SL_2(\mathbb{F}_q), \mathbb{C})$ came from the fact that

$$\mathbb{1}_{\mathrm{SL}_2(\mathbb{F}_q)} * \mathbb{1}_{\mathrm{SL}_2(\mathbb{F}_q)} = (1+q)\mathbb{1}_{\mathrm{SL}_2(\mathbb{F}_q)}.$$

Why was this again? Well, the multiplication map mult : $SL_2 \times_B SL_2 \to SL_2$ factors through $SL_2 / B \times SL_2 = \mathbb{P}^1 \times SL_2$:



Here the horizontal arrow is the map $(g,h) \mapsto (gB,gh)$. So we obtain the formula above by pushing forward the constant sheaf on either side of the isomorphism, and the (1+q) comes from $\#\mathbb{P}^1(\mathbb{F}_q) = 1+q$.

In $D^b_{B\times B}(\mathrm{SL}_2,\mathbb{Q})$, the same diagram shows

$$\mathbb{Q}_{\mathrm{SL}_2} * \mathbb{Q}_{\mathrm{SL}_2} = p_* \mathbb{Q}_{\mathbb{P}^1 \times \mathrm{SL}_2} = H^*(\mathbb{P}^1) \otimes \mathbb{Q}_{\mathrm{SL}_2} = \mathbb{Q}_{\mathrm{SL}_2} \oplus \mathbb{Q}_{\mathrm{SL}_2}[-2]$$

In the Grothendieck group, this gives

$$[\mathbb{Q}_{SL_2}]^2 = 2[\mathbb{Q}_{SL_2}]. \tag{24.2}$$

Exercise 24.9. The map

$$\mathbb{Z}[W] \to \left[D^b_{B \times B}(\mathrm{SL}_2, \mathbb{Q}) \right]$$
$$s \mapsto \left[\mathbb{Q}_{\mathrm{SL}_2} \right] - \left[\mathbb{Q}_{B/B} \right]$$

is an isomorphism of algebras.

We see from the exercise that the trivial group wasn't just an anomaly - with this definition of \mathcal{H} , the Grothendieck group really is too small. Also notice that we want to replace the 2 in (24.2) by (1 + q). With the Grothendieck-Lefschetz trace formula in mind, we wish to replace an Euler characteristic (2) by the trace of Frobenius (1 + q). Therefore we are led to the following conclusion:

The moral: Somehow we need to remember weights!

24.4 Third try

Again, let G/\mathbb{F}_q . This time, we set

$$\mathcal{H} = \langle IC_x \mid x \in W_f \rangle_{[\mathbb{Z}],(\mathbb{Z}),\Delta} \subset D^b_{B \times B}(G, \mathbb{Q}_\ell).$$

In this incarnation of \mathcal{H} , we are keeping track of weight by introducing "Tate twists," denoted above by (\mathbb{Z}) .

What is a Tate twist? Define

$$\overline{\mathbb{Q}}_{\ell}(-1) := H_c^2(\mathbb{A}^1) = \overline{\mathbb{Q}}_{\ell} \in \operatorname{Sh}_{\mathrm{\acute{e}t}}(\operatorname{Spec} \mathbb{F}_q).$$

On $\overline{\mathbb{Q}}_{\ell}(-1)$, Frob acts via multiplication by q. Let $p: X \to \operatorname{Spec} \mathbb{F}_q$. For $\mathcal{F} \in D^b_c(X, \overline{\mathbb{Q}}_{\ell})$, set

$$\mathcal{F}(m) := \mathcal{F} \otimes \left(p^* (\mathbb{Q}_{\ell}(-1)^{\otimes (-m)}) \right).$$

(Here we are using that $\mathbb{Q}_{\ell}(-1)$ is invertible, as an étale sheaf on a point. Thus it makes sense to take any integral tensor power.)

Very nontrivial fact: This definition of \mathcal{H} is closed under convolution. (For the experts: This is a consequence of the Decomposition Theorem and the fact that the objects are "already semi-simple over \mathbb{F}_q ".)

Remark 24.10. (*Technical point*) In the Hecke algebra, it is very useful to introduce a square root of q. To reflect this, we often choose a square root of q in $\overline{\mathbb{Q}}_{\ell}$ and use it to define $\overline{\mathbb{Q}}_{\ell}(-1/2)$, the "square root of Tate twist". This can also be done purely formally if one prefers.

Now we are in good shape: If G is the trivial group,


$$\mathcal{H} = \left\langle \left. \overline{\mathbb{Q}}_{\ell} \left(\frac{m}{2} \right) \right| m \in \mathbb{Z} \right\rangle_{\Delta} \subset D^{b}(\operatorname{Spec} \mathbb{F}_{q}).$$

Hence the Grothendieck group is

$$[\mathcal{H}] \xrightarrow{\sim} \mathbb{Z}[v^{\pm 1}], v \mapsto \overline{\mathbb{Q}}_{\ell}(-1/2).$$

So this \mathcal{H} passes our first test. Great!

Using Grothendieck's theory of weights, there is an alternative version of this third try. Recall (see "Scholze's motivic plane" in Lecture 10):



Under the red arrow above, we have a second incarnation of our third try:

$$\mathcal{H} = \langle \boldsymbol{I}\boldsymbol{C}_x \mid x \in W_f \rangle_{[\mathbb{Z}], (\frac{1}{2}\mathbb{Z}), \Delta} \subset D^b_{B \times B} \left(\begin{array}{c} \text{mixed Hodge} \\ \text{modules} \end{array} \right)$$

This version of \mathcal{H} also has the correct Grothendieck group. It is closed under convolution by the Decomposition Theorem.

A subtlety: These two incarnations $\mathcal{H}_{\text{étale}}$ and \mathcal{H}_{mHm} of our third attempt are really different. If G is the trivial group, then

• in $\mathcal{H}_{\text{étale}}$,

$$\operatorname{Ext}^{1}\left(\overline{\mathbb{Q}}_{\ell}, \overline{\mathbb{Q}}_{\ell}\left(\frac{m}{2}\right)\right) = \begin{cases} \overline{\mathbb{Q}}_{\ell} & \text{if } m = 0; \\ 0 & \text{otherwise.} \end{cases}$$

A non-trivial self-extension of $\overline{\mathbb{Q}}_{\ell}$ is given by $\overline{\mathbb{Q}}_{\ell}^2$, with Frobenius acting by Jordan block $\begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}$. Hence, there are extensions between objects of the same weights, but no extensions between objects of different weights.

• However, in \mathcal{H}_{mHm} , there are no extensions between objects of the same weight, but there are extensions between objects of different weights²⁸.

Remark 24.11. Under Bezrukavnikov's equivalence, we expect

weight \leftrightarrow weight of \mathbb{C}^{\times} -action.

So there should be no Exts at all for the trivial group!

A historical incarnation of this issue: Let \mathcal{O}_0 be the principal block of category \mathcal{O} . There is a finite-dimensional \mathbb{C} -algebra A such that $\mathcal{O}_0 \simeq A$ -mod_{f.g.}. In 1990, Soergel showed that A admits a \mathbb{Z} -grading \widetilde{A} defined over \mathbb{Q} , and \widetilde{A} -gmod_{f.g.} provides a grading on category \mathcal{O} . This grading explains the q in the Kazhdan–Lusztig polynomials. Hence we have the following diagram:

$$\begin{array}{cccc} A\operatorname{-mod}_{\mathrm{f.g.}} & \longleftarrow & A\operatorname{-gmod}_{\mathrm{f.g.}} \\ & & & \\ & & & \\ & \mathcal{O}_0 & \underbrace{\qquad \sim \qquad }_{\text{\mathcal{D}-modules}} & \underbrace{\qquad \sim \qquad }_{\text{\mathcal{P}-rv}_{(B)}(G/B)} \end{array}$$

What is the geometric meaning of the grading on \mathcal{O}_0 coming from \widetilde{A} ? This was explained in [BGS96]:

$$\widetilde{A}_{\overline{\mathbb{Q}}_{\ell}}\text{-gmod} \simeq \begin{array}{c} \text{some geometric} \\ \text{category} \end{array} \leftrightarrow D^b_B(G/B, \overline{\mathbb{Q}}_{\ell}), \\ \widetilde{A}_{\mathbb{C}}\text{-gmod} \simeq \begin{array}{c} \text{some geometric} \\ \text{category} \end{array} \leftrightarrow D^b_B(MHM_{G/B}). \end{array}$$

In each case, the graded version is explained by some category related, but not equal, to mixed sheaves. In both cases, some cooking is involved to remove the problematic Exts from earlier. The "cooking" is different in each case.

Remark 24.12. Recent motivic versions (Soergel-Wendt [SW18] and Soergel-Wendt-Virk, [SVW18]) explain how to remove the cooking.

²⁸Stating exactly what these extension groups are would require us to be more precise about what version of mixed Hodge structures we are using (integral, real, complex, etc.). However, the basic phenomenon that different weights extend persists in all of them.

24.5 Fourth (and final) try

We have finally converged on the correct formulation. Let

$$\mathcal{H}_{\mathrm{s.s}} = \langle IC_x \mid x \in W_f \rangle_{\oplus, [\mathbb{Z}]} \subset D^b_{B \times B}(G, \mathbb{Q})$$

This is an additive category, but it is *not* triangulated. Thus it consists of all semi-simple complexes in $D^b_{B\times B}(G,\mathbb{Q})$. It is closed under convolution by the Decomposition Theorem. We need to make make one necessary change of notation.

Redefine: (1) := [1] on $\mathcal{H}_{s.s.}$

Definition 24.13. The geometric Hecke category is

$$\mathcal{H} := K^b(\mathcal{H}_{\mathbf{s},\mathbf{s}}).$$

This category has two shift functors: (n) denotes the shift in $\mathcal{H}_{s.s.}$ and [n] denotes the shift in K^b .

Before declaring victory, we should verify that this satisfies all of our desired properties for the trivial group. If G is the trivial group, then

$$\mathcal{H}_{\text{s.s.}} = D_c^b(\text{pt}) = D^b \left(\begin{array}{c} \text{finite-dimensional} \\ \text{vector spaces} \end{array} \right) = \begin{array}{c} \text{finite-dimensional} \\ \text{graded vector spaces.} \end{array}$$

(Here the final equality is as *additive* categories.) Hence

$$\mathcal{H} = K^b \left(\begin{array}{c} \text{finite-dimensional} \\ \text{graded vector spaces} \end{array} \right).$$


Note that this category has no extensions, so our woes of our third attempt have disappeared. Moreover, we have

$$[\mathcal{H}] = \mathbb{Z}[v^{\pm 1}],$$

with v corresponding to the shift of grading. Moreover, there are no q's or ℓ 's in sight. Hurrah!

Remark 24.14. 1. We have $[\mathcal{H}] = [\mathcal{H}_{s.s.}] = H_f$.

- 2. A similar definition over $\overline{\mathbb{F}}_q$ with $\overline{\mathbb{Q}}_\ell$ -sheaves leads to an equivalent category (after extensions of scalars to $\overline{\mathbb{Q}}_\ell$). This proves independence of q and ℓ . The proof is by showing that $\mathcal{H}_{\text{s.s.}} \simeq \mathbb{S}\text{Bim}$, the category of Soergel bimodules²⁹.
- 3. With difficulty, one can construct monoidal realisation functors:



 $^{^{29}}$ We hope to explain this sometime in the future!

This provides considerable evidence that ${\mathcal H}$ is the right object.

Remark 24.15. It is possible that although \mathcal{H} is the right object, it may not yet have the right definition. Recent motivic versions [SVW18] probably provide the "correct" definition.

Lecture 25: The categorical anti-spherical module and its symmetries

Recall that we are working towards proving the following theorem of Arkhipov–Bezrukavnikov:

Theorem 25.1. [AB09] There exists an equivalence of triangulated categories

$$D^{G^{\vee} \times \mathbb{G}_m}(\widetilde{\mathcal{N}}) \stackrel{\sim}{\longrightarrow} \mathcal{M}^{\mathrm{asph}}.$$

This equivalence categorifies the isomorphism

$$K^{G^{\vee} \times \mathbb{G}_m}(\widetilde{\mathcal{N}}) \xrightarrow{\sim} H \otimes_{H_f} \operatorname{sgn.}$$

Today we will define the category \mathcal{M}^{asph} and discuss the philosophy of higher representation theory.

25.1 The affine Hecke category

For the remainder of this course, $k \in \{\mathbb{Q}, \mathbb{R}, \mathbb{C}\}$. Last lecture in our quest to find the correct definition of the geometric Hecke category, we defined

$$\mathcal{H}_f^{\rm ss} := \langle \boldsymbol{I}\boldsymbol{C}_x \mid x \in W_f \rangle_{\oplus, [\mathbb{Z}]} \subset D_{B \times B}(G, k).$$

Here "ss" stands for semi-simple. This category is additive, but not triangulated.

Remark 25.2. (Technical point) It is convenient to normalize IC_x so that it corresponds to the usual IC sheaf on G/B under the equivalence

$$D_{B \times B}(G, k) \simeq D_B(G/B, k);$$

e.g.

$$IC_{id} = k_B, IC_s = k_{P_s}[1], \ldots, IC_x|_{BxB} = k_{BxB}[\ell(x)]$$

We used \mathcal{H}_{f}^{ss} to define the geometric Hecke category:

$$\mathcal{H}_f := K^b(\mathcal{H}_f^{\mathrm{ss}}).$$

This is a triangulated category (in contrast to \mathcal{H}_f^{ss}). It has two natural shift functors:

[1] = triangulated shift functor, and $(1) = \text{ functor induced by the shift functor on } \mathcal{H}_{f}^{ss}.$

A picture of these shifts on an object $\mathcal{F} \in \mathcal{H}_f$ is

Here each $\mathcal{F}^i \in \mathcal{H}_f^{\mathrm{ss}}$.

Theorem 25.3. The map $b_s \mapsto [IC_s]$ induces an isomorphism of $\mathbb{Z}[v^{\pm 1}]$ -algebras

$$H_f \xrightarrow{\sim} [\mathcal{H}_f^{\mathrm{ss}}]_{\oplus} \simeq [\mathcal{H}_f]_{\Delta}.$$

Here the subscript \oplus denotes the split Grothendieck group of an additive category, and the subscript Δ denotes the triangulated Grothendieck group of a triangulated category. Under this isomorphism, the Kazhdan–Lusztig basis element b_x corresponds to the class of an intersection cohomology complex, in formulas

$$b_x \mapsto [IC_x].$$

The same construction works in the affine case. Define an additive category

$$\mathcal{H}^{\rm ss} := \langle IC_x \mid x \in W \rangle_{\oplus, [\mathbb{Z}]} \subset D^b_I(\mathcal{F}\ell, k)$$

for $\mathcal{F}\ell := G((t))/I$, and a triangulated category

$$\mathcal{H} := K^b(\mathcal{H}^{\mathrm{ss}}).$$

The analogue of Theorem 25.3 holds:

Theorem 25.4.

$$H \simeq [\mathcal{H}^{ss}]_{\oplus} \simeq [\mathcal{H}]_{\Delta}, b_x \mapsto [IC_x].$$

25.2 The categorical (anti-)spherical module

In the Hecke algebra, we have the quadratic relation for each $s \in S$:

$$\delta_s^2 = (v^{-1} - v)\delta_s + 1$$
, or reformulated, $(\delta_s + v)(\delta_x - v^{-1}) = 0$.

This leads to the existence of two H_f -modules, sgn and triv, where δ_s acts by -v and v^{-1} , respectively, and two corresponding induced modules for H:

 $M^{\mathrm{sph}} := \mathrm{triv} \otimes_{H_f} H$ the spherical module, and $M^{\mathrm{asph}} := \mathrm{sgn} \otimes_{H_f} H$ the antispherical module.

Remark 25.5. We have made a notational switch from denoting the anti-spherical module by N (c.f. Lecture 21) to M^{asph} . This is because we wish to denote categorifications by the corresponding script letters and the symbol \mathcal{N} has already been assigned to the nilpotent cone.

Remark 25.6. We have also made a switch from left to right modules. From now on, the spherical and anti-spherical module will be considered as right *H*-modules to align with [AB09].

In Lecture 20, we discussed how as right modules over the lattice part $\mathcal{L} = \bigoplus_{\lambda \in X^{\vee}} \mathbb{Z}[v^{\pm 1}] \theta_{\lambda}$ of H, M^{sph} is a free module of rank 1:

$$M^{\mathrm{sph}} = \mathrm{triv} \otimes_{H_f} H = \mathrm{triv} \otimes_{H_f} H_f \otimes \mathcal{L} \simeq \mathcal{L}.$$

A similar statement holds for M^{asph} . This description of the (anti-)spherical module, which follows from the Bernstein presentation of the affine Hecke algebra, is the "coherent perspective" of $M^{(a)\text{sph}}$. On the other hand, using the Coxeter structure of W, we can view $M^{(a)\text{sph}}$ from a "constructible perspective" as follows. Any $w \in W$ can be written as $w_{\text{fn}}{}^{f}w$ for $w_{\text{fn}} \in W_{f}$ and ${}^{f}w \in {}^{f}W$, where ${}^{f}W$ is the set of minimal coset representatives in $W_{f} \setminus W$. Hence we have an isomorphism

$$H_f \otimes \left(\bigoplus_{x \in {}^{f}W} \mathbb{Z}[v^{\pm 1}] \delta_x \right) \xrightarrow{\sim} H,$$

and $M^{(a)sph}$ has a "standard basis"

$$\{\delta_x^{(a)sph} := 1 \otimes \delta_x \mid x \in {}^fW\}.$$

Exercise 25.7. (Fun!) Compute the bijection

$$\begin{array}{l} \text{dominant} \\ \text{alcoves} \end{array} \stackrel{\sim}{\leftrightarrow} {}^{f}W \stackrel{\sim}{\leftrightarrow} W_{f} \backslash W = W_{f} \backslash (W_{f} \ltimes \mathbb{Z}\Phi^{\vee}) = \mathbb{Z}\Phi^{\vee}$$

in a few examples (e.g. for SL_2, SL_3, \ldots).

Using Kazhdan–Lusztig combinatorics, we can give descriptions of the spherical and antispherical module in terms of the Kazhdan–Lusztig basis of H. We dedicate the next part of the lecture to doing this.

Exercise 25.8. Prove the following:

- 1. If $x \in W$ and $s \in S$ are such that xs < x, then $b_x b_s = (v + v^{-1})b_x$.
- 2. If $t \in S$ and $x \in W$ are such that tx < x and $s \in S$ is arbitrary, then in the decomposition

$$b_x b_s = \sum n_y b_y,$$

 $n_y \neq 0$ implies ty < y. Conclude from this that $\{b_x \mid tx < x\}$ span a right ideal in H.

Lemma 25.9. (realisations of M^{sph} and M^{asph} via Kazhdan–Lusztig theory) Let w_f be the longest element in W_f .

1. The map $1 \otimes 1 \mapsto b_{w_f}$ induces an isomorphism of right *H*-modules

$$M^{\mathrm{sph}} \xrightarrow{\sim} b_{w_f} H.$$

2. The $\mathbb{Z}[v^{\pm 1}]$ -span of $\{b_s \mid s \notin {}^{f}W\}$ is a right ideal. Moreover,

$$M^{\operatorname{asph}} \simeq H/\langle b_x \mid x \notin {}^fW \rangle.$$

Proof. By Exercise 25.8.1, $b_{w_f}b_s = (v + v^{-1})b_{w_f}$ for all $s \in S_f$, so

$$b_{w_f}\delta_s = b_{w_f}(b_s - v\delta_{id}) = v^{-1}b_{w_f}$$

for all $s \in S_f$. By Frobenius reciprocity, we have a map

$$M^{\mathrm{sph}} \xrightarrow{\phi} b_{w_f} H, \ \delta^{\mathrm{sph}}_{id} \mapsto b_{w_f}.$$

For $x \in {}^{f}W$

$$\delta_x^{\mathrm{sph}} \stackrel{\phi}{\mapsto} b_{w_f} \delta_x = \delta_{w_f x} + \text{ lower terms.}$$

By upper triangularity, we conclude that ϕ is an isomorphism. This proves 1.

By Exercise 25.8.2, $\langle b_x \mid x \notin {}^f W \rangle$ is a right ideal in H. Hence $H/\langle b_x \mid x \notin {}^f W \rangle$ is a right H-module. In $H/\langle b_x \mid x \notin {}^f W \rangle$,

$$0 = 1 \cdot b_s = \delta_s + v$$

for $s \in S_f$, so

 $1 \cdot \delta_s = -v \cdot 1.$

By Frobenius reciprocity, we have a map

$$M^{\mathrm{asph}} \to H/\langle B_x \mid x \notin {}^fW \rangle,$$

and an analogous argument to the one above shows that this map is an isomorphism. This proves 2. $\hfill \Box$

Remark 25.10. One can use the other Kazhdan–Lusztig basis $\{b'_x\}$ to realise M^{sph} as a quotient and M^{asph} as a submodule. This is categorified by Koszul duality.

Now it makes sense to define the categorical versions of M^{sph} and M^{asph} :

$$\mathcal{M}^{\mathrm{sph,ss}} := \langle IC_{w_f} * \mathcal{H}^{\mathrm{ss}} \rangle_{\ominus} \subset \mathcal{H}^{\mathrm{ss}},$$

 $\mathcal{M}^{\mathrm{sph}} := \langle IC_{w_f} * \mathcal{H} \rangle_{\Delta} \subset \mathcal{H},$
 $\mathcal{M}^{\mathrm{asph,ss}} := \mathcal{H}^{\mathrm{ss}} / \langle IC_x \mid x \notin {}^{f}W \rangle_{\oplus,(\mathbb{Z})},$
 $\mathcal{M}^{\mathrm{asph}} := \mathcal{H} / \langle IC_x \mid x \notin {}^{f}W \rangle_{\Delta,(\mathbb{Z})}.$

Here the quotients are quotients of additive categories, and \ominus denotes the closure under direct summands.

Lemma 25.11.

$$\mathcal{M}^{\mathrm{asph}} = K^b(\mathcal{M}^{\mathrm{asph,ss}}).$$

Theorem 25.12. The category $\mathcal{M}^{(a)sph}$ is a right \mathcal{H} -module. The map $\delta_{id}^{asph} \mapsto IC_{id}$ induces an isomorphism of right H-modules

$$M^{(\mathrm{a})\mathrm{sph}} \simeq [\mathcal{M}^{(a)\mathrm{sph}}]_{\oplus} = [\mathcal{M}^{(\mathrm{a})\mathrm{sph}}]_{\Delta}.$$

Remark 25.13. In future lectures, we will discuss in more detail what it means for a category to be a module over a monoidal category like \mathcal{H} .

25.3 Symmetries of categories

The remainder of this lecture will be a discussion of higher representation theory. We start at the beginning: **Representation theory** is the study of linear symmetry (of groups, algebras, etc.). This field stems from the underlying observation:

Group actions are difficult, but they become easier once we linearize.

- **Example 25.14.** 1. Linearizing the action of $S^1 \circlearrowright S^1$ (resp. $SO(3) \circlearrowright S^2$) leads to the theory of Fourier series (resp. spherical harmonics).
 - 2. If X is a variety over \mathbb{Q} , studying $X(\mathbb{Q})$ is hard. This can be expressed as a question about the action of the absolute Galois group of \mathbb{Q} on all points of $X(\overline{\mathbb{Q}})$. A central technique in modern number theory is instead to study the linear problem $\operatorname{Gal}(\overline{\mathbb{Q}}/\mathbb{Q}) \circlearrowright$ $H^i(X, \overline{\mathbb{Q}}_{\ell})$.

2-representation theory is the study of symmetries of linear categories (additive, abelian, triangulated, etc.). These symmetries take the form of monoidal categories.

Philosophy: (learned from Manin) In differential geometry, if we have a group acting on a manifold $G \circlearrowright M$, we get a lot of leverage by linearizing and studying $G \circlearrowright L^2(M, \mathbb{C})$. In algebraic geometry, if we want to do the same thing for a group G acting on an algebraic variety X, a useful technique is to study is $G \circlearrowright \operatorname{Coh}(X), \operatorname{QCoh}(X), D^b(\operatorname{Coh}(X))$. The categories $\operatorname{QCoh}(X), \operatorname{Coh}(X), D^b(\operatorname{Coh}(X))$ can be thought of as 2-linearizations of X.

Remark 25.15. (Side remark, c.f. [Man14]) Let \mathcal{F} be a sheaf on X. Early approaches to studying such objects (~ 1950s) emphasized Čech coverings. Later, Grothendieck shifted the emphasis to injective resolutions. In sheaf cohomology

 $H^i(X,\mathcal{F})$

there is a non-linear variable (X), and a linear variable (\mathcal{F}) . We can interpret Grothendieck's shift as a movement from the non-linear variable to the linear variable.

Long range hope: From the perspective of 2-representation theory, our current goal is to show that the natural symmetries of

$$D^{G^{\vee} \times \mathbb{G}_m}(\widetilde{\mathcal{N}})$$
 and $\mathcal{M}^{\operatorname{asph}}$

agree, and yield the Hecke category.

A priori the symmetries on each side look rather different:

RHS
$$\circlearrowleft \mathcal{H}$$

LHS $\circlearrowright \operatorname{Rep} G^{\lor}, \operatorname{Rep} \mathbb{G}_m, \operatorname{Pic}^{G^{\lor} \times \mathbb{G}_m}(\widetilde{\mathcal{N}}), \ldots$

25.4 A notational interlude:

We will use the language of stacks.

- All stacks we study will be of the form Y/H, where Y is a quasi-projective variety and H is an affine algebraic group.
- See Emily's IFS talk "Sheaves on Stacks" for an excellent introduction to coherent sheaves on stacks.
- The most important fact (which will be used repeatedly) is

$$\operatorname{QCoh}(Y/H) \simeq \operatorname{QCoh}^H(Y),$$

 $\operatorname{Coh}(Y/H) \simeq \operatorname{Coh}^H(Y).$

- **Basic observation:** If Y is a stack as above, then
 - 1. $\operatorname{QCoh}(Y)$ and the category of vector bundles on Y are symmetric monoidal categories.
 - 2. If $X \xrightarrow{f} Y$ is a map of stacks, then QCoh is a module over QCoh(Y) via

$$\mathcal{F} * \mathcal{G} := f^*(\mathcal{F}) \otimes_{\mathcal{O}_X} \mathcal{G}$$

for $\mathcal{F} \in \operatorname{QCoh}(Y)$, $\mathcal{G} \in \operatorname{QCoh}(X)$. (Note that here * denotes the action map of $\operatorname{QCoh}(X)$ and not convolution.)

25.5 The basic approach of Arkhipov-Bezrukavnikov

The basic approach of [AB09] is the following. We wish to show that

$$D^{G^{\vee} \times \mathbb{G}_m}(\widetilde{\mathcal{N}}) \simeq \mathcal{M}^{\operatorname{asph}}$$

(For the moment, we ignore \mathbb{G}_m .) We have the following maps of stacks:

$$\begin{array}{ccc} \widetilde{\mathcal{N}}/G^{\vee} & \operatorname{QCoh}(\widetilde{\mathcal{N}}/G^{\vee}) \\ \downarrow^{q} & q^{*} & \uparrow \\ \mathfrak{g}^{\vee}/G^{\vee} & \rightsquigarrow & \operatorname{QCoh}(\mathfrak{g}^{\vee}/G^{\vee}) \\ \downarrow^{p} & p^{*} & \uparrow \\ \operatorname{pt}/G^{\vee} & \operatorname{QCoh}(\operatorname{pt}/G^{\vee}) \end{array}$$

The maps of stacks in the column on the left induce the increasing chain of symmetric monoidal categories in the column on the right. Step by step, we will construct an action of the symmetric monoidal categories in this chain on \mathcal{M}^{asph} . The action of $\operatorname{QCoh}(\operatorname{pt}/G^{\vee})$ will come from Gaitsgory's central functors, the action of $\operatorname{QCoh}(\mathfrak{g}^{\vee}/G^{\vee})$ from the monodromy endomorphism of nearby cycles, and the action of $\operatorname{QCoh}(\widetilde{\mathcal{N}}/G^{\vee})$ from "Wakimoto arrows" (which are yet to be defined). This procedure yields a functor

$$\operatorname{QCoh}(\widetilde{\mathcal{N}}/G^{\vee}) \to \mathcal{M}^{\operatorname{asph}},$$

which we will argue is an equivalence.

Lecture 26: Monoidal categories and their actions

In representation theory, we study

$$A$$
 \circlearrowright M

algebra vector space

This process lets us draw conclusions about M (composition series, semi-simplicity, etc.) In 2-representation theory, we study

 \mathcal{A} \circlearrowright \mathcal{M}

monoidal category category

What does this process let us conclude about \mathcal{M} ? Today we'll examine this question.

A theme of this lecture: There are two approaches to higher algebra:

- (a) Generators and relations: carry "just enough" coherence data, or
- (b) *Holistic*: carry "all" coherence data.

Approach (a) is rigid, but (sometimes) computable; whereas approach (b) is flexible but (often) incomputable. Historically, (a) is usually developed first, but (b) proves itself to be more powerful in the long run. The following familiar example from algebraic topology illustrates this.

Example 26.1. Given a reasonable topological space X, there are two approaches to computing homology $H_*(X)$:

- (a) triangulate your space as efficiently as possible and compute homology using a "small" complex (simplicial homology), or
- (b) consider all possible singular *n*-simplices and build an enormous chain complex from the spaces $\text{Hom}(\Delta_n, X)$ (singular homology).

The first approach is by "generators and relations," whereas the second is "holistic".

Remark 26.2. (For those who know about simplicial sets) The two approaches in Example 26.1 underly two approaches to studying the homotopy theory of X.

26.1 What is a monoidal category?

There are two approaches.

Generators and relations: A monoidal category is a category \mathcal{A} equipped with a bifunctor

$$\otimes:\mathcal{A}\times\mathcal{A}\to\mathcal{A},$$


a unit

 $\mathbb{1}\in\mathcal{A},$

and natural transformations

$$\alpha_{X,Y,Z} : (X \otimes Y) \otimes Z \xrightarrow{\sim} X \otimes (Y \otimes Z)$$
$$\lambda_X : \mathbb{1} \otimes X \xrightarrow{\sim} X, \text{ and}$$
$$\rho_X : X \otimes \mathbb{1} \xrightarrow{\sim} X$$

such that the diagrams



commute for all objects X, Y, Z, W. We refer to the first diagram as the *pentagon* and the second as the *unit*. With this set-up, we can also formulate the notion of a *monoidal functor* between monoidal categories, though we won't state this definition precisely today.

Basic claim: ("MacLane's coherence theorem") Any two maps consisting of associators and units agree.

Holistic approach: It's possible to give an alternative (and perhaps better?) definition of a monoidal category using the holistic approach, it can be found in [Lur07, §1.1]. This definition is complicated so we won't state it precisely here, but the rough idea is the following.

Let $\mathcal{A}^{\otimes n}$ be the category of sequences of *n*-elements in \mathcal{A} . A monoidal category is equivalence to a whole host of functors of the flavor

$$\mathcal{A}^{\otimes 5} \to \mathcal{A}^{\otimes 3}$$
$$(X_1, X_2, X_3, X_4, X_5) \mapsto (X_1 \otimes X_2, \mathbb{1}, X_3 \otimes X_4 \otimes X_5),$$

together with many many compatibilities.

26.2 Modules for monoidal categories

Similarly, we can formulate the notion of a module for a monoidal category either in terms of generators and relations, or holistically.

Generators and relations: A left module M for a monoidal category \mathcal{A} consists of a functor

$$\otimes: \mathcal{A} \times \mathcal{M} \to \mathcal{M},$$

together with natural transformations

$$\beta_{X,Y,M} : (X \otimes Y) \otimes M \xrightarrow{\sim} X \otimes (Y \otimes M),$$
$$\lambda_M : \mathbb{1} \otimes M \xrightarrow{\sim} M, \text{ and}$$
$$\rho_M : M \otimes \mathbb{1} \xrightarrow{\sim} M$$

such that the analogous pentagon and unit diagrams commute for all objects $X, Y \in \mathcal{A}$ and $M \in \mathcal{M}$.

Holistic: A similar approach is possible, see [Lur07, §2].

In representation theory, we are familiar with the fact that representations of an algebra are equivalent to modules over that algebra. An analogous statement holds in 2representation theory. A representation of \mathcal{A} is a pair (\mathcal{M}, ϕ) of a category \mathcal{M} and a \otimes functor $\phi : \mathcal{A} \to \operatorname{End}(\mathcal{M})$.

Proposition 26.3.

 $\{\mathcal{A}\text{-modules}\} \simeq \{\mathcal{A}\text{-representations}\}.$

Exercise 26.4. (Which Geordie assumes is true but hasn't done.) Check that this is an equivalence of 2-categories.

26.3 Examples of modules over monoidal categories

In representation theory, we started by studying groups, then realized that many of our questions about linear group symmetry could be formulated in terms of representations of algebras. In a similar vein, we can start 2-representation theory by building categories from groups.

Given a discrete group G, construct a category \mathcal{A}_G as follows:

- **Objects:** $\{r_g \mid g \in G\}$ satisfying $r_g \otimes r_h = r_{gh}$, $\mathbb{1} = r_{id}$, and α, λ, ρ are all the identity.
- Morphisms: $\operatorname{Hom}(r_g, r_h) = \emptyset$ for $g \neq h$, and $\operatorname{End}(r_g) = \{id_{r_g}\}$.

What is a module over \mathcal{A}_G ? Given a category \mathcal{M} and a \otimes -functor $F : \mathcal{A}_G \to \text{End}(\mathcal{M})$, define $F_g := F(r_g)$. Then the \otimes -functor F gives us natural isomorphisms

$$\mu_{gh}: F_g F_h \to F_{gh} \text{ and } \epsilon: F_{id} \to id_{\mathcal{M}}$$

such that

commute for all $g, h, k \in G$. This is called a *strict action* of G on \mathcal{M} .

Exercise 26.5. Show that giving an action of \mathbb{Z} on \mathcal{M} is the same as giving an autoequivalence E of \mathcal{M} .

Example 26.6. (An example to urge caution) What does it mean to give an action of $\mathbb{Z}/2\mathbb{Z}$ on \mathcal{M} ?


• First guess: A category \mathcal{M} with an autoequivalence $E : \mathcal{M} \to \mathcal{M}$ together with a natural isomorphism $m : E^2 \xrightarrow{\sim} id_{\mathcal{M}}$.

This is wrong! Here's why: It follows from the definition of a categorical action that the diagram

commutes. (This is a consequence of the previously displayed commutative square.) But we can cook up an example of data as above where it doesn't. Let \mathcal{M} be the category with two objects X and Y and morphisms

$$\operatorname{Hom}(X, Y) = \operatorname{Hom}(Y, X) = 0, \ \operatorname{End}(X) = \operatorname{End}(Y) = k,$$

where k is a field. Define $E : \mathcal{M} \to \mathcal{M}$ by E(X) = Y, E(Y) = X, and $id_X \xrightarrow{E} id_Y \xrightarrow{E} id_X$. A picture of this category:



Note that $E^2 = id_{\mathcal{M}}$, so any $m : E^2 \xrightarrow{\sim} id_{\mathcal{M}}$ is a natural transformation $m : id_{\mathcal{M}} \to id_{\mathcal{M}}$; i.e. $m \in Z(\mathcal{M})$. Hence

$$m = \begin{cases} a & \text{on End}(X) \\ b & \text{on End}(Y) \end{cases}$$

for some $a, b \in k$. Now in this example, the diagram (26.1) on X becomes

$$\begin{array}{ccc} Y & \stackrel{a}{\longrightarrow} Y \\ \downarrow & \swarrow \\ Y & & & \\ Y & & & \\ \end{array}$$

This diagram commutes if and only if a = b. However, any choice of a and b defines a natural transformation $m : E^2 \to id_{\mathcal{M}}$, so our first guess must be wrong!

• Revised guess (Exercise): Show that the data of our first guess along with the requirement that (26.1) commutes does determine an action of $\mathbb{Z}/2\mathbb{Z}$ on \mathcal{M} .

26.4 What is going on here?


We will briefly take a small diversion and discuss a beautiful dictionary related to these concepts. As we discussed in the beginning of this lecture, in the generators and relations approach to categorical actions, we want to include "just enough" coherence data. Example 26.6 illustrates that sometimes it can be subtle to determine how much coherence data is enough. The reader might be wondering:

How do we determine the necessary coherences?

It turns out that they are determined by the "cells of BG". We will illustrate what we mean by this through a series of examples.

Example 26.7. Associated to G is its classifying space BG^{30} .


1. For $G = \mathbb{Z}$, $BG = S^1$. This is a CW complex consisting of one 0-cell and one 1-cell. Recall that in Exercise 26.5 we showed that an action of \mathbb{Z} on a category \mathcal{M} is given by an autoequivalence $E : \mathcal{M} \to \mathcal{M}$:



Hence we have one piece of "coherence data" for each cell of BG.

2. For $G = \mathbb{Z}/2\mathbb{Z}$, $BG = \mathbb{RP}^{\infty} = S^{\infty}/\{\pm 1\}$. The sphere S^{∞} has a CW complex structure with two *n*-cells for each $n \in \mathbb{Z}_{>0}$:


Moreover, the action of $\mathbb{Z}/2\mathbb{Z}$ on this cell complex is cellular. Hence \mathbb{RP}^{∞} is a CW complex with one *n*-cell for each $n \in \mathbb{Z}_{\geq 0}$. We saw in Example 26.6 that an action of $\mathbb{Z}/2\mathbb{Z}$ on \mathcal{M} is determined by $(\mathcal{M}, E, m : E^2 \simeq id_{\mathcal{M}}, (26.1))$:



 $^{^{30}}$ Recall that BG is only defined up to homotopy. We will ignore this point below.

Again, for each *n*-cell with $0 \le n \le 3$, we have a corresponding piece of "coherence data".

3. For any group G, the Milnor construction of BG yields a CW complex strucure with one 0-cell, G 1-cells, $G \times G$ 2-cells, $G \times G \times G$ 3-cells, etc. Comparing this to the definition of a G-action on \mathcal{M} in Section 26.3, we see:



So each of piece of coherence data in the definition of a G-action corresponds to an n-cell of BG.

In each example above, the CW complex structure gives us the necessary coherence data to determine the action of G. In parts 1 and 2, we used a CW complex with a small number of cells to obtain "just enough" coherence data, illustrating the "generators and relations" approach. In part 3, we used a CW complex with many cells to capture "all" coherence data of the G-action, illustrating the "holistic" approach.

This pattern is very pretty, but why do we stop at 3-cells? This is because we are acting on 1-categories. In general,

26.5 Example: Representations of the Verlinde category

For some reason, an action of the category \mathcal{A}_G is not as powerful as a *G*-action on a vector space.

Reasons: (Speculative)

1. We need more structure to categorify linear algebra. For example, what is the eigenvalue of a functor? If $\mathbb{Z} \oslash \mathcal{M}$, how do we take its logarithm? In examples of categorical actions (e.g. of braid groups in highest weight representation theory), there are often interesting morphisms $F_g \to F_h$ and these need to be studied.

2. It is possible that this isn't the right generalization. In this course, a much more important action will be

$$\operatorname{Rep} G \circlearrowright \mathcal{M},$$

where G is a linear algebraic group. In other words, we will study "representations of representations of G".

Next lecture we will discuss the general case; today we will give a simple and beautiful example. To do so we need to introduce the Verlinde category.

Recall that for each $m \geq 0$, $\mathfrak{sl}_2(\mathbb{C})$ has a unique simple module V_m of highest weight mand dimension m + 1. In the category Rep $\mathfrak{sl}_2(\mathbb{C})$, the module V_0 is the monoidal unit, and other representations multiply according to the rule

$$V_1 \otimes V_m \simeq V_m \otimes V_1 \simeq V_{m+1} \oplus V_{m-1}.$$
(26.2)

For any positive integer $\ell \geq 1$, there exists a monoidal category C_{ℓ} which can be seen as a "finite" version of Rep $\mathfrak{sl}_2(\mathbb{C})$. It has $\ell + 1$ simple objects, $V_0, \ldots, V_{\ell}, V_0$ is the unit, and multiplication³¹ is given by

$$V_1 \otimes V_m = V_m \otimes V_1 = V_{m+1} \oplus V_{m-1}$$
 for $1 \le m < \ell$, and $V_\ell \otimes V_1 = V_1 \otimes V_\ell \simeq V_{\ell-1}$. (26.3)

Remark 26.8. There are two realizations of C_{ℓ} (see [Ost03], [Kac90]). The first is as the category of level ℓ representations of the affine Lie algebra $\widehat{\mathfrak{sl}}_2(\mathbb{C})$ with its fusion product. The second is as the semi-simplification of representations of the Lusztig form of the quantum group of \mathfrak{sl}_2 at a root of unity. In both settings, C_{ℓ} is braided but (in contrast to Rep $\mathfrak{sl}(2, \mathbb{C})$) not symmetric. The braiding will not play a role below.

Theorem 26.9. Let $\mathcal{A} = C_{\ell}$. Then

semi-simple abelian
$$\mathcal{A}$$
-modules \mathcal{M} such that
1. \mathcal{M} has finitely many simple objects, and
2. \mathcal{M} is generated by a single simple object $\Big|_{\simeq} \stackrel{\sim}{\longrightarrow} \left\{ \begin{array}{c} \text{simply laced} \\ \text{Dynkin diagrams} \\ \text{with Coxeter number } \ell + 2 \end{array} \right\}.$

Example 26.10. The category C_{10} admits 3 module categories on the right hand side, corresponding to the A_{11}, D_7 , and E_6 Dynkin diagrams.

Proof. (Sketch) Let us explain how to go from the LHS to RHS. Starting with \mathcal{M} on the LHS, we can build a graph as follows (cf. McKay correspondence):

- Vertices: One vertex for every simple module in \mathcal{M} .
- Edges: An edge between M and M' if M is a summand of $V_1 \otimes M'$.

Just as in $\operatorname{Rep} \mathfrak{sl}_2(\mathbb{C})$, V_1 is self-biadjoint in \mathcal{A} , so

M is a summand of $V_1 \otimes M' \iff M'$ is a summand of $V_1 \otimes M$.

³¹One can remember (26.3) by noticing that it is the same as (26.2), except that all classes V_m for $m > \ell$ are zero.

Hence the graph associated to any \mathcal{A} -module \mathcal{M} in this way is undirected.

From this graph, one can use (26.3) to deduce how V_m acts on the Grothendieck group $[\mathcal{M}]$ for all m. For example, if D denotes the adjacency matrix of the graph associated to \mathcal{M} , then the action of V_2 on \mathcal{M} is described by $D^2 - id$ because $[V_2] = [V_1^{\otimes 2}] - [V_0]$ in C_{ℓ} . Now some linear algebra allows one to deduce that the graph we construct from \mathcal{M} must either

be a simply laced affine Dynkin diagram or a "tadpole" $\begin{pmatrix} \\ \\ \\ \\ \end{pmatrix}$. Then we can work harder to

rule out the tadpole and show that the graph determines \mathcal{M} up to equivalence.

Remark 26.11. In lectures, it was incorrectly stated that one can classify module categories over Rep $\mathfrak{sl}_2(\mathbb{C})$ in a similar fashion. It turns out that the answer here is much more complicated. For example, in this setting, graphs of the form \mathcal{C} with any $n \ge 2$ loops appear. For more details³², the reader is referred to [EO03].

 $^{^{32}}$ An explanation of why these flower-shaped graphs appear can also be found in the notes from Anna's talk "Representations of representations of $\mathfrak{sl}_2(\mathbb{C})$ " at Macquarie University, available on her website www.maths.usyd.edu.au/u/romanova.

Lecture 27: Abelian categories over stacks

27.1 Modules over algebras in monoidal categories

General goal: Get a feeling for module categories (i.e. categories acted on by monoidal categories).

One of the first surprises in representation theory is that what appears to be external search (the search for all representations of a group) turns out to be an internal search (all representations occur in the regular representation). A similar phenomenon happens in the representation theory of monoidal categories.

Given a monoidal category (\mathcal{A}, \otimes) , an algebra A in \mathcal{A} is an object A together with morphisms $\mathbb{1} \to A$ and $m : A \otimes A \to A$ satisfying the axioms that one might expect. Given an algebra, let

 $\operatorname{Mod}_{\mathcal{A}}(A) = \operatorname{category} \operatorname{of} \operatorname{right} A \operatorname{-modules} \operatorname{in} \mathcal{A},$

where a right A-module in \mathcal{A} is an object M and a morphism $a : M \otimes A \to M$ satisfying the usual right module axioms. For objects $X \in \mathcal{A}$ and $M \in Mod_{\mathcal{A}}(A)$, $X \otimes M$ is a right A-module via $id_X \otimes a$, so the category $Mod_{\mathcal{A}}(A)$ is a left \mathcal{A} -module.

Meta Theorem: " \mathcal{A} -modules are A-modules."

The Meta Theorem captures the external-becomes-internal phenomenon we see in representation theory: the external search for \mathcal{A} -modules becomes an internal search for algebras in \mathcal{A} . For a precise version of the Meta Theorem in the case of a rigid semisimple category \mathcal{A} with finitely many simple objects, see [Ost03].

Example 27.1. Module categories for finite groups.

Let G be a finite group and let k be an algebraically closed field, such that k[G] is semi-simple. Let $H \subset G$ be a subgroup. The restriction functor

$$\operatorname{Rep} G \to \operatorname{Rep} H$$

is a \otimes -functor, and it gives Rep *H* the structure of a Rep *G*-module. The following exercise shows that this \mathcal{A} -module is an *A*-module (for $\mathcal{A} = \operatorname{Rep} G$, A = k[G/H]).

Exercise 27.2. Show that k[G/H] is an algebra in Rep G, and Rep $H \simeq \operatorname{Mod}_{\operatorname{Rep} G} - k[G/H]$.

Question: Are these all semisimple indecomposible Rep *G*-modules?

Theorem 27.3. [Ost03]

$$\left\{ \begin{array}{c} \text{indecomposible} \\ \text{module categories} \\ \text{over } \operatorname{Rep} G \end{array} \right\}_{\simeq} \stackrel{\sim}{\leftrightarrow} \left\{ (H, \omega) \left| \begin{array}{c} H \subset G \text{ subgroup} \\ \omega \in H^2(H, k^{\times}) \end{array} \right\}_{G\text{-conjugacy}} \right.$$

Under this correspondence, the pair (H, ω) corresponds to the category $\operatorname{Rep}^1(\widetilde{H})$ of representations of the central extension

$$k^{\times} \to \tilde{H} \to H$$

on which k^{\times} acts via the identity character.

We can sum up this example with the following motto:

"Module categories over $\operatorname{Rep} G$ are the same thing as subgroups with sauce."

Remark 27.4. 1. The proof of Theorem 27.3 is easy once one has the Meta Theorem.

- 2. Objects on the LHS are not always monoidal categories as they are when the central extension is trivial.
- 3. Question: In stacks language, Rep H as a Rep G-module corresponds to $pt/H \rightarrow pt/G$. Can one interpret $H^2(H, k^{\times})$ as twistings of QCoh(pt/H)?

27.2 Linear algebra over a stack

For the remainder of this lecture, we will summarize a short note of Gaitsgory on categories with actions of algebraic stacks [Gai05]. Recall our motivation: In lecture 25, we introduced the "constructable" anti-spherical category \mathcal{M}^{asp} . We wish to show that $\mathcal{M}^{asp} \simeq D^{G^{\vee} \times \mathbb{G}_m}(\widetilde{\mathcal{N}})$, and our strategy for doing so is to view \mathcal{M}^{asp} progressively as a module over $\operatorname{QCoh}(\operatorname{pt}/G^{\vee})$, then $\operatorname{QCoh}(\mathfrak{g}^{\vee}/G^{\vee})$, then $\operatorname{QCoh}(\widetilde{\mathcal{N}}/G^{\vee})$. The following language will be useful.

The notion of a category over an algebraic stack:

Let \mathcal{C} be a k-linear category. Assume that \mathcal{C} is closed under inductive limits. (E.g. Vect and $\operatorname{QCoh}(X)$ are okay, $\operatorname{Vect}_{f.d.}$ and $\operatorname{Coh}(X)$ are not.) Let \mathcal{Y} be a stack. Our goal is to address the following question:

What does it mean for \mathcal{C} to be linear over \mathcal{Y} ?

Step 1: Start by assuming \mathcal{Y} is an affine scheme; i.e. $\mathcal{Y} = \operatorname{Spec} A$ for some k-algebra A. In this case, \mathcal{C} being linear over \mathcal{Y} should mean that \mathcal{C} is A-linear; i.e. we have a map

$$A \to Z(\mathcal{C}) := \operatorname{End}(id_{\mathcal{C}}).$$

Example 27.5. If $X \xrightarrow{f} \operatorname{Spec} A$, then $\operatorname{QCoh}(X)$ is A-linear.

Claim: If \mathcal{C} is A-linear, then $\operatorname{QCoh}(\mathcal{Y})$ acts on \mathcal{C} ; i.e. we have

$$\operatorname{QCoh}(\mathcal{Y}) \times \mathcal{C} \to \mathcal{C}.$$

Example 4.5. (continued) QCoh(Spec A) acts on QCoh(X) via $f^*(-) \otimes (-)$.

Proof. We want to define $M \otimes X$ for $M \in \operatorname{QCoh}(\mathcal{Y}), X \in \mathcal{C}$. If $M = A^I$ (recall that $\mathcal{Y} = \operatorname{Spec} A$), define $M \otimes X := X^I$. In general, choose a presentation

$$A^I \to A^J \twoheadrightarrow M$$

of M, and define $M \otimes X := \operatorname{coker}(A^I \otimes X \to A^J \otimes X)$, where the ij^{th} -matrix coefficient is given by the a_{ij} action on $\operatorname{Hom}(X, X)$.

Exercise 27.6. Show that this is independent of presentation.


We conclude that $\operatorname{QCoh}(\mathcal{Y}) \circlearrowright \mathcal{C}$.

On the other hand, given $\operatorname{QCoh}(\operatorname{Spec} A) \circ \mathcal{C}$, we can also go back; that is, we can recover the A-linear structure on \mathcal{C} . Indeed, because $\mathbb{1}_{\operatorname{QCoh}(\operatorname{Spec} A)} = A$, for any $M \in \mathcal{C}$ we have a map

$$\operatorname{End}(A) = A \to \operatorname{End}(\mathbb{1} \otimes M) = \operatorname{End}(M).$$


Hence we obtain a map $A \to Z(\mathcal{C})$; i.e. an A-linear structure on \mathcal{C} .

We would like to have some notion of base change in this setting. But first we need a pull-back. For a map $A' \leftarrow A$ of k-algebras, we want functors f_*, f^* fitting into a diagram



We define the category $\mathcal{C}' = \mathcal{C} \times_S S'$ as follows:

• **Objects:** objects of C, equipped with an additional action of A' such that



commutes.

• Morphisms: morphisms in C compatible with the A'-action.

Exercise 27.7. (If you are stuck, ask Emily!) Show that $\mathcal{C}' = \mathcal{C} \times_S S'$ is abelian.

Example 27.8. Let C = QCoh(S) = A-Mod. Then $C \times_S S'$ is the category of A-modules with A'-actions; that is,

$$\mathcal{C} \times_S S' = A' \operatorname{-Mod} = \operatorname{QCoh}(S').$$

Returning to the general setting, we see that our desired functors

$$\begin{array}{c} \mathcal{C} \times_S S' & \xleftarrow{f^*} \mathcal{C} \\ & \downarrow & & \downarrow \\ & S' & \longrightarrow S \end{array}$$

are given by

$$f_* = \text{ forget } A'\text{-structure}, \\ f^* : X \mapsto A' \otimes X.$$

It is not difficult to see that they form an adjoint pair (f^*, f_*) .

27.3 A brief review of descent

The idea: Often in mathematics, we can construct an object by constructing it "locally".

More specifically, the principle of descent is that for some mathematical object Z,

giving
$$Z$$
 on $X \longleftrightarrow$ giving pieces of Z on
a "cover" of $X +$ glue.

Remark 27.9. One of Grothendieck's great insights is that here "cover" can be interpreted *very* generally.


Different types of objects require different glue:

• Function: Define a function f on X by giving functions f_i on U_i such that

$$f_i|_{U_{ij}} = f_j|_{U_{ij}}.$$


In this setting the glue is a "truth value" (i.e. the functions agree or don't). If you want to be more fancy, you can interpret a function as a "sheaf of truth values".

• Sheaf: Define a sheaf \mathcal{F} by giving sheaves \mathcal{F}_i on U_i and morphisms $\alpha_{ij} : \mathcal{F}|_{U_{ij}} \to \mathcal{F}|_{U_{ij}}$ such that



commutes. In this setting, the morphisms are the glue, which is subject to the restriction imposed by the diagram.

• Category: Define a category \mathcal{C} by giving categories \mathcal{C}_i on U_i , functors $\alpha_{ij} : \mathcal{C}_i|_{U_{ij}} \to \mathcal{C}_j|_{U_{ij}}$, and natural transformations $\beta_{ijk} : \alpha_{jk} \circ \alpha_{ij} \xrightarrow{\sim} \alpha_{ik}$ such that




commutes. In this setting, both the functors and natural transformations make up the glue.

Topological approach: Again coherence conditions come from a topological space. Assume for simplicity $X = \bigcup_{i \in I} U_i$ is a cover (in the classical sense). Then there is a bijection

simplicial complexes with I vertices \leftrightarrow the indexing set I of the cover

 $n\text{-simplices} \leftrightarrow \begin{array}{c} n\text{-element subsets } K \text{ of } I \\ \text{such that } \cap_{i \in K} U_i \neq \emptyset. \end{array}$


Example 27.10.



Coherence conditions come from cells:


- Function: 0 and 1 cells.
- Sheaf: 0, 1, and 2 cells.
- **Category:** 0, 1, 2, and 3 cells.
- Etc.

Example 27.11. Coherence condition (27.1) above corresponds to the 3-cell in the 3-simplex:



27.4 Back to sheaves over stacks

Return to the setting of the end of Section 27.2 and the diagram



Lemma 27.12. (Key Lemma) Sheaves of categories satisfy descent with respect to flat covers of affine schemes; that is, if $S' \to S$ is a flat cover of affine schemes, then,

 $\begin{array}{ccc} \text{sheaf of categories} & \underset{\text{over } S}{\to} & \xrightarrow{\text{sheaf of categories over } \mathcal{C}'} \\ & + \text{ descent data} \end{array}.$

Now let \mathcal{Y} be an algebraic stack in the faithfully-flat sense, with affine diagonal. This means that any morphism

$$\operatorname{Spec} A \to \mathcal{Y}$$

with source an affine scheme is affine. Recall from Emily's IFS talk *Sheaves on stacks*, that we define a quasi-coherent sheaf on \mathcal{Y} by imagining that there is a quasi-coherent sheaf " \mathcal{F} " on \mathcal{Y} and axiomatising what this would mean for f^* " \mathcal{F} " for all $f : \operatorname{Spec} A \to \mathcal{Y}$.

Given an algebraic stack \mathcal{Y} as above, let $\operatorname{Sch}_{\mathcal{Y}}^{\operatorname{aff}}$ denote the category of affine schemes over \mathcal{Y} . We define a *sheaf of categories over* \mathcal{Y} to be an assignment $\mathcal{C}^{\operatorname{sh}}$:

- 1. $S \in \operatorname{Sch}_{\mathcal{V}}^{\operatorname{aff}} \mapsto \mathcal{C}_S$, linear over S;
- 2. for each $S' \to S$ in Sch^{aff}_{\mathcal{Y}}, an S-linear functor $f^* : \mathcal{C}_S \to \mathcal{C}_{S'}$ inducing an equivalence $\mathcal{C}_S \times_S S' \xrightarrow{\sim} \mathcal{C}_{S'}$;
- 3. for $S'' \xrightarrow{g} S' \xrightarrow{f} S$ in $\operatorname{Sch}_{\mathcal{Y}}^{\operatorname{aff}}$, an isomorphism $g^* \circ f^* \simeq (f \circ g)^*$ such that 3-way compatibility holds.

Remark 27.13. This is holistic definition. We could also give a "generators and relations" definition via a fixed flat cover.

Example 27.14. If $\mathcal{Y} = \text{pt}$, then the assignment $S \mapsto \text{QCoh}(S)$ is a sheaf of categories on \mathcal{Y} .

Exercise 27.15. Given \mathcal{C}^{sh} on \mathcal{Y} , define $\Gamma(\mathcal{Y}, \mathcal{C}^{sh})$ and show that it is a Vect_{\mathcal{Y}}-module. (Here Vect_{\mathcal{Y}} is the category of vector bundles on \mathcal{Y} .)

Suppose that \mathcal{Y} satisfies:

- 1. \mathcal{Y} is locally Noetherian and every quasicoherent sheaf is a limit of coherent sheaves, and
- 2. every coherent sheaf is a quotient of a vector bundle (e.g. "enough ample line bundles").

Then we have the following theorem. (See Gaitsgory's note [Gai05] for a proof.)

Theorem 27.16.

sheaves of categories
over
$$\mathcal{Y} \leftrightarrow \operatorname{Vect}_{\mathcal{Y}}$$
-modules
 $\mathcal{C}^{\operatorname{sh}} \mapsto \Gamma(\mathcal{Y}, \mathcal{C}^{\operatorname{sh}}).$

Example 27.17. (Key example) Let G be a linear algebraic group. Then

$$\begin{array}{c} \text{sheaf of categories} \\ \text{on } \mathrm{pt}/G \end{array} \leftrightarrow \mathrm{Vect}_{\mathrm{pt}/G} \simeq \mathrm{Rep}^{\mathrm{f.d.}} \, G\text{-module}. \end{array}$$

To see this, recall Emily's approach to understanding sheaves on pt/G. She started with a prestack $(pt/G)^{triv}$ given by

$$(\mathrm{pt}/G)^{\mathrm{triv}}(S) = \mathrm{trivial}\ G$$
-bundle on $S \circlearrowright G(S)$,

then applied stackification to obtain the stack pt/G. A coherent sheaf on $(pt/G)^{triv}$ is the data

$$V + G(A) \circlearrowright V \otimes A$$

for every affine k-scheme S = Spec A. In other words, an algebraic representation of G. Similarly, a category over $(\text{pt}/G)^{\text{triv}}$ is the data

k-linear category $\mathcal{C} + G(A) \circlearrowright \mathcal{C}_S$

for all affine $S = \operatorname{Spec} A$.

Consider the case of a G-scheme X. Then the assignment

S/k affine \mapsto QCoh $(X_S) \circlearrowleft G(S)$

is a sheaf of categories over pt/G. On the other hand,

$$\operatorname{QCoh}^G(X) = \operatorname{QCoh}(X/G)$$

is a RepG-module. Under the above equivalence, these correspond to one another.

Remark 27.18. We will have more to say about this correspondence in a slightly different language in the next lecture.

Lecture 28: (De)equivariantisation


We explained last week that if \mathcal{Y} is an algebraic stack satisfying certain assumptions, then

sheaf of abelian categories $\mathcal{C}^{\mathrm{sh}}$ on $\mathcal{Y} \xrightarrow{}$ (i.e. category with an action of vector bundles on \mathcal{Y})

The most basic example of this is the following. Let G be a linear algebraic group, and $\mathcal{Y} = \text{pt}/G$. Recall that pt/G is the stackification of the prestack $(\text{pt}/G)^{\text{triv}}$, which is given by the assignment

$$S \mapsto G \times S$$

of a scheme S to the trivial G-bundle on S. As a stack in groupoids, this assignment sends



Recall that a quasi-coherent sheaf on pt/G is the data of a vector space V (the value of this sheaf on a point), together with an action of the S-points G(S) on $V \times S$ for all test schemes S. This is known as an algebraic representation.

Similarly, a sheaf of categories on pt/G (an object on the left hand side of the correspondence above) is the data of a category \mathcal{C} (the value of this sheaf on a point), together with an action of the S-points G(S) on \mathcal{C}_S for all test schemes S.

Hence for the stack pt/G, the correspondence above matches


$$\left(S \mapsto \mathcal{C}_S = \begin{array}{c} \text{abelian category} \\ \text{with } G(S) \text{-action} \end{array}\right) \longleftrightarrow \operatorname{Rep}_{f.d.} G \text{-module.}$$

28.1 A simple example of this phenomenon

Remark 28.1. This example should have come earlier, but Geordie didn't discover the very clear reference [DGNO10] providing this perspective until last week.

Let G be a finite group and \mathcal{C} an additive monoidal category.

Theorem 28.2. There are functors "equivariantisation" and "deequivariantisation" which allow us to move between categories with *G*-actions and categories with Rep *G*-actions:



On Karoubian k-linear categories, the upper two arrows provide equivalences of categories. The lower arrows can be made into equivalences too, with additional minor assumptions.

How do these functors work?

28.2 Equivariantisation

Suppose G acts on a category \mathcal{M} ; i.e. we have morphisms $F_g : \mathcal{M} \to \mathcal{M}$ for each element $g \in G$ and natural isomorphisms $\mu_{gh} : F_g \circ F_h \xrightarrow{\sim} F_{gh}$ for each pair $g, h \in G$. (See lecture 26 for a review of this notion.)

An equivariant object in \mathcal{M} is a tuple $(X, \{u_g\}_{g \in G})$ where $X \in \mathcal{M}$ and $u_g : F_g(X) \xrightarrow{\sim} X$ such that the diagram

$$F_{g}(F_{h}(X)) \xrightarrow{F_{g}(u_{h})} F_{g}(X)$$

$$\downarrow^{\mu_{gh}(X)} \qquad \qquad \downarrow^{u_{g}}$$

$$F_{gh}(X) \xrightarrow{u_{gh}} X$$

commutes.

An equivariant morphism in \mathcal{M} is a morphism $X \xrightarrow{f} Y$ in \mathcal{M} which commutes with all u_g ; that is, a morphism such that the diagram

$$F_g(X) \xrightarrow{u_g} X$$

$$F_g(f) \downarrow \qquad \qquad \downarrow f$$

$$F_g(Y) \xrightarrow{u_g} Y$$

commutes for all u_q .

The equivariantisation \mathcal{M}^G of \mathcal{M} is the category whose objects are equivariant objects in \mathcal{M} and whose morphisms are equivariant morphisms in \mathcal{M} .

Remark 28.3. The category \mathcal{M}^G is a Rep *G*-module. Indeed, given $V \in \operatorname{Rep} G$ and $(X, \{u_g\}_{g \in G}) \in \mathcal{M}^G$, define $V \otimes X := (V \otimes X, u_{g_{g \in G}}^{\vee})$, where

$$u_g^{\vee}: F_g(V \otimes X) \simeq V \otimes F_g(X) \xrightarrow{\rho_g \otimes u_g} V \otimes X.$$

Here $\rho: G \to \operatorname{Aut}(V)$ is the *G*-representation structure on *V*.

Exercise 28.4. If X is a G-scheme, show that

- 1. G acts on Coh X, and
- 2. the equivariantisation of Coh(X) is equivalent to the category of G-equivariant sheaves on X:

$$(\operatorname{Coh}(X))^G \simeq \operatorname{Coh}^G(X).$$

28.3 Deequivariantisation

Let A be the algebra of functions from G to k. The algebra A carries commuting left and right G-actions:

- Left action: $(g \cdot f)(x) = f(g^{-1}x)$
- Right action: $(f \cdot g)(x) = f(xg)$.

The left action of G gives A the structure of a G-representation, and it is an algebra object in the category Rep G. The right G-action means that G acts on A, preserving its Rep Galgebra structure.

If \mathcal{N} is a Rep *G*-module, define a category \mathcal{N}_G , the *deequivariantisation* of \mathcal{N} , by

- **Objects:** A-modules in \mathcal{N} ; i.e. objects Y together with a morphism $a : A \otimes Y \to Y$ satisfying the appropriate module conditions
- Morphisms: A-linear morphisms in \mathcal{N} .

Example 28.5. Let \mathcal{N} be the category of *G*-equivariant coherent sheaves on *X*, and A = k[G] as above. Then

$$\mathcal{N}_G = A \text{-modules in } \operatorname{Coh}^G(X)$$
$$= \mathcal{O}_X[G] \text{-modules in } \operatorname{Coh}^G(X)$$
$$= \operatorname{Coh}^G(G \times X)$$
$$= \operatorname{Coh}(X).$$

28.4 Rep G enrichments

Both equivariantisation and deequivariantisation can be realised as a two-step process in which the first step passes through a Rep G-enrichment (i.e. a category whose Hom spaces are objects in Rep G) of our original category. We describe this perspective now.

Let \mathcal{M} be a category with an action of G, and let M, M' be equivariant objects in \mathcal{M} .

Claim 28.6. Hom_{\mathcal{M}}(M, M') is a *G*-module, and

$$\operatorname{Hom}_{\mathcal{M}^G}(M, M') = \operatorname{Hom}_{\mathcal{M}}(M, M')^G.$$

Proof. The action is given by

$$(M \xrightarrow{f} M') \xrightarrow{g_{\cdot}} \left(\begin{array}{cc} F_g(M) \xrightarrow{F_g(f)} F_g(M') \\ \downarrow_{\sim} & \downarrow_{\sim} \\ M \xrightarrow{g \cdot f} M' \end{array} \right).$$

The second statement follows from the definitions.

From \mathcal{M} , we can construct a Rep *G*-enriched category by considering just the equivariant objects of \mathcal{M} , but keeping all morphisms between those objects. This is a category where *G* fixes all objects but moves morphisms. Claim 28.6 shows that from this Rep *G*-enrichment of \mathcal{M} , we can then pass to the equivariantisation of \mathcal{M} by only keeping morphisms which are fixed by *G*.

On the other hand, given a Rep G-module \mathcal{N} , we can also construct a Rep G-enriched category. For $X, Y \in \mathcal{N}$, consider the functor

$$\operatorname{Rep} G \to k : V \mapsto \operatorname{Hom}(V \otimes X, Y).$$

Because $\operatorname{Rep} G$ is semisimple, any k-linear functor on $\operatorname{Rep} G$ is representable. The functor above is represented by


$$\underline{\operatorname{Hom}}(X,Y) \in \operatorname{Rep} G;$$

i.e.

$$\operatorname{Hom}_{\operatorname{Rep} G}(V, \operatorname{\underline{Hom}}(X, Y)) \simeq \operatorname{Hom}_{\mathcal{N}}(V \otimes X, Y).$$

Hence the category consisting of the objects of \mathcal{N} with morphisms given by $\underline{\operatorname{Hom}}(X,Y)$ is a Rep *G*-enriched category. Forgetting the Rep *G*-enrichment and taking the Karoubi completion of this category results in the deequivariantisation \mathcal{N}_G of \mathcal{N} . Note that $\operatorname{Hom}_{\mathcal{N}}(X,Y) = \underline{\operatorname{Hom}}(X,Y)^G$.

Attempt at a big picture:



Remark 28.7. The fact that non-equivariant objects are consequences of equivariant ones is not so surprising in the case of a finite group G: Any X is a summand of $\text{Ind}(X) = \bigoplus F_g(X)$, which has a canonical equivariant structure.

28.5 Deequivariantisation principle

Now we return to the setting of the beginning of the lecture. Let S = Spec A be an affine scheme with an action of a linear algebraic group H (e.g. $S = \mathfrak{g}^{\vee}$, $H = G^{\vee}$). A key tool in


[AB09, Bez16] is the *deequivariantisation principle*, which answers the question:

How can we make a category C linear over S/H?

The deequivariantisation principle states that it is enough to give:

- 1. a Rep *H*-module structure on C; and
- 2. an *H*-equivariant *A*-linear structure on \mathcal{C}_{deeq}^{33} (which is equivalent to an *H*-equivariant *A*-action on Hom_{\mathcal{C}}(*X*, $\mathcal{O}_H \otimes Y$)).

Big diagram:



Recall that since C is a Rep *H*-module, C_{deeq} consists of \mathcal{O}_H -modules in C, and \mathcal{O}_H is an ind-object in Rep_{f.d.} *H*. Hence the deequivariantisation functor sends

$$\mathcal{C} \to \mathcal{C}_{\text{deeq}} : X \mapsto \mathcal{O}_H \otimes X.$$

Exercise 28.8. The functor $\mathcal{O}_H \otimes (-)$ is left adjoint to the forgetful functor $\mathcal{C}_{deeq} \to \mathcal{C}$.

Moreover,

 $\operatorname{Hom}_{\mathcal{C}_{\operatorname{deeg}}}(\mathcal{O}_H \otimes X, \mathcal{O}_H \otimes Y) = \operatorname{Hom}_{\mathcal{C}}(X, \mathcal{O}_H \otimes Y).$

Example 28.9. What is a sheaf of categories on $\mathbb{A}^1/\mathbb{G}_m$? By the deequivariantisation principle, we need to give:

- 1. A Rep \mathbb{G}_m -action on \mathcal{C} ; i.e. an auto-equivalence $M \mapsto M(1) := \operatorname{nat} \otimes M$ of \mathcal{C} . (Recall that Rep \mathbb{G}_m is freely generated³⁴ by the natural representation nat.)
- 2. A \mathbb{G}_m -equivariant $\mathcal{O}_{\mathbb{A}^1} = k[X]$ -linear structure on $\mathcal{C}_{\text{deeq}}$; i.e. a k[X]-graded module structure on

$$\operatorname{Hom}_{\mathcal{C}_{\operatorname{deeq}}}(\mathcal{O}_{\mathbb{G}_m} \otimes M, \mathcal{O}_{\mathbb{G}_m} \otimes N) = \operatorname{Hom}_{\mathcal{C}}(M, \bigoplus_{m \in \mathbb{Z}} N(m))$$
$$= \bigoplus_{m \in \mathbb{Z}} \operatorname{Hom}_{\mathcal{C}}(M, N(m))$$

³³Here C_{deeq} is the deequivariantisation of C, the category we denoted in the previous section by C_G . We are making this notational switch to align with Bezrukavnikov's notation in what is coming.

 $^{^{34}}$ as a k-linear additive tensor category

with M in degree 1. In other words, this is the same data as a morphisms of functors $id \rightarrow (1)$.


We conclude that a sheaf of categories on $\mathbb{A}^1/\mathbb{G}_m$ consists of: (1) a category \mathcal{C} , an autoequivalence $M \mapsto M(1)$ of \mathcal{C} , and a natural transformation $X : id \to (1)$.

The point of this example:

sheaf over $\mathbb{A}^1/\mathbb{G}_m \longleftrightarrow$ auto-equivalence (1) + morphism $id \to (1)$ "geometric" "combinatorial"

This pattern is a feature in [AB09] and [Bez16].

Exercise 28.10. (Worthwhile). For each \mathcal{Y} in the diagram



calculate $C_{\mathcal{Y}}$ in terms of the above data.

A bigger fish: What does it mean to give a sheaf of categories on \mathfrak{g}/G ?

Based on the deequivariantisation principle, this is

 $\operatorname{Rep} G\operatorname{-module} \mathcal{C} + \frac{G\operatorname{-equivariant} \mathcal{O}_{\mathfrak{g}}\operatorname{-linear}}{\operatorname{structure} \text{ on } \mathcal{C}_{\operatorname{deeq}}}.$

Key observation: (To be explained below.) Via Tannakian formalism, the $\mathcal{O}_{\mathfrak{g}}$ -linear structure on \mathcal{C}_{deeq} is simply an "endomorphism".

28.6 Tannakian formalism

Let G/k be a group scheme, and For : Rep $G \to k$ the forgetful functor. We have a k-group functor:

$$A \mapsto \operatorname{Aut}^{\otimes}(\operatorname{For} \otimes A) = \otimes$$
-automorphisms of $A \otimes \operatorname{For}$.

(A \otimes -automorphism of $A \otimes$ For is a collection of functors $F_v : V \otimes A \to V \otimes A$ for all $V \in \operatorname{Rep}_{f.d.} G$ such that

commutes.)

We discover that "G can be recovered from its category of representations":

Theorem 28.11.

$$G(A) \simeq \operatorname{Aut}^{\otimes}(A \otimes \operatorname{For}).$$

How can we think about this theorem? The A-points of G clearly give \otimes -automorphisms, so we have a map

$$G(A) \to \operatorname{Aut}^{\otimes}(\operatorname{For} \otimes A).$$

The miracle is that this map is an isomorphism.

Question: How can we recover the Lie algebra $\mathfrak{g} = \operatorname{Lie} G$ from Tannakian formalism?

Definition 28.12. A \otimes -derivation of For \otimes *A* is an endomorphism

$$N_V: V \otimes A \to V \otimes A$$

for all $V \in \operatorname{Rep} G$ such that $N_{V \otimes V'} = N_V \otimes 1 + 1 \otimes N_{V'}$.

Theorem 28.13.

$$\mathfrak{g} \otimes A \simeq \operatorname{End}^{\otimes}(\operatorname{For} \otimes A).$$

Exercise 28.14. Deduce Theorem 28.13 from Theorem 28.11 using

$$\mathfrak{g} = \ker \left(G\left(k[\epsilon]/(\epsilon^2)\right) \right) \to G(k).$$

A rather startling consequence of this is the following. If A is a k-algebra with a G-action, then

$$\begin{array}{l} G\text{-equivariant}\\ \otimes\text{-derivation}\\ V\mapsto V\otimes A \end{array} = \begin{array}{c} \text{element}\\ x\in (\mathfrak{g}\otimes A)^G \end{array} \iff \operatorname{Hom}_{\mathrm{v.s.}}^G(\mathfrak{g}^*,A)\\ \Leftrightarrow \begin{array}{c} G\text{-equivariant}\\ \operatorname{Hom}_{\mathrm{alg.}}(\mathcal{O}_{\mathfrak{g}},A)\\ \Leftrightarrow \end{array} \\ \text{Spec } A/G \to \mathfrak{g}/G. \end{array}$$

The moral is that, for a G-equivariant algebra, equipping the functor $V \mapsto V \otimes A$ with a G-equivariant tensor derivation is the same as equipping its category of equivariant sheaves with a \mathfrak{g}/G -linear structure.

Lecture 29: Coherent sheaves on base affine space

Today we will discuss coherent sheaves on base affine space. But before doing so, we'll start

with a brief reminder on where we're going. Let $\operatorname{Coh}_{\operatorname{free}}^{G^{\vee}}(\widetilde{\mathcal{N}}) \subset \operatorname{Coh}^{G^{\vee}}(\widetilde{\mathcal{N}})$ be the full subcategory consisting of $V \otimes \mathcal{O}_{\widetilde{\mathcal{N}}}(\lambda)$ for $V \in \operatorname{Rep} G^{\vee}$ and λ a character of T^{\vee} . As in Lectures 24 and 25, let \mathcal{H} be the affine Hecke category and \mathcal{M}^{asph} the categorical anti-spherical module. Our goal is to give an equivalence of categories $\operatorname{Coh}^{G^{\vee}}(\widetilde{\mathcal{N}}) \xrightarrow{\sim} \widetilde{\mathcal{M}}^{\operatorname{asph}}$. To do so, we will construct a monoidal functor F fitting into the following diagram.

$$\operatorname{Coh}^{G^{\vee}}(\widetilde{\mathcal{N}}) \longleftrightarrow \operatorname{Coh}_{\operatorname{free}}^{G^{\vee}}(\widetilde{\mathcal{N}}) \xrightarrow{\operatorname{construct}}_{F} \mathcal{H} \longrightarrow \mathcal{M}^{\operatorname{asph}}$$

Once we have constructed $F : \operatorname{Coh}_{\operatorname{free}}^{G^{\vee}}(\widetilde{\mathcal{N}}) \to \mathcal{M}^{\operatorname{asph}}$, we can extend F to a functor from $\operatorname{Coh}^{G^{\vee}}(\widetilde{\mathcal{N}})$ using the fact that any coherent sheaf admits a resolution via vector bundles. (This is analogous to the extension from an A-linear structure to an action of QCoh(Spec A)) which occurred at the start of Lecture 27.)

How might we go about constructing such a functor F? The key philosophical observation is that F can be built "softly" once one has a functor $\operatorname{Rep} G^{\vee} \to \mathcal{H}$. More precisely, one of the main technical lemmas in [AB09] (in a somewhat diluted form, to aid comprehensibility at this point) is the following.

Theorem 29.1. Let \mathcal{C} be an additive monoidal category, and $F : \operatorname{Rep} G^{\vee} \to \mathcal{C}$ a \otimes -functor.

1. If N is a \otimes -derivation of F (cf. Lecture 28), then F extends to a \otimes -functor

$$F: \operatorname{Coh}_{\operatorname{free}}(\mathfrak{g}^{\vee}/G^{\vee}) \to \mathcal{C}.$$

2. Suppose that we can upgrade F to a \otimes -functor Rep $(G^{\vee} \times T^{\vee}) \to \mathcal{C}$. Moreover, suppose that we have arrows

$$b_{\lambda}: F(V_{\lambda}) \to F(k_{\lambda})$$

satisfying the "Plücker relations" (and an "acyclicity condition"). Then F extends to a functor

$$F: \operatorname{Coh}_{\operatorname{free}}^{G^{\vee}}(\mathcal{B}^{\vee}) \to \mathcal{C}.$$

(Here \mathcal{B}^{\vee} denotes the flag variety of G^{\vee} .) Combining 1 and 2 we obtain a functor

$$F: \operatorname{Coh}_{\operatorname{free}}^{G^{\vee}}(\mathcal{B}^{\vee} \times \mathfrak{g}^{\vee}) \to \mathcal{C}.$$

3. Suppose that for all λ , the arrow $b_{\lambda} \circ N_{V_{\lambda}} : F(V_{\lambda}) \to F(V_{\lambda}) \to F(k_{\lambda})$ is equal to zero. Then F extends to

$$F: \operatorname{Coh}_{\operatorname{free}}^{G^{\vee}}(\widetilde{\mathcal{N}}) \to \mathcal{C}.$$

For the remainder of the lecture we will work entirely on the Langland's dual side, so from here on, we drop all checks from our notation.

29.1 Serre's description of coherent sheaves on \mathbb{P}^n

Serre's paper [Ser55] was very influential, and is recommended reading for any readers who have not done so. It was one of the first papers dealing in detail with coherent sheaves and their cohomology. The last part gives a beautiful description of coherent sheaves on projective space, which we now recall.

The variety \mathbb{P}^n has a standard affine cover

$$\mathbb{P}^n = \bigcup_{i=0}^n U_i, \ U_i \simeq \mathbb{A}^n.$$

Using this cover, one way to view a coherent sheaf on \mathbb{P}^n is as

$$\begin{array}{c} \text{collection of } n+1 \text{ modules} \\ \text{over } k[x_1, \dots, x_n] \end{array} + \text{glue.}$$

But this is not very practical. Instead, we can make the observation that

$$\operatorname{Coh} \mathbb{P}^n = \operatorname{Coh}^{\mathbb{G}_m}(\mathbb{A}^{n+1} \setminus \{0\}) \simeq \operatorname{Coh}^{\mathbb{G}_m}(\mathbb{A}^{n+1}) / \operatorname{Coh}^{\mathbb{G}_m}(\{0\}).$$

If you haven't thought about this before, the second equivalence shouldn't be entirely obvious. Now,

$$\operatorname{Coh}^{\mathbb{G}_m}(\mathbb{A}^{n+1}) =$$

$$finitely generated graded$$

$$S = k[x_0, \dots, x_n] \text{-modules}$$

$$finite-dimensional$$

$$\operatorname{modules}$$

Hence

$$\operatorname{Coh} \mathbb{P}^{n} = \left(\begin{array}{c} \text{finitely generated} \\ \text{graded } S \text{-modules} \end{array} \right) \middle/ \left\langle \begin{array}{c} \text{finite dimensional} \\ S \text{-modules} \end{array} \right\rangle$$

Here $\langle - \rangle$ denotes a "Serre subcategory", and the quotient is a "Serre quotient". (Perhaps this language shouldn't be a surprise!) If this is new to you, the following exercise is recommended.

Exercise 29.2. 1. Convince yourself that it gives the right answer for $\operatorname{Coh}(\mathbb{P}^0)$.

2. The functor $\operatorname{Coh} \mathbb{P}^n \to S$ -grmod is given by

$$\Gamma := \operatorname{Hom}(\bigoplus_{n \in \mathbb{Z}} \mathcal{O}(n), -)$$

Hence describe $\Gamma(\mathcal{O}(n))$ and $\Gamma(\text{skyscraper})$.

Remark 29.3. We can view these descriptions through the lens of descent. The "classical" cover $\mathbb{P}^n = \bigcup U_i$ leads to our not-very-useful description. The cover

$$\mathbb{A}^{n+1} \setminus \{0\} \to \mathbb{P}^n$$

leads to Serre's description. Note that $\mathbb{A}^{n+1}\setminus\{0\}$ is "almost" affine.

29.2 How can we describe coherent shaves on G/B?

If $G = \mathrm{SL}_2$, $G/B \simeq \mathbb{P}^1$, and we can use Serre's description of coherent sheaves on \mathbb{P}^n to describe coherent sheaves on G/B. For other groups, a similar philosophy can be employed. At first glance, three options present themselves for describing coherent sheaves on G/B:

- 1. via covers (not very useful);
- 2. using $\operatorname{Coh}^G(G/B) \simeq \operatorname{Coh}(\operatorname{pt}/B) \simeq \operatorname{Rep} B$;
- 3. via base affine space (analogue of Serre's description).

What is **base affine**³⁵ **space**? Let $U \subset B$ be the unipotent radical. Then G/U is a $G \times T$ -variety via the *T*-action

$$gU \cdot t = gtU.$$

This is well-defined because T normalizes U. Hence we have a quotient map

$$G/U \to G/B \simeq (G/U)/T.$$

Example 29.4. Let $G = SL_2$. Then $G \circlearrowright \mathbb{C}^2$ and

$$\operatorname{stab}\begin{pmatrix}1\\0\end{pmatrix} = \begin{pmatrix}1 & *\\0 & 1\end{pmatrix} = U$$

Hence $G/U \simeq \text{orbit of} \begin{pmatrix} 1 \\ 0 \end{pmatrix} \simeq \mathbb{C}^2 \setminus \{0\}$. The *T*-action is given by scaling: $x \cdot \lambda = \lambda x$, and $\left(\mathbb{C}^2 \setminus \{0\}\right) / \mathbb{C}^{\times} = \mathbb{P}^1 \mathbb{C} = G/B.$

In general, G/U is quasi-affine, meaning that it is an open set inside an affine variety.

Why? Denote by V_{λ} a simple highest weight module of G and $v_{\lambda} \in V_{\lambda}$ a fixed highest weight vector. We can choose $\{\lambda_1, \ldots, \lambda_n\}$ such that the stabiliser of

$$v := (v_{\lambda_1}, \dots, v_{\lambda_n}) \in V := V_{\lambda_1} \oplus \dots \oplus V_{\lambda_n}$$

in G is U. Hence

$$\begin{array}{c} G/U \hookrightarrow V\\ g \mapsto g \imath \end{array}$$

gives a map $G/U \xrightarrow{\text{open}} \overline{G/U} \subset V$ and $\overline{G/U}$ is affine.

Ring of functions: Assume chark = 0. Then we have the Peter-Weyl theorem:

$$k[G] = \bigoplus_{\lambda \in X_+} V_{\lambda} \oplus V_{\lambda}^*$$
 as $G \times G$ -representations.

³⁵Note that it is almost never affine, so the nomenclature is a little strange!

Hence,

$$k[G/U] \simeq k[G]^U \simeq \bigoplus_{\lambda \in X_+} V_\lambda \otimes (V_\lambda^*)^U \simeq \bigoplus_{\lambda \in X_+} V_\lambda.$$

The first isomorphism follows from the fact that the right action of U on G is free, and the final isomorphism follows from $(V_{\lambda}^*)^U \simeq \mathbb{C}$. This gives us a description of k[G/U] in terms of representations of G, but how do we describe multiplication from this perspective?

The torus T acts on $(V_{\lambda}^*)^U$ via the character λ , so T acts on $V_{\lambda} \simeq V_{\lambda} \otimes (V_{\lambda}^*)^U$ with character λ . The multiplication map

$$m: \left(\bigoplus_{\lambda \in X_+} V_{\lambda}\right) \otimes \left(\bigoplus_{\lambda \in X_+} V_{\lambda}\right) \to \bigoplus_{\lambda \in X_+} V_{\lambda}$$

is T-equivariant, so it must map

$$V_{\lambda} \otimes V_{\mu} \to V_{\lambda+\mu}$$

The space $\operatorname{Hom}_G(V_\lambda \otimes V_\mu, V_{\lambda+\mu})$ is one-dimensional, hence there is a unique map

$$m_{\lambda,\mu}: V_{\lambda} \otimes V_{\mu} \to V_{\lambda+\mu}: v_{\lambda} \otimes v_{\mu} \mapsto v_{\lambda+\mu}.$$

(Note that the multiplication map could also be zero on this component, but this can be ruled out.) This gives the multiplication in k[G/U].

Remark 29.5. This is sometimes referred to as *Chevalley multiplication* on k[G/U].

Example 29.6. If $G = SL_2$, $k[G/U] = k[x, y] = \bigoplus_{m \in \mathbb{Z}} k[x, y]_m$, and multiplication

$$k[x,y]_m \otimes k[x,y]_{m'} \to k[x,y]_{m+m'}$$

is the unique SL₂-equivariant morphism which sends $x^m \otimes x^{m'}$ to $x^{m+m'}$.

Important Notation:

$$G/U \xrightarrow{\text{open dense}} \overline{G/U} = \operatorname{Spec} k[G/U]$$

base affine space affine closure

We should think that these two spaces are "almost the same", like \mathbb{A}^{n+1} and $\mathbb{A}^{n+1} \setminus \{0\}$.

Exercise 29.7. (Well worth thinking about!) Describe the ideal of the boundary of G/U inside $\overline{G/U}$ in terms of the above.

29.3 Motivational interlude

The algebra $k[G/U] = \bigoplus_{\lambda \in X_+} V_{\lambda}$ is a *T*-algebra. The deequivariantisation principle of Lecture 28 tells us that to make a category \mathcal{M} linear over G/B = (G/U)/T is the same as giving:

1. a Rep *T*-module $\mathcal{M} \iff$ an action of the character lattice *X* of *T* on a category \mathcal{M} . We will denote this action via $\lambda \cdot M = M(\lambda)$. 2. a *T*-equivariant k[G/U]-linear structure on \mathcal{M}_{deeq} ; i.e. for all $M, M' \in \mathcal{M}$, a *T*-equivariant map

$$k[G/U] \xrightarrow{\phi_{M,M'}} \operatorname{Hom}(M, \mathcal{O}_T \otimes M') = \operatorname{Hom}(M, \bigoplus M'(\lambda)).$$

By the description of k[G/U] above, this is the same data as a collection of maps

 $V_{\lambda} \to \operatorname{Hom}(M, M'(\lambda))$

for each $\lambda \in X_+$ (the λ -isotypic components of $\phi_{M,M'}$); or, equivalently (as we will explain momentarily), a collection of "highest weight arrows"

$$b_{\lambda} \in \operatorname{Hom}(V_{\lambda} \otimes M, M'(\lambda))$$

such that the associated map $\phi_{M,M'}$ is K[G/U]-linear ("Drinfeld-Plücker relations").

3. with the property that sheaves on the complement

$$\overline{G/U} \backslash G/U$$

act by zero.

Remark 29.8. This explains the arrows in Theorem 29.1.2. Note that C from Theorem 29.1 is End \mathcal{M} here.

29.4 Arrows between coherent sheaves on G/U

Now we will examine the "highest weight arrows" in the previous section more carefully. Set

$$\mathcal{O} := k[G/U] = \bigoplus_{\lambda \in X_+} V_{\lambda}$$

with Chevalley multiplication and X-grading. Then

$$\operatorname{Hom}_{\operatorname{vect\ bndls}}^{G \times T\operatorname{-eqvt}}(V_{\lambda} \otimes \mathcal{O}, \mathcal{O}(\lambda)) = \operatorname{Hom}_{G \times T\operatorname{-eqvt}}(V_{\lambda}, \mathcal{O}(\lambda))$$
$$= \operatorname{Hom}_{G\operatorname{-eqvt}}(V_{\lambda}, V_{\lambda})$$
$$= \mathbb{C}.$$

is spanned by

$$B_{\lambda}: V_{\lambda} \otimes \mathcal{O} \to \mathcal{O}(\lambda),$$


with the property that

$$B_{\lambda}|_{V_{\lambda}\otimes V_{\mu}} = m_{\lambda,\mu} : V_{\lambda}\otimes V_{\mu} \to V_{\lambda+\mu}$$

Lemma 29.9. These arrows satisfy *Plücker relations*:

$$B_{\lambda} \otimes B_{\mu} = B_{\lambda+\mu} \circ (m_{\lambda,\mu} \otimes id_{\mathcal{O}}).$$

That is, the diagram



commutes.

Proof. It is enough to check that the restriction to $V_{\lambda} \otimes V_{\mu}$ commutes. This becomes the definition of Chevalley multiplication.

Remark 29.10. Recall that the classical Plücker relations describe the homogeneous equations defining the Grassmannian of k-planes in \mathbb{C}^n in its "Plücker" embedding inside $\mathbb{P}(\Lambda^k \mathbb{C}^n)$. Analogously, the above theory can be used to describe the defining relations of the flag variety in its embedding inside a product of Grassmannians. For more on this (and the connection to moduli of vector bundles and Drinfeld's compactification), see [FGV02, §4].

We will end today's lecture by describing the analogue in this setting of the $pt/G \rightsquigarrow \mathfrak{g}/G$ upgrade that we discussed last lecture. Here, we have an upgrade

$$\operatorname{pt}/G \rightsquigarrow G \setminus (\overline{G/U})/T.$$

Proposition 29.11. Let A be a $G \times T$ -algebra. Suppose for all $\lambda \in X_+$, we are given a G-equivariant morphism

$$b_{\lambda}: V_{\lambda} \otimes A \to A(\lambda)$$

satisfying the Plücker relations. Then there exists a unique $G \times T$ -equivariant homomorphism

$$\phi: \mathcal{O} \to A$$

such that $b_{\lambda} = \phi_*(B_{\lambda}) := id_A \otimes B_{\lambda}$.

Proof. What does the last condition mean? It means that

$$V_{\lambda} \otimes \mathcal{O} \longrightarrow V_{\lambda} \otimes A$$
$$\begin{array}{c} B_{\lambda} \downarrow & \downarrow b_{\lambda} \\ \mathcal{O}(\lambda) \longrightarrow A(\lambda) \end{array}$$

commutes; i.e.

$$\phi|_{V_{\lambda}} = b_{\lambda}|_{V_{\lambda} \otimes 1}.$$

This determines ϕ uniquely: define $\phi_{\lambda} := b_{\lambda}|_{V_{\lambda} \otimes 1} : V_{\lambda} \to A$ and $\phi = \bigoplus \phi_{\lambda}$. This is clearly $G \times T$ -equivariant. One can check that ϕ is a homomorphism, which amounts to the Plücker relations holding.

Lecture 30: Constructible sheaves on finite flag varieties and braid groups


Today we return to the constructible world. In this lecture and the following one, we will see how the formalism of the last two lectures is used to produce a \otimes -functor

$$\operatorname{Coh}_{\operatorname{free}}^{G^{\vee}}(\widetilde{\mathcal{N}}) \to P_I \subset D^b_I(\mathcal{F}\ell)$$

The key tool is provided by BGK central sheaves and certain easier sheaves called *Wakimoto sheaves*. Wakimoto sheaves are special to the affine setting; however, to warm up (it's been a few weeks since we've been on this side!), we'll spend today's lecture in the finite case.

30.1 A fun calculation

Choose points $x \neq y$ on \mathbb{P}^1 :




There are two ways to include these points into \mathbb{P}^1 :

Exercise 30.1. Show that

$$j_{x!}j_{y*}\mathbb{Q}_U \simeq j_{y*}j_{x!}\mathbb{Q}_U =: \mathbb{Q}_{x!,y*}$$

(*Hint*: construct a map and then show that it is an isomorphism.)

A picture:



Claim 30.2. $H^*(\mathbb{P}^1, \mathbb{Q}_{x!,y*}) = 0.$

Proof. Let $i_x : \{x\} \hookrightarrow \mathbb{P}^1$ be the inclusion. By applying the functorial distinguished triangle


$$j_{x!}j_x^! \to id \to i_{x*}i_x^* \xrightarrow{+1}$$

to $\mathbb{Q}_{y*} := j_{y*}\mathbb{Q}_{\mathbb{P}^1 \setminus \{y\}}$, we obtain a distinguished triangle

$$j_{x!}j_x^! \mathbb{Q}_{y*} \to \mathbb{Q}_{y*} \to i_{x*}i_x^* \mathbb{Q}_{y*} \xrightarrow{+1} \cdot$$

By base change, $j_{x!}j_x^! \mathbb{Q}_{y*} \simeq \mathbb{Q}_{x!,y*}$, so the long exact sequence in cohomology

$$H^*(\mathbb{Q}_{x!,y*}) \qquad H^*(\mathbb{Q}_{y*}) = H^*(\mathbb{C}) \qquad H^*(i_{x*}i_x^*\mathbb{Q}_{y*})$$



implies the claim.

Exercise 30.3. • Prove the claim again as follows: use Artin vanishing (which we will state later) and the previous exercise to show that $H^i(\mathbb{P}^1, \mathbb{Q}_{x!,y*}[1]) = 0$ for $i \neq 0$, then compute that

$$\chi(\mathbb{P}^1, \mathbb{Q}_{x!, y*}) = 0$$

to deduce the claim.

• Find yet another proof of the claim by interpreting $H^*(\mathbb{P}^1, \mathbb{Q}_{x!,y*})$ as certain locally finite chains.

30.2 The finite Hecke category

Fix

$$G \supset B \supset T$$

as usual, taken over \mathbb{C} . In what follows we will take our coefficients in \mathbb{Q} . To simplify the discussion, we will work in

$$D^b_{B \times B}(G, \mathbb{Q}) \simeq D^b_B(G/B, \mathbb{Q}).$$

Remark 30.4. As we learned in Lecture 24, this is an approximation to the Hecke category. Everything we do below may be lifted to the Hecke category \mathcal{H} with a little more effort. However, we choose to focus on $D^b_{B\times B}(G,\mathbb{Q})$ in today's lecture to illustrate the ideas more clearly.

Recall that $D_B^b(G/B)$ comes with a convolution structure

$$*: D^b_B(G/B, \mathbb{Q}) \times D^b_B(G/B, \mathbb{Q}) \to D^b_B(G/B, \mathbb{Q}).$$

The *B*-orbits on G/B stratify G/B by the Bruhat decomposition

$$G/B = \bigsqcup_{x \in W_f} BxB/B$$

Hence we have two related (but not equal!) derived categories, each containing a corresponding category of perverse sheaves: the construcible (with respect to the Bruhat stratification) derived category

$$D^b_{(B)}(G/B,\mathbb{Q}) \supset P_{(B)},$$

and the B-equivariant derived category

$$D^b_B(G/B, \mathbb{Q}) \supset P_B.$$

Convolution gives a right action

$$D^b_{(B)}(G/B,\mathbb{Q}) \circlearrowleft D^b_B(G/B,\mathbb{Q}).$$

Notation 30.1. For $\mathcal{F} \in D_B(G/B)$, denote by $\dot{\mathcal{F}} := \mathbb{Q}_{B/B} * \mathcal{F} = For(\mathcal{F}) \in D^b_{(B)}(G/B)$ the object we get by "forgetting equivariance". This gives a map

$$P_B \rightarrow P_{(B)}$$
.

Remark 30.5. The map $\mathcal{F} \mapsto \dot{\mathcal{F}}$ is fully faithful, but the image is not closed under extensions.

Let $j_x : BxB/B \hookrightarrow G/B$. Define

$$\Delta_x := j_{x!} \mathbb{Q}_{BxB/B}[\ell(x)], \quad \nabla_x := j_{x*} \mathbb{Q}_{BxB/B}[\ell(x)]$$

in P_B .

Remark 30.6. Recall that in Lecture 20, we showed that $\{\dot{\Delta}_x\}$ and $\{\dot{\nabla}_x\}$ are standard and costandard objects in a highest weight structure on $P_{(B)}$. In general, the category P_B does not admit a highest weight structure.

Example 30.7. Let $G = SL_2$. As we discussed in Lecture 16, we can use Beilinson's description

$$\operatorname{Perv}_{(B)}(\mathbb{P}^1) \simeq \operatorname{Rep}\left(\bullet \xleftarrow{f}{e} \bullet \middle| fe = 0 \right)$$

of the category $P_{(B)}$ to see that it has five indecomposible objects:

1. $\operatorname{IC}_{id} \longleftrightarrow \left(\begin{array}{c} 0 \\ & \underbrace{\longleftarrow} \end{array} \right),$

2.
$$\operatorname{I\dot{C}}_{s} \longleftrightarrow \left(\begin{array}{c} k & \xleftarrow{} 0 \end{array} \right),$$

3. $\dot{\Delta}_{s} \longleftrightarrow \left(\begin{array}{c} k & \xleftarrow{} 0 \\ & \xrightarrow{} \end{array} k \end{array} \right),$
4. $\dot{\nabla}_{s} \longleftrightarrow \left(\begin{array}{c} k & \xleftarrow{} 0 \\ & \xrightarrow{} 0 \end{array} k \end{array} \right),$

5. $T_s \longleftrightarrow (k \longleftrightarrow k \oplus k)$, where the arrow \rightarrow embeds k into the first factor of $k \oplus k$ and the arrow \leftarrow projects onto the second factor.

The object T_s is called the "big projective" or "big tilting sheaf".

We can also use a modified version Beilinson's description to describe the indecomposible objects in the category P_B . The *B*-equivariance implies that the micolocal monodromy is zero, so we have an extra condition on our quiver representations:

$$\operatorname{Perv}_B(\mathbb{P}^1) \simeq \operatorname{Rep}\left(\bullet \xleftarrow{f}{e} \bullet \left| fe = ef = 0 \right) \right).$$

Hence there are four indecomposible objects in P_B : IC_{id}, IC_s, Δ_s , and ∇_s , which correspond to the quiver representations 1-4 above. The quiver representation 5 corresponding to the big projective does not satisfy the condition that ef = 0, so we have no analogue to T_s in P_B . With this description, it is easy to see that P_B is not a highest weight category. For instance, minimal projective resolutions of IC_{id} and IC_s are

$$\cdots \to \Delta_s \to \nabla_s \to \Delta_s \to \nabla_s \to \Delta_s \twoheadrightarrow \mathrm{IC}_s, \\ \cdots \to \nabla_s \to \Delta_s \to \nabla_s \to \Delta_s \to \nabla_s \twoheadrightarrow \mathrm{IC}_{id},$$

so P_B does not have finite homological dimension, and hence cannot admit a highest weight structure.

30.3 The Braid group

Recall that W_f has a presentation

$$W_f = \langle s \in S_f \mid s^2 = id, \underbrace{st \dots}_{m_{st}} = \underbrace{ts \dots}_{m_{st}} \rangle.$$

The braid group associated to W_f is

$$B_{W_f} := \langle \sigma_s, s \in S_f \mid \underbrace{\sigma_s \sigma_t \dots}_{m_{st}} = \underbrace{\sigma_t \sigma_s \dots}_{m_{st}} \rangle.$$

Let \underline{B}_{W_f} be the corresponding monoidal category³⁶. We have the following important and beautiful theorem.

³⁶See Lecture 26, where this was denoted $\mathcal{A}_{B_{W_{\ell}}}$.

Theorem 30.8. The assignment $\sigma_s \mapsto \Delta_s$ extends to a monoidal functor

$$\underline{B}_{W_f} \to D^b_B(G/B, \mathbb{Q}).$$

Corollary 30.9. There is a (strict) action of \underline{B}_{W_f} on $D^b_{(B)}(G/B)$.

Proof. (of Theorem 30.8) We will use the following "generators and relations" theorem of Deligne [Del97].

Theorem 30.10. Let \mathcal{A} be a monoidal category. Giving a monoidal functor

$$F:\underline{B}_{W_f}\to\mathcal{A}$$

is the same as giving

- a collection $\{F_x \in \mathcal{A}\}_{x \in W_f}$ of invertible objects, and
- a collection $\{F_x F_y \xrightarrow{\sim} F_{xy}\}_{x,y \in W_f \text{ with } \ell(xy) = \ell(x) + \ell(y)}$ of isomorphisms,

such that

commutes whenever $\ell(xyz) = \ell(x) + \ell(y) + \ell(z)$.

We will show that there exists a monoidal functor

$$F:\underline{B}_{W_f}\to (D^b_{(B)}(G/B),*)$$

sending $\sigma_s \mapsto \Delta_s$ by providing the data of Theorem 30.10. Most of this data can be obtained from the following lemma:

Lemma 30.11. If $\ell(xy) = \ell(x) + \ell(y)$, then $\Delta_x * \Delta_y \simeq \Delta_{xy}$ canonically.

Proof. Recall that $\Delta_x := j_{x!}(\mathbb{Q}_{BxB/B}[\ell(x)])$. Unpacking definitions, we have

$$\Delta_x * \Delta_y = m_* \left(j_! \mathbb{Q}_{BxB \times ByB/B} [\ell(x) + \ell(y)] \right).$$

If $\ell(x) + \ell(y) = \ell(xy)$, then we have a commutative diagram:

$$BxB \times_B ByB/B \xrightarrow{m'} BxyB/B$$

$$\downarrow^j \qquad \qquad \qquad \downarrow^{j_{xy}}$$

$$G \times_B G/B \xrightarrow{m} G/B$$

Hence,

$$\Delta_x * \Delta_y = j_{xy!} \left(\mathbb{Q}_{BxyB/B}[\ell(xy)] \right) = \Delta_{xy}. \quad \Box$$

Lemma 30.11 provides the canonical isomorphisms $F_x F_y \xrightarrow{\sim} F_{xy}$ of Deligne's theorem, and it is easy to check that (30.1) commutes if $\ell(xyz) = \ell(x) + \ell(y) + \ell(z)$. Hence to complete the proof, it remains to show that Δ_x are invertible. Because

$$\Delta_x = \Delta_{s_1} * \cdots * \Delta_{s_n}$$

for $x = s_1 \cdots s_m$, it is enough to show that Δ_s is invertible for all $s \in S_f$. Lemma 30.12. For $s \in S_f$,

$$\Delta_s * \nabla_s \simeq \nabla_s * \Delta_s \simeq \Delta_{id} = \nabla_{id}.$$

Proof. (Sketch #1, a "hygenic" proof) First, one can easily reduce to SL_2 . Here we have a distinguished triangle of perverse sheaves

$$\mathbb{Q}_{\mathbb{P}^1}[1] = \mathrm{IC}_s \to \nabla_s \to \mathrm{IC}_{id} = \mathbb{Q}_{id} \xrightarrow{+1} \mathbb{Q}_{id}$$

Now, using the fact that convolution with Δ_s preserves distinguished triangles, we have another distinguished triangle

$$\Delta_s * IC_s = \mathbb{Q}_{\mathbb{P}^1} \to \Delta_s * \nabla_s \to \Delta_s \xrightarrow{+1} .$$

From this, we conclude that $\Delta_s * \nabla_s = \ker(\Delta_s \to \mathrm{IC}_s) = \mathbb{Q}_{id}$, using that the connecting map is actually a map between perverse sheaves to make sense of the kernel.


Exercise 30.13. Fill in the details of this argument.

Proof. (Sketch # 2, a more intuitive proof) We have an isomorphism

$$SL_2 \times_B SL_2 / B \xrightarrow{\sim} \mathbb{P}^1 \times \mathbb{P}^1$$
$$(g, g'B) \mapsto (gB, gg'B)$$

Under this isomorphism, $B \times_B \operatorname{SL}_2/B$ corresponds to Z := x-axis, and $\operatorname{SL}_2 \times_B B/B$ corre-

sponds to $\Delta :=$ diagonal in the following picture:



After forgetting equivariance,

$$\Delta_s * \nabla_s = p_* \left((j_{\Delta *} \mathbb{Q}_{\mathbb{P}^1 \times \mathbb{P}^1 \setminus \Delta}) \otimes (j_{Z!} \mathbb{Q}_{\mathbb{P}^1 \otimes \mathbb{P}^1 \setminus Z}[2]) \right), \tag{30.2}$$

where $j_{\Delta} : \mathbb{P}^1 \times \mathbb{P}^1 \setminus \Delta \hookrightarrow \mathbb{P}^1 \times \mathbb{P}^1$ and $j_Z : \mathbb{P}^1 \times \mathbb{P}^1 \setminus Z \hookrightarrow \mathbb{P}^1 \times \mathbb{P}^1$. By proper base change, we can compute stalks:

$$(\Delta_s * \nabla_s)_x = \begin{cases} H^*(\mathbb{P}^1, \mathbb{Q}_{0!, x*}) & \text{if } x \neq 0; \\ H^{*+2}(\mathbb{P}^1, j_{0!}\mathbb{Q}) & \text{if } x = 0. \end{cases}$$

In the second case, we have $H^{*+2}(\mathbb{P}^1, j_{0!}\mathbb{Q}) = \mathbb{Q}$, and by the fun computation from earlier, we have $H^*(\mathbb{P}^1, \mathbb{Q}_{0!,x*}) = 0$. Hence,

$$\Delta_s * \nabla_s \simeq \mathbb{Q}_{B/B}$$

which completes the proof.

Exercise 30.14. Fill in the details of this argument.

This completes the proof of Theorem 30.8.

30.4 Further properties of the braid group action

We will spend the rest of the lecture exploring the properties of the elements associated to braid group elements. All of this is in preparation for the affine case next week.

The $\{\Delta_y\}$ and $\{\nabla_y\}$ form "exceptional collections":

$$\operatorname{Hom}^{\bullet}(\Delta_x, \Delta_y) = 0 \text{ unless } x \leq y;$$

$$\operatorname{Hom}^{\bullet}(\nabla_x, \nabla_y) = 0 \text{ unless } x \geq y;$$

$$\operatorname{Hom}^{\bullet}(\Delta_x, \nabla_y) = 0 \text{ if } x \neq y.$$

Indeed, ignoring shifts,

$$\operatorname{Hom}^{\bullet}(\nabla_x, \nabla_y) = \operatorname{Hom}^{\bullet}(j_{x*}\mathbb{Q}_{X_x}, j_{y*}\mathbb{Q}_{X_y}) = \operatorname{Hom}^{\bullet}(j_y^* j_{x*}\mathbb{Q}_{X_x}, \mathbb{Q}_{X_y}) = 0 \text{ unless } y \le x,$$

and similarly for $\operatorname{Hom}^{\bullet}(\Delta_x, \Delta_y)$. (Here $X_x = BxB/B$ is the Bruhat cell associated to $x \in W_f$.)

Remark 30.15. In Lecture 20, we showed that

$$\operatorname{Hom}^{i}(\dot{\Delta}_{x}, \dot{\nabla}_{y}) = \begin{cases} \mathbb{Q} & \text{if } i = 0 \text{ and } x = y; \\ 0 & \text{otherwise.} \end{cases}$$

This was how we established the necessary Ext^2 vanishing property of the highest weight structure on $P_{(B)}$. In contrast,

$$\operatorname{Hom}^{\bullet}(\Delta_x, \nabla_y) = \begin{cases} H_B^{\bullet}(\mathrm{pt}) & \text{if } x = y; \\ 0 & \text{if } x \neq y. \end{cases}$$

Note that in this case, it is not true that the derived category of P_B agrees with the equivariant derived category, so we cannot use this computation to conclude that Ext^2 vanishing fails in P_B . However, it is an indication that P_B might not be highest weight, and indeed it is not in general. (See Remark 30.6.)

Remark 30.16. In \underline{B}_{W_f} , Hom spaces are very easy. By contrast, morphisms between Δ_x 's are complicated and quite mysterious. For example, $\text{Ext}^{\bullet}(\text{Vermas})$ is an unsolved problem.

Recall that convolution gives a right action

$$D^b_{(B)}(G/B) \circlearrowleft D^b_B(G/B) \supset P_B.$$

Within $D_{(B)}^b$ we have the highest weight category $P_{(B)} \subset D_{(B)}^b(G/B)$. The convolution of two perverse sheaves is not generally perverse, so the subcategory $P_{(B)}$ is not preserved under the action of P_B .

Definition 30.17. A perverse sheaf $\mathcal{F} \in P_{(B)}$ is *convolution exact* if $\mathcal{F} * \mathcal{G} \in P_{(B)}$ for any $\mathcal{G} \in P_B$.

Mirkovic's observation: A perverse sheaf $\mathcal{F} \in P_{(B)}$ is convolution exact if and only if \mathcal{F} is tilting.

Proof. (\implies) By highest weight formalism,

$$\mathcal{F}$$
 is tilting $\iff \operatorname{Hom}^{>0}(\mathcal{F}, \dot{\nabla}_y) \stackrel{(*)}{=} 0 \stackrel{(!)}{=} \operatorname{Hom}^{>0}(\dot{\Delta}_x, \mathcal{F}).$

We will check that equality (!) holds for \mathcal{F} convolution exact, (*) is similar.

Because \mathcal{F} is convolution exact, $\mathcal{F} * \Delta_{w_0}$ is perverse. This implies that

$$\mathcal{F} * \Delta_{w_0} \in {}^p D^{\leq 0} = \langle \dot{\Delta}_y[d] \mid y \in W_f, d \in \mathbb{Z}_{\geq 0} \rangle_{ext},$$

where the subscript "ext" denotes the closure under extensions. Hence

$$\mathcal{F} = \mathcal{F} * \Delta_{w_0} * \nabla_{w_0} \in \langle \dot{\Delta}_y * \nabla_{w_0}[d] = \dot{\nabla}_{yw_0} \mid y \in W_f, d \in \mathbb{Z}_{\geq 0} \rangle_{ext}.$$

Recall that $w_0 = y^{-1} \cdot y w_0$ and $\ell(w_0) = \ell(y^{-1}) + \ell(y w_0)$, so $\nabla_{w_0} = \nabla_{y^{-1}} * \nabla_{y w_0}$. Now any object in $\langle \dot{\Delta}_y * \nabla_{w_0}[d] = \dot{\nabla}_{y w_0} \mid y \in W_f, d \in \mathbb{Z}_{\geq 0} \rangle_{ext}$ satisfies (!), so we are done.

For the (\Leftarrow) direction, we need a very important property of affine morphisms.

Theorem 30.18. Let $f: X \to Y$ be affine. Then f_* preserves ${}^pD^{\geq 0}$, and $f_!$ preserves ${}^pD^{\geq 0}$. **Remark 30.19.** Recall Artin vanishing: if \mathcal{F}/X is perverse and X is affine,

$$H^{>0}(f_*\mathcal{F}) = H^{>0}(X,\mathcal{F}) = 0 = H_c^{<0}(X,\mathcal{F}) = H^{<0}(f_!\mathcal{F}).$$

This is exactly Theorem 30.18 when Y = pt.

Exercise 30.20. Can you prove the theorem for \mathcal{D} -modules?

Now back to our setting.

Proposition 30.21. The maps

$$BxB \times_B G/B \xrightarrow{\lambda_x} G/B$$
 and $G \times_B BsB/B \xrightarrow{\rho_x} G/B$

are affine.

Proof. (Idea) We can write $BxB \simeq U_x \cdot \{x\} \cdot B$, hence we have a commutative diagram

$$BxB \times_B G \xrightarrow{\text{mult}} G$$
$$\begin{vmatrix} \sim \\ | \sim \\ U_x \times G \xrightarrow{\text{mult}} G \end{vmatrix}$$

The space $U_x \times G$ is affine, so mult : $BxB \times_B G \to G$ is affine, hence λ_x is affine (being affine is local in the flat topology).

Exercise 30.22. Show that

$$\Delta_x * (-) = \lambda_{x!} \lambda_x^* (-) [\ell(x)],$$

$$\nabla_x * (-) = \lambda_{x*} \lambda_x^* (-) [\ell(x)],$$

$$(-) * \Delta_x = \rho_{x!} \rho_x^* (-) [\ell(x)],$$

$$(-) * \nabla_x = \rho_{x*} \rho_x^* (-) [\ell(x)].$$

(Note: $\lambda_x^! \simeq \lambda_x^* [2\ell(x)]$ because λ_x is smooth of relative dimension $\ell(x)$.)

Corollary 30.23. $\Delta_x * (-), (-) * \Delta_x$ preserve ${}^p D^{\geq 0}$, and $\nabla_x * (-), (-) * \nabla_x$ preserve ${}^p D^{\leq 0}$, whenever this makes sense.

Corollary 30.24. $\Delta_x * \nabla_y$ and $\nabla_y * \Delta_x$ are perverse for all $x, y \in W_f$.

With this we can complete the proof of Mirkovic's observation. Recall that

$${}^{p}D_{B}^{\leq 0} = \langle \Delta_{x}[d] \mid x \in W_{f}, d \in \mathbb{Z}_{\geq 0} \rangle.$$

By Corollary 30.24,

$$\langle \dot{\nabla}_y \mid y \in W \rangle * {}^p D_B^{\leq 0} \subset {}^p D^{\leq 0}.$$

Hence

tilting
$$* {}^p D_B^{\leq 0} \subset {}^p D^{\leq 0}$$
.

Similary, ${}^{p}D^{\geq 0} = \langle \nabla_x[-d] \mid x \in W_f, d \in \mathbb{Z}_{\geq 0} \rangle$, so

tilting
$$* {}^p D_B^{\geq 0} \subset {}^p D^{\geq 0}$$
.

This lets us conclude that

tilting
$$* P_B \subset P_{(B)}$$
.

This completes the proof of Mirkovic's observation.

Exercise 30.25. 1. Use these ideas to show that if $\mathcal{F} \in P_B$ is convolution exact, then $\mathcal{F} = \Delta_{id}$, (i.e. there are no interesting convolution exact sheaves in P_B).

Remark 30.26. We will see next week that there are many convolution exact $\mathcal{F} \in P_I \subset D_I(\mathcal{F}\ell)$ in the affine case.

2. Let T_x be indecomposable tilting. Show that

$$T_x * \mathbb{Q}_{P_s/B}[1] = 0$$

unless x = id. (This is related to the fact that $H^*(G/B, T_x) = 0$ unless x = id.) Use this to give another proof of the (\Leftarrow) direction of Mirkovic's observation.

Lecture 31: Affine flags, affine braids, and Wakimoto sheaves

Last week we established the existence of a monoidal functor

$$\underline{B}_{W_f} \to (D^b_B(G/B), *)$$
$$\sigma_x \mapsto \Delta_x$$

and described several of its nice properties. Today we'll describe the situation in the affine case.

As in previous lectures, let $\mathcal{F}\ell = G((t))/I$, where $I \subset G((t))$ is an Iwahori subgroup. Denote by $W = X^{\vee} \rtimes W_f$ the affine Weyl group. For simplicity, assume that G is simply connected, so W is a Coxeter group³⁷.

The same construction as we described last week in the finite case yields a monoidal functor

$$\underline{B}_W \to (D^b_B(G/B, \mathbb{Q}), *)$$

$$\sigma_x \mapsto \Delta_x.$$

31.1 Wakimoto sheaves

In Lecture 12, we described Bernstein's presentation of the extended affine Hecke algebra. We briefly recall this construction now. In Lecture 11, we proved the Iwahori-Matsumoto Lemma (Lemma 11.8): for $w \in W_f, \lambda \in X^{\vee}$,

$$\ell(t_{\lambda}w_f) = \sum_{\substack{\alpha \in R_+\\ w_f(\alpha) > 0}} |\langle \lambda, \alpha \rangle| + \sum_{\substack{\alpha \in R_+\\ w_f(\alpha) < 0}} |\langle \lambda, \alpha \rangle - 1|.$$

A consequence of this is that if $\lambda', \lambda'' \in X^{\vee}$ are dominant, then $\ell(t_{\lambda'}t_{\lambda''}) = \ell(t_{\lambda'}) + \ell(t_{\lambda''})$. (This is also intuitively clear because $t_{\lambda'}$ and $t_{\lambda''}$ translate in the same direction.) Hence in the Hecke algebra H^{38} ,

$$\delta_{t_{\lambda'}}\delta_{t_{\lambda''}}=\delta_{t_{\lambda'+\lambda''}}=\delta_{t_{\lambda''}}\delta_{t_{\lambda}'}.$$

We can write any $\lambda \in X^{\vee}$ as $\lambda = \lambda' - \lambda''$ with $\lambda', \lambda'' \in X^{\vee}$ dominant. Hence we have have a well-defined map

$$\mathbb{Z}[v^{\pm 1}][X^{\vee}] \hookrightarrow H : \lambda \mapsto \theta_{\lambda} := \delta_{t_{\lambda'}} \delta_{t_{\lambda''}}^{-1}.$$

Bernstein used this map to describe the center of H.

Wakimoto sheaves categorify this construction; i.e. given $\lambda \in X^{\vee}$, write $\lambda = \lambda' - \lambda''$ for $\lambda', \lambda'' \in X_+^{\vee}$ and define

$$J_{\lambda} := \Delta_{t_{\lambda'}} * \nabla_{t_{-\lambda''}} = \Delta_{t_{\lambda'}} * (\Delta_{t_{\lambda''}})^{-1}.$$

The J_{λ} are called *Wakimoto sheaves*.

 $^{^{37}}$ If we drop this assumption, W is a quasi-Coxeter group. This adds a few complications to proofs, but doesn't change the story in any major way.

³⁸Here we are aligning notation with Lecture 23, which differs slightly from Lecture 12.

Example 31.1. If λ is dominant, then $J_{\lambda} = \Delta_{t_{\lambda}}$. If λ is anti-dominant, then $J_{\lambda} = \nabla_{t_{\lambda}}$.

Lemma 31.2. The map $\lambda \mapsto J_{\lambda}$ extends to a monoidal functor $\underline{X}^{\vee} \mapsto (D_I^b(\mathcal{F}\ell), *)$, and hence to a monoidal functor

$$(\operatorname{Rep} T^{\vee}, \otimes) \mapsto (D^b_I(\mathcal{F}\ell), *)$$

We saw last week that $\Delta_x * \nabla_y$ is perverse for all $x, y \in W_f$. An easy adaptation of this to the affine case implies the following corollary.

Corollary 31.3. For all $\lambda \in X^{\vee}$, J_{λ} is perverse; i.e. $J_{\lambda} \in P_I \subset D_I^b(\mathcal{F}\ell)$.

Notation 31.1. Let

 $\mathcal{A} = \text{ Wakimoto-filtered objects in } P_I,$ gr $\mathcal{A} = \text{ image of } \operatorname{Rep} T^{\vee} \text{ in } D^b_I(\mathcal{F}\ell).$

Note that $\operatorname{gr} \mathcal{A}$ is not a full subcategory in $D_I^b(\mathcal{F}\ell)$. This notation will make sense in a moment.

Remark 31.4. It is immediate from the definitions that \mathcal{A} and $\operatorname{gr}\mathcal{A}$ are monoidal subcategories in P_I . Note that this is in contrast to the finite case, where P_B has no interesting monoidal subcategories.

Recall that in the finite case,

 $\operatorname{Hom}^{\bullet}(\Delta_x, \Delta_y) = 0 \text{ unless } x \leq y,$ $\operatorname{Hom}^{\bullet}(\nabla_x, \nabla_y) = 0 \text{ unless } y \leq x.$

On X^{\vee} , we have the *periodic order:*

 $\lambda \leq \mu$ if and only if $\mu - \lambda \in \mathbb{Z} X_+^{\vee}$.

Lemma 31.5. With respect to the periodic order,

$$\operatorname{Hom}^{\bullet}(J_{\lambda}, J_{\mu}) = 0 \text{ unless } \lambda \leq \mu,$$

$$\operatorname{Hom}^{\bullet}(J_{\lambda}, J_{\lambda}) = \mathbb{Q}.$$

Proof. For γ sufficiently dominant,


$$\operatorname{Hom}^{\bullet}(J_{\lambda}, J_{\mu}) = \operatorname{Hom}^{\bullet}(J_{\lambda} * J_{\gamma}, J_{\mu} * J_{\gamma})$$
$$= \operatorname{Hom}^{\bullet}(\Delta_{t_{\lambda+\gamma}}, \Delta_{t_{\mu+\gamma}})$$
$$= 0$$

unless $t_{\lambda+\gamma} \leq t_{\mu+\gamma}$. It is a known fact about the Bruhat order in affine type that for sufficiently dominant weights we have

$$t_{\lambda+\gamma} \leq t_{\mu+\gamma} \iff \lambda+\gamma \leq \mu+\gamma.$$

Hence $\operatorname{Hom}^{\bullet}(J_{\lambda}, J_{\mu}) = 0$ unless $\lambda \leq \mu$.

Example 31.6. Let $G = SL_2$, so $X^{\vee} = \mathbb{Z}\alpha^{\vee}$. The Homs between Wakimoto sheaves only go in one direction:



Lemma 31.5 implies:

Exercise 31.7. Every $\mathcal{F} \in \mathcal{A}$ admits a functorial³⁹ filtration (called a "Wakimoto filtration") \mathcal{F}_I indexed by upper closet subsets $I \subset X^{\vee}$ such that if $\lambda \in I$ is minimal, then

$$\mathcal{F}_I/\mathcal{F}_{I\setminus\{\lambda\}}\simeq \bigoplus m J_{\lambda}^{\oplus m_{\lambda}}$$

Remark 31.8. This exercise is an analogue of a standard (but nonetheless beautiful) feature of highest weight categories: (co)standard filtrations are unique when they exist.

The exercise gives us a functor

$$\operatorname{gr}: \mathcal{A} \mapsto \operatorname{gr} \mathcal{A}$$

 $\mathcal{F} \mapsto \bigoplus \mathcal{F}_{\{\geq \lambda\}} / \mathcal{F}_{\{>\lambda\}},$

which sends an object to its "associated graded under the Wakimoto filtration".

Exercise 31.9. The Wakimoto filtration is compatible with *, and hence gr is a \otimes -functor.

31.2 Motivational interlude

We'll take a brief motivational interlude to explain how Wakimoto sheaves fit into our bigger picture.


Move back to the coherent side, and let $G^{\vee} \supset B^{\vee}$, R^{\vee}_+ be as usual. Any G^{\vee} -module V has a B^{\vee} -filtration indexed by upper closed subsets $I \subset X^{\vee}$, namely

$$V_I = \bigoplus_{\lambda \in I} V_\lambda \subset V,$$

and the associated graded of this filtration is isomorphic to a direct sum of one-dimensional modules k_{λ} .

Example 31.10. If $G = SL_2$, a picture of this filtration is:





³⁹in particular unique!

In fact, $\operatorname{Hom}^{\bullet}(k_{\lambda}, k_{\mu}) = 0$ unless $\lambda \leq \mu$. Similarly on $\widetilde{\mathcal{N}}$, we have

$$\operatorname{Hom}_{\operatorname{Coh}^{G^{\vee}}(\widetilde{\mathcal{N}})}(\mathcal{O}(\lambda),\mathcal{O}(\mu))=0 \text{ unless } \lambda \leq \mu.$$

Our equivalence will end up mapping

$$\mathcal{O}(\lambda) \mapsto J_{\lambda}.$$

31.3 Beilinson–Gaitsgory–Kottwitz (BGK) central sheaves

Let $\mathcal{G}r = G((t))/G[[t]]$ be the affine Grassmannian and $\mathcal{F}\ell = G((t))/I$ the affine flag variety.

Theorem 31.11. There exists a central functor

$$Z: P_{G(\mathcal{O})}(\mathcal{G}r) \to P_I = P_I(\mathcal{F}\ell) \subset D_I(\mathcal{F}\ell).$$

Moreover, Z is equipped with a \otimes -derivation N.

Remark 31.12. The necessary compatibilities are checked in the appendix "braiding compatabilities" by Gaitsgory to Bezrukavnikov's [Bez04] and in more detail in a book in progress by Achar-Riche.

For the purposes of stating the next theorem, we (temporarily) extend the definition of Wakimoto sheaves: write $w \in W$ as $w = t_{\lambda}w_f$ for $t_{\lambda} \in X^{\vee}, w_f \in W_f$, and let

$$J_w = J_\lambda * \nabla_{w_f}$$

The affine analogue of Mirkovic's observation from last lecture is the following.

Theorem 31.13. [AB09]

- 1. Any convolution exact $\mathcal{F} \in P_I$ is $\{J_w \mid w \in W\}$ -filtered.
- 2. If in addition \mathcal{F} is "weakly central" (i.e. $\mathcal{F} * \mathcal{G} \simeq \mathcal{G} * \mathcal{F}$ for all \mathcal{G}), then \mathcal{F} is Wakimotofiltered (i.e. only J_{λ} for $\lambda \in X^{\vee}$ occur in the filtration above).

Corollary 31.14. $Z(\mathcal{G})$ is Wakimoto-filtered! i.e.,

$$Z: P_{G(\mathcal{O})}(\mathcal{G}r) \to \mathcal{A}$$

We omit the proof of Theorem 31.13, but note that it is not much more difficult than the proof of Mirkovic's observation that we described last week.

The following is beautiful and bears a striking resemblance to Mirkivic-Vilonen's theorem on weight functors. (In fact, according to [AB09, Remark 6], they are "equivalent".)

Theorem 31.15. 1. The following diagram commutes up to canonical isomorphism.

2. We have

$$H_c^i(\iota_w^*(Z(V))) = \begin{cases} V_\nu & \text{if } w = \nu \in X^{\vee}, i = \ell(\nu); \\ 0 & \text{otherwise.} \end{cases}$$

Remark 31.16. Part 2. is included for experts. The notation involved is the following: The $N_{\mathbb{C}((t))}$ -orbits on $\mathcal{F}\ell$ are indexed by W. Given $w \in W$, let $S_w = N_{\mathbb{C}((t))} \cdot wI/I$ and let $\iota_w : S_w \hookrightarrow \mathcal{F}\ell$ denote its inclusion.

We now have all of the ingredients that we need to define our functor:


- 1. $Z : \operatorname{Rep} G^{\vee} \to \mathcal{A};$
- 2. $F : \operatorname{Rep} T^{\vee} \xrightarrow{\sim} \operatorname{gr} \mathcal{A} \subset \mathcal{A} \text{ (not full)};$
- 3. N = nilpotent \otimes -derivation of Z, coming from monodromy of vanishing cycles;
- 4. $b_{\lambda}: J_{\lambda} \to Z(V_{\lambda})$ highest weight arrow coming from Wakimoto filtration.

Now we need to check:

(a) **Plücker relations:** Thus we need to check that the following diagram commutes:

This is immediate from the multiplicativity of the Wakimoto functor.

(b) Compatibility between b_{λ} and N: (relates \mathfrak{g}^{\vee} and G^{\vee}/N^{\vee}) We want a commutative diagram



Proof. $Z(V_{\lambda})$ has a Wakimoto filtration $\begin{pmatrix} \overline{J_{\lambda'}} \\ A < \lambda \\ \hline{J_{\lambda}} \end{pmatrix}$. Because $\operatorname{Hom}(J_{\lambda}, J_{\mu}) = 0$ for $\lambda > \mu$,

 $N_{V_{\lambda}} \circ b_{\lambda}$ maps J_{λ} into $J_{\lambda} \subset Z(V_{\lambda})$, hence $N_{V_{\lambda}}$ induces an endomorphism of J_{λ} , which has to be zero because $\operatorname{End}(J_{\lambda}) = \mathbb{Q}$ and N is nilpotent.

Now we have the following set-up:

$$\begin{array}{ccc} \widehat{\widetilde{\mathcal{N}}}_{\mathrm{aff}} & \longleftrightarrow & \mathfrak{g}^{\vee} \times \overline{G^{\vee}/U^{\vee}} \\ & \uparrow^{\mathrm{open}} & & \uparrow^{\mathrm{open}} \\ & \widehat{\widetilde{\mathcal{N}}} & \longleftrightarrow & \mathfrak{g}^{\vee} \times G^{\vee}/U^{\vee} \\ & \downarrow_{T^{\vee}\text{-bundle}} & & \downarrow_{T^{\vee}\text{-bundle}} \\ & \widetilde{\mathcal{N}} & \longleftarrow & \mathfrak{g}^{\vee} \times \mathcal{B} \end{array}$$

(For the definition of $\widehat{\widetilde{\mathcal{N}}}$, see [AB09].) By a variant of the Tannakian formalism discussed in previous lectures, the above data gives a functor

$$\operatorname{Coh}_{\operatorname{free}}^{G^{\vee} \times T^{\vee}}(\widehat{\widetilde{\mathcal{N}}}) \to \mathcal{A}$$

From this we obtain a functor

$$K^{b}\left(\operatorname{Coh}_{\operatorname{free}}^{G^{\vee}\times T^{\vee}}(\widehat{\widetilde{\mathcal{N}}})\right) \to K^{b}(\mathcal{A}).$$

The functor passes to derived categories, complexes on $\partial \widehat{\widetilde{\mathcal{N}}}$ go to zero⁴⁰, and we arrive at our desired functor:


$$\widetilde{F}: D^{G^{\vee}}(\widetilde{\mathcal{N}}) \to D^b(P_I).$$

All that remains is to see that \widetilde{F} induces an equivalence with the anti-spherical module.

Example 31.17. We'll end today's lecture with an illustrative example. Recall that we described Z(nat) explicitly for $G = \text{GL}_2$ in Lecture 23. We arrived at the following picture.

⁴⁰This fact must sadly remain a black box here, due to time constraints.

(Here the notation aligns with the exercises of Lecture 23.)



Lecture 32: Whittaker sheaves and Arkhipov-Bezrukavnikov theorem

We'll start today with a simple and beautiful example. But first we need to introduce some notation.

32.1 Averaging functors

Let $N \odot X$ be the action of an algebraic group, and denote by m and p the corresponding action and projection maps:

$$N \times X \xrightarrow{m} X$$

$$\downarrow^{p} \qquad X$$

$$X$$

We define two functors

$$D^b_c(X) \underbrace{\overbrace{Av_{N*}}^{Av_{N!}}}_{Av_{N*}} D^b_N(X)$$

by

$$Av_{N*} := m_*(\mathbb{Q}_N \boxtimes (-)[\dim N]) \simeq m_* p^*[\dim N] \simeq m_* p^![-\dim N],$$
$$Av_{N!} := m_!(\mathbb{Q}_N \boxtimes (-)[\dim N]).$$

One can imagine these functors as "smearing out" (integrating) our sheaf over the N-orbits to make it equivariant.

Remark 32.1. 1. We have

$$\mathbb{D}Av_{N*} \simeq Av_{N!}\mathbb{D}.$$

2. The functors Av_{N*} and $Av_{N!}$ fit into adjoint pairs

(For $[-\dim N]$, Av_{N*}) and $(Av_{N!}$, For $[\dim N]$).


3. The functor Av_{N*} preserves ${}^{p}D^{\leq 0}$ and $Av_{N!}$ preserves ${}^{p}D^{\geq 0}$.

Remark 32.2. These are a special case of the induction and restriction functors of Bernstein–Lunts.

32.2 Another fun calculation


Let $X = \mathbb{P}^1$ and $N = \begin{pmatrix} 1 & * \\ 0 & 1 \end{pmatrix}$. Then one can show that N-equivariant perverse sheaves are equivalent to perverse sheaves with respect to the B-orbit stratification: $P_N = P_{(B)}$. (This follows from the fact that the "forget equivariance" functor is fully faithful when the group is unipotent.) As we've discussed in previous lectures, there are 5 indecomposable objects

in $P_{(B)}$: $\Delta_s, \nabla_s, \text{IC}_s, \text{IC}_{id}$, and $T_s = P_{id}$. Their relationship is captured by the following diagram:



We have described geometric constructions for Δ_s , ∇_s , IC_s , IC_{id} , but we established the existence of T_s via formal properties of highest weight categories. Is there a geometric construction of $T_s = P_{id}$ as well?

The answer to this question is yes, and it can be realised using averaging functors. Recall the perverse sheaf $\mathbb{Q}_{x!,y*}[1]$ for $x, y \neq 0$ that we saw in Lecture 30:



Lemma 32.3.

$$Av_{N*}(\mathbb{Q}_{x!,y*}[1]) \simeq Av_{N!}(\mathbb{Q}_{x!,y*}[1]) \simeq T_s.$$

For $x, y \neq 0$, we have

Proof. Step 1: Let $j : \mathbb{C} \hookrightarrow \mathbb{P}^1$ be inclusion of the open Bruhat cell. Averaging functors commute with equivariant inclusions, so we have

$$j^*Av_{N*}(\mathbb{Q}_{x!,y*}[1]) \simeq Av_{N*}(j^*\mathbb{Q}_{x!,y*}[1]).$$

This is an N-equivariant sheaf on \mathbb{C} , so it is determined by its global sections.

Exercise 32.4. $H^i(\mathbb{C}, \mathbb{Q}_{x!,y*}[1]) = \begin{cases} \mathbb{Q} & \text{ for } i = 0, \\ 0 & \text{ otherwise.} \end{cases}$

(Hint: See the "fun calculation" from Lecture 30.)

The shift by [1] in the definition of Av produces a shift by [2] above, hence by Exercise 32.4, we have:

$$j^*Av_{N*}(\mathbb{Q}_{x!,y*}[1]) \simeq Av_{N*}(j^*\mathbb{Q}_{x!,y*}[1]) = \mathbb{Q}_{\mathbb{C}}[1].$$

Step 2: Let $i: \{0\} \hookrightarrow \mathbb{P}^1$ be inclusion of the closed Bruhat cell. Then by adjunction,

$$\operatorname{Hom}_{D_{N}^{b}}^{\bullet}(i_{*}\mathbb{Q}_{0}, Av_{N*}\mathbb{Q}_{x^{!},y^{*}}[1]) \simeq \operatorname{Hom}^{\bullet}(\operatorname{For}(i_{*}\mathbb{Q}_{0})[-1], \mathbb{Q}_{x^{!},y^{*}}[1])$$
$$\simeq \operatorname{Hom}^{\bullet}(\mathbb{Q}_{0}, i^{!}\mathbb{Q}_{x^{!},y^{*}}[2])$$
$$= \begin{cases} \mathbb{Q} & \text{if } \bullet = 0, \\ 0 & \text{otherwise.} \end{cases}$$

Step 3: By Remark 32.1, we know that $Av_{N*}\mathbb{Q}_{x!,y*}[1] \in {}^{p}D^{\leq 0}$. Moreover, by steps 1 and 2, $Av_{N*}\mathbb{Q}_{x!,y*} \in {}^{p}D^{\geq 0}$, so we conclude that $Av_{N*}\mathbb{Q}_{x!,y*} \in P_N(\mathbb{P}^1)$. Hence $Av_{N*}\mathbb{Q}_{x!,y*}$ decomposes into a direct sum of indecomposable perverse sheaves:

$$Av_{N*}\mathbb{Q}_{x!,y*}[1] \simeq \mathrm{IC}_{id}^{\oplus m_{id}} \oplus \cdots \oplus T_s^{\oplus m_{id}}$$

Step 4: Finally, by the fun calculation of last week, we have

$$H^*(Av_{N*}\mathbb{Q}_{x!,y*}) = H^*(\mathbb{Q}_{x!,y*}) = 0.$$


This implies that only T_s can occur in the decomposition of $Av_{N*}\mathbb{Q}_{x!,y*}[1]$ into indecomposables, and by steps 1 and 2, there must be only a single copy. We conclude that

$$Av_{N*}\mathbb{Q}_{x!,y*}[1] = T_s,$$

as desired.


Another construction: (which Geordie learned from K. Vilonen)

Rough idea: Perhaps, it is easy to calculate the composition series of $\mathbb{Q}_{x!,y*}$. In this case, one could dream of taking a limit as $x, y \to 0$ to obtain the big tilting sheaf:



Question: How do we formalize "taking a limit"?

Answer: Nearby cycles!



Let

$$\mathcal{F} = (j_! \mathbb{Q}_{(\mathbb{P}^1 \times \mathbb{C}) \setminus \Delta}) \otimes (j_* \mathbb{Q}_{(\mathbb{P}^1 \times \mathbb{C}) \setminus Z})[2].$$

Then

$$\mathcal{F}|_{\mathbb{P}^1 \times \{x\}} \simeq (\mathbb{Q}_{x!,0*}[1])[1]$$

is a "family of $\mathbb{Q}_{x!,0*}$ for varying x".

Exercise 32.5. (Beautiful!) $T_s \simeq \psi_f \mathcal{F}$.

Remark 32.6. (Spoiler alert): It's perverse and *B*-constructible . . . now what are its global sections?!

32.3 Lightning introduction to the Whittaker world

Let $N \subset B \subset G$ be (for a moment) finite groups of Lie type. Fix $\chi : N \to \mathbb{C}^{\times}$ a character. Given a representation V of G, consider

$$V^{(N,\chi)} = \{ v \in V \mid n \cdot v = \chi(n)v \}.$$

This is the space of *Whittaker vectors* in the representation. By Frobenius reciprocity,

$$\operatorname{Hom}_{N}(\mathbb{C}_{\chi}, V) = \operatorname{Hom}_{G}(\operatorname{Ind}_{N}^{G} \mathbb{C}_{\chi}, V),$$

so Whittaker vectors detect whether a representation is "seen" by a 1-dimensional character of N. Put another way, what irreducible representations can one get by inducing from a character of N? Precisely those that admit a non-zero Whittaker vector!

Suppose $V = Fun(X, \mathbb{C})$ for some G-space X. In this setting, what do Whittaker vectors look like?

Fix an N-orbit $U \subset X$. This is a homogeneous space for N, so $U \simeq N/K$ for some $K \subset N$. Then

$$\operatorname{Fun}(U) = \operatorname{Ind}_{K}^{N} \mathbb{C} = \bigoplus_{\lambda \in \widehat{N}} V_{\lambda} \otimes (V_{\lambda}^{*})^{K}.$$

Hence

 $\operatorname{Fun}(U)^{(N,\chi)} = \begin{cases} \mathbb{C} & \text{if } \chi \text{ is trivial on } K, \\ 0 & \text{otherwise.} \end{cases}$

So up to scaling, we have "either 0 or 1 Whittaker vectors per N-orbit" and we can tell exactly which orbits admit non-zero Whittaker vectors.

Exercise 32.7. (Important) Let X = G/B and N^- be the unipotent radical of the opposite Borel. Then

$$N^-/[N^-, N^-] \simeq \bigoplus_{ ext{simple} \atop ext{roots}} \mathbb{F}_q^{lpha}.$$

Hence a character $\chi: N^- \to \mathbb{C}$ determines a subset of simple reflections:

$$I = \{ s_{\alpha} \in S \mid \chi \neq 0 \text{ on } \mathbb{F}_q^{\alpha} \}.$$

Show that the orbit N^-xB/B supports a Whittaker vector if and only if $x \in {}^{I}W$ (the set of minimal coset representatives of $W_I \setminus W$).

Example 32.8. If χ is nondegenerate (i.e. I = S), then $\operatorname{Fun}(G/B, \mathbb{C})^{(N^{-}, \chi)} = \mathbb{C}$.

Categorifying Whittaker functions: Now let G, B, N, N^- , etc. be defined over $\overline{\mathbb{F}}_q$. Just as

 $\operatorname{Fun}(X)^{N^{-}}$ is categorified by $D_{N^{-}}(X)$,

we will see that

$$\operatorname{Fun}(X)^{(N^-,\chi)}$$
 is categorified by $D_{(N^-,\mathcal{L})}(X)$

But what is this category $D_{(N^-,\mathcal{L})}(X)$? Our first step is to find an appropriate way to categorify the character χ . We do this using the **Artin-Schreier sheaf**.

Before defining this sheaf, we give some motivation. Recall that

$$\mathbb{C}/\mathbb{Z} \xrightarrow{\sim} \mathbb{C}^{\times}$$

via $z \mapsto \exp(2\pi i z)$. The exponential map is the fundamental additive character of \mathbb{C} . Analogously,

$$\mathbb{A}^1/\mathbb{F}_p \xrightarrow{\sim} \mathbb{A}^1$$

via the Artin-Schreier map $a: x \mapsto x^p - x$. (The map $a: \mathbb{A}^1 \to \mathbb{A}^1$ has kernel \mathbb{F}_p , so provides the isomorphism above.) Moreover, we have

$$a_*(\mathbb{Q}_\ell)_{\mathbb{A}^1} \simeq \bigoplus_{\chi: \mathbb{F}_p \to \overline{\mathbb{Q}}_\ell^{\times}} \mathcal{L}_{\chi}.$$

The sheaves \mathcal{L}_{χ} are examples of *character sheaves* on \mathbb{A}^1 .

Define a map p by the following diagram:

$$N^{-} \xrightarrow{p} N^{-} / [N^{-}, N^{-}] \xrightarrow{\sim} \prod_{\alpha \text{ simple}} \mathbb{G}_{a}^{\alpha} \xrightarrow{\Sigma} \mathbb{G}_{a}$$

Fix a non-trivial additive character χ of \mathbb{A}^1 , and define

$$\mathcal{L} := p^* \mathcal{L}_{\chi}.$$

Exercise 32.9. The sheaves \mathcal{L}_{χ} and \mathcal{L} are "multiplicative"; i.e. we have $m^*\mathcal{L} \simeq \mathcal{L} \boxtimes \mathcal{L}$, where $m : N^- \times N^- \to N^-$ is multiplication. Multiplicative sheaves are "one-dimensional character sheaves" or "categorified characters".

Now let $N^- \circlearrowright X$. A (N^-, \mathcal{L}) -equivariant complex on X is a pair (\mathcal{F}, β) , where $\mathcal{F} \in D^b_c(X)$ and

$$\beta: a^* \mathcal{F} \xrightarrow{\sim} \mathcal{L} \boxtimes \mathcal{F}$$

is an isomorphism satisfying the usual cocycle condition. Let

 $D_{(N^{-},\mathcal{L})}(X) =$ category of (N^{-},\mathcal{L}) -equivariant complexes on X.

Morphisms in this category are morphisms in $D_c^b(X)$ which commute with β . We can define averaging functors

$$Av_{\mathcal{L}*}, Av_{\mathcal{L}!}: D^b_c(V) \to D^b_{(N^-,\mathcal{L})}(X)$$

as we did in the beginning of this lecture,

$$Av_{\mathcal{L}*} = m_*(\mathcal{L} \boxtimes (-))[\dim N^-], \text{ etc.}$$

Important fact: The forgetful functor

For :
$$D_{(N^-,\mathcal{L})}(X) \to D^b_c(X)$$

is fully faithful.

Exercise 32.10. Let

$$\begin{array}{ccc} \mathbb{A}^1 & \longrightarrow & \mathbb{P}^1 \\ \downarrow^a & & \downarrow \\ \mathbb{A}^1 & \longrightarrow & \mathbb{P}^1 \end{array}$$

- Show explicitly that the map $a: x \mapsto x^p x$ extends to \mathbb{P}^1 . (Find a formula!)
- Show that a breaks all the rules you know about coverings of Riemann surfaces. (The map a at ∞ is the simplest example of "wild ramification".)
- Show that $j_! \mathcal{L}_{\chi} \xrightarrow{\sim} j_* \mathcal{L}_{\chi}$.
- Show that $H^*(\mathbb{P}^1, j_!\mathcal{L}_{\chi}) = 0$. (We can interpret this as saying that \mathcal{L}_{χ} is a bit like our friend $\mathbb{Q}_{x!,y*}$ from earlier.)
- Show that $j_! \mathcal{L}_{\chi}$ is $(N^-, \mathcal{L}_{\chi})$ -equivariant.
- Show that $Av_{N*}(j_!\mathcal{L}_{\chi}) \simeq Av_{N!}(j_*\mathcal{L}_{\chi}) = T_s.$

32.4 Bird's eye view of the rest of the proof

We return to our usual setting: fix G, $\mathcal{F}\ell$ the corresponding affine flag variety, $I \subset G((t))$ Iwahori subgroup, I^- Iwahori for the opposite Borel, $I_u \subset I$, $I_u^- \subset I^-$ pro-unipotent radicals. Let

$$N_K = \begin{pmatrix} 1 & * & * \\ 0 & \ddots & * \\ 0 & 0 & 1 \end{pmatrix} \subset G((t)), \quad N_K^- = \begin{pmatrix} 1 & 0 & 0 \\ * & \ddots & 0 \\ * & * & 1 \end{pmatrix}.$$

We have seen that the antispherical module can be realized as

$$M_{\mathrm{asph}} \simeq H/\langle b_x \mid x \notin {}^fW \rangle.$$

Another realization is important in *p*-adic groups:

$$M_{\text{asph}} = \text{Fun}(\mathcal{F}\ell)^{(N_K^-,\psi)},$$

the "Whittaker vectors in the principal series". We might hope that on the level of categories, we would also have

$$\mathcal{M}_{asph} \simeq D_{(N_{\kappa}^{-},\mathcal{L})}(\mathcal{F}\ell).$$

Problem: N_K^- orbits on $\mathcal{F}\ell$ are " $\infty/2$ -dimensional" (i.e. they have neither finite dimension nor finite co-dimension). It is difficult to work with sheaves on an ∞ -dimensional space.

One solution: Use Drinfeld compactification. This approach is described in [FGV01]. This is not yet understood by Geordie.

Another solution: Use "Iwahori-Whittaker" or "baby Whittaker" techniques. The idea is to replace N_K^- by I_u^- .

Lemma 32.11. For nondegenerate characters ψ of N_K^- and χ of I_u^- ,

$$\operatorname{Fun}(\mathcal{F}\ell(\mathbb{F}_q))^{(N_K^-,\psi)} = \operatorname{Fun}(\mathcal{F}\ell(\mathbb{F}_q))^{(I_u^-,\chi)}.$$

The I_u^- -orbits on $\mathcal{F}\ell$ are just (opposite) Bruhat cells, so passing to I_u^- -orbits resolves our problem of infinte-dimensional orbits.

Define

 $P_{IW} \subset D_{IW} = D_{(I_u^-, \mathcal{L})}(\mathcal{F}\ell),$ "Iwahori-Whittaker sheaves".

The irreducible objects in P_{IW} are $\mathrm{IC}_{\chi}^{\mathcal{L}_{\chi}}$ for $x \in {}^{f}W$. Last week we constructed a functor

$$\widetilde{F}: D^b(\operatorname{Coh}^{G^\vee}(\widetilde{\mathcal{N}})) \to D_I$$

induced by

 $\begin{aligned} \operatorname{Coh}_{\operatorname{free}}^{G^{\vee}}(\widetilde{\mathcal{N}}) &\to P_{I} \\ V \otimes \mathcal{O} &\mapsto Z(\operatorname{Sat}(V)) \quad \text{central sheaf} \\ \mathcal{O}(\lambda) &\mapsto J_{\lambda} & \text{Wakimoto sheaf} \end{aligned}$

Let

 ${}^{f}P_{I} = P_{I} / \langle \mathrm{IC}_{x} \mid x \notin {}^{f}W \rangle.$

The main theorem is the following.


Theorem 32.12. \widetilde{F} induces an equivalence

$$D^b(\operatorname{Coh}^{G^{\vee}}(\widetilde{N})) \xrightarrow{\sim} D^b({}^fP_I).$$

The averaging functor

$$Av_{\mathcal{L}}: P_I \to D_{IW}$$

factors over ${}^{f}P_{I}$. Now



Theorem 32.13. 1. ${}^{f}P_{I} \simeq P_{IW}$ (reasonably easy consequence of part 3. below)

- 2. $D^b(P_{IW}) \simeq D_{IW}$ (same argument as Corollary 20.12)
- 3. $D^b(\operatorname{Coh}^{G^{\vee}}(\widetilde{\mathcal{N}})) \xrightarrow[F_{IW}]{\sim} D_{IW}$ (main difficulty)

As usual, fully-faithfulness of F_{IW} is the main issue.

Step 1: F_{IW} is faithful

A game with induction and restriction functors (using the G/B version of the fun calculation at the start of this lecture) implies that Av_{ψ} is faithful. Hence it is enough to show that \tilde{F} is faithful. The key idea is the following. We have

$$\widetilde{\mathcal{N}} \to \mathcal{N} \subset \mathfrak{g}^{\vee},$$

and $\mathcal{N}_{\text{reg}} \subset \mathcal{N}$ is open and dense. For $V, V' \in \operatorname{Coh}_{\text{free}}^{G^{\vee}}(\widetilde{\mathcal{N}})$,

$$\operatorname{Hom}(V, V') \to \operatorname{Hom}(V|_{\mathcal{N}_{\operatorname{reg}}}, V'|_{\mathcal{N}_{\operatorname{reg}}})$$

is injective. We have a diagram

What corresponds to (*)?

$$D_I \supset D_I^{\neq id} = \langle \mathrm{IC}_x \mid x \neq id \rangle_\Delta.$$

Using this, we can define a \otimes -category

]

$$D_I^{id} := D_I / D_I^{\neq id}.$$

Moreover, if P_I^{id} is the image of P_I in D_I^{id} , then P_I^{id} is a \otimes -category with one simple object. Hence we have

$$\operatorname{Rep} G^{\vee} \xrightarrow{Z} P_I \to P_I^{id}.$$

Using the central functor + derivation, Tannakian formalism (+ a bit of sauce) gives us the following diagram

Because res is faithful, \tilde{F} is faithful.

Remark 32.14. The above is an instance of an important theme in Bezrukavnikov's work: from a two-sided cell \underline{c} , Lusztig constructed a semi-simple abelian tensor category $J_{\underline{c}}$. Bezrukavnikov observed that Z provides a central functor $\operatorname{Rep} G^{\vee} \to J_{\underline{c}}$. Bezrukavnikov then uses Z to identify J_c . The theorem is the case when $\underline{c} = \{id\}$. Step 2: F_{IW} is full:

This can be shown using Beilinson's lemma and a generation argument. It reduces to checking

$$\operatorname{Ext}^{i}_{\operatorname{Coh}^{G^{\vee}}(\widetilde{\mathcal{N}})}(V \otimes \mathcal{O}, \mathcal{O}(\lambda)) \xrightarrow{F_{IW}} \operatorname{Ext}^{i}(Av_{\mathcal{L}}(\operatorname{Sat}(V)), Av_{\mathcal{L}}(J_{\lambda}))$$

for $V \in \operatorname{Rep} G^{\vee}$ and λ dominant. The left side is

$$\operatorname{Hom}_{G^{\vee}}(V, H^{i}(\widetilde{\mathcal{N}}, \mathcal{O}(\lambda))) = \begin{cases} 0 & \text{if } i \neq 0 \text{ (Frobenius splitting of } T^{*}\mathcal{B}), \\ V_{\lambda} & \text{if } i = 0. \end{cases}$$

We can chek that the right side has the same dimension and we are done.

Potential moral of the proof: It is "obvious" that P_{IW} is highest weight. Hence one can do calculations more more easily here than in ${}^{f}P_{I}$.

Lecture 33: Soergel bimodules, Soergel calculus, and BGK central sheaves

We will begin today's lecture by describing a more combinatorial approach to the Hecke category using Soergel bimodules.

33.1 The Hecke category and Soergel bimodules

We start with some motivation. Recall that given G a (split) finite group of Lie type and B a Borel subgroup, we can define a \mathbb{C} -algebra

$$H_q := (\operatorname{Fun}_{B \times B}(G, \mathbb{C}), *).$$

This is the first incarnation of the Hecke algebra. Iwahori showed that H_q admits a presentation which is "independent" of q, in that it only depends on the Coxeter system (W, S)determined by $B \subset G$. This leads us to define a $\mathbb{Z}[v^{\pm 1}]$ -algebra using this presentation, which allows us study "all q at once". This construction now makes sense for any Coxeter system (W, S). Can we do a similar thing for the Hecke category?

More specifically, recall from Lecture 24 that given an algebraic group G/\mathbb{C} , we defined a monoidal⁴¹ category

$$\mathcal{H}_{s.s.} := \left\langle \begin{array}{c} \text{additive category of} \\ \text{semi-simple complexes} \end{array} \right\rangle \subset D^b_{B \times B}(G, \mathbb{Q}).$$

From this category, we built our final incarnation of the Hecke category:

$$\mathcal{H} := K^b(\mathcal{H}_{s.s.}).$$

Questions: Can we present \mathcal{H} or $\mathcal{H}_{s.s.}$ by generators and relations? Does \mathcal{H} makes sense for any Coxeter system? We can also define \mathcal{H} using $G/\overline{\mathbb{F}}_q$ and étale sheaves, do we get equivalent categories after extending scalars?

To answer these questions, we need to introduce the notion of equivariant cohomology. Let K be a group acting on a space X. Define

$$H_K^*(X) := H^*(X \times_K EK),$$

where EK is a classifying space for K (i.e. a path-connected, contractible space with a free K-action). This construction is sometimes referred to as the "Borel construction". There is a natural map

$$X \times_K EK \to \text{pt} \times_K EK,$$

which gives $H_K^*(X)$ the structure of a graded $H_K^*(\text{pt})$ -module. Similarly, given a complex $\mathcal{F} \in D_K^b(X)$, $H_K^*(X, \mathcal{F})$ is a graded module over $H_K^*(\text{pt})$.

 $^{^{41}}$ by the Decomposition Theorem

Equivariant cohomology for tori: Let $T = \mathbb{C}^{\times}$. Then a classifying space for T is

$$ET = \mathbb{C}^{\infty} \setminus \{\infty\} = \lim_{n \to \infty} \mathbb{C}^n \setminus \{0\}.$$

Hence

$$H^*_{\mathbb{C}^{\times}}(\mathrm{pt}) = H^*(\mathrm{pt} \times_{\mathbb{C}^{\times}} \mathbb{C}^{\infty} \setminus \{0\}) = H^*(\mathbb{P}^{\infty}) = \mathbb{C}[x],$$

where x is in degree 2. Similarly, if $T \simeq (\mathbb{C}^{\times})^n$, then $ET = (\mathbb{C}^{\infty} \setminus \{0\})^n$, and

$$H_T^*(\mathrm{pt}) \simeq H^*((\mathbb{P}^\infty)^n) = \mathbb{C}[x_1, \dots, x_n].$$
(33.1)

Note that the isomorphism (33.1) depends on the isomorphism $T \simeq (\mathbb{C}^{\times})^n$. We can give a more canonical description as follows. Let $\chi : T \to \mathbb{C}^{\times}$ be a character. We have an associated \mathbb{C}^{\times} -bundle

$$L_{\chi} := \mathbb{C}^{\times} \times_{T,\chi} ET \to \mathrm{pt} \times_T ET$$

Borel showed that there is an isomorphism

$$\begin{array}{l} X \xrightarrow{\sim} H_T^2(\mathrm{pt}, \mathbb{Z}) \\ x \mapsto c_1(L_{\chi}), \end{array}$$

where X is the character lattice of T and $c_1(L_{\chi})$ is the first Chern class of L_{χ} . This leads to a canonical isomorphism

$$S^{\bullet}(X) \simeq H^*_T(\mathrm{pt}, \mathbb{C}),$$

where $S^{\bullet}(X)$ is the symmetric algebra of the character lattice. This is the **Borel isomorphism**.

Remark 33.1. Over \mathbb{C} , this can be further simplified. Given $\chi : T \to \mathbb{C}^{\times}$, we can differentiate to get a linear functional $d\chi : \text{Lie } T \to \mathbb{C}$. Then the Borel isomorphism becomes

$$\mathcal{O}(\operatorname{Lie} T) \simeq H_T^*(\mathrm{pt}, \mathbb{C}).$$

This is the version which we will use today.

Useful trick: If $K \subset G$ is a subgroup, then any model of EG is also a model of EK via restriction. Examples:

1. $T \subset B$: The map

 $T \backslash EB \rightarrow B \backslash EB$

is a $B/T \simeq \mathbb{C}^n$ -bundle, so $H_B^*(\text{pt}) \simeq H_T^*(\text{pt})$.

2. $T \subset G$: The map

 $T \backslash EG \xrightarrow{p} G \backslash EG$

induces a map $H^*_G(\text{pt}) \xrightarrow{p^*} H^*_T(\text{pt})$.

Theorem 33.2. p^* is injective, and the image is in $(\mathcal{O}(\operatorname{Lie} T))^W$.

By taking equivariant cohomology, we can give an algebraic description of the Hecke category.

$$\mathcal{H}_{s.s.} \subset D^b_{B \times B}(G) \xrightarrow{H^*_{B \times B}} H^*_{B \times B}(\text{pt})\text{-graded modules}$$
$$= H^*_{T \times T}(\text{pt})\text{-graded modules}$$
$$= R\text{-gbim},$$

where $R = \mathcal{O}(\text{Lie } T)$. The category R-gbim is a monoidal category via $- \otimes_R -$. In most situations, taking cohomology loses a lot of information, so the following theorem is especially remarkable.

Theorem 33.3. (Soergel) $H^*_{B\times B}$ is fully faithful and monoidal on $\mathcal{H}_{s.s.}$.

TT*

Hence,

$$\mathcal{H}_{s.s.} \hookrightarrow R$$
-gbim.

How can we describe the image? To start, observe that $\mathcal{H}_{s.s.}$ is generated by $IC_s = \mathbb{C}_{\overline{PsP}}[1]$ for $s \in S$ under $*, \oplus, \ominus, [1]$; i.e.

$$\mathcal{H}_{s.s.} = \langle IC_s \mid s \in S \rangle_{*,[\mathbb{Z}], \oplus, \ominus}$$

This gives us a way of describing the image:

$$\text{Sbim} := \langle B_s \mid s \in S \rangle_{*, \oplus, (1), \ominus},$$

where $B_s = H^*_{B \times B}(IC_s) = R \otimes_{R^s} R(1)$. (Exercise: prove this!) This is the category of **Soergel bimodules**. Unpacking definitions, we obtain equivalences of monoidal categories:

Remarkable consequence: To define $\mathcal{H}_{s.s.}$, we need G, B, hundreds of pages of sheaf theory, the decomposition theorem⁴², and more. To define Sbim, all we need is $W \circlearrowright$ Lie T and a bit of algebra!

- **Remark 33.4.** 1. This can be seen as a first step toward freeing \mathcal{H} from its concrete realization as a category of sheaves. By replacing Lie T with a reflection representation \mathfrak{h} of W, the definition makes sense for *any* Coxeter group. This led to the proof by Soergel and Elias-W. that the Kazhdan-Lusztig polynomials have non-negative coefficients.
 - 2. One of the main goals of this course is to approach Bezrukavnikov's equivalence:

 $\mathcal{H}^{\text{affine}} \simeq$ "coherent sheaves on Steinberg".

Soergel's theorem tells us that the Hecke category always has a coherent description in terms of Soergel bimodules. One way of understanding Bezrukavnikov's theorem is that "Soergel bimodules have another name" in the affine setting.

⁴²can be avoided via the theory of parity sheaves

3. One can see Soergel bimodules as "half way" towards a generators and relations description. A generators and relations description was obtained by Elias-W., following Elias-Khovanov, Libedinsky, and Elias. See [EMTW20]. An interesting recent take has been given by Abe in [Abe19].

33.2 BGK central sheaves for $G = SL_2$

Goal: Describe $Z : \operatorname{Rep} \operatorname{SL}_2 \to \mathcal{H}_{ext}$ by "generators and relations".

We start with the left hand side, $\operatorname{Rep} \operatorname{SL}_2$, which we can describe via the **Temperly-Lieb** category TL:

- Objects: $\mathbb{Z}_{\geq 0}$
- Morphisms: crossingless matchings, up to isotopy, () = -2

Let $V = \mathbb{C}^2$ be the natural representation of SL_2 .

Theorem 33.5. $\langle V^{\otimes n} \rangle \simeq TL$, hence $(TL)_{\oplus,\ominus} \simeq \operatorname{Rep} \operatorname{SL}_2$.

Remark 33.6. This description of Rep SL₂ is very useful. For example, an immediate consequence of the theorem is that giving a functor Rep SL₂ \rightarrow (\mathcal{C}, \otimes) with \mathcal{C} additive Karoubian is the same as giving a self-dual object $X \in \mathcal{C}$ of dimension 2.


Giving a generators and relations description of the right hand side \mathcal{H}_{ext} is more complicated. We'll start with the unextended case and describe \mathcal{H} , the Hecke category corresponding to the affine Weyl group $W = \langle s, t \rangle$. (Here s is the finite simple reflection.) Fix a "realisation" of W; that is, fix the data of

- a complex C-vector space \mathfrak{h} ,
- vectors $\alpha_s, \alpha_t \in \mathfrak{h}^*$, and
- vectors $\alpha_s^{\vee}, \alpha_t^{\vee} \in \mathfrak{h}$, such that the pairing between the α and the α^{\vee} is given by the Cartan matrix

$$\begin{pmatrix} 2 & -2 \\ -2 & 2 \end{pmatrix}.$$

With such a realisation, W acts via automorphisms of \mathfrak{h} according to the usual formulas. Most important choices of realisations:

1.
$$\mathfrak{h} = \mathbb{C}, \, \alpha_s = -\alpha_t, \, \alpha_s^{\vee} = -\alpha_t^{\vee}.$$
 (Realises canonical quotient $W \twoheadrightarrow W_f$.)




Remark 33.7. In geometry, \mathfrak{h} arises from I, whereas \mathfrak{h}_{loop} arises from $I \ltimes \mathbb{C}^{\times}$ (loop rotation). Now we can describe a diagrammatic version of the Hecke category. Let

 $\mathcal{H}_{BS} =$ monoidal category generated by B_s and B_t .


Morphisms in \mathcal{H}_{BS} are isotopy classes of diagrams generated by



subject to a collection of relations. The most important of these relations are:



along with the same relations (and generators) obtained by swapping red \leftrightarrow blue and $s \leftrightarrow t$. Example 33.8. The diagram



is a morphism $B_s B_t B_s B_s \rightarrow B_t B_s B_t$.

From here, we can obtain a version of $\mathcal{H}_{s.s.}$:

$$\mathcal{H}_{BS} \longrightarrow \begin{array}{cc} \text{formally add} & \text{add formal} \\ \text{shifts, keep} & \text{sums and} \\ \text{only degree} & \text{take Karoubi} \\ \text{zero maps} & \text{envelope} \end{array} =: \mathcal{H}_{s.s.}^{\text{diag}}$$


Theorem 33.9. (Elias-W.) Using \mathfrak{h}_{loop} ,

$$\mathcal{H}_{s.s.}^{\mathrm{diag}} \simeq \mathrm{Sbim} \simeq \mathcal{H}_{s.s.}$$

As a corollary to this theorem, we obtain a rather explicit version of the Hecke category \mathcal{H} where we can do calculations.

Corollary 33.10. $K^b(\mathcal{H}_{s,s}^{\text{diag}}) \simeq \mathcal{H}.$

How do we get \mathcal{H}_{ext} ? We can extend this constuction to \mathcal{H}_{ext} by adding



Now we can translate what we've done in previous lectures to the language of diagrammatic Soergel bimodules.

33.3 Braid group categorification

In Soergel bimodule world,

 $\begin{array}{lll} \Delta_s\longmapsto & 0\to \underline{\underline{B_s}} \xrightarrow{\uparrow} R(1) \to 0 \\ \Delta_w\longmapsto & \text{``Rouquier complex'' with many intriguing properties} \\ \Delta_\omega\longmapsto & 0\to\underline{\underline{\omega}}\to 0 \end{array}$

Here a double underline indicates degree zero. Recall that for the affine simple reflection t, we have

translation by $\varpi = t_{\omega}$.

Hence the Wakimoto sheaves

$$J_{\varpi} = \underline{\underline{B}_t \omega} \xrightarrow{\dagger} \omega(1), \quad J_{-\varpi} = \omega(-1) \xrightarrow{\dagger} \underline{\underline{B}_s \omega},$$

and

$$J_{2\varpi} = J_{\varpi}^2 = \underbrace{\underline{B_t \omega B_t \omega}}_{\omega B_t \omega} \to \underbrace{\begin{array}{c} B_t \omega \omega(1) \\ \oplus \\ \omega B_t \omega \end{array}}_{\omega B_t \omega} \to \mathbb{1}(2) \simeq \underbrace{\underline{B_t B_s}}_{\omega B_s} \xrightarrow{(|\mathbf{1},\mathbf{1}|)} \underbrace{\begin{array}{c} B_{ts} \omega(1) \\ \oplus \\ B_{st} \omega(1) \end{array}}_{\omega B_s \omega(1)} \xrightarrow{(\mathbf{1},\mathbf{1})} \mathbb{1}(2).$$

Exercise 33.11. Show that

$$J_{3\varpi} = \underbrace{\underline{B}_{tst}}_{B_{ts}\omega} \to \begin{array}{cc} B_{ts}\omega(1) & B_{s}\omega(2) \\ \oplus & \to & \oplus \\ B_{st}\omega(1) & B_{t}\omega(2) \end{array} \to B_{id}\omega$$

and

$$J_{-2\varpi} = \mathbb{1}(-2) \to \begin{array}{c} B_s(-1) \\ \oplus \\ B_t(-1) \end{array} \to \underline{\underline{B_s B_t}}.$$

Consider

$$\mathcal{F} = \mathbb{1}(-1) \xrightarrow{\begin{pmatrix} \mathbf{l} \\ \mathbf{l} \end{pmatrix}} \begin{array}{c} B_s \\ \oplus \\ B_t \end{array} \xrightarrow{(\uparrow,\uparrow)} \mathbb{1}(1).$$

Note that $\alpha^2 = \mathbf{1} \mathbf{+} \mathbf{1} = \begin{cases} \delta & \text{ in } \mathfrak{h}_{\text{loop}}, \\ 0 & \text{ in } \mathfrak{h}. \end{cases}$

We have

$$\mathcal{F}B_s \simeq B_s(-1) \rightarrow \begin{array}{ccc} B_sB_s & \oplus \\ \oplus \\ B_tB_s \end{array} \rightarrow B_s(1) \simeq B_s(-1) \rightarrow \begin{array}{ccc} B_s(-1) \\ \oplus \\ B_tB_s \end{array} \rightarrow B_s(1) \simeq B_tB_s \end{array}$$

Similarly,

$$\mathcal{F}B_t \simeq B_s B_t, \quad B_s \mathcal{F} \simeq B_s B_t, \text{ and } \quad B_t \mathcal{F} \simeq B_t B_s.$$

Exercise 33.12. Checek this.

Theorem 33.13. (Elias) $z_1 := \mathcal{F}\omega$ has a canonical central structure. Moreover, there exist maps


$$\mathbb{1} \to z_1^2, \quad z_1^2 \to \mathbb{1}$$

satisfying the Temperly-Lieb relations.

Hence we obtain


$$Z : \operatorname{Rep} \operatorname{SL}_2 \to \mathcal{H}_{ext}.$$

Example 33.14.



$$\simeq \mathbf{1}^{(-1)} \xrightarrow{B_{s}(-1)}_{\substack{\mathfrak{B}_{t}(-1)\\\mathfrak{B}_{t}(-1)}} \xrightarrow{B_{s}t}_{\substack{\mathfrak{B}_{s}t\\\mathfrak{B}_{t}(-1)}} \xrightarrow{\mathfrak{B}_{s}t}_{\substack{\mathfrak{B}_{s}t\\\mathfrak{B}_{t}(-1)}} \xrightarrow{\mathfrak{B}_{s}t}_{\substack{\mathfrak{B}_{s}(1)\\\mathfrak{B}_{t}(-1)}} \xrightarrow{\mathfrak{B}_{s}t}_{\substack{\mathfrak{B}_{s}(1)\\\mathfrak{B}_{t}(-1)}} \xrightarrow{\mathfrak{B}_{s}t}_{\substack{\mathfrak{B}_{s}(1)\\\mathfrak{B}_{t}(-1)}} \xrightarrow{\mathfrak{B}_{s}t}_{\substack{\mathfrak{B}_{s}(1)\\\mathfrak{B}_{t}(-1)}} \xrightarrow{\mathfrak{B}_{s}t}_{\substack{\mathfrak{B}_{s}(1)\\\mathfrak{B}_{t}(-1)}} \xrightarrow{\mathfrak{B}_{s}t}_{\substack{\mathfrak{B}_{s}(1)\\\mathfrak{B}_{t}(-1)}} \xrightarrow{\mathfrak{B}_{s}t}_{\substack{\mathfrak{B}_{s}(1)\\\mathfrak{B}_{t}(-1)}} \xrightarrow{\mathfrak{B}_{s}t}_{\substack{\mathfrak{B}_{s}(1)\\\mathfrak{B}_{t}(-1)}} \xrightarrow{\mathfrak{B}_{s}t}_{\substack{\mathfrak{B}_{s}(1)\\\mathfrak{B}_{t}(-1)}} \xrightarrow{\mathfrak{B}_{s}t}_{\substack{\mathfrak{B}_{s}(1)\\\mathfrak{B}_{s}(1)}} \xrightarrow{\mathfrak{B}_{s}t}_{\substack{\mathfrak{B}_{s}(1)}} \xrightarrow{\mathfrak{B}_{s}t}_{\substack{\mathfrak{B}_{s}t}} \xrightarrow{\mathfrak{B}_{s}t}_{\substack{\mathfrak{B}_{s}(1)}} \xrightarrow{\mathfrak{B}_{s}t}_{\substack{\mathfrak{B}_{s}(1)}} \xrightarrow{\mathfrak{B}_{s}t}_{\substack{\mathfrak{B}_{s}t}} \xrightarrow{\mathfrak{B}_{s}t}_{\substack{\mathfrak{B}_{s}t}} \xrightarrow{\mathfrak{B}_{s}t}_{\substack{\mathfrak{B}_{s}t}} \xrightarrow{\mathfrak{B}_{s}t}_{\substack{\mathfrak{B}_{s}t}} \xrightarrow{\mathfrak{B}_{s}t}_{\substack{\mathfrak{B}_{s}t}} \xrightarrow{\mathfrak{B}_{s}t}_{\substack{\mathfrak{B}$$

(compare $(\mathbb{C}^{n})^{\otimes L} \mathbb{C}^{3} \otimes \mathbb{C}^{\otimes}$ los Rep SL2).



Lecture 34: Bezrukavnikov's equivalence

This is the final lecture of the course!

34.1 Potted history of Koszul duality

A more thorough description of the material in this section can be found in [Wil18, §3]. Let \mathcal{O}_0 be the principal block of category \mathcal{O} for \mathfrak{g} , and $\Delta_x, \nabla_x, L_x, T_x, P_x, I_x$ the standard, costandard, simple, tilting, projective, and injective objects in \mathcal{O}_0 corresponding to $x \in W$. Let \mathcal{O}_0^{\vee} be the principal block of category \mathcal{O} for the Langlands dual Lie algebra \mathfrak{g}^{\vee} , and $\Delta_x^{\vee}, \nabla_x^{\vee}$, etc. the corresponding objects in \mathcal{O}_0^{\vee} .

Beilinson-Ginzburg ('86) conjectured that:

- 1. \mathcal{O}_0 admits a graded version $\widetilde{\mathcal{O}}_0$ with a "shift of grading functor" $\langle m \rangle$, and all "canonical" objects above admit lifts $\widetilde{\Delta}_x, \ldots, \widetilde{I}_x$.
- 2. There exists an equivalence of triangulated categories

$$\kappa: D^b(\widetilde{\mathcal{O}}_0) \xrightarrow{\sim} D^b(\widetilde{\mathcal{O}}_0^{\vee})$$

such that $\kappa \circ \langle -1 \rangle [1] \simeq \langle 1 \rangle \circ \kappa$, and κ sends

$$\widetilde{\Delta}_x \mapsto \widetilde{\Delta}_{x^{-1}w_0}^{\vee}, \quad \widetilde{L}_x \mapsto \widetilde{I}_{x^{-1}w_0}^{\vee}, \quad \widetilde{P}_x \mapsto \widetilde{L}_{x^{-1}w_0}^{\vee}$$

The conjecture was proved by Beilinson-Ginzburg-Soergel in [BGS96].

Exercise 34.1. Decategorify κ and deduce something remarkable about Kazhdan–Lusztig polynomials! (See the first chapter on Koszul duality chapter in [EMTW20].)

Beilinson-Ginzburg ('99): If we compose κ with the Radon transform $(*\Delta_{w_0})$ + inversion $(g \mapsto g^{-1})$ and pass to a geometric setting, we obtain a functor

$$\widetilde{\kappa}: D^{\mathrm{mix}}_{(B)}(G/B) \xrightarrow{\sim} D^{\mathrm{mix}}_{(B^{\vee})}(B^{\vee} \backslash G^{\vee})$$

is an equivalence with $\widetilde{\kappa} \circ \langle -1 \rangle [1] \simeq \langle 1 \rangle \circ \widetilde{\kappa}$, and

$$\widetilde{IC}_x \mapsto \widetilde{T}_x^{\vee}, \quad \widetilde{\Delta}_x \mapsto \widetilde{\Delta}_x^{\vee}, \quad \widetilde{\nabla}_x \mapsto \widetilde{\nabla}_x^{\vee}, \quad \widetilde{T}_x^{\vee} \mapsto \widetilde{IC}_x^{\vee}.$$

After seeing this more symmetric version, one starts to start to dream a little.

Dream:

- 1. It should work for any Kac-Moody group. (We've removed w_0 from the formulas!)
- 2. It should be "monoidal"; i.e. we should roughly have

 $(\text{semisimple}, *) \simeq (\text{tiltings}, *^{\vee}).$

Unfortunately, this is very much in dreamland.

Basic problem: $T_x \in D_{(B)}(G/B)$ lifts to $D_B(G/B)$ if and only if x = id.

Remark 34.2. We have seen this a number of times for \mathbb{P}^1 (e.g. Example 30.7): T_s does not have an equivariant structure.

One way we could try to resolve this problem is to work in $D_U(G/B)$ instead, because T_x does lift to this category. However, in $D_U(G/B)$ we no longer have convolution. To fix this, we can go one step further and work in $D_U(G/U)$, which gives us access to T_x and convolution. But unfortunately, this introduces new problems.

Second problem: * is not exact on $D_U(G/U)$.

To illustrate this problem, we can consider a toy example

34.2 A toy example

Let $G = S^1$ (if you like Lie groups) or \mathbb{G}_m (if you prefer the algebraic setting).

Exercise 34.3. Given $\mathcal{L}_1, \mathcal{L}_2 \in \operatorname{Loc}(S^1)$,

$$(\mathcal{L}_1 * \mathcal{L}_2)_1 \simeq H^*(\mathcal{L}_1 \otimes \mathcal{L}_2^*).$$

The subscript 1 denotes the stalk at the identity. As an additional exercise, describe the monodromy of $\mathcal{L}_1 * \mathcal{L}_2$ in terms of the monodromy on \mathcal{L}_1 and \mathcal{L}_2 .

The basic issue is the following. We have seen (Example 17.6) that for a local system \mathcal{L}/S^1 given by monodromy $\mu \circlearrowright V$,

$$H^0(S^1, \mathcal{L}) = V^{\mu}$$
 and $H^1(S^1, \mathcal{L}) = V_{\mu}$.

For V finite dimensional, $V^{\mu} \neq 0 \iff V_{\mu} \neq 0$. Hence for any finite-dimensional local system \mathcal{L} on S^1 , either $H^*(S^1, \mathcal{L}) = 0$ or $H^*(S^1, \mathcal{L}) \neq 0$ in both degree 0 and degree 1. This implies (by Exercise 34.3) that * is either zero or *not exact*.

Solution: Infinite-dimensional local systems! Take \mathcal{L} to be the local system corresponding to

$$\mathbb{C}[[t]] \circlearrowleft \mu = 1 + t.$$

Note that $\mathbb{C}[[t]]^{\mu} = 0$, $\mathbb{C}[[t]]_{\mu} = \mathbb{C}[[t]]/(1-\mu) = \mathbb{C}$.

Remark 34.4. Roughly one can think of \mathcal{L} as an infinite-dimensional Jordan block

which has one-dimensional coinvariants and no invariants.


Exercise 34.5. $m_!(\mathcal{L} \boxtimes \mathcal{L})[1] \simeq \mathcal{L}$, hence we have an exact *-product!

34.3 Torus monodromic sheaves á la Bezrukavnikov-Yun

The ideas we encountered in the toy example above were formalized by Bezrukavnikov-Yun in [BY13] (c.f. Beilinson's 1983 ICM talk). Let

$$D(B \searrow G \swarrow B) := \langle q^* D(B \backslash G/B) \rangle$$
$$= \langle p^* D(U \backslash G/B) \rangle,$$

where



Example 34.6. 1. If $G = \mathbb{C}^{\times}$,

$$D(B \searrow G \swarrow B) = \langle p^* D(\text{pt}) \rangle$$

= full subcategory of $D^b_c(\mathbb{C}^{\times})$ generated by $\mathbb{Q}_{\mathbb{C}}$
= $\{\mathcal{F} \mid \mathcal{H}^i(\mathcal{F}) \text{ local systems with unipotent monodromy} \}.$


2. If $G = \operatorname{SL}_2$, $G/U \simeq \mathbb{C}^2 \setminus \{0\}$, so $D(U \setminus G/B) \simeq (\operatorname{IC}_0, \operatorname{IC}_{\mathbb{P}^1})$. Hence

 $D(B \searrow G \swarrow B) = \langle \mathbb{Q}_{\mathbb{C}^{\times}}, \mathbb{Q}_{\mathbb{C}^{2} \setminus \{0\}} \rangle \subset D_{U}(G/U).$

By replacing B with U, we get a convolution product

$${}^{U}_{*}: D(B \searrow G \swarrow B) \times D(B \searrow G \swarrow B) \to D(B \searrow G \swarrow B)$$

from the diagram



using the same definition as previously, except that $m : G \times_U G/U$ is not proper, so we need to make a choice of which pushforward m_* or $m_!$ to use. We choose $m_!$.

Very technical point: (cf. appendix by Yun to [BY13])

1. One can complete $D(B \searrow G \swarrow B)$ to the "free monodromic completion" $\widehat{D}(B \searrow G \swarrow B)$ to allow pro-local systems (like $\mathbb{C}[[t]] \bigcirc$ from earlier) along the fibres of p.

2. The convolution product $\overset{U}{*}$ extends to $\widehat{D}(B \searrow G \swarrow B)$. Moreover, we can define a category

$$\widehat{\mathrm{Tilt}} \subset \widehat{D}(B \searrow G \swarrow B)$$

of "free monodromic tilting sheaves," and $\overset{U}{*}$ is exact on $\widehat{\text{Tilt}}$.

Theorem 34.7. [BY13]

$$(\widehat{\mathrm{Tilt}}, \overset{U}{*}) \simeq (\widehat{\mathrm{SBim}}, \underset{\widehat{R}}{\otimes})$$

The category on the right side of this equivalence consists of Soergel bimodules for $R = \mathcal{O}(\text{Lie }T^{\vee}) = \mathcal{O}((LieT)^*)$ (note the dual!), completed along the grading.

Idea of proof: Suppose that $\bigvee_{f}^{X} p$ is a *T*-torsor. Define a category Y $D^{b}(X \swarrow T) = \langle p^{*}D_{c}^{b}(Y) \rangle_{\Delta} \subset D^{b}(X)$

of "unipotently monodromic sheaves". The fundamental observation of Verdier is that every $\mathcal{F} \in D^b(X \swarrow T)$ has a canonical monodromy action of $\pi_1(T)$. The action is unipotent. In other words, $D^b(X \swarrow T)$ is linear over $\mathbb{Q}[\pi_1(T)]$. Taking logs, we obtain that $D^b(X \swarrow T)$ is \widehat{R} -linear.

Theorem 34.8. [BY13] The functor $\mathbb{V} : \widehat{\text{Tilt}} \to \widehat{\text{SBim}}$ is monoidal and fully-faithful.

Adding weights, one gets monoidal equivalences

$$(\widehat{\mathrm{Tilt}}^{\mathrm{mixed}}, *) \simeq (\mathrm{SBim}, \bigotimes_R) \simeq (\mathcal{H}_{s.s.}^{G^{\vee}}, *),$$

and hence taking homotopy categories,

$$D^{\mathrm{mixed}}(B \searrow G \swarrow B) \simeq \mathcal{H} \simeq D^{\mathrm{mixed}}(B^{\vee} \backslash G^{\vee} / B^{\vee}).$$

This achieves the dream of a "monoidal Koszul duality".

A beautiful feature: Let $\pi: U \setminus G/U \to U \setminus G/B$ be projection. Then

$$\pi_*(T_x)[\operatorname{rank} T] = T_x \in D_U(G/B).$$

In other words, taking coinvariants of free monodromic tilting sheaves gives indecomposible tilting sheaves downstairs.

Remark 34.9. With enough⁴³ homological algebra, one can do all of the above algebraically (and also mod p), c.f. [AMRW19]. Some of the constructions there are very adhoc. Recent foundational work of Hogencamp and Makisumi explains much more satisfactorily what is going on.

⁴³approximately 250 pages

34.4 Bezrukavnikov's equivalence

The constructible side: Associated to our choice $T \subset B \subset G$, we have the familiar zoo of characters:

- the pro-unipotent radical $I^0 \subset I \subset G((t))$,
- the extended affine flag variety (aka "affine base affine space") $\widetilde{\mathcal{F}\ell} = G((t))/I^0$,
- the affine flag variety $\mathcal{F}\ell = G((t))/I$,
- the affine Grassmannian $\mathcal{G}r = G((t))/G[[t]],$
- projections

$$\widetilde{\mathcal{F}\ell} \xrightarrow{T\text{-torsor}} \mathcal{F}\ell \to \mathcal{G}r,$$

• derived categories

 $D_{II} := D(I \setminus G((t))/I),$ $D_{I^0I} := D(I^0 \setminus G((t))/I) \rightsquigarrow \widehat{D}_{I^0I}, \quad \text{free monodromic completion}$ $D_{I^0I^0} := D(I^0 \searrow G((t)) \swarrow I^0) \rightsquigarrow \widehat{D}_{I^0I^0}, \quad \text{free monodromic completion}$

• and actions

$$\widehat{D}_{I^0I^0} \circlearrowright \widehat{D}_{I^0I} \circlearrowright \widehat{D}_{II} D_{I^0I^0} \circlearrowright D_{I^0I} \circlearrowright D_{II}.$$

The coherent side: Associated to G^{\vee} , we have

$$\begin{array}{ccc} \widetilde{\mathcal{N}}^{\vee} & \longrightarrow & \widetilde{\mathfrak{g}}^{\vee} \\ \downarrow & & \downarrow \\ \mathcal{N}^{\vee} & \longrightarrow & \mathfrak{g}^{\vee}, \end{array}$$

where the left vertical arrow is the Springer resolution and the right vertical arrow is the Grothendieck-Springer resolution sending $(x, \mathfrak{b}) \in \tilde{\mathfrak{g}}^{\vee} = \{(x, \mathfrak{b}) \subset \mathfrak{g}^{\vee} \times \mathcal{B}^{\vee} \mid x \in \mathfrak{b}\}$ to $x \in \mathfrak{g}^{\vee}$.

Recall: Given $X \xrightarrow{f} Y$ proper, one can construct a convolution structure on $(D^b(X \times_Y X), *)$ (e.g. if X is a finite set and Y is a point, then $D^b(X \times_Y X) \simeq X \times X$ matrices of chain complexes of vector spaces). Earlier we used this construction to give a coherent realization of the affine Hecke algebra via the Kazhdan–Lusztig isomorphism:

$$H^{\mathrm{aff}} \simeq K^{G^{\vee} \times \mathbb{C}^{\times}} (\widetilde{\mathcal{N}}^{\vee} \times_{\mathcal{N}} \widetilde{\mathcal{N}}^{\vee}).$$

We need a derived version of this (which will mostly disappear in a moment).

Recall that $X \times_Y X$ is constructed by gluing $\operatorname{Spec}(B \otimes_A B)$ for an affine cover $\operatorname{Spec} B \to$ Spec A of f. Similarly, we can construct $X \times_Y^L X$ by gluing the dg-schemes $\operatorname{Spec}(B \otimes_A^L B)$ for an affine cover $\operatorname{Spec} B \to \operatorname{Spec} A$ of f.

Key points:

- 1. $D^b(X \stackrel{L}{\times_Y} X)$ is still triangulated and monoidal.
- 2. If $\operatorname{Tor}_{>0}^{\mathcal{O}_Y}(\mathcal{O}_X, \mathcal{O}_X) = 0$, then $X \stackrel{L}{\times_Y} X = X \times_Y X$.

From this we define our coherent characters:

- $St_{\mathfrak{g}^{\vee}} := \widetilde{\mathfrak{g}}^{\vee} \overset{L}{\times}_{\mathfrak{g}^{\vee}} \widetilde{\mathfrak{g}}^{\vee} = \widetilde{\mathfrak{g}}^{\vee} \times_{\mathfrak{g}^{\vee}} \widetilde{\mathfrak{g}}^{\vee},$
- $St_{\mathcal{N}^{\vee}} := \widetilde{\mathfrak{g}}^{\vee} \overset{L}{\times}_{\mathfrak{g}^{\vee}} \widetilde{\mathcal{N}}^{\vee} = \widetilde{\mathfrak{g}}^{\vee} \times_{\mathfrak{g}^{\vee}} \widetilde{\mathcal{N}}^{\vee}$, and
- $St^L := \widetilde{\mathcal{N}}^{\vee} \overset{L}{\times}_{\mathfrak{g}^{\vee}} \widetilde{\mathcal{N}}^{\vee} \qquad (\neq \widetilde{\mathcal{N}}^{\vee} \times_{\mathfrak{g}^{\vee}} \widetilde{\mathcal{N}}^{\vee} = \widetilde{\mathcal{N}} \times_{\mathcal{N}^{\vee}} \widetilde{\mathcal{N}}^{\vee}).$

Theorem 34.10. (Bezrukavnikov) There exist vertical equivalences

$$\begin{array}{ccccc} D^b \operatorname{Coh}^{G^{\vee}}(St_{\mathfrak{g}^{\vee}}) & \circlearrowright & D^b \operatorname{Coh}^{G^{\vee}}(St_{\mathcal{N}^{\vee}}) & \circlearrowleft & D^b \operatorname{Coh}^{G^{\vee}}(St^L) \\ & & & & & \\ & & & & & \\ & & & & & \\ D_{I^0I^0} & \circlearrowright & D_{I^0I} & \circlearrowright & D_{II} \end{array}$$

making the diagram of module categories commute.

Remarks on the proof:

- 1. Equivalences (b) and (c) and compatibility are reasonably straightforward consequences of (a), which is where Bezrukavnikov spends most of the paper.
- 2. It is technically convenient to instead prove

$$\widehat{D}_{I^0I^0} \simeq D(\operatorname{Coh}^{G^{\vee}}(\widehat{St}_{\mathfrak{g}^{\vee}})),$$

where \widehat{St} denotes the formal completion of $St_{\mathfrak{g}^{\vee}}$ along the preimage of \mathcal{N}^{\vee} .

3. The proof relies heavily on ideas in [AB09].

The theorem has several remarkable consequences, but exploring them will have to wait until the next course!

References

- [AB09] S. Arkhipov and R. Bezrukavnikov. Perverse sheaves on affine flags and Langlands dual group. *Israel Journal of Mathematics*, 170(1):135, 2009.
- [Abe19] Noriyuki Abe. On Soergel bimodules. *arXiv:1901.02336*, 2019.
- [AMRW19] Pramod Achar, Shotaro Makisumi, Simon Riche, and Geordie Williamson. Koszul duality for Kac–Moody groups and characters of tilting modules. *Journal* of the American Mathematical Society, 32(1):261–310, 2019.
- [Art24] Emil Artin. Über eine neue art von L-reihen. In Abhandlungen aus dem Mathematischen Seminar der Universität Hamburg, volume 3, pages 89–108. Springer, 1924.
- [Bei87] Alexander A Beilinson. On the derived category of perverse sheaves. In *K-theory, Arithmetic and Geometry*, pages 27–41. Springer, 1987.
- [Bez04] Roman and Bezrukavnikov. On tensor categories attached to cells in affine weyl groups. In *Representation theory of algebraic groups and quantum groups*, pages 69–90. Mathematical Society of Japan, 2004.
- [Bez16] Roman Bezrukavnikov. On two geometric realizations of an affine Hecke algebra. *Publications mathématiques de l'IHÉS*, 123(1):1–67, 2016.
- [BGS96] Alexander Beilinson, Victor Ginzburg, and Wolfgang Soergel. Koszul duality patterns in representation theory. *Journal of the American Mathematical Society*, 9(2):473–527, 1996.
- [Bum98] Daniel Bump. Automorphic Forms and Representations, volume 55. Cambridge University Press, 1998.
- [Buz17] Kevin Buzzard. Automorphic forms and the Langlands program, MSRI summer school. Available at https://www.youtube.com/playlist?list= PLhsb6tmzSpiysoRR0bZozub-MM0k3mdFR, 2017.
- [BY13] Roman Bezrukavnikov and Zhiwei Yun. On koszul duality for Kac-Moody groups. *Representation Theory of the American Mathematical Society*, 17(1):1–98, 2013.
- [CF67] JWS Cassels and A Fröhlich. *Algebraic Number Theory*. Thompson Book Company, Inc., 1967.
- [CG09] Neil Chriss and Victor Ginzburg. *Representation theory and complex geometry*. Springer Science & Business Media, 2009.
- [Clo06] L Clozel. The Sato-Tate conjecture. Current Developments in Mathematics, 2006(1):1–34, 2006.

- [Con01] Keith Conrad. History of class field theory. This unpublished essay is available online as a PDF file at www.math.uconn.edu/~kconrad/blurbs/gradnumthy/cfthistory.pdf, 2001.
- [DBB83] Pierre Deligne, Alexander A Beilinson, and Joseph Bernstein. Faisceaux pervers. Astérisque, 100, 1983.
- [Del97] Pierre Deligne. Action du groupe des tresses sur une catégorie. *Inventiones mathematicae*, 128(1):159–175, 1997.
- [DGNO10] Vladimir Drinfeld, Shlomo Gelaki, Dmitri Nikshych, and Victor Ostrik. On braided fusion categories i. *Selecta Mathematica*, 16(1):1–119, 2010.
- [EMTW20] Ben Elias, Shotaro Makisumi, Ulrich Thiel, and Geordie Williamson. Introduction to Soergel bimdules. Springer International Publishing, 2020.
- [EO03] Pavel Etingof and Viktor Ostrik. Module categories over representations of $_q(2)$ and graphs. arXiv preprint math/0302130, 2003.
- [Far18] Laurent Fargues. La courbe. In *Proc. Int. Cong. of Math*, volume 1, pages 261–290. World Scientific, 2018.
- [FGV01] Edward Frenkel, Dennis Gaitsgory, and Kari Vilonen. Whittaker patterns in the geometry of moduli spaces of bundles on curves. *Annals of Mathematics*, pages 699–748, 2001.
- [FGV02] Edward Frenkel, Dennis Gaitsgory, and Kari Vilonen. On the geometric Langlands conjecture. Journal of the American Mathematical Society, 15(2):367–417, 2002.
- [Gai05] Dennis Gaitsgory. The notion of category over an algebraic stack. arXiv:math/0507192 [math.AG], 2005.
- [Gro11] Benedict Gross. Representation Theory and Number Theory, Eilenberg lectures at Columbia University. Available at https://www.youtube.com/playlist? list=PL5E0D6DC4BCD8309D, 2011.
- [Kac90] Victor G Kac. Infinite-dimensional Lie algebras. Cambridge University Press, 1990.
- [KL87] David Kazhdan and George Lusztig. Proof of the Deligne-Langlands conjecture for Hecke algebras. *Inventiones mathematicae*, 87(1):153–215, 1987.
- [Kle93] Felix Klein. Vorlesungen über das Ikosaeder und die Auflösung der Gleichungen vom fünften Grade. Birkhäuser Verlag, Basel; B. G. Teubner, Stuttgart, 1993.
 Reprint of the 1884 original, Edited, with an introduction and commentary by Peter Slodowy.

[Kri05] Martin H. Krieger. A 1940 letter of André Weil on analogy in mathematics. Notices Amer. Math. Soc., 52(3):334-341, 2005. Excerpted from it Doing mathematics [World Scientific Publishing Co., Inc., River Edge, NJ, 2003; MR1961400]. [Lan90] R. P. Langlands. Representation theory: its rise and its role in number theory. In Proceedings of the Gibbs Symposium (New Haven, CT, 1989), pages 181–210. Amer. Math. Soc., Providence, RI, 1990. Jacob Lurie. Derived algebraic geometry ii: Noncommutative algebra. arXiv [Lur07] preprint math/0702299, 2007. [Man14] Yuri I. Manin. Forgotten Motives: the Varieties of Scientific Experience, 2014. [Mil97] James S Milne. Class Field Theory. Lecture notes available at https://www. jmilne.org/math/CourseNotes/cft.html, 1997. [Miy11] Katsuya Miyake. Takagi's class field theory: From where? and to where? RIMS Kôkyûroku Bessatsu, B25:125–160, 2011. [Mor18] Sophie Morel. Beilinson's construction of nearby cycles and gluing. *preprint*, available at https://web.math.princeton.edu/~smorel/gluing.pdf, 2018.[MS16] Barry Mazur and William Stein. Prime Numbers and the Riemann Hypothesis. Cambridge University Press, 2016. [Ost03]Victor Ostrik. Module categories, weak Hopf algebras and modular invariants. Transformation Groups, 8(2):177–206, 2003. [Sch17] Peter Scholze. p-adic geometry. ICM Lecture, arXiv:1712.03708, 2017. [Ser 55]Jean-Pierre Serre. Faisceaux algébriques cohérents. Annals of Mathematics, pages 197–278, 1955. [Ser79] Jean-Pierre Serre. Local fields, volume 67 of Graduate Texts in Mathematics. Springer-Verlag, New York-Berlin, 1979. Translated from the French by Marvin Jay Greenberg. [SVW18] Wolfgang Soergel, Rahbar Virk, and Matthias Wendt. Equivariant motives and geometric representation theory. (with an appendix by f. hörmann and m. wendt). arXiv:1809.05480, 2018. [SW18] Wolfgang Soergel and Matthias Wendt. Perverse motives and graded derived category \mathcal{O} . Journal of the Institute of Mathematics of Jussieu, 17(2):347–395, 2018.

- [Tat79] J. Tate. Number theoretic background. In Automorphic forms, representations and L-functions (Proc. Sympos. Pure Math., Oregon State Univ., Corvallis, Ore., 1977), Part 2, Proc. Sympos. Pure Math., XXXIII, pages 3–26. Amer. Math. Soc., Providence, R.I., 1979.
- [TT90] Robert W Thomason and Thomas Trobaugh. Higher algebraic K-theory of schemes and of derived categories. In *The Grothendieck festschrift*, pages 247– 435. Springer, 1990.
- [Wey16] Hermann Weyl. Über die gleichverteilung von zahlen mod. eins. *Mathematische Annalen*, 77(3):313–352, 1916.
- [Wil18] Geordie Williamson. Parity sheaves and the Hecke category. *Proc. Int. Cong.* of Math., Rio de Janeiro, Vol. 1:979 – 1016, 2018.
- [You16] Alexander Youcis. Weil-Deligne representations and p-adic Hodge theory: motivation. Available at: https://ayoucis.files.wordpress.com/2016/11/ weil-deligne-representations-2.pdf, 2016.