

Singular limit in Hopf bifurcation for doubly diffusive convection equations II: bifurcation and stability

Chun-Hsiung Hsia¹, Yoshiyuki Kagei², Takaaki Nishida³
and Yuka Teramoto⁴

Abstract

A singular perturbation problem from the artificial compressible system to the incompressible system is considered for a doubly diffusive convection when a Hopf bifurcation from the motionless state occurs in the incompressible system. It is proved that the Hopf bifurcation also occurs in the artificial compressible system for small singular perturbation parameter, called the artificial Mach number. The time periodic solution branch of the artificial compressible system is shown to converge to the corresponding bifurcating branch of the incompressible system in the singular limit of vanishing artificial Mach number.

1 Introduction

This paper studies a singular limit problem for Hopf bifurcation in the artificial compressible system for thermal convection equations in the presence of the diffusion of solute concentration. The system of equations under consideration is written as

$$\varepsilon^2 \partial_t \phi + \operatorname{div} \mathbf{w} = 0, \quad (1.1)$$

$$\partial_t \mathbf{w} - \operatorname{Pr} \Delta \mathbf{w} + \operatorname{Pr} \nabla \phi - \operatorname{Pr} \mathcal{R}_1 \theta \mathbf{e}_2 + \operatorname{Pr} \mathcal{R}_2 \psi \mathbf{e}_2 + \mathbf{w} \cdot \nabla \mathbf{w} = \mathbf{0}, \quad (1.2)$$

¹Department of Mathematics, National Taiwan University, Taipei 10617, Taiwan

²Department of Mathematics, Tokyo Institute of Technology, Tokyo 152-8551, Japan

³Department of Advanced Mathematical Sciences, Kyoto University, Kyoto 606-8317, Japan

⁴Research Institute for Interdisciplinary Science, Okayama University, Okayama 700-8530, Japan

$$\partial_t \theta - \Delta \theta - \mathcal{R}_1 \mathbf{w} \cdot \mathbf{e}_2 + \mathbf{w} \cdot \nabla \theta = 0, \quad (1.3)$$

$$\partial_t \psi - d \Delta \psi - \mathcal{R}_2 \mathbf{w} \cdot \mathbf{e}_2 + \mathbf{w} \cdot \nabla \psi = 0. \quad (1.4)$$

Here $\phi = \phi(x, t)$, $\mathbf{w} = {}^\top(w^1(x, t), w^2(x, t))$, $\theta = \theta(x, t)$ and $\psi = \psi(x, t)$ denote the perturbation of the pressure, velocity field, temperature and solute concentration, respectively, at position $x \in \mathbb{R}^2$ and time $t \in \mathbb{R}$, from their values of the motionless state in a thermal convection; \mathcal{R}_1 , \mathcal{R}_2 , Pr and d are non-dimensional positive parameters; \mathcal{R}_1^2 and \mathcal{R}_2^2 are the thermal and salinity Rayleigh numbers, respectively; Pr and d are called the Prandtl and Lewis numbers, respectively. The parameter $\varepsilon > 0$ is called the artificial Mach number.

The system (1.1)–(1.4) is considered on the two dimensional infinite layer $\{x = (x_1, x_2) \in \mathbb{R}^2; x_1 \in \mathbb{R}, 0 < x_2 < 1\}$ under the following boundary condition on the boundary $\{x_2 = 0, 1\}$:

$$\frac{\partial w^1}{\partial x_2} = w^2 = \theta = \psi = 0 \text{ on } \{x_2 = 0, 1\} \quad (1.5)$$

and the periodic boundary condition in x_1 for ϕ , \mathbf{w} , θ and ψ with period $\frac{2\pi}{\alpha}$, where $\alpha > 0$. The system (1.1)–(1.4) is thus considered in the domain

$$\Omega = \mathbb{T}_{\frac{2\pi}{\alpha}} \times (0, 1)$$

under the boundary condition (1.5). Here \mathbb{T}_a denotes $\mathbb{T}_a = \mathbb{R}/a\mathbb{Z}$. We write $u = {}^\top(\phi, \mathbf{w}, \theta, \psi)$ for the unknown functions of (1.1)–(1.4). Here and in what follows, the superscript ${}^\top \cdot$ denotes the transposition.

When $\varepsilon = 0$ one obtains the thermal convection equations for a viscous incompressible fluid under the Oberbeck-Boussinesq approximation. A rough explanation of the structure of (1.1)–(1.4) with $\varepsilon = 0$ is as follows. The terms including \mathcal{R}_1 have a symmetric structure which may cause instabilities against the dissipativity by the Laplacians when \mathcal{R}_1 increases; the terms including \mathcal{R}_2 have a skew-symmetric structure which may cause oscillatory behavior. Indeed, when $\varepsilon = 0$, it was proved in [1] that, for some range of \mathcal{R}_2 , there exists a critical number $\mathcal{R}_{1,c}$ such that if $\mathcal{R}_1 < \mathcal{R}_{1,c}$ then the motionless state $u = 0$ is asymptotically stable, while if $\mathcal{R}_1 > \mathcal{R}_{1,c}$ then $u = 0$ is unstable and nontrivial time periodic solutions bifurcate from $u = 0$.

On the other hand, when $\varepsilon > 0$, the system (1.1)–(1.4) is a hyperbolic-parabolic system and the limit $\varepsilon \rightarrow 0$ is a singular limit from a hyperbolic-parabolic system to a parabolic system. One of mathematical questions

is thus whether the artificial compressible system (1.1)–(1.4) gives a good approximation of the incompressible system in the singular limit $\varepsilon \rightarrow 0$.

The aim of this paper is to consider whether the artificial compressible system approximates well the incompressible system when a Hopf bifurcation, i.e., a time periodic bifurcation, occurs in the incompressible system. In this paper we shall investigate the Hopf bifurcation problem of the singularly perturbed system (1.1)–(1.4) with $0 < \varepsilon \ll 1$ for \mathcal{R}_1 near a critical value $\mathcal{R}_{1,c}$, where a Hopf bifurcation occurs from the motionless state $u = 0$ in the incompressible system (1.1)–(1.4) with $\varepsilon = 0$ for $\mathcal{R}_1 \sim \mathcal{R}_{1,c}$ as was shown in [1].

The artificial compressible system (1.1)–(1.4) with $\varepsilon > 0$ was introduced by Chorin ([2, 3, 4]) and Temam ([17, 18]) to avoid the computational difficulties in the incompressible system due to the constraint $\operatorname{div} \mathbf{w} = 0$. By using the artificial compressible system (1.1)–(1.4) with small $\varepsilon > 0$, Chorin [2, 3, 4] computed stationary convective bifurcating patterns near the onset of convection when $\mathcal{R}_2 = 0$, which suggests the artificial compressible system (1.1)–(1.4) would give a good approximation of the incompressible system. Temam ([17, 18, 19]) introduced the artificial compressible system with an additional stabilizing nonlinear term (in the context of (1.1)–(1.4), the corresponding stabilizing terms are to be $+\frac{1}{2}(\operatorname{div} \mathbf{w})\mathbf{w}$, $+\frac{1}{2}(\operatorname{div} \mathbf{w})\theta$ and $+\frac{1}{2}(\operatorname{div} \mathbf{w})\psi$ which are added on the left-hand side of (1.2), (1.3) and (1.4), respectively); and for such a system with stabilizing nonlinear term, the global in time solutions of the initial boundary value problem for the artificial compressible system on a two-dimensional bounded domain D converge in the limit $\varepsilon \rightarrow 0$ to the one for the incompressible system on time interval $(0, T)$ for all $T > 0$. Convergence result was also established in the framework of weak solutions on three-dimensional bounded domains. Donatelli [5, 6] and Donatelli and Marcati [7, 8] proved similar convergence results in the case of unbounded domains on any finite time interval by using the wave equation structure of the pressure and the dispersive estimates.

In this paper we shall show that a time periodic bifurcation occurs in (1.1)–(1.4) with $0 < \varepsilon \ll 1$ for \mathcal{R}_1 near the criticality $\mathcal{R}_{1,c}$ and the bifurcating solution branch converges as $\varepsilon \rightarrow 0$ to the time periodic bifurcating solution branch of the incompressible system obtained in [1].

To prove these results, we shall employ the Lyapunov-Schmidt method in a time periodic function space \mathcal{X}_a . For this purpose, we need to investigate the spectral properties of the operator $\partial_t + L_{\mathcal{R}_1}^\varepsilon$ in \mathcal{X}_a for \mathcal{R}_1 near the criticality $\mathcal{R}_{1,c}$, where $L_{\mathcal{R}_1}^\varepsilon$ denotes the linearized operator around $u = 0$ for (1.1)–(1.4),

which was studied in [11] in detail. Based on the spectral properties of the operator $\partial_t + L_{\mathcal{R}_1}^\varepsilon$, we shall show that (1.1)–(1.4) has a nontrivial bifurcating time periodic solution u^ε for sufficiently small $\varepsilon > 0$ and that u^ε converges to the bifurcating time periodic solution of the incompressible system (1.1)–(1.4) with $\varepsilon = 0$ as $\varepsilon \rightarrow 0$.

One of the key points in the proof is to use the norms with ε -weights, which enables us to establish the uniform estimates in the nonlinear problem as well as in the linearized problem. Another key is the fact that the limiting solution of the incompressible system and the eigenfunctions for the critical eigenvalues of $L_{\mathcal{R}_1}^\varepsilon$ for $\mathcal{R}_1 \sim \mathcal{R}_{1,c}$ are smooth, which compensate the loss of uniform estimates in ε . There is one more thing to be mentioned. At criticality $\mathcal{R}_1 = \mathcal{R}_{1,c}^\varepsilon$, the linearized problem has nontrivial time periodic solutions and the bifurcating solutions of the nonlinear problem are obtained as a perturbation of a time periodic solution of the linearized problem. The period of the bifurcating solution of the nonlinear problem is also a perturbation of the period of the time periodic solution of the linearized problem. This perturbation of the period is transformed into the equation as a perturbation of the time derivative of the unknown. In the case of the incompressible problem (i.e., in a classical setting), this perturbation term can be regraded as a regular perturbation. In contrast to the classical case, in the case of the artificial compressible problem, this perturbation term is not a regular perturbation. To overcome this difficulty, we put it into the principal part of the linearized operator and establish uniform estimates for small ε .

In the proof, we make use of the two-dimensional aspect of the problem to estimate the nonlinearity uniformly in small ε . It is not straightforward to extend the argument to the three-dimensional problem.

We shall also show that the bifurcating time periodic solutions of (1.1)–(1.4) for small $\varepsilon > 0$ is stable under perturbations with the same symmetries as those of the time periodic solutions, if the corresponding bifurcating time periodic solutions of the incompressible system is stable. This, in particular, implies that, by using the artificial compressible system with small $\varepsilon > 0$, one can numerically compute bifurcating time periodic solutions which are close to the bifurcating ones of the incompressible system near the onset of convection. See [12, 13, 20] for the stability of stationary bifurcating solutions.

This paper is organized as follows. In section 2 we introduce notation used in this paper. In section 3 we state the results on the Hopf bifurcation for the incompressible problem obtained in [1]. In section 4 we state the

result on the occurrence of the Hopf bifurcation in the artificial compressible system with small ε and also the result on the singular limit $\varepsilon \rightarrow 0$ of the time periodic bifurcating solutions. In section 5 we summarize the results on the spectral properties of the linearized problem obtained in [11]. Section 6 is devoted to a proof of the existence and singular limit. In section 7 we state the result on the stability of the bifurcating solutions and give a proof of the stability.

2 Preliminaries

In this section we introduce notation used in this paper. Let $\Omega = \mathbb{T}_{\frac{2\pi}{\alpha}} \times (0, 1)$. We denote by $L^p(\Omega)$ the usual L^p space on Ω with norm $\|\cdot\|_p$. We also denote by $\mathbf{L}^p(\Omega)$ the space of all L^p vector fields on Ω with norm $\|\cdot\|_p$. The inner product of $L^2(\Omega)$ is denoted by $(\cdot, \cdot)_{L^2}$. We also denote by $(\cdot, \cdot)_{\mathbf{L}^2}$ the inner product of $\mathbf{L}^2(\Omega)$.

The inner product and the norm of $\mathbf{L}^2(\Omega) \times L^2(\Omega) \times L^2(\Omega)$ are defined by

$$(\mathbf{u}_1, \mathbf{u}_2) = \text{Pr}^{-1}(\mathbf{w}_1, \mathbf{w}_2)_{L^2} + (\theta_1, \theta_2)_{L^2} + (\psi_1, \psi_2)_{L^2}$$

for $\mathbf{u}_j = {}^\top(\mathbf{w}_j, \theta_j, \psi_j) \in \mathbf{L}^2(\Omega) \times L^2(\Omega) \times L^2(\Omega)$

$$\|\mathbf{u}\|_2 = \sqrt{(\mathbf{u}, \mathbf{u})} = \sqrt{\text{Pr}^{-1}\|\mathbf{w}\|_2^2 + \|\theta\|_2^2 + \|\psi\|_2^2}$$

for $\mathbf{u} = {}^\top(\mathbf{w}, \theta, \psi) \in \mathbf{L}^2(\Omega) \times L^2(\Omega) \times L^2(\Omega)$, respectively.

The symbols $H^k(\Omega)$ and $\mathbf{H}^k(\Omega)$ stand for the k th order L^2 Sobolev spaces for scalar functions and for vector fields, respectively.

We define the spaces of L^2 and H^k functions with vanishing mean value on Ω by

$$L_*^2(\Omega) = \{\phi \in L^2(\Omega); \int_{\Omega} \phi(x) dx = 0\}$$

and

$$H_*^k(\Omega) = H^k(\Omega) \cap L_*^2(\Omega),$$

respectively.

We shall employ the following function spaces with symmetries:

$$L_{sym}^2(\Omega) = \{\theta \in L^2(\Omega); \theta \text{ is even in } x_1\},$$

$$L_{*,sym}^2(\Omega) = L_{sym}^2(\Omega) \cap L_*^2(\Omega),$$

$$\begin{aligned}
H_{*,sym}^k(\Omega) &= H^k(\Omega) \cap L_{*,sym}^2(\Omega), \\
\mathbf{L}_{sym}^2(\Omega) &= \{\mathbf{v} = {}^\top(v^1, v^2) \in \mathbf{L}^2(\Omega); v^1 \text{ is odd in } x_1, v^2 \text{ is even in } x_1\}.
\end{aligned}$$

It is known ([9, 16, 19]) that $\mathbf{L}_{sym}^2(\Omega)$ admits the Helmholtz decomposition:

$$\mathbf{L}_{sym}^2(\Omega) = \mathbf{L}_{\sigma,sym}^2(\Omega) \oplus \mathbf{G}_{sym}^2(\Omega) \quad (\text{orthogonal decomposition}),$$

where

$$\begin{aligned}
\mathbf{L}_{\sigma,sym}^2(\Omega) &= \{\mathbf{w} \in \mathbf{L}_{sym}^2(\Omega); \operatorname{div} \mathbf{w} = 0 \text{ in } \Omega, \mathbf{w} \cdot \mathbf{n}|_{\partial\Omega} = 0\}, \\
\mathbf{G}_{sym}^2(\Omega) &= \{\nabla\phi; \phi \in H_{*,sym}^1(\Omega)\}.
\end{aligned}$$

We denote by \mathbb{P}_σ the orthogonal projection on $\mathbf{L}_{\sigma,sym}^2(\Omega)$.

We define the function spaces \mathbf{X} , \mathbf{Y} , \mathbb{X}_σ , \mathbb{Y}_σ , X and Y by

$$\begin{aligned}
\mathbf{X} &= \mathbf{L}_{sym}^2(\Omega) \times \mathbf{L}_{sym}^2(\Omega) \times \mathbf{L}_{sym}^2(\Omega), \\
\mathbf{Y} &= [\mathbf{H}_b^2(\Omega) \times (H^2 \cap H_0^1)(\Omega) \times (H^2 \cap H_0^1)(\Omega)] \cap \mathbf{X}, \\
\mathbb{X}_\sigma &= \mathbf{L}_{\sigma,sym}^2(\Omega) \times \mathbf{L}_{sym}^2(\Omega) \times \mathbf{L}_{sym}^2(\Omega), \\
\mathbb{Y}_\sigma &= [\mathbf{H}_b^2(\Omega) \times (H^2 \cap H_0^1)(\Omega) \times (H^2 \cap H_0^1)(\Omega)] \cap \mathbb{X}_\sigma, \\
X &= H_{*,sym}^1(\Omega) \times \mathbf{L}_{sym}^2(\Omega) \times \mathbf{L}_{sym}^2(\Omega) \times \mathbf{L}_{sym}^2(\Omega), \\
Y &= [H_{*,sym}^1(\Omega) \times \mathbf{H}_b^2(\Omega) \times (H^2 \cap H_0^1)(\Omega) \times (H^2 \cap H_0^1)(\Omega)] \cap X,
\end{aligned}$$

respectively, where

$$\mathbf{H}_b^2(\Omega) = \left\{ \mathbf{w} = {}^\top(w^1, w^2) \in \mathbf{H}^2(\Omega); \frac{\partial w^1}{\partial x_2} = w^2 = 0 \text{ on } \{x_2 = 0, 1\} \right\}$$

$$H_0^1(\Omega) = \{\theta \in H^1(\Omega); \theta|_{x_2=0,1} = 0\}.$$

We will also employ the space \mathbf{X}^1 defined by

$$\mathbf{X}^1 = [\mathbf{H}_b^1(\Omega) \times H_0^1(\Omega) \times H_0^1(\Omega)] \cap \mathbf{X}$$

with norm

$$\|\mathbf{u}\|_{\mathbf{X}^1} = \sqrt{\operatorname{Pr}^{-1} \|\nabla \mathbf{w}\|_2^2 + \|\nabla \theta\|_2^2 + \|\nabla \psi\|_2^2}$$

for $\mathbf{u} = {}^\top(\mathbf{w}, \theta, \psi) \in \mathbf{X}^1$. Here

$$\mathbf{H}_b^1(\Omega) = \{\mathbf{w} = {}^\top(w^1, w^2) \in \mathbf{H}^1(\Omega); w^2|_{x_2=0,1} = 0\}.$$

Observe that the Poincaré inequality $\|\mathbf{w}\|_2 \leq C\|\nabla\mathbf{w}\|_2$ holds for $\mathbf{w} = {}^\top(w^1, w^2) \in \mathbf{H}_b^1(\Omega) \cap \mathbf{L}_{sym}^2(\Omega)$, which follows from the fact $\int_\Omega w^1 dx = 0$ due to the oddness of w^1 in x_1 and the fact $w^2|_{x_2=0,1} = 0$.

We next introduce $\|\mathbf{u}\|_{(\mathbf{X}^1)^*}$. We expand $\mathbf{u} = {}^\top(\mathbf{w}, \theta, \psi) \in \mathbf{X}$ with $\mathbf{w} = {}^\top(w^1, w^2)$ in the Fourier series as

$$\begin{aligned} w^1 &= \sum_{j \geq 1, k \geq 0} w_{jk}^1 \sin \alpha j x_1 \cos k \pi x_2, \\ w^2 &= \sum_{j \geq 0, k \geq 1} w_{jk}^2 \cos \alpha j x_1 \sin k \pi x_2, \\ \theta &= \sum_{j \geq 0, k \geq 1} \theta_{jk} \cos \alpha j x_1 \sin k \pi x_2, \\ \psi &= \sum_{j \geq 0, k \geq 1} \psi_{jk} \cos \alpha j x_1 \sin k \pi x_2, \end{aligned}$$

and define $\|\mathbf{u}\|_{(\mathbf{X}^1)^*}$ by

$$\|\mathbf{u}\|_{(\mathbf{X}^1)^*} = \left[\sum_{j \geq 0, k \geq 0} (\alpha^2 j^2 + k^2 \pi^2)^{-1} (\text{Pr}^{-1} \{(w_{jk}^1)^2 + (w_{jk}^2)^2\} + \theta_{jk}^2 + \psi_{jk}^2) \right]^{\frac{1}{2}},$$

where $w_{0k}^1 = w_{j0}^2 = \theta_{j0} = \psi_{j0} = 0$ for $k, j \geq 0$. It is easily verified that

$$\|\mathbf{u}\|_{(\mathbf{X}^1)^*} \leq C\|\mathbf{u}\|_2$$

and

$$|(\mathbf{u}_1, \mathbf{u}_2)| \leq \|\mathbf{u}_1\|_{\mathbf{X}^1} \|\mathbf{u}_2\|_{(\mathbf{X}^1)^*}.$$

We set

$$X^1 = H_{*,sym}^1(\Omega) \times \mathbf{X}^1.$$

We next introduce function spaces on time intervals. Let $t_1 < t_2$. We define $\mathcal{X}(t_1, t_2)$ and $\mathcal{Y}(t_1, t_2)$ by

$$\begin{aligned} \mathcal{X}(t_1, t_2) &= L^2(t_1, t_2; X), \\ \mathcal{Y}(t_1, t_2) &= L^2(t_1, t_2, T; Y) \cap H^1(t_1, t_2, T; X), \end{aligned}$$

with norms $\|u\|_{\mathcal{X}(t_1, t_2)}$ and $\|u\|_{\mathcal{Y}(t_1, t_2)}$, respectively.

We define the inner product $\langle \mathbf{u}_1, \mathbf{u}_2 \rangle$ by

$$\langle \mathbf{u}_1, \mathbf{u}_2 \rangle = \frac{a}{2\pi} \int_0^{\frac{2\pi}{a}} (\mathbf{u}_1(t), \mathbf{u}_2(t)) dt$$

for $\mathbf{u}_j = {}^\top(\mathbf{w}_j, \theta_j, \psi_j) \in L^2(\mathbb{T}_{\frac{2\pi}{a}}; \mathbf{L}^2(\Omega) \times L^2(\Omega) \times L^2(\Omega))$ ($j = 1, 2$)

Let ε be a given positive number. For $u_j = {}^\top(\phi_j, \mathbf{u}_j) \in L^2(\Omega) \times \mathbf{L}^2(\Omega) \times L^2(\Omega) \times L^2(\Omega)$ ($j = 1, 2$), we define the inner product $(u_1, u_2)_\varepsilon$ and the norm $|||u|||_\varepsilon$ with ε -weights by

$$(u_1, u_2)_\varepsilon = \varepsilon^2(\phi_1, \phi_2)_{L^2} + (\mathbf{u}_1, \mathbf{u}_2)$$

for $u_j = {}^\top(\phi_j, \mathbf{u}_j) \in L^2(\Omega) \times \mathbf{L}^2(\Omega) \times L^2(\Omega) \times L^2(\Omega)$ ($j = 1, 2$) and

$$|||u|||_\varepsilon = \sqrt{(u, u)_\varepsilon} = \sqrt{\varepsilon^2 \|\phi\|_2^2 + \|\mathbf{u}\|_2^2}$$

for $u = {}^\top(\phi, \mathbf{u}) \in L^2(\Omega) \times \mathbf{L}^2(\Omega) \times L^2(\Omega) \times L^2(\Omega)$, and, likewise, for $\mathbf{u}_j = {}^\top(\mathbf{w}_j, \theta_j, \psi_j) \in L^2(\mathbb{T}_{\frac{2\pi}{a}}; \mathbf{L}^2(\Omega) \times \mathbf{L}^2(\Omega) \times L^2(\Omega) \times L^2(\Omega))$ ($j = 1, 2$), the inner product $\langle u_1, u_2 \rangle_\varepsilon$ is defined by

$$\langle u_1, u_2 \rangle_\varepsilon = \frac{a}{2\pi} \int_0^{\frac{2\pi}{a}} (u_1(t), u_2(t))_\varepsilon dt.$$

We shall also employ the norms $|||u|||_{\varepsilon, X^1}$, $|||u|||_{\varepsilon, \mathcal{X}(t_1, t_2)}$ and $|||u|||_{\varepsilon, \mathcal{Y}(t_1, t_2)}$ of $u = {}^\top(\phi, \mathbf{u})$ with ε weights defined by

$$\begin{aligned} |||u|||_{\varepsilon, X^1} &= \left\{ |||u|||_\varepsilon^2 + \varepsilon^2 |||\partial_x u|||_\varepsilon^2 \right\}^{\frac{1}{2}}, \\ |||u|||_{\varepsilon, \mathcal{X}(t_1, t_2)} &= \left\{ \int_{t_1}^{t_2} \left(\varepsilon^2 \|\phi\|_2^2 + \|\mathbf{u}\|_{(\mathbf{X}^1)^*}^2 + \varepsilon^6 \|\partial_x \phi\|_2^2 + \varepsilon^2 \|\mathbf{u}\|_2^2 \right) dt \right\}^{\frac{1}{2}}, \\ |||u|||_{\varepsilon, \mathcal{Y}(t_1, t_2)} &= \left\{ \sup_{t_1 \leq t \leq t_2} |||u(t)|||_{\varepsilon, X^1}^2 \right. \\ &\quad \left. + \int_{t_1}^{t_2} \left(|||\partial_x u|||_\varepsilon^2 + \varepsilon^2 |||\partial_t u|||_\varepsilon^2 + \varepsilon^2 \|\partial_x^2 \mathbf{u}\|_2^2 + \varepsilon^6 \|\partial_x \partial_t \phi\|_2^2 \right) dt \right\}^{\frac{1}{2}}. \end{aligned}$$

Time periodic function spaces \mathcal{X}_a and \mathcal{Y}_a with period $\frac{2\pi}{a}$ are defined by

$$\mathcal{X}_a = L^2(\mathbb{T}_{\frac{2\pi}{a}}; X),$$

$$\mathcal{Y}_a = L^2(\mathbb{T}_{\frac{2\pi}{a}}; Y) \cap H^1(\mathbb{T}_{\frac{2\pi}{a}}; X),$$

respectively. We denote the norm of $u \in \mathcal{X}_a$ (resp. $u \in \mathcal{Y}_a$) by $\|u\|_{\mathcal{X}_a} = \|u\|_{\mathcal{X}(0, \frac{2\pi}{a})}$ (resp. $\|u\|_{\mathcal{Y}_a} = \|u\|_{\mathcal{Y}(0, \frac{2\pi}{a})}$), and likewise, the norm of $u \in \mathcal{X}_a$ (resp. $u \in \mathcal{Y}_a$) by $\|u\|_{\varepsilon, \mathcal{X}_a} = \|u\|_{\varepsilon, \mathcal{X}(0, \frac{2\pi}{a})}$ (resp. $\|u\|_{\varepsilon, \mathcal{Y}_a} = \|u\|_{\varepsilon, \mathcal{Y}(0, \frac{2\pi}{a})}$).

The resolvent set and spectrum of an operator A are denoted by $\rho(A)$ and $\sigma(A)$, respectively. We denote by $\mathfrak{B}(E_1, E_2)$ the space of all bounded linear operators from E_1 to E_2 .

3 Hopf bifurcation in the incompressible system

In this section we summarize the results on the Hopf bifurcation for the incompressible system (1.1) obtained in [1] and introduce the associated pressure of the velocity field of the bifurcating time periodic solution.

Setting $\varepsilon = 0$ in (1.1)–(1.4), we obtain the incompressible system:

$$\begin{cases} \operatorname{div} \mathbf{w} = 0, \\ \partial_t \mathbf{w} - \operatorname{Pr} \Delta \mathbf{w} + \operatorname{Pr} \nabla \phi - \operatorname{Pr} \mathcal{R}_1 \theta \mathbf{e}_2 + \operatorname{Pr} \mathcal{R}_2 \psi \mathbf{e}_2 + \mathbf{w} \cdot \nabla \mathbf{w} = \mathbf{0}, \\ \partial_t \theta - \Delta \theta - \mathcal{R}_1 \mathbf{w} \cdot \mathbf{e}_2 + \mathbf{w} \cdot \nabla \theta = 0, \\ \partial_t \psi - d \Delta \psi - \mathcal{R}_2 \mathbf{w} \cdot \mathbf{e}_2 + \mathbf{w} \cdot \nabla \psi = 0. \end{cases} \quad (3.1)$$

The boundary condition on the boundary $\{x_2 = 0, 1\}$ is given as

$$\frac{\partial w^1}{\partial x_2} = w^2 = \theta = \psi = 0 \text{ on } \{x_2 = 0, 1\}. \quad (3.2)$$

The problem (3.1)–(3.2) then has a trivial stationary solution $u = 0$ which corresponds to the motionless state. See [1, 11] for the derivation of the non-dimensional perturbation equations (3.1).

The result on the Hopf bifurcation for (3.1)–(3.2) by Bona, Hsia, Ma and Wang [1] is summarized as follows. There are positive numbers \mathcal{R}_{2*} and \mathcal{R}_2^* such that if $\mathcal{R}_2 \in [\mathcal{R}_{2*}, \mathcal{R}_2^*)$, $\operatorname{Pr} > 1$ and $0 < d < 1$, then there exists a critical number $\mathcal{R}_{1,c}$ such that the basic state $u = 0$ is stable when $\mathcal{R}_1 < \mathcal{R}_{1,c}$; and if $\mathcal{R}_1 > \mathcal{R}_{1,c}$, then $u = 0$ loses its stability and time periodic solutions of the incompressible problem (3.1)–(3.2) bifurcate from $u = 0$.

To state the bifurcating result in [1] more precisely, we take $\eta = \mathcal{R}_1 - \mathcal{R}_{1,c}$ as a bifurcation parameter. The linearized operator $\mathbb{L}_{\mathcal{R}_{1,c} + \eta}$ on \mathbb{X}_σ is then

given by

$$D(\mathbb{L}_{\mathcal{R}_{1,c}+\eta}) = \mathbb{Y}_\sigma,$$

$$\mathbb{L}_{\mathcal{R}_{1,c}+\eta} = \begin{pmatrix} -\text{Pr}\mathbb{P}_\sigma\Delta & -\text{Pr}(\mathcal{R}_{1,c} + \eta)\mathbb{P}_\sigma\mathbf{e}_2 & \text{Pr}\mathcal{R}_2\mathbb{P}_\sigma\mathbf{e}_2 \\ -(\mathcal{R}_{1,c} + \eta)^\top\mathbf{e}_2 & -\Delta & 0 \\ -\mathcal{R}_2^\top\mathbf{e}_2 & 0 & -d\Delta \end{pmatrix}.$$

We write problem (3.1)–(3.2) in the form

$$\partial_t \mathbf{u} + \mathbb{L}_{\mathcal{R}_{1,c}} \mathbf{u} + \eta \mathbb{P} \mathbf{K} \mathbf{u} + \mathbb{P} \mathbf{N}(\mathbf{u}) = \mathbf{0}, \quad (3.3)$$

where $\mathbf{u} = {}^\top(\mathbf{w}, \theta, \psi)$,

$$\mathbb{P} = \begin{pmatrix} \mathbb{P}_\sigma & \mathbf{0} & \mathbf{0} \\ {}^\top\mathbf{0} & 1 & 0 \\ {}^\top\mathbf{0} & 0 & 1 \end{pmatrix}, \quad \mathbf{K} = \begin{pmatrix} \mathbf{0} & -\text{Pr} \mathbf{e}_2 & \mathbf{0} \\ -{}^\top\mathbf{e}_2 & 0 & 0 \\ {}^\top\mathbf{0} & 0 & 0 \end{pmatrix},$$

and

$$\mathbf{N}(\mathbf{u}) = \mathbf{N}(\mathbf{u}, \mathbf{u})$$

with

$$\mathbf{N}(\mathbf{u}_1, \mathbf{u}_2) = \begin{pmatrix} \mathbf{w}_1 \cdot \nabla \mathbf{w}_2 \\ \mathbf{w}_1 \cdot \nabla \theta_2 \\ \mathbf{w}_1 \cdot \nabla \psi_2 \end{pmatrix} \text{ for } \mathbf{u}_j = \begin{pmatrix} \mathbf{w}_j \\ \theta_j \\ \psi_j \end{pmatrix} \quad (j = 1, 2).$$

The adjoint operator $\mathbb{L}_{\mathcal{R}_{1,c}+\eta}^*$ on \mathbb{X}_σ is given by

$$\mathbb{L}_{\mathcal{R}_{1,c}+\eta}^* = \begin{pmatrix} -\text{Pr}\mathbb{P}_\sigma\Delta & -\text{Pr}(\mathcal{R}_{1,c} + \eta)\mathbb{P}_\sigma\mathbf{e}_2 & -\text{Pr}\mathcal{R}_2\mathbb{P}_\sigma\mathbf{e}_2 \\ -(\mathcal{R}_{1,c} + \eta)\mathbf{e}_2 & -\Delta & 0 \\ \mathcal{R}_2\mathbf{e}_2 & 0 & -d\Delta \end{pmatrix}$$

with domain $D(\mathbb{L}_{\mathcal{R}_{1,c}+\eta}^*) = D(\mathbb{L}_{\mathcal{R}_{1,c}+\eta}) = \mathbb{Y}_\sigma$.

Bona, Hsia, Ma and Wang in [1] proved the following result on the spectrum of $\mathbb{L}_{\mathcal{R}_{1,c}+\eta}$.

Proposition 3.1 ([1]) (i) *There exist positive numbers \mathcal{R}_{2*} and \mathcal{R}_2^* such that if $\mathcal{R}_2 \in [\mathcal{R}_{2*}, \mathcal{R}_2^*]$, $\text{Pr} > 1$ and $0 < d < 1$, then the following assertions hold. There exist positive constants η_0 , b_0 and Λ_0 such that if $|\eta| \leq \eta_0$ then it holds that*

$$\Sigma \setminus \{\lambda_+(\eta), \lambda_-(\eta)\} \subset \rho(-\mathbb{L}_{\mathcal{R}_{1,c}+\eta}),$$

where $\Sigma = \{\lambda \in \mathbb{C}; \operatorname{Re}\lambda \geq -b_0|\operatorname{Im}\lambda|^2 - \Lambda_0\}$; $\lambda_+(\eta)$ and $\lambda_-(\eta)$ are simple eigenvalues of $-\mathbb{L}_{\mathcal{R}_{1,c}+\eta}$ satisfying $\lambda_-(\eta) = \overline{\lambda_+(\eta)}$ and

$$\lambda_+(0) = ia, \quad \frac{d\operatorname{Re}\lambda_+}{d\eta}(0) > 0.$$

Here a is a positive constant.

(ii) Let \mathbf{u}_\pm be eigenfunctions for the eigenvalues $\pm ia$ of $-\mathbb{L}_{\mathcal{R}_{1,c}}$ and let \mathbf{u}_\pm^* be eigenfunctions for the eigenvalues $\mp ia$ of the adjoint operator $-\mathbb{L}_{\mathcal{R}_{1,c}}^*$ satisfying $(\mathbf{u}_j, \mathbf{u}_k^*) = \delta_{jk}$, where $j, k \in \{+, -\}$. Then $\mathbf{u} = \overline{\mathbf{u}_+}$ and $\mathbf{u}^* = \overline{\mathbf{u}_+^*}$; and the eigenprojections \mathbf{P}_\pm for the eigenvalues $\pm ia$ of $-\mathbb{L}_{\mathcal{R}_{1,c}}$ are given by

$$\mathbf{P}_\pm \mathbf{u} = (\mathbf{u}, \mathbf{u}_\pm^*) \mathbf{u}_\pm.$$

Furthermore, it holds that

$$\frac{d\operatorname{Re}\lambda_+}{d\eta}(0) = -\operatorname{Re}(\mathbf{K}\mathbf{u}_+, \mathbf{u}_+^*) > 0.$$

We next define the operators \mathbb{B} and \mathbb{B}^* on $L^2(\mathbb{T}_{\frac{2\pi}{a}}; \mathbb{X}_\sigma)$ by

$$\mathbb{B} = \partial_t + \mathbb{L}_{\mathcal{R}_{1,c}} \quad \text{and} \quad \mathbb{B}^* = -\partial_t + \mathbb{L}_{\mathcal{R}_{1,c}}^*$$

with domain $D(\mathbb{B}) = D(\mathbb{B}^*) = L^2(\mathbb{T}_{\frac{2\pi}{a}}; D(\mathbb{L}_0)) \cap H^1(\mathbb{T}_{\frac{2\pi}{a}}; \mathbb{X}_\sigma)$.

One can then see by a direct computation that the functions

$$\mathbf{z}_\pm = e^{\pm iat} \mathbf{u}_\pm, \quad \mathbf{z}_\pm^* = e^{\pm iat} \mathbf{u}_\pm^*$$

are eigenfunctions for the eigenvalue 0 of \mathbb{B} and \mathbb{B}^* with properties

$$\langle \mathbf{z}_\pm, \mathbf{z}_\pm^* \rangle = 1, \quad \langle \mathbf{z}_\mp, \mathbf{z}_\pm^* \rangle = 0.$$

We set

$$[\mathbf{z}]_\pm = \langle \mathbf{z}, \mathbf{z}_\pm^* \rangle.$$

The operators $\hat{\mathcal{P}}_\pm$ defined by

$$\hat{\mathcal{P}}_\pm \mathbf{z} = [\mathbf{z}]_\pm \mathbf{z}_\pm \quad (\mathbf{z} \in L^2(\mathbb{T}_{\frac{2\pi}{a}}; \mathbb{X}_\sigma))$$

are then projections which satisfy $\hat{\mathcal{P}}_j \hat{\mathcal{P}}_k = \delta_{jk} \hat{\mathcal{P}}_j$ for $j, k \in \{+, -\}$. As for the operator \mathbb{B} , we then have the following proposition.

Proposition 3.2 0 is a semisimple eigenvalue of \mathbb{B} and it holds that

$$L^2(\mathbb{T}_{\frac{2\pi}{a}}; \mathbb{X}_\sigma) = \text{Ker}(\mathbb{B}) \oplus R(\mathbb{B}).$$

Set

$$\hat{\mathcal{P}}_0 = \hat{\mathcal{P}}_+ + \hat{\mathcal{P}}_-, \quad \hat{\mathcal{Q}}_0 = I - \hat{\mathcal{P}}_0.$$

Then $\hat{\mathcal{P}}_0$ is an eigenprojection for the eigenvalue 0 of \mathbb{B} and $\hat{\mathcal{Q}}_0$ is a projection on $R(\mathbb{B})$ along $N(\mathbb{B})$. There holds that $\mathbf{u} \in R(\mathbb{B})$ if and only if $\hat{\mathcal{P}}_0 \mathbf{u} = 0$, i.e., $[\mathbf{z}]_+ = [\mathbf{z}]_- = 0$.

We note that if \mathbf{u} is a real valued function, then $[\mathbf{u}]_- = \overline{[\mathbf{u}]_+}$, and hence,

$$\hat{\mathcal{P}}_0 \mathbf{u} = 2\text{Re}([\mathbf{u}]_+ \mathbf{z}_+).$$

Based on Proposition 3.1, a Hopf bifurcation would occur when $\mathcal{R}_1 > \mathcal{R}_{1,c}$. It is expected that (3.3) has a nontrivial time periodic solution of period $\frac{2\pi}{a(1+\omega)}$ with small ω for sufficiently small η . We thus change the variable $t \mapsto (1 + \omega)t$. The problem is then reduced to finding a nontrivial time periodic solution of period $\frac{2\pi}{a}$ to the equation

$$(1 + \omega)\partial_t \mathbf{u} + \mathbb{L}_{\mathcal{R}_{1,c}} \mathbf{u} + \eta \mathbb{P} \mathbf{K} \mathbf{u} + \mathbb{P} \mathbf{N}(\mathbf{u}) = \mathbf{0}. \quad (3.4)$$

The following result on the Hopf bifurcation was shown by Bona, Hsia, Ma and Wang in [1].

Proposition 3.3 ([1]) (i) *There exists a positive constant δ_0 such that (3.4) has a nontrivial time periodic solution $\mathbf{u}_\delta \in L^2(\mathbb{T}_{\frac{2\pi}{a}}; \mathbb{Y}_\sigma) \cap H^1(\mathbb{T}_{\frac{2\pi}{a}}; \mathbb{X}_\sigma)$ for $\eta = \eta_\delta$ and $\omega = \omega_\delta$, where $\{\eta_\delta, \omega_\delta, \mathbf{u}_\delta\}$ with a parameter δ takes the form*

$$\begin{aligned} \eta_\delta &= \tilde{\eta}_0 \delta^2 + \tilde{\eta}_1(\delta) \delta^3, \\ \omega_\delta &= \tilde{\omega}_0 \delta^2 + \tilde{\omega}_1(\delta) \delta^3, \\ \mathbf{u}_\delta &= \delta(\mathbf{z}_0 + \delta \mathbf{U}_\delta) \end{aligned}$$

for $|\delta| \leq \delta_0$. Here $\tilde{\eta}_0$ and $\tilde{\omega}_0$ are constants and $\tilde{\eta}_0 > 0$; $\tilde{\eta}_1(\delta)$ and $\tilde{\omega}_1(\delta)$ are analytic in δ satisfying $\tilde{\eta}_1(\delta) = O(1)$ and $\tilde{\omega}_1(\delta) = O(1)$ as $\delta \rightarrow 0$; $\mathbf{z}_0 = {}^\top(\mathbf{w}_0, \theta_0, \psi_0) = 2\text{Re}(\mathbf{z}_+)$; and \mathbf{U}_δ is in $R(\mathbb{B}) \cap [L^2(\mathbb{T}_{\frac{2\pi}{a}}; \mathbb{Y}_\sigma) \cap H^1(\mathbb{T}_{\frac{2\pi}{a}}; \mathbb{X}_\sigma)]$ and is analytic in δ .

(ii) *There exists a neighborhood \mathbf{O}_0 of ${}^\top(\eta, \omega, \mathbf{u}) = {}^\top(0, 0, \mathbf{0})$ in $\mathbb{R} \times \mathbb{R} \times [L^2(\mathbb{T}_{\frac{2\pi}{a}}; \mathbb{Y}_\sigma) \cap H^1(\mathbb{T}_{\frac{2\pi}{a}}; \mathbb{X}_\sigma)]$ such that if ${}^\top(\eta, \omega, \mathbf{u}) \in \mathbf{O}_0$ is a solution of (3.4), then*

$${}^\top(\eta, \omega, \mathbf{u}) \in \{{}^\top(\eta, \omega, \mathbf{0}) \in \mathbf{O}_0\} \cup \{{}^\top(\eta_\delta, \omega_\delta, \mathbf{u}_\delta(\cdot + \tau)); |\delta| \leq \delta_1, -\frac{\pi}{a} \leq \tau < \frac{\pi}{a}\}$$

where $\{\eta_\delta, \omega_\delta, \mathbf{u}_\delta\}$ is the solution branch obtained in (i).

We next introduce the associated pressure ϕ_δ of the time periodic bifurcating solution \mathbf{u}_δ of (3.4). It is known [9, 16, 19] that there exists the associated pressure $\phi_\delta \in L^2(\mathbb{T}_{\frac{2\pi}{a}}; H_{*,sym}^1(\Omega))$ of $\mathbf{u}_\delta = {}^\top(\mathbf{w}_\delta, \theta_\delta, \psi_\delta)$, namely, there exists a unique $\phi_\delta \in L^2(\mathbb{T}_{\frac{2\pi}{a}}; H_{*,sym}^1(\Omega))$ such that $u_\delta = {}^\top(\phi_\delta, \mathbf{u}_\delta) \in \mathcal{Y}_a$ is a time periodic solution of

$$(1 + \omega)\partial_t \begin{pmatrix} 0 \\ \mathbf{u} \end{pmatrix} + L_{\mathcal{R}_1, c}^\varepsilon u + \eta K u + N(u) = 0 \quad (3.5)$$

for $\eta = \eta_\delta$ and $\omega = \omega_\delta$. Here $L_{\mathcal{R}_1}^\varepsilon$, K and $N(\cdot)$ are the maps defined as follows; $L_{\mathcal{R}_1}^\varepsilon$ is the linearized operator around $u = 0$ on the space X with domain

$$D(L_{\mathcal{R}_1}^\varepsilon) = Y$$

and it is given in the form

$$L_{\mathcal{R}_1}^\varepsilon = \begin{pmatrix} 0 & \frac{1}{\varepsilon^2} \operatorname{div} & 0 & 0 \\ \operatorname{Pr} \nabla & -\operatorname{Pr} \Delta & -\operatorname{Pr} \mathcal{R}_1 \mathbf{e}_2 & \operatorname{Pr} \mathcal{R}_2 \mathbf{e}_2 \\ 0 & -\mathcal{R}_1^\top \mathbf{e}_2 & -\Delta & 0 \\ 0 & -\mathcal{R}_2^\top \mathbf{e}_2 & 0 & -d\Delta \end{pmatrix};$$

K is the linear operator given by

$$K = \left(\begin{array}{c|ccc} 0 & \top \mathbf{0} & 0 & 0 \\ \hline \mathbf{0} & & & \\ 0 & & \mathbf{K} & \\ 0 & & & \end{array} \right);$$

and $N(\cdot)$ is the nonlinear map given by

$$N(u) = N(u, u),$$

where

$$N(u_1, u_2) = \begin{pmatrix} 0 \\ \mathbf{N}(\mathbf{u}_1, \mathbf{u}_2) \end{pmatrix}$$

for $u_j = {}^\top(\phi_j, \mathbf{u}_j)$ ($j = 1, 2$).

The associated pressure ϕ_δ takes the form

$$\phi_\delta = \delta(p_0 + \delta\Phi_\delta),$$

where $p_0 = \text{Re}(e^{iat}\phi_+)$ and $\Phi_\delta \in L^2(\mathbb{T}_{\frac{2\pi}{a}}; H_{*,sym}^1(\Omega))$. Here $\phi_+ \in H_{*,sym}^1(\Omega)$ is the associated pressure of $\mathbf{u}_+ = {}^\top(\mathbf{w}_+, \theta_+, \psi_+)$, i.e., $u_+ = {}^\top(\phi_+, \mathbf{u}_+)$ satisfies

$$ia \begin{pmatrix} 0 \\ \mathbf{u}_+ \end{pmatrix} + L_{\mathcal{R}_{1,c}}^\varepsilon u_+ = 0,$$

and Φ_δ is the associated pressure of $\mathbf{U}_\delta = {}^\top(\mathbf{W}_\delta, \Theta_\delta, \Psi_\delta)$, i.e., $U_\delta = {}^\top(\Phi_\delta, \mathbf{U}_\delta)$ satisfies

$$\partial_t \begin{pmatrix} 0 \\ \mathbf{U}_\delta \end{pmatrix} + \delta\tilde{\omega}_\delta \partial_t \begin{pmatrix} 0 \\ \mathbf{z}_0 + \delta\mathbf{U}_\delta \end{pmatrix} + L_{\mathcal{R}_{1,c}}^\varepsilon U_\delta + \delta\tilde{\eta}_\delta K(z_0 + \delta U_\delta) + N(z_0 + \delta U_\delta) = 0,$$

where $z_0 = {}^\top(p_0, \mathbf{z}_0)$, $\tilde{\eta}_\delta = \delta^{-2}\eta_\delta$ and $\tilde{\omega}_\delta = \delta^{-2}\omega_\delta$. Furthermore, Φ_δ is in $H^1(\mathbb{T}_{\frac{2\pi}{a}}; H_{*,sym}^1(\Omega))$ and satisfies $\|\Phi_\delta\|_{H^1(\mathbb{T}_{\frac{2\pi}{a}}; H_*^1(\Omega))} \leq C$ uniformly for $|\delta| \leq \delta_0$ if δ_0 is suitably small.

The above decomposition of ϕ_δ into p_0 and Φ_δ parts can be seen by decomposing (3.4) into $\hat{\mathcal{P}}_0$ and $\hat{\mathcal{Q}}_0$ parts. To see the boundedness of $\{\Phi_\delta\}_{|\delta| \leq \delta_0}$, we first observe that \mathbf{U}_δ is a time periodic solution of

$$(1 + \delta^2\tilde{\omega}_\delta)\partial_t \mathbf{U}_\delta + \mathbb{L}_{\mathcal{R}_{1,c}} \mathbf{U}_\delta + \delta\tilde{\eta}_\delta \hat{\mathcal{Q}}_0 \mathbb{P}\mathbf{K}(z_0 + \delta\mathbf{U}_\delta) + \hat{\mathcal{Q}}_0 \mathbb{P}\mathbf{N}(z_0 + \delta\mathbf{U}_\delta) = \mathbf{0}.$$

Since z_0 is a smooth function and $\{\mathbf{U}_\delta\}_{|\delta| \leq \delta_0}$ is bounded in $L^2(\mathbb{T}_{\frac{2\pi}{a}}; \mathbb{Y}_\sigma) \cap H^1(\mathbb{T}_{\frac{2\pi}{a}}; \mathbb{X}_\sigma)$, we see that $\{\partial_t \mathbf{U}_\delta\}_{|\delta| \leq \delta_0}$ is bounded in $L^2(\mathbb{T}_{\frac{2\pi}{a}}; \mathbb{Y}_\sigma) \cap H^1(\mathbb{T}_{\frac{2\pi}{a}}; \mathbb{X}_\sigma)$ if δ_0 is sufficiently small. It then follows that $\{\Phi_\delta\}_{|\delta| \leq \delta_0}$ is bounded in $H^1(\mathbb{T}_{\frac{2\pi}{a}}; H_*^1(\Omega))$.

In summary we have the following proposition.

Proposition 3.4 (i) *There exists a positive constant δ_0 such that (3.5) has a nontrivial time periodic solution $u_\delta \in \mathcal{Y}_a$ for $\eta = \eta_\delta$ and $\omega = \omega_\delta$, where $\{\eta_\delta, \omega_\delta, u_\delta\}$ with a parameter δ takes the form*

$$\begin{aligned} \eta_\delta &= \tilde{\eta}_0 \delta^2 + \tilde{\eta}_1(\delta) \delta^3, \\ \omega_\delta &= \tilde{\omega}_0 \delta^2 + \tilde{\omega}_1(\delta) \delta^3, \\ u_\delta &= \delta(z_0 + \delta\mathbf{U}_\delta) \end{aligned}$$

for $|\delta| \leq \delta_0$. Here η_δ and ω_δ are the numbers given in Proposition 3.3 and $U_\delta \in \mathcal{Y}_a$ is analytic in δ . Furthermore, $\|U_\delta\|_{\mathcal{Y}_a} \leq C$ uniformly for $|\delta| \leq \delta_0$.

(ii) There exists a neighborhood \mathcal{O}_0 of ${}^\top(\eta, \omega, u) = {}^\top(0, 0, 0)$ in $\mathbb{R} \times \mathbb{R} \times \mathcal{Y}_a$ such that if ${}^\top(\eta, \omega, u) \in \mathcal{O}_0$ is a solution of (3.5), then

$${}^\top(\eta, \omega, u) \in \{{}^\top(\eta, \omega, 0) \in \mathcal{O}_0\} \cup \{{}^\top(\eta_\delta, \omega_\delta, u_\delta(\cdot + \tau)); |\delta| \leq \delta_1, -\frac{\pi}{a} \leq \tau < \frac{\pi}{a}\}$$

where $\{\eta_\delta, \omega_\delta, u_\delta\}$ is the solution branch given in (i).

We close this section by introducing notation related to Proposition 3.4. The associated pressure p_- of $\mathbf{u}_- = {}^\top(\mathbf{w}_-, \theta_-, \psi_-)$ is given by $\phi_- = \overline{\phi_+}$, and it holds that $u_- = {}^\top(\phi_-, \mathbf{u}_-)$ satisfies

$$-ia \begin{pmatrix} 0 \\ \mathbf{u}_- \end{pmatrix} + L_{\mathcal{R}_1, c}^{\varepsilon*} u_- = 0.$$

where $L_{\mathcal{R}_1}^{\varepsilon*} : X \rightarrow X$ is the adjoint operator of $L_{\mathcal{R}_1}^\varepsilon$ and it is defined by

$$D(L_{\mathcal{R}_1}^{\varepsilon*}) = Y,$$

$$L_{\mathcal{R}_1}^{\varepsilon*} = \begin{pmatrix} 0 & -\frac{1}{\varepsilon^2} \operatorname{div} & 0 & 0 \\ -\operatorname{Pr} \nabla & -\operatorname{Pr} \Delta & -\operatorname{Pr} \mathcal{R}_1 \mathbf{e}_2 & -\operatorname{Pr} \mathcal{R}_2 \mathbf{e}_2 \\ 0 & -\mathcal{R}_1^\top \mathbf{e}_2 & -\Delta & 0 \\ 0 & \mathcal{R}_2^\top \mathbf{e}_2 & 0 & -d\Delta \end{pmatrix}.$$

Similarly, we have the associated pressures ϕ_\pm^* of $\mathbf{u}_\pm^* = {}^\top(\mathbf{w}_\pm^*, \theta_\pm^*, \psi_\pm^*)$, and it holds that $u_\pm^* = {}^\top(\phi_\pm^*, \mathbf{u}_\pm^*)$ satisfies

$$\mp ia \begin{pmatrix} 0 \\ \mathbf{u}_\pm^* \end{pmatrix} + L_{\mathcal{R}_1, c}^{\varepsilon*} u_\pm^* = 0.$$

We define functions u_\pm , u_\pm^* , z_\pm and z_\pm^* by

$$u_\pm = {}^\top(\phi_\pm, \mathbf{u}_\pm), \quad u_\pm^* = {}^\top(\phi_\pm^*, \mathbf{u}_\pm^*), \quad z_\pm = e^{\pm iat} u_\pm, \quad z_\pm^* = e^{\pm iat} u_\pm^*, \quad (3.6)$$

respectively. We finally define the operators $\mathcal{P}_\pm : \mathcal{X}_a \rightarrow \mathcal{X}_a$ by

$$\mathcal{P}_\pm u = [\mathbf{u}]_\pm z_\pm$$

for $u = {}^\top(\phi, \mathbf{u}) \in \mathcal{X}_a$, and $\mathcal{P}_0, \mathcal{Q}_0 : \mathcal{X}_a \rightarrow \mathcal{X}_a$ by

$$\mathcal{P}_0 = \mathcal{P}_+ + \mathcal{P}_-, \quad \mathcal{Q}_0 = I - \mathcal{P}_0. \quad (3.7)$$

Observe that if $u = {}^\top(\phi, \mathbf{u})$ is real valued, then $\mathcal{P}_0 u = 2\operatorname{Re}([\mathbf{u}]_+ z_+)$.

4 Hopf bifurcation in the artificial compressible system

In this section we state the result on the occurrence of a Hopf bifurcation in the artificial compressible system (1.1)–(1.4) under (1.5) and the result on the singular limit $\varepsilon \rightarrow 0$ for the time periodic bifurcating solutions.

We fix the parameters Pr , d and \mathcal{R}_2 in such a way that these parameters satisfies the assumption of Proposition 3.1.

The time periodic problem for the artificial compressible system (1.1)–(1.5) is formulated as

$$\partial_t u + L_{\mathcal{R}_1}^\varepsilon u + N(u) = 0. \quad (4.1)$$

Here $u = {}^\top(\phi, \mathbf{u})$ with $\mathbf{u} = {}^\top(\mathbf{w}, \theta, \psi)$ is periodic in t , and $L_{\mathcal{R}_1, c}^\varepsilon$, K and $N(\cdot)$ are the maps given in (3.5).

As for the spectrum of the linearized operator $-L_{\mathcal{R}_1}^\varepsilon$ for \mathcal{R}_1 close to the criticality $\mathcal{R}_{1, c}$, we obtained the following result in [11]. (Cf., [14].)

Theorem 4.1 ([11]) (i) *There exist positive constants Λ_1 , ε_1 and η_1 such that for each $0 < \varepsilon \leq \varepsilon_1$ there exists a critical value $\mathcal{R}_{1, c}^\varepsilon = \mathcal{R}_{1, c} + O(\varepsilon^2)$ such that if $|\eta| \leq \eta_1$ with $\eta = \mathcal{R}_1 - \mathcal{R}_{1, c}^\varepsilon$, then*

$$\{\lambda \in \mathbb{C}; \text{Re } \lambda \geq -\Lambda_1\} \setminus \{\lambda_+^\varepsilon(\eta), \lambda_-^\varepsilon(\eta)\} \subset \rho(-L_{\mathcal{R}_{1, c}^\varepsilon + \eta}^\varepsilon),$$

where $\lambda_\pm^\varepsilon(\eta)$ are simple eigenvalues of $-L_{\mathcal{R}_{1, c}^\varepsilon + \eta}^\varepsilon$ satisfying $\lambda_-^\varepsilon(\eta) = \overline{\lambda_+^\varepsilon(\eta)}$,

$$\lambda_\pm^\varepsilon(\eta) = \lambda_\pm(\eta) + O(\varepsilon^2),$$

and

$$\lambda_+^\varepsilon(0) = ia^\varepsilon, \quad \frac{\partial \text{Re} \lambda_+^\varepsilon}{\partial \eta}(0) = -\text{Re}(\mathbf{K} \mathbf{u}_+, \mathbf{u}_+^*) + O(\varepsilon^2) > 0.$$

Here a^ε is a constant satisfying $a^\varepsilon = a + O(\varepsilon^2)$.

The eigenspaces for $\lambda_\pm^\varepsilon(0)$ are spanned by u_\pm^ε , respectively, where u_\pm^ε satisfy $\overline{u_-^\varepsilon} = u_+^\varepsilon$ and $u_\pm^\varepsilon = u_\pm + O(\varepsilon^2)$.

(ii) For $j = \pm$, the eigenprojections P_j^ε for $\lambda_j^\varepsilon(0)$ ($j = \pm$) satisfies

$$P_j^\varepsilon u = (u, u_j^{\varepsilon*})_\varepsilon u_j^\varepsilon,$$

where u_{\pm}^{ε} are eigenfunction for the eigenvalues $\lambda_{\pm}^{\varepsilon}$ of $-L_{\mathcal{R}_{1,c}^{\varepsilon}}$ satisfying $\overline{u_{-}^{\varepsilon}} = u_{+}^{\varepsilon}$ and $u_{+}^{\varepsilon} = u_{+} + O(\varepsilon^2)$ in Y ; $u_{\pm}^{\varepsilon*}$ are the adjoint eigenfunctions for the eigenvalues $\overline{\lambda_j^{\varepsilon}(0)}$ of $-L_{\mathcal{R}_{1,c}^{\varepsilon*}}$ satisfying $\overline{u_{-}^{\varepsilon*}} = u_{+}^{\varepsilon*}$, $(u_j^{\varepsilon}, u_k^{\varepsilon*})_{\varepsilon} = \delta_{jk}$ and $u_{+}^{\varepsilon*} = u_{+}^* + O(\varepsilon^2)$ in Y ; and P_{\pm}^0 are projections defined by $P_{\pm}^0 u = (\mathbf{u}, \mathbf{u}_{\pm}^*) u_{\pm}$ for $u = {}^{\top}(\phi, \mathbf{u})$. Here u_{\pm} and u_{\pm}^* are functions defined in (3.6).

Furthermore, if $k \in \mathbb{Z}$ with $k \geq 0$ and $1 \leq p \leq 2$, there exists a positive constant $\tilde{\varepsilon}_1 = \tilde{\varepsilon}_1(k, p, \Omega)$ such that

$$\begin{aligned} \|u_{\pm}^{\varepsilon}\|_{H^k \times H^k} + \|u_{\pm}^{\varepsilon*}\|_{H^k \times H^k} &\leq C, \\ \|u_{\pm}^{\varepsilon} - u_j\|_{H^k \times H^k} + \|u_{\pm}^{\varepsilon*} - u_j^*\|_{H^k \times H^k} &\leq C\varepsilon^2, \\ \|P_{\pm}^{\varepsilon} u\|_{H^k \times H^k} &\leq C\|u\|_{L^p \times L^p}, \\ \|(P_{\pm}^{\varepsilon} - P_{\pm}^0)u\|_{H^k \times H^k} &\leq C\varepsilon^2\|u\|_{L^p \times L^p} \end{aligned}$$

uniformly for $\varepsilon \in (0, \tilde{\varepsilon}_1]$.

By Theorem 4.1, a Hopf bifurcation is expected to occur when \mathcal{R}_1 passes the critical value $\mathcal{R}_{1,c}^{\varepsilon}$. In fact, we have the following bifurcation result.

To fix the time interval, we change the variables t and u into \tilde{t} and \tilde{u} , respectively, by $\frac{a^{\varepsilon}}{a}t = \tilde{t}$ and $u(x, t) = \tilde{u}(x, \tilde{t})$. Omitting the tildes $\tilde{\cdot}$ of \tilde{t} and \tilde{u} , we see from (4.1) that the problem is transformed into

$$\frac{a^{\varepsilon}}{a} \partial_t u + L_{\mathcal{R}_{1,c}^{\varepsilon}} u + \eta K u + N(u) = 0. \quad (4.2)$$

As in the incompressible problem, we look for a nontrivial time periodic solution of (4.2) whose period is $\frac{2\pi}{a(1+\omega)}$ with small ω for sufficiently small ε and η . We thus again change the variable $t \mapsto (1 + \omega)t$. The problem is then reduced to finding a nontrivial time periodic solution of period $\frac{2\pi}{a}$ of the following equation

$$\frac{a^{\varepsilon}}{a} (1 + \omega) \partial_t u + L_{\mathcal{R}_{1,c}^{\varepsilon}} u + \eta K u + N(u) = 0. \quad (4.3)$$

To formulate the time periodic problem in a functional analytic setting, we introduce the operator B^{ε} on \mathcal{X}_a defined by

$$B^{\varepsilon} = \frac{a^{\varepsilon}}{a} \partial_t + L_{\mathcal{R}_{1,c}^{\varepsilon}}$$

with domain $D(B^\varepsilon) = \mathcal{Y}_a$. The time periodic problem (4.3) is then formulated as

$$B^\varepsilon u + \frac{a^\varepsilon}{a} \omega \partial_t u + \eta K u + N(u) = 0, \quad u \in \mathcal{Y}_a. \quad (4.4)$$

Let $z_\pm^\varepsilon = e^{\pm iat} u_\pm^\varepsilon$. It is easily verified that $B^\varepsilon z_\pm^\varepsilon = 0$. Let

$$[u]_{\pm, \varepsilon} = \langle u, z_\pm^{\varepsilon*} \rangle_\varepsilon$$

with $z_\pm^{\varepsilon*} = e^{\pm iat} u_\pm^{\varepsilon*}$ and define the operators $\mathcal{P}_\pm^\varepsilon$ by

$$\mathcal{P}_\pm^\varepsilon u = [u]_{\pm, \varepsilon} z_\pm^\varepsilon.$$

The operators $\mathcal{P}_\pm^\varepsilon$ are projections onto the spaces $\text{span}\{z_\pm^\varepsilon\}$, respectively. As was shown in [11] (see Theorem 5.4 below), 0 is a semisimple eigenvalue of B^ε and the eigenprojection for the eigenvalue 0 is given by

$$\mathcal{P}^\varepsilon = \mathcal{P}_+^\varepsilon + \mathcal{P}_-^\varepsilon.$$

We set $\mathcal{Q}^\varepsilon = I - \mathcal{P}^\varepsilon$. Then

$$u \in R(\mathcal{Q}^\varepsilon) \text{ if and only if } [u]_{+, \varepsilon} = [u]_{-, \varepsilon} = 0.$$

If u is real valued, then $[u]_{+, \varepsilon} = \overline{[u]_{-, \varepsilon}}$, from which we see that $u \in R(\mathcal{Q}^\varepsilon)$ if and only if $[u]_{+, \varepsilon} = 0$, when u is a real valued function.

We state the result on the Hopf bifurcation for the artificial compressible system (1.1)–(1.4).

Theorem 4.2 (i) *There exist positive numbers ε_2 and δ_2 such that if $0 < \varepsilon \leq \varepsilon_2$, then there exists a nontrivial solution $u_\delta^\varepsilon \in \mathcal{Y}_a$ of (4.4) for $\eta = \eta_\delta^\varepsilon$ and $\omega = \omega_\delta^\varepsilon$, where*

$$\begin{aligned} \eta_\delta^\varepsilon &= \tilde{\eta}_0^\varepsilon \delta^2 + \tilde{\eta}_1^\varepsilon(\delta) \delta^3, \\ \omega_\delta^\varepsilon &= \tilde{\omega}_0^\varepsilon \delta^2 + \tilde{\omega}_1^\varepsilon(\delta) \delta^3, \\ u_\delta^\varepsilon &= \delta(z_0^\varepsilon + \delta U_\delta^\varepsilon) \end{aligned}$$

for $|\delta| \leq \delta_2$. Here $z_0^\varepsilon = \text{Re}(e^{iat} u_+^\varepsilon)$ and $[U_\delta^\varepsilon]_{\varepsilon, +} = 0$; and $\tilde{\eta}_0^\varepsilon, \tilde{\omega}_0^\varepsilon, \tilde{\eta}_1^\varepsilon(\delta), \tilde{\omega}_1^\varepsilon(\delta)$ and U_δ^ε satisfy $|\tilde{\eta}_0^\varepsilon| + |\tilde{\omega}_0^\varepsilon| \leq C$ and $|\tilde{\eta}_1^\varepsilon(\delta)| + |\tilde{\omega}_1^\varepsilon(\delta)| \leq C$; and, furthermore, U_+^ε satisfies $\partial_t U_\delta^\varepsilon \in \mathcal{Y}_a$ with $[\partial_t U_\delta^\varepsilon]_{+, \varepsilon} = 0$ and $|||U_\delta^\varepsilon|||_{\varepsilon, \mathcal{Y}_a} + |||\partial_t U_\delta^\varepsilon|||_{\varepsilon, \mathcal{Y}_a} \leq C$ uniformly in $0 < \varepsilon \leq \varepsilon_2$ and $|\delta| \leq \delta_2$.

(ii) Let $u_\delta = {}^\top(\phi_\delta, \mathbf{u}_\delta) = {}^\top(\delta(p_0 + \delta\Phi_\delta), \delta(\mathbf{z}_0 + \delta\mathbf{U}_\delta))$ be the time periodic bifurcating solution of (3.5) for $\eta = \eta_\delta$ and $\omega = \omega_\delta$ given in Proposition 3.4. Then

$$\begin{aligned}\tilde{\eta}_0^\varepsilon &= \tilde{\eta}_0 + O(\varepsilon), & \tilde{\eta}_1^\varepsilon(\delta) &= \tilde{\eta}_1(\delta) + O(\varepsilon), \\ \tilde{\omega}_0^\varepsilon &= \tilde{\omega}_0 + O(\varepsilon), & \tilde{\omega}_1^\varepsilon(\delta) &= \tilde{\omega}_1(\delta) + O(\varepsilon)\end{aligned}$$

and

$$\|U_\delta^\varepsilon - U_\delta\|_{\varepsilon, \mathcal{Y}_a} \leq C\varepsilon$$

uniformly for $0 < \varepsilon \leq \varepsilon_2$ and $|\delta| \leq \delta_2$, where $\tilde{\eta}_0, \tilde{\omega}_0, \tilde{\eta}_1(\delta)$ and $\tilde{\omega}_1(\delta)$ are the numbers given in Proposition 3.4. In particular,

$$\begin{aligned}|\eta_\delta^\varepsilon - \eta_\delta| + |\omega_\delta^\varepsilon - \omega_\delta| &\leq C\varepsilon\delta^2, \\ \|\phi_\delta^\varepsilon - \phi_\delta\|_{C(\mathbb{T}_{\frac{2\pi}{a}}; L^2)} + \|\phi_\delta^\varepsilon - \phi_\delta\|_{L^2(\mathbb{T}_{\frac{2\pi}{a}}; H^1)} &\leq C|\delta| \\ \|\mathbf{u}_\delta^\varepsilon - \mathbf{u}_\delta\|_{C(\mathbb{T}_{\frac{2\pi}{a}}; \mathbf{X})} + \|\mathbf{u}_\delta^\varepsilon - \mathbf{u}_\delta\|_{L^2(\mathbb{T}_{\frac{2\pi}{a}}; \mathbf{X}^1)} &\leq C\varepsilon|\delta| \\ \|\partial_x^2(\mathbf{u}_\delta^\varepsilon - \mathbf{u}_\delta)\|_{L^2(\mathbb{T}_{\frac{2\pi}{a}}; \mathbf{X})} + \|\partial_t(\mathbf{u}_\delta^\varepsilon - \mathbf{u}_\delta)\|_{L^2(\mathbb{T}_{\frac{2\pi}{a}}; \mathbf{X})} &\leq C|\delta|\end{aligned}$$

uniformly for $0 < \varepsilon \leq \varepsilon_2$ and $|\delta| \leq \delta_2$.

(iii) There exists a neighborhood \mathcal{O} of ${}^\top(\eta, \omega, u) = {}^\top(0, 0, 0)$ in $\mathbb{R} \times \mathbb{R} \times \mathcal{Y}_a$ such that if ${}^\top(\eta, \omega, u) \in \mathcal{O}$ is a solution of (4.4), then

$${}^\top(\eta, \omega, u) \in \{{}^\top(\eta, \omega, 0) \in \mathcal{O}\} \cup \{{}^\top(\eta_\delta^\varepsilon, \omega_\delta^\varepsilon, u_\delta^\varepsilon(\cdot + \tau)); |\delta| \leq \delta_1, -\frac{\pi}{a} \leq \tau < \frac{\pi}{a}\}$$

where $\{\eta_\delta^\varepsilon, \omega_\delta^\varepsilon, u_\delta^\varepsilon\}$ is the solution branch obtained in (i).

The stability of the bifurcating solution u_δ^ε is also an important question; we shall investigate it in section 7.

To prove Theorem 4.2, we employ the spectral properties of $-B^\varepsilon$ which was investigated in [11]. Some of them will be summarized in section 5 which will then be used in sections 6 and 7.

In what follows we assume that \mathcal{R}_1 is in the interval $[\frac{1}{2}\mathcal{R}_{1,c}, \frac{3}{2}\mathcal{R}_{1,c}]$.

5 Spectrum of B^ε

In this section we summarize the results on the spectrum of B^ε obtained in [11].

We begin with the following lemma.

Lemma 5.1 ([11, Lemma 5.1]) *Let β and T be positive constants. Suppose that $u_0 \in X^1$ and let $F = {}^\top(f, \mathbf{g}, h, k) \in L^2(0, T; X)$. Then there is a unique solution $u(t) \in L^2(0, T; Y) \cap H^1(0, T; X)$ of the problem*

$$\beta \partial_t u + L_{\mathcal{R}_{1,c}^\varepsilon}^\varepsilon u = F, \quad (5.1)$$

$$u(0) = u_0 \quad (5.2)$$

with estimate

$$\begin{aligned} & \beta \|\|u(t)\|\|_{\varepsilon, X^1}^2 + \int_0^t (\|\|\partial_x u\|\|_\varepsilon^2 + \beta^2 \varepsilon^2 \|\|\partial_t u\|\|_\varepsilon^2 + \varepsilon^2 \|\|\partial_x^2 u\|\|_2^2 + \beta^2 \varepsilon^6 \|\|\partial_x \partial_t \phi\|\|_2^2) ds \\ & \leq C e^{\frac{C}{\beta} t} \left\{ \|\|u_0\|\|_{\varepsilon, X^1}^2 + \int_0^t (\varepsilon^2 \|f\|_2^2 + |\operatorname{Re}(\mathbf{F}, \mathbf{u})| + \varepsilon^2 \|\mathbf{F}\|_2^2 + \varepsilon^6 \|\partial_x f\|_2^2) ds \right\} \end{aligned}$$

uniformly for $0 \leq t \leq T$ and $0 < \varepsilon \leq \varepsilon_1$ and β . Furthermore, if $\beta \geq \beta_1$, then the estimate

$$\|\|u\|\|_{\varepsilon, \mathcal{Y}(0,T)} \leq C \left\{ \|\|u_0\|\|_{\varepsilon, X^1} + \|\|F\|\|_{\varepsilon, \mathcal{X}(0,T)} \right\}$$

holds uniformly for $0 < \varepsilon \leq \varepsilon_1$ with a constant depending only on β_1 .

We denote by $\mathcal{V}_\omega^\varepsilon(t)$ the restriction of the semigroup $e^{-\frac{at}{a^\varepsilon(1+\omega)} L_{\mathcal{R}_{1,c}^\varepsilon}^\varepsilon}$ to the space X^1 :

$$\mathcal{V}_\omega^\varepsilon(t) = e^{-\frac{at}{a^\varepsilon(1+\omega)} L_{\mathcal{R}_{1,c}^\varepsilon}^\varepsilon} |_{X^1}.$$

As for the decay property of $\mathcal{V}_\omega^\varepsilon(t)$, we have the following result.

We define the projections P^ε and Q^ε by

$$P^\varepsilon = P_+^\varepsilon + P_-^\varepsilon \text{ and } Q^\varepsilon = I - P^\varepsilon, .$$

respectively. Since $\pm ia^\varepsilon$ are simple eigenvalues of $-L_{\mathcal{R}_{1,c}^\varepsilon}^\varepsilon$, we have

$$P_\pm^\varepsilon \mathcal{V}_\omega^\varepsilon(t) = \mathcal{V}_\omega^\varepsilon(t) P_\pm^\varepsilon = e^{\pm \frac{iat}{1+\omega}} P_\pm^\varepsilon.$$

The Q^ε part of $\mathcal{V}_\omega^\varepsilon(t)$ has the following decay property.

Lemma 5.2 ([11, Lemma 4.3]) *There exist positive constants ε_1 and κ_1 such that the estimate*

$$\|\|\mathcal{V}_0^\varepsilon(t) Q^\varepsilon u_0\|\|_{\varepsilon, X^1} \leq C e^{-\kappa_1 t} \|\|u_0\|\|_{\varepsilon, X^1}$$

holds uniformly in $t \geq 0$ and $0 < \varepsilon \leq \varepsilon_1$. Furthermore, if $\operatorname{Re} \lambda > -\kappa_1$, then $(I - e^{-\frac{2\pi}{a}\lambda} \mathcal{V}_0^\varepsilon(\frac{2\pi}{a}))Q^\varepsilon$ has a bounded inverse on $Q^\varepsilon X^1$ and its inverse $[(I - e^{-\frac{2\pi}{a}\lambda} \mathcal{V}_0^\varepsilon(\frac{2\pi}{a}))Q^\varepsilon]^{-1}$ satisfies

$$\|[(I - e^{-\frac{2\pi}{a}\lambda} \mathcal{V}_0^\varepsilon(\frac{2\pi}{a}))Q^\varepsilon]^{-1}\| \leq \frac{C}{1 - e^{-\frac{2\pi}{a}(\operatorname{Re} \lambda + \kappa_1)}}$$

uniformly in $\varepsilon \in (0, \varepsilon_1]$.

We also have the following spectral property of the monodromy operator $\mathcal{V}_\omega^\varepsilon(\frac{2\pi}{a})$.

Lemma 5.3 ([11, Remark 5.6]) *For any $r > 0$, there exists a positive constant $\omega_r = O(r)$ as $r \rightarrow 0$ such that if $|\omega| \leq \omega_r$ then $\mu - \mathcal{V}_\omega^\varepsilon(\frac{2\pi}{a})$ has a bounded inverse on X^1 for $\mu \in \mathbb{C}$ satisfying $|\mu - 1| \geq r$ and $|\mu| \geq e^{-\frac{3}{4}\kappa_1 \frac{2\pi}{a}}$ and $(\mu - \mathcal{V}_\omega^\varepsilon(\frac{2\pi}{a}))^{-1}$ satisfies the estimate*

$$\|(\mu - \mathcal{V}_\omega^\varepsilon(\frac{2\pi}{a}))^{-1}\|_{\varepsilon, X^1} \leq C \left(\frac{1}{r} + \frac{1}{|\mu|} \right) \|F\|_{\varepsilon, X^1}$$

uniformly in $\varepsilon \in (0, \varepsilon_1]$.

As for the operator B^ε , we have the following result on its spectrum ([11]).

Theorem 5.4 ([11, Theorem 4.2]) (i) *If $\varepsilon \in (0, \varepsilon_1]$, then*

$$\rho(-B^\varepsilon) \supset \Sigma_1 \setminus (\cup_{k \in \mathbb{Z}} \{ika^\varepsilon\}).$$

Here $\Sigma_1 = \{\lambda \in \mathbb{C}; \operatorname{Re} \lambda > -\frac{a^\varepsilon}{a} \kappa_1\}$ with κ_1 being the positive number given in Lemma 5.2; and each ika^ε is a semisimple eigenvalue of B^ε and the corresponding eigenspace is spanned by $e^{i(1-k)at} u_+^{(\varepsilon)}$ and $e^{-i(k+1)at} u_-^{(\varepsilon)}$.

(ii) *If $\lambda \in \Sigma_1 \setminus (\cup_{k \in \mathbb{Z}} \{ika^\varepsilon\})$, then*

$$\begin{aligned} & (\lambda + B^\varepsilon)^{-1} F \\ &= \frac{2\pi e^{-\frac{a}{a^\varepsilon} \lambda t}}{a^\varepsilon (1 - e^{-\frac{2\pi}{a^\varepsilon} \lambda})} \left(\left[e^{-\frac{a}{a^\varepsilon} \lambda (\frac{2\pi}{a} - s)} F \right]_{+, \varepsilon} z_+^\varepsilon + \left[e^{-\frac{a}{a^\varepsilon} \lambda (\frac{2\pi}{a} - s)} F \right]_{-, \varepsilon} z_-^\varepsilon \right) \\ &+ \frac{a}{a^\varepsilon} e^{-\frac{a}{a^\varepsilon} \lambda t} \mathcal{V}_0^\varepsilon(t) \left[(I - e^{-\frac{2\pi}{a^\varepsilon} \lambda} \mathcal{V}_0^\varepsilon(\frac{2\pi}{a})) Q^\varepsilon \right]^{-1} \\ &\quad \cdot Q^\varepsilon \int_0^{\frac{2\pi}{a}} e^{-\frac{a}{a^\varepsilon} \lambda (\frac{2\pi}{a} - s)} \mathcal{V}_0^\varepsilon(\frac{2\pi}{a} - s) F(s) ds \end{aligned}$$

$$+ \frac{a}{a^\varepsilon} \int_0^t e^{-\frac{a}{a^\varepsilon} \lambda(t-s)} \mathcal{V}_0^\varepsilon(t-s) F(s) ds.$$

Furthermore,

$$\|(\lambda + B^\varepsilon)^{-1} F\|_{\mathcal{Y}_a} \leq \frac{C}{|1 - e^{-\frac{2\pi}{a^\varepsilon} \lambda}|} \sum_{j=+,-} \left| \left[e^{-\frac{a}{a^\varepsilon} \lambda(\frac{2\pi}{a} - s)} F \right]_{j,\varepsilon} \right| + C \|F\|_{\mathcal{X}_a}.$$

(iii) 0 is a semisimple eigenvalue of B^ε and

$$\mathcal{X}_a = N(B^\varepsilon) \oplus R(B^\varepsilon), \quad N(B^\varepsilon) = \text{span} \{z_+^\varepsilon, z_-^\varepsilon\}.$$

The projection \mathcal{P}^ε is an eigenprojection for the eigenvalue 0 and \mathcal{Q}^ε is a projection on $R(B^\varepsilon)$ along $N(B^\varepsilon)$.

(iv) Let $F = {}^\top(f, \mathbf{F}) \in \mathcal{X}_a$. Then the equation

$$B^\varepsilon u = F, \quad u \in \mathcal{Y}_a$$

is solvable if and only if $F \in \mathcal{Q}^\varepsilon \mathcal{X}_a$, i.e., F satisfies $[F]_{+,\varepsilon} = [F]_{-,\varepsilon} = 0$. If this condition for F is satisfied, then the problem $B^\varepsilon u = F$ with $u \in \mathcal{Q}^\varepsilon \mathcal{Y}_a$ is uniquely solvable and the solution $u \in \mathcal{Q}^\varepsilon \mathcal{Y}_a$ is given by

$$\begin{aligned} u(t) &= \frac{a}{a^\varepsilon} \mathcal{P}^\varepsilon(sF(s)) \\ &+ \frac{a}{a^\varepsilon} \mathcal{V}_0^\varepsilon(t) \left[(I - \mathcal{V}_0^\varepsilon(\frac{a}{a^\varepsilon})) \mathcal{Q}^\varepsilon \right]^{-1} \int_0^{\frac{2\pi}{a}} \mathcal{V}_0^\varepsilon(\frac{2\pi}{a} - s) F(s) ds \\ &+ \frac{a}{a^\varepsilon} \int_0^t \mathcal{V}_0^\varepsilon(t-s) F(s) ds. \end{aligned}$$

Furthermore, u satisfies

$$\|u\|_{\varepsilon, \mathcal{Y}_a} \leq C \left\{ \int_0^{\frac{2\pi}{a}} (\varepsilon^2 \|f\|_2^2 + |\text{Re}(\mathbf{F}, \mathbf{u})| + \varepsilon^2 \|\mathbf{F}\|_2^2 + \varepsilon^6 \|\partial_x f\|_2^2) dt \right\}^{\frac{1}{2}}$$

and

$$\|u\|_{\varepsilon, \mathcal{Y}_a} \leq C \|F\|_{\varepsilon, \mathcal{X}_a}$$

uniformly in $\varepsilon \in (0, \varepsilon_1]$.

To prove the Hopf bifurcation in (4.4), Theorem 5.4 is insufficient; the term $\omega\partial_t u$ on the left-hand side of the \mathcal{Q}^ε part of the equation (4.4) cannot be regarded as a perturbation of $B^\varepsilon u$. To overcome this we shall put the term $\omega\partial_t$ into the principal part, so we define the operator $B^\varepsilon(\omega)$ on \mathcal{X}_a by

$$D(B^\varepsilon(\omega)) = \mathcal{Y}_a, \quad B^\varepsilon(\omega)u = \frac{a^\varepsilon}{a}(1 + \omega)\partial_t u + L_{\mathcal{R}_{1,c}^\varepsilon}^\varepsilon u \quad (u \in D(B^\varepsilon(\omega))).$$

As for $B^\varepsilon(\omega)$, we have the following estimate.

Lemma 5.5 ([11, Lemma 5.6]) *There exist a positive constant ε_1 such that the following assertion holds. For a given $F \in \mathcal{Q}^\varepsilon \mathcal{X}_a$, there exists a unique solution $u \in \mathcal{Q}^\varepsilon \mathcal{Y}_a$ of $B^\varepsilon(\omega)u = F$, and the solution u satisfies the estimate*

$$\|u\|_{\varepsilon, \mathcal{Y}_a} \leq C \left\{ \int_0^{\frac{2\pi}{a}} (\varepsilon^2 \|f\|_2^2 + |\operatorname{Re}(\mathbf{F}, \mathbf{u})| + \varepsilon^2 \|\mathbf{F}\|_2^2 + \varepsilon^6 \|\partial_x f\|_2^2) dt \right\}^{\frac{1}{2}}$$

and

$$\|u\|_{\varepsilon, \mathcal{Y}_a} \leq C \|F\|_{\varepsilon, \mathcal{X}_a}$$

uniformly for $0 < \varepsilon \leq \varepsilon_1$ and $|\omega| \leq \frac{1}{4}$.

6 Proof of Theorem 4.2

In this section we prove Theorem 4.2, i.e., we shall prove that a nontrivial solution branch of (4.4) bifurcates from $\{\eta, u\} = \{0, 0\}$. In what follows, we will use the same letters ε_0 and δ_0 for bounds of the ranges of ε and δ , even when they should be taken suitably smaller than those of the previous ones if no confusion will occur from the context.

6.1 Proof of Theorem 4.2 (i)

We look for a solution u of (4.4) in the form

$$u = \delta(z_0^\varepsilon + \delta U^\varepsilon), \tag{6.1}$$

where δ is a small parameter; $z_0^\varepsilon = 2\operatorname{Re}z_+^\varepsilon$; and U^ε is a real valued function in \mathcal{Y}_a satisfying $[U^\varepsilon]_{+, \varepsilon} = 0$.

We note that if U is real valued, then $[U]_{-, \varepsilon} = \overline{[U]_{+, \varepsilon}}$. Therefore, the condition $[U]_{+, \varepsilon} = 0$ automatically implies the condition $[U]_{-, \varepsilon} = 0$, which in turn implies that $U = \mathcal{Q}^\varepsilon U \in R(B^\varepsilon) = \mathcal{Q}^\varepsilon \mathcal{X}_a$ by Theorem 5.4.

We first collect basic facts concerning \mathcal{P}^ε .

Lemma 6.1 (i) *It holds that $[z_0^\varepsilon]_{+, \varepsilon} = 1$, $[\partial_t z_0^\varepsilon]_{+, \varepsilon} = ia$ and $\mathcal{P}^\varepsilon z_0^\varepsilon = z_0^\varepsilon$.*

(ii) *It holds that $\mathcal{P}_\pm^\varepsilon \partial_t \subset \partial_t \mathcal{P}_\pm^\varepsilon = \pm ia \mathcal{P}_\pm^\varepsilon$ and $\mathcal{P}^\varepsilon \partial_t \subset \partial_t \mathcal{P}^\varepsilon$.*

(iii) *If u is real valued, then $\mathcal{P}^\varepsilon u = 0$ is equivalent to $[u]_{+, \varepsilon} = 0$ that is equivalent to $\operatorname{Re}[u]_{+, \varepsilon} = \operatorname{Im}[u]_{+, \varepsilon} = 0$.*

Lemma 6.1 can be proved by straightforward computations. We omit the proof.

The following estimates will be used to estimate nonlinear terms. We define the bilinear form $M(\cdot, \cdot)$ by

$$M(u_1, u_2) = \begin{pmatrix} 0 \\ \mathbf{M}(\mathbf{u}_1, \mathbf{u}_2) \end{pmatrix}, \quad \mathbf{M}(\mathbf{u}_1, \mathbf{u}_2) = \mathbf{N}(\mathbf{u}_1, \mathbf{u}_2) + \mathbf{N}(\mathbf{u}_2, \mathbf{u}_1)$$

for $u_j = {}^\top(\phi_j, \mathbf{u}_j)$ ($j = 1, 2$).

Lemma 6.2 *The following inequalities hold for $u_j = {}^\top(\phi_j, \mathbf{u}_j) \in \mathcal{Y}_a$ ($j = 1, 2, 3$) uniformly in $0 < \varepsilon \leq \varepsilon_0$:*

(i) $\| [u]_{+, \varepsilon} \| \leq C \| \|u\| \|_{\varepsilon, \mathcal{X}_a},$

(ii) $\| [N(\mathbf{u}_1, \mathbf{u}_2)]_{+, \varepsilon} \| \leq C \| \|u_1\| \|_{\varepsilon, \mathcal{Y}_a} \| \|u_2\| \|_{\varepsilon, \mathcal{Y}_a},$

(iii) $\| \|M(z_0^\varepsilon, u)\| \|_{\varepsilon, \mathcal{X}_a} \leq C \| \|u\| \|_{\varepsilon, \mathcal{Y}_a},$

(iv) $\int_0^{\frac{2\pi}{a}} |(\mathbf{N}(\mathbf{u}_1, \mathbf{u}_2), \mathbf{u}_3)| dt \leq C \| \|u_1\| \|_{\varepsilon, \mathcal{Y}_a} \| \|u_2\| \|_{\varepsilon, \mathcal{Y}_a} \| \|u_3\| \|_{\varepsilon, \mathcal{Y}_a},$

(v) $\int_0^{\frac{2\pi}{a}} \varepsilon^2 \| \mathbf{N}(\mathbf{u}_1, \mathbf{u}_2) \|_2^2 dt \leq C \| \|u_1\| \|_{\varepsilon, \mathcal{Y}_a}^2 \| \|u_2\| \|_{\varepsilon, \mathcal{Y}_a}^2.$

We look for a solution u in the form (6.1) of the equation (4.4) for $\eta = \delta^2 \tilde{\eta}^\varepsilon$ and $\omega = \delta^2 \tilde{\omega}^\varepsilon$. This scaling of η and ω is due to the fact $[N(z_0^\varepsilon)]_{+, \varepsilon} = 0$. We substitute $\{\eta, \omega, u\} = \{\delta^2 \tilde{\eta}^\varepsilon, \delta^2 \tilde{\omega}^\varepsilon, \delta(z_0^\varepsilon + \delta U^\varepsilon)\}$ into (4.4) and divide the resulting equation by δ^2 . We then obtain

$$B^\varepsilon U^\varepsilon + \frac{a^\varepsilon}{a} \delta \tilde{\omega}^\varepsilon (\partial_t z_0^\varepsilon + \delta \partial_t U^\varepsilon) + \delta \tilde{\eta}^\varepsilon K(z_0^\varepsilon + \delta U^\varepsilon) + N(z_0^\varepsilon + \delta U^\varepsilon) = 0. \quad (6.2)$$

We decompose (6.2) into its \mathcal{P}^ε and \mathcal{Q}^ε parts. Applying \mathcal{P}^ε and \mathcal{Q}^ε to (6.2) and noting that $[N(z_0^\varepsilon)]_{+, \varepsilon} = 0$, we see from Lemma 6.1 that

$$\tilde{\eta}^\varepsilon \operatorname{Re}[K z_0^\varepsilon]_{+, \varepsilon} + \delta \tilde{\eta}^\varepsilon \operatorname{Re}[K U^\varepsilon]_{+, \varepsilon} + \operatorname{Re}[M(z_0^\varepsilon, U^\varepsilon) + \delta N(U^\varepsilon)]_{+, \varepsilon} = 0, \quad (6.3)$$

$$a^\varepsilon \tilde{\omega}^\varepsilon + \tilde{\eta}^\varepsilon \operatorname{Im}[K z_0^\varepsilon]_{+, \varepsilon} + \delta \tilde{\eta}^\varepsilon \operatorname{Im}[K U^\varepsilon]_{+, \varepsilon} + \operatorname{Im}[M(z_0^\varepsilon, U^\varepsilon) + \delta N(U^\varepsilon)]_{+, \varepsilon} = 0, \quad (6.4)$$

$$B^\varepsilon(\delta^2 \tilde{\omega}^\varepsilon) U^\varepsilon + \mathcal{Q}^\varepsilon(\delta \tilde{\eta}^\varepsilon K(z_0^\varepsilon + \delta U^\varepsilon) + N(z_0^\varepsilon + \delta U^\varepsilon)) = 0. \quad (6.5)$$

Here, as explained in section 5, we regard the term $\frac{a^\varepsilon}{a} \delta^2 \tilde{\omega}^\varepsilon \partial_t U^\varepsilon$ in (6.5) as a part of the principal part of the equation (6.5).

We define the map $\mathcal{B}^\varepsilon(\omega) : \mathbb{R} \times \mathbb{R} \times \mathcal{Q}^\varepsilon \mathcal{Y}_a \rightarrow \mathbb{R} \times \mathbb{R} \times \mathcal{Q}^\varepsilon \mathcal{X}_a$ by

$$\mathcal{B}^\varepsilon(\omega) = \begin{pmatrix} \operatorname{Re}[K z_0^\varepsilon]_{+, \varepsilon} & 0 & \operatorname{Re}[M(z_0^\varepsilon, \cdot)]_{+, \varepsilon} \\ \operatorname{Im}[K z_0^\varepsilon]_{+, \varepsilon} & a^\varepsilon & \operatorname{Im}[M(z_0^\varepsilon, \cdot)]_{+, \varepsilon} \\ 0 & 0 & B^\varepsilon(\omega) \mathcal{Q}^\varepsilon \end{pmatrix}.$$

The problem (6.3)–(6.5) is then written as

$$\mathcal{B}^\varepsilon(\delta^2 \tilde{\omega}^\varepsilon) \mathcal{U}^\varepsilon = -\mathcal{N}^\varepsilon(\delta; \mathcal{U}^\varepsilon), \quad (6.6)$$

where $\mathcal{U}^\varepsilon = {}^\top(\tilde{\eta}^\varepsilon, \tilde{\omega}^\varepsilon, U^\varepsilon) \in \mathbb{R} \times \mathbb{R} \times \mathcal{Q}^\varepsilon \mathcal{Y}_a$ and

$$\mathcal{N}^\varepsilon(\delta; \mathcal{U}^\varepsilon) = \begin{pmatrix} \delta \tilde{\eta}^\varepsilon \operatorname{Re}[K U^\varepsilon]_{+, \varepsilon} + \delta \operatorname{Re}[N(U^\varepsilon)]_{+, \varepsilon} \\ \delta \tilde{\eta}^\varepsilon \operatorname{Im}[K U^\varepsilon]_{+, \varepsilon} + \delta \operatorname{Im}[N(U^\varepsilon)]_{+, \varepsilon} \\ \delta \tilde{\eta}^\varepsilon \mathcal{Q}^\varepsilon K(z_0^\varepsilon + \delta U^\varepsilon) + \mathcal{Q}^\varepsilon N(z_0^\varepsilon + \delta U^\varepsilon) \end{pmatrix}$$

Let us suppose that $|\tilde{\eta}^\varepsilon| \leq 1$ and $|\tilde{\omega}^\varepsilon| \leq 1$. By Theorem 4.1 we see that $\operatorname{Re}[K z_0^\varepsilon]_{+, \varepsilon} = \operatorname{Re}(\mathbf{K} \mathbf{u}_+, \mathbf{u}_+^*) + O(\varepsilon^2) < 0$ and $a^\varepsilon = a + O(\varepsilon^2) > 0$ if $\varepsilon > 0$ is sufficiently small. Furthermore, Lemma 5.5 implies that $B^\varepsilon(\delta^2 \tilde{\omega}^\varepsilon) : \mathcal{Q}^\varepsilon \mathcal{Y}_a \rightarrow \mathcal{Q}^\varepsilon \mathcal{X}_a$ has a bounded inverse if $\varepsilon > 0$ and δ are sufficiently small. As a result, $\mathcal{B}^\varepsilon(\delta^2 \tilde{\omega}^\varepsilon)$ has a bounded inverse if $0 < \varepsilon \leq \varepsilon_0$ and $\delta^2 \leq \frac{1}{4}$ for some small $\varepsilon_0 > 0$. Therefore, if $0 < \varepsilon \leq \varepsilon_0$ and $|\delta| \leq \delta_0$ for some small $\varepsilon_0, \delta_0 > 0$, then (6.6) is rewritten as

$$\mathcal{U}^\varepsilon = -\mathcal{B}^\varepsilon(\delta^2 \tilde{\omega}^\varepsilon)^{-1} \mathcal{N}^\varepsilon(\delta; \mathcal{U}^\varepsilon). \quad (6.7)$$

We note that if $\tilde{\mathcal{U}} = {}^\top(\tilde{\eta}, \tilde{\omega}, U)$ with $U = \mathcal{Q}^\varepsilon U$, then

$$\mathcal{B}^\varepsilon(\omega)^{-1} \tilde{\mathcal{U}} = \begin{pmatrix} (\mathcal{B}^\varepsilon(\omega)^{-1} \tilde{\mathcal{U}})^1 \\ (\mathcal{B}^\varepsilon(\omega)^{-1} \tilde{\mathcal{U}})^2 \\ (\mathcal{B}^\varepsilon(\omega)^{-1} \tilde{\mathcal{U}})^3 \end{pmatrix},$$

where

$$\begin{aligned}
(\mathcal{B}^\varepsilon(\omega)^{-1}\tilde{\mathcal{U}})^1 &= \frac{1}{\operatorname{Re}[Kz_0^\varepsilon]_{+, \varepsilon}} \left\{ \tilde{\eta} - \operatorname{Re}[M(z_0^\varepsilon, B^\varepsilon(\omega)^{-1}U)]_{+, \varepsilon} \right\}, \\
(\mathcal{B}^\varepsilon(\omega)^{-1}\tilde{\mathcal{U}})^2 &= \frac{1}{a^\varepsilon} \tilde{\omega} - \frac{\operatorname{Im}[Kz_0^\varepsilon]_{+, \varepsilon}}{a^\varepsilon} (\mathcal{B}^\varepsilon(\omega)^{-1}\tilde{\mathcal{U}})^1 \\
&\quad - \frac{1}{a^\varepsilon} \operatorname{Im}[M(z_0^\varepsilon, B^\varepsilon(\omega)^{-1}U)]_{+, \varepsilon}, \\
(\mathcal{B}^\varepsilon(\omega)^{-1}\tilde{\mathcal{U}})^3 &= (B^\varepsilon(\omega)\mathcal{Q}^\varepsilon)^{-1}U.
\end{aligned}$$

To solve (6.6), we introduce a norm of $\mathbb{R} \times \mathbb{R} \times \mathcal{Y}_a$. We define the norm $\|\mathcal{U}\|_{\varepsilon, \mathbb{R} \times \mathbb{R} \times \mathcal{Y}_a}$ of $\mathcal{U} = {}^\top(\hat{\eta}, \hat{\omega}, U) \in \mathbb{R} \times \mathbb{R} \times \mathcal{Y}_a$ by

$$\|\mathcal{U}\|_{\varepsilon, \mathbb{R} \times \mathbb{R} \times \mathcal{Y}_a} = |\hat{\eta}| + |\hat{\omega}| + \|U\|_{\varepsilon, \mathcal{Y}_a}.$$

We construct approximate solutions $\{\mathcal{U}_n^\varepsilon\}_{n=0}^\infty$ in the following way. Consider first the problem

$$B^\varepsilon U_0^\varepsilon + N(z_0^\varepsilon) = 0. \quad (6.8)$$

Since $[N(z_0^\varepsilon)]_{+, \varepsilon} = 0$, i.e., $N(z_0^\varepsilon) = \mathcal{Q}^\varepsilon N(z_0^\varepsilon)$, it follows from Theorems 4.1 and 5.4 and Lemma 6.2 that (6.8) has a unique solution $U_0^\varepsilon \in \mathcal{Q}^\varepsilon \mathcal{Y}_a$ with estimate $\|U_0^\varepsilon\|_{\varepsilon, \mathcal{Y}_a} \leq C \|N(z_0^\varepsilon)\|_{\varepsilon, \mathcal{X}_a} \leq C$ uniformly for $0 < \varepsilon \leq \varepsilon_0$. We also note that $\|\partial_t U_0^\varepsilon\|_{\varepsilon, \mathcal{Y}_a} \leq C \|M(\partial_t z_0^\varepsilon, z_0^\varepsilon)\|_{\varepsilon, \mathcal{X}_a} \leq C$ uniformly for $0 < \varepsilon \leq \varepsilon_0$.

We now define ${}^\top(\tilde{\eta}_0^\varepsilon, \tilde{\omega}_0^\varepsilon)$ by the solution of

$$\tilde{\eta}_0^\varepsilon \operatorname{Re}[Kz_0^\varepsilon]_{+, \varepsilon} + \operatorname{Re}[M(z_0^\varepsilon, U_0^\varepsilon)]_{+, \varepsilon} = 0, \quad (6.9)$$

$$a^\varepsilon \tilde{\omega}_0^\varepsilon + \tilde{\eta}_0^\varepsilon \operatorname{Im}[Kz_0^\varepsilon]_{+, \varepsilon} + \operatorname{Im}[M(z_0^\varepsilon, U_0^\varepsilon)]_{+, \varepsilon} = 0. \quad (6.10)$$

Using the estimate of U_0^ε , we have $|\tilde{\eta}_0^\varepsilon| + |\tilde{\omega}_0^\varepsilon| \leq C$ uniformly for $0 < \varepsilon \leq \varepsilon_0$. We thus obtain $\mathcal{U}_0^\varepsilon = {}^\top(\tilde{\eta}_0^\varepsilon, \tilde{\omega}_0^\varepsilon, U_0^\varepsilon)$ which is the solution of

$$\mathcal{B}^\varepsilon(0)\mathcal{U}_0^\varepsilon = -\mathcal{N}^\varepsilon(0; 0) = - \begin{pmatrix} 0 \\ 0 \\ N(z_0^\varepsilon) \end{pmatrix}.$$

We set

$$D_1 = \sup \left\{ \|\mathcal{B}^\varepsilon(\omega)^{-1}\mathcal{N}^\varepsilon(0; 0)\|_{\varepsilon, \mathbb{R} \times \mathbb{R} \times \mathcal{Y}_a} : |\omega| \leq \frac{1}{4}, 0 < \varepsilon \leq \varepsilon_0 \right\}. \quad (6.11)$$

By Theorem 4.1, Lemmas 5.5 and 6.2, we see that $D_1 < \infty$ and $\mathcal{U}_0^\varepsilon$ satisfies

$$\|\mathcal{U}_0^\varepsilon\|_{\varepsilon, \mathbb{R} \times \mathbb{R} \times \mathcal{Y}_a} \leq D_1$$

uniformly in $0 < \varepsilon \leq \varepsilon_0$.

We now define $\mathcal{U}_n^\varepsilon = {}^\top(\tilde{\eta}_n^\varepsilon, \tilde{\omega}_n^\varepsilon, U_n^\varepsilon)$ ($n = 1, 2, \dots$) by

$$\mathcal{U}_n^\varepsilon = -\mathcal{B}^\varepsilon(\delta^2 \tilde{\omega}_{n-1}^\varepsilon)^{-1} \mathcal{N}^\varepsilon(\delta; \mathcal{U}_{n-1}^\varepsilon) \quad (n = 1, 2, \dots).$$

We shall prove that the sequence $\{\mathcal{U}_n^\varepsilon\}_{n=0}^\infty$ converges to a solution $\mathcal{U}^\varepsilon = {}^\top(\tilde{\eta}^\varepsilon, \tilde{\omega}^\varepsilon, U^\varepsilon)$ of (6.6) satisfying $\partial_t U^\varepsilon \in \mathcal{Q}^\varepsilon \mathcal{Y}_a$ with estimates for \mathcal{U}^ε and $\partial_t U^\varepsilon$ uniformly in $0 < \varepsilon \leq \varepsilon_0$ and $|\delta| \leq \delta_0$.

We claim that there exists a positive constant δ_0 such that if $|\delta| \leq \delta_0$, then

$$\|\mathcal{U}_n^\varepsilon\|_{\varepsilon, \mathbb{R} \times \mathbb{R} \times \mathcal{Y}_a} \leq 2D_1 \quad (6.12)$$

for all $n = 0, 1, 2, \dots$.

Indeed, (6.12) can be proved by induction. We have already seen that (6.12) holds for $n = 0$. Suppose that (6.12) holds with n replaced by $n - 1$ ($n \geq 1$). We shall show that (6.12) also holds for n .

We take δ so that $|\delta| \leq 1$ and $D_1 \delta^2 \leq \frac{1}{4}$. Writing

$$\mathcal{U}_n^\varepsilon = -\mathcal{B}^\varepsilon(\delta^2 \tilde{\omega}_{n-1}^\varepsilon)^{-1} \{ \mathcal{N}^\varepsilon(0; 0) + (\mathcal{N}^\varepsilon(\delta; \mathcal{U}_{n-1}^\varepsilon) - \mathcal{N}^\varepsilon(0; 0)) \},$$

we deduce from Theorem 5.4, Lemmas 5.5 and 6.2 that

$$\|\mathcal{U}_n^\varepsilon\|_{\varepsilon, \mathbb{R} \times \mathbb{R} \times \mathcal{Y}_a} \leq D_1 + C|\delta| \left\{ D_1 + (1 + |\delta|)D_1^2 + |\delta|D_1 \|\mathcal{U}_{n-1}^\varepsilon\|_{\varepsilon, \mathcal{Y}_a}^{\frac{1}{2}} \right\},$$

and hence,

$$\|\mathcal{U}_n^\varepsilon\|_{\varepsilon, \mathbb{R} \times \mathbb{R} \times \mathcal{Y}_a} \leq \frac{4}{3}D_1 + C|\delta|D_1 \{ 1 + (1 + |\delta| + |\delta|^3)D_1 \}.$$

Taking δ in such a way that $|\delta| \leq \min \left\{ 1, \frac{1}{4D_1}, \frac{2}{3C(1+3D_1)} \right\}$, we have

$$\|\mathcal{U}_n^\varepsilon\|_{\varepsilon, \mathbb{R} \times \mathbb{R} \times \mathcal{Y}_a} \leq 2D_1.$$

This proves that (6.12) holds for n . We thus conclude by induction that (6.12) holds for all $n = 0, 1, 2, \dots$.

We next estimate $|||\partial_t U_n^\varepsilon|||_{\varepsilon, \mathcal{Y}_a}$. One can verify that $\partial_t U_n^\varepsilon$ satisfies the linear problem

$$\begin{aligned} B^\varepsilon(\delta^2 \tilde{\omega}_{n-1}^\varepsilon) U_n^\varepsilon &= -M(z_0^\varepsilon, \partial_t z_0^\varepsilon) - \delta \tilde{\eta}_{n-1}^\varepsilon \mathcal{Q}^\varepsilon K(\partial_t z_0^\varepsilon + \delta \partial_t U_{n-1}^\varepsilon) \\ &\quad - \delta \mathcal{Q}^\varepsilon (M(z_0^\varepsilon, \partial_t U_{n-1}^\varepsilon) + M(U_{n-1}^\varepsilon, \partial_t z_0^\varepsilon) + \delta M(U_{n-1}^\varepsilon, \partial_t U_{n-1}^\varepsilon)) \end{aligned}$$

and this linear problem has a unique solution in $\mathcal{Q}^\varepsilon \mathcal{Y}_a$. Note that $\mathcal{Q}^\varepsilon M(z_0^\varepsilon, \partial_t z_0^\varepsilon) = M(z_0^\varepsilon, \partial_t z_0^\varepsilon)$ since $[M(z_0^\varepsilon, \partial_t z_0^\varepsilon)]_{+, \varepsilon} = 0$.

We set

$$D_2 = \max\{D_1, \tilde{D}_2\}, \quad (6.13)$$

where $\tilde{D}_2 = \sup \{|||B^\varepsilon(\omega)^{-1} M(z_0^\varepsilon, \partial_t z_0^\varepsilon)|||_{\varepsilon, \mathcal{Y}_a} : |\omega| \leq \frac{1}{4}, 0 < \varepsilon \leq \varepsilon_0\}$.

As above, one can prove

$$|||\partial_t U_n^\varepsilon|||_{\varepsilon, \mathcal{Y}_a} \leq 2D_2 \quad (6.14)$$

for all $n = 0, 1, 2, \dots$. Indeed, this is true for $n = 0$. Suppose that (6.14) holds for n replaced by $n-1$ ($n \geq 1$). We then see from (6.12), Lemmas 5.5, 6.1 and 6.2 that

$$|||\partial_t U_n^\varepsilon|||_{\varepsilon, \mathcal{Y}_a} \leq D_2 + C|\delta| \left\{ D_2 + |\delta| D_2^2 + D_2^{\frac{1}{2}} \left(1 + D_2^{\frac{1}{2}}\right) |||\partial_t U_n^\varepsilon|||_{\varepsilon, \mathcal{Y}_a} \right\},$$

and hence, by the same argument as above, we see that (6.14) also holds for n if $|\delta| \leq \delta_0$ for some small $\delta_0 > 0$. We thus conclude that (6.14) holds for all $n = 0, 1, 2, \dots$.

We next show that $\{\mathcal{U}_n^\varepsilon\}_{n \geq 0}$ is a Cauchy sequence in $\mathbb{R} \times \mathbb{R} \times \mathcal{Q}^\varepsilon \mathcal{Y}_a$. Since $\mathcal{U}_{n+1}^\varepsilon - \mathcal{U}_n^\varepsilon$ satisfies

$$\begin{aligned} \mathcal{B}^\varepsilon(\delta^2 \tilde{\omega}_n^\varepsilon)(\mathcal{U}_{n+1}^\varepsilon - \mathcal{U}_n^\varepsilon) &= -\frac{a^\varepsilon}{a} \delta^2 (\tilde{\omega}_n^\varepsilon - \tilde{\omega}_{n-1}^\varepsilon) \begin{pmatrix} 0 \\ 0 \\ \partial_t U_n^\varepsilon \end{pmatrix} \\ &\quad - (\mathcal{N}^\varepsilon(\delta; \mathcal{U}_n^\varepsilon) - \mathcal{N}^\varepsilon(\delta; \mathcal{U}_{n-1}^\varepsilon)), \end{aligned}$$

we see from (6.12) and (6.14) that

$$\begin{aligned} &|||\mathcal{U}_{n+1}^\varepsilon - \mathcal{U}_n^\varepsilon|||_{\varepsilon, \mathbb{R} \times \mathbb{R} \times \mathcal{Y}_a} \\ &\leq C|\delta|(1 + |\delta|D_1 + |\delta|D_2) |||\mathcal{U}_n^\varepsilon - \mathcal{U}_{n-1}^\varepsilon|||_{\varepsilon, \mathbb{R} \times \mathbb{R} \times \mathcal{Y}_a} \\ &\quad + C\delta^2 D_1^{\frac{1}{2}} |||\mathcal{U}_n^\varepsilon - \mathcal{U}_{n-1}^\varepsilon|||_{\varepsilon, \mathbb{R} \times \mathbb{R} \times \mathcal{Y}_a}^{\frac{1}{2}} |||\mathcal{U}_{n+1}^\varepsilon - \mathcal{U}_n^\varepsilon|||_{\varepsilon, \mathbb{R} \times \mathbb{R} \times \mathcal{Y}_a}^{\frac{1}{2}}, \end{aligned}$$

from which we have

$$|||\mathcal{U}_{n+1}^\varepsilon - \mathcal{U}_n^\varepsilon|||_{\varepsilon, \mathbb{R} \times \mathbb{R} \times \mathcal{Y}_a} \leq C|\delta| |||\mathcal{U}_n^\varepsilon - \mathcal{U}_{n-1}^\varepsilon|||_{\varepsilon, \mathbb{R} \times \mathbb{R} \times \mathcal{Y}_a}$$

uniformly in $0 < \varepsilon \leq \varepsilon_0$, $|\delta| \leq \delta_0$ and $n = 1, 2, \dots$. This implies that $\{\mathcal{U}_n^\varepsilon\}_{n \geq 0}$ is a Cauchy sequence in $\mathbb{R} \times \mathbb{R} \times \mathcal{Q}^\varepsilon \mathcal{Y}_a$ if δ_0 is taken so that $0 < \delta_0 \leq 1/(2C)$.

Consequently, we deduce that there exists $\mathcal{U}^\varepsilon = {}^\top(\tilde{\eta}^\varepsilon, \tilde{\omega}^\varepsilon, U^\varepsilon) \in \mathbb{R} \times \mathbb{R} \times \mathcal{Q}^\varepsilon \mathcal{Y}_a$ with $\partial_t U^\varepsilon \in \mathcal{Q}^\varepsilon \mathcal{Y}_a$ such that

$$\begin{aligned} \mathcal{U}_n^\varepsilon &\rightarrow \mathcal{U}^\varepsilon \text{ strongly in } \mathbb{R} \times \mathbb{R} \times \mathcal{Q}^\varepsilon \mathcal{Y}_a, \\ \partial_t \mathcal{U}_n^\varepsilon &\rightarrow \partial_t \mathcal{U}^\varepsilon \text{ weakly in } \mathcal{Q}^\varepsilon \mathcal{Y}_a. \end{aligned}$$

Furthermore, this \mathcal{U}^ε is a solution of (6.6) and $\mathcal{U}^\varepsilon = {}^\top(\tilde{\eta}^\varepsilon, \tilde{\omega}^\varepsilon, U^\varepsilon)$ satisfies the estimates

$$|||\mathcal{U}^\varepsilon|||_{\varepsilon, \mathbb{R} \times \mathbb{R} \times \mathcal{Y}_a} \leq 2D_1, \quad |||\partial_t \mathcal{U}^\varepsilon|||_{\varepsilon, \mathcal{Y}_a} \leq 2D_2 \quad (6.15)$$

uniformly for $0 < \varepsilon \leq \varepsilon_0$ and $|\delta| \leq \delta_0$.

Similarly to above, one can show

$$|||\mathcal{U}^\varepsilon - \mathcal{U}_0^\varepsilon|||_{\varepsilon, \mathbb{R} \times \mathbb{R} \times \mathcal{Y}_a} \leq C|\delta| \quad (6.16)$$

uniformly in $0 < \varepsilon \leq \varepsilon_0$ and $|\delta| \leq \delta_0$. This implies that $\tilde{\eta}^\varepsilon$ and $\tilde{\omega}^\varepsilon$ take the forms

$$\tilde{\eta}^\varepsilon = \tilde{\eta}_0^\varepsilon + \tilde{\eta}_1^\varepsilon(\delta)\delta, \quad \tilde{\omega}^\varepsilon = \tilde{\omega}_0^\varepsilon + \tilde{\omega}_1^\varepsilon(\delta)\delta \quad (6.17)$$

for some $\tilde{\eta}_1^\varepsilon(\delta)$ and $\tilde{\omega}_1^\varepsilon(\delta)$ satisfying $|\tilde{\eta}_1^\varepsilon(\delta)| + |\tilde{\omega}_1^\varepsilon(\delta)| \leq C$ uniformly for $0 < \varepsilon \leq \varepsilon_0$ and $|\delta| \leq \delta_0$. This completes the proof of Theorem 4.2 (i). \square

6.2 Proof of Theorem 4.2 (ii)

We next prove Theorem 4.2 (ii). In this subsection we denote the solution branch obtained in the previous subsection by $\{\eta_\delta^\varepsilon, \omega_\delta^\varepsilon, u_\delta^\varepsilon\}$ with

$$\eta_\delta^\varepsilon = \delta^2 \tilde{\eta}_\delta^\varepsilon, \quad \omega_\delta^\varepsilon = \delta^2 \tilde{\omega}_\delta^\varepsilon, \quad u_\delta^\varepsilon = \delta(z_0^\varepsilon + \delta U_\delta^\varepsilon),$$

where $\{\tilde{\eta}_\delta^\varepsilon, \tilde{\omega}_\delta^\varepsilon, U_\delta^\varepsilon\}$ is the solution of (6.3)–(6.5) obtained in the previous subsection. Therefore, $\tilde{\eta}_\delta^\varepsilon$ and $\tilde{\omega}_\delta^\varepsilon$ are written in the forms in (6.17).

We first consider the behavior of the time periodic solution branch for the incompressible problem (3.5) as $\delta \rightarrow 0$.

Let $\{\eta_\delta, \omega_\delta, u_\delta\}$ be the time periodic solution branch for (3.5) given in Proposition 3.4 which takes the form

$$\eta_\delta = \tilde{\eta}_0 \delta^2 + \tilde{\eta}_1(\delta) \delta^3, \quad \omega_\delta = \tilde{\omega}_0 \delta^2 + \tilde{\omega}_1(\delta) \delta^3, \quad u_\delta = \delta(z_0 + \delta U_\delta).$$

Since $[\mathbf{N}(z_0)]_+ = 0$, there exists a unique solution $\mathbf{U}_0 \in L^2(\mathbb{T}_{\frac{2\pi}{a}}; \mathbb{Y}_\sigma) \cap H^1(\mathbb{T}_{\frac{2\pi}{a}}; \mathbb{X}_\sigma)$ of

$$\partial_t \mathbf{U}_0 + \mathbb{L}_{\mathcal{R}_{1,c}} \mathbf{U}_0 + \mathbb{P} \mathbf{N}(z_0) = \mathbf{0} \quad (6.18)$$

with $[\mathbf{U}_0]_+ = 0$. Let $U_0 = {}^\top(\Phi_0, \mathbf{U}_0) \in \mathcal{Y}_a$ with Φ_0 being the associated pressure of \mathbf{U}_0 . Then $U_0 = {}^\top(\Phi_0, \mathbf{U}_0)$ is the unique solution of

$$\partial_t \begin{pmatrix} 0 \\ \mathbf{U}_0 \end{pmatrix} + L_{\mathcal{R}_{1,c}}^\varepsilon U_0 + N(z_0) = 0 \quad (6.19)$$

with $[\mathbf{U}_0]_+ = 0$.

Similarly to the proof of Theorem 4.2 (i), one can show the following proposition.

Proposition 6.3 *Let $\{\eta_\delta, \omega_\delta, u_\delta\}$ be the time periodic solution branch for (3.5) given in Proposition 3.4 and let $U_0 = {}^\top(\Phi_0, \mathbf{U}_0) \in \mathcal{Y}_a$ be the unique solution of (6.19) with $[\mathbf{U}_0]_+ = 0$. Then $\{\eta_0, \omega_0, U_0\}$ satisfies*

$$\tilde{\eta}_0 \operatorname{Re}[\mathbf{K} z_0]_+ + \operatorname{Re}[\mathbf{M}(z_0, \mathbf{U}_0)]_+ = 0, \quad (6.20)$$

$$a \tilde{\omega}_0 + \tilde{\eta}_0 \operatorname{Im}[\mathbf{K} z_0]_+ + \operatorname{Im}[\mathbf{M}(z_0, \mathbf{U}_0)]_+ = 0 \quad (6.21)$$

and

$$|\tilde{\eta}_\delta - \tilde{\eta}_0| + |\tilde{\omega}_\delta - \tilde{\omega}_0| + \|U_\delta - U_0\|_{\mathcal{Y}_a} \leq C|\delta|$$

uniformly for $|\delta| \leq \delta_0$.

We now estimate the difference between the solution branches for the incompressible system and the artificial compressible system.

Let $\{\eta_\delta, \omega_\delta, u_\delta\}$ be the time periodic solution branch of the incompressible problem (3.5) given in Proposition 3.4. Then $\{\eta_\delta, \omega_\delta, u_\delta\}$ satisfies

$$(1 + \omega_\delta) \partial_t \begin{pmatrix} 0 \\ \mathbf{u}_\delta \end{pmatrix} + L_{\mathcal{R}_{1,c}}^\varepsilon u_\delta + \eta_\delta K u_\delta + N(u_\delta) = 0. \quad (6.22)$$

We set $\eta_\delta = \delta^2 \tilde{\eta}_\delta$ and $\omega_\delta = \delta^2 \tilde{\omega}_\delta$. Then $\{\tilde{\eta}_\delta, \tilde{\omega}_\delta, U_\delta\}$ satisfies the equation

$$(1 + \delta^2 \tilde{\omega}_\delta) \partial_t \begin{pmatrix} 0 \\ \mathbf{U}_\delta \end{pmatrix} + L_{\mathcal{R}_{1,c}}^\varepsilon U_\delta + \delta \tilde{\omega}_\delta \partial_t \begin{pmatrix} 0 \\ \mathbf{z}_0 \end{pmatrix} + \delta \tilde{\eta}_\delta K(z_0 + \delta U_\delta) + N(z_0 + \delta U_\delta) = 0. \quad (6.23)$$

As in the proof of Theorem 4.2 (i), we see from (6.23) that

$$\tilde{\eta}_\delta \operatorname{Re}[\mathbf{K} \mathbf{z}_0]_+ + \delta \tilde{\eta}_\delta \operatorname{Re}[\mathbf{K} \mathbf{U}_\delta]_+ + \operatorname{Re}[\mathbf{M}(z_0, \mathbf{U}_\delta) + \delta \mathbf{N}(U_\delta)]_+ = 0, \quad (6.24)$$

$$a \tilde{\omega}_\delta + \tilde{\eta}_\delta \operatorname{Im}[\mathbf{K} \mathbf{z}_0]_+ + \delta \tilde{\eta}_\delta \operatorname{Im}[\mathbf{K} \mathbf{U}_\delta]_+ + \operatorname{Im}[\mathbf{M}(z_0, \mathbf{U}_\delta) + \delta \mathbf{N}(U_\delta)]_+ = 0. \quad (6.25)$$

Applying \mathcal{Q}^ε to (6.23), we have

$$B^\varepsilon(\delta^2 \tilde{\omega}_\delta^\varepsilon) \mathcal{Q}^\varepsilon U_\delta + \mathcal{Q}^\varepsilon H^\varepsilon(\delta; \tilde{\omega}_\delta^\varepsilon, \tilde{\omega}_\delta, U_\delta) + \mathcal{Q}^\varepsilon (\delta \tilde{\eta}_\delta K(z_0 + \delta U_\delta) + N(z_0 + \delta U_\delta)) = 0. \quad (6.26)$$

Here $H^\varepsilon(\delta; \tilde{\omega}_\delta^\varepsilon, \tilde{\omega}_\delta, U_\delta)$ is a function of $\tilde{\omega}_\delta^\varepsilon$, $\tilde{\omega}_\delta$ and $U_\delta = {}^\top(\Phi_\delta, \mathbf{U}_\delta)$ with $\hat{\mathcal{Q}}_0 \mathbf{U}_\delta = \mathbf{0}$ given by

$$H^\varepsilon(\delta; \tilde{\omega}_\delta^\varepsilon, \tilde{\omega}_\delta, U_\delta) = -\frac{a^\varepsilon - a}{a} (1 + \delta^2 \tilde{\omega}_\delta^\varepsilon) \begin{pmatrix} 0 \\ \partial_t \mathbf{U}_\delta \end{pmatrix} - \delta^2 (\tilde{\omega}_\delta^\varepsilon - \tilde{\omega}_\delta) \begin{pmatrix} 0 \\ \partial_t \mathbf{U}_\delta \end{pmatrix} - \frac{a^\varepsilon}{a} (1 + \delta^2 \tilde{\omega}_\delta^\varepsilon) \begin{pmatrix} \partial_t \Phi_\delta \\ \mathbf{0} \end{pmatrix} + \delta \tilde{\omega}_\delta \begin{pmatrix} 0 \\ \partial_t \mathbf{z}_0 \end{pmatrix} - (\mathcal{R}_{1,c}^\varepsilon - \mathcal{R}_{1,c}) K U_\delta.$$

We set $\mathcal{U}_\delta = {}^\top(\tilde{\eta}_\delta, \tilde{\omega}_\delta, U_\delta)$. It then follows from (6.3)–(6.5) and (6.24)–(6.26) that

$$\mathcal{B}^\varepsilon(\delta^2 \tilde{\omega}_\delta^\varepsilon) (\mathcal{U}_\delta^\varepsilon - \mathcal{U}_\delta) = -\mathcal{M}^\varepsilon(\delta; \mathcal{U}_\delta^\varepsilon, \mathcal{U}_\delta),$$

Here

$$\mathcal{M}^\varepsilon(\delta; \mathcal{U}^\varepsilon, \mathcal{U}_\delta) = \begin{pmatrix} \mathcal{M}^{\varepsilon,1}(\delta; \mathcal{U}_\delta^\varepsilon, \mathcal{U}_\delta) \\ \mathcal{M}^{\varepsilon,2}(\delta; \mathcal{U}_\delta^\varepsilon, \mathcal{U}_\delta) \\ \mathcal{Q}^\varepsilon H^\varepsilon(\delta; \tilde{\omega}_\delta^\varepsilon, \tilde{\omega}_\delta, U_\delta) + \mathcal{M}^{\varepsilon,3}(\delta; \mathcal{U}_\delta^\varepsilon, \mathcal{U}_\delta) \end{pmatrix},$$

where

$$\begin{aligned} \mathcal{M}^{\varepsilon,1}(\delta; \mathcal{U}_\delta^\varepsilon, \mathcal{U}_\delta) &= \tilde{\eta}_\delta \operatorname{Re}\{[K z_0^\varepsilon]_{+,\varepsilon} - [\mathbf{K} \mathbf{z}_0]_{+}\} + \delta \operatorname{Re}\{\tilde{\eta}_\delta^\varepsilon [K U_\delta^\varepsilon]_{+,\varepsilon} - \tilde{\eta}_\delta [\mathbf{K} \mathbf{U}_\delta]_{+}\} \\ &\quad + \operatorname{Re}\{[M(z_0^\varepsilon, U_\delta)]_{+,\varepsilon} - [\mathbf{M}(z_0, \mathbf{U}_\delta)]_{+}\} \\ &\quad + \delta \operatorname{Re}\{[N(U_\delta^\varepsilon)]_{+,\varepsilon} - [\mathbf{N}(U_\delta)]_{+}\} \end{aligned}$$

$$\begin{aligned}
\mathcal{M}^{\varepsilon,2}(\delta; \mathcal{U}^\varepsilon, \mathcal{U}) &= (a^\varepsilon - a)\tilde{\omega}_\delta \\
&\quad + \tilde{\eta}_\delta \text{Im}\{[Kz_0^\varepsilon]_{+,\varepsilon} - [\mathbf{K}z_0]_{+}\} + \delta \text{Im}\{\tilde{\eta}_\delta^\varepsilon [KU_\delta^\varepsilon]_{+,\varepsilon} - \tilde{\eta}_\delta [\mathbf{K}U_\delta]_{+}\} \\
&\quad + \text{Im}\{[M(z_0^\varepsilon, U_\delta)]_{+,\varepsilon} - [\mathbf{M}(z_0, \mathbf{U}_\delta)]_{+}\} \\
&\quad + \delta \text{Im}\{[N(U_\delta^\varepsilon)]_{+,\varepsilon} - [\mathbf{N}(\mathbf{U}_\delta)]_{+}\} \\
\mathcal{M}^{\varepsilon,3}(\delta; \mathcal{U}^\varepsilon, \mathcal{U}) &= \mathcal{Q}^\varepsilon \left\{ \delta(\tilde{\eta}_\delta^\varepsilon Kz_0^\varepsilon - \tilde{\eta}_\delta Kz_0) + \delta^2(\tilde{\eta}_\delta^\varepsilon KU_\delta^\varepsilon - \tilde{\eta}_\delta KU_\delta) \right. \\
&\quad \left. + N(z_0^\varepsilon + \delta U_\delta^\varepsilon) - N(z_0 + \delta U_\delta) \right\}.
\end{aligned}$$

To estimate $\mathcal{U}_\delta^\varepsilon - \mathcal{U}_\delta$, we use the following lemma.

Lemma 6.4 *Let $u_j = {}^\top(\phi_j, \mathbf{u}_j)$, $\tilde{u}_j = {}^\top(\tilde{\phi}_j, \tilde{\mathbf{u}}_j)$, $j = 1, 2$. Then the following estimates hold uniformly for $0 < \varepsilon \leq \varepsilon_0$:*

- (i) $||[u_1]_{+,\varepsilon} - [\mathbf{u}_2]_{+}| \leq C \left\{ \varepsilon^2 \|\phi_1\|_{L^2(\mathbb{T}_{\frac{2\pi}{a}}; L^2(\Omega))} + \varepsilon^2 \|\mathbf{u}_2\|_{L^2(\mathbb{T}_{\frac{2\pi}{a}}; (\mathbf{X}^1)^*)} \right. \\ \left. + \|\mathbf{u}_1 - \mathbf{u}_2\|_{L^2(\mathbb{T}_{\frac{2\pi}{a}}; (\mathbf{X}^1)^*)} \right\},$
- (ii) $||[N(u_1, u_2)]_{+,\varepsilon} - [\mathbf{N}(\tilde{\mathbf{u}}_1, \tilde{\mathbf{u}}_2)]_{+}| \\ \leq C \left\{ \varepsilon^2 \|\|u_1\|\|_{\varepsilon, \mathcal{Y}_a} \|\|u_2\|\|_{\varepsilon, \mathcal{Y}_a} + \|\|u_1\|\|_{\varepsilon, \mathcal{Y}_a} \|\|u_2 - \tilde{u}_2\|\|_{\varepsilon, \mathcal{Y}_a} \right. \\ \left. + \|\|u_1 - \tilde{u}_1\|\|_{\varepsilon, \mathcal{Y}_a} \|\|\tilde{u}_2\|\|_{\varepsilon, \mathcal{Y}_a} \right\},$
- (iii) $|||\mathcal{Q}^\varepsilon H^\varepsilon(\delta; \tilde{\omega}_\delta^\varepsilon, \tilde{\omega}_\delta, U_\delta)|||_{\varepsilon, \mathcal{X}_a} \leq C \left\{ \varepsilon(1 + |\delta| |\tilde{\omega}_\delta| + \delta^2 |\tilde{\omega}_\delta^\varepsilon|) + \delta^2 |\tilde{\omega}_\delta^\varepsilon - \tilde{\omega}_\delta| \right\}.$

We will give a proof of Lemma 6.4 in the end of this section.

Noting that $Ku = {}^\top(0, \mathbf{K}u)$ for $u = {}^\top(\phi, \mathbf{u})$, we apply Theorems 4.1, 4.2 (i), Lemmas 5.5, 6.1, 6.2 and 6.4 to obtain

$$\begin{aligned}
&|\tilde{\eta}_\delta^\varepsilon - \tilde{\eta}_\delta| + |\tilde{\omega}_\delta^\varepsilon - \tilde{\omega}_\delta| \\
&\leq C \left\{ \varepsilon^2 + |\delta| (|\tilde{\eta}_\delta^\varepsilon - \tilde{\eta}_\delta| + |\tilde{\omega}_\delta^\varepsilon - \tilde{\omega}_\delta|) + \|\|U_\delta^\varepsilon - U_\delta\|\|_{\varepsilon, \mathcal{Y}_a} \right\}, \tag{6.27}
\end{aligned}$$

and

$$\|\|\mathcal{Q}^\varepsilon(U_\delta^\varepsilon - U_\delta)\|\|_{\varepsilon, \mathcal{Y}_a} \leq C \left\{ \varepsilon + |\delta| (|\tilde{\eta}_\delta^\varepsilon - \tilde{\eta}_\delta| + |\tilde{\omega}_\delta^\varepsilon - \tilde{\omega}_\delta|) + |\delta| \|\|U_\delta^\varepsilon - U_\delta\|\|_{\varepsilon, \mathcal{Y}_a} \right\} \tag{6.28}$$

uniformly in $0 < \varepsilon \leq \varepsilon_0$ and $|\delta| \leq \delta_0$.

Since $[\mathbf{U}_\delta]_+ = (\mathbf{U}_\delta, \mathbf{z}_+^*) = 0$, $[U_\delta]_{+, \varepsilon} = \varepsilon^2(\Phi_\delta, p_+^{\varepsilon*}) + (\mathbf{U}_\delta, \mathbf{z}_+^{\varepsilon*})$ and $\mathbf{z}_+^{\varepsilon*} = \mathbf{z}_+^* + O(\varepsilon^2)$, we have

$$|[U_\delta]_{+, \varepsilon}| \leq C\{\varepsilon^2\|\Phi_\delta\|_{L^2(\mathbb{T}_{\frac{2\pi}{a}}; L^2)} + \varepsilon^2\|\mathbf{U}_\delta\|_{L^2(\mathbb{T}_{\frac{2\pi}{a}}; (\mathbf{X}^1)^*)}\} \leq C\varepsilon^2\|\mathbf{U}_\delta\|_{\mathcal{X}_a}$$

uniformly for $0 < \varepsilon \leq \varepsilon_0$ and $|\delta| \leq \delta_0$, and hence,

$$|\mathcal{P}^\varepsilon U_\delta| = |2\operatorname{Re}([U_\delta]_{+, \varepsilon} z_+^\varepsilon)| \leq C\varepsilon^2\|\mathbf{U}_\delta\|_{\mathcal{X}_a}|z_+^\varepsilon|.$$

Combining this with $\mathcal{P}^\varepsilon U_\delta^\varepsilon = 0$, we see that

$$\begin{aligned} \||U_\delta^\varepsilon - U_\delta\||_{\varepsilon, \mathcal{Y}_a} &\leq \||\mathcal{Q}^\varepsilon(U_\delta^\varepsilon - U_\delta)\||_{\varepsilon, \mathcal{Y}_a} + \||\mathcal{P}^\varepsilon(U_\delta^\varepsilon - U_\delta)\||_{\varepsilon, \mathcal{Y}_a} \\ &= \||\mathcal{Q}^\varepsilon(U_\delta^\varepsilon - U_\delta)\||_{\varepsilon, \mathcal{Y}_a} + \||\mathcal{P}^\varepsilon U_\delta\||_{\varepsilon, \mathcal{Y}_a} \\ &\leq \||\mathcal{Q}^\varepsilon(U_\delta^\varepsilon - U_\delta)\||_{\varepsilon, \mathcal{Y}_a} + C\varepsilon^2\|\mathbf{U}_\delta\|_{\mathcal{X}_a} \\ &\leq \||\mathcal{Q}^\varepsilon(U_\delta^\varepsilon - U_\delta)\||_{\varepsilon, \mathcal{Y}_a} + C\varepsilon^2. \end{aligned}$$

This, together with (6.27) and (6.28), implies that the estimate

$$\||\mathcal{U}_\delta^\varepsilon - \mathcal{U}_\delta\||_{\varepsilon, \mathbb{R} \times \mathbb{R} \times \mathcal{Y}_a} \leq C\varepsilon \quad (6.29)$$

holds uniformly for $0 < \varepsilon \leq \varepsilon_0$ and $|\delta| \leq \delta_0$.

Before proceeding further, we make an observation. It is not difficult to see that $\|\partial_t^2 U_0\|_{\mathcal{X}_a} \leq C$, from which one can also prove, in a similar manner to above, that

$$\||\partial_t(U_0^\varepsilon - U_0)\||_{\varepsilon, \mathcal{Y}_a} \leq C\varepsilon. \quad (6.30)$$

It remains to show that

$$|\tilde{\eta}_1^\varepsilon(\delta) - \tilde{\eta}_1(\delta)| + |\tilde{\omega}_1^\varepsilon(\delta) - \tilde{\omega}_1(\delta)| \leq C\varepsilon. \quad (6.31)$$

To this end, we shall estimate $\mathcal{V}_\delta^\varepsilon = (\mathcal{U}_\delta^\varepsilon - \mathcal{U}_0^\varepsilon) - (\mathcal{U}_\delta - \mathcal{U}_0)$. Since $\mathcal{V}_\delta^\varepsilon = (\mathcal{U}_\delta^\varepsilon - \mathcal{U}_\delta) - (\mathcal{U}_0^\varepsilon - \mathcal{U}_0)$, we have

$$\begin{aligned} \mathcal{B}^\varepsilon(\delta^2 \tilde{\omega}_\delta^\varepsilon) \mathcal{Q}^\varepsilon \mathcal{V}_\delta^\varepsilon &= -\frac{a^\varepsilon}{a} \delta^2 \tilde{\omega}_\delta^\varepsilon \begin{pmatrix} 0 \\ 0 \\ \mathcal{Q}^\varepsilon \partial_t(U_0^\varepsilon - U_0) \end{pmatrix} \\ &\quad - (\mathcal{M}^\varepsilon(\delta; \mathcal{U}_\delta^\varepsilon, \mathcal{U}_\delta) - \mathcal{M}^\varepsilon(0; \mathcal{U}_0^\varepsilon, \mathcal{U}_0)). \end{aligned}$$

By using (6.30), we have $|||\frac{a^\varepsilon}{a}\delta^2\tilde{\omega}_\delta^\varepsilon\partial_t(U_0^\varepsilon - U_0)|||_{\varepsilon,\mathcal{Y}_a} \leq C\varepsilon|\delta|$. Based on this, together with (6.21) and (6.29), we have, in a similar manner to above,

$$\begin{aligned} & |(\tilde{\eta}_\delta^\varepsilon - \tilde{\eta}_0^\varepsilon) - (\tilde{\eta}_\delta - \tilde{\eta}_0)| + |(\tilde{\omega}_\delta^\varepsilon - \tilde{\omega}_0^\varepsilon) - (\tilde{\omega}_\delta - \tilde{\omega}_0)| \\ & \leq C\{\varepsilon|\delta| + |||(U_\delta^\varepsilon - U_0^\varepsilon) - (U_\delta - U_0)|||_{\varepsilon,\mathcal{Y}_a}\}, \end{aligned}$$

and

$$|||\mathcal{Q}^\varepsilon[(U_\delta^\varepsilon - U_0^\varepsilon) - (U_\delta - U_0)]|||_{\varepsilon,\mathcal{Y}_a} \leq C\varepsilon|\delta|.$$

Since $[\mathbf{U}_\delta - \mathbf{U}_0]_+ = 0$, we see, as above, that

$$\begin{aligned} & |||(U_\delta^\varepsilon - U_0^\varepsilon) - (U_\delta - U_0)|||_{\varepsilon,\mathcal{Y}_a} \\ & \leq C\{|||\mathcal{Q}^\varepsilon[(U_\delta^\varepsilon - U_0^\varepsilon) - (U_\delta - U_0)]|||_{\varepsilon,\mathcal{Y}_a} + \varepsilon^2\|U_\delta - U_0\|_{\mathcal{X}_a}\} \\ & \leq C\{|||\mathcal{Q}^\varepsilon[(U_\delta^\varepsilon - U_0^\varepsilon) - (U_\delta - U_0)]|||_{\varepsilon,\mathcal{Y}_a} + \varepsilon^2|\delta|\}. \end{aligned}$$

We thus conclude that $|||\mathcal{V}^\varepsilon|||_{\varepsilon,\mathbb{R}\times\mathbb{R}\times\mathcal{Y}_a} \leq C\varepsilon|\delta|$ uniformly for $0 < \varepsilon \leq \varepsilon_0$ and $|\delta| \leq \delta_0$. In particular, we obtain (6.31). This completes the proof. \square

6.3 Proof of Theorem 4.2 (iii)

We next prove Theorem 4.2 (iii), i.e., the uniqueness of time periodic solutions up to translations in the time variable t .

We introduce notation. We set $z_0^{\varepsilon,\tau} = 2\text{Re}(e^{ia\tau}z_+^\varepsilon)$ and $(z_0^{\varepsilon*})^\tau = 2\text{Re}(e^{ia\tau}z_+^{\varepsilon*})$ for a constant $\tau \in [-\frac{\pi}{a}, \frac{\pi}{a})$ and define $[u]_{+,\varepsilon}^{(\tau)}$ by $[u]_{+,\varepsilon}^{(\tau)} = \langle u, e^{ia\tau}z_+^{\varepsilon*} \rangle_\varepsilon = e^{-ia\tau}[u]_{+,\varepsilon}$. We note that $\mathcal{P}_\pm^\varepsilon$ can be expressed as

$$\mathcal{P}_\pm^\varepsilon u = \langle u, e^{\pm ia\tau}z_\pm^{\varepsilon*} \rangle_\varepsilon e^{\pm ia\tau}z_\pm^\varepsilon.$$

Proof of Theorem 4.2 (iii). Let $u^\varepsilon \in \mathcal{Y}_a$ be a solution of (4.4) with $\eta = \eta^\varepsilon$ and $\omega = \omega^\varepsilon$. Then, by Theorem 5.4, u^ε is written as

$$u^\varepsilon = u_0^\varepsilon + U^\varepsilon, \quad u_0^\varepsilon = \mathcal{P}^\varepsilon u^\varepsilon \in N(B^\varepsilon), \quad U^\varepsilon \in R(B^\varepsilon) \quad (6.32)$$

Assume first that $u_0^\varepsilon = 0$ which is equivalent to $[u^\varepsilon]_{+,\varepsilon} = 0$. We then find that

$$B^\varepsilon U^\varepsilon + \frac{a^\varepsilon}{a}\omega^\varepsilon\partial_t U^\varepsilon + \eta^\varepsilon K U^\varepsilon + N(U^\varepsilon) = 0,$$

and hence,

$$B^\varepsilon(\omega^\varepsilon)U^\varepsilon + \eta^\varepsilon \mathcal{Q}^\varepsilon K U^\varepsilon + \mathcal{Q}^\varepsilon N(U^\varepsilon) = 0.$$

By using Lemmas 5.5 and 6.1, we see that if $|\omega^\varepsilon| \leq \frac{1}{4}$, then

$$\|U^\varepsilon\|_{\varepsilon, \mathcal{Y}_a} \leq C\{|\eta^\varepsilon| \|U^\varepsilon\|_{\varepsilon, \mathcal{Y}_a} + \|U^\varepsilon\|_{\varepsilon, \mathcal{Y}_a}^{\frac{3}{2}} + \|U^\varepsilon\|_{\varepsilon, \mathcal{Y}_a}^2\}.$$

This implies that there exists a positive constant r_0 such that if $|\eta^\varepsilon| + \|U^\varepsilon\|_{\varepsilon, \mathcal{Y}_a} \leq r_0$, then $\|U^\varepsilon\|_{\varepsilon, \mathcal{Y}_a} \leq \frac{1}{2}\|U^\varepsilon\|_{\varepsilon, \mathcal{Y}_a}$, and therefore, $U^\varepsilon = 0$. We thus conclude that if $|\eta^\varepsilon| + \|u^\varepsilon\|_{\varepsilon, \mathcal{Y}_a} \leq r_0$, $|\omega^\varepsilon| \leq \frac{1}{4}$ and $u^\varepsilon = U^\varepsilon \in R(B^\varepsilon)$, then $u^\varepsilon = 0$.

Assume next that $u_0^\varepsilon \neq 0$ in (6.32). Then there exists a constant $\tau \in [-\frac{\pi}{a}, \frac{\pi}{a})$ such that $u_0^\varepsilon = \mathcal{P}^\varepsilon u^\varepsilon$ is written as $u_0^\varepsilon = \delta z_0^{\varepsilon, \tau}$, where $\delta = |[u^\varepsilon]_{+, \varepsilon}|$ and $z_0^{\varepsilon, \tau} = 2\text{Re}(e^{ia\tau} z_+^\varepsilon)$. It then follows that u^ε is written as

$$u^\varepsilon = \delta z_0^{\varepsilon, \tau} + U^\varepsilon, \quad U^\varepsilon \in R(B^\varepsilon).$$

We substitute this into (4.4) with $\eta = \eta^\varepsilon$ and $\omega = \omega^\varepsilon$. Since $[Kz_0^{\varepsilon, \tau}]_{+, \varepsilon}^{(\tau)} = [Kz_0^\varepsilon]_{+, \varepsilon}$, as in the proof of Theorem 4.2 (i), we obtain

$$\begin{aligned} \delta \eta^\varepsilon \text{Re}[Kz_0^\varepsilon]_{+, \varepsilon} + \eta^\varepsilon \text{Re}[KU^\varepsilon]_{+, \varepsilon}^{(\tau)} + \text{Re}[N(\delta z_0^{\varepsilon, \tau} + U^\varepsilon)]_{+, \varepsilon}^{(\tau)} &= 0, \\ \delta a^\varepsilon \omega^\varepsilon + \delta \eta^\varepsilon \text{Im}[Kz_0^\varepsilon]_{+, \varepsilon} + \eta^\varepsilon \text{Im}[KU^\varepsilon]_{+, \varepsilon}^{(\tau)} + \text{Im}[N(\delta z_0^{\varepsilon, \tau} + U^\varepsilon)]_{+, \varepsilon}^{(\tau)} &= 0, \\ B^\varepsilon(\omega^\varepsilon)U^\varepsilon + \delta \eta^\varepsilon \mathcal{Q}^\varepsilon Kz_0^{\varepsilon, \tau} + \eta^\varepsilon \mathcal{Q}^\varepsilon KU^\varepsilon + \mathcal{Q}^\varepsilon N(\delta z_0^{\varepsilon, \tau} + U^\varepsilon) &= 0. \end{aligned}$$

Setting $\hat{\mathcal{V}}_\delta^\varepsilon = {}^\top(\delta \eta^\varepsilon, \delta \omega^\varepsilon, U^\varepsilon)$, we have

$$\hat{\mathcal{B}}^\varepsilon(\omega^\varepsilon) \hat{\mathcal{V}}_\delta^\varepsilon = - \begin{pmatrix} \eta^\varepsilon \text{Re}[KU^\varepsilon]_{+, \varepsilon}^{(\tau)} + \text{Re}[N(\delta z_0^{\varepsilon, \tau} + U^\varepsilon)]_{+, \varepsilon}^{(\tau)} \\ \eta^\varepsilon \text{Im}[KU^\varepsilon]_{+, \varepsilon}^{(\tau)} + \text{Im}[N(\delta z_0^{\varepsilon, \tau} + U^\varepsilon)]_{+, \varepsilon}^{(\tau)} \\ \eta^\varepsilon \mathcal{Q}^\varepsilon KU^\varepsilon + \mathcal{Q}^\varepsilon N(\delta z_0^{\varepsilon, \tau} + U^\varepsilon) \end{pmatrix},$$

where $\hat{\mathcal{B}}^\varepsilon(\omega) : \mathbb{R} \times \mathbb{R} \times \mathcal{Q}^\varepsilon \mathcal{Y}_a \rightarrow \mathbb{R} \times \mathbb{R} \times \mathcal{Q}^\varepsilon \mathcal{X}_a$ is defined by

$$\hat{\mathcal{B}}^\varepsilon(\omega^\varepsilon) = \begin{pmatrix} \text{Re}[Kz_0^\varepsilon]_{+, \varepsilon} & 0 & 0 \\ \text{Im}[Kz_0^\varepsilon]_{+, \varepsilon} & a^\varepsilon & 0 \\ \mathcal{Q}^\varepsilon Kz_0^{\varepsilon, \tau} & 0 & B^\varepsilon(\omega^\varepsilon) \mathcal{Q}^\varepsilon \end{pmatrix}.$$

Applying Lemmas 5.5 and 6.2, we see that

$$\|\hat{\mathcal{V}}_\delta^\varepsilon\|_{\varepsilon, \mathbb{R} \times \mathbb{R} \times \mathcal{Y}_a} \leq C\{|\delta|^2 + |\delta| \|U^\varepsilon\|_{\varepsilon, \mathcal{Y}_a} + |\eta^\varepsilon| \|U^\varepsilon\|_{\varepsilon, \mathcal{Y}_a} + \|U^\varepsilon\|_{\varepsilon, \mathcal{Y}_a}^{\frac{3}{2}} + \|U^\varepsilon\|_{\varepsilon, \mathcal{Y}_a}^2\}.$$

This implies that there exists a positive number r_1 such that if $|\eta^\varepsilon| + \|U^\varepsilon\|_{\varepsilon, \mathcal{Y}_a} \leq r_1$ and $|\omega^\varepsilon| \leq \frac{1}{4}$, then

$$\|\hat{\mathcal{Y}}_\delta^\varepsilon\|_{\varepsilon, \mathbb{R} \times \mathbb{R} \times \mathcal{Y}_a} \leq C_1 |\delta|^2. \quad (6.33)$$

This suggests us to write

$$\eta^\varepsilon = \delta \hat{\eta}^\varepsilon, \quad \omega^\varepsilon = \delta \hat{\omega}^\varepsilon \quad (6.34)$$

and

$$u = \delta(z_0^{\varepsilon, \tau} + \delta \hat{U}^\varepsilon), \quad \hat{U}^\varepsilon \in R(B^\varepsilon). \quad (6.35)$$

We note that, by (6.33), $\hat{\mathcal{U}}^\varepsilon = {}^\top(\hat{\eta}^\varepsilon, \hat{\omega}^\varepsilon, \hat{U}^\varepsilon)$ with $\hat{\eta}^\varepsilon, \hat{\omega}^\varepsilon$ and \hat{U}^ε given in (6.34) and (6.35) satisfy

$$\|\hat{\mathcal{U}}^\varepsilon\|_{\varepsilon, \mathbb{R} \times \mathbb{R} \times \mathcal{Y}_a} \leq C_1.$$

As in the proof of Theorem 4.2 (i), we also find that $\partial_t \hat{U}^\varepsilon \in \mathcal{Y}_a$ with estimate $\|\partial_t \hat{U}^\varepsilon\|_{\mathcal{Y}_a} \leq C_2$ uniformly in $0 < \varepsilon \leq \varepsilon_0$ and $|\delta| \leq \delta_0$.

We thus find that if $u^\varepsilon = \delta z_0^{\varepsilon, \tau} + U^\varepsilon$, $|\delta| \leq \delta_0$, $|\eta^\varepsilon| + \|U^\varepsilon\|_{\varepsilon, \mathcal{Y}_a} \leq r_1$ and $|\omega^\varepsilon| \leq \frac{1}{4}$, then ${}^\top(\eta^\varepsilon, \omega^\varepsilon, U^\varepsilon)$ belongs to the set

$$\{{}^\top(\delta \hat{\eta}^\varepsilon, \delta \hat{\omega}^\varepsilon, \delta^2 \hat{U}^\varepsilon); \|\hat{\eta}^\varepsilon, \hat{\omega}^\varepsilon, \hat{U}^\varepsilon\|_{\varepsilon, \mathbb{R} \times \mathbb{R} \times \mathcal{Y}_a} \leq C_1, \|\partial_t \hat{U}^\varepsilon\|_{\mathcal{Y}_a} \leq C_2\}.$$

We next claim that if $\hat{\mathcal{U}}_j^\varepsilon = {}^\top(\hat{\eta}_j^\varepsilon, \hat{\omega}_j^\varepsilon, \hat{U}_j^\varepsilon)$, $j = 1, 2$, are solutions of (6.2) with z_0^ε replaced by $z_0^{\varepsilon, \tau}$ for $0 < |\delta| \leq \delta_0$ and $\hat{\mathcal{U}}_j^\varepsilon = {}^\top(\hat{\eta}_j^\varepsilon, \hat{\omega}_j^\varepsilon, \hat{U}_j^\varepsilon)$ satisfy $\|\hat{\mathcal{U}}_j^\varepsilon\|_{\varepsilon, \mathcal{Y}_a} \leq C_1$ and $\|\partial_t \hat{U}_j^\varepsilon\|_{\mathcal{Y}_a} \leq C_2$ ($j = 1, 2$), then it holds $\hat{\mathcal{U}}_1^\varepsilon = \hat{\mathcal{U}}_2^\varepsilon$.

Indeed, we see that $\hat{\mathcal{U}}_j^\varepsilon$ ($j = 1, 2$) satisfy

$$\hat{\mathcal{B}}^\varepsilon(\delta \hat{\omega}_j^\varepsilon) \hat{\mathcal{U}}_j^\varepsilon = -\hat{\mathcal{N}}^\varepsilon(\delta; \hat{\eta}_j^\varepsilon, \hat{U}_j^\varepsilon),$$

where

$$\hat{\mathcal{N}}^\varepsilon(\delta; \hat{\eta}^\varepsilon, \hat{U}^\varepsilon) = \begin{pmatrix} \delta \hat{\eta}^\varepsilon \operatorname{Re}[K \hat{U}^\varepsilon]_{+, \varepsilon}^{(\tau)} + \operatorname{Re}[N(z_0^{\varepsilon, \tau} + \delta \hat{U}^\varepsilon)]_{+, \varepsilon}^{(\tau)} \\ \delta \hat{\eta}^\varepsilon \operatorname{Im}[K \hat{U}^\varepsilon]_{+, \varepsilon}^{(\tau)} + \operatorname{Im}[N(z_0^{\varepsilon, \tau} + \delta \hat{U}^\varepsilon)]_{+, \varepsilon}^{(\tau)} \\ \delta \hat{\eta}^\varepsilon \mathcal{Q}^\varepsilon K \hat{U}^\varepsilon + \mathcal{Q}^\varepsilon N(z_0^{\varepsilon, \tau} + \delta \hat{U}^\varepsilon) \end{pmatrix}$$

It follows that

$$\hat{\mathcal{B}}^\varepsilon(\delta \hat{\omega}_1^\varepsilon)(\hat{\mathcal{U}}_1^\varepsilon - \hat{\mathcal{U}}_2^\varepsilon) = \frac{a^\varepsilon}{a} \delta(\hat{\omega}_2^\varepsilon - \hat{\omega}_1^\varepsilon) \begin{pmatrix} 0 \\ 0 \\ \partial_t \hat{U}_2^\varepsilon \end{pmatrix} + \hat{\mathcal{N}}^\varepsilon(\delta; \hat{\eta}_1^\varepsilon, \hat{U}_1^\varepsilon) - \hat{\mathcal{N}}^\varepsilon(\delta; \hat{\eta}_2^\varepsilon, \hat{U}_2^\varepsilon).$$

By using Lemmas 5.5 and 6.1–6.4, we find that

$$\begin{aligned}
\|\hat{\mathcal{U}}_1^\varepsilon - \hat{\mathcal{U}}_2^\varepsilon\|_{\varepsilon, \mathbb{R} \times \mathbb{R} \times \mathcal{Y}_a} &\leq C\{|\delta|\|\hat{\omega}_1^\varepsilon - \hat{\omega}_2^\varepsilon\|C_2 + C_1|\delta|(|\hat{\omega}_1^\varepsilon - \hat{\omega}_2^\varepsilon| + |\hat{\eta}_1^\varepsilon - \hat{\eta}_2^\varepsilon|) \\
&\quad + C_1|\delta|\|\hat{U}_1^\varepsilon - \hat{U}_2^\varepsilon\|_{\varepsilon, \mathcal{Y}_a} + |\delta|\|\hat{U}_1^\varepsilon - \hat{U}_2^\varepsilon\|_{\varepsilon, \mathcal{Y}_a} \\
&\quad + (C_1^{\frac{1}{2}} + C_1)|\delta|^2\|\hat{U}_1^\varepsilon - \hat{U}_2^\varepsilon\|_{\varepsilon, \mathcal{Y}_a}\} \\
&\leq C|\delta|\|\hat{\mathcal{U}}_1^\varepsilon - \hat{\mathcal{U}}_2^\varepsilon\|_{\varepsilon, \mathbb{R} \times \mathbb{R} \times \mathcal{Y}_a}.
\end{aligned}$$

We thus conclude that there exists a positive number δ_0 such that if $|\delta| \leq \delta_0$, then $\|\hat{\mathcal{U}}_1^\varepsilon - \hat{\mathcal{U}}_2^\varepsilon\|_{\varepsilon, \mathbb{R} \times \mathbb{R} \times \mathcal{Y}_a} = 0$, i.e., $\hat{\mathcal{U}}_1^\varepsilon = \hat{\mathcal{U}}_2^\varepsilon$.

Let ${}^\top(\eta_\delta^\varepsilon, \omega_\delta^\varepsilon, u_\delta^\varepsilon)$ be the solution branch obtained in Theorem 4.2 (i). We denote $\eta_\delta^\varepsilon = \delta\hat{\eta}_\delta^\varepsilon$, $\omega_\delta^\varepsilon = \delta\hat{\omega}_\delta^\varepsilon$ and $u_\delta^\varepsilon = \delta(z_0^\varepsilon + \delta U_\delta^\varepsilon)$. Set $\hat{\mathcal{U}}_\delta^\varepsilon = {}^\top(\hat{\eta}_\delta^\varepsilon, \hat{\omega}_\delta^\varepsilon, \hat{U}_\delta^\varepsilon)$. We then see that $|\eta_\delta^\varepsilon| + \|U_\delta^\varepsilon\|_{\varepsilon, \mathcal{Y}_a} \leq r_1$ and $|\omega_\delta^\varepsilon| \leq \frac{1}{4}$ by taking δ_0 in the proof of Theorem 4.2 (i) suitably smaller if necessary. The above argument is then applicable to $\hat{\mathcal{U}}_\delta^\varepsilon$ to obtain $\|\hat{\mathcal{U}}_\delta^\varepsilon\|_{\varepsilon, \mathbb{R} \times \mathbb{R} \times \mathcal{Y}_a} \leq C_1$ and $\|\partial_t \hat{U}_\delta^\varepsilon\|_{\varepsilon, \mathcal{Y}_a} \leq C_2$. We set $\hat{\mathcal{U}}_\delta^{\varepsilon, \tau} = {}^\top(\hat{\eta}_\delta^\varepsilon, \hat{\omega}_\delta^\varepsilon, U_\delta^\varepsilon(\cdot + \tau))$. Since $u_\delta^\varepsilon(\cdot + \tau)$ takes the form

$$u_\delta^\varepsilon(\cdot + \tau) = \delta(z_0^{\varepsilon, \tau}(\cdot) + \delta U_\delta^\varepsilon(\cdot + \tau))$$

and $\hat{\mathcal{U}}_\delta^{\varepsilon, \tau}$ satisfies $\|\hat{\mathcal{U}}_\delta^{\varepsilon, \tau}\|_{\varepsilon, \mathbb{R} \times \mathbb{R} \times \mathcal{Y}_a} \leq C_1$ and $\|\partial_t \hat{U}_\delta^{\varepsilon, \tau}\|_{\varepsilon, \mathcal{Y}_a} \leq C_2$, we see that $\hat{\mathcal{U}}^\varepsilon = \hat{\mathcal{U}}_\delta^{\varepsilon, \tau}$, and hence, we have ${}^\top(\eta^\varepsilon, \omega^\varepsilon, u^\varepsilon(\cdot)) = {}^\top(\eta_\delta^\varepsilon, \omega_\delta^\varepsilon, u_\delta^\varepsilon(\cdot + \tau))$. This completes the proof. \square

We finally give proofs of Lemmas 6.2 and 6.4.

Proof of Lemma 6.2. By the definition of $[u]_{+, \varepsilon}$, we have

$$\begin{aligned}
|[u]_{+, \varepsilon}| &\leq \frac{a}{2\pi} \int_0^{\frac{2\pi}{a}} \varepsilon^2 |(\phi, p_+^{\varepsilon*})_{L^2} + |(\mathbf{u}, \mathbf{z}_+^{\varepsilon*})| dt \\
&\leq C\{\varepsilon^2 \|\phi\|_{L^2(\mathbb{T}_{\frac{2\pi}{a}}; L^2(\Omega))} + \|\mathbf{u}\|_{L^2(\mathbb{T}_{\frac{2\pi}{a}}; (\mathbf{X}^1)^*)}\} \\
&\leq C\|u\|_{\varepsilon, \mathcal{X}_a}.
\end{aligned}$$

This proves (i).

As for (ii), by Theorem 4.1, we have $\sup_{0 \leq t \leq \frac{2\pi}{a}} \|z_+^\varepsilon(t)\|_{L^\infty(\Omega)} \leq C\|u_+^\varepsilon\|_{H^2(\Omega)} \leq C$ uniformly in $0 < \varepsilon \leq \varepsilon_0$. It follows that

$$|[N(u_1, u_2)]_{+, \varepsilon}| \leq \frac{a}{2\pi} \int_0^{\frac{2\pi}{a}} \|N(\mathbf{u}_1, \mathbf{u}_2)\|_1 \|z_+^\varepsilon\|_\infty dt$$

$$\begin{aligned}
&\leq C \int_0^{\frac{2\pi}{a}} \|\mathbf{u}_1\|_2 \|\nabla \mathbf{u}_2\|_2 dt \\
&\leq C \left(\sup_{0 \leq t \leq \frac{2\pi}{a}} \|\mathbf{u}_1(t)\|_2^2 \right)^{\frac{1}{2}} \left(\int_0^{\frac{2\pi}{a}} \|\nabla \mathbf{u}_2\|_2^2 dt \right)^{\frac{1}{2}} \\
&\leq C \| \mathbf{u}_1 \|_{\varepsilon, \mathcal{Y}_a} \| \mathbf{u}_2 \|_{\varepsilon, \mathcal{Y}_a}.
\end{aligned}$$

This shows (ii). Similarly, by using $\sup_{0 \leq t \leq \frac{2\pi}{a}} \|\mathbf{z}_+^\varepsilon(t)\|_\infty + \|\nabla \mathbf{z}_+^\varepsilon(t)\|_\infty \leq C$, we have

$$\int_0^{\frac{2\pi}{a}} \|\mathbf{M}(\mathbf{z}_0^\varepsilon, \mathbf{u})\|_{(\mathbf{X}_1)^*}^2 + \varepsilon^2 \|\mathbf{M}(\mathbf{z}_0^\varepsilon, \mathbf{u})\|_2^2 dt \leq C \int_0^{\frac{2\pi}{a}} \|\nabla \mathbf{u}\|_2^2 dt,$$

which gives (iii).

As for (iv), by the Gagliardo-Nirenberg inequality, we have

$$\begin{aligned}
|(\mathbf{N}(\mathbf{u}_1, \mathbf{u}_2), \mathbf{u}_3)| &\leq \|\mathbf{u}_1\|_4 \|\nabla \mathbf{u}_2\|_2 \|\mathbf{u}_3\|_4 \\
&\leq \|\mathbf{u}_1\|_2^{\frac{1}{2}} \|\nabla \mathbf{u}_1\|_2^{\frac{1}{2}} \|\nabla \mathbf{u}_2\|_2 \|\mathbf{u}_3\|_2^{\frac{1}{2}} \|\nabla \mathbf{u}_3\|_2^{\frac{1}{2}},
\end{aligned}$$

and hence,

$$\begin{aligned}
&\int_0^{\frac{2\pi}{a}} |(\mathbf{N}(\mathbf{u}_1, \mathbf{u}_2), \mathbf{u}_3)| dt \\
&\leq C \left(\sup_{0 \leq t \leq \frac{2\pi}{a}} \|\mathbf{u}_1(t)\|_2^2 \int_0^{\frac{2\pi}{a}} \|\nabla \mathbf{u}_1\|_2^2 dt \right)^{\frac{1}{4}} \left(\int_0^{\frac{2\pi}{a}} \|\nabla \mathbf{u}_2\|_2^2 dt \right)^{\frac{1}{2}} \\
&\quad \cdot \left(\sup_{0 \leq t \leq \frac{2\pi}{a}} \|\mathbf{u}_3(t)\|_2^2 \int_0^{\frac{2\pi}{a}} \|\nabla \mathbf{u}_3\|_2^2 dt \right)^{\frac{1}{4}} \\
&\leq C \| \mathbf{u}_1 \|_{\varepsilon, \mathcal{Y}_a} \| \mathbf{u}_2 \|_{\varepsilon, \mathcal{Y}_a} \| \mathbf{u}_3 \|_{\varepsilon, \mathcal{Y}_a}.
\end{aligned}$$

This proves (iv). Similarly, we have

$$\|\mathbf{N}(\mathbf{u}_1, \mathbf{u}_2)\|_2^2 dt \leq \|\mathbf{u}_1\|_4^2 \|\nabla \mathbf{u}_2\|_4^2 \leq C \|\mathbf{u}_1\|_2 \|\nabla \mathbf{u}_1\|_2 \|\nabla \mathbf{u}_2\|_2 \|\partial_x^2 \mathbf{u}_2\|_2,$$

and hence,

$$\int_0^{\frac{2\pi}{a}} \varepsilon^2 \|\mathbf{N}(\mathbf{u}_1, \mathbf{u}_2)\|_2^2 dt \leq C \int_0^{\frac{2\pi}{a}} \varepsilon^2 \|\mathbf{u}_1\|_2 \|\nabla \mathbf{u}_1\|_2 \|\nabla \mathbf{u}_2\|_2 \|\partial_x^2 \mathbf{u}_2\|_2 dt$$

$$\begin{aligned}
&\leq C \left(\sup_{0 \leq t \leq \frac{2\pi}{a}} \|\mathbf{u}_1(t)\|_2^2 \int_0^{\frac{2\pi}{a}} \|\nabla \mathbf{u}_1\|_2^2 dt \right)^{\frac{1}{2}} \\
&\quad \cdot \left(\sup_{0 \leq t \leq \frac{2\pi}{a}} \varepsilon^2 \|\nabla \mathbf{u}_2(t)\|_2^2 \int_0^{\frac{2\pi}{a}} \varepsilon^2 \|\partial_x^2 \mathbf{u}_2\|_2 dt \right)^{\frac{1}{2}} \\
&\leq C \| |u_1| \|_{\varepsilon, \mathcal{Y}_a}^2 \| |u_2| \|_{\varepsilon, \mathcal{Y}_a}^2,
\end{aligned}$$

and (v) is obtained. This completes the proof. \square

Proof of Lemma 6.4. As for (i), we have

$$\begin{aligned}
|[u_1]_{+, \varepsilon} - [u_2]_{+}] &\leq \frac{a}{2\pi} \int_0^{\frac{2\pi}{a}} |\varepsilon^2 (\phi_1, p_+^{\varepsilon*}) + (\mathbf{u}_1 - \mathbf{u}_2, \mathbf{z}_+^{\varepsilon*}) + (\mathbf{u}_2, \mathbf{z}_+^{\varepsilon*} - \mathbf{z}_+^*)| dt \\
&\leq C \left\{ \varepsilon^2 \|\phi_1\|_{L^2(\mathbb{T}_{\frac{2\pi}{a}}; L^2(\Omega))} + \varepsilon^2 \|\mathbf{u}_2\|_{L^2(\mathbb{T}_{\frac{2\pi}{a}}; (\mathbf{X}^1)^*)} \right. \\
&\quad \left. + \|\mathbf{u}_1 - \mathbf{u}_2\|_{L^2(\mathbb{T}_{\frac{2\pi}{a}}; (\mathbf{X}^1)^*)} \right\}.
\end{aligned}$$

We thus obtain (i). Similarly, we have

$$\begin{aligned}
&|[N(u_1, u_2)]_{+, \varepsilon} - [N(\tilde{u}_1, \tilde{u}_2)]_{+}] \\
&= |\langle N(\mathbf{u}_1, \mathbf{u}_2), \mathbf{z}_+^{\varepsilon*} \rangle - \langle N(\tilde{\mathbf{u}}_1, \tilde{\mathbf{u}}_2), \mathbf{z}_+^* \rangle| \\
&= |\langle N(\mathbf{u}_1, \mathbf{u}_2), \mathbf{z}_+^{\varepsilon*} - \mathbf{z}_+^* \rangle + \langle N(\mathbf{u}_1, \mathbf{u}_2) - N(\tilde{\mathbf{u}}_1, \tilde{\mathbf{u}}_2), \mathbf{z}_+^* \rangle| \\
&\leq C \varepsilon^2 \int_0^{\frac{2\pi}{a}} \|N(\mathbf{u}_1, \mathbf{u}_2)\|_1 dt + C \int_0^{\frac{2\pi}{a}} \|N(\mathbf{u}_1, \mathbf{u}_2) - N(\tilde{\mathbf{u}}_1, \tilde{\mathbf{u}}_2)\|_1 dt \\
&\leq C \left\{ \varepsilon^2 \| |u_1| \|_{\varepsilon, \mathcal{Y}_a} \| |u_2| \|_{\varepsilon, \mathcal{Y}_a} + \| |u_1| \|_{\varepsilon, \mathcal{Y}_a} \| |u_2 - \tilde{u}_2| \|_{\varepsilon, \mathcal{Y}_a} \right. \\
&\quad \left. + \| |u_1 - \tilde{u}_1| \|_{\varepsilon, \mathcal{Y}_a} \| |\tilde{u}_2| \|_{\varepsilon, \mathcal{Y}_a} \right\}.
\end{aligned}$$

This proves (ii).

The inequality (iii) is proved as follows. We first consider the term $\mathcal{Q}^{\varepsilon \top}(0, \partial_t \mathbf{z}_0)$. Since $\hat{\mathcal{Q}}_0 \partial_t \mathbf{z}_0 = \partial_t \mathbf{z}_0 - 2\text{Re}([\partial_t \mathbf{z}_0]_{+} \mathbf{z}_+) = \mathbf{0}$, we have

$$\begin{aligned}
\mathcal{Q}^{\varepsilon \top}(0, \partial_t \mathbf{z}_0) &= - \left(\begin{array}{c} 2\text{Re}(\langle \partial_t \mathbf{z}_0, \mathbf{z}_+^{\varepsilon*} \rangle p_+^{\varepsilon}) \\ 2\text{Re}(\langle \partial_t \mathbf{z}_0, \mathbf{z}_+^{\varepsilon*} \rangle \mathbf{z}_+^{\varepsilon} - \langle \partial_t \mathbf{z}_0, \mathbf{z}_+^* \rangle \mathbf{z}_+) \end{array} \right) \\
&= - \left(\begin{array}{c} 2\text{Re}(\langle \partial_t \mathbf{z}_0, \mathbf{z}_+^{\varepsilon*} \rangle p_+^{\varepsilon}) \\ 2\text{Re}(\langle \partial_t \mathbf{z}_0, \mathbf{z}_+^{\varepsilon*} - \mathbf{z}_+^* \rangle \mathbf{z}_+^{\varepsilon} + \langle \partial_t \mathbf{z}_0, \mathbf{z}_+^* \rangle (\mathbf{z}_+^{\varepsilon} - \mathbf{z}_+)) \end{array} \right).
\end{aligned}$$

We thus obtain $||| \mathcal{Q}^{\varepsilon \top}(0, \partial_t \mathbf{z}_0) |||_{\varepsilon, \mathcal{X}_a} \leq C\varepsilon$. We also note that $||| \top(\partial_t \Phi_\delta, \mathbf{0}) |||_{\varepsilon, \mathcal{X}_a} \leq C\varepsilon ||| \top(\partial_t \Phi_\delta, \mathbf{0}) |||_{\mathcal{X}_a} \leq C\varepsilon$. It then follows that

$$||| \mathcal{Q}^\varepsilon H^\varepsilon(\delta; \tilde{\omega}_\delta^\varepsilon, \tilde{\omega}_\delta, U_\delta) |||_{\varepsilon, \mathcal{X}_a} \leq C \left\{ \varepsilon(1 + |\delta| |\tilde{\omega}_\delta| + \delta^2 |\tilde{\omega}_\delta^\varepsilon|) + \delta^2 |\tilde{\omega}_\delta^\varepsilon - \tilde{\omega}_\delta| \right\}.$$

This completes the proof. \square

7 Stability of bifurcating solutions

Throughout this section we denote $T_a = \frac{2\pi}{a}$. In what follows, we will use the same letters ε_2 and δ_2 for bounds of the ranges of ε and δ , even when they should be taken suitably smaller than those of the previous ones if no confusion will occur from the context.

We investigate the linearized problem around $u_\delta^\varepsilon(t)$. The linearized problem around $u_\delta^\varepsilon(t)$ then takes the form

$$\frac{a^\varepsilon}{a} \partial_t u + \frac{1}{1 + \omega_\delta^\varepsilon} L_{\mathcal{R}_{1,c}^\varepsilon} u + \frac{\delta}{1 + \omega_\delta^\varepsilon} M_\delta^\varepsilon(t) u = F, \quad (7.1)$$

$$u|_{t=s} = v, \quad (7.2)$$

where $v \in X^1$ and $F \in \mathcal{X}(s, s + T_a)$ are given functions. Here and in what follows, we denote

$$\begin{aligned} \hat{u}_\delta^\varepsilon(t) &= \frac{1}{\delta} u_\delta^\varepsilon(t) = z_0^\varepsilon(t) + \delta U_\delta^\varepsilon(t), \\ M_\delta^\varepsilon(t) u &= \delta \tilde{\eta}_\delta^\varepsilon K u + M(\hat{u}_\delta^\varepsilon(t), u). \end{aligned}$$

Recall that $\eta_\delta^\varepsilon = \tilde{\eta}_0^\varepsilon \delta^2 + \tilde{\eta}_1^\varepsilon(\delta) \delta^3$.

By Theorem 4.2, we have $||| \hat{u}_\delta^\varepsilon |||_{\varepsilon, \mathcal{Y}_a} \leq C$ uniformly in $0 < \varepsilon \leq \varepsilon_1$ and $|\delta| \leq \delta_1$. Based on this, we see, by a perturbation argument, that there are positive constants ε_2 and δ_2 such that if $0 < \varepsilon \leq \varepsilon_2$ and $|\delta| \leq \delta_2$, then the problem (7.1)–(7.2) has a unique solution $u \in \mathcal{Y}(s, s + T_a)$ and it satisfies the estimate

$$||| u |||_{\varepsilon, \mathcal{Y}(s, s + T_a)} \leq C \{ ||| v |||_{X^1} + ||| F |||_{\varepsilon, \mathcal{X}(s, s + T_a)} \}$$

uniformly for $s \in \mathbb{R}$, $0 < \varepsilon \leq \varepsilon_2$ and $|\delta| \leq \delta_2$.

Let $\mathcal{U}_\delta^\varepsilon(t, s)$ be the solution operator for (7.1)–(7.2) with $F = 0$ and let $\tilde{\mathcal{W}}_\delta^\varepsilon(t, s)$ be the solution operator for (7.1)–(7.2) with $v = 0$. The solution u of (7.1)–(7.2) is then written as

$$u(t) = \mathcal{U}_\delta^\varepsilon(t, s) v + \tilde{\mathcal{W}}_\delta^\varepsilon(t, s) F \quad (7.3)$$

and $\mathcal{U}_\delta^\varepsilon(t, s)$ and $\widetilde{\mathcal{U}}_\delta^\varepsilon(t, s)$ satisfy the uniform estimates

$$\|\|\mathcal{U}_\delta^\varepsilon(t, s)v\|\|_{\varepsilon, \mathcal{Y}(s, s+T_a)} \leq C\|\|v\|\|_{X^1}, \quad (7.4)$$

$$\|\|\widetilde{\mathcal{U}}_\delta^\varepsilon(t, s)F\|\|_{\varepsilon, \mathcal{Y}(s, s+T_a)} \leq C\|\|F\|\|_{\varepsilon, \mathcal{X}(s, s+T_a)} \quad (7.5)$$

uniformly for $s \in \mathbb{R}$, $0 < \varepsilon \leq \varepsilon_2$ and $|\delta| \leq \delta_2$.

Since the bifurcating solution $u_\delta^\varepsilon(t) = \delta(z_0^\varepsilon(t) + \delta U^\varepsilon(t))$ is a time periodic function of period T_a , its stability is determined by the spectrum of $\mathcal{U}_\delta^\varepsilon(T_a, 0)$. We shall investigate the spectrum of $\mathcal{U}_\delta^\varepsilon(T_a, 0)$ as a perturbation of that of $\mathcal{V}_{\omega_\delta^\varepsilon}^\varepsilon(T_a)$. Here recall that $\mathcal{V}_{\omega_\delta^\varepsilon}^\varepsilon(t) = e^{-\frac{at}{a^\varepsilon(1+\omega_\delta^\varepsilon)}L_{\mathcal{R}_{1,c}}^\varepsilon}|_{X^1}$. We shall prove the following result on the spectrum of $\mathcal{U}_\delta^\varepsilon(T_a, 0)$ for $0 < |\delta| \ll 1$.

Theorem 7.1 *There exist positive constants ε_2 , δ_2 and Λ_2 such that if $0 < \varepsilon \leq \varepsilon_2$ and $|\delta| \leq \delta_2$, then*

$$\sigma(\mathcal{U}_\delta^\varepsilon(T_a, 0)) \subset \{1, e^{\frac{a}{a^\varepsilon}T_a\lambda_\delta^\varepsilon}\} \cup \{\mu \in \mathbb{C}; |\mu| \leq e^{-T_a\Lambda_2}\}.$$

Here 1 and $e^{T_a\lambda_\delta^\varepsilon}$ are simple eigenvalues of $\mathcal{U}_\delta^\varepsilon(T_a, 0)$ and $\lambda_\delta^\varepsilon$ satisfies

$$\lambda_\delta^\varepsilon = 2\delta^2\tilde{\eta}_0^\varepsilon \operatorname{Re}[Kz_0^\varepsilon]_{+, \varepsilon} + O(\delta^3)$$

uniformly for $0 < \varepsilon \leq \varepsilon_2$ and $|\delta| \leq \delta_2$. The eigenspace of the eigenvalue 1 is spanned by $\partial_t u_\delta^\varepsilon$.

To prove Theorem 7.1, we first show the following lemma.

Lemma 7.2 *There exist positive constants Λ_2 and r_0 such that for all $0 < r \leq r_0$ there exists a positive constant $\delta_2 = \delta_2(r)$ such that for $0 < \varepsilon \leq \varepsilon_2$ and $|\delta| \leq \delta_2$, it holds that*

$$\rho(\mathcal{U}_\delta^\varepsilon(T_a, 0)) \supset \{\mu; |\mu - 1| \geq r\} \cap \{\mu; |\mu| \geq e^{-T_a\Lambda_2}\},$$

and $\sigma(\mathcal{U}_\delta^\varepsilon(T_a, 0)) \cap \{\mu; |\mu - 1| < r\}$ consists of eigenvalues and the spectral projection associated with the set of these eigenvalues is a finite rank operator of rank 2.

To prove Lemma 7.2, we observe that $\mathcal{U}_\delta^\varepsilon(t, \tau)$ is written as

$$\mathcal{U}_\delta^\varepsilon(t, \tau) = \mathcal{V}_{\omega_\delta^\varepsilon}^\varepsilon(t - \tau) - \frac{a\delta}{a^\varepsilon(1 + \omega_\delta^\varepsilon)} \int_\tau^t \mathcal{V}_{\omega_\delta^\varepsilon}^\varepsilon(t - s)M_\delta^\varepsilon(s)\mathcal{U}_\delta^\varepsilon(s, \tau) ds.$$

See (7.6) below. Using this formula, we shall investigate the spectrum of $\mathcal{W}_\delta^\varepsilon(T_a, 0)$ by a perturbation argument. We thus introduce an operator $\mathcal{S}_{\omega, \delta}^\varepsilon$ on $\mathfrak{B}(X^1, \mathcal{Y}(0, T_a))$ defined by

$$(\mathcal{S}_{\omega, \delta}^\varepsilon \mathcal{W}v)(t) = -\frac{a}{a^\varepsilon(1+\omega)} \int_0^t \mathcal{V}_\omega^\varepsilon(t-s) M_\delta^\varepsilon(s) \mathcal{W}v(s) ds$$

for $\mathcal{W} \in \mathfrak{B}(X^1, \mathcal{Y}(0, T_a))$ and $v \in X^1$. The following estimates hold for $\mathcal{S}_{\omega, \delta}^\varepsilon$.

Lemma 7.3 (i) *Let $|\omega| \leq \frac{1}{4}$. Then the operator $\mathcal{S}_{\omega, \delta}^\varepsilon$ is a bounded operator on $\mathfrak{B}(X^1, \mathcal{Y}(0, T_a))$ and satisfies*

$$\|\mathcal{S}_{\omega, \delta}^\varepsilon \mathcal{W}\|_{\mathfrak{B}(X^1, \mathcal{Y}(0, T_a))} \leq C \|\mathcal{W}\|_{\mathfrak{B}(X^1, \mathcal{Y}(0, T_a))}$$

and

$$\|\mathcal{S}_{\omega, \delta}^\varepsilon\|_{\mathfrak{B}(\mathfrak{B}(X^1, \mathcal{Y}(0, T_a)))} \leq C$$

uniformly for $0 < \varepsilon \leq \varepsilon_2$, $|\omega| \leq \frac{1}{4}$ and $|\delta| \leq \delta_2$.

(ii) *There exists a positive constant δ_2 such that if $|\delta| \leq \delta_2$, then $I - \delta \mathcal{S}_{\omega, \delta}^\varepsilon$ has the bounded inverse $(I - \delta \mathcal{S}_{\omega, \delta}^\varepsilon)^{-1} \in \mathfrak{B}(\mathfrak{B}(X^1, \mathcal{Y}(0, T_a)))$ with estimate*

$$\|(I - \delta \mathcal{S}_{\omega, \delta}^\varepsilon)^{-1}\|_{\mathfrak{B}(\mathfrak{B}(X^1, \mathcal{Y}(0, T_a)))} \leq 2$$

uniformly for $0 < \varepsilon \leq \varepsilon_2$, $|\omega| \leq \frac{1}{4}$ and $|\delta| \leq \delta_2$.

Proof. Applying Lemmas 5.1 and 6.2, we obtain

$$\begin{aligned} \|\|\mathcal{S}_{\omega, \delta}^\varepsilon \mathcal{W}v\|\|_{\varepsilon, \mathcal{Y}(0, T_a)} &\leq \frac{1}{2} \|\|\mathcal{S}_{\omega, \delta}^\varepsilon \mathcal{W}v\|\|_{\varepsilon, \mathcal{Y}(0, T_a)} \\ &\quad + C \{ \|\|\mathcal{W}v\|\|_{\varepsilon, \mathcal{X}(0, T_a)} + \|\|\hat{u}_\delta^\varepsilon\|\|_{\varepsilon, \mathcal{Y}_a} \|\|\mathcal{W}(\cdot)v\|\|_{\varepsilon, \mathcal{Y}(0, T_a)} \} \\ &\leq \frac{1}{2} \|\|\mathcal{S}_{\omega, \delta}^\varepsilon \mathcal{W}v\|\|_{\varepsilon, \mathcal{Y}(0, T_a)} + C \|\|\mathcal{W}\|\|_{\mathfrak{B}(X^1, \mathcal{Y}(0, T_a))} \|\|v\|\|_{\varepsilon, X^1}, \end{aligned}$$

and hence,

$$\|\|\mathcal{S}_{\omega, \delta}^\varepsilon \mathcal{W}v\|\|_{\varepsilon, \mathcal{Y}(0, T_a)} \leq C \|\|\mathcal{W}\|\|_{\mathfrak{B}(X^1, \mathcal{Y}(0, T_a))} \|\|v\|\|_{\varepsilon, X^1}$$

uniformly for $0 < \varepsilon \leq \varepsilon_2$, $|\omega| \leq \frac{1}{4}$ and $|\delta| \leq \delta_2$. This implies that

$$\|\|\mathcal{S}_{\omega, \delta}^\varepsilon \mathcal{W}\|\|_{\mathfrak{B}(X^1, \mathcal{Y}(0, T_a))} \leq C \|\|\mathcal{W}\|\|_{\mathfrak{B}(X^1, \mathcal{Y}(0, T_a))},$$

and hence,

$$\|\mathcal{S}_{\omega,\delta}^\varepsilon\|_{\mathfrak{B}(\mathfrak{B}(X^1, \mathcal{Y}(0, T_a)))} \leq C$$

uniformly for $0 < \varepsilon \leq \varepsilon_2$, $|\omega| \leq \frac{1}{4}$ and $|\delta| \leq \delta_2$. It then follows that if $|\delta| \leq \frac{1}{2C}$, then there exists $(I - \delta \mathcal{S}_{\omega,\delta}^\varepsilon)^{-1} \in \mathfrak{B}(\mathfrak{B}(X^1, \mathcal{Y}(0, T_a)))$ with estimate $\|(I - \delta \mathcal{S}_{\omega,\delta}^\varepsilon)^{-1}\|_{\mathfrak{B}(\mathfrak{B}(X^1, \mathcal{Y}(0, T_a)))} \leq 2$. This completes the proof. \square

We next give a proof of Lemma 7.2.

Proof of Lemma 7.2. Let $u(t)$ be a solution of (7.1)–(7.2) with $F = 0$. Since $u(t) = \mathcal{U}_\delta^\varepsilon(t, \tau)v$ and $\partial_t u + \frac{a}{a^\varepsilon} L_{\omega_\delta^\varepsilon}^\varepsilon u = -\frac{a\delta}{a^\varepsilon(1+\omega_\delta^\varepsilon)} M_\delta^\varepsilon(t)u$, the solution operator $\mathcal{U}_\delta^\varepsilon(t, \tau)$ for (7.1) satisfies

$$\begin{aligned} \mathcal{U}_\delta^\varepsilon(t, \tau)v &= \mathcal{V}_{\omega_\delta^\varepsilon}^\varepsilon(t - \tau)v - \frac{a\delta}{a^\varepsilon(1 + \omega_\delta^\varepsilon)} \int_\tau^t \mathcal{V}_{\omega_\delta^\varepsilon}^\varepsilon(t - s) M_\delta^\varepsilon(s) u(s) ds \\ &= \mathcal{V}_{\omega_\delta^\varepsilon}^\varepsilon(t - \tau)v - \frac{a\delta}{a^\varepsilon(1 + \omega_\delta^\varepsilon)} \int_\tau^t \mathcal{V}_{\omega_\delta^\varepsilon}^\varepsilon(t - s) M_\delta^\varepsilon(s) \mathcal{U}_\delta^\varepsilon(s, \tau)v ds. \end{aligned} \quad (7.6)$$

We set $\mathcal{W}_\delta^\varepsilon(t, s) = \mathcal{U}_\delta^\varepsilon(t, s) - \mathcal{V}_{\omega_\delta^\varepsilon}^\varepsilon(t - s)$. It then follows that

$$\begin{aligned} \mathcal{W}_\delta^\varepsilon(t, \tau)v &= -\frac{a\delta}{a^\varepsilon(1 + \omega_\delta^\varepsilon)} \int_\tau^t \mathcal{V}_{\omega_\delta^\varepsilon}^\varepsilon(t - s) M_\delta^\varepsilon(s) \mathcal{U}_\delta^\varepsilon(s, \tau)v ds \\ &= -\frac{a\delta}{a^\varepsilon(1 + \omega_\delta^\varepsilon)} \int_\tau^t \mathcal{V}_{\omega_\delta^\varepsilon}^\varepsilon(t - s) M_\delta^\varepsilon(s) \mathcal{V}_{\omega_\delta^\varepsilon}^\varepsilon(s - \tau)v ds \\ &\quad - \frac{a\delta}{a^\varepsilon(1 + \omega_\delta^\varepsilon)} \int_\tau^t \mathcal{V}_{\omega_\delta^\varepsilon}^\varepsilon(t - s) M_\delta^\varepsilon(s) \mathcal{W}_\delta^\varepsilon(s, \tau)v ds. \end{aligned} \quad (7.7)$$

We claim that

$$\|\mathcal{W}_\delta^\varepsilon(T_a, 0)\|_{\mathfrak{B}(X^1)} \leq 2C|\delta| \quad (7.8)$$

uniformly for $0 < \varepsilon \leq \varepsilon_2$ and $|\delta| \leq \delta_2$. Indeed, we see from (7.7) that $\mathcal{W}_\delta^\varepsilon(t, 0)$ is written as

$$\mathcal{W}_\delta^\varepsilon(t, 0)v = -\delta(\mathcal{S}_{\omega_\delta^\varepsilon, \delta}^\varepsilon \mathcal{V}_{\omega_\delta^\varepsilon}^\varepsilon(\cdot)v)(t) - \delta(\mathcal{S}_{\omega_\delta^\varepsilon, \delta}^\varepsilon \mathcal{W}_\delta^\varepsilon(\cdot, 0)v)(t).$$

This implies

$$(I + \delta \mathcal{S}_{\omega_\delta^\varepsilon, \delta}^\varepsilon) \mathcal{W}_\delta^\varepsilon(\cdot, 0) = -\delta \mathcal{S}_{\omega_\delta^\varepsilon, \delta}^\varepsilon \mathcal{V}_{\omega_\delta^\varepsilon}^\varepsilon(\cdot).$$

By Lemmas 5.1 and 7.3, we have

$$\|\mathcal{S}_{\omega_\delta^\varepsilon, \delta}^\varepsilon \mathcal{V}_{\omega_\delta^\varepsilon}^\varepsilon(\cdot)v\|_{\varepsilon, \mathcal{Y}(0, T_a)} \leq C\|v\|_{\varepsilon, X^1}$$

uniformly for $0 < \varepsilon \leq \varepsilon_2$ and $|\delta| \leq \delta_2$. We also find from Lemma 7.3 that

$$\mathcal{W}_\delta^\varepsilon(\cdot, 0) = -\delta(I + \delta \mathcal{S}_{\omega_\delta^\varepsilon, \delta}^\varepsilon)^{-1} \mathcal{S}_{\omega_\delta^\varepsilon, \delta}^\varepsilon \mathcal{V}_{\omega_\delta^\varepsilon}^\varepsilon(\cdot)$$

and

$$\|\mathcal{W}_\delta^\varepsilon(\cdot, 0)v\|_{\varepsilon, \mathcal{Y}(0, T_a)} \leq 2|\delta| \|\mathcal{S}_{\omega_\delta^\varepsilon, \delta}^\varepsilon \mathcal{V}_{\omega_\delta^\varepsilon}^\varepsilon(\cdot)v\|_{\varepsilon, \mathcal{Y}(0, T_a)} \leq 2C|\delta| \|v\|_{\varepsilon, X^1}.$$

In particular, we have $\|\mathcal{W}_\delta^\varepsilon(T_a, 0)v\|_{\varepsilon, X^1} \leq 2C|\delta| \|v\|_{\varepsilon, X^1}$, which yields $\|\mathcal{W}_\delta^\varepsilon(T_a, 0)\|_{\mathfrak{B}(X^1)} \leq 2C|\delta|$ uniformly for $0 < \varepsilon \leq \varepsilon_2$ and $|\delta| \leq \delta_2$. This proves (7.8).

We next consider the resolvent of $\mathcal{U}_\delta^\varepsilon(T_a, 0)$. For a fixed positive constant r we set

$$\Sigma_r = \{\mu \in \mathbb{C}; |\mu - 1| \geq r, |\mu| \geq e^{-\frac{3}{4}\kappa_1 T_a}\}.$$

We see from Lemma 5.3 that if $|\omega_\delta^\varepsilon| \leq C \min\{r, 1\}$, then

$$\|(\mu - \mathcal{V}_{\omega_\delta^\varepsilon}^\varepsilon(T_a))^{-1}\|_{\mathfrak{B}(X^1)} \leq C \left(\frac{1}{r} + \frac{1}{|\mu|} \right).$$

Since $\omega_\delta^\varepsilon = \tilde{\omega}_\delta^\varepsilon \delta^2$, there exists a positive constant $\delta_2 = O(r)$ ($r \rightarrow 0$) such that if $|\delta| \leq \delta_2$, then

$$\|\mathcal{W}_\delta^\varepsilon(T_a, 0)(\mu - \mathcal{V}_{\omega_\delta^\varepsilon}^\varepsilon(T_a))^{-1}\|_{\mathfrak{B}(X^1)} \leq \frac{1}{2}.$$

We thus conclude that $\Sigma_r \subset \rho(\mathcal{U}_\delta^\varepsilon(T_a, 0))$ and

$$\begin{aligned} & (\mu - \mathcal{U}_\delta^\varepsilon(T, 0))^{-1} \\ &= (\mu - \mathcal{V}_{\omega_\delta^\varepsilon}^\varepsilon(T_a))^{-1} (I - \mathcal{W}_\delta^\varepsilon(T_a, 0)(\mu - \mathcal{V}_{\omega_\delta^\varepsilon}^\varepsilon(T_a))^{-1})^{-1} \end{aligned}$$

for $\mu \in \Sigma_r$. Furthermore, it holds

$$\|(\mu - \mathcal{U}_\delta^\varepsilon(T, 0))^{-1}\|_{\mathfrak{B}(X^1)} \leq 2C \left(\frac{1}{r} + \frac{1}{|\mu|} \right)$$

for $\mu \in \Sigma_r$ uniformly in $0 < \varepsilon \leq \varepsilon_2$ and $|\delta| \leq \delta_2$. This completes the proof. \square

We next investigate a part of $\sigma(\mathcal{U}_\delta^\varepsilon(T_a, 0))$ in $\{\mu \in \mathbb{C}; |\mu - 1| < r\}$. To specify it, we consider the spectrum of the linearized operator B_δ^ε around the bifurcating time periodic solution u_δ^ε of (4.3) which is given by

$$B_\delta^\varepsilon = B^\varepsilon(\omega_\delta^\varepsilon) + \delta M_\delta^\varepsilon, \tag{7.9}$$

where M_δ^ε is the operator defined by

$$(M_\delta^\varepsilon u)(t) = M_\delta^\varepsilon(t)u(t) \quad \text{for } t \in \mathbb{T}_{T_a}$$

with $M_\delta^\varepsilon(t)$ given in (7.1). Note that $B_0^\varepsilon = B^\varepsilon$.

Theorem 7.4 *There exists a positive constant r_0 such that*

$$\sigma(-B_\delta^\varepsilon) \cap \{\lambda \in \mathbb{C}; |\lambda| \leq r_0\} = \{0, \lambda_\delta^\varepsilon\},$$

where 0 and $\lambda_\delta^\varepsilon$ are simple eigenvalues of $-B_\delta^\varepsilon$. Furthermore, $\lambda_\delta^\varepsilon$ satisfies

$$\lambda_\delta^\varepsilon = 2\delta^2 \tilde{\eta}_0^\varepsilon \operatorname{Re}[K z_0^\varepsilon]_{+, \varepsilon} + O(\delta^3)$$

uniformly for $0 < \varepsilon \leq \varepsilon_2$ and $|\delta| \leq \delta_2$.

To give a proof of Theorem 7.4, we introduce notation. We change the bases of $N(B_0^\varepsilon)$ from $\{z_+^\varepsilon, z_-^\varepsilon\}$ to $\{z_0^\varepsilon, z_1^\varepsilon\}$, where

$$z_0^\varepsilon = 2\operatorname{Re}z_+^\varepsilon, \quad z_1^\varepsilon = 2\operatorname{Im}z_+^\varepsilon,$$

and likewise, the dual bases from $\{z_+^{\varepsilon*}, z_-^{\varepsilon*}\}$ to $\{z_0^{\varepsilon*}, z_1^{\varepsilon*}\}$, where

$$z_0^{\varepsilon*} = \operatorname{Re}z_+^{\varepsilon*}, \quad z_1^{\varepsilon*} = \operatorname{Im}z_+^{\varepsilon*}.$$

We set

$$[[u]]_{j, \varepsilon} = \langle u, z_j^{\varepsilon*} \rangle_\varepsilon \quad (j = 0, 1)$$

and define the operators $\tilde{\mathcal{P}}^\varepsilon$ and $\tilde{\mathcal{Q}}^\varepsilon$ by

$$\tilde{\mathcal{P}}^\varepsilon u = [[u]]_{0, \varepsilon} z_0^\varepsilon + [[u]]_{1, \varepsilon} z_1^\varepsilon$$

and

$$\tilde{\mathcal{Q}}^\varepsilon = I - \tilde{\mathcal{P}}^\varepsilon,$$

respectively. It then follows that $\tilde{\mathcal{P}}^\varepsilon$ is an eigenprojection for the eigenvalue 0 of $-B_0^\varepsilon$. Observe also that

$$[[\partial_t u]]_{0, \varepsilon} = a[[u]]_{1, \varepsilon}, \quad [[\partial_t u]]_{1, \varepsilon} = -a[[u]]_{0, \varepsilon}.$$

Furthermore, it holds that

$$[[u]]_{0, \varepsilon} = \operatorname{Re}[u]_{+, \varepsilon} \text{ if } u \text{ is a real valued function.}$$

We see from Lemma 5.5 that if $\operatorname{Re}\lambda \geq -\frac{3}{4}\kappa_1 T_a$, then the problem

$$(\lambda + B^\varepsilon(\omega))u = F \quad (7.10)$$

has a unique solution $u \in \tilde{\mathcal{Q}}^\varepsilon \mathcal{Y}_a$ for any given $F \in \tilde{\mathcal{Q}}^\varepsilon \mathcal{X}_a$ with estimate

$$\|u\|_{\varepsilon, \mathcal{Y}_a} \leq C \|F\|_{\varepsilon, \mathcal{X}_a}$$

uniformly for $0 < \varepsilon \leq \varepsilon_2$, $|\omega| \leq \frac{1}{4}$ and λ with $\operatorname{Re}\lambda \geq -\frac{3}{4}\kappa_1 T_a$ and $|\operatorname{Im}\lambda| \leq \frac{a}{2}$. We denote the solution operator for problem (7.10) by $\mathcal{S}^\varepsilon(\lambda, \omega)$, namely, the operator $\mathcal{S}^\varepsilon(\lambda, \omega) : \tilde{\mathcal{Q}}^\varepsilon \mathcal{X}_a \rightarrow \tilde{\mathcal{Q}}^\varepsilon \mathcal{Y}_a$ is defined by

$$u = \mathcal{S}^\varepsilon(\lambda, \omega)F,$$

where $u \in \tilde{\mathcal{Q}}^\varepsilon \mathcal{Y}_a$ is the unique solution of (7.10).

We now consider the resolvent problem

$$(\lambda + B_\delta^\varepsilon)u = F. \quad (7.11)$$

We shall employ the Lyapunov-Schmidt method to investigate problem (7.11).

We decompose u in (7.11) into its $\tilde{\mathcal{P}}^\varepsilon$ and $\tilde{\mathcal{Q}}^\varepsilon$ parts as

$$u = \zeta_0 z_0^\varepsilon + \zeta_1 z_1^\varepsilon + U,$$

where $\zeta_j = \llbracket u \rrbracket_{j, \varepsilon}$ ($j = 0, 1$) and $U = \tilde{\mathcal{Q}}^\varepsilon u \in \tilde{\mathcal{Q}}^\varepsilon \mathcal{Y}_a$. Applying $\tilde{\mathcal{P}}^\varepsilon$ and $\tilde{\mathcal{Q}}^\varepsilon$ to (7.11), we see that (7.11) is reduced to

$$\lambda \zeta_0 + a^\varepsilon \omega_\delta^\varepsilon \zeta_1 + \eta_\delta^\varepsilon \llbracket Ku \rrbracket_{0, \varepsilon} + \delta \llbracket M(\hat{u}_\delta^\varepsilon, u) \rrbracket_{0, \varepsilon} = \llbracket F \rrbracket_{0, \varepsilon}, \quad (7.12)$$

$$\lambda \zeta_1 - a^\varepsilon \omega_\delta^\varepsilon \zeta_0 + \eta_\delta^\varepsilon \llbracket Ku \rrbracket_{1, \varepsilon} + \delta \llbracket M(\hat{u}_\delta^\varepsilon, u) \rrbracket_{1, \varepsilon} = \llbracket F \rrbracket_{1, \varepsilon}, \quad (7.13)$$

$$(\lambda + \tilde{\mathcal{Q}}^\varepsilon B_\delta^\varepsilon)U + \delta \tilde{\mathcal{Q}}^\varepsilon \sum_{j=0,1} \zeta_j M_\delta^\varepsilon z_j^\varepsilon = \tilde{\mathcal{Q}}^\varepsilon F, \quad (7.14)$$

where $u = \zeta_0 z_0^\varepsilon + \zeta_1 z_1^\varepsilon + U$ with $\zeta = {}^\top(\zeta_0, \zeta_1) \in \mathbb{C}^2$ and $U \in \tilde{\mathcal{Q}}^\varepsilon \mathcal{Y}_a$.

We shall reduce (7.12)–(7.14) to a two-dimensional problem by solving (7.14) for U in terms of ζ_j ($j = 0, 1$) and F and then substituting U into (7.12) and (7.13). To this end, we next consider the problem

$$(\lambda + \tilde{\mathcal{Q}}^\varepsilon B_\delta^\varepsilon)u = \tilde{\mathcal{Q}}^\varepsilon F. \quad (7.15)$$

Proposition 7.5 *There exist positive constants ε_2 and δ_2 such that if $0 < \varepsilon \leq \varepsilon_2$, $|\delta| \leq \delta_2$, $\operatorname{Re}\lambda \geq -\frac{3}{4}\kappa_1 T_a$ and $|\operatorname{Im}\lambda| \leq \frac{a}{2}$, then for any given $F \in \mathcal{X}_a$, problem (7.15) has a unique solution $U \in \tilde{\mathcal{Q}}^\varepsilon \mathcal{Y}_a$ and U satisfies the estimate*

$$\|U\|_{\varepsilon, \mathcal{Y}_a} \leq C \|F\|_{\varepsilon, \mathcal{X}_a}$$

uniformly for $0 < \varepsilon \leq \varepsilon_2$, $|\delta| \leq \delta_2$ and λ with $\operatorname{Re}\lambda \geq -\frac{3}{4}\kappa_1 T_a$ and $|\operatorname{Im}\lambda| \leq \frac{a}{2}$.

Proof. We regard $\tilde{\mathcal{Q}}^\varepsilon B_\delta^\varepsilon \tilde{\mathcal{Q}}^\varepsilon = B^\varepsilon(\omega_\delta^\varepsilon) + \delta \tilde{\mathcal{Q}}^\varepsilon M_\delta^\varepsilon$ as a perturbation of $B^\varepsilon(\omega_\delta^\varepsilon)$. By Lemma 5.5 and Lemma 6.2, we see that if $\operatorname{Re}\lambda \geq -\frac{3}{4}\kappa_1 T_a$ and $|\operatorname{Im}\lambda| \leq \frac{a}{2}$, then

$$\begin{aligned} \|\mathcal{S}^\varepsilon(\lambda, \omega_\delta^\varepsilon) \tilde{\mathcal{Q}}^\varepsilon \tilde{M}_\delta^\varepsilon u\|_{\varepsilon, \mathcal{Y}_a} &\leq \frac{1}{2} \|\mathcal{S}^\varepsilon(\lambda, \omega_\delta^\varepsilon) \tilde{\mathcal{Q}}^\varepsilon \tilde{M}_\delta^\varepsilon u\|_{\varepsilon, \mathcal{Y}_a} \\ &\quad + C \{ \|u\|_{\varepsilon, \mathcal{Y}_a} + \|\hat{u}_\delta^\varepsilon\|_{\varepsilon, \mathcal{Y}_a} \|u\|_{\varepsilon, \mathcal{Y}_a} \} \end{aligned}$$

uniformly for $u \in \tilde{\mathcal{Q}}^\varepsilon \mathcal{Y}_a$, $0 < \varepsilon \leq \varepsilon_2$, $|\delta| \leq \delta_2$ and $\operatorname{Re}\lambda \geq -\frac{3}{4}\kappa_1 T_a$. This implies that there exists a positive constant δ_2 such that if $|\delta| \leq \delta_2$, then $I + \delta \mathcal{S}^\varepsilon(\lambda, \omega_\delta^\varepsilon) \tilde{\mathcal{Q}}^\varepsilon M_\delta^\varepsilon \tilde{\mathcal{Q}}^\varepsilon$ has a bounded inverse $(I + \delta \mathcal{S}^\varepsilon(\lambda, \omega_\delta^\varepsilon) \tilde{\mathcal{Q}}^\varepsilon M_\delta^\varepsilon \tilde{\mathcal{Q}}^\varepsilon)^{-1}$ on $\tilde{\mathcal{Q}}^\varepsilon \mathcal{Y}_a$ with estimate

$$\|(I + \delta \mathcal{S}^\varepsilon(\lambda, \omega_\delta^\varepsilon) \tilde{\mathcal{Q}}^\varepsilon M_\delta^\varepsilon \tilde{\mathcal{Q}}^\varepsilon)^{-1} u\|_{\varepsilon, \mathcal{Y}_a} \leq C \|u\|_{\varepsilon, \mathcal{Y}_a}$$

uniformly for $0 < \varepsilon \leq \varepsilon_2$, $|\delta| \leq \delta_2$ and $\operatorname{Re}\lambda \geq -\frac{3}{4}\kappa_1 T_a$. It then follows that (7.15) has a unique solution $U = (I + \delta \mathcal{S}^\varepsilon(\lambda, \omega_\delta^\varepsilon) \tilde{\mathcal{Q}}^\varepsilon M_\delta^\varepsilon \tilde{\mathcal{Q}}^\varepsilon)^{-1} \mathcal{S}^\varepsilon(\lambda, \omega_\delta^\varepsilon) F \in \tilde{\mathcal{Q}}^\varepsilon \mathcal{Y}_a$ and U satisfies the estimate

$$\|U\|_{\varepsilon, \mathcal{Y}_a} \leq C \|\mathcal{S}^\varepsilon(\lambda, \omega_\delta^\varepsilon) \tilde{\mathcal{Q}}^\varepsilon F\|_{\varepsilon, \mathcal{Y}_a} \leq C \|F\|_{\varepsilon, \mathcal{X}_a}$$

uniformly for $0 < \varepsilon \leq \varepsilon_2$, $|\delta| \leq \delta_2$, $\operatorname{Re}\lambda \geq -\frac{3}{4}\kappa_1 T_a$ and $|\operatorname{Im}\lambda| \leq \frac{a}{2}$. This completes the proof. \square

Let $0 < \varepsilon \leq \varepsilon_2$ and $|\delta| \leq \delta_2$. For λ satisfying $\operatorname{Re}\lambda \geq -\frac{3}{4}\kappa_1 T_a$, we denote the solution operator for (7.15) by $\tilde{\mathcal{S}}_\delta^\varepsilon(\lambda)$.

In terms of $\tilde{\mathcal{S}}_\delta^\varepsilon(\lambda)$, (7.14) is written as

$$U = \tilde{\mathcal{S}}_\delta^\varepsilon(\lambda) \tilde{\mathcal{Q}}^\varepsilon F - \delta \sum_{j=0,1} \zeta_j \tilde{\mathcal{S}}_\delta^\varepsilon(\lambda) \tilde{\mathcal{Q}}^\varepsilon M_\delta^\varepsilon z_j^\varepsilon.$$

Substituting this into (7.12) and (7.13) and using the fact that $\llbracket M(z_0^\varepsilon, z_j^\varepsilon) \rrbracket_{k,\varepsilon} = 0$, ($j, k \in \{0, 1\}$), we have

$$\lambda \zeta + \delta^2 \Gamma_\delta^\varepsilon(\lambda) \zeta = \mathcal{F}_\delta^\varepsilon(\lambda) F. \quad (7.16)$$

Here $\zeta = {}^\top(\zeta_0, \zeta_1) \in \mathbb{C}^2$; $\Gamma_\delta^\varepsilon(\lambda)$ is a 2×2 matrix given by

$$\Gamma_\delta^\varepsilon(\lambda) = \begin{pmatrix} \tilde{\eta}_\delta^\varepsilon \llbracket K z_0^\varepsilon \rrbracket_{0,\varepsilon} & a^\varepsilon \tilde{\omega}_\delta^\varepsilon + \tilde{\eta}_\delta^\varepsilon \llbracket K z_1^\varepsilon \rrbracket_{0,\varepsilon} \\ + \llbracket M(U_\delta^\varepsilon, z_0^\varepsilon) \rrbracket_{0,\varepsilon} & + \llbracket M(U_\delta^\varepsilon, z_1^\varepsilon) \rrbracket_{0,\varepsilon} \\ -a^\varepsilon \tilde{\omega}_\delta^\varepsilon + \tilde{\eta}_\delta^\varepsilon \llbracket K z_0^\varepsilon \rrbracket_{1,\varepsilon} & \tilde{\eta}_\delta^\varepsilon \llbracket K z_1^\varepsilon \rrbracket_{1,\varepsilon} \\ + \llbracket M(U_\delta^\varepsilon, z_0^\varepsilon) \rrbracket_{1,\varepsilon} & + \llbracket M(U_\delta^\varepsilon, z_1^\varepsilon) \rrbracket_{1,\varepsilon} \end{pmatrix} \\ - \begin{pmatrix} \llbracket M_\delta^\varepsilon \tilde{S}_\delta^\varepsilon(\lambda) \tilde{Q}^\varepsilon M_\delta^\varepsilon z_0^\varepsilon \rrbracket_{0,\varepsilon} & \llbracket M_\delta^\varepsilon \tilde{S}_\delta^\varepsilon(\lambda) \tilde{Q}^\varepsilon M_\delta^\varepsilon z_1^\varepsilon \rrbracket_{0,\varepsilon} \\ \llbracket M_\delta^\varepsilon \tilde{S}_\delta^\varepsilon(\lambda) \tilde{Q}^\varepsilon M_\delta^\varepsilon z_0^\varepsilon \rrbracket_{1,\varepsilon} & \llbracket M_\delta^\varepsilon \tilde{S}_\delta^\varepsilon(\lambda) \tilde{Q}^\varepsilon M_\delta^\varepsilon z_1^\varepsilon \rrbracket_{1,\varepsilon} \end{pmatrix}$$

and

$$\mathcal{F}_\delta^\varepsilon(\lambda)F = \begin{pmatrix} \llbracket F \rrbracket_{0,\varepsilon} \\ \llbracket F \rrbracket_{1,\varepsilon} \end{pmatrix} - \delta \begin{pmatrix} \llbracket M_\delta^\varepsilon \tilde{S}_\delta^\varepsilon(\lambda) \tilde{Q}^\varepsilon F \rrbracket_{0,\varepsilon} \\ \llbracket M_\delta^\varepsilon \tilde{S}_\delta^\varepsilon(\lambda) \tilde{Q}^\varepsilon F \rrbracket_{1,\varepsilon} \end{pmatrix}.$$

Proposition 7.6 *There exists a positive constant c_0 such that if $0 < \varepsilon \leq \varepsilon_2$ and $|\delta| \leq \delta_2$ then $\{\lambda \in \mathbb{C}; |\lambda| > c_0 \delta^2, \operatorname{Re} \lambda \geq -\frac{3}{4} \kappa_1 T_a, |\operatorname{Im} \lambda| \leq \frac{a}{2}\} \subset \rho(-B_\delta^\varepsilon)$ and*

$$\|(\lambda + B_\delta^\varepsilon)^{-1} F\|_{\varepsilon, \mathcal{Y}_a} \leq C \left(\frac{1}{|\lambda| - c_0 |\delta|^2} + 1 \right) \|F\|_{\varepsilon, \mathcal{X}_a}$$

uniformly for $0 < \varepsilon \leq \varepsilon_2$, $|\delta| \leq \delta_2$ and λ with $|\lambda| > c_0 |\delta|^2$, $\operatorname{Re} \lambda \geq -\frac{3}{4} \kappa_1 T_a$ and $|\operatorname{Im} \lambda| \leq \frac{a}{2}$.

Proof. By using Proposition 7.5, we see that

$$|\Gamma_\delta^\varepsilon(\lambda) \zeta| \leq c_0 |\zeta|$$

uniformly for $0 < \varepsilon \leq \varepsilon_2$, $|\delta| \leq \delta_2$ and λ with $\operatorname{Re} \lambda \geq -\frac{3}{4} \kappa_1 T_a$. It then follows that if $|\lambda| > c_0 |\delta|^2$, $\operatorname{Re} \lambda \geq -\frac{3}{4} \kappa_1 T_a$ and $|\operatorname{Im} \lambda| \leq \frac{a}{2}$, then (7.16) has a unique solution $\zeta \in \mathbb{C}^2$ with estimate

$$|\zeta| \leq \frac{C}{|\lambda| - c_0 |\delta|^2} \|F\|_{\varepsilon, \mathcal{X}_a},$$

from which we deduce that $\{\lambda; |\lambda| > c_0 |\delta|^2, \operatorname{Re} \lambda \geq -\frac{3}{4} \kappa_1 T_a, |\operatorname{Im} \lambda| \leq \frac{a}{2}\} \subset \rho(-B_\delta^\varepsilon)$ and

$$(\lambda + B_\delta^\varepsilon)^{-1} F = (z_0^\varepsilon \ z_1^\varepsilon)(\lambda + \delta^2 \Gamma_\delta^\varepsilon(\lambda))^{-1} \mathcal{F}_\delta^\varepsilon(\lambda) F + \tilde{S}_\delta^\varepsilon(\lambda) \tilde{Q}^\varepsilon F,$$

$$\|(\lambda + \tilde{B}_\delta^\varepsilon)^{-1} F\|_{\varepsilon, \mathcal{Y}_a} \leq C \left(\frac{1}{|\lambda| - c_0 |\delta|^2} + 1 \right) \|F\|_{\varepsilon, \mathcal{X}_a}.$$

This completes the proof. \square

We take a positive constant δ_2 so that $c_0\delta_2^2 < \frac{3}{4}\kappa_1 T_a$. It then follows from Proposition 7.6 that

$$\sigma(-B_\delta^\varepsilon) \cap \{\lambda \in \mathbb{C}; \operatorname{Re}\lambda \geq -\frac{3}{4}\kappa_1 T_a, |\operatorname{Im}\lambda| \leq \frac{a}{2}\} \subset \{\lambda \in \mathbb{C}; |\lambda| < c_0\delta^2\}$$

for $0 < \varepsilon \leq \varepsilon_2$ and $|\delta| \leq \delta_2$. We shall, therefore, investigate

$$\sigma(-B_\delta^\varepsilon) \cap \{\lambda \in \mathbb{C}; |\lambda| \leq c_0\delta^2\}$$

for $0 < \varepsilon \leq \varepsilon_2$ and $|\delta| \leq \delta_2$. From the argument above, we see that

$$\lambda \in \sigma(-B_\delta^\varepsilon) \cap \{\lambda \in \mathbb{C}; |\lambda| \leq c_0\delta^2\} \text{ if and only if } \det(\lambda I + \delta^2 \Gamma_\delta^\varepsilon(\lambda)) = 0.$$

We thus consider zeros of $\det(\lambda I + \delta^2 \Gamma_\delta^\varepsilon(\lambda))$. To this end, we regard $\Gamma_\delta^\varepsilon(\lambda)$ as a perturbation of $\Gamma_0^\varepsilon(0)$.

Proposition 7.7 *If $|\lambda| \leq c_1|\delta|$, then*

$$|\Gamma_\delta^\varepsilon(\lambda) - \Gamma_0^\varepsilon(0)| \leq C|\delta|$$

uniformly for $0 < \varepsilon \leq \varepsilon_2$, $|\delta| \leq \delta_2$ and $|\lambda| \leq c_1|\delta|$.

Proof. We first prove

$$\|\tilde{\mathcal{S}}_\delta^\varepsilon(\lambda) \tilde{\mathcal{Q}}^\varepsilon M_\delta^\varepsilon z_j^\varepsilon - \mathcal{S}^\varepsilon(0, 0) \tilde{\mathcal{Q}}^\varepsilon M_0^\varepsilon z_j^\varepsilon\|_{\varepsilon, \mathcal{Y}_a} \leq C|\delta| \quad (j = 0, 1) \quad (7.17)$$

uniformly for $0 < \varepsilon \leq \varepsilon_2$, $|\delta| \leq \delta_2$ and $|\lambda| \leq c_1|\delta|$. To show (7.17), we write

$$\tilde{\mathcal{S}}_\delta^\varepsilon(\lambda) \tilde{\mathcal{Q}}^\varepsilon M_\delta^\varepsilon z_j^\varepsilon - \mathcal{S}^\varepsilon(0, 0) \tilde{\mathcal{Q}}^\varepsilon M_0^\varepsilon z_j^\varepsilon = I_1 + I_2 + I_3 + I_4,$$

where

$$\begin{aligned} I_1 &= \tilde{\mathcal{S}}_\delta^\varepsilon(\lambda) \tilde{\mathcal{Q}}^\varepsilon (M_\delta^\varepsilon z_j^\varepsilon - M_0^\varepsilon z_j^\varepsilon), & I_2 &= (\tilde{\mathcal{S}}_\delta^\varepsilon(\lambda) - \tilde{\mathcal{S}}_\delta^\varepsilon(0)) \tilde{\mathcal{Q}}^\varepsilon M_0^\varepsilon z_j^\varepsilon, \\ I_3 &= (\tilde{\mathcal{S}}_\delta^\varepsilon(0) - \mathcal{S}^\varepsilon(0, \omega_\delta^\varepsilon)) \tilde{\mathcal{Q}}^\varepsilon M_0^\varepsilon z_j^\varepsilon, & I_4 &= (\mathcal{S}^\varepsilon(0, \omega_\delta^\varepsilon) - \mathcal{S}^\varepsilon(0, 0)) \tilde{\mathcal{Q}}^\varepsilon M_0^\varepsilon z_j^\varepsilon. \end{aligned}$$

As for I_1 , since $M_\delta^\varepsilon z_j^\varepsilon - M_0^\varepsilon z_j^\varepsilon = \delta \tilde{\eta}_\delta^\varepsilon K z_j^\varepsilon + \delta M(U_\delta^\varepsilon, z_j^\varepsilon)$, we have $\|I_1\|_{\varepsilon, \mathcal{Y}_a} \leq C|\delta|$. To estimate I_2 , we note that

$$\tilde{\mathcal{S}}_\delta^\varepsilon(\lambda) = \tilde{\mathcal{S}}_\delta^\varepsilon(0) \sum_{N=0}^{\infty} (-1)^N \lambda^N \tilde{\mathcal{S}}_\delta^\varepsilon(0)^N.$$

This, together with Proposition 7.5, implies that

$$\|I_2\|_{\varepsilon, \mathcal{Y}_a} \leq C|\lambda| \|\tilde{\mathcal{S}}_\delta^\varepsilon(0)M_0^\varepsilon z_j^\varepsilon\|_{\varepsilon, \mathcal{Y}_a} \leq C|\delta|.$$

As for I_3 , we see from the proof of Proposition 7.5 that

$$\|I_3\|_{\varepsilon, \mathcal{Y}_a} \leq C|\delta| \|\mathcal{S}^\varepsilon(0, \omega_\delta^\varepsilon)\tilde{\mathcal{Q}}^\varepsilon M_0^\varepsilon z_j^\varepsilon\|_{\varepsilon, \mathcal{Y}_a} \leq C|\delta|.$$

Since $(\mathcal{S}^\varepsilon(0, \omega_\delta^\varepsilon) - \mathcal{S}^\varepsilon(0, 0))\tilde{\mathcal{Q}}^\varepsilon = (B^\varepsilon(\omega_\delta^\varepsilon)\tilde{\mathcal{Q}}^\varepsilon)^{-1} - (B^\varepsilon\tilde{\mathcal{Q}}^\varepsilon)^{-1}$ and $M_0^\varepsilon z_j^\varepsilon = M(z_0^\varepsilon, z_j^\varepsilon)$, we estimate I_4 as in the proof of Theorem 4.2 (i) to obtain $\|I_4\|_{\varepsilon, \mathcal{Y}_a} \leq C|\delta|$. The estimate (7.17) is thus proved.

It follows from (7.17) that

$$|\|M_\delta^\varepsilon \tilde{\mathcal{S}}_\delta^\varepsilon(\lambda)\tilde{\mathcal{Q}}^\varepsilon M_\delta^\varepsilon z_j^\varepsilon\|_{k, \varepsilon} - \|M_\delta^\varepsilon \mathcal{S}^\varepsilon(0, 0)\tilde{\mathcal{Q}}^\varepsilon M_\delta^\varepsilon z_j^\varepsilon\|_{k, \varepsilon}| \leq C|\delta|, \quad (j, k \in \{0, 1\})$$

uniformly for $0 < \varepsilon \leq \varepsilon_2$, $|\delta| \leq \delta_2$ and $\lambda \in \Sigma$. Since $\tilde{\eta}_\delta^\varepsilon = \tilde{\eta}_0^\varepsilon + O(\delta)$ and $\tilde{\omega}_\delta^\varepsilon = \tilde{\omega}_0^\varepsilon + O(\delta)$, we conclude that $|\Gamma_\delta^\varepsilon(\lambda) - \Gamma_0^\varepsilon(0)| \leq C|\delta|$ uniformly for $0 < \varepsilon \leq \varepsilon_2$, $|\delta| \leq \delta_2$ and $|\lambda| \leq c_1|\delta|$. This completes the proof. \square

We are now in a position to prove Theorem 7.4. We set

$$D_\delta^\varepsilon(\tilde{\lambda}) = \det\left(\tilde{\lambda} + \Gamma_\delta^\varepsilon(\delta^2\tilde{\lambda})\right).$$

If $D_\delta^\varepsilon(\tilde{\lambda}) \neq 0$, then $\lambda = \delta^2\tilde{\lambda} \in \rho(-B_\delta^\varepsilon)$ and

$$(\lambda + B_\delta^\varepsilon)^{-1}F = \begin{pmatrix} z_0^\varepsilon & z_1^\varepsilon \end{pmatrix} (\lambda + \delta^2\Gamma_\delta^\varepsilon(\lambda))^{-1}\mathcal{F}_\delta^\varepsilon(\lambda)F + \tilde{\mathcal{S}}_\delta^\varepsilon(\lambda)\tilde{\mathcal{Q}}^\varepsilon F.$$

If $D_\delta^\varepsilon(\tilde{\lambda}) = 0$, then $\lambda = \delta^2\tilde{\lambda} \in \sigma(-B_\delta^\varepsilon)$. Therefore, we investigate zeros of $D_\delta^\varepsilon(\tilde{\lambda})$ in $\{\tilde{\lambda} \in \mathbb{C}; |\tilde{\lambda}| \leq \tilde{c}_0\}$, where $\tilde{c}_0 = \frac{c_0}{\delta_2}$. We shall prove the following proposition.

Proposition 7.8 *There exists a positive constant δ_2 such that if $0 < \varepsilon \leq \varepsilon_2$ and $|\delta| \leq \delta_2$ then $\tilde{D}_\delta^\varepsilon(\tilde{\lambda})$ has two zeros $\tilde{\lambda} = 0$ and $\tilde{\lambda}_\delta^\varepsilon$ in $\{\tilde{\lambda} \in \mathbb{C}; |\tilde{\lambda}| \leq \tilde{c}_0\}$ and both zeros are of order 1. Here $\tilde{\lambda}_\delta^\varepsilon$ satisfies $\tilde{\lambda}_\delta^\varepsilon = \tilde{\lambda}_0^\varepsilon + O(\delta)$, $\tilde{\lambda}_0^\varepsilon = 2\tilde{\eta}_0^\varepsilon \operatorname{Re}[Kz_0^\varepsilon]_{+, \varepsilon}$.*

It follows from Proposition 7.8 that

$$\sigma(-B_\delta^\varepsilon) \cap \{\lambda \in \mathbb{C}; |\lambda| \leq c_0\delta^2\} = \{0, \lambda_\delta^\varepsilon\},$$

where 0 and $\lambda_\delta^\varepsilon = \delta^2\tilde{\lambda}_\delta^\varepsilon$ are simple eigenvalues of $-B_\delta^\varepsilon$. This proves Theorem 7.4. Furthermore, we shall see from the proof of Proposition 7.8 below that

the eigenspaces for the eigenvalues 0 and $\lambda_\delta^\varepsilon$ are spanned by $\partial_t u_\delta^\varepsilon$ and $z_0^\varepsilon + O(\delta)$, respectively.

Let us prove Proposition 7.8.

Proof of Proposition 7.8. We first observe that

$$\Gamma_\delta^\varepsilon(0) = \begin{pmatrix} \gamma_{11,\delta}^\varepsilon(0) & 0 \\ \gamma_{21,\delta}^\varepsilon(0) & 0 \end{pmatrix}, \quad (7.18)$$

where

$$\begin{aligned} \gamma_{11,\delta}^\varepsilon(\lambda) &= \tilde{\eta}_\delta^\varepsilon \llbracket K z_0^\varepsilon \rrbracket_{0,\varepsilon} + \llbracket M(U_\delta^\varepsilon, z_0^\varepsilon) \rrbracket_{0,\varepsilon} - \llbracket M_\delta^\varepsilon \tilde{\mathcal{S}}_\delta^\varepsilon(\lambda) \tilde{\mathcal{Q}}^\varepsilon M_\delta^\varepsilon z_0^\varepsilon \rrbracket_{0,\varepsilon}, \\ \gamma_{21,\delta}^\varepsilon(\lambda) &= -a^\varepsilon \tilde{\omega}_\delta^\varepsilon + \tilde{\eta}_\delta^\varepsilon \llbracket K z_0^\varepsilon \rrbracket_{1,\varepsilon} + \llbracket M(U_\delta^\varepsilon, z_0^\varepsilon) \rrbracket_{1,\varepsilon} - \llbracket M_\delta^\varepsilon \tilde{\mathcal{S}}_\delta^\varepsilon(\lambda) \tilde{\mathcal{Q}}^\varepsilon M_\delta^\varepsilon z_0^\varepsilon \rrbracket_{1,\varepsilon}. \end{aligned}$$

Indeed, differentiating the equation

$$B^\varepsilon(\omega_\delta^\varepsilon)u_\delta^\varepsilon + \eta_\delta^\varepsilon K u_\delta^\varepsilon + N(u_\delta^\varepsilon) = 0 \quad (7.19)$$

in t , we have

$$B_\delta^\varepsilon \partial_t U_\delta^\varepsilon + \delta \frac{a^\varepsilon}{a} \tilde{\omega}_\delta^\varepsilon \partial_t^2 z_0^\varepsilon + M_\delta^\varepsilon \partial_t z_0^\varepsilon = 0.$$

This, together with $\partial_t z_0^\varepsilon = -a z_1^\varepsilon$ and Proposition 7.6, gives (7.18). It then follows that $\tilde{\lambda} = 0$ is a zero of $D_\delta^\varepsilon(\tilde{\lambda})$.

We next claim that

$$D_0^\varepsilon(\tilde{\lambda}) = \tilde{\lambda}(\tilde{\lambda} + \tilde{\lambda}_0^\varepsilon). \quad (7.20)$$

This can be verified by using (7.19). Indeed, we see from (7.19) that

$$B^\varepsilon(\omega_\delta^\varepsilon)U_\delta^\varepsilon + \frac{a^\varepsilon}{a} \delta \tilde{\omega}_\delta^\varepsilon \partial_t z_0^\varepsilon + \delta \tilde{\eta}_\delta^\varepsilon K(z_0^\varepsilon + \delta U_\delta^\varepsilon) + N(z_0^\varepsilon + \delta U_\delta^\varepsilon) = 0. \quad (7.21)$$

Noting that $\llbracket M(z_0^\varepsilon, z_0^\varepsilon) \rrbracket_{j,\varepsilon} = 0$ ($j = 0, 1$), we have

$$\tilde{\eta}_\delta^\varepsilon \llbracket K z_0^\varepsilon \rrbracket_{0,\varepsilon} + \delta \tilde{\eta}_\delta^\varepsilon \llbracket K U_\delta^\varepsilon \rrbracket_{0,\varepsilon} + \llbracket M(U_\delta^\varepsilon, z_0^\varepsilon) \rrbracket_{0,\varepsilon} + \frac{\delta}{2} \llbracket M(U_\delta^\varepsilon, U_\delta^\varepsilon) \rrbracket_{0,\varepsilon} = 0$$

and

$$U_\delta^\varepsilon = -\mathcal{S}^\varepsilon(0, \omega_\delta^\varepsilon) \tilde{\mathcal{Q}}^\varepsilon \{ \delta \tilde{\eta}_\delta^\varepsilon K(z_0^\varepsilon + \delta U_\delta^\varepsilon) + N(z_0^\varepsilon + \delta U_\delta^\varepsilon) \}.$$

Letting $\delta \rightarrow 0$, we obtain

$$\tilde{\eta}_0^\varepsilon \llbracket K z_0^\varepsilon \rrbracket_{0,\varepsilon} + \llbracket M(U_0^\varepsilon, z_0^\varepsilon) \rrbracket_{0,\varepsilon} = 0 \quad (7.22)$$

and

$$U_0^\varepsilon = -\frac{1}{2}\tilde{\mathcal{S}}_0^\varepsilon(0)\tilde{\mathcal{Q}}^\varepsilon M(z_0^\varepsilon, z_0^\varepsilon). \quad (7.23)$$

We deduce from (7.22) and (7.23) that

$$\gamma_{11,0}^\varepsilon(0) = -\llbracket M(z_0^\varepsilon, \tilde{\mathcal{S}}_0^\varepsilon(0)M(z_0^\varepsilon, z_0^\varepsilon)) \rrbracket_{0,\varepsilon} = -2\tilde{\eta}_0^\varepsilon \llbracket K z_0^\varepsilon \rrbracket_{0,\varepsilon} = -\tilde{\lambda}_0^\varepsilon,$$

and hence,

$$\Gamma_0^\varepsilon(0) = \begin{pmatrix} -\tilde{\lambda}_0^\varepsilon & 0 \\ \gamma_0^\varepsilon(0) & 0 \end{pmatrix}.$$

This implies (7.20).

We take δ_2 so small that $\tilde{c}_0 = \frac{c_0}{\delta_2} \leq 2|\tilde{\lambda}_0^\varepsilon|$. By Proposition 7.6, we obtain

$$|D_\delta^\varepsilon(\tilde{\lambda}) - D_0^\varepsilon(\tilde{\lambda})| \leq c_1|\delta|$$

for $0 < \varepsilon \leq \varepsilon_2$, $|\delta| \leq \delta_2$ and $|\tilde{\lambda}| \leq \tilde{c}_0$. Therefore, there exists a positive constant δ_2 such that if $|\delta| \leq \delta_2$, then $|D_0^\varepsilon(\tilde{\lambda})| > |D_\delta^\varepsilon(\tilde{\lambda}) - D_0^\varepsilon(\tilde{\lambda})|$ on $|\tilde{\lambda}| = \tilde{c}_0$. Applying the Rouché theorem, we find that $D_\delta^\varepsilon(\tilde{\lambda})$ has two zeros in $\{\tilde{\lambda} \in \mathbb{C}; |\tilde{\lambda}| \leq \tilde{c}_0\}$; one of them is $\tilde{\lambda} = 0$. Furthermore, we see that, by taking δ_2 smaller if necessary, that there exists a positive constant c_3 such that if $|\delta| \leq \delta_2$, then $|D_0^\varepsilon(\tilde{\lambda})| > |D_\delta^\varepsilon(\tilde{\lambda}) - D_0^\varepsilon(\tilde{\lambda})|$ on $|\tilde{\lambda} - \tilde{\lambda}_0^\varepsilon| = c_3|\delta|$. The Rouché theorem then implies that $D_\delta^\varepsilon(\tilde{\lambda})$ has a zero $\tilde{\lambda}_\delta^\varepsilon$ of order 1 in $\{\tilde{\lambda} \in \mathbb{C}; |\tilde{\lambda} - \tilde{\lambda}_0^\varepsilon| < c_3|\delta|\}$. This completes the proof. \square

Proof of Theorem 7.1 Let μ_0 and μ_1 be eigenvalues of $\mathcal{U}_\delta^\varepsilon(T_a, 0)$ with $|\mu_j - 1| < r$ ($j = 0, 1$). We know that one of μ_j 's, say μ_0 , is equal to 1. On the other hand, as in the proof of Theorem 5.4, i.e., the proof of [11, Theorem 4.2], we see from (7.3)–(7.5) that if $e^{\frac{a}{\alpha^\varepsilon}T_a\lambda} \in \rho(\mathcal{U}_\delta^\varepsilon(T_a, 0))$, then $\lambda \in \rho(-B_\delta^\varepsilon)$. Therefore, since $\sigma(\mathcal{U}_\delta^\varepsilon(T_a, 0)) \cap \{\mu; |\mu - 1| < r\} = \{\mu_0, \mu_1\}$, by Theorem 7.4, we conclude that $\mu_0 = 1$, $\mu_1 = e^{\frac{a}{\alpha^\varepsilon}T_a\lambda_\delta^\varepsilon}$. This completes the proof. \square

Acknowledgements. Y. Kagei was partly supported by JSPS KAKENHI Grant Numbers 16H03947, 16H06339 and 20H00118. T. Nishida is partly supported by JSPS KAKENHI Grant Number 20K03699.

References

- [1] J.-L. Bona, C.-H. Hsia, T. Ma and S. Wang, Hopf bifurcation for two-dimensional doubly diffusive convection, *Appl. Anal.*, **90** (2011), pp. 5–30.

- [2] A. Chorin, The numerical solution of the Navier-Stokes equations for an incompressible fluid, *Bull. Amer. Math. Soc.*, **73** (1967), pp. 928–931.
- [3] A. Chorin, A numerical method for solving incompressible viscous flow problems, *J. Comput. Phys.*, **2** (1967), pp. 12–26.
- [4] A. Chorin, Numerical solution of the Navier-Stokes equations, *Math. Comp.*, **22** (1968), pp. 745–762.
- [5] D. Donatelli, On the artificial compressibility method for the Navier-Stokes-Fourier system, *Quart. Appl. Math.*, **68** (2010), pp. 469–485.
- [6] D. Donatelli, The artificial compressibility approximation for MHD equations in unbounded domain, *J. Hyperbolic Differential Equations*, **10** (2013), pp. 181–198.
- [7] D. Donatelli and P. Marcati, A dispersive approach to the artificial compressibility approximations of the Navier-Stokes equations in 3D, *J. Hyperbolic Differential Equations*, **3** (2006), pp. 575–588.
- [8] D. Donatelli and P. Marcati, Leray weak solutions of the incompressible Navier Stokes system on exterior domains via the artificial compressibility method, *Indiana Univ. Math. J.*, **59** (2010), pp. 1831–1852.
- [9] G. P. Galdi, *An Introduction to the Mathematical Theory of the Navier-Stokes Equations*, Vol. 1, Springer-Verlag New York (1994).
- [10] C.-H. Hsia, T. Ma, and S. Wang, Bifurcation and stability of two-dimensional double diffusive convection, *Commun. Pure Appl. Anal.*, **7** (2008), pp. 23–48.
- [11] C.-H. Hsia, Y. Kagei, T. Nishida and Y. Teramoto, Singular limit in Hopf bifurcation for doubly diffusive convection equations I: linearized analysis at criticality, preprint.
- [12] Y. Kagei and T. Nishida, On Chorin’s method for stationary solutions of the Oberbeck-Boussinesq equation, *J. Math. Fluid Mech.*, **19** (2017), pp. 345–365.
- [13] Y. Kagei, T. Nishida and Y. Teramoto, On the spectrum for the artificial compressible system, *J. Differential Equations*, **264** (2018), pp. 897–928.

- [14] Y. Kagei and Y. Teramoto, On the spectrum of the linearized operator around compressible Couette flows between two concentric cylinders, *J. Math. Fluid Mech.*, **22** (2020), no. 2, Paper No. 21, 23 pp.
- [15] M. Murata, On a maximal L_p - L_q approach to the compressible viscous fluid flow with slip boundary condition, *Nonlinear Anal.*, **106** (2014), pp. 86–109.
- [16] H. Sohr, *The Navier-Stokes equations: an elementary functional analytic approach*, Birkhäuser, Basel (2001).
- [17] R. Témam, Sur l’approximation de la solution des équations de Navier-Stokes par la méthode des pas fractionnaires. I, *Arch. Rational Mech. Anal.*, **32** (1969), pp. 135–153.
- [18] R. Témam, Sur l’approximation de la solution des équations de Navier-Stokes par la méthode des pas fractionnaires. II, *Arch. Rational Mech. Anal.* **33** (1969), pp. 377–385.
- [19] R. Temam, *Navier-Stokes equations. Theory and numerical analysis*, reprint of the 1984 edition, AMS Chelsea Publishing, Providence, RI, 2001.
- [20] Y. Teramoto, Stability of bifurcating stationary solutions of the artificial compressible system, *J. Math. Fluid Mech.*, **20** (2018), pp. 1213–1228.