

RATIONALITY OVER NON-CLOSED FIELDS OF FANO THREEFOLDS WITH HIGHER GEOMETRIC PICARD RANK

ALEXANDER KUZNETSOV AND YURI PROKHOROV

ABSTRACT. We prove rationality criteria over algebraically non-closed fields of characteristic 0 for five out of six types of geometrically rational Fano threefolds of Picard number 1 and geometric Picard number bigger than 1. For the last type of such threefolds we provide a unirationality criterion and prove stable non-rationality under additional assumptions.

1. INTRODUCTION

1.1. The results. The goal of this paper is to discuss rationality of smooth Fano threefolds over non-closed fields of characteristic 0. In [KP19] we considered the case of geometrically rational Fano threefolds with geometric Picard number $\rho(X_{\bar{k}}) = 1$ and here we switch the focus to the case of geometrically rational Fano threefolds X with Picard numbers

$$(1.1.1) \quad \rho(X) = 1 \quad \text{and} \quad \rho(X_{\bar{k}}) > 1.$$

In fact, Fano threefolds satisfying (1.1.1) have been classified in [Pro13b] and [AB92] explains which of these are geometrically rational. A combination of these results gives the following

Theorem 1.1 ([Pro13b, Theorem 1.2], [AB92]). *There are exactly six families of geometrically rational Fano threefolds satisfying (1.1.1) as listed in the following table.*

	$\iota(X_{\bar{k}})$	$\rho(X_{\bar{k}})$	$-K_X^3$	$g(X)$	$h^{1,2}(X_{\bar{k}})$	$X_{\bar{k}}$
$X_{(3,3)}$	1	2	20	11	3	<i>an intersection of three divisors of bidegree $(1, 1)$ in $\mathbb{P}_{\bar{k}}^3 \times \mathbb{P}_{\bar{k}}^3$</i>
$X_{(1,1,1,1)}$	1	4	24	13	1	<i>a divisor of multidegree $(1, 1, 1, 1)$ on $(\mathbb{P}_{\bar{k}}^1)^4$</i>
$X_{(4,4)}$	1	2	28	15	0	<i>the blow-up of a smooth quadric $Q_{\bar{k}} \subset \mathbb{P}_{\bar{k}}^4$ along a linearly normal smooth rational quartic curve</i>
$X_{(2,2,2)}$	1	3	30	16	0	<i>an intersection of three divisors of multidegrees $(0, 1, 1)$, $(1, 0, 1)$, $(1, 1, 0)$ in $\mathbb{P}_{\bar{k}}^2 \times \mathbb{P}_{\bar{k}}^2 \times \mathbb{P}_{\bar{k}}^2$</i>
$X_{(2,2)}$	2	2	48	25	0	<i>a divisor of bidegree $(1, 1)$ on $\mathbb{P}_{\bar{k}}^2 \times \mathbb{P}_{\bar{k}}^2$</i>
$X_{(1,1,1)}$	2	3	48	25	0	$\mathbb{P}_{\bar{k}}^1 \times \mathbb{P}_{\bar{k}}^1 \times \mathbb{P}_{\bar{k}}^1$

TABLE 1. Geometrically rational Fano threefolds X satisfying (1.1.1)

The paper was partially supported by the HSE University Basic Research Program.

The first column of Table 1 contains the name for the family we use in this paper, the next columns contain the index $\iota(X_{\bar{k}})$, defined as

$$\iota(X_{\bar{k}}) = \max \left\{ i \mid \frac{1}{i} K_{X_{\bar{k}}} \in \text{Pic}(X_{\bar{k}}) \right\},$$

the geometric Picard number $\rho(X_{\bar{k}})$, the anticanonical degree $(-K_X)^3$, the genus $g(X)$, defined by

$$(-K_X)^3 = 2g(X) - 2.$$

and the Hodge number $h^{1,2}(X_{\bar{k}})$ of the threefold, while the last column provides a geometric description of these varieties over an algebraic closure \bar{k} of the field.

We discuss some geometric properties of threefolds from Table 1 in §2. In particular, we describe their extremal contractions over \bar{k} and identify their Hilbert schemes of lines and conics, as well as the subschemes of the Hilbert schemes of twisted cubic curves passing through a general point.

The next theorem is the main result of the paper.

Theorem 1.2. *Let X be a Fano threefold from Table 1; in particular we assume $\rho(X) = 1$.*

- (i) *X is unirational if and only if $X(\mathbf{k}) \neq \emptyset$.*
- (ii) *If X has type $X_{(4,4)}$, $X_{(2,2,2)}$, $X_{(2,2)}$, or $X_{(1,1,1)}$ then X is \mathbf{k} -rational if and only if $X(\mathbf{k}) \neq \emptyset$.*
- (iii) *If X has type $X_{(3,3)}$ then X is never \mathbf{k} -rational.*

Note that over an algebraically closed field threefolds of types $X_{(4,4)}$, $X_{(2,2,2)}$, $X_{(2,2)}$, and $X_{(1,1,1)}$ have $h^{1,2} = 0$, hence trivial intermediate Jacobians, while the intermediate Jacobians of threefolds of types $X_{(3,3)}$ and $X_{(1,1,1,1)}$ are Jacobians of curves of genus 3 and 1, respectively; this explains the difference in the behavior.

It is a classical fact that the existence of a \mathbf{k} -point is necessary for rationality or unirationality, so the major part of the proof of the theorem consists of proving rationality or unirationality under this assumption. We use for this a case-by-case analysis (see §1.2 for a description of our approach). The theorem is thus a combination of the following results:

- rationality for threefolds of type $X_{(1,1,1)}$ is proved in Corollary 3.4;
- rationality for threefolds of type $X_{(2,2)}$ is proved in Proposition 4.1,
- rationality for threefolds of type $X_{(2,2,2)}$ is proved in Proposition 4.3,
- rationality for threefolds of type $X_{(4,4)}$ is proved in Proposition 5.5;
- unirationality for threefolds of type $X_{(1,1,1,1)}$ is proved in Proposition 4.5;
- unirationality for threefolds of type $X_{(3,3)}$ is proved in Proposition 6.9;
- non-rationality for threefolds of type $X_{(3,3)}$ is proved in Corollary 6.11.

Theorem 1.2 provides nice criteria of rationality for the five out of six types of Fano threefolds listed in Table 1. For the remaining type $X_{(1,1,1,1)}$ we have a conjecture and a partial result.

Conjecture 1.3. *If X has type $X_{(1,1,1,1)}$ and $\rho(X) = 1$ then X is never \mathbf{k} -rational.*

To explain the partial result we need to introduce some notation. Let X be a Fano threefold of type $X_{(1,1,1,1)}$. As we show in Lemma 2.5, the action of the Galois group $G(\bar{\mathbf{k}}/\mathbf{k})$ on $\text{Pic}(X_{\bar{k}})$ factors through the group \mathfrak{S}_4 that acts by permutations of the pullbacks of the point classes of the factors of the ambient $(\mathbb{P}_{\bar{\mathbf{k}}}^1)^4$, and the assumption $\rho(X) = 1$ means that the subgroup

$$G_X := \text{Im}(G(\bar{\mathbf{k}}/\mathbf{k}) \longrightarrow \mathfrak{S}_4) \subset \mathfrak{S}_4$$

is *transitive*, hence belongs to the following list of (conjugacy classes) of transitive subgroups of \mathfrak{S}_4 :

$$G_X \in \{\mathfrak{S}_4, \mathfrak{A}_4, D_4, V_4, C_4\},$$

where \mathfrak{A}_4 is the alternating subgroup, D_4 is the dihedral group of order 8 (a Sylow 2-subgroup in \mathfrak{S}_4), V_4 is the Klein group of order 4, and C_4 is the cyclic group of order 4. Note that all of these groups contain V_4 except for C_4 .

Theorem 1.4. *Assume the base field k is uncountable. Then a very general X of type $X_{(1,1,1,1)}$ with $\rho(X) = 1$ and $X(k) \neq \emptyset$ such that the group G_X contains V_4 is not stably k -rational.*

1.2. The proofs. For (uni)rationality constructions it is natural to use k -Sarkisov links:

$$(1.2.1) \quad \begin{array}{ccccc} & & \tilde{X} & \overset{\psi}{\dashrightarrow} & \tilde{X}^+ \\ & \swarrow \sigma & & \searrow \phi & \\ X & & & & X^+ \\ & \searrow \phi & \tilde{X} & \swarrow \phi_+ & \\ & & & & \end{array}$$

where σ is the blowup of a k -irreducible subvariety, ϕ and ϕ_+ are small crepant birational contractions, ψ is a flop, and σ_+ is a Mori extremal contraction. Note that such a link is completely determined by the center of the blowup σ — the contractions and the flop are obtained by the Minimal Model Program applied to \tilde{X} (note that $\rho(\tilde{X}) = 2$, so the output of the MMP is unambiguous); in particular the link is defined over k . For our purpose it is enough to consider two types of Sarkisov links:

- Sarkisov links where σ is the blowup of a k -point;
- Sarkisov links where σ is the blowup of a reduced k -irreducible singular conic.

We construct the corresponding links accurately for threefolds of type $X_{(4,4)}$ in §5 (see Theorem 5.1) by using standard MMP arguments. Of course, a similar construction could be given for other types of Fano threefolds from Table 1, but to make the argument less tedious we use the fact that all other among these threefolds are k -forms of complete intersections in products of projective spaces and deduce the required (uni)rationality constructions from an appropriate birational transformation for a product of projective spaces.

With this goal in mind we construct in §3 a toric birational transformation between the product $(\mathbb{P}^n)^r$ of projective spaces and a \mathbb{P}^r -bundle over the product $(\mathbb{P}^{n-1})^r$ of smaller projective spaces, see Theorem 3.1 (in fact, we construct a birational transformation in a slightly more general situation, but the setup described above is the only one that we need for applications of the paper). This theorem has a consequence of independent interest, Corollary 3.3, saying that a k -form of a product of projective spaces is k -rational if and only if it has a k -point. This corollary immediately gives the required rationality construction for Fano threefolds of type $X_{(1,1,1)}$ (Corollary 3.4), and with a bit of more work provides rationality constructions for threefolds of types $X_{(2,2)}$ (Proposition 4.1) and $X_{(2,2,2)}$ (Proposition 4.3) as well as a unirationality construction for threefolds of type $X_{(1,1,1,1)}$ (Proposition 4.5).

In the case of a variety X of type $X_{(3,3)}$ with a k -point x we again use the toric transformation of Theorem 3.1 to construct a birational equivalence of X with a divisor X^+ of bidegree $(2, 2)$ in a k -form of $\mathbb{P}^2 \times \mathbb{P}^2$. If x lies on a line in X , we check that X^+ contains a k -form of the quadric surface $\mathbb{P}^1 \times \mathbb{P}^1$ and use this to deduce unirationality of X (Proposition 6.9). If x does not lie

on a line, we check in Proposition 6.6 that X^+ described above is, in fact, the mid-point of a Sarkisov link, that ends with a conic bundle over \mathbb{P}^2 which has a smooth quartic curve $\Gamma \subset \mathbb{P}^2$ as discriminant. We also check that the discriminant double covering $\tilde{\Gamma} \rightarrow \Gamma$ obtained from this conic bundle is trivial over a quadratic extension k' of the base field k but nontrivial over k and that the conic bundle has a rational section over k' . We check in Theorem 6.10 that these geometric properties characterize the non-rational conic bundles constructed by Benoist and Wittenberg in [BW20] and deduce in Corollary 6.11 non-rationality of X from [BW20, Proposition 3.4].

In the last part of the paper, §7, we discuss Fano threefolds of type $X_{(1,1,1,1)}$. To prove Theorem 1.4 we use degeneration technique. Namely we observe that one can degenerate these varieties to a singular toric threefold (with ordinary double points) which is well-known not to be stably rational. Since stable rationality is specialization-closed by a result of Nicaise and Shinder [NS19], we conclude that a very general threefold of type $X_{(1,1,1,1)}$ is not stably rational.

Acknowledgements. We would like to thank Sergey Gorchinskiy and Costya Shramov for useful discussions.

2. EXTREMAL CONTRACTIONS AND HILBERT SCHEMES OF CURVES

In this section we describe the geometry of Fano threefolds of index 1 from Table 1. In particular, we describe their extremal contractions over \bar{k} as well as their Hilbert schemes of lines, conics, and twisted cubic curves.

To start with, recall that for most Fano threefolds the anticanonical linear system is very ample and the anticanonical image is an intersection of quadrics; in fact Fano threefolds which do not enjoy these nice properties (hyperelliptic and trigonal ones) have been classified and listed in [Isk80]. It is easy to check that Fano threefolds from Table 1 are not in this list; therefore we obtain

Theorem 2.1 ([Isk80, Chapter 2, Theorems 2.2 and 3.4]). *Let X be a Fano threefold from Table 1. The anticanonical class $-K_X$ is very ample and the anticanonical image*

$$X = X_{2g-2} \subset \mathbb{P}^{g+1}$$

is an intersection of quadrics (as a scheme), where $g = g(X)$.

2.1. Contractions over \bar{k} . Assume X is a Fano threefold of index 1 from Table 1, i.e., a threefold of either of types $X_{(2,2,2)}$, $X_{(4,4)}$, $X_{(3,3)}$, $X_{(1,1,1,1)}$. Then there is an embedding

$$(2.1.1) \quad X_{\bar{k}} \subset Y \cong (\mathbb{P}^n)^r,$$

(we will see in Lemma 2.5 that $r = \rho(X_{\bar{k}})$, hence the notation), where

$$(n, r) = (2, 3), (4, 2), (3, 2), \text{ or } (1, 4).$$

Indeed, for types $X_{(2,2,2)}$, $X_{(3,3)}$, $X_{(1,1,1,1)}$ this holds by definition and for type $X_{(4,4)}$ this follows from the following

Lemma 2.2. *Let $\Gamma_1 \subset Q_1 \subset \mathbb{P}^4$ be a linearly normal smooth rational quartic curve in a smooth quadric threefold. If H_1 is the hyperplane class of Q_1 then the linear system $|2H_1 - \Gamma_1|$ of quadrics through Γ_1 defines a birational morphism $\pi_2: \text{Bl}_{\Gamma_1} Q_1 \rightarrow Q_2 \subset \mathbb{P}^4$ onto another smooth quadric*

threefold Q_2 and this morphism is itself a blowup of a linearly normal smooth rational quartic curve $\Gamma_2 \subset Q_2$, so that

$$\mathrm{Bl}_{\Gamma_1}(Q_1) \cong \mathrm{Bl}_{\Gamma_2}(Q_2).$$

In particular, if X is a Fano threefold of type $X_{(4,4)}$ there is a natural embedding

$$X_{\bar{k}} \hookrightarrow Q_1 \times Q_2 \subset \mathbb{P}_{\bar{k}}^4 \times \mathbb{P}_{\bar{k}}^4$$

such that $-K_{X_{\bar{k}}}$ is the sum of the pullbacks of the hyperplane classes of the factors.

Proof. The curve Γ_1 is an intersection of five quadrics. Hence, if E_1 is the exceptional divisor of the blowup $\pi_1: X_{\bar{k}} \rightarrow Q_1$ and H_1 is the pullback of the hyperplane class of Q_1 , the linear system $|2H_1 - E_1|$ is 4-dimensional and base point free. Therefore, this linear system defines a morphism $\pi_2: \mathrm{Bl}_{\Gamma_1}(Q_1) \rightarrow \mathbb{P}^4$; moreover, $(2H_1 - E_1)^3 = 2$, hence the image of π_2 is a quadric $Q_2 \subset \mathbb{P}^4$ and π_2 is birational. Since $-K_{\mathrm{Bl}_{\Gamma_1}(Q_1)}$ is ample on the fibers of π_2 and $\rho(\mathrm{Bl}_{\Gamma_1}(Q_1)) = 2$, we see that π_2 is an extremal Mori contraction. By [Mor82] the quadric Q_2 is smooth and π_2 is the blowup of a curve which must be a linearly normal smooth rational quartic curve. For the last statement just note that $H_1 + (2H_1 - E_1) = 3H_1 - E_1$ is the anticanonical class of $\mathrm{Bl}_{\Gamma_1}(Q_1)$. \square

We denote by H_i , $1 \leq i \leq r$, the pullbacks to $Y = (\mathbb{P}^n)^r$ of the hyperplane classes of the factors and, abusing the notation, also their restrictions to $X_{\bar{k}}$ via the embedding (2.1.1).

Lemma 2.3. *The Picard group $\mathrm{Pic}(X_{\bar{k}})$ is freely generated by the classes H_i :*

$$\mathrm{Pic}(X_{\bar{k}}) = \bigoplus_{i=1}^r \mathbb{Z}H_i.$$

Moreover,

$$(2.1.2) \quad -K_{X_{\bar{k}}} = H := H_1 + \cdots + H_r;$$

Proof. For type $X_{(4,4)}$ this follows from Lemma 2.2, and for other types the first statement follows from the Lefschetz Hyperplane Theorem and the second from adjunction. \square

For each subset $I \subset \{1, \dots, r\}$ we consider the projection

$$(2.1.3) \quad \pi_I: X_{\bar{k}} \hookrightarrow Y \longrightarrow \prod_{i \in I} \mathbb{P}^n \cong (\mathbb{P}^n)^{|I|}.$$

Especially useful are the morphisms π_I for I of cardinality $r - 1$, so we introduce the notation

$$\widehat{i} := \{1, \dots, r\} \setminus \{i\}$$

and write

$$(2.1.4) \quad \pi_{\widehat{i}}: X_{\bar{k}} \rightarrow (\mathbb{P}^n)^{r-1}$$

for the corresponding morphisms. Note that in the case $r = 2$ we have $\widehat{i} = \{3 - i\}$, so these morphisms are the same as morphisms π_{3-i} . The next lemma describes $X_{\bar{k}}$ in terms of the $\pi_{\widehat{i}}$.

Lemma 2.4. *The morphism $\pi_{\widehat{i}}$ is birational onto its image and the exceptional divisor $E_{\widehat{i}}$ is irreducible. More precisely, the morphism $\pi_{\widehat{i}}$ identifies $X_{\bar{k}}$ as follows:*

- (i) if X has type $X_{(2,2,2)}$ the map $\pi_{\widehat{i}}$ is the blowup of a smooth divisor $W_{\widehat{i}} \subset \mathbb{P}^2 \times \mathbb{P}^2$ of bidegree $(1, 1)$ along a smooth rational curve $\Gamma_{\widehat{i}} \subset W_{\widehat{i}}$ of bidegree $(2, 2)$ whose projections to the factors \mathbb{P}^2 are closed embeddings; the divisor class H_i is equal to $\sum_{j \neq i} H_j - E_{\widehat{i}}$;

- (ii) if X has type $X_{(4,4)}$ the map π_i is the blowup of a 3-dimensional quadric Q_i along a smooth rational curve $\Gamma_i \subset Q_i$ of degree 4; the divisor class $H_{\hat{\tau}}$ is equal to $2H_i - E_i$;
- (iii) if X has type $X_{(3,3)}$ the map π_i is the blowup of \mathbb{P}^3 along a smooth curve $\Gamma_i \subset \mathbb{P}^3$ of genus 3 and degree 6; the divisor class $H_{\hat{\tau}}$ is equal to $3H_i - E_i$;
- (iv) if X has type $X_{(1,1,1,1)}$ the map $\pi_{\hat{\tau}}$ is the blowup of $\mathbb{P}^1 \times \mathbb{P}^1 \times \mathbb{P}^1$ along a smooth elliptic curve $\Gamma_{\hat{\tau}} \subset (\mathbb{P}^1)^3$ of multidegree $(2, 2, 2)$; the divisor class H_i is equal to $\sum_{j \neq i} H_j - E_{\hat{\tau}}$.

Proof. Part (ii) is proved in Lemma 2.2. So, assume X is a variety of either of types $X_{(2,2,2)}$, $X_{(3,3)}$, or $X_{(1,1,1,1)}$. Birationality of the projection $\pi_{\hat{\tau}}$ is clear from the descriptions of Table 1; it also follows that all fibers of $\pi_{\hat{\tau}}$ are linear subspaces in \mathbb{P}^n and $-K_{X_{\bar{k}}}$ restricts to each of them as the hyperplane class by (2.1.2). Also, it is easy to see that the image of $\pi_{\hat{\tau}}$ is smooth in all cases (for type $X_{(2,2,2)}$ if $W_{\hat{\tau}} \subset \mathbb{P}^2 \times \mathbb{P}^2$ is singular then its preimage in $\mathbb{P}^2 \times \mathbb{P}^2 \times \mathbb{P}^2$ is singular along a plane, hence $X_{\bar{k}}$, which is the intersection of this preimage with two other divisors, must be singular; and for types $X_{(3,3)}$ and $X_{(1,1,1,1)}$ the image is just \mathbb{P}^3 or $\mathbb{P}^1 \times \mathbb{P}^1 \times \mathbb{P}^1$, respectively).

By Lemma 2.3 the relative Picard number of $\pi_{\hat{\tau}}$ is 1 and $-K_{X_{\bar{k}}}$ is ample, hence $\pi_{\hat{\tau}}$ is an extremal Mori contraction. Since both the source and target of $\pi_{\hat{\tau}}$ are smooth, it follows from [Mor82] that the morphism $\pi_{\hat{\tau}}$ is either the blowup of a smooth curve or the blowup of a smooth point. In the latter case the restriction of $-K_{X_{\bar{k}}}$ to the nontrivial fiber \mathbb{P}^2 of $\pi_{\hat{\tau}}$ would be isomorphic to $\mathcal{O}_{\mathbb{P}^2}(2)$, contradicting to the above observation, hence $\pi_{\hat{\tau}}$ is the blowup of a smooth curve.

The remaining assertions are easy and left to the reader (see also [MM82]). \square

Lemma 2.5. *The classes H_i are semiample and generate the nef cone of $X_{\bar{k}}$. The Galois group $G(\bar{k}/k)$ permutes these classes in a transitive way. In other words, the natural homomorphism $G(\bar{k}/k) \rightarrow \text{Aut}(\text{Pic}(X_{\bar{k}}))$ factors through the subgroup $\mathfrak{S}_r \subset \text{Aut}(\text{Pic}(X_{\bar{k}}))$ and its image*

$$(2.1.5) \quad G_X := \text{Im}(G(\bar{k}/k) \longrightarrow \mathfrak{S}_r)$$

is a transitive subgroup of \mathfrak{S}_r .

Proof. The classes H_i are pullbacks of ample classes on \mathbb{P}^n , hence semiample, and they generate $\text{Pic}(X_{\bar{k}})$ by Lemma 2.3. If Λ_i is the classes of a non-trivial fiber of $\pi_{\hat{\tau}}$, we have

$$H_j \cdot \Lambda_i = \delta_{ij},$$

therefore H_j generate the rays of the nef cone. It follows that the Galois group permutes the H_i , hence its action on $\text{Pic}(X_{\bar{k}})$ factors through the permutation group. Transitivity of the subgroup $G_X \subset \mathfrak{S}_r$ follows from the equality $\rho(X) = 1$. \square

2.2. Lines. By a line on X we understand a curve (defined over \bar{k}) of anticanonical degree 1. We denote by $F_1(X)$ the Hilbert scheme of lines on X . Note that $F_1(X)_{\bar{k}} \cong F_1(X_{\bar{k}})$.

Lemma 2.6. *Let X be a Fano threefold of types $X_{(2,2,2)}$, $X_{(4,4)}$, $X_{(3,3)}$, or $X_{(1,1,1,1)}$. A line on X is a fiber of the exceptional divisor of one of the projections (2.1.4). In particular*

$$F_1(X_{\bar{k}}) \cong \bigsqcup_{i=1}^r \Gamma_{\hat{\tau}},$$

where the smooth curves $\Gamma_{\hat{\tau}}$ have been described in Lemma 2.4. The normal bundle of each line is

$$(2.2.1) \quad \mathcal{N}_{L/X_{\bar{k}}} \cong \mathcal{O}_L \oplus \mathcal{O}_L(-1).$$

Finally, the action of the Galois group $G(\bar{k}/k)$ on the set of connected components of the Hilbert scheme of lines factors through the group G_X and is transitive.

Proof. Since the classes H_i are semiample, it follows from (2.1.2) that for each \bar{k} -line L on X there is a unique i such that $L \cdot H_i = 1$ and $L \cdot H_j = 0$ for $j \neq i$ (i.e., $[L] = \Lambda_i$ in the notation of Lemma 2.5). Thus, L is contracted by the projection $\pi_{\hat{\tau}}$, hence it is equal to a fiber of the exceptional divisor of this projection. Taking into account the description of the projections $\pi_{\hat{\tau}}$, we obtain the description of $F_1(X_{\bar{k}})$.

Further, the description of the normal bundle of L follows from the exact sequence

$$0 \longrightarrow \mathcal{N}_{L/E_{\hat{\tau}}} \longrightarrow \mathcal{N}_{L/X_{\bar{k}}} \longrightarrow \mathcal{N}_{E_{\hat{\tau}}/X_{\bar{k}}}|_L \longrightarrow 0$$

because the first term is trivial and the last is $\mathcal{O}_L(-1)$. Finally, factorization of the Galois action on the components of $F_1(X_{\bar{k}})$ and its transitivity follow from Lemma 2.5. \square

For a \bar{k} -point $x \in X$ we denote by $F_1(X, x) \subset F_1(X)$ the subscheme parameterizing lines passing through x . We will need the following observation.

Lemma 2.7. *Let X be a Fano threefold of types $X_{(2,2,2)}$, $X_{(4,4)}$, $X_{(3,3)}$, or $X_{(1,1,1,1)}$. If $x \in X(\bar{k})$, the scheme $F_1(X_{\bar{k}}, x)$ is a finite reduced scheme of length at most $r = \rho(X_{\bar{k}})$. If, moreover, $x \in X(k)$ then either $F_1(X_{\bar{k}}, x) = \emptyset$, or $F_1(X_{\bar{k}}, x)$ is a reduced scheme of length r and the Galois group $G(\bar{k}/k)$ action on $F_1(X_{\bar{k}}, x)$ factors through the group G_X and is transitive.*

Proof. By Lemma 2.6 for each \bar{k} -point x of X there is at most one line from each of the connected components of the Hilbert scheme $F_1(X_{\bar{k}})$ passing through x . This proves that $F_1(X_{\bar{k}}, x)$ is finite and reduced and gives the bound for its length.

Now assume x is a point defined over k and let L be a \bar{k} -line through x . Then for any $g \in G(\bar{k}/k)$ the line $g(L)$ also passes through x . Transitivity of the Galois action on the set of components of $F_1(X_{\bar{k}})$ then implies that there is a unique line of each type through x , hence the length of $F_1(X_{\bar{k}}, x)$ is r , and the $G(\bar{k}/k)$ -action on $F_1(X_{\bar{k}}, x)$ factors through G_X and is transitive. \square

2.3. Conics. By a conic on X we understand a curve (defined over \bar{k}) of anticanonical degree 2. We denote by $F_2(X)$ the Hilbert scheme of conics on X . Note that $F_2(X)_{\bar{k}} \cong F_2(X_{\bar{k}})$.

Lemma 2.8. *Let X be a Fano threefold of types $X_{(2,2,2)}$, $X_{(4,4)}$, $X_{(3,3)}$, or $X_{(1,1,1,1)}$. We have the following descriptions of the Hilbert schemes of conics $F_2(X_{\bar{k}})$:*

$$\begin{aligned} F_2((X_{(2,2,2)})_{\bar{k}}) &\cong \mathbb{P}_{\bar{k}}^2 \sqcup \mathbb{P}_{\bar{k}}^2 \sqcup \mathbb{P}_{\bar{k}}^2, \\ F_2((X_{(4,4)})_{\bar{k}}) &\cong \Gamma_1 \times \Gamma_2, \\ F_2((X_{(3,3)})_{\bar{k}}) &\cong \text{Sym}^2 \Gamma_1 \cong \text{Sym}^2 \Gamma_2, \\ F_2((X_{(1,1,1,1)})_{\bar{k}}) &\cong \bigsqcup_6 (\mathbb{P}_{\bar{k}}^1 \times \mathbb{P}_{\bar{k}}^1), \end{aligned}$$

where Γ_i are the curves described in Lemma 2.4.

Moreover, the morphism from each component of the universal conic to $X_{\bar{k}}$ is dominant.

Proof. First, note that no conic on X is contracted by the projections $\pi_{\hat{\tau}}$, since by Lemma 2.4 any reduced connected curve contracted by $\pi_{\hat{\tau}}$ is a line and lines do not support nonreduced conics

by (2.2.1) and [KPS18, Remark 2.1.7]. Therefore, again from (2.1.2) we deduce that for each \bar{k} -conic $C \subset X_{\bar{k}}$ there is a pair of indices $1 \leq i_1 < i_2 \leq r$ such that

$$(2.3.1) \quad H_{i_1} \cdot C = H_{i_2} \cdot C = 1 \quad \text{and} \quad H_j \cdot C = 0 \quad \text{for } j \notin \{i_1, i_2\}.$$

If $r \geq 3$, i.e., if X is of type $X_{(2,2,2)}$ or $X_{(1,1,1,1)}$, such C is contracted by the projection

$$(2.3.2) \quad \pi_i: X_{\bar{k}} \longrightarrow \mathbb{P}_{\bar{k}}^2 \quad \text{or} \quad \pi_{i_1, i_2}: X_{\bar{k}} \longrightarrow \mathbb{P}_{\bar{k}}^1 \times \mathbb{P}_{\bar{k}}^1.$$

respectively. It is easy to see that the maps (2.3.2) are flat conic bundles, hence C is a fiber of one of them, and therefore $F_2(X_{\bar{k}})$ is the disjoint union of $\mathbb{P}_{\bar{k}}^2$, or of $\mathbb{P}_{\bar{k}}^1 \times \mathbb{P}_{\bar{k}}^1$, respectively.

Assume X is of type $X_{(4,4)}$. Applying Corollary A.2 twice we obtain a morphism

$$\varphi = (\varphi_1, \varphi_2): F_2(X_{\bar{k}}) \longrightarrow \Gamma_1 \times \Gamma_2$$

that takes a conic $C \subset X_{\bar{k}}$ to the pair of lines (L_2, L_1) of different types such that $C \cap L_i \neq \emptyset$. We will show that φ is an isomorphism.

First, note that by (2.3.1), Lemma 2.4(ii), and Corollary A.2 for $[C] \in \varphi_1^{-1}(x_1)$ with $x_1 \in \Gamma_1$ the image of C with respect to the blowup $\pi_1: X_{\bar{k}} \rightarrow Q_1$ is a line intersecting the curve $\Gamma_1 \subset Q_1$ at the point x_1 . Since any line on Q_1 through x_1 lies in the embedded tangent space to Q_1 at x_1 , and the intersection of this tangent space with Q_1 is a 2-dimensional quadratic cone with vertex at x_1 , it follows that

$$\varphi_1^{-1}(x_1) \cong \mathbb{P}^1$$

for any $x_1 \in \Gamma_1$. Since also $\Gamma_2 \cong \mathbb{P}^1$, the morphism φ is a morphism of \mathbb{P}^1 -bundles over Γ_1 , and to show that it is an isomorphism, it is enough to check that it is generically injective.

So, consider a general pair (L_2, L_1) of lines on X of different types. It follows from (2.3.1) and Lemma 2.4(ii) that $\bar{L}_1 := \pi_1(L_1)$ is a line on Q_1 bisecant to Γ_1 , $x_1 := \pi_1(L_2)$ is a point on Γ_1 , and $\varphi^{-1}(L_2, L_1)$ is the Hilbert scheme of lines $L \subset Q_1$ passing through x_1 and intersecting \bar{L}_1 . By genericity we may assume $x_1 \notin \bar{L}_1$ (i.e., that the lines L_1 and L_2 do not intersect). Then any line L as above is contained in the intersection of the plane spanned by \bar{L}_1 and x_1 with Q_1 , which is equal to the union of the line \bar{L}_1 with a residual line. Therefore, L must be equal to the residual line, hence the scheme $\varphi^{-1}(L_2, L_1)$ consists of a single point, and φ is an isomorphism.

Since the embedded tangent space to Q_1 at a general point $x \in Q_1$ intersects the quartic curve Γ_1 at 4 points, the universal conic is dominant over $X_{\bar{k}}$ and its degree over $X_{\bar{k}}$ is 4.

Finally, assume X is of type $X_{(3,3)}$. By (2.3.1) and Lemma 2.4(iii) the image of C with respect to the blowup $\pi_i: X_{\bar{k}} \rightarrow \mathbb{P}^3$ is a line intersecting the curve $\Gamma_i \subset \mathbb{P}^3$ at two points. This defines a morphism

$$F_2(X_{\bar{k}}) \longrightarrow \text{Sym}^2 \Gamma_i,$$

and it is easy to see that it is an isomorphism. It is also easy to see that for a general point $x \in \mathbb{P}^3$ there are three lines passing through x and bisecant to Γ_1 ; therefore the universal conic on $X_{\bar{k}}$ is dominant of degree 3 over $X_{\bar{k}}$. \square

Remark 2.9. Let X be a threefold of type $X_{(4,4)}$. Clearly a general line on the quadric Q_1 passing through a point $x \in \Gamma_1$ is not bisecant to Γ_1 and its strict transform in X intersects the line $L_2 = \pi_1^{-1}(x)$ transversally. This means that a general conic intersecting L_2 is smooth and intersects L_2 transversally.

For a given curve $\Theta \subset X$ we denote by $F_2(X, \Theta)$ the subscheme of the Hilbert scheme $F_2(X)$ that parameterizes conics intersecting the curve Θ . Furthermore, we denote by $R_\Theta \subset X$ the image of the universal conic intersecting Θ .

Lemma 2.10. *If X is of type $X_{(4,4)}$ and Θ is a singular conic then $F_2(X_{\bar{k}}, \Theta) \cong \Gamma_1 \cup \Gamma_2$ is the union of the two rulings of the surface $F_2(X_{\bar{k}}) \cong \Gamma_1 \times \Gamma_2$ passing through the point $[\Theta]$. Moreover, the surface $R_\Theta \subset X$ is an anticanonical divisor passing through each component of the curve Θ with multiplicity 3.*

Proof. Let L_1 and L_2 be the irreducible components (over \bar{k}) of the conic Θ . Then, of course, L_i are lines of two different types and

$$F_2(X_{\bar{k}}, \Theta) = F_2(X_{\bar{k}}, L_1) \cup F_2(X_{\bar{k}}, L_2).$$

Recall that by Lemma 2.6 the curves Γ_1 and Γ_2 can be identified with the two connected components of $F_1(X_{\bar{k}})$ and the isomorphism $F_2(X_{\bar{k}}) \cong \Gamma_1 \times \Gamma_2$ of Lemma 2.8 is defined by taking a conic C to the unique pair of lines of different types intersecting C . This means that

$$F_2(X_{\bar{k}}, \Theta) \cong (\Gamma_1 \times [L_1]) \cup ([L_2] \times \Gamma_2) \subset \Gamma_1 \times \Gamma_2.$$

Furthermore, it follows from the description of Lemma 2.8 that conics parameterized by the curve $F_2(X_{\bar{k}}, L_2)$ sweep on Q_1 the hyperplane section tangent to Q_1 at the point $\pi_1(L_2)$; it contains the line $\pi_1(L_1)$ with multiplicity 1 and has multiplicity 2 at the point $\pi_1(L_2)$. Similarly, conics parameterized by the curve $F_2(X_{\bar{k}}, L_1)$ sweep on Q_2 the hyperplane section containing the line $\pi_2(L_2)$ with multiplicity 1 and having multiplicity 2 at the point $\pi_2(L_1)$. The divisor they sweep together has the class $(H_1 - L_1 - 2L_2) + (H_2 - L_2 - 2L_1) = H - 3\Theta$. \square

2.4. Twisted cubic curves. Finally, we describe the Hilbert scheme $F_3(X, x)$ of rational normal cubic curves passing through a point x . We denote by $R_x \subset X$ the image of the universal rational normal cubic curve through x . Recall the curves $\Gamma_{\hat{i}}$ described in Lemma 2.4.

Lemma 2.11. *Let X be a Fano threefold of types $X_{(2,2,2)}$, $X_{(4,4)}$, $X_{(3,3)}$, or $X_{(1,1,1,1)}$. If x is a k -point on X not lying on a \bar{k} -line, one has the following descriptions of the schemes $F_3(X_{\bar{k}}, x)$:*

$$\begin{aligned} F_3((X_{(2,2,2)})_{\bar{k}}, x) &\cong \Gamma_{1,2} \cong \Gamma_{1,3} \cong \Gamma_{2,3}, \\ F_3((X_{(4,4)})_{\bar{k}}, x) &\cong \mathbb{P}_{\bar{k}}^1 \sqcup \mathbb{P}_{\bar{k}}^1, \\ F_3((X_{(3,3)})_{\bar{k}}, x) &\cong \Gamma_1 \sqcup \Gamma_2, \\ F_3((X_{(1,1,1,1)})_{\bar{k}}, x) &\cong \bigsqcup_8 \mathbb{P}_{\bar{k}}^1. \end{aligned}$$

Moreover, for threefolds of type $X_{(4,4)}$ the surface R_x is an anticanonical divisor passing through the point x with multiplicity 4.

Proof. First, consider a threefold X of type $X_{(2,2,2)}$. If C is a rational normal cubic curve and $H_i \cdot C = 0$ for some i then C is contracted by one of the conic bundles (2.3.2), hence C is supported on a fiber of (2.3.2). But the conormal bundle of any such fiber is trivial, hence it cannot support a nonreduced rational curve of degree more than 2. This means that we have $H_i \cdot C = 1$ for each i , and we conclude from this and Lemma 2.4 that the image of C under the map $\pi_{1,2}: X_{\bar{k}} \rightarrow W_{1,2}$ is a rational curve of bidegree $(1, 1)$ intersecting the curve $\Gamma_{1,2}$

and passing through x . The argument analogous to that of Corollary A.2 shows that there is a morphism

$$\varphi_{1,2}: F_3(X, x) \longrightarrow \Gamma_{1,2}$$

that takes a twisted cubic curve C to the unique point $x_{1,2} \in \Gamma_{1,2}$ such that $C \cap \pi_{1,2}^{-1}(x_{1,2}) \neq \emptyset$. This morphism is an isomorphism, because on $W_{1,2}$ there is a unique curve of bidegree $(1, 1)$ through a given pair of points (unless they lie on a fiber of either of the projections $W_{1,2} \rightarrow \mathbb{P}_{\bar{k}}^2$, in which case x lies on a line in X). The same argument proves isomorphisms of $F_3(X, x)$ with the curves $\Gamma_{1,3}$ and $\Gamma_{2,3}$.

Next, consider a threefold of type $X_{(4,4)}$. If $H_i \cdot C = 0$ for some i then C is contracted by π_i , hence is supported on a line. But the conormal bundle of a line is globally generated by (2.2.1), hence a line cannot support a nonreduced rational curve of degree more than 1. This means that C has bidegree $(1, 2)$ or $(2, 1)$. In the first case the image of C under π_1 is a line on the quadric Q_1 passing through x ; hence the corresponding component of $F_3(X, x)_{\bar{k}}$ is isomorphic to $\mathbb{P}_{\bar{k}}^1$. These lines sweep on Q_1 the hyperplane section tangent at x , i.e., a divisor of class H_1 passing through x with multiplicity 2. The second component is described analogously. The total divisor class of the surface R_x is $H_1 + H_2 - 4x$, i.e., the anticanonical class passing through x with multiplicity 4.

Next, consider a threefold of type $X_{(3,3)}$. The same argument as above shows that C has bidegree $(1, 2)$ or $(2, 1)$. In the first case the image of C under π_1 is a line on \mathbb{P}^3 passing through x and intersecting the curve Γ_1 . Since for any point of Γ_1 there is a unique line through it and x , the corresponding component of $F_3(X_{\bar{k}}, x)$ is isomorphic to Γ_1 . Analogously, the second component is isomorphic to Γ_2 .

Finally, consider a threefold of type $X_{(1,1,1,1)}$. Then, of course, $H_i \cdot C = 0$ for some i . The argument used for threefolds of type $X_{(2,2,2)}$ shows this cannot hold for two distinct i . So, assume this holds for $i = 1$. By Lemma 2.4(iv) the image of C under the map $\pi_{1,2,3}$ is a curve of multidegree $(0, 1, 1)$ on $\mathbb{P}_{\bar{k}}^1 \times \mathbb{P}_{\bar{k}}^1 \times \mathbb{P}_{\bar{k}}^1$ intersecting the curve $\Gamma_{1,2,3}$ and passing through x . In other words, it is a curve of bidegree $(1, 1)$ on the surface $\mathbb{P}_{\bar{k}}^1 \times \mathbb{P}_{\bar{k}}^1$ passing through x and either of the two points of intersection of $\Gamma_{1,2,3}$ with this surface (note that these points cannot collide because otherwise x would lie on a line in X). Therefore, there are two pencils of such curves. Using the same argument for other i we see that altogether there are 8 pencils of twisted cubic curves on X passing through x . \square

Remark 2.12. Let X be a threefold of type $X_{(4,4)}$. Since a general line on Q_1 passing through a point $x \notin \Gamma_1$ does not intersect Γ_1 , it follows that a general twisted cubic curve on X passing through x is smooth.

3. A BIRATIONAL TRANSFORMATION FOR A PRODUCT OF PROJECTIVE SPACES

In this section we construct a birational transformation for a product of projective spaces and deduce a consequence for rationality of its k -forms.

3.1. Product of projective spaces. Consider the product

$$Y = \mathbb{P}^{n_1} \times \mathbb{P}^{n_2} \times \cdots \times \mathbb{P}^{n_r} = \mathbb{P}(V_1) \times \mathbb{P}(V_2) \times \cdots \times \mathbb{P}(V_r)$$

of projective spaces. Assume that $r = p + q$ and

$$(3.1.1) \quad n_1 \geq n_2 \geq \cdots \geq n_p \geq 2, \quad n_{p+1} = \cdots = n_{p+q} = 1.$$

Let $y \in Y$ be a point, and let (v_1, v_2, \dots, v_r) , $0 \neq v_i \in V_i$, be the corresponding collection of vectors. Consider the blowup

$$\tilde{Y} = \text{Bl}_y(Y)$$

and let $E \subset \tilde{Y}$ be its exceptional divisor.

Let $\text{PGL}(V_i)_{v_i} \subset \text{PGL}(V_i)$ be the stabilizer of the point $[v_i] \in \mathbb{P}(V_i)$. The group

$$G = \prod_{i=1}^r \text{PGL}(V_i)_{v_i}$$

acts naturally on \tilde{Y} and has finitely many orbits, which can be described as follows. First, for each $1 \leq i \leq r$ let

$$(3.1.2) \quad \tilde{Y}_i := \text{Bl}_y \left(\mathbb{P}(V_1) \times \cdots \times \mathbb{P}(V_{i-1}) \times [v_i] \times \mathbb{P}(V_{i+1}) \times \cdots \times \mathbb{P}(V_r) \right) \subset \tilde{Y}.$$

Furthermore, for any subset $I \subsetneq \{1, \dots, r\}$ denote

$$(3.1.3) \quad \tilde{Y}_I := \bigcap_{i \in I} \tilde{Y}_i \quad \text{and} \quad E_I := E \cap \tilde{Y}_I.$$

Finally, set

$$(3.1.4) \quad \tilde{Y}_I^\circ := \tilde{Y}_I \setminus \left(E_I \cup \bigcup_{I \subsetneq J} \tilde{Y}_J \right) \quad \text{and} \quad E_I^\circ := E_I \setminus \left(\bigcup_{I \subsetneq J} E_J \right).$$

Then $\tilde{Y}_\emptyset^\circ$ is the open orbit, E_\emptyset° and \tilde{Y}_i° , $p+1 \leq i \leq q$, are the orbits of codimension 1, and all other orbits have higher codimension.

To describe the other side of the transformation, denote

$$\bar{V}_i := V_i / kv_i$$

and choose splittings $V_i = kv_i \oplus \bar{V}_i$. They induce a direct sum decomposition

$$V_1 \otimes \cdots \otimes V_r = \bigoplus_{I \subset \{1, \dots, r\}} \bar{V}_I, \quad \text{where} \quad \bar{V}_I := \bigotimes_{i \in I} \bar{V}_i.$$

Note that the point y corresponds to the summand $\bar{V}_\emptyset = \mathbf{k}$, and the tangent space to Y at y corresponds to the sum of the summands \bar{V}_I with $|I| = 1$.

Note also that for $i \geq p+1$ one has $\mathbb{P}(\bar{V}_i) \cong \text{Spec}(\mathbf{k})$. Let

$$Y^+ := \prod_{i=1}^r \mathbb{P}(\bar{V}_i) = \mathbb{P}(\bar{V}_1) \times \mathbb{P}(\bar{V}_2) \times \cdots \times \mathbb{P}(\bar{V}_p) \cong \mathbb{P}^{n_1-1} \times \mathbb{P}^{n_2-1} \times \cdots \times \mathbb{P}^{n_p-1}.$$

Denote by h_i the pullback to Y^+ of the hyperplane class of the i -th factor (note that $h_i = 0$ for $i \geq p+1$ because, as we noticed above, $\mathbb{P}(\bar{V}_i)$ is just a point) and for $I \subset \{1, \dots, r\}$ set

$$h_I := \sum_{i \in I} h_i.$$

Consider the vector bundle \mathcal{E} of rank $r+1$ on Y^+ defined by

$$(3.1.5) \quad \mathcal{E} := \bigoplus_{|I| \geq r-1} \mathcal{O}(-h_I).$$

Denote by

$$s_i: Y^+ \longrightarrow \mathbb{P}_{Y^+}(\mathcal{E})$$

the section of (3.1.5) corresponding to the summand with $I = \{1, \dots, i-1, i+1, \dots, r\}$. Set

$$(3.1.6) \quad \hat{Y}^+ := \mathbb{P}_{Y^+}(\mathcal{E}), \quad \tilde{Y}^+ := \text{Bl}_{s_{p+1}(Y^+) \sqcup \dots \sqcup s_{p+q}(Y^+)}(\hat{Y}^+).$$

and let $E_i \subset \tilde{Y}^+$, $p+1 \leq i \leq p+q$, be the exceptional divisors. The group G acts transitively on Y^+ , the vector bundle \mathcal{E} is G -equivariant, and its summands $\mathcal{O}(-h_I)$ with $|I| = r-1$ are G -invariant. Therefore, the action of G lifts naturally to \hat{Y}^+ and \tilde{Y}^+ . Moreover, the action of G on \tilde{Y}^+ still has a finite number of orbits, which can be described as follows.

For a subset $J \subsetneq \{1, \dots, r\}$ denote

$$(3.1.7) \quad \bar{\mathcal{E}}_J = \bigoplus_{J \subset I, |I|=r-1} \mathcal{O}(-h_I);$$

this is a subbundle in \mathcal{E} of corank $1 + |J|$. Let $\tilde{Y}_J^+ \subset \tilde{Y}^+$ denote the strict transform of $\mathbb{P}_{Y^+}(\bar{\mathcal{E}}_J)$. Then the G -orbits are

$$\begin{aligned} (\tilde{Y}^+)^{\circ} &= \tilde{Y}^+ \setminus \left(\tilde{Y}_{\emptyset}^+ \cup \bigcup_{i=p+1}^q E_i \right), & (\tilde{Y}_J^+)^{\circ} &= \tilde{Y}_J^+ \setminus \left(\bigcup_{i=p+1}^q E_i \right), \\ E_i^{\circ} &= E_i \setminus \tilde{Y}_{\emptyset}^+, & E_{i,J}^{\circ} &= (E_i \cap \tilde{Y}_J^+) \setminus \left(\bigcup_{J \subsetneq K} E_i \cap \tilde{Y}_K^+ \right), \end{aligned}$$

where in the last formula we assume $i \notin J$. Note that $(\tilde{Y}^+)^{\circ}$ is the open orbit, $(\tilde{Y}_{\emptyset}^+)^{\circ}$ and E_i° are the orbits of codimension 1, and all other orbits have higher codimension.

The linear projection out of $[v_i]$ defines a $\text{PGL}(V_i)_{v_i}$ -equivariant rational map $\mathbb{P}(V_i) \dashrightarrow \mathbb{P}(\bar{V}_i)$ which is regular if $i \geq p+1$. The product of these maps is a G -equivariant rational map, which we denote by $\psi_0: Y \dashrightarrow Y^+$. It gives rise to the following birational transformation.

Theorem 3.1. *There is a small birational G -equivariant isomorphism $\psi: \tilde{Y} \dashrightarrow \tilde{Y}^+$ that fits into the commutative diagram*

$$(3.1.8) \quad \begin{array}{ccc} \tilde{Y} & \overset{\psi}{\dashrightarrow} & \tilde{Y}^+ \\ \downarrow \sigma & \searrow \hat{\psi} & \downarrow \tilde{\sigma}_+ \\ & & \hat{Y}^+ \\ & \searrow \hat{\psi}_0 & \downarrow \hat{\sigma}_+ \\ Y & \overset{\psi_0}{\dashrightarrow} & Y^+ \end{array} \quad \begin{array}{c} \curvearrowright \\ \sigma_+ \end{array}$$

where $\hat{\sigma}_+: \hat{Y}^+ = \mathbb{P}_{Y^+}(\mathcal{E}) \rightarrow Y^+$ is the projection, $\tilde{\sigma}_+: \tilde{Y}^+ = \text{Bl}_{s_{p+1}(Y^+) \sqcup \dots \sqcup s_{p+q}(Y^+)}(\hat{Y}^+) \rightarrow \hat{Y}^+$ is the blowup, and $\sigma_+ = \hat{\sigma}_+ \circ \tilde{\sigma}_+$, and such that ψ induces isomorphisms of G -orbits

$$\tilde{Y}_{\emptyset}^{\circ} \cong (\tilde{Y}^+)^{\circ}, \quad E_{\emptyset}^{\circ} \cong (\tilde{Y}_{\emptyset}^+)^{\circ}, \quad \text{and} \quad \tilde{Y}_i^{\circ} \cong E_i^{\circ}$$

of codimension 0 and 1. Moreover, if

- H_i , $1 \leq i \leq r$, are the hyperplane classes of $\mathbb{P}(V_i)$ and $H = H_1 + \dots + H_r$,
- E is the exceptional divisor of σ ,
- h is the relative hyperplane class of the projective bundle $\hat{\sigma}_+$,
- h_i , $1 \leq i \leq p$, are the hyperplane classes of $\mathbb{P}(\bar{V}_i)$, and
- e_i , $p+1 \leq i \leq p+q$, are the exceptional divisor classes of the blowup $\tilde{\sigma}_+$,

then in the Picard group $\text{Pic}(\tilde{Y}) = \text{Pic}(\tilde{Y}^+)$ there are the following equalities

$$(3.1.9) \quad \begin{aligned} h_i &= H_i - E, & 1 \leq i \leq p, \\ e_i &= H_i - E, & p+1 \leq i \leq p+q, \\ h &= H - (r-1)E. \end{aligned}$$

Conversely, one has

$$(3.1.10) \quad E = h - \sum_{i=1}^p h_i - \sum_{j=p+1}^{p+q} e_j, \quad H_i = \begin{cases} h_i + E, & \text{for } 1 \leq i \leq p, \\ e_i + E, & \text{for } p+1 \leq i \leq p+q. \end{cases}$$

Proof. For each $u_i \in V_i$ denote by $\bar{u}_i \in \bar{V}_i$ the linear projection of u_i from the fixed vector $v_i \in V_i$. Then the rational map $\psi_0: Y \dashrightarrow Y^+$ is given by the formula

$$(u_1, \dots, u_r) \mapsto (\bar{u}_1, \dots, \bar{u}_r).$$

This map is well defined on the open orbit $Y^\circ \subset Y$ (given by the conditions $\bar{u}_i \neq 0$ for all $1 \leq i \leq r$) and it extends regularly to the orbits $Y_i^\circ \subset Y$ of codimension 1 (given by the condition $\bar{u}_i = 0$ for some $p+1 \leq i \leq p+q$ and $\bar{u}_j \neq 0$ for all $j \neq i$).

Now consider the rational G -equivariant map

$$(3.1.11) \quad \hat{\psi}_0: Y \dashrightarrow \hat{Y}^+, \quad (u_1, \dots, u_r) \mapsto \left((\bar{u}_1, \dots, \bar{u}_r), \sum_{|I| \geq r-1} \bigotimes_{i \in I} \bar{u}_i \right).$$

Here we consider the summand $\bigotimes_{i \in I} \bar{u}_i$ as a point in the fiber of the line bundle $\mathcal{O}(-h_I)$ and their sum for $|I| \geq r-1$ as a point in the fiber (of the projectivization) of \mathcal{E} . The map $\hat{\psi}_0$ induces an isomorphism of the open orbit $Y^\circ \subset Y$ onto the open orbit $\mathbb{P}_{Y^+}(\mathcal{E}) \setminus \mathbb{P}_{Y^+}(\bar{\mathcal{E}}_\emptyset)$ in \hat{Y}^+ and contracts each orbit Y_i° of codimension 1 to the section $s_i(Y^+) \subset \hat{Y}^+$.

Now consider the composition $\hat{\psi} = \hat{\psi}_0 \circ \sigma: \tilde{Y} \dashrightarrow \hat{Y}^+$. The restriction of $\hat{\psi}$ to the exceptional divisor E is given by

$$(3.1.12) \quad E = \mathbb{P}(\bar{V}_1 \oplus \dots \oplus \bar{V}_r) \dashrightarrow \hat{Y}^+, \quad (\bar{u}_1 + \dots + \bar{u}_r) \mapsto \left((\bar{u}_1, \dots, \bar{u}_r), \sum_{|I|=r-1} \bigotimes_{i \in I} \bar{u}_i \right).$$

It maps the orbit $E_\emptyset^\circ \subset \tilde{Y}$ isomorphically onto the G -orbit $(\tilde{Y}_\emptyset^+)^{\circ} = \mathbb{P}_{Y^+}(\bar{\mathcal{E}}) \setminus (\bigcup_{i=1}^r \mathbb{P}_{Y^+}(\bar{\mathcal{E}}_i))$ of codimension 1. By the above arguments it also gives an isomorphism of open G -orbits and contracts the orbits $\tilde{Y}_i^\circ \cong Y_i^\circ$, $p+1 \leq i \leq p+q$, to the sections $s_i(Y^+) \subset \hat{Y}^+$. Therefore, $\hat{\psi}$ induces a birational isomorphism

$$\psi: \tilde{Y} \dashrightarrow \text{Bl}_{s_{p+1}(Y^+) \sqcup \dots \sqcup s_{p+q}(Y^+)}(\mathbb{P}_{Y^+}(\mathcal{E})) = \tilde{Y}^+.$$

Finally, it is easy to see that the induced map $\tilde{Y}_i^\circ \rightarrow E_i^\circ$ is an isomorphism for all $p+1 \leq i \leq p+q$. This gives the commutative diagram (3.1.8) and proves that ψ is small.

The first two lines in (3.1.9) follow easily from the formulas (3.1.11), (3.1.12), and (3.1.2). The last line follows from the equality of the canonical classes of \tilde{Y} and \tilde{Y}^+ expressed in terms of H_i and E on the one hand, and h_i , h , and e_i on the other hand.

Finally, (3.1.10) follows from (3.1.9). □

Remark 3.2. Alternatively, one can use the fact that the varieties \tilde{Y} and \tilde{Y}^+ , as well as the birational isomorphism ψ are toric. Thus, to check that ψ is small, it is enough to identify the generators of rays of the corresponding fans. Moreover, comparing the other cones in the fans one can check that the map ψ factors as the composition

$$\tilde{Y} \xrightarrow{\psi_1} \tilde{Y}' \xrightarrow{\psi_2} \dots \xrightarrow{\psi_{r-2}} \tilde{Y}^{(r-2)} \xrightarrow{\psi_{r-1}} \tilde{Y}^+$$

of standard (anti)flips ψ_l in the strict transforms of \tilde{Y}_I for $|I| = l$, $1 \leq l \leq r-2$, and for $|I| = r-1$ with $\{p+1, \dots, p+q\} \subset I$, respectively.

3.2. Rationality of forms of products of projective spaces. Here we apply the birational transformation of the previous subsection to deduce the following corollary (see [Zak07] for a different proof).

Corollary 3.3. *Let Y be a \mathbf{k} -form of $\mathbb{P}^{n_1} \times \mathbb{P}^{n_2} \times \dots \times \mathbb{P}^{n_r}$. For any $y \in Y(\mathbf{k})$ the diagram (3.1.8) is defined over \mathbf{k} and if $Y(\mathbf{k}) \neq \emptyset$ then Y is \mathbf{k} -rational.*

Proof. First, we prove that for any $y \in Y(\mathbf{k})$ the diagram (3.1.8) is defined over \mathbf{k} . The divisor classes $H = \sum_{i=1}^r H_i$ and $H' := \sum_{i=1}^p H_i$ on $Y_{\bar{\mathbf{k}}}$ are Galois-invariant, and since we have $Y(\mathbf{k}) \neq \emptyset$ by assumption, we conclude that they are defined over \mathbf{k} . Also \tilde{Y} and E are defined over \mathbf{k} as y is a \mathbf{k} -point. Therefore, the divisor classes

$$\sum_{i=1}^p h_i = H' - pE, \quad h = H - (r-1)E, \quad \text{and} \quad - \sum_{i=p+1}^{p+q} e_i = H - H' - qE$$

(which by Theorem 3.1 are equal to the strict transforms of the classes that are ample on Y^+ , relatively ample for $\hat{Y}^+ \rightarrow Y^+$ and for $\tilde{Y}^+ \rightarrow \hat{Y}^+$, respectively) are defined over \mathbf{k} , hence the varieties Y^+ , \hat{Y}^+ , and \tilde{Y}^+ , equal to the images of \tilde{Y} under the maps given by their appropriate linear combinations, are defined over \mathbf{k} , as well as the remaining arrows in the diagram.

Now to prove \mathbf{k} -rationality of Y we argue by induction in $\dim(Y) = \sum n_i$. If the dimension is zero, there is nothing to prove. So, assume the dimension is positive and consider the diagram (3.1.8). By Theorem 3.1 the variety Y^+ is a \mathbf{k} -form of $Y_{\bar{\mathbf{k}}}^+ = \mathbb{P}_{\bar{\mathbf{k}}}^{n_1-1} \times \mathbb{P}_{\bar{\mathbf{k}}}^{n_2-1} \times \dots \times \mathbb{P}_{\bar{\mathbf{k}}}^{n_r-1}$. By Nishimura lemma we have $Y^+(\mathbf{k}) \neq \emptyset$, hence Y^+ is \mathbf{k} -rational by the induction assumption. Furthermore, the morphism $\hat{Y}^+ \rightarrow Y^+$ is a \mathbf{k} -form of a projective bundle, and by (3.1.10) the strict transform of the exceptional divisor E of \tilde{Y} provides for it a relative hyperplane section. But E is defined over \mathbf{k} , therefore \tilde{Y}^+ is rational over Y^+ , hence it is \mathbf{k} -rational. It remains to note that the morphisms σ , ψ , and $\tilde{\sigma}_+$ in (3.1.8) are birational, hence Y is \mathbf{k} -rational as well. \square

Applying this to the case of a \mathbf{k} -form of $(\mathbb{P}^1)^3$ we obtain

Corollary 3.4. *If X is a Fano threefold of type $X_{(1,1,1)}$ with $X(\mathbf{k}) \neq \emptyset$ then X is \mathbf{k} -rational.*

For other applications of the theorem we will often use the following observation. Recall the definitions (3.1.2) and (3.1.7) of the subvarieties $Y_i \subset Y$ of codimension n_i and subbundles $\bar{\mathcal{E}}_i \subset \mathcal{E}$ of corank 2.

Proposition 3.5. *Let Y be a \mathbf{k} -form of $(\mathbb{P}^n)^r$ where $n \geq 2$ and assume Y has a \mathbf{k} -point $y \in Y(\mathbf{k})$. Let $X \subset Y$ be a closed \mathbf{k} -subvariety containing the point y such that*

$$X_{\bar{\mathbf{k}}} = \bigcap_{\alpha=1}^c D_{\alpha} \subset (\mathbb{P}_{\bar{\mathbf{k}}}^n)^r$$

is a complete intersection of divisors $D_{\alpha} \subset (\mathbb{P}_{\bar{\mathbf{k}}}^n)^r$, $1 \leq \alpha \leq c$. Let $\tilde{D}_{\alpha} \subset \tilde{Y}_{\bar{\mathbf{k}}}$ and $\tilde{D}_{\alpha}^+ \subset \tilde{Y}_{\bar{\mathbf{k}}}^+ = \hat{Y}_{\bar{\mathbf{k}}}^+$ be the strict transforms of D_{α} and set

$$\tilde{X}_{\bar{\mathbf{k}}}^+ := \bigcap_{\alpha=1}^c \tilde{D}_{\alpha}^+ \subset \tilde{Y}_{\bar{\mathbf{k}}}^+ = \hat{Y}_{\bar{\mathbf{k}}}^+ = \mathbb{P}_{(\mathbb{P}_{\bar{\mathbf{k}}}^{n-1})^r}(\mathcal{E}).$$

If X is smooth at y and for each $1 \leq i \leq r$ one has

$$(3.2.1) \quad \dim(X_{\bar{\mathbf{k}}} \cap (Y_i)_{\bar{\mathbf{k}}}) < \dim(X_{\bar{\mathbf{k}}}) \quad \text{and} \quad \dim(\tilde{X}_{\bar{\mathbf{k}}}^+ \cap \mathbb{P}_{Y_{\bar{\mathbf{k}}}^+}(\bar{\mathcal{E}}_i)) < \dim(X_{\bar{\mathbf{k}}})$$

then the strict transform $\tilde{X}^+ = \psi_(\text{Bl}_y(X))$ of X in \tilde{Y}^+ is a \mathbf{k} -form of the complete intersection $\tilde{X}_{\bar{\mathbf{k}}}^+$.*

Proof. By Corollary 3.3 the diagram (3.1.8) is defined over \mathbf{k} , so it is enough to check that the complete intersection $\tilde{X}_{\bar{\mathbf{k}}}^+$ is equal to the strict transform of $\text{Bl}_y(X_{\bar{\mathbf{k}}})$. First, note that the assumption that y is a smooth point of X implies that the strict transform $\text{Bl}_y(X_{\bar{\mathbf{k}}})$ of $X_{\bar{\mathbf{k}}}$ in \tilde{Y} is the complete intersection of the divisors \tilde{D}_{α} . Furthermore, the first part of the assumptions (3.2.1) implies that the intersection of $\text{Bl}_y(X_{\bar{\mathbf{k}}})$ with the open G -orbit in $\tilde{Y}_{\bar{\mathbf{k}}}$ is dense in $\text{Bl}_y(X_{\bar{\mathbf{k}}})$. Therefore, the strict transform of $\text{Bl}_y(X_{\bar{\mathbf{k}}})$ in $\tilde{Y}_{\bar{\mathbf{k}}}^+$ is contained in $\tilde{X}_{\bar{\mathbf{k}}}^+$. So, it remains to check that $\tilde{X}_{\bar{\mathbf{k}}}^+$ is irreducible of dimension $\dim(X_{\bar{\mathbf{k}}})$. This is definitely true for the intersection of $\tilde{X}_{\bar{\mathbf{k}}}^+$ with the complement of the union of projective subbundles $\mathbb{P}_{Y_{\bar{\mathbf{k}}}^+}(\bar{\mathcal{E}}_i)$, because the map ψ defines an isomorphism of this complement with an open subset of $\tilde{Y}_{\bar{\mathbf{k}}}$. On the other hand, the second part of the assumptions (3.2.1) gives a bound for the dimension of the intersections with these projective subbundles, which implies the irreducibility. \square

4. RATIONALITY AND UNIRATIONALITY OF TYPES $X_{(2,2)}$, $X_{(2,2,2)}$, AND $X_{(1,1,1,1)}$

In this section we prove rationality of Fano threefolds of types $X_{(2,2)}$ and $X_{(2,2,2)}$ as well as unirationality of threefolds of type $X_{(1,1,1,1)}$ under the assumption $X(\mathbf{k}) \neq \emptyset$.

4.1. Rationality of $X_{(2,2)}$. To start with we deal with threefolds of type $X_{(2,2)}$.

Proposition 4.1. *Let X be a Fano threefold of type $X_{(2,2)}$. If $X(\mathbf{k}) \neq \emptyset$ then X is \mathbf{k} -rational.*

Proof. Let x be a \mathbf{k} -point of X . By definition X is a smooth divisor of bidegree $(1, 1)$ in a \mathbf{k} -form Y of $\mathbb{P}^2 \times \mathbb{P}^2$. By Theorem 3.1 it is birational to a \mathbf{k} -form of a divisor

$$\tilde{X}_{\bar{\mathbf{k}}}^+ \subset \mathbb{P}_{\mathbb{P}^1 \times \mathbb{P}^1}(\mathcal{E}) = \mathbb{P}_{\mathbb{P}^1 \times \mathbb{P}^1}(\mathcal{O}(-1, -1) \oplus \mathcal{O}(-1, 0) \oplus \mathcal{O}(0, -1))$$

of type $H_1 + H_2 - E = h$; any such divisor corresponds to a morphism

$$\xi: \mathcal{O}(-1, -1) \oplus \mathcal{O}(-1, 0) \oplus \mathcal{O}(0, -1) \longrightarrow \mathcal{O}.$$

Furthermore, the divisor $\tilde{X}^+ \subset \tilde{Y}^+$ comes with a morphism $\sigma_+: \tilde{X}^+ \rightarrow Y^+$ defined over \mathbf{k} . By Nishimura lemma we have $\tilde{X}^+(\mathbf{k}) \neq \emptyset$, hence $Y^+(\mathbf{k}) \neq \emptyset$, and since Y^+ is a \mathbf{k} -form of $\mathbb{P}^1 \times \mathbb{P}^1$, it is \mathbf{k} -rational by Corollary 3.3. Finally, the general fiber of the morphism $\sigma_+: \tilde{X}^+ \rightarrow Y^+$ is a

1-dimensional linear section of a form of a projective plane, hence it is isomorphic to \mathbb{P}^1 , hence \tilde{X}^+ is rational over Y^+ , hence is \mathbf{k} -rational, hence so is X . \square

Remark 4.2. One can check that the birational isomorphism $\psi: \tilde{X} \dashrightarrow \tilde{X}^+$ is a flop in the union of the strict transforms of two $\bar{\mathbf{k}}$ -lines passing through the point $x \in X$ and that $\sigma_+: \tilde{X}^+ \rightarrow Y^+$ is the projectivization of the rank-2 vector bundle $\text{Ker}(\xi)$ over Y^+ , and that these maps provide a Sarkisov link (1.2.1).

4.2. Rationality of $X_{(2,2,2)}$. A similar argument works for threefolds of type $X_{(2,2,2)}$.

Proposition 4.3. *Let X be a Fano threefold of type $X_{(2,2,2)}$. If $X(\mathbf{k}) \neq \emptyset$ then X is \mathbf{k} -rational.*

Proof. Let x be a \mathbf{k} -point of X . By definition $X_{\bar{\mathbf{k}}}$ is a complete intersection of three divisors in $Y_{\bar{\mathbf{k}}} = \mathbb{P}(V_1) \times \mathbb{P}(V_2) \times \mathbb{P}(V_3) \cong \mathbb{P}^2 \times \mathbb{P}^2 \times \mathbb{P}^2$ of multidegree $(1, 1, 0)$, $(1, 0, 1)$, and $(0, 1, 1)$, respectively. Denote by

$$F_{12} \in V_1^\vee \otimes V_2^\vee, \quad F_{13} \in V_1^\vee \otimes V_3^\vee, \quad F_{23} \in V_2^\vee \otimes V_3^\vee,$$

their equations. To apply Proposition 3.5 consider the intersection

$$\tilde{X}_{\bar{\mathbf{k}}}^+ \subset \mathbb{P}_{\mathbb{P}^1 \times \mathbb{P}^1 \times \mathbb{P}^1}(\mathcal{E}) = \mathbb{P}_{\mathbb{P}^1 \times \mathbb{P}^1 \times \mathbb{P}^1}(\mathcal{O}(-1, -1, -1) \oplus \mathcal{O}(-1, -1, 0) \oplus \mathcal{O}(-1, 0, -1) \oplus \mathcal{O}(0, -1, -1))$$

of three divisors which by (3.1.10) have types

$$H_1 + H_2 - E = h - h_3, \quad H_1 + H_3 - E = h - h_2, \quad H_2 + H_3 - E = h - h_1,$$

hence correspond to a morphism of vector bundles

$$\begin{aligned} \xi: \mathcal{O}(-1, -1, -1) \oplus \mathcal{O}(-1, -1, 0) \oplus \mathcal{O}(-1, 0, -1) \oplus \mathcal{O}(0, -1, -1) &\longrightarrow \\ &\longrightarrow \mathcal{O}(0, 0, -1) \oplus \mathcal{O}(0, -1, 0) \oplus \mathcal{O}(-1, 0, 0). \end{aligned}$$

It is easy to see that ξ is given by the matrix

$$(4.2.1) \quad \xi = \begin{pmatrix} \bar{F}_{12} & 0 & F_{12}(-, v_2) & F_{12}(v_1, -) \\ \bar{F}_{13} & F_{13}(-, v_3) & 0 & F_{13}(v_1, -) \\ \bar{F}_{23} & F_{23}(-, v_3) & F_{23}(v_2, -) & 0 \end{pmatrix},$$

where we write $x = (v_1, v_2, v_3)$, choose splittings $V_i = \bar{\mathbf{k}}v_i \oplus \bar{V}_i$, write \bar{F}_{ij} for the restriction of F_{ij} to $\bar{V}_i \otimes \bar{V}_j$, and consider $F_{ij}(v_i, -)$ and $F_{ij}(-, v_j)$ as linear functions on V_j and V_i , vanishing at v_j and v_i , hence as linear functions on \bar{V}_j and \bar{V}_i , respectively.

Let us check the dimension conditions (3.2.1). Since $(Y_i)_{\bar{\mathbf{k}}}$ is a fiber of the projection

$$\pi_i: Y_{\bar{\mathbf{k}}} = \mathbb{P}^2 \times \mathbb{P}^2 \times \mathbb{P}^2 \longrightarrow \mathbb{P}^2,$$

it follows from Lemma 2.8 that $X_{\bar{\mathbf{k}}} \cap (Y_i)_{\bar{\mathbf{k}}}$ is a conic, hence the first part of the dimension conditions is satisfied. To check the second part, we need to show that the restriction

$$\xi_{2,3}: \mathcal{O}(-1, 0, -1) \oplus \mathcal{O}(0, -1, -1) \longrightarrow \mathcal{O}(0, 0, -1) \oplus \mathcal{O}(0, -1, 0) \oplus \mathcal{O}(-1, 0, 0)$$

of ξ to the last two summands of \mathcal{E} (given by the last two columns of (4.2.1)) cannot be everywhere degenerate, and similarly for the restrictions $\xi_{1,3}$ and $\xi_{1,2}$. Assuming that $\xi_{2,3}$ is everywhere degenerate we conclude from (4.2.1) that

$$F_{12}(v_1, -) = F_{13}(v_1, -) = 0 \quad \text{or} \quad F_{12}(-, v_2) = F_{23}(v_2, -) = 0 \quad \text{or} \quad F_{13}(v_1, -) = F_{23}(v_2, -) = 0.$$

In any case it would follow that at least two of the divisors $W_{i,j} \subset \mathbb{P}(V_i) \times \mathbb{P}(V_j)$ (defined by the equation $F_{i,j}$) are singular, which contradicts Lemma 2.4(i).

Thus, the conditions (3.2.1) are satisfied, and we conclude from Proposition 3.5 that X is \mathbf{k} -birational to a \mathbf{k} -form of the complete intersection $\tilde{X}_{\bar{\mathbf{k}}}^+$.

Finally, the subvariety $\tilde{X}^+ \subset \tilde{Y}^+$ comes with a morphism $\sigma_+ : \tilde{X}^+ \rightarrow Y^+$ defined over \mathbf{k} . By Nishimura lemma we have $\tilde{X}^+(\mathbf{k}) \neq \emptyset$, hence $Y^+(\mathbf{k}) \neq \emptyset$, and since Y^+ is a \mathbf{k} -form of $\mathbb{P}^1 \times \mathbb{P}^1 \times \mathbb{P}^1$, it is \mathbf{k} -rational by Corollary 3.3. Moreover, the general fiber of the morphism $\sigma_+ : \tilde{X}^+ \rightarrow Y^+$ is a 0-dimensional linear section of a form of a projective space, hence this morphism is birational, hence \tilde{X}^+ is \mathbf{k} -rational, hence so is X . \square

Remark 4.4. If the point x does not lie on a $\bar{\mathbf{k}}$ -line, i.e., $F_1(X, x) = \emptyset$, one can check that the birational isomorphism $\psi : \tilde{X} \dashrightarrow \tilde{X}^+$ is a flop in the union of the strict transforms of three smooth $\bar{\mathbf{k}}$ -conics passing through the point $x \in X$, that $\sigma_+ : \tilde{X}^+ \rightarrow Y^+$ is the blowup of a smooth geometrically rational curve of multidegree $(2, 2, 2)$ and that these maps provide a Sarkisov link (1.2.1).

4.3. Unirationality of $X_{(1,1,1,1)}$. Finally, we deal with threefolds of type $X_{(1,1,1,1)}$.

Proposition 4.5. *Let X be a Fano threefold of type $X_{(1,1,1,1)}$. If $X(\mathbf{k}) \neq \emptyset$ then X is \mathbf{k} -unirational.*

Proof. Let x be a \mathbf{k} -point of X . By definition X is a smooth divisor of multidegree $(1, 1, 1, 1)$ in a \mathbf{k} -form Y of $\mathbb{P}^1 \times \mathbb{P}^1 \times \mathbb{P}^1 \times \mathbb{P}^1$. By Theorem 3.1 it is birational to a \mathbf{k} -form of a divisor

$$\tilde{X}_{\bar{\mathbf{k}}}^+ \subset \text{Bl}_{s_1, s_2, s_3, s_4}(\mathbb{P}^4)$$

which by (3.1.10) has type $H_1 + H_2 + H_3 + H_4 - E = 3h - 2 \sum_{i=1}^4 e_i$. Moreover, the exceptional divisor $E \cap \text{Bl}_x(X) \subset \tilde{Y} = \text{Bl}_x(Y)$ of the blowup $\text{Bl}_x(X) \rightarrow X$ is an irreducible \mathbf{k} -rational surface birational to a \mathbf{k} -form of the complete intersection of $\tilde{X}_{\bar{\mathbf{k}}}^+$ with the linear span $\mathbb{P}^3 \subset \mathbb{P}^4$ of the points s_i .

Let us prove that $\tilde{X}_{\bar{\mathbf{k}}}^+$ is not a cone. Indeed, $\tilde{X}_{\bar{\mathbf{k}}}^+$ is smooth away from the linear span \mathbb{P}^3 of the s_i , because the map ψ from Theorem 3.1 is an isomorphism over it and $X_{\bar{\mathbf{k}}}$ is smooth, so if $\tilde{X}_{\bar{\mathbf{k}}}^+$ is a cone, its vertex belongs to the \mathbb{P}^3 . But then its intersection with the \mathbb{P}^3 (which has been shown to be an irreducible \mathbf{k} -rational surface) is itself a cone and has singular points at all the s_i . But it is easy to see that any such cone is reducible; this contradiction proves the claim. Now we conclude that $\tilde{X}_{\bar{\mathbf{k}}}^+$ is \mathbf{k} -unirational by [Kol02]. \square

5. RATIONALITY OF TYPE $X_{(4,4)}$

In this section we prove rationality of Fano threefolds of type $X_{(4,4)}$.

5.1. Sarkisov links. We start with a construction of two Sarkisov links. Recall that $F_1(X, x)$ denotes the Hilbert scheme of lines on X passing through x , see §2.2; and that by Lemma 2.7 if $x \in X(\mathbf{k})$ and this scheme is not empty, it is the union of two reduced $\bar{\mathbf{k}}$ -points swapped by the Galois action. If L_1 and L_2 are the corresponding $\bar{\mathbf{k}}$ -lines on X (passing through x), then

$$\Theta(x) := L_1 \cup L_2$$

is a singular \mathbf{k} -conic on X irreducible over \mathbf{k} with $\text{Sing}(\Theta(x)) = \{x\}$.

Recall that a quintic del Pezzo threefold is a Fano threefold of index 2 and half-anticanonical degree 5. Over an algebraically closed field it can be realized as a linear section of the Grassmannian $\text{Gr}(2, 5)$ by a subspace of codimension 3, see [Isk80, Chapter 2, Theorem 1.1].

Theorem 5.1. *Let X be a Fano threefold of type $X_{(4,4)}$ and let $x \in X(\mathbf{k})$ be a \mathbf{k} -point.*

- (i) *If $F_1(X, x) = \emptyset$ then there exists a Sarkisov link (1.2.1) defined over \mathbf{k} , where*
 - σ is the blowup of the point x ,
 - X^+ is a smooth quintic del Pezzo threefold, and
 - σ_+ is the blowup of a smooth \mathbf{k} -irreducible curve $B^+ \subset X^+$ of degree 4 with two geometrically rational components.
- (ii) *If $F_1(X, x) \neq \emptyset$ then there exists a Sarkisov link (1.2.1) defined over \mathbf{k} , where*
 - σ is the blowup of the singular \mathbf{k} -irreducible conic $\Theta(x)$,
 - X^+ is a smooth Fano threefold of type $X_{(2,2)}$, and
 - σ_+ is the blowup of a singular \mathbf{k} -irreducible curve $B^+ \subset X^+$ of degree 6 with two geometrically rational components.

The proof of the theorem takes §5.1 and §5.2: in the rest of §5.1 we prove the existence of the links, and in §5.2 we describe their details. The proofs of cases (i) and (ii) are completely analogous, so to carry them on simultaneously we introduce the following convenient notation

$$(5.1.1) \quad m = m(x) := \begin{cases} 2, & \text{if } F_1(X, x) = \emptyset, \\ 1, & \text{if } F_1(X, x) \neq \emptyset. \end{cases}$$

The proof of the existence of the links is analogous to the first parts of [KP19, Theorem 5.18 and Theorem 5.10], so we use below some results from [KP19]. Note that although in [KP19] we work under the assumption $\rho(X_{\bar{\mathbf{k}}}) = 1$, the arguments in the proofs do not use it.

Let

$$\sigma: \tilde{X} \longrightarrow X$$

be the blowup of X at x or at $\Theta(x)$, respectively. We denote by H (the pullback to \tilde{X} of) the anticanonical class of X and by E (the class of) the exceptional divisor of σ .

First, note that the anticanonical linear system

$$|-K_{\tilde{X}}| = |H - mE|$$

is base-point free by Theorem 2.1 and [KP19, Lemma 5.6 and 5.5]. Moreover, combining [KP19, (5.1.9) and (5.1.7)], we can uniformly write

$$(5.1.2) \quad H^3 = 2g - 2, \quad H^2 \cdot E = 0, \quad H \cdot E^2 = 2(m - 2), \quad E^3 = m - 1,$$

where recall from Table 1 that $g = g(X) = 15$. We will also need the following observation.

Lemma 5.2. *The linear system $\mathcal{M} := |H - (m+1)E|$ on the blowup \tilde{X} of X has positive dimension*

$$(5.1.3) \quad \dim \mathcal{M} \geq g - m - 7 \geq 6,$$

and has no fixed components.

Proof. The dimension is estimated in [KP19, Lemma 5.4(iii) and (i)]. To prove that \mathcal{M} has no fixed components, note that the divisor E is rigid (since it is the exceptional divisor of a blowup),

hence the only possibility for a fixed component of \mathcal{M} is the divisor E with some multiplicity. So, assume

$$|H - (m + 1)E| = (a - m - 1)E + |H - aE|,$$

where $a \geq m + 2$ and E is not a fixed component of the linear system $|H - aE|$. Then a general member of the linear system $|H - aE|$ is irreducible with $\dim |H - aE| = \dim \mathcal{M} > 0$, and since the linear system $|H - mE|$ is base-point free, using (5.1.2) we obtain

$$0 \leq (H - aE)^2 \cdot (H - mE) = 2g - 2 - a^2(m^2 - 3m + 4) + 4am(m - 2).$$

When $m = 2$ this gives $a^2 \leq 14$, hence $a \leq 3$, and when $m = 1$ this gives $(a + 1)^2 \leq 15$, hence $a \leq 2$. In both cases this contradicts the assumption $a \geq m + 2$. \square

Now we can deduce the existence of the Sarkisov links.

Proposition 5.3. *Let X be a Fano threefold of type $X_{(4,4)}$ with a \mathbf{k} -point x .*

- (i) *If $F_1(X, x) = \emptyset$, there exists a Sarkisov link (1.2.1), where σ is the blowup of x .*
- (ii) *If $F_1(X, x) \neq \emptyset$, there exists a Sarkisov link (1.2.1), where σ is the blowup of $\Theta(x)$.*

In both cases the link is defined over \mathbf{k} .

Proof. We use notation (5.1.1). Recall that the anticanonical class $H = -K_X$ is very ample and the image of the anticanonical embedding $X \subset \mathbb{P}^{g+1}$ is an intersection of quadrics (see Theorem 2.1). The anticanonical morphism $\phi: \tilde{X} \rightarrow \mathbb{P}^{g-m-1}$ cannot contract a divisor D , because by [KP19, Lemma 5.6 and 5.5] this divisor is then a fixed component of \mathcal{M} , but by Lemma 5.2 this linear system has no fixed components. Therefore, the required link exists by [KP19, Lemma 5.6 and 5.5]. \square

5.2. The second contraction. By Proposition 5.3 we have the diagram (1.2.1), so to finish the proof of Theorem 5.1 it remains to describe the extremal contraction σ_+ . During this step we systematically use the classification of extremal contractions from [Mor82] and [Cut88].

We denote by $H^+, E^+ \in \text{Pic}(\tilde{X}^+)$ the strict transforms of the classes $H, E \in \text{Pic}(\tilde{X})$. Note that

$$-K_{\tilde{X}^+} = H^+ - mE^+$$

because the birational isomorphism ψ is small. Consider also the strict transform

$$\mathcal{M}^+ := |H^+ - (m + 1)E^+|$$

of the linear system \mathcal{M} .

We denote by $\Upsilon_i \subset \tilde{X}$ the flopping curves and by $\Upsilon_i^+ \subset \tilde{X}^+$ the corresponding flopped curves. Finally, when $F_1(X, x) = \emptyset$ we denote by C a general twisted cubic curve on X passing through x and otherwise we denote by C a general conic meeting $\Theta(x)$ (recall Lemma 2.11 and Lemma 2.10 for the description of the corresponding Hilbert schemes). Note that

$$(5.2.1) \quad H \cdot C = m + 1.$$

We denote by \tilde{C} the strict transform of C in \tilde{X} and by \tilde{C}^+ the strict transform of \tilde{C} in \tilde{X}^+ . Note that by Remark 2.9 and Remark 2.12 the curve \tilde{C} is smooth and

$$(5.2.2) \quad E \cdot \tilde{C} = 1;$$

in particular \tilde{C} does not contain the curves Υ_i .

Lemma 5.4. *The nef cone of \tilde{X}^+ is generated by the anticanonical class $-K_{\tilde{X}^+}$ and $M^+ \in \mathcal{M}^+$, and the Mori cone of \tilde{X}^+ is generated by the class of the curves Υ_i^+ and the class of \tilde{C}^+ . The extremal contraction σ_+ is given by a multiple of the linear system \mathcal{M}^+ and contracts the extremal ray generated by \tilde{C}^+ .*

Proof. Since ϕ is crepant and ψ is a flop, the morphism ϕ_+ is crepant as well. Moreover, the anticanonical linear system $| -K_{\tilde{X}} |$ is base-point free by [KP19, Lemma 5.6], hence $-K_{\tilde{X}}^+$ is nef.

On the other hand, we have $(H - mE) \cdot \Upsilon_i = -K_X \cdot \Upsilon_i = 0$, and since $H \cdot \Upsilon_i > 0$, we conclude that $E \cdot \Upsilon_i > 0$. Therefore, for $M \in \mathcal{M}$ we have

$$M \cdot \Upsilon_i = (H - (m + 1)E) \cdot \Upsilon_i = -E \cdot \Upsilon_i < 0.$$

If $M^+ \in \mathcal{M}^+$ is the strict transform of M , this implies that $M^+ \cdot \Upsilon_i^+ > 0$ by definition of a flop, see e.g. [KM98, Definition 6.10]. Now if M^+ is not nef, it is negative on the extremal ray R corresponding to the contraction $\sigma_+ : \tilde{X}^+ \rightarrow X^+$. Since the canonical class is also negative on R , the contraction σ_+ cannot be small (see [Ben85, Theorem 0] or [MP19, Corollary 6.3.4]), hence curves in R sweep a subvariety of \tilde{X}^+ of dimension ≥ 2 , hence the base locus of \mathcal{M}^+ is at least 2-dimensional, which contradicts Lemma 5.2. This proves that M^+ is nef.

Now we have $-K_{\tilde{X}} \cdot \tilde{C} = (H - mE) \cdot \tilde{C} = 1$ by (5.2.1) and (5.2.2), hence $-K_{\tilde{X}^+} \cdot \tilde{C}^+ = 1$. On the other hand, since a general divisor H meets C away from the indeterminacy locus of the map $X \dashrightarrow \tilde{X}^+$, we have $H^+ \cdot \tilde{C}^+ \geq H \cdot C = m + 1$ and so $E^+ \cdot \tilde{C}^+ \geq 1$. Thus $M^+ \cdot \tilde{C}^+ \leq 0$. Since M^+ is nef, $M^+ \cdot \tilde{C}^+ = 0$.

Combining the above computations we conclude that the nef cone of \tilde{X}^+ is generated by $-K_{\tilde{X}^+}$ and M^+ and the Mori cone is generated by Υ_i^+ and \tilde{C}^+ . The rest of the lemma follows from the Mori contraction theorem. \square

Now we can finally prove Theorem 5.1.

Proof of Theorem 5.1. Let X be a Fano threefold of type $X_{(4,4)}$. Since σ_+ is an extremal contraction, we have $\rho(X^+) = \rho(\tilde{X}^+) - 1 = \rho(\tilde{X}) - 1 = \rho(X)$, hence $\rho(X^+) = 1$. Similarly, we have $\rho(X_{\bar{k}}^+) \leq \rho(\tilde{X}_{\bar{k}}^+) - 1 = \rho(\tilde{X}_{\bar{k}}) - 1 = \rho(X_{\bar{k}}) = 2$, hence

$$(5.2.3) \quad \rho(X_{\bar{k}}^+) \leq 2.$$

On the other hand, in the case $F_1(X, x) \neq \emptyset$, the varieties \tilde{X} and \tilde{X}^+ are not smooth and arguing as in the proof of [KP19, Theorem 5.10] we obtain

$$(5.2.4) \quad \text{rk Cl}(\tilde{X}^+) = 2, \quad \text{rk Cl}(\tilde{X}_{\bar{k}}^+) = 5 - m.$$

Since ϕ and ϕ_+ are crepant morphisms, the projection formula implies that a triple intersection product of divisor classes on \tilde{X}^+ which includes $K_{\tilde{X}^+}$ is equal to the analogous triple product on \tilde{X} , so using (5.1.2) we compute (recall that $g = g(X) = 15$)

$$(-K_{\tilde{X}^+})^3 = 2(g - m - 3) = 24 - 2m, \quad (-K_{\tilde{X}^+})^2 \cdot E^+ = 4, \quad (-K_{\tilde{X}^+}) \cdot (E^+)^2 = -2.$$

On the other hand, by Lemma 5.4 and primitivity of $H^+ - (m + 1)E^+$ we have

$$H^+ - (m + 1)E^+ = \sigma_+^* A^+,$$

where A^+ is the ample generator of the Picard group of X^+ . We have

$$(5.2.5) \quad (\sigma_+^* A^+)^2 \cdot (-K_{\tilde{X}^+}) = (H^+ - (m + 1)E^+)^2 \cdot (-K_{X^+}) = 2(g - m - 8) = 14 - 2m > 0,$$

therefore σ_+ is not a del Pezzo fibration. Similarly, if σ_+ is a conic bundle, it follows that

$$(A^+)^2 = 7 - m,$$

hence X^+ is a smooth quintic or sextic del Pezzo surface, which of course contradicts the inequality (5.2.3). Therefore, the morphism σ_+ is birational.

By Lemma 5.4 the morphism σ_+ contracts the strict transform R^+ of the divisor swept by curves C , i.e., the strict transform of the divisor $R_x \subset X$ if $F_1(X, x) = \emptyset$, or of the divisor $R_{\Theta(x)}$ otherwise. In both cases Lemma 2.11 and 2.10 show that R^+ has over \bar{k} two irreducible components swapped by the Galois group. Therefore, it follows from (5.2.4) that

$$\text{rk Cl}(X^+) = 1, \quad \text{rk Cl}(X_{\bar{k}}^+) = 3 - m,$$

and σ_+ is the blowup of two \bar{k} -curves or two \bar{k} -points. Furthermore, by Lemmas 2.11 and 2.10 we have

$$R^+ \sim H^+ - (m + 2)E^+.$$

Denoting by i_+ the index of X^+ and by a_+ the discrepancy of the exceptional divisor R^+ of σ_+ , and computing the anticanonical class of \tilde{X}^+ in two ways we obtain the equality

$$H^+ - mE^+ = i_+(H^+ - (m + 1)E^+) - a_+(H^+ - (m + 2)E^+).$$

Solving this equation, we obtain $i_+ = 2$ and $a_+ = 1$. Thus, X^+ is a Fano threefold of index 2 and σ_+ is either the blowup of a k -irreducible curve B^+ , or of rational double points on X^+ swapped by the Galois action. Moreover, using the equality from (5.2.5) we obtain

$$14 - 2m = (\sigma_+^* A^+)^2 \cdot (-K_{\tilde{X}^+}) = (\sigma_+^* A^+)^2 \cdot (2\sigma_+^* A^+ - R^+) = 2(\sigma_+^* A^+)^3,$$

hence X^+ is a quintic or sextic del Pezzo threefold, respectively. Finally, if X^+ is singular, its class group $\text{Cl}(X^+)$ has rank greater than 1 (see [Pro13a, Theorem 1.7]), which contradicts to the equality $\text{rk Cl}(X^+) = 1$ obtained above. Thus, X^+ is smooth and σ_+ is the blowup of a curve B^+ such that $B_{\bar{k}}^+$ has two irreducible \bar{k} -components swapped by the Galois group. To finally compute the degree of B^+ , recall that R^+ is the exceptional divisor of σ_+ . Note that on the one hand

$$(\sigma_+^* A^+) \cdot (-K_{\tilde{X}^+})^2 = 2g - 2m - 10 = 20 - 2m,$$

and on the other hand, this expression is equal to

$$(\sigma_+^* A^+) \cdot (2\sigma_+^* A^+ - R^+)^2 = 4(\sigma_+^* A^+)^3 + (\sigma_+^* A^+) \cdot (R^+)^2 = 4(7 - m) - \deg(B^+).$$

Thus,

$$\deg(B^+) = 8 - 2m,$$

hence B^+ is a quartic or sextic curve with two connected \bar{k} -components (swapped by the Galois action), i.e., a union of two conics or two cubic curves. \square

5.3. Rationality. Now we use the constructed links to prove rationality of threefolds of type $X_{(4,4)}$.

Proposition 5.5. *Let X be a Fano threefold of type $X_{(4,4)}$. If $X(\mathbf{k}) \neq \emptyset$ then X is \mathbf{k} -rational.*

Proof. Let $x \in X(\mathbf{k})$ be a \mathbf{k} -point. First, assume that $F_1(X, x) = \emptyset$. Then by Theorem 5.1(i) the variety X is birational to a smooth quintic del Pezzo threefold X^+ . But X^+ is \mathbf{k} -rational by [KP19, Theorem 3.3], hence so is X .

Now assume that $F_1(X, x) \neq \emptyset$. Then by Theorem 5.1(ii) the variety X is birational to a smooth Fano threefold X^+ of type $X_{(2,2)}$. Moreover, by Nishimura lemma we have $X^+(\mathbf{k}) \neq \emptyset$. Therefore, X^+ is \mathbf{k} -rational by Proposition 4.1, hence so is X . \square

6. FANO THREEFOLDS OF TYPE $X_{(3,3)}$

In this section we prove that a Fano threefold X of type $X_{(3,3)}$ is \mathbf{k} -unirational if $X(\mathbf{k}) \neq \emptyset$, but not \mathbf{k} -rational if $\rho(X) = 1$.

6.1. The discriminant curve. Let X be a Fano threefold of type $X_{(3,3)}$ with $X(\mathbf{k}) \neq \emptyset$. Recall from Lemma 2.5 that the image G_X of the Galois group $G(\bar{\mathbf{k}}/\mathbf{k})$ in $\text{Aut}(\text{Pic}(X_{\bar{\mathbf{k}}}))$ is a group of order 2 swapping the generators H_1 and H_2 of $\text{Pic}(X_{\bar{\mathbf{k}}})$. The homomorphism $G(\bar{\mathbf{k}}/\mathbf{k}) \rightarrow G_X$ therefore defines a quadratic extension \mathbf{k}'/\mathbf{k} such that H_1 and H_2 are defined on $X_{\mathbf{k}'}$, hence

$$X_{\mathbf{k}'} \cong \left(\mathbb{P}(V_1) \times \mathbb{P}(V_2) \right) \cap \mathbb{P}(A^\perp),$$

where V_i are \mathbf{k}' -vector spaces of dimension 4 and $A \subset V_1^\vee \otimes V_2^\vee$ is the 3-dimensional subspace of linear equations of $X_{\mathbf{k}'}$. Note that the \mathbf{k}' -spaces $V_1 \otimes V_2$ and A are defined over \mathbf{k} , as well as the inclusion $A \subset V_1^\vee \otimes V_2^\vee$. We think of vectors $a \in A$ as of bilinear forms on $V_1 \otimes V_2$ and denote by

$$\Gamma \xrightarrow{\alpha} \mathbb{P}(A)$$

the discriminant curve parameterizing degenerate bilinear forms; it is also defined over \mathbf{k} .

Lemma 6.1. *The curve Γ is a smooth plane quartic curve; in particular it is a non-hyperelliptic curve of genus 3.*

Proof. The discriminant divisor in $\mathbb{P}(V_1^\vee \otimes V_2^\vee)$ is a quartic hypersurface, hence Γ is a quartic curve or the entire plane. To prove that Γ is a smooth curve we can work over \mathbf{k}' , and it is enough to show that the tangent space to Γ at any point is 1-dimensional. Assume to the contrary that the tangent space at a point $[a] \in \mathbb{P}(A)$ is 2-dimensional; then

- (i) either the bilinear form $a(-, -) \in V_1^\vee \otimes V_2^\vee$ has corank at least 2,
- (ii) or a has corank 1 and if the vectors $v_1 \in V_1$ and $v_2 \in V_2$ generate its left and right kernels, respectively, the point $[(v_1, v_2)] \in \mathbb{P}(V_1) \times \mathbb{P}(V_2)$ belongs to $X_{\mathbf{k}'}$.

In case (i), if $K_1 \subset V_1$ and $K_2 \subset V_2$ are the left and right kernels of a (they have dimension ≥ 2 by assumption), the form a vanishes on $\mathbb{P}(K_1) \times \mathbb{P}(K_2)$, hence the intersection $\left(\mathbb{P}(K_1) \times \mathbb{P}(K_2) \right) \cap X_{\mathbf{k}'}$ is a codimension-2 linear section of $\mathbb{P}(K_1) \times \mathbb{P}(K_2)$, hence it is non-empty. Therefore, in case (i), similarly to the case (ii), there is a point $[(v_1, v_2)] \in X_{\mathbf{k}'}$ such that v_1 and v_2 belong to the left and right kernels of some a . Then the hyperplane section of $\mathbb{P}(V_1) \times \mathbb{P}(V_2)$ by the hyperplane corresponding to a is singular at $[(v_1, v_2)]$, hence $X_{\mathbf{k}'}$ is also singular at this point. \square

As explained in Lemma 2.4(iii) the projections $\pi_1: X_{\mathbf{k}'} \rightarrow \mathbb{P}(V_1)$ and $\pi_2: X_{\mathbf{k}'} \rightarrow \mathbb{P}(V_2)$ defined over \mathbf{k}' , but not over \mathbf{k} , are the blowups of curves $\Gamma_i \subset \mathbb{P}(V_i)$ of genus 3 and degree 6 also defined over \mathbf{k}' . The next lemma relates the curves Γ_i to the discriminant curve Γ .

Lemma 6.2. *There is a natural isomorphism $\Gamma_i \cong \Gamma_{\mathbf{k}'}$ of curves over \mathbf{k}' .*

Proof. The fiber of the projection π_1 over a point $[v_1] \in \mathbb{P}(V_1)$ is the intersection of the orthogonals of v_1 with respect to all bilinear forms $a \in A$. Therefore, it has positive dimension if and only if v_1 belongs to the left kernel of one of the forms a . This means that the morphism

$$\gamma_i: \Gamma_{k'} \longrightarrow \mathbb{P}(V_i), \quad a \longmapsto \text{Ker}_i(a),$$

where Ker_1 and Ker_2 denote the left and right kernels of the bilinear form a , respectively, is an isomorphism $\Gamma_{k'} \rightarrow \Gamma_i$. \square

Remark 6.3. It is also easy to check that if $H_i|_\Gamma$ are the pullbacks of the hyperplane classes of $\Gamma_i \subset \mathbb{P}(V_i)$ to $\Gamma_{k'}$ under the isomorphism of Lemma 6.2 then $H_1|_\Gamma + H_2|_\Gamma = 3K_\Gamma$ and that the divisor classes $H_i|_\Gamma - K_\Gamma$ are non-effective and swapped by the $G(k'/k)$ -action. Conversely, given two such classes on a curve $\Gamma_{k'}$ one can reconstruct the variety X .

We denote by $E_i \subset X$ the exceptional divisors of the projections π_i . As explained in §2.2 every line on X is a fiber of one the projections $E_i \rightarrow \Gamma_i$ induced by π_1 and π_2 , respectively. The Galois group G_X swaps the E_i , while their sum stays Galois-invariant, hence defined over k .

6.2. The double projection from a point. Recall the quadratic extension k'/k defined in §6.1. Recall also the canonical embedding $X \subset Y$, where Y is a k -form of $\mathbb{P}^3 \times \mathbb{P}^3$. We consider the birational transformation of Theorem 3.1 for the variety $Y_{k'} = \mathbb{P}(V_1) \times \mathbb{P}(V_2)$ associated with a k -point

$$x_0 = (v_1, v_2) \in X \subset Y.$$

As in 3.1 we denote $\bar{V}_i := V_i/k'v_i$ and choose a splitting $V_i = k'v_i \oplus \bar{V}_i$. The transformation in this case looks as follows

$$(6.2.1) \quad \begin{array}{ccc} \tilde{Y} & \xrightarrow{\psi} & \tilde{Y}^+ \\ \sigma \downarrow & & \downarrow \sigma_+ \\ Y & & Y^+ \end{array}$$

where σ is the blowup of x_0 , $Y_{k'}^+ \cong \mathbb{P}(\bar{V}_1) \times \mathbb{P}(\bar{V}_2)$, σ_+ is the projectivization of the vector bundle

$$(6.2.2) \quad \mathcal{E} = \mathcal{O}(-h_1 - h_2) \oplus \mathcal{O}(-h_1) \oplus \mathcal{O}(-h_2)$$

(here h_i stand for the hyperplane classes of $\mathbb{P}(\bar{V}_i)$) over $Y_{k'}^+$ and the map ψ is a small birational isomorphism. Note that all varieties and maps in (6.2.1) are defined over k .

Recall also the relations (3.1.9) in $\text{Pic}(\tilde{Y}_{k'}) = \text{Pic}(\tilde{Y}_{k'}^+)$ between the hyperplane classes H_i of the factors $\mathbb{P}(V_i)$ of $Y_{k'}$, the class E of the exceptional divisor of σ , the hyperplane classes h_i , and the relative hyperplane class h of $\tilde{Y}_{k'}^+ = \mathbb{P}_{Y_{k'}^+}(\mathcal{E})$:

$$(6.2.3) \quad \begin{cases} h_1 = H_1 - E, \\ h_2 = H_2 - E, \\ h = H_1 + H_2 - E \end{cases} \quad \begin{cases} H_1 = h - h_2, \\ H_2 = h - h_1, \\ E = h - h_1 - h_2. \end{cases}$$

Since X is a smooth linear section of Y , containing the point x_0 , it is a complete intersection of three divisors D_α , $1 \leq \alpha \leq 3$, in the linear system $|H_1 + H_2|$ whose strict transforms on \tilde{Y} belong

to the linear system $|H_1 + H_2 - E|$. Now it follows from (6.2.3) that their strict transforms \tilde{D}_α^+ on \tilde{Y}^+ belong to the linear system $|h|$. As in Proposition 3.5 we consider the complete intersection

$$\tilde{X}_{k'}^+ := \tilde{D}_1^+ \cap \tilde{D}_2^+ \cap \tilde{D}_3^+ \subset \mathbb{P}_{Y_{k'}^+}(\mathcal{E}).$$

It follows that $\tilde{X}_{k'}^+$ is determined by a morphism of vector bundles

$$\xi: \mathcal{E} \longrightarrow A^\vee \otimes \mathcal{O},$$

and if we choose a basis a_1, a_2, a_3 in A , it is easy to see that ξ is given by the matrix

$$(6.2.4) \quad \xi = \begin{pmatrix} \bar{a}_1(-, -) & a_1(-, v_2) & a_1(v_1, -) \\ \bar{a}_2(-, -) & a_2(-, v_2) & a_2(v_1, -) \\ \bar{a}_3(-, -) & a_3(-, v_2) & a_3(v_1, -) \end{pmatrix},$$

where \bar{a}_i denotes the restriction of the bilinear form a_i to $\bar{V}_1 \otimes \bar{V}_2$, while $a_i(-, v_2)$ and $a_i(v_1, -)$ are considered as linear functions on \bar{V}_1 and \bar{V}_2 , respectively.

Proposition 6.4. *The threefold X is \mathbf{k} -birational to a \mathbf{k} -form \tilde{X}^+ of the threefold $\tilde{X}_{k'}^+$ defined by (6.2.4) and to a \mathbf{k} -form X^+ of its image in Y^+*

$$X_{k'}^+ = \sigma_+(\tilde{X}_{k'}^+) = \{\det(\xi) = 0\} \subset \mathbb{P}(\bar{V}_1) \times \mathbb{P}(\bar{V}_2),$$

which is a geometrically irreducible divisor of bidegree $(2, 2)$. Moreover,

- if $F_1(X, x_0) = \emptyset$ then $\tilde{X}^+ \cong \tilde{X} = \text{Bl}_x(X)$ is smooth, the morphism $\sigma_+: \tilde{X}^+ \rightarrow X^+$ is induced by the double projection from x_0 , and it is a small resolution of singularities;
- if $F_1(X, x_0) \neq \emptyset$ then X^+ contains a \mathbf{k} -form of a quadric surface $\mathbb{P}^1 \times \mathbb{P}^1 \subset \mathbb{P}(\bar{V}_1) \times \mathbb{P}(\bar{V}_2)$.

Proof. To prove birationality of \tilde{X} and the \mathbf{k} -form $\tilde{X}^+ = \psi_*(\tilde{X})$ we apply Proposition 3.5, so we need to verify the dimension conditions (3.2.1). We have $(Y_1)_{k'} = [v_1] \times \mathbb{P}(V_2)$, hence

$$X_{k'} \cap (Y_1)_{k'} = ([v_1] \times \mathbb{P}(V_2)) \cap \mathbb{P}(A^\perp)$$

is a fiber of the projection $\pi_1: X_{k'} \rightarrow \mathbb{P}(V_1)$. By Lemma 2.4 it is a point or a line. A similar argument for $X_{k'} \cap (Y_2)_{k'}$ shows that the first part of (3.2.1) holds. Moreover, the same argument shows that in the case $F_1(X, x_0) = \emptyset$, the blowup $\tilde{X}_{k'}$ of $X_{k'}$ has empty intersection with the indeterminacy locus $(\tilde{Y}_1)_{k'} \sqcup (\tilde{Y}_2)_{k'}$ of the map ψ .

On the other hand, the subbundle $\bar{\mathcal{E}}_1 \subset \mathcal{E}$ is just the summand $\mathcal{O}(-h_2)$, hence the corresponding intersection $\tilde{X}_{k'}^+ \cap \mathbb{P}_{Y_{k'}^+}(\bar{\mathcal{E}}_1)$ is the zero locus of the morphism

$$\xi_2: \mathcal{O}(-h_2) \longmapsto A^\vee \otimes \mathcal{O}$$

given by the last column of (6.2.4). It is easy to see that this is empty, if $F_1(X, x_0) = \emptyset$, or isomorphic to a line otherwise. A similar argument works for $\tilde{X}_{k'}^+ \cap \mathbb{P}_{Y_{k'}^+}(\bar{\mathcal{E}}_2)$; therefore, the second part of (3.2.1) also holds. This proves that $\tilde{X}^+ = \psi(\tilde{X})$ is a \mathbf{k} -form of $\tilde{X}_{k'}^+$, which is \mathbf{k} -birational to X , and if $F_1(X, x_0) = \emptyset$, it is isomorphic to \tilde{X} , and in particular in this case it is smooth.

By definition, $\tilde{X}_{k'}^+$ parameterizes points in the projectivization of kernel spaces of ξ ; therefore, its image in $Y_{k'}^+ = \mathbb{P}(\bar{V}_1) \times \mathbb{P}(\bar{V}_2)$ is the degeneracy locus $X_{k'}^+$ of ξ given by the equation $\det(\xi) = 0$. Since $\det(\mathcal{E}) \cong \mathcal{O}(-2h_1 - 2h_2)$ by (6.2.2), this is a divisor of bidegree $(2, 2)$, which is geometrically irreducible because $\tilde{X}_{k'}^+$ is. Moreover, the fibers of the morphism $\sigma_+: \tilde{X}_{k'}^+ \rightarrow X_{k'}^+$ are linear spaces, so since both the source and the target are 3-dimensional, the morphism is birational.

Now assume $F_1(X, x_0) = \emptyset$. In this case the pullback along σ_+ of the ample class $h_1 + h_2$ on $\mathbb{P}(\bar{V}_1) \times \mathbb{P}(\bar{V}_2)$ by (6.2.3) equals to $H_1 + H_2 - 2E$, the anticanonical class of $\tilde{X}^+ \cong \tilde{X}$, hence the morphism σ_+ is the double projection from a point. Consequently, it is small by the argument of [KP19, Theorem 5.18] — by [KP19, Lemma 5.4(iii)] we have $\dim |H_1 + H_2 - 3E| \geq g - 9 = 2$ (recall that $g = g(X) = 11$, see Table 1), hence by [KP19, Lemma 5.6(ii)] any divisor D contracted by σ_+ must be a fixed component of $|H_1 + H_2 - 3E|$, and at the same time by [KP19, (5.1.8)] its class should be a multiple of $H_1 + H_2 - 5E$, and these two conclusions are incompatible.

Finally, assume that $F_1(X, x_0) \neq \emptyset$. As it was explained in Lemma 6.2 this means that (for appropriate k' -basis in A) we have

$$a_1(v_1, -) = 0 \quad \text{and} \quad a_2(-, v_2) = 0$$

as linear functions on \bar{V}_2 and \bar{V}_1 , respectively; moreover, a_1 and a_2 as above are unique and swapped by the Galois action. Consider the surface

$$\{a_1(-, v_2) = 0, a_2(v_1, -) = 0\} \subset \mathbb{P}(\bar{V}_1) \times \mathbb{P}(\bar{V}_2) = Y_{k'}^+.$$

(isomorphic to $\mathbb{P}^1 \times \mathbb{P}^1$). The equations, defining it are Galois-conjugate, hence it comes from a k -surface in Y^+ (geometrically, this surface is the image of the exceptional divisor E of σ). It is clear from (6.2.4) that this surface is contained in the degeneracy locus X^+ of ξ . \square

6.3. A conic bundle structure. In this section we work under the assumption $F_1(X, x_0) = \emptyset$ and show that in this case X admits a nice conic bundle structure.

We will need a general result about what we call **Springer resolutions**. Let M be a variety and let $\xi: \mathcal{E}_1 \rightarrow \mathcal{E}_2^\vee$ and $\xi^\vee: \mathcal{E}_2 \rightarrow \mathcal{E}_1^\vee$ be a morphism of vector bundles on M of the same rank and its dual morphism. Assume the degeneracy locus $Z \subset M$ of ξ is a geometrically integral divisor. Let $Z_1 \subset \mathbb{P}_M(\mathcal{E}_1)$ and $Z_2 \subset \mathbb{P}_M(\mathcal{E}_2)$ be the zero loci of the morphisms

$$\mathcal{O}(-h_{\mathcal{E}_1}) \hookrightarrow p_1^* \mathcal{E}_1 \xrightarrow{p_1^* \xi} p_1^* \mathcal{E}_2^\vee \quad \text{and} \quad \mathcal{O}(-h_{\mathcal{E}_2}) \hookrightarrow p_2^* \mathcal{E}_2 \xrightarrow{p_2^* \xi^\vee} p_2^* \mathcal{E}_1^\vee,$$

where $p_i: \mathbb{P}_M(\mathcal{E}_i) \rightarrow M$ are the projections, $h_{\mathcal{E}_i}$ are their relative hyperplane classes, and the first arrows are the tautological embeddings.

Lemma 6.5. *If one of the morphisms*

$$p_1|_{Z_1}: Z_1 \longrightarrow Z \quad \text{or} \quad p_2|_{Z_2}: Z_2 \longrightarrow Z$$

is birational then so is the other. Moreover, if one of them is small then so is the other, and there is an equality

$$(6.3.1) \quad (h_{\mathcal{E}_1} + c_1(p_1^* \mathcal{E}_1)) + (h_{\mathcal{E}_2} + c_1(p_2^* \mathcal{E}_2)) = 0$$

in the group $\text{Cl}(Z_1) \cong \text{Cl}(Z_2)$, where the isomorphism of the class groups is induced by the birational isomorphisms.

Proof. Let $Z^{\geq c} \subset Z$ be the locus of points where the corank of ξ is at least c (so that $Z = Z^{\geq 1}$). Then the morphisms $p_i|_{Z_i}$ are \mathbb{P}^{c-1} -fibrations over $Z^{\geq c} \setminus Z^{\geq c+1}$. In particular if one of the morphisms is birational then $\dim Z^{\geq c} \leq \dim Z - c$ for $c \geq 2$ and then the other morphism is also birational. Similarly, if one of the morphisms is small then $\dim Z^{\geq c} \leq \dim Z - c - 1$ for $c \geq 2$ and then the other morphism is also small. Finally, assuming that the morphisms are small, we have

$$\text{Cl}(Z_1) = \text{Cl}(Z_1 \setminus p_1^{-1}(Z^{\geq 2})) = \text{Cl}(Z \setminus Z^{\geq 2}) = \text{Cl}(Z_2 \setminus p_2^{-1}(Z^{\geq 2})) = \text{Cl}(Z_2),$$

and when restricted to $Z \setminus Z^{\geq 2}$ the morphism ξ has constant corank 1, the summands in (6.3.1) are equal to $c_1(\text{Im}(\xi))$ and $c_1(\text{Im}(\xi^\vee))$, respectively, and (6.3.1) follows from the natural duality isomorphism $\text{Im}(\xi^\vee) \cong \text{Im}(\xi)^\vee$. \square

Now, coming back to the threefold X of type $X_{(3,3)}$ and assuming $F_1(X, x_0) = \emptyset$ we note that $X^+ \subset Y^+$ is the degeneracy locus of $\xi: \mathcal{E} \rightarrow A^\vee \otimes \mathcal{O}$ and that $\tilde{X}^+ \subset \mathbb{P}_{Y^+}(\mathcal{E})$ is one of its Springer resolutions. Consider the other Springer resolution

$$\tilde{X}^{++} \subset \mathbb{P}_{Y^+}(A \otimes \mathcal{O}) \cong Y^+ \times \mathbb{P}(A),$$

which by definition is the zero locus of the morphism

$$\mathcal{O}(-h_A) \hookrightarrow A \otimes \mathcal{O} \xrightarrow{\xi^\vee} \mathcal{E}^\vee,$$

where h_A is the hyperplane class of $\mathbb{P}(A)$ and we suppress the pullback in the notation. In view of (6.2.2) over k' this is just a complete intersection of divisors of types $h_1 + h_2 + h_A$, $h_1 + h_A$, and $h_2 + h_A$ in $\mathbb{P}(\bar{V}_1) \times \mathbb{P}(\bar{V}_2) \times \mathbb{P}(A)$ that correspond to the columns of (6.2.4).

Proposition 6.6. *If $F_1(X, x_0) = \emptyset$ there is a commutative diagram defined over k*

$$(6.3.2) \quad \begin{array}{ccccc} \tilde{X} & \xrightarrow{\psi} & \tilde{X}^+ & \overset{\psi_+}{\dashrightarrow} & \tilde{X}^{++} \\ \sigma \downarrow & & \searrow \sigma_+ & & \swarrow \sigma_{++} \\ X & & X^+ & & \mathbb{P}(A), \end{array} \quad \begin{array}{c} \downarrow f \\ \mathbb{P}(A), \end{array}$$

where \tilde{X}^+ and \tilde{X}^{++} are the Springer resolutions of the degeneracy locus $X^+ \subset Y^+$ of ξ , the morphisms σ_+ and σ_{++} are small birational contractions, and $\psi_+ = \sigma_{++}^{-1} \circ \sigma_+$ is a flop.

Moreover, \tilde{X}^{++} is smooth and f is a flat conic bundle with discriminant curve Γ defined in §6.1.

Proof. The morphism σ_+ is small by Proposition 6.4, hence σ_{++} is small by Lemma 6.5; moreover, it follows that both morphisms are crepant. Now, the relation (6.3.1) implies that the σ_+ -antiample class $-h$ is σ_{++} -ample, hence ψ_+ is a flop. Since $\tilde{X}^+ \cong \tilde{X}$ is smooth and ψ_+ is a flop, \tilde{X}^{++} is smooth as well, see [Kol89, Theorem 2.4].

It remains to show that f is a conic bundle and compute its discriminant. For this note that by definition the fiber of f over a point $[a] \in \mathbb{P}(A)$ is given in $\mathbb{P}(\bar{V}_1) \times \mathbb{P}(\bar{V}_2)$ by the equations

$$a(-, v_2) = a(v_1, -) = \bar{a}(-, -) = 0.$$

The first is a linear function on \bar{V}_1 , the second is a linear function on \bar{V}_2 , and both are non-zero because $F_1(X, x_0) = \emptyset$, so their common zero locus is $\mathbb{P}^1 \times \mathbb{P}^1 \subset \mathbb{P}(\bar{V}_1) \times \mathbb{P}(\bar{V}_2)$. The last equation $\bar{a}(-, -) = 0$ cuts a divisor of bidegree $(1, 1)$ on this $\mathbb{P}^1 \times \mathbb{P}^1$, i.e., a conic; and if $\bar{a}(-, -)$ vanishes identically, then the corresponding bilinear form a vanishes on $\mathbb{P}^2 \times \mathbb{P}^2 \subset \mathbb{P}(V_1) \times \mathbb{P}(V_2)$, hence has corank 2, which is impossible by the argument of Lemma 6.1 as X is smooth. This shows that f is a flat conic bundle. Finally, note that if $\bar{v}'_i, \bar{v}''_i, \bar{v}'''_i$ are bases of vector spaces \bar{V}_i such that $a(v_1, \bar{v}'_2) = a(v_1, \bar{v}''_2) = a(\bar{v}'_1, v_2) = a(\bar{v}''_1, v_2) = 0$ then the matrix of a has the form

$$\begin{pmatrix} 0 & 0 & 0 & a(v_1, \bar{v}'''_2) \\ 0 & \bar{a}(\bar{v}'_1, \bar{v}'_2) & \bar{a}(\bar{v}'_1, \bar{v}''_2) & * \\ 0 & \bar{a}(\bar{v}''_1, \bar{v}'_2) & \bar{a}(\bar{v}''_1, \bar{v}''_2) & * \\ a(\bar{v}'''_1, v_2) & * & * & * \end{pmatrix}$$

with non-zero $a(v_1, \bar{v}_2''')$ and $a(\bar{v}_1''', v_2)$, and with the 2-by-2 matrix in the middle giving the equation of the conic $f^{-1}(a)$ in $\mathbb{P}^1 \times \mathbb{P}^1$. Therefore, the conic is singular if and only if $\det(a) = 0$, i.e., if and only if $[a] \in \Gamma$. Thus, the discriminant curve of f equals Γ . \square

If $f: \mathcal{X} \rightarrow S$ is a flat conic bundle over a surface S (not necessarily proper) with a smooth discriminant curve $\Delta \subset S$, consider the preimage $\mathcal{X}_\Delta := f^{-1}(\Delta)$, its normalization $\mathcal{X}_\Delta^\nu \rightarrow \mathcal{X}_\Delta$, and the Stein factorization

$$\mathcal{X}_\Delta^\nu \longrightarrow \tilde{\Delta} \longrightarrow \Delta.$$

Then the first arrow is a \mathbb{P}^1 -bundle and the second arrow is an étale double covering (because Δ was assumed to be smooth). We will say that the étale covering $\tilde{\Delta} \rightarrow \Delta$ is the **discriminant double covering** of the conic bundle f .

Lemma 6.7. *The discriminant double covering of the conic bundle $f: \tilde{X}^{++} \rightarrow \mathbb{P}(A)$ has the form*

$$\tilde{\Gamma} \cong \Gamma \times_{\mathbf{k}} \mathbf{k}' \longrightarrow \Gamma.$$

Proof. If the conic $f^{-1}([a])$ is singular, it is a union of two components that correspond to the two factors $\mathbb{P}(\bar{V}_i)$ in $Y_{\mathbf{k}'}^+$ and each of them is contracted by appropriate projection $Y_{\mathbf{k}'}^+ \rightarrow \mathbb{P}(\bar{V}_i)$. Therefore, the discriminant double covering $\tilde{\Gamma} \rightarrow \Gamma$ becomes trivial after the extension of scalars to \mathbf{k}' , while it is non-trivial over \mathbf{k} , hence the claim. \square

6.4. Unirationality. In this section we prove unirationality of X assuming that $X(\mathbf{k}) \neq \emptyset$. We start with the following observation which might be useful in other situations.

Lemma 6.8. *Let Y be a \mathbf{k} -form of $\mathbb{P}^2 \times \mathbb{P}^2$ and let $W \subset Y$ be a \mathbf{k} -form of $\mathbb{P}^1 \times \mathbb{P}^1 \subset \mathbb{P}^2 \times \mathbb{P}^2$. Any geometrically irreducible divisor $Z \subset Y$ of bidegree $(2, 2)$ such that $W \subset Z$ and $Z(\mathbf{k}) \neq \emptyset$ is \mathbf{k} -unirational.*

Proof. Consider the toric birational isomorphism

$$\begin{array}{ccc} \text{Bl}_{\mathbb{P}^1 \times \mathbb{P}^1}(\mathbb{P}^2 \times \mathbb{P}^2) & \xlongequal{\quad} & \text{Bl}_{\mathbb{P}^1 \sqcup \mathbb{P}^1}(\mathbb{P}^4) \\ \swarrow & & \searrow \\ \mathbb{P}^2 \times \mathbb{P}^2 & \xrightarrow{\quad \chi \quad} & \mathbb{P}^4. \end{array}$$

analogous to the birational transformation of Theorem 3.1 (see also [Zak07, Proposition 3]). Here the map χ is the projection from the linear span of W under the Segre embedding $\mathbb{P}^2 \times \mathbb{P}^2 \subset \mathbb{P}^8$, and the right arrow is the blowup of two skew lines in \mathbb{P}^4 swapped by the action of $G(\mathbf{k}'/\mathbf{k})$. Denoting by \hat{e} the class of the exceptional divisor of the left blowup, by \hat{h} the hyperplane class of \mathbb{P}^4 and by e_1 and e_2 the classes of the exceptional divisors of the right blowup, it is easy to check that we have the relations

$$\begin{cases} \hat{h} = h_1 + h_2 - \hat{e}, \\ e_1 = h_1 - \hat{e}, \\ e_2 = h_2 - \hat{e}, \end{cases} \quad \text{and} \quad \begin{cases} h_1 = \hat{h} - e_2, \\ h_2 = \hat{h} - e_1, \\ \hat{e} = \hat{h} - e_1 - e_2. \end{cases}$$

In particular, the map χ is given by the linear system $|h_1 + h_2 - \hat{e}|$, hence it is defined over \mathbf{k} . Furthermore, we have $2h_1 + 2h_2 - \hat{e} = 3\hat{h} - e_1 - e_2$, hence the strict transform of Z under the map χ is a cubic threefold $\hat{Z} \subset \mathbb{P}^4$ passing through the pair of skew lines $\mathbb{P}^1 \sqcup \mathbb{P}^1$. Moreover, we

have $\hat{e} = \hat{h} - e_1 - e_2$, hence the image of \hat{E} is the hyperplane section of this cubic threefold (by the linear span of these lines). Finally, we have $\hat{Z}(\mathbf{k}) \neq \emptyset$ by Nishimura lemma.

Now, if \hat{Z} is not a cone, it is \mathbf{k} -unirational by Kollár's theorem [Kol02]. Otherwise, if \hat{Z} is a cone and its vertex lies away from the hyperplane spanned by the two skew lines, then the base of the cone is a cubic surface containing the pair of skew lines swapped by the $G(\mathbf{k}'/\mathbf{k})$ -action. This surface is birational to the product of these lines, which is a quadrics surface with a point, hence it is \mathbf{k} -rational. Therefore, the cone \hat{Z} is also \mathbf{k} -rational. Finally, if the vertex of the cone lies on \hat{E} , then \hat{E} itself must be a cubic cone in \mathbb{P}^3 , and since it also contains two skew lines, it is not geometrically irreducible, which is absurd. \square

Now we can deduce unirationality of X .

Proposition 6.9. *If X is a Fano threefold of type $X_{(3,3)}$ with $X(\mathbf{k}) \neq \emptyset$ then X is \mathbf{k} -unirational.*

Proof. Let x_0 be a \mathbf{k} -point on X .

First, assume $F_1(X, x_0) = \emptyset$. By Proposition 6.6 we have a \mathbf{k} -birational isomorphism $X \sim_{\text{bir}} \tilde{X}^{++}$, where $f: \tilde{X}^{++} \rightarrow \mathbb{P}(A)$ is a conic bundle. Moreover, the \mathbf{k} -rational surface $E \cong \mathbb{P}(T_{x_0}X) \subset \tilde{X}$ dominates the base of this conic bundle. Therefore, \tilde{X} is \mathbf{k} -unirational (and hence so is X).

Now assume that $F_1(X, x_0) \neq \emptyset$. By Proposition 6.4 we have a birational isomorphism $X \sim_{\text{bir}} X^+$ where X^+ is a geometrically irreducible divisor of bidegree $(2, 2)$ in a \mathbf{k} -form of $\mathbb{P}^2 \times \mathbb{P}^2$ that contains a \mathbf{k} -form of a quadric surface $\mathbb{P}^1 \times \mathbb{P}^1$. Moreover, by Nishimura lemma we have $X^+(\mathbf{k}) \neq \emptyset$. Therefore, X^+ is \mathbf{k} -unirational by Lemma 6.8, hence so is X . \square

6.5. Non-rationality. In this section we prove non-rationality of Fano threefolds of type $X_{(3,3)}$. We will use the following reformulation of a result of Benoist–Wittenberg from [BW20].

Theorem 6.10. *Let $\mathcal{X} \rightarrow S$ be a flat conic bundle over a smooth \mathbf{k} -rational surface S with smooth connected discriminant curve $\Delta \subset S$. Assume the discriminant double covering takes the form*

$$\tilde{\Delta} \cong \Delta \times_{\mathbf{k}} \mathbf{k}' \longrightarrow \Delta$$

where \mathbf{k}'/\mathbf{k} is a quadratic extension of the base field. If the conic bundle $\mathcal{X}_{\mathbf{k}'} \rightarrow S_{\mathbf{k}'}$ admits a rational section and the curve Δ is not hyperelliptic then \mathcal{X} is not \mathbf{k} -rational.

Note that we do not require neither the surface S nor the curve Δ to be proper; moreover, during the proof we will further shrink S but keep (the generic point of) the curve Δ in S .

Proof. Since S is normal and f is proper any rational section of f extends to codimension 1 points, hence defines a regular section over the complement of a finite subscheme of S . Moreover, over the complement of this finite subscheme the section does not pass through singular points of fibers of f , hence it defines a section of the morphism $\mathcal{X}_{\Delta}^{\nu} \rightarrow \Delta$, where recall that $\mathcal{X}_{\Delta}^{\nu}$ is the normalization of $\mathcal{X}_{\Delta} = f^{-1}(\Delta)$. Therefore it also gives a section of the discriminant double covering $\tilde{\Delta} \rightarrow \Delta$. This contradicts to the isomorphism $\tilde{\Delta} \cong \Delta \times_{\mathbf{k}} \mathbf{k}'$, hence the morphism f has no rational sections defined over \mathbf{k} .

Now consider a rational section of $f: \mathcal{X}_{\mathbf{k}'} \rightarrow S_{\mathbf{k}'}$. Removing if necessary a finite subscheme from S we may assume that this section is regular. Its intersection with the conjugate (with respect to the $G(\mathbf{k}'/\mathbf{k})$ -action) section projects to a curve in S which is disjoint from Δ (because

a regular section does not pass through singular points of fibers). So, shrinking S further we may assume that the section and its conjugate do not intersect. Then the union

$$Z \subset \mathcal{X}$$

of the section and its conjugate is a 2-section of f defined over k ; moreover, $Z \cong S \times_k k'$ and in particular Z is étale over S .

Consider the rank-3 and rank-2 bundles $\mathcal{V} := (f_*\omega_{\mathcal{X}}^{-1})^\vee$ and $\mathcal{V}_Z := (f_*\omega_{\mathcal{X}}^{-1}|_Z)^\vee$ on S . The restriction morphism $\omega_{\mathcal{X}}^{-1} \rightarrow \omega_{\mathcal{X}}^{-1}|_Z$ induces an embedding of vector bundles $\mathcal{V}_Z \hookrightarrow \mathcal{V}$ and a Cartesian square

$$\begin{array}{ccc} Z & \longrightarrow & \mathbb{P}_S(\mathcal{V}_Z) \\ \downarrow & & \downarrow \\ \mathcal{X} & \longrightarrow & \mathbb{P}_S(\mathcal{V}), \end{array}$$

where all arrows are the natural embeddings.

Shrinking the surface S again but keeping an open part of the curve Δ in it we may assume that the bundles \mathcal{V} and \mathcal{V}_Z are trivial and that the subvarieties $\mathcal{X} \subset \mathbb{P}_S(\mathcal{V})$ and $Z \subset \mathbb{P}_S(\mathcal{V}_Z)$ are given by a quadratic form $q \in \text{Sym}^2 \mathcal{V}^\vee$ and its restriction $q_Z \in \text{Sym}^2 \mathcal{V}_Z^\vee$ to \mathcal{V}_Z , respectively. Since Z is étale over S , the form q_Z is everywhere non-degenerate and takes the form

$$q_Z = x^2 - \alpha y^2,$$

where (x, y) are homogeneous coordinates in the fiber of $\mathbb{P}_S(\mathcal{V}_Z) \cong S \times \mathbb{P}^1$ and $\alpha \in k^\times$ is such that $k' = k(\sqrt{\alpha})$. Now, considering the orthogonal complement to \mathcal{V}_Z in \mathcal{V} we see that q takes the form

$$q = x^2 - \alpha y^2 - Fz^2,$$

where F is an equation of Δ on S . Thus, the conic bundle $\mathcal{X} \rightarrow S$ is birational to conic bundles considered in [BW20, §3.3.1], hence \mathcal{X} is not k -rational by [BW20, Proposition 3.4]. \square

Now we apply this to prove non-rationality of threefolds of type $X_{(3,3)}$.

Corollary 6.11. *If X is a Fano threefold of type $X_{(3,3)}$ then X is not k -rational.*

Proof. If X is not k -unirational, there is nothing to prove. So, assume X is k -unirational. Then there exists a k -point $x_0 \in X$ such that $F_1(X, x_0) = \emptyset$. Consider the conic bundle $\tilde{X}^{++} \rightarrow \mathbb{P}^2$ constructed in Proposition 6.6. By Lemma 6.1 the discriminant curve of f is the smooth non-hyperelliptic curve Γ defined in 6.1 and by Lemma 6.7 the discriminant double covering has the form $\tilde{\Gamma} \cong \Gamma \times_k k'$. Finally, the description of Proposition 6.6 shows that f admits a rational section after base change to k' . Therefore, Theorem 6.10 applies and proves that \tilde{X}^{++} is not k -rational, hence X is not k -rational as well. \square

7. FANO THREEFOLDS OF TYPE $X_{(1,1,1,1)}$

In this section we prove Theorem 1.4. Recall the definition (2.1.5) of the Galois group $G_X \subset \mathfrak{S}_4$ of X . Recall also that $V_4 \subset \mathfrak{S}_4$ denotes the Klein group. To prove that a very general X with $G_X \supset V_4$ is not stably rational, we use degeneration technique. Explicitly, we consider a singular toric degeneration of X .

7.1. Toric degeneration. To start with we consider $Y_0 = (\mathbb{P}^1)^4$ and denote by $(u_i : v_i)$ the homogeneous coordinates on the i -th factor. This is a toric variety with respect to the action of the split torus \mathbb{G}_m^4 that rescales the v_i . We also consider the action of \mathfrak{S}_4 on Y_0 that permutes the factors. It normalizes the torus action, and together they generate an action of the group $\mathbb{G}_m^4 \rtimes \mathfrak{S}_4$.

Lemma 7.1. *The subvariety*

$$X_0 := \{u_1 u_2 u_3 u_4 - v_1 v_2 v_3 v_4 = 0\} \subset Y_0$$

is a toric variety with 6 ordinary double points at the \mathfrak{S}_4 -orbit of the point $(0, 0, \infty, \infty) \in Y_0$.

Proof. In the affine chart $v_1 \neq 0, v_2 \neq 0, u_3 \neq 0, u_4 \neq 0$ we can set $v_1 = v_2 = u_3 = u_4 = 1$ and use u_1, u_2, v_3, v_4 as coordinates. Then the equation of X_0 takes the form

$$u_1 u_2 - v_3 v_4 = 0,$$

which means that the origin of the chart, i.e., the point $(0, 0, \infty, \infty)$ is an ordinary double point. Considering similarly the other charts, we see that the singular locus of X_0 is the \mathfrak{S}_4 -orbit of this point; in particular each singular point of X_0 is an ordinary double point. Moreover, we see that the hypersurface X_0 is normal and irreducible.

On the other hand, the intersection of X_0 with the torus $\mathbb{G}_m^4 \subset Y_0$ (identified with the set of points such that $u_i \neq 0$ and $v_i \neq 0$ for all i) is the subtorus

$$(7.1.1) \quad T_0 := \{(t_1, t_2, t_3, t_4) \in \mathbb{G}_m^4 \mid t_1 t_2 t_3 t_4 = 1\}.$$

This torus acts on X_0 , hence X_0 is toric with respect to the torus T_0 . □

Remark 7.2. Note that the toric variety X_0 contains the point

$$(7.1.2) \quad x_0 = (1, 1, 1, 1) \in Y_0$$

and is equal to the closure in Y_0 of the T_0 -orbit of x_0 .

Now we describe all varieties of type $X_{(1,1,1,1)}$ with a \mathbf{k} -point. For a field extension \mathbf{k}'/\mathbf{k} we denote by $\text{Res}_{\mathbf{k}'/\mathbf{k}}: \text{Sch}_{\mathbf{k}'} \rightarrow \text{Sch}_{\mathbf{k}}$ the Weil restriction of scalars functor, the right adjoint functor to the extension of scalars $-\otimes_{\mathbf{k}} \mathbf{k}': \text{Sch}_{\mathbf{k}} \rightarrow \text{Sch}_{\mathbf{k}'}$.

Proposition 7.3. *If X is a Fano threefold of type $X_{(1,1,1,1)}$ with $X(\mathbf{k}) \neq \emptyset$ then there is a quartic field extension \mathbf{k}'/\mathbf{k} and an embedding*

$$(7.1.3) \quad X \subset Y := \text{Res}_{\mathbf{k}'/\mathbf{k}}(\mathbb{P}_{\mathbf{k}'}^1)$$

such that $X_{\bar{\mathbf{k}}} \subset Y_{\bar{\mathbf{k}}} \cong \mathbb{P}_{\bar{\mathbf{k}}}^1 \times \mathbb{P}_{\bar{\mathbf{k}}}^1 \times \mathbb{P}_{\bar{\mathbf{k}}}^1 \times \mathbb{P}_{\bar{\mathbf{k}}}^1$ is a smooth hypersurface of multidegree $(1, 1, 1, 1)$ containing the point (7.1.2).

Proof. By definition of threefolds of type $X_{(1,1,1,1)}$, there is an embedding $X_{\bar{\mathbf{k}}} \subset \mathbb{P}_{\bar{\mathbf{k}}}^1 \times \mathbb{P}_{\bar{\mathbf{k}}}^1 \times \mathbb{P}_{\bar{\mathbf{k}}}^1 \times \mathbb{P}_{\bar{\mathbf{k}}}^1$ as a smooth hypersurface of multidegree $(1, 1, 1, 1)$. By Lemma 2.5 there is a homomorphism

$$G(\bar{\mathbf{k}}/\mathbf{k}) \twoheadrightarrow G_X \subset \mathfrak{S}_4$$

that corresponds to the action of $G(\bar{\mathbf{k}}/\mathbf{k})$ on the set $\{H_1, H_2, H_3, H_4\}$ of the generators of the nef cone of $X_{\bar{\mathbf{k}}}$. Consider the stabilizer of H_1 in the Galois group $G(\bar{\mathbf{k}}/\mathbf{k})$ (this is a subgroup of index 4) and the corresponding quartic field extension \mathbf{k}'/\mathbf{k} . Then the divisor class H_1 on $X_{\mathbf{k}'}$ is Galois invariant, and since $X(\mathbf{k}) \neq \emptyset$, we have $X_{\mathbf{k}'}(\mathbf{k}') \neq \emptyset$, hence H_1 is defined over \mathbf{k}' . Moreover,

it defines a morphism $p_1: X_{k'} \rightarrow \mathbb{P}_{k'}^1$, and if we fix a k -point $x \in X$, we can choose coordinates on $\mathbb{P}_{k'}^1$ in such a way that

$$(7.1.4) \quad p_1(x) = 1 \in \mathbb{P}_{k'}^1.$$

By adjunction, we obtain a morphism $X \rightarrow Y := \text{Res}_{k'/k}(\mathbb{P}_{k'}^1)$. Over the field \bar{k} , where all classes H_i are defined, it becomes the morphism

$$X_{\bar{k}} \rightarrow (\text{Res}_{k'/k}(\mathbb{P}_{k'}^1))_{\bar{k}} \cong (\mathbb{P}_{\bar{k}}^1)^4$$

induced by the four projections $p_i: X_{\bar{k}} \rightarrow \mathbb{P}_{\bar{k}}^1$, defined by the H_i , hence it is the natural embedding. Moreover, by definition of the Weil restriction we have $Y(k) = \mathbb{P}_{k'}^1(k')$, so if $x_0 \in Y(k)$ is the k -point corresponding to the k' -point (7.1.4), then $x_0 \in X$. Finally, the above isomorphism $(\text{Res}_{k'/k}(\mathbb{P}_{k'}^1))_{\bar{k}} \cong (\mathbb{P}_{\bar{k}}^1)^4$ can be chosen in such a way, that the \bar{k} -point corresponding to the k -point x_0 is the point (7.1.2). \square

Now we fix the quartic field extension k'/k and the embedding $X \subset Y := \text{Res}_{k'/k}(\mathbb{P}_{k'}^1)$ defined in Proposition 7.3 so that X contains the point (7.1.2). We denote by $H = H_1 + H_2 + H_3 + H_4$ the Segre class of Y . Note that H is defined over k . Let

$$\mathfrak{P} := |H - x_0| \cong \mathbb{P}^{14}$$

be the linear system of all hyperplane sections of Y containing the point (7.1.2). The open subset

$$\mathfrak{P}_0 \subset \mathfrak{P}$$

corresponding to smooth hyperplane sections is a parameter space for Fano threefolds X of type $X_{(1,1,1,1)}$ with a k -point. In other words, for each X with $X(k) \neq \emptyset$ there is a k -point of \mathfrak{P}_0 such that X is isomorphic to the corresponding hyperplane section of Y , and conversely, each k -point of \mathfrak{P}_0 corresponds to a Fano threefold X of type $X_{(1,1,1,1)}$ with $X(k) \neq \emptyset$.

Now we can prove the last result of this paper.

Proof of Theorem 1.4. The above discussion shows that it is enough to check that a very general k -point of \mathfrak{P}_0 corresponds to a Fano threefold which is not stably rational. For this we use the results [NS19] of Nicaise and Shinder on degeneration of stable rationality.

First, we note that there is a point in \mathfrak{P} such that the corresponding hyperplane section X of Y is isomorphic over \bar{k} to the toric variety X_0 of Lemma 7.1. Indeed, the \mathbb{G}_m -action on $\mathbb{P}_{k'}^1$ induces an action of the torus $\text{Res}_{k'/k} \mathbb{G}_m$ on Y . Consider the subtorus

$$T_X := \text{Ker}(\text{Res}_{k'/k} \mathbb{G}_m \rightarrow \mathbb{G}_m)$$

where the arrow is the norm map, and define X to be the closure in Y of the T_X -orbit of x_0 . Clearly, $(T_X)_{\bar{k}} = (T_0)_{\bar{k}}$, hence the observation of Remark 7.2 shows that $X_{\bar{k}}$ is isomorphic to $(X_0)_{\bar{k}}$.

The Galois group of the extension k'/k coincides by definition with the group G_X , hence contains the Klein group V_4 . Therefore, the torus T_X and the corresponding toric variety X are not stably rational by [Vos98, §2.4.8].

Next, we apply [NS19, Theorem 4.2.11]. Since by Lemma 7.1 the toric variety X_0 has only ordinary double points, we conclude that the geometric general fiber of the universal family $\mathfrak{X} \rightarrow \mathfrak{P}$ of hyperplane sections of Y parameterized by \mathfrak{P} is not stably rational, and hence by [NS19, Corollary 4.1.5] a very general Fano threefold of type $X_{(1,1,1,1)}$ is not stably rational. \square

APPENDIX A. CONSTRUCTING MORPHISMS OF HILBERT SCHEMES

In this section we show how one can use technique of derived categories to construct morphisms of Hilbert schemes. For smooth projective varieties X and Y we denote by π_X and π_Y the projections from $X \times Y$ to the factors, and for an object $\mathcal{K} \in \mathbf{D}(X \times Y)$ we denote by

$$\Phi_{\mathcal{K}}: \mathbf{D}(X) \rightarrow \mathbf{D}(Y), \quad \mathcal{F} \mapsto \mathbf{R}\pi_{Y*}(\mathbf{L}\pi_X^*(\mathcal{F}) \otimes^{\mathbf{L}} \mathcal{K})$$

the corresponding Fourier–Mukai functor from the bounded derived category of coherent sheaves on X to that of Y . For an integral valued polynomial $p \in \mathbb{Q}[t]$ we denote by $\text{Hilb}_p(X)$ the Hilbert scheme of subschemes in X with Hilbert polynomials p .

Proposition A.1. *Let X and Y be smooth projective varieties. If $\mathcal{K} \in \mathbf{D}(X \times Y)$ is an object such that for any subscheme $Z \subset X$ with Hilbert polynomial p the object $\Phi_{\mathcal{K}}(\mathcal{O}_Z) \in \mathbf{D}(Y)$ is isomorphic to the structure sheaf of a point $y(Z) \in Y$ then there is a morphism of schemes*

$$\varphi: \text{Hilb}_p(X) \longrightarrow Y$$

such that $\varphi([Z]) = y(Z)$.

Proof. Let $Z \subset X \times S$ be an S -flat family of subschemes in X with Hilbert polynomial p . Let

$$\mathcal{F} := \Phi_{\pi_{XY}^* \mathcal{K}}(\mathcal{O}_Z) = \mathbf{R}\pi_{SY*}(\mathbf{L}\pi_{XS}^*(\mathcal{O}_Z) \otimes^{\mathbf{L}} \pi_{XY}^* \mathcal{K}) \in \mathbf{D}(S \times Y)$$

be the image of the structure sheaf of Z under the induced Fourier–Mukai functor from $\mathbf{D}(X \times S)$ to $\mathbf{D}(S \times Y)$, where π_{XS} , π_{SY} and π_{XY} are the projections of $X \times S \times Y$ to the pairwise products of factors. By base change and the projection formula, for each point $s \in S$ we have

$$i_s^* \mathcal{F} \cong \Phi_{\mathcal{K}}(\mathcal{O}_{Z_s}),$$

where $i_s: \{s\} \times Y \hookrightarrow S \times Y$ is the natural embedding and $Z_s \subset X$ is the fiber of Z over $s \in S$. Thus, we have $i_s^* \mathcal{F} \cong \mathcal{O}_{y(Z_s)}$ by assumption, therefore by [Kuz19, Lemma 4.4(iii)] there is a morphism $\varphi_S: S \rightarrow Y$ such that \mathcal{F} is isomorphic up to twist to the structure sheaf of the graph of φ_S . Now applying this argument to $S = \text{Hilb}_p(X)$ and Z the universal subscheme, we obtain the required morphism φ . \square

In §2.3 we apply Proposition A.1 to the Hilbert scheme of conics on the threefold $X \cong \text{Bl}_{\Gamma_1}(Q_1)$, where $Q_1 \subset \mathbb{P}^4$ is a smooth quadric and $\Gamma_1 \subset Q_1$ is a linearly normal smooth rational quartic curve. Recall that $F_1(X)$ and $F_2(X)$ denote the Hilbert schemes of (anticanonical) lines and conics on X , and that there is a natural embedding $\Gamma_1 \subset F_1(X)$.

Corollary A.2. *There is a morphism $\varphi_1: F_2(X) \rightarrow \Gamma_1$ such that for a smooth conic $C \subset X$ one has $\varphi_1([C]) = [L]$, where $L \subset X$ is the unique line corresponding to a point of Γ_1 such that $C \cap L \neq \emptyset$. Moreover, if $C = L_1 \cup L_2$ is a reducible conic, so that $L_1 \cap L_2 \neq \emptyset$ and if L'_2 is the other line corresponding to a point of Γ_1 such that $L_1 \cap L'_2 \neq \emptyset$, then $\varphi_1([C]) = [L'_2]$.*

Proof. Let $\pi_1: X \rightarrow Q_1$ be the blowup morphism and let $E_1 \subset X$ be its exceptional divisor; note that E_1 is the universal family of lines on X over the connected component $\Gamma_1 \subset F_1(X)$ of the Hilbert scheme of lines. Let $\varepsilon: E_1 \rightarrow X \times \Gamma_1$ be the corresponding embedding and consider

$$\mathcal{K} := \varepsilon_* \mathcal{O}_{E_1}(E_1) \in \mathbf{D}(X \times \Gamma_1).$$

Let us check that the assumption of Proposition A.1 is satisfied for $Y = \Gamma_1$.

If $C \subset X$ is a smooth conic, then $C \cdot E_1 = 1$, and $C \not\subset E_1$, therefore $C \cap E_1 = \{x\}$ is a single point and the intersection is transverse. Therefore $\mathcal{O}_C \otimes^{\mathbf{L}} \mathcal{O}_{E_1}(E_1) \cong \mathcal{O}_x$, hence $\Phi_{\mathcal{X}}(\mathcal{O}_C) \cong \mathcal{O}_{\pi_1(x)}$.

If $C = L_1 \cup L_2$ is a reducible conic, so that $L_2 \subset E_1$ and $L_1 \cap E_1 = \{x, x'\}$ with $L_1 \cap L_2 = \{x\}$ (if L_1 is tangent to E_1 we take $x' = x$), then using the exact sequences

$$0 \longrightarrow \mathcal{O}_{L_1}(-1) \longrightarrow \mathcal{O}_C \longrightarrow \mathcal{O}_{L_2} \longrightarrow 0 \quad \text{and} \quad 0 \longrightarrow \mathcal{O}_{L_2}(-1) \longrightarrow \mathcal{O}_C \longrightarrow \mathcal{O}_{L_1} \longrightarrow 0$$

it is easy to check that $\mathcal{O}_C \otimes^{\mathbf{L}} \mathcal{O}_{E_1}(E_1)$ fits into a distinguished triangle

$$\mathcal{O}_C \otimes^{\mathbf{L}} \mathcal{O}_{E_1}(E_1) \longrightarrow \mathcal{O}_{L_2}(-1) \oplus \mathcal{O}_{x'} \longrightarrow \mathcal{O}_{L_2}(-1)[2]$$

(if L_1 is tangent to E_1 the middle term should be replaced by an extension of $\mathcal{O}_{L_2}(-1)$ by $\mathcal{O}_{x'}$). Since the pushforward functor $\mathbf{R}\pi_{1*}$ kills the sheaf $\mathcal{O}_{L_2}(-1)$, it follows that $\Phi_{\mathcal{X}}(\mathcal{O}_C) \cong \mathcal{O}_{\pi_1(x)}$.

Now, applying Proposition A.1 we conclude that there is a morphism $\varphi_1: F_2(X) \rightarrow \Gamma_1$ such that $\varphi_1([C]) = \pi_1(x)$ if C is smooth and $\varphi_1([L_1 \cup L_2]) = \pi_1(x')$, with the notation for points x and x' introduced above. \square

REFERENCES

- [AB92] Alberto Alzati and Marina Bertolini. Sulla razionalità delle 3-varietà di Fano con $B_2 \geq 2$. *Matematiche*, 47(1):63–74, 1992.
- [Ben85] X. Benveniste. Sur le cone des 1-cycles effectifs en dimension 3. *Math. Ann.*, 272:257–265, 1985.
- [BW20] Olivier Benoist and Olivier Wittenberg. The Clemens–Griffiths method over non-closed fields. *Algebraic Geometry*, 7(6):696–721, 2020.
- [Cut88] Steven Cutkosky. Elementary contractions of Gorenstein threefolds. *Math. Ann.*, 280(3):521–525, 1988.
- [Isk80] V. A. Iskovskikh. Anticanonical models of three-dimensional algebraic varieties. *J. Sov. Math.*, 13:745–814, 1980.
- [KM98] János Kollár and Shigefumi Mori. *Birational geometry of algebraic varieties*, volume 134 of *Cambridge Tracts in Mathematics*. Cambridge University Press, Cambridge, 1998. With the collaboration of C. H. Clemens and A. Corti, Translated from the 1998 Japanese original.
- [Kol89] János Kollár. Flops. *Nagoya Math. J.*, 113:15–36, 1989.
- [Kol02] János Kollár. Unirationality of cubic hypersurfaces. *J. Inst. Math. Jussieu*, 1(3):467–476, 2002.
- [KP19] Alexander Kuznetsov and Yuri Prokhorov. Rationality of Fano threefolds over non-closed fields. *arXiv e-print*, 1911.08949, 2019.
- [KPS18] Alexander Kuznetsov, Yuri Prokhorov, and Constantin Shramov. Hilbert schemes of lines and conics and automorphism groups of Fano threefolds. *Japanese J. Math.*, 13(1):109–185, 2018.
- [Kuz19] Alexander Kuznetsov. Derived categories of families of sextic del Pezzo surfaces. *Int. Math. Res. Not. IMRN*, 2019. published online.
- [MM82] Shigefumi Mori and Shigeru Mukai. Classification of Fano 3-folds with $B_2 \geq 2$. *Manuscripta Math.*, 36(2):147–162, 1981/82. Erratum: *Manuscripta Math.* 110 (2003), 407.
- [Mor82] S. Mori. Threefolds whose canonical bundles are not numerically effective. *Ann. Math.*, 115:133–176, 1982.
- [MP19] S. Mori and Yu. Prokhorov. Threefold extremal curve germs with one non-Gorenstein point. *Izv. Math.*, 83(3):565–612, 2019.
- [NS19] Johannes Nicaise and Evgeny Shinder. The motivic nearby fiber and degeneration of stable rationality. *Invent. Math.*, 217(2):377–413, 2019.
- [Pro13a] Yuri Prokhorov. G-Fano threefolds, I. *Adv. Geom.*, 13(3):389–418, 2013.
- [Pro13b] Yuri Prokhorov. G-Fano threefolds, II. *Adv. Geom.*, 13(3):419–434, 2013.
- [Vos98] V. E. Voskresenskii. *Algebraic groups and their birational invariants*, volume 179 of *Translations of Mathematical Monographs*. American Mathematical Society, Providence, RI, 1998. Translated from the Russian manuscript by Boris Kunyavski.
- [Zak07] N. F. Zak. Quasi-triviality of forms of Segre varieties. *Russ. Math. Surv.*, 62(5):1018–1020, 2007.

STEKLOV MATHEMATICAL INSTITUTE OF RUSSIAN ACADEMY OF SCIENCES, MOSCOW, RUSSIA
LABORATORY OF ALGEBRAIC GEOMETRY, NRU HSE, MOSCOW, RUSSIA

Email address: akuznet@mi-ras.ru

STEKLOV MATHEMATICAL INSTITUTE OF RUSSIAN ACADEMY OF SCIENCES, MOSCOW, RUSSIA
LABORATORY OF ALGEBRAIC GEOMETRY, NRU HSE, MOSCOW, RUSSIA
DEPARTMENT OF ALGEBRA, MOSCOW STATE UNIVERSITY, MOSCOW, RUSSIA

Email address: prokhoro@mi-ras.ru