

CHARACTERIZATION OF COMMUTATIVE ALGEBRAS EMBEDDED INTO THE ALGEBRA OF SMOOTH OPERATORS

TOMASZ CIAŚ

ABSTRACT. The paper deal with the noncommutative Fréchet $*$ -algebra $\mathcal{L}(s', s)$ of the so-called smooth operators, i.e. linear and continuous operators acting from the space s' of slowly increasing sequences to the Fréchet space s of rapidly decreasing sequences. By a canonical identification, this algebra of smooth operators can be also seen as the algebra of the rapidly decreasing matrices. We give a full description of closed commutative $*$ -subalgebras of this algebra and we show that every closed subspace of s with basis is isomorphic (as a Fréchet space) to some closed commutative $*$ -subalgebra of $\mathcal{L}(s', s)$. As a consequence, we give some equivalent formulation of the long-standing Quasi-equivalence Conjecture for closed subspaces of s .

1. INTRODUCTION

In this paper we consider some specific noncommutative Fréchet algebra with involution – known, for instance, as the algebra $\mathcal{L}(s', s)$ of *smooth operators* or as the algebra

$$\mathcal{K}_\infty := \{(x_{j,k})_{j,k \in \mathbb{N}} \in \mathbb{C}^{\mathbb{N}^2} : \sup_{j,k \in \mathbb{N}} |x_{j,k}| j^q k^q < \infty \text{ for all } q \in \mathbb{N}_0\}$$

of *rapidly decreasing matrices*. Our main goal is to solve the following problems:

- (A) characterize Fréchet $*$ -algebras isomorphic to closed commutative $*$ -subalgebras of $\mathcal{L}(s', s)$ (Theorem 4);
- (B) characterize the underlying Fréchet spaces for the class of closed commutative $*$ -subalgebras of $\mathcal{L}(s', s)$ (Theorem 7).

By definition, $\mathcal{L}(s', s)$ is the Fréchet $*$ -algebra consisting of all linear and continuous operators from the space

$$(1) \quad s' := \left\{ (\xi_j)_{j \in \mathbb{N}} \in \mathbb{C}^{\mathbb{N}} : \sup_{j \in \mathbb{N}} |\xi_j| j^{-q} < \infty \text{ for some } q \in \mathbb{N}_0 \right\}$$

of *slowly increasing sequences* (this an inductive limit of Banach spaces) to the Fréchet space

$$(2) \quad s := \left\{ (\xi_j)_{j \in \mathbb{N}} \in \mathbb{C}^{\mathbb{N}} : \sup_{j \in \mathbb{N}} |\xi_j| j^q < \infty \text{ for all } q \in \mathbb{N}_0 \right\}$$

of *rapidly decreasing sequences*. Since there is a natural inclusion $\mathcal{L}(s', s)$ into $\mathcal{B}(\ell_2)$, the space $\mathcal{L}(s', s)$ is indeed a $*$ -algebra with composition of operators and hilbertain involution taken from $\mathcal{B}(\ell_2)$. Furthermore, $\mathcal{L}(s', s)$ with its natural topology of the uniform convergence on bounded subsets of s' is a Fréchet space.

We prove, in particular, that the class of closed commutative $*$ -subalgebras of $\mathcal{L}(s', s)$ is – in the sense of an isomorphism of Fréchet $*$ -algebras – exactly the class of Köthe sequence algebras which are closed under taking square roots (Definition 3) and at the same time isomorphic (as Fréchet spaces) to closed subspaces of s (Theorem 4). This can be easily expressed by conditions on the corresponding Köthe matrices. Moreover, from the solution of Problem (A) we get the solution of Problem (B): the

¹2010 *Mathematics Subject Classification*. Primary: 46A04, 46A11, 46A45. Secondary: 46A63, 46J40.

Key words and phrases: Köthe sequence algebras, nuclear Fréchet spaces, property (DN), structure of Fréchet algebras, smooth operators.

The research of the author was supported by the National Center of Science, grant no. 2013/09/N/ST1/04410.

underlying Fréchet spaces for closed commutative $*$ -subalgebras of $\mathcal{L}(s', s)$ are just closed subspaces of s with basis (Theorem 7).

It has been already shown in [2, Th. 4.8] that every infinite-dimensional closed commutative $*$ -subalgebra X of $\mathcal{L}(s', s)$ is isomorphic as a Fréchet $*$ -algebra to a Köthe sequence algebra

$$(3) \quad \lambda^\infty(\|P_j\|_q) := \left\{ (\xi_j)_{j \in \mathbb{N}} \in \mathbb{C}^{\mathbb{N}} : \sup_{j \in \mathbb{N}} |\xi_j| \|P_j\|_q < \infty \text{ for all } q \in \mathbb{N}_0 \right\},$$

where $(P_j)_{j \in \mathbb{N}}$ is a suitable sequence of pairwise orthogonal, nonzero, self-adjoint projections belonging to X and $(\|\cdot\|_q)_{q \in \mathbb{N}_0}$ is a sequence of continuous norms on $\mathcal{L}(s', s)$ which determines the Fréchet space topology of $\mathcal{L}(s', s)$. Hence, in order to solve problems (A) and (B), we needed to “understand” matrices $(\|P_j\|_q)_{j \in \mathbb{N}, q \in \mathbb{N}_0}$, where $(P_j)_{j \in \mathbb{N}}$ are pairwise orthogonal, nonzero, self-adjoint projections belonging to $\mathcal{L}(s', s)$. The key step in solutions of our problems was to show that for every nuclear Fréchet space with the so-called dominating Hilbert norm $\|\cdot\|$ there is a topological embedding $V: X \hookrightarrow s$ such that $\|Vx\|_{\ell_2} = \|x\|$ for every $x \in X$ (Theorem 14).

In this context, it is worth mentioning that, by [4, Th. 6.2], a Köthe sequence algebra of the form (3) is isomorphic to some closed $*$ -subalgebra of the algebra s if and only if it is isomorphic as a Fréchet space to a complemented subspace of s . One would think that every closed commutative $*$ -subalgebra of $\mathcal{L}(s', s)$ is of this type. Theorem [4, Th. 6.9] shows that this is not true (in the proof we give a concrete counterexample) and the present paper shows that indeed – taking into account the representation (3) – the class of the underlying Fréchet spaces for closed commutative $*$ -subalgebras of $\mathcal{L}(s', s)$ is as big as possible, i.e. as we mentioned above, it consists of all closed subspaces of s with basis.

The algebra of smooth operators is a quite natural object. It can be seen as a kind of a noncommutative analogon of the commutative Fréchet algebra s (look at the definition of \mathcal{K}_∞ above) and, in particular, it is isomorphic as a Fréchet space to s (see [12, Lemma 31.1]) – one of the most significant Fréchet spaces. On the level of Fréchet spaces, the space s is isomorphic to many classical spaces of smooth functions e.g. to the Schwartz space $\mathcal{S}(\mathbb{R})$ of smooth rapidly decreasing functions on the real line (which is a natural domain for the Fourier transform and differential operators with smooth coefficients), the space $C^\infty[-1, 1]$ of smooth functions on the interval $[-1, 1]$, the space $\mathcal{D}(K)$ of smooth functions with support contained in a compact set $K \subset \mathbb{R}^n$, $\text{int}(K) \neq \emptyset$, or the space $C^\infty(M)$ of smooth functions on an arbitrary compact smooth manifold M . Finally, the space s carries all the information about nuclear locally convex spaces. Indeed, the Kōmura-Kōmura theorem states that a locally convex space is nuclear if and only if it is isomorphic to some closed subspace of a suitable big cartesian product of the space s (see [12, Th. 29.8]). Consequently, a Fréchet space is nuclear (i.e. a Fréchet space in which every unconditionally convergent series is already absolutely convergent) if and only if it is isomorphic to some closed subspace of $s^{\mathbb{N}}$ (see [12, Cor. 29.9]).

Replacing s' with the space $\mathcal{S}'(\mathbb{R})$ of tempered distributions and s with $\mathcal{S}(\mathbb{R})$, we still end up with the same Fréchet $*$ -algebra – this time consisting of all linear and continuous operators acting from $\mathcal{S}'(\mathbb{R})$ to $\mathcal{S}(\mathbb{R})$, where $*$ -algebra operations are inherited from $L^2(\mathbb{R})$. In fact, for every Fréchet space E isomorphic to s and its topological dual E' , the space $\mathcal{L}(E', E)$ (with appropriate $*$ -algebra operations) of linear and continuous operators from E' to E is a Fréchet $*$ -algebra isomorphic to $\mathcal{L}(s', s)$ (for details see [3, Th. 1.10 & Ex. 1.13] and [8, Th. 2.1]).

The algebra $\mathcal{L}(s', s)$ appears and plays a significant role in K -theory of Fréchet algebras (see Bhatt, Inoue and Ogi [1, Ex. 2.12], Cuntz [6, p. 144], [7, p. 64–65], Glöckner and Langkamp [10], Phillips [14, Def. 2.1]) and in C^* -dynamical systems (Elliot, Natsume and Nest [9, Ex. 2.6]). Recently, Piszczek obtained several results concerning closed ideals, automatic continuity, amenability, Jordan decomposition and Grothendieck-type inequality in \mathcal{K}_∞ (see Piszczek [16, 17, 18, 19] and his forthcoming paper “The noncommutative Schwartz space is weakly amenable”).

Recall that two bases $(x_j)_{j \in \mathbb{N}}$ and $(y_j)_{j \in \mathbb{N}}$ of a Fréchet space X are called *quasi-equivalent* if there is a permutation $\sigma: \mathbb{N} \rightarrow \mathbb{N}$, a sequence $(\lambda_j)_{j \in \mathbb{N}}$ of nonzero scalars and a Fréchet space isomorphism

$T: X \rightarrow X$ such that $Tx_j = \lambda_j y_{\sigma(j)}$ for $j \in \mathbb{N}$. It is still an open question – the so-called Quasi-equivalence Conjecture – whether all bases in a nuclear Fréchet space are quasi-equivalent. This conjecture was posed in 1961 by Mityagin [13, §8.3]. There were many attempts to solve this problem (see [26] and references therein), e.g. Crone and Robinson [5] showed that in every nuclear Fréchet space with the so-called regular basis all bases are quasi-equivalent. In particular, all bases in s are quasi-equivalent but the conjecture remains open for closed subspaces of s . As a consequence of Theorem 4, we obtain some characterization of closed subspaces of s which have all bases quasi-equivalent (Theorem 15). We hope that our characterization will shed some light on this long-standing conjecture.

2. PRELIMINARIES

In what follows, \mathbb{N} will denote the set of natural numbers $\{1, 2, \dots\}$ and $\mathbb{N}_0 := \mathbb{N} \cup \{0\}$.

By e_j we denote the vector in $\mathbb{C}^{\mathbb{N}}$ whose j -th coordinate equals 1 and the others equal 0.

By a *Fréchet space* we mean a complete metrizable locally convex space over \mathbb{C} (we will not use locally convex spaces over \mathbb{R}). A *Fréchet algebra* is a Fréchet space which is an algebra with continuous multiplication. A *Fréchet *-algebra* is a Fréchet algebra with continuous involution.

We use the standard notation and terminology. All the notions from functional analysis are explained in [12].

We define the *space of rapidly decreasing sequences* as the Fréchet space

$$s := \left\{ \xi = (\xi_j)_{j \in \mathbb{N}} \in \mathbb{C}^{\mathbb{N}} : |\xi|_q := \left(\sum_{j=1}^{\infty} |\xi_j|^2 j^{2q} \right)^{1/2} < \infty \text{ for all } q \in \mathbb{N}_0 \right\}$$

with the topology corresponding to the system $(|\cdot|_q)_{q \in \mathbb{N}_0}$ of norms. The strong dual of s (i.e. the space of all continuous linear functionals on s with the topology of uniform convergence on bounded subsets of s , see e.g. [12, Definition on p. 267]) can be identified with the *space of slowly increasing sequences*

$$s' := \left\{ \xi = (\xi_j)_{j \in \mathbb{N}} \in \mathbb{C}^{\mathbb{N}} : |\xi|'_q := \left(\sum_{j=1}^{\infty} |\xi_j|^2 j^{-2q} \right)^{1/2} < \infty \text{ for some } q \in \mathbb{N}_0 \right\}.$$

It is an easy exercise to show that replacing in the above definition of s and s' the ℓ_2 -norms by the corresponding sup-norms – as in (1) and (2) – or ℓ_p -norms ($1 \leq p < \infty$), we end up with the same sequence spaces.

Closed subspaces of the space s can be characterized by the so-called property (DN) (see [12, Prop. 31.5]).

Definition 1. A Fréchet space X with a fundamental system $(\|\cdot\|_q)_{q \in \mathbb{N}_0}$ of seminorms has the *property (DN)* (see [12, Def. on p. 359]) if there is a continuous norm $\|\cdot\|$ on X such that for all $q \in \mathbb{N}_0$ there is $r \in \mathbb{N}_0$ and $C > 0$ such that

$$\|x\|_q^2 \leq C \|x\| \|x\|_r$$

for all $x \in X$. The norm $\|\cdot\|$ is called a *dominating norm*.

We will also use the condition (DN) in the following equivalent form (see [12, Lemma 29.10]): there is a continuous norm $\|\cdot\|$ on X such that for any $q \in \mathbb{N}_0$ and $\theta \in (0, 1)$ there is $r \in \mathbb{N}_0$ and $C > 0$ such that

$$\|x\|_q \leq C \|x\|^{1-\theta} \|x\|_r^\theta$$

for all $x \in X$.

The *algebra of smooth operators* is defined as the Fréchet *-algebra $\mathcal{L}(s', s)$ of continuous linear operators from s' to s with the topology of uniform convergence on bounded sets in s' . The *-algebra operations on $\mathcal{L}(s', s)$ are inherited from $\mathcal{B}(\ell_2)$ (note that every smooth operator is bounded on ℓ_2).

The algebra $\mathcal{L}(s', s)$ can be also seen as the algebra of matrices which are sometimes more handy; it is isomorphic, via the map $x \mapsto ((xe_k, ej))_{j,k \in \mathbb{N}}$, to the Fréchet $*$ -algebra

$$\mathcal{K}_\infty := \{(x_{j,k})_{j,k \in \mathbb{N}} \in \mathbb{C}^{\mathbb{N}^2} : \sup_{j,k \in \mathbb{N}} |x_{j,k}| j^q k^q < \infty \text{ for all } q \in \mathbb{N}_0\}$$

of *rapidly decreasing matrices* (with matrix multiplication and matrix complex conjugation). Therefore – when reading this paper – one can replace each “ $\mathcal{L}(s', s)$ ” with “ \mathcal{K}_∞ ”. More details concerning $\mathcal{L}(s', s)$ can be find in the introductions of [2, 3, 4].

Definition 2. A matrix $A = (a_{j,q})_{j \in \mathbb{N}, q \in \mathbb{N}_0}$ of nonnegative numbers such that

- (i) for each $j \in \mathbb{N}$ there is $q \in \mathbb{N}_0$ such that $a_{j,q} > 0$
- (ii) $a_{j,q} \leq a_{j,q+1}$ for $j \in \mathbb{N}$ and $q \in \mathbb{N}_0$

is called a *Köthe matrix*. For $1 \leq p < \infty$ and a Köthe matrix A we define the *Köthe space*

$$\lambda^p(A) := \left\{ \xi = (\xi_j)_{j \in \mathbb{N}} \in \mathbb{C}^{\mathbb{N}} : |\xi|_{\lambda^p(A), q} := \sum_{j=1}^{\infty} |\xi_j|^p a_{j,q}^p < \infty \text{ for all } q \in \mathbb{N}_0 \right\}$$

and for $p = \infty$

$$\lambda^\infty(A) := \left\{ \xi = (\xi_j)_{j \in \mathbb{N}} \in \mathbb{C}^{\mathbb{N}} : |\xi|_{\lambda^\infty(A), q} := \sup_{j \in \mathbb{N}} |\xi_j| a_{j,q} < \infty \text{ for all } q \in \mathbb{N}_0 \right\}$$

with the locally convex topology given by the seminorms $(|\cdot|_{\lambda^p(A), q})_{q \in \mathbb{N}_0}$ (see e.g. [12, Definition p. 326]). For simplicity, we will also write $\lambda^p(a_{j,q})$ (i.e. only the entries of the matrix) instead of $\lambda^p(A)$.

All spaces $\lambda^p(A)$ are Fréchet spaces ([12, Lemma 27.1]). By the Dynin-Mityagin Theorem (see [12, Th. 28.12]) if $\lambda^p(A)$ is nuclear (i.e. for all q there is r such that $\sum_{j=1}^{\infty} \frac{a_{j,q}}{a_{j,r}} < \infty$), then the sequence of vectors $(e_j)_{j \in \mathbb{N}}$ is an absolute Schauder basis of $\lambda^p(A)$. We will also use the following result: $\lambda^p(A)$ is nuclear for some $1 \leq p \leq \infty$ if and only if $\lambda^p(A) = \lambda^q(A)$ as Fréchet spaces for all $1 \leq p, q \leq \infty$ ([12, Prop. 28.16]).

For Köthe matrices $A = (a_{j,q})_{j \in \mathbb{N}, q \in \mathbb{N}_0}$ and $B = (b_{j,q})_{j \in \mathbb{N}, q \in \mathbb{N}_0}$ such that

$$\forall q \in \mathbb{N}_0 \exists r \in \mathbb{N}_0 \exists C > 0 \forall j \in \mathbb{N} \quad a_{j,q} \leq C b_{j,r},$$

we write $A \prec B$. If $A \prec B$ and $B \prec A$, then we write $A \sim B$. For a bijection $\sigma: \mathbb{N} \rightarrow \mathbb{N}$, A_σ denotes the Köthe matrix $(a_{\sigma(j),q})_{j \in \mathbb{N}, q \in \mathbb{N}_0}$ and, moreover, A^2 is by definition the Köthe matrix $(a_{j,q}^2)_{j \in \mathbb{N}, q \in \mathbb{N}_0}$.

By [15, Prop. 3.1], $\lambda^1(A)$ with pointwise multiplication is an algebra (and, clearly, if $\lambda^1(A)$ is nuclear so is $\lambda^p(A)$ for $1 < p \leq \infty$) if and only if $A \prec A^2$; in this case $\lambda^p(A)$ is called a *Köthe algebra*.

Definition 3. We say that a Köthe space $\lambda^p(A)$ is *closed under taking square root* if it is a Köthe algebra and for each sequence $(\xi_j)_{j \in \mathbb{N}}$ in $\lambda^p(A)$ of nonnegative scalars, the sequence $(\sqrt{\xi_j})_{j \in \mathbb{N}}$ belongs to $\lambda^p(A)$ as well.

3. COMMUTATIVE SUBALGEBRAS OF $\mathcal{L}(s', s)$

Our main result – solving Problem (A) from Introduction – reads as follows:

Theorem 4. For a Köthe matrix $A = (a_{j,q})_{j \in \mathbb{N}, q \in \mathbb{N}_0}$ the following assertions are equivalent:

- (1) $\lambda^2(A)$ is isomorphic as a Fréchet $*$ -algebra to some closed commutative $*$ -subalgebra of $\mathcal{L}(s', s)$;
- (2) there is a sequence $(P_j)_{j \in \mathbb{N}}$ of nonzero pairwise orthogonal self-adjoint projections belonging to $\mathcal{L}(s', s)$ such that $A \sim (\|P_j\|_q)_{j \in \mathbb{N}, q \in \mathbb{N}_0}$;
- (3) there is a basic sequence $(f_j)_{j \in \mathbb{N}}$ of s which is at the same time an orthonormal sequence in ℓ_2 such that $A \sim (\|f_j\|_q)_{j \in \mathbb{N}, q \in \mathbb{N}_0}$;
- (4) the matrix A satisfies the following conditions:
 - (i) $\forall q \in \mathbb{N}_0 \exists r \in \mathbb{N}_0 \quad \sum_{j=1}^{\infty} \frac{a_{j,q}}{a_{j,r}} < \infty$,
 - (ii) $\exists p \in \mathbb{N}_0 \exists C > 0 \forall j \in \mathbb{N} \quad a_{j,p} \geq \frac{1}{C}$,

- (iii) $A^2 \prec A$;
- (5) *the matrix A satisfies the following conditions:*
 - (i) $\forall q \in \mathbb{N}_0 \exists r \in \mathbb{N}_0 \sum_{j=1}^{\infty} \frac{a_{j,q}}{a_{j,r}} < \infty$,
 - (ii) $\exists p \in \mathbb{N}_0 \forall q \in \mathbb{N}_0 \exists r \in \mathbb{N}_0, C > 0 \forall j \in \mathbb{N} \quad a_{j,q}^2 \leq C a_{j,p} a_{j,r}$,
 - (iii) $A \sim A^2$;
- (6) $\lambda^2(A)$ *is a nuclear biprojective Köthe algebra with the property (DN);*
- (7) $\lambda^2(A)$ *is isomorphic as a Fréchet space to a closed subspace of s and it is closed under taking square roots;*
- (8) $\lambda^2(A)$ *is a nuclear Fréchet space with $\|\cdot\|_{\ell_2}$ as a dominating norm.*

Remark 5. By [2, Th. 4.8], every infinite-dimensional closed commutative $*$ -subalgebra of $\mathcal{L}(s', s)$ is isomorphic as a Fréchet $*$ -algebra to some Köthe sequence algebra, and thus the above theorem characterizes in fact all infinite-dimensional closed commutative $*$ -subalgebras of $\mathcal{L}(s', s)$.

Remark 6. Homology theory is out of the mainstream of this paper, and therefore we only recall that the a Fréchet algebra A is called *biprojective* if the product map $\pi: A \hat{\otimes} A \rightarrow A, a \otimes b \mapsto ab$, has a right inverse in the category of A -bimodules (see also [11, Def. IV.5.1]). The characterization of biprojectivity for Köthe algebras in terms of the corresponding Köthe matrices is due to Pirkovskii [15, Th. 5.9].

The following theorem leads to the solution of Problem (B) (see Introduction).

Theorem 7. *For a Köthe matrix $A = (a_{j,q})_{j \in \mathbb{N}, q \in \mathbb{N}_0}$ the following assertions are equivalent:*

- (1) $\lambda^2(A)$ *is isomorphic as a Fréchet space to some closed commutative $*$ -subalgebra of $\mathcal{L}(s', s)$;*
- (2) *the matrix A satisfies the following conditions:*
 - (i) $\forall q \in \mathbb{N}_0 \exists r \in \mathbb{N}_0 \sum_{j=1}^{\infty} \frac{a_{j,q}}{a_{j,r}} < \infty$,
 - (ii) $\exists p \in \mathbb{N}_0 \forall q \in \mathbb{N}_0 \exists r \in \mathbb{N}_0, C > 0 \forall j \in \mathbb{N} \quad a_{j,q}^2 \leq C a_{j,p} a_{j,r}$;
- (3) $\lambda^2(A)$ *is a nuclear Fréchet space with the property (DN).*
- (4) $\lambda^2(A)$ *is a Fréchet space isomorphic to a closed subspace of s .*

Proof. It is well known that a Köthe space $\lambda^2(A)$ is nuclear and has the property (DN) if and only if A satisfies conditions (i) and (ii), respectively. This shows (1) \Rightarrow (2) \Rightarrow (3), because $\mathcal{L}(s', s) \cong s$ as Fréchet spaces and nuclearity together with the property (DN) are inherited by closed subspaces.

In order to prove (3) \Rightarrow (1), choose $p_0 \in \mathbb{N}_0$ for which $|\cdot|_{\lambda^2(A), p_0}$ is dominating norm on $\lambda^2(A)$ and define the Köthe matrix $B := (a_{j,q}/a_{j,p_0})_{j \in \mathbb{N}, q \in \mathbb{N}_0}$. Let $T: \lambda^2(B) \rightarrow \lambda^2(A)$, $T\xi := (\xi_j/a_{j,p_0})_{j \in \mathbb{N}}$. Then, for $\xi \in \lambda^2(B)$ and $q \in \mathbb{N}_0$, we have $|T\xi|_{\lambda^2(A), q} = |\xi|_{\lambda^2(B), q}$, and thus T is an isomorphism of Fréchet spaces. Moreover, for $q \in \mathbb{N}_0$ there is $r \in \mathbb{N}_0$ and $C > 0$ such that

$$|\xi|_{\lambda^2(B), q}^2 = |T\xi|_{\lambda^2(A), q}^2 \leq C |T\xi|_{\lambda^2(A), p_0} |T\xi|_{\lambda^2(A), r} = C \|\xi\|_{\ell_2} |\xi|_{\lambda^2(B), r}$$

$\xi \in \lambda^2(B)$. Therefore, $\|\cdot\|_{\ell_2}$ is a dominating norm on $\lambda^2(B)$, and the conclusion follows from Theorem 4 (implication (8) \Rightarrow (1)).

Finally, equivalence (3) \Leftrightarrow (4) follows directly from [12, Prop. 31.5]. □

As a direct consequence of Theorems 4, 7 and [12, Cor. 28.13, Prop. 28.16, Prop. 31.5] we also get the following characterization.

Corollary 8. *For a Fréchet space E the following assertions are equivalent:*

- (1) *E is isomorphic as a Fréchet space to some closed commutative $*$ -subalgebra of $\mathcal{L}(s', s)$;*
- (2) *E is isomorphic as a Fréchet space to a Köthe space $\lambda^2(|f_j|_q)$ for some basic sequence $(f_j)_{j \in \mathbb{N}}$ of s which is at the same time an orthonormal sequence in ℓ_2 ;*
- (3) *E is nuclear, has a basis and the property (DN);*
- (4) *E is isomorphic as a Fréchet space to some closed subspace of s with basis.*

Now, we shall provide the proof of Theorem 4. We start with some lemmas involving short exact sequences of Fréchet-Hilbert spaces.

Lemma 9. *Let*

$$0 \longrightarrow E \xrightarrow{j} F \xrightarrow{q} G \longrightarrow 0$$

be a short exact sequence of Fréchet-Hilbert spaces. Let $\|\cdot\|_E$ be a continuous Hilbert norm on E and let $\|\cdot\|_G$ be a continuous norm on G . Then there is a continuous Hilbert norm $\|\cdot\|_F$ on F with

- (α) $\|x\|_E \sim \|j(x)\|_F$ for $x \in E$;
- (β) $\|q(x)\|_G \leq \inf_{y \in \ker q} \|x - y\|_F$ for $x \in F$.

Proof. Without loss of generality we may assume that E is a closed subspace of F and $j(x) = x$ for $x \in E$. By [12, Remark 22.8], there is a continuous Hilbert norm $\|\cdot\|_1$ on F with $\|x\|_E \leq \|x\|_1$ for $x \in E$. Moreover, by the continuity of q , there is a continuous Hilbert norm $\|\cdot\|_2$ on F such that

$$\|q(x)\|_G \leq \inf_{z \in E} \|x - z\|_2$$

for $x \in F$. Then $\|\cdot\| := (\|\cdot\|_1^2 + \|\cdot\|_2^2)^{1/2}$ is a continuous Hilbert norm on F such that

- (i) $\|x\|_E \leq \|x\|$ for $x \in E$;
- (ii) $\|q(x)\|_G \leq \inf_{z \in E} \|x - z\|$ for $x \in F$.

Define $\|\cdot\|_F$, by

$$\|x\|_F^2 = \inf_{z \in E} (\|z\|_E^2 + \|x - z\|^2)$$

for $x \in F$.

To show condition (α), let us fix $x \in E$. Taking $z := x$ in the infimum, we get

$$\|x\|_F \leq \|x\|_E.$$

Moreover, for $z \in E$, we obtain

$$\|x\|_E^2 \leq (\|z\|_E + \|x - z\|_E)^2 = \|z\|_E^2 + \|x - z\|_E^2 + 2\|z\|_E \|x - z\|_E.$$

If $\|z\|_E \leq \|x - z\|_E$, then $\|z\|_E \|x - z\|_E \leq \|x - z\|_E^2$. Otherwise, $\|z\|_E \|x - z\|_E \leq \|z\|_E^2$ so, by (i),

$$\|x\|_E^2 \leq 3(\|z\|_E^2 + \|x - z\|_E^2) \leq 3(\|z\|_E^2 + \|x - z\|^2).$$

In consequence,

$$(4) \quad \|x\|_E \leq \sqrt{3} \inf_{z \in E} (\|z\|_E^2 + \|x - z\|^2)^{1/2} = \sqrt{3} \|x\|_F,$$

and thus condition (α) is fulfilled. By (ii), for $x \in F$, we get

$$\begin{aligned} \|q(x)\|_G^2 &\leq \inf_{z \in E} \|x - z\|^2 = \inf_{z_1 \in E} \inf_{z_2 \in E} (\|z_2\|_E^2 + \|x - z_1\|^2) = \inf_{z_1, z_2 \in E} (\|z_2\|_E^2 + \|x - (z_1 + z_2)\|^2) \\ &= \inf_{z_1 \in E} \inf_{z_2 \in E} (\|z_2\|_E^2 + \|(x - z_1) - z_2\|^2) = \inf_{z \in E} \|x - z\|_F^2, \end{aligned}$$

which yields condition (β).

Now define a Hilbert norm $\|(\cdot, \cdot)\|_H$ on $H := E \times F$ by

$$\|(x, y)\|_H^2 := \|x\|_E^2 + \|y\|_F^2.$$

Let

$$R: (H, \|(\cdot, \cdot)\|_H) \rightarrow F, \quad (x, y) \mapsto x + y$$

and let

$$Q: (H, \|(\cdot, \cdot)\|_H) \rightarrow (H/\ker R, \|\cdot\|_{H/\ker R})$$

be the quotient map. Then

$$\begin{aligned}
\|Q(x, y)\|_{H/\ker R} &= \inf_{(x', y') \in \ker R} \|(x, y) - (x', y')\|_H = \inf_{x' \in E} \|(x, y) - (x', -x')\|_H \\
(5) \quad &= \inf_{x' \in E} (\|x - x'\|_E^2 + \|y + x'\|_E^2)^{1/2} = \inf_{z \in E} (\|z\|_E^2 + \|(x + y) - z\|_E^2)^{1/2} \\
&= \|x + y\|_F.
\end{aligned}$$

In particular

$$(6) \quad \|\cdot\|_F = \|Q(0, \cdot)\|_{H/\ker R},$$

hence $\|\cdot\|_F$ is a seminorm. Moreover, since $\|\cdot\|_F \leq \|\cdot\|$, it follows that $\|\cdot\|_F$ is continuous.

Now assume that $\|x\|_F = 0$ for some $x \in F$. Then, by condition (β) , $\|q(x)\|_G \leq \|x\|_F = 0$, and since $\|\cdot\|_G$ is a norm, we have $x \in \ker q = E$. Moreover, by (4), $\|x\|_E \leq \sqrt{3}\|x\|_F = 0$, but $\|\cdot\|_E$ is a norm, so $x = 0$. This shows that $\|\cdot\|_F$ is a norm, and therefore, by (5), $\|\cdot\|_{H/\ker R}$ is a norm as well (i.e. $\ker R$ is a closed subspace of H – see e.g. [12, Lemma 5.10]). Since the quotient norm induced by a Hilbert norm is a Hilbert norm, the norm $\|\cdot\|_{H/\ker R}$ is hilbertian (see e.g. [20, p. 44]). Therefore, by (6), $\|\cdot\|_F$ satisfies the parallelogram law, hence $\|\cdot\|_F$ is a Hilbert norm. \square

An easy proof of our next lemma is left to the reader.

Lemma 10. *Let us consider the diagram*

$$\begin{array}{ccc}
(Y, \|\cdot\|_Y) & \xleftarrow{\iota} & (X, \|\cdot\|_X) \\
\downarrow j_Y & & \downarrow j_X \\
(\tilde{Y}, \|\cdot\|_{\tilde{Y}}) & \xrightarrow{\tilde{\iota}} & (\tilde{X}, \|\cdot\|_{\tilde{X}})
\end{array}$$

where

- $(X, \|\cdot\|_X)$ and $(Y, \|\cdot\|_Y)$ are normed spaces such that Y is a closed subspace of $(X, \|\cdot\|_X)$ and $\|\cdot\|_Y \sim \|\cdot\|_{X|_Y}$;
- $(\tilde{Y}, \|\cdot\|_{\tilde{Y}})$ and $(\tilde{X}, \|\cdot\|_{\tilde{X}})$ are the completions of $(Y, \|\cdot\|_Y)$ and $(X, \|\cdot\|_X)$, respectively;
- ι, j_Y and j_X are the canonical inclusions;
- $\tilde{\iota}$ is the continuous linear extensions of the map $j_X \circ \iota: Y \rightarrow \tilde{X}$.

Then $\tilde{\iota}(\tilde{Y})$ is a closed subspace of $(\tilde{X}, \|\cdot\|_{\tilde{X}})$.

Lemma 11. *Let*

$$0 \longrightarrow E \xrightarrow{j} F \xrightarrow{q} G \longrightarrow 0$$

be a short exact sequence of Fréchet-Hilbert spaces. Let $\|\cdot\|_E$ be a continuous Hilbert norm on E and let us assume that G has a continuous norm. Then there is a continuous Hilbert norm $\|\cdot\|$ on F such that $\|j(x)\| = \|x\|_E$ for $x \in E$.

Proof. Without loss of generality we may assume that E is a closed subspace of F and $j(x) = x$ for $x \in E$. Let $\|\cdot\|_G$ be a continuous norm on G . By Lemma 9, there is a continuous Hilbert norm $\|\cdot\|_F$ on F such that

- (α) $\|\cdot\|_E \sim \|\cdot\|_{F|_E}$ for $x \in E$;
- (β) $\|q(x)\|_G \leq \inf_{y \in \ker q} \|x - y\|_F =: \|x\|_G$ for $x \in F$.

Consider the diagram

$$\begin{array}{ccccccc}
0 & \longrightarrow & (E, \|\cdot\|_E) & \xrightarrow{j} & (F, \|\cdot\|_F) & \xrightarrow{q} & (G, \|\cdot\|_G) \longrightarrow 0 \\
& & \downarrow j_E & & \downarrow j_F & & \\
& & (\tilde{E}, \|\cdot\|_{\tilde{E}}) & \xrightleftharpoons[\pi]{\tilde{j}} & (\tilde{F}, \|\cdot\|_{\tilde{F}}) & & \\
& & \uparrow \kappa & \swarrow \pi_0 & & & \\
& & (\hat{E}, \|\cdot\|_{\hat{E}}) & & & &
\end{array}$$

where

- $(\tilde{E}, \|\cdot\|_{\tilde{E}})$ and $(\tilde{F}, \|\cdot\|_{\tilde{F}})$ are the completions of $(E, \|\cdot\|_E)$ and $(F, \|\cdot\|_F)$, respectively;
- j_E and j_F are the canonical inclusions;
- \tilde{j} is the continuous extensions of the map $j_F \circ j: E \rightarrow \tilde{F}$.

Since, by (β) , $\|\cdot\|_G$ is a norm, E is a closed subspace of $(F, \|\cdot\|_F)$, and thus, by Lemma 10, $\hat{E} := \tilde{j}(\tilde{E})$ is a closed subspace of the Hilbert space $(\tilde{F}, \|\cdot\|_{\tilde{F}})$. Then π_0 is defined to be the orthogonal projections onto \hat{E} , κ is the inverse of $\tilde{\iota}$ on \hat{E} and $\pi := \kappa \circ \pi_0$.

Let us define

$$\|x\|^2 := \|(\pi \circ j_F)(x)\|_{\tilde{E}}^2 + \|q(x)\|_G^2$$

for $x \in F$. Clearly, $\|\cdot\|$ is a continuous seminorm on F . Next, for $x \in E$, we get

$$\|x\| = \|(\pi \circ j_F)(x)\|_{\tilde{E}} = \|j_E(x)\|_{\tilde{E}} = \|x\|_E,$$

hence $\|\cdot\|_E = \|\cdot\|$. For $x \in F$ with $\|x\| = 0$ we have $\|q(x)\|_G = 0$, and thus $x \in E$. Then we have also $\|x\|_E = \|x\| = 0$ so $x = 0$. This shows that $\|\cdot\|$ is a norm.

Finally, since $\|\cdot\|_{\tilde{E}}$ and $\|\cdot\|_G$ are Hilbert norms, $\|\cdot\|$ is a Hilbert norm as well. \square

The proof of the following lemma is a slight modification of the Vogt's proof of [23, Lemma 3.1]. For the convenience of the reader we present here the full argument.

Lemma 12. *Let*

$$0 \longrightarrow E \xrightarrow{j} F \xrightarrow{q} G \longrightarrow 0$$

be a short exact sequence of Fréchet-Hilbert spaces. Let $\|\cdot\|_E$ be dominating Hilbert norm on E and let us assume that G has the property (DN). Then there is a dominating Hilbert norm $\|\cdot\|$ on F such that $\|j(x)\| = \|x\|_E$ for $x \in E$.

Proof. Without loss of generality we may assume that E is a closed subspace of F and $j(x) = x$ for $x \in E$. By Lemma 11, there is continuous Hilbert norm $\|\cdot\|_F$ on F such that $\|\cdot\|_F|_E = \|\cdot\|_E$ and, by assumption, there is a dominating Hilbert norm $\|\cdot\|_G$ on G . Define

$$\|x\|^2 := \|x\|_F^2 + \|qx\|_G^2$$

for $x \in F$ and let $U_0 := \{x \in E: \|x\| < 1\}$. Then, clearly, $\|\cdot\|$ is a continuous Hilbert norm on F . We shall show that $\|\cdot\|$ is a dominating norm.

For an absolutely convex zero neighborhood U in F , let $\|\cdot\|_U$ denote the Minkowski functional for U and let $\|\cdot\|_{\hat{U}}: G \rightarrow [0, \infty)$ be defined by $\|qx\|_{\hat{U}} := \inf\{\|x+y\|_U: y \in E\}$ for $x \in F$.

Since $\|\cdot\|_E$ is a dominating norm on E , for each absolutely convex zero neighborhood U in F there is an absolutely convex zero neighborhood W in F with $W \subseteq U \cap U_0$ such that

$$(7) \quad \|z\|_U \leq \|z\|_E^{2/3} \|z\|_W^{1/3},$$

for all $z \in E$. Since $\|\cdot\|_G$ is a dominating norm on G , there is an absolutely convex zero neighborhood V in F with $V \subseteq W$ such that

$$(8) \quad \|qx\|_{\widehat{W}} \leq \|qx\|_G^{3/4} \|qx\|_{\widehat{V}}^{1/4} \leq \|x\|^{3/4} \|x\|_V^{1/4}.$$

for all $x \in F$.

Let us fix $x \in F$ and $\varepsilon > 0$. By the very definition of the norm $\|\cdot\|_{\widehat{W}}$, there is $y \in F$ such that

$$qx = qy \quad \text{and} \quad \|y\|_W \leq \|qx\|_{\widehat{W}} + \varepsilon,$$

and thus (8) yields

$$(9) \quad \|y\|_W \leq \|x\|^{3/4} \|x\|_V^{1/4} + \varepsilon.$$

Hence, for $z := x - y$, we have

$$\|z\|_E \leq \|x\|_F + \|y\|_F \leq \|x\| + \|y\|_W \leq \|x\| + \|x\|^{3/4} \|x\|_V^{1/4} + \varepsilon,$$

and since

$$\|x\| = \|x\|^{3/4} \|x\|_V^{1/4} \leq \|x\|^{3/4} \|x\|_V^{1/4},$$

it follows that

$$(10) \quad \|z\|_E \leq 2\|x\|^{3/4} \|x\|_V^{1/4} + \varepsilon.$$

Moreover, by (9),

$$\|z\|_W \leq \|x\|_W + \|y\|_W \leq \|x\|_V + \|x\|^{3/4} \|x\|_V^{1/4} + \varepsilon \leq 2\|x\|_V + \varepsilon.$$

Now, since for $a, b \geq 0$ we have $(a+b)^{2/3} \leq 3(a^{2/3} + b^{2/3})$, it follows from (7) and (10) that

$$(11) \quad \begin{aligned} \|z\|_U &\leq \left(2\|x\|^{3/4} \|x\|_V^{1/4} + \varepsilon\right)^{2/3} (2\|x\|_V + \varepsilon)^{1/3} \\ &\leq \left[3 \left(2\|x\|^{3/4} \|x\|_V^{1/4}\right)^{2/3} + \varepsilon^{2/3}\right] \left[(2\|x\|_V)^{1/3} + \varepsilon^{1/3}\right] \\ &= \left(3\sqrt[3]{4}\|x\|^{1/2} \|x\|_V^{1/6} + 3\varepsilon^{2/3}\right) \left(\sqrt[3]{2}\|x\|_V^{1/3} + \varepsilon^{1/3}\right) = 6\|x\|^{1/2} \|x\|_V^{1/2} + h(\varepsilon), \end{aligned}$$

where $h(\varepsilon) \searrow 0$ when $\varepsilon \searrow 0$.

Next, by (9),

$$(12) \quad \|y\|_U \leq \|y\|_W \leq \|x\|^{3/4} \|x\|_V^{1/4} + \varepsilon = \|x\|^{1/2} \|x\|_V^{1/2} \left(\frac{\|x\|}{\|x\|_V}\right)^{1/4} + \varepsilon \leq \|x\|^{1/2} \|x\|_V^{1/2} + \varepsilon.$$

Consequently, by (11),

$$\|x\|_U \leq \|y\|_U + \|z\|_U \leq 7\|x\|^{1/2} \|x\|_V^{1/2} + h(\varepsilon) + \varepsilon,$$

and finally, since $h(\varepsilon) + \varepsilon$ can be chosen arbitrary small, we have

$$\|x\|_U^2 \leq 49\|x\| \|x\|_V,$$

which shows that $\|\cdot\|$ is a dominating norm on F . □

Corollary 13. *Let E be a closed subspace of s such that s/E has a dominating norm. Then for every dominating Hilbert norm $\|\cdot\|_E$ on E there is a dominating Hilbert norm $\|\cdot\|$ on s such that the restriction of $\|\cdot\|$ to E is equal to $\|\cdot\|_E$.*

Proof. The results follows by applying Lemma 12 to the canonical short exact sequence

$$0 \longrightarrow E \hookrightarrow s \longrightarrow s/E \longrightarrow 0.$$

□

Corollary 13 together with some results of Vogt ([25, Cor. 7.6], [22, Prop. 3.3(a)]) allows to give a stronger version of the well-known theorem of Vogt characterizing closed subspaces of s in terms of nuclearity and the property (DN) (see [21, Satz 1.7] or [12, Prop. 31.5]).

Theorem 14. *For a Fréchet space X and a seminorm $\|\cdot\|: X \rightarrow [0, \infty)$ the following assertions are equivalent:*

- (1) *there is a topological embedding $V: X \hookrightarrow s$ such that $\|Vx\|_{\ell_2} = \|x\|$ for all $x \in X$;*
- (2) *X is nuclear and $\|\cdot\|$ is a dominating Hilbert norm on X .*

Proof. (\Rightarrow) By assumption, X is isomorphic to a closed subspace of s , so it is nuclear. Moreover, since $\|\cdot\|_{\ell_2}$ is a dominating Hilbert norm on s , it is easily seen that $\|\cdot\|$ is a dominating Hilbert norm on X .

(\Leftarrow) Since X is a nuclear Fréchet space with the property (DN), by Theorem [12, Prop. 31.5], it is isomorphic to some closed subspace E of s . Without loss of generality we may assume that s/E has (DN). Indeed, by [22, Prop. 3.3(a)], for an arbitrary closed subspace E_0 of s isomorphic to X , there is a closed subspace F_0 of s and an exact sequence

$$0 \longrightarrow E_0 \xrightarrow{\iota_0} s \xrightarrow{q_0} F_0 \longrightarrow 0$$

with continuous linear maps. Then q_0 induces an isomorphism $\overline{q_0}: s/\ker q_0 \rightarrow F_0$ (see [12, Prop. 22.11]), and thus

$$s/\iota_0(E_0) = s/\ker q_0 \cong F_0.$$

Hence $E := \iota_0(E_0)$ is a closed subspace of s such that s/E has (DN), and by the open mapping theorem $E \cong E_0$.

Let $T: X \rightarrow E$ be an isomorphism and set $\|x\|_E := \|T^{-1}x\|$ for $x \in E$. An easy calculation shows that $\|\cdot\|_E$ is a dominating Hilbert norm on E . Hence, by Corollary 13, there is a dominating Hilbert norm $\|\cdot\|_s$ on s such that $\|\cdot\|_s|_E = \|\cdot\|_E$. Moreover, by [25, Cor. 7.6], there is an automorphism U of s such that $\|Ux\|_{\ell_2} = \|x\|_s$ for $x \in s$. Now, for $V := U \circ T$ and $x \in X$, we get

$$\|Vx\|_{\ell_2} = \|U(Tx)\|_{\ell_2} = \|Tx\|_s = \|Tx\|_E = \|x\|,$$

which proves the theorem. □

Finally, we are ready to prove our main result.

Proof of Theorem 4. (3) \Rightarrow (1): We should only note that, by [4, Prop. 4.2], $\lambda^2(A) \cong \lambda^\infty(|f_j|_q)$ as a Fréchet *-algebras and $\lambda^\infty(|f_j|_q)$ is isomorphic as a Fréchet *-algebra to the closed *-subalgebra of $\mathcal{L}(s', s)$ generated by the sequence $(f_j \otimes f_j)_{j \in \mathbb{N}}$.

(1) \Rightarrow (2): Let E be a closed commutative *-subalgebra of $\mathcal{L}(s', s)$ isomorphic to the Fréchet *-algebra $\lambda^2(A)$. By [2, Th. 4.8] and the nuclearity of E , there is a sequence $(Q_j)_{j \in \mathbb{N}}$ of nonzero pairwise orthogonal self-adjoint projections belonging to $\mathcal{L}(s', s)$ such that $\lambda^2(A) \cong E \cong \lambda^2(\|Q_j\|_q)$ as Fréchet *-algebras. Therefore, according to [4, Prop. 4.2], there is a bijection $\sigma: \mathbb{N} \rightarrow \mathbb{N}$ such that for $P_j := Q_{\sigma(j)}$ ($j \in \mathbb{N}$), we have $A \sim (\|P_j\|_q)_{j \in \mathbb{N}, q \in \mathbb{N}_0}$.

(2) \Rightarrow (4): By [2, Prop. 4.7] and [12, Lemma 27.25], $\lambda^2(\|P_j\|_q)$ is isomorphic to the closed *-subalgebra of $\mathcal{L}(s', s)$ generated by the sequence $(P_j)_{j \in \mathbb{N}}$. Therefore, the space $\lambda^2(A)$ (which is isomorphic as a Fréchet *-algebra to $\lambda^2(\|P_j\|_q)$) is nuclear, and consequently, the matrix A satisfies condition (i). Moreover, it is easy to check that the matrix $(\|P_j\|_q)_{j \in \mathbb{N}, q \in \mathbb{N}_0}$ fulfils conditions (ii) and (iii), and thus (ii) and (iii) are also satisfied by the matrix A (apply [4, Prop. 4.2]).

(4) \Rightarrow (5): By conditions (ii) and (iii), we get

$$a_{j,r} \leq C a_{j,p} a_{j,r} \leq C a_{j, \max\{p,r\}}^2$$

and

$$a_{j,q}^2 \leq C_q a_{j,r} \leq CC_q a_{j,p} a_{j,r}$$

with appropriate quantifiers and constants.

(5) \Leftrightarrow (6): It is well known that conditions (i) and (ii) are equivalent to nuclearity and the property (DN), respectively. By [15, Prop. 3.1, Th. 5.9], $\lambda^\infty(A)$ is a biprojective Köthe algebra if and only if A satisfies condition (iii).

(5) \Leftrightarrow (7): By [12, Prop. 31.5], conditions (i) and (ii) are equivalent to the fact that $\lambda^2(A)$ is isomorphic to a closed subspace of s and, by [15, Prop. 3.1], $\lambda^\infty(A)$ is an algebra if and only if $A \prec A^2$. Moreover, by [15, Lemmas 7.7 and 7.9], conditions (i) and (iii) imply that the square root of each nonnegative element of $\lambda^\infty(A)$ belongs to $\lambda^\infty(A)$. On the other hand, if $\lambda^2(A)$ is closed under taking square root and nuclear (note that closed subspaces of s are nuclear) then, by [12, Prop. 28.16], $\lambda^2(A) \subset \lambda^2(A^2)$. Consequently, by the closed graph theorem, the identity map from $\lambda^2(A)$ to $\lambda^2(A^2)$ is continuous, and hence $A^2 \prec A$.

(5) \Rightarrow (8): Nuclearity is guaranteed by condition (i). For an index p from condition (ii) take, according to (i), an index r and a constant $C_0 > 0$ such that $\sum_{j=1}^{\infty} \frac{a_{j,p}}{a_{j,r}} < C_0$. By (ii) and (iii) (more precisely by $A \prec A^2$), there are $q_1, q_2, q_3 \in \mathbb{N}_0$ and constants $C_1, C_2, C_3 > 0$ such that

$$a_{j,r} \leq C_1 a_{j,q_1}^2 \leq C_2 a_{j,p} a_{j,q_2} \leq C_3 a_{j,p} a_{j,q_3}^2.$$

Then for $\xi \in \lambda^2(A)$ we obtain

$$\|\xi\|_{\ell_2} \leq C_3 \sum_{j=1}^{\infty} |\xi_j|^2 \frac{a_{j,p} a_{j,q_3}^2}{a_{j,r}} \leq C_3 \sum_{j=1}^{\infty} \frac{a_{j,p}}{a_{j,r}} \cdot \sup_{j \in \mathbb{N}} |\xi_j|^2 a_{j,q_3}^2 \leq C_0 C_3 |\xi|_{\lambda^2(A), q_3}^2 < \infty,$$

and thus $\|\cdot\|_{\ell_2}$ is a norm on $\lambda^2(A)$.

Moreover, the Cauchy-Schwartz inequality and condition (iii) (more precisely relation $A^2 \prec A$) imply that for all $q \in \mathbb{N}_0$ there are $r \in \mathbb{N}_0$ and $C > 0$ such that

$$\begin{aligned} |\xi|_{\lambda^2(A), q}^2 &:= \sum_{j \in \mathbb{N}} |\xi_j|^2 a_{j,q}^2 \leq C \left(\sum_{j \in \mathbb{N}} |\xi_j|^2 \right)^{1/2} \cdot \left(\sum_{j \in \mathbb{N}} |\xi_j|^2 a_{j,q}^4 \right)^{1/2} \leq C \|\xi\|_{\ell_2} \left(\sum_{j \in \mathbb{N}} |\xi_j|^2 a_{j,r}^2 \right)^{1/2} \\ &= C \|\xi\|_{\ell_2} |\xi|_{\lambda^2(A), r}, \end{aligned}$$

which shows that $\|\cdot\|_{\ell_2}$ is a dominating norm on $\lambda^2(A)$.

(8) \Rightarrow (3): By Theorem 14, there is a topological embedding $V: \lambda^2(A) \hookrightarrow s$ such that $\|V\xi\|_{\ell_2} = \|\xi\|_{\ell_2}$ for all $\xi \in \lambda^2(A)$. Set $f_j := Ve_j$ for $j \in \mathbb{N}$. Then $(f_j)_{j \in \mathbb{N}} \subset s$ is an orthonormal sequence in ℓ_2 and a Schauder basis of $\text{im } V$. Therefore, $\Phi: \text{im } V \rightarrow \lambda^2(|f_j|_q)$ defined by $\Phi f_j := e_j$ for $j \in \mathbb{N}$ is a Fréchet space isomorphism (see [12, Lemma 27.25]), and so is $\Phi \circ V: \lambda^2(A) \rightarrow \lambda^2(|f_j|_q)$. But $(\Phi \circ V)e_j = e_j$ for $j \in \mathbb{N}$, hence $\lambda^2(A) = \lambda^2(|f_j|_q)$ as Fréchet $*$ -algebras, whence $A \sim (|f_j|_q)_{j \in \mathbb{N}, q \in \mathbb{N}_0}$. \square

4. QUASI-EQUIVALENCE CONJECTURE

In the last section we shall show how commutative subalgebras of $\mathcal{L}(s', s)$ are connected with the so-called Quasi-equivalence Conjecture stating that all bases in a nuclear Fréchet space are quasi-equivalent (see [13, §8.3], [26] and references therein). Here we only consider the case (still unsolved) of closed subspaces of s , i.e. nuclear Fréchet spaces with the property (DN). Let us recall that two bases $(f_j)_{j \in \mathbb{N}}$ and $(g_j)_{j \in \mathbb{N}}$ of a Fréchet space X are called *quasi-equivalent* if there is a bijection $\sigma: \mathbb{N} \rightarrow \mathbb{N}$ and a sequence $(\lambda_j)_{j \in \mathbb{N}}$ of nonzero scalars such that the operator $T: X \rightarrow X$ defined by $Tf_j = \lambda_j g_{\sigma(j)}$ is a Fréchet space isomorphism.

Theorem 15. *Let X be an infinite-dimensional closed subspace of s . Then the following assertions are equivalent.*

- (i) For each two closed commutative $*$ -subalgebras E, F of $\mathcal{L}(s', s)$ we have: if $E \cong F \cong X$ as Fréchet spaces then $E \cong F$ as Fréchet $*$ -algebras.
- (ii) For each two basic sequences $(f_j)_{j \in \mathbb{N}}$ and $(g_j)_{j \in \mathbb{N}}$ of s which are at the same time orthonormal sequences in ℓ_2 we have: if $\lambda^2(|f_j|_q) \cong \lambda^2(|g_j|_q) \cong X$ as Fréchet spaces then there is a bijection $\sigma: \mathbb{N} \rightarrow \mathbb{N}$ such that $(|f_j|_q)_{j \in \mathbb{N}, q \in \mathbb{N}_0} \sim (|g_{\sigma(j)}|_q)_{j \in \mathbb{N}, q \in \mathbb{N}_0}$.
- (iii) For each two Köthe matrices A and B such that $A \sim A^2$ and $B \sim B^2$ we have: if $\lambda^2(A) \cong \lambda^2(B) \cong X$ as Fréchet spaces then there is a bijection $\sigma: \mathbb{N} \rightarrow \mathbb{N}$ such that $A \sim B_\sigma$.
- (iv) In X all bases are quasi-equivalent.

Proof. (i) \Leftrightarrow (ii): If E and F are closed commutative $*$ -subalgebras of $\mathcal{L}(s', s)$ then, by Theorem 4, there are basic sequences $(f_j)_{j \in \mathbb{N}}$ and $(g_j)_{j \in \mathbb{N}}$ of s which are at the same time orthonormal in ℓ_2 such that $E \cong \lambda^2(|f_j|_q)$ and $F \cong \lambda^2(|g_j|_q)$ as Fréchet $*$ -algebras. Moreover, for each such a sequence $(f_j)_{j \in \mathbb{N}}$, $\lambda^2(|f_j|_q)$ is isomorphic as a Fréchet $*$ -algebra to a closed commutative $*$ -subalgebra of $\mathcal{L}(s', s)$. Therefore, assuming $\lambda^2(|f_j|_q) \cong \lambda^2(|g_j|_q) \cong X$ as Fréchet spaces, it follows from (i) and [4, Prop. 4.2] that $(|f_j|_q)_{j \in \mathbb{N}, q \in \mathbb{N}_0} \sim (|g_{\sigma(j)}|_q)_{j \in \mathbb{N}, q \in \mathbb{N}_0}$. Conversely, if we assume that $E \cong F \cong X$ as Fréchet spaces then, by (ii) and again by [4, Prop. 4.2], we get that $E \cong F$ as Fréchet $*$ -algebras.

(i) \Leftrightarrow (iii): This is an immediate consequence of Theorem 4 and [4, Prop. 4.2].

(iii) \Rightarrow (iv): Let $(x_j)_{j \in \mathbb{N}}, (y_j)_{j \in \mathbb{N}}$ be two bases in X . Then

$$\lambda^2(A) \cong \lambda^2(B) \cong X$$

as Fréchet spaces, where

$$A := \left(\frac{|x_j|_q}{\|x_j\|_{\ell_2}} \right)_{j \in \mathbb{N}, q \in \mathbb{N}_0} \quad \text{and} \quad B := \left(\frac{|y_j|_q}{\|y_j\|_{\ell_2}} \right)_{j \in \mathbb{N}, q \in \mathbb{N}_0}.$$

Then for all $q \in \mathbb{N}_0$ we have

$$\left(\frac{|x_j|_q}{\|x_j\|_{\ell_2}} \right)^2 \leq \frac{\|x_j\|_{\ell_2} |x_j|_{2q}}{\|x_j\|_{\ell_2}^2} = \frac{|x_j|_{2q}}{\|x_j\|_{\ell_2}}$$

so $A^2 \prec A$. Moreover, since

$$\frac{|x_j|_q}{\|x_j\|_{\ell_2}} \geq 1$$

for all $q \in \mathbb{N}_0$, we have also $A \prec A^2$, and thus $A \sim A^2$. Similarly, we show that $B \sim B^2$. Therefore, by (iii), there is a bijection $\sigma: \mathbb{N} \rightarrow \mathbb{N}$ such that $A \sim B_\sigma$. This means that

$$\forall q \in \mathbb{N}_0 \exists r \in \mathbb{N}_0 \exists C > 0 \forall j \in \mathbb{N} \quad |x_j|_q \leq C \lambda_j |y_{\sigma(j)}|_r$$

and

$$\forall r' \in \mathbb{N}_0 \exists q' \in \mathbb{N}_0 \exists C' > 0 \forall j \in \mathbb{N} \quad \lambda_j |y_{\sigma(j)}|_{r'} \leq C' |x_j|_{q'},$$

where $\lambda_j := \frac{\|x_j\|_{\ell_2}}{\|y_{\sigma(j)}\|_{\ell_2}}$. Hence $x_j \mapsto \lambda_j y_{\sigma(j)}$, $j \in \mathbb{N}$, defines an isomorphism of X , i.e. the bases $(x_j)_{j \in \mathbb{N}}$ and $(y_j)_{j \in \mathbb{N}}$ are quasi-equivalent.

(iv) \Rightarrow (ii): Assume that in X all bases are quasi-equivalent. Let $(f_j)_{j \in \mathbb{N}}$ and $(g_j)_{j \in \mathbb{N}}$ be basic sequences of s which are at the same time orthonormal sequences of ℓ_2 and such that $\lambda^2(|f_j|_q) \cong \lambda^2(|g_j|_q) \cong X$ as Fréchet spaces. Let F and G be closed linear span in s of $(f_j)_{j \in \mathbb{N}}$ and $(g_j)_{j \in \mathbb{N}}$, respectively. Then $F \cong G \cong X$ as Fréchet spaces; let $T: G \rightarrow F$ be an isomorphism. Clearly, $(F_j)_{j \in \mathbb{N}}$, $F_j := Tg_j$, is a basis in F which is, by assumption quasi-equivalent to $(f_j)_{j \in \mathbb{N}}$. Consequently, there is a sequence $(\lambda_j)_{j \in \mathbb{N}}$ of nonzero scalars and a bijection $\sigma: \mathbb{N} \rightarrow \mathbb{N}$ such that $S: F \rightarrow F$ defined by sending $F_{\sigma(j)}$ to $\lambda_j^{-1} f_j$ is an isomorphism of Fréchet spaces.

Now, let $V := ST: G \rightarrow F$ and define $\|\cdot\|_V: F \rightarrow [0, \infty)$, by $\|V\xi\|_V := \|\xi\|_{\ell_2}$. Since $\|\cdot\|_{\ell_2}$ is a dominating Hilbert norm on G , $\|\cdot\|_V$ is a dominating Hilbert norm on F and, consequently,

$(\lambda_j^{-1} f_j)_{j \in \mathbb{N}} = (V g_{\sigma(j)})_{j \in \mathbb{N}}$ is an orthonormal sequence in F with respect to the Hilbert norm $\|\cdot\|_V$. In particular, $\|f_j\|_V = |\lambda_j|$ for $j \in \mathbb{N}$ and thus, $\|\cdot\|_V$ being a dominating norm on F ,

$$(13) \quad \forall q \in \mathbb{N}_0 \exists r \in \mathbb{N}_0, C > 0 \forall j \quad |f_j|_q \leq C \|f_j\|_V |f_j|_r = C |\lambda_j| |f_j|_r.$$

Moreover, since $\|\cdot\|_V$ is a continuous norm on F ,

$$\exists r_0 \in \mathbb{N}_0, C_0 > 0 \forall j \quad |\lambda_j| = \|f_j\|_V \leq C_0 |f_j|_{r_0}.$$

Now, since

$$\forall q \in \mathbb{N}_0 \exists r \in \mathbb{N}_0, C > 0 \forall j \in \mathbb{N} \quad |f_j|_q^2 \leq C |f_j|_r,$$

it follows that

$$(14) \quad \forall q \in \mathbb{N}_0 \exists r \in \mathbb{N}_0, C > 0 \forall j \quad |\lambda_j| |f_j|_q \leq C_0 |f_j|_{r_0} |f_j|_q \leq C_0 |f_j|_{\max\{r_0, q\}}^2 \leq C |f_j|_r.$$

Finally, let $W: F \rightarrow F$ be defined by $f_j \mapsto \lambda_j f_j$ for $j \in \mathbb{N}$. Then, by (13) and (14), W is an automorphism of the Fréchet space F , and thus WV is an isomorphism of Fréchet spaces which sends $g_{\sigma(j)}$ to f_j . This, clearly, implies that $(|f_j|_q)_{j \in \mathbb{N}, q \in \mathbb{N}_0} \sim (|g_{\sigma(j)}|_q)_{j \in \mathbb{N}, q \in \mathbb{N}_0}$. \square

For a monotonically increasing sequence $\alpha = (\alpha_j)_{j \in \mathbb{N}}$ in $[0, \infty)$ such that $\lim_{j \rightarrow \infty} \alpha_j = \infty$ we define the *power series space of infinite type*

$$\Lambda_\infty(\alpha) := \{(\xi_j)_{j \in \mathbb{N}} \subset \mathbb{C}^{\mathbb{N}} : \sum_{j=1}^{\infty} |\xi_j|^2 e^{2q\alpha_j} < \infty \text{ for all } q \in \mathbb{N}_0\}.$$

It appears that the space $\Lambda_\infty(\alpha)$ is nuclear if and only if $\sup_{j \in \mathbb{N}} \frac{\log j}{\alpha_j} < \infty$ (see [12, Prop. 29.6]). As a consequence, of Theorem 15, we get another proof of the following well-known result of Mityagin [13, Th. 12].

Corollary 16 (Mityagin 1961). *In nuclear power series space of infinite type $\Lambda_\infty(\alpha)$ all bases are quasi-equivalent.*

Proof. Let E, F be a closed commutative *-subalgebras of $\mathcal{L}(s', s)$ isomorphic as Fréchet spaces to $\Lambda_\infty(\alpha)$. Then, by [4, Cor. 6.10], E and F are isomorphic to $\Lambda_\infty(\alpha)$ as a Fréchet *-algebra, and thus, by Theorem 15, all bases in $\Lambda_\infty(\alpha)$ are quasi-equivalent. \square

Since, by Corollary 8, every closed subspace of s with basis is isomorphic as a Fréchet space to some closed commutative *-subalgebra of $\mathcal{L}(s', s)$, we can rewrite Theorem 15 in the following way.

Corollary 17. *The following assertions are equivalent.*

- (i) *For each two closed commutative *-subalgebras E, F of $\mathcal{L}(s', s)$ we have: if $E \cong F$ as Fréchet spaces then $E \cong F$ as Fréchet *-algebras.*
- (ii) *For each two basic sequences $(f_j)_{k \in \mathbb{N}}$ and $(g_j)_{k \in \mathbb{N}}$ of s which are at the same time orthonormal sequences in ℓ_2 we have: if $\lambda^2(|f_j|_q) \cong \lambda^2(|g_j|_q)$ as Fréchet spaces then there is a bijection $\sigma: \mathbb{N} \rightarrow \mathbb{N}$ such that $(|f_j|_q)_{j \in \mathbb{N}, q \in \mathbb{N}_0} \sim (|g_{\sigma(j)}|_q)_{j \in \mathbb{N}, q \in \mathbb{N}_0}$.*
- (iii) *For each two Köthe matrices A and B such that $A \sim A^2$ and $B \sim B^2$ we have: if $\lambda^2(A) \cong \lambda^2(B)$ as Fréchet spaces then there is a bijection $\sigma: \mathbb{N} \rightarrow \mathbb{N}$ such that $A \sim B_\sigma$.*
- (iv) *In each closed subspace of s all bases are quasi-equivalent.*

Acknowledgements. I cordially thank Leonhard Frerick for many valuable suggestions and encouraging me to solve the problems raised in the present paper.

REFERENCES

- [1] S. J. Bhatt, A. Inoue, H. Ogi, *Spectral invariance, K-theory isomorphism and an application to the differential structure of C^* -algebras*. J. Operator Theory **49** (2003), no. 2, 389–405.
- [2] T. Ciaś, *On the algebra of smooth operators*. Studia Math. **218** (2013), no. 2, 145–166.
- [3] T. Ciaś, *Algebra of smooth operators*. PhD dissertation, A. Mickiewicz University, Poznań, 2014. Available at https://repozytorium.amu.edu.pl/jspui/bitstream/10593/10958/1/phd_thesis_TCias.pdf
- [4] T. Ciaś, *Commutative subalgebras of the algebra of smooth operators*. Monatsh. Math. **179** (2016), no. 4, 1–23.
- [5] L. Crone, W.B. Robinson, *Every nuclear Fréchet space with a regular basis has the quasi-equivalence property*. Studia Math. **52** (1974/75), 203–207.
- [6] J. Cuntz, *Bivariante K-Theorie für lokalkonvexe Algebren und der Chern-Connes-Charakter*. Doc. Math. **2** (1997), 139–182.
- [7] J. Cuntz, *Cyclic theory and the bivariant Chern-Connes character*. Noncommutative geometry, Lecture Notes in Math., 1831, Springer, Berlin, 2004, 73–135.
- [8] P. Domański, *Algebra of smooth operators*. Unpublished note available at www.staff.amu.edu.pl/~domanski/salgebra1.pdf.
- [9] G. A. Elliot, T. Natsume, R. Nest, *Cyclic cohomology for one-parameter smooth crossed products*. Acta Math. **160** (1998), 285–305.
- [10] H. Glöckner, B. Langkamp, *Topological algebras of rapidly decreasing matrices and generalizations*. Topology Appl. **159** (2012), no. 9, 2420–2422.
- [11] A.Yu. Helemskii, *The Homology of Banach and Topological Algebras*. Mathematics and its Applications (Soviet Series), **41**. Kluwer Academic Publishers, Dordrecht, 1989.
- [12] R. Meise, D. Vogt, *Introduction to functional analysis*. Oxford University Press, New York 1997.
- [13] B.S. Mityagin, *Approximate dimension and bases in nuclear spaces*. (Russian) Uspehi Mat. Nauk **16** (1961), no. 4 (100), 63–132.
- [14] N.C. Phillips, *K-theory for Fréchet algebras*. Internat. J. Math. **2** (1991), no. 1, 77–129.
- [15] A. Yu. Pirkovskii, *Homological dimensions and approximate contractibility for Köthe algebras*. Banach algebras 2009, 261–278, Banach Center Publ., **91**, Polish Acad. Sci. Inst. Math., Warsaw, 2010.
- [16] K. Piszczyk, *One-sided ideals of the non-commutative Schwartz space*. Monatsh. Math. **178** (2015), no.4, 599–610.
- [17] K. Piszczyk, *A Jordan-like decomposition in the noncommutative Schwartz space*. Bull. Aust. Math. Soc. **91** (2015), no. 2, 322–330.
- [18] K. Piszczyk, *Automatic continuity and amenability in the non-commutative Schwartz space*. J. Math. Anal. Appl. **432** (2015), no. 2, 954–964.
- [19] K. Piszczyk, *Corrigendum to “Automatic continuity and amenability in the non-commutative Schwartz space”* [J. Math. Anal. Appl. **432** (2015), no. 2, 954–964]. J. Math. Anal. Appl. **435** (2016), no. 1, 1015–1016.
- [20] H. H. Schaefer, *Topological vector spaces*. Third printing corrected. Graduate Texts in Mathematics, Vol. 3. Springer-Verlag, New York-Berlin, 1971.
- [21] D. Vogt, *Charakterisierung der Unterräume von s* . Math. Z. **155** (1977), 109–117.
- [22] D. Vogt, *Subspaces and quotient spaces of (s)* . Functional analysis: surveys and recent results (Proc. Conf., Paderborn, 1976), North-Holland Math. Studies, Vol. **27**; Notas de Mat., No. 63, North-Holland, Amsterdam, 1977, 167–187.
- [23] D. Vogt *Eine Charakterisierung der Potenzreihenräume von endlichem Typ und ihre Folgerungen*. Manuscripta Math. **37** (1982), no. 3, 269–301.
- [24] D. Vogt, *On the functors $\text{Ext}^1(E, F)$ for Fréchet spaces*. Studia Math. **85** (1987), no. 2, 163–197.
- [25] D. Vogt, *Unitary endomorphisms of power series spaces*. Math. Forum, Vol. 7, Studies on Mathematical Analysis, Vladikavkaz, (2013), 220–239.
- [26] N. Zobin, *Some remarks on quasi-equivalence of bases in Fréchet spaces*. Linear Algebra Appl. **307** (2000), no. 1-3, 47–67.

T. Ciaś
 Faculty of Mathematics and Computer Science
 Adam Mickiewicz University in Poznań
 Umultowska 87
 61-614 Poznań, POLAND
 e-mail: tcias@amu.edu.pl