

PHASE TRANSITIONS OF COMPOSITION SCHEMES: MITTAG-LEFFLER AND MIXED POISSON DISTRIBUTIONS

CYRIL BANDERIER, MARKUS KUBA, AND MICHAEL WALLNER

Multitudinous combinatorial structures are counted by generating functions satisfying a composition scheme $F(z) = G(H(z))$. The corresponding asymptotic analysis becomes challenging when this scheme is critical (i.e., G and H are simultaneously singular). The singular exponents appearing in the Puiseux expansions of G and H then dictate the asymptotics.

In this work, we first complement results of Flajolet et al. for a full family of singular exponents of G and H . Motivated by many examples (random mappings, planar maps, directed lattice paths), we consider a natural extension of this scheme, namely $F(z, u) = G(uH(z))M(z)$. We also consider a variant of this scheme, which allows us to analyse the number of H -components of a given size in F .

These two models lead to a rich world of limit laws, where we identify the key rôle played by a new universal three-parameter law: the beta-Mittag-Leffler distribution, which is essentially the product of a beta and a Mittag-Leffler distribution. We prove (double) phase transitions, additionally involving Boltzmann and mixed Poisson distributions, with a unified explanation of the associated thresholds. We also obtain moment convergence and local limit theorems. We end with extensions of the critical composition scheme to a cycle scheme and to the multivariate case, leading to product distributions. Applications are presented for random walks, trees (supertrees of trees, increasingly labelled trees, preferential attachment trees), triangular Pólya urns, and the Chinese restaurant process.

*** This article is kindly devoted to Alois Panholzer, on the occasion of his 50th birthday. ***

CONTENTS

1	Introduction	2
2	New main results	4
2.1	Composition schemes analysed in this article	4
2.2	Phase transitions	5
2.3	Plan of the paper	6
3	Singularity analysis, stable laws, and mixed Poisson distributions	7
3.1	Singularity analysis and asymptotic expansions	7
3.2	Probability distribution melting pot	10
4	Extended composition scheme	17
5	Size-refined composition scheme	23
6	Applications and examples	26
6.1	Core size of supertrees and a confluent example	26
6.2	Root degree and branching structure in bilabelled increasing trees	29
6.3	Returns to zero: walks and bridges with drift zero	32
6.4	Initial returns in coloured bridges	33
6.5	Sign changes in walks	35
6.6	Tables in the Chinese restaurant process	37
6.7	Triangular urn models and beta-Mittag-Leffler distributions	41
7	Further extensions	43
7.1	Critical cycle scheme	43
7.2	Multivariate critical composition schemes	44
8	Conclusion	50
	Acknowledgments	51
	References	52

MSC2020 subject classifications: Primary 60C05; secondary 60E07, 60G50, 05A15.

Keywords and phrases: Mixed Poisson distributions, Mittag-Leffler distributions, stable laws, Boltzmann distributions, analytic combinatorics, generating functions, singularity analysis, critical composition schemes.

1. Introduction. Many combinatorial structures are an assemblage of more basic building blocks, and this situation is ubiquitous in many different fields, such as combinatorics, probability theory, and statistical mechanics. It appears for example in permutations, random walks, random mappings, random forests, parking functions, Pólya–Eggenberger urn models, (Bienaymé)–Galton–Watson processes, destruction procedures in simply generated trees, inversions in labelled tree families, generalized plane-oriented recursive trees (scale-free trees), set or integer partitions with some constraints, sequences of words, tilings, different families of graphs or maps, etc.; see, e.g., [5–7, 11–13, 28, 29, 37, 40–42, 49, 83, 99, 100, 102, 112, 118]. In the language of generating functions, one then has a functional composition scheme such as

$$F(z) = G(H(z)).$$

Let us illustrate this composition scheme with some examples (each of them being in fact the starting point of many theorems in the literature): A random forest is a set of random trees; a permutation is a set of cycles; a bridge (a random walk on \mathbb{Z}) is a sequence of arches; functional mappings are cycles of Cayley trees; supertrees are trees in which each leaf is replaced by another family of trees; an integer partition is a sequence of parts; the factorization of a polynomial in a finite field is a multiset of irreducible factors; following the work of Tutte, several important families of planar maps can be seen as a “simple core” in which each node is replaced by some “simple map”, etc. The reader can find many other examples illustrating the universality of the scheme $F(z) = G(H(z))$ in the wonderful book by Flajolet and Sedgewick on analytic combinatorics [40]. Structurally, this composition scheme is at the heart of many fascinating phase transition phenomena (analytically corresponding, e.g., to coalescing saddle points or to confluence of singularities). More precisely, let $G(z) = \sum_{n \geq 0} g_n z^n$ and $H(z) = \sum_{n \geq 0} h_n z^n$ be analytic functions at the origin with nonnegative coefficients and $H(0) = 0$. Let ρ_G and ρ_H be the radii of convergence of $G(z)$ and $H(z)$, respectively. Then, following [6, 40], we focus on *critical composition schemes*.

DEFINITION 1.1 (Critical composition scheme). The composition scheme $F(z) = G(H(z))$ is *critical* if it satisfies $H(\rho_H) = \rho_G$.

In other words, $G(z)$ and $H(z)$ are concomitantly singular. We will assume throughout this work that we are always in the critical case (the asymptotic analysis is straightforward otherwise). Note that this terminology is a generalization of the notion of critical/supercritical/subcritical Galton–Watson processes, initially popularized by Harris for neutron branching processes [52].

Often, $G(z)$ and $H(z)$ are the counting series of certain combinatorial families \mathcal{G} and \mathcal{H} such that $\mathcal{F} = \mathcal{G}(\mathcal{H})$. We refer to the first part of [40] for a more detailed presentation of this combinatorial approach, starting from the so-called atoms, and then assembling them into more elaborate structured blocks via combinatorial constructors. Some important subclasses of such structures were also subject of more probabilistic approaches; see, e.g., [3, 49, 72].

Now, our goal is to analyse probabilistic properties of critical compositions like

$$F(z, u) = G(uH(z)),$$

where u marks each occurrence of objects of \mathcal{H} . From a combinatorial perspective, $F(z, u)$ enumerates \mathcal{F} -structures of size n made of k “building blocks from \mathcal{H} ” (also simply called \mathcal{H} -components), i.e., \mathcal{G} -structures made of k blocks, where each block is then replaced by an \mathcal{H} -block (which is itself a structured set of atoms). For any combinatorial structure in \mathcal{F} , its corresponding \mathcal{G} -structure is sometimes called its *core*¹ (or “skeleton”, or “backbone”).

¹The word “core” comes from the theory of graphs and maps, where this composition scheme is natural and was, e.g., analysed in [6, 67].

A first natural question is what is the typical size of this core, i.e., what is the *typical number of \mathcal{H} -components*? Such insight helps, for example, to make many algorithms on combinatorial structures more efficient, as Knuth shows in [71] (see also [6], where this insight is used to design faster random generation algorithms). To answer this question, one considers the discrete random variable X_n associated with this core size in a uniformly chosen object of size n . Its probability mass function is obtained by extraction of coefficients:

$$\mathbb{P}\{X_n = k\} = \frac{[z^n u^k]F(z, u)}{[z^n]F(z, 1)} = g_k \frac{[z^n]H(z)^k}{f_n},$$

where $[z^n]$ denotes the extraction of coefficient operator: $[z^n]\sum_n f_n z^n = f_n$. As $H(z)$ has typically a singular expansion of the type

$$H(z) = \tau_H + c_H \left(1 - \frac{z}{\rho_H}\right)^{\lambda_H} + \dots,$$

this implies that the asymptotic behaviour of $\mathbb{P}\{X_n = k\}$ depends on the exponent λ_H (which is called the *singular exponent* of H). Actually, this exponent even plays a key rôle, as it entails four types of asymptotic behaviour for X_n :

- For $\lambda_H > 2$ the limit law is related to a Gaussian law.
- For $1 < \lambda_H < 2$ the limit law is related to a stable law of parameter λ_H (this distribution is supported on \mathbb{R} and possesses moments up to order λ_H ; e.g., for $\lambda_H = 3/2$ this gives the map-Airy distribution).
- For $0 < \lambda_H < 1$ we will show that the limit law is related to a stable law of parameter λ_H , or more precisely to a generalized Mittag-Leffler distribution (this distribution is supported on \mathbb{R}^+ and has moments of any order; e.g., for $\lambda = 1/2$, this gives the Rayleigh distribution).
- For $\lambda_H < 0$ the scheme is not critical because the function $H(z)$ diverges at $z = \rho_H$, and thus leads to a singularity of $G(H(z))$ at some $z < \rho_H$. Such a scheme is called supercritical and typically leads to a Gaussian limit law.

The case $\lambda_H < 0$ is analysed by Flajolet and Sedgewick [40], building on the seminal work of Bender [15]. The three cases $0 < \lambda_H < 1$, $1 < \lambda_H < 2$, and $\lambda_H > 2$ were partially analysed by Flajolet et al. [6], but without a precise statement for the limit laws, the right renormalizations, etc. It is partly due to the fact that the initial motivation of the authors of [6] was to analyse the core of planar maps, so they focused on the subcase $\lambda_H = \frac{3}{2}$, which corresponds to the map-Airy distribution. Thus, a more complete analysis of the composition scheme in these three regions of λ_H remained to be done.

For sure, we expected that the different possible analytic behaviours of G introduce further subcases, but we were surprised that the detailed analyses were much more challenging than expected: As we shall see, they require several new ingredients. Our identification of the limit laws involves *moment-tilted distributions*, *product distributions*, and *Boltzmann distributions* (see Section 3.2 for a formal presentation of these three types of distributions and their key properties). Therefore, the first main objective of our work is to give a complete landscape of the limit laws associated with critical composition schemes. We analyse the case $0 < \lambda_H < 1$ in this article, and the cases $1 < \lambda_H < 2$ and $\lambda_H > 2$ in our companion article [9].

Our second main objective is to explain the phase transitions observed for the *number of \mathcal{H} -components of a given size*. This builds on the work of Panholzer and the second author [83], in which they started to unify the diversity of limit laws encountered in these phase transitions under the umbrella of *mixed Poisson distributions*, and relies on the study of a size-refined composition scheme that we detail in the next section.

2. New main results.

2.1. *Composition schemes analysed in this article.* In this work we complete the analysis of composition schemes with exponent $0 < \lambda_H < 1$. We identify the corresponding limit laws as generalized Mittag-Leffler distributions and product distributions; this unifies and refines many previous studies. Motivated by models associated with directed lattice paths [5, 11–13, 118] and triangular Pólya urns [37, 64, 66], we also relax and extend the scheme by adding another component $M(z)$, which allows us to apply our results to various examples in Section 6.

First, we consider the following *extended composition scheme*

$$F(z) = G(H(z)) \cdot M(z), \quad (1)$$

for some functions $F/G/H/M$ analytic at the origin, with nonnegative coefficients. Such schemes are *critical* if $\rho_G = H(\rho_H)$ (like in Definition 1.1) and additionally satisfy $\rho_M \geq \rho_H$, where ρ_M is the radius of convergence of $M(z)$ (the analysis is straightforward if the extended composition scheme is not critical). In order to enumerate the family \mathcal{F} according to the occurrences of \mathcal{H} -components, we consider

$$F(z, u) = G(uH(z)) \cdot M(z), \quad (2)$$

which from now on we will refer to as the extended composition scheme. Equivalently, $[z^n u^k]F(z, u)$ is the number of \mathcal{F} -structures of size n having k \mathcal{H} -components. The corresponding random variable X_n has a probability mass function given by

$$\mathbb{P}\{X_n = k\} = \frac{[z^n u^k]F(z, u)}{[z^n]F(z, 1)} = g_k \frac{[z^n]H(z)^k \cdot M(z)}{f_n}. \quad (3)$$

Our first main result is Theorem 4.1, in which we give explicit expressions for the asymptotics of the factorial moments of X_n , the limit distribution of X_n (suitably normalized), and its density function. It appears that this limit distribution differs depending on some relationship between the singular exponents of $G(z)$, $H(z)$, and $M(z)$, as summarized in Table 1 (where we write $X_n \sim c_n \cdot \mathcal{D}$ when $c_n^{-1} \cdot X_n \rightarrow \mathcal{D}$ in distribution for $n \rightarrow \infty$). In addition to these convergences in distribution, we prove moment convergence for the continuous limit laws.




Singular exponent	$\lambda_M > \lambda_G \lambda_H$ (pure scheme)	$\lambda_M = \lambda_G \lambda_H$ (confluent scheme)	$\lambda_M < \lambda_G \lambda_H$ (degenerate scheme)
Limit law	continuous (gen. Mittag-Leffler ML)	linear combination (ML + \mathcal{B})	discrete (Boltzmann \mathcal{B})
Example			
	$X_n \sim Cn^{\lambda_H} \text{ML}$	$X_n \sim \text{LinComb}(n^{\lambda_H} \text{ML}, \mathcal{B})$	$\mathbb{P}\{X_n = k\} \sim \frac{g_k \rho_G^k}{G(\rho_G)}$

TABLE 1

Three asymptotic behaviours according to the singular exponents $\lambda_G/\lambda_H/\lambda_M$ of $G/H/M$: the number X_n of \mathcal{H} -components in \mathcal{F} -structures of size n in the extended scheme (1) is given by three completely different types of limit laws, depending on whether the scheme is analytically pure/confluent/degenerate (Theorems 4.1, 4.3, and 4.4).

Secondly, we consider a *size-refined composition scheme* which allows us to capture some threshold phenomenon via a bivariate generating function $F(z, v)$ ²:

$$F(z, v) = G(H(z) - (1 - v)h_j z^j) \cdot M(z), \quad j \in \mathbb{N}. \quad (4)$$

In this scheme, $[z^n v^k]F(z, v)$ is therefore the number of \mathcal{F} -structures of size n having k \mathcal{H} -components each of size j ; this is combinatorially summarized by:

$$\mathcal{F} = \mathcal{G}(\mathcal{H}_{\neq j} + v\mathcal{H}_{=j}) \times \mathcal{M}.$$

Given $n, j \in \mathbb{N}$, let $X_{n,j}$ denote the random variable counting the number of \mathcal{H} -components of size j inside \mathcal{F} -structures³ of size n . Note that the random variables $X_{n,j}$ naturally refine the distribution of the core size X_n given in (3), since

$$\sum_{j \in \mathbb{N}} X_{n,j} = X_n.$$

Formula (4) implies that one has

$$\mathbb{P}\{X_{n,j} = k\} = \frac{[z^n v^k]F(z, v)}{[z^n]F(z, 1)} = \frac{h_j^k [z^{n-jk}]G^{(k)}(H(z) - h_j z^j)M(z)}{f_n}. \quad (5)$$

Our second main result is Theorem 5.1, in which we prove that the factorial moments of $X_{n,j}$ are asymptotically of mixed Poisson type, and establish a convergence in distribution, with convergence of all moments, towards explicit limit distributions.

We extend these two main results to the composition schemes involving a logarithmic singularity in Theorem 7.1 and 7.3, leading to Mittag-Leffler distributions.

Then, our third main result is Theorem 7.4 in which we give a multivariate generalization with arbitrary many variables, leading to Dirichlet product distributions.

2.2. Phase transitions. Analogously to what can happen in physics or chemistry for some small change of temperature, pressure, or concentration, a phase transition in mathematics corresponds to a sudden non-smooth change of properties under smooth variation of the parameters. Such non-smooth changes are thus analytically reflected by a singularity of some function associated with these properties. For combinatorial structures, generating function methods were successfully used to analyse such phase transitions, e.g., for random graphs or planar maps [6, 47, 48, 67], for satisfiability problems [23], and for many other problems [40, 44, 45, 55–57].

Usually, the main problems are to locate the phase transition, to properly describe the phase transition via special functions in terms of the involved parameters, and to give intuitive explanations of the observed phenomena. Some cases even exhibit two successive phase transitions. In probability theory, such a double phase transition occurs for example with the binomial distribution $X_n \stackrel{d}{=} B(n, p)$ when p depends on n , with p bounded away from one. It indeed leads to the following trichotomy involving a continuous to discrete to degenerate phase transition: First, when both $\mathbb{E}(X_n) = p(n) \cdot n$ and the variance $\mathbb{V}(X_n)$ tend to infinity, then a central limit theorem for X_n follows. Second, if $p(n) \cdot n \rightarrow \lambda > 0$, then X_n is asymptotically Poisson distributed. Third, if $p(n) \cdot n \rightarrow 0$, then X_n degenerates to a Dirac distribution with all mass at 0.

The analysis of the size-refined composition scheme unravels, though more subtly, such a double phase transition: the random variable $X_{n,j}$ given by Equation (5) has three phases, each leading to its own limit law, visualized in Table 2.

²Note that we still use the letter F to denote the main function. The auxiliary variable is now v and no more u as we are marking a different parameter. This choice avoids more cumbersome notation and is also motivated by the fact that we always have $F(z, 1) = F(z)$.

³Flajolet and Sedgewick call the corresponding distribution the *profile* of the combinatorial object, by analogy with the profile of integer compositions; see [40, p. 169, p. 451, p. 632].




Scale	$j \ll n^{\frac{\lambda_H}{1+\lambda_H}}$	$j = \Theta\left(n^{\frac{\lambda_H}{1+\lambda_H}}\right)$	$j \gg n^{\frac{\lambda_H}{1+\lambda_H}}$
Limit law	continuous (gen. Mittag-Leffler ML)	discrete (mixed Poisson MPo(ξ ML))	discrete (Dirac)
Example			
	$X_{n,j} \sim C h_j \rho_H^j n^{\lambda_H}$ ML	$X_{n,j} \sim \text{MPo}(\xi \text{ ML})$	$\mathbb{P}\{X_{n,j} \geq 1\} \sim 0$

TABLE 2

Three consecutive régimes for the number $X_{n,j}$ of \mathcal{H} -components of size j in \mathcal{F} -structures of size n in the critical size-refined scheme (4), depending on the relation between j and n ; this double phase transition is proven in Theorem 5.1.

Table 2 also motivates the following important remark.

REMARK 2.1 (Ubiquity of the exponent 1/3). Generating functions often have a dominant singularity of the square-root type (i.e., $\lambda_H = 1/2$). This phenomenon is explained by the Drmota–Lalley–Woods theorem: Whenever $H(z)$ can be defined by a strongly connected set of polynomial equations with nonnegative coefficients, it has a singular exponent 1/2 (see, e.g., [4, 27, 40]). Accordingly, in conjunction with Table 2, this explains why one observes a threshold at $j = n^{1/3}$ in many phase transitions; see Section 6.

One pleasant consequence of our work is that it gives a unified explanation of phase transitions from continuous to discrete observed in many examples: descendants in increasing trees [75], node degrees in increasing trees [76], block sizes in k -Stirling permutations [78], stopping times in urn models [80], death processes [81], inversions in labelled tree families [102], ancestors and descendants in evolving k -tree models [103].

These case by case studies lacked a proper comprehensive and uniform description of the arising phase transitions. So, instead of treating these combinatorial structures individually, we directly study the size-refined composition scheme (4). As summarized in Table 2, we show how the phase transitions for the random variable $X_{n,j}$ depend on the growth of $j = j(n)$ with respect to the size n . We prove that the distribution of $X_{n,j(n)}$ is *continuous* for small values of j (a three-parameter generalization of the Mittag-Leffler distribution), or *discrete* for some threshold values of j (a Poisson distribution mixed with the previous Mittag-Leffler distribution), or a *Dirac distribution* for large values of j . We further exemplify these results on different processes, like the Chinese restaurant process, sign changes and returns to zero in random walks, and the branching structure of random trees.

2.3. *Plan of the paper.* In Section 3 we collect results from analytic combinatorics. We also present our basic assumptions on the generalized composition scheme and collect properties of various distributions that appear later in our main results. Section 4 is devoted to our results on the random variable X_n corresponding to the extended composition scheme (2) involving $F(z, u)$. Section 5 contains the results for the random variable $X_{n,j}$ corresponding to the size-refined composition scheme (4) involving $F(z, v)$ and exhibiting phase transitions. We also give the covariance and the correlation coefficient of X_{n,j_1}, X_{n,j_2} , observing again some phase transitions. In Section 6 we discuss various examples to which we apply our results. Finally, in Section 7, we analyse further extensions for a cycle scheme and for a multivariate critical composition scheme. We also present new examples for these two extensions.

3. Singularity analysis, stable laws, and mixed Poisson distributions. In this section, we first present a few important notions from analytic combinatorics [40] which we use to identify the radius of convergence and the singular exponents in our composition schemes. Then, we present a few results on the family of moment-tilted stable laws, based on James [59–61] and Janson [66]. We also collect properties of mixed Poisson distributions and their factorial moments [51, 83]. All of this allows us to identify in Sections 4 and 5 the distribution of the \mathcal{H} -components in our composition schemes.




FIG 1. As visually summarized by Flajolet in [36], three fundamental methods of asymptotic analysis require information on the function in different parts (shown here in red) of the complex plane. Flajolet and Odlyzko’s singularity analysis [39] offers more powerful results, but requires analyticity in a Δ -domain (tastefully also sometimes called “camembert domain” or “Pac-Man” by Flajolet himself!). This is the domain inside the blue curve, defined by $\Delta = \{z \in \mathbb{C} \text{ such that } |z| < \rho + \varepsilon \text{ and } \arg(z - \rho) > \theta\}$, for some $\varepsilon > 0$ and $0 < \theta < \pi/2$. This analyticity is agreeably typically granted for most combinatorial constructions (e.g., for the ones leading to meromorphic, algebraic-logarithmic, hypergeometric, or D -finite functions).

3.1. *Singularity analysis and asymptotic expansions.* Let $F(z) = \sum_{n \geq 0} f_n z^n$ be a function with nonnegative coefficients f_n that is analytic in a Δ -domain (see Figure 1 for this notion) with a finite radius of convergence ρ and singular expansion

$$F(z) = P\left(1 - \frac{z}{\rho}\right) + c_F \cdot \left(1 - \frac{z}{\rho}\right)^{\lambda_F} (1 + o(1)), \quad (6)$$

where $\lambda_F \in \mathbb{R} \setminus \{0, 1, 2, \dots\}$ is called the *singular exponent* (of $F(z)$ at $z = \rho$), and where $P(x) \in \mathbb{C}[x]$ is a polynomial (of degree ≥ 1 for $\lambda_F > 1$, of degree 0 for $0 < \lambda_F < 1$, and $P = 0$ for $\lambda_F < 0$). Then, by standard singularity analysis [39], if ρ is the unique singularity of $F(z)$ in $|z| \leq \rho$, the Taylor series coefficients of $F(z)$ satisfy

$$[z^n]F(z) = f_n = \frac{c_F}{\rho^n} \cdot \frac{n^{-\lambda_F-1}}{\Gamma(-\lambda_F)} \cdot (1 + o(1)). \quad (7)$$

As the f_n ’s are nonnegative, this implies the sign property

$$\text{sgn}(c_F) = \text{sgn}(\Gamma(-\lambda_F)), \quad (8)$$

i.e., due to the sign change of the gamma function at each negative integer we have $c_F < 0$ for $0 < \lambda_F < 1$, $c_F > 0$ for $\lambda_F < 0$, and $\text{sgn}(c_F) = (-1)^{\lceil \lambda_F \rceil}$ for $\lambda_F > 1$.

Note that it is in general easy to get more asymptotic terms in (6), and that singularity analysis directly translates them into more asymptotic terms in (7). What is more, if one has several dominant singularities, one just has to sum the contributions of the local expansions at each singularity to get the asymptotics of the coefficients (this standard process is well presented in [40, Chapter IV.6]; see also the rotation law in [14] for walks or trees, which involve multiple dominant singularities as soon as their offspring distribution has a periodic support).

Note that algebraic functions constitute one of the main sources of functions satisfying the conditions of the expansion (6); indeed, they admit a Puiseux expansion

$$F(z) = \sum_{k \geq k_0} c_k \cdot (1 - z/\rho)^{k/r},$$

for $k_0 \in \mathbb{Z}$ and an integer $r \geq 1$. For example, in Section 6.1 we will encounter the Catalan generating function $1/2 - \sqrt{1-4z}/2$, for which one has $\rho = 1/4$, $P(x) = 1/2$, $k_0 = 0$, and $r = 2$. This Catalan example is pleasantly simple, and thus obviously not generic, as here the Puiseux expansion contains only two terms. In full generality it can involve an infinite number of terms whose sum is converging. Let us now present this general case.

LEMMA 3.1 (Singular expansion). *Let $F(z)$ be a power series with nonnegative coefficients satisfying (6). Then $F(z)$ has the following singular expansion*

$$F(z) = \begin{cases} c_F \left(1 - \frac{z}{\rho}\right)^{\lambda_F} (1 + o(1)) & \text{if } \lambda_F < 0, \\ \tau_F + c_F \left(1 - \frac{z}{\rho}\right)^{\lambda_F} (1 + o(1)) & \text{if } 0 < \lambda_F < 1, \\ \tau_F + \sum_{i=1}^{\lfloor \lambda_F \rfloor} p_i \left(1 - \frac{z}{\rho}\right)^i + c_F \left(1 - \frac{z}{\rho}\right)^{\lambda_F} (1 + o(1)) & \text{if } \lambda_F > 1, \end{cases}$$

where $\tau_F = F(\rho) > 0$ for $\lambda_F > 0$ and $p_1 = -\rho F'(\rho) < 0$ for $\lambda_F > 1$.

PROOF. First, if $\lambda_F < 0$, then the lowest order of the Puiseux expansion is λ_F . Second, for $\lambda_F > 0$, we rewrite (6) into

$$F(z) = \sum_{i=0}^k p_i (1 - z/\rho)^i + c_F (1 - z/\rho)^{\lambda_F} + \dots$$

Then, we have $p_0 = P(0) = F(\rho)$ which we define to be τ_F . We get $\tau_F > 0$ as it is an infinite convergent sum of nonnegative not-all-zero terms. Next, observe that the nonnegative coefficients of $F(z)$ imply that $F'(\rho) > 0$. Thus, taking the derivative in the expansion of $F(z)$ we get $\lim_{z \rightarrow \rho} F'(z) = +\infty$ for $0 < \lambda_F < 1$ and $p_1 = -\rho F'(\rho) \neq 0$ for $\lambda_F > 1$. \square

We now consider the critical scheme $F(z) = G(H(z))M(z)$ where we assume that each of the functions $G(z)$, $H(z)$, and $M(z)$ has a finite radius of convergence with a unique dominant singularity (i.e., the one of smallest modulus). By Pringsheim's theorem [40, p. 240] applied to each of these functions, the nonnegativity of its coefficients implies that its dominant singularity lies on the positive real axis and corresponds therefore to its radius of convergence denoted by ρ_G , ρ_H , and ρ_M , respectively.

Note that if $M(z)$ has an infinite radius of convergence (denoted by $\rho_M = +\infty$) or if $\rho_M \neq \rho_H$, then the asymptotics are easily obtained via

$$[z^n]F(z) \sim \begin{cases} G(H(\rho_M))[z^n]M(z) & \text{if } \rho_M < \rho_H, \\ M(\rho_H)[z^n]G(H(z)) & \text{if } \rho_M > \rho_H. \end{cases} \quad (9a)$$

$$[z^n]F(z) \sim \begin{cases} G(H(\rho_M))[z^n]M(z) & \text{if } \rho_M < \rho_H, \\ M(\rho_H)[z^n]G(H(z)) & \text{if } \rho_M > \rho_H. \end{cases} \quad (9b)$$

Now, as in (6), we define for each function:

- the singular exponents λ_G , λ_H , and λ_M ,
- the constant terms τ_G , τ_H , and τ_M ,
- and the singular coefficients c_G , c_H , and c_M .

Thus, thanks to Equations (9a) and (9b), we can now focus (without loss of generality, or rather “without loss of difficulty!”) on the case $\rho_M = \rho_H$ which is more involved as here $G(z)$, $H(z)$, and $M(z)$ are *all contributing* to the asymptotics in a nontrivial way. Then, one gets different régimes (depending on λ_H) for the asymptotics of the coefficient of $F(z)$. In this article we focus on the range $0 < \lambda_H < 1$, while we treat the other range $\lambda_H > 1$ in a companion article [9].

Note that, as our work extends to some interesting combinatorial cases where $M(z)$ has a radius of convergence $\rho_M > \rho_H$, we will also encompass this case, for which it is then convenient to set $\lambda_M = +\infty$ (the singular exponent of $M(z)$ at $z = \rho_H$ is infinite whenever M is analytical there). The case $\lambda_M = +\infty$ is thus archetypal of cases where $M(z)$ only affects the asymptotics of f_n by a multiplicative constant like in Equation (9b).

We can now express the singular exponent of F in terms of those of $G/H/M$.

LEMMA 3.2. *Let $F(z) = G(H(z))M(z)$ be a critical composition scheme that is singular at ρ_H . Then, the singular exponent λ_H of $H(z)$ satisfies $\lambda_H > 0$. Moreover, for the range $0 < \lambda_H < 1$, the singular exponent λ_F of $F(z)$ satisfies*

$$\lambda_F = \min(\lambda_G \lambda_H, \lambda_H, \lambda_M, \lambda_G \lambda_H + \lambda_M).$$

For $\lambda_H > 1$, the singular exponent λ_F of $F(z)$ satisfies

$$\lambda_F = \min(\lambda_G, \lambda_H, \lambda_M, \lambda_G + \lambda_M).$$

PROOF. The claim $\lambda_H > 0$ follows from $H(\rho_H) = \rho_G \in (0, \infty)$ as one would have $H(\rho_H) = +\infty$ if $H(z)$ had a pole (or any algebraic singularity of negative singular exponent) at $z = \rho_H$.

Now, we plug the singular expansions from Lemma 3.1 at $z = \rho_H$ for $G(z)$, $H(z)$, and $M(z)$ into $F(z) = G(H(z))M(z)$. When $0 < \lambda_H < 1$ we get the following expansions (in which we omit the terms not contributing to the first-order asymptotics):

$$F(z) = \begin{cases} c_M c_G \left(\frac{-c_H}{\rho_G} \right)^{\lambda_G} \left(1 - \frac{z}{\rho_H} \right)^{\lambda_G \lambda_H + \lambda_M} + \dots & \text{if } \lambda_G < 0, \lambda_M < 0, \quad (10a) \\ \tau_M c_G \left(\frac{-c_H}{\rho_G} \right)^{\lambda_G} \left(1 - \frac{z}{\rho_H} \right)^{\lambda_G \lambda_H} + \dots & \text{if } \lambda_G < 0, \lambda_M > 0, \quad (10b) \\ \tau_M c_G \left(\frac{-c_H}{\rho_G} \right)^{\lambda_G} \left(1 - \frac{z}{\rho_H} \right)^{\lambda_G \lambda_H} + c_M \tau_G \left(1 - \frac{z}{\rho_H} \right)^{\lambda_M} + \dots & \text{if } 0 < \lambda_G < 1, \quad (10c) \\ G'(\rho_G) \tau_M c_H \left(1 - \frac{z}{\rho_H} \right)^{\lambda_H} + c_M \tau_G \left(1 - \frac{z}{\rho_H} \right)^{\lambda_M} + \dots & \text{if } \lambda_G > 1, \quad (10d) \end{cases}$$

where, if $\lambda_M = +\infty$, $\tau_M = c_M = M(\rho_H)$ and $\left(1 - \frac{z}{\rho_H} \right)^{\lambda_M} = 0$.

When $\lambda_H > 1$ the linear term of $H(z)$ is nonzero and therefore λ_G plays the rôle of $\lambda_G \lambda_H$ in the above expansions. So, for $\lambda_H > 1$, one gets (omitting the constants for simplicity):

$$F(z) = \begin{cases} C_1 \left(1 - \frac{z}{\rho_H} \right)^{\lambda_G + \lambda_M} + \dots & \text{if } \lambda_G < 0, \lambda_M < 0, \\ C_2 \left(1 - \frac{z}{\rho_H} \right)^{\lambda_G} + \dots & \text{if } \lambda_G < 0, \lambda_M > 0, \\ C_{3a} \left(1 - \frac{z}{\rho_H} \right)^{\lambda_G} + C_{3b} \left(1 - \frac{z}{\rho_H} \right)^{\lambda_M} + \dots & \text{if } 0 < \lambda_G < 1, \\ C_{4a} \left(1 - \frac{z}{\rho_H} \right)^{\lambda_G} + C_{4b} \left(1 - \frac{z}{\rho_H} \right)^{\lambda_M} + C_{4c} \left(1 - \frac{z}{\rho_H} \right)^{\lambda_H} + \dots & \text{if } \lambda_G > 1. \end{cases}$$

Finally, the singular exponent λ_F is equal to the minimal exponent in each of these Puiseux expansions. \square

This lemma motivates the following definition.




FIG 2. The three different régimes (pure, confluent, degenerate) for extended or size-refined composition schemes: The Puiseux expansions of $G/H/M$ go into resonance (or not), thus leading to these three cases.

DEFINITION 3.3 (Pure/confluent/degenerate composition schemes). Consider an extended or size-refined composition scheme (1) or (4) with a unique dominant singularity $\rho_F = \rho_H$, and with $0 < \lambda_H < 1$. It is either analytically

- *pure* if $\begin{cases} \lambda_G < 0 \text{ or} \\ 0 < \lambda_G < 1 \text{ and } \lambda_M > \min(\lambda_G \lambda_H, \lambda_H); \end{cases}$
- *confluent* if $0 < \lambda_G < 1 \text{ and } \lambda_M = \min(\lambda_G \lambda_H, \lambda_H);$
- *degenerate* if $\begin{cases} \lambda_G > 1 \text{ or} \\ 0 < \lambda_G < 1 \text{ and } \lambda_M < \min(\lambda_G \lambda_H, \lambda_H). \end{cases}$

REMARK 3.4 (Dominant asymptotics and degenerate/confluent/pure composition schemes). Consider an extended or size-refined composition scheme (1) or (4) with a unique dominant singularity $\rho_F = \rho_H$.

- In the subcases (10a), (10b), and (10c) with $\lambda_M > \lambda_G \lambda_H$, one gets a *pure* composition scheme. This adjective stresses the fact that the singular exponent of F is expressible, as expected, in terms of λ_G and λ_H .
- In the subcase (10c) with $\lambda_M = \lambda_G \lambda_H$, one gets a *confluent* composition scheme.
- In the subcases (10c) with $\lambda_M < \lambda_G \lambda_H$ and (10d) the first-order asymptotics are dictated by $M(z)$ or $H(z)$ only, leading to a *degenerate* composition scheme. This adjective stresses the fact that the singular exponent of F is not expressible in terms of λ_G .

We characterize the distributions associated with critical composition schemes in the analytically pure/confluent/degenerate cases in Theorems 4.1, 4.3, and 4.4. We now present some probabilistic results on the distributions which will appear in these theorems.

3.2. Probability distribution melting pot. First, we discuss properties of tilted probability distributions and study in particular positive stable distributions. Then, we collect properties of a family of discrete distributions called mixed Poisson distributions [51, 70, 83, 120], and we will end this section with a brief introduction to Boltzmann distributions.

For a random variable X of density $f(x)$, the tilt of $f(x)$ by a nonnegative integrable function $g(x)$ is the following density

$$\frac{g(x)}{\mathbb{E}(g(X))} \cdot f(x).$$

An important class of tilted densities are the *polynomially tilted densities*, where one tilts by a polynomial $g(x) = x^c$ (with c being any real value such that $\mathbb{E}(X^c)$ is well defined). We then use the notation

$$\text{tilt}_c(f(x)) = \frac{x^c}{\mathbb{E}(X^c)} \cdot f(x).$$

Such tilted densities occur in many places: in the degree distribution in preferential attachment trees [60, 61], in Lambert-type laws [59], in triangular urn schemes [64, 66], in node-degrees in plane-oriented recursive trees [76], and in table sizes in the Chinese restaurant process [2, 83, 108, 109]. Note that many classes of distributions like the beta distribution, generalized gamma distribution [10], the F -distribution, the beta-prime distribution, and distributions with gamma-type moments [66] are closed under the tilting operation.

The following lemma shows that the operator tilt_c admits in fact several equivalent definitions using the density, the moments, or the Laplace transform (see also [66, Remark 2.11]).

LEMMA 3.5 (Polynomially tilted density functions and moment shifts). *Consider a random variable X with moment sequence $(\mu_s)_{s \geq 0}$ and density $f(x)$ with support $[0, \infty)$. Now consider a random variable X_c with $c \in \mathbb{N}$, having a distribution uniquely determined by its moments. Then the following properties are equivalent:*

1. *Tilted density: X_c is a random variable with density $f_c(x) = \frac{x^c}{\mu_c} \cdot f(x)$.*
2. *Shifted moments: X_c is a random variable with moments $\mathbb{E}(X_c^s) = \frac{\mu_{s+c}}{\mu_c}$.*
3. *Differentiated moment generating function: X_c is such that*

$$\mathbb{E}(e^{tX_c}) = \frac{1}{\mu_c} \frac{d^c}{dt^c} \mathbb{E}(e^{tX}).$$

REMARK 3.6 (Tilt with $c \in \mathbb{R}$). For the properties (1) and (2) of Lemma 3.5, it is possible to extend their equivalence to $c \in \mathbb{R}$, assuming that the corresponding moments exist. More generally, the equivalence between properties (1) and (2) stays valid for any random variable with density $f(x)$ such that only moments μ_1, \dots, μ_n exist up to a certain value $n \geq 1$.

REMARK 3.7 (Densities with support \mathbb{R}). In Lemma 3.5, if c is even, then one can drop the restriction that the support of $f(x)$ is in $[0, \infty)$.

REMARK 3.8 (The tilt operator for densities/moments/random variables). This lemma justifies a slight abuse of notation: Starting with the densities of X and X_c linked by $\text{tilt}_c(f(x)) = f_c(x)$, the operator tilt_c is also used to denote the corresponding tilted random variable $\text{tilt}_c(X) := X_c$ and the corresponding tilted moments $\text{tilt}_c(\mu_s) := \frac{\mu_{s+c}}{\mu_c}$.

PROOF OF LEMMA 3.5. For (1) \Rightarrow (2), first observe that $f_c(x)$ is indeed a density: One has $f_c(x) \geq 0$ on $[0, \infty)$ and $\int_0^\infty f_c(x) dx = \frac{\mu_c}{\mu_c} = 1$. Then, one checks that

$$\mathbb{E}(X_c^s) = \int_0^\infty x^s f_c(x) dx = \int_0^\infty x^{s+c} \frac{f(x)}{\mu_c} dx = \frac{\mu_{s+c}}{\mu_c}.$$

The fact that X_c is uniquely determined by its moments then implies (2) \Rightarrow (1).

For (2) \Leftrightarrow (3), observe that

$$\frac{d^c}{dt^c} \mathbb{E}(e^{tX}) = \frac{d^c}{dt^c} \sum_{s \geq 0} \frac{\mu_s t^s}{s!} = \sum_{s \geq c} \frac{\mu_s t^{s-c}}{(s-c)!} = \sum_{s \geq 0} \frac{\mu_{s+c} t^s}{s!},$$

and, on the other hand, (1) and (2) imply

$$\mathbb{E}(e^{tX_c}) = \int_0^\infty f_c(x) e^{tx} dx = \frac{1}{\mu_c} \sum_{s \geq 0} \frac{t^s}{s!} \int_0^\infty x^{s+c} f(x) dx = \frac{1}{\mu_c} \sum_{s \geq 0} \frac{t^s}{s!} \mu_{s+c},$$

which proves the claim. □

Let us now introduce *positive stable laws* (also called *one-sided stable laws*, as their density has support $(0, +\infty)$).

DEFINITION 3.9 (Positive stable laws and their negative powers). We say that a positive random variable S_α follows a stable law of parameter $\alpha \in (0, 1)$ if its Laplace transform is $\mathbb{E}(e^{-tS_\alpha}) = e^{-t^\alpha}$ (see [117] for a general presentation involving skewness, scale, and location parameters; they are respectively always 0, 1, and 0 in our work). The density of S_α is⁴

$$f_{S_\alpha}(x) = \sum_{n=1}^{\infty} \frac{(-1)^n}{n! \Gamma(-n\alpha)} x^{-n\alpha-1} \quad (12)$$

$$= \frac{1}{\pi} \sum_{n=1}^{\infty} (-1)^{n+1} \frac{\Gamma(n\alpha + 1) \sin(\pi n\alpha)}{n!} x^{-n\alpha-1} \quad (13)$$

$$= \frac{1}{\pi} \frac{\alpha}{1-\alpha} \int_0^\pi \frac{K(\phi)}{x^{1/(1-\alpha)}} \exp\left(-\frac{K(\phi)}{x^{\alpha/(1-\alpha)}}\right) d\phi, \quad (14)$$

where

$$K(\phi) = \left(\frac{\sin(\alpha\phi)}{\sin(\phi)}\right)^{1/(1-\alpha)} \frac{\sin((1-\alpha)\phi)}{\sin(\alpha\phi)}. \quad (15)$$

Formula (12) was first obtained by Humbert [53], and then rigorously proven by Pollard [111] (see also Feller [33, Chapter XVII.6, Lemma 1], with the parameter $\gamma = -\alpha$ therein). Formula (14) is due, up to a typo that we corrected here, to Ibragimov and Chernin [58].

Now, let $\beta > 0$ and define $S_{\alpha,\beta} = (S_\alpha)^{-\beta}$. Since $\mathbb{P}\{S_{\alpha,\beta} \leq x\} = 1 - \mathbb{P}\{S_\alpha < x^{-1/\beta}\}$, we directly obtain from (12) the density of $S_{\alpha,\beta}$ on its support $(0, +\infty)$:

$$f_{S_{\alpha,\beta}}(x) = \frac{1}{\beta} \sum_{n=1}^{\infty} \frac{(-1)^n}{n! \Gamma(-n\alpha)} x^{n\alpha/\beta-1} = \frac{x^{-1/\beta-1}}{\beta} f_{S_\alpha}(x^{-1/\beta}). \quad (16)$$

Its moments are given by (see, e.g., Janson's survey on moments of Gamma type [66]):

$$\mathbb{E}(S_{\alpha,\beta}^s) = \frac{\Gamma(\frac{s\beta}{\alpha} + 1)}{\Gamma(s\beta + 1)}, \quad s > -\frac{\alpha}{\beta}.$$

We will encounter composition schemes leading to powers of stable laws, an important subcase of it being the Mittag-Leffler distribution.⁵

EXAMPLE 3.10 (Mittag-Leffler distribution). We say that a random variable M_α follows a *Mittag-Leffler distribution* $ML(\alpha)$ if $M_\alpha \stackrel{d}{=} S_{\alpha,\alpha}$. Its moment generating function $\mathbb{E}(e^{xM_\alpha})$ is the Mittag-Leffler function $E_\alpha(x) = \sum_{k \geq 0} \frac{x^k}{\Gamma(1+\alpha k)}$. An important special case is $M_{\frac{1}{2}}$, the *half-normal distribution* $|\mathcal{N}(0, \sigma^2)|$ with $\sigma = \sqrt{2}$; see Example 3.17 hereafter.

For $c > -\frac{\alpha}{\beta}$ consider the moment-tilted random variable $X_c = \text{tilt}_c(S_{\alpha,\beta})$. Then,

$$\mathbb{E}(X_c^s) = \frac{\Gamma(\frac{(s+c)\beta}{\alpha} + 1)}{\Gamma((s+c)\beta + 1)} \frac{\Gamma(\beta c + 1)}{\Gamma(\frac{c\beta}{\alpha} + 1)} = \frac{\Gamma(\frac{(s+c)\beta}{\alpha})}{\Gamma((s+c)\beta)} \frac{\Gamma(\beta c)}{\Gamma(\frac{c\beta}{\alpha})}; \quad (17)$$

see James [59–61]. By (16) and Lemma 3.5, the density of X_c is given by

$$f_{X_c}(x) = \frac{\Gamma(\beta c + 1)}{\beta \Gamma(\frac{c\beta}{\alpha} + 1)} \sum_{n=1}^{\infty} \frac{(-1)^n}{n! \Gamma(-n\alpha)} x^{n\alpha/\beta+c-1}.$$

E.g., $\text{tilt}_1(M_{\frac{1}{2}})$ is the *Rayleigh distribution* of parameter $\sqrt{2}$; see Example 3.18 hereafter. ■

⁴Throughout this article, we use that, by analytical continuation, $1/\Gamma(m) = 0$ whenever m is an integer ≤ 0 .

⁵In the literature, there are unfortunately two distinct distributions which are called *Mittag-Leffler distribution*. Both of them are defined in terms of the function $E_\alpha(x)$ introduced in 1903 by Mittag-Leffler [94, 95]. The first distribution (which we use in this article) was popularized by Feller [32] (with a slight change of variable) and by Darling and Kac [22] for the study of the local time of Markov processes. It has an exponentially bounded tail. The second one, which has a heavy tail, was introduced by Pillai [107].

Some probabilistic processes are related to a two-parameter generalization of this Mittag-Leffler distribution (see, e.g., [49, 61, 109]). We will establish in the next section that some critical composition schemes lead to this generalized Mittag-Leffler distribution.

DEFINITION 3.11 (Generalized Mittag-Leffler distribution). For $\alpha \in (0, 1)$ and $\theta > -\alpha$, we say that a random variable X follows a *generalized Mittag-Leffler distribution* $\text{ML}(\alpha, \theta)$ if $X \stackrel{d}{=} (\text{tilt}_{-\theta}(S_\alpha))^{-\alpha}$; see James [61]. This distribution is uniquely defined by its moments

$$\mathbb{E}(X^s) = \frac{\Gamma\left(s + \frac{\theta}{\alpha} + 1\right) \Gamma(\theta + 1)}{\Gamma(\alpha s + \theta + 1) \Gamma\left(\frac{\theta}{\alpha} + 1\right)} = \frac{\Gamma\left(s + \frac{\theta}{\alpha}\right) \Gamma(\theta)}{\Gamma(\alpha s + \theta) \Gamma\left(\frac{\theta}{\alpha}\right)}. \quad (18)$$

Comparing these moments with (17) we directly get that for $\beta = \alpha$ and $c = \theta/\alpha$ we have

$$\text{tilt}_{\theta/\alpha}(\text{ML}(\alpha)) \stackrel{d}{=} \text{ML}(\alpha, \theta), \quad \text{i.e.,} \quad \text{tilt}_{\theta/\alpha}(S_\alpha^{-\alpha}) = (\text{tilt}_{-\theta}(S_\alpha))^{-\alpha}. \quad (19)$$

In other words, the permutation of the tilt and the power creates a change of the tilt parameter.

Next, we discuss product distributions. First, we recall properties of the beta distribution.

DEFINITION 3.12 (Beta distribution). A *beta-distributed* random variable $B \stackrel{d}{=} \text{Beta}(\alpha, \beta)$ with parameters $\alpha, \beta > 0$ has a probability density function defined on $(0, 1)$ by

$$f(x) = \frac{\Gamma(\alpha + \beta)}{\Gamma(\alpha)\Gamma(\beta)} x^{\alpha-1} (1-x)^{\beta-1}.$$

The moments of B are given by

$$\mathbb{E}(B^s) = \frac{\Gamma(s + \alpha)\Gamma(\alpha + \beta)}{\Gamma(s + \alpha + \beta)\Gamma(\alpha)}, \quad s > 0, \quad (20)$$

and the beta distribution is uniquely determined by the sequence of its moments. Furthermore, let the reader be convinced of the convenient convention $\text{Beta}(\alpha, 0) \stackrel{d}{=} 1$.

We now have all the ingredients to present the main properties of the distribution which will play a key rôle in the next sections, namely, the beta-Mittag-Leffler distribution.

DEFINITION 3.13 (Beta-Mittag-Leffler distribution). We define the *beta-Mittag-Leffler distribution* $\text{BML}(\alpha, \theta, \beta)$ as the distribution of the following product of independent random variables

$$Z \stackrel{d}{=} Y \cdot B^\alpha$$

where $Y \stackrel{d}{=} \text{tilt}_{\theta/\alpha}(S_{\alpha, \alpha}) \stackrel{d}{=} \text{ML}(\alpha, \theta)$ and $B \stackrel{d}{=} \text{Beta}(\theta, \beta)$ are respectively distributed like a Mittag-Leffler and a beta distribution, with $0 < \alpha < 1$, $\theta > 0$, and $\beta \geq 0$.

LEMMA 3.14. *The beta-Mittag-Leffler distribution $\text{BML}(\alpha, \theta, \beta)$ has the following moments of order s (for $s > 0$):*

$$\mathbb{E}(Z^s) = \frac{\Gamma\left(s + \frac{\theta}{\alpha}\right) \Gamma(\theta + \beta)}{\Gamma(\alpha s + \theta + \beta) \Gamma\left(\frac{\theta}{\alpha}\right)}. \quad (21)$$

One has the following identity

$$Z \stackrel{d}{=} \text{ML}(\alpha, \theta) \text{Beta}(\theta, \beta)^\alpha \stackrel{d}{=} \text{ML}(\alpha, \theta + \beta) \text{Beta}\left(\frac{\theta}{\alpha}, \frac{\beta}{\alpha}\right). \quad (22)$$

PROOF. Due to the independence of the random variables we have $\mathbb{E}(Z^s) = \mathbb{E}(Y^s) \cdot \mathbb{E}(B^{\alpha s})$. Then, using (17) and (20), one gets (21). With the relation (19), this gives (22). \square

One thus has the relations $\text{BML}(\alpha, \theta, 0) = \text{ML}(\alpha, \theta)$, $\text{BML}(\alpha, \alpha, 1 - \alpha) = \text{ML}(\alpha)$, and more generally $\text{BML}(\alpha, \alpha k, 1 - \alpha) = \text{ML}(\alpha, \alpha(k - 1))$.

Note that the beta-Mittag-Leffler distribution is one important instance of a “distribution with moments of Gamma type”, a class of distributions popularized by Janson in his nice thorough survey [66]. Therein, amongst several properties of these distributions, he obtains the asymptotics of their tail; applied to $\text{BML}(\alpha, \theta, \beta)$, this implies that its density $f(x)$ satisfies

$$f(x) \sim Cx^{d-1} \exp(-cx^{\frac{1}{1-\alpha}}) \quad \text{for } x \sim +\infty, \text{ with}$$

$$c = (1 - \alpha)\alpha^{\frac{\alpha}{1-\alpha}}, \quad d = \frac{\theta/\alpha - \theta - \beta + 1/2}{1 - \alpha}, \quad \text{and} \quad C = \frac{\Gamma(\theta + \beta)}{\Gamma(\frac{\theta}{\alpha})} \frac{\alpha^{\frac{1-2\beta}{2(1-\alpha)}}}{\sqrt{2\pi(1-\alpha)}}.$$

Let us end this short introduction on the beta-Mittag-Leffler distribution by mentioning that an explicit series representation of its density is given later (in Formula (25) in the next section); we establish it via an inverse Mellin transform.

Another important ingredient of this article will be the mixed Poisson distributions. These distributions were first introduced by Dubourdieu in 1938 for actuarial mathematics/insurance modelling [30], and then also studied by Lundberg and others (sometimes under the name “compound Poisson processes”, a term that has a different meaning nowadays); they were also used by Neyman for applications in bacteriology [97], or for the analysis of some point processes in [51]. Their unimodality properties are studied in [91], and their tail asymptotics are analysed in [121].

DEFINITION 3.15 (Mixed Poisson distributions). Let X denote a nonnegative random variable with cumulative distribution function U . We say that the discrete random variable Y has a *mixed Poisson distribution with mixing distribution U* and scale parameter $\xi \geq 0$, if its probability mass function is given for $\ell \geq 0$ by

$$\mathbb{P}\{Y = \ell\} = \frac{\xi^\ell}{\ell!} \int_{\mathbb{R}^+} X^\ell e^{-\xi X} dU = \frac{\xi^\ell}{\ell!} \mathbb{E}(X^\ell e^{-\xi X}).$$

This is summarized by the notation $Y \stackrel{d}{=} \text{MPo}(\xi U)$, or, indifferently, $Y \stackrel{d}{=} \text{MPo}(\xi X)$.

Note that mixed Poisson distributions provide a common generalization of three major discrete laws ubiquitous in combinatorics (see [40, Figure IX.5]), namely the Poisson, the geometric, and the negative binomial distributions, making them of great importance per se.

We emphasize that the *factorial moments*⁶ of a mixed Poisson distribution are closely related to the classical *raw moments* of its mixing distribution:

$$\mathbb{E}(Y^s) = \xi^s \mathbb{E}(X^s), \quad s \geq 1.$$

Additionally, like for any distribution, the factorial and raw moments of Y are related via the Stirling set partition numbers $\left\{ \begin{smallmatrix} s \\ k \end{smallmatrix} \right\}$ (also called Stirling numbers of the second kind):

$$\mathbb{E}(Y^s) = \sum_{k=0}^s \left\{ \begin{smallmatrix} s \\ k \end{smallmatrix} \right\} \mathbb{E}(Y^k).$$

We note in passing that such general relations are called Stirling transforms [17]. We refer to [83] for more properties of the Stirling transformation and mixed Poisson distributions; therein, Panholzer and the second author also gave the following useful expression for the probability mass function of $Y \stackrel{d}{=} \text{MPo}(\xi X)$ in terms of its *factorial* moments.

⁶Throughout this work we denote by $x^{\underline{n}}$ the n^{th} falling factorial, $x^{\underline{n}} = x(x-1)\cdots(x-n+1)$, $n \geq 0$, with $x^{\underline{0}} = 1$. It will be used for $\mathbb{E}(X^{\underline{n}})$, the factorial moment of order n of a random variable X .

PROPOSITION 3.16. *Let X denote a random variable with moment sequence given by $(\mu_s)_{s \in \mathbb{N}}$ such that $\mathbb{E}(e^{zX})$ exists in a neighbourhood of zero, including the value $z = -\xi$. If a random variable Y has factorial moments given by $\mathbb{E}(Y^{\underline{s}}) = \xi^s \mu_s$, then $Y \stackrel{d}{=} \text{MPo}(\xi X)$. What is more, the sequence of moments of Y is the Stirling transform of the moment sequence $(\mu_s)_{s \in \mathbb{N}}$, and the probability mass function of Y is given by*

$$\mathbb{P}\{Y = \ell\} = \sum_{s \geq \ell} (-1)^{s-\ell} \binom{s}{\ell} \mu_s \frac{\xi^s}{s!}, \quad \ell \geq 0.$$

Let us give two short examples of mixed Poisson distributions, which, as we shall later see, correspond to ubiquitous cases in combinatorics.

EXAMPLE 3.17 (Mixed Poisson half-normal distribution). A half-normally distributed random variable $X \stackrel{d}{=} \text{HN}(\sigma)$ with parameter σ is the absolute value of a normally distributed random variable $Y \stackrel{d}{=} \mathcal{N}(0, \sigma^2)$: $X \stackrel{d}{=} |Y|$. Consequently, X has the probability density function

$$f(x; \sigma) = \frac{\sqrt{2}}{\sigma\sqrt{\pi}} e^{-\frac{x^2}{2\sigma^2}}, \quad x > 0;$$

alternatively, it is fully characterized by its moment sequence

$$\mathbb{E}(X^s) = \sigma^s 2^{s/2} \frac{\Gamma(\frac{s+1}{2})}{\Gamma(\frac{1}{2})}.$$

Thus, a discrete random variable Y with probability mass function

$$\mathbb{P}\{Y = \ell\} = \frac{\xi^\ell}{\ell!} \cdot \frac{\sqrt{2}}{\sigma\sqrt{\pi}} \int_0^\infty x^\ell e^{-\xi x - \frac{x^2}{2\sigma^2}} dx, \quad \ell \geq 0,$$

has a mixed Poisson distribution: $Y \stackrel{d}{=} \text{MPo}(\xi X)$ with $X \stackrel{d}{=} \text{HN}(\sigma)$. Note that we can readily expand the exponential function and obtain various series representations of $\mathbb{P}\{Y = \ell\}$. ■

EXAMPLE 3.18 (Mixed Poisson Rayleigh distribution). A Rayleigh distributed random variable $X \stackrel{d}{=} \text{Rayleigh}(\sigma)$ with parameter σ has the probability density function

$$f(x; \sigma) = \frac{x}{\sigma^2} e^{-\frac{x^2}{2\sigma^2}}, \quad x \geq 0;$$

alternatively, it is fully characterized by its moment sequence

$$\mathbb{E}(X^s) = \sigma^s 2^{s/2} \Gamma\left(\frac{s}{2} + 1\right).$$

Thus, a discrete random variable Y with probability mass function

$$\mathbb{P}\{Y = \ell\} = \frac{\xi^\ell}{\ell! \sigma^2} \int_0^\infty x^{\ell+1} e^{-\xi x - \frac{x^2}{2\sigma^2}} dx, \quad \ell \geq 0,$$

has a mixed Poisson distribution: $Y \stackrel{d}{=} \text{MPo}(\xi X)$ with $X \stackrel{d}{=} \text{Rayleigh}(\sigma)$. Another representation valid for all $\xi > 0$ can be stated in terms of the incomplete gamma function $\Gamma(s, x) = \int_x^\infty t^{s-1} e^{-t} dt$:

$$\mathbb{P}\{Y = \ell\} = \frac{(\xi\sigma)^\ell}{\ell!} e^{-\frac{(\xi\sigma)^2}{2}} \sum_{i=0}^{\ell+1} \binom{\ell+1}{i} (-\xi\sigma)^{\ell+1-i} 2^{\frac{i-1}{2}} \Gamma\left(\frac{i+1}{2}, \frac{(\xi\sigma)^2}{2}\right). \quad \blacksquare$$

In our results, we shall also encounter another important family of discrete distributions: the *Boltzmann distributions*.

Boltzman distributions were introduced in combinatorics by Duchon, Flajolet, Louchard, and Schaeffer in order to perform sampling of combinatorial structures [31]. The starting point of these authors was the idea to give, like in statistical mechanics, a Gibbs measure/Boltzmann weight x^n (for some fixed real number x) to each combinatorial object of size n . Note that objects of the same size then follow a uniform distribution. It was then a nice surprise that, if one deals with an assemblage of combinatorial objects, the corresponding Boltzmann weights are given by very simple probabilistic laws (similarly to the symbolic method [40] which directly gives the generating functions of unions/products/cycles of objects). This led to an outstanding generic linear time sampling algorithm: Its astonishing efficiency is partially due to the fact that, thanks to these Boltzmann weights, the sampling of a product of two combinatorial structures is simply obtained by *two* independent recursive subsamplings. The sampling algorithm is thus essentially based on the following definition:

DEFINITION 3.19 (Boltzmann distribution). For any generating function $G(z) = \sum_{n \geq 0} g_n z^n$, and for any parameter $x > 0$ inside the radius of convergence of G , a random variable X follows a *Boltzmann distribution (associated with G) of parameter x* , denoted by $\mathcal{B}_G(x)$, if

$$\mathbb{P}\{X = n\} = \frac{g_n x^n}{G(x)}, \quad n \geq 0.$$

Then, the key idea behind Boltzmann sampling (of objects of size n) is to choose x adequately to maximize $\mathbb{P}\{X = n\}$. If the object generated is not of size n , one rejects it and restarts the sampling. This leads to a uniform sampling algorithm of optimal efficiency when x is the unique real root of the equation $xG'(x) = nG(x)$. This equation is reminiscent of many probabilistic results with mean $\mu = xG'(x)/G(x)$ (e.g., when G encodes the offspring of a Galton–Watson process). This is no coincidence: By design, Boltzmann sampling “reverse-engineers” these results [98, 115].

As we shall see, these Boltzmann distributions also occur in our *critical* composition schemes. Retrospectively, it explains and puts in a unified framework earlier sporadic occurrences of such distributions for the limit law of the degree of a random node in simply-generated trees, the root degree in simply-generated trees, as well as in *subcritical* composition schemes; see [40, pages 460, 629–633].




FIG 3. Boltzmann distributions $\mathcal{B}_G(x)$ have different shapes⁸, related to the singular exponent λ_G of the function G : It goes from an asymptotic Gaussian shape if $\lambda_G \ll -1$ (and for entire functions) to a spread shape if $\lambda_G \gg -1$. The above distributions are drawn with a value of the parameter x such that $\mathbb{E}(\mathcal{B}_G(x)) = xG'(x)/G(x) = n$ (with $n = 200$). It is interesting that they appear in the context of critical schemes, independently of uniform random generation motivations, but, as explained above, both aspects are in fact intimately related.

⁸See <https://lipn.fr/~cb/Papers/CriticalSchemes/> for several animations of the different limit laws occurring in this article.

4. Extended composition scheme. In the following we state and prove our main theorem on the extended critical composition scheme (2) for pure schemes. For the terms critical and pure, we refer to Definitions 1.1 and 3.3, respectively. This theorem shows the universality of the beta-Mittag-Leffler distribution (introduced in Definition 3.13).

THEOREM 4.1 (Extended composition scheme: pure case). *In a pure extended critical composition scheme $F(z, u) = G(uH(z))M(z)$, the core size X_n , rescaled, converges in distribution and in moments⁹ to a random variable X distributed like a beta-Mittag-Leffler distribution:*

$$\frac{X_n}{\kappa \cdot n^{\lambda_H}} \xrightarrow{d} X, \quad \text{with} \quad X \stackrel{d}{=} \text{BML}(\alpha, \theta, \beta), \quad (23)$$

where

$$\alpha = \lambda_H, \quad \theta = -\lambda_G \lambda_H, \quad \beta = -\lambda_M^- = -\min(0, \lambda_M), \quad \text{and} \quad \kappa = \frac{\tau_H}{-c_H}.$$

What is more, one has a local limit theorem

$$\mathbb{P}\{X_n = x \cdot \kappa n^{\lambda_H}\} \sim \frac{1}{\kappa n^{\lambda_H}} \cdot f_X(x), \quad (24)$$

where $f_X(x)$ is the density of X :

$$f_X(x) = \frac{\Gamma(\theta + \beta)}{\Gamma(\theta/\alpha)} \sum_{j \geq 0} \frac{(-1)^j}{j! \Gamma(\beta - j\alpha)} x^{\theta/\alpha + j - 1}. \quad (25)$$

REMARK 4.2 (Two simplifications of the beta-Mittag-Leffler distribution). In the above theorem, if $\lambda_M \geq 0$ (which includes the critical scheme $F(z, u) = G(uH(z))$ as $\lambda_M = +\infty$), then $\beta = 0$ and the beta-Mittag-Leffler distribution (23) simplifies into a generalized Mittag-Leffler distribution:

$$X \stackrel{d}{=} \text{tilt}_{-\lambda_G}(\text{ML}(\lambda_H)) \stackrel{d}{=} \text{ML}(\lambda_H, -\lambda_G \lambda_H).$$

In particular, for $\lambda_H = \frac{1}{2}$ and $\lambda_G = -1$ the random variable X follows a Rayleigh distribution of parameter $\sigma = \sqrt{2}$; see Example 3.10.

Another noteworthy simplification occurs for $\lambda_M < 0$, in the special case $\lambda_G = -1$ and $\lambda_H - \lambda_M = 1$: We then obtain a Mittag-Leffler distribution of parameter λ_H :

$$X \stackrel{d}{=} \text{ML}(\lambda_H).$$

In particular, for $\lambda_H = \frac{1}{2}$ the random variable X follows a half-normal distribution of parameter $\sigma = \sqrt{2}$; see again Example 3.10.

Note that the cases with $\lambda_G = -1$ occur in a great many places in applied probability theory; they indeed correspond to a natural combinatorial framework where \mathcal{F} -objects are essentially *sequences* of \mathcal{H} -components (see [40] and examples in our Section 6).

PROOF OF THEOREM 4.1. The factorial moments satisfy¹⁰:

$$\mathbb{E}(X_n^s) = \frac{[z^n] \partial_u^s (F)(z, 1)}{[z^n] F(z, 1)} = \frac{[z^n] H(z)^s G^{(s)}(H(z)) M(z)}{[z^n] G(H(z)) M(z)}.$$

⁹We write $X_n \xrightarrow{d} X$ to denote that X_n converges in distribution to X , with convergence of all moments, i.e., $\mathbb{E}(X_n^s) \rightarrow \mathbb{E}(X^s)$ for all s . This notion of convergence in moments was, e.g., used in [46, 69]. It is also indirectly used in [113], which deals with more constrained models that offer a convergence in L^p , for all $p > 1$. Note that in all our results involving convergence in distribution or in moments, we omit the speed of convergence, which is in fact easily obtained by considering the Puiseux expansions of order 2 of $G/H/M$.

¹⁰Throughout this work, we denote by ∂_u the differentiation operator with respect to the variable u . Accordingly, we use the shorthand notation $\partial_u(F)(z, 1) = (\partial_u F(z, u))|_{u=1}$.

In the following we use the notation ρ_F , τ_F , λ_F , and c_F from Section 3.1 for the singular expansions of $F = G/H/M$. As we are in a pure critical scheme (Definition 3.3), the unique singularity of $F(z) = G(H(z))M(z)$ is at $z = \rho_H$. Then, we unify the three cases (10a), (10b), and (10c) by using $\lambda_M^- = \min(0, \lambda_M)$ and choosing C_M according to the specific case (C_M is for $\lambda_M \neq \lambda_G \lambda_H$ either τ_M or c_M ; this quantity C_M will anyway cancel in the end). Note that the case (10d) does not hold in a pure scheme. This gives

$$F(z) \sim C_M c_G \left(\frac{-c_H}{\rho_G} \right)^{\lambda_G} \left(1 - \frac{z}{\rho_H} \right)^{\lambda_G \lambda_H + \lambda_M^-}.$$

Therefore, using singularity analysis we get

$$f_n = [z^n]F(z, 1) \sim \frac{C_M c_G (-c_H/\rho_G)^{\lambda_G}}{\rho_H^n} \cdot \frac{n^{-\lambda_G \lambda_H - \lambda_M^- - 1}}{\Gamma(-\lambda_G \lambda_H - \lambda_M^-)}. \quad (26)$$

Using singular differentiation [35, 40] for $G(z)$, we get the following singular expansion of the higher-order derivatives $G^{(s)}(z)$, for integer $s \geq 1$:

$$G^{(s)}(z) \sim (-1)^s \frac{c_G}{\rho_G^s} \lambda_G^s \left(1 - \frac{z}{\rho_G} \right)^{\lambda_G - s}.$$

From this we get

$$G^{(s)}(H(z)) \sim (-1)^s c_G \rho_G^{-\lambda_G} \lambda_G^s (-c_H)^{\lambda_G - s} \left(1 - \frac{z}{\rho_H} \right)^{\lambda_G \lambda_H - s \lambda_H}.$$

Next, from the singular expansion of $H(z)$ we directly get

$$H(z)^s \sim \tau_H^s - s \tau_H^{s-1} \cdot (-c_H) \left(1 - \frac{z}{\rho_H} \right)^{\lambda_H}.$$

Combining these expansions with the one of $M(z)$ gives the required expansion

$$H(z)^s G^{(s)}(H(z)) M(z) \sim (-1)^s \tau_H^s C_M c_G \rho_G^{-\lambda_G} \lambda_G^s (-c_H)^{\lambda_G - s} \left(1 - \frac{z}{\rho_H} \right)^{\lambda_M^- + \lambda_H \lambda_G - s \lambda_H}.$$

Next we rewrite $(-1)^s \lambda_G^s$ using the gamma function:

$$(-1)^s \lambda_G^s = \frac{\Gamma(s - \lambda_G)}{\Gamma(-\lambda_G)}.$$

Hence, we obtain by extraction of coefficients and singularity analysis

$$[z^n] H(z)^s G^{(s)}(H(z)) M(z) \sim \left(\frac{\tau_H}{-c_H} \right)^s \frac{C_M c_G (-c_H/\rho_G)^{\lambda_G}}{\rho_H^n} \times \frac{\Gamma(s - \lambda_G)}{\Gamma(-\lambda_G)} \cdot \frac{n^{-\lambda_G \lambda_H - \lambda_M^- - 1 + s \lambda_H}}{\Gamma(s \lambda_H - \lambda_G \lambda_H - \lambda_M^-)}. \quad (27)$$

Combining this expression with (26) gives

$$\mathbb{E}(X_n^s) \sim n^{s \lambda_H} \kappa^s \cdot \frac{\Gamma(s - \lambda_G) \Gamma(-\lambda_G \lambda_H - \lambda_M^-)}{\Gamma(-\lambda_G) \Gamma(s \lambda_H - \lambda_G \lambda_H - \lambda_M^-)},$$

with $\kappa = \frac{\tau_H}{-c_H}$. What is more, one has $\mathbb{E}(X_n^s) \sim \mathbb{E}(X_n^{\bar{s}})$ since we can express the raw moments by using the factorial moments and the Stirling numbers of the second kind:

$$\mathbb{E}(X_n^s) = \mathbb{E} \left(\sum_{j=0}^s \left\{ \begin{matrix} s \\ j \end{matrix} \right\} X_n^j \right) = \sum_{j=0}^s \left\{ \begin{matrix} s \\ j \end{matrix} \right\} \mathbb{E}(X_n^j). \quad (28)$$

Consequently, we obtain the moment convergence to the stated moment sequence:

$$\frac{\mathbb{E}(X_n^s)}{n^{s\lambda_H} \kappa^s} \rightarrow \mu_s.$$

By Stirling's formula for the gamma function, one has

$$\Gamma(z) = \left(\frac{z}{e}\right)^z \frac{\sqrt{2\pi}}{\sqrt{z}} \left(1 + \mathcal{O}\left(\frac{1}{z}\right)\right); \quad (29)$$

this entails that for $s \rightarrow \infty$ the moments satisfy

$$(\mu_s)^{-1/(2s)} \sim \sqrt{e^{1-\lambda_H} \lambda_H^{\lambda_H} s^{\frac{H-1}{2}}} \left(1 + \mathcal{O}\left(\frac{1}{s}\right)\right).$$

As $0 < \lambda_H < 1$, the divergence in Carleman's criterion [21, pp. 189–220] is satisfied:

$$\sum_{s=0}^{\infty} \mu_s^{-1/(2s)} = +\infty; \quad (30)$$

consequently, the moment sequence $(\mu_s)_{s \in \mathbb{N}}$ characterizes a unique distribution. Thus, by the Fréchet–Shohat theorem [43], we obtain the weak convergence of the normalized random variable $\frac{X_n}{\kappa \cdot n^{\lambda_H}}$ to a random variable X with moment sequence $(\mu_s)_{s \in \mathbb{N}}$.

Concerning the local limit theorem, we have to analyse $\mathbb{P}\{X_n = k\} = \frac{g_k [z^n] H(z)^k M(z)}{f_n}$, for $k = x \cdot \kappa \cdot n^{\lambda_H}$, with x in a compact subinterval of $(0, \infty)$. First, we note that by (7) applied to $G(z)$ we directly obtain

$$g_k \sim \frac{c_G}{\rho_G^k} \cdot \frac{(\kappa x)^{-\lambda_G - 1} n^{-\lambda_H \lambda_G - \lambda_H}}{\Gamma(-\lambda_G)}. \quad (31)$$

It remains to determine the asymptotics of $[z^n] H(z)^k M(z)$. We have

$$[z^n] H(z)^k M(z) = \frac{1}{2\pi i} \oint \frac{H(z)^k M(z)}{z^{n+1}} dz.$$

Introducing the point A of coordinates $(\frac{1}{n}, \rho_H(1 + \frac{\log^2 n}{n}))$, this Cauchy integral can be transformed into an integral over a larger contour in the Delta-domain (in blue in Figures 1 and 4). Then, setting $z = \rho_H(1 + t/n)$ leads to an integral asymptotically concentrated on the Hankel contour \mathcal{C} (which starts and ends at $+\infty$):

$$[z^n] H(z)^k M(z) \sim \frac{\tau_H^k C_M}{\rho_H^n n^{1+\lambda_M}} \cdot \frac{1}{2\pi i} \int_{\mathcal{C}} (-t)^{\lambda_M} e^{-t-x(-t)^{\lambda_H}} dt.$$




FIG 4. The integration contour of the Cauchy integral (left) is transformed into a larger contour (in red in the middle image) inside a Delta-domain (in blue), becoming in the limit the Hankel contour \mathcal{C} (right), allowing the identification of the gamma function.

Then, expanding $e^{-x(-t)^{\lambda_H}}$ leads to

$$[z^n]H(z)^k M(z) \sim \frac{\tau_H^k C_M}{\rho_H^n n^{1+\lambda_M}} \cdot \frac{1}{2\pi i} \int_{\mathcal{C}} e^{-t} \sum_{j \geq 0} \frac{(-x)^j}{j!} (-t)^{j\lambda_H + \lambda_M} dt, \quad (32)$$

in which we recognize the following Hankel contour representation [119, Section 12.22]:

$$\frac{1}{\Gamma(-z)} = \frac{1}{2i\pi} \int_{\mathcal{C}} (-t)^z e^{-t} dt.$$

Recall that we use, by analytic continuation, $1/\Gamma(m) = 0$ for any integer $m < 0$; this allows us to avoid more cumbersome expressions which make use of Euler's reflection formula $\frac{1}{\Gamma(-z)} = -\frac{1}{\pi} \sin(\pi z) \Gamma(1+z)$. Now, combining the expansions of f_n from (26), g_k from (31), and term-wise integration in (32), we get the desired local limit theorem:

$$\begin{aligned} \mathbb{P}\{X_n = k\} &= \frac{g_k}{f_n} [z^n]H(z)^k M(z) \\ &\sim \frac{\Gamma(-\lambda_G \lambda_H - \lambda_M)}{\kappa n^{\lambda_H} \Gamma(-\lambda_G)} \sum_{j \geq 0} \frac{(-1)^j}{j! \Gamma(-j\lambda_H - \lambda_M)} x^{j-\lambda_G-1}. \end{aligned} \quad (33)$$

It remains to verify that $f_X(x)$ is the density function of X with moments

$$\mu_s = \frac{\Gamma(s - \lambda_G) \Gamma(-\lambda_G \lambda_H - \lambda_M)}{\Gamma(-\lambda_G) \Gamma(s \lambda_H - \lambda_G \lambda_H - \lambda_M)}.$$

Note that the gamma function is never zero and that its only singularities are simple poles at the negative integers. Therefore μ_s (considered as a complex function of s) has simple poles at $\rho_0 = \lambda_G$ (if $\lambda_M \neq 0$, which entails $\lambda_G < 0$ as we have a pure scheme) and at $\rho_j = \lambda_G - j$ (for $j \in \{1, 2, \dots\}$). Then, as μ_{s-1} is the Mellin transform of the density $f_X(x)$ of X , an inverse Mellin computation¹¹ implies that $f_X(x)$ is expressible in terms of the residues of μ_s :

$$f_X(x) = \frac{1}{2\pi i} \int_{\sigma-i\infty}^{\sigma+i\infty} \mu_{s-1} x^{-s} ds = \sum_{\substack{\sigma_j = 1 + \rho_j \\ \sigma_j < \sigma}} \text{Res}_{s=\sigma_j}(\mu_{s-1}) x^{-\sigma_j},$$

which is valid for $x > 0$, and where $\sigma = 1 + \rho_0 + \epsilon$ is in the fundamental strip of μ_{s-1} , and

$$\text{Res}_{s=\sigma_j}(\mu_{s-1}) = \text{Res}_{s=\rho_j}(\mu_s) = \frac{(-1)^j}{j!} \cdot \frac{\Gamma(-\lambda_G \lambda_H - \lambda_M)}{\Gamma(-\lambda_G) \Gamma(-j\lambda_H - \lambda_M)}.$$

Summing for $j \geq 0$ (we can include ρ_0 , even if it is not a pole, as the corresponding residue is then 0) gives the same density function $f_X(x)$ as in the local limit theorem (33).

Now, thanks to our probability distribution melting pot section 3.2, we identify the corresponding limit law by matching the parameters in the moments of the beta-Mittag-Leffler product given in Lemma 3.14. \square

Above, we have established the limit laws occurring for analytically *pure* composition schemes. Next we deal with the analytically *confluent* and *degenerate* cases of Definition 3.3. They lead either to discrete distributions, or, more interestingly, to a mixture of discrete and continuous distributions.

This generalizes the phenomenon observed in [6], where for $\lambda_H = \frac{3}{2}$ the limit law also consists of a discrete part *plus* a continuous part, a map-Airy distribution (this phenomenon also occurs for variants of 3-connected graphs; see [48]). The following two theorems explain the connection between this discrete part and Boltzmann distributions (usually used for random sampling; see, e.g., [19, 31]).

¹¹See [40, Appendix B.7] for more on this method; see also [66, Theorem 5.4] with $\gamma = \gamma' = 1 - \lambda_H > 0$, and [66, Equation (6.12)] for similar results on the class of functions with moments of Gamma type.

THEOREM 4.3 (Extended composition scheme: degenerate case). *Let a degenerate extended critical composition scheme $F(z, u) = G(uH(z))M(z)$ with $0 < \lambda_H < 1$ be given.*

For $0 < \lambda_G < 1$ and $\lambda_M < \lambda_G \lambda_H$, the core size X_n converges for $k \geq 0$ and $n \rightarrow \infty$ to a Boltzmann distribution $\mathcal{B}_G(\rho_G)$ (see Definition 3.19):

$$\mathbb{P}\{X_n = k\} \rightarrow \mathbb{P}\{\mathcal{B}_G(\rho_G) = k\} = \frac{g_k \rho_G^k}{G(\rho_G)}.$$

For $\lambda_G > 1$, the core size X_n has for $k \geq 0$ and $n \rightarrow \infty$ the following behaviour:

- *For $\lambda_M < \lambda_H$, the random variable X_n converges to a Boltzmann distribution $\mathcal{B}_G(\rho_G)$:*

$$\mathbb{P}\{X_n = k\} \rightarrow \frac{g_k \rho_G^k}{G(\rho_G)}.$$

- *For $\lambda_M = \lambda_H$, the random variable X_n converges to a convex combination of two discrete Boltzmann distributions:*

$$X_n \xrightarrow{d} \text{Be}(p) \cdot \mathcal{B}_G(\rho_G) + (1 - \text{Be}(p)) \cdot \mathcal{B}_{G'}(\rho_{G'}),$$

with $p = \frac{c_M G(\rho_G)}{c_M G(\rho_G) + c_H \tau_M G'(\rho_G)}$ and where the Bernoulli distribution $\text{Be}(p)$ is independent of the other random variables. Therefore, we have

$$\mathbb{P}\{X_n = k\} \rightarrow p \cdot \frac{g_k \rho_G^k}{G(\rho_G)} + (1 - p) \cdot \frac{k g_k \rho_G^{k-1}}{G'(\rho_G)}.$$

- *For $\lambda_M > \lambda_H$, the random variable X_n converges to a Boltzmann distribution $\mathcal{B}_{G'}(\rho_{G'})$:*

$$\mathbb{P}\{X_n = k\} \rightarrow \frac{k g_k \rho_G^{k-1}}{G'(\rho_G)}.$$

PROOF. Let us start with the case $0 < \lambda_G < 1$. We study for arbitrary but fixed $k \in \mathbb{N}$ the probability $\mathbb{P}\{X_n = k\}$, as n tends to infinity. From (3) we obtain

$$\begin{aligned} \mathbb{P}\{X_n = k\} &= g_k \frac{[z^n] H(z)^k \cdot M(z)}{f_n} \\ &\sim \frac{g_k}{f_n} [z^n] \left(\left(\tau_H^k + k \tau_H^{k-1} c_H \left(1 - \frac{z}{\rho_H}\right)^{\lambda_H} \right) \left(\tau_M + c_M \left(1 - \frac{z}{\rho_H}\right)^{\lambda_M} \right) \right), \end{aligned} \quad (34)$$

where we apply the Singular-expansion Lemma 3.1 to the function $H(z)$. From the expansion (10c), we get by standard singularity analysis (7), for $\lambda_M < \lambda_G \lambda_H$, that f_n satisfies

$$f_n \sim \tau_G c_M \frac{1}{\rho_H^n} \frac{n^{-\lambda_M-1}}{\Gamma(-\lambda_M)},$$

as the singular exponent λ_M dominates the asymptotics. Thus we obtain from (34)

$$\mathbb{P}\{X_n = k\} \sim \frac{g_k \tau_H^k}{\tau_G} = \frac{g_k \rho_G^k}{G(\rho_G)},$$

because $\tau_H = \rho_G$ and $\tau_G = G(\rho_G)$ by our assumptions on criticality.

Let us continue with the case $\lambda_G > 1$. From expansion (10d) we see that we now have to distinguish three subcases: $\lambda_M < \lambda_H$, $\lambda_M = \lambda_H$, or $\lambda_H < \lambda_M$. The case $\lambda_M < \lambda_H$ is exactly the same as before. The case $\lambda_M > \lambda_H$ is different, yet the results are derived analogously: the asymptotics of f_n depend now on λ_H and are given by

$$f_n \sim G'(\rho_G) \tau_M c_H \frac{1}{\rho_H^n} \frac{n^{-\lambda_H-1}}{\Gamma(-\lambda_H)}.$$

Then, we obtain from (34) where now again the contribution from $H(z)$ dominates

$$\mathbb{P}\{X_n = k\} \sim \frac{k g_k \tau_H^{k-1}}{G'(\rho_G)} = \frac{k g_k \rho_G^{k-1}}{G'(\rho_G)}.$$

Finally, in the case $\lambda_M = \lambda_H$ the previous two contributions are mixed as the first two terms in (10d) contribute: The asymptotics of f_n are given by

$$f_n \sim (G'(\rho_G)\tau_M c_H + c_M \tau_G) \frac{1}{\rho_H^n} \frac{n^{-\lambda_H-1}}{\Gamma(-\lambda_H)}.$$

Note that the coefficient is not zero, as both terms are negative: $G'(\rho_g), \tau_M, \tau_M > 0$ and $c_H, c_M < 0$ due to the sign property (8) for $0 < \lambda_H < 1$. Then, we obtain from (34) where again both contributions have to be taken into account

$$\mathbb{P}\{X_n = k\} \sim \frac{c_M g_k \rho_G^k + \tau_M c_H k g_k \rho_G^{k-1}}{G'(\rho_G)\tau_M c_H + c_M \tau_G} = p \cdot \frac{g_k \rho_G^k}{G(\rho_G)} + (1-p) \cdot \frac{k g_k \rho_G^{k-1}}{G'(\rho_G)},$$

where $p = \frac{c_M G(\rho_G)}{c_M G(\rho_G) + c_H \tau_M G'(\rho_G)} \in (0, 1)$ by the sign property (8). We thus get a linear combination of two Boltzmann distributions, weighted by a Bernoulli random variable $\text{Be}(p)$. \square

THEOREM 4.4 (Extended composition scheme: confluent case). *Let a confluent extended critical composition scheme $F(z, u) = G(uH(z))M(z)$ with $0 < \lambda_H < 1$ be given (i.e., $0 < \lambda_G < 1$ and $\lambda_M = \lambda_G \lambda_H$). Then the core size X_n is a convex combination of a Boltzmann distribution $\mathcal{B}_G(\rho_G)$ and an asymptotically continuous random variable Z_n :*

$$X_n \sim \text{Be}(p) \cdot \mathcal{B}_G(\rho_G) + (1 - \text{Be}(p)) \cdot Z_n, \quad \frac{Z_n}{\kappa \cdot n^{\lambda_H}} \xrightarrow{d} X,$$

where κ and the limit law $X \stackrel{d}{=} \text{ML}(\lambda_H, -\lambda_G \lambda_H)$ are the same as in Remark 4.2, and where $\text{Be}(p)$ (with $p = \frac{c_M G(\rho_G)}{c_M G(\rho_G) + \tau_M c_G (-c_H/\rho_G)^{\lambda_G}}$) is independent of $\mathcal{B}_G(\rho_G)$, Z_n , and X .

PROOF. The proof follows the same lines as the one of Theorem 4.3. We start from expansion (10c). Due to $\lambda_M = \lambda_G \lambda_H$ both terms contribute, and we get

$$f_n \sim \frac{\tau_M c_G (-c_H/\rho_G)^{\lambda_G} + \tau_G c_M}{\rho_H^n} \frac{n^{-\lambda_M-1}}{\Gamma(-\lambda_M)}. \quad (35)$$

Then we extract the asymptotics from (34), where now the contributions from $M(z)$ dominate, as $\lambda_M < \lambda_G$:

$$\mathbb{P}\{X_n = k\} = \frac{c_M g_k \tau_H^k}{\tau_M c_G (-c_H/\rho_G)^{\lambda_G} + \tau_G c_M} + o(1) = p \cdot \frac{g_k \rho_G^k}{G(\rho_G)} + o(1), \quad (36)$$

Using the sign property (8) we get $p \in (0, 1)$. So, for large n , X_n behaves with probability p like $\mathcal{B}_G(\rho_G)$, but what happens with probability $1-p$? Where is this missing mass in (36)? For sure, it is sneakily spread in $\sum_k o(1)$: It turns out that more and more mass is distributed in the range $k \sim \Theta(\kappa n^{\lambda_H})$, leading to an asymptotic continuous distribution therein.

In order to identify this distribution, we compute the factorial moments of X_n like in the proof of Theorem 4.1. We again use the singular expansions of $G^{(s)}(H(z))$, $H(z)^s$, and $H(z)^s G^{(s)}(H(z))M(z)$. Formula (27) holds verbatim with $C_M = \tau_M$. The big difference lies now in the asymptotics of f_n : It is given by (26) in the pure case and by (35) in the confluent case. Thus, after rescaling (27) by f_n , the factorial moments have the same shape, but with an additional prefactor $1-p$: $\mathbb{E}(X_n^{\underline{s}}) \sim (1-p) \cdot \mathbb{E}(Z_n^{\underline{s}})$, which proves the claim. \square

REMARK 4.5 (Physical interpretation of the bimodal behaviour). As illustrated by Table 1 on page 4, the confluent case gives a bimodal distribution. The first mode is dictated by small values of k , where a Boltzmann distribution is dominant, while for larger values of $k \approx n^{\lambda_H}$, a second mode appears, which is associated with a continuous Mittag-Leffler distribution. This phenomenon explains the following behaviour: If one picks an atom at random in a structure of size n , then with probability p it lies in a large \mathcal{H} -block of size $\Theta(n)$ and with probability $1-p$ in a smaller \mathcal{H} -block of size $\Theta(n^{1-\lambda_H})$.

In the next section, we will refine these considerations by having a closer look at the distribution of the \mathcal{H} -blocks of any given size.

5. Size-refined composition scheme. In this section, we give the limit laws for the profile of combinatorial structures given by a critical composition scheme. We focus here on schemes which are analytically pure (see Definition 3.3), while we handle the confluent and degenerate cases in our companion article [9] (as they require additional technical details and different families of limit laws, which also pop up for $\lambda_H > 1$). The profile is captured by the size-refined composition scheme (4). As we see in the theorem below, we get three distinct asymptotic régimes, each leading to its own limit law. Two of these limit laws are expressible in terms of the generalized Mittag-Leffler distribution of Theorem 4.1 (see also Definition 3.11 and Definition 3.13).

THEOREM 5.1 (Mixed Poisson limit behaviour for the size-refined scheme). *Consider a size-refined pure critical composition scheme $F(z, v) = G(H(z) - (1 - v)h_j z^j)M(z)$, with $j \in \mathbb{N}$. Let $\xi_{n,j} = \frac{\rho_H^j}{-c_H} h_j n^{\lambda_H}$, and X be the beta-Mittag-Leffler distribution of Theorem 4.1:*

$$X \stackrel{d}{=} \text{BML}(\alpha, \theta, \beta),$$

where $\alpha = \lambda_H$, $\theta = -\lambda_G \lambda_H$, and $\beta = -\min(0, \lambda_M)$.

Then, the random variable $X_{n,j}$, which counts the number of \mathcal{H} -components of size j possesses three distinct asymptotic régimes, with a phase transition at $j = \Theta(n^{\frac{\lambda_H}{1+\lambda_H}})$:

(i) For $j \ll n^{\frac{\lambda_H}{1+\lambda_H}}$, we have $\xi_{n,j} \rightarrow +\infty$ and convergence in distribution and in moments:

$$\frac{X_{n,j}}{\xi_{n,j}} \xrightarrow[m]{d} X.$$

(ii) For $j \sim r \cdot n^{\frac{\lambda_H}{1+\lambda_H}}$, $r \in (0, \infty)$, we have $\xi_{n,j} \rightarrow \xi$ with $\xi = r^{-\frac{\lambda_H}{1+\lambda_H}} \cdot \frac{1}{-\Gamma(-\lambda_H)}$ and convergence in distribution and in moments towards a mixed Poisson distribution:

$$X_{n,j} \xrightarrow[m]{d} \text{MPo}(\xi X).$$

(iii) For $j \gg n^{\frac{\lambda_H}{1+\lambda_H}}$, we have $\xi_{n,j} \rightarrow 0$, and $X_{n,j}$ converges to a Dirac distribution at 0.

REMARK 5.2 (Phase transition I). The intuition behind the phase transition is as follows: In the limit $n \rightarrow \infty$, there are many small ($j \ll n^{\frac{\lambda_H}{1+\lambda_H}}$), some giant ($j \sim r n^{\frac{\lambda_H}{1+\lambda_H}}$), and no super-giant ($j \gg n^{\frac{\lambda_H}{1+\lambda_H}}$) \mathcal{H} -components of size j . It is interesting to compare this situation with the birth of the giant component in Erdős–Rényi random graphs; see [67]. Note that the case $j \in \mathbb{N}$ fixed (i.e., independent of n as n tends to infinity) falls into the case $j \ll n^{\frac{\lambda_H}{1+\lambda_H}}$.

REMARK 5.3 (Phase transition II). For the often observed case of a square-root singularity of $H(z)$ (i.e., $\lambda_H = \frac{1}{2}$), we reobtain the critical range $j = \Theta(n^{1/3})$, which was already observed in the mixed Poisson Rayleigh distributions in [83]. Furthermore, as one has

$$\mathbb{E}(X_{n,j}^s) = \xi_{n,j}^s \cdot \mathbb{E}(X^s) \cdot (1 + o(1)),$$

this offers en passant a link between the $\xi_{n,j}$'s and the rescaling factor κn^{λ_H} in Theorem 4.1:

$$\sum_{j \geq 1} \xi_{n,j} = \sum_{j \geq 1} \frac{\rho_H^j}{-c_H} h_j n^{\lambda_H} = \frac{H(\rho_H)}{-c_H} n^{\lambda_H} = \frac{\tau_H}{-c_H} n^{\lambda_H} = \kappa n^{\lambda_H}.$$

This link can be seen as an asymptotic avatar of the combinatorial relation $\sum_{j \geq 1} X_{n,j} = X_n$, implying

$$\sum_{j \geq 1} \mathbb{E}(X_{n,j}) = \mathbb{E}(X_n).$$

In order to prove Theorem 5.1, we first need the following lemma concerning the convergence of mixed Poisson distributions.

LEMMA 5.4 (Factorial moments and limit laws of mixed Poisson type [83]). *Let $(X_n)_{n \in \mathbb{N}}$ denote a sequence of random variables, whose factorial moments are asymptotically of mixed Poisson type, i.e., they satisfy for $n \rightarrow \infty$ the asymptotic expansion*

$$\mathbb{E}(X_n^s) = \xi_n^s \cdot \mu_s \cdot (1 + o(1)), \quad s \geq 1,$$

with $\mu_s \geq 0$ and $\xi_n > 0$. Furthermore, assume that the moment sequence $(\mu_s)_{s \in \mathbb{N}}$ determines a unique distribution X satisfying Carleman's condition. Then, the following limit distribution results hold:

- (i) If $\xi_n \rightarrow \infty$, the random variable $\frac{X_n}{\xi_n}$ converges in distribution, with convergence of all moments, to X .
- (ii) If $\xi_n \rightarrow \xi \in (0, \infty)$, the random variable X_n converges in distribution, with convergence of all moments, to a mixed Poisson distributed random variable $Y \stackrel{d}{=} \text{MPo}(\xi X)$.
- (iii) If $\xi_n \rightarrow 0$, the random variable X_n converges to a Dirac distribution: $X_n \xrightarrow{d} 0$.

REMARK 5.5. The second and third case could be grouped together, since for $\xi = 0$ we have $Y \stackrel{d}{=} \text{MPo}(0) \stackrel{d}{=} 0$. Furthermore, in the third case, for positive random variables $(X_n)_{n \in \mathbb{N}}$ the assumptions can be relaxed to simply $\mathbb{E}(X_n) \rightarrow 0$, without requiring the specific structure of the moments. Note further that the discrete random variable $Y \stackrel{d}{=} \text{MPo}(\xi X)$ converges, after scaling, to its mixing distribution X : One has $\frac{Y}{\xi} \xrightarrow[\xi \rightarrow \infty]{d} X$, with convergence of all moments.

PROOF OF THEOREM 5.1. The factorial moments $\mathbb{E}(X_{n,j}^s) = \sum_{k \geq 0} \mathbb{P}\{X_{n,j} = k\} k^s$ of $X_{n,j}$ are obtained from $F(z, v)$ by repeated differentiation and evaluation at $s = 1$:

$$\mathbb{E}(X_{n,j}^s) = \frac{[z^n] \partial_v^s (F)(z, 1)}{[z^n] F(z, 1)} = h_j^s \frac{[z^{n-j-s}] G^{(s)}(H(z)) M(z)}{f_n}.$$

We already know the asymptotics of f_n from (26). For fixed j we can proceed by extraction of coefficients, while for $j = j(n)$ tending to infinity, the asymptotic expansion of h_j follows by singularity analysis applied to $H(z)$ (see Equation (7)):

$$h_j = \frac{c_H}{\rho_H^j} \cdot \frac{j^{-\lambda_H - 1}}{\Gamma(-\lambda_H)} \cdot (1 + o(1)). \quad (37)$$

What is more, one has

$$G^{(s)}(H(z)) M(z) \sim (-1)^s c_M c_G \rho_G^{-\lambda_G} \lambda_G^s (-c_H)^{\lambda_G - s} \left(1 - \frac{z}{\rho_H}\right)^{\lambda_H \lambda_G - s \lambda_H + \lambda_M}. \quad (38)$$

This implies that $X_{n,j}$ has factorial moments of mixed Poisson type:

$$\mathbb{E}(X_{n,j}^s) \sim h_j^s (-c_H)^{-s} \rho_H^{j s} \cdot \mu_s \cdot n^{s \lambda_H} \quad \text{with} \quad \mu_s := \frac{\Gamma(s - \lambda_G) \Gamma(-\lambda_G \lambda_H - \lambda_M)}{\Gamma(-\lambda_G) \Gamma(-\lambda_H \lambda_G + s \lambda_H - \lambda_M)}.$$

We already observed that the moment sequence $(\mu_s)_{s \in \mathbb{N}}$ determines a unique distribution, by Carleman's criterion (30). Thus, the limit laws follow by using Lemma 5.4.

Finally, the critical growth range is obtained via the closed-form expression for $\xi_{n,j}$ (in which one inserts the expansion (37)): Indeed, $\xi_{n,j}$ is of growth order $\frac{n^{\lambda_H}}{j^{1+\lambda_H} \Gamma(-\lambda_H)}$ and converges to a nonzero constant if and only if $j(n) \sim r \cdot n^{\frac{\lambda_H}{1+\lambda_H}}$, therefore the critical growth range is $\Theta(n^{\frac{\lambda_H}{1+\lambda_H}})$. Collecting all contributions from $\xi_{n,j}$ (for $j = o(n)$ tending to infinity) gives the constant ξ . The Dirac case is now finally obtained by an additional analysis of the expected value in the remaining range $j \gg n^{\frac{\lambda_H}{1+\lambda_H}}$. There, we directly obtain $\mathbb{E}(X_{n,j}) \rightarrow 0$, which proves the stated result. \square

Next we turn to the dependence between the number of \mathcal{H} -components of size j_1 and j_2 , determining the covariance and the correlation coefficient.

THEOREM 5.6. *In a size-refined pure critical composition scheme, the covariance of the random variables X_{n,j_1} and X_{n,j_2} , counting the number of \mathcal{H} -components of size j_1 and j_2 (with $j_1, j_2 = o(n)$), satisfies*

$$\text{Cov}(X_{n,j_1}, X_{n,j_2}) \sim \xi_{n,j_1} \xi_{n,j_2} \cdot \mathbb{V}(X), \quad (39)$$

where $\xi_{n,j} = \frac{\rho_H^j}{c_H} h_j n^{\lambda_H} > 0$ and $X \stackrel{d}{=} \text{BML}(\alpha, \theta, \beta)$ denotes the beta-Mittag-Leffler distribution from Theorem 4.1. Furthermore, the correlation coefficient between X_{n,j_1} and X_{n,j_2} satisfies

$$\rho(X_{n,j_1}, X_{n,j_2}) \sim \frac{1}{\sqrt{1 + \frac{\mathbb{E}(X)}{\xi_1}}} \frac{1}{\sqrt{1 + \frac{\mathbb{E}(X)}{\xi_2}}},$$

where, for $k = 1, 2$, one has $\xi_k := \lim_n \xi_{n,j_k}$ and $\sqrt{1 + \frac{\mathbb{E}(X)}{\xi_k}} \sim 1$ if $j_k \ll n^{\frac{\lambda_H}{1+\lambda_H}}$.

REMARK 5.7. We observe that for small j_1, j_2 (e.g., if $j_1, j_2 = O(1)$), the random variables are asymptotically highly correlated: $\rho(X_{n,j_1}, X_{n,j_2}) \sim 1$.

PROOF OF THEOREM 5.6. The combinatorial scheme $\mathcal{F} = \mathcal{G}(\mathcal{H}_{\neq j_1, j_2} + v_1 \mathcal{H}_{=j_1} + v_2 \mathcal{H}_{=j_2}) \times \mathcal{M}$ (where one takes distinct sizes $1 \leq j_1 < j_2$) directly translates into

$$F(z; v_1, v_2) = G(H(z) - (1 - v_1)h_{j_1}z^{j_1} - (1 - v_2)h_{j_2}z^{j_2})M(z).$$

Accordingly, the random variables X_{n,j_1} and X_{n,j_2} have the joint distribution

$$\mathbb{P}\{X_{n,j_1} = k_1, X_{n,j_2} = k_2\} = \frac{[z^n v_1^{k_1} v_2^{k_2}] F(z; v_1, v_2)}{[z^n] F(z; 1, 1)}.$$

We already know the asymptotics of $f_n = [z^n] F(z; 1, 1)$, given in (26). We get by differentiation and evaluation at $v_1 = v_2 = 1$

$$\mathbb{E}(X_{n,j_1} X_{n,j_2}) = \frac{[z^n] \partial_{v_1} \partial_{v_2} (F)(z; 1, 1)}{[z^n] F(z; 1, 1)} = h_{j_1} h_{j_2} \frac{[z^{n-j_1-j_2}] G''(H(z)) M(z)}{f_n}.$$

The asymptotics of h_{j_1} and h_{j_2} are given in (37). The singular expansion of $G''(H(z))M(z)$ is a special case of (38), so we obtain for $j_1, j_2 = o(n)$:

$$\mathbb{E}(X_{n,j_1} X_{n,j_2}) \sim h_{j_1} h_{j_2} c_H^{-2} \rho_H^{j_1+j_2} \cdot \mathbb{E}(X^2) \cdot n^{2\lambda_H}.$$

Hence, using the explicit form of $\mathbb{E}(X)$ in (21), we obtain for the covariance:

$$\begin{aligned} \text{Cov}(X_{n,j_1}, X_{n,j_2}) &= \mathbb{E}(X_{n,j_1} X_{n,j_2}) - \mathbb{E}(X_{n,j_1}) \mathbb{E}(X_{n,j_2}) \\ &\sim \frac{h_{j_1} h_{j_2} \rho_H^{j_1+j_2}}{c_H^2} (\mathbb{E}(X^2) - \mathbb{E}(X)^2) n^{2\lambda_H}. \end{aligned}$$

By $\mathbb{V}(X) = \mathbb{E}(X^2) - \mathbb{E}(X)^2$ and the definition of $\xi_{n,j}$, this implies (39). For the correlation coefficient, we observe that Theorem 5.1 together with Lemma 5.4 implies $\mathbb{V}(X_{n,j}) \sim \xi_{n,j}^2 \mathbb{V}(X) + \xi_{n,j} \mathbb{E}(X)$. Collecting all contributions, this implies that

$$\rho(X_{n,j_1}, X_{n,j_2}) = \frac{\text{Cov}(X_{n,j_1}, X_{n,j_2})}{\sqrt{\mathbb{V}(X_{n,j_1})} \sqrt{\mathbb{V}(X_{n,j_2})}} \sim \frac{1}{\sqrt{1 + \frac{\mathbb{E}(X)}{\xi_{n,j_1}}} \sqrt{1 + \frac{\mathbb{E}(X)}{\xi_{n,j_2}}}}.$$

Taking the limit of this expression for n going to infinity concludes the proof. \square

Before to present in Section 7 a multivariate generalization of these results, we now list several applications to a variety of combinatorial structures.

6. Applications and examples. Our main results, Theorems 4.1 and 5.1, can be readily applied to the problems considered by Drmota and Soria [29], Flajolet and Sedgewick [40], Dumas, Flajolet and Puyhaubert [37], Janson [65], Meir and Moon [92], Panholzer and Seitz [102] (see also [83] for many additional pointers to the literature) once the required singular expansions of the involved generating functions are established. This includes returns to record-subtrees in Cayley trees, edge-cutting in Cayley trees, returns to zero in Dyck paths, cyclic points and trees in graphs of random mappings, all leading to (mixed Poisson) Rayleigh laws, as well as block sizes in k -Stirling permutations.

In the following we discuss several new results for the distribution of different parameters such as returns to zero and sign changes in walks and bridges with arbitrary steps, the number of subtrees satisfying some constraint in different fundamental families of trees, as well as the table sizes in the Chinese restaurant process, and the evolution of the number of balls in triangular urn models. These examples illustrate that composition schemes $\mathcal{F} = \mathcal{G}(\mathcal{H}) \times \mathcal{M}$ lead universally to beta-Mittag-Leffler distributions (Definition 3.13), which simplify into generalized Mittag-Leffler distributions if $M(z)$ has a nonnegative singular exponent.

6.1. Core size of supertrees and a confluent example. Let \mathcal{C} denote the family of plane trees (i.e., trees with all arities allowed and embedded into the plane) defined by

$$\mathcal{C} = \mathcal{Z} \times \text{SEQ}(\mathcal{C}), \quad \text{which translates to} \quad C(z) = \frac{1 - \sqrt{1 - 4z}}{2}.$$

Then, following [40, pp. 412–414, 714], we consider supertrees, or “trees of trees”, defined by

$$\mathcal{K} = \mathcal{C}\left((\mathcal{Z}_{\text{red}} + \mathcal{Z}_{\text{blue}}) \times \mathcal{C}\right), \quad \text{which translates to} \quad K(z) = C(2zC(z)).$$

Note that this is a critical scheme as one has $\rho_{\mathcal{C}} = 1/4$ and $\tau_{\mathcal{C}} = C(\rho_{\mathcal{C}}) = 1/2$.

The tree family \mathcal{K} corresponds to trees where onto each node we graft a red or blue tree; one can also draw them like done in Figure 5. By Lagrange inversion, these supertrees \mathcal{K} are thus enumerated by a neat combinatorial sum:

$$K_n = \sum_{k=1}^{\lfloor n/2 \rfloor} \frac{2^k}{n-k} \binom{2k-2}{k-1} \binom{2n-3k-1}{n-k-1},$$

thus the sequence K_n for $n \geq 2$ starts like 2, 2, 8, 18, 64, 188, 656, 2154, \dots , constituting sequence A168506 in the OEIS¹².




FIG 5. A bicoloured supertree is a “tree of trees”: Each node (here drawn in a potato shape) of some initial plane tree is replaced by a red or a blue node to which one attaches another plane tree.

¹²OEIS stands for the On-Line Encyclopedia of Integer Sequences, accessible via <https://oeis.org>.

By the Laplace method or by singularity analysis, this directly leads to the asymptotic expansion

$$K_n \sim \frac{4^n}{8\Gamma(3/4)n^{5/4}}.$$

(See also DeVries [24, 106] for an alternative approach to this expansion via multivariate analysis.) This asymptotic behaviour is noteworthy, because one sees here an unusual occurrence of the exponent $-\frac{5}{4}$, while most tree-like structures in combinatorics usually involve the exponent $-\frac{3}{2}$. In fact, one could similarly define super-supertrees, super-super-supertrees, and so on, by further iterations of the critical scheme: $\mathcal{C}_{k+1} = \mathcal{C}_k(2\mathcal{Z}\mathcal{C}_k)$ with $\mathcal{C}_0 = \mathcal{C}$. This gives an interesting family of combinatorial structures whose asymptotic enumeration involves a dyadic exponent $-1 - 1/2^{k+1}$; see [4] for a complete characterization of the possible singular exponents for \mathbb{N} -algebraic functions (i.e., generating functions of any structure which can be generated by a context-free language).

With respect to supertrees, the critical scheme is

$$K(z) = G(H(z)), \quad G(z) = C(z), \quad H(z) = 2zC(z),$$

where $H(z)$ has the following Puiseux expansion at $z \sim \frac{1}{4}$:

$$H(z) \sim \frac{1}{4} - \frac{1}{4}\sqrt{1-4z}.$$

Now, we can study the core size X_n via the bivariate generating function

$$K(z, u) = C(u \cdot 2zC(z)),$$

as well as the number of \mathcal{H} -components of size j , as captured by

$$K(z, v) = C(2zC(z) + (v-1)2c_{j-1}z^j).$$

We can then apply our main Theorems 4.1 and 5.1 (with $\lambda_G = \lambda_H = 1/2$, $\tau_H = \frac{1}{4}$, $c_H = -\frac{1}{4}$, and $\kappa = 1$). This directly gives the following corollaries.

COROLLARY 6.1. *The core size X_n in supertrees of size n has factorial moments given by*

$$\mathbb{E}(X_n^s) \sim n^{s/2} \cdot \mu_s, \quad \mu_s = \frac{\Gamma(s - \frac{1}{2})\Gamma(-\frac{1}{4})}{\Gamma(-\frac{1}{2})\Gamma(\frac{s}{2} - \frac{1}{4})}.$$

The scaled random variable $X_n/n^{1/2}$ converges in distribution with convergence of all moments to a $c = -1/2$ moment-tilted stable distribution, more precisely, a generalized Mittag-Leffler distribution of index $1/2$:

$$\frac{X_n}{n^{1/2}} \xrightarrow{d} X, \quad \text{with} \quad X \stackrel{d}{=} \text{tilt}_{-\frac{1}{2}}\left(\text{ML}\left(\frac{1}{2}\right)\right) \stackrel{d}{=} \text{ML}\left(\frac{1}{2}, -\frac{1}{4}\right).$$

Moreover, we have the local limit theorem

$$\mathbb{P}\{X_n = x \cdot n^{1/2}\} \sim \frac{1}{n^{1/2}} \cdot f_X(x),$$

with $f_X(x)$ denoting the density of the random variable X .

Note that by Legendre's duplication formula one has

$$\mu_s = 2^s \cdot \frac{\Gamma(\frac{s}{2} + \frac{1}{4})}{\Gamma(\frac{1}{4})},$$

so the random variable X can also be seen as equal in law to the chi distribution $\chi(\frac{1}{2})$ of parameter $\frac{1}{2}$, which is itself a generalized gamma distribution [10].

COROLLARY 6.2. *The number of coloured trees of size $j - 1$ in supertrees of size n has factorial moments of mixed Poisson type given by*

$$\mathbb{E}(X_{n,j}^s) = \xi_{n,j}^s \cdot \mu_s(1 + o(1)),$$

with $\xi_{n,j} = 2 \cdot (\frac{1}{4})^{j-1} c_{j-1} \cdot n^{1/2}$ and mixing distribution $X = \chi(\frac{1}{2})$ with $\mathbb{E}(X^s) = \mu_s$.

Furthermore, the random variable $X_{n,j}$ possesses the three distinct asymptotic régimes of Theorem 5.1, with a phase transition at $j = \Theta(n^{1/3})$.

We note in passing that a very similar result holds for the family of binary supertrees \mathcal{S} , occurring in Bousquet-Mélou's study [20] of the integrated super-Brownian excursion. This family is defined in terms of the family of complete binary trees \mathcal{B} :

$$\mathcal{S} = \mathcal{B}(\mathcal{Z} \times \mathcal{B}), \quad \mathcal{B} = \mathcal{Z} + \mathcal{Z} \times \mathcal{B} \times \mathcal{B},$$

with initial values of S_n given by 1, 1, 3, 8, 25, 80, 267, 911, ..., constituting sequence [A101490](#) in the OEIS. These functional equations indeed lead to $B(z) = \frac{1 - \sqrt{1 - 4z^2}}{2z}$, and to the following Puiseux expansion for $\tilde{S}(z) = S(\sqrt{z})$:

$$\tilde{S}(z) \sim 1 - \sqrt{2}(1 - 4z)^{1/4} + (1 - 4z)^{1/2} + \dots;$$

thus leading to limit laws similar to the ones of supertrees in Corollaries 6.1 and 6.2.

A confluent example. Next, we consider pairs of supertrees in which we mark the number of \mathcal{H} -trees in the first part of the pair. This translates to the scheme $F(z, u) = G(uH(z))K(z)$ (with G , H , and K defined as in (6.1)), which is confluent because we have $\lambda_H \lambda_G = \lambda_M = \frac{1}{4}$. This case is interesting as $\mathbb{P}\{X_n = k\}$ (the probability that a pair of supertrees has k \mathcal{H} -trees in its first part) converges to the sum of a discrete law and a continuous law. More precisely, as given by Theorem 4.4, and illustrated in Figure 6, the limit is a linear combination of a Boltzmann distribution, namely $\mathcal{B}_C(\frac{1}{4})$ ¹³, and a generalized Mittag-Leffler distribution, namely $\text{ML}(\frac{1}{2}, -\frac{1}{4})$.




FIG 6. The empirical distribution $\mathbb{P}\{X_n = k\}$ (drawn with red dots), and its theoretical limiting distribution (in blue). This blue curve is a linear combination of a discrete and a continuous distribution (the middle and right curves drawn in black): $\frac{1}{2} \mathcal{B}_C(\frac{1}{4}) + \frac{1}{2} \sqrt{n} \text{ML}(\frac{1}{2}, -\frac{1}{4})$. The blue theoretical limit curve agrees almost perfectly with the red empirical distribution even for small values of n (here, $n = 500$).

¹³ C denoting the Catalan generating function, it could be natural to call this Boltzmann distribution $\mathcal{B}_C(\frac{1}{4})$ the *Catalan distribution*, but this name is already used for some other distributions popping up in random matrix theory, and having moments related to the Catalan numbers, like the Marchenko–Pastur distribution.




FIG 7. An increasing diamond and the bijectively equivalent bilabelled increasing plane-oriented recursive tree.

6.2. *Root degree and branching structure in bilabelled increasing trees.* Bilabelled increasing trees are a natural generalization of increasing trees [77] where every node is assigned two labels instead of just one. General families of bilabelled trees are in bijection with increasing diamonds, which are a natural type of directed acyclic graphs; see [84] for the general statement and Figure 7 for a concrete example. Increasing diamonds model partial orders and their linear extensions, as well as computational processes and their executions in parallel computing [18]. They possess nice combinatorial properties and are enumerated by variants of hook-length formulas [79, 82, 84].

Given a *degree-weight sequence* $(\varphi_j)_{j \geq 0}$, the corresponding *degree-weight generating function* is defined as $\varphi(t) = \sum_{j \geq 0} \varphi_j t^j$. The family \mathcal{T} of bilabelled increasing trees can be described by the following symbolic equation:

$$\mathcal{T} = \mathcal{Z}^\square * \left(\mathcal{Z}^\square * \varphi(\mathcal{T}) \right),$$

where \mathcal{Z} denotes single unilabelled nodes, $\mathcal{A}^\square * \mathcal{B}$ denotes the *boxed product* (i.e., the smallest label is constrained to lie in the \mathcal{A} component) of the combinatorial classes \mathcal{A} and \mathcal{B} , and $\varphi(\mathcal{A}) = \varphi_0 \cdot \{\epsilon\} + \varphi_1 \cdot \mathcal{A} + \varphi_2 \cdot \mathcal{A}^2 + \dots$ denotes the class containing all *weighted finite labelled sequences* of objects of \mathcal{A} (i.e., each sequence of length k is weighted by φ_k ; ϵ denotes the *neutral object* of size 0); see [40]. Note that increasing diamonds are associated with $\varphi(t) = \frac{1}{1-t}$. For the exponential generating function $T(z) = \sum_{n \geq 1} T_n \frac{z^n}{n!}$ where n counts the number of labels, the above construction translates into

$$T''(z) = \varphi(T(z)), \quad T(0) = 0, \quad T'(0) = 0. \tag{40}$$

We now focus on the case of 3-bundled bilabelled increasing trees; see Figure 8.




FIG 8. An increasing plane-oriented recursive tree and the bijectively equivalent 3-bundled bilabelled increasing plane-oriented recursive tree. (The grey ellipses are just here to sketch the bijective correspondences.)

The family of 3-bundled trees is defined by Equation (40) with the following degree-weight generating function

$$\varphi(t) = \frac{1}{(1-t)^3} = \sum_{k \geq 0} \binom{k+2}{2} t^k. \quad (41)$$

In other words, each node may have any number k of children, and the binomial indicates two bars between these children, thus creating 3 (possibly empty) sequences (or *bundles*) of children. From [82] we get the remarkably simple closed form

$$T(z) = 1 - \sqrt{1-z^2} = \sum_{n \geq 1} (2n-1)!!(2n-3)!! \frac{z^{2n}}{(2n)!}, \quad (42)$$

where the double factorials are defined as

$$(2n-1)!! = \prod_{k=1}^n (2k-1) = \frac{(2n)!}{n! \cdot 2^n},$$

with initial values of T_{2n} given by 1, 3, 45, 1575, 99225, 9823275, 1404728325, \dots , constituting sequence [A079484](#) in the OEIS.

We are interested in the random variable X_n counting the root degree of these 3-bundled bilabelled increasing trees of size n , under the uniform random tree model. Note that by definition there are no bilabelled trees with an odd number of labels, so $T_{2n+1} = 0$ and, consequently, $X_{2n+1} = 0$. In the following we use the notation $R_n = X_{2n}$ for the root degree.

By (40) and (41), the generating function

$$T(z, u) = \sum_{n \geq 1} T_n \mathbb{E}(u^{X_n}) \frac{z^n}{n!}$$

satisfies

$$\frac{\partial^2}{\partial z^2} T(z, u) = \varphi(uT(z)) = \frac{1}{(1-u(1-\sqrt{1-z^2}))^3}.$$

Therefore, the Taylor expansion of $T(z, u)$ starts as follows

$$T(z, u) = 1 + 3u \frac{z^2}{2!} + (3u + 9u^2) \frac{z^4}{4!} + (36u + 135u^2 + 540u^3) \frac{z^6}{6!} + \dots$$

Since a derivative with respect to z is simply a shift in the coefficient sequence of exponential generating functions, we obtain

$$\mathbb{E}(u^{R_{n+1}}) = \frac{(2n)!}{T_{2n+2}} [z^n] \frac{1}{(1-u(1-\sqrt{1-z}))^3}.$$

Now we observe, by Stirling's formula for the gamma function (29) and singularity analysis (7) applied to (42), that

$$\frac{T_{2n+2}}{(2n)!} \sim \frac{2\sqrt{n}}{\sqrt{\pi}} \sim [z^n] \frac{1}{(1-z)^{3/2}}. \quad (43)$$

This implies that, except for the non-standard shift in the random variable, the problem is equivalent (for first-order asymptotics) to the composition scheme $(1-u(1-\sqrt{1-z}))^{-3}$, i.e., $\rho_H = 1$, $\lambda_H = \frac{1}{2}$, $\lambda_G = -3$, and $\lambda_M = +\infty$ (as $M(z) = 1$ is entire). We note in passing that in this special case, it is also possible to obtain a quite simple closed-form expression for the probability mass function, as well as the (factorial) moments, due to the explicit expressions for the involved generating functions:

$$\mathbb{E}(X_{2n+2}^s) = \frac{(2n)!}{T_{2n+2}} [z^n] \partial_u^s \frac{1}{(1-u(1-\sqrt{1-z}))^3} \Big|_{u=1}.$$

However, here we use our general scheme and Theorem 4.1. We apply Legendre's duplication formula and obtain

$$\frac{\Gamma(s+3)\Gamma(\frac{3}{2})}{\Gamma(3)\Gamma(\frac{s+3}{2})} = 2^s \cdot \Gamma\left(\frac{s+4}{2}\right).$$

This leads to the following result.

COROLLARY 6.3. *The random variable R_n , counting the root degree in a random strict bilabelled increasing three-bundled tree with $2n$ labels, with tree generating function given by $\varphi(t) = (1-t)^{-3}$, satisfies*

$$\mathbb{E}(R_n^s) \sim n^{s/2} \cdot 2^s \cdot \Gamma\left(\frac{s+4}{2}\right).$$

The random variable $R_n/n^{1/2}$ converges (in distribution and in moments) to a multiple of a chi-distributed random variable $X \stackrel{d}{=} \chi(4)$, with four degrees of freedom:

$$\frac{R_n}{n^{1/2}} \xrightarrow[m]{d} \sqrt{2} \cdot X, \quad X \stackrel{d}{=} \chi(4).$$

We can refine the root degree by looking at the branching structure. We denote by $R_{n,j} = X_{2n,j}$ the random variable counting the number of branches (i.e., subtrees) with $2j$ labels, $1 \leq j \leq n$, attached to the root:

$$R_n = \sum_{j \geq 1} R_{n,j}.$$

Such random variables naturally arise in the context of the Chinese restaurant process [83, 108, 109] and generalized plane-oriented recursive trees [74]. See also Feng et al. [116] for the analysis of the branching structure of recursive trees. The generating function $T(z, v) = \sum_{n \geq 1} T_n \mathbb{E}(v^{X_{n,j}}) \frac{z^n}{n!}$ satisfies

$$\frac{\partial^2}{\partial z^2} T(z, v) = \varphi \left(T(z) - (1-v) \frac{T_{2j}}{(2j)!} z^{2j} \right).$$

Consequently,

$$\mathbb{E}(u^{R_{n+1,j}}) = \frac{(2n)!}{T_{2n+2}} [z^{2n}] \frac{1}{\left(1 - \left((1 - \sqrt{1-z^2}) - (1-v) \frac{T_{2j}}{(2j)!} z^{2j}\right)\right)^3}.$$

We can use the asymptotics (43) and apply Theorem 5.1 to obtain the following result.

COROLLARY 6.4. *The random variable $R_{n,j}$ counting the number of size $2j$ branches attached to the root in a random strict bilabelled increasing three-bundled tree with $2n$ labels, with tree generating function given by $\varphi(t) = (1-t)^{-3}$, has factorial moments of mixed Poisson type,*

$$\mathbb{E}(R_{n,j}^s) = \xi_{n,j}^s \cdot \mathbb{E}(X^s)(1 + o(1)),$$

with $\xi_{n,j} = \sqrt{2} \cdot \frac{T_{2j}}{(2j)!} \cdot n^{1/2}$ and mixing distribution $X = \chi(4)$.

Furthermore, the random variable $R_{n,j}$ possesses the three distinct asymptotic régimes of Theorem 5.1, with a phase transition at $j = \Theta(n^{1/3})$.




FIG 9. A walk consists of an initial bridge which contains all returns to zero (red dots) and a final walk never returning to zero. The bridge is further decomposed into minimal bridges touching zero only twice.

6.3. Returns to zero: walks and bridges with drift zero. A lattice path of length n is a sequence (s_1, \dots, s_n) of steps $s_i \in \mathcal{S}$ for a fixed finite subset $\mathcal{S} \subseteq \mathbb{Z}$ called step set. Geometrically, we fix the starting point 0 and consider the partial sums $\sum_{i=1}^k s_i$ which can be interpreted as appending the steps one after another. Each step s_i gets a weight $p_i > 0$ and the weight of a path is the product of all steps. The *step polynomial* $P(u) = \sum_i p_i u^i$ connects the weights and the steps. We call a step set *periodic* if there exist integers $b, p \in \mathbb{Z}$, $p \geq 2$ such that $P(u) = u^b P(u^p)$; otherwise we call it *aperiodic*. Here and in the next section, we assume that the step set is aperiodic. Note that this is no major constraint as the asymptotics of walks with periodic steps can be deduced from the ones with aperiodic ones; see [14]. We call a lattice path a walk if it is unconstrained, and a bridge if it ends at zero, i.e., $\sum_{i=1}^n s_i = 0$. A *return to zero* is a point in the path such that $\sum_{i=1}^k s_i = 0$; see Figure 9.

Generalizing results from Feller [33, Problems 9–10] and Barton [114, Discussion p. 115], it was shown in [118, Section 3.2] that for drift $P'(1) = 0$ the law of the number of returns to zero follows a Rayleigh distribution for bridges, while it follows a half-normal distribution for walks. The proof utilized a general theorem on the singular structure of the generating functions. However, the situation is also amenable to Theorem 4.1 and we can give an alternative proof next.

Let $w_{n,k}$ be the number of walks of length n with k returns to zero. The bivariate generating function of walks $W(z, u) = \sum_{n,k \geq 0} w_{n,k} z^n u^k$ is given by

$$W(z, u) = \frac{1}{1 - u \left(1 - \frac{1}{B(z)}\right)} \frac{W(z)}{B(z)}, \quad (44)$$

where $B(z)$ and $W(z) = 1/(1 - zP(1))$ are the generating functions of bridges and walks, respectively; see [118, Equation (3.3)]. To explain (44), observe that every bridge is a sequence of *minimal bridges*, which are bridges that never return to the x -axis between the start- and endpoint; see Figure 9. Therefore, minimal bridges are enumerated by $1 - 1/B(z)$. Hence, this is exactly the situation of the extended composition scheme (2) with $G(z) = 1/(1 - z)$, $H(z) = 1 - 1/B(z)$, and $M(z) = W(z)/B(z)$. Now, for zero drift, [118, Equation (3.4)] shows that

$$B(z) = \frac{c_B}{\sqrt{1 - zP(1)}} + \mathcal{O}(1) \quad \text{with} \quad c_B = \sqrt{\frac{P(1)}{2P''(1)}}. \quad (45)$$

Hence, one has $\lambda_G = -1$, $\lambda_H = \frac{1}{2}$, and $\lambda_M^- = -\frac{1}{2}$, therefore we are in the pure régime; see Definition 3.3. This is exactly the situation in Remark 4.2 and the number of returns to zero in walks thus follows a half-normal distribution with parameter $\sigma = \sqrt{2}c_B = \sqrt{P(1)/P''(1)}$.

Now, the generating function for bridges is nearly the same as $W(z, u)$ from (44) except that the last factor $W(z)/B(z)$ is omitted. So, by Corollary 4.2, the number of returns to zero here follows a Rayleigh distribution with the same parameter $\sigma = \sqrt{P(1)/P''(1)}$.

We can refine this result for the random variable $X_{n,j}$ counting the number of distance- j -zeroes (which were introduced in [83]). These are the number of returns to zero which have a distance of exactly j steps to the *previous* zero contact. The union over j of distance- j -zeroes gives all returns to zero, and they therefore clearly represent a partition of all returns to zero. Using Theorem 5.1, we then get the following limit theorem.

COROLLARY 6.5. *Let $X_{n,j}$ be the number of distance- j -zeroes in lattice paths of length n . For walks (resp. bridges) with zero drift (i.e., $P'(1) = 0$), $X_{n,j}$ has factorial moments of mixed Poisson half-normal type (resp. mixed Poisson Rayleigh type)*

$$\mathbb{E}(X_{n,j}^s) = \xi_{n,j}^s \cdot \mathbb{E}(X^s) (1 + o(1)), \tag{46}$$

with $\xi_{n,j} = \sqrt{\frac{P(1)}{2P''(1)}} \frac{h_j}{P(1)^j} \cdot n^{1/2}$, where X is given by

$$X = \begin{cases} \text{HN}(\sigma) & \text{for walks,} \\ \text{Rayleigh}(\sigma) & \text{for bridges,} \end{cases} \quad \sigma = \sqrt{\frac{P(1)}{P''(1)}}.$$

Furthermore, the random variable $X_{n,j}$ possesses the three distinct asymptotic régimes of Theorem 5.1, with a phase transition at $j = \Theta(n^{1/3})$.

REMARK 6.6 (Universality of the rescaling factor). Note that the rescaling factor $\xi_{n,j}$ in (46) is the same for walks and bridges, while the moment sequence μ_s changes. This independence of $\xi_{n,j}$ can be explained: In Figure 9, the last factor of the walk is a walk not touching zero and is encoded by $M(z) = W(z)/B(z)$. Then by Theorem 5.1 we know that $\xi_{n,j}$ is independent of this factor, and thus has the same value for walks and bridges.

Furthermore, this factor $M(z)$ is also responsible for the often observed dichotomy between half-normal and Rayleigh distributions in the extended composition scheme which we will also observe in the next examples of initial returns and sign changes.

Note that Formula (46) offers a neat factorization for the moments: One can regroup in one factor the quantities with a probabilistic flavour (involving the variance $P''(1)$ of the allowed steps, and μ_s), while the remaining factor (h_j , the number of minimal bridges of length j) corresponds to a quantity with a combinatorial flavour. This could also be explained using renewal theory.

6.4. Initial returns in coloured bridges. We generalize the previous model by introducing m -coloured bridges B_m (see Figure 10): We append m non-empty bridges one after the other (and each one with a different colour): $B_m = (B - 1)^m$. Then, we are interested in the number of returns to zero in the first bridge, i.e., the initial one that we uniformly coloured. We call such returns the *initial returns*.




FIG 10. A 4-coloured bridge, with all its initial returns to zero marked by red dots.

Reusing the combinatorial constructions of the previous section, this gives for the bivariate generating function $B_m(z, u)$ the following decomposition

$$B_m(z, u) = \left(\frac{1}{1 - u \left(1 - \frac{1}{B(z)}\right)} - 1 \right) (B(z) - 1)^{m-1}. \quad (47)$$

For $m = 1$ this is (44) except the factor $W(z)/B(z)$ and the constraint to be non-empty. Asymptotically, and therefore for the law, the non-emptiness is negligible. The generating function $W_m(z, u)$ of m -coloured walks (m -tuples of bridges with a few more steps coloured in the same colour as the final bridge) is given by

$$W_m(z, u) = (1 + B_m(z, u)) \frac{W(z)}{B(z)}.$$

Now, we can directly apply Theorem 4.1. From the reasoning above we see that $\lambda_G = -1$, $\lambda_H = \frac{1}{2}$, and $\lambda_M = -\frac{m-1}{2}$ for bridges and $\lambda_M = -\frac{m}{2}$ for walks.

COROLLARY 6.7. *The random variable X_n counting the number of initial returns in a m -coloured walk (resp. bridge) of length n satisfies*

$$\mathbb{E}(X_n^s) \sim n^{s/2} \left(\frac{\sigma}{\sqrt{2}} \right)^n \mu_s, \quad \sigma = \sqrt{\frac{P(1)}{P''(1)}}, \quad \mu_s = \begin{cases} \frac{\Gamma(s+1)\Gamma((m+1)/2)}{\Gamma((m+s+1)/2)}, & \text{for walks,} \\ \frac{\Gamma(s+1)\Gamma(m/2)}{\Gamma((m+s)/2)}, & \text{for bridges.} \end{cases}$$

The scaled random variable $X_n/n^{1/2}$ converges in distribution with convergence of all moments to a Rayleigh distribution and a scaled beta distribution (see Definition 3.12 and Example 3.18):

$$\frac{X_n}{n^{1/2}} \xrightarrow{d} X, \quad X \stackrel{d}{=} \text{Rayleigh}(\sigma) \cdot B^{1/2},$$

with independent

$$\text{Rayleigh}(\sigma) \quad \text{and} \quad B = \begin{cases} \text{Beta}\left(\frac{1}{2}, \frac{m}{2}\right), & \text{for walks,} \\ \text{Beta}\left(\frac{1}{2}, \frac{m-1}{2}\right), & \text{for bridges,} \end{cases}$$

where $\text{Beta}(\alpha, 0) = 1$.

Moreover, we have the local limit theorem

$$\mathbb{P}\{X_n = x \cdot n^{1/2}\} \sim n^{-1/2} \cdot f_X(x),$$

with density $f_X(x)$ of the random variable X for bridges (for walks one replaces m by $m+1$) given by

$$f_X(x) = \sqrt{\frac{2}{\pi\sigma^2}} \Gamma\left(\frac{m}{2}\right) e^{-\frac{x^2}{2\sigma^2}} U\left(\frac{m}{2} - 1, \frac{1}{2}, \frac{x^2}{2\sigma^2}\right),$$

where $U(a, b, x)$ is the confluent hypergeometric function of the second kind which is the solution of $zy'' + (b-z)y' - ay = 0$ such that $U(a, b, x) \sim z^{-a}$ for $z \rightarrow \infty$ and $|\arg(z)| < 3\pi/2$; see [25, Section 13.2].

Observe the special cases $U(-1/2, 1/2, x) = \sqrt{x}$ and $U(0, 1/2, x) = 1$ which nicely give the density functions of a Rayleigh (see Example 3.18) and a half-normal distribution (see Example 3.17). Hence, for $m = 1$ we recover the results of the previous section and uncover a large family of connected probability distributions. It is interesting that this family also appears in the context of preferential attachments in graphs [104, Formula (1.1)].

It is also interesting to consider multicoloured bridges, where we allow any number of colours. We still mark by u the initial returns. The corresponding bivariate generating function $B(z, u)$ is given by

$$B(z, u) = \sum_{m \geq 1} B_m(z, u) = \left(\frac{1}{1 - u \left(1 - \frac{1}{B(z)}\right)} - 1 \right) \frac{1}{2 - B(z)}.$$

The generating function for the number of multicoloured bridges is thus $B(z, 1) = \frac{1}{2 - B(z)} - 1$. From (45) we see that $B(z)$ possesses a singularity of order $-1/2$ at $z = 1/P(1)$, and hence $B(z, 1)$ becomes singular at some $z_0 > 0$ which is the unique solution of $B(z_0) = 2$. Hence, the probability generating function reveals a geometric distribution of parameter $1/2$:

$$\frac{[z^n]B(z, u)}{[z^n]B(z, 1)} \sim \frac{1}{1 - u \left(1 - \frac{1}{B(z_0)}\right)} - 1 = \frac{u/2}{1 - u/2}.$$

As the truncated sum $\sum_{m=1}^{m_0} B_m(z, u)$ behaves asymptotically like $B_{m_0}(z, u)$, we see here a phase transition from a continuous law (for any finite m_0) to a discrete law (when m_0 goes to infinity). Note that this phenomenon holds verbatim for walks.

Finally, let us apply the size-refined scheme Theorem 5.1, counting initial returns in m -coloured bridges (or walks) which are a certain distance apart:

COROLLARY 6.8. *Let $X_{n,j}$ be the number of initial returns at distance j from the previous zero in m -coloured walks or bridges of length n . Then, $X_{n,j}$ has moments of mixed Poisson type*

$$\mathbb{E}(X_{n,j}^s) = \xi_{n,j}^s \cdot \mathbb{E}(X^s) (1 + o(1)),$$

with $\xi_{n,j} = \sqrt{\frac{P(1)}{2P^n(1)}} \frac{h_j}{P(1)^j} \cdot n^{1/2}$, $h_j = [z^j](1 - 1/B(z))$, and where X is the product of a beta and a Rayleigh distribution given in Corollary 6.7. Furthermore, $X_{n,j}$ possesses the three distinct asymptotic régimes of Theorem 5.1, with a phase transition at $j = \Theta(n^{1/3})$.

These results also hold for other variants of paths, in bijection with sequences that already appeared in the literature; see Table 3.

6.5. Sign changes in walks. Using the same notation as in Example 6.3, we now define the *sign* of the path (s_1, \dots, s_n) after k steps as $\text{sgn}(\sum_{i=1}^k s_i) \in \{-1, 0, 1\}$. Thereby every lattice path is associated with a sequence of signs. A *sign change* is therein any subsequence $(-1, 0^*, 1)$ or $(1, 0^*, -1)$ where 0^* denotes a (possibly empty) sequence of 0s; see Figure 11.

In this section we consider *Motzkin paths*. They are composed of up steps $+1$, down steps -1 , and horizontal steps 0 ; see again Figure 11. Their step polynomial is therefore given by $P(u) = \frac{p_{-1}}{u} + p_0 + p_1 u$ (with $p_{-1}p_0p_1 \neq 0$). In the case of zero drift, let us show how to apply our results to get that the number of sign changes follows asymptotically a Rayleigh distribution for bridges and a half-normal distribution for walks, while for nonzero drift it follows a geometric distribution; see [118].

Steps	GF	Sequence	OEIS
$\{U, D\}$	$\frac{8z^2 - 2 - \sqrt{1 - 4z^2}}{16z^2 - 3}$	1, 0, 2, 0, 10, 0, 52, 0, 274, 0, 1452, 0, 7716, ...	A075436
$\{U, D, H_1\}$	$\frac{z + \sqrt{1 - 4z^2}}{1 - 5z^2}$	1, 1, 3, 5, 13, 25, 61, 125, 295, 625, 1447, ...	A098615
$\{U, D, H_2\}$	$\frac{z^2 + \sqrt{1 - 4z^2}}{1 - 4z^2 - z^4}$	1, 0, 3, 0, 11, 0, 43, 0, 173, 0, 707, 0, 2917, ...	A026671

TABLE 3

Multicoloured bridge models: they end at 0 and use up steps $U = (1, 1)$, down steps $D = (1, -1)$, and horizontal steps $H_i = (i, 0)$ allowed only at altitude 0. The limit laws of initial returns to zero in these models are all the same and special cases of Corollaries 6.7 and 6.8.




FIG 11. A Motzkin walk (i.e., step set $S = \{-1, 0, 1\}$) with 4 sign changes marked in red.

Combinatorially, we see that the bivariate generating function of bridges is

$$B(z, u) = S(z) \left(1 + \frac{2H(z)}{1 - uH(z)} \right), \quad \text{where } S(z) = \frac{1}{1 - p_0 z} \quad \text{and} \quad H(z) = \frac{E(z)}{S(z)} - 1.$$

Here, $S(z)$ counts sequences of horizontal steps, $E(z)$ counts excursions (bridges constrained to be nonnegative; see [5]), and $H(z)$ counts excursions which start with an up or a down step (and not with a horizontal step).

We now give the main corresponding Puiseux expansions. First one has

$$H(z) = 1 - 2\sqrt{\frac{2P(1)}{P''(1)}}\sqrt{1 - zP(1)} + \mathcal{O}(1 - zP(1)).$$

Then, as the radius of convergence $1/p_0$ of $S(z)$ is strictly larger than $1/P(1)$, which is the one of $H(z)$, we see that the additive term $S(z)$ is negligible for the limit law. Thus, we have a composition scheme (2) where $M(z) = 2S(z)H(z)$ has the asymptotic expansion

$$M(z) = 2E\left(\frac{1}{P(1)}\right) + \mathcal{O}\left(\sqrt{1 - zP(1)}\right).$$

Hence, we have $\lambda_M^- = 0$, which means that the factor $M(z)$ is asymptotically negligible for the law. The asymptotic dominant part arises from $\frac{1}{1 - uH(z)}$ and we get from Corollary 4.2 the expected convergence to a Rayleigh distribution with parameter $\sigma = -\sqrt{2}\frac{\tau_H}{c_H} = \frac{1}{2}\sqrt{\frac{P''(1)}{P(1)}}$.

A similar reasoning (and Remark 4.2) allows us to prove that the number of sign changes in walks asymptotically follows a half-normal distribution with the same parameter σ . We now refine the analysis by counting sign changes which are j steps apart. Then we can apply Theorem 5.1 to get the following refined result which strongly depends on $H(z) = \sum_{j \geq 0} h_j z^j$. Note that a statement analogous to Remark 6.6 also applies here.

COROLLARY 6.9. *For walks of length n of Motzkin paths, let the random variable $X_{n,j}$ be the number of sign changes at distance j from the previous sign change or the origin. For walks (resp. bridges) with zero drift (i.e., $P'(1) = 0$), $X_{n,j}$ has factorial moments of mixed Poisson half-normal type (resp. mixed Poisson Rayleigh type)*

$$\mathbb{E}(X_{n,j}^s) = \xi_{n,j}^s \cdot \mathbb{E}(X^s) (1 + o(1)),$$

with $\xi_{n,j} = \frac{1}{2}\sqrt{\frac{P''(1)}{2P(1)}}\frac{h_j}{P(1)^j} \cdot n^{1/2}$ and mixing distributions

$$X \stackrel{d}{=} \begin{cases} \text{HN}(\sigma) & \text{for walks,} \\ \text{Rayleigh}(\sigma) & \text{for bridges,} \end{cases} \quad \sigma = \frac{1}{2}\sqrt{\frac{P''(1)}{P(1)}}.$$

Furthermore, the random variable $X_{n,j}$ (for walks and for bridges) possesses the three distinct asymptotic régimes of Theorem 5.1, with a phase transition at $j = \Theta(n^{1/3})$.




FIG 12. A plane-oriented recursive tree of size 15, where the root node labelled zero has one size-one, one size-two, one size-three, and two size-four branches, as well as the corresponding table structure in the Chinese restaurant model.

6.6. *Tables in the Chinese restaurant process.* Following Aldous, Pitman, and Dubins (see [2,108,109]), we now consider the Chinese restaurant process. This a discrete-time stochastic process having as value at time n one of the B_n partitions of the set $[n] = \{1, 2, \dots, n\}$ (where B_n denotes the Bell numbers found as sequence A000110 in the OEIS). One fancifully imagines a Chinese restaurant with an infinite number of tables, where each table has a possibly infinite number of seats. In the beginning the first customer takes a seat at the first table. At each discrete time step a new customer arrives and either joins one of the existing tables, or takes a seat at the next empty table. Each table corresponds to a block of a random partition. The process thus starts at time $n = 1$ with the partition $\{\{1\}\}$. Now, given a partition $T = \{t_1, \dots, t_k\}$ of $[n]$ with $|T| = k$ parts t_i , at time $n + 1$ the element $n + 1$ is either added to one of the existing parts $t_i \in T$ with probability

$$\mathbb{P}\{n + 1 \prec t_i\} = \frac{|t_i| - \alpha}{n + \theta}, \quad 1 \leq i \leq k,$$

where $n + 1 \prec t_i$ denotes that $n + 1$ is a customer sitting at table t_i , or as a new singleton block with probability

$$\mathbb{P}\{n + 1 \prec t_{|T|+1}\} = \frac{\theta + k \cdot \alpha}{n + \theta}.$$

This model (parametrized by the two parameters $0 < \alpha < 1$ and $\theta > -\alpha$) thus assigns a probability to any particular partition T of $[n]$. We are interested in the random variable C_n , counting the total number of tables in the Chinese restaurant process, as well as the random variable $C_{n,j}$, counting the number of parts of size j in a partition of $[n]$. As pointed out in [83], this process can be embedded into a variant of the growth process of generalized plane-oriented recursive trees with two different connectivity parameters $a > 0$ and $b > -1$. This allows us to study properties of the Chinese restaurant process using analytic combinatorial tools. For the reader's convenience, we restate this embedding below.

We collect the results of [83] and complement them by extending the constraint $b > 0$ to the full range $b > -1$ as well as by providing the missing identification of the limit law as a (moment-shifted) stable law.

Combinatorially, we consider a family $\mathcal{T}_{a,b}$ of generalized plane-oriented recursive trees, where the degree-weight generating function $\psi(t) = \frac{1}{(1-t)^b}$, $b > 0$, associated with the root of the tree, is different to the one for non-root nodes in the tree, $\varphi(t) = \frac{1}{(1-t)^a}$, $a > 0$. Then, the family $\mathcal{T}_{a,b}$ is closely related to the corresponding family \mathcal{T} of generalized plane-oriented recursive trees with degree-weight generating $\varphi(t) = \frac{1}{(1-t)^a}$, $a > 0$, via the following formal recursive equations (see Section 6.2 for the definition of the boxed product):

$$\mathcal{T}_{a,b} = \mathcal{Z}^\square * \psi(\mathcal{T}), \quad \mathcal{T} = \mathcal{Z}^\square * \varphi(\mathcal{T}).$$

The weight $w(T)$ of a tree $T \in \mathcal{T}_{a,b}$ is then defined by

$$w(T) = \psi_{d(\text{root})} \prod_{v \in T \setminus \{\text{root}\}} \varphi_{d(v)},$$

where $d(v)$ denotes the outdegree of node v . Thus, the generating functions

$$T_{a,b}(z) = \sum_{n \geq 1} T_{a,b;n} \frac{z^n}{n!} \quad \text{and} \quad T(z) = \sum_{n \geq 1} T_n \frac{z^n}{n!}$$

of the total weight of size- n trees in $\mathcal{T}_{a,b}$ and \mathcal{T} , respectively, satisfy the differential equations

$$T'_{a,b}(z) = \psi(T(z)), \quad T'(z) = \varphi(T(z)).$$

The ordinary tree evolution process to generate a random tree of arbitrary given size in the family \mathcal{T} (see [101] for a detailed discussion) can be extended in the following way to generate a random tree in the family $\mathcal{T}_{a,b}$. The process, evolving in discrete time, starts with the root labelled zero. At step $n + 1$, with $n \geq 0$, the node with label $n + 1$ is attached as a new child to any previous node v (this is denoted by $n + 1 \prec v$) with probabilities

$$\mathbb{P}\{n + 1 \prec v\} = \begin{cases} \frac{d(v)+b}{b+(a+1)n} & \text{if } v \text{ is the root,} \\ \frac{d(v)+a}{b+(a+1)n} & \text{if } v \text{ is not the root.} \end{cases}$$

We recall the following result from [83].

PROPOSITION 6.10 (Chinese restaurant process and generalized plane-oriented recursive trees). *A random partition of $\{1, \dots, n\}$ generated by the Chinese restaurant process with parameters $\alpha > 0$ and $\theta > 0$ can be generated equivalently by the growth process of the family of generalized plane-oriented recursive trees $\mathcal{T}_{a,b}$ when generating such a tree of size $n + 1$. The parameters α, θ and $a, b > 0$, respectively, are related via*

$$\alpha = \frac{1}{1+a}, \quad \theta = \frac{b}{1+a}.$$

The random variable C_n is distributed as the outdegree X_{n+1} of the root of a random generalized plane-oriented recursive trees of size $n + 1$ from the family $\mathcal{T}_{a,b}$:

$$C_n \stackrel{d}{=} X_{n+1}.$$

The random variable $C_{n,j}$ is distributed as $X_{n+1,j}$, which corresponds to the number of branches of size j attached to the root of a random tree of size $n + 1$ from the family $\mathcal{T}_{a,b}$:

$$C_{n,j} \stackrel{d}{=} X_{n+1,j}.$$

Note that in the above relation, θ cannot be negative, since b is assumed to be positive. As already observed in [83], the correspondence can be extended to the full range $0 < \alpha < 1$ and $\theta > -a$, where one has $a = \frac{1}{\alpha} - 1 > 0$ and $b = \frac{\theta}{\alpha} > -1$. For $-1 < b \leq 0$, we cannot directly use the degree-weight generating function $\psi(t) = (1-t)^{-b}$. Indeed, for $-1 < b < 0$ we would have $\psi(t) = 1 + bt + \dots$, involving a negative weight; while for $b = 0$ we would have $\psi(t) = 1$, a degenerate case. However, we can use a modified generating function, leading to a correct model of the Chinese restaurant process in the range $-1 < b \leq 0$ (see [83, 101] for more details on the growth process):

$$\psi(t) = 1 + \int_0^t \frac{1}{(1-x)^{1+b}} dx = 1 + \frac{1}{b} \left(\frac{1}{(1-t)^b} - 1 \right) = 1 + \sum_{k \geq 1} \binom{b+k}{k-1} \frac{t^k}{k},$$

for $-1 < b < 0$, while for $b = 0$ one uses

$$\psi(t) = 1 - \log(1-t) = 1 + \sum_{k \geq 1} \frac{t^k}{k}.$$

Thus, we have some generalized plane-oriented recursive trees attached to a root with a different tree-weight generating function $\psi(t)$. Summarizing, we have

$$\psi(t) = \begin{cases} \frac{1}{(1-t)^b} & \text{if } b > 0, \\ 1 - \log(1-t) & \text{if } b = 0, \\ 1 + \frac{1}{b} \left(\frac{1}{(1-t)^b} - 1 \right) & \text{if } -1 < b < 0. \end{cases}$$

Here (except for the special case $b = 0$, which is handled by a slightly different approach, detailed later in Theorem 7.1), we can directly apply our results from Theorem 4.1 to

$$T'_{a,b}(z, u) = \sum_{n \geq 1} T_n \mathbb{E}(u^{X_n}) \frac{z^{n-1}}{(n-1)!} = \psi(u \cdot T(z)), \quad T'(z) = \varphi(T(z)),$$

for the total number of tables, and from Theorem 5.1 to

$$R'_{a,b}(z, v) = \sum_{n \geq 1} T_n \mathbb{E}(v^{X_{n,j}}) \frac{z^{n-1}}{(n-1)!} = \psi(T(z) - (1-v)z^j \frac{T_j}{j!}),$$

for the number of tables of size j . This allows us to extend the corresponding result of [83] to the full range of $b > -1$, also providing the missing identification of the limit law as a (moment-tilted) stable law:

THEOREM 6.11. *Let $a > 0$, $b > -1$. The random variable $X_{n,j}$ counting the number of branches of size j in a random $\mathcal{T}_{a,b}$ tree of size n (or, equivalently, the number of tables with j seated customers in a Chinese restaurant process of parameter $\alpha = 1/(1+a)$ and $\theta = b/(1+a)$, with a total of $n-1$ customers) possesses the three distinct asymptotic régimes of Theorem 5.1, with a phase transition at $j = \Theta(n^{1/(a+2)})$:*

- (i) *For $j \ll n^{\frac{1}{a+2}}$ we have $\xi_{n,j} = \frac{\alpha n^\alpha}{j} \binom{j-1-\alpha}{j-1} \rightarrow \infty$ and $\frac{X_{n,j}}{\xi_{n,j}}$ converges in distribution, with convergence of all moments, to a generalized Mittag-Leffler distribution:*

$$\frac{X_{n,j}}{\xi_{n,j}} \xrightarrow[m]{d} X \quad \text{with} \quad X \stackrel{d}{=} \text{ML}(\alpha, \theta).$$

- (ii) *For $j \sim r \cdot n^{\frac{1}{a+2}}$, $r \in (0, \infty)$, we have $\xi_{n,j} \rightarrow \xi$, and the random variable $X_{n,j}$ converges in distribution, with convergence of all moments, to a mixed Poisson distribution:*

$$X_{n,j} \xrightarrow[m]{d} \text{MPo}(\xi X).$$

- (iii) *For $j \gg n^{\frac{1}{a+2}}$ we have $\xi_{n,j} \rightarrow 0$, so $X_{n,j}$ converges to a Dirac distribution at 0.*

REMARK 6.12. Our result above implies that there are only a few giant tables in the Chinese restaurant process (a mixed-Poisson number of tables with a number of customers proportional to $n^{\frac{1}{a+2}}$). In contrast, there are much more tables with a smaller number of customers, and an asymptotically negligible number of tables of size $\gg n^{\frac{1}{a+2}}$; see Figure 13.

REMARK 6.13. Closed formulas for the factorial moments $\mathbb{E}(X_{n,j}^{\underline{s}})$, as well as a formula for the probability mass function of $X_{n,j}$ are readily obtained from the generating functions by extraction of coefficients.

REMARK 6.14. Our results also allow recovering the limit theorem in [109] for the total number of tables C_n in the Chinese restaurant process (via X_n), albeit with a totally different proof, as the normalized random variable X_n/n^α converges in distribution with convergence of all moments to a random variable X , with X given in the theorem before. For the reader's




FIG 13. Schematic representation of the tables in the Chinese restaurant model: When the number of customers goes to infinity, one observes, with probability one, many tables with a small number of customers (coloured teal), only a few larger tables of size $\Theta(n^{\frac{1}{a+2}})$ (coloured purple), and no super-giant tables.

convenience, we state the moments in terms of θ and α , compare with [109, Theorem 3.8]:

$$\mathbb{E}(X^s) = \frac{\Gamma(s + \frac{\theta}{\alpha})\Gamma(\theta)}{\Gamma(\theta + s \cdot \alpha)\Gamma(\frac{\theta}{\alpha})}.$$

PROOF OF THEOREM 6.11. We follow very closely [83] and sketch the remaining steps. We solve the differential equation $T'(z) = \varphi(T(z))$ and get

$$T(z) = 1 - (1 - (a+1)z)^{\frac{1}{a+1}}.$$

Thus, the probability generating function is given by

$$\mathbb{E}(v^{X_{n+1,j}}) = \frac{n!}{T_{n+1}} [z^n] \psi \left(T(z) - (1-v)z^j \frac{T_j}{j!} \right),$$

where the coefficient

$$\frac{T_{n+1}}{n!} = [z^n] \psi(T(z))$$

is computed by standard singularity analysis. Therefore, except for the non-standard shift, we can readily apply our scheme to the generating function

$$\psi \left(T(z) - (1-v)z^j \frac{T_j}{j!} \right).$$

More precisely, since $a > 0$ and $b > -1$, we have $\lambda_H = \frac{1}{a+1}$ and $\lambda_G = -b$ for $b \neq 0$. Hence, the critical range is given by

$$j(n) = \Theta \left(n^{\frac{\lambda_H}{1+\lambda_H}} \right) = \Theta \left(n^{\frac{1}{a+2}} \right).$$

Here, no additional factor $M(z)$ is present, so $\lambda_M^- = 0$. In the case of $b = 0$ we apply the cycle scheme of Theorem 7.3. This gives

$$\mathbb{E}(X^s) = \mu_s = \begin{cases} \frac{\Gamma(s+b)\Gamma(\frac{b}{a+1})}{\Gamma(b)\Gamma(\frac{b+s}{a+1})} & \text{if } \beta \neq 0, \\ \frac{\Gamma(s+1)}{\Gamma(\frac{s}{a+1}+1)} & \text{if } \beta = 0. \end{cases}$$

Finally, we unify both expressions by simply using $\Gamma(x+1) = x\Gamma(x)$. □

6.7. *Triangular urn models and beta-Mittag-Leffler distributions.* Two-colour triangular urns are instances of generalized Pólya urn models [37, 62, 86]. At each time step $n \geq 1$, a ball is drawn uniformly at random, reinserted, and depending on the observed colour, balls of both colours are added to the urn: If a white ball was drawn, we add α white and β black balls, whereas, if a black ball was drawn, we add γ white and δ black balls. The addition/replacement of balls can be described by the so-called *ball replacement matrix*

$$M = \begin{pmatrix} \alpha & \beta \\ \gamma & \delta \end{pmatrix},$$

where for *balanced* urn models it holds that $\alpha + \beta = \gamma + \delta$, such that the total number $\sigma = \alpha + \beta$ of added balls in each step is independent of the observed colour. The initial configuration of the urn consists of w_0 white balls and b_0 black balls, and the random variable \mathcal{W}_n counts the number of white balls in the urn after n draws. For balanced triangular urns with replacement matrix

$$M = \begin{pmatrix} \alpha & \beta \\ 0 & \delta \end{pmatrix}, \quad \alpha, \beta > 0, \quad \delta = \sigma = \alpha + \beta,$$

it was shown by Flajolet, Dumas, and Puyhaubert [37] (and also by Janson [64, 66] via different analytic methods) that $\frac{\mathcal{W}_n}{\alpha n^{\alpha/\sigma}} \xrightarrow{d} \mathcal{W}$, for a random variable with moments

$$\mathbb{E}(\mathcal{W}^s) = \frac{\Gamma(\frac{b_0+w_0}{\sigma})}{\Gamma(\frac{w_0}{\alpha})} \cdot \frac{\Gamma(s + \frac{w_0}{\alpha})}{\Gamma(s \cdot \frac{\alpha}{\sigma} + \frac{b_0+w_0}{\sigma})}.$$

Figure 14 shows the evolution of the first three draws of a triangular urn.

In the special case $(w_0, b_0) = (\alpha, \beta)$, and thus $w_0 + b_0 = \sigma$, this simplifies to a Mittag-Leffler distribution with parameter $0 < \alpha/\sigma < 1$. As it is often the case for urn models, the limit law strongly depends on the initial composition of the urn. For $b_0 > 0$ and either $w_0 = 0$ or $w_0 = \beta$, Janson observed a moment-tilted stable law, leaving the other cases open; see [64, Theorem 1.8 and Problem 1.15].

We can directly obtain the limit law \mathcal{W} using our extended scheme. The key tool is the history generating function $F(z, u) = \sum_{n,k \geq 0} f_{n,k} u^k \frac{z^n}{n!}$ where $f_{n,k}$ is equal to the number of transitions (or histories) leading to a configuration with k white balls after n steps; see Figure 14.




FIG 14. The evolution of a balanced triangular urn with replacement matrix $M = \begin{pmatrix} 1 & 1 \\ 0 & 2 \end{pmatrix}$ and initially $w_0 = 2$ white balls and $b_0 = 1$ black ball. Dashed arrows indicate that a white ball was drawn, solid arrows indicate that a black ball was drawn. The number below each node corresponds to the number of possible transitions (or histories) to reach this state.

The closed form of this history generating function was derived in [37, Proposition 14]:

$$F(z, u) = u^{w_0} (1 - \sigma z)^{-b_0/\sigma} \left(1 - u^\alpha (1 - (1 - \sigma z)^{\alpha/\sigma}) \right)^{-w_0/\alpha}.$$

Putting aside the prefactor u^{w_0} , and after a change of variable $u^\alpha \mapsto u$, this equation can be interpreted as an extended critical composition scheme

$$F(z, u) = M(z) \cdot G(uH(z)), \quad (48)$$

involving the exponential generating functions with nonnegative integer coefficients

$$M(z) = (1 - \sigma z)^{-b_0/\sigma}, \quad G(z) = (1 - \alpha z)^{-w_0/\alpha}, \quad \text{and} \quad H(z) = (1 - (1 - \sigma z)^{\alpha/\sigma})/\alpha. \quad (49)$$

The fact that the singular exponents depend on b_0 and w_0 explains en passant why the limit distribution of \mathcal{W}_n differs according to the initial composition of the urn. Indeed, as the number of white balls at time n satisfies

$$\mathbb{P}\{\mathcal{W}_n = \alpha k + w_0\} = \frac{n! [z^n u^k] F(z, u)}{n! [z^n] F(z, 1)} = \frac{g_k [z^n] H(z)^k M(z)}{k! [z^n] F(z, 1)},$$

we can apply Theorem 4.1 and we then get the following limit distributions of \mathcal{W}_n for balanced triangular urn models, completing and extending earlier results [64, Theorem 1.8]:

COROLLARY 6.15. *Let \mathcal{W}_n be the random variable for the number of white balls in a balanced triangular Pólya urn with initially $w_0 > 0$ white and $b_0 \geq 0$ black balls. Then, we have a convergence in distribution, with convergence of all moments, towards a beta-Mittag-Leffler distribution (see Definition 3.13)*

$$\frac{\mathcal{W}_n}{\alpha n^{\alpha/\sigma}} \xrightarrow[m]{d} \text{BML} \left(\frac{\alpha}{\sigma}, \frac{w_0}{\alpha}, \frac{b_0}{\alpha} \right).$$

REMARK 6.16 (Almost sure convergence and beyond). This limit was also recently identified by Goldschmidt et al. [49] for urns with non-integer weights: A link with the Chinese restaurant model (for $b_0 = 0$) leads to a Mittag-Leffler distribution, then they show that the impact of $b_0 > 0$ on the process leads to a distribution with an additional beta law factor. It is interesting to stress that their approach implies an almost sure convergence. Note that the fluctuations around this almost sure limit are known: A second-order central limit theorem (that is, the random variable minus its almost sure limit converges, rescaled, to a Gaussian distribution), as well as a law of the iterated logarithm was obtained using discrete martingale [85]. What is more, following Gouet [50], a continuous-time reparametrization leads to a functional second-order limit theorem for balanced urn models. There is currently no systematic way to obtain an almost sure convergence for all combinatorial models covered by our composition schemes; however, in a few cases (e.g., for walks, trees, and maps), some ad-hoc clever constructions entail this almost sure convergence [88–90, 93, 113].

Note that applying Theorem 5.1 to the size-refined version of the composition scheme (48), we get factorial moments of mixed Poisson type for a size-refined random variable $X_{n,j}$ and the corresponding limit laws. However, the combinatorial interpretation of the random variable(s) $X_{n,j}$ is more involved and will be given elsewhere.

We stress the fact that the methods and results presented in this article are thus holding both for ordinary generating functions (typically used for unlabelled structures) and for exponential generating functions (typically used for labelled structures); see [40].

This concludes the list of applications for our results on the extended and size-refined composition schemes. We now give some extensions of our work to other schemes.

7. Further extensions.

7.1. *Critical cycle scheme.* Many combinatorial structures are cycles of more basic building blocks (e.g., cyclic permutations or functional applications are cycles of Cayley trees). If one marks the number of such basic building blocks, this corresponds to

$$\mathcal{F} = \mathcal{G}(\mathcal{H}) = \text{CYC}(\mathcal{H}) \implies F(z, u) = -\log(1 - uH(z)),$$

where $\mathcal{G} = \text{CYC}$ denotes the cycle operator. This scheme is analysed in Flajolet and Sedgewick's magnum opus [40, page 414] in the supercritical case, and we now extend this analysis to the critical case (i.e., by Definition 1.1 for $H(\rho_H) = 1$). Note that the previous sections were assuming Puiseux-like expansions for the generating function $F(z, 1)$ at its dominant singularity $z = \rho = \rho_H$. Now, for critical cycle schemes, F does not have a Puiseux expansion, so the previous results need to be adapted.

Let us begin with an example: For $H(z) = 1 - \sqrt{1 - 2z}$ we get the sequence

$$n![z^n]F(z) = n![z^n]\frac{1}{2}\log\left(\frac{1}{1-2z}\right) = (n-1)!2^{n-1} = (2n-2)!!, \quad n \geq 1,$$

which starts with 1, 2, 8, 48, 384, 3840, \dots , and constitutes the entry [A000165](#) in the OEIS. Here, the moments $\mathbb{E}(X_n^s)$ are of order $n^{s/2}$, so the scaling with $1/\sqrt{n}$ leads directly to moment convergence. This is just one instance of the following more general result.

THEOREM 7.1 (Critical schemes with a log). *In a critical cycle composition scheme*

$$F(z, u) = -\log(1 - uH(z)), \tag{50}$$

if $H(z)$ has a singular exponent $0 < \lambda_H < 1$, the core size X_n (i.e., the number of \mathcal{H} -components in structures of size n) has factorial moments given by

$$\mathbb{E}(X_n^s) \sim \kappa n^{s\lambda_H} \mu_s, \quad \text{with} \quad \kappa = \frac{1}{-c_H} \quad \text{and} \quad \mu_s = \frac{\Gamma(s+1)}{\Gamma(s\lambda_H+1)}.$$

The scaled random variable $X_n/(\kappa n^{\lambda_H})$ converges in distribution with convergence of all moments to a Mittag-Leffler distributed random variable $X \stackrel{d}{=} M_{\lambda_H}$.

REMARK 7.2. Observe that this scheme leads to a distribution similar (except for a shift in the moments) to the one obtained for the scheme involving the sequence operator $\mathcal{G} = \text{SEQ}$, i.e., $G(z) = \frac{1}{1-z}$, for which one has $c_G = 1$, $\lambda_G = -1$, and $\rho_G = 1$. Alternatively, we may think of this cycle scheme as the limit case of Theorem 4.1 when $\lambda_G \rightarrow 0$. Indeed, for $\lambda_M = 0$ the moments (21) of the extended composition scheme can be rewritten into

$$\mathbb{E}(X^s) = \frac{\Gamma(s - \lambda_G + 1)\Gamma(-\lambda_G\lambda_H + 1)}{\Gamma(s\lambda_H - \lambda_G\lambda_H + 1)\Gamma(-\lambda_G + 1)}.$$

Thus, for $\lambda_G \rightarrow 0$ the moments of the random variable X converge to the moments of an ordinary Mittag-Leffler distribution; see Definition 3.9. Similarly, taking the limit $\lambda_G \rightarrow 0$ in Remark 4.2 gives $\lambda_G = 0$ as the tilting parameter, resulting again in the ordinary Mittag-Leffler distribution.

Now, for $j \in \mathbb{N}$, we can also look at the size-refined scheme

$$\mathcal{F} = \text{CYC}(v\mathcal{H}_{=j} + \mathcal{H}_{\neq j}),$$

for which we get the following theorem.

THEOREM 7.3 (Size-refined critical schemes with a log). *In the size-refined critical cycle composition scheme*

$$F(z, v) = -\log\left(1 - (H(z) - (1-v)h_j z^j)\right), \quad j \in \mathbb{N}, \quad (51)$$

if $H(z)$ has a singular exponent $0 < \lambda_H < 1$, then the number $X_{n,j}$ of \mathcal{H} -components of size j in structures of size n has factorial moments of mixed Poisson type,

$$\mathbb{E}(X_{n,j}^s) = \xi_{n,j}^s \cdot \mu_s \cdot (1 + o(1)),$$

with $\xi_{n,j} = \frac{\rho_H^j}{c_H} h_j n^{\lambda_H}$ and Mittag-Leffler mixing distribution $X \stackrel{d}{=} M_{\lambda_H}$. The random variable $X_{n,j}$ converges to one of the three limit laws given in Theorem 5.1, depending on whether $j = j(n)$ is smaller, equal, or larger than the critical growth range $j = \Theta(n^{\frac{\lambda_H}{1+\lambda_H}})$.

PROOFS OF THEOREM 7.1 AND 7.3. The proofs are analogous to those of Theorems 4.1 and 5.1, and we only point out the differences next. Let us start with the proof of Theorem 7.1. The factorial moments of order s satisfy

$$\mathbb{E}(X_n^s) = \frac{[z^n] \partial_u^s (F)(z, 1)}{[z^n] F(z, 1)},$$

where F is defined by Equation (50). Hence, we first compute

$$F(z, 1) = -\log(1 - H(z)) = -\log\left((1 - z/\rho_H)^{\lambda_H} (1 + o(1))\right) \sim -\lambda_H \cdot \log\left(1 - z/\rho_H\right).$$

Using the transfer theorems of [40] we directly obtain $[z^n] F(z, 1) \sim \lambda_H \frac{\rho_H^{-n}}{n}$. It remains to compute $\partial_u^s F$. Note that the log function can be replaced by a quasi-inverse using

$$\partial_u^s \log\left(\frac{1}{1-u}\right) = \partial_u^{s-1} \frac{1}{1-u}.$$

Thus, the s th factorial moment is obtained from the asymptotics in (27) computed for $s-1$ and with $G(z) = \frac{1}{1-z}$. Then, we obtain the final result: the normalized moments converge to the moments of a Mittag-Leffler distribution.

For the proof of Theorem 7.3 one replaces (50) by (51) and $\partial_u^s F$ by $\partial_v^s F$. \square

7.2. Multivariate critical composition schemes. It is possible to generalize the critical composition scheme by looking at combinatorial constructions of the form

$$\mathcal{F} = \mathcal{M} \times \mathcal{G}_1(\mathcal{H}_1) \times \mathcal{G}_2(\mathcal{H}_2) \times \cdots \times \mathcal{G}_m(\mathcal{H}_m) = \mathcal{M} \times \prod_{\ell=1}^m \mathcal{G}_\ell(\mathcal{H}_\ell).$$

We measure the size of the \mathcal{G}_ℓ component by the variable u_ℓ ; accordingly this gives

$$F(z, u_1, \dots, u_m) = M(z) \cdot \prod_{\ell=1}^m G_\ell(u_\ell H_\ell(z)). \quad (52)$$

The random vector $\mathbf{X}_n = (X_{n,1}, \dots, X_{n,m})$ measures the sizes of the \mathcal{G}_ℓ -components,

$$\mathbb{P}\{X_{n,1} = k_1, \dots, X_{n,m} = k_m\} = \frac{[z^n u_1^{k_1} \cdots u_m^{k_m}] F(z, u_1, \dots, u_m)}{[z^n] F(z, 1, \dots, 1)}.$$

We now introduce a suitable extension of the terms *critical* (Definition 1.1) and *pure* (Definition 3.3) for multivariate schemes. We call a multivariate scheme *critical* if all functions $H_\ell(z)$ have the identical radius of convergence $\rho_{H_\ell} = \rho_H$ such that $\tau_\ell := H_\ell(\rho_H) = \rho_{G_\ell}$ and $M(z)$ has radius of convergence $\rho_M \geq \rho_H$. We call a multivariate scheme *pure* if

- $H_\ell(z)$ has a singular exponent $0 < \lambda_{H_\ell} < 1$ for $1 \leq \ell \leq m$;
- $G_\ell(z)$ has a singular exponent $\lambda_{G_\ell} < 0$ for $1 \leq \ell \leq m$;
- $M(z)$ has a singular exponent $\lambda_M \leq 0$ or $M(z)$ is analytic at ρ_H .

We can now state our multivariate result.

THEOREM 7.4. *In a multivariate pure extended critical composition scheme (52), the joint moments of the random vector $\mathbf{X}_n = (X_{n,1}, \dots, X_{n,m})$ satisfy*

$$\mathbb{E}(X_{n,1}^{s_1} \dots X_{n,m}^{s_m}) \sim \mu_{s_1, \dots, s_m} \prod_{\ell=1}^m n^{s_\ell \lambda_{H_\ell}} \kappa_\ell^{s_\ell},$$

with $\kappa_\ell = \frac{\tau_{H_\ell}}{-c_{H_\ell}}$ and μ_{s_1, \dots, s_m} given by

$$\mu_{s_1, \dots, s_m} = \frac{\Gamma(-\sum_{\ell=1}^m \lambda_{G_\ell} \lambda_{H_\ell} - \lambda_M)}{\Gamma(\sum_{\ell=1}^m s_\ell \lambda_{H_\ell} - \sum_{\ell=1}^m \lambda_{G_\ell} \lambda_{H_\ell} - \lambda_M)} \prod_{\ell=1}^m \frac{\Gamma(s_\ell - \lambda_{G_\ell})}{\Gamma(-\lambda_{G_\ell})}. \quad (53)$$

Consequently, one gets a convergence in distribution and in moments

$$\left(\frac{X_{n,1}}{\kappa_1 n^{\lambda_1}}, \dots, \frac{X_{n,m}}{\kappa_m n^{\lambda_m}} \right) \xrightarrow[m]{d} \mathbf{X}, \quad (54)$$

where \mathbf{X} is determined by its joint moment sequence $\mu_{\mathbf{s}} = \mu_{s_1, \dots, s_m}$. Moreover, the random vector \mathbf{X} has a scaled Dirichlet-stable product distribution,

$$\mathbf{X} = (X_1, \dots, X_m) \stackrel{d}{=} (V_1 \cdot W_1^{\lambda_{H_1}}, \dots, V_m \cdot W_m^{\lambda_{H_m}}), \quad (55)$$

where $\mathbf{W} = (W_1, \dots, W_m, W_{m+1})$ follows a Dirichlet distribution

$$\mathbf{W} \stackrel{d}{=} \text{Dir}(-\lambda_{G_1} \lambda_{H_1}, \dots, -\lambda_{G_m} \lambda_{H_m}, -\lambda_M), \quad (56)$$

and where the V_ℓ 's, for $1 \leq \ell \leq m$, are m independent generalized Mittag-Leffler distributions $V_\ell \stackrel{d}{=} \text{ML}(\lambda_{H_\ell}, -\lambda_{G_\ell} \lambda_{H_\ell})$, also independent of \mathbf{W} .

PROOF. We proceed similarly to the proof of the first part of Theorem 4.1. First, the mixed factorial moments of \mathbf{X}_n , which are obtained by differentiation and extraction of coefficients:

$$\mathbb{E}(X_{n,1}^{s_1} \dots X_{n,m}^{s_m}) = \frac{[z^n] \partial_{u_1}^{s_1} \dots \partial_{u_m}^{s_m} (F)(z, 1, \dots, 1)}{[z^n] F(z, 1, \dots, 1)}.$$

The differentiation with respect to u_ℓ only affects the factor $G_\ell(u_\ell H_\ell(z))$, leading to a singular expansion covered in Section 3.1. Extraction of coefficients then gives an asymptotic expansion of the factorial moments. Converting all the factorial moments into moments using (28) gives the desired asymptotics and moments in (53).

It remains to identify the distribution. To this aim, note that a Dirichlet distributed random vector $(W_1, \dots, W_{m+1}) \stackrel{d}{=} \text{Dir}(a_1, \dots, a_{m+1})$ with positive parameters a_1, \dots, a_{m+1} has a density function supported on the m simplex $\{(x_1, \dots, x_{m+1}) \in \mathbb{R}_{\geq 0}^{m+1} \mid \sum_{j=1}^{m+1} x_j = 1\}$:

$$f(x_1, \dots, x_{m+1}) = \frac{\Gamma(\sum_{j=1}^{m+1} a_j)}{\prod_{j=1}^{m+1} \Gamma(a_j)} \prod_{j=1}^{m+1} x_j^{a_j-1}.$$

Accordingly, the joint moments are given by

$$\mathbb{E}(W_1^{s_1} \dots W_{m+1}^{s_{m+1}}) = \frac{\Gamma(\sum_{j=1}^{m+1} a_j)}{\Gamma(\sum_{j=1}^{m+1} (s_j + a_j))} \frac{\prod_{j=1}^{m+1} \Gamma(s_j + a_j)}{\prod_{j=1}^{m+1} \Gamma(a_j)}.$$

Now, consider a random vector (Z_1, \dots, Z_m) satisfying

$$(Z_1, \dots, Z_m) \stackrel{d}{=} (V_1 \cdot W_1^{\alpha_1}, \dots, V_m \cdot W_m^{\alpha_m})$$

with generalized Mittag-Leffler distributions $V_\ell \stackrel{d}{=} \text{ML}(\alpha_\ell, a_\ell)$ for $\ell = 1, \dots, m$, such that all random variables are mutually independent. Using the closed form (18) we get

$$\mathbb{E}(Z_1^{s_1} \dots Z_m^{s_m}) = \frac{\Gamma(\sum_{\ell=1}^m a_\ell)}{\Gamma(a_{m+1} + \sum_{\ell=1}^m (\alpha_\ell s_\ell + a_\ell))} \prod_{\ell=1}^m \frac{\Gamma(s_\ell + a_\ell/\alpha_\ell)}{\Gamma(a_\ell/\alpha_\ell)}.$$

Comparing this expression with the moments (53), the claim follows. \square

REMARK 7.5. The marginals X_ℓ of the random vector \mathbf{X} are also covered by Theorem 4.1. The random vector \mathbf{X} is closely related to Poisson–Dirichlet distributions $\text{PD}(\alpha, \theta)$, [60, 61, 110] and the joint limit law of node degrees in preferential attachment trees or generalized plane-oriented recursive trees [96]; see also the subsequent example.

Now, if one considers the multivariate size-refined scheme

$$\mathcal{F} = \mathcal{M} \times \prod_{\ell=1}^m \mathcal{G}_\ell(\mathcal{H}_{\ell, \neq j_\ell} + v_\ell \mathcal{H}_{\ell, =j_\ell}),$$

one gets the following multivariate version of Theorem 5.1.

THEOREM 7.6 (Multivariate pure size-refined critical scheme). *In a multivariate pure size-refined critical composition scheme*

$$F(z, v_1, \dots, v_m) = M(z) \cdot \prod_{\ell=1}^m G_\ell(H_\ell(z) - (1 - v_\ell)h_{\ell, j_\ell} z^{j_\ell})$$

the random variables X_{n, ℓ, j_ℓ} , which count the number of H_ℓ -components of size j_ℓ , have joint factorial moments of mixed Poisson type:

$$\mathbb{E}(X_{n, 1, j_1}^{s_1} \dots X_{n, m, j_m}^{s_m}) = \mu_{s_1, \dots, s_m} \cdot \prod_{\ell=1}^m \xi_{n, \ell, j_\ell}^{s_\ell} \cdot (1 + o(1)),$$

with $\xi_{n, \ell, j_\ell} = \frac{\rho_{H_\ell}^{j_\ell}}{-c_{H_\ell}} h_{\ell, j_\ell} n^{\lambda_{H_\ell}}$ and joint mixing distribution $\mathbf{X} = (X_1, \dots, X_m)$ as in Equation (55).

Let X_ℓ , for $1 \leq \ell \leq m$, denote the marginal distribution of the ℓ th coordinate of $\mathbf{X} = (X_1, \dots, X_m)$. For $n \rightarrow \infty$, the limiting distributions of X_{n, ℓ, j_ℓ} jointly undergo mixed Poisson type phase transitions with mixing distributions X_ℓ . The phase transitions depend on the growth of $j_\ell = j_\ell(n)$, with critical growth ranges given by $j_\ell = j_\ell(n) = \Theta(n^{\frac{\lambda_{H_\ell}}{1+\lambda_{H_\ell}}})$.

In particular, for $j_\ell(n) \sim \xi_\ell \cdot n^{\frac{\lambda_{H_\ell}}{1+\lambda_{H_\ell}}}$, the random vector $\mathbf{X}_{n, \mathbf{j}} = (X_{n, 1, j_1}, \dots, X_{n, m, j_m})$ converges in distribution with convergence of all (factorial) moments to a multivariate distribution $\text{MPo}(\boldsymbol{\xi}, \mathbf{X})$.

For properties of multivariate mixed Poisson distributions we refer to [34] or [83]. The key tool for proving Theorem 7.6 is the following multivariate extension of Lemma 5.4.

LEMMA 7.7 (Joint factorial moments and limit laws of mixed Poisson type). *Let $(\mathbf{X}_n)_{n \in \mathbb{N}}$ denote a sequence of m -dimensional random vectors, whose factorial moments are asymptotically of mixed Poisson type, i.e., they satisfy for $n \rightarrow \infty$ the asymptotic expansion*

$$\mathbb{E}(\mathbf{X}_n^{\mathbf{s}}) = \mathbb{E}(X_{n, 1}^{s_1} \dots X_{n, m}^{s_m}) = \mu_{s_1, \dots, s_m} \cdot \prod_{\ell=1}^m \xi_{n, \ell}^{s_\ell} \cdot (1 + o(1)), \quad s \geq 1,$$

with $\mu_{s_1, \dots, s_m} \geq 0$, and $\xi_{n, \ell} > 0$ for $1 \leq \ell \leq m$. Furthermore, assume that the sequence of joint moments $(\mu_{\mathbf{s}})_{\mathbf{s} \in \mathbb{N}^m}$ determines a unique distribution $\mathbf{L} = (L_1, \dots, L_m)$. Then, for $n \rightarrow \infty$, one has the following joint limit distributions:

- (i) If $\xi_{n, \ell} \rightarrow \infty$, the random variable $\frac{X_{n, \ell}}{\xi_{n, \ell}}$ converges in distribution, with convergence of all moments, to L_ℓ .
- (ii) If $\xi_{n, \ell} \rightarrow \xi \in (0, \infty)$, the random variable $X_{n, \ell}$ converges in distribution, with convergence of all moments, to $Y \stackrel{d}{=} \text{MPo}(\xi L_\ell)$.
- (iii) If $\xi_{n, \ell} \rightarrow 0$, $X_{n, \ell}$ converges to a Dirac distribution: $X_{n, \ell} \xrightarrow{d} 0$.

PROOF. We follow the proof of the univariate case [83, Lemma 2]. Assume that the set $\{1, \dots, m\}$ decomposes into two disjoint sets C and D such that for indices $k \in C$ we have $\lambda_{n,k} \rightarrow \infty$, whereas for an index $k \in D$ it holds $\lambda_{n,k} \rightarrow \rho_k$ for $n \rightarrow \infty$. We observe that the mixture of joint raw moments and joint factorial moments converge:

$$\mathbb{E} \left(\left(\prod_{k \in C} \frac{X_{n,k}^{s_k}}{\lambda_{n,k}^{s_k}} \right) \cdot \left(\prod_{k \in D} X_{n,k}^{s_k} \right) \right) \rightarrow \mu_{s_1, \dots, s_m} \cdot \prod_{k \in D} \rho_k^{s_k}.$$

The latter joint moment sequence, both raw and factorial moments, is exactly the joint moment sequence of a random vector $\mathbf{Z} = (Z_1, \dots, Z_m)$, with

$$\forall k \in C: Z_k \stackrel{d}{=} L_k, \quad \forall k \in D: Z_k \stackrel{d}{=} \text{MPo}(\rho_k L_k),$$

such that

$$\mathbb{E} \left(\left(\prod_{k \in C} Z_k^{s_k} \right) \cdot \left(\prod_{k \in D} Z_k^{s_k} \right) \right) = \mu_{s_1, \dots, s_m} \cdot \prod_{k \in D} \rho_k^{s_k}. \quad \square$$

PROOF OF THEOREM 7.6. The proof is very similar to proofs of Theorems 5.1 and 7.4, so we will be brief again. The mixed factorial moments of \mathbf{X}_{n, j_ℓ} are obtained by differentiation and extraction of coefficients:

$$\mathbb{E}(X_{n,1,j_1}^{s_1} \cdots X_{n,m,j_m}^{s_m}) = \frac{[z^n] \partial_{v_1}^{s_1} \cdots \partial_{v_m}^{s_m} (F)(z, 1, \dots, 1)}{[z^n] F(z, 1, \dots, 1)}.$$

The differentiation with respect to v_ℓ only affects the factor

$$G_\ell(H_\ell(z) - (1 - v_\ell)h_{\ell, j_\ell} z^{j_\ell}),$$

leading to singular expansions covered in Section 3.1 and additional factors $h_{\ell, j_\ell}^{s_\ell}$, $1 \leq \ell \leq m$. The asymptotics of these factors for $j_\ell \rightarrow \infty$ are all governed by (37). The individual singular expansions are similar to (38). Thus, extracting the coefficient $z^{n-s_1 j_1 - \dots - s_m j_m}$ from

$$M(z) \cdot \prod_{\ell=1}^m G_\ell^{(s_\ell)}(H_\ell(z))$$

leads to the asymptotic expansion of the joint factorial moments. Lemma 7.7 then yields the stated limit law. \square

We end this section with four examples covered by the multivariate scheme.

EXAMPLE 7.8 (m -bundled plane-oriented recursive trees). We have previously encountered 3-bundled trees in Section 6.2 in the framework of bilabelled trees. In the following we study ordinary increasing trees [16, 27, 68, 76, 87, 101], where each node has only one label. As before, given a degree-weight sequence $(\varphi_j)_{j \geq 0}$, the corresponding degree-weight generating function is defined by $\varphi(t) = \sum_{j \geq 0} \varphi_j t^j$. The associated family \mathcal{T} of increasing trees can be described by the following symbolic equation using the boxed product (see Section 6.2):

$$\mathcal{T} = \mathcal{Z}^\square * \varphi(\mathcal{T}).$$

For the exponential generating function

$$T(z) = \sum_{n \geq 1} T_n \frac{z^n}{n!},$$

we thus have the following equation

$$T'(z) = \varphi(T(z)), \quad T(0) = 0.$$

We are interested in families of generalized plane-oriented recursive trees with degree-weight generating function $\varphi(t) = 1/(1-t)^m$, with $m \in \mathbb{N}$.

For $m = 1$ we get the ordinary plane-oriented recursive tree, whereas for $m > 1$ we get the so-called m -bundled trees, with generating function

$$T(z) = 1 - (1 - (m + 1)z)^{1/(m+1)}.$$

One may think of each node holding $m - 1$ additional separation bars [68], which can be regarded as a special edge type. Naturally, this refines the root degree X_n in a random tree of size n , since we may look at the number of nodes attached to the root in a specific cluster, induced by the $m - 1$ bars: $X_n = \sum_{\ell=1}^m X_{n,\ell}$.

By construction, the random variables $X_{n,\ell}$ are exchangeable, but not independent. Writing $\mathbf{u}^{\mathbf{X}_n} = u_1^{X_{n,1}} \dots u_m^{X_{n,m}}$, the generating function

$$T(z, \mathbf{u}) = \sum_{n \geq 1} T_n \mathbb{E}(\mathbf{u}^{\mathbf{X}_n}) \frac{z^n}{n!}$$

of the random vector $\mathbf{X}_n = (X_{n,1}, \dots, X_{n,m})$ is given by

$$\frac{\partial}{\partial z} T(z, \mathbf{u}) = \prod_{\ell=1}^m \frac{1}{1 - u_\ell T(z)}.$$

This refinement is covered by the multivariate scheme (52) (with a shift: $X_{n+1,\ell}$, instead of $X_{n,\ell}$). Similarly, one may study the outdegree of a node labelled j , as well as multiple nodes, leading to closely related generating functions [61, 76, 96]. ■

EXAMPLE 7.9 (Bilabelled increasing trees and refined root degree). Continuing from Section 6.2, we can refine the root-degree X_n in 3-bundled bilabelled increasing trees of size $2n$, $X_n = X_{n,1} + X_{n,2} + X_{n,3}$, where the $X_{n,\ell}$ are exchangeable. The corresponding generating function

$$T(z, u_1, u_2, u_3) = \sum_{n \geq 1} T_n \mathbb{E}(u_1^{X_{n,1}} u_2^{X_{n,2}} u_3^{X_{n,3}}) \frac{z^n}{n!}$$

then satisfies

$$\frac{\partial^2}{\partial z^2} T(z, u_1, u_2, u_3) = \prod_{\ell=1}^3 \frac{1}{1 - u_\ell (1 - \sqrt{1 - z^2})}.$$

Except for the non-standard shift in the random variable, the problem corresponds directly to a multivariate pure critical composition scheme. ■

EXAMPLE 7.10 (Returns in coloured bridges and walks). Consider k -coloured bridges B_k as defined in Section 6.4: A bridge is coloured in exactly k colours, where each colour consists of a non-empty bridge. Combinatorially, we append k non-empty bridges one after the other. We are interested in the individual number of returns to zero in each of the first k_1 bridges, followed by k_2 additional bridges, such that $k_1 + k_2 = k$. Reusing the combinatorial construction (47), the multivariate generating function $B_k(z, \mathbf{u})$ satisfies

$$B_k(z, u_1, \dots, u_{k_1}) = \left(\prod_{j=1}^{k_1} \left(\frac{1}{1 - u_j \left(1 - \frac{1}{B(z)}\right)} - 1 \right) \right) (B(z) - 1)^{k_2}.$$

By our previous reasoning we see that the corresponding random variables are exchangeable and the multivariate scheme directly applies. Moreover, we can also consider walks and a refined generating function $W_k(z, \mathbf{u})$ of k -coloured walks with the tail coloured in the same colour as the final bridge, keeping track of the individual returns to zero of the first k_1 bridges,

$$W_k(z, \mathbf{u}) = (1 + B_k(z, \mathbf{u})) \frac{W(z)}{B(z)}.$$

Again, the corresponding random variables are exchangeable and the multivariate scheme directly applies again. ■

EXAMPLE 7.11 (Triangular urn models and node degrees in generalized recursive trees). We discuss a specific balanced triangular urn model with $k + 1$ colours, which generalizes the case with 2 colours from Section 6.7. Our urn model is specified by the following $(k + 1) \times (k + 1)$ balanced ball replacement matrix M with $\alpha_r, \beta_r \in \mathbb{N}$ and $\alpha_r + \beta_r = \sigma$, $1 \leq r \leq k + 1$:

$$M = \begin{pmatrix} \alpha_1 & 0 & \dots & 0 & \beta_1 \\ 0 & \alpha_2 & \dots & 0 & \beta_2 \\ \vdots & \ddots & \ddots & & \vdots \\ 0 & \dots & 0 & \alpha_k & \beta_k \\ 0 & 0 & \dots & 0 & \alpha_{k+1} + \beta_{k+1} \end{pmatrix} \quad (57)$$

We assume that there are initially $A_{0,r} = a_r$ balls of type r for $1 \leq r \leq k + 1$ and that the random variables $A_{n,r}$ count the number of balls of type r after n draws. As before, we use the history-counting approach [37, 38] to analyse the urn model. For this purpose we will first derive the history generating function $H(x_1, \dots, x_k, x_{k+1}; z)$. Due to the balance condition, it suffices to study the function $H(x_1, \dots, x_k, 1; z)$. In the associated differential system, the functions $x_\ell(t)$ satisfy

$$\dot{x}_r(z) = x_r^{\alpha_r+1} x_{k+1}^{\beta_r}(z) \text{ (for } 1 \leq r \leq k) \text{ and } \dot{x}_{k+1}(z) = x_{k+1}^{\sigma+1}(z),$$

with initial conditions $x_\ell(0) = x_{0,\ell}$. Now, by separation of variables, we directly obtain¹⁴

$$x_{k+1}(z) = x_{0,k+1} \cdot (1 - \sigma x_{0,k+1}^\sigma z)^{-1/\sigma}.$$

By integration, we readily get the other functions $x_r(z)$ for $1 \leq r \leq k$:

$$x_r(z) = x_{0,r} \cdot \left(1 - x_{0,r}^{\alpha_r} x_{0,k+1}^{-\alpha_r} (1 - (1 - \sigma x_{0,k+1}^\sigma z)^{\alpha_r/\sigma})\right)^{-1/\alpha_r}.$$

Finally, using the basic isomorphism between differential systems and history generating functions [37], we obtain the desired closed form for the generating function of urn histories, which falls into the class of our multivariate pure critical composition scheme Theorem 7.4.

PROPOSITION 7.12. *The history generating function $H(x_1, \dots, x_k, 1; z)$, associated with the balanced triangular urn model with ball replacement matrix M in (57) and initial conditions $A_{0,r} = a_r$, $1 \leq r \leq k + 1$, is given by*

$$H(x_1, \dots, x_k, 1; z) = (1 - \sigma z)^{-\frac{a_{k+1}}{\sigma}} \cdot \prod_{r=1}^k x_r^{a_r} \left(1 - x^{\alpha_r} (1 - (1 - \sigma z)^{\alpha_r/\sigma})\right)^{-\frac{a_r}{\alpha_r}}.$$

It is well known that node degrees in generalized plane-oriented recursive trees can be modelled by such urns [63, 96]. As an application, we set $\alpha_r = 1$ and $\beta_r = k$ for $1 \leq r \leq k + 1$ and $a_r = 1$, $1 \leq r \leq k$ and $a_{k+1} = 0$ for the initial values. Then, the random variables $A_{n,1}$ up to $A_{n,k}$ are exchangeable and count the refined root degree of k -bundled plane-oriented recursive trees, i.e., the number of children of the root in each bundle.

Moreover, the urn model M can be used to study the joint degree distribution of the nodes labelled $1, 2, \dots, k$ in generalized plane-oriented recursive trees; this will be discussed in a forthcoming paper [73]. ■

¹⁴There is a small sign typo in [37, page 36] where, in the corresponding result for the 2×2 model, $y_0^{-\sigma}$ should be replaced by y_0^σ .

8. Conclusion. This work makes explicit the limit laws for combinatorial structures counted by schemes like $G(H(z))M(z)$, where we additionally mark either the number of \mathcal{H} -components, or the number of \mathcal{H} -components of a given size. We focused on the technically more delicate and mathematically richer case where G , H , and M are simultaneously singular. We proved that when these functions have algebraic dominant singularities with exponents between 0 and 1, the limit laws of the schemes are moment-tilted stable distributions and product distributions involving Mittag-Leffler laws. In Table 4 we give an overview of the covered schemes and the corresponding limit laws, where we have convergence in distribution *and* convergence of all moments.

In the extended scheme, in which one follows the total number of \mathcal{H} -components, we proved the appearance of three régimes for the corresponding limit law (continuous, Boltzmann, and linear combination of a continuous and a Boltzmann distribution), depending on the relation of the singular exponents of G , H , and M ; see Table 1. In the size-refined scheme, in which one follows the number of \mathcal{H} -components of a given size, we proved the appearance of mixed Poisson distributions (see Definition 3.15) and a double phase transition for the limit law from continuous to discrete to degenerate, with explicit threshold sizes depending on the exponents; see Table 2.

We also presented several extensions (logarithmic singularities, multivariate cases) and a variety of applications to important probabilistic and combinatorial objects. This allowed us to obtain new results for the core size of supertrees, the number of subtrees in different increasing trees, the returns to zero and sign changes in walks and bridges, the table sizes in the Chinese restaurant process, and the number of balls in some urn models.

Composition scheme	Symbolic form	Limit law	Thm.
Ordinary	$F(z, u) = G(uH(z))$	generalized Mittag-Leffler	4.2
Extended	$F(z, u) = M(z)G(uH(z))$	beta-Mittag-Leffler and Boltzmann distribution	4.1 4.3 4.4
Cyclic	$F(z, u) = -\log(1 - uH(z))$	Mittag-Leffler	7.1
Multivariate extended	$F(z, \mathbf{u}) = M(z) \prod_{\ell=1}^m G_{\ell}(u_{\ell}H_{\ell}(z))$	multivariate product distribution	7.4
Size-refined	$F(z, v) = M(z)G(H(z) - z^j h_j(1 - v))$	mixed Poisson type phase transition	5.1
Size-refined cyclic	$F(z, v) = -\log(1 - (H(z) - (1 - v)h_j z^j / j!))$	mixed Poisson type phase transition	7.3
Multivariate size-refined	$F(z, \mathbf{v}) = M(z) \prod_{\ell=1}^m G_{\ell}(H_{\ell}(z) - z^{j_{\ell}} h_{\ell, j_{\ell}}(1 - v_{\ell}))$	mv. mixed Poisson type phase transition	7.6

TABLE 4

Overview of our results on critical composition schemes, where $\mathbf{u} = (u_1, \dots, u_m)$ and $\mathbf{v} = (v_1, \dots, v_m)$.

The results of Table 4 hold for functions $H(z)$ having a dominant singularity of algebraic type. Our methods can also deal with schemes having other types of singularities. For example, one could allow algebraic-logarithmic behaviours of the form

$$H(z) \sim \left(1 - \frac{z}{\rho_H}\right)^{\lambda_H} \left(\frac{1}{z} \log\left(1 - \frac{z}{\rho_H}\right)\right)^{\psi_H}.$$

This would cover some instances of 3-colour balanced triangular urn models [37, 112], whose complete analysis, however, remains a challenge as other types of singularities appear, thus leading to new families of limit laws whose nature is unclear. Compare also with the related open problem by Janson [66], asking for a more detailed description of the limit law of unbalanced 2-colour triangular urn models.

To keep this article readable, we eluded the question of the speed of convergence of X_n , properly normalized, to its limit law X . Recently, a few articles addressed this for some preferential attachment rules [105] and for the Chinese restaurant process [26] (see Remark 6.16). In fact, following the Puiseux expansions at order 2 of $G/H/M$ allow us to capture this speed of convergence, but this leads to many subcases. In a future work we plan to give uniform bounds on the moments, leading to Berry–Esseen-like inequalities for the speed of convergence of all the limit laws associated with critical composition schemes; see also [1, 54].

Last but not least, in our companion article [9], we further extend our analysis to schemes with algebraic singularities, where $\lambda_H > 1$: We analyse a generalization of the composition scheme analysed in [6], leading to stable laws (e.g., the map-Airy distribution), Gaussian laws, as well as bimodal distributions. In extended composition schemes we observe a new behaviour, obtaining for example beta limit laws. In size-refined schemes we anticipate further continuous to discrete phase transitions. This will achieve the exhaustive exploration of the landscape of critical composition schemes with algebraic singularities and associated phase transitions.

Acknowledgments. We are pleased to dedicate this article to our colleague/mentor Alois Panholzer, on the occasion of his 50th birthday. In fact, Alois played a central rôle in the birth of this article in several ways:

- As a byproduct of Alois’ analyses of combinatorial structures and stochastic processes (such as permutations, parking functions, lattice paths, urns, and trees), several phase transitions involving the mixed Poisson distribution were uncovered [83, 102]. This guided the second author to study a general framework for these phase transitions, and to join forces with the other authors to extend the results of [6, 118] involving Rayleigh, half-normal, and stable distributions. This led us to the composition schemes analysed in this work.
- Alois also connected the first and second authors via a French-Austrian international project (PHC Amadeus) in 2005/06, perpetuating a long tradition of exchanges in combinatorics between Austria and France. A part of this project led to the article [8], which established Gaussian limit laws for some composition schemes related to trees.
- Alois was also a constant source of inspiration and guidance for the second author. Without him, we would have never started this work.

Funding. Michael Wallner was funded by the FWF Erwin Schrödinger Fellowship J 4162 and the FWF Stand-alone Project P 34142.

References.

- [1] José A. Adell and Jesús de la Cal. [On the uniform convergence of normalized Poisson mixtures to their mixing distribution](#). *Statistics & Probability Letters*, 18:227–232, 1993.
- [2] David J. Aldous. [Exchangeability and related topics](#), volume 1117 of *Lecture Notes in Mathematics*. Springer, 1985.
- [3] Richard Arratia, Andrew D. Barbour, and Simon Tavaré. [Logarithmic Combinatorial Structures: a Probabilistic Approach](#). EMS Monographs in Mathematics. European Mathematical Society, 2003.
- [4] Cyril Banderier and Michael Drmota. [Formulae and asymptotics for coefficients of algebraic functions](#). *Combin. Probab. Comput.*, 24(1):1–53, 2015.
- [5] Cyril Banderier and Philippe Flajolet. [Basic analytic combinatorics of directed lattice paths](#). *Theoret. Comput. Sci.*, 281(1-2):37–80, 2002.
- [6] Cyril Banderier, Philippe Flajolet, Gilles Schaeffer, and Michèle Soria. [Random maps, coalescing saddles, singularity analysis, and Airy phenomena](#). *Random Struct. Algorithms*, 19(3-4):194–246, 2001.
- [7] Cyril Banderier and Paweł Hitczenko. [Enumeration and asymptotics of restricted compositions having the same number of parts](#). *Discrete Appl. Math.*, 160(18):2542–2554, 2012.
- [8] Cyril Banderier, Markus Kuba, and Alois Panholzer. [Analysis of three graph parameters for random trees](#). *Random Struct. Algorithms*, 35(1):42–69, 2009.
- [9] Cyril Banderier, Markus Kuba, and Michael Wallner. Phase transitions of composition schemes: stable laws and Boltzmann distributions. *In preparation*, 2022.
- [10] Cyril Banderier, Philippe Marchal, and Michael Wallner. [Periodic Pólya urns, the density method and asymptotics of Young tableaux](#). *Ann. Probab.*, 48(4):1921–1965, 2020.
- [11] Cyril Banderier and Michael Wallner. [Some reflections on directed lattice paths](#). In *Proceedings of the 25th International Conference on Probabilistic, Combinatorial and Asymptotic Methods for the Analysis of Algorithms (AofA'14)*, pages 25–36. Discrete Math. Theor. Comput. Sci., 2014.
- [12] Cyril Banderier and Michael Wallner. [Lattice paths with catastrophes](#). *Discrete Math. Theor. Comput. Sci.*, Vol. 19 no. 1, 2017.
- [13] Cyril Banderier and Michael Wallner. [Local time for lattice paths and the associated limit laws](#). In *Proceedings of GASCom 2018*, volume 2113, pages 69–78. CEUR Workshop Proceedings, 2018.
- [14] Cyril Banderier and Michael Wallner. [The kernel method for lattice paths below a rational slope](#). In *Lattice paths combinatorics and applications*, volume 58 of *Developments in Mathematics Series*, pages 119–154. Springer, 2019.
- [15] Edward A. Bender. [Central and local limit theorems applied to asymptotic enumeration](#). *J. Comb. Theory, Ser. A*, 15:91–111, 1973.
- [16] François Bergeron, Philippe Flajolet, and Bruno Salvy. [Varieties of increasing trees](#). In *Colloque sur les arbres en algèbre et programmation (CAAP'92)*, volume 581 of *Lect. Notes Comput. Science*, pages 24–48. Springer, 1992.
- [17] Mira Bernstein and Neil J. A. Sloane. [Some canonical sequences of integers](#). *Linear Algebra Appl.*, 226-228:57–72, 1995.
- [18] Olivier Bodini, Matthieu Dien, Xavier Fontaine, Antoine Genitrini, and Hsien-Kuei Hwang. [Increasing diamonds](#). In *LATIN 2016*, volume 9644 of *Lect. Notes Comput. Science*, pages 207–219. Springer, 2016.
- [19] Olivier Bodini, Olivier Roussel, and Michèle Soria. [Boltzmann samplers for first-order differential specifications](#). *Discrete Appl. Math.*, 160(18):2563–2572, 2012.

- [20] Mireille Bousquet-Mélou. [Limit laws for embedded trees: applications to the integrated superBrownian excursion](#). *Random Struct. Algorithms*, 29(4):475–523, 2006.
- [21] Torsten Carleman. *Sur les équations intégrales singulières à noyau réel et symétrique*. Uppsala Universitets Årsskrift, 1923.
- [22] Donald A. Darling and Mark Kac. [On occupation times for Markoff processes](#). *Trans. Amer. Math. Soc.*, 84:444–458, 1957.
- [23] Hervé Daudé and Vlady Ravelomanana. [Random 2-XORSAT phase transition](#). *Algorithmica*, 59:48–65, 2011.
- [24] Timothy DeVries. *Algorithms for Bivariate Singularity Analysis*. PhD thesis, Univ. of Pennsylvania, 2011.
- [25] *NIST Digital Library of Mathematical Functions*. <http://dlmf.nist.gov/>, release 1.1.2 of 2021-06-15. F. W. J. Olver, A. B. Olde Daalhuis, D. W. Lozier, B. I. Schneider, R. F. Boisvert, C. W. Clark, B. R. Miller, B. V. Saunders, H. S. Cohl, and M. A. McClain, eds.
- [26] Emanuele Dolera and Stefano Favaro. [A Berry–Esseen theorem for Pitman’s \$\alpha\$ -diversity](#). *Ann. Appl. Probab.*, 30(2):847–869, 2020.
- [27] Michael Drmota. *Random Trees*. Springer, 2009.
- [28] Michael Drmota and Michèle Soria. [Marking in combinatorial constructions: generating functions and limiting distributions](#). *Theoret. Comput. Sci.*, 144(1-2):67–99, 1995.
- [29] Michael Drmota and Michèle Soria. [Images and preimages in random mappings](#). *SIAM J. Discrete Math.*, 10(2):246–269, 1997.
- [30] Jules Dubourdieu. *Les principes fondamentaux du calcul des probabilités et la théorie de l’assurance-maladie*. Gauthier-Villars, 1939.
- [31] Philippe Duchon, Philippe Flajolet, Guy Louchard, and Gilles Schaeffer. [Boltzmann samplers for the random generation of combinatorial structures](#). *Combin. Probab. Comput.*, 13(4-5):577–625, 2004.
- [32] William Feller. [Fluctuation theory of recurrent events](#). *Trans. Amer. Math. Soc.*, 67:98–119, 1949.
- [33] William Feller. *An Introduction to Probability Theory and its Applications. Vol. II*. Second edition (1st ed.: 1966). John Wiley & Sons, 1971.
- [34] Anthony Ferrari, Gérard Letac, and Jean-Yves Tourneret. [Multivariate mixed Poisson distributions](#). In *12th European Signal Processing Conference, IEEE*, pages 1067–1070, 2004.
- [35] James Allen Fill, Philippe Flajolet, and Nevin Kapur. [Singularity analysis, Hadamard products, and tree recurrences](#). *J. Comput. Appl. Math.*, 174(2):271–313, 2005.
- [36] Philippe Flajolet. [Some exactly solvable models of urn process theory](#). Slides for the conference Mathinfo in Nancy, 2006.
- [37] Philippe Flajolet, Philippe Dumas, and Vincent Puyhaubert. [Some exactly solvable models of urn process theory](#). In *Fourth Colloquium on Mathematics and Computer Science Algorithms, Trees, Combinatorics and Probabilities*, pages 59–118. Discrete Math. Theor. Comput. Sci., 2006.
- [38] Philippe Flajolet, Joaquim Gabarró, and Helmut Pekari. [Analytic urns](#). *Ann. Probab.*, 33(3):1200–1233, 2005.
- [39] Philippe Flajolet and Andrew Odlyzko. [Singularity analysis of generating functions](#). *SIAM J. Discrete Math.*, 3(2):216–240, 1990.
- [40] Philippe Flajolet and Robert Sedgewick. *Analytic Combinatorics*. Cambridge University Press, 2009.
- [41] Philippe Flajolet and Michèle Soria. [Gaussian limiting distributions for the number of components in combinatorial structures](#). *J. Combin. Theory Ser. A*, 53(2):165–182, 1990.

- [42] Philippe Flajolet and Michèle Soria. [General combinatorial schemas: Gaussian limit distributions and exponential tails](#). *Discrete Math.*, 114(1-3):159–180, 1993.
- [43] Maurice Fréchet and James Shohat. [A proof of the generalized second limit theorem in the theory of probability](#). *Trans. Amer. Math. Soc.*, 33(2):533–543, 1931.
- [44] Michael Fuchs. [Subtree sizes in recursive trees and binary search trees: Berry–Esseen bounds and Poisson approximations](#). *Combin. Probab. Comput.*, 17(5):661–680, 2008.
- [45] Michael Fuchs. [Limit theorems for subtree size profiles of increasing trees](#). *Combin. Probab. Comput.*, 21(3):412–441, 2012.
- [46] Michael Fuchs, Hsien-Kuei Hwang, and Ralph Neininger. [Profiles of random trees: Limit theorems for random recursive trees and binary search trees](#). *Algorithmica*, 46(3-4):367–407, 2006.
- [47] Omer Giménez and Marc Noy. [Asymptotic enumeration and limit laws of planar graphs](#). *J. Am. Math. Soc.*, 22(2):309–329, 2009.
- [48] Omer Giménez, Marc Noy, and Juanjo Rué. [Graph classes with given 3-connected components: asymptotic enumeration and random graphs](#). *Random Struct. Algorithms*, 42(4):438–479, 2013.
- [49] Christina Goldschmidt, Bénédicte Haas, and Delphin Sénizergues. [Stable graphs: distributions and line-breaking construction](#). *arXiv*, 2020.
- [50] Raúl Gouet. [Martingale Functional Central Limit Theorems for a Generalized Pólya Urn](#). *Ann. Probab.*, 21:1624–1639, 1993.
- [51] Jan Grandell. *Mixed Poisson Processes*. Chapman & Hall, 1997.
- [52] Theodore Edward Harris. *The Theory of Branching Processes*. Springer, 1963.
- [53] Pierre Humbert. [Nouvelles correspondances symboliques](#). *Bull. Sci. Math., II. Sér.*, 69:121–129, 1945.
- [54] Hsien-Kuei Hwang. [Second phase changes in random m-ary search trees and generalized quicksort: convergence rates](#). *Ann. Probab.*, 31(2):609–629, 1998.
- [55] Hsien-Kuei Hwang. [Asymptotics of Poisson approximation to random discrete distributions: an analytic approach](#). *Advances in Applied Probability*, 31(2):448–491, 1999.
- [56] Hsien-Kuei Hwang. [Phase changes in random recursive structures and algorithms](#). In *Proceedings of the Workshop on Probability with Applications to Finance and Insurance*, pages 82–97. World Scientific, 2004.
- [57] Hsien-Kuei Hwang and Vytas Zacharovas. [Uniform asymptotics of Poisson approximation to the Poisson-binomial distribution](#). *Theory of Probability and Its Applications*, 55(2):198–224, 2011.
- [58] Ildar Abdulovich Ibragimov and Kalman Eremeevich Chernin. [On the unimodality of stable laws](#). *Theory Probab. Appl.*, 4:417–419, 1959.
- [59] Lancelot Fitzgerald James. [Lamperti-type laws](#). *Ann. Appl. Probab.*, 20(4):1303–1340, 2010.
- [60] Lancelot Fitzgerald James. [Stick-breaking \$PG\(\alpha, \zeta\)\$ -generalized gamma processes](#). *arXiv*, 2013.
- [61] Lancelot Fitzgerald James. [Generalized Mittag Leffler distributions arising as limits in preferential attachment models](#). *arXiv*, 2015.
- [62] Svante Janson. [Functional limit theorems for multitype branching processes and generalized Pólya urns](#). *Stochastic Process. Appl.*, 110(2):177–245, 2004.
- [63] Svante Janson. [Asymptotic degree distribution in random recursive trees](#). *Random Struct. Algorithms*, 26(1-2):69–83, 2005.
- [64] Svante Janson. [Limit theorems for triangular urn schemes](#). *Probab. Theory Related Fields*, 134(3):417–452, 2006.

- [65] Svante Janson. [Random cutting and records in deterministic and random trees](#). *Random Struct. Algorithms*, 29:139–179, 2006.
- [66] Svante Janson. [Moments of Gamma type and the Brownian supremum process area](#). *Probab. Surv.*, 7:1–52, 2010. With addendum on pages 207–208.
- [67] Svante Janson, Donald E. Knuth, Tomasz Łuczak, and Boris Pittel. [The birth of the giant component](#). *Random Struct. Algorithms*, 4(3):233–358, 1993.
- [68] Svante Janson, Markus Kuba, and Alois Panholzer. [Generalized Stirling permutations, families of increasing trees and urn models](#). *J. Combin. Theory Ser. A*, 118(1):94–114, 2011.
- [69] Svante Janson and Nicolas Pouyanne. [Moment convergence of balanced Pólya processes](#). *Electron. J. Probab.*, 23:13, 2018. Id/No 34.
- [70] Dimitris Karlis. [EM algorithm for mixed Poisson and other discrete distributions](#). *Astin Bull.*, 35(1):3–24, 2005.
- [71] Donald E. Knuth. *The Art of Computer Programming. Vol. 3: Sorting and searching*. Addison-Wesley, 1998. Second edition (1st ed: 1973).
- [72] Valentin F. Kolchin. *Random Graphs*, volume 53. Cambridge University Press, 1999.
- [73] Markus Kuba. A note on central limit theorem analogues for node degree preferential attachment trees and Pitman’s α -diversity. *In preparation*, 2021.
- [74] Markus Kuba and Alois Panholzer. [Analysis of label-based parameters in increasing trees](#). In *Fourth Colloquium on Mathematics and Computer Science Algorithms, Trees, Combinatorics and Probabilities*, pages 321–330. Discrete Math. Theor. Comput. Sci., 2006.
- [75] Markus Kuba and Alois Panholzer. [Descendants in increasing trees](#). *Electronic Journal of Combinatorics*, 13 (1)(Paper 8), 2006.
- [76] Markus Kuba and Alois Panholzer. [On the degree distribution of the nodes in increasing trees](#). *J. Combin. Theory Ser. A*, 114(4):597–618, 2007.
- [77] Markus Kuba and Alois Panholzer. [A combinatorial approach to the analysis of bucket recursive trees](#). *Theoret. Comput. Sci.*, 411(34-36):3255–3273, 2010.
- [78] Markus Kuba and Alois Panholzer. [Analysis of statistics for generalized Stirling permutations](#). *Discrete Math.*, 20:875–910, 2011.
- [79] Markus Kuba and Alois Panholzer. [Bilabelled increasing trees and hook-length formulae](#). *European J. Combin.*, 33(2):248–258, 2012.
- [80] Markus Kuba and Alois Panholzer. [Limiting distributions for a class of diminishing urn models](#). *Advances in Applied Probability*, 44(4):1–31, 2012.
- [81] Markus Kuba and Alois Panholzer. [On death processes and urn models](#). In *Proceeding of the 23rd International Conference on Probabilistic, Combinatorial, and Asymptotic Methods in the Analysis of Algorithms (AofA’12)*, pages 29–42. Discrete Math. Theor. Comput. Sci., 2012.
- [82] Markus Kuba and Alois Panholzer. [Combinatorial families of multilabelled increasing trees and hook-length formulas](#). *Discrete Math.*, 339(1):227–254, 2016.
- [83] Markus Kuba and Alois Panholzer. [On moment sequences and mixed Poisson distributions](#). *Probab. Surv.*, 13:89–155, 2016.
- [84] Markus Kuba and Alois Panholzer. [On bucket increasing trees, clustered increasing trees and increasing diamonds](#). *To appear in Combinatorics, Probability and Computing*, 2021.
- [85] Markus Kuba and Henning Sulzbach. [On martingale tail sums in affine two-color urn models with multiple drawings](#). *J. Appl. Probab.*, 54:1–21, 2017.
- [86] Hosam M. Mahmoud. *Pólya Urn Models*. CRC Press, 2009.
- [87] Hosam M. Mahmoud, Robert Thomas Smythe, and Jerzy Szymański. [On the structure of random plane-oriented recursive trees and their branches](#). *Random Struct. Algorithms*, 4(2):151–176, 1993.

- [88] Philippe Marchal. [Constructing a sequence of random walks strongly converging to Brownian motion](#). In *Proceedings of the conference Discrete random walks (DRW'03), Paris, France, September 1–5, 2003*, pages 181–190. Discrete Mathematics & Theoretical Computer Science, 2003.
- [89] Philippe Marchal. [A note on the fragmentation of a stable tree](#). In *Proceedings of the fifth colloquium on mathematics and computer science. Blaubeuren, Germany, September 22–26, 2008*, pages 489–500. Discrete Mathematics & Theoretical Computer Science, 2008.
- [90] Jean-François Marckert. [Growing random uniform \$d\$ -ary trees](#). *arXiv*, 2021.
- [91] Jean-Claude Massé and Radu Theodorescu. [Neyman type A distribution revisited](#). *Statistica Neerlandica*, 59:206–213, 2005.
- [92] Amram Meir and John W. Moon. [Cutting down random trees](#). *Journal of the Australian Mathematical Society*, 11:313–324, 1970.
- [93] Grégory Miermont. [Random maps and their scaling limits](#). In *Fractal geometry and stochasticity IV. Proceedings of the 4th conference, Greifswald, Germany, September 8–12, 2008*, pages 197–224. Basel: Birkhäuser, 2009.
- [94] Gösta Mittag-Leffler. [Une généralisation de l'intégrale de Laplace–Abel](#). *C. R. Acad. Sci., Paris*, 136:537–539, 1903.
- [95] Gösta Mittag-Leffler. [Sur la nouvelle fonction \$E_\alpha\(x\)\$](#) . *C. R. Acad. Sci., Paris*, 137:554–558, 1903.
- [96] Tamás F. Móri. [The maximum degree of the Barabási–Albert random tree](#). *Combin. Probab. Comput.*, 14(3):339–348, 2005.
- [97] Jerzy Neyman. [On a new class of “contagious” distributions applicable in entomology and bacteriology](#). *Statistica Neerlandica*, 10:35–57, 1939.
- [98] Konstantinos Panagiotou, Leon Ramzews, and Benedikt Stuffer. [Exact-size sampling of combinatorial structures in linear time](#). *Preprint*, 2021.
- [99] Alois Panholzer. [Non-crossing trees revisited: cutting down and spanning subtrees](#). In *Discrete random walks (Paris, 2003)*, pages 265–276. Discrete Math. Theor. Comput. Sci., 2003.
- [100] Alois Panholzer. [Cutting down very simple trees](#). *Quaest. Math.*, 29(2):211–227, 2006.
- [101] Alois Panholzer and Helmut Prodinger. [Level of nodes in increasing trees revisited](#). *Random Struct. Algorithms*, 31(2):203–226, 2007.
- [102] Alois Panholzer and Georg Seitz. [Limiting distributions for the number of inversions in labelled tree families](#). *Ann. Comb.*, 16(4):847–870, 2012.
- [103] Alois Panholzer and Georg Seitz. [Ancestors and descendants in evolving \$k\$ -tree models](#). *Random Struct. Algorithms*, 44:465–489, 2014.
- [104] Erol A. Peköz, Adrian Röllin, and Nathan Ross. [Degree asymptotics with rates for preferential attachment random graphs](#). *Ann. Appl. Probab.*, 23(3):1188–1218, 2013.
- [105] Erol A. Peköz, Adrian Röllin, and Nathan Ross. [Generalized gamma approximation with rates for urns, walks and trees](#). *Ann. Probab.*, 44(3):1776–1816, 2016.
- [106] Robin Pemantle and Mark C. Wilson. *Analytic Combinatorics in Several Variables*. Cambridge University Press, 2013.
- [107] Raman N. Pillai. [On Mittag-Leffler functions and related distributions](#). *Ann. Inst. Statist. Math.*, 42(1):157–161, 1990.
- [108] Jim Pitman. [Exchangeable and partially exchangeable random partitions](#). *Probab. Theory Relat. Fields*, 102(2):145–158, 1995.
- [109] Jim Pitman. *Combinatorial Stochastic Processes*, volume 1875 of *Lecture Notes in Mathematics*. Springer, 2006. Lectures from the 32nd Summer School on Probability Theory held in Saint-Flour, July 2002.
- [110] Jim Pitman and Marc Yor. [The two-parameter Poisson–Dirichlet distribution derived from a stable subordinator](#). *Ann. Probab.*, 25(2):855–900, 1997.

- [111] Harry Pollard. [The representation of \$e^{-x^\lambda}\$ as a Laplace integral](#). *Bull. Amer. Math. Soc.*, 52:908–910, 1946.
- [112] Vincent Puyhaubert. [Modèles d’urnes et phénomènes de seuils en combinatoire analytique](#). PhD thesis, École polytechnique, Palaiseau, 2005.
- [113] Uwe Rösler and Ludger Rüschemdorf. [The contraction method for recursive algorithms](#). *Algorithmica*, 29(1-2):3–33, 2001.
- [114] John G. Skellam and Leonard R. Shenton. [Distributions associated with random walk and recurrent events](#). *J. Roy. Statist. Soc. Ser. B.*, 19:64–111 (discussion 111–118), 1957.
- [115] Andrea Sportiello. [Boltzmann sampling of irreducible context-free structures in linear time](#). *arXiv*, 2021.
- [116] Chun Su, Qunqiang Feng, and Zhishui Hu. [Uniform recursive trees: branching structure and simple random downward walk](#). *J. Math. Anal. Appl.*, 315(1):225–243, 2006.
- [117] Vladimir Vasilevich Uchaikin and Vladimir Mikhailovich Zolotarev. [Chance and stability. Stable distributions and their applications](#). De Gruyter, 1999.
- [118] Michael Wallner. [A half-normal distribution scheme for generating functions](#). *European J. Combin.*, 87:103138, 21, 2020.
- [119] Edmund Taylor Whittaker and George Neville Watson. [A Course of Modern Analysis](#). Fifth edition (1st ed.: 1902). Cambridge University Press, 2021.
- [120] Gordon E. Willmot. [Mixed compound Poisson distributions](#). *ASTIN Bulletin*, 16(S1):S59–S79, 1986.
- [121] Gordon E. Willmot and X. Sheldon Lin. [Lundberg Approximations for Compound Distributions with Insurance Applications](#). Springer, 2001.

Authors’ URLs:

Cyril Banderier <https://lipn.fr/~banderier>  0000-0003-0755-3022
Laboratoire d’Informatique de Paris Nord, Univ. Sorbonne Paris Nord, France.

Markus Kuba <https://dmg.tuwien.ac.at/kuba>  0000-0001-7188-6601
Department of Applied Mathematics and Physics, FH Technikum Wien, Austria.

Michael Wallner <https://dmg.tuwien.ac.at/mwallner>  0000-0001-8581-449X
Institute of Discrete Mathematics and Geometry, TU Wien, Austria.