

# DIAMETER TWO PROPERTIES IN SOME VECTOR-VALUED FUNCTION SPACES

HAN JU LEE AND HYUNG-JOON TAG

**ABSTRACT.** We introduce a vector-valued version of a uniform algebra, called the vector-valued function space over a uniform algebra. The diameter two properties of the vector-valued function space over a uniform algebra on an infinite compact Hausdorff space are investigated. Every nonempty relatively weakly open subset of the unit ball of a vector-valued function space  $A(K, (X, \tau))$  over an infinite dimensional uniform algebra has the diameter two, where  $\tau$  is a locally convex Hausdorff topology on a Banach space  $X$  compatible to a dual pair. Under the assumption on  $X$  being uniformly convex with norm topology  $\tau$  and the additional condition that  $A \otimes X \subset A(K, X)$ , it is shown that Daugavet points and  $\Delta$ -points on  $A(K, X)$  over a uniform algebra  $A$  are the same, and they are characterized by the norm-attainment at a limit point of the Shilov boundary of  $A$ . In addition, a sufficient condition for the convex diametral local diameter two property of  $A(K, X)$  is also provided. As a result, the similar results also hold for an infinite dimensional uniform algebra.

## 1. INTRODUCTION

In this article, we study the diameter two properties of a uniform algebra and the vector-valued function space  $A(K, (X, \tau))$  for a complex Banach space  $X$  with a locally convex Hausdorff topology  $\tau$  compatible with a dual pair by using the Urysohn-type lemma. The diameter two properties have gained a lot of attention recently and been studied in various Banach spaces. Exploring these properties involves weakly open subsets and slices of the unit ball  $B_X$  of a Banach space  $X$ . Let  $S_X$  be the unit sphere of a Banach space  $X$  and  $X^*$  be the dual space of  $X$ . For  $x^* \in S_{X^*}$  and  $\epsilon > 0$ , a slice  $S(x^*, \epsilon)$  of the unit ball  $B_X$  is defined by  $S(x^*, \epsilon) = \{x \in B_X : \operatorname{Re} x^*x > 1 - \epsilon\}$ . Here we recall the definitions of the “classical” diameter two properties introduced in [3].

**Definition 1.1.** (“Classical” D2Ps)

- (i) A Banach space  $X$  has the strong diameter two property (SD2P) when every convex combination of slices has diameter two.
- (ii) A Banach space  $X$  has the diameter two property (D2P) when every nonempty relatively weakly open subset of the unit ball  $B_X$  has diameter two.
- (iii) A Banach space  $X$  has the local diameter two property (LD2P) when every slice of the unit ball  $B_X$  has diameter two.

Since every weakly open subset of the unit ball contains a convex combination of slices [12, Lemma II.1], the SD2P implies the D2P. The D2P implies the LD2P because every slice is a relatively weakly open subset of the unit ball. It is well-known that these properties are weaker than the Daugavet property [3, Theorem 4.4] and that these properties are the other extremes of the Radon-Nikodým property which states, geometrically, that there exists a nonempty slice of the unit ball with arbitrarily small diameter. The classical D2Ps are examined in various spaces. It is well-known that  $c_0$  and  $\ell_\infty$

---

*Date:* 2021. February 20.

*2010 Mathematics Subject Classification.* Primary 46B04; Secondary 46B20, 46E30, 47B38.

*Key words and phrases.* diameter two property, uniform algebra, Urysohn-type lemma, Shilov boundary.

The first author was supported by Basic Science Research Program through the National Research Foundation of Korea(NRF) funded by the Ministry of Education, Science and Technology [NRF-2020R1A2C1A01010377]. The second author was supported by Basic Science Research Program through the National Research Foundation of Korea(NRF) funded by the Ministry of Education, Science and Technology [NRF-2020R1A2C1A01010377].

has the SD2P. The infinite dimensional uniform algebra over a compact Hausdorff space also has the SD2P [3, 20], in fact, even stronger property called the symmetric strong diameter two property (SSD2P) [4]. Other known examples with the D2P are interpolation spaces  $L_1 + L_\infty$  and  $L_1 \cap L_\infty$  equipped with certain norms [2]. These spaces do not satisfy the Daugavet property but the D2P. For a Banach space  $X$ , the space  $C(K, (X, \tau))$  of  $X$ -valued continuous function over a compact Hausdorff space  $K$ , where  $\tau$  a locally convex Hausdorff topology compatible to a dual pair, is also known to have the D2P [5].

Not only the ‘‘classical’’ D2Ps mentioned above, but we also explore the ‘‘diametral’’ version of the D2Ps. Studying these properties requires us to have a firm grasp on the  $\Delta$ -points and the Daugavet points. Let  $\Delta_\epsilon(x) = \{y \in B_X : \|x - y\| \geq 2 - \epsilon\}$ . Then a point  $x \in S_X$  is said to be a  $\Delta$ -point if  $x \in \overline{\text{conv}}\Delta_\epsilon(x)$  for every  $\epsilon > 0$ . We denote the set of all  $\Delta$ -points by  $\Delta_X$ . Similarly, a point  $x \in S_X$  is said to be a Daugavet point if  $B_X = \overline{\text{conv}}\Delta_\epsilon(x)$  for every  $\epsilon > 0$ . Notice that every Daugavet point is a  $\Delta$ -point. We recall the definitions of the diametral diameter two properties introduced in [1, 6].

**Definition 1.2.** (*‘‘Diametral’’ D2Ps*)

- (i) A Banach space  $X$  has the convex diametral local diameter two property (convex-DLD2P) if  $B_X = \overline{\text{conv}}\Delta_X$ , that is, the unit ball is the closed convex hull of the set of all  $\Delta$ -points.
- (ii) A Banach space  $X$  has the diametral local diameter two property (DLD2P) if for every slice  $S$  of  $B_X$ , every  $x \in S_X \cap S$ , and every  $\epsilon > 0$  there exists  $y \in S$  such that  $\|x - y\| \geq 2 - \epsilon$ .
- (iii) A Banach space  $X$  has the diametral diameter two property (DD2P) if for every relative weakly open subset  $U$  of  $B_X$ , every  $x \in S_X \cap U$ , and every  $\epsilon > 0$ , there exists  $y \in U$  such that  $\|x - y\| \geq 2 - \epsilon$ .

There is also a stronger version of the DD2P called the diametral strong diameter two property (DSD2P), which uses a convex combination of slices, but this property is now known to be equivalent to the Daugavet property [14]. It was shown in [6] that the Daugavet property implies the DD2P and that the DD2P implies the DLD2P. The convex-DLD2P is stronger than the LD2P and weaker than DLD2P [1]. Moreover, we can see that the D2P is weaker than the DD2P from their definitions. If one wishes to see a clearer picture, we refer to a nice visualization of the relationship between various diameter two properties in the dissertation of Pirk (see [23, pg 82-85]). These properties have been also examined on several Banach spaces. For example, the spaces  $c$  of convergent sequences and  $l_\infty$  do not have the DLD2P but the convex-DLD2P [1, Corollary 5.4, Remark 5.5]. Moreover,  $c_0$  even fails to have the convex-DLD2P [1, pg 18].

The DLD2P and the Daugavet property have equivalent definitions in terms of  $\Delta$ -points and Daugavet points. A Banach space  $X$  having the DLD2P is equivalent to say that every point in  $S_X$  is a  $\Delta$ -point (see [13, Theorem 1.4] and [24, Open problem (7)]). Similarly, a Banach space  $X$  has the Daugavet property if and only if every point in  $S_X$  is a Daugavet point [24, Corollary 2.3].

The organization of this article consists of four parts. In section 2, we provide necessary information about the uniform algebra  $A$ , the space  $A(K, (X, \tau))$ , and the Urysohn-type lemma, which is the key ingredient to prove our results throughout this article. In section 3, we make a few remarks on the symmetric strong diameter two property of a scalar-valued function algebra. In section 4, we show that the vector-valued function space  $A(K, (X, \tau))$  over a uniform algebra also satisfies the D2P. By the Gelfand transform we show that a function space  $A(\Omega, X)$  on a Hausdorff space  $\Omega$  is isometrically isomorphic to a function space  $A(M_A, (X^{**}, w^*))$  over a uniform algebra, so it is shown that  $A(\Omega, X)$  also has the D2P if the base (function) algebra is infinite dimensional. This result shows D2P of some function spaces like  $A_b(B_X, Y)$  and (resp.  $A_u(B_X, Y)$ ) consisting of  $Y$ -valued functions which are bounded (resp. uniformly continuous) on  $B_X$  and are holomorphic on the interior of  $B_X$  when  $X$  and  $Y$  are Banach spaces. In section 5, we present some results on the Daugavet and  $\Delta$ -points in a vector-valued function space  $A(K, X)$  over a uniform algebra. Under the condition of  $X$  being uniformly convex and the additional assumption that  $A \otimes X \subset A(K, X)$ , we show that these points are the same and characterized by norm-attainment at a limit point of the Shilov boundary. Also under the same assumptions, a sufficient condition for the convex-DLD2P of  $A(K, X)$  is provided.

## 2. PRELIMINARIES

In this article, we assume that  $K$  is a compact Hausdorff space and  $X, Y$  are nontrivial complex Banach spaces unless specified. Let  $X^*$  be the dual space of  $X$ . We work with a Banach space  $X$  equipped with a locally convex Hausdorff (LCH) topology  $\tau$  compatible to the dual pair  $(X, X^*)$  or  $(X^*, X)$ . Recall that  $(X, \tau)$  is said to be compatible to a dual pair  $(X, X^*)$  if  $(X, \tau)^* = X^*$  and a LCH topology  $\tau$  on  $X^*$  is said to be compatible to a dual pair  $(X^*, X)$  if  $(X^*, \tau)^* = X$ . In this paper, we say that a LCH topology  $\tau$  on a Banach space  $X$  is compatible to a dual pair if  $(X, \tau)^* = X^*$  or  $(X, \tau)^* = X_*$ , where  $X_*$  is the predual of  $X$ .

The weak topology on  $X$  is denoted by  $(X, w)$ , the weak\* topology on  $X^*$  by  $(X^*, w^*)$ , and the norm topology on  $X$  by  $(X, \|\cdot\|)$ . The Mackey topology for  $X$  (resp.  $X^*$ ) is the finest topology where every linear functional  $x^* \in X^*$  (resp. linear functional defined by  $x \in X$ ) is continuous. The Mackey-Arens theorem [19, Theorem 8.7.4] says that a locally convex topology  $\tau$  for  $X$  (resp.  $X^*$ ) compatible to a dual pair includes the weak (resp. weak\*-) topology and is included in the Mackey topology on  $X$  (resp.  $X^*$ ) and that is known to be coarser than the norm topology on  $X$  [22, IV.3.4] (resp. the norm topology on  $X^*$  [19, Example 8.5.5 and Example 8.8.9]). Hence for any LCH topology  $\tau$  compatible to a dual pair  $(X, X^*)$  (resp.  $(X^*, X)$ ), we have  $(X, w) \subset \tau \subset (X, \|\cdot\|)$  (resp.  $(X^*, w^*) \subset \tau \subset (X^*, \|\cdot\|)$ ). Also for such topology  $\tau$  on  $X$  compatible to  $(X, X^*)$ ,  $\tau$ -bounded sets are weakly bounded and so norm-bounded. Similarly, for a LCH topology  $\tau$  on  $X^*$  compatible to  $(X^*, X)$ ,  $\tau$ -bounded sets are weakly\*-bounded and so norm-bounded in  $X^*$ . For more details on the theory of topological vector spaces, we refer to [19, 22].

Now we show that the space  $C(K, (X, \tau))$ , which consists of all continuous functions from a compact Hausdorff space  $K$  to  $(X, \tau)$ , equipped with the supremum norm is a Banach space. Even though this fact is already known in [5], we include the proof for completeness.

**Proposition 2.1.** *Let  $K$  be a compact Hausdorff space, let  $X$  be either a Banach space and let  $\tau$  be a locally convex Hausdorff topology compatible to a dual pair. Then the space  $C(K, (X, \tau))$  equipped with the supremum norm is a Banach space.*

*Proof.* Let us consider when  $\tau$  is a LCH topology compatible to either a dual pair  $(X, X^*)$  or  $(X, X_*)$ . Since  $f(K)$  is  $\tau$ -compact for every  $f \in C(K, (X, \tau))$ , the range  $f(K)$  is bounded in the norm topology of  $X$ . This makes the norm  $\|f\| = \sup\{\|f(t)\|_X : t \in K\}$  to be well-defined.

To show the completeness of  $C(K, (X, \tau))$ , let  $(f_n)_{n=1}^\infty \subset C(K, (X, \tau))$  be a Cauchy sequence. Then for any  $\epsilon > 0$ , there exists  $N$  such that for every  $n, m \geq N$ , we have  $\|f_n - f_m\| < \epsilon$ . Notice that for every  $t \in K$ ,

$$(1) \quad \|f_n(t) - f_m(t)\|_X \leq \|f_n - f_m\| < \epsilon,$$

and so  $(f_n(t))_{n=1}^\infty$  is a Cauchy sequence in  $X$  with respect to the norm topology. Since  $X$  is complete with the norm topology,  $(f_n(t))_{n=1}^\infty$  converges in norm. So for each  $t \in K$ , let  $f(t) = \lim_{n \rightarrow \infty} f_n(t)$ . Then by (1), for any  $\epsilon > 0$  there exists  $N$  such that for all  $n \geq N$ ,  $\sup_{t \in K} \|f_n(t) - f(t)\|_X < \frac{\epsilon}{2}$ .

Now we have to show that  $f \in C(K, (X, \tau))$ . First, let  $V$  be a  $\tau$ -neighborhood of 0. Choose a balanced, convex  $\tau$ -neighborhood  $W$  of 0 such that  $W + W + W \subset V$ . Since  $W$  is also open in the norm topology, there exists  $\epsilon B_X \subset W$  and  $N \in \mathbb{N}$  such that for all  $n \geq N$  and  $t \in K$ ,  $f(t) - f_n(t) \in \epsilon B_X$ . Hence

$$\begin{aligned} f(t) - f(t_0) &= (f(t) - f_N(t)) + (f_N(t) - f_N(t_0)) + (f_N(t_0) - f(t_0)) \\ &\in \epsilon B_X + (f_N(t) - f_N(t_0)) + \epsilon B_X \\ &\subset W + (f_N(t) - f_N(t_0)) + W. \end{aligned}$$

Moreover,  $f_N$  is continuous with respect to  $\tau$ . So there exists a neighborhood  $U$  of  $t_0$  such that for every  $t \in U$ ,  $f_N(t) - f_N(t_0) \in W$ . Therefore,  $f(t) - f(t_0) \in W + W + W \subset V$  for all  $t \in U$ . This shows that  $f \in C(K, (X, \tau))$ . Therefore,  $C(K, (X, \tau))$  is a Banach space.  $\square$

Let  $\Omega$  be a Hausdorff space and  $C_b(\Omega)$  be the Banach algebra of all bounded continuous functions over  $\Omega$  equipped with the supremum norm. A function algebra  $A(\Omega)$  on  $\Omega$  is a closed subalgebra of  $C_b(\Omega)$  that separates the points of  $\Omega$  and contains constant functions. Here separating points means for each pair  $(s, t) \in \Omega \times \Omega$  with  $s \neq t$ , there exists  $f \in A(\Omega)$  such that  $f(s) \neq f(t)$ . If  $\Omega$  is a compact Hausdorff space, a function algebra is called a uniform algebra.

We introduce  $(X, \tau)$ -valued function space over a uniform algebra for a Banach space  $X$  as follows.

**Definition 2.2.** *Let  $X$  be a Banach space and  $\tau$  be a LCH topology on  $X$  compatible to a dual pair. A closed subspace  $A(K, (X, \tau))$  of  $C(K, (X, \tau))$  is said to be the  $(X, \tau)$ -valued function space over a uniform algebra  $A$  if the following two conditions are satisfied:*

- (i) *The base algebra, defined by  $A := \{y^* \circ f : f \in A(K, (X, \tau)), y^* \in (X, \tau)^*\}$ , is a uniform algebra over  $K$ .*
- (ii)  *$\phi f \in A(K, (X, \tau))$  for every  $\phi \in A$  and  $f \in A(K, (X, \tau))$  where  $(\phi f)(t) = \phi(t)f(t)$  for  $t \in K$ .*

Notice that if  $X = \mathbb{C}$ , we have the usual uniform algebra  $A$  on  $K$ . When  $\tau$  is the norm topology, we call the space  $A(K, (X, \tau))$  an  $X$ -valued function space over a uniform algebra and denote the space by  $A(K, X)$ . Similar vector-valued version of a uniform algebra has been studied in [7].

A point  $t_0 \in K$  is said to be a strong boundary point for a uniform algebra  $A$  if for every neighborhood  $U$  containing  $t_0$ , there exists  $f \in A$  such that  $f(t_0) = \|f\| = 1$  and  $\sup_{t \in K \setminus U} |f(t)| < 1$ . A subset  $S \subset K$  is said to be a boundary if for each  $f \in A$ , there exists an element  $t \in S$  such that  $|f(t)| = \|f\|_\infty$ . The Shilov boundary  $\Gamma(A)$  of  $A$  is the smallest closed boundary of  $A$  and it is known that it is the intersection of all closed boundaries of  $A$ . For a uniform algebra, let  $K_A = \{\lambda \in A^* : \|\lambda\| = \lambda(1_A) = 1\}$  and denote the set of its extreme points by  $\text{ex } K_A$ . The set  $\text{ex } K_A$  is called the Choquet boundary of  $A$ . It is well-known that each  $\lambda \in \text{ex } K_A$  are associated with some elements  $x \in K$  and uniquely represented by the Dirac measure  $\delta_x$  [9, Lemma 4.3.2]. For this reason, we use  $\Gamma_0(A)$  to denote the set of such elements  $x \in K$  corresponding to the elements in the Choquet boundary. Also, the set of strong boundary points and  $\Gamma_0(A)$  coincide with each other when we consider a uniform algebra  $A$  over a compact Hausdorff space  $K$  [9, Theorem 4.3.5]. It is known that the Shilov boundary  $\Gamma(A)$  of a uniform algebra  $A$  is the closure of its Choquet boundary  $\Gamma_0(A)$  [9, Corollary 4.3.7.a].

Now we recall the Urysohn-type lemma that is extensively used throughout this article.

**Lemma 2.3.** [8, Lemma 2.5] *Let  $A \subset C(K)$  be a uniform algebra and  $\Gamma_0$  be its Choquet boundary. Then, for every open set  $U \subset K$  with  $U \cap \Gamma_0 \neq \emptyset$  and  $0 < \epsilon < 1$ , there exists  $f \in A$  and  $t_0 \in U \cap \Gamma_0$  such that  $f(t_0) = \|f\|_\infty = 1$ ,  $|f(t)| < \epsilon$  for every  $t \in K \setminus U$  and*

$$(2) \quad |f(t)| + (1 - \epsilon)|1 - f(t)| \leq 1 \text{ for all } t \in K.$$

The following observation is useful for later.

**Lemma 2.4.** *Let  $X$  be a Banach space and let  $\tau$  be a LCH topology compatible to a dual pair. Suppose that  $A(K, (X, \tau))$  is a  $(X, \tau)$ -valued function space over the base algebra  $A$  and  $L$  is a closed boundary for  $A$ . The space of restrictions of elements of  $A(K, (X, \tau))$  to  $L$  is denoted by  $A(L, (X, \tau))$  and the restrictions of elements of  $A$  to  $L$  is denoted by  $A(L)$ . Then  $A(L, (X, \tau))$  is a  $(X, \tau)$ -valued function space over the base algebra  $A(L)$  and it is isometrically isomorphic to  $A(K, (X, \tau))$ .*

*Proof.* Given  $f \in A(K, (X, \tau))$ , let  $f|_L$  be the restriction of  $f$  to  $L$ . Then we have

$$\begin{aligned} \|f\| &= \sup\{|f(t)| : t \in K\} = \sup\{|x^* f(t)| : t \in K, x^* \in B_{(X, \tau)^*}\} \\ &= \sup\{|x^* f(t)| : t \in L, x^* \in B_{(X, \tau)^*}\} = \sup\{\|f(t)\| : t \in L\} = \|f|_L\|. \end{aligned}$$

So  $A(L, (X, \tau))$  is a surjective isometric image of  $A(K, (X, \tau))$  by the map  $f \mapsto f|_L$  and it is a closed subspace of  $C(L, (X, \tau))$ . It is easy to see that  $A(L, (X, \tau))$  satisfies the condition (i) and (ii) in the Definition 2.2 and its base algebra is  $A(L)$ . This completes the proof.  $\square$

Since the Shilov boundary  $\Gamma$  of a uniform algebra is a closed boundary of  $K$ , by Lemma 2.4 or [17, Theorem 4.1.6]), a uniform algebra is isometric to a uniform algebra of its restrictions to  $\Gamma$ . We need another useful lemma on the isolated point of the Shilov boundary.

**Lemma 2.5.** *Let  $A$  be a uniform algebra on a compact Hausdorff space  $K$  and let  $t_0$  be an isolated point of the Shilov boundary  $\Gamma$  of  $A$ . Then there exists a function  $\phi \in A$  such that  $\phi(t_0) = \|\phi\| = 1$  and  $\phi(t) = 0$  for  $t \in \Gamma \setminus \{t_0\}$ .*

*Proof.* Since  $t_0$  is an isolated point of  $\Gamma$ , there is an open set such that  $U \cap \Gamma = \{t_0\}$ . The set of strong boundary points  $\Gamma_0$  is dense in  $\Gamma$ . So the point  $t_0 \in \Gamma$  is in fact a strong boundary point of  $A$ . By definition of a strong boundary point, there exists a function  $\psi \in A$  such that  $\psi(t_0) = \|\psi\| = 1$  and  $\sup_{t \in K \setminus U} |\psi(t)| < 1$ . Note that  $\{(\psi|_{\Gamma})^n\}_{n=1}^{\infty}$  converges uniformly in  $C(L)$ . By the isometry of the map from  $f \in A$  to its restriction  $f|_L$  to  $L$ ,  $\{\psi^n\}_{n=1}^{\infty}$  is Cauchy in  $A$  and converges uniformly to  $\phi$  in  $A$ . Now it is clear that  $\phi(t) = 0$  for  $t \in \Gamma \setminus \{t_0\}$  and  $\phi(t_0) = \|\phi\| = 1$ . Thus, we obtain the desired function.  $\square$

### 3. A SHORT REMARK ON THE SSD2P OF FUNCTION ALGEBRAS

First we start with what is known as the symmetric strong diameter two property (SSD2P) that is stronger than the SD2P.

**Definition 3.1.** [4, Definition 1.3] *A Banach space  $X$  has the symmetric strong diameter two property (SSD2P) if whenever  $n \in \mathbb{N}$ ,  $S_1, \dots, S_n$  are slices of  $B_X$ , and  $\epsilon > 0$ , there exists  $x_i \in S_i$ ,  $i = 1, 2, 3, \dots, n$ , and  $y \in B_X$  such that  $x_i \pm y \in S_i$  for every  $i \in \{1, 2, 3, \dots, n\}$  and  $\|y\| > 1 - \epsilon$ .*

It has been already known from [3, Theorem 4.2] that the infinite dimensional uniform algebra over a compact Hausdorff space has the SSD2P and their proof method is extended to show the SSD2P of somewhat regular subspaces of the space  $C_0(L)$  of continuous functions over a locally compact Hausdorff space  $L$  vanishing at infinity [4, Theorem 2.2].

**Definition 3.2.** [4, Definition 2.1] *Let  $L$  be a locally compact Hausdorff space and  $C_0(L)$  be the space of continuous functions over  $L$  vanishing at infinity. A linear subspace  $Y$  of  $C_0(L)$  is somewhat regular, if whenever  $V$  is a non-empty open subset of  $L$  and  $0 < \epsilon < 1$ , there is an  $f \in Y$  such that*

$$\|f\| = 1 \text{ and } |f(x)| \leq \epsilon \text{ for every } x \in L \setminus V.$$

We make a few remarks here. In view of Lemma 2.3 we can easily verify that the infinite dimensional uniform algebra  $A$  over a compact Hausdorff space  $K$  is an example of somewhat regular subspace of  $C(\Gamma)$  due to the fact that  $A$  is isometric to  $A(\Gamma)$  and that  $\Gamma_0 = \Gamma_0(A)$  is dense in  $\Gamma = \Gamma(A)$  because  $K$  is compact. In addition, Lemma 2.3 is a special case of the Urysohn-type lemma for somewhat regular subspaces (see [4, Lemma 2.4]) that targets on uniform algebras.

**Theorem 3.3.** [4, Theorem 2.2] *The infinite dimensional uniform algebra  $A$  over a compact Hausdorff space  $K$  has the SSD2P.*

Now let  $C_b(B_X)$  be the space of bounded, complex-valued, continuous functions on the unit ball  $B_X$  of a Banach space  $X$ . The space  $A_b(B_X)$  is a closed subalgebra of  $C_b(B_X)$  that consists of holomorphic functions on the interior of  $B_X$ . The space  $A_u(B_X)$  is a closed subalgebra of  $C_b(B_X)$  that consists of functions in  $A_b(B_X)$  that are uniformly continuous on  $B_X$ . From the fact that function algebras  $A_b(B_X)$  and  $A_u(B_X)$  are isometric to a uniform algebra on a compact Hausdorff space via the Gelfand transformation (see [16, Proposition 2] or Theorem 4.5), we can deduce the following fact on the SSD2Ps of  $C_b(B_X)$ ,  $A_b(B_X)$  and  $A_u(B_X)$ .

**Corollary 3.4.** *Let  $X$  be a Banach space. The function algebras  $C_b(B_X)$ ,  $A_b(B_X)$  and  $A_u(B_X)$  have the SSD2P.*

## 4. THE D2P OF VECTOR-VALUED FUNCTION SPACES OVER A UNIFORM ALGEBRA

Now we consider the space  $A(K, (X, \tau))$  where  $X$  is a Banach space with a locally convex Hausdorff topology  $\tau$  compatible with a dual pair. Here the norm of the space  $A = A(K)$  is denoted by  $\|\cdot\|_\infty$ . First we recall a useful fact about the D2P of a direct sum of two Banach spaces.

**Lemma 4.1.** [5, Lemma 2.2] *Let  $X$  be a Banach space satisfying the diameter two property. Then for any arbitrary Banach space  $Y$ ,  $X \oplus_\infty Y$  has the diameter two property.*

Here we present the main result of this section.

**Theorem 4.2.** *Let  $K$  be a compact Hausdorff space and  $X$  be a Banach space endowed with a locally convex Hausdorff topology  $\tau$  compatible to a dual pair. If the base algebra  $A$  is infinite dimensional, then the space  $A(K, (X, \tau))$  has the diameter two property, i.e. every nonempty weakly open subset of the unit ball has the diameter two.*

*Proof.* In view of Lemma 2.4, we only have to show the D2P for  $A(\Gamma, (X, \tau))$  where  $\Gamma = \Gamma(A)$  is a Shilov boundary of a uniform algebra. Since  $A$  is infinite dimensional,  $\Gamma$  is infinite. We write  $A(\Gamma, X)$  instead of  $A(\Gamma, (X, \tau))$  for convenience.

Now let  $W$  be a nonempty weakly open subset of  $A(\Gamma, X)$ . Three cases will be considered.

- (i) Assume that  $\Gamma$  is perfect. Let  $a \in W \cap S_{A(\Gamma, X)}$ . Then for every  $\delta > 0$ , there exists  $t_0 \in \Gamma$  such that  $\|a(t_0)\|_X > 1 - \delta$ . Also we can find  $f \in S_{(X, \tau)^*}$  such that  $\operatorname{Re} f(a(t_0)) > 1 - \delta$  because  $(X, \tau)^* = X^*$  or  $(X, \tau)^* = X_*$ , where  $X_*$  is a predual of  $X$ .

Define  $U = \{t \in \Gamma : \operatorname{Re} f(a(t)) > 1 - \delta\}$ . Notice that  $U$  contains  $t_0$ . Since  $t_0$  is a limit point of  $\Gamma$ , we can construct pairwise disjoint open subsets  $U_n \subset U$  for  $n \in \mathbb{N}$ . From the fact that  $\Gamma_0 = \Gamma_0(A)$  is dense in  $\Gamma$ , we know  $U_n \cap \Gamma_0 \neq \emptyset$  for all  $n \in \mathbb{N}$ . So by Lemma 2.3, for each  $n \in \mathbb{N}$  there exist  $\phi_n \in A$  and  $t_n \in U_n \cap \Gamma_0$  such that  $\phi_n(t_n) = \|\phi_n\|_\infty = 1$ ,  $|\phi_n(t)| < \frac{\delta}{2^{n+2}}$  for  $t \in \Gamma \setminus U_n$ , and

$$|\phi_n(t)| + \left(1 - \frac{\delta}{2^{n+2}}\right) |1 - \phi_n(t)| \leq 1 \text{ for } t \in \Gamma.$$

Define  $g_n = 1 - 2\phi_n$ . First we want to show that  $\|g_n\| \leq 1 + \frac{\delta}{2^n}$ . For every  $t \in \Gamma$ ,

$$\begin{aligned} |1 - 2\phi_n(t)| &= \left| 1 - 2\phi_n(t) + \frac{\delta}{2^{n+2}} - \frac{\delta}{2^{n+2}} + \frac{\delta}{2^{n+1}}\phi_n(t) - \frac{\delta}{2^{n+1}}\phi_n(t) \right| \\ &\leq \left| \left(1 - \frac{\delta}{2^{n+2}}\right) (1 - 2\phi_n(t)) \right| + \frac{\delta}{2^{n+2}} + \left| \frac{\delta}{2^{n+1}}\phi_n(t) \right| \\ &\leq \left| \left(1 - \frac{\delta}{2^n}\right) (1 - \phi_n(t)) - \left(1 - \frac{\delta}{2^n}\right) \phi_n(t) \right| + \frac{\delta}{2^n} \\ &\leq 1 - |\phi_n(t)| + |\phi_n(t)| + \frac{\delta}{2^n} \\ &\leq 1 + \frac{\delta}{2^n}. \end{aligned}$$

Hence  $\|g_n\|_\infty \leq 1 + \frac{\delta}{2^n}$ . Define  $h_n = \frac{g_n}{1 + \delta/2^n}$ . Notice that  $h_n \in B_A$  for all  $n \in \mathbb{N}$  and that  $h_n(t)$  converges to 1 pointwise.

Now, let  $F \in A(\Gamma, X)^*$  be a bounded linear functional on  $A(\Gamma, X)$ . If we define  $\varphi(h) = F(ha)$  for  $h \in A$ ,  $\varphi$  is a bounded linear functional on  $A$  because  $|\varphi(h)| \leq \|F\|_{A(\Gamma, X)^*} \|h\|_\infty \|a\| < \|h\|_\infty \|F\|_{A(\Gamma, X)^*} < \infty$  for all  $h \in A$ . By the Hahn-Banach extension theorem and the Riesz representation theorem,  $\varphi$  is represented by a regular complex Borel measure  $\mu$  and  $\varphi(h_n) = \int_\Gamma h_n d\mu$ . Moreover,  $\varphi(h_n) = \int_\Gamma h_n d\mu \rightarrow \int_\Gamma 1 d\mu = \varphi(1)$  by the bounded convergence theorem. Hence  $F(h_n a) = \varphi(h_n) \rightarrow \varphi(1) = F(a)$  and so the functions  $h_n a$  converges to  $a$  weakly in  $A(\Gamma, X)$ . Furthermore, we have

$$\begin{aligned}
\|a - h_n a\| &\geq \|a(t_n) - h_n(t_n)a(t_n)\|_X = |1 - h_n(t_n)| \|a(t_n)\|_X \\
&\geq \frac{2 + \delta/2^n}{1 + \delta/2^n} \cdot \operatorname{Re} f(a(t_n)) \\
&\geq \left(2 - \frac{\delta/2^n}{1 + \delta/2^n}\right) (1 - \delta) \\
&\geq (2 - \delta)(1 - \delta) \\
&\geq 2 - 3\delta.
\end{aligned}$$

Since  $\delta > 0$  is arbitrary, the diameter of  $W \cap B_{A(\Gamma, X)}$  is 2. The proof is finished.

- (ii) Now we consider when  $\Gamma$  has infinitely many isolated points. Let  $(t_n)_{n=1}^\infty$  be a sequence of isolated points and  $t_0$  be the accumulation point of the sequence. Let  $a \in W \cap S_{A(\Gamma, X)}$ . Let  $U$  be a  $\tau$ -open neighborhood around 0 and let  $V$  be an  $\tau$ -open neighborhood around  $a(t_0) \neq 0$  such that  $U \cap V = \emptyset$ . Notice that every  $\tau$ -open neighborhood is also an open neighborhood in the norm topology on  $X$ . So by the continuity of  $a$ ,  $a^{-1}(V)$  contains infinitely many  $t_n$ 's. Since  $U$  is norm-open, for each  $n \in \mathbb{N}$  there is  $\delta > 0$  such that  $\delta B_X \subset U$ . Hence  $a(t_n) \in V \subset X \setminus \delta B_X$ . That is,  $\|a(t_n)\|_X > \delta$  infinitely many  $n \in \mathbb{N}$ . So we may assume that  $\|a(t_n)\|_X > \delta$  for all  $n \in \mathbb{N}$ . Since  $\Gamma_0$  is dense in  $\Gamma$ , every isolated point  $t_n \in \Gamma$  is a strong boundary point. Hence by Lemma 2.5, there exists  $\phi_n \in A$  such that  $\phi_n(t_n) = \|\phi_n\| = 1$  and  $\phi_n(t) = 0$  for all  $t \in \Gamma \setminus \{t_n\}$ .

Define  $g_n = 1 + \left(\frac{1}{\|a(t_n)\|_X} - 1\right) \phi_n + \left(\frac{-1}{\|a(t_{n+1})\|_X} - 1\right) \phi_{n+1}$ . We see that  $g_n(t)$  converges to 1 pointwise,  $\sup_{n \in \mathbb{N}} \|g_n\|_\infty < 1 + \frac{1}{\delta}$  and  $\|g_n a\| = 1$  for every  $n \in \mathbb{N}$ . Hence  $g_n \rightarrow 1$  weakly in  $A$ . Moreover,  $g_n a \rightarrow a$  weakly in  $A(\Gamma, X)$  by the same argument in (i). When we compute the diameter of the relatively weakly open set of  $B_{A(\Gamma, X)}$ , we have

$$\begin{aligned}
\operatorname{diam}(W \cap B_{A(\Gamma, X)}) &\geq \|g_{n+1}a - g_n a\| \geq \|g_{n+1}(t_{n+1})a(t_{n+1}) - g_n(t_{n+1})a(t_{n+1})\|_X \\
&= \frac{2}{\|a(t_{n+1})\|_X} \|a(t_{n+1})\|_X = 2.
\end{aligned}$$

- (iii) Finally, let us consider when  $\Gamma$  has only finitely many isolated points  $t_1, t_2, \dots, t_n$ . Denote  $\Gamma_1 = \{t_1, t_2, \dots, t_n\}$  and the perfect subset of  $\Gamma$  by  $\Gamma_2 = \Gamma \setminus \Gamma_1$ . Since the set of strong boundary points are dense in  $\Gamma$ , we see that  $t_1, t_2, \dots, t_n$  are also strong boundary points and for each  $i = 1, 2, \dots, n$ , once again by Lemma 2.5, there exist  $\phi_i \in A(\Gamma)$  such that  $\phi_i(t_i) = 1$  and  $\phi_i(t) = 0$  for all  $t \neq t_i$ . Define a bounded linear operator  $P : A(\Gamma, X) \rightarrow A(\Gamma, X)$  by  $Pf = \sum_{i=1}^n \phi_i f$ . Then it is easy to check that  $P$  is a norm-one projection. Let  $B = P(A(\Gamma, X))$  and  $C$  be the restrictions of  $f \in A(\Gamma, X)$  to  $\Gamma_2$ . Then  $C$  is an  $(X, \tau)$ -valued function space over an infinite dimensional uniform algebra on  $\Gamma_2$ . We will show that  $A(\Gamma, X)$  is isometrically isomorphic to  $B \oplus_\infty C$ . Then by (i),  $C$  has diameter two property and Lemma 4.1 completes the proof.

For  $f \in A(\Gamma, X)$ , define  $\Phi(f) = (P(f), f|_{\Gamma_2})$ . Then  $\Phi : A(\Gamma, X) \rightarrow B \oplus_\infty C$  is well-defined and an isometry as follows

$$\|\Phi(f)\| = \max\{\|P(f)\|, \|f|_{\Gamma_2}\|\} = \max\left\{\sup_{t \in \Gamma_1} \|f(t)\|_X, \sup_{t \in \Gamma_2} \|f(t)\|_X\right\} = \sup_{t \in \Gamma} \|f(t)\|_X = \|f\|.$$

Given  $(f, g) \in B \oplus_\infty C$ , there exist  $f_1, f_2 \in A(\Gamma, X)$  such that  $f = P(f_1)$  and  $g = f_2|_{\Gamma_2}$ . Let  $h = P(f_1) + f_2 - P(f_2) \in A(\Gamma, X)$  and it is clear that  $\Phi(h) = (P(f_1), f_2|_{\Gamma_2}) = (f, g)$ . So  $\Phi$  is surjective. This shows that  $A(\Gamma, X)$  is isometrically isomorphic to  $B \oplus_\infty C$  and completes the proof.  $\square$

The norm, weak and weak-\* topologies are compatible to a dual pair and we get the following.

**Corollary 4.3.** *Let  $X$  be a Banach space and let  $K$  be an infinite compact Hausdorff space. Then the spaces  $A(K, X)$ ,  $A(K, (X, w))$ , and  $A(K, (X^*, w^*))$  have the D2P if their corresponding base algebras are infinite dimensional.*

Now we introduce a Banach space  $X$ -valued function space (over a function space) on a Hausdorff space and we show that this space is isometrically isomorphic to a  $(X^{**}, w^*)$ -valued function space over a uniform algebra. Let  $\Omega$  be a Hausdorff space and let  $X$  be a Banach space equipped with the norm topology. The space  $C_b(\Omega, X)$  is the Banach space of all bounded  $X$ -valued continuous functions over  $\Omega$  equipped with the supremum norm.

**Definition 4.4.** *Let  $X$  be a Banach space. An  $X$ -valued function space over a function algebra on  $\Omega$ , denoted by  $A(\Omega, X)$ , is a closed subspace of  $C_b(\Omega, X)$  satisfying*

- (i) *the base algebra  $A := \{x^* \circ f : f \in A(\Omega, X)\}$  is a function algebra on  $\Omega$ .*
- (ii)  *$fg \in A(\Omega, X)$  for every  $f \in A$  and  $g \in A(\Omega, X)$ .*

Given a function algebra  $A$  on a Hausdorff space  $\Omega$ , let  $M_A$  be its maximal ideal space consisting of all nonzero algebraic homomorphisms from  $A$  to  $\mathbb{C}$ .  $M_A$  is a compact Hausdorff space with the Gelfand topology [21, Theorem 11.9]. The Gelfand transform  $\hat{f} : M_A \rightarrow \mathbb{C}$  of  $f \in A$  is defined by  $\hat{f}(\phi) = \phi(f)$ . Now for  $g \in A(\Omega, X)$  and  $x^* \in X^*$  let  $\hat{g}(\phi)(x^*) = \phi(x^* \circ g)$ . From the fact that

$$|\hat{g}(\phi)(x^*)| = |\phi(x^* \circ g)| \leq \|x^* \circ g\|_\infty = \sup_{t \in \Omega} \{|x^*(g(t))|\} \leq \|g\| \|x^*\|_{X^*},$$

the mapping  $\hat{g}(\phi)$  is a bounded linear functional on  $X^*$ . Moreover,  $\hat{g} : M_A \rightarrow X^{**}$  is continuous on  $M_A$  if we consider the weak\*-topology on  $X^{**}$ . Let  $A(M_A, (X^{**}, w^*))$  be the set of such  $\hat{g}$ 's.

**Theorem 4.5.** *Let  $X$  be a Banach space and  $A(\Omega, X)$  be an  $X$ -valued function space over a base algebra  $A$  on a Hausdorff space  $\Omega$ . Then  $A(\Omega, X)$  is isometrically isomorphic to an  $(X^{**}, w^*)$ -valued function space over a base algebra  $\hat{A}$  on the compact Hausdorff space  $M_A$ . In fact,  $M_A$  is the maximal ideal space of  $A$  with the Gelfand topology.*

We split the proof of Theorem 4.5 into several parts for readability. First we show that the mapping  $g \mapsto \hat{g}$  is an isometry.

**Proposition 4.6.** *Let  $X$  be a Banach space, let  $A(\Omega, X)$  be an  $X$ -valued function space over a base algebra  $A$  on  $\Omega$  and let  $M_A$  be the maximal ideal space of  $A$ . Then for every  $g \in A(\Omega, X)$  and  $x^* \in X^*$ , the mapping  $g \mapsto \hat{g}$  where  $\hat{g} \in A(M_A, (X^{**}, w^*))$  is defined by  $\hat{g}(\phi)(x^*) = \phi(x^* \circ g)$  is an isometry.*

*Proof.* By the fact that  $\|\phi\|_\infty = 1$  we obtain

$$\begin{aligned} \|\hat{g}\| &= \sup_{\phi \in M_A} \{\|\hat{g}(\phi)\|\} = \sup_{\phi \in M_A} \sup_{x^* \in B_{X^*}} \{|\hat{g}(\phi)(x^*)|\} = \sup_{\phi \in M_A} \sup_{x^* \in B_{X^*}} \{|\phi(x^* \circ g)|\} \\ &\leq \sup_{x^* \in B_{X^*}} \{\|x^* \circ g\|_\infty\} \\ &= \sup_{x^* \in B_{X^*}} \sup_{t \in \Omega} \{|(x^* \circ g)(t)|\} \\ &\leq \sup_{t \in \Omega} \{\|g(t)\|_X\} = \|g\|. \end{aligned}$$

To show the reverse inequality, let  $\phi_t \in M_A$  be an evaluation functional at  $t \in \Omega_0 = \{t \in \Omega : g(t) \neq 0\}$ . Then we have

$$\begin{aligned} \|g\| &= \sup_{t \in \Omega_0} \{\|g(t)\|_X\} = \sup_{t \in \Omega_0} \sup_{x^* \in B_{X^*}} \{|\phi_t(x^* \circ g)|\} = \sup_{t \in \Omega_0} \sup_{x^* \in B_{X^*}} \{|\phi_t(x^* \circ g)|\} \\ &\leq \sup_{\phi \in M_A} \sup_{x^* \in B_{X^*}} \{|\phi(x^* \circ g)|\} \\ &= \sup_{\phi \in M_A} \sup_{x^* \in B_{X^*}} \{|\hat{g}(\phi)(x^*)|\} \\ &\leq \sup_{\phi \in M_A} \{\|\hat{g}(\phi)\|\} = \|\hat{g}\|, \end{aligned}$$



so the mapping  $g \mapsto \hat{g}$  is an isometry.  $\square$

Now we show that the space  $A(M_A, (X^{**}, w^*))$  satisfies the condition (i) and (ii) in the Definition 2.2 for a  $(X, \tau)$ -valued function space over a uniform algebra on a compact Hausdorff space.

**Proposition 4.7.** *Let  $X$  be a Banach space, let  $A(\Omega, X)$  be a  $X$ -valued function space over a base algebra  $A$  on  $\Omega$ , and let  $M_A$  be the maximal ideal space of  $A$ . Then the space*

$$A_1 = \{x^* \circ \hat{g} : \hat{g} \in A(M_A, (X^{**}, w^*)) \text{ and } x^* \in X^*\}$$

*is a uniform algebra over  $M_A$ .*

*Proof.* From the definition of the Gelfand transformation, notice that  $x^* \circ \hat{g}(\phi) = \phi(x^* \circ g) = \widehat{x^* \circ g}(\phi)$ . Hence we can rewrite  $A_1 = \{\widehat{x^* \circ g} : g \in A(\Omega, X) \text{ and } x^* \in X^*\}$ . We see that the set  $A_1$  is, in fact, the image of the Gelfand transformation on  $A$ . Thus  $A$  is isometrically isomorphic to  $A_1$  and so  $A_1$  is a closed subalgebra of  $C(M_A)$  that separates the points of  $M_A$  and contains constant functions. This shows that  $A_1$  is a uniform algebra over  $M_A$ .  $\square$

**Proposition 4.8.** *Suppose that  $X$  is a Banach space and  $A(\Omega, X)$  is an  $X$ -valued function space over a base algebra  $A$  on  $\Omega$ . Let  $M_A$  be the maximal ideal space of  $A$  and let*

$$A_1 = \{x^* \circ \hat{g} : \hat{g} \in A(M_A, (X^{**}, w^*)) \text{ and } x^* \in X^*\}.$$

*Then  $\phi \cdot \hat{g} \in A(M_A, (X^{**}, w^*))$  for every  $\phi \in A_1$  and  $\hat{g} \in A(M_A, (X^{**}, w^*))$ .*

*Proof.* Let  $\hat{f}, \hat{g} \in A(M_A, (X^{**}, w^*))$  and let  $x^* \circ \hat{f} \in A_1$  where  $x^* \in (X^{**}, w^*)^* = X^*$ . We claim that  $(x^* \circ \hat{f}) \cdot \hat{g} \in A(M_A, (X^{**}, w^*))$ . Note first that  $h = (x^* \circ f) \cdot g \in A(\Omega, X)$ . We need to show that  $\hat{h} = \widehat{(x^* \circ f) \cdot g}$ .

Since  $x^* \circ \hat{f}(\phi) = x^* \circ f(\phi) = \phi(x^* \circ f)$  for every nonzero algebra homomorphism  $\phi \in M_A$ , we have

$$\hat{h}(\phi)(y^*) = \widehat{(x^* \circ f \cdot g)}(\phi)(y^*) = (x^* \circ \hat{f})(\phi) \cdot \hat{g}(\phi)(y^*) = ((x^* \circ \hat{f}) \cdot \hat{g})(\phi)(y^*)$$

for  $y^* \in X^*$ . Therefore,  $\hat{h} = (x^* \circ \hat{f}) \cdot \hat{g}$  and this proves our claim.  $\square$

Theorem 4.2 and Theorem 4.5 prove the following.

**Corollary 4.9.** *Suppose that  $X$  is a Banach space and  $A(\Omega, X)$  is an  $X$ -valued function space over a function algebra  $A$  on a Hausdorff space  $\Omega$ . If the base algebra  $A$  is infinite dimensional, then  $A(\Omega, X)$  has the D2P.*

For Banach spaces  $X$  and  $Y$ , let  $C_b(B_X, Y)$  be the bounded  $Y$ -valued continuous functions on the unit ball  $B_X$ . The space  $A_b(B_X, Y)$  is defined to be a closed subspace of  $C_b(B_X, Y)$  that consists of holomorphic functions on the interior of  $B_X$  and the space  $A_u(B_X, Y)$  is a closed subspace of  $A_b(B_X, Y)$  consisting of uniformly continuous functions on  $B_X$ . It is easy to check that  $C_b(B_X, Y)$ ,  $A_b(B_X, Y)$  and  $A_u(B_X, Y)$  are  $Y$ -valued function spaces in Definition 4.4. Hence we prove the following.

**Corollary 4.10.** *Let  $X$  and  $Y$  be Banach spaces. Then  $C_b(B_X, Y)$ ,  $A_b(B_X, Y)$  and  $A_u(B_X, Y)$  have the D2P.*

## 5. DAUGAVET POINTS AND $\Delta$ -POINTS ON VECTOR-VALUED FUNCTION SPACES OVER A UNIFORM ALGEBRA

In this section, we present another application of Lemma 2.3 to the Daugavet points and  $\Delta$ -points of  $A(K, X)$  with respect to its Shilov boundary of the base algebra. Recall that a Banach space  $X$  has the Daugavet property if for every rank-one operator  $T : X \rightarrow X$  satisfies  $\|I + T\| = 1 + \|T\|$ . Well-known examples are  $L_1(\mu)$  and  $L_\infty(\mu)$  where  $\mu$  is a nonatomic measure [11, 18] and  $C(K)$  where  $K$  does not have isolated points [10]. An infinite dimensional uniform algebra over a compact Hausdorff space satisfies the Daugavet property if the Shilov boundary does not contain isolated points [25, 26]. More generally, if a locally compact Hausdorff space  $L$  does not have isolated points, somewhat regular

subspaces of  $C_0(L)$  also have the Daugavet property [4]. Similar results are extended to vector-valued spaces. The space  $C(K, X)$  has the Daugavet property if  $K$  does not have isolated points [24]. A subspace  $A^X$  of  $C(K, X)$ , defined by  $A^X = \{f \in C(K, X) : x^* \circ f \in A\}$  where  $A$  is a uniform algebra, is known to satisfy the Daugavet property from the fact that it satisfies the polynomial Daugavet property [7].

We mentioned earlier that we can investigate the Daugavet property and the DLD2P of Banach spaces through the Daugavet and  $\Delta$ -points. So it is clear that the nonexistence of isolated points in the Shilov boundary of an infinite dimensional uniform algebra implies that every point on the unit sphere  $S_A$  is a Daugavet point. However, a characterization of Daugavet points and  $\Delta$ -points on an infinite dimensional uniform algebra has not been known. In [1], the  $\Delta$ -points and the Daugavet points for  $C(K)$  are determined by norm-attainment at a limit point of  $K$ . Motivated by this result, we provide a characterization for the Daugavet points and the  $\Delta$ -points on the vector-valued function space  $A(K, X)$  over a uniform algebra when  $X$  is uniformly convex. First we have the following observation on the  $\Delta$ -points.

**Lemma 5.1.** [1, Lemma 2.1] *Let  $X$  be a Banach space. The following are equivalent:*

- (i)  $x \in S_X$  is a  $\Delta$ -point.
- (ii) for every  $x^* \in X^*$  with  $x^*x = 1$ , the projection  $P = x^* \otimes x$  satisfies  $\|I - P\| \geq 2$ .

We show that if  $f \in S_{A(K, (X, \tau))}$  attains its norm at a limit point on the Shilov boundary of its base algebra with the additional condition  $A \otimes X \subset A(K, (X, \tau))$ , it is a Daugavet point. For a function algebra  $A$  on  $\Omega$  and a Banach space  $X$ ,  $A \otimes X$  means the set of all functions  $f \otimes x$  ( $f \in A$ ,  $x \in X$ ) defined by  $(f \otimes x)(t) = f(t)x$  ( $t \in \Omega$ ).

**Theorem 5.2.** *Let  $K$  be a compact Hausdorff space and let  $X$  be a Banach space endowed with a locally convex Hausdorff topology  $\tau$  compatible to a dual pair. Suppose that a function space  $A(K, (X, \tau))$  over a base algebra  $A$  satisfies the additional condition that  $A \otimes X \subset A(K, (X, \tau))$  and let  $\Gamma$  be the Shilov boundary of  $A$ . If  $f$  is a norm-one element of  $A(K, (X, \tau))$  and there is a limit point  $t_0$  of  $\Gamma$  such that  $\|f\| = \|f(t_0)\|_X$ , then  $f$  is a Daugavet point.*

*Proof.* Suppose that  $\|f\| = \|f(t_0)\|_X = 1$  for a limit point  $t_0$  of  $\Gamma$ . Since  $\Gamma$  is closed,  $t_0$  is an element of  $\Gamma$ . Let  $A(\Gamma, (X, \tau))$  be set of all restrictions of  $f \in A(K, (X, \tau))$  to the Shilov boundary  $\Gamma$ . By Lemma 2.4, the mapping, defined by  $g \in A(K, (X, \tau)) \mapsto g|_\Gamma$ , is an isometry. So it is enough to show that  $f|_\Gamma$  is a Daugavet point of  $A(\Gamma, (X, \tau))$  and we may assume that  $K = \Gamma$ . Denote the base algebra of  $A(K, (X, \tau))$  as  $A$ .

Now fix  $g \in B_{A(K, (X, \tau))}$  and  $\epsilon > 0$ . Let  $U$  be an open set containing  $t_0$  such that  $\|f(t) - f(t_0)\|_X < \epsilon$  for every  $t \in U$ . By the Hausdorff condition and the fact that  $t_0$  is a limit point of  $\Gamma$ , we can find pairwise disjoint nonempty open subsets  $U_1, U_2, \dots$  such that  $U_n \subset U$  for each  $n \in \mathbb{N}$ . Since the set of strong boundary points is dense in  $\Gamma$ , each  $U_n$  contains a strong boundary point. By Lemma 2.3, for each  $n \in \mathbb{N}$ , there exist  $\phi_n \in A$  and  $t_n \in U_n \cap \Gamma_0$  such that  $\phi_n(t_n) = \|\phi_n\|_\infty = 1$ ,  $|\phi_n(t)| < \frac{\epsilon}{3 \cdot 2^n}$  for all  $t \in K \setminus U_n$ , and

$$|\phi_n(t)| + \left(1 - \frac{\epsilon}{3 \cdot 2^n}\right) |1 - \phi_n(t)| \leq 1 \text{ for all } t \in K.$$

Now for each  $n \in \mathbb{N}$ , define a function  $g_n(t) = (1 - \frac{\epsilon}{3 \cdot 2^n})(1 - \phi_n(t))g(t) - \phi_n(t)f(t_0)$ . Then  $g_n$  is an element of  $A(K, (X, \tau))$  and we show that  $g_n \in B_{A(K, (X, \tau))}$  for all  $n \in \mathbb{N}$ . It is easy to see that for all  $t \in K$ ,

$$\|g_n(t)\|_X = \left\| \left(1 - \frac{\epsilon}{3 \cdot 2^n}\right)(1 - \phi_n(t))g(t) - \phi_n(t)f(t_0) \right\|_X \leq \left(1 - \frac{\epsilon}{3 \cdot 2^n}\right) |1 - \phi_n(t)| + |\phi_n(t)| \leq 1.$$

Hence  $\|g_n\| \leq 1$ . Notice that  $g_n(t_n) = -f(t_0)$ . This implies that

$$\|f - g_n\| \geq \|f(t_n) - g_n(t_n)\|_X = \|f(t_n) + f(t_0)\|_X > 2\|f(t_0)\|_X - \|f(t_n) - f(t_0)\|_X \geq 2 - \epsilon,$$

and so  $g_n \in \Delta_\epsilon(f)$ . Furthermore, for every  $t \in U_n$  we see that

$$\begin{aligned} \|g(t) - g_n(t)\|_X &= \left\| g(t) - \left(1 - \frac{\epsilon}{3 \cdot 2^n}\right) (1 - \phi_n(t))g(t) + \phi_n(t)f(t_0) \right\|_X \\ &= \left\| \frac{\epsilon}{3 \cdot 2^n} g(t) + \left(1 - \frac{\epsilon}{3 \cdot 2^n}\right) \phi_n(t)g(t) + \phi_n(t)f(t_0) \right\|_X \\ &\leq \frac{\epsilon}{3 \cdot 2^n} + \left(1 - \frac{\epsilon}{3 \cdot 2^n}\right) + 1 = 2. \end{aligned}$$

Since  $|\phi_n(t)| < \frac{\epsilon}{3 \cdot 2^n}$  for  $t \in K \setminus U_n$ , we also have

$$\begin{aligned} \|g(t) - g_n(t)\|_X &= \left\| \frac{\epsilon}{3 \cdot 2^n} g(t) + \left(1 - \frac{\epsilon}{3 \cdot 2^n}\right) \phi_n(t)g(t) + \phi_n(t)f(t_0) \right\|_X \\ &\leq \frac{\epsilon}{3 \cdot 2^n} + \left(1 - \frac{\epsilon}{3 \cdot 2^n}\right) \frac{\epsilon}{3 \cdot 2^n} + \frac{\epsilon}{3 \cdot 2^n} < \frac{\epsilon}{2^n}. \end{aligned}$$

From the fact that each  $U_n$ 's are pairwise disjoint, we obtain, for each  $t \in \cup_{n=1}^m U_n$ ,

$$\left\| g(t) - \frac{1}{m} \sum_{n=1}^m g_n(t) \right\|_X \leq \frac{1}{m} \left( 2 + \left( \frac{\epsilon}{2} + \frac{\epsilon}{4} + \dots + \frac{\epsilon}{2^m} \right) \right) = \frac{1}{m} \left( 2 + \left( 1 - \frac{1}{2^m} \right) \epsilon \right)$$

and for each  $t \in K \setminus \cup_{n=1}^m U_n$ , we have

$$\left\| g(t) - \frac{1}{m} \sum_{n=1}^m g_n(t) \right\|_X \leq \frac{1}{m} \left( \frac{\epsilon}{2} + \frac{\epsilon}{4} + \dots + \frac{\epsilon}{2^m} \right) = \frac{1}{m} \left( 1 - \frac{1}{2^m} \right) \epsilon.$$

Hence, we get

$$\left\| g - \frac{1}{m} \sum_{n=1}^m g_n \right\| \leq \frac{1}{m} \left( 2 + \left( 1 - \frac{1}{2^m} \right) \epsilon \right).$$

By taking the limit  $m \rightarrow \infty$ , we show that  $g \in \overline{\text{conv}}\Delta_\epsilon(f)$ . Hence  $B_X = \overline{\text{conv}}\Delta_\epsilon(f)$ , in other words,  $f$  is a Daugavet point.  $\square$

From this, we obtain a sufficient condition for the Daugavet property of  $A(K, X)$  since the map  $t \mapsto \|f(t)\|$  is continuous on a compact space and attains its maximum on  $K$  if  $\tau$  is norm topology and  $f \in A(K, X)$ .

**Corollary 5.3.** *Let  $K$  be a compact Hausdorff space and let  $X$  be a Banach space. Suppose that a function space  $A(K, X)$  over a base algebra  $A$  satisfies the additional condition that  $A \otimes X \subset A(K, X)$ . If the Shilov boundary  $\Gamma$  of the base algebra  $A$  does not have isolated points, then  $A(K, X)$  has the Daugavet property.*

Furthermore, when  $X$  is uniformly convex, Daugevet points and  $\Delta$ -points are the same in  $A(K, X)$  and characterized by the norm-attainment at a limit point of  $\Gamma$ . Recall that the modulus of convexity  $\delta_X(\epsilon)$  of a Banach space  $X$  is defined by for  $0 < \epsilon < 2$

$$\delta_X(\epsilon) = \inf \left\{ 1 - \left\| \frac{x+y}{2} \right\| : x, y \in B_X, \|x-y\| \geq \epsilon \right\}.$$

A Banach space  $X$  is said to be uniformly convex if  $\delta_X(\epsilon) > 0$  for each  $0 < \epsilon < 2$ .

**Theorem 5.4.** *For a uniformly convex Banach space  $X$  and a compact Hausdorff space  $K$ , let  $\Gamma$  be the Shilov boundary of the base algebra  $A$  of a function space  $A(K, X)$  and let  $f \in S_{A(K, X)}$ . Then (i)  $\implies$  (ii)  $\implies$  (iii) holds.*

(i)  $f$  is a Daugavet point.

(ii)  $f$  is a  $\Delta$ -point.

(iii) there is a limit point  $t_0$  of  $\Gamma$  such that  $\|f\| = \|f(t_0)\|_X$ .

Moreover, if we assume the additional condition that  $A \otimes X \subset A(K, X)$ , then (i), (ii) and (iii) are equivalent.

*Proof.* In view of Lemma 2.4,  $A(K, X)$  is isometric to  $A(\Gamma, X)$  and so it is enough to prove our claim with respect to  $f|_{\Gamma} \in S_{A(\Gamma, X)}$ . So we may assume that  $K = \Gamma$ . (i)  $\implies$  (ii) is clear from their definitions. (iii)  $\implies$  (i) is already shown in Theorem 5.2 with the additional assumption that  $A \otimes X \subset A(K, X)$ . Hence we only need to show (ii)  $\implies$  (iii).

Assume to the contrary that  $f \in S_{A(K, X)}$  is a  $\Delta$ -point but  $\|f\| \neq \|f(t)\|_X$  for every limit point of  $K$ . Let  $F = \{t \in K : \|f(t)\|_X = 1\}$ . Then  $F$  only contains isolated points of  $K$  by the assumption and is nonempty. Since  $K$  is compact, we also see that  $|F| < \infty$ . For each  $t \in F$  we can always find  $x_t^* \in S_{X^*}$  such that  $x_t^*(f(t)) = 1$ .

Since  $X$  is uniformly convex, let  $\delta = \delta_X \left( \frac{1}{2|F|} \right) > 0$ , where  $|F|$  is the number of elements of  $F$ . If  $\operatorname{Re} x_t^*(x) \geq 1 - \delta$  and  $x \in B_X$ , then  $\left\| \frac{x+f(t)}{2} \right\| \geq \operatorname{Re} x_t^* \left( \frac{x+f(t)}{2} \right) \geq 1 - \frac{1}{2}\delta > 1 - \delta_X \left( \frac{1}{2|F|} \right)$ . This means that  $\|x - f(t)\| < \frac{1}{2|F|}$  by the definition of  $\delta_X$ . Hence

$$(3) \quad \text{if } \operatorname{Re} x_t^*(x) \geq 1 - \delta \text{ and } x \in B_X \text{ then } \|x - f(t)\|_X \leq \frac{1}{2|F|} \text{ for every } t \in F.$$

Let  $\epsilon = 1 - \max_{t \in K \setminus F} \|f(t)\|_X > 0$ . Define a bounded linear functional  $\psi$  on  $A(K, X)$  by

$$\psi(g) = \frac{1}{|F|} \sum_{t \in F} x_t^*(g(t)),$$

where  $g \in A(K, X)$ . Notice that  $\|\psi\| = \psi(f) = 1$  and this defines a bounded projection  $P : A(K, X) \rightarrow A(K, X)$  by  $P(g) = \psi(g)f$  for  $g \in A(K, X)$ . It is clear that  $\|P\| = \|P(f)\| = \|f\| = 1$ . We see that  $P$  is in fact a norm-one projection on  $A(K, X)$ . Then in view of Lemma 5.1, we should have  $\|I - P\| \geq 2$ . But we show that in fact  $\|I - P\| < 2$ .

For a given  $g \in B_{A(K, X)}$ , if  $t \in K \setminus F$  we have

$$(4) \quad \|g(t) - Pg(t)\|_X = \|g(t) - \psi(g)f(t)\|_X \leq 1 + \|f(t)\|_X \leq 2 - \epsilon.$$

Now we divide  $F$  into two parts. Let  $t_0 \in F_1 = \{t \in F : |x_{t_0}^*(g(t_0))| \geq 1 - \delta\}$ . Then there exists a scalar  $\lambda_0 \in B_{\mathbb{C}}$  such that  $|x_{t_0}^*(g(t_0))| = \lambda_0 x_{t_0}^*(g(t_0)) = x_{t_0}^*(\lambda_0 g(t_0)) \geq 1 - \delta$ . Then by (3), we have  $\|\lambda_0 g(t_0) - f(t_0)\|_X \leq \frac{1}{2|F|}$ . Then

$$\begin{aligned} \|g(t_0) - Pg(t_0)\|_X &= \left\| g(t_0) - \frac{1}{|F|} \sum_{t \in F} x_t^*(g(t))f(t) \right\|_X \leq \left\| g(t_0) - \frac{1}{|F|} x_{t_0}^*(g(t_0))f(t_0) \right\|_X + \frac{|F| - 1}{|F|} \\ &= \left\| \lambda_0 g(t_0) - \frac{1}{|F|} x_{t_0}^*(\lambda_0 g(t_0))f(t_0) \right\|_X + \frac{|F| - 1}{|F|}. \end{aligned}$$

From the fact that  $x_{t_0}^*(\lambda_0 g(t_0)) \leq 1$ , we have  $1 - \frac{1}{|F|} x_{t_0}^*(\lambda_0 g(t_0)) \geq 0$ . Hence

$$\begin{aligned} \left\| \lambda_0 g(t_0) - \frac{1}{|F|} x_{t_0}^*(\lambda_0 g(t_0))f(t_0) \right\|_X &\leq \|\lambda_0 g(t_0) - f(t_0)\|_X + \|f(t_0)\|_X \left( 1 - \frac{1}{|F|} x_{t_0}^*(\lambda_0 g(t_0)) \right) \\ &\leq \frac{1}{2|F|} + 1 - \frac{1}{|F|}(1 - \delta), \end{aligned}$$

and so

$$(5) \quad \|g(t_0) - Pg(t_0)\|_X \leq \frac{1}{2|F|} + 1 - \frac{1}{|F|}(1 - \delta) + \frac{|F| - 1}{|F|} \leq 2 - \frac{1}{2|F|}.$$

When  $t_0 \in F_2 = F \setminus F_1$ , we have  $|x_{t_0}^*(g(t_0))| < 1 - \delta$ . So we see that

$$\begin{aligned} \|g(t_0) - Pg(t_0)\|_X &\leq \left\| \lambda_0 g(t_0) - \frac{1}{|F|} x_{t_0}^*(\lambda_0 g(t_0)) f(t_0) \right\|_X + \frac{|F| - 1}{|F|}. \\ &\leq 1 + \frac{1}{|F|} (1 - \delta) + \frac{|F| - 1}{|F|} \leq 2 - \frac{\delta}{|F|}. \end{aligned}$$

Hence, with (4) and (5) we obtain  $\|g - Pg\| \leq \max\{2 - \frac{1}{2|F|}, 2 - \frac{\delta}{|F|}, 2 - \epsilon\} < 2$ . Since  $g \in B_{A(K,X)}$  is chosen arbitrarily,  $\|I - P\| < 2$ , and we get a contradiction.  $\square$

Since  $\mathbb{C}$  is uniformly convex, we have the following consequences for a scalar-valued uniform algebra.

**Corollary 5.5.** *Let  $K$  be a compact Hausdorff space and let  $\Gamma$  be the Shilov boundary of  $A(K)$ . Then for  $f \in S_{A(K,X)}$ , the following are equivalent:*

- (i)  $f$  is a Daugavet point.
- (ii)  $f$  is a  $\Delta$ -point.
- (iii) there is a limit point  $t_0$  of  $\Gamma$  such that  $\|f\|_\infty = |f(t_0)|$ .

By the characterization given in Theorem 5.4, we show when the Daugavet property, DD2P, and DLD2P of  $A(K, X)$  are equivalent to each other.

**Theorem 5.6.** *Let  $X$  be a uniformly convex Banach space, let  $K$  be a compact Hausdorff space, and let  $\Gamma$  be the Shilov boundary of the base algebra of  $A(K, X)$ . Then (i)  $\implies$  (ii)  $\implies$  (iii)  $\implies$  (iv) holds:*

- (i)  $A(K, X)$  has the Daugavet property.
- (ii)  $A(K, X)$  has the DD2P.
- (iii)  $A(K, X)$  has the DLD2P.
- (iv) The Shilov boundary  $\Gamma$  does not have isolated points.

Moreover, if we assume the additional condition that  $A \otimes X \subset A(K, X)$ , then (i), (ii), (iii) and (iv) are equivalent.

*Proof.* Since  $A(K, X)$  is isometric to  $A(\Gamma, X)$  by Lemma 2.4, showing the equivalence for  $A(\Gamma, X)$  is enough. It is a well-known that (i)  $\implies$  (ii)  $\implies$  (iii) [6]. By Corollary 5.3, we have (iv)  $\implies$  (i) with the additional assumption that  $A \otimes X \subset A(K, X)$ .

Finally we prove (iii)  $\implies$  (iv). Assume that  $A(\Gamma, X)$  has the DLD2P but the Shilov boundary  $\Gamma$  of the base algebra of  $A(K, X)$  has an isolated point  $t_0$ . Since the set of strong boundary points  $\Gamma_0$  is dense in  $\Gamma$ , this  $t_0$  is also in  $\Gamma_0$ . Then by Lemma 2.5 there is  $\phi \in A$  such that  $\phi(t_0) = 1 = \|\phi\|$  and  $\phi(t) = 0$  for all  $t \in \Gamma \setminus \{t_0\}$ . Since  $\phi$  is in the base algebra  $A$ , there is  $g \in A(\Gamma, X)$  such that  $\phi = x^* \circ g$  for some  $x^* \in X^*$ . Then  $g(t_0) \neq 0$  and let  $h = \frac{\phi}{\|g(t_0)\|_X} g$ . So  $h$  is an element of  $A(K, X)$ . Since  $\|h\| = \|h(t_0)\| = 1$  and  $\|h(t)\| = 0$  for every point  $t$  of  $\Gamma$  other than  $t_0$ . So  $\|h(t)\| = 0$  for every limit point  $t$  of  $\Gamma$ . Hence in view of Theorem 5.4, we see that  $h \in S_{A(K,X)}$  is not a  $\Delta$ -point, and this is a contradiction to our assumption that  $A(K, X)$  has the DLD2P, which states that every function in  $S_{A(K,X)}$  is a  $\Delta$ -point. Therefore (iii)  $\implies$  (iv) is proved.  $\square$

As we mentioned before, it is well-known that the nonexistence of isolated points on the Shilov boundary implies that the scalar-valued uniform algebra has the Daugavet property [25, 26]. Here we have something even further that the Daugavet property, DD2P, and DLD2P are equivalent to each other.

**Corollary 5.7.** *Let  $K$  be a compact Hausdorff space and let  $\Gamma$  be the Shilov boundary of a uniform algebra  $A$ . Then the following are equivalent:*

- (i)  $A$  has the Daugavet property.
- (ii)  $A$  has the DD2P.
- (iii)  $A$  has the DLD2P.

(iv) *The Shilov boundary  $\Gamma$  does not have isolated points.*

We conclude this article with showing when the space  $A(K, X)$  has the convex-DLD2P. But here we take a slightly different approach. To use Lemma 2.3 we need to find a limit point of  $\Gamma$  that is also a strong boundary point, but the existence of such point does not seem to be guaranteed in general. So here we use a different version of the Urysohn-type lemma for strong peak points. For a uniform algebra  $A$  over a compact Hausdorff space  $K$ , an element  $t_0 \in K$  is said to be a strong peak point if there exists a function  $f \in A$  such that  $|f(t_0)| = \|f\|_\infty$  and  $\|f\|_\infty > \sup\{|f(t)| : t \in K \setminus U\}$  for every open neighborhood  $U$  of  $t_0$ . Let us denote the set of all strong peak points for  $A$  by  $\rho(A)$ .

**Lemma 5.8.** [15, Lemma 3] *Let  $\Omega$  be a Hausdorff space and  $A$  be a subalgebra of  $C_b(\Omega)$ . Suppose that  $U$  is an open subset of  $\Omega$  and  $t_0 \in U \cap \rho(A)$ . Then given  $0 < \epsilon < 1$ , there exists  $\phi \in A$  such that  $\|\phi\| = 1 = \phi(t_0)$ ,  $\sup_{t \in K \setminus U} |\phi(t)| < \epsilon$ , and for all  $t \in \Omega$ ,*

$$|\phi(t)| + (1 - \epsilon)|1 - \phi(t)| \leq 1.$$

The original statement of Lemma 5.8 is given with  $\|\phi\| = 1 = |\phi(t_0)|$ . However, if we observe the proof of Lemma 3 in [15] carefully, it is easy to see that the statement without the absolute value also holds.

**Theorem 5.9.** *For a compact Hausdorff space  $K$  and a uniformly convex Banach space  $X$ , suppose that a function space  $A(K, X)$  over a base algebra  $A$  satisfies the additional condition that  $A \otimes X \subset A(K, X)$ . Let  $\Gamma'$  be the set of limit points in the Shilov boundary  $\Gamma$  of the base algebra  $A$  and let  $\rho(A)$  the set of all strong peak points for  $A$ . If  $\rho(A) \cap \Gamma' \neq \emptyset$ , then the space  $A(K, X)$  has the convex-DLD2P.*

*Proof.* By the isometry in Lemma 2.4, we may assume that  $K = \Gamma$ . We need to show that  $S_{A(K, X)} \subset \overline{\text{conv}}\Delta$ , where  $\Delta$  is the set of all  $\Delta$ -points of  $A(K, X)$ .

Fix  $f \in S_{A(K, X)}$ . Pick  $t_0 \in \rho(A) \cap K'$  and let  $\lambda = \frac{1 + \|f(t_0)\|_X}{2}$ . For  $\epsilon > 0$ , let  $U$  be an open neighborhood containing  $t_0$  such that  $\|f(t) - f(t_0)\|_X < \epsilon$ . By Lemma 5.8, there exists a function  $\phi \in A$  such that  $\phi(t_0) = \|\phi\|_\infty = 1$ ,  $\sup_{t \in K \setminus U} |\phi(t)| < \epsilon$  and

$$|\phi(t)| + (1 - \epsilon)|1 - \phi(t)| \leq 1.$$

for every  $t \in K$ . Choose a norm-one vector  $v_0 \in X$  and set

$$x_0 = \begin{cases} \frac{f(t_0)}{\|f(t_0)\|_X} & \text{if } f(t_0) \neq 0 \\ v_0 & \text{if } f(t_0) = 0. \end{cases}$$

Now define two functions

$$\begin{aligned} f_1(t) &= (1 - \epsilon)(1 - \phi(t))f(t) + \phi(t)x_0 \\ f_2(t) &= (1 - \epsilon)(1 - \phi(t))f(t) - \phi(t)x_0, \quad t \in K. \end{aligned}$$

Since  $A \otimes X \subset A(K, X)$ , we see that the functions  $f_1, f_2 \in A(K, X)$ . Notice that

$$\begin{aligned} \|f_1(t)\|_X &= \|(1 - \epsilon)(1 - \phi(t))f(t) + \phi(t)x_0\|_X \\ &\leq (1 - \epsilon)|1 - \phi(t)| + |\phi(t)| \leq 1, \end{aligned}$$

for every  $t \in K$ , in particular,  $\|f_1(t_0)\|_X = 1$ . Hence  $\|f_1\| = \|f_1(t_0)\|_X = 1$ . We can also show  $\|f_2\| = \|f_2(t_0)\|_X = 1$  by the same argument. Thus by Theorem 5.4, we see that  $f_1, f_2 \in \Delta$ . Let  $g(t) = \lambda f_1(t) + (1 - \lambda)f_2(t)$ . We consider two cases.

(i) First, consider when  $f(t_0) \neq 0$ . Then  $g = (1 - \epsilon)(1 - \phi(t))f(t) + \phi(t)f(t_0)$ . We see that

$$\begin{aligned} \|g(t) - f(t)\|_X &= \|(1 - \epsilon)(1 - \phi(t))f(t) + \phi(t)f(t_0) - f(t)\|_X \\ &= \|(1 - \epsilon)(1 - \phi(t))f(t) + \phi(t)f(t_0) - (1 - \epsilon)f(t) - \epsilon f(t)\|_X \\ &= \|(1 - \epsilon)(-\phi(t))f(t) + (1 - \epsilon)\phi(t)f(t_0) + \epsilon\phi(t)f(t_0) - \epsilon f(t)\|_X \\ &= \|(1 - \epsilon)\phi(t)(f(t_0) - f(t)) + \epsilon\phi(t)f(t_0) - \epsilon f(t)\|_X \\ &\leq (1 - \epsilon)|\phi(t)| \cdot \|f(t) - f(t_0)\|_X + \epsilon|\phi(t)| \cdot \|f(t_0)\|_X + \epsilon\|f(t)\|_X \\ &\leq (1 - \epsilon)|\phi(t)| \cdot \|f(t) - f(t_0)\|_X + 2\epsilon. \end{aligned}$$

For  $t \in U$ , we have  $(1 - \epsilon)|\phi(t)| \cdot \|f(t) - f(t_0)\|_X \leq (1 - \epsilon)\epsilon < \epsilon$ . Since  $|\phi(t)| < \epsilon$  for  $t \in K \setminus U$ , we have  $(1 - \epsilon)|\phi(t)| \cdot \|f(t) - f(t_0)\|_X \leq 2(1 - \epsilon)\epsilon < 2\epsilon$  by the triangle inequality. Therefore  $\|g - f\| < 4\epsilon$ , and this implies that  $f \in \overline{\text{conv}}\Delta$ .

(ii) Now consider when  $f(t_0) = 0$ . Then for all  $t \in U$ ,  $\|f(t)\|_X < \epsilon$ . Notice that  $\lambda = \frac{1}{2}$  and  $g(t) = (1 - \epsilon)(1 - \phi(t))f(t)$ . Hence

$$\begin{aligned} \|g(t) - f(t)\|_X &= \|(1 - \epsilon)(1 - \phi(t))f(t) - (1 - \epsilon)f(t) - \epsilon f(t)\|_X \\ &\leq (1 - \epsilon)|\phi(t)| \cdot \|f(t)\|_X + \epsilon\|f(t)\|_X \leq (1 - \epsilon)|\phi(t)| \cdot \|f(t)\|_X + \epsilon. \end{aligned}$$

For  $t \in U$ , we have  $(1 - \epsilon)|\phi(t)| \cdot \|f(t)\|_X \leq (1 - \epsilon)\epsilon < \epsilon$ . In view of Lemma 5.8, since  $\sup_{t \in K \setminus U} |\phi(t)| < \epsilon$  for  $t \in K \setminus U$ , we have  $(1 - \epsilon)|\phi(t)| \cdot \|f(t)\|_X \leq (1 - \epsilon)\epsilon < \epsilon$ . Therefore  $\|g - f\| < 2\epsilon$ , and this also implies that  $f \in \overline{\text{conv}}\Delta$ .

Since  $f \in S_{A(K,X)}$  is arbitrary, we see that  $S_{A(K,X)} \subset \overline{\text{conv}}(\Delta)$ , and so  $A(K,X)$  has the convex-DLD2P.  $\square$

**Corollary 5.10.** *Let  $K$  be a compact Hausdorff space and  $\Gamma'$  be the set of limit points in the Shilov boundary  $\Gamma$  of a uniform algebra  $A$ . If  $\rho(A) \cap \Gamma' \neq \emptyset$ , then the uniform algebra  $A$  has the convex-DLD2P.*

**Remark 5.11.** The sufficient condition  $\rho(A) \cap \Gamma' \neq \emptyset$  in Corollary 5.10 does not guarantee the DLD2P. For example, let  $K = \{\frac{1}{n} : n \in \mathbb{N}\} \cup \{0\}$  in the real line and let  $A = C(K)$  be the space of continuous functions over  $K$  equipped with the supremum norm. Notice that  $K = \rho(A)$  is its Shilov boundary and  $\rho(A) \cap \Gamma' = \{0\}$ . So in view of Theorem 5.7 and Theorem 5.10, the space  $C(K)$  has the convex-DLD2P, while it does not have DLD2P. In fact,  $C(K)$  is isometric to the space  $c$  of convergent sequences. The same results in the space  $c$  were shown in [1, Corollary 5.4, Remark 5.5].

## REFERENCES

- [1] T. Abrahamsen, R. Haller, V. Lima, and K. Pirk, *Delta- and Daugavet points in Banach Spaces*, Proc. Edin. Math. Soc., Vol. 63, Issue 2, 2020 pp. 475–496.
- [2] M. D. Acosta and A. Kamińska, *Weak neighborhoods of the unit ball in interpolation spaces  $L^1 + L^\infty$  and  $L^1 \cap L^\infty$* , Indiana Univ. Math. J., **57**, No. 1 (2008), 77–96.
- [3] T. Abrahamsen, V. Lima and O. Nygaard, *Remarks on Diameter 2 Properties*, J. Convex Analysis, **20** (2013), pp. 439–452.
- [4] T. Abrahamsen, O. Nygaard, and Märt Põldvere, *New applications of extremely regular function spaces*, Pacific Journal of Mathematics, Vol. **301**, No.2 (2019), pg 385–394.
- [5] J. Bercerra Guerrero and G. López Pérez, *Relatively Weakly Open Subsets of the Unit Ball in Function Spaces*, J. Math. Anal. Appl. **315** (2006), no.2, 544–554.
- [6] J. Bercerra Guerrero, G. López Pérez, and A. Rueda Zoca, *Diametral Diameter Two Properties in Banach Spaces*, J. Convex Anal. **25** (2018), 817–840.
- [7] Y. S. Choi, D. García, S. K. Kim and M. Maestre, *Some geometric properties of disk algebras*, J. Math. Anal. Appl. **409** (2014), no. 1, 147–157.
- [8] B. Cascales, A. Guirao, and V. Kadets, *A Bishop-Phelps-Bollobás type Theorem for Uniform Algebras*, Advances in Mathematics, vol 240, 2013, pp. 370–382
- [9] H.G. Dales, *Banach Algebras and Automatic Continuity*, vol.24, London Mathematical Society Monographs. New Series, the Clarendon Press, Oxford University Press, New York, NY, USA, 2000

- [10] I.K Daugavet, *On a property of completely continuous operators in the space  $C$* , Uspekhi Mat. Nauk **18.5** (1963) 157–158 (Russian)
- [11] C.Foiaş and I. Singer *Points of diffusion of linear operators and almost diffuse operators in spaces of continuous functions*, Math. Z. **87** (1965), 434–450.
- [12] N. Ghoussoub, G. Godefroy, B. Maurey, W. Schachermayer, *Some Topological and Geometrical Structures in Banach Spaces*, Mem. Amer. Math. Soc. **70** (1987), no.378, iv +116. MR 912637 (89h:46024)
- [13] Y. Ivakhno and V. Kadets, *Unconditional Sums of Spaces with Bad Projections*, Visn. Khark. Univ., Ser. Mat. Pryki. Mat. Mekh **645** (2004), 30–35.
- [14] V. Kadets, *The Diametral Strong Diameter 2 Property of Banach spaces Is the Same as the Daugavet Property*, Proc. Amer. Math. Soc, 2020, <https://doi.org/10.1090/proc/15448>
- [15] S.K Kim, H.J Lee, *A Urysohn-type Theorem and the Bishop-Phelps-Bollobás Theorem for Holomorphic Functions*, J. Math. Anal. and Appl, Volume 480, Issue 2, 2019, 123393, ISSN 0022-247X, <https://doi.org/10.1016/j.jmaa.2019.123393>.
- [16] H.J Lee, *Generalized Numerical Index of Function Algebras*, Journal of Function Spaces, vol. 2019, Article ID 9080867, 6 pages, 2019. <https://doi.org/10.1155/2019/9080867>
- [17] G.M Leibowitz, *Lectures on Complex Function Algebras*. Scott, Foresman and Company 1970
- [18] G. Ya, Lozanovskii, *On almost integral operators in  $KB$ -spaces*, Vestnik Leningrad Univ. Mat. Mekh. Astr, **21.7** (1966), 35–44 (Russian)
- [19] L. Narici, E. Beckenstein, *Topological Vector Spaces*, 2nd ed., CRC Press, 2011.
- [20] O. Nygaard and D. Werner, *Slices in the Unit Ball of a Uniform Algebra*, Arch. Math. (Basel) **76** (2001), no.6, pp 441-444.
- [21] W. Rudin, *Functional Analysis*, 2nd ed, McGraw Hill, 1991.
- [22] H. Schaefer, M. Wolff, *Topological Vector spaces*, 2nd ed., Springer-Verlag, 1999.
- [23] K. Pirk, *Diametral Diameter Two Properties, Daugavet-, and  $\Delta$ -points in Banach Spaces*, PhD dissertation, Dissertationes Mathematicae Universitatis Tartuensis, **133**, 2020
- [24] D. Werner, *Recent Progress on the Daugavet property*, Irish Math. Soc. Bulletin, **46**, 2001, pg 77–97
- [25] D. Werner, *The Daugavet equation for operators not fixing a copy of  $C[0, 1]$* , J. Operator Theory **39** (1998), 89–98.
- [26] P. Wojtaszczyk, *Some remarks on the Daugavet equation*, Proc. Amer. Math. Soc **115** (1992), 1047–1052.

DEPARTMENT OF MATHEMATICS EDUCATION, DONGGUK UNIVERSITY - SEOUL, 04620 (SEOUL), REPUBLIC OF KOREA  
 Email address: [hanjulee@dgu.ac.kr](mailto:hanjulee@dgu.ac.kr)

DEPARTMENT OF MATHEMATICS EDUCATION, DONGGUK UNIVERSITY - SEOUL, 04620 (SEOUL), REPUBLIC OF KOREA  
 Email address: [htag@dongguk.edu](mailto:htag@dongguk.edu)