## EXISTENCE OF CO-ROTATING AND TRAVELLING VORTEX PATCHES WITH DOUBLY CONNECTED COMPONENTS FOR ACTIVE SCALAR EQUATIONS

#### DAOMIN CAO, GUOLIN QIN, WEICHENG ZHAN, CHANGJUN ZOU

ABSTRACT. By applying implicit function theorem on contour dynamics, we prove the existence of co-rotating and travelling patch solutions for both Euler and the generalized surface quasi-geostrophic equation. The solutions obtained constitute a desingularization of points vortices when the size of patch support vanishes. In particular, solutions constructed in this paper consist of doubly connected components, which is essentially different from all known results.

#### 1. INTRODUCTION

We consider the following active scalar equation

$$\begin{cases} \partial_t \theta + \mathbf{u} \cdot \nabla \theta = 0 & \text{in } \mathbb{R}^2 \times (0, T) \\ \mathbf{u} = \nabla^{\perp} (-\Delta)^{-1 + \frac{\alpha}{2}} \theta & \text{in } \mathbb{R}^2 \times (0, T) \\ \theta \big|_{t=0} = \theta_0 & \text{in } \mathbb{R}^2, \end{cases}$$
(1.1)

where  $0 \leq \alpha < 2$ ,  $\theta(\boldsymbol{x},t) : \mathbb{R}^2 \times (0,T) \to \mathbb{R}$  is the active scalar being transported by the velocity field  $\mathbf{u}(\boldsymbol{x},t) : \mathbb{R}^2 \times (0,T) \to \mathbb{R}^2$  generated by  $\theta$ , and  $(x_1,x_2)^{\perp} = (x_2,-x_1)$ . The operator  $(-\Delta)^{-1+\frac{\alpha}{2}}$  is given by the expression

$$(-\Delta)^{-1+\frac{\alpha}{2}}\theta(\boldsymbol{x}) = \int_{\mathbb{R}^2} K_{\alpha}(\boldsymbol{x}-\boldsymbol{y})\theta(\boldsymbol{y})d\boldsymbol{y},$$

where  $K_{\alpha}$  is the fundamental solution of  $(-\Delta)^{-1+\frac{\alpha}{2}}$  in  $\mathbb{R}^2$  given by

$$K_{\alpha}(\boldsymbol{x}) = \begin{cases} \frac{1}{2\pi} \ln \frac{1}{|\boldsymbol{x}|}, & \text{if } \alpha = 0;\\ \frac{C_{\alpha}}{2\pi} \frac{1}{|\boldsymbol{x}|^{\alpha}}, & C_{\alpha} = \frac{\Gamma(\alpha/2)}{2^{1-\alpha}\Gamma(\frac{2-\alpha}{2})}, & \text{if } 0 < \alpha < 2 \end{cases}$$

with  $\Gamma$  the Euler gamma function.

When  $\alpha = 0$ , (1.1) corresponds to the vorticity formulation of 2D incompressible Euler equation. In the case  $\alpha = 1$ , (1.1) becomes the surface quasi-geostrophic (SQG) equation, which models geophysical flows and is applied to study the atmosphere circulation and ocean dynamics [29, 34, 37]. The SQG equation also shares mathematical analogy with the three-dimensional incompressible Euler equation [15]. As a generalization of the Euler equation and the SQG equation, the gSQG model (1.1) with  $0 < \alpha < 2$  was proposed by Córdoba et al. in [18].

Mathematical study of (1.1) has attracted more and more attention in the past years. For the case  $\alpha = 0$ , global well-posedness of solutions has already been solved in Sobolev spaces [40]. While Yudovich [49] proved the existence and uniqueness of solutions to (1.1) with the initial data in  $L^1 \cap L^{\infty}$  in 1963. It is natural to ask whether it is possible to extend theories of the Euler equation to the general case  $0 < \alpha < 2$  and a lot of effort has been made in recent years. Constantin et al. [15] established local well-posedness of the gSQG equation for classical solutions with sufficiently regular initial data. Local existence of solutions was also studied in different function spaces, see [13, 38, 47, 48]. Uniqueness and non-uniqueness of weak solutions under certain assumptions was recently shown in [17] and [2] respectively. However, the problem whether the gSQG equation has global solutions or not remains unsolved. In [36], Kiselev and Nazarov constructed solutions of the gSQG equations to the SQG equations with any initial data in  $L^2$ , which was improved by Marchand [39] to any initial data belonging to  $L^p$  with p > 4/3. To give concrete examples for global existence, various kind of global solutions to (1.1) were also constructed.

Due to the structure of the nonlinear term, all radially symmetric functions are stationary solutions to Euler and the gSQG equation. The first explicit non-trivial rotating solution of 2-fold symmetry for  $\alpha = 0$  is constructed by Kirchhoff [35], which is an elliptic patch of semi-axes a and b subjected to a perpetual rotation with uniform angular velocity  $ab/(a+b)^2$ . Deem and Zabusky [20] then carried out a series of numerical simulations, which provided evidence for existence of various N-fold symmetric solutions with N > 2. In 1980s, Burbea [3] outlined a new way of constructing N-fold symmetric patch solutions by bifurcation theory. Following Burbea's framework, Hassainia and Hmidi [28] used the contour dynamics equation to give a rigorous proof of existence for simply-connected Nfold symmetric V-states with  $\alpha \in [0,1)$ , which are bifurcated from the unit disk. Their construction relies on the structure of linearization of contour dynamics equation, and the uniform angular velocity is selected appropriately such that the transversality assumption of Crandall-Rabinowitz's theorem is satisfied. By changing the function spaces from Hölder space to  $H^k$  space on torus, Castro et al. [9] extended this result for the remaining open cases  $\alpha \in [1,2)$  and showed the  $C^{\infty}$  regularity of patch boundaries for  $0 < \alpha < 2$ . In [16, 21, 22, 23], the existence of doubly connected V-states are established by studying the bifurcation from annulli with special angular velocity. However, there are two coupled nonlinear equations in this situation, and the estimate for the spectral of linearized operator is much more complicated. In [10, 32], 2-fold symmetric V-states bifurcated from Kirchhoff elliptic vortices were considered for  $\alpha = 0$ . Recently, [11, 12, 24] successfully applied the same method to construct travelling and rotating smooth solutions. Using a different method which is based on variational principle, Gravejat and Smets [27] and Godard-Cadillac [25] constructed smooth traveling solutions for the SQG and gSQG equations respectively.

It is well-known that the Euler equation and the gSQG equation exhibit point vortex solutions [43]. Of special interest are highly concentrated solutions which desingularize the point vortices. There are abundant literatures on the desingularized N-vortices problem for  $\alpha = 0$  and we refer to [4, 6, 7, 8, 19, 44, 45, 46] and references therein. As for  $0 < \alpha < 2$ , the results are relatively less. The first result was obtained by Hmidi and Mateu [33], where they showed the existence of concentrating travelling and co-rotating vortex pairs for  $0 \leq \alpha < 1$  using the contour dynamics equation and implicit function theorem. Recently, Ao et al. [1] gave the construction of concentrating smooth solutions for all  $\alpha \in (0, 2)$  by the Lyapunov-Schmidt reduction method. Godard-Cadillac et al. [26] constructed concentrating co-rotating *m*-fold symmetric vortex patches for  $\alpha \in (0, 2)$  via vorticity method.

It is noteworthy that all known concentrating solutions have supports composed of components which are perturbations of small disks. The aim of our paper is to construct co-rotating and travelling patch solutions for  $0 \le \alpha < 2$ , whose supports consist of doubly connected components perturbed from small annuli. Recall that a planar domain  $D_0$  is called doubly connected if  $D_0 = D_2 \setminus D_1$  for simply connected bounded domains  $D_1$  and  $D_2$  satisfying  $\overline{D}_1 \subset D_2$ . Our solutions provide a brand new phenomenon for the behavior of quasi-geostrophic flow.

The first kind of global solutions we are concerned with in this paper are co-rotating solutions. Suppose that the initial data  $\theta_{0,\varepsilon}$  is of N-fold symmetric patch type, that is

$$\theta_{0,\varepsilon}(x) = \frac{1}{\pi(1-b^2+\gamma b^2)\varepsilon^2} \sum_{n=0}^{N-1} \omega_n^{\varepsilon}, \qquad (1.2)$$

where  $\omega_0^{\varepsilon} = \boldsymbol{\chi}_{D_{out}^{\varepsilon}} + (\gamma - 1) \boldsymbol{\chi}_{D_{in}^{\varepsilon}}$  for some  $b \in (0, 1), \ \gamma \neq \frac{b^2 - 1}{b^2}$  and  $\omega_n^{\varepsilon}$  satisfy

 $\omega_n^{\varepsilon}(\boldsymbol{x}) = \omega_0^{\varepsilon}(Q_{\frac{2\pi n}{N}}(\boldsymbol{x} - d\boldsymbol{e}_1) + d\boldsymbol{e}_1), \quad \forall n = 1, 2, \dots N - 1$ 

with some d > 0 fixed,  $e_1 = (1,0)$  the unit vector in  $x_1$  direction and  $Q_\beta$  to denote the counterclockwise rotation operator of angle  $\beta$  with respect to the origin. In addition,  $D_{\text{out}}^{\varepsilon}$  and  $D_{\text{in}}^{\varepsilon}$  are simply connected domains satisfying  $\overline{D}_{\text{in}}^{\varepsilon} \subset D_{\text{out}}^{\varepsilon}$  and are perturbations of small disks  $\varepsilon B_1(0)$  and  $\varepsilon B_b(0)$  respectively. Notice that if  $\gamma = 0$ , then  $\omega_0^{\varepsilon} = \chi_{D_{\text{out}}^{\varepsilon} \setminus D_{\text{in}}^{\varepsilon}}$  and  $D_{\text{out}}^{\varepsilon} \setminus D_{\text{in}}^{\varepsilon}$  is a doubly connected domain. For  $\gamma \ll 0$ , an interesting phenomenon appears where  $\omega_1^{\varepsilon} < 0$  in  $D_{\text{out}} \setminus D_{\text{in}}$ , and  $\omega_1^{\varepsilon} > 0$  in  $D_{\text{in}}$ . The case  $\gamma = 1$  corresponds to simply connected cases, which have been studied as mentioned above. We intend to prove the existence of a series of co-rotating N-fold symmetric solutions to (1.1) about (d, 0), which take the form

$$\theta_{\varepsilon}(\boldsymbol{x},t) = \theta_{0,\varepsilon} \left( Q_{\Omega t}(\boldsymbol{x} - d\boldsymbol{e}_1) + d\boldsymbol{e}_1 \right)$$

with  $\Omega$  some fixed angular velocity.

As mentioned above, Hmidi and Mateu [33] obtained the existence of concentrating travelling and co-rotating vortex pairs with simply connected components for  $0 \le \alpha < 1$ using the contour dynamics equation and implicit function theorem. To sketch their proof, we denote  $G(\varepsilon, \Omega, f) = 0$  as the contour dynamics equation that the patch boundary satisfies, where f parameterizes the boundary. The first step is to extend  $G(\varepsilon, \Omega, f)$  such that  $\varepsilon \le 0$  is allowed. One can verify the case  $\varepsilon = 0$ , f = 0 corresponds to the point vortices and hence  $G(0, \Omega, 0) = 0$  holds for some special  $\Omega = \Omega_{sing}$ . To apply the implicit function theorem, the Gautex derivative  $\partial_f G(0, \Omega, 0) = 0$  should be an isomorphism, which only holds true for the space where the first Fourier coefficient vanishes. Consequently, the condition  $\int G(\varepsilon, \Omega, f) \sin(x) dx = 0$  is needed, which can be achieved by choosing proper  $\Omega(\varepsilon, f) = \Omega_{sing} + O(\varepsilon)$ . Then the existence of solutions follows from the implicit function theorem.

However, for the doubly connected components case, there are two coupled equations  $G_1(\varepsilon, \Omega, f_1, f_2) = 0$  and  $G_2(\varepsilon, \Omega, f_1, f_2) = 0$  to be solved. It seems impossible to choose suitable  $\Omega(\varepsilon)$  such that  $\int G_1(\varepsilon, \Omega, f_1, f_2) \sin(x) dx = 0$  and  $\int G_2(\varepsilon, \Omega, f_1, f_2) \sin(x) dx = 0$  hold simultaneously. Fortunately, we find that letting

$$\int G_1(\varepsilon, \Omega, f_1, f_2) \sin(x) dx = (1 - \gamma)b^2 \int G_2(\varepsilon, \Omega, f_1, f_2) \sin(x) dx$$

is enough for our purpose if we take appropriate function spaces. By choosing  $\Omega$  such that this assumption is satisfied, we successfully apply implicit function theorem to prove the following result of existence for the Euler equation.

**Theorem 1.1.** Suppose  $\alpha = 0$ ,  $N \geq 2$  and  $\gamma \in \mathbb{R} \setminus \{0\}$ . Then there exist  $\varepsilon_0 > 0$  and  $b \in (0,1)$ , such that for any  $\varepsilon \in (0,\varepsilon_0)$ , equation (1.1) has a global co-rotating solution  $\theta_{\varepsilon}(\boldsymbol{x} - d\boldsymbol{e}_1, t) = \theta_{0,\varepsilon}(Q_{\Omega^0 t}(\boldsymbol{x} - d\boldsymbol{e}_1))$  centered at (d, 0), where  $\theta_{0,\varepsilon}$  is of the form (1.2), and  $\Omega^0$  satisfies

$$\Omega^0 = \Omega^0_* + O(\varepsilon)$$

with  $\Omega^0_*$  given by

$$\Omega^{0}_{*} := \sum_{n=1}^{N-1} \frac{1 - \cos(\frac{2\pi n}{N})}{2\pi \left( \left( -1 + \cos(\frac{2\pi n}{N}) \right)^{2} + \sin^{2}(\frac{2\pi n}{N}) \right) d^{2}}.$$
(1.3)

Moreover, it holds in the sense of measures

$$\theta_{0,\varepsilon}(x) \to \sum_{n=0}^{N-1} \delta_{Q_{\frac{2n\pi}{N}}(d,0)} \quad as \quad \varepsilon \to 0^+.$$

Remark 1.2. If we take  $\mathcal{B}$  as the set of all b such that Theorem 1.1 holds true. Then  $(0,1) \setminus \mathcal{B}$  composed of at most countable many points. Moreover, for  $\gamma \in (0,1)$  it holds  $(0,\frac{\sqrt{2}}{2}) \subset \mathcal{B}$  and if  $\gamma \geq 1$  then  $\mathcal{B} = (0,1)$ . Theorem 1.1 does not cover the case  $\gamma = 0$  due to the fail of the invertibility of derivative matrices, see Lemma 3.5 for details. However, since the bifurcation method does not rely on this invertibility, in [23] the authors let  $\gamma = 0$  and constructed doubly connected vorticity to the Euler equation.

We also construct co-rotating solutions of the form (1.2) for the gSQG equation. Since the linearized operator has better property than the Euler case ( $\alpha = 0$ ), we can take  $\gamma = 0$  here, which means that the solutions constructed have supports composed of doubly connected components.

**Theorem 1.3.** Suppose  $\alpha \in (0,2)$ ,  $N \geq 2$  and  $\gamma \in [0,1)$ . Then there exist  $\varepsilon_0 > 0$  and  $b_0 \in (0,1)$  such that for any  $\varepsilon \in (0,\varepsilon_0)$  and  $b \in (0,b_0)$ , equation (1.1) has a global corotating solution  $\theta_{\varepsilon}(\boldsymbol{x} - d\boldsymbol{e}_1, t) = \theta_{0,\varepsilon}(Q_{\Omega^{\alpha}t}(\boldsymbol{x} - d\boldsymbol{e}_1))$  centered at (d,0), where  $\theta_{0,\varepsilon}$  is of the form (1.2), and  $\Omega^{\alpha}$  satisfies

$$\Omega^{\alpha} = \Omega^{\alpha}_* + O(\varepsilon^{1+\alpha})$$

with  $\Omega^{\alpha}_{*}$  given by

$$\Omega_*^{\alpha} := \sum_{n=1}^{N-1} \frac{\alpha C_{\alpha} (1 - \cos(\frac{2\pi n}{N}))}{2\pi \left( (-1 + \cos(\frac{2\pi n}{N}))^2 + \sin^2(\frac{2\pi n}{N}) \right)^{1+\frac{\alpha}{2}} d^{2+\alpha}}, \quad 0 < \alpha < 2.$$
(1.4)

Moreover, it holds in the sense of measures

$$\theta_{0,\varepsilon}(x) \to \sum_{n=0}^{N-1} \delta_{Q_{\frac{2n\pi}{N}}(d,0)} \quad as \quad \varepsilon \to 0^+.$$

Our next result concerns the travelling patch pairs. In this situation, the initial data  $\theta_{0,\varepsilon}$  is given by

$$\theta_{0,\varepsilon}(\boldsymbol{x}) = \frac{1}{\pi(1 - b^2 + \gamma b^2)\varepsilon^2} (\omega_0^{\varepsilon} - \omega_1^{\varepsilon}).$$
(1.5)

where  $\omega_0^{\varepsilon} = \chi_{D_{\text{out}}^{\varepsilon}} + (\gamma - 1)\chi_{D_{\text{in}}^{\varepsilon}}$  for some  $\gamma \in [0, 1]$ ,  $b \in (0, 1)$  and  $\omega_1^{\varepsilon}$  satisfies

$$\omega_1^{\varepsilon}(x_1, x_2) = \omega_0^{\varepsilon}(-x_1 + 2d, x_2)$$

with some d > 0 fixed, and  $e_1$  the unit vector in  $x_1$  direction. Furthermore,  $D_{\text{out}}^{\varepsilon}$  and  $D_{\text{in}}^{\varepsilon}$  are simply connected domains satisfying  $\overline{D}_{\text{in}}^{\varepsilon} \subset D_{\text{out}}^{\varepsilon}$  and are perturbations of the small disks  $\varepsilon B_1(0)$  and  $\varepsilon B_b(0)$  respectively. A travelling patch pair centered at (d, 0) takes the form

$$\theta_{\varepsilon}(x,t) = \theta_{0,\varepsilon}(x - tW\boldsymbol{e}_2)$$

with W some fixed speed, and  $e_2$  the unit vector in  $x_2$  direction. Using the same method as the proof of Theorems 1.1 and 1.3, we obtain the following result.

**Theorem 1.4.** Suppose  $\alpha \in [0,2)$ ,  $\gamma \in \mathbb{R} \setminus \{0\}$  if  $\alpha = 0$  and  $\gamma \in [0,1)$  if  $\alpha \in (0,2)$ . Then there exist  $\varepsilon_0 > 0$  and  $b \in (0,1)$  such that for any  $\varepsilon \in (0,\varepsilon_0)$ , equation (1.1) has a global travelling solution  $\theta_{\varepsilon}(\boldsymbol{x},t) = \theta_{0,\varepsilon}(\boldsymbol{x} - tW^{\alpha}\boldsymbol{e}_2)$ , where  $\theta_{0,\varepsilon}$  is of the form (1.5), and  $W^{\alpha}$ satisfies

$$W^{\alpha} = W^{\alpha}_* + O(|\varepsilon|^{1+\alpha})$$

with  $W^{\alpha}_{*}$  given by

$$W_*^{\alpha} := \begin{cases} \frac{1}{4\pi d}, & \alpha = 0, \\ \frac{\alpha C_{\alpha}}{2\pi (2d)^{1+\alpha}}, & 0 < \alpha < 2. \end{cases}$$
(1.6)

Moreover, it holds in the sense of measures

$$\theta_{0,\varepsilon}(x) \to \delta_{(0,0)} - \delta_{(2d,0)} \quad as \quad \varepsilon \to 0^+.$$

Remark 1.5. If we take  $\mathcal{B}$  as the set of all b such that Theorem 1.4 holds true. Then for  $\alpha = 0, (0, 1) \setminus \mathcal{B}$  composed of at most countable many points. Moreover, for  $\gamma \in (0, 1)$  it holds  $(0, \frac{\sqrt{2}}{2}) \subset \mathcal{B}$  and if  $\gamma \geq 1$  then  $\mathcal{B} = (0, 1)$ . For  $\alpha \in (0, 2)$ , there exists  $b_0 \in (0, 1)$  such that  $(0, b_0) \subset \mathcal{B}$ .

Remark 1.6. Calculating the signed curvature as the Corollary at the end of [30] and Proposition 4.12 in [9], one can prove that the doubly connected domain  $D_0^{\varepsilon}$  in Theorem 1.1, 1.3 or 1.4 is the difference of two convex sets. We also expect that the regularity of the boundaries can be improved to be analytic using related spaces defined in [10]. Some other interesting prosperities for doubly connected vortices with single component have been investigated in [31].

This paper is organized as follows. In section 2, we derive the contour dynamics equation and define the function spaces used later. In Section 3, we show the existence of N-fold symmetric co-rotating patch solutions for  $\alpha = 0$ . Section 4 is devoted to the existence of N-fold symmetric co-rotating patch solutions with doubly connected components for  $0 < \alpha < 2$ . In Section 5, we sketch the proof of existence of travelling patch pair solutions with doubly connected components.

### 2. The contour dynamics equation and function spaces

The main tool used in this paper is the contour dynamics equation for  $\alpha$ -patch. An  $\alpha$ -patch is a solution of (1.1) whose initial data is given by  $\theta_0 = \kappa \chi_D$ , with  $\kappa \in \mathbb{R}$ ,  $D \subset \mathbb{R}^2$  a bounded domain and  $\chi_D$  its characteristic function. Due to the transport formula  $\partial_t \theta + \mathbf{u} \cdot \nabla \theta = 0$ , the solution will preserve its patch structure and can be written as  $\theta_t = \kappa \chi_{D_t}$ . When  $0 \leq \alpha < 1$ , by using Biot-Savart law and Green-Stokes formula, the velocity can be recovered by

$$\boldsymbol{u}(\boldsymbol{x},t) = \begin{cases} \frac{\kappa}{2\pi} \int_{\partial D_t} \log\left(\frac{1}{|\boldsymbol{x}-\boldsymbol{y}|}\right) d\boldsymbol{y}, & \alpha = 0, \\\\ \frac{\kappa C_{\alpha}}{2\pi} \int_{\partial D_t} \frac{1}{|\boldsymbol{x}-\boldsymbol{y}|^{\alpha}} d\boldsymbol{y}, & 0 < \alpha < 1. \end{cases}$$

Thus if the patch boundary  $\partial D_t$  is parameterized as  $\boldsymbol{z}(t,\sigma)$  with  $\sigma \in [0, 2\pi)$ , then  $\boldsymbol{z}(t,\sigma)$  satisfies

$$\partial_t \boldsymbol{z}(t,\sigma) = \begin{cases} \frac{\kappa}{2\pi} \int_0^{2\pi} \log\left(\frac{1}{|\boldsymbol{z}(t,\sigma) - \boldsymbol{z}(t,\tau)|}\right) \partial_\tau \boldsymbol{z}(t,\tau) d\tau, & \alpha = 0, \\ \\ \frac{\kappa C_\alpha}{2\pi} \int_0^{2\pi} \frac{\partial_\tau \boldsymbol{z}(t,\tau)}{|\boldsymbol{z}(t,\sigma) - \boldsymbol{z}(t,\tau)|^\alpha} d\tau & 0 < \alpha < 1, \end{cases}$$

which is known as the contour dynamics equation. This equation is locally well-posed if the boundary of the initial  $\alpha$ -patch is composed with finite number of disjoint smooth Jordan curves. For  $1 \leq \alpha < 2$  the above integral is divergent. To eliminate the singularity, one can change the velocity at the boundary by subtracting a tangential vector and define

$$\partial_t \boldsymbol{z}(t,\sigma) = \frac{\kappa C_\alpha}{2\pi} \int_0^{2\pi} \frac{\partial_\tau \boldsymbol{z}(t,\tau) - \partial_\sigma \boldsymbol{z}(t,\sigma)}{|\boldsymbol{z}(t,\sigma) - \boldsymbol{z}(t,\tau)|^\alpha} d\tau.$$

The interfaces of the  $\alpha$ -patch under our consideration can be parameterized as two  $2\pi$  periodic curves  $\boldsymbol{z}_i(x)$  with i = 1, 2 and  $x \in [0, 2\pi)$ . For 0 < b < 1, we assume that  $\boldsymbol{z}_i$  are of the following form:

$$\boldsymbol{z}_i(x,t) = \left(\varepsilon R_i(x,t)\cos(x), \varepsilon R_i(x,t)\sin(x)\right),\,$$

with

$$R_1(x,t) = 1 + \varepsilon^{1+\alpha} f_1(x,t),$$
  

$$R_2(x,t) = b + \varepsilon^{1+\alpha} f_2(x,t).$$

For simplicity, let

$$\int g(\tau)d\tau := \frac{1}{2\pi} \int_0^{2\pi} g(\tau)d\tau$$

be the mean value of integral on the unit circle and denote  $\mathbf{f} = (f_1, f_2)$ .

For co-rotating patches

$$\theta_{\varepsilon}(\boldsymbol{x},t) = \theta_{0,\varepsilon} \left( Q_{\Omega t}(\boldsymbol{x} - d\boldsymbol{e}_1) + d\boldsymbol{e}_1 \right), \quad \boldsymbol{x} \in \mathbb{R}^2$$

with  $\Omega$  some uniform angular velocity. Inserting this equality into (1.1), we derive

$$\left(\mathbf{u}_0(\boldsymbol{x}) + \Omega(\boldsymbol{x} - d\boldsymbol{e}_1)^{\perp}\right) \cdot \nabla \theta_{0,\varepsilon}(\boldsymbol{x}) = 0.$$

Then we use the patch structure to obtain

$$\left(\mathbf{u}_0(\boldsymbol{x}) + \Omega(\boldsymbol{x} - d\boldsymbol{e}_1)^{\perp}\right) \cdot \mathbf{n}(\boldsymbol{x}) = 0, \quad \forall \, \boldsymbol{x} \in \bigcup_{n=0}^{N-1} \partial D_n^{\varepsilon},$$

where  $D_n^{\varepsilon}$  is the rotation of annuler  $D_{out}^{\varepsilon} \setminus \overline{D}_{in}^{\varepsilon}$  and  $\mathbf{n}(\boldsymbol{x})$  is the normal vector to the boundary. According to Biot-Savart law and Green-Stokes formula, the interfaces satisfies the following equation for  $\alpha = 0$ 

$$\begin{aligned} 0 &= G_{i}^{0}(\varepsilon,\Omega,\mathbf{f}) \end{aligned} \tag{2.1} \\ &= \pi (1-b^{2}+\gamma b^{2})\Omega \left(\varepsilon R_{i}'(x) - \frac{dR_{i}'(x)\cos(x)}{R_{i}(x)} + d\sin(x)\right) \\ &+ \frac{1}{2R_{i}(x)\varepsilon} \int \log \left(\frac{1}{(R_{i}(x)-R_{1}(y))^{2} + 4R_{i}(x)R_{1}(y)\sin^{2}\left(\frac{x-y}{2}\right)}\right) \\ &\times \left[(R_{i}(x)R_{1}(y) + R_{i}'(x)R_{1}'(y))\sin(x-y) + (R_{i}(x)R_{1}'(y) - R_{i}'(x)R_{1}(y))\cos(x-y)\right]dy \\ &+ \frac{\gamma-1}{2R_{i}(x)\varepsilon} \int \log \left(\frac{1}{(R_{i}(x)-R_{2}(y))^{2} + 4R_{i}(x)R_{2}(y)\sin^{2}\left(\frac{x-y}{2}\right)}\right) \\ &\times \left[(R_{i}(x)R_{2}(y) + R_{i}'(x)R_{2}'(y))\sin(x-y) + (R_{i}(x)R_{2}'(y) - R_{i}'(x)R_{2}(y))\cos(x-y)\right]dy \\ &+ \sum_{n=1}^{N-1} \frac{1}{R_{i}(x)\varepsilon} \int \log \left(\frac{1}{\left|\mathbf{z}_{i}(x) - d\mathbf{e}_{1} - Q_{\frac{2\pi n}{N}}(\mathbf{z}_{1}(y) - d\mathbf{e}_{1})\right|}\right) \\ &\times \left[(R_{i}(x)R_{1}(y) + R_{i}'(x)R_{1}'(y))\cos(x-y - \frac{2\pi n}{N}) \\ &+ (R_{i}(x)R_{1}'(y) - R_{i}'(x)R_{1}(y))\cos(x-y - \frac{2\pi n}{N})\right]dy \\ &+ \sum_{n=1}^{N-1} \frac{\gamma-1}{R_{i}(x)\varepsilon} \int \log \left(\frac{1}{\left|\mathbf{z}_{i}(x) - d\mathbf{e}_{1} - Q_{\frac{2\pi n}{N}}(\mathbf{z}_{2}(y) - d\mathbf{e}_{1})\right|}\right) \end{aligned}$$

$$\times \left[ (R_i(x)R_2(y) + R'_i(x)R'_2(y))\sin(x - y - \frac{2\pi n}{N}) + (R_i(x)R'_2(y) - R'_i(x)R_2(y))\cos(x - y - \frac{2\pi n}{N}) \right] dy$$
  
=  $G_{i1} + G_{i2} + G_{i3} + G_{i4} + G_{i5}.$ 

When  $0 < \alpha < 2$ , the interfaces obey the following equation  $0 = G_i^{\alpha}(\varepsilon, \Omega, \mathbf{f})$ 

$$\begin{aligned} 0 &= G_{i}^{\alpha}(\varepsilon,\Omega,\mathbf{f}) \end{aligned} \tag{2.2} \\ &= \pi(1-b^{2}+\gamma b^{2})\Omega\left(\varepsilon R_{i}'(x) - \frac{dR_{i}'(x)\cos(x)}{R_{i}(x)} + d\sin(x)\right) \\ &+ \frac{C_{\alpha}}{R_{i}(x)\varepsilon|\varepsilon|^{\alpha}} \int \frac{1}{|(R_{i}(x) - R_{1}(y))^{2} + 4R_{i}(x)R_{1}(y)\sin^{2}\left(\frac{x-y}{2}\right)|^{\frac{\alpha}{2}}} \\ &\times \left[(R_{i}(x)R_{1}(y) + R_{i}'(x)R_{1}'(y))\sin(x-y) + (R_{i}(x)R_{1}'(y) - R_{i}'(x)R_{1}(y)\cos(x-y))\right] dy \\ &+ \frac{(\gamma-1)C_{\alpha}}{R_{i}(x)\varepsilon|\varepsilon|^{\alpha}} \int \frac{1}{|(R_{i}(x) - R_{2}(y))^{2} + 4R_{i}(x)R_{2}(y)\sin^{2}\left(\frac{x-y}{2}\right)|^{\frac{\alpha}{2}}} \\ &\times \left[(R_{i}(x)R_{2}(y) + R_{i}'(x)R_{2}'(y))\sin(x-y) + (R_{i}(x)R_{2}'(y) - R_{i}'(x)R_{2}(y))\cos(x-y)\right] dy \\ &+ \sum_{n=1}^{N-1} \frac{C_{\alpha}}{R_{i}(x)\varepsilon} \int \frac{1}{|\mathbf{z}_{i}(x) - d\mathbf{e}_{1} - Q_{\frac{2\pi n}{N}}(\mathbf{z}_{1}(y) - d\mathbf{e}_{1})|^{\alpha}} \\ &\times \left[(R_{i}(x)R_{1}(y) + R_{i}'(x)R_{1}(y))\cos(x-y - \frac{2\pi n}{N}) \\ &+ (R_{i}(x)R_{1}'(y) - R_{i}'(x)R_{1}(y))\cos(x-y - \frac{2\pi n}{N})\right] dy \\ &+ \sum_{n=1}^{N-1} \frac{(\gamma-1)C_{\alpha}}{R_{i}(x)\varepsilon} \int \frac{1}{|\mathbf{z}_{i}(x) - d\mathbf{e}_{1} - Q_{\frac{2\pi n}{N}}(\mathbf{z}_{2}(y) - d\mathbf{e}_{1})|^{\alpha}} \\ &\times \left[(R_{i}(x)R_{2}(y) + R_{i}'(x)R_{2}'(y))\sin(x-y - \frac{2\pi n}{N}) \\ &+ (R_{i}(x)R_{2}(y) - R_{i}'(x)R_{2}(y))\cos(x-y - \frac{2\pi n}{N})\right] dy \end{aligned}$$

For  $0 \leq \alpha < 1$ , the function spaces used in the current paper are

$$X^{k} = \left\{ g \in H^{k}, \ g(x) = \sum_{j=1}^{\infty} a_{j} \cos(jx) \right\},$$
$$X^{k}_{b} := \left\{ \left( f_{1}, f_{2} \right) \in X^{k} \times X^{k} \middle| \oint f_{1}(x) \cos(x) dx = (1 - \gamma) b^{2} \oint f_{2}(x) \cos(x) dx \right\},$$

$$Y^{k} = \left\{ g \in H^{k}, \ g(x) = \sum_{j=1}^{\infty} a_{j} \sin(jx) \right\},$$

and

$$Y_b^k := \left\{ (g_1, g_2) \in Y^k \times Y^k \middle| \oint g_1(x) \sin(x) dx = (1 - \gamma) b^2 \oint g_2(x) \sin(x) dx \right\}.$$

The following spaces are also in order for the case  $\alpha = 1$ 

$$X^{k+\log} = \left\{ g \in H^k, \ g(x) = \sum_{j=2}^{\infty} a_j \cos(jx), \ \left\| \int_0^{2\pi} \frac{\partial^k g(x-y) - \partial^k g(x)}{|\sin(\frac{y}{2})|} dy \right\|_{L^2} < \infty \right\},$$

and

$$X_{b}^{k+\log} := \left\{ (f_{1}, f_{2}) \in X^{k+\log} \times X^{k+\log} \middle| \oint f_{1}(x) \cos(x) dx = (1-\gamma)b^{2} \oint f_{2}(x) \cos(x) dx \right\}.$$

For  $1 < \alpha < 2$ , we will need the following fractional spaces

$$X^{k+\alpha-1} = \left\{ g \in H^k, \ g(x) = \sum_{j=2}^{\infty} a_j \cos(jx), \ \left\| \int_0^{2\pi} \frac{\partial^k g(x-y) - \partial^k g(x)}{|\sin(\frac{y}{2})|^{\alpha}} dy \right\|_{L^2} < \infty \right\},$$

and

$$X_{b}^{k+\alpha-1} := \left\{ \left(f_{1}, f_{2}\right) \in X^{k+\alpha-1} \times X^{k+\alpha-1} \middle| \oint f_{1}(x) \cos(x) dx = (1-\gamma)b^{2} \oint f_{2}(x) \cos(x) dx \right\}.$$

The norm for  $X^k$  and  $Y^k$  is the  $H^k$ -norm. For  $X^{k+\log}$  and  $X^{k+\alpha-1}$ , the norms are given by the sum of the  $H^k$ -norm and the integral in their definitions respectively. Note that for every  $\beta > 0$ , the embedding  $X^{k+\beta} \subset X^{k+\log} \subset X^k$  holds.

Throughout this paper, we always assume  $k \geq 3$ .

### 3. Existence of co-rotating solutions: the case $\alpha = 0$

We will split up our proof into several lemmas, which correspond to checking the hypotheses of the implicit function theorem. Set  $V := \{f \in X_b^k \mid ||g||_{H^k} < 1\}$ . In what follows,  $\varepsilon_0$  denotes a small positive number determined in the process of proof.

We first prove that  $G_i^0$  is continuous and can be extended to  $\varepsilon \leq 0$ .

**Lemma 3.1.** 
$$G_i^0(\varepsilon, \Omega, f): (-\varepsilon_0, \varepsilon_0) \times \mathbb{R} \times V \to Y^{k-1}$$
 is continuous for  $i = 1, 2$ .

*Proof.* We only prove the continuity of  $G_1^0$ .

For the first term  $G_{11}$ , it is easy to show that  $G_{11}: \left(-\frac{1}{2}, \frac{1}{2}\right) \times \mathbb{R} \times V \to Y^{k-1}$  is continuous by defining  $G_{11}(0, \Omega, \mathbf{f}) = \pi (1 - b^2 + \gamma b^2) \Omega d \sin(x)$ . And we can rewrite  $G_{11}$  as

$$G_{11} = \pi (1 - b^2 + \gamma b^2) \Omega \left( d \sin(x) + \varepsilon \mathcal{R}_{11}(\varepsilon, f_1) \right), \qquad (3.1)$$

where  $\mathcal{R}_{11}(\varepsilon, \Omega, f_1) : \left(-\frac{1}{2}, \frac{1}{2}\right) \times \mathbb{R} \times X^k \to Y^{k-1}$  is continuous.

Now, consider the term  $G_{12}$ . We will prove that  $\partial^l G_{12} \in L^2$  for  $l = 0, 1, \ldots, k-1$ . Since  $R_1(x) = 1 + \varepsilon f_1(x)$ , the possible singularity caused by  $\varepsilon = 0$  may occur only when we take zeroth order derivative of  $G_{12}$ . Thus, we first show that  $G_{12} \in L^2$ .

Firstly, we consider the terms in  $G_{12}$ ,

$$\begin{aligned} &\frac{1}{2R_1(x)\varepsilon} \oint \log\left(\frac{1}{(R_1(x) - R_1(y))^2 + 4R_1(x)R_1(y)\sin^2\left(\frac{x-y}{2}\right)}\right) \\ &\times [R'_1(x)R'_1(y)\sin(x-y) + (R_1(x)R'_1(y) - R'_1(x)R_1(y))\cos(x-y)]\,dy \\ &= \frac{1}{2R_1(x)} \oint \log\left(\frac{1}{(R_1(x) - R_1(y))^2 + 4R_1(x)R_1(y)\sin^2\left(\frac{x-y}{2}\right)}\right) \\ &\times [\varepsilon f'_1(x)f'_1(y)\sin(x-y) + (R_1(x)f'_1(y) - f'_1(x)R_1(y))\cos(x-y)]\,dy \\ &= \frac{1}{2} \oint \log\left(\frac{1}{(R_1(x) - R_1(y))^2 + 4R_1(x)R_1(y)\sin^2\left(\frac{x-y}{2}\right)}\right) (f'_1(y) - f'_1(x))\cos(x-y)dy \\ &+ \frac{\varepsilon}{2R_1(x)} \oint \log\left(\frac{1}{(R_1(x) - R_1(y))^2 + 4R_1(x)R_1(y)\sin^2\left(\frac{x-y}{2}\right)}\right) \\ &\times [f'_1(x)f'_1(y)\sin(x-y) + f'_1(x)(f_1(x) - f_1(y))\cos(x-y)]\,dy. \end{aligned}$$

Since  $c|x-y|^2 \leq \left| (R_1(x) - R_1(y))^2 + 4R_1(x)R_1(y)\sin^2\left(\frac{x-y}{2}\right) \right| \leq C|x-y|^2$  for some positive constants c and C, we deduce that the kernel

$$\log\left(\frac{1}{\left(R_{1}(x) - R_{1}(y)\right)^{2} + 4R_{1}(x)R_{1}(y)\sin^{2}\left(\frac{x-y}{2}\right)}\right) \in L^{1}.$$
(3.2)

Hence, these terms belong to  $L^2$ .

To prove that the remaining term in  $G_{12}$  belongs to  $L^2$ , we will use the following Taylor's formula:

$$\log\left(\frac{1}{A+B}\right) = \log\left(\frac{1}{A}\right) - \int_0^1 \frac{B}{A+tB} dt.$$
(3.3)

and

$$\frac{1}{(A+B)^{\beta}} = \frac{1}{A^{\beta}} - \beta \int_0^1 \frac{B}{(A+tB)^{1+\beta}} dt.$$
 (3.4)

Denote

$$A := 4\sin^2\left(\frac{x-y}{2}\right),\,$$

and

$$B := \varepsilon (f_1(x) - f_1(y))^2 + \sin^2 \left(\frac{x - y}{2}\right) \left(4f_1(x) + 4f_1(y) + \varepsilon f_1(x)f_1(y)\right).$$
re helde

Then, there holds

$$(R_1(x) - R_1(y))^2 + 4R_1(x)R_1(y)\sin^2\left(\frac{x-y}{2}\right) = A + \varepsilon B.$$

Using (3.3), (3.4) and the fact that  $\sin(\cdot)$  is an odd function, we have

$$\begin{aligned} \frac{1}{\varepsilon} \oint \log\left(\frac{1}{A+\varepsilon B}\right) \sin(x-y) dy &+ \int \log\left(\frac{1}{A+\varepsilon B}\right) f_1(y) \sin(x-y) dy \\ &= \frac{1}{\varepsilon} \int \log\left(\frac{1}{A}\right) \sin(x-y) dy - \frac{1}{\varepsilon} \oint \int_0^1 \frac{\varepsilon B}{A+t\varepsilon B} dt \sin(x-y) dy \\ &+ \int \log\left(\frac{1}{A}\right) f(y) \sin(x-y) dy - \int \int_0^1 \frac{\varepsilon B}{A+t\varepsilon B} dt f_1(y) \sin(x-y) dy \\ &= -\int \int_0^1 \frac{B}{A+t\varepsilon B} dt \sin(x-y) dy + \int \log\left(\frac{1}{A}\right) f_1(y) \sin(x-y) dy \\ &- \varepsilon \oint \int_0^1 \frac{B}{A+t\varepsilon B} dt f(y) \sin(x-y) dy \\ &= -\int \int_0^1 \frac{B}{A} dt \sin(x-y) dy + \int \int_0^1 \int_0^1 \frac{t\varepsilon B^2}{(A+t\varepsilon B)^2} d\tau dt \sin(x-y) dy \\ &+ \int \log\left(\frac{1}{A}\right) f(y) \sin(x-y) dy - \varepsilon \oint \int_0^1 \frac{B}{A+t\varepsilon B} dt f_1(y) \sin(x-y) dy \\ &= -\int f_1(y) \sin(x-y) dy + \int \log\left(\frac{1}{A}\right) f_1(y) \sin(x-y) dy + \varepsilon \mathcal{R}_{121}(\varepsilon, f_1), \end{aligned}$$

where  $\mathcal{R}_{121}(\varepsilon, f_1)$  is not singular with respect to  $\varepsilon$  and belongs to  $L^2$  due to  $|B| \leq C|x-y|^2$ . In view of (3.2), it is easy to see  $\int \log(\frac{1}{A}) f_1(y) \sin(x-y) dy \in L^2$ , and hence  $G_{12} \in L^2$  for all  $\varepsilon \neq 0$ .

Applying (3.3) to  $\frac{1}{2} \oint \log \left( \frac{1}{(R_1(x) - R_1(y))^2 + 4R_1(x)R_1(y)\sin^2\left(\frac{x-y}{2}\right)} \right) (f_1'(y) - f_1'(x))\cos(x-y)dy$  we obtain

$$\frac{1}{2} \int \log\left(\frac{1}{(R_1(x) - R_1(y))^2 + 4R_1(x)R_1(y)\sin^2\left(\frac{x-y}{2}\right)}\right) (f_1'(y) - f_1'(x))\cos(x-y)dy$$

$$= \frac{1}{2} \int \log\left(\frac{1}{A + \varepsilon B}\right) (f_1'(y) - f_1'(x))\cos(x-y)dy$$

$$= \frac{1}{2} \int \log\left(\frac{1}{A}\right) (f_1'(y) - f_1'(x))\cos(x-y)dy$$

$$- \frac{1}{2} \int \int_0^1 \frac{\varepsilon B}{A + t\varepsilon B} dt (f_1'(y) - f_1'(x))\cos(x-y)dy$$

$$= \frac{1}{2} \int \log\left(\frac{1}{A}\right) (f_1'(y) - f_1'(x))\cos(x-y)dy + \varepsilon \mathcal{R}_{122}(\varepsilon, f_1).$$

Define

$$G_{12}(0, f_1) = \frac{1}{2} \int \log\left(\frac{1}{A}\right) \left(f_1(y)\sin(x-y) + \left(f_1'(y) - f_1'(x)\right)\cos(x-y)\right) dy$$

$$\begin{aligned} &-\frac{1}{2} \oint f_1(y) \sin(x-y) dy \\ &= \frac{1}{2} \oint \log\left(\frac{1}{4\sin^2\left(\frac{x-y}{2}\right)}\right) (f_1(y) \sin(x-y) + (f_1'(y) - f_1'(x)) \cos(x-y)) \, dy \\ &-\frac{1}{2} \oint f_1(y) \sin(x-y) dy \\ &= \frac{1}{2} \oint \log\left(\frac{1}{\sin^2\left(\frac{y}{2}\right)}\right) (f_1(x-y) \sin(y) + (f_1'(x-y) - f_1'(x)) \cos(y)) \, dy \\ &-\frac{1}{2} \oint f_1(y) \sin(x-y) dy. \end{aligned}$$

Then  $G_{12} \in L^2$  for all  $(\varepsilon, \mathbf{f}) \in \left(-\frac{1}{2}, \frac{1}{2}\right) \times V$ . We are going to show that  $\partial^{k-1}G_{12} \in L^2$ . Splitting the term  $R_1(x)R'_1(y) - R'_1(x)R_1(y)$ into  $R_1(x)(R'_1(y) - R'_1(x)) + R'_1(x)(R_1(x) - R_1(y))$  and taking (k-1)th derivatives of  $G_{12}$ , the most singular term is

$$\begin{aligned} &-\frac{\partial^{k-1}f_1(x)}{R_1(x)}\varepsilon G_{12} \\ &+ \int \log\left(\frac{1}{(R_1(x) - R_1(y))^2 + 4R_1(x)R_1(y)\sin^2\left(\frac{x-y}{2}\right)}\right) \\ &\times \left((\partial^k f(x)f'(y) + f'(x)\partial^k f(y)) + (\partial^k f(y) - \partial^k f(x))\cos(x-y)\right)dy \\ &- \int \frac{(f(x) - f(y))(\partial^{k-1}f(x) - \partial^{k-1}f(y)) + 2(f(x)\partial^{k-1}f(y) + \partial^{k-1}f(x)f(y))\sin^2\left(\frac{x-y}{2}\right)}{(R_1(x) - R_1(y))^2 + 4R_1(x)R_1(y)\sin^2\left(\frac{x-y}{2}\right)} \\ &\times \left[(R_1(x)R_1(y) + R_1'(x)R_1'(y))\sin(x-y) + (R_1(x)R_1'(y) - R_1'(x)R_1(y))\cos(x-y)\right]dy \\ &- \int \frac{(f(x) - f(y))(f'(x) - f'(y)) + 2(f(x)f'(y) + f'(x)f(y))\sin^2\left(\frac{x-y}{2}\right)}{(R_1(x) - R_1(y))^2 + 4R_1(x)R_1(y)\sin^2\left(\frac{x-y}{2}\right)} \\ &\times \left[\varepsilon(\partial^{k-1}f_1(x)R_1'(y) + f_1'(x)\partial^{k-1}f_1(y))\sin(x-y) \\ &+ (R_1(x)\partial^{k-1}f_1(y) - \partial^{k-1}f_1(x)R_1(y))\cos(x-y)\right]dy. \end{aligned}$$

Since  $f \in H^k$  and  $k \geq 3$ , we have  $\|\partial^l f\|_{L^{\infty}} < C\|f\|_{H^k}$  for l = 0, 1, 2. By mean value theorem, we can deduce that

$$\begin{aligned} |(f(x) - f(y))(\partial^{k-1}f(x) - \partial^{k-1}f(y)) + 2(f(x)\partial^{k-1}f(y) + \partial^{k-1}f(x)f(y))\sin^2(\frac{x-y}{2})| \\ &\leq C|x-y|(|\partial^{k-1}f(x)| + |\partial^{k-1}f(y)|), \end{aligned}$$

$$|(R_1(x)R_1(y) + R'_1(x)R'_1(y))\sin(x-y)| \le C|x-y|,$$

$$|(R_1(x)R'_1(y) - R'_1(x)R_1(y))\cos(x-y)| \le C|x-y|,$$

$$\left| (f(x) - f(y))(f'(x) - f'(y)) + 2(f(x)f'(y) + f'(x)f(y))\sin^2(\frac{x - y}{2}) \right| \le C|x - y|^2.$$

Therefore, by Hölder's inequality, we have  $\partial^{k-1}G_{12} \in L^2$ , which implies  $G_{12} \in H^{k-1}$ . Notice that  $f_1(x) \in V$  is an even function, and  $f'_1(x)$  is odd. By changing y to -y in  $G_{12}$ , we can deduce that  $G_{12}(\varepsilon, \mathbf{f})$  is odd, and hence  $G_{12} \in Y^{k-1}$ .

Next, we prove the continuity of  $G_{12}$ . By the definition of  $G_{12}(0, \mathbf{f})$  and above calculations, one can easily check that  $G_{12}$  is continuous with respect to  $\varepsilon$  at  $\varepsilon = 0$  and  $G_{12}(0, \mathbf{f})$ is continuous with respect to  $\mathbf{f}$ . Thus, we only need to prove that  $G_{12}$  is continuous for  $\varepsilon \neq 0$ . We deal with the most singular term

$$G_{122} = \frac{1}{2} \oint \log \left( \frac{1}{\left(R_1(x) - R_1(y)\right)^2 + 4R_1(x)R_1(y)\sin^2\left(\frac{x-y}{2}\right)} \right) (f_1'(y) - f_1'(x))\cos(x-y)dy$$
$$= \frac{1}{2} \oint \log \left( \frac{1}{\varepsilon^2 \left(f_1(x) - f_1(y)\right)^2 + 4(1+\varepsilon f_1(x))(1+f_1(y))\sin^2\left(\frac{x-y}{2}\right)} \right)$$
$$\times (f_1'(y) - f_1'(x))\cos(x-y)dy.$$

The following notations will be used for convenience. For a general function h, denote

$$\Delta g = g(x) - g(y), \quad g = g(x), \quad \tilde{g} = g(y),$$

and

$$D(h) = \varepsilon^2 (\Delta h)^2 + 4(1 + \varepsilon h)(1 + \varepsilon \tilde{h}) \sin^2(\frac{x - y}{2}).$$

Then for  $f_1, f_2 \in V$ , it holds

$$\begin{aligned} G_{122}(\varepsilon, f_1) &- G_{22}(\varepsilon, f_2) \\ &= -\frac{1}{2} \int \log\left(\frac{1}{D(f_1)}\right) \Delta f_1' \cos(x-y) dy + \frac{1}{2} \int \log\left(\frac{1}{D(f_2)}\right) \Delta f_2' \cos(x-y) dy \\ &= -\frac{1}{2} \int \log\left(\frac{1}{D(f_1)}\right) (\Delta f_1' - \Delta f_2') \cos(x-y) dy \\ &+ \frac{1}{2} \int \left(\log\left(\frac{1}{D(f_2)}\right) - \log\left(\frac{1}{D(f_1)}\right)\right) \Delta f_2' \cos(x-y) dy \\ &= I_1 + I_2. \end{aligned}$$

Similar to  $G_{12}$ , it is easy to show that  $||I_1||_{H^{k-1}} \leq C||f_1 - f_2||_{H^k}$ . To show  $||I_2||_{H^{k-1}} \leq C||f_1 - f_2||_{H^k}$ , it is enough to prove

$$\left\| \partial^l \left( \log \left( \frac{1}{D(f_2)} \right) - \log \left( \frac{1}{D(f_1)} \right) \right) \cdot |x - y| \right\|_{L^2} \le C \|f_1 - f_2\|_{H^k},$$

for l = 0, ..., k - 1, which can be verified by direct calculations. Indeed, we consider the most singular term

$$\begin{pmatrix} \frac{\partial^{k-1}D(f_1)}{D(f_1)} - \frac{\partial^{k-1}D(f_2)}{D(f_2)} \end{pmatrix} |x-y|$$
  
=  $\frac{\partial^{k-1}D(f_1) - \partial^{k-1}D(f_2)}{D(f_1)} |x-y| + \left(\frac{1}{D(f_1)} - \frac{1}{D(f_2)}\right) |x-y| \partial^{k-1}D(f_2)$   
=  $I_{21} + I_{22}.$ 

Obviously, it holds  $||I_{21}||_{H^{k-1}} \leq C ||f_1 - f_2||_{H^k}$ . Using mean value theorem, we have

$$I_{22} = \left(\frac{1}{D(f_1)} - \frac{1}{D(f_2)}\right) |x - y| \partial^{k-1} D(f_2)$$
  
=  $\frac{1}{\xi^2} (D(f_2) - D(f_1)) |x - y| \partial^{k-1} D(f_2)$ 

where  $\xi$  lies between  $D(f_1)$  and  $D(f_2)$  and hence  $c_1|x-y|^2 \leq \xi \leq c_2|x-y|^2$  for some constant  $c_1, c_2 > 0$ . Noticing that  $|D(f_2)-D(f_1)| \leq C|x-y|(|f_1(x)-f_2(x)|+|f_1(y)-f_2(y)|)$ , it follows that  $||I_{22}||_{H^{k-1}} \leq C||f_1-f_2||_{H^k}$ . Hence we have proved  $G_{12}(\varepsilon, \mathbf{f}) : (-\varepsilon_0, \varepsilon_0) \times V \to Y^{k-1}$  is continuous. And  $G_{12}$  takes the form

$$G_{12} = \frac{1}{2} \int \log\left(\frac{1}{\sin^2\left(\frac{y}{2}\right)}\right) (f_1(x-y)\sin(y) + (f_1'(x-y) - f_1'(x))\cos(y)) \, dy - \frac{1}{2} \int f_1(y)\sin(x-y) \, dy + \varepsilon \mathcal{R}_{12}(\varepsilon, \mathbf{f}),$$
(3.5)

where  $\mathcal{R}_{12}(\varepsilon, \mathbf{f}) : (-\varepsilon_0, \varepsilon_0) \times V \to Y^{k-1}$  is continuous by previous discussion.

Then we turn to consider  $G_{13}$ . Since  $(R_1(x) - R_2(y))^2 + 4R_1(x)R_2(y)\sin^2\left(\frac{x-y}{2}\right) = (1-b)^2 + O(\varepsilon)$ , the term  $G_{13}$  is more regular than  $G_{12}$  for  $\varepsilon$  small and can be proved to be bounded in  $Y^{k-1}$  and continuous as above. Applying the Taylor's formula (3.3) and (3.4) to  $G_{13}$ , one has

$$G_{13} = \frac{\gamma - 1}{2} \int \log\left(\frac{1}{(1 - b)^2 + 4b\sin^2\left(\frac{y}{2}\right)}\right) \left(f_2(x - y)\sin(y) + \left(f_2'(x - y) - bf_1'(x)\right)\cos(y)\right) dy + \frac{1 - \gamma}{2} \int \frac{b\sin(y)f_2(y)\left(-2(1 - b) + 4\sin^2\left(\frac{y}{2}\right)\right)}{(1 - b)^2 + 4b\sin^2\left(\frac{y}{2}\right)} dy + \varepsilon \mathcal{R}_{13}(\varepsilon, \mathbf{f}),$$
(3.6)

where  $\mathcal{R}_{13}(\varepsilon, f) : (-\varepsilon_0, \varepsilon_0) \times V \to Y^{k-1}$  is continuous.

Similarly, one has that the denominators of  $G_{14}$  and  $G_{15}$  are bounded from below by a positive constant for  $\varepsilon$  small. Hence  $G_{14}$  and  $G_{15}$  are more regular than  $G_{12}$  and can be proved to be bounded in  $Y^{k-1}$  and continuous more easily than  $G_{12}$ . Thus, the proof of continuity of  $G_1^0$  is complete. For later use, we compute the expansion of  $G_1^0$  at  $\varepsilon = 0$ .

Denote

$$A_n = \left(-1 + \cos\left(\frac{2\pi n}{N}\right)\right)^2 d^2 + \sin^2\left(\frac{2\pi n}{N}\right) d^2 > 0,$$
  

$$B_n = 2d(-1 + \cos\left(\frac{2\pi n}{N}\right))(\cos(x) - \cos(y + \frac{2\pi n}{N})) + 2d\sin\left(\frac{2\pi n}{N}\right)(\sin(x) - \sin(x + \frac{2\pi n}{N})),$$
  

$$\tilde{B}_n = 2d(-1 + \cos\left(\frac{2\pi n}{N}\right))(\cos(x) - b\cos(y + \frac{2\pi n}{N})) + 2d\sin\left(\frac{2\pi n}{N}\right)(\sin(x) - b\sin(x + \frac{2\pi n}{N})).$$
  
Then, we have

$$\left|\boldsymbol{z}_{1}(x) - d\boldsymbol{e}_{1} - Q_{\frac{2\pi n}{N}}\left(\boldsymbol{z}_{1}(y) - d\boldsymbol{e}_{1}\right)\right|^{2} = A_{n} + \varepsilon B_{n} + O(\varepsilon^{2}),$$

and

$$\left|\boldsymbol{z}_{1}(x) - d\boldsymbol{e}_{1} - Q_{\frac{2\pi n}{N}} \left(\boldsymbol{z}_{2}(y) - d\boldsymbol{e}_{1}\right)\right|^{2} = A_{n} + \varepsilon \tilde{B}_{n} + O(\varepsilon^{2}).$$

Applying the Taylor's formula (3.3) and (3.4), we have

$$\begin{aligned} G_{14} &= \sum_{n=1}^{N-1} \frac{1}{\varepsilon} \int \log \left( \frac{1}{|z_1(x) - de_1 - Q_{\frac{2\pi n}{N}}(z_1(y) - de_1)|} \right) \sin(x - y - \frac{2\pi n}{N}) dy \end{aligned} (3.7) \\ &+ \sum_{n=1}^{N-1} \int \log \left( \frac{1}{|(z_1(x) - de_1 - Q_{\frac{2\pi n}{N}}(z_1(y) - de_1))||} \right) \\ &\times \left( f_1(y) \sin(x - y - \frac{2\pi n}{N}) + (f_1'(y) - f_1'(x)) \cos(x - y - \frac{2\pi n}{N}) \right) dy + \varepsilon \mathcal{R}_{14} \\ &= \sum_{n=1}^{N-1} \frac{1}{2\varepsilon} \int \log \left( \frac{1}{A_n + \varepsilon B_n + O(\varepsilon^2)} \right) \sin(x - y - \frac{2\pi n}{N}) dy \\ &+ \sum_{n=1}^{N-1} \int \log \left( \frac{1}{A_n} \right) \left( f_1(y) \sin(x - y - \frac{2\pi n}{N}) + (f_1'(y) - f_1'(x)) \cos(x - y - \frac{2\pi n}{N}) \right) dy \\ &+ \varepsilon \mathcal{R}_{14} \\ &= \sum_{n=1}^{N-1} \frac{1}{2\varepsilon} \int \log \left( \frac{1}{A_n} \right) \sin(x - y - \frac{2\pi n}{N}) dy \\ &= \sum_{n=1}^{N-1} \frac{1}{2\varepsilon} \int \log \left( \frac{1}{A_n} \right) \sin(x - y - \frac{2\pi n}{N}) dy \\ &- \sum_{n=1}^{N-1} \frac{1}{2\varepsilon} \int \log \left( \frac{1}{A_n} \right) \sin(x - y - \frac{2\pi n}{N}) dy \\ &- \sum_{n=1}^{N-1} \frac{1}{2} \int \frac{B_n}{A_n + \varepsilon t B_n + tO(\varepsilon^2)} \sin(x - y - \frac{2\pi n}{N}) dy + \varepsilon \mathcal{R}_{14} \\ &= -\sum_{n=1}^{N-1} \frac{1}{2} \int \frac{B_n}{A_n} \sin(x - y - \frac{2\pi n}{N}) dy + \varepsilon \mathcal{R}_{14} \end{aligned}$$

$$= -\sum_{n=1}^{N-1} \frac{(1 - \cos(\frac{2\pi n}{N}))\sin(x)}{2d\left((-1 + \cos(\frac{2\pi n}{N}))^2 + \sin^2(\frac{2\pi n}{N})\right)} + \varepsilon \mathcal{R}_{14}(\varepsilon, \boldsymbol{f}),$$

where  $\mathcal{R}_{14}(\varepsilon, f)$  is regular and may be different from line to line.

Applying the Taylor's formula (3.3) and (3.4), we obtain

$$G_{15} = (3.8)$$

$$\sum_{n=1}^{N-1} \frac{\gamma - 1}{\varepsilon} \int \log\left(\frac{1}{\left|\boldsymbol{z}_{1}(x) - d\boldsymbol{e}_{1} - Q_{\frac{2\pi n}{N}}\left(\boldsymbol{z}_{2}(y) - d\boldsymbol{e}_{1}\right)\right|}\right) b\sin(x - y - \frac{2\pi n}{N}) dy \quad (3.9)$$

$$\begin{aligned} &+ \sum_{n=1}^{N-1} (\gamma - 1) \int \log \left( \frac{1}{|z_1(x) - de_1 - Q_{\frac{2\pi n}{N}}(z_2(y) - de_1)|} \right) \\ &\times \left( f_2(y) \sin(x - y - \frac{2\pi n}{N}) + (f_2'(y) - bf_1'(x)) \cos(x - y - \frac{2\pi n}{N}) \right) dy + \varepsilon \mathcal{R}_{15} \\ &= \sum_{n=1}^{N-1} \frac{(\gamma - 1)b}{2\varepsilon} \int \log \left( \frac{1}{A_n + \varepsilon \tilde{B}_n + O(\varepsilon^2)} \right) \sin(x - y - \frac{2\pi n}{N}) dy \\ &+ \sum_{n=1}^{N-1} (\gamma - 1) \int \log \left( \frac{1}{A_n} \right) \left( f_2(y) \sin(x - y - \frac{2\pi n}{N}) + (f_2'(y) - f_1'(x)) \cos(x - y - \frac{2\pi n}{N}) \right) dy \\ &+ \varepsilon \mathcal{R}_{15} \end{aligned}$$

$$= -\sum_{n=1}^{N-1} \frac{(\gamma - 1)b^2}{2} \int \frac{\pi}{A_n} \sin(x - y - \frac{\pi}{N}) dy + \varepsilon \mathcal{R}_{15}$$
$$= -\sum_{n=1}^{N-1} \frac{(\gamma - 1)b^2(1 - \cos(\frac{2\pi n}{N}))\sin(x)}{2d\left((-1 + \cos(\frac{2\pi n}{N}))^2 + \sin^2(\frac{2\pi n}{N})\right)} + \varepsilon \mathcal{R}_{15}(\varepsilon, \boldsymbol{f}),$$

where  $\mathcal{R}_{15}(\varepsilon, \Omega, f) : (-\varepsilon_0, \varepsilon_0) \times \mathbb{R} \times V \to Y^{k-1}$  is continuous and may be different from line to line.

Combining (3.1), (3.5), (3.6), (3.7) and (3.8) we conclude  $G_1^0 = \pi (1 - b^2 + \gamma b^2) \Omega d \sin(x)$ (3.10)

$$\begin{split} &+ \frac{1}{2} \int \log\left(\frac{1}{\sin^2\left(\frac{y}{2}\right)}\right) \left(f_1(x-y)\sin(y) + \left(f_1'(x-y) - f_1'(x)\right)\cos(y)\right) dy \\ &- \frac{1}{2} \int f_1(y)\sin(x-y)dy \\ &\frac{\gamma-1}{2} \int \log\left(\frac{1}{(1-b)^2 + 4b\sin^2\left(\frac{y}{2}\right)}\right) \left(f_2(x-y)\sin(y) + \left(f_2'(x-y) - bf_1'(x)\right)\cos(y)\right) dy \\ &+ \frac{1-\gamma}{2} \int \frac{b\sin(y)f_2(y)\left(-2(1-b) + 4\sin^2\left(\frac{y}{2}\right)\right)}{(1-b)^2 + 4b\sin^2\left(\frac{y}{2}\right)} \\ &- \sum_{n=1}^{N-1} \frac{(1-b^2 + \gamma b^2)(1-\cos(\frac{2\pi n}{N}))\sin(x)}{2d\left((-1+\cos(\frac{2\pi n}{N}))^2 + \sin^2(\frac{2\pi n}{N})\right)} + \varepsilon \mathcal{R}_1(\varepsilon, \Omega, \boldsymbol{f}), \end{split}$$

where  $\mathcal{R}_1(\varepsilon, \Omega, \mathbf{f}) : (-\varepsilon_0, \varepsilon_0) \times \mathbb{R} \times V \to Y^{k-1}$  is continuous. Similarly, we can prove that  $G_2^0(\varepsilon, \Omega, \mathbf{f}) : (-\varepsilon_0, \varepsilon_0) \times \mathbb{R} \times V \to Y^{k-1}$  is continuous and is of the form

$$\begin{aligned} G_{2}^{0} &= \pi (1 - b^{2} + \gamma b^{2}) \Omega d \sin(x) \end{aligned}$$

$$&+ \frac{1}{2b} \oint \log \left( \frac{1}{(1 - b)^{2} + 4b \sin^{2}\left(\frac{y}{2}\right)} \right) (bf_{1}(x - y) \sin(y) + (bf_{1}'(x - y) - f_{2}'(x)) \cos(y)) \, dy \\ &- \frac{1}{2} \int \frac{\sin(y) f_{1}(y) \left(2(1 - b) + 4b \sin^{2}\left(\frac{y}{2}\right)\right)}{(1 - b)^{2} + 4b \sin^{2}\left(\frac{y}{2}\right)} \\ &+ \frac{\gamma - 1}{2} \int \log \left( \frac{1}{\sin^{2}\left(\frac{y}{2}\right)} \right) (f_{2}(x - y) \sin(y) + (f_{2}'(x - y) - f_{2}'(x)) \cos(y)) \, dy \\ &+ \frac{1 - \gamma}{2} \int f_{2}(y) \sin(x - y) dy \\ &- \sum_{n=1}^{N-1} \frac{(1 - b^{2} - \gamma b^{2})(1 - \cos\left(\frac{2\pi n}{N}\right)) \sin(x)}{2d \left((-1 + \cos\left(\frac{2\pi n}{N}\right))^{2} + \sin^{2}\left(\frac{2\pi n}{N}\right)} + \varepsilon \mathcal{R}_{2}(\varepsilon, \Omega, \mathbf{f}), \end{aligned}$$

where  $\mathcal{R}_2(\varepsilon, \Omega, f): (-\varepsilon_0, \varepsilon_0) \times \mathbb{R} \times V \to Y^{k-1}$  is continuous. The proof of Lemma 3.1 is thus complete.

For  $(\varepsilon, \Omega, \mathbf{f}) \in (-\varepsilon_0, \varepsilon_0) \times \mathbb{R} \times V$  and  $h \in X^k$ , denote  $\partial_{f_i} G^0_i(\varepsilon, \Omega, f) h$  to be the Gateaux derivatives. We have following lemma:

**Lemma 3.2.** The Gautex derivative  $\partial_{f_i} G_j^0(\varepsilon, \Omega, f) : (-\varepsilon_0, \varepsilon_0) \times \mathbb{R} \times V \to L(X^k, Y^{k-1})$ exist and is continuous for i, j = 1, 2.

Proof. Set

$$B_{ijn} := 2\left(-1 + \cos\left(\frac{2\pi n}{N}\right)\right) d\left(R_i(x)\cos(x) - R_j(y)\cos\left(y + \frac{2\pi n}{N}\right)\right)$$

$$+ 2\sin\left(\frac{2\pi n}{N}\right) \left(R_i(x)\sin(x) - R_j(y)\sin\left(y + \frac{2\pi n}{N}\right)\right)$$
$$+ \varepsilon \left(R_i(x)^2 + R_j^2 + R_i(x)R_j(y)\cos\left(x - y - \frac{2\pi n}{N}\right)\right).$$

Then

$$\left|z_i(x) - d\boldsymbol{e}_1 - Q_{\frac{2\pi n}{N}}\left(z_j(y) - d\boldsymbol{e}_1\right)\right|^2 = A_n + \varepsilon B_{ijn}.$$

We are going to prove that the derivative of  $G_1^0$  with respect to  $f_1$  is as follows,

$$\partial_{f_1} G_1^0(\varepsilon, \Omega, \boldsymbol{f}) h = F_1 + F_2 + F_3 + F_4 + F_5,$$

where  $F_1, F_2, F_3, F_4, F_5$  are given by

$$F_{1} = \pi (1 - b^{2} + \gamma b^{2}) \Omega \left( \varepsilon^{2} h'(x) - \frac{d\varepsilon h'(x) \cos(x)}{R_{1}(x)} + \frac{d\varepsilon R_{1}'(x) h'(x) \cos(x)}{R_{1}(x)^{2}} \right),$$

$$F_2 =$$

$$\begin{split} &-\frac{h(x)}{2R_{1}(x)^{2}} \oint \log\left(\frac{1}{(R_{1}(x)-R_{1}(y))^{2}+4R_{1}(x)R_{1}(y)\sin^{2}\left(\frac{x-y}{2}\right)}\right) \\ &\times \left[(R_{1}(x)R_{1}(y)+R_{1}'(x)R_{1}'(y))\sin(x-y)+(R_{1}(x)R_{1}'(y)-R_{1}'(x)R_{1}(y))\cos(x-y)\right]dy \\ &+\frac{1}{2R_{1}(x)} \oint \log\left(\frac{1}{(R_{1}(x)-R_{1}(y))^{2}+4R_{1}(x)R_{1}(y)\sin^{2}\left(\frac{x-y}{2}\right)}\right) \\ &\times \left[(h(x)R_{1}(y)+R_{1}(x)h(y)+h'(x)R_{1}'(y)+R_{1}'(x)h'(y))\sin(x-y) \\ &+(h(x)R_{1}'(y)+R_{1}(x)h'(y)-h'(x)R_{1}(y)-R_{1}'(x)h(y))\cos(x-y)\right]dy \\ &-\frac{1}{2R_{1}(x)} \int \frac{(R_{1}(x)R_{1}(y)+R_{1}'(x)R_{1}'(y))\sin(x-y)+(R_{1}(x)R_{1}(y)-R_{1}'(x)R_{1}(y))\cos(x-y)}{(R_{1}(x)-R_{1}(y))^{2}+4R_{1}(x)R_{1}(y)\sin^{2}\left(\frac{x-y}{2}\right)} \\ &\times \left[2\left(R_{1}(x)-R_{1}(y)\right)(h(x)-h(y))+4(h(x)R_{1}(y)+R_{1}(x)h(y))\sin^{2}\left(\frac{x-y}{2}\right)\right]dy, \\ F_{3} &=\frac{(1-\gamma)h(x)}{2R_{1}(x)^{2}} \int \log\left(\frac{1}{(R_{1}(x)-R_{2}(y))^{2}+4R_{1}(x)R_{2}(y)\sin^{2}\left(\frac{x-y}{2}\right)}\right) \\ &\times \left[(R_{1}(x)R_{2}(y)+R_{1}'(x)R_{2}'(y))\sin(x-y)+(R_{1}(x)R_{2}(y)-R_{1}'(x)R_{2}(y))\cos(x-y)\right]dy \\ &+\frac{\gamma-1}{2R_{1}(x)} \int \log\left(\frac{1}{(R_{1}(x)-R_{2}(y))^{2}+4R_{1}(x)R_{2}(y)\sin^{2}\left(\frac{x-y}{2}\right)}\right) \end{split}$$

$$\times \left[ (h(x)R_{2}(y) + h'(x)R'_{2}(y))\sin(x-y) + (h(x)R'_{2}(y) - h'(x)R_{2}(y))\cos(x-y) \right] dy + \frac{1-\gamma}{2R_{1}(x)} \int \frac{\left[ R_{1}(x)R_{2}(y) + R'_{1}(x)R'_{2}(y) \right]\sin(x-y) + \left[ R_{1}(x)R'_{2}(y) - R'_{1}(x)R_{2}(y) \right]\cos(x-y)}{\left( R_{1}(x) - R_{2}(y) \right)^{2} + 4R_{1}(x)R_{2}(y)\sin^{2}\left(\frac{x-y}{2}\right)}$$

$$\times \left[2\left(R_1(x) - R_2(y)\right)h(x) + 4h(x)R_2(y)\sin^2\left(\frac{x-y}{2}\right)\right]dy,$$

$$\begin{split} F_4 &= \\ &- \sum_{n=1}^{N-1} \frac{h(x)}{R_1(x)^2} \int \log \left( \frac{1}{\left| \mathbf{z}_1(x) - d\mathbf{e}_1 - Q_{\frac{2\pi n}{N}} \left( \mathbf{z}_1(y) - d\mathbf{e}_1 \right) \right|} \right) \\ &\times \left[ (R_1(x)R_1(y) + R_1'(x)R_1'(y)) \sin(x - y - \frac{2\pi n}{N}) \right] \\ &+ (R_1(x)R_1'(y) - R_1'(x)R_1(y)) \cos(x - y - \frac{2\pi n}{N}) \right] dy \\ &+ \sum_{n=1}^{N-1} \frac{1}{R_1(x)} \int \log \left( \frac{1}{\left| \mathbf{z}_1(x) - d\mathbf{e}_1 - Q_{\frac{2\pi n}{N}} \left( \mathbf{z}_1(y) - d\mathbf{e}_1 \right) \right|} \right) \\ &\times \left[ (h(x)R_1(y) + R_1(x)h(y) + h'(x)R_1'(y) + R_1'(x)h'(y)) \sin(x - y - \frac{2\pi n}{N}) \right. \\ &+ (h(x)R_1'(y) + R_1(x)h'(y) - h'(x)R_1(y) - R_1'(x)h(y)) \cos(x - y - \frac{2\pi n}{N}) \right] dy \\ &- \sum_{n=1}^{N-1} \frac{1}{2R_1(x)} \int \frac{\partial_{f_1}B_{11n}h}{\left| \mathbf{z}_1(x) - d\mathbf{e}_1 - Q_{\frac{2\pi n}{N}} \left( \mathbf{z}_1(y) - d\mathbf{e}_1 \right) \right|^2} \left[ \left( R_1(x)R_1(y) + R_1'(x)R_1'(y) \right) \sin(x - y - \frac{2\pi n}{N}) \right. \\ &+ \left( R_1(x)R_1'(y) - R_1'(x)R_1(y) \right) \cos(x - y - \frac{2\pi n}{N}) \right] dy \end{split}$$

and

$$F_{5} = \sum_{n=1}^{N-1} \frac{(1-\gamma)h(x)}{R_{1}(x)^{2}} \int \log\left(\frac{1}{\left|\boldsymbol{z}_{1}(x) - d\boldsymbol{e}_{1} - Q_{\frac{2\pi n}{N}}\left(\boldsymbol{z}_{2}(y) - d\boldsymbol{e}_{1}\right)\right|}\right)$$

$$\left[\left(R_{1}(x)R_{2}(y) + R'_{1}(x)R'_{2}(y)\right)\sin(x-y-\frac{2\pi n}{N}) + \left(R_{1}(x)R'_{2}(y) - R'_{1}(x)R_{2}(y)\right)\cos(x-y-\frac{2\pi n}{N})\right]dy$$

$$+ \sum_{n=1}^{N-1} \frac{\gamma-1}{R_{1}(x)} \int \log\left(\frac{1}{\left|\boldsymbol{z}_{1}(x) - d\boldsymbol{e}_{1} - Q_{\frac{2\pi n}{N}}\left(\boldsymbol{z}_{2}(y) - d\boldsymbol{e}_{1}\right)\right|}\right)$$

$$\times \left[\left(h(x)R_{2}(y) + h'(x)R'_{2}(y)\right)\sin(x-y-\frac{2\pi n}{N}) + \left(h(x)R'_{2}(y) - h'(x)R_{2}(y)\right)\cos(x-y-\frac{2\pi n}{N})\right]dy$$

$$+\sum_{n=1}^{N-1} \frac{1-\gamma}{2R_1(x)} \oint \frac{\partial_{f_1} B_{12n} h}{\left| \boldsymbol{z}_1(x) - d\boldsymbol{e}_1 - Q_{\frac{2\pi n}{N}} \left( \boldsymbol{z}_2(y) - d\boldsymbol{e}_1 \right) \right|^2} \left[ \left( R_1(x) R_2(y) + R_1'(x) R_2'(y) \right) \sin(x-y-\frac{2\pi n}{N}) + \left( R_1(x) R_2'(y) - R_1'(x) R_2(y) \right) \cos(x-y-\frac{2\pi n}{N}) \right] dy.$$

To this end, we need to show for l = 1, 2, 3, 4, 5,

$$\lim_{t \to 0} \left\| \frac{G_{1l}^0(\varepsilon, \Omega, f_1 + th, f_2) - G_{1l}^0(\varepsilon, \Omega, f_1, f_2)}{t} - F_l(\varepsilon, \Omega, f)(\varepsilon, \boldsymbol{f}, h) \right\|_{Y^{k-1}} = 0$$

The derivative of  $G_{11}^0$  is obvious. Next, we only consider the most singular term

$$\begin{aligned} G_{12} = &\frac{1}{2R_1(x)\varepsilon} \oint \log\left(\frac{1}{(R_1(x) - R_1(y))^2 + 4R_1(x)R_1(y)\sin^2\left(\frac{x-y}{2}\right)}\right) \\ &\times \left[(R_1(x)R_1(y) + R_1'(x)R_1'(y))\sin(x-y) + (R_1(x)R_1'(y) - R_1'(x)R_1(y))\cos(x-y)\right]dy \\ &= &\frac{1}{2R_1(x)\varepsilon} \oint \log\left(\frac{1}{(R_1(x) - R_1(y))^2 + 4R_1(x)R_1(y)\sin^2\left(\frac{x-y}{2}\right)}\right) \\ &\times (R_1(x)R_1(y) + R_1'(x)R_1'(y))\sin(x-y)dy \\ &+ &\frac{1}{2} \oint \log\left(\frac{1}{(R_1(x) - R_1(y))^2 + 4R_1(x)R_1(y)\sin^2\left(\frac{x-y}{2}\right)}\right) (f_1'(y) - f_1'(x))\cos(x-y)dy \\ &+ &\frac{f_1'(x)}{2R_1(x)} \oint \log\left(\frac{1}{(R_1(x) - R_1(y))^2 + 4R_1(x)R_1(y)\sin^2\left(\frac{x-y}{2}\right)}\right) (f_1(x) - f_1(y))\cos(x-y)dy \\ &= &G_{121} + G_{122} + G_{123}. \end{aligned}$$

The proof for other terms is much easier, so we omit it.

Recall that  $R_1 = 1 + \varepsilon f_1$ , thus the  $\varepsilon$  in the denominator of  $G_{12}$  will be eliminated after taking derivatives. Thus, the  $\varepsilon$  in  $G_{12}$  causes no singularity. For simplicity, we consider the most singular term in  $G_{12}$ 

$$G_{122} = \frac{1}{2} \oint \log\left(\frac{1}{\left(R_1(x) - R_1(y)\right)^2 + 4R_1(x)R_1(y)\sin^2\left(\frac{x-y}{2}\right)}\right) \left(f_1'(y) - f_1'(x)\right)\cos(x-y)dy.$$

We need to prove

$$\lim_{t \to 0} \left\| \frac{G_{122}(\varepsilon, f_1 + th) - G_{122}(\varepsilon, f_1)}{t} - F_{22}(\varepsilon, f)(\varepsilon, f, h) \right\|_{Y^{k-1}} = 0,$$
(3.12)

where  $F_{22}$  is

$$F_{22} = \frac{1}{2} \int \log\left(\frac{1}{\left(R_1(x) - R_1(y)\right)^2 + 4R_1(x)R_1(y)\sin^2\left(\frac{x-y}{2}\right)}\right) \left(h'(y) - h'(x)\right)\cos(x-y)dy$$

$$-\frac{1}{2} \int \frac{(f_1'(y) - f_1'(x))\cos(x - y)}{(R_1(x) - R_1(y))^2 + 4R_1(x)R_1(y)\sin^2\left(\frac{x - y}{2}\right)} \times \left[2\varepsilon \left(R_1(x) - R_1(y)\right)(h(x) - h(y)) + 4\varepsilon(h(x)R_1(y) + R_1(x)h(y))\sin^2\left(\frac{x - y}{2}\right)\right] dy.$$

Using the notations given in Lemma 3.1, we deduce

$$\begin{aligned} \frac{G_{122}(\varepsilon, f_1 + th) - G_{122}(\varepsilon, f_1)}{t} &- F_{22}(\varepsilon, f_1, h) \\ &= \frac{1}{t} \int (f_1'(x) - f_1'(y)) \cos(x - y) \\ &\times \left( \log\left(\frac{1}{D(f_1 + th)}\right) - \log\left(\frac{1}{D(f_1)}\right) + t\frac{2\varepsilon^2 \Delta f_1 \Delta h + 4\varepsilon(\tilde{R}_1 h + \tilde{h}R_1) \sin^2(\frac{x - y}{2})}{D(f_1)} \right) dy \\ &+ \int (h'(x) - h'(y)) \cos(x - y) \left( \log\left(\frac{1}{D(f_1 + th)}\right) - \left(\frac{1}{D(f_1)}\right) \right) dy \\ &= F_{221} + F_{222}. \end{aligned}$$

By taking k - 1th partial derivatives of  $F_{221}$ , we find the most singular term is

$$\frac{1}{t} \oint \left( \log\left(\frac{1}{D(f_1+th)}\right) - \log\left(\frac{1}{D(f_1)}\right) + t \frac{2\varepsilon^2 \Delta f_1 \Delta h + 4\varepsilon (\tilde{R}_1 h + \tilde{h}R_1) \sin^2(\frac{x-y}{2})}{D(f_1)} \right) \times (\partial^k f(x) - \partial^k f(y)) \cos(x-y) dy.$$

Using the mean value theorem, we have

$$\left|\log\left(\frac{1}{D(f_1+th)}\right) - \log\left(\frac{1}{D(f_1)}\right) + t\frac{2\varepsilon^2\Delta f_1\Delta h + 4\varepsilon(\tilde{R}_1h + \tilde{h}R_1)\sin^2(\frac{x-y}{2})}{D_1(f)}\right| \le Ct^2\zeta(\varepsilon, f, h),$$

where  $\|\zeta(\varepsilon, f, h)\|_{L^{\infty}} < \infty$ . It follows that

$$\|F_{221}\|_{Y^{k-1}} \le Ct^2 \left\| \int \partial^k f(x) - \partial^k f(y) dy \right\|_{L^2} \le Ct^2 \|f\|_{X^k}.$$

Taking k - 1th partial derivatives of  $F_{222}$ , the most singular term is

$$\int (\partial^k h(x) - \partial^k h(y)) \cos(x - y) \left( \log \left( \frac{1}{D(f_1 + th)} \right) - \left( \frac{1}{D(f_1)} \right) \right) dy.$$

Using the mean value theorem, we have

$$\log\left(\frac{1}{D(f_1+th)}\right) - \left(\frac{1}{D(f_1)}\right)$$
  
=  $-\frac{1}{\xi(\varepsilon, f_1, h)} \left(D(f_1+th) - D(f_1)\right)$   
=  $-\frac{t}{\xi(\varepsilon, f_1, h)} \left(2\varepsilon^2 \Delta f_1 \Delta h + t(\Delta h)^2 + 4\varepsilon (h\tilde{R}_1 + R_1\tilde{h}\sin^2\left(\frac{x-y}{2}\right)\right),$ 

where  $\xi(\varepsilon, f_1, h)$  lies between  $D(f_1+th)$  and  $D(f_1)$  and hence  $0 < c_1|x-y|^2 \le |\xi(\varepsilon, f_1, h)| \le c_2|x-y|^2$  for some positive constants  $c_1$  and  $c_2$ . Noticing that

$$\left|2\varepsilon^2\Delta f_1\Delta h + t(\Delta h)^2 + 4\varepsilon(h\tilde{R}_1 + R_1\tilde{h}\sin^2\left(\frac{x-y}{2}\right)\right| \le C|x-y|^2,$$

we derive

$$\left|\log\left(\frac{1}{D(f_1+th)}\right) - \left(\frac{1}{D(f_1)}\right)\right| \le Ct,$$

which implies

$$\|F_{222}\|_{Y^{k-1}} \le Ct \left\| \oint \partial^k h(x) - \partial^k h(y) dy \right\|_{L^2} \le Ct \|h\|_{X^k}$$

Letting  $t \to 0$ , we obtain (3.12). And hence, we obtain the existence of Gateaux derivative of  $G_{22}$ . To prove the continuity of  $\partial_{f_1}G_{22}(\varepsilon, f)h$ , one just need to verify by the definition. Since there is no other new idea than the proof of continuity for  $G_1^0(\varepsilon, \Omega, f)$ , we omit it. Similarly, one can prove the existence and continuity of all derivatives  $\partial_{f_i}G_j^0$ , we omit the details and give them directly.

$$\begin{split} &\partial_{f_2} G_1^0(\varepsilon,\Omega,\boldsymbol{f})h \\ &= \frac{\gamma-1}{2R_1(x)} \int \log\left(\frac{1}{(R_1(x)-R_2(y))^2 + 4R_1(x)R_2(y)\sin^2\left(\frac{x-y}{2}\right)}\right) \\ &\times \left[(R_1(x)h(y) + R_1'(x)h'(y))\sin(x-y) + (R_1(x)h'(y) - R_1'(x)h(y))\cos(x-y)\right]dy \\ &+ \frac{1-\gamma}{2R_1(x)} \int \frac{-2\left(R_1(x)-R_2(y)\right)h(y) + 4R_1(x)h(y)\sin^2\left(\frac{x-y}{2}\right)}{(R_1(x)-R_2(y))^2 + 4R_1(x)R_2(y)\sin^2\left(\frac{x-y}{2}\right)} \\ &\times \left[(R_1(x)R_2(y) + R_1'(x)R_2'(y))\sin(x-y) + (R_1(x)R_2'(y) - R_1'(x)R_2(y))\cos(x-y)\right]dy \\ &+ \sum_{n=1}^{N-1} \frac{\gamma-1}{R_1(x)} \int \log\left(\frac{1}{\left|\boldsymbol{z}_1(x) - d\boldsymbol{e}_1 - Q_{\frac{2\pi n}{N}}\left(\boldsymbol{z}_2(y) - d\boldsymbol{e}_1\right)\right|}\right) \\ &\times \left[(R_1(x)h(y) + R_1'(x)h'(y))\sin(x-y - \frac{2\pi n}{N}) \\ &+ (R_1(x)h'(y) - R_1'(x)h(y))\cos(x-y - \frac{2\pi n}{N})\right]dy \\ &+ \sum_{n=1}^{N-1} \frac{\gamma-1}{2R_1(x)} \int \frac{\partial_{f_2}B_{12n}h}{\left|\boldsymbol{z}_1(x) - d\boldsymbol{e}_1 - Q_{\frac{2\pi n}{N}}\left(\boldsymbol{z}_2(y) - d\boldsymbol{e}_1\right)\right|^2} \\ &\times \left[(R_1(x)R_2(y) + R_1'(x)R_2'(y))\sin(x-y - \frac{2\pi n}{N}) \\ &+ (R_1(x)R_2(y) - R_1'(x)R_2(y))\cos(x-y - \frac{2\pi n}{N})\right]dy, \end{split}$$

$$\begin{split} &\partial_{f_1} G_2^0(\varepsilon,\Omega,f)h \\ &= \frac{1}{2R_2(x)} \int \log\left(\frac{1}{(R_2(x) - R_1(y))^2 + 4R_2(x)R_1(y)\sin^2\left(\frac{x-y}{2}\right)}\right) \\ &\times \left[(R_2(x)h(y) + R_2'(x)h'(y))\sin(x-y) + (R_2(x)h'(y) - R_2'(x)h(y))\cos(x-y)\right]dy \\ &- \frac{1}{2R_2(x)\varepsilon} \int \frac{-2(R_2(x) - R_1(y))h(y) + 4R_2(x)h(y)\sin^2\left(\frac{x-y}{2}\right)}{(R_2(x) - R_1(y))^2 + 4R_2(x)R_1(y)\sin^2\left(\frac{x-y}{2}\right)} \\ &\times \left[(R_2(x)R_1(y) + R_2'(x)R_1'(y))\sin(x-y) + (R_2(x)R_1'(y) - R_2'(x)R_1(y))\cos(x-y)\right]dy \\ &+ \sum_{n=1}^{N-1} \frac{1}{R_2(x)} \int \log\left(\frac{1}{\left|z_2(x) - de_1 - Q_{\frac{2\pi n}{N}}(z_1(y) - de_1)\right|}\right) \\ &\times \left[(R_2(x)h(y) + R_2'(x)h'(y))\sin(x-y - \frac{2\pi n}{N}) \\ &+ (R_2(x)h'(y) - R_2'(x)h(y))\cos(x-y - \frac{2\pi n}{N})\right]dy \\ &- \sum_{n=1}^{N-1} \frac{1}{R_2(x)} \int \frac{\partial_{f_1}B_{21n}h}{\left|z_2(x) - de_1 - Q_{\frac{2\pi n}{N}}(z_1(y) - de_1)\right|} \\ &\times \left[(R_2(x)R_1(y) + R_2'(x)R_1'(y))\sin(x-y - \frac{2\pi n}{N}) \\ &+ (R_2(x)R_1'(y) - R_2'(x)R_1(y))\cos(x-y - \frac{2\pi n}{N})\right]dy, \end{split}$$

$$\begin{aligned} \partial_{f_2} G_2^0(\varepsilon, \Omega, \mathbf{f}) h \\ &= (1 - b^2 - \gamma b^2) \Omega \left( \varepsilon^2 h'(x) - \frac{d\varepsilon h'(x) \cos(x)}{R_2(x)} + \frac{d\varepsilon R'_2(x) h'(x) \cos(x)}{R_2(x)^2} \right) \\ &- \frac{h(x)}{2R_2(x)^2} \int \log \left( \frac{1}{(R_2(x) - R_1(y))^2 + 4R_2(x)R_1(y) \sin^2\left(\frac{x-y}{2}\right)} \right) \\ &\times \left[ (R_2(x)R_1(y) + R'_2(x)R'_1(y)) \sin(x-y) + (R_2(x)R'_1(y) - R'_2(x)R_1(y)) \cos(x-y) \right] dy \\ &+ \frac{1}{2R_2(x)} \int \log \left( \frac{1}{(R_2(x) - R_1(y))^2 + 4R_2(x)R_1(y) \sin^2\left(\frac{x-y}{2}\right)} \right) \end{aligned}$$

$$\begin{split} & \times \left[ (h(x)R_{1}(y) + h'(x)R'_{1}(y))\sin(x-y) + (h(x)R'_{1}(y) - h'(x)R_{1}(y))\cos(x-y) \right] dy \\ & - \frac{1}{2R_{2}(x)} \int \frac{2(R_{2}(x) - R_{1}(y))h(x) + 4h(x)R_{1}(y)\sin^{2}\left(\frac{x-y}{2}\right)}{(R_{2}(x) - R_{1}(y))^{2} + 4R_{2}(x)R_{1}(y)\sin^{2}\left(\frac{x-y}{2}\right)} \\ & \times \left[ (R_{2}(x)R_{1}(y) + R'_{2}(x)R'_{1}(y))\sin(x-y) + (R_{2}(x)R'_{1}(y) - R'_{2}(x)R_{1}(y))\cos(x-y) \right] dy \\ & + \frac{(1-\gamma)h(x)}{2R_{2}(x)^{2}} \int \log \left( \frac{1}{(R_{2}(x) - R_{2}(y))^{2} + 4R_{2}(x)R_{2}(y)\sin^{2}\left(\frac{x-y}{2}\right)} \right) \\ & \times \left[ (R_{2}(x)R_{2}(y) + R'_{2}(x)R'_{2}(y))\sin(x-y) + (R_{2}(x)R'_{2}(y) - R'_{2}(x)R_{2}(y))\cos(x-y) \right] dy \\ & + \frac{\gamma-1}{2R_{2}(x)} \int \log \left( \frac{1}{(R_{2}(x) - R_{2}(y))^{2} + 4R_{2}(x)R_{2}(y)\sin^{2}\left(\frac{x-y}{2}\right)} \right) \\ & \times \left[ (h(x)R_{2}(y) + R'_{2}(x)h(y) + h'(x)R'_{2}(y) + R'_{2}(x)h'(y))\sin(x-y) \\ & + (h(x)R'_{2}(y) + R_{2}(x)h(y) - h'(x)R_{2}(y) - R'_{2}(x)h(y))\cos(x-y) \right] dy \\ & + \frac{1-\gamma}{2R_{2}(x)} \int \frac{2(R_{2}(x) - R_{2}(y))(h(x) - h(y)) + 4(h(x)R_{2}(y) + R_{2}(x)h(y))\sin^{2}\left(\frac{x-y}{2}\right)}{(R_{2}(x) - R_{2}(y))^{2} + 4R_{2}(x)R_{2}(y)\sin^{2}\left(\frac{x-y}{2}\right)} \\ & \times \left[ (R_{2}(x)R_{2}(y) + R'_{2}(x)R'_{2}(y))\sin(x-y) + (R_{2}(x)R'_{2}(y) - R'_{2}(x)R_{2}(y))\cos(x-y) \right] dy \\ & + \frac{\sum_{n=1}^{N-1} \frac{(\gamma-1)h(x)}{R_{2}(x)^{2}} \int \log \left( \frac{1}{\left|z_{2}(x) - de_{1} - Q_{\frac{2\pi n}{N}}(z_{1}(y) - de_{1})\right|} \right) \\ & \times \left[ (R_{2}(x)R_{1}(y) + R'_{2}(x)R'_{1}(y))\cos(x-y - \frac{2\pi n}{N}) \\ & + (h(x)R'_{1}(y) - R'_{2}(x)R_{1}(y))\cos(x-y - \frac{2\pi n}{N}) \\ & + (h(x)R'_{1}(y) - h'(x)R_{1}(y))\cos(x-y - \frac{2\pi n}{N}) \right] dy \\ & + \sum_{n=1}^{N-1} \frac{\gamma-1}{2R_{2}(x)} \int \frac{\partial_{f_{2}}B_{2nn}h}{\left|z_{2}(x) - de_{1} - Q_{\frac{2\pi n}{N}}(z_{1}(y) - de_{1})\right|^{2} \\ & \times \left[ (R_{2}(x)R_{1}(y) + R'_{2}(x)R'_{1}(y))\sin(x-y - \frac{2\pi n}{N}) \right] dy \\ & + \sum_{n=1}^{N-1} \frac{\gamma-1}{2R_{2}(x)} \int \frac{\partial_{f_{2}}B_{2nn}h}{\left|z_{2}(x) - de_{1} - Q_{\frac{2\pi n}{N}}(z_{1}(y) - de_{1})\right|^{2}} \\ & \times \left[ (R_{2}(x)R_{1}(y) + R'_{2}(x)R'_{1}(y))\sin(x-y - \frac{2\pi n}{N}) \right] dy \\ \end{aligned}$$

$$\begin{split} &+ (R_{2}(x)R_{1}'(y) - R_{2}'(x)R_{1}(y))\cos(x - y - \frac{2\pi n}{N}) \bigg] dy \\ &+ \sum_{n=1}^{N-1} \frac{(1 - \gamma)h(x)}{R_{2}(x)^{2}} \int \log \left( \frac{1}{\left| z_{2}(x) - de_{1} - Q_{\frac{2\pi n}{N}}(z_{2}(y) - de_{1}) \right|} \right) \\ &\times \left[ (R_{2}(x)R_{2}(y) + R_{2}'(x)R_{2}'(y))\sin(x - y - \frac{2\pi n}{N}) \\ &+ (R_{2}(x)R_{2}'(y) - R_{2}'(x)R_{2}(y))\cos(x - y - \frac{2\pi n}{N}) \right] dy \\ &+ \sum_{n=1}^{N-1} \frac{\gamma - 1}{R_{2}(x)} \int \log \left( \frac{1}{\left| z_{2}(x) - de_{1} - Q_{\frac{2\pi n}{N}}(z_{2}(y) - de_{1}) \right|} \right) \\ &\times \left[ (h(x)R_{2}(y) + R_{2}(x)h(y) + h'(x)R_{2}'(y) + R_{2}'(x)h'(y))\sin(x - y - \frac{2\pi n}{N}) \\ &+ (h(x)R_{2}'(y) + R_{2}(x)h'(y) - h'(x)R_{2}(y) - R_{2}'(x)h(y))\cos(x - y - \frac{2\pi n}{N}) \right] dy \\ &+ \sum_{n=1}^{N-1} \frac{1 - \gamma}{R_{2}(x)} \int \frac{\partial_{f_{2}}B_{22n}h}{\left| z_{2}(x) - de_{1} - Q_{\frac{2\pi n}{N}}(z_{2}(y) - de_{1}) \right|^{2}} \\ &\times \left[ (R_{2}(x)R_{2}(y) + R_{2}'(x)R_{2}'(y))\sin(x - y - \frac{2\pi n}{N}) \\ &+ (R_{2}(x)R_{2}'(y) - R_{2}'(x)R_{2}(y))\cos(x - y - \frac{2\pi n}{N}) \right] dy. \end{split}$$

Taking  $\varepsilon = 0$  and  $\mathbf{f} = 0$  in the derivatives  $\partial_{f_i} G_j^0$ , using integration by parts and changing x - y to y, we obtain

$$\partial_{f_1} G_1^0(0,\Omega,0)h$$

$$= \frac{1}{2} \int \log\left(\frac{1}{4\sin^2\left(\frac{y}{2}\right)}\right) \left[h(x-y)\sin(y) + (h'(x-y) - h'(x))\cos(y)\right] dy$$

$$- \frac{1}{2} \int \sin(y)h(x-y)dy$$

$$+ \frac{\gamma - 1}{2} \int \log\left(\frac{1}{(1-b)^2 + 4b\sin^2\left(\frac{y}{2}\right)}\right) \left[-h'(x)b\cos(y)\right] dy$$

$$+ \sum_{n=1}^{m-1} \int \log\left(\frac{1}{A_n}\right) \left[h(x-y)\sin(y - \frac{2\pi n}{m}) + h'(x-y)\cos(y - \frac{2\pi n}{m})\right] dy$$
(3.13)

$$= \frac{1}{2} \int \log\left(\frac{1}{\sin^2\left(\frac{y}{2}\right)}\right) \left[h(x-y)\sin(y) + (h'(x-y) - h'(x))\cos(y)\right] dy$$
$$- \frac{1}{2} \int \sin(y)h(x-y)dy$$
$$+ \frac{(1-\gamma)bh'(x)}{2} \int \log\left(\frac{1}{(1-b)^2 + 4b\sin^2\left(\frac{y}{2}\right)}\right)\cos(y)dy,$$

$$\partial_{f_2} G_1^0(0, \Omega, 0)h \tag{3.14}$$

$$= \frac{\gamma - 1}{2} \int \log\left(\frac{1}{(1 - b)^2 + 4b\sin^2\left(\frac{x - y}{2}\right)}\right) [h(y)\sin(x - y) + h'(y)\cos(x - y)] dy$$

$$+ \frac{1 - \gamma}{2} \int \frac{b\sin(x - y)\left(-2\left(1 - b\right)h(y) + 4h(y)\sin^2\left(\frac{x - y}{2}\right)\right)}{(1 - b)^2 + 4b\sin^2\left(\frac{x - y}{2}\right)}$$

$$+ \sum_{n=1}^{N-1} \frac{\gamma - 1}{2} \int \log\left(\frac{1}{A_n}\right) \left[h(y)\sin(x - y - \frac{2\pi k}{m}) + h'(y)\cos(x - y - \frac{2\pi n}{N})\right] dy$$

$$= \frac{\gamma - 1}{2} \int \log\left(\frac{1}{(1 - b)^2 + 4b\sin^2\left(\frac{x - y}{2}\right)}\right) h(y)\sin(x - y) dy$$

$$+ \frac{1 - \gamma}{2} \int \partial_y \left(\log\left(\frac{1}{(1 - b)^2 + 4b\sin^2\left(\frac{x - y}{2}\right)}\right) \cos(x - y)\right) h(y) dy$$

$$+ \frac{1 - \gamma}{2} \int \frac{2b\sin(x - y) \left(b - \cos(y)\right)}{(1 - b)^2 + 4b\sin^2\left(\frac{x - y}{2}\right)} h(y) dy$$

$$= (1 - \gamma)b^2 \int \frac{\sin(x - y)h(y)}{(1 - b)^2 + 4b\sin^2\left(\frac{x - y}{2}\right)} dy,$$

$$\partial_{f_1} G_2^0(0, \Omega, 0)h$$

$$= \frac{1}{2} \int \log\left(\frac{1}{(1-b)^2 + 4b\sin^2\left(\frac{x-y}{2}\right)}\right) [h(y)\sin(x-y) + h'(y)\cos(x-y)] dy$$

$$- \frac{1}{2} \int \frac{-2(b-1)h(y) + 4bh(y)\sin^2\left(\frac{x-y}{2}\right)}{(1-b)^2 + 4b\sin^2\left(\frac{x-y}{2}\right)} \sin(x-y) dy$$

$$+ \sum_{n=1}^{m-1} \frac{1}{2} \int \log\left(\frac{1}{A_n}\right) \left[h(y)\sin(x-y-\frac{2\pi k}{m}) + h'(y)\cos(x-y-\frac{2\pi k}{m})\right] dy$$

$$= -\int \frac{\sin(x-y)h(y)}{(1-b)^2 + 4b\sin^2\left(\frac{x-y}{2}\right)} dy.$$
(3.15)

$$\partial_{f_2} G_2^0(0,\Omega,0)h \tag{3.16}$$

$$= -\frac{1}{2b} \int \log\left(\frac{1}{(1-b)^2 + 4b\sin^2\left(\frac{x-y}{2}\right)}\right) h'(x)\cos(x-y)dy$$

$$\frac{\gamma - 1}{2} \int \log\left(\frac{1}{4b^2\sin^2\left(\frac{x-y}{2}\right)}\right) [h(y)\sin(x-y) + (h'(y) - h'(x))\cos(x-y)]dy$$

$$+ \frac{1-\gamma}{2} \int h(y)\sin(x-y)dy.$$

Take  $h_1, h_2 \in H^k$  with

$$h_1(x) = \sum_{j=1}^{\infty} a_j \cos(jx), \quad h_2(x) = \sum_{j=1}^{\infty} b_j \cos(jx).$$

To compute the precise values of  $\partial_{f_i} G_j^0(0,\Omega,0)$ , we will need the following integral identities taken from [10].

Lemma 3.3. For 0 < b < 1, there hold

$$\int e^{imy} \log\left(\frac{1}{\sin^2\left(\frac{y}{2}\right)}\right) = \begin{cases} \frac{1}{|m|}, & m \in \mathbb{Z}, m \neq 0, \\ 2\log 2, & m = 0, \end{cases}$$
(3.17)

$$\int e^{imy} \log\left(\frac{1}{(1-b)^2 + 4b\sin^2\left(\frac{y}{2}\right)}\right) = \begin{cases} \frac{1}{|m|}b^{|m|}, & m \in \mathbb{Z}, m \neq 0, \\ 2\log(2b), & m = 0, \end{cases}$$
(3.18)

$$\int \frac{e^{imy}}{(1-b)^2 + 4b\sin^2\left(\frac{y}{2}\right)} = \frac{b^{|m|}}{1-b^2}.$$
(3.19)

*Proof.* (3.17) is exactly Lemma 3.3 [10]. (3.18) and (3.19) follows from Lemma 3.4 and Lemma 3.1 [10] respectively by taking  $r = \frac{1-b}{1+b}$ .

We denote  $G^0 = (G_1^0, G_2^0)$ , then the linearized operator of  $G^0$  is as follows.

### Lemma 3.4.

$$DG^{0}(0,\Omega,0)(h_{1},h_{2}) = \begin{pmatrix} \partial_{f_{1}}G_{1}^{0}(0,\Omega,0)h_{1} + \partial_{f_{2}}G_{1}^{0}(0,\Omega,0)h_{2} \\ \partial_{f_{1}}G_{2}^{0}(0,\Omega,0)h_{1} + \partial_{f_{2}}G_{2}^{0}(0,\Omega,0)h_{2} \end{pmatrix} = \sum_{j=1}^{\infty} \frac{1}{2}M_{j} \begin{pmatrix} a_{j} \\ b_{j} \end{pmatrix} \sin(jx),$$
(3.20)

where

$$M_j = \begin{pmatrix} j - 1 + (\gamma - 1)jb^2 & (1 - \gamma)b^{j+1} \\ -b^{j-1} & (\gamma - 1)(j - 1) + j \end{pmatrix}.$$

*Proof.* We first compute  $\partial_{f_1} G_1^{\alpha}(0,\Omega,0)h_1$ . Using (3.17), we have

$$\begin{split} &\frac{1}{2} \int \log\left(\frac{1}{\sin^2\left(\frac{y}{2}\right)}\right) [h_1(x-y)\sin(y) + (h'_1(x-y) - h'_1(x))\cos(y)] \, dy \\ &= \sum_{j=1}^{\infty} \frac{a_j}{2} \int \log\left(\frac{1}{\sin^2\left(\frac{y}{2}\right)}\right) [\cos(j(x-y))\sin(y) - j\sin(j(x-y))\cos(y) + j\sin(jx)\cos(y)] \, dy \\ &= \sum_{j=1}^{\infty} \frac{a_j}{2} \int \log\left(\frac{1}{\sin^2\left(\frac{y}{2}\right)}\right) [\cos(jx)\cos(jy)\sin(y) + \sin(jx)\sin(jy)\sin(y) \\ &- j\sin(jx)\cos(jy)\cos(y) + j\cos(jx)\sin(jy)\cos(y) + j\sin(jx)\cos(y)] \, dy \\ &= \sum_{j=1}^{\infty} \frac{a_j}{2} \int \log\left(\frac{1}{\sin^2\left(\frac{y}{2}\right)}\right) [\sin(jx)\sin(jy) - j\sin(jx)\cos(jy)\cos(y) + j\sin(jx)\cos(y)] \, dy \\ &= \frac{a_1\sin(x)}{2} \int \log\left(\frac{1}{\sin^2\left(\frac{y}{2}\right)}\right) [\sin(y)\sin(y) - \cos(y)\cos(y) + \cos(y)] \, dy \\ &+ \sum_{j=2}^{\infty} \frac{a_j\sin(jx)}{2} \int \log\left(\frac{1}{\sin^2\left(\frac{y}{2}\right)}\right) [\sin(jy)\sin(y) - j\cos(jy)\cos(y) + j\cos(y)] \, dy \\ &= \frac{a_1\sin(x)}{2} \int \log\left(\frac{1}{\sin^2\left(\frac{y}{2}\right)}\right) [-\cos(2y) + \cos(y)] \, dy \\ &+ \sum_{j=2}^{\infty} \frac{a_j\sin(jx)}{4} \int \log\left(\frac{1}{\sin^2\left(\frac{y}{2}\right)}\right) [(1 - j)\cos((j - 1)y) - (j + 1)\cos((j + 1)y) + 2j\cos(y)] \, dy \\ &= \frac{a_1\sin(x)}{2} \left(-\frac{1}{2} + 1\right) + \sum_{j=2}^{\infty} \frac{a_j\sin(jx)}{4} (-1 - 1 + 2j) \\ &= \frac{a_1\sin(x)}{4} + \sum_{j=2}^{\infty} \frac{(j - 1)a_j\sin(jx)}{2}. \end{split}$$

By trigonometric formula, we drive

$$-\frac{1}{2} \int \sin(y) h_1(x-y) dy$$
  
=  $\sum_{j=1}^{\infty} \frac{-a_j}{2} \int \sin(y) \cos(j(x-y)) dy$   
=  $\sum_{j=1}^{\infty} \frac{-a_j}{4} \int (\sin(jx+(1-j)y) - \sin((j+1)y - jx)) dy$ 

$$=\frac{-a_1\sin(x)}{4}.$$

The remaining term follows by using (3.18)

$$\frac{bh_1'(x)}{2} \oint \log\left(\frac{1}{(1-b)^2 + 4b\sin^2\left(\frac{y}{2}\right)}\right)\cos(y)dy = \frac{b^2h_1'(x)}{2}.$$

Thus, if we combine the above identities, we arrive at

$$\partial_{f_1} G_1^{\alpha}(0,\Omega,0) h_1 = \sum_{j=1}^{\infty} \frac{j-1+(\gamma-1)jb^2}{2} a_j \sin(jx).$$

Now, we compute  $\partial_{f_2} G_1^0(0,\Omega,0)h_2$ . By (3.19), we have

$$\begin{aligned} &\int \frac{\sin(y)h_2(x-y)}{(1-b)^2 + 4b\sin^2\left(\frac{y}{2}\right)} dy \\ &= \sum_{j=1}^{\infty} b_j \int \frac{\sin(y)\cos(jx-jy)}{(1-b)^2 + 4b\sin^2\left(\frac{y}{2}\right)} dy \\ &= \sum_{j=1}^{\infty} b_j \int \frac{\cos(jx)\sin(y)\cos(jy) + \sin(jx)\sin(y)\sin(jy)}{(1-b)^2 + 4b\sin^2\left(\frac{y}{2}\right)} dy \\ &= \sum_{j=1}^{\infty} b_j\sin(jx) \int \frac{\sin(y)\sin(jy)}{(1-b)^2 + 4b\sin^2\left(\frac{y}{2}\right)} dy \\ &= \sum_{j=1}^{\infty} \frac{b_j\sin(jx)}{2} \int \frac{\cos((j-1)y) - \cos((j+1)y)}{(1-b)^2 + 4b\sin^2\left(\frac{y}{2}\right)} dy \\ &= \sum_{j=1}^{\infty} \frac{b_j\sin(jx)}{2} \frac{1}{1-b^2} \left(b^{j-1} - b^{j+1}\right) \\ &= \sum_{j=1}^{\infty} \frac{b^{j-1}}{2} b_j\sin(jx). \end{aligned}$$

Thus, we obtain

$$\partial_{f_2} G_1^{\alpha}(0,\Omega,0) h_2 = \sum_{j=1}^{\infty} \frac{(1-\gamma)b^{j+1}}{2} b_j \sin(jx).$$

Using the above results, we deduce

$$\partial_{f_2} G_1^{\alpha}(0,\Omega,0) h_2 = \sum_{j=1}^{\infty} \frac{-b^{j-1}}{2} b_j \sin(jx).$$

and

$$\partial_{f_2} G_2^{\alpha}(0,\Omega,0) h_2 = \sum_{j=1}^{\infty} \frac{(\gamma-1)(j-1) + j}{2} b_j \sin(jx).$$

We choose b such that  $DG^0(0,\Omega,0)$  is a isomorphism from  $X_b^k \to Y_b^{k-1}$ .

**Lemma 3.5.** For  $\gamma \neq 0$ , there exists  $b \in (0,1)$  such that  $DG^0(0,\Omega,0) : X_b^k \to Y_b^{k-1}$  is an isomorphism.

*Proof.* By Lemma 3.4 and  $M_1 = \begin{pmatrix} -(1-\gamma)b^2 & (1-\gamma)b^2 \\ -1 & 1 \end{pmatrix}$ , it is easy to see  $DG^0(0,\Omega,0)$ 

maps  $X_b^k$  to  $Y_b^{k-1}$  for any  $k \ge 3$ . Next, we show the invertibility of  $DG^0$ .

Case i):  $\gamma \in (0, 1)$ . We first consider the case  $\gamma \in (0, 1)$ , where we assume  $b \in (0, \frac{\sqrt{2}}{2})$ . The determinant of the matrix  $M_j$  is

det 
$$M_j = \gamma (1 + (\gamma - 1)b^2)j^2 + ((1 - \gamma)(1 + (\gamma - 1)b^2) - \gamma)j + \gamma - 1 + (1 - \gamma)b^{2j},$$

for each  $j \ge 2$ . We define the auxiliary function

$$D(x) = \gamma (1 + (\gamma - 1)b^2)x^2 + ((1 - \gamma)(1 + (\gamma - 1)b^2) - \gamma)x + \gamma - 1.$$

Suppose  $0 < b < \frac{\sqrt{2}}{2}$ , then for  $x \ge 2$ , we have

$$D'(x) = 2\gamma(1 + (\gamma - 1)b^2)x + (1 - \gamma)(1 + (\gamma - 1)b^2) - \gamma$$
  

$$\geq 2\gamma(1 - b^2)x + (1 - \gamma)(1 - b^2) - \gamma$$
  

$$\geq 4\gamma(1 - b^2) + (1 - \gamma)(1 - b^2) - \gamma$$
  

$$\geq \gamma > 0,$$

which implies that D(x) is strict increasing and

$$D(x) \ge D(2) = 4\gamma(1 + (\gamma - 1)b^2) + 2((1 - \gamma)(1 + (\gamma - 1)b^2) - \gamma) + \gamma - 1$$
  
>  $4\gamma(1 - b^2) + 2((1 - \gamma)(1 - b^2) - \gamma) + \gamma - 1$   
>  $2\gamma + 1 - \gamma - 2\gamma + \gamma - 1 = 0,$ 

for  $x \ge 2$ . Thus,  $\det(M_j) > D(j) \ge D(2) > 0$  for  $j \ge 2$  and hence  $M_j$  is invertible for  $\gamma \in (0,1)$  and  $b \in (0, \frac{\sqrt{2}}{2})$ .

Case ii):  $\gamma > 1$ . In this case, we assume  $b \in (0, 1)$ . Noticing that for all  $\gamma > 1$  and 0 < b < 1, it holds

$$\begin{cases} \gamma(1+(\gamma-1)b^2) > 0\\ \frac{(\gamma-1)(1+(\gamma-1)b^2)+\gamma}{2\gamma(1+(\gamma-1)b^2)} \leq 1\\ \gamma-1+(1-\gamma)b^{2j} = (\gamma-1)(1-b^{2j}) > 0. \end{cases}$$

Thus, one easily infer from the properties of quadratic function that  $det(M_j) > 0$  for  $j \ge 2$ , which implies the invertibility of  $M_j$ .

Case iii):  $\gamma < 0$ . In this case, it is hard to give a precise interval, where  $M_j$  is invertible for all  $j \geq 2$  and b belonging to this interval. Instead, we will give a abstract proof of the existence of  $b \in (0,1)$  such that  $M_j$  is invertible. We claim that for all  $\gamma \in \mathbb{R}$  with  $\gamma \neq 0$ , there exists a set  $S_{\gamma}$  composed of at most countable points such that as long as  $b \in (0,1) \setminus S_{\gamma}, M_j$  is invertible for any  $j \geq 2$ . Indeed, for  $\gamma \neq 0$ , we have  $\gamma(1+(\gamma-1)b^2) \neq 0$ for all  $b \in (0,1)$  except for at most one point  $b_{\gamma}$ . Thus, D(x) is a quadratic polynomial of x for  $b \neq b_{\gamma}$ . Noticing that  $b^{2x} \to 0$  as  $x \to +\infty$ , for  $b \in (0,1) \setminus \{b_{\gamma}\}$ , there exists  $x_{\gamma}(b)$ sufficiently large such that  $\det(M_j) \neq 0$  for all  $j \geq x_{\gamma}(b)$ . Moreover,  $x_{\gamma}(b)$  is contious with respect to b. Take arbitrary closed interval  $I \subset ((0,1) \setminus \{b_{\gamma}\})$ . By the continuity of  $x_{\gamma}(b)$ , one can choose a universal  $x_{\gamma,I}$  satisfying  $\det(M_j) \neq 0$  for all  $j \geq x_{\gamma,I}$  and  $b \in I$ . For each fixed j,  $\det(M_j)$  is a 2jth degree polynomial of b and hence the number of b such that  $\det(M_j) = 0$  is at most 2j. Therefore, there are finite many b such that  $\det(M_j)$  is not invertible for  $2 \leq j \leq x_{\gamma,I}$ . This is,  $\det(M_j)$  is invertible for all  $j \geq 2$  and  $b \in I$  except for finite many b. Since  $(0,1) \setminus \{b_{\gamma}\}$  is a union of countable closed intervals, the claim above follows easily.

Now, we fix b such that  $det(M_j)$  is invertible for all  $j \ge 2$ . Then,  $M_j^{-1}$  is given by

$$M_j^{-1} = \frac{1}{\det(M_j)} \begin{pmatrix} (\gamma - 1)(j - 1) + j & b^{j-1} \\ -(1 - \gamma)b^{j+1} & j - 1 + (\gamma - 1)jb^2, \end{pmatrix}.$$

For  $g = (g_1, g_2) \in Y_b^{k-1}$  with  $g_1 = (1-\gamma)b^2d_1\sin(x) + \sum_{j=1}^{\infty} c_j\sin(jx)$  and  $g_2 = \sum_{j=1}^{\infty} d_j\sin jx$ , we define its inverse  $f = (f_1, f_2)$  with

$$f^{t} = 2 \begin{pmatrix} (1-\gamma)b^{2} \\ 1 \end{pmatrix} \frac{d_{1}}{1 - (1-\gamma)b^{2}} \cos(x) + \sum_{j=2}^{\infty} 2M_{j}^{-1} \begin{pmatrix} c_{j} \\ d_{j} \end{pmatrix} \cos(jx)$$

Noticing that  $\det(M_j) = O(j^2)$ , thus the elements of  $M_j^{-1}$  is of order  $j^{-1}$  and hence  $f \in X_b^k$ .

Remark 3.6. Lemma 3.5 fails for  $\gamma = 0$ . In this case  $M_j = \begin{pmatrix} j - 1 - jb^2 & b^{j+1} \\ -b^{j-1} & 1 \end{pmatrix}$ . Though  $M_j$  is still invertible for  $j \ge 2$  and  $0 < b < \frac{\sqrt{2}}{2}$ , the lower right element of  $M_j^{-1}$  is of order O(1) and hence f defined as above does not belong to  $X_b^k$ .

Our next step is to choose  $\Omega$  such that  $G^0 = (G_1^0, G_2^0)$  maps  $X_b^k$  to  $Y_b^{k-1}$  for all  $\varepsilon$  small. According to Lemma 3.1, we conclude that

$$G_1^0(0,\Omega,0) = G_2^0 = (1-b^2+\gamma b^2) \left(\pi\Omega d - \sum_{n=1}^{N-1} \frac{1-\cos(\frac{2\pi n}{N})}{2d\left((-1+\cos(\frac{2\pi n}{N}))^2 + \sin^2(\frac{2\pi n}{N})\right)}\right) \sin(x) + \frac{1-\cos(\frac{2\pi n}{N})}{2d\left((-1+\cos(\frac{2\pi n}{N}))^2 + \sin^2(\frac{2\pi n}{N})\right)}$$

Therefore, if we take  $\Omega$  to be

$$\Omega^{0}_{*} := \sum_{n=1}^{N-1} \frac{1 - \cos(\frac{2\pi n}{N})}{2\pi d^{2} \left( (-1 + \cos(\frac{2\pi n}{N}))^{2} + \sin^{2}(\frac{2\pi n}{N}) \right)},$$
(3.21)

then  $G_1^0(0, \Omega^0_*, 0) = G_2^0(0, \Omega^0_*, 0) = 0$  and hence  $G^0(0, \Omega^0_*, 0) \in Y_b^{k-1}$ .

For general  $\varepsilon$  and f, we can take  $\Omega$  properly such that the range of  $G^0 = (G_1^0, G_2^0)$  belongs to  $Y_b^{k-1}$ .

## Lemma 3.7. There exists

$$\Omega^0(\varepsilon, \boldsymbol{f}) := \Omega^0_* + \varepsilon \mathcal{R}_{\Omega}(\varepsilon, \boldsymbol{f})$$

with  $\Omega^0_*$  given by (3.21) and continuous function  $\mathcal{R}_{\Omega}(\varepsilon, \mathbf{f})$  such that  $\tilde{G}^0(\varepsilon, \mathbf{f}) := G^0(\varepsilon, \Omega^0(\varepsilon, \mathbf{f}), \mathbf{f})$ maps  $(-\varepsilon_0, \varepsilon_0) \times V$  to  $Y^{k-1}_b$ . Moreover,  $\partial_{\mathbf{f}} \mathcal{R}_{\Omega}(\varepsilon, \mathbf{f}) : X^k \times X^k \to \mathbb{R}^2$  is continuous.

*Proof.* It suffices to find  $\Omega$  such that

$$\oint G_1^0(\varepsilon, \Omega, \boldsymbol{f}) \sin(x) dx = (1 - \gamma) b^2 \oint G_2^0(\varepsilon, \Omega, \boldsymbol{f}) \sin(x) dx.$$
(3.22)

For the sake of convenience, set  $\Omega = \Omega^0_* + \Omega_{\varepsilon}$  and assume  $f_1 = \sum_{j=1}^{\infty} a_j \cos(jx)$  and  $f_2 = \sum_{j=1}^{\infty} b_j \cos(jx)$  with  $a_1 = (1 - \gamma)b^2b_1$ . By (3.1), (3.10), (3.11) and the calculations in Lemma 3.4, we have

$$\begin{aligned} & \oint G_1^0(\varepsilon, \Omega, \boldsymbol{f}) \sin(x) dx = \pi (1 - b^2 + \gamma b^2) \Omega \left( \frac{d}{2} + \varepsilon \oint \mathcal{R}_{11}(\varepsilon, \boldsymbol{f}) \sin(x) dx \right) + (\gamma - 1) b^2 (a_1 - b_1) \\ & - \frac{(1 - b^2 + \gamma b^2) \Omega_*^0 d}{2} + \int \varepsilon \mathcal{R}_1(\varepsilon, \boldsymbol{f}) \sin(x) dx \\ &= \pi (1 - b^2 + \gamma b^2) \Omega_\varepsilon \left( \frac{d}{2} + \varepsilon \oint \mathcal{R}_{11}(\varepsilon, \boldsymbol{f}) \sin(x) dx \right) + (\gamma - 1) b^2 (a_1 - b_1) \\ & + \varepsilon \oint \left( \pi (1 - b^2 + \gamma b^2) \Omega_*^0 \mathcal{R}_{11}(\varepsilon, \boldsymbol{f}) + \mathcal{R}_1(\varepsilon, \boldsymbol{f}) \right) \sin(x) dx, \end{aligned}$$

and

$$\begin{aligned} \oint G_2^0(\varepsilon,\Omega,\boldsymbol{f})\sin(x)dx &= \pi(1-b^2+\gamma b^2)\Omega\left(\frac{d}{2}+\varepsilon\int\mathcal{R}_{21}(\varepsilon,\boldsymbol{f})\sin(x)dx\right) + (-a_1+b_1)\\ &-\frac{(1-b^2+\gamma b^2)\Omega_*^0d}{2} + \int\varepsilon\mathcal{R}_2(\varepsilon,\boldsymbol{f})\sin(x)dx\\ &= \pi(1-b^2+\gamma b^2)\Omega_\varepsilon\left(\frac{d}{2}+\varepsilon\int\mathcal{R}_{21}(\varepsilon,\boldsymbol{f})\sin(x)dx\right) + (-a_1+b_1)\\ &+\varepsilon\int\left(\pi(1-b^2+\gamma b^2)\Omega_*^0\mathcal{R}_{21}(\varepsilon,\boldsymbol{f}) + \mathcal{R}_2(\varepsilon,\boldsymbol{f})\right)\sin(x)dx.\end{aligned}$$

Then, (3.22) is equivalent to

$$\pi (1 - b^2 + \gamma b^2) \Omega_{\varepsilon} \left( \frac{d}{2} + \varepsilon \int \mathcal{R}_{11}(\varepsilon, \mathbf{f}) \sin(x) dx \right) + (\gamma - 1) b^2 (a_1 - b_1)$$
  
+  $\varepsilon \int (\pi (1 - b^2 + \gamma b^2) \Omega_*^0 \mathcal{R}_{11}(\varepsilon, \mathbf{f}) + \mathcal{R}_1(\varepsilon, \mathbf{f})) \sin(x) dx$   
=  $(1 - \gamma) b^2 \left( \pi (1 - b^2 + \gamma b^2) \Omega_{\varepsilon} \left( \frac{d}{2} + \varepsilon \int \mathcal{R}_{21}(\varepsilon, \mathbf{f}) \sin(x) dx \right) + (-a_1 + b_1) \right)$ 

$$+\varepsilon \int \left(\pi(1-b^2+\gamma b^2)\Omega^0_*\mathcal{R}_{21}(\varepsilon,\boldsymbol{f})+\mathcal{R}_2(\varepsilon,\boldsymbol{f})\right)\sin(x)dx\right),$$

which implies

$$\Omega_{\varepsilon} = \varepsilon \frac{-\oint \left(\pi (1-b^2+\gamma b^2) \Omega_*^0 \mathcal{R}_{21}(\varepsilon, \boldsymbol{f}) + \mathcal{R}_2(\varepsilon, \boldsymbol{f})\right) \sin(x) dx + (1-\gamma) b^2 \oint \left(\pi (1-b^2+\gamma b^2) \Omega_*^0 \mathcal{R}_{21}(\varepsilon, \boldsymbol{f}) + \mathcal{R}_2(\varepsilon, \boldsymbol{f})\right) \sin(x) dx}{\pi (1-b^2+\gamma b^2) \left((1-(1-\gamma) b^2) d/2 + \varepsilon \left(\int \mathcal{R}_{11}(\varepsilon, \boldsymbol{f}) \sin(x) dx - (1-\gamma) b^2 \int \mathcal{R}_{21}(\varepsilon, \boldsymbol{f}) \sin(x) dx\right)\right)}$$

Hence, the desired  $\mathcal{R}_{\Omega}(\varepsilon, f)$  is given by  $\mathcal{R}_{\Omega}(\varepsilon, f) = \frac{\Omega_{\varepsilon}}{\varepsilon}$  and the regularity of  $\mathcal{R}_{\Omega}(\varepsilon, f)$  follows from Lemmas 3.1 and 3.2.

The proof is thus complete.

Now, we are in the position to prove Theorem 1.1. **Proof of Theorem 1.1:** We first prove  $\partial_f \tilde{G}^0(0,0) : X_b^k \to Y_b^{k-1}$  is an isomorphism. By chain rule, it holds

$$\partial_f \tilde{G}^0(0,0)h = \partial_\Omega G^0(0,\Omega^0_*,0)\partial_f \Omega^0(0,0)h + \partial_f G^0(0,\Omega^0_*,0).$$

From Lemma 3.7, we know  $\partial_f \Omega^0(0,0) = 0$ . Hence  $\partial_f \tilde{G}^0(0,0) = \partial_f G^0(0,\Omega^0_*,0)$ , and we achieve the desired result by Lemma 3.5.

According to Lemma 3.1–Lemma 3.7, we can apply implicit function theorem to prove that there exists  $\varepsilon_0 > 0$  such that

$$\left\{ (\varepsilon, \boldsymbol{f}) \in [-\varepsilon_0, \varepsilon_0] \times V : \tilde{G}_i^0(\varepsilon, \boldsymbol{f}) = 0, \quad \text{for } i = 1, 2 \right\}$$

is parameterized by one-dimensional curve  $\varepsilon \in [-\varepsilon_0, \varepsilon_0] \to (\varepsilon, f_{\varepsilon})$ .

Let  $\tilde{\boldsymbol{f}}(x) = \boldsymbol{f}(-x)$ . Our last step is to show that if  $(\varepsilon, \boldsymbol{f})$  is a solution to  $\tilde{G}_i^0(\varepsilon, f) = 0$ for i = 1, 2, then so is  $(-\varepsilon, \tilde{\boldsymbol{f}})$ . By changing y to -y in the integral representation of  $G^0(\varepsilon, \Omega, f)$  and using the fact that f is an even function, we obtain  $\Omega^0(\varepsilon, \boldsymbol{f}) = \Omega^0(-\varepsilon, \tilde{\boldsymbol{f}})$ . Then we can insert it into  $G^0(\varepsilon, \Omega, \boldsymbol{f})$  and derive  $\tilde{G}^0(-\varepsilon, \tilde{\boldsymbol{f}}) = 0$  by a similar substitution of variables as before.

Hence the proof of Theorem 1.1 is finished.

# 4. Existence of co-rotating solutions: the case $0 < \alpha < 2$

The proof for the case  $0 < \alpha < 2$  follows the same argument as the case  $\alpha = 0$ , so we will be sketch and omit the details.

Set

$$V := \begin{cases} \{f \in X_b^k \mid ||g||_{X^k} < 1\}, & 0 < \alpha < 1, \\ \{f \in X_b^{k+\log} \mid ||g||_{X^{k+\log}} < 1\}, & \alpha = 1, \\ \{f \in X_b^{k+\alpha-1} \mid ||g||_{X^{k+\alpha-1}} < 1\}, & 1 < \alpha < 2. \end{cases}$$

We have the following two lemmas concerning the regularity of  $G_i^{\alpha}$ .

**Lemma 4.1.**  $G_i^{\alpha}(\varepsilon, \Omega, f): (-\varepsilon_0, \varepsilon_0) \times \mathbb{R} \times V \to Y^{k-1}$  is continuous for i = 1, 2.

*Proof.* For  $0 < \alpha < 1$ , the proof is almost the same as the case  $\alpha = 0$  since the kernel  $\frac{1}{|(R_1(x)-R_1(y))^2+4R_1(x)R_1(y)\sin^2(\frac{x-y}{2})|^{\frac{\alpha}{2}}} \in L^1$ . For the case  $1 \le \alpha < 2$ , the most singular terms

has been treated in [5] and hence no additional difficulty arises in the proof of this case. Further more, by Taylor's formula (3.4), we have

$$\begin{aligned} G_{1}^{\alpha} &= \pi (1 - b^{2} + \gamma b^{2}) \Omega d \sin(x) \end{aligned}$$
(4.1)  
+  $C_{\alpha} (1 - \frac{\alpha}{2}) \int \frac{f_{1}(x - y) \sin(y) dy}{(4 \sin^{2}(\frac{y}{2}))^{\frac{\alpha}{2}}} + C_{\alpha} \int \frac{(f_{1}'(x - y) - f_{1}'(x)) \cos(y) dy}{(4 \sin^{2}(\frac{y}{2}))^{\frac{\alpha}{2}}} \\ &+ \frac{(1 - \gamma) \alpha C_{\alpha}}{2} \int \frac{b \left(-2(1 - b) f_{2}(y) + 4 f_{2}(y) \sin^{2}\left(\frac{x - y}{2}\right)\right) \sin(x - y)}{|(1 - b)^{2} + 4b \sin^{2}\left(\frac{x - y}{2}\right)|^{\frac{\alpha}{2} + 1}} dy \\ &+ (1 - \gamma) C_{\alpha} \int \frac{f_{2}(y) \sin(x - y) + (f_{2}'(y) - b f_{1}'(x)) \cos(x - y))}{|(1 - b)^{2} + 4b \sin^{2}\left(\frac{x - y}{2}\right)|^{\frac{\alpha}{2}}} dy \\ &+ (b^{2} - \gamma b^{2} - 1) \sum_{n=1}^{N-1} \frac{\alpha C_{\alpha} (1 - \cos(\frac{2\pi n}{N})) \sin(x)}{2 \left((-1 + \cos(\frac{2\pi n}{N}))^{2} + \sin^{2}(\frac{2\pi n}{N})\right)^{1 + \frac{\alpha}{2}} d^{1 + \alpha}} + \varepsilon |\varepsilon|^{\alpha} \mathcal{R}_{1}^{\alpha}(\varepsilon, \Omega, \boldsymbol{f}), \end{aligned}$ 

and

$$\begin{aligned} G_{2}^{\alpha} &= \pi (1 - b^{2} + \gamma b^{2}) \Omega d \sin(x) \end{aligned}$$
(4.2)  
 
$$&= -(1 - \gamma) b^{-\alpha} \left( C_{\alpha} (1 - \frac{\alpha}{2}) \int \frac{f_{2}(x - y) \sin(y) dy}{(4 \sin^{2}(\frac{y}{2}))^{\frac{\alpha}{2}}} + C_{\alpha} \int \frac{(f_{2}'(x - y - )f_{2}'(x)) \cos(y) dy}{(4 \sin^{2}(\frac{y}{2}))^{\frac{\alpha}{2}}} \right) \\ &- \frac{\alpha C_{\alpha}}{2} \int \frac{(2(1 - b)f_{1}(y) + 4bf_{1}(y) \sin^{2}(\frac{x - y}{2})) \sin(x - y)}{|(1 - b)^{2} + 4b \sin^{2}(\frac{x - y}{2})|^{\frac{\alpha}{2} + 1}} dy \\ &+ \frac{C_{\alpha}}{b} \int \frac{bf_{1}(y) \sin(x - y) + (bf_{1}'(y) - f_{2}'(x)) \cos(x - y))}{|(1 - b)^{2} + 4b \sin^{2}(\frac{x - y}{2})|^{\frac{\alpha}{2}}} dy \\ &+ (b^{2} - \gamma b^{2} - 1) \sum_{n=1}^{N-1} \frac{\alpha C_{\alpha} (1 - \cos(\frac{2\pi n}{N})) \sin(x)}{2 \left((-1 + \cos(\frac{2\pi n}{N}))^{2} + \sin^{2}(\frac{2\pi n}{N})\right)^{1 + \frac{\alpha}{2}} d^{1 + \alpha}} + \varepsilon |\varepsilon|^{\alpha} \mathcal{R}_{2}^{\alpha}(\varepsilon, \Omega, \boldsymbol{f}). \end{aligned}$$

**Lemma 4.2.** The Gautex derivative  $\partial_{f_i} G_j^{\alpha}(\varepsilon, \Omega, f) : (-\varepsilon_0, \varepsilon_0) \times \mathbb{R} \times V \to L(X^k, Y^{k-1})$ exists and is continuous for i, j = 1, 2.

*Proof.* For the case  $0 < \alpha < 1$ , the proof is almost the same as the case  $\alpha = 0$ . For the case  $1 \leq \alpha < 2$ , the most singular terms are treated in [5] and hence the proof of this case contains no additional difficulty.

To calculate the exact expression of the Gautex derivatives, we will follow the notations of [21]. Let F(a, b, c, z) be the hypergeometric function, whose definition and properties can be found in [21, 41].

Denote

$$\Theta_j = \begin{cases} 2^{\alpha - 1} \frac{\Gamma(1 - \alpha)}{(\Gamma(1 - \frac{\alpha}{2}))^2} \left( \frac{\Gamma(1 + \frac{\alpha}{2})}{\Gamma(2 - \frac{\alpha}{2})} - \frac{\Gamma(j + \frac{\alpha}{2})}{\Gamma(1 + j - \frac{\alpha}{2})} \right), & \alpha \neq 1 \\ \frac{2}{\pi} \sum_{l=2}^{j}, & \alpha = 1, \end{cases}$$

$$(x)_j := \begin{cases} 1, & j = 0\\ x(x+1)(x+2)\cdots(x+j-1), & j \ge 1. \end{cases}$$

and

$$\Lambda_j(b) := \frac{1}{b} \int_0^\infty \frac{J_n(bt)J_n(t)}{t^{1-\alpha}} dt,$$

where  $J_n$  represents the Bessel function of the first kind. The relationship between  $\Lambda_j$  and the hypergeometric function is (see (5.24) [21])

$$\Lambda_j = \frac{\Gamma\left(\frac{\alpha}{2}\right)}{2^{1-\alpha}\Gamma\left(1-\frac{\alpha}{2}\right)} \frac{\left(\frac{\alpha}{2}\right)_j}{j!} b^{j-1} F\left(\left(\frac{\alpha}{2}\right), j+\frac{\alpha}{2}, j+1, b^2\right).$$

Take  $h_1, h_2$  with

$$h_1(x) = \sum_{j=1}^{\infty} a_j \cos(jx), \quad h_2(x) = \sum_{j=1}^{\infty} b_j \cos(jx).$$

The linearized operator of  $G^{\alpha}$  is as follows.

## Lemma 4.3.

$$DG^{\alpha}(0,\Omega,0)(h_1,h_2) = \begin{pmatrix} \partial_{f_1} G_1^{\alpha}(0,\Omega,0)h_1 + \partial_{f_2} G_1^{\alpha}(0,\Omega,0)h_2\\ \partial_{f_1} G_2^{\alpha}(0,\Omega,0)h_1 + \partial_{f_2} G_2^{\alpha}(0,\Omega,0)h_2 \end{pmatrix} = \sum_{j=1}^{\infty} jM_j \begin{pmatrix} a_j\\ b_j \end{pmatrix} \sin(jx),$$
(4.3)

where

$$M_j = \begin{pmatrix} \Theta_j + (\gamma - 1)b^2\Lambda_1(b) & (1 - \gamma)b^2\Lambda_j(b) \\ -\Lambda_j(b) & (\gamma - 1)b^{-\alpha}\Theta_j + \Lambda_1(b) \end{pmatrix}.$$

*Proof.* By direct calculations, we obtain

$$\partial_{f_{1}}G_{1}^{\alpha}(0,\Omega,0)h \tag{4.4}$$

$$= \left(1 - \frac{\alpha}{2}\right)C_{\alpha}\int \frac{h(x-y)\sin(y)}{|4\sin^{2}(\frac{y}{2})|^{\frac{\alpha}{2}}}dy + C_{\alpha}\int \frac{(h'(x-y) - h'(x))\cos(y)}{|4\sin^{2}(\frac{y}{2})|^{\frac{\alpha}{2}}}dy$$

$$+ (1 - \gamma)C_{\alpha}\int \frac{bh'(x)\cos(y)}{|(1 - b)^{2} + 4b\sin^{2}(\frac{y}{2})|^{\frac{\alpha}{2}}}dy$$

$$\partial_{f_{2}}G_{1}^{\alpha}(0,\Omega,0)h = (\gamma - 1)C_{\alpha}\int \frac{h(x-y)\sin(y) + h'(x-y)\cos(y)}{|(1 - b)^{2} + 4b\sin^{2}(\frac{y}{2})|^{\frac{\alpha}{2}}}dy$$

$$+ (1 - \gamma)\alpha C_{\alpha}\int \frac{bh(x-y)\sin(y)(b - \cos(y))}{|(1 - b)^{2} + 4b\sin^{2}(\frac{y}{2})|^{\frac{\alpha}{2}+1}}dy$$

$$\partial_{f_{1}}G_{2}^{\alpha}(0,\Omega,0)h = C_{\alpha}\int \frac{h(x-y)\sin(y) + h'(x-y)\cos(y)}{|(1 - b)^{2} + 4b\sin^{2}(\frac{y}{2})|^{\frac{\alpha}{2}}}dy \tag{4.6}$$

$$-\alpha C_{\alpha} \int \frac{h(x-y)\sin(y)\left(1-b\cos(y)\right)dy}{\left|(1-b)^{2}+4b\sin^{2}\left(\frac{y}{2}\right)\right|^{\frac{\alpha}{2}+1}}dy$$

$$= (1-\gamma)\left(\frac{\alpha}{2}-1\right)C_{\alpha} \int \frac{\sin(y)h(x-y)}{\left|4b^{2}\sin^{2}\left(\frac{y}{2}\right)\right|^{\frac{\alpha}{2}}}dy + (\gamma-1)C_{\alpha} \int \frac{(h'(x-y)-h'(x))\cos(y)}{\left|4b^{2}\sin^{2}\left(\frac{y}{2}\right)\right|^{\frac{\alpha}{2}}}dy$$

$$- \frac{C_{\alpha}h'(x)}{b} \int \frac{\cos(y)}{\left|(1-b)^{2}+4b\sin^{2}\left(\frac{y}{2}\right)\right|^{\frac{\alpha}{2}}}dy$$
(4.7)

We first calculate  $\partial_{f_1} G_1^{\alpha}(0,\Omega,0)h_1$ . The first two terms has been obtained in [28] for  $0 < \alpha < 1$  and [9] for  $1 \le \alpha < 2$ ,

$$\begin{split} \left(1-\frac{\alpha}{2}\right)C_{\alpha} & \int \frac{h_{1}(x-y)\sin(y)}{\left|4\sin^{2}\left(\frac{y}{2}\right)\right|^{\frac{\alpha}{2}}}dy + C_{\alpha} \int \frac{(h_{1}'(x-y)-h_{1}'(x))\cos(y)}{\left|4\sin^{2}\left(\frac{y}{2}\right)\right|^{\frac{\alpha}{2}}}dy \\ &= \sum_{j=1}^{\infty} j\Theta_{j}a_{j}\sin(jx), \\ \text{where } \Theta_{j} = \begin{cases} 2^{\alpha-1}\frac{\Gamma(1-\alpha)}{(\Gamma(1-\frac{\alpha}{2}))^{2}} \left(\frac{\Gamma(1+\frac{\alpha}{2})}{\Gamma(2-\frac{\alpha}{2})} - \frac{\Gamma(j+\frac{\alpha}{2})}{\Gamma(1+j-\frac{\alpha}{2})}\right), & \alpha \neq 1 \\ \frac{2}{\pi}\sum_{l=2}^{j}, & \alpha = 1. \end{cases}$$

Using (3.19) in [21],

$$C_{\alpha}h'_{1}(x)b \int \frac{\cos(y)}{\left|(1-b)^{2}+4b\sin^{2}\left(\frac{y}{2}\right)\right|^{\frac{\alpha}{2}}}dy$$
  
=  $C_{\alpha}h'_{1}(x)b^{2}\frac{\alpha}{2}F\left(\frac{\alpha}{2},1+\frac{\alpha}{2},2,b^{2}\right)$   
=  $b^{2}\Lambda_{1}(b)h'_{1}(x).$ 

$$\partial_{f_1} G_1^{\alpha}(0,\Omega,0) h_1 \tag{4.8}$$
$$= \sum_{j=1}^{\infty} j \left( \Theta_j + (\gamma - 1) b^2 \Lambda_1(b) \right) a_j \sin(jx).$$

Next, we consider the second term  $\partial_{f_2} G_1^{\alpha}(0,\Omega,0)h_2$ . By (3.19) in [21], we have

$$-C_{\alpha} \oint \frac{h_2(x-y)\sin(y)}{\left|(1-b)^2 + 4b\sin^2\left(\frac{y}{2}\right)\right|^{\frac{\alpha}{2}}} dy$$
  
=  $-C_{\alpha} \sum_{j=1}^{\infty} b_j \oint \frac{\cos(jx-jy)\cos(y)}{\left|(1-b)^2 + 4b\sin^2\left(\frac{y}{2}\right)\right|^{\frac{\alpha}{2}}} dy$ 

$$= -C_{\alpha} \sum_{j=1}^{\infty} b_j \int \frac{\cos(jx)\cos(jy)\sin(y) + \sin(jx)\sin(jy)\sin(y)}{\left|(1-b)^2 + 4b\sin^2\left(\frac{y}{2}\right)\right|^{\frac{\alpha}{2}}} dy$$
$$= -C_{\alpha} b_j \sum_{j=1}^{\infty} \sin(jx) \int \frac{\sin(jy)\sin(y)}{\left|(1-b)^2 + 4b\sin^2\left(\frac{y}{2}\right)\right|^{\frac{\alpha}{2}}} dy.$$

Using integration by parts, we obtain

$$C_{\alpha} \oint \frac{-h'_{2}(x-y)\cos(y)}{\left|(1-b)^{2}+4b\sin^{2}\left(\frac{y}{2}\right)\right|^{\frac{\alpha}{2}}} dy$$

$$= C_{\alpha} \sum_{j=1}^{\infty} b_{j} \int \frac{j\sin(jx-jy)\cos(y)}{\left|(1-b)^{2}+4b\sin^{2}\left(\frac{y}{2}\right)\right|^{\frac{\alpha}{2}}} dy$$

$$= C_{\alpha} \sum_{j=1}^{\infty} b_{j} \int \frac{j\sin(jx)\cos(jy)\cos(y)-j\cos(jx)\sin(jy)\cos(y)}{\left|(1-b)^{2}+4b\sin^{2}\left(\frac{y}{2}\right)\right|^{\frac{\alpha}{2}}} dy$$

$$= C_{\alpha} \sum_{j=1}^{\infty} b_{j}j\sin(jx) \int \frac{\cos(jy)\cos(y)}{\left|(1-b)^{2}+4b\sin^{2}\left(\frac{y}{2}\right)\right|^{\frac{\alpha}{2}}} dy$$

$$= C_{\alpha} \sum_{j=1}^{\infty} b_{j}\sin(jx) \int \frac{\cos(y)}{\left|(1-b)^{2}+4b\sin^{2}\left(\frac{y}{2}\right)\right|^{\frac{\alpha}{2}}} d\sin(jy)$$

$$= C_{\alpha} \sum_{j=1}^{\infty} b_{j}\sin(jx) \int \left(\frac{\sin(jy)\sin(y)}{\left|(1-b)^{2}+4b\sin^{2}\left(\frac{y}{2}\right)\right|^{\frac{\alpha}{2}}} + \frac{\alpha b}{2} \frac{\sin(jy)\sin(2y)}{\left|(1-b)^{2}+4b\sin^{2}\left(\frac{y}{2}\right)\right|^{\frac{\alpha}{2}+1}} \right) dy.$$

By direct calculation and (3.19) in [21], we derive

$$-\alpha bC_{\alpha} \oint \frac{h_{2}(x-y)\sin(y)\cos(y)}{\left|(1-b)^{2}+4b\sin^{2}\left(\frac{y}{2}\right)\right|^{\frac{\alpha}{2}+1}}dy$$

$$=-\frac{\alpha bC_{\alpha}}{2}\sum_{j=1}^{\infty}b_{j} \oint \frac{\cos(j(x-y))\sin(2y)}{\left|(1-b)^{2}+4b\sin^{2}\left(\frac{y}{2}\right)\right|^{\frac{\alpha}{2}+1}}dy$$

$$=-\frac{\alpha bC_{\alpha}}{2}\sum_{j=1}^{\infty}b_{j} \oint \frac{\cos(jx)\cos(jy)\sin(2y)+\sin(jx)\sin(jy)\sin(2y)}{\left|(1-b)^{2}+4b\sin^{2}\left(\frac{y}{2}\right)\right|^{\frac{\alpha}{2}+1}}dy$$

$$=-\frac{\alpha bC_{\alpha}}{2}\sum_{j=1}^{\infty}b_{j}\sin(jx) \oint \frac{\sin(jy)\sin(2y)}{\left|(1-b)^{2}+4b\sin^{2}\left(\frac{y}{2}\right)\right|^{\frac{\alpha}{2}+1}}dy.$$

Thus, we conclude by combining the above equalities that

$$\partial_{f_2} G_1^{\alpha}(0,\Omega,0) h_2 = (1-\gamma)\alpha b^2 C_{\alpha} \int \frac{h_2(x-y)\sin(y)}{\left|(1-b)^2 + 4b\sin^2\left(\frac{y}{2}\right)\right|^{\frac{\alpha}{2}+1}}$$

$$= (1 - \gamma)\alpha b^{2}C_{\alpha}\sum_{j=1}^{\infty} b_{j}\sin(jx) \oint \frac{\sin(jy)\sin(y)}{\left|(1 - b)^{2} + 4b\sin^{2}\left(\frac{y}{2}\right)\right|^{\frac{\alpha}{2}+1}} dy$$

$$= (1 - \gamma)\alpha b^{2}\frac{C_{\alpha}}{2}\sum_{j=1}^{\infty} b_{j}\sin(jx) \oint \frac{\cos((j - 1)y) - \cos((j + 1)y)}{\left|(1 - b)^{2} + 4b\sin^{2}\left(\frac{y}{2}\right)\right|^{\frac{\alpha}{2}+1}} dy$$

$$= (1 - \gamma)\alpha b^{2}\frac{C_{\alpha}}{2}\sum_{j=1}^{\infty} b_{j}\sin(jx) \left[b^{j-1}\frac{\left(\frac{\alpha}{2}+1\right)_{(j-1)}}{(j - 1)!}F\left(\frac{\alpha}{2} + 1, j + \frac{\alpha}{2}, j, b^{2}\right)\right]$$

$$-b^{j+1}\frac{\left(\frac{\alpha}{2}+1\right)_{(j+1)}}{(j + 1)!}F\left(\frac{\alpha}{2} + 1, j + 2 + \frac{\alpha}{2}, j + 2, b^{2}\right)\right]$$

$$= (1 - \gamma)\alpha b^{2}\frac{C_{\alpha}}{2}\sum_{j=1}^{\infty} b_{j}\sin(jx)b^{j-1}\frac{\left(\frac{\alpha}{2}+1\right)_{(j-1)}}{(j - 1)!}\left[F\left(\frac{\alpha}{2} + 1, j + \frac{\alpha}{2}, j, b^{2}\right)\right]$$

$$-b^{2}\frac{\left(\frac{\alpha}{2}+j\right)\left(\frac{\alpha}{2}+j + 1\right)}{j(j + 1)}F\left(\frac{\alpha}{2} + 1, j + 2 + \frac{\alpha}{2}, j + 2, b^{2}\right)\right].$$

Using (3.11), (3.14) and (3.14)-(3.15) in [21], we have

$$\begin{split} b^2 \frac{\left(\frac{\alpha}{2}+j\right)\left(\frac{\alpha}{2}+j+1\right)}{j(j+1)} F\left(\frac{\alpha}{2}+1,j+2+\frac{\alpha}{2},j+2,b^2\right) \\ &= \frac{\frac{\alpha}{2}+j}{j} \left[F\left(\frac{\alpha}{2}+1,j+1+\frac{\alpha}{2},j+1,b^2\right) - F\left(\frac{\alpha}{2},j+1+\frac{\alpha}{2},j+1,b^2\right)\right] \\ &= \frac{\frac{\alpha}{2}+j}{j} \left[\frac{j}{\frac{\alpha}{2}+j} F\left(\frac{\alpha}{2}+1,j+\frac{\alpha}{2},j,b^2\right) + \frac{\frac{\alpha}{2}}{j+\frac{\alpha}{2}} F\left(\frac{\alpha}{2}+1,j+\frac{\alpha}{2},j+1,b^2\right)\right] \\ &- \frac{\frac{\alpha}{2}+j}{j} F\left(\frac{\alpha}{2},j+1+\frac{\alpha}{2},j+1,b^2\right) \\ &= F\left(\frac{\alpha}{2}+1,j+\frac{\alpha}{2},j,b^2\right) \\ &+ \frac{\frac{\alpha}{2}}{j} F\left(\frac{\alpha}{2}+1,j+\frac{\alpha}{2},j+1,b^2\right) - \frac{\frac{\alpha}{2}+j}{j} F\left(\frac{\alpha}{2},j+1+\frac{\alpha}{2},j+1,b^2\right) \\ &= F\left(\frac{\alpha}{2}+1,j+\frac{\alpha}{2},j,b^2\right) - F\left(\frac{\alpha}{2},j+\frac{\alpha}{2},j+1,b^2\right). \end{split}$$

Hence, we finally arrive at

$$\partial_{f_2} G_1^{\alpha}(0,\Omega,0) h_2 \tag{4.9}$$

$$= (1-\gamma)\alpha b^2 \frac{C_{\alpha}}{2} \sum_{j=1}^{\infty} b_j \sin(jx) b^{j-1} \frac{\left(\frac{\alpha}{2}+1\right)_{(j-1)}}{(j-1)!} F(\frac{\alpha}{2}, j+\frac{\alpha}{2}, j+1, b^2)$$

$$= (1 - \gamma)b^2 \sum_{j=1}^{\infty} j\Lambda_j(b)b_j \sin(jx)$$

Using the above calculations, one easily obtain

$$\partial_{f_1} G_2^{\alpha}(0,\Omega,0) h_1 = -\sum_{j=1}^{\infty} j \Lambda_j(b) a_j \sin(jx),$$
(4.10)

and

$$\partial_{f_2} G_2^{\alpha}(0,\Omega,0) h_2 \tag{4.11}$$
$$= -\sum_{j=1}^{\infty} j \left( (1-\gamma) b^{-\alpha} \Theta_j - \Lambda_1(b) \right) b_j \sin(jx).$$

The proof of this lemma is therefore complete.

We can choose b such that  $DG^{\alpha}(0,\Omega,0)$  is an isomorphism.

**Lemma 4.4.** There exists  $b_0 \in (0, 1)$  such that for  $b \in (0, b_0)$ ,

$$DG^{\alpha}(0,\Omega,0): \begin{cases} X_{b}^{k} \to Y_{b}^{k-1}, & 0 < \alpha < 1, \\ X_{b}^{k+\log} \to Y_{b}^{k-1}, & \alpha = 1, \\ X_{b}^{k+\alpha-1} \to Y_{b}^{k-1}, & 1 < \alpha < 2, \end{cases}$$

is an isomorphism.

*Proof.* We choose  $b_0$  such that for  $0 < b < b_0$ ,  $DG^{\alpha}(0, \Omega, 0)$  is invertible. The determinant of  $M_j$  is

det $(M_j) = -(1-\gamma)b^{-\alpha}\Theta_j \left(\Theta_j + (\gamma-1)b^2\Lambda_1(b)\right) + \Lambda_1(b) \left(\Theta_j + (\gamma-1)b^2\Lambda_1(b)\right) + b^2\Lambda_j(b)^2$ . We need the following properties of  $\Theta_j$  and  $\Lambda_j$ , whose proof can be found in [9], [21] and [28]:

> $\Theta_j > 0 \text{ for } j \ge 2,$   $\Theta_j(b) \text{ is increasing with respect to } j,$   $\Lambda_j(b) \text{ is decreasing with respect to } j,$   $\Lambda_j(b) \text{ is increasing with respect to } b,$   $\int O(1), \qquad 0 < \alpha < 1,$ O(1) = 0 = 0

$$\Theta_j = \begin{cases} O(1), & 0 < \alpha < 1, \\ O(\log(j)), & \alpha = 1, \\ O(j^{\alpha - 1}), & 1 < \alpha < 2. \end{cases}$$

In view of the above properties, we choose  $b_0 \in (0, 1)$  such that

$$\frac{\Theta_2}{2} \ge b_0^2 \Lambda_1(b_0), \quad \frac{(1-\gamma)b_0^{-\alpha}\Theta_2}{4} \ge \Lambda_1(b_0), \quad \frac{(1-\gamma)b_0^{-\alpha}\Theta_2^2}{8} \ge 3b_0^2 \Lambda_1(b_0)^2.$$

Then, for  $0 < b < b_0$ , we have for  $j \ge 2$ ,

$$\begin{aligned} \det(M_j) &= -(1-\gamma)b^{-\alpha}\Theta_j \left(\Theta_j + (\gamma-1)b^2\Lambda_1(b)\right) + \Lambda_1(b) \left(\Theta_j + (\gamma-1)b^2\Lambda_1(b)\right) + b^2\Lambda_j(b)^2 \\ &\leq -(1-\gamma)b^{-\alpha}\Theta_j \left(\Theta_j - b^2\Lambda_1(b)\right) + \Lambda_1(b) \left(\Theta_j + \gamma b^2\Lambda_1(b)\right) + b^2\Lambda_j(b)^2 \\ &\leq -(1-\gamma)b^{-\alpha}\frac{\Theta_j^2}{2} + \Lambda_1(b) \left(\Theta_j + \gamma b^2\Lambda_1(b)\right) + b^2\Lambda_j(b)^2 \\ &\leq -(1-\gamma)b^{-\alpha}\frac{\Theta_j^2}{4} + \gamma b^2\Lambda_1(b)^2 + b^2\Lambda_j(b)^2 \\ &\leq -(1-\gamma)b^{-\alpha}\frac{\Theta_j^2}{8} \leq -(1-\gamma)b^{-\alpha}\frac{\Theta_j^2}{4}. \end{aligned}$$

Hence,  $M_j$  is invertible for  $j \ge 2$ . Notice that

$$M_{1} = \begin{pmatrix} -(1-\gamma)b^{2}\Lambda_{1}(b) & (1-\gamma)b^{2}\Lambda_{1}(b) \\ -\Lambda_{1}(b) & \Lambda_{1}(b) \end{pmatrix} = \Lambda_{1}(b) \begin{pmatrix} -(1-\gamma)b^{2} & (1-\gamma)b^{2} \\ -1 & 1 \end{pmatrix}$$

is of rank 1. For  $g = (g_1, g_2) \in Y_b^{k-1}$  with  $g_1 = b^2 d_1 \sin(x) + \sum_{j=1}^{\infty} c_j \sin(jx)$  and  $g_2 = b^2 d_1 \sin(x) + \sum_{j=1}^{\infty} c_j \sin(jx)$  $\sum_{j=1}^{\infty} d_j \sin(jx)$ , we define its inverse  $f = (f_1, f_2)$  with

$$f^{t} = \binom{(1-\gamma)b^{2}}{1} \frac{d_{1}}{(1-(1-\gamma)b^{2})\Lambda_{1}(b)}\cos(x) + \sum_{j=2}^{\infty}\frac{1}{j}M_{j}^{-1}\binom{c_{j}}{d_{j}}\cos(jx).$$

Then, by the asymptotic behavior of  $\Theta_j$ , the fact  $\Lambda_j \leq \Lambda_1$  and the characterization of  $X^{k+\log}$  and  $X^{k+\alpha-1}$  (see Proposition 1.1 and Remark 3.2 [10]), it is easy to verify that  $f \in X_b^k$  if  $0 < \alpha < 1$ ,  $f \in X_b^{k+\log}$  if  $\alpha = 1$  and  $f \in X_b^{k+\alpha-1}$  if  $1 < \alpha < 2$ .

Next, we adjust  $\Omega$  such that the range of  $G^{\alpha}$  belongs to  $Y_b^{k-1}$ .

Lemma 4.5. There exists

$$\Omega^{\alpha}(\varepsilon, \boldsymbol{f}) := \Omega^{\alpha}_* + \varepsilon \mathcal{R}_{\Omega}(\varepsilon, \boldsymbol{f})$$

with

$$\Omega_*^{\alpha} = \sum_{n=1}^{N-1} \frac{\alpha C_{\alpha} (1 - \cos(\frac{2\pi n}{N}))}{2\pi \left( (-1 + \cos(\frac{2\pi n}{N}))^2 + \sin^2(\frac{2\pi n}{N}) \right)^{1+\frac{\alpha}{2}} d^{2+\alpha}}$$

and a continuous function  $\mathcal{R}_{\Omega}(\varepsilon, \boldsymbol{f}) : (-\varepsilon_0, \varepsilon_0) \times V \to \mathbb{R}$ , such that  $\tilde{G}^{\alpha}(\varepsilon, \boldsymbol{f}) := G^{\alpha}(\varepsilon, \Omega^{\alpha}(\varepsilon, \boldsymbol{f}), \boldsymbol{f})$  $\begin{array}{l} maps \ (-\varepsilon_0, \varepsilon_0) \times V \ to \ Y_b^{k-1}. \\ Moreover, \ \partial_{\boldsymbol{f}} \mathcal{R}_{\Omega}(\varepsilon, \boldsymbol{f})h : X^k \times X^k \to \mathbb{R}^2 \ is \ continuous. \end{array}$ 

*Proof.* The proof is the same as Lemma 3.7, so we omit it.

Theorem 1.3 follows from combining the above lemmas and applying the implicit function theorem.

### 5. EXISTENCE OF TRAVELING SOLUTIONS

The travelling patch pair centered at (d, 0) takes the form

$$\theta_{\varepsilon}(\boldsymbol{x},t) = \theta_{0,\varepsilon}(\boldsymbol{x} - tW\boldsymbol{e}_2)$$

with W some fixed speed, and  $e_2$  the unit vector in  $x_2$  direction. According to (1.1), we derive

$$(\mathbf{u}_0(\boldsymbol{x}) - W\boldsymbol{e}_2) \cdot \nabla \theta_{0,\varepsilon}(\boldsymbol{x}) = 0,$$

which yields

$$(\mathbf{u}_0(\boldsymbol{x}) - W\boldsymbol{e}_2) \cdot \mathbf{n}(\boldsymbol{x}) = 0,$$

for all  $\boldsymbol{x}$  lies on the interfaces. According to Biot-Savart law and Green-Stokes formula, when  $\alpha = 0$ , the interfaces satisfies the following equation

$$0 = H_i^0(\varepsilon, \Omega, \mathbf{f}) \tag{5.1}$$

$$\begin{split} &= -\pi (1 - b^2 + \gamma b^2) W \left( \sin(x) - \frac{R_i'(x)}{R_i(x)} \cos(x) \right) \\ &+ \frac{1}{2R_i(x)\varepsilon} \int \log \left( \frac{1}{(R_i(x) - R_1(y))^2 + 4R_i(x)R_1(y)\sin^2\left(\frac{x-y}{2}\right)} \right) \\ &\times \left[ (R_i(x)R_1(y) + R_i'(x)R_1'(y))\sin(x-y) + (R_i(x)R_1'(y) - R_i'(x)R_1(y))\cos(x-y) \right] dy \\ &- \frac{1 - \gamma}{2R_i(x)\varepsilon} \int \log \left( \frac{1}{(R_i(x) - R_2(y))^2 + 4R_i(x)R_2(y)\sin^2\left(\frac{x-y}{2}\right)} \right) \\ &\times \left[ (R_i(x)R_2(y) + R_i'(x)R_2'(y))\sin(x-y) + (R_i(x)R_2'(y) - R_i'(x)R_2(y))\cos(x-y) \right] dy \\ &- \frac{1}{R_i(x)\varepsilon} \int \log \left( \frac{1}{|z_i(x) - de_1 - Q_\pi\left(z_1(y) - de_1\right)|} \right) \left[ (R_i(x)R_1(y) + R_i'(x)R_1'(y))\sin(x-y-\pi) + (R_i(x)R_1'(y) - R_i'(x)R_1(y))\cos(x-y-\pi) \right] dy \\ &+ \frac{1 - \gamma}{R_i(x)\varepsilon} \int \log \left( \frac{1}{|z_i(x) - de_1 - Q_\pi\left(z_2(y) - de_1\right)|} \right) \left[ (R_i(x)R_2(y) + R_i'(x)R_2'(y))\sin(x-y-\pi) + (R_i(x)R_2'(y) - R_i'(x)R_2(y))\cos(x-y-\pi) \right] dy \end{split}$$

When  $0 < \alpha < 2$ , the interfaces obey the following equation  $0 = H_i^{\alpha}(\varepsilon, \Omega, \mathbf{f})$ 

$$= -\pi (1 - b^{2} + \gamma b^{2}) W \left( \sin(x) - \frac{R'_{i}(x)}{R_{i}(x)} \cos(x) \right) + \frac{C_{\alpha}}{R_{i}(x)\varepsilon|\varepsilon|^{\alpha}} \int \frac{1}{\left| (R_{i}(x) - R_{1}(y))^{2} + 4R_{i}(x)R_{1}(y)\sin^{2}\left(\frac{x-y}{2}\right) \right|^{\frac{\alpha}{2}}} \times \left[ (R_{i}(x)R_{1}(y) + R'_{i}(x)R'_{1}(y))\sin(x-y) + (R_{i}(x)R'_{1}(y) - R'_{i}(x)R_{1}(y)) \right] \cos(x-y)dy - \frac{(1 - \gamma)C_{\alpha}}{R_{i}(x)\varepsilon|\varepsilon|^{\alpha}} \int \frac{1}{\left| (R_{i}(x) - R_{2}(y))^{2} + 4R_{i}(x)R_{2}(y)\sin^{2}\left(\frac{x-y}{2}\right) \right|^{\frac{\alpha}{2}}}$$

(5.2)

$$\times \left[ (R_{i}(x)R_{2}(y) + R'_{i}(x)R'_{2}(y))\sin(x-y) + (R_{i}(x)R'_{2}(y) - R'_{i}(x)R_{2}(y)) \right]\cos(x-y)dy \\ - \frac{C_{\alpha}}{R_{i}(x)\varepsilon} \int \frac{1}{|\boldsymbol{z}_{i}(x) - d\boldsymbol{e}_{1} - Q_{\pi}\left(\boldsymbol{z}_{1}(y) - d\boldsymbol{e}_{1}\right)|^{\alpha}} \left[ (R_{i}(x)R_{1}(y) + R'_{i}(x)R'_{1}(y))\sin(x-y-\pi) + (R_{i}(x)R'_{1}(y) - R'_{i}(x)R_{1}(y))\cos(x-y-\pi) \right] dy \\ + \frac{(1-\gamma)C_{\alpha}}{R_{i}(x)\varepsilon} \int \frac{1}{|\boldsymbol{z}_{i}(x) - d\boldsymbol{e}_{1} - Q_{\pi}\left(\boldsymbol{z}_{2}(y) - d\boldsymbol{e}_{1}\right)|^{\alpha}} \left[ (R_{i}(x)R_{2}(y) + R'_{i}(x)R'_{2}(y))\sin(x-y-\pi) + (R_{i}(x)R'_{2}(y) - R'_{i}(x)R_{2}(y))\cos(x-y-\pi) \right] dy.$$

**Proof of Theorem 1.4:** We only sketch the proof. Notice that  $H^{\alpha}$  and  $G^{\alpha}$  with N = 2 are almost identical except for the first term and the signs before the fourth and fifth terms. Thus we can use a similar method as in Sections 3 and 4 to prove the continuity of  $H^{\alpha}(\varepsilon, W, f)$  and  $\partial_{f} H^{\alpha}(\varepsilon, W, f)$ . Moreover, one can verify that for each  $W \in \mathbb{R}$ ,  $\partial_{f} H^{\alpha}(0, W, 0)$  exactly equals  $\partial_{f} G^{\alpha}(0, \Omega, 0)$  and hence  $DH^{\alpha}(0, W, 0)$  is an isomorphism under the same assumptions as previous sections. Then we just need to select the appropriate W and apply the implicit function theorem. The  $W^{\alpha}_{*}$  chosen is given by

$$W_*^{\alpha} := \begin{cases} \frac{1}{4\pi d}, & \alpha = 0, \\ \frac{\alpha C_{\alpha}}{2\pi (2d)^{1+\alpha}}, & 0 < \alpha < 2. \end{cases}$$

We omit the details.

#### References

- W. Ao, J. Dávila, L.D. Pino, M. Musso and J. Wei, Travelling and rotating solutions to the generalized inviscid surface quasi-geostrophic equation, to appear in Transaction of AMS, https://doi.org/10.1090/tran/8406.
- [2] T. Buckmaster, S. Shkoller and V. Vicol, Nonuniqueness of weak Solutions to the SQG Equation, Comm. Pure Appl. Math., 72 (2019), 1809-1874.
- [3] J. Burbea, Motions of vortex patches, Lett. Math. Phys., 6(1982), 1–16.
- [4] G. R. Burton, Steady symmetric vortex pairs and rearrangements, Proc. R. Soc. Edinb., Sect. A, 108(1988), 269-290.
- [5] D. Cao, G. Qin, W. Zhan and C. Zou, Existence and regularity of co-rotating and travelling global solutions for the generalized SQG equation, preprint, 2021.
- [6] D. Cao, S. Peng and S. Yan, Planar vortex patch problem in incompressible steady flow, Adv. Math, 270(2015), 2741-2785.
- [7] D. Cao, S. Lai and W. Zhan, Traveling vortex pairs for 2D incompressible Euler equations, Preprint arXiv:2012.10918.
- [8] D. Cao, G. Wang and W. Zhan, Desingularization of vortices for 2D steady Euler flows via the vorticity method, SIAM J. Math. Anal., 52(2020), no. 6, 5363–5388.
- [9] A. Castro, D. Córdoba and J. Gómez-Serraon, Existence and regularity of rotating global solutions for the generalized surface quasi-geostrophic equations, *Duke Math. J.*, 165(2016), no. 5, 935–984.
- [10] A. Castro, D. Córdoba and J. Gómez-Serraon, Uniformly rotating analytic global patch solutions for active scalars, Ann. PDE, 2(2016), no. 1, Art. 1, 34.
- [11] A. Castro, D. Córdoba, and J. Gómez-Serrano, Uniformly rotating smooth solutions for the incompressible 2D Euler equations, Arch. Ration. Mech. Anal., 231(2019), no. 2, 719–785.

- [12] A. Castro, D. Córdoba and J. Gómez-Serrano, Global smooth solutions for the inviscid SQG equation, Mem. Amer. Math. Soc., 266(2020), no. 1292.
- [13] D. Chae, The quasi-geostrophic equation in the Triebel-Lizorkin spaces, Nonlinearity, 16(2003), no. 2, 479–495.
- [14] D. Chae, P. Constantin, D. Cordoba, F. Gancedo and J. Wu, Generalized surface quasi-geostrophic equations with singular velocities, *Comm. Pure Appl. Math.*, 65(2012), no. 8, 1037–1066.
- [15] P. Constantin, A.J. Majda and E. Tabak, Formation of strong fronts in the 2-D quasigeostrophic thermal active scalar, *Nonlinearity*, 7(1994), no. 6, 1495–1533.
- [16] R. Coralie, Relative equilibria with holes for the surface quasi-geostrophic equations, J. Differential Equations, 263(2017), 567-614.
- [17] A. Córdoba, D. Córdoba and F. Gancedo. Uniqueness for SQG patch solutions. Trans. Amer. Math. Soc. Ser. B, 5 (2018), 1–31
- [18] D. Córdoba, M. A. Fontelos, A. M. Mancho and J. L. Rodrigo, Evidence of singularities for a family of contour dynamics equations, *Proc. Natl. Acad. Sci. USA* 102(2005), 5949–5952.
- [19] J. Davila, M. D. Pino, M. Musso and J. Wei, Gluing Methods for Vortex Dynamics in Euler Flows, Arch. Ration. Mech. Anal., 235(2020), 1467–1530.
- [20] G. S. Deem and N. J. Zabusky, Vortex waves: stationary "V-states", interactions, recurrence, and breaking, Phys. Rev. Lett., 40(13)(1978), 859–862.
- [21] F. de la Hoz, Z. Hassainia and T. Hmidi, Doubly connected V-states for the generalized surface quasigeostrophic equations, Arch. Ration. Mech. Anal., 220(3)(2016), 1209–1281.
- [22] F. de la Hoz, Z. Hassainia, T. Hmidi and J. Mateu, An analytical and numerical study of steady patches in the disc, Anal. PDE, 9(2016), no. 7, 1609–1670.
- [23] F. de la Hoz, T. Hmidi, J. Mateu and J. Verdera, Doubly connected V-states for the planar Euler equations, SIAM J. Math. Anal., 48(3)(2016), 1892–1928.
- [24] C. García, T.Hmidi and J.Soler, Non uniform Rotating Vortices and Periodic Orbits for the Two-Dimensional Euler Equations, Arch. Ration. Mech. Anal., 238 (2020), 929–1085.
- [25] L. Godard-Cadillac, Smooth traveling-wave solutions to the inviscid surface quasi-geostrophic equations, Preprint arXiv:2010.09289.
- [26] L. Godard-Cadillac, P. Gravejat and D. Smets, Co-rotating vortices with N fold symmetry for the inviscid surface quasi-geostrophic equation, arXiv: 2010.08194.
- [27] P. Gravejat and D. Smets, Smooth travelling-wave solutions to the inviscid surface quasigeostrophic equation, Int. Math. Res. Not., (6)2019, 1744–1757.
- [28] Z. Hassainia and T. Hmidi, On the V-states for the generalized quasi-geostrophic equations, Comm. Math. Phys., 337(2015), no. 1, 321–377.
- [29] I. Held, R. Pierrehumbert, S. Garner and K. Swanson, Surface quasi-geostrophic dynamics, J. Fluid Mech., 282 (1995), 1–20.
- [30] T. Hmidi, J.Mateu and J. Verdera, Boundary regularity of rotating vortex patches, Arch. Ration. Mech. Anal., 209(1)(2013), 171–208.
- [31] T. Hmidi, J. Mateu and J. Verdera, On rotating doubly connected vortices, J. Differential Equations, 258(4)(2015), 1395–1429.
- [32] T. Hmidi and J.Mateu, Bifurcation of rotating patches from Kirchhoff vortices, Discret, Contin. Dyn. Syst., 36(10)(2016), 5401–5422.
- [33] T. Hmidi and J. Mateu, Existence of corotating and counter-rotating vortex pairs for active scalar equations, Comm. Math. Phys., 350(2017), 699–747.
- [34] M. Juckes, Quasigeostrophic dynamics of the tropopause, J. Armos. Sci., (1994) 2756–2768.
- [35] G. Kirchhoff, Vorlesungen uber mathematische Physik, Leipzig, 1874.
- [36] A. Kiselev and F. Nazarov, A simple energy pump for the surface quasi-geostrophic equation, Nonlinear partial differential equations, volume 7 of Abel Symposia, Springer-Verlag, Berlin, Heidelberg, 2012, 175–179.

- [37] G. Lapeyre and P. Klein, Dynamics of the upper oceanic layers in terms of surface quasigeostrophic theory, J. Phys. Oceanogr., 36(2006), 165–176.
- [38] D. Li, Existence theorems for the 2D quasi-geostrophic equation with plane wave initial conditions, *Nonlinearity*, 22(2009), no. 7, 1639–1651.
- [39] F. Marchand, Existence and regularity of weak solutions to the quasi-geostrophic equations in the spaces  $L^p$  or  $\dot{H}^{-\frac{1}{2}}$ , Comm. Math. Phys., 277(2008), no. 1, 45–67.
- [40] A. J. Majda and A. L. Bertozzi, Vorticity and incompressible flow, Cambridge Texts in Applied Mathematics, Vol. 27., Cambridge University Press, 2002.
- [41] E.D. Rainville, Special Functions. Macmillan, New York, 1960
- [42] S. Resnick, Dynamical problems in non-linear advective partial differential equations. Ph.D. thesis, The University of Chicago, 1995.
- [43] M. Rosenzweig, Justification of the point vortex approximation for modified surface quasi-geostrophic equations, SIAM J. Math. Anal., 52(2020), no. 2, 1690–1728.
- [44] D. Smets and J. Van Schaftingen, Desingularization of vortices for the Euler equation, Arch. Rational Mech. Anal., 198(2010), 869–925.
- [45] B. Turkington, On steady vortex flow in two dimensions. I, II, Comm. Partial Differential Equations, 8(1983), 999–1030, 1031–1071.
- [46] B. Turkington, Corotating steady vortex flows with N-fold symmetry, Nonlinear Anal., 9(1985), no. 4, 351–369.
- [47] J. Wu, Quasi-geostrophic-type equations with initial data in Morrey spaces, Nonlinearity, 10(1997), no. 6, 1409–1420.
- [48] J. Wu, Solutions of the 2D quasi-geostrophic equation in Hölder spaces, Nonlinear Anal., 62(2005), no. 4, 579–594.
- [49] V. I. Yudovich, Non-stationnary flows of an ideal incompressible fluid, Zhurnal Vych Matematika, 3(1963), 1032–1106.

INSTITUTE OF APPLIED MATHEMATICS, CHINESE ACADEMY OF SCIENCES, BEIJING 100190, AND UNIVERSITY OF CHINESE ACADEMY OF SCIENCES, BEIJING 100049, P.R. CHINA *Email address*: dmcao@amt.ac.cn

INSTITUTE OF APPLIED MATHEMATICS, CHINESE ACADEMY OF SCIENCES, BEIJING 100190, AND UNIVERSITY OF CHINESE ACADEMY OF SCIENCES, BEIJING 100049, P.R. CHINA *Email address*: qinguolin18@mails.ucas.edu.cn

INSTITUTE OF APPLIED MATHEMATICS, CHINESE ACADEMY OF SCIENCES, BEIJING 100190, AND UNIVERSITY OF CHINESE ACADEMY OF SCIENCES, BEIJING 100049, P.R. CHINA *Email address*: zhanweicheng16@mails.ucas.ac.cn

INSTITUTE OF APPLIED MATHEMATICS, CHINESE ACADEMY OF SCIENCES, BEIJING 100190, AND UNIVERSITY OF CHINESE ACADEMY OF SCIENCES, BEIJING 100049, P.R. CHINA *Email address*: zouchangjun17@mails.ucas.ac.cn