

ON THE REGULARITY OF SCALAR TYPE SPECTRAL C_0 -SEMIGROUPS

MARAT V. MARKIN

ABSTRACT. We show that, for the C_0 -semigroups of *scalar type spectral operators*, a well-known necessary condition for the generation of *eventually norm-continuous* C_0 -semigroups, formulated exclusively in terms of the location of the spectrum of the semigroup's generator in the complex plane, is also sufficient and, in fact, characterizes the generators of *immediately norm-continuous* such semigroups.

Combining characterizations of the *immediate differentiability* and the *Gevrey ultradifferentiability* of scalar type spectral C_0 -semigroups with the *generation theorem*, found earlier by the author, we arrive at respective characterizations of the generation of such semigroups.

We further establish characterizations of the generation of *eventually differentiable* and *immediately compact* scalar type spectral C_0 -semigroups also in terms of the generator's spectrum and show that, for such semigroups, eventual compactness implies immediate.

All the obtained results are instantly transferred to the C_0 -semigroups of *normal operators*.

1. INTRODUCTION

While a Hille–Yosida type characterization of the generation of *eventually norm-continuous* C_0 -semigroups (see Preliminaries) appears to be still unestablished (see, e.g., [3, 11–14], for C_0 -semigroups on Hilbert spaces, see [10, 34] and [12]), well-known is the following necessary condition formulated exclusively in terms of the location of the spectrum of the semigroup's generator in the complex plane.

Theorem 1.1 (Necessary Condition for the Generation of Eventually Norm-Continuous C_0 -Semigroups [13, Theorem II.5.3]). *If A is a generator of an eventually norm continuous C_0 -semigroup on a complex Banach space, then, for any $b \in \mathbb{R}$, the set*

$$\{\lambda \in \sigma(A) \mid \operatorname{Re} \lambda \geq b\}$$

($\sigma(\cdot)$ is the spectrum of an operator) is bounded.

To a significant extent, the importance of eventually norm-continuous C_0 -semigroups is attributed the fact, that such a semigroup $\{T(t)\}_{t \geq 0}$ with generator A is subject to a *weak spectral mapping theorem*

$$(WSMT) \quad \sigma(T(t)) \setminus \{0\} = \overline{e^{t\sigma(A)}} \setminus \{0\}, \quad t \geq 0,$$

2020 *Mathematics Subject Classification.* Primary 47B40, 47D03, 47D60; Secondary 47B15, 47D06.

Key words and phrases. Scalar type spectral operator, C_0 -semigroup of linear operators.

($\bar{\cdot}$ is the *closure* of a set) [13, Examples V.2.2], in fact, a stronger *spectral mapping theorem*

$$(SMT) \quad \sigma(T(t)) \setminus \{0\} = e^{t\sigma(A)}, \quad t \geq 0,$$

[13, Theorem V.2.8, Proposition V.2.3], and with that the following *spectral bound equal growth bound condition*

$$(SBeGB) \quad s(A) = \omega_0,$$

where

$$s(A) := \sup \{\operatorname{Re} \lambda \mid \lambda \in \sigma(A)\} \quad (s(A) := -\infty \text{ if } \sigma(A) = \emptyset)$$

is the *spectral bound* of the generator and

$$(1.1) \quad \omega_0 := \inf \{ \omega \in \mathbb{R} \mid \exists M = M(\omega) \geq 1 : \|T(t)\| \leq M e^{\omega t}, \quad t \geq 0 \}$$

is the *growth bound* of the semigroup [13, Proposition V.2.3, Corollary V.2.10]. The asymptotic behavior of a C_0 -semigroup satisfying the *spectral bound equal growth bound condition* (SBeGB) is governed by the spectral bound of its generator and, in particular, is subject to a *generalized Lyapunov stability theorem* [13, Theorem V.3.6].

Eventually norm-continuous are C_0 -semigroups with certain regularity properties such as *eventual differentiability*, in particular *analyticity* or *uniform continuity*, or *eventual compactness* [29, 33] (see also [13]).

As is shown in [27], a *scalar type spectral* C_0 -semigroup $\{T(t)\}_{t \geq 0}$ with generator A is subject to the following *weak spectral mapping theorem*

$$\sigma(T(t)) = \overline{e^{t\sigma(A)}}, \quad t \geq 0,$$

[27, Theorem 4.1] (cf. [13, Corollary V.2.12]) and with that *spectral bound equal growth bound condition* (SBeGB) without any regularity requirement.

We show that, for the C_0 -semigroups of *scalar type spectral operators*, the above necessary condition is also sufficient and, in fact, characterizes the generators of *immediately norm continuous* such semigroups.

Combining the characterizations of the *immediate differentiability* and *Gevrey ultradifferentiability* of scalar type spectral C_0 -semigroups found in [17, 18, 23] (see also [21]) with the *generation theorem* [17, Proposition 3.1], we arrive at respective characterizations of the generation of such semigroups.

We further establish characterizations of the generation of *eventually differentiable* and *immediately compact* scalar type spectral C_0 -semigroups also in terms of the generator's spectrum and show that, for such semigroups, eventual compactness implies immediate.

All the obtained results are immediately transferred to the C_0 -semigroups of *normal operators*.

This work continues the series of papers [17, 18, 21, 23] (see also [19, 22]) on the generation of differentiable and Carleman, in particular Gevrey, ultradifferentiable scalar type spectral C_0 -semigroups.

2. PRELIMINARIES

Here, we concisely outline preliminaries necessary for our subsequent discourse.

2.1. C_0 -Semigroups.

Definition 2.1 (Norm-Continuity).

A C_0 -semigroup $\{T(t)\}_{t \geq 0}$ on a Banach space $(X, \|\cdot\|)$ is said to be

- *eventually norm-continuous* if there exists a $t_0 > 0$ such that the operator function

$$[t_0, \infty) \ni t \mapsto T(t) \in L(X)$$

($L(X)$ is the space of bounded linear operators on X) is continuous relative to the operator norm;

- *immediately norm-continuous* if the operator function

$$(0, \infty) \ni t \mapsto T(t) \in L(X)$$

is continuous relative to the operator norm;

- *uniformly continuous* if the operator function

$$[0, \infty) \ni t \mapsto T(t) \in L(X)$$

is continuous relative to the operator norm.

Definition 2.2 (Differentiability).

A C_0 -semigroup $\{T(t)\}_{t \geq 0}$ on a Banach space $(X, \|\cdot\|)$ is said to be

- *eventually differentiable* if there exists a $t_0 > 0$ such that the semigroup is strongly differentiable on $[t_0, \infty)$, i.e., the orbit maps

$$[0, \infty) \ni t \mapsto T(t)f \in X$$

are differentiable on $[t_0, \infty)$ for all $f \in X$;

- *immediately differentiable* if the semigroup is strongly differentiable on $(0, \infty)$, i.e., the orbit maps

$$[0, \infty) \ni t \mapsto T(t)f \in X$$

are differentiable on $(0, \infty)$ for all $f \in X$.

Remarks 2.1.

- If a C_0 -semigroup is strongly differentiable for $t \geq t_0$, it is *norm-continuous* for $t \geq t_0$ [29, Lemma 2.1] (see also [15, Theorem 10.3.5]).
- An immediately differentiable C_0 -semigroup $\{T(t)\}_{t \geq 0}$, often referred to as a C^∞ -semigroup, is *immediately norm-continuous*, and furthermore, *infinitely differentiable* on $(0, \infty)$ relative to the operator norm with

$$T^{(n)}(t) = A^n T(t), \quad t > 0,$$

[29, Lemma 2.1].

Definition 2.3 (Compactness).

A C_0 -semigroup $\{T(t)\}_{t \geq 0}$ on a Banach space $(X, \|\cdot\|)$ is said to be

- *eventually compact* if there exists a $t_0 > 0$ such that the operator $T(t_0)$ is compact, and hence, $T(t)$ is compact for all $t \geq t_0$ (see, e.g., [26]);

- *immediately compact* if the operator $T(t)$ is compact for all $t > 0$.

Remarks 2.2.

- If a C_0 -semigroup is compact for $t \geq t_0$, it is *norm-continuous* for $t \geq t_0$ [29, Theorem 3.1].

Thus, an immediately compact C_0 -semigroup is *immediately norm-continuous*.

- Observe that a *bounded* operator A in a complex infinite-dimensional Banach space cannot generate a *compact* C_0 -semigroup $\{T(t)\}_{t \geq 0}$, which would be *uniformly continuous*, and hence, the identity operator I would be *compact* as the uniform limit of $T(t)$ as $t \rightarrow 0+$. The latter would contradict the infinite-dimensionality of X (see, e.g., [26]).

For a linear operator A generating a C_0 -semigroup $\{T(t)\}_{t \geq 0}$ in a complex Banach space $(X, \|\cdot\|)$ the *abstract Cauchy problem*

$$(ACP) \quad \begin{cases} y'(t) = Ay(t), & t \geq 0, \\ y(0) = f \in X, \end{cases}$$

is *well-posed*, in particular *uniquely solvable*, in which case the *weak solutions* (also called *mild solutions*) of the associated *abstract evolution equation*

$$(AEE) \quad y'(t) = Ay(t), \quad t \geq 0,$$

are the orbit maps

$$y(t) = T(t)f, \quad t \geq 0,$$

with $f \in X$ [1, 12, 13], whereas the *classical* ones are the orbit maps with $f \in D(A)$.

2.2. Scalar Type Spectral Operators.

A *scalar type spectral operator* is a densely defined closed linear operator A in a complex Banach space $(X, \|\cdot\|)$ with strongly σ -additive *spectral measure* (the *resolution of the identity*) $E_A(\cdot)$, assigning to Borel sets of the complex plane \mathbb{C} projection operators on X and having the operator's *spectrum* $\sigma(A)$ as its *support*, and the associated *Borel operational calculus*, assigning to each Borel measurable function $F : \sigma(A) \rightarrow \overline{\mathbb{C}}$ ($\overline{\mathbb{C}} := \mathbb{C} \cup \{\infty\}$ is the extended complex plane) with $E_A(\{\lambda \in \mathbb{C} \mid F(\lambda) = \infty\}) = 0$ (here and whenever appropriate, 0 designates the *zero operator* on X) a scalar type spectral operator in X

$$F(A) := \int_{\sigma(A)} F(\lambda) dE_A(\lambda)$$

with

$$A = \int_{\sigma(A)} \lambda dE_A(\lambda)$$

[5, 6, 9].

In a complex finite-dimensional space, the scalar type spectral operators are those linear operators on the space, which furnish an *eigenbasis* for it (see, e.g., [6, 9])

and, in a complex Hilbert space, the scalar type spectral operators are those that are similar to the *normal operators* [32].

Due to its strong σ -additivity, the spectral measure is uniformly bounded, i.e.,

$$(2.2) \quad \exists M \geq 1 \forall \delta \in \mathcal{B}(\mathbb{C}) : \|E_A(\delta)\| \leq M$$

($\mathcal{B}(\mathbb{C})$ is the *Borel σ -algebra* on \mathbb{C}) (see, e.g., [7]).

Remark 2.3. Here and henceforth, we use the same notation $\|\cdot\|$ for the operator norm and adhere to this rather common economy of symbols also for the functional norm in the *dual space* X^* .

By [9, Theorem XVIII.2.11 (c)], for a Borel measurable function $F : \sigma(A) \rightarrow \overline{\mathbb{C}}$, the operator $F(A)$ is *bounded* iff

$$E_{A\text{-ess sup}} |F(\lambda)| < \infty,$$

in which case, we have the estimates

$$(2.3) \quad E_{A\text{-ess sup}} |F(\lambda)| \leq \|F(A)\| \leq 4M E_{A\text{-ess sup}} |F(\lambda)|,$$

where $M \geq 1$ is from (2.2).

In particular, this implies that a scalar type spectral operator is *bounded* iff its spectrum $\sigma(A)$ is a *bounded* set.

For arbitrary Borel measurable function $F : \mathbb{C} \rightarrow \mathbb{C}$, $f \in D(F(A))$, $g^* \in X^*$, and Borel set $\delta \subseteq \mathbb{C}$,

$$(2.4) \quad \int_{\delta} |F(\lambda)| dv(f, g^*, \lambda) \leq 4M \|E_A(\delta)F(A)f\| \|g^*\|,$$

where $v(f, g^*, \cdot)$ is the *total variation measure* of the complex-valued Borel measure $\langle E_A(\cdot)f, g^* \rangle$, for which

$$(2.5) \quad v(f, g^*, \mathbb{C}) = v(f, g^*, \sigma(A)) \leq 4M \|f\| \|g^*\|,$$

where $M \geq 1$ in (2.4) and (2.5) is from (2.2) (see, e.g., [18, 19]).

The following statement allows to characterize the domains of the Borel measurable functions of a scalar type spectral operator in terms of positive Borel measures.

Proposition 2.1 ([16, Proposition 3.1]).

Let A be a scalar type spectral operator in a complex Banach space X with spectral measure $E_A(\cdot)$ and $F : \sigma(A) \rightarrow \mathbb{C}$ be a Borel measurable function. Then $f \in D(F(A))$ iff

$$(i) \text{ for each } g^* \in X^*, \int_{\sigma(A)} |F(\lambda)| dv(f, g^*, \lambda) < \infty \text{ and}$$

$$(ii) \sup_{\{g^* \in X^* \mid \|g^*\|=1\}} \int_{\{\lambda \in \sigma(A) \mid |F(\lambda)| > n\}} |F(\lambda)| dv(f, g^*, \lambda) \rightarrow 0, \quad n \rightarrow \infty,$$

where X^* is the dual space and $v(f, g^*, \cdot)$ is the total variation measure of $\langle E_A(\cdot)f, g^* \rangle$.

In [17], it is shown that, for scalar type spectral C_0 -semigroups, the classical *Hille-Yosida-Feller-Miyadera-Phillips generation theorem* (see, e.g., [13, Theorems II.3.5 and II.3.8]) acquires the following inherently qualitative form exclusively in terms of the location of the generator's spectrum in the complex plane, void of any estimates on the generator's resolvent.

Theorem 2.1 (Generation Theorem [17, Proposition 3.1]).

A scalar type spectral operator A in a complex Banach space with spectral measure $E_A(\cdot)$ generates a C_0 -semigroup (of scalar type spectral operators) $\{T(t)\}_{t \geq 0}$ iff there exists an $\omega \in \mathbb{R}$ such that

$$\sigma(A) \subseteq \{\lambda \in \mathbb{C} \mid \operatorname{Re} \lambda \leq \omega\}$$

in which case

$$T(t) = e^{tA} := \int_{\sigma(A)} e^{t\lambda} dE_A(\lambda), \quad t \geq 0,$$

Furthermore, scalar type spectral C_0 -semigroups are generated by scalar type spectral operators [2, 28].

In [17, 18, 21, 23], characterizations of the *analyticity*, *immediate differentiability*, and *Carleman*, in particular *Gevrey ultradifferentiability*, of scalar type spectral C_0 -semigroups are found also purely in terms of the location of the generator's spectrum in the complex plane. In particular, the characterizations *analyticity* and *immediate differentiability*, similarly to the above *generation theorem*, are obtained from their general Hille–Yosida type counterparts [29, 33] (see also [4]) by dropping the corresponding estimates on the generator's resolvent.

2.3. Finer Spectrum Structure.

The spectrum $\sigma(A)$ of a *closed linear operator* A in a complex Banach space $(X, \|\cdot\|)$ is partitioned into the following pairwise disjoint subsets:

$$\begin{aligned} \sigma_p(A) &:= \{\lambda \in \mathbb{C} \mid A - \lambda I \text{ is not one-to-one, i.e., } \lambda \text{ is an eigenvalue of } A\}, \\ \sigma_c(A) &:= \left\{ \lambda \in \mathbb{C} \mid A - \lambda I \text{ is one-to-one and } R(A - \lambda I) \neq X, \text{ but } \overline{R(A - \lambda I)} = X \right\}, \\ \sigma_r(A) &:= \left\{ \lambda \in \mathbb{C} \mid A - \lambda I \text{ is one-to-one and } \overline{R(A - \lambda I)} \neq X \right\} \end{aligned}$$

($R(\cdot)$ is the *range* of an operator), called the *point*, *continuous*, and *residual spectrum* of A , respectively (see, e.g., [26]).

For a scalar type spectral operator A ,

$$(2.6) \quad \sigma_r(A) = \emptyset$$

[24, Corollary 3.1] (see also [20]).

3. NORM CONTINUITY

Theorem 3.1 (Characterization of the Generation of Immediately Norm-Continuous Scalar Type Spectral C_0 -Semigroups). *A scalar type spectral operator A in a complex Banach space generates an immediately norm continuous C_0 -semigroup (of scalar type spectral operators) iff, for any $b \in \mathbb{R}$, the set*

$$(3.7) \quad \{\lambda \in \sigma(A) \mid \operatorname{Re} \lambda \geq b\}$$

is bounded.

Proof.

“Only if” part immediately follows from the *Necessary Condition for the Generation of Eventually Norm-Continuous C_0 -Semigroups* (Theorem 1.1).

“If” part. Suppose that a scalar type spectral operator A in a complex Banach space with spectral measure $E_A(\cdot)$ satisfies condition (3.7). Then there exists an $\omega \in \mathbb{R}$ such that

$$\sigma(A) \subseteq \{\lambda \in \mathbb{C} \mid \operatorname{Re} \lambda \leq \omega\},$$

and hence, by the *Generation Theorem* (Theorem 2.1), A generates a scalar type spectral C_0 -semigroup

$$T(t) = e^{tA} := \int_{\sigma(A)} e^{t\lambda} dE_A(\lambda), \quad t \geq 0.$$

The premise also implies that, for each $n \in \mathbb{N}$ ($\mathbb{N} := \{1, 2, \dots\}$ is the set of *natural numbers*), by the properties of the Borel operational calculus [9, Theorem XVIII.2.11 (f)],

$$A_n := AE_A(\{\lambda \in \sigma(A) \mid \operatorname{Re} \lambda \geq -n\}) = \int_{\sigma(A)} \lambda \chi_{\{\lambda \in \sigma(A) \mid \operatorname{Re} \lambda \geq -n\}}(\lambda) dE_A(\lambda)$$

($\chi_\delta(\cdot)$ is the *characteristic function* of a set $\delta \subseteq \mathbb{C}$) is a *bounded* scalar type spectral operator since its spectrum $\sigma(A_n)$ is a bounded set (see Preliminaries). Indeed, by the *A.E. Weak Spectral Inclusion and Mapping Theorem* ([27, Theorem 3.1]), we have the inclusion

$$\sigma(A_n) \subseteq \{\lambda \in \sigma(A) \mid \operatorname{Re} \lambda \geq -n\} \cup \{0\}$$

(see also [25]).

Thus, for each $n \in \mathbb{N}$, A_n generates the *uniformly continuous* semigroup $\{e^{tA_n}\}_{t \geq 0}$ of its exponentials (see Preliminaries).

By the properties of the Borel operational calculus [9, Theorem XVIII.2.11],

$$\begin{aligned} e^{tA_n} &= \int_{\sigma(A)} e^{t\lambda \chi_{\{\lambda \in \sigma(A) \mid \operatorname{Re} \lambda \geq -n\}}(\lambda)} dE_A(\lambda) = \int_{\sigma(A)} e^{t\lambda} \chi_{\{\lambda \in \sigma(A) \mid \operatorname{Re} \lambda \geq -n\}}(\lambda) dE_A(\lambda) \\ &+ \int_{\sigma(A)} \chi_{\{\lambda \in \sigma(A) \mid \operatorname{Re} \lambda < -n\}}(\lambda) dE_A(\lambda) \\ &= e^{tA} E_A(\{\lambda \in \sigma(A) \mid \operatorname{Re} \lambda \geq -n\}) + E_A(\{\lambda \in \sigma(A) \mid \operatorname{Re} \lambda < -n\}), \quad t \geq 0. \end{aligned}$$

Hence, for each $n \in \mathbb{N}$, by the properties of the Borel operational calculus [9, Theorem XVIII.2.11 (g)], the operator function

$$\begin{aligned} [0, \infty) \ni t \mapsto E_A(\{\lambda \in \sigma(A) \mid \operatorname{Re} \lambda \geq -n\}) e^{tA_n} \\ = E_A(\{\lambda \in \sigma(A) \mid \operatorname{Re} \lambda \geq -n\}) e^{tA} E_A(\{\lambda \in \sigma(A) \mid \operatorname{Re} \lambda \geq -n\}) \\ = e^{tA} E_A(\{\lambda \in \sigma(A) \mid \operatorname{Re} \lambda \geq -n\}) \in L(X) \end{aligned}$$

is *continuous* relative to the *operator norm* on $[0, \infty)$.

For arbitrary $t_0 > 0$ and $n \in \mathbb{N}$, we have:

$$\begin{aligned} & \sup_{t \geq t_0} \|T(t) - E_A(\{\lambda \in \sigma(A) \mid \operatorname{Re} \lambda \geq -n\}) e^{tA_n}\| \\ &= \sup_{t \geq t_0} \|e^{tA} - E_A(\{\lambda \in \sigma(A) \mid \operatorname{Re} \lambda \geq -n\}) e^{tA_n}\| \\ &= \sup_{t \geq t_0} \|e^{tA} - e^{tA} E_A(\{\lambda \in \sigma(A) \mid \operatorname{Re} \lambda \geq -n\})\| \\ &= \sup_{t \geq t_0} \|e^{tA} E_A(\{\lambda \in \sigma(A) \mid \operatorname{Re} \lambda < -n\})\| \end{aligned}$$

by (2.3);

$$\begin{aligned} &= \sup_{t \geq t_0} \left\| \int_{\{\lambda \in \sigma(A) \mid \operatorname{Re} \lambda < -n\}} e^{t\lambda} dE_A(\lambda) \right\| \leq 4M \sup_{t \geq t_0} \sup_{\{\lambda \in \sigma(A) \mid \operatorname{Re} \lambda < -n\}} |e^{t\lambda}| \\ &= 4M \sup_{t \geq t_0} \sup_{\{\lambda \in \sigma(A) \mid \operatorname{Re} \lambda < -n\}} e^{t \operatorname{Re} \lambda} = 4M e^{-t_0 n} \rightarrow 0, \quad n \rightarrow \infty, \end{aligned}$$

where the constant $M \geq 1$ is from (2.2).

Thus, for any $t_0 > 0$, the sequence of operator functions

$$(E_A(\{\lambda \in \sigma(A) \mid \operatorname{Re} \lambda \geq -n\}) e^{tA_n})_{n \in \mathbb{N}},$$

continuous relative to the operator norm on $[0, \infty)$, converges *uniformly* to $T(\cdot)$ in the operator norm on $[t_0, \infty)$. This implies *immediate norm continuity* for the semigroup $\{T(t)\}_{t \geq 0}$ and completes the proof. \square

We directly obtain the following

Corollary 3.1 (Characterization of Immediately Norm-Continuous Scalar Type Spectral C_0 -Semigroups). *A C_0 -semigroup (of scalar type spectral operators) on a complex Banach space generated by a scalar type spectral operator A is immediately norm-continuous iff, for any $b \in \mathbb{R}$, the set*

$$\{\lambda \in \sigma(A) \mid \operatorname{Re} \lambda \geq b\}$$

is bounded.

By the *Necessary Condition for the Generation of Eventually Norm-Continuous C_0 -Semigroups* (Theorem 1.1), we also instantly arrive at

Corollary 3.2 (Equivalence of Eventual and Immediate Norm-Continuity).

If a scalar type spectral C_0 -semigroup in a complex Banach space is eventually norm-continuous, it is immediately norm-continuous.

4. DIFFERENTIABILITY

Combining the characterizations of the *immediate differentiability* and *Gevrey ultradifferentiability*, in particular *analyticity*, of scalar type spectral C_0 -semigroups found in [17, 18, 23] (see also [21]) with the *Generation Theorem* (Theorem 2.1) for such semigroups, we instantly obtain the subsequent characterizations of the generation of such semigroups.

Theorem 4.1 (Characterization of the Generation of Immediately Differentiable Scalar Type Spectral C_0 -Semigroups). *A scalar type spectral operator A in a complex Banach space generates an immediately differentiable C_0 -semigroup (of scalar type spectral operators) iff there exists an $\omega \in \mathbb{R}$ and, for any $b > 0$, there exists an $a \in \mathbb{R}$ such that*

$$\sigma(A) \subseteq \{\lambda \in \mathbb{C} \mid \operatorname{Re} \lambda \leq \min(\omega, a - b \ln |\operatorname{Im} \lambda|)\}.$$

See [18] for details (cf. [29]).

Theorem 4.2 (Characterization of the Generation of Roumieu-type Gevrey Ultradifferentiable Scalar Type Spectral C_0 -Semigroups). *Let $1 \leq \beta < \infty$. A scalar type spectral operator A in a complex Banach space generates a β th-order Roumieu-type Gevrey ultradifferentiable on $(0, \infty)$ C_0 -semigroup (of scalar type spectral operators) iff there exist $b > 0$ and $a \in \mathbb{R}$ such that*

$$\sigma(A) \subseteq \left\{ \lambda \in \mathbb{C} \mid \operatorname{Re} \lambda \leq a - b |\operatorname{Im} \lambda|^{1/\beta} \right\}.$$

See [18, 21] (cf. [4]).

In particular, for $\beta = 1$, we arrive at

Corollary 4.1 (Characterization of the Generation of Analytic Scalar Type Spectral C_0 -Semigroups). *A scalar type spectral operator A in a complex Banach space generates an analytic C_0 -semigroup (of scalar type spectral operators) iff there exist $b > 0$ and $a \in \mathbb{R}$ such that*

$$\sigma(A) \subseteq \{\lambda \in \mathbb{C} \mid \operatorname{Re} \lambda \leq a - b |\operatorname{Im} \lambda|\}.$$

See [17] (cf. [33]).

A similar characterization of the generation of the β th-order Beurling-type Gevrey ultradifferentiable on $(0, \infty)$ scalar type spectral C_0 -semigroups ($1 < \beta < \infty$) is found in [23].

For characterizations of the generation of the *Carleman ultradifferentiable* scalar type spectral C_0 -semigroups, see [21, 23].

We obtain the following characterization of the generation of *eventually differentiable* scalar type spectral C_0 -semigroups dropping the estimate on the resolvent of the semigroup's generator in its general Hille–Yosida type counterpart [29, Theorem 2.5].

Theorem 4.3 (Characterization of the Generation of Eventual Differentiable Scalar Type Spectral C_0 -Semigroups). *A scalar type spectral operator A in a complex Banach space generates an eventually differentiable C_0 -semigroup $\{T(t)\}_{t \geq 0}$ (of scalar type spectral operators) iff there exist $\omega \in \mathbb{R}$, $b > 0$, and $a \in \mathbb{R}$ such that*

$$(4.8) \quad \sigma(A) \subseteq \{\lambda \in \mathbb{C} \mid \operatorname{Re} \lambda \leq \min(\omega, a - b \ln |\operatorname{Im} \lambda|)\},$$

in which case the orbit maps

$$[0, \infty) \ni t \mapsto T(t)f \in X$$

are differentiable on $[b^{-1}, \infty)$ for all $f \in X$.

Proof.

“Only if” part immediately follows from [29, Theorem 2.5].

“If” part. Suppose that, for a scalar type spectral operator A in a complex Banach space $(X, \|\cdot\|)$ with spectral measure $E_A(\cdot)$, inclusion (4.8) holds with some $\omega \in \mathbb{R}$, $b > 0$, and $a \in \mathbb{R}$.

To prove that the C_0 -semigroup of exponentials $\{e^{tA}\}_{t \geq 0}$ generated by A (see Preliminaries) is eventually differentiable is to show that

$$(4.9) \quad \exists t_0 > 0 : e^{t_0 A} X \subseteq D(A)$$

[29].

For arbitrary $f \in X$ and $g^* \in X^*$,

$$\begin{aligned} \int_{\sigma(A)} |\lambda| e^{b^{-1} \operatorname{Re} \lambda} dv(f, g^*, \lambda) &= \int_{\{\lambda \in \sigma(A) \mid \operatorname{Re} \lambda < 0\}} |\lambda| e^{b^{-1} \operatorname{Re} \lambda} dv(f, g^*, \lambda) \\ &+ \int_{\{\lambda \in \sigma(A) \mid \operatorname{Re} \lambda \geq 0\}} |\lambda| e^{b^{-1} \operatorname{Re} \lambda} dv(f, g^*, \lambda). \end{aligned}$$

Indeed,

$$\int_{\{\lambda \in \sigma(A) \mid \operatorname{Re} \lambda \geq 0\}} |\lambda| e^{b^{-1} \operatorname{Re} \lambda} dv(f, g^*, \lambda)$$

due to the boundedness of the set

$$\{\lambda \in \sigma(A) \mid \operatorname{Re} \lambda \geq 0\},$$

which follows from inclusion (4.8), the continuity of the integrated function on \mathbb{C} , and the finiteness of the measure $v(f, g^*, \cdot)$.

Further, for any $f \in X$ and $g^* \in X^*$,

$$(4.10) \quad \int_{\{\lambda \in \sigma(A) \mid \operatorname{Re} \lambda < 0\}} |\lambda| e^{b^{-1} \operatorname{Re} \lambda} dv(f, g^*, \lambda) \leq \int_{\sigma(A)} (|\operatorname{Re} \lambda| + |\operatorname{Im} \lambda|) e^{b^{-1} \operatorname{Re} \lambda} dv(f, g^*, \lambda)$$

since, by (4.8), for $\lambda \in \{\lambda \in \sigma(A) \mid \operatorname{Re} \lambda < 0\}$,
 $\operatorname{Re} \lambda < 0$ and $|\operatorname{Im} \lambda| \leq e^{b^{-1}(a - \operatorname{Re} \lambda)}$;

$$\begin{aligned}
&\leq \int_{\{\lambda \in \sigma(A) \mid \operatorname{Re} \lambda < 0\}} \left[-\operatorname{Re} \lambda + e^{b^{-1}(a - \operatorname{Re} \lambda)} \right] e^{b^{-1} \operatorname{Re} \lambda} dv(f, g^*, \lambda) \\
&\hspace{15em} \text{since } x \leq e^x, x \geq 0; \\
&\leq \int_{\{\lambda \in \sigma(A) \mid \operatorname{Re} \lambda < 0\}} \left[b e^{b^{-1}(-\operatorname{Re} \lambda)} + e^{ab^{-1}} e^{b^{-1}(-\operatorname{Re} \lambda)} \right] e^{b^{-1} \operatorname{Re} \lambda} dv(f, g^*, \lambda) \\
&= \left[b + e^{ab^{-1}} \right] \int_{\{\lambda \in \sigma(A) \mid \operatorname{Re} \lambda < 0\}} 1 dv(f, g^*, \lambda) \leq \left[b + e^{ab^{-1}} \right] \int_{\sigma(A)} 1 dv(f, g^*, \lambda) \\
&= \left[b + e^{ab^{-1}} \right] v(f, g^*, \sigma(A)) \\
&\hspace{15em} \text{by (2.5);} \\
&\leq 4M \left[b + e^{ab^{-1}} \right] \|f\| \|g^*\|.
\end{aligned}$$

Also, for any $f \in X$,

$$\begin{aligned}
(4.11) \quad &\sup_{\{g^* \in X^* \mid \|g^*\|=1\}} \int_{\{\lambda \in \sigma(A) \mid |\lambda| e^{b^{-1} \operatorname{Re} \lambda} > n\}} |\lambda| e^{b^{-1} \operatorname{Re} \lambda} dv(f, g^*, \lambda) \\
&\leq \sup_{\{g^* \in X^* \mid \|g^*\|=1\}} \int_{\{\lambda \in \sigma(A) \mid \operatorname{Re} \lambda < 0, |\lambda| e^{b^{-1} \operatorname{Re} \lambda} > n\}} |\lambda| e^{b^{-1} \operatorname{Re} \lambda} dv(f, g^*, \lambda) \\
&\quad + \sup_{\{g^* \in X^* \mid \|g^*\|=1\}} \int_{\{\lambda \in \sigma(A) \mid \operatorname{Re} \lambda \geq 0, |\lambda| e^{b^{-1} \operatorname{Re} \lambda} > n\}} |\lambda| e^{b^{-1} \operatorname{Re} \lambda} dv(f, g^*, \lambda) \\
&\rightarrow 0, \quad n \rightarrow \infty.
\end{aligned}$$

Indeed, since, due to the boundedness of the set

$$\{\lambda \in \sigma(A) \mid \operatorname{Re} \lambda \geq 0\}$$

and the continuity of the integrated function on \mathbb{C} , the set

$$\left\{ \lambda \in \sigma(A) \mid \operatorname{Re} \lambda \geq 0, |\lambda| e^{b^{-1} \operatorname{Re} \lambda} > n \right\}$$

is *empty* for all sufficiently large $n \in \mathbb{N}$, we immediately infer that

$$\lim_{n \rightarrow \infty} \sup_{\{g^* \in X^* \mid \|g^*\|=1\}} \int_{\{\lambda \in \sigma(A) \mid \operatorname{Re} \lambda \geq 0, |\lambda| e^{b^{-1} \operatorname{Re} \lambda} > n\}} |\lambda| e^{b^{-1} \operatorname{Re} \lambda} dv(f, g^*, \lambda) = 0.$$

Finally, as in (4.10), for an arbitrary $f \in X$, we have:

$$\begin{aligned}
&\sup_{\{g^* \in X^* \mid \|g^*\|=1\}} \int_{\{\lambda \in \sigma(A) \mid \operatorname{Re} \lambda < 0, |\lambda| e^{b^{-1} \operatorname{Re} \lambda} > n\}} |\lambda| e^{b^{-1} \operatorname{Re} \lambda} dv(f, g^*, \lambda) \\
&\leq \sup_{\{g^* \in X^* \mid \|g^*\|=1\}} \left[b + e^{ab^{-1}} \right] \int_{\{\lambda \in \sigma(A) \mid \operatorname{Re} \lambda < 0, |\lambda| e^{b^{-1} \operatorname{Re} \lambda} > n\}} 1 dv(f, g^*, \lambda) \\
&\hspace{15em} \text{by (2.4);}
\end{aligned}$$

$$\begin{aligned}
&\leq \left[b + e^{ab^{-1}} \right] \cdot \\
&\quad \cdot \sup_{\{g^* \in X^* \mid \|g^*\|=1\}} 4M \|E_A(\{\lambda \in \sigma(A) \mid \operatorname{Re} \lambda < 0, |\lambda|e^{b^{-1}\operatorname{Re} \lambda} > n\})f\| \|g^*\| \\
&= 4M \left[b + e^{ab^{-1}} \right] \|E_A(\{\lambda \in \sigma(A) \mid \operatorname{Re} \lambda < 0, |\lambda|e^{b^{-1}\operatorname{Re} \lambda} > n\})f\| \\
&\quad \text{by the strong continuity of the spectral measure;} \\
&\rightarrow 4M \left[b + e^{ab^{-1}} \right] \|E_A(\emptyset)f\| = 0, \quad n \rightarrow \infty.
\end{aligned}$$

By Proposition 2.1 and the properties of the *operational calculus* (see [9, Theorem XVIII.2.11 (f)]), (4.10) and (4.11) jointly imply that,

$$f \in D\left(Ae^{b^{-1}A}\right),$$

and hence,

$$e^{b^{-1}A}f \in D(A),$$

i.e., inclusion (4.9) holds for $t_0 = b^{-1} > 0$, which implies that the orbit maps

$$[0, \infty) \ni t \mapsto T(t)f \in X$$

are differentiable on $[b^{-1}, \infty)$ for all $f \in X$. \square

Instantly, we obtain the following

Corollary 4.2 (Characterization of Eventually Differentiable Scalar Type Spectral C_0 -Semigroups). *A C_0 -semigroup $\{T(t)\}_{t \geq 0}$ (of scalar type spectral operators) on a complex Banach space generated by a scalar type spectral operator A is eventually differentiable iff there exist $b > 0$ and $a \in \mathbb{R}$ such that*

$$\sigma(A) \subseteq \{\lambda \in \mathbb{C} \mid \operatorname{Re} \lambda \leq a - b \ln |\operatorname{Im} \lambda|\},$$

in which case the orbit maps

$$[0, \infty) \ni t \mapsto T(t)f \in X$$

are differentiable on $[b^{-1}, \infty)$ for all $f \in X$.

Since an eventually differentiable C_0 -semigroup is eventually norm-continuous, by the *Equivalence of Eventual and Immediate Norm-Continuity* (Corollary 3.2), we also arrive at

Corollary 4.3 (Eventual Differentiability Implies Immediate Norm-Continuity). *If a scalar type spectral C_0 -semigroup is eventually differentiable, it is immediately norm-continuous.*

5. COMPACTNESS

Let us first prove the following characterization of compactness for the scalar type spectral operators emerging directly from the classical *Riesz-Schauder Theorem* (see, e.g., [26]).

Lemma 5.1 (Characterization of Compactness for Scalar Type Spectral Operators). *A scalar type spectral operator $A \neq 0$ in a complex infinite-dimensional Banach space is compact iff the spectrum $\sigma(A)$ of A consists of 0 and a nonempty countable set of nonzero eigenvalues with finite geometric multiplicities.*

If the set $\sigma(A) \setminus \{0\}$ is countably infinite, for its arbitrary arrangement $\{\lambda_n\}_{n=1}^\infty$ (i.e., $\mathbb{N} \ni n \mapsto \lambda_n \in \sigma(A) \setminus \{0\}$ is a bijection),

$$\lambda_n \rightarrow 0, \quad n \rightarrow \infty.$$

Proof.

“Only if” part follows directly from the *Riesz-Schauder Theorem* (see, e.g., [26]) in view of the fact that, for a scalar type spectral operator $A \neq 0$ with spectral measure $E_A(\cdot)$, due to the integral representation

$$A = \int_{\sigma(A)} \lambda dE_A(\lambda)$$

(see Preliminaries),

$$\sigma(A) \setminus \{0\} \neq \emptyset.$$

“If” part. Suppose that, a scalar type spectral operator $A \neq 0$ in a complex Banach space $(X, \|\cdot\|)$ with spectral measure $E_A(\cdot)$ is subject to the conditions of the “if” part. Then the spectrum $\sigma(A)$ is a *bounded* set, which implies that A is *bounded* (see Preliminaries).

If the set

$$\sigma(A) \setminus \{0\} \subseteq \sigma_p(A)$$

is finite, then

$$\sigma(A) \setminus \{0\} = \{\lambda_1, \dots, \lambda_n\}$$

with some $n \in \mathbb{N}$. This implies, that

$$A = \int_{\sigma(A)} \lambda dE_A(\lambda) = \sum_{k=1}^n \lambda_k E_A(\{\lambda_k\}).$$

Whence, in view of the fact that each λ_k , $k = 1, \dots, n$, is an eigenvalue with a finite geometric multiplicity, i.e., the subspace $E_A(\{\lambda_k\})X$, $k = 1, \dots, n$, is *finite-dimensional*, we infer that the operator A is of *finite rank*, and hence, compact.

Now suppose that the set

$$\sigma(A) \setminus \{0\} \subseteq \sigma_p(A)$$

is countably infinite and let $\{\lambda_n\}_{n=1}^\infty$ is its arbitrary arrangement. Then, by the premise,

$$\lambda_n \rightarrow 0, \quad n \rightarrow \infty.$$

Whence, by the properties of the Borel operational calculus [9], we infer that, for any $n \in \mathbb{N}$,

$$\left\| A - \sum_{k=1}^n \lambda_k E_A(\{\lambda_k\}) \right\| = \left\| \int_{\sigma(A) \setminus \{\lambda_1, \dots, \lambda_n\}} \lambda dE_A(\lambda) \right\|$$

by (2.3);

$$\leq 4M \sup_{k \geq n+1} |\lambda_k| \rightarrow 0, \quad n \rightarrow \infty.$$

Therefore, the operator A is *compact* as the *uniform limit* of the sequence

$$\left(\sum_{k=1}^n \lambda_k E_A(\{\lambda_k\}) \right)_{n \in \mathbb{N}}$$

of finite-rank operators (see, e.g., [26]).

With all possible cases considered, the proof of the “if” part, and hence, of the entire statement, is complete. \square

Theorem 5.1 (Characterization of the Generation of Immediately Compact Scalar Type Spectral C_0 -Semigroups). *A scalar type spectral operator $A \neq 0$ in a complex infinite-dimensional Banach space generates an immediately compact C_0 -semigroup iff $\sigma(A)$ is a countably infinite set of eigenvalues with finite geometric multiplicities, which has no finite limit points and is such that, for its arbitrary arrangement $\{\lambda_n\}_{n=1}^\infty$,*

$$\operatorname{Re} \lambda_n \rightarrow -\infty, \quad n \rightarrow \infty.$$

Proof.

“Only if” part. Suppose that, a scalar type spectral operator $A \neq 0$ in a complex Banach space $(X, \|\cdot\|)$ with spectral measure $E_A(\cdot)$ generates an *immediately compact* C_0 -semigroup $\{T(t)\}_{t \geq 0}$. Then A is necessarily *unbounded* (see Remarks 2.2), and hence, its spectrum $\sigma(A)$ is an unbounded set (see Preliminaries), which, in particular, is *infinite*.

Then, by [29, Theorem 3.3], the semigroup is *immediately norm-continuous* and the resolvent

$$R(\lambda, A) := (A - \lambda I)^{-1} = \int_{\sigma(A)} \frac{1}{\mu - \lambda} dE_A(\mu)$$

is *compact* for all $\lambda \in \rho(A) := \mathbb{C} \setminus \sigma(A)$, i.e., the operator A is *regular* in the sense of [31, Definition 1].

By [31, Lemma 1] and [13, Corollary V.1.15],

$$\sigma(A) = \sigma_p(A)$$

is a countable set with no finite limit points. This, in view of the fact that the spectrum $\sigma(A)$ is an *infinite* set, implies that, for its arbitrary arrangement $\{\lambda_n\}_{n=1}^\infty$,

$$\lambda_n \rightarrow \infty, \quad n \rightarrow \infty.$$

Since $T(t) = e^{tA}$, $t > 0$, (see Preliminaries), in particular $T(1) = e^A$, is a *compact* operator and by the *Point Spectrum Spectral Mapping Theorem* ([27, Theorem 4.2])

$$\sigma_p(T(1)) = e^{\sigma_p(A)}, \quad t \geq 0,$$

(cf. [13, Theorem V.2.6]), by the *Characterization of Compactness for Scalar Type Spectral Operators* (Lemma 5.1), we infer that

$$e^{\operatorname{Re} \lambda_n} = |e^{\lambda_n}| \rightarrow 0, \quad n \rightarrow \infty,$$

and hence,

$$\operatorname{Re} \lambda_n \rightarrow -\infty, \quad n \rightarrow \infty.$$

Furthermore, for an arbitrary $\mu \in \rho(A)$,

$$(5.12) \quad \lambda \in \sigma_p(A) \Leftrightarrow (\lambda - \mu)^{-1} \in \sigma_p(R(\mu, A)),$$

in which case

$$(5.13) \quad \ker(A - \lambda I) = \ker(R(\mu, A) - (\lambda - \mu)^{-1} I)$$

(see, e.g., [26]).

This, by the *Characterization of Compactness for Scalar Type Spectral Operators* (Lemma 5.1), implies that each eigenvalue of A has a finite geometric multiplicity.

“If” part. Suppose that, a scalar type spectral operator $A \neq 0$ in a complex Banach space $(X, \|\cdot\|)$ with spectral measure $E_A(\cdot)$ is subject to the conditions of the “if” part.

Since the spectrum $\sigma(A)$ is unbounded, the operator A is *unbounded* (see Preliminaries).

By the *Characterization of the Generation of Immediately Norm-Continuous Scalar Type Spectral C_0 -Semigroups* (Theorem 3.1), we infer that A generates an *immediately norm-continuous C_0 -semigroup*.

Furthermore, by the *Spectral Mapping Theorem for the Resolvent* ([13, Theorem V.1.13]), for an arbitrary $\lambda \in \rho(A)$,

$$\sigma(R(\lambda, A)) \setminus \{0\} = (\sigma(A) - \lambda)^{-1},$$

and hence, the spectrum $\sigma(R(\lambda, A))$ of $R(\lambda, A)$ consists of 0 (because A is unbounded) and a nonempty countably infinite set $\sigma(R(\lambda, A)) \setminus \{0\}$ such that, for its arbitrary arrangement $\{\lambda_n\}_{n=1}^\infty$,

$$\lambda_n \rightarrow 0, \quad n \rightarrow \infty.$$

By (5.12) and (5.13), we also infer that the set $\sigma(R(\lambda, A)) \setminus \{0\}$ consists of the eigenvalues of $R(\lambda, A)$ with finite geometric multiplicities.

By the *Characterization of Compactness for Scalar Type Spectral Operators* (Lemma 5.1), we conclude that the resolvent $R(\lambda, A)$ is *compact* for all $\lambda \in \rho(A)$, which along with the *immediate norm-continuity* of the C_0 -semigroup generated by A , by [29, Theorem 3.3], implies that the latter is *immediately compact*. \square

Instantly, we obtain

Corollary 5.1 (Characterization of Immediately Compact Scalar Type Spectral C_0 -Semigroups). *A C_0 -semigroup (of scalar type spectral operators) on complex infinite-dimensional Banach space generated by a scalar type spectral operator A is immediately compact iff $\sigma(A)$ is a countably infinite set of eigenvalues with finite geometric multiplicities, which has no finite limit points and is such that, for its arbitrary arrangement $\{\lambda_n\}_{n=1}^\infty$,*

$$\operatorname{Re} \lambda_n \rightarrow -\infty, \quad n \rightarrow \infty.$$

We also have the following analogue of the *Equivalence of Eventual and Immediate Norm-Continuity* (Corollary 3.2).

Corollary 5.2 (Equivalence of Eventual and Immediate Compactness).

If a scalar type spectral C_0 -semigroup in a complex infinite-dimensional Banach space is eventually compact, its is immediately compact.

Proof. Suppose that a C_0 -semigroup $\{T(t)\}_{t \geq 0}$ on a complex infinite-dimensional Banach space generated by a scalar type spectral operator $A \neq 0$ with spectral measure $E_A(\cdot)$ is *compact* for $t \geq t_0$ with some $t_0 > 0$.

Since $T(t) = e^{tA}$, $t \geq t_0$, (see Preliminaries) is a *compact operator*, in view of the *infinite dimensionality* of the underlying space, its inverse

$$T^{-1}(t) = e^{-tA} := \int_{\sigma(A)} e^{-t\lambda} dE_A(\lambda)$$

is necessarily *unbounded* (see, e.g., [26]), which implies that the scalar type spectral operator A has an *unbounded*, in particular *infinite*, spectrum $\sigma(A)$, and hence, is *unbounded* (see Preliminaries).

Since, by the *Weak Spectral Mapping Theorem* ([27, Theorem 4.1]),

$$(5.14) \quad \sigma(T(t)) = \overline{e^{t\sigma(A)}}, \quad t \geq 0,$$

we infer that $\sigma(A)$ has no finite limit points. Otherwise, by the *continuity* of the exponential function $e^{t\cdot}$, the spectrum $\sigma(T(t_0))$ of the *compact operator* $T(t_0)$ would have a *nonzero finite limit point*, which is impossible by the *Characterization of Compactness for Scalar Type Spectral Operators* (Lemma 5.1).

Being an infinite set in \mathbb{C} with no finite limit points, the spectrum $\sigma(A)$ is *countably infinite*.

Further, since $\sigma(A)$ is the *support* for the spectral measure $E_A(\cdot)$ (see Preliminaries) and consist of *isolated points*,

$$\forall \lambda \in \sigma(A) : E_A(\lambda) \neq 0,$$

and hence,

$$\sigma(A) = \sigma_p(A),$$

[9].

Since the operator $T(t) = e^{tA}$ is *compact* for each $t \geq t_0$, considering the $2\pi i$ -*periodicity* (i is the *imaginary unit*) of the exponential function e^{\cdot} , we infer that there exists a $t_1 \geq t_0$ such that the spectrum $\sigma(T(t_1))$ is an *infinite* set. Indeed, assuming that the set

$$\sigma(T(t_0)) = \overline{e^{t_0\sigma(A)}}$$

(see (5.14)) is *finite*, we conclude that the *infinite set* $\sigma(A)$ contains an infinitely subset consisting of the points of the form

$$\lambda_0 + it_0^{-1}2\pi n, \quad n \in \mathbb{Z},$$

with some $\lambda_0 \in \mathbb{C}$, mapped under $e^{t_0\cdot}$ into the same point of $\sigma(T(t_0))$. Then, for any $t > t_0$, these points are mapped under $e^{t\cdot}$ into infinitely many points of

$$\sigma(T(t)) = \overline{e^{t\sigma(A)}}$$

(see (5.14)).

Let $\{\lambda_n\}_{n=1}^\infty$ be an arbitrary arrangement of $\sigma(A)$. Then, by the *Characterization of Compactness for Scalar Type Spectral Operators* (Lemma 5.1),

$$e^{t_1 \operatorname{Re} \lambda_n} = |e^{t_1 \lambda_n}| \rightarrow 0, \quad n \rightarrow \infty.$$

Therefore,

$$\operatorname{Re} \lambda_n \rightarrow -\infty, \quad n \rightarrow \infty.$$

Furthermore, since, for arbitrary $\lambda \in \sigma_p(A)$ and $f \in \ker(A - \lambda I)$,

$$y(t) := e^{t\lambda} f, \quad t \geq 0,$$

is the *unique* eigenvalue solution of the Cauchy problem

$$\begin{cases} y'(t) = Ay(t), & t \geq 0, \\ y(0) = f, \end{cases}$$

(see Preliminaries), we infer that

$$T(t)f = e^{t\lambda} f, \quad t \geq 0,$$

and hence,

$$f \in \ker(T(t) - e^{t\lambda} I), \quad t \geq 0,$$

in particular, $f \in \ker(T(t_0) - e^{t_0\lambda} I)$, which implies that

$$\ker(A - \lambda I) \subseteq \ker(T(t_0) - e^{t_0\lambda} I).$$

Whence, since, by the *Characterization of Compactness for Scalar Type Spectral Operators* (Lemma 5.1), $\ker(T(t_0) - e^{t_0\lambda} I)$ is *finite-dimensional*, we conclude that each eigenvalue of A has a *finite geometric multiplicity*.

Thus, by the *Characterization of the Generation of Immediately Compact Scalar Type Spectral C_0 -Semigroups* (Theorem 5.1), the C_0 -semigroup generated by A is *immediately compact*. \square

6. THE CASE OF NORMAL C_0 -SEMIGROUPS

For the important particular case of the C_0 -semigroups of *normal operators*, we instantly obtain the following counterparts of the corresponding characterizations established for C_0 -semigroups of *scalar type spectral operators*.

Corollary 6.1 (Characterization of the Generation of Immediately Norm-Continuous Normal C_0 -Semigroups). *A normal operator A in a complex Hilbert space generates an immediately norm continuous C_0 -semigroup (of normal operators) iff, for any $b \in \mathbb{R}$, the set*

$$\{\lambda \in \sigma(A) \mid \operatorname{Re} \lambda \geq b\}$$

is bounded.

Remark 6.1. In view of the fact that, for a normal operator A in a complex Hilbert space,

$$\|R(\lambda, A)\| = \frac{1}{\operatorname{dist}(\lambda, \sigma(A))}, \quad \lambda \in \rho(A),$$

where

$$\text{dist}(\lambda, \sigma(A)) := \inf_{\mu \in \sigma(A)} |\mu - \lambda| > 0,$$

(see, e.g., [8, 30]), the prior corollary is consistent with the known characterizations of immediate and eventual norm continuity of C_0 -semigroups in complex Hilbert spaces ([34, Theorem 1] and [10, Theorem 4], respectively).

Corollary 6.2 (Characterization of Immediately Norm-Continuous Normal C_0 -Semigroups). *A C_0 -semigroup (of normal operators) on a complex Hilbert space generated by a normal operator A is immediately norm-continuous iff, for any $b \in \mathbb{R}$, the set*

$$\{\lambda \in \sigma(A) \mid \text{Re } \lambda \geq b\}$$

is bounded.

Corollary 6.3 (Equivalence of Eventual and Immediate Norm-Continuity). *If a normal C_0 -semigroup in a complex Hilbert space is eventually norm-continuous, it is immediately norm-continuous.*

Corollary 6.4 (Characterization of the Generation of Roumieu-type Gevrey Ultradifferentiable Normal C_0 -Semigroups). *Let $1 \leq \beta < \infty$. A normal operator A in a complex Hilbert space generates a β th-order Roumieu-type Gevrey ultradifferentiable on $(0, \infty)$ C_0 -semigroup (of normal operators) iff there exist $b > 0$ and $a \in \mathbb{R}$ such that*

$$\sigma(A) \subseteq \left\{ \lambda \in \mathbb{C} \mid \text{Re } \lambda \leq a - b |\text{Im } \lambda|^{1/\beta} \right\}.$$

Corollary 6.5 (Characterization of the Generation of Beurling-type Gevrey Ultradifferentiable Normal C_0 -Semigroups). *Let $1 < \beta < \infty$. A normal operator A in a complex Hilbert space generates a β th-order Beurling-type Gevrey ultradifferentiable on $(0, \infty)$ C_0 -semigroup (of normal operators) iff, for any $b > 0$, there exists an $a \in \mathbb{R}$ such that*

$$\sigma(A) \subseteq \left\{ \lambda \in \mathbb{C} \mid \text{Re } \lambda \leq a - b |\text{Im } \lambda|^{1/\beta} \right\}.$$

Cf. [23, Corollary 4.1].

Corollary 6.6 (Characterization of the Generation of Eventually Differentiable Normal C_0 -Semigroups). *A normal operator A in a complex Hilbert space generates an eventually differentiable C_0 -semigroup $\{T(t)\}_{t \geq 0}$ (of normal operators) iff there exist $\omega \in \mathbb{R}$, $a > 0$, and $b > 0$ such that*

$$\sigma(A) \subseteq \{\lambda \in \mathbb{C} \mid \text{Re } \lambda \leq \min(\omega, a - b \ln |\text{Im } \lambda|)\},$$

in which case the orbits

$$[0, \infty) \ni t \mapsto T(t)f \in X$$

are strongly differentiable on $[b^{-1}, \infty)$ for all $f \in X$.

Corollary 6.7 (Characterization of Eventually Differentiable Normal C_0 -Semigroups). *A C_0 -semigroup $\{T(t)\}_{t \geq 0}$ (of normal operators) on a complex Hilbert space generated by a normal operator A is eventually differentiable iff there exist $a > 0$ and $b > 0$ such that*

$$\sigma(A) \subseteq \{\lambda \in \mathbb{C} \mid \text{Re } \lambda \leq a - b \ln |\text{Im } \lambda|\},$$

in which case the orbits

$$[0, \infty) \ni t \mapsto T(t)f \in X$$

are strongly differentiable on $[b^{-1}, \infty)$ for all $f \in X$.

Corollary 6.8 (Characterization of Compactness for Normal Operators).

A normal operator $A \neq 0$ in a complex infinite-dimensional Hilbert space is compact iff the spectrum $\sigma(A)$ of A consists of 0 and a nonempty countable set of nonzero eigenvalues with finite geometric multiplicities.

If the set $\sigma(A) \setminus \{0\}$ is countably infinite, for its arbitrary arrangement $\{\lambda_n\}_{n=1}^\infty$ (i.e., $\mathbb{N} \ni n \mapsto \lambda_n \in \sigma(A) \setminus \{0\}$ is a bijection),

$$\lambda_n \rightarrow 0, \quad n \rightarrow \infty.$$

Corollary 6.9 (Characterization of the Generation of Immediately Compact Normal C_0 -Semigroups). A normal operator $A \neq 0$ in a complex infinite-dimensional Hilbert space generates an immediately compact C_0 -semigroup (of normal operators) iff $\sigma(A)$ is a countably infinite set of eigenvalues with finite geometric multiplicities, which has no finite limit points and is such that, for its arbitrary arrangement $\{\lambda_n\}_{n=1}^\infty$,

$$\operatorname{Re} \lambda_n \rightarrow -\infty, \quad n \rightarrow \infty.$$

Corollary 6.10 (Characterization of Immediately Compact Normal C_0 -Semigroups).

A C_0 -semigroup (of normal operators) on a complex infinite-dimensional Hilbert space generated by a normal operator A is immediately compact iff $\sigma(A)$ is a countably infinite set of eigenvalues with finite geometric multiplicities, which has no finite limit points and is such that, for its arbitrary arrangement $\{\lambda_n\}_{n=1}^\infty$,

$$\operatorname{Re} \lambda_n \rightarrow -\infty, \quad n \rightarrow \infty.$$

Corollary 6.11 (Equivalence of Eventual and Immediate Compactness).

If a normal C_0 -semigroup in a complex infinite-dimensional Hilbert space is eventually compact, it is immediately compact.

REFERENCES

- [1] J.M. Ball, *Strongly continuous semigroups, weak solutions, and the variation of constants formula*, Proc. Amer. Math. Soc. **63** (1977), no. 2, 101–107.
- [2] E. Berkson, *Semi-groups of scalar type operators and a theorem of Stone*, Illinois J. Math. **10** (1966), no. 2, 345–352.
- [3] O. Blasco and J. Martínez, *Norm continuity and related notions for semigroups on Banach spaces*, Arch. Math. **66** (1996), no. 6, 470–478.
- [4] M.G. Crandall and A. Pazy, *On the differentiability of weak solutions of a differential equations in Banach space*, J. Math. Mech. **18** (1969), no. 10, 1007–1016.
- [5] N. Dunford, *Spectral operators*, Pacific J. Math. **4** (1954), 321–354.
- [6] N. Dunford, *A survey of the theory of spectral operators*, Bull. Amer. Math. Soc. **64** (1958), 217–274.
- [7] N. Dunford and J.T. Schwartz with the assistance of W.G. Bade and R.G. Bartle, *Linear Operators. Part I: General Theory*, Interscience Publishers, New York, 1958.
- [8] ———, *Linear Operators. Part II: Spectral Theory. Self Adjoint Operators in Hilbert Space*, Interscience Publishers, New York, 1963.
- [9] ———, *Linear Operators. Part III: Spectral Operators*, Interscience Publishers, New York, 1971.

- [10] O. El-Mennaoui and K.-J. Engel, *On the characterization of eventually norm continuous semigroups in Hilbert spaces*, Arch. Math. **63** (1994), no. 5, 437–440.
- [11] ———, *Towards a characterization of eventually norm continuous semigroups on Banach spaces*, Quaestiones Math. **19** (1996), no. 1-2, 183–190.
- [12] K.-J. Engel and R. Nagel, *One-Parameter Semigroups for Linear Evolution Equations*, Graduate Texts in Mathematics, vol. 194, Springer-Verlag, New York, 2000.
- [13] ———, *A Short Course on Operator Semigroups*, Universitext, Springer, New York, 2006.
- [14] V. Goersmeyer and L. Weis, *Norm continuity of C_0 -semigroups*, Studia Math. **134** (1999), no. 2, 169–178.
- [15] E. Hille and R.S. Phillips, *Functional Analysis and Semi-groups*, American Mathematical Society Colloquium Publications, vol. 31, Amer. Math. Soc., Providence, RI, 1957.
- [16] M.V. Markin, *On an abstract evolution equation with a spectral operator of scalar type*, Int. J. Math. Math. Sci. **32** (2002), no. 9, 555–563.
- [17] ———, *A note on the spectral operators of scalar type and semigroups of bounded linear operators*, Ibid. **32** (2002), no. 10, 635–640.
- [18] ———, *On scalar type spectral operators, infinite differentiable and Gevrey ultradifferentiable C_0 -semigroups*, Ibid. **2004** (2004), no. 45, 2401–2422.
- [19] ———, *On the Carleman classes of vectors of a scalar type spectral operator*, Ibid. **2004** (2004), no. 60, 3219–3235.
- [20] ———, *A note on one decomposition of Banach spaces*, Methods Funct. Anal. Topology **12** (2006), no. 3, 254–257.
- [21] ———, *On scalar type spectral operators and Carleman ultradifferentiable C_0 -semigroups*, Ukrainian Math. J. **60** (2008), no. 9, 1418–1436.
- [22] ———, *On the Carleman ultradifferentiable vectors of a scalar type spectral operator*, Methods Funct. Anal. Topology **21** (2015), no. 4, 361–369.
- [23] ———, *On the generation of Beurling type Carleman ultradifferentiable C_0 -semigroups by scalar type spectral operators*, Ibid., **22** (2016), no. 2, 169–183.
- [24] ———, *On certain spectral features inherent to scalar type spectral operators*, Ibid. **23** (2017), no. 1, 60–65.
- [25] ———, *On the Gevrey ultradifferentiability of weak solutions of an abstract evolution equation with a scalar type spectral operator on the real axis*, Open Math. **18** (2020), no. 1, 1952–1976.
- [26] ———, *Elementary Operator Theory*, De Gruyter Graduate, Walter de Gruyter GmbH, Berlin/Boston, 2020.
- [27] ———, *On spectral mapping theorems and asymptotics of scalar type spectral C_0 -semigroups*, [arXiv:2002.09087](https://arxiv.org/abs/2002.09087).
- [28] T.V. Panchapagesan, *Semi-groups of scalar type operators in Banach spaces*, Pacific J. Math. **30** (1969), no. 2, 489–517.
- [29] A. Pazy, *On the differentiability and compactness of semi-groups of linear operators*, J. Math. Mech. **17** (1968), no. 12, 1131–1141.
- [30] A.I. Plesner, *Spectral Theory of Linear Operators*, Nauka, Moscow, 1965 (Russian).
- [31] J. Schwartz, *Perturbations of spectral operators, and applications. I. Bounded perturbations*, Pacific J. Math. **4** (1954), 415–458.
- [32] J. Wermer, *Commuting spectral measures on Hilbert space*, Ibid. **4** (1954), 355–361.
- [33] K. Yosida, *On the differentiability of semi-groups of linear operators*, Proc. Japan Acad. **34** (1958), no. 6, 337–340.
- [34] P. You, *Characteristic conditions for a C_0 -semigroup with continuity in the uniform operator topology for $t > 0$ in Hilbert space*, Proc. Amer. Math. Soc. **116** (1992), no. 4, 991–997.

DEPARTMENT OF MATHEMATICS
 CALIFORNIA STATE UNIVERSITY, FRESNO
 5245 N. BACKER AVENUE, M/S PB 108
 FRESNO, CA 93740-8001, USA

Email address: mmarkin@csufresno.edu