

WICK AND ANTI-WICK CHARACTERIZATIONS OF LINEAR OPERATORS ON SPACES OF POWER SERIES EXPANSIONS

JOACHIM TOFT

ABSTRACT. We study links between Wick, anti-Wick and analytic kernel operators on the Bargmann transform side. We consider classes of kernels, whose corresponding operators agree with the sets of linear and continuous operators on spaces of power series expansions, which are Bargmann images of Pilipović spaces. We show that in several situations, the sets of Wick and kernel operators with symbols and kernels in such classes agree. We also find suitable subclasses to these kernel classes, whose corresponding sets of Wick and anti-Wick operators agree. We also show ring, module and composition properties for such classes.

0. INTRODUCTION

A convenient way to represent linear operators acting on functions defined on the configuration space \mathbf{R}^d , is to apply the Bargmann transform. In this process, these operators are transformed into (analytic) integral operators T_K with kernels K , or as analytic pseudo-differential operators $\text{Op}_{\mathfrak{B}}(a)$ with symbols a , given by

$$T_K F(z) = \pi^{-d} \int_{\mathbf{C}^d} K(z, w) F(w) e^{-|w|^2} d\lambda(w) \quad (0.1)$$

and

$$\text{Op}_{\mathfrak{B}}(a) F(z) = \pi^{-d} \int_{\mathbf{C}^d} a(z, w) F(w) e^{(z-w, w)} d\lambda(w), \quad (0.2)$$

respectively, when F is a suitable analytic function on \mathbf{C}^d , where $d\lambda$ is the Lebesgue measure and (\cdot, \cdot) is the scalar product on \mathbf{C}^d . Here $K(z, w)$ and $a(z, w)$ are analytic functions or, more generally, power series expansions with respect to $(z, \bar{w}) \in \mathbf{C}^d \times \mathbf{C}^d \simeq \mathbf{C}^{2d}$. (See [19] or Section 1 for notations.) The operator $\text{Op}_{\mathfrak{B}}(a)$ is often called the Wick or Berezin operator with (Wick) symbol a .

An important subclass of Wick operators are the anti-Wick operators, $\text{Op}_{\mathfrak{B}}^{\text{aw}}(a)$ with suitable symbols $a(z, w)$ being analytic with respect to $(z, \bar{w}) \in \mathbf{C}^{2d}$, and given by

$$\text{Op}_{\mathfrak{B}}^{\text{aw}}(a) F(z) = \pi^{-d} \int_{\mathbf{C}^d} a(w, w) F(w) e^{(z-w, w)} d\lambda(w), \quad (0.3)$$

where F is a suitable analytic function on \mathbf{C}^d . An important feature for anti-Wick operators is that $\text{Op}_{\mathfrak{B}}^{\text{aw}}(a)$ is positive semi-definite when $a(z, z) \geq 0$ for every $z \in \mathbf{C}^d$, which is essential when performing certain types of energy estimates in quantum physics. Any similar transitions on positivity from symbols to Wick operators are in general not

2010 *Mathematics Subject Classification*. Primary: 32W25, 35S05, 32A17, 46F05, 42B35 Secondary: 32A25, 32A05.

Key words and phrases. Bargmann transform, Wick operators, anti-Wick operators, pseudo-differential operators, Toeplitz operators, Pilipović spaces, Gelfand-Shilov spaces.

true. On the other hand for suitable a we have the expansion formula

$$\text{Op}_{\mathfrak{W}}(a) = \sum_{|\alpha| < N} \frac{(-1)^{|\alpha|}}{\alpha!} \text{Op}_{\mathfrak{W}}^{\text{aw}}(b_\alpha) + \text{Op}_{\mathfrak{W}}(c_N) \quad \text{where} \quad b_\alpha = \partial_z^\alpha \bar{\partial}_w^\alpha a, \quad (0.4)$$

of Wick operators into a superposition of anti-Wick operators with error term $\text{Op}_{\mathfrak{W}}(c_N)$, deduced in [32]. Here the first term $\text{Op}_{\mathfrak{W}}^{\text{aw}}(b_0) = \text{Op}_{\mathfrak{W}}^{\text{aw}}(a)$ on the right-hand side, possess the convenient positivity transition from symbols to anti-Wick operators, while c_N is a linear combination of expressions of the form

$$\int_0^1 (1-t)^{|\alpha|-1} \partial_z^\alpha \bar{\partial}_w^\alpha a(w+t(z-w), w) dt, \quad |\alpha| = N.$$

In several situations, this term is dominating, while the reminder term $\text{Op}_{\mathfrak{W}}(c_N)$ is small compared to the other operators in the expansion. (See e. g. [28, 32].)

Wick and anti-Wick operators act between suitable spaces of analytic functions, which are the Bargmann transforms of spaces of functions or distributions on the configuration space \mathbf{R}^d . Some ideas on this approach were performed by G. C. Wick in [39]. Later on, F. Berezin improved and extended the theory [6, 7], where more general classes of operators were considered, compared to [39]. (See also [13, 28] for some other earlier results on Wick and anti-Wick operators.) An important part of the investigations in [6, 7, 39] concerns the links between Wick and anti-Wick operators. Especially in [6, 7] convenient formulae were established which explain such links in the case when the Wick and anti-Wick symbols are polynomials. These formulae are in some sense related to (0.4).

Some recent continuity properties for (analytic) integral, Wick and anti-Wick operators are obtained in [31, 32]. In [31], such continuity properties are obtained in the framework of the spaces of power series expansions,

$$\mathcal{A}_{0,s}(\mathbf{C}^d), \quad \mathcal{A}_s(\mathbf{C}^d), \quad \text{with duals} \quad \mathcal{A}'_s(\mathbf{C}^d), \quad \mathcal{A}'_{0,s}(\mathbf{C}^d), \quad (0.5)$$

respectively. These spaces are the Bargmann images of the so-called Pilipović spaces and their distribution (dual) spaces

$$\mathcal{H}_{0,s}(\mathbf{R}^d), \quad \mathcal{H}_s(\mathbf{R}^d), \quad \mathcal{H}'_s(\mathbf{R}^d) \quad \text{and} \quad \mathcal{H}'_{0,s}(\mathbf{R}^d), \quad (0.6)$$

a family of function and (ultra-)distribution spaces, which contains the Fourier invariant Gelfand-Shilov spaces and their ultra-distribution spaces, and which are thoroughly investigated in [35]. Some early ideas to some of the spaces $\mathcal{H}_s(\mathbf{R}^d)$, $\mathcal{H}_{0,s}(\mathbf{R}^d)$ and their duals is given by S. Pilipović in [24]. For example in [24] it is proved that $\mathcal{H}_s(\mathbf{R}^d)$ and $\mathcal{H}_{0,t}(\mathbf{R}^d)$ agree with the Fourier invariant Gelfand-Shilov spaces $\mathcal{S}_s(\mathbf{R}^d)$ and $\Sigma_t(\mathbf{R}^d)$, respectively, when $s \geq \frac{1}{2}$ and $t > \frac{1}{2}$. (See also Proposition 1.12 in Section 1 for a more complete view.)

In [31] it is remarked that the usual distribution kernel results for linear operators from $\mathcal{H}_s(\mathbf{R}^d)$ or $\mathcal{H}_{0,s}(\mathbf{R}^d)$ to their duals, give one-to-one correspondence of linear and continuous operators between $\mathcal{A}_s(\mathbf{C}^d)$ and $\mathcal{A}'_s(\mathbf{C}^d)$ and analytic integral operators with kernels in $\widehat{\mathcal{A}}_s(\mathbf{C}^{2d})$ or in $\widehat{\mathcal{A}}'_s(\mathbf{C}^{2d})$. Here $K(z, w) \in \widehat{\mathcal{A}}_s(\mathbf{C}^{2d})$, if and only if $K(z, \bar{w}) \in \mathcal{A}_s(\mathbf{C}^{2d})$, and similarly for other spaces. It is then proved that for suitable s , the set of Wick operators with symbols in $\widehat{\mathcal{A}}_s(\mathbf{C}^{2d})$ agrees with the set of integral operators with kernels in $\widehat{\mathcal{A}}'_s(\mathbf{C}^{2d})$. Consequently, for such s , there is a one-to-one correspondence of linear and continuous operators between $\mathcal{A}_s(\mathbf{C}^d)$ and $\mathcal{A}'_s(\mathbf{C}^d)$ and Wick operators with symbols in $\widehat{\mathcal{A}}_s(\mathbf{C}^{2d})$. Similar facts hold true with $\mathcal{A}_{0,s}$ in place of \mathcal{A}_s at each occurrence. (See Theorems 2.7 and 2.8 in [31].)

In the paper we find characterizations of the kernels to (analytic) integral operators and to symbols of Wick and anti-Wick operators in order for these operators should be continuous on the spaces in (0.5). This leads to introduction of certain classes of power series expansions

$$\widehat{\mathcal{B}}_{0,s}(\mathbf{C}^{2d}), \quad \widehat{\mathcal{B}}_s(\mathbf{C}^{2d}), \quad \widehat{\mathcal{B}}_s^*(\mathbf{C}^{2d}) \quad \text{and} \quad \widehat{\mathcal{B}}_{0,s}^*(\mathbf{C}^{2d}), \quad (0.7)$$

and slightly smaller subclasses

$$\widehat{\mathcal{C}}_{0,s}(\mathbf{C}^{2d}), \quad \widehat{\mathcal{C}}_s(\mathbf{C}^{2d}), \quad \widehat{\mathcal{C}}_s^*(\mathbf{C}^{2d}) \quad \text{and} \quad \widehat{\mathcal{C}}_{0,s}^*(\mathbf{C}^{2d}), \quad (0.8)$$

which are defined in similar and convenient ways as $\widehat{\mathcal{A}}_{0,s}(\mathbf{C}^{2d})$, $\widehat{\mathcal{A}}'_s(\mathbf{C}^{2d})$ and their duals (see Definition 1.5). In Section 2 we prove that $K \mapsto T_K$ is bijective from the spaces in (0.7) into the sets of continuous mappings on the respective spaces in (0.5). In particular,

$$\{T_K; K \in \widehat{\mathcal{B}}_s(\mathbf{C}^{2d})\} = \mathcal{L}(\mathcal{A}_s(\mathbf{C}^d))$$

(see Propositions 2.4 and 2.5). For $s < \frac{1}{2}$ we extend the last equality in Section 4 into

$$\begin{aligned} \text{Op}_{\mathfrak{W}}^{\text{aw}}(\mathcal{C}_s(\mathbf{C}^{2d})) &= \text{Op}_{\mathfrak{W}}(\mathcal{C}_s(\mathbf{C}^{2d})) \subseteq \text{Op}_{\mathfrak{W}}(\mathcal{B}_s(\mathbf{C}^{2d})) \\ &= \{T_K; K \in \widehat{\mathcal{B}}_s(\mathbf{C}^{2d})\} = \mathcal{L}(\mathcal{A}_s(\mathbf{C}^d)), \end{aligned} \quad (0.9)$$

and similarly for the other spaces in (0.5), (0.7) and (0.8) (see Theorems 4.8 and 4.9).

For certain s , the spaces in (0.7) and (0.8) are subspaces of $\widehat{A}(\mathbf{C}^{2d})$ and have simple characterizations in terms of convenient estimates on involved symbols. For example, if $s = b_\sigma$ with $\sigma > 1$, then

$$\mathcal{A}_s(\mathbf{C}^d) = \left\{ F \in A(\mathbf{C}^d); \exists r_0 > 0, |F(z)| \lesssim e^{r_0|z|^{\frac{2\sigma}{\sigma+1}}} \right\}$$

and

$$\widehat{\mathcal{B}}_s(\mathbf{C}^{2d}) = \left\{ a \in \widehat{A}(\mathbf{C}^{2d}); \forall r > 0, \exists r_0 > 0, |a(z, w)| \lesssim e^{r_0|z|^{\frac{2\sigma}{\sigma+1}} + r|w|^{\frac{2\sigma}{\sigma-1}}} \right\}.$$

(Cf. [3, 35].)

In several situations it is straight-forward to carry over the characterizations above to characterize linear and continuous operators acting on the spaces in (0.6), by applying the Bargmann transform and its inverse in suitable ways. Here the anti-Wick operator $\text{Op}_{\mathfrak{W}}^{\text{aw}}(a)$ corresponds to the Toeplitz operator $\text{Tp}_{\mathfrak{W}}(a)$, in the sense that

$$\text{Op}_{\mathfrak{W}}^{\text{aw}}(a) = \mathfrak{W}_d \circ \text{Tp}_{\mathfrak{W}}(a) \circ \mathfrak{W}_d^{-1},$$

when \mathfrak{W}_d is the Bargmann transform. Here $\text{Tp}_{\mathfrak{W}}(a)$ is defined by the formula

$$(\text{Tp}_{\mathfrak{W}}(a)f, g)_{L^2(\mathbf{R}^d)} = (\mathbf{a} \cdot V_\phi f, V_\phi g)_{L^2(\mathbf{R}^{2d})}, \quad \mathbf{a}(\sqrt{2}x, -\sqrt{2}\xi) = a(z, z), \quad z = x + i\xi,$$

where $V_\phi f$ is the short-time Fourier transform of f with respect to the window function

$$\phi(x) = \pi^{-\frac{d}{4}} e^{-\frac{1}{2}|x|^2}$$

(see Subsection 1.5). Especially it follows from (0.9) that

$$\text{Tp}_{\mathfrak{W}}(\mathcal{C}_s(\mathbf{C}^{2d})) \subseteq \mathcal{L}(\mathcal{H}_s(\mathbf{R}^d)), \quad s < \frac{1}{2}, \quad (0.10)$$

(see Theorem 5.1). Here we remark that in (0.10), the former class of operators is, in some sense close to the latter class. In particular several linear and continuous operators on $\mathcal{H}_s(\mathbf{R}^d)$ can be described as Toeplitz operators.

If instead $s \geq \frac{1}{2}$, then the closed links (0.9) and (0.10) between Wick, anti-Wick and kernel operators are violated. For example, it is shown in Subsection 4.4 that estimates, which seems rather natural, can not be performed when finding an anti-Wick operator to a Wick operator (see Remark 4.24).

The analysis behind (0.9) and the other related results are based on some continuity properties for linear and bilinear binomial type operators of independent interests, which acts on coefficients of power series expansions, given in [31] and in Section 3. The linear binomial operators are given by

$$(\mathcal{T}_{0,t}c)(\alpha, \beta) = \sum_{\gamma \leq \alpha, \beta} \left(\binom{\alpha}{\gamma} \binom{\beta}{\gamma} \right)^{1/2} t^{|\gamma|} c(\alpha - \gamma, \beta - \gamma),$$

and their formal adjoints

$$(\mathcal{T}_{0,t}^*c)(\alpha, \beta) = \sum_{\gamma \in \mathbf{N}^d} \left(\binom{\alpha + \gamma}{\gamma} \binom{\beta + \gamma}{\gamma} \right)^{1/2} t^{|\gamma|} c(\alpha + \gamma, \beta + \gamma),$$

when c is a suitable sequence on \mathbf{N}^{2d} and $t \in \mathbf{C}$ is fixed. Let $T_{\mathcal{A}}$ be the linear and bijective map which takes the sequence $\{c(\alpha, \beta)\}_{\alpha, \beta \in \mathbf{N}^d}$ into the power series expansion

$$\sum_{\alpha, \beta \in \mathbf{N}^d} c(\alpha, \beta) e_{\alpha}(z) e_{\beta}(\bar{w}), \quad e_{\alpha}(z) = \frac{z^{\alpha}}{\sqrt{\alpha!}}.$$

Also let

$$\ell'_{\mathcal{A},s}(\mathbf{N}^{2d}) \quad \text{and} \quad \ell'_{\mathcal{A},0,s}(\mathbf{N}^{2d})$$

be the counter images of $\widehat{\mathcal{A}}'_s(\mathbf{C}^{2d})$ and $\widehat{\mathcal{A}}'_{0,s}(\mathbf{C}^{2d})$, respectively, of $T_{\mathcal{A}}$, and let

$$\ell_{\mathcal{B},0,s}(\mathbf{N}^{2d}), \quad \ell_{\mathcal{B},s}(\mathbf{N}^{2d}), \quad \ell_{\mathcal{B},s}^*(\mathbf{N}^{2d}) \quad \text{and} \quad \ell_{\mathcal{B},0,s}^*(\mathbf{N}^{2d}) \quad (0.11)$$

and

$$\ell_{\mathcal{C},0,s}(\mathbf{N}^{2d}), \quad \ell_{\mathcal{C},s}(\mathbf{N}^{2d}), \quad \ell_{\mathcal{C},s}^*(\mathbf{N}^{2d}) \quad \text{and} \quad \ell_{\mathcal{C},0,s}^*(\mathbf{N}^{2d}) \quad (0.12)$$

be the counter images of the respective spaces in (0.7) and (0.8) under $T_{\mathcal{A}}$. Then it is proved in [31] that $\mathcal{T}_{0,t}$ is homeomorphic on $\ell'_{\mathcal{A},s}(\mathbf{N}^{2d})$ when $s < \frac{1}{2}$ and on $\ell'_{\mathcal{A},0,s}(\mathbf{N}^{2d})$ when $s \leq \frac{1}{2}$.

In Sections 3 and 4 we complete these mapping properties and prove that for similar s one has that $\mathcal{T}_{0,t}$ is homeomorphic on each of the spaces in (0.11), and that $\mathcal{T}_{0,t}^*$ is homeomorphic on each of the spaces in (0.12). By letting

$$\mathcal{T}_t = T_{\mathcal{A}} \circ \mathcal{T}_{0,t} \circ T_{\mathcal{A}}^{-1} \quad \text{and} \quad \mathcal{T}_t^* = T_{\mathcal{A}} \circ \mathcal{T}_{0,t}^* \circ T_{\mathcal{A}}^{-1}, \quad (0.13)$$

the relations of the form (0.9) then follows from

$$T_K = \text{Op}_{\mathfrak{W}}(a), \quad \text{Op}_{\mathfrak{W}}(\mathcal{T}_1^*a) = \text{Op}_{\mathfrak{W}}^{\text{aw}}(a) \quad \text{and} \quad \text{Op}_{\mathfrak{W}}^{\text{aw}}(\mathcal{T}_{-1}^*a) = \text{Op}_{\mathfrak{W}}(a),$$

when $K = \mathcal{T}_1 a$, or equivalently, when $a = \mathcal{T}_{-1} K$. Here we observe that the inverses of the operators in (0.13) are given by

$$\mathcal{T}_t^{-1} = \mathcal{T}_{-t} \quad \text{and} \quad (\mathcal{T}_t^*)^{-1} = \mathcal{T}_{-t}^*.$$

(See also (3.5) and (3.10) in Section 3 for exact formulae for \mathcal{T}_t and \mathcal{T}_t^* .)

The paper is organized as follows. In Section 1 we set the stage by providing necessary background notions and fixing the notation. It contains useful properties for weight

functions, Pilipović spaces, the Bargmann transform, Toeplitz operators, Wick and anti-Wick operators. Here we also introduce the spaces (0.7) and (0.8), and explain some of their basic properties.

In Section 2 we deduce kernel results. Especially we show that the spaces of linear and continuous operators on the spaces in (0.5) agree with sets of operators with kernels belonging to the spaces in (0.7).

In Section 3 we discuss continuity and bijectivity properties of the binomial operators $\mathcal{T}_{0,t}$ and $\mathcal{T}_{0,t}^*$, especially on the spaces in (0.11) and (0.12). We link these binomial operators to transitions between kernel, Wick and anti-Wick operators. Here we also introduce certain bilinear binomial operators which are linked to multiplications and compositions of Wick symbols, and prove continuity properties for such operators.

In Section 4 we apply the results and analysis from Section 3 to find continuity and identifications between kernel, Wick and anti-Wick operators, as well as continuity properties for compositions of Wick and anti-Wick symbols. For example, here we show relationships like (0.9).

In Section 5 we present some consequences of our investigations concerning linear operators which acts on functions and distributions on \mathbf{R}^d . For example we show that for suitable s , the sets of Toeplitz operators with symbols in (0.8) is in some sense close to the set of linear and continuous operators on $\mathcal{A}_s(\mathbf{C}^d)$, $\mathcal{A}_{0,s}(\mathbf{C}^d)$ and their duals.

In Section 6 we deduce mapping properties for $\mathcal{A}_s(\mathbf{C}^d)$, $\mathcal{A}_{0,s}(\mathbf{C}^d)$ and their duals, as well as the spaces in (0.7) and (0.8) under linear pullbacks and trace mappings. These results can be used to get alternative proofs of the multiplication properties in Section 4.

Finally some background analyses are presented in Appendices A and B. In Appendix A we present some identifications of the spaces in (0.5), (0.7) and some other spaces in terms of spaces of analytic functions. In Appendix B we deduce some basic formulae in the transition between Wick and anti-Wick symbols.

ACKNOWLEDGEMENT

The author was supported by Vetenskapsrådet (Swedish Science Council) within the project 2019-04890.

1. PRELIMINARIES

In this section we recall some facts on involved function and distribution spaces as well as on pseudo-differential operators. In Subsections 1.1 and 1.2 we give definitions and review some basic properties for Gelfand-Shilov spaces, Pilipović spaces and the spaces in (0.7). Thereafter we discuss in Subsection 1.2 the Bargmann transform and recall some topological spaces of entire functions or power series expansions on \mathbf{C}^d . The section is concluded with a review of some facts on pseudo-differential operators, Toeplitz operators, Wick and anti-Wick operators.

1.1. Gelfand-Shilov spaces. Let $0 < s \in \mathbf{R}$ be fixed. Then the (Fourier invariant) Gelfand-Shilov space $\mathcal{S}_s(\mathbf{R}^d)$ ($\Sigma_s(\mathbf{R}^d)$) of Roumieu type (Beurling type) consists of all $f \in C^\infty(\mathbf{R}^d)$ such that

$$\|f\|_{\mathcal{S}_{s,h}} \equiv \sup \left(\frac{|x^\alpha \partial^\beta f(x)|}{h^{|\alpha+\beta|} (\alpha! \beta!)^s} \right) \quad (1.1)$$

is finite for some $h > 0$ (for every $h > 0$). Here the supremum should be taken over all $\alpha, \beta \in \mathbf{N}^d$ and $x \in \mathbf{R}^d$. The semi-norms $\|\cdot\|_{\mathcal{S}_{s,h}^\sigma}$ induce an inductive limit topology for

the space $\mathcal{S}_s(\mathbf{R}^d)$ and projective limit topology for $\Sigma_s(\mathbf{R}^d)$, and the latter space becomes a Fréchet space under this topology.

The space $\mathcal{S}_s(\mathbf{R}^d) \neq \{0\}$ ($\Sigma_s(\mathbf{R}^d) \neq \{0\}$), if and only if $s \geq \frac{1}{2}$ ($s > \frac{1}{2}$).

The *Gelfand-Shilov distribution spaces* $\mathcal{S}'_s(\mathbf{R}^d)$ and $\Sigma'_s(\mathbf{R}^d)$ are the (strong) duals of $\mathcal{S}_s(\mathbf{R}^d)$ and $\Sigma_s(\mathbf{R}^d)$, respectively.

Let $(\cdot, \cdot)_{L^2}$ be the scalar product in $L^2(\mathbf{R}^d)$. Then the duality between $\mathcal{S}_s(\mathbf{R}^d)$ and $\mathcal{S}'_s(\mathbf{R}^d)$ can be obtained by straight-forward extensions of the restriction of $(\cdot, \cdot)_{L^2}$ to $\mathcal{S}_{1/2}(\mathbf{R}^d)$. In this setting we have

$$\begin{aligned} \mathcal{S}_{1/2}(\mathbf{R}^d) &\hookrightarrow \Sigma_s(\mathbf{R}^d) \hookrightarrow \mathcal{S}_s(\mathbf{R}^d) \hookrightarrow \Sigma_t(\mathbf{R}^d) \\ &\hookrightarrow \mathcal{S}(\mathbf{R}^d) \hookrightarrow \mathcal{S}'(\mathbf{R}^d) \hookrightarrow \Sigma'_t(\mathbf{R}^d) \\ &\hookrightarrow \mathcal{S}'_s(\mathbf{R}^d) \hookrightarrow \Sigma'_s(\mathbf{R}^d) \hookrightarrow \mathcal{S}'_{1/2}(\mathbf{R}^d), \quad \frac{1}{2} < s < t. \end{aligned} \tag{1.2}$$

Here and in what follows we use the notation $A \hookrightarrow B$ when the topological spaces A and B satisfy $A \subseteq B$ with continuous embeddings.

A convenient family of functions concerns the Hermite functions

$$h_\alpha(x) = \pi^{-\frac{d}{4}} (-1)^{|\alpha|} (2^{|\alpha|} \alpha!)^{-\frac{1}{2}} e^{\frac{1}{2}|x|^2} (\partial^\alpha e^{-|x|^2}), \quad \alpha \in \mathbf{N}^d.$$

The set of Hermite functions on \mathbf{R}^d is an orthonormal basis for $L^2(\mathbf{R}^d)$. It is also a basis for the Schwartz space and its distribution space, and for any Σ_s when $s > \frac{1}{2}$, \mathcal{S}_s when $s \geq \frac{1}{2}$ and their distribution spaces. They are also eigenfunctions to the Harmonic oscillator $H = H_d \equiv |x|^2 - \Delta$ and to the Fourier transform \mathcal{F} , given by

$$(\mathcal{F}f)(\xi) = \hat{f}(\xi) \equiv (2\pi)^{-\frac{d}{2}} \int_{\mathbf{R}^d} f(x) e^{-i\langle x, \xi \rangle} dx, \quad \xi \in \mathbf{R}^d,$$

when $f \in L^1(\mathbf{R}^d)$. Here $\langle \cdot, \cdot \rangle$ denotes the usual scalar product on \mathbf{R}^d . In fact, we have

$$H_d h_\alpha = (2|\alpha| + d) h_\alpha.$$

The Fourier transform \mathcal{F} extends uniquely to homeomorphisms on $\mathcal{S}'(\mathbf{R}^d)$, $\mathcal{S}'_s(\mathbf{R}^d)$ and on $\Sigma'_s(\mathbf{R}^d)$. Furthermore, \mathcal{F} restricts to homeomorphisms on $\mathcal{S}(\mathbf{R}^d)$, $\mathcal{S}_s(\mathbf{R}^d)$ and on $\Sigma_s(\mathbf{R}^d)$, and to a unitary operator on $L^2(\mathbf{R}^d)$. Similar facts hold true when the Fourier transform is replaced by a partial Fourier transform.

Gelfand-Shilov spaces and their distribution spaces can also be characterized by estimates of short-time Fourier transforms, (see e. g. [17, 29, 35]). They also obey convenient mapping properties under short-time Fourier transforms. More precisely, let $\phi \in \mathcal{S}(\mathbf{R}^d)$ be fixed. Then the *short-time Fourier transform* $V_\phi f$ of $f \in \mathcal{S}'(\mathbf{R}^d)$ with respect to the *window function* ϕ is the Gelfand-Shilov distribution on \mathbf{R}^{2d} , defined by

$$V_\phi f(x, \xi) = \mathcal{F}(f \overline{\phi(\cdot - x)})(\xi), \quad x, \xi \in \mathbf{R}^d. \tag{1.3}$$

If $f, \phi \in \mathcal{S}(\mathbf{R}^d)$, then it follows that

$$V_\phi f(x, \xi) = (2\pi)^{-\frac{d}{2}} \int_{\mathbf{R}^d} f(y) \overline{\phi(y - x)} e^{-i\langle y, \xi \rangle} dy, \quad x, \xi \in \mathbf{R}^d.$$

By [33, Theorem 2.3] it follows that the definition of the map $(f, \phi) \mapsto V_\phi f$ from $\mathcal{S}(\mathbf{R}^d) \times \mathcal{S}(\mathbf{R}^d)$ to $\mathcal{S}(\mathbf{R}^{2d})$ is uniquely extendable to a continuous map from $\mathcal{S}'_s(\mathbf{R}^d) \times \mathcal{S}'_s(\mathbf{R}^d)$ to $\mathcal{S}'_s(\mathbf{R}^{2d})$, and restricts to a continuous map from $\mathcal{S}_s(\mathbf{R}^d) \times \mathcal{S}_s(\mathbf{R}^d)$ to $\mathcal{S}_s(\mathbf{R}^{2d})$. The same conclusions hold with Σ_s in place of \mathcal{S}_s , at each place.

1.2. **Spaces of sequences, Hermite series and power series expansions.** Next we recall the definitions of topological vector spaces of Hermite series expansions, given in [35]. As in [35], it is convenient to use suitable extensions of \mathbf{R}_+ when indexing our spaces.

Definition 1.1. The sets \mathbf{R}_b , $\overline{\mathbf{R}}_b$, $\mathbf{R}_{b,\infty}$ and $\overline{\mathbf{R}}_{b,\infty}$ are given by

$$\mathbf{R}_b = \mathbf{R}_+ \bigcup_{\sigma>0} \{b_\sigma\}, \quad \overline{\mathbf{R}}_b = \mathbf{R}_b \cup \{0\}, \quad \mathbf{R}_{b,\infty} = \mathbf{R}_b \cup \{\infty\} \quad \text{and} \quad \overline{\mathbf{R}}_{b,\infty} = \overline{\mathbf{R}}_b \cup \{\infty\}.$$

Beside the usual ordering in $\overline{\mathbf{R}}_+ \cup \{\infty\}$, the elements b_σ in these subsets of $\overline{\mathbf{R}}_{b,\infty}$ are ordered by the relations $x_1 < b_{\sigma_1} < b_{\sigma_2} < x_2$, when $\sigma_1, \sigma_2, x_1, x_2 \in \mathbf{R}_+$ satisfy $x_1 < \frac{1}{2}$, $x_2 \geq \frac{1}{2}$ and $\sigma_1 < \sigma_2$.

In order for defining the sequence spaces we shall make use of the weight

$$\vartheta_{r,s}(\alpha) \equiv \begin{cases} 1 & \text{when } s = 0, |\alpha| \leq r, \\ \infty & \text{when } s = 0, |\alpha| > r, \\ e^{r|\alpha|^{\frac{1}{2s}}} & \text{when } s \in \mathbf{R}_+, \\ r^{|\alpha|} (\alpha!)^{\frac{1}{2\sigma}} & \text{when } s = b_\sigma, \\ \langle \alpha \rangle^r & \text{when } s = \infty, \quad \alpha \in \mathbf{N}^d, \end{cases} \quad (1.4)$$

and we observe that such weights obey estimates given in the following lemma.

Lemma 1.2. Let $\alpha, \beta \in \mathbf{N}^d$ and $s, \sigma \in \mathbf{R}_+$. Then

$$\begin{aligned} \vartheta_{c_1 r, s}(\alpha) \vartheta_{c_1 r, s}(\beta) &\leq \vartheta_{r, s}(\alpha + \beta) \leq \vartheta_{c_2 r, s}(\alpha) \vartheta_{c_2 r, s}(\beta), \\ c_1 &= \min(2^{\frac{1}{2s}-1}, 1), \quad c_2 = \max(2^{\frac{1}{2s}-1}, 1), \end{aligned} \quad (1.5)$$

and

$$\begin{aligned} \vartheta_{r, b_\sigma}(\alpha) \vartheta_{r, s}(\beta) &\leq \vartheta_{r, b_\sigma}(\alpha + \beta) \leq C \vartheta_{c r, b_\sigma}(\alpha) \vartheta_{c r, s}(\beta), \\ c &= 2^{\frac{1}{2\sigma}}, \quad C = \begin{cases} 1, & |\alpha + \beta| = 0, \\ 2^{-\frac{1}{2\sigma}}, & |\alpha + \beta| > 0. \end{cases} \end{aligned} \quad (1.6)$$

If $R \geq 0$ and in addition $s < \frac{1}{2}$, then it also holds

$$R^{|\alpha|} \lesssim \vartheta_{r, s}(\alpha) \quad \text{and} \quad R^{|\alpha|} \lesssim \vartheta_{r, b_\sigma}(\alpha), \quad \alpha \in \mathbf{N}^d. \quad (1.7)$$

Proof. Let $\theta = \frac{1}{2s}$. Then the inequalities in (1.5) follows from

$$c_1(s^\theta + t^\theta) \leq (s + t)^\theta \leq c_2(s^\theta + t^\theta), \quad s, t \geq 0.$$

The inequalities in (1.6) follows from

$$\vartheta_{r, b_\sigma}(\alpha + \beta) = \vartheta_{r, b_\sigma}(\alpha) \vartheta_{r, b_\sigma}(\beta) \binom{\alpha + \beta}{\alpha}^{\frac{1}{2\sigma}},$$

and that

$$1 \leq \binom{\alpha + \beta}{\alpha} \leq 2^{|\alpha + \beta| - 1}, \quad \text{when } |\alpha + \beta| \geq 1.$$

The estimates in (1.7) follows from

$$R^{|\alpha|} \lesssim \alpha!^{\frac{1}{2\sigma}} \lesssim R^{|\alpha|^{\frac{1}{2s}}}$$

when $R > 1$ and $s < \frac{1}{2}$, and the result follows. \square

Definition 1.3. Let $p \in (0, \infty]$, $s \in \overline{\mathbf{R}_{b,\infty}}$, $r \in \mathbf{R}$, ϑ be a map from \mathbf{N}^d to $\mathbf{R}_+ \cup \infty$, and let $\vartheta_{r,s}$ be as in (1.4).

(1) The set $\ell_{[\vartheta]}^p(\mathbf{N}^d)$ consists of all sequences $\{c(\alpha)\}_{\alpha \in \mathbf{N}^d} \subseteq \mathbf{C}$ such that

$$\|\{c(\alpha)\}_{\alpha \in \mathbf{N}^d}\|_{\ell_{[\vartheta]}^p} \equiv \|\{c(\alpha)\vartheta(\alpha)\}_{\alpha \in \mathbf{N}^d}\|_{\ell^p} < \infty;$$

(2) if $s < \infty$, then

$$\ell_s(\mathbf{N}^d) \equiv \text{ind} \lim_{r>0} \ell_{[\vartheta_{r,s}]}^p(\mathbf{N}^d) \quad \text{and} \quad \ell_s^*(\mathbf{N}^d) \equiv \text{proj} \lim_{r>0} \ell_{[1/\vartheta_{r,s}]}^p(\mathbf{N}^d); \quad (1.8)$$

(3) if $s > 0$, then

$$\ell_{0,s}(\mathbf{N}^d) \equiv \text{proj} \lim_{r>0} \ell_{[\vartheta_{r,s}]}^p(\mathbf{N}^d) \quad \text{and} \quad \ell_{0,s}^*(\mathbf{N}^d) \equiv \text{ind} \lim_{r>0} \ell_{[1/\vartheta_{r,s}]}^p(\mathbf{N}^d). \quad (1.9)$$

We observe that $\ell_{[\vartheta_{r,s}]}^p(\mathbf{N}^d)$ in Definition 1.3 is a quasi-Banach space under the quasi-norm $\|\cdot\|_{\ell_{[\vartheta_{r,s}]}^p}$ when $s > 0$. If in addition $p \geq 1$, then $\ell_{[\vartheta_{r,s}]}^p(\mathbf{N}^d)$ is a Banach space with norm $\|\cdot\|_{\ell_{[\vartheta_{r,s}]}^p}$.

In several situations we deal with two-parameter version of the spaces in Definition 1.3, which are denoted by

$$\ell_{\mathcal{A},0,(s_2,s_1)}(\Lambda), \quad \ell_{\mathcal{A},(s_2,s_1)}(\Lambda), \quad \ell_{\mathcal{A},0,(s_2,s_1)}^*(\Lambda), \quad \ell_{\mathcal{A},(s_2,s_1)}^*(\Lambda), \quad (1.10)$$

$$\ell_{\mathcal{B},0,(s_2,s_1)}(\Lambda), \quad \ell_{\mathcal{B},(s_2,s_1)}(\Lambda), \quad \ell_{\mathcal{B},0,(s_2,s_1)}^*(\Lambda), \quad \ell_{\mathcal{B},(s_2,s_1)}^*(\Lambda), \quad (1.11)$$

$$\ell_{\mathcal{C},0,(s_2,s_1)}(\Lambda), \quad \ell_{\mathcal{C},(s_2,s_1)}(\Lambda), \quad \ell_{\mathcal{C},0,(s_2,s_1)}^*(\Lambda) \quad \text{and} \quad \ell_{\mathcal{C},(s_2,s_1)}^*(\Lambda). \quad (1.12)$$

Definition 1.4. Let $\vartheta_{r,s}$ be as in (1.4), $s_1, s_2 \in \overline{\mathbf{R}_{b,\infty}}$, $\Lambda = \mathbf{N}^{d_2} \times \mathbf{N}^{d_1}$, $p \in (0, \infty]$ and let

$$\vartheta_{r,(s_2,s_1)}(\alpha) = \vartheta_{r,s_1}(\alpha_1)\vartheta_{r,s_2}(\alpha_2), \quad \alpha = (\alpha_2, \alpha_1), \quad \alpha_1 \in \mathbf{N}^{d_1}, \quad \alpha_2 \in \mathbf{N}^{d_2}.$$

Then the spaces in (1.10) are given by

$$\ell_{\mathcal{A},0,(s_2,s_1)}(\Lambda) \equiv \text{proj} \lim_{r>0} \ell_{[\vartheta_{r,(s_2,s_1)}]}^p(\Lambda), \quad s_1, s_2 > 0, \quad (1.13)$$

$$\ell_{\mathcal{A},(s_2,s_1)}(\Lambda) \equiv \text{ind} \lim_{r>0} \ell_{[\vartheta_{r,(s_2,s_1)}]}^p(\Lambda), \quad s_1, s_2 < \infty, \quad (1.14)$$

$$\ell_{\mathcal{A},0,(s_2,s_1)}^*(\Lambda) \equiv \text{ind} \lim_{r>0} \ell_{[1/\vartheta_{r,(s_2,s_1)}]}^p(\Lambda), \quad s_1, s_2 > 0, \quad (1.15)$$

and

$$\ell_{\mathcal{A},(s_2,s_1)}^*(\Lambda) \equiv \text{proj} \lim_{r>0} \ell_{[1/\vartheta_{r,(s_2,s_1)}]}^p(\Lambda), \quad s_1, s_2 < \infty. \quad (1.16)$$

Definition 1.5. Let $\vartheta_{r,s}$ be as in Definition 1.3, $s_1, s_2 \in \overline{\mathbf{R}_{b,\infty}}$, $\Lambda = \mathbf{N}^{d_2} \times \mathbf{N}^{d_1}$, $p \in (0, \infty]$ and let

$$\omega_{s_2,s_1;r_2,r_1}(\alpha) \equiv \frac{\vartheta_{r_2,s_2}(\alpha_2)}{\vartheta_{r_1,s_1}(\alpha_1)}, \quad \alpha = (\alpha_2, \alpha_1), \quad \alpha_1 \in \mathbf{N}^{d_1}, \quad \alpha_2 \in \mathbf{N}^{d_2}.$$

Then the spaces in (1.11) are given by

$$\ell_{\mathcal{B},0,(s_2,s_1)}(\Lambda) = \text{proj lim}_{r_2>0} \left(\text{ind lim}_{r_1>0} \left(\ell_{[\omega_{s_2,s_1};r_2,r_1]}^p(\Lambda) \right) \right), \quad s_1, s_2 > 0, \quad (1.17)$$

$$\ell_{\mathcal{B},(s_2,s_1)}(\Lambda) = \text{proj lim}_{r_1>0} \left(\text{ind lim}_{r_2>0} \left(\ell_{[\omega_{s_2,s_1};r_2,r_1]}^p(\Lambda) \right) \right), \quad s_1, s_2 < \infty, \quad (1.18)$$

$$\ell_{\mathcal{B},0,(s_2,s_1)}^*(\Lambda) = \text{proj lim}_{r_1>0} \left(\text{ind lim}_{r_2>0} \left(\ell_{[1/\omega_{s_2,s_1};r_2,r_1]}^p(\Lambda) \right) \right), \quad s_1, s_2 > 0. \quad (1.19)$$

and

$$\ell_{\mathcal{B},(s_2,s_1)}^*(\Lambda) = \text{proj lim}_{r_2>0} \left(\text{ind lim}_{r_1>0} \left(\ell_{[1/\omega_{s_2,s_1};r_2,r_1]}^p(\Lambda) \right) \right), \quad s_1, s_2 < \infty. \quad (1.20)$$

The spaces in (1.12) for admissible s_1, s_2 are given by the right-hand sides of (1.17)–(1.20), after the orders of inductive and projective limits are swapped.

Remark 1.6. By playing with r, r_1 and r_2 it follows that the topological vector spaces in (1.8), (1.9), (1.13)–(1.16), (1.11) and (1.12) are independent of $p \in (0, \infty]$.

For conveniency we also complete the spaces in Definitions 1.3, 1.4 and 1.5 with the following.

Definition 1.7. Let $s_1, s_2 \in \overline{\mathbf{R}_{b,\infty}}$ and $\Lambda = \mathbf{N}^{d_2} \times \mathbf{N}^{d_1}$.

- (1) $\ell_\infty(\mathbf{N}^d)$ and $\ell_\infty^*(\mathbf{N}^d)$ are given by (1.8) with $s = \infty$ and $p = 2$;
- (2) if $s_1 = 0$ or $s_2 = 0$, then

$$\begin{aligned} \ell_{0,0}(\Lambda) &= \ell_{0,0}^*(\Lambda) = \ell_{\mathcal{A},0,(s_2,s_1)}(\Lambda) = \ell_{\mathcal{B},0,(s_2,s_1)}(\Lambda) = \ell_{\mathcal{C},0,(s_2,s_1)}(\Lambda) \\ &= \ell_{\mathcal{A},0,(s_2,s_1)}^*(\Lambda) = \ell_{\mathcal{B},0,(s_2,s_1)}^*(\Lambda) = \ell_{\mathcal{C},0,(s_2,s_1)}^*(\Lambda) \equiv \{0\} = \{\{0\}_{\alpha \in \Lambda}\}; \end{aligned}$$

- (3) if $s_1 = \infty$ or $s_2 = \infty$, then

$$\ell_{\mathcal{A},(s_2,s_1)}(\Lambda), \quad \ell_{\mathcal{A},(s_2,s_1)}^*(\Lambda), \quad \ell_{\mathcal{B},(s_2,s_1)}(\Lambda) \quad \text{and} \quad \ell_{\mathcal{B},(s_2,s_1)}^*(\Lambda)$$

are defined by (1.14), (1.16), (1.18) and (1.19) with $p = 2$, and $\ell_{\mathcal{C},s_1,s_2}(\Lambda)$ and $\ell_{\mathcal{C},s_1,s_2}^*(\Lambda)$ are defined by (1.18) and (1.19) with $p = 2$ after the orders of inductive and projective limits are swapped.

Remark 1.8. By the definitions it follows that (1.8), (1.9), (1.13)–(1.16) and (1.17)–(1.19) hold true with \bigcup and \bigcap in place of ind lim and proj lim , respectively, at each occurrence.

We observe that the following holds true:

- (1) the space $\ell_0^*(\mathbf{N}^d)$ is the set of all sequences $\{c(\alpha)\}_{\alpha \in \mathbf{N}^d} \subseteq \mathbf{C}$ on \mathbf{N}^d , and that $\ell_0(\mathbf{N}^d)$ is the set of all such sequences such that $c(\alpha) \neq 0$ for at most finite numbers of α . In similar ways, the condition $s_1 = 0$ or $s_2 = 0$ impose support restrictions of the elements in the spaces (1.10)–(1.12);
- (2) the spaces in (1.10) are complete Hausdorff topological vector spaces, and that $\ell_{\mathcal{A},0,(s_2,s_1)}(\Lambda)$ and $\ell_{\mathcal{A},(s_2,s_1)}^*(\Lambda)$ are Fréchet spaces. It holds that $\ell_{\mathcal{A},(s,s)} = \ell_s$, $\ell_{\mathcal{A},0,(s,s)} = \ell_{0,s}$;
- (3) $\ell_0(\Lambda)$ is dense in all of the spaces in (1.10)–(1.12).

Remark 1.9. Let $s_1, s_2 \in \overline{\mathbf{R}}_b$ and $\Lambda = \mathbf{N}^{d_2} \times \mathbf{N}^{d_1}$. Then it follows by straight-forward computations that the $\ell^2(\Lambda)$ form $(\cdot, \cdot)_{\ell^2(\Lambda)}$ on $\ell_0(\Lambda)$ is uniquely extendable to continuous mappings from

$$\begin{aligned} \ell_{\mathcal{A},(s_2,s_1)}^*(\Lambda) \times \ell_{\mathcal{A},(s_2,s_1)}(\Lambda), & \quad \ell_{\mathcal{C},(s_2,s_1)}^*(\Lambda) \times \ell_{\mathcal{B},(s_2,s_1)}(\Lambda) \\ \text{and} & \quad \ell_{\mathcal{B},(s_2,s_1)}^*(\Lambda) \times \ell_{\mathcal{C},(s_2,s_1)}(\Lambda) \end{aligned} \quad (1.21)$$

to \mathbf{C} . These mappings constitutes dualities in the sense that they are non-degenerate, i. e. if $c_1 \in \ell_{\mathcal{A},(s_2,s_1)}^*(\Lambda) \setminus 0$ and $c_2 \in \ell_{\mathcal{A},(s_2,s_1)}(\Lambda) \setminus 0$, then

$$\{(c_1, c)_{\ell^2}; c \in \ell_{\mathcal{A},(s_2,s_1)}(\Lambda)\} \neq 0 \quad \text{and} \quad \{(c, c_2)_{\ell^2}; c \in \ell_{\mathcal{A},(s_2,s_1)}^*(\Lambda)\} \neq 0,$$

and similarly for the other pairs of spaces in (1.21). We also observe that $\ell_{\mathcal{A},(s_2,s_1)}^*(\Lambda)$ and $\ell_{\mathcal{A},(s_2,s_1)}(\Lambda)$ are (strong) duals to each others, through the form $(\cdot, \cdot)_{\ell^2(\Lambda)}$.

If instead $s_1, s_2 \in \mathbf{R}_{b,\infty}$, then the same holds true with $\ell_{\mathcal{A},0,(s_2,s_1)}$, $\ell_{\mathcal{B},0,(s_2,s_1)}$ and $\ell_{\mathcal{C},0,(s_2,s_1)}$ in place of $\ell_{\mathcal{A},(s_2,s_1)}$, $\ell_{\mathcal{B},(s_2,s_1)}$ and $\ell_{\mathcal{C},(s_2,s_1)}$, respectively, at each occurrence.

Next we consider, in similar ways as in [35], spaces of formal Hermite series expansions

$$f = \sum_{\alpha \in \mathbf{N}^d} c(f; \alpha) h_\alpha, \quad \{c(\alpha)\}_{\alpha \in \mathbf{N}^d} \in \ell_0^*(\mathbf{N}^d). \quad (1.22)$$

and spaces of formal power series expansions

$$F = \sum_{\alpha \in \mathbf{N}^d} c(F; \alpha) e_\alpha, \quad \{c(\alpha)\}_{\alpha \in \mathbf{N}^d} \in \ell_0^*(\mathbf{N}^d). \quad (1.23)$$

which correspond to

$$\ell_{0,s}(\mathbf{N}^d), \quad \ell_s(\mathbf{N}^d), \quad \ell_{0,s}^*(\mathbf{N}^d) \quad \text{and} \quad \ell_s^*(\mathbf{N}^d). \quad (1.24)$$

Here

$$e_\alpha(z) \equiv \frac{z^\alpha}{\sqrt{\alpha!}}, \quad z \in \mathbf{C}^d, \quad \alpha \in \mathbf{N}^d. \quad (1.25)$$

We consider the mappings

$$T_{\mathcal{H}} : \{c(\alpha)\}_{\alpha \in \mathbf{N}^d} \mapsto \sum_{\alpha \in \mathbf{N}^d} c(\alpha) h_\alpha \quad \text{and} \quad T_{\mathcal{A}} : \{c(\alpha)\}_{\alpha \in \mathbf{N}^d} \mapsto \sum_{\alpha \in \mathbf{N}^d} c(\alpha) e_\alpha \quad (1.26)$$

between sequences, and formal Hermite series and power series expansions.

Definition 1.10. If $s \in \overline{\mathbf{R}}_{b,\infty}$, then

$$\mathcal{H}_{0,s}(\mathbf{R}^d), \quad \mathcal{H}_s(\mathbf{R}^d), \quad \mathcal{H}_{0,s}^*(\mathbf{R}^d) \quad \text{and} \quad \mathcal{H}_s^*(\mathbf{R}^d), \quad (1.27)$$

and

$$\mathcal{A}_{0,s}(\mathbf{C}^d), \quad \mathcal{A}_s(\mathbf{C}^d), \quad \mathcal{A}_{0,s}^*(\mathbf{C}^d) \quad \text{and} \quad \mathcal{A}_s^*(\mathbf{C}^d), \quad (1.28)$$

are the images of $T_{\mathcal{H}}$ and $T_{\mathcal{A}}$ respectively in (1.26) of corresponding spaces in (1.24). The topologies of the spaces in (1.27) and (1.28) are inherited from the corresponding spaces in (1.24).

By Remark 1.9 it follows that the latter spaces in (1.27) are the (strong) duals of the former spaces with respect to unique extensions of the form $(\cdot, \cdot)_{L^2}$ on $\mathcal{H}_0(\mathbf{R}^d)$. That is,

$$\mathcal{H}'_s(\mathbf{R}^d) = \mathcal{H}_s^*(\mathbf{R}^d) \quad \text{and} \quad \mathcal{H}'_{0,s}(\mathbf{R}^d) = \mathcal{H}_{0,s}^*(\mathbf{R}^d). \quad (1.29)$$

In the same way as in (1.28) we let

$$\mathcal{A}_{0,(s_2,s_1)}(W), \quad \mathcal{A}_{(s_2,s_1)}(W), \quad \mathcal{A}_{0,(s_2,s_1)}^*(W) \quad \text{and} \quad \mathcal{A}_{(s_2,s_1)}^*(W) \quad (1.28)'$$

be the images of the spaces (1.10) under the map $T_{\mathcal{A}}$ when $W = \mathbf{C}^{d_2} \times \mathbf{C}^{d_1}$, $\Lambda = \mathbf{N}^{d_2} \times \mathbf{N}^{d_1}$ and $s_1, s_2 \in \overline{\mathbf{R}}_{b, \infty}$.

Since locally absolutely convergent power series expansions can be identified with entire functions, several of the spaces in (1.28) are identified with topological vector spaces contained in $A(\mathbf{C}^d)$ or $A(W)$. (See [35], Theorem A.1 in Appendix A and the introduction.) Here $A(\Omega)$ is the set of all (complex valued) functions which are analytic in the open set $\Omega \subseteq \mathbf{C}^d$. If $\Omega_0 \subseteq \mathbf{C}^d$ is arbitrary, then $A(\Omega_0) = \cup A(\Omega)$, where the union is taken over all open sets $\Omega \subseteq \mathbf{C}^d$ such that $\Omega_0 \subseteq \Omega$.

We recall that $f \in \mathcal{S}(\mathbf{R}^d)$ if and only if (1.22) holds with

$$|c(\alpha)| \lesssim \langle \alpha \rangle^{-N},$$

for every $N \geq 0$. That is, $\mathcal{S}(\mathbf{R}^d) = \mathcal{H}_{0, \infty}(\mathbf{R}^d)$. In the same way, $\mathcal{S}'(\mathbf{R}^d) = \mathcal{H}_{0, \infty}^*(\mathbf{R}^d)$, and $f \in L^2(\mathbf{R}^d)$, if and only if $\{c(\alpha)\}_{\alpha \in \mathbf{N}^d} \in \ell^2(\mathbf{N}^d)$ (cf. e. g. [26]). In particular it follows from the definitions that the inclusions

$$\begin{aligned} \mathcal{H}_0(\mathbf{R}^d) &\hookrightarrow \mathcal{H}_{0, s}(\mathbf{R}^d) \hookrightarrow \mathcal{H}_s(\mathbf{R}^d) \hookrightarrow \mathcal{H}_{0, t}(\mathbf{R}^d) \\ &\hookrightarrow \mathcal{H}_{0, t}^*(\mathbf{R}^d) \hookrightarrow \mathcal{H}_s^*(\mathbf{R}^d) \hookrightarrow \mathcal{H}_{0, s}^*(\mathbf{R}^d) \hookrightarrow \mathcal{H}_0^*(\mathbf{R}^d), \quad s, t \in \mathbf{R}_{b, \infty}, \quad s < t, \end{aligned} \quad (1.30)$$

are dense.

Remark 1.11. By the definition it follows that $T_{\mathcal{H}}$ in (1.26) is a homeomorphism between any of the spaces in (1.24) and corresponding space in (1.27), and that $T_{\mathcal{A}}$ in (1.26) is a homeomorphism between any of the spaces in (1.24) and corresponding space in (1.28).

The next results give some characterizations of $\mathcal{H}_s(\mathbf{R}^d)$ and $\mathcal{H}_{0, s}(\mathbf{R}^d)$ when s is a non-negative real number. See also [3] for similar characterizations of $\mathcal{H}_{b, \sigma}(\mathbf{R}^d)$ and $\mathcal{H}_{0, b, \sigma}(\mathbf{R}^d)$.

Proposition 1.12. *Let $0 \leq s, s_1, s_2 \in \mathbf{R}$ and let $f \in \mathcal{H}_0^*(\mathbf{R}^d)$. Then $f \in \mathcal{H}_s(\mathbf{R}^d)$ ($f \in \mathcal{H}_{0, s}(\mathbf{R}^d)$), if and only if $f \in C^\infty(\mathbf{R}^d)$ and satisfies*

$$\|H_d^N f\|_{L^\infty} \lesssim h^N N!^{2s}, \quad (1.31)$$

for some (every) $h > 0$. If $s_1 < \frac{1}{2}$ and $s_2 \geq \frac{1}{2}$ ($0 < s_1 \leq \frac{1}{2}$ and $s_2 > \frac{1}{2}$), then

$$\begin{aligned} \mathcal{H}_{s_1}(\mathbf{R}^d) \neq \mathcal{S}_{s_1}(\mathbf{R}^d) = \{0\}, & \quad (\mathcal{H}_{0, s_1}(\mathbf{R}^d) \neq \Sigma_{s_1}(\mathbf{R}^d) \neq \{0\}), \\ \mathcal{H}_{s_2}(\mathbf{R}^d) = \mathcal{S}_{s_2}(\mathbf{R}^d) \neq \{0\} & \quad \text{and} \quad (\mathcal{H}_{0, s_2}(\mathbf{R}^d) = \Sigma_{s_2}(\mathbf{R}^d) \neq \{0\}). \end{aligned}$$

We refer to [35] for the proof of Proposition 1.12.

Due to the pioneering investigations related to Proposition 1.12 by Pilipović in [24], we call the spaces $\mathcal{H}_s(\mathbf{R}^d)$ and $\mathcal{H}_{0, s}(\mathbf{R}^d)$ *Pilipović spaces of Roumieu and Beurling types*, respectively. In fact, in the restricted case $s, s_1, s_2 \geq \frac{1}{2}$, Proposition 1.12 was proved already in [24].

In what follows we let $F(z_2, \bar{z}_1)$ be the formal power series

$$\sum c(\alpha_2, \alpha_1) e_{\alpha_2}(z_2) e_{\alpha_1}(\bar{z}_1), \quad z_j \in \mathbf{C}^{d_j}, \quad j = 1, 2, \quad (1.32)$$

when $F(z_2, z_1)$ is the formal power series

$$\sum c(\alpha_2, \alpha_1) e_{\alpha_2}(z_2) e_{\alpha_1}(z_1). \quad (1.33)$$

Here $z_j \in \mathbf{C}^{d_j}$, $j = 1, 2$, and the sums should be taken over all $(\alpha_2, \alpha_1) \in \mathbf{N}^{d_2} \times \mathbf{N}^{d_1}$.

Definition 1.13. Let $s, s_1, s_2 \in \overline{\mathbf{R}_{b,\infty}}$, Θ_C , $W = \mathbf{C}^{d_2} \times \mathbf{C}^{d_1}$ and be the operator

$$(\Theta_C F)(z_2, z_1) = F(z_2, \bar{z}_1) \quad (1.34)$$

between formal power series in (1.32) and (1.33), $z_j \in \mathbf{C}^{d_j}$, $j = 1, 2$. Then

$$\widehat{\mathcal{A}}_{0,s}(W), \quad \widehat{\mathcal{A}}_s(W), \quad \widehat{\mathcal{A}}_s^*(W), \quad \widehat{\mathcal{A}}_{0,s}^*(W) \quad (1.35)$$

and

$$\widehat{\mathcal{A}}_{0,(s_2,s_1)}(W), \quad \widehat{\mathcal{A}}_{(s_2,s_1)}(W), \quad \widehat{\mathcal{A}}_{(s_2,s_1)}^*(W), \quad \widehat{\mathcal{A}}_{0,(s_2,s_1)}^*(W) \quad (1.36)$$

are the images of (1.28) respectively (1.28)' under Θ_C , and $\widehat{A}(W)$ is the image of $A(W)$ under Θ_C . The topologies of the spaces in (1.35), (1.36) and $\widehat{A}(W)$ are inherited from the topologies of the spaces (1.28), (1.28)' and $A(W)$ respectively, through the map Θ_C .

Remark 1.14. Let $W = \mathbf{C}^{d_2} \times \mathbf{C}^{d_1}$. By letting $d_2 = d$ and $d_1 = 0$, it follows that $A(\mathbf{C}^d)$ and the spaces in (1.28) can be considered as special cases of $\widehat{A}(W)$ and the spaces in (1.35).

Since $\mathcal{A}_{b_1}^*(\mathbf{C}^d) = A(\mathbf{C}^d)$ and $\mathcal{A}_{0,b_1}^*(\mathbf{C}^d) = A_d(\{0\})$, it follows that

$$\widehat{\mathcal{A}}_{b_1}^*(W) = \widehat{A}(W) \quad \text{and} \quad \widehat{\mathcal{A}}_{0,b_1}^*(W) = \widehat{A}_{d_2,d_1}(\{0\}). \quad (1.37)$$

The subspaces of $\widehat{A}(W)$ in the following definition are important when considering analytic kernel operators which are continuous on the spaces in (1.28).

Definition 1.15. Let $s_1, s_2 \in \overline{\mathbf{R}_{b,\infty}}$, $W = \mathbf{C}^{d_2} \times \mathbf{C}^{d_1}$, T_A and Θ_C be given by (1.26) and (1.34). Then

$$\widehat{\mathcal{B}}_s(W), \quad \widehat{\mathcal{B}}_{0,s}(W), \quad \widehat{\mathcal{B}}_s^*(W), \quad \widehat{\mathcal{B}}_{0,s}^*(W), \quad s = (s_2, s_1), \quad (1.38)$$

and

$$\widehat{\mathcal{C}}_s(W), \quad \widehat{\mathcal{C}}_{0,s}(W), \quad \widehat{\mathcal{C}}_s^*(W), \quad \widehat{\mathcal{C}}_{0,s}^*(W), \quad s = (s_2, s_1), \quad (1.39)$$

are the images of the spaces in (1.11) and (1.12) under the map $\Theta_C \circ T_A$. The topologies of the spaces in (1.38) and (1.39) are inherited from the topologies of the corresponding spaces in (1.11) and (1.12).

For conveniency we set

$$\begin{aligned} \ell_{\mathcal{B},s} &= \ell_{\mathcal{B},(s,s)}, & \ell_{\mathcal{B},0,s} &= \ell_{\mathcal{B},0,(s,s)}, & \ell_{\mathcal{C},s} &= \ell_{\mathcal{C},(s,s)}, & \ell_{\mathcal{C},0,s} &= \ell_{\mathcal{C},0,(s,s)}, \\ \widehat{\mathcal{B}}_s &= \widehat{\mathcal{B}}_{(s,s)}, & \widehat{\mathcal{B}}_{0,s} &= \widehat{\mathcal{B}}_{0,(s,s)}, & \widehat{\mathcal{C}}_s &= \widehat{\mathcal{C}}_{(s,s)} & \text{and} & \widehat{\mathcal{C}}_{0,s} &= \widehat{\mathcal{C}}_{0,(s,s)}. \end{aligned}$$

In particular, if $W = \mathbf{C}^d \times \mathbf{C}^d \simeq \mathbf{C}^{2d}$, then the spaces in (1.38) and (1.39) becomes

$$\widehat{\mathcal{B}}_s(\mathbf{C}^{2d}), \quad \widehat{\mathcal{B}}_{0,s}(\mathbf{C}^{2d}), \quad \widehat{\mathcal{B}}_s^*(\mathbf{C}^{2d}), \quad \widehat{\mathcal{B}}_{0,s}^*(\mathbf{C}^{2d}) \quad (1.38)'$$

and

$$\widehat{\mathcal{C}}_s(\mathbf{C}^{2d}), \quad \widehat{\mathcal{C}}_{0,s}(\mathbf{C}^{2d}), \quad \widehat{\mathcal{C}}_s^*(\mathbf{C}^{2d}), \quad \widehat{\mathcal{C}}_{0,s}^*(\mathbf{C}^{2d}). \quad (1.39)'$$

In Proposition 2.1 in Section 2 we present some duality properties of the spaces in Definitions 1.13 and 1.15. In Appendix A we identify the spaces in (1.38) and (1.39) with convenient subspaces of $\widehat{A}(W)$, for suitable $s_1, s_2 \in \overline{\mathbf{R}_b}$.

1.3. The Bargmann transform and spaces of analytic functions. Let $p \in [1, \infty]$. Then the Bargmann transform $\mathfrak{B}_d f$ of $f \in L^p(\mathbf{R}^d)$ is the entire function given by

$$(\mathfrak{B}_d f)(z) = \pi^{-\frac{d}{4}} \int_{\mathbf{R}^d} \exp\left(-\frac{1}{2}(\langle z, z \rangle + |y|^2) + 2^{1/2} \langle z, y \rangle\right) f(y) dy, \quad z \in \mathbf{C}^d.$$

We have

$$(\mathfrak{B}_d f)(z) = \int_{\mathbf{R}^d} \mathfrak{A}_d(z, y) f(y) dy, \quad z \in \mathbf{C}^d,$$

or

$$(\mathfrak{B}_d f)(z) = \langle f, \mathfrak{A}_d(z, \cdot) \rangle, \quad z \in \mathbf{C}^d, \quad (1.40)$$

where the Bargmann kernel \mathfrak{A}_d is given by

$$\mathfrak{A}_d(z, y) = \pi^{-\frac{d}{4}} \exp\left(-\frac{1}{2}(\langle z, z \rangle + |y|^2) + 2^{1/2} \langle z, y \rangle\right), \quad z \in \mathbf{C}^d, y \in \mathbf{R}^d.$$

(Cf. [4, 5].) Here

$$\langle z, w \rangle = \sum_{j=1}^d z_j w_j \quad \text{and} \quad (z, w) = \langle z, \bar{w} \rangle$$

when

$$z = (z_1, \dots, z_d) \in \mathbf{C}^d \quad \text{and} \quad w = (w_1, \dots, w_d) \in \mathbf{C}^d,$$

and otherwise $\langle \cdot, \cdot \rangle$ denotes the duality between test function spaces and their corresponding duals which is clear from the context. We note that the right-hand side of (1.40) makes sense when $f \in \mathcal{S}'_{1/2}(\mathbf{R}^d)$ and defines an element in $A(\mathbf{C}^d)$, since $y \mapsto \mathfrak{A}_d(z, y)$ can be interpreted as an element in $\mathcal{S}_{1/2}(\mathbf{R}^d)$ with values in $A(\mathbf{C}^d)$.

It was proved by Bargmann in [4] that $f \mapsto \mathfrak{B}_d f$ is a bijective and isometric map from $L^2(\mathbf{R}^d)$ to the Hilbert space $A^2(\mathbf{C}^d)$, the set of entire functions F on \mathbf{C}^d which fullfils

$$\|F\|_{A^2} \equiv \left(\int_{\mathbf{C}^d} |F(z)|^2 d\mu(z) \right)^{1/2} < \infty. \quad (1.41)$$

Here $d\mu(z) = \pi^{-d} e^{-|z|^2} d\lambda(z)$, where $d\lambda(z)$ is the Lebesgue measure on \mathbf{C}^d . The scalar product on $A^2(\mathbf{C}^d)$ is given by

$$(F, G)_{A^2} \equiv \int_{\mathbf{C}^d} F(z) \overline{G(z)} d\mu(z), \quad F, G \in A^2(\mathbf{C}^d). \quad (1.42)$$

For future references we note that the latter scalar product induces the bilinear form

$$(F, G) \mapsto \langle F, G \rangle_{A^2} = \langle F, G \rangle_{A^2(\mathbf{C}^d)} \equiv \int_{\mathbf{C}^d} F(z) G(z) d\mu(z) \quad (1.43)$$

on $A^2(\mathbf{C}^d) \times \overline{A^2(\mathbf{C}^d)}$.

In [4] it was proved that the Bargmann transform maps the Hermite functions to monomials as

$$\mathfrak{B}_d h_\alpha = e_\alpha, \quad z \in \mathbf{C}^d, \quad \alpha \in \mathbf{N}^d \quad (1.44)$$

(cf. (1.25)). The orthonormal basis $\{h_\alpha\}_{\alpha \in \mathbf{N}^d} \subseteq L^2(\mathbf{R}^d)$ is thus mapped to the orthonormal basis $\{e_\alpha\}_{\alpha \in \mathbf{N}^d} \subseteq A^2(\mathbf{C}^d)$.

In particular it follows that the definition of the Bargmann transform from $\mathcal{H}_0(\mathbf{R}^d)$ to $A^2(\mathbf{C}^d)$ is uniquely extendable to a homeomorphism from $\mathcal{H}_0^*(\mathbf{R}^d) = \mathcal{H}'_0(\mathbf{R}^d)$ to $\mathcal{A}_0^*(\mathbf{C}^d)$, by letting

$$\mathfrak{B}_d = T_{\mathcal{A}} \circ T_{\mathcal{H}}^{-1}, \quad (1.45)$$

where $T_{\mathcal{A}}$ and $T_{\mathcal{H}}$, where $T_{\mathcal{H}}$ and $T_{\mathcal{A}}$ are given by (1.26). From the definitions of $\mathcal{H}_s(\mathbf{R}^d)$ to $\mathcal{A}_s(\mathbf{C}^d)$ and their duals it follows that the Bargmann transform restricts to homeomorphisms from $\mathcal{H}_s(\mathbf{R}^d)$ to $\mathcal{A}_s(\mathbf{C}^d)$ and from $\mathcal{H}_s^*(\mathbf{R}^d)$ to $\mathcal{A}_s^*(\mathbf{C}^d)$. Similar facts hold true

with $\mathcal{H}_{0,s}$ and $\mathcal{A}_{0,s}$ in place of \mathcal{H}_s and \mathcal{A}_s , respectively, at each occurrence. (Cf. [35].) In particular, Remark 1.9, (1.46) and (1.44) show that the latter spaces in (1.28) are the (strong) duals of the former spaces with respect to unique extensions of the form $(\cdot, \cdot)_{A^2}$ on $\mathcal{A}_0(\mathbf{C}^d)$. That is,

$$\mathcal{A}'_s(\mathbf{C}^d) = \mathcal{A}^*_s(\mathbf{C}^d) \quad \text{and} \quad \mathcal{A}'_{0,s}(\mathbf{C}^d) = \mathcal{A}^*_{0,s}(\mathbf{C}^d). \quad (1.46)$$

Bargmann also proved that there is a reproducing formula for $A^2(\mathbf{C}^d)$. In fact, let Π_A be the operator from $L^2(d\mu)$ to $A(\mathbf{C}^d)$, given by

$$(\Pi_A F)(z) = \int_{\mathbf{C}^d} F(w) e^{(z,w)} d\mu(w), \quad z \in \mathbf{C}^d. \quad (1.47)$$

Then it is proved in [4] that Π_A is an orthonormal projection from $L^2(d\mu)$ to $A^2(\mathbf{C}^d)$.

From now on we assume that ϕ in short-time Fourier transforms (1.3) is given by

$$\phi(x) = \pi^{-\frac{d}{4}} e^{-|x|^2/2}, \quad x \in \mathbf{R}^d, \quad (1.48)$$

if nothing else is stated. For such ϕ , it follows by straight-forward computations that the relationship between the Bargmann transform and the short-time Fourier transform is given by

$$\mathfrak{B}_d = U_{\mathfrak{B}} \circ V_\phi, \quad \text{and} \quad U_{\mathfrak{B}}^{-1} \circ \mathfrak{B}_d = V_\phi, \quad (1.49)$$

where $U_{\mathfrak{B}}$ is the linear, continuous and bijective operator from $\mathcal{D}'(\mathbf{R}^{2d})$ to $\mathcal{D}'(\mathbf{C}^d)$, given by

$$(U_{\mathfrak{B}} F)(x + i\xi) = (2\pi)^{\frac{d}{2}} e^{\frac{1}{2}(|x|^2 + |\xi|^2)} e^{-i\langle x, \xi \rangle} F(\sqrt{2}x, -\sqrt{2}\xi), \quad x, \xi \in \mathbf{R}^d. \quad (1.50)$$

We notice that the inverse of $U_{\mathfrak{B}}$ is given by

$$(U_{\mathfrak{B}}^{-1} F)(x, \xi) = (2\pi)^{-\frac{d}{2}} e^{-\frac{1}{4}(|x|^2 + |\xi|^2)} e^{-\frac{i}{2}\langle x, \xi \rangle} F\left(\frac{x - i\xi}{\sqrt{2}}\right), \quad x, \xi \in \mathbf{R}^d \quad (1.51)$$

(cf. [33]).

If $W = \mathbf{C}^{d_2} \times \mathbf{C}^{d_1}$, then $\widehat{A}^2(W)$ consists of all $K \in C(W)$ such that $(z_2, z_1) \mapsto K(z_2, \bar{z}_1)$ belongs to $A^2(W)$. Then $\widehat{A}^2(W)$ is a Hilbert space under the scalar product

$$(K_1, K_2)_{\widehat{A}^2} = (K_1, K_2)_{\widehat{A}^2(W)} \equiv (K_{0,1}, K_{0,2})_{A^2(W)}, \quad (1.52)$$

when

$$K_j \in \widehat{A}^2(W), \quad K_{0,j}(z_2, z_1) = K_j(z_2, \bar{z}_1), \quad j = 1, 2. \quad (1.53)$$

1.4. Wick and anti-Wick operators. Next we recall the definition of analytic pseudo-differential operators, so-called Wick operators. Suppose that $a(z, w) \in \widehat{A}(\mathbf{C}^{2d})$ and $F \in A(\mathbf{C}^d)$ satisfy

$$w \mapsto a(z, w) F(w) e^{r|w| - |w|^2} \in L^1(\mathbf{C}^d) \quad (1.54)$$

for every $r > 0$ and $z \in \mathbf{C}^d$. Then the *analytic pseudo-differential operator*, or *Wick operator* $\text{Op}_{\mathfrak{B}}(a)$ with symbol a , is the linear and continuous operator from $\mathcal{A}_0(\mathbf{C}^d)$ to $A(\mathbf{C}^d)$, defined by the formula

$$\text{Op}_{\mathfrak{B}}(a)F(z) = \int_{\mathbf{C}^d} a(z, w) F(w) e^{(z,w)} d\mu(w), \quad z \in \mathbf{C}^d, \quad (1.55)$$

when $F \in \mathcal{A}_0(\mathbf{C}^d)$. (Cf. e.g. [6, 13, 31, 33, 35].) Here we remark that $\text{Op}_{\mathfrak{B}}(a)F$ is extendable in several ways, allowing a and F to belong to different spaces where (1.54) is violated. For example, in [31] it is proved that the definition of $\text{Op}_{\mathfrak{B}}(a)$ is uniquely extendable to any $a \in \widehat{\mathcal{A}}'_0(\mathbf{C}^{2d})$, and then $\text{Op}_{\mathfrak{B}}(a)$ is continuous from $\mathcal{A}_0(\mathbf{C}^d)$ to $\mathcal{A}'_0(\mathbf{C}^d)$.

The definition of the Wick operator in (1.55) resembles on the definition of the classical Kohn-Nirenberg pseudo-differential operators on \mathbf{R}^d . Let A be a real $d \times d$ matrix. Then the *pseudo-differential operator* $\text{Op}_A(\mathbf{a})$ with *symbol* $\mathbf{a} \in \mathcal{S}_{1/2}(\mathbf{R}^{2d})$ is the linear and continuous operator on $\mathcal{S}_{1/2}(\mathbf{R}^d)$, given by

$$(\text{Op}_A(\mathbf{a})f)(x) = (2\pi)^{-d} \iint \mathbf{a}(x - A(x - y), \xi) f(y) e^{i\langle x-y, \xi \rangle} dy d\xi, \quad x \in \mathbf{R}^d.$$

The definition of $\text{Op}_A(\mathbf{a})$ extends to any $\mathbf{a} \in \mathcal{S}'_{1/2}(\mathbf{R}^{2d})$, and then $\text{Op}_A(\mathbf{a})$ is continuous from $\mathcal{S}_{1/2}(\mathbf{R}^d)$ to $\mathcal{S}'_{1/2}(\mathbf{R}^d)$. (See e. g. [19, 34] and the references therein.)

The normal (Kohn-Nirenberg) representation and the Weyl quantization are obtained by choosing $A = 0$ and $A = \frac{1}{2}I$ respectively, where $I = I_d$ is the $d \times d$ identity matrix.

In the literature it is also common to consider anti-Wick operators. Suppose that $a \in \widehat{A}(\mathbf{C}^{2d})$ satisfies

$$w \mapsto a(w, w) e^{r|w| - |w|^2} \in L^1(\mathbf{C}^d) \quad (1.56)$$

for every $r > 0$. Then the *anti-Wick operator* $\text{Op}_{\mathfrak{W}}^{\text{aw}}(a)$ with symbol a is the linear and continuous operator from $\mathcal{A}_0(\mathbf{C}^d)$ to $A(\mathbf{C}^d)$, given by

$$\text{Op}_{\mathfrak{W}}^{\text{aw}}(a)F(z) = \int_{\mathbf{C}^d} a(w, w) F(w) e^{(z, w)} d\mu(w), \quad z \in \mathbf{C}^d, \quad (1.57)$$

when $F \in \mathcal{A}_0(\mathbf{C}^d)$.

Berezin established in [6] convenient links between Wick and anti-Wick operators (see page 587 in [6]). More precisely, if $a \in \widehat{\mathcal{A}}_0(\mathbf{C}^{2d})$, then

$$\text{Op}_{\mathfrak{W}}^{\text{aw}}(a) = \text{Op}_{\mathfrak{W}}(a^{\text{aw}}) \quad (1.58)$$

when a^{aw} is given by

$$a^{\text{aw}}(z, w) = \pi^{-d} \int_{\mathbf{C}^d} a(w_1, w_1) e^{-(z-w_1, w-w_1)} d\lambda(w_1). \quad (1.59)$$

If instead $a^{\text{aw}} \in \widehat{\mathcal{A}}_0(\mathbf{C}^{2d})$, then (1.58) holds true when $a^{\text{aw}} \in \widehat{A}(\mathbf{C}^{2d})$ is given by

$$a(z, -w) = \pi^{-d} \int_{\mathbf{C}^d} a^{\text{aw}}(w_1, -w_1) e^{-(z-w_1, w-w_1)} d\lambda(w_1). \quad (1.60)$$

In fact, if $F \in \mathcal{A}_0(\mathbf{C}^d)$ and a^{aw} is given by (1.59), then

$$\text{Op}_{\mathfrak{W}}(a^{\text{aw}})F(z) = \pi^{-d} \int_{\mathbf{C}^d} a(w_1, w_1) e^{(z, w_1) - |w_1|^2} \left(\int_{\mathbf{C}^d} F(w) e^{(w_1, w)} d\mu(w) \right) d\lambda(w_1),$$

and by applying the reproducing formula (1.47) on the inner integral, it follows that the right-hand side becomes $\text{Op}_{\mathfrak{W}}^{\text{aw}}(a)F(z)$ in (1.57). This shows that (1.58) holds true when a^{aw} is given by (1.59). In Section 3 and 4 we show the equivalence between (1.59) and (1.60), when a or a^{aw} are allowed to belong to larger classes than $\widehat{\mathcal{A}}_0(\mathbf{C}^{2d})$.

Since any element $a(z, w)$ in $\widehat{A}(\mathbf{C}^{2d})$ is uniquely determined on its values at the diagonal $z = w$ or anti-diagonal $z = -w$, it follows that (1.59) and (1.60) are equivalent to

$$a^{\text{aw}}(z, z) = \pi^{-d} \int_{\mathbf{C}^d} a(w_1, w_1) e^{-|z-w_1|^2} d\lambda(w_1) \quad (1.59)'$$

and

$$a(z, -z) = \pi^{-d} \int_{\mathbf{C}^d} a^{\text{aw}}(w_1, -w_1) e^{-|z-w_1|^2} d\lambda(w_1). \quad (1.60)'$$

These formulae were especially emphasized on page 587 in [6].

Let $a_1, a_2 \in \widehat{\mathcal{A}}'_0(\mathbf{C}^{2d})$. For suitable additional conditions on a_1 and a_2 , it is possible to compose the Wick operators $\text{Op}_{\mathfrak{W}}(a_1)$ and $\text{Op}_{\mathfrak{W}}(a_2)$, and the resulting operator is again a Wick operator. The (complex) twisted product $a_1 \#_{\mathfrak{W}} a_2$ is then defined by

$$\text{Op}_{\mathfrak{W}}(a_1) \circ \text{Op}_{\mathfrak{W}}(a_2) = \text{Op}_{\mathfrak{W}}(a_1 \#_{\mathfrak{W}} a_2), \quad (1.61)$$

provided the composition on the left-hand side is well-defined as a continuous operator from $\mathcal{A}_0(\mathbf{C}^d)$ to $\mathcal{A}'_0(\mathbf{C}^d)$. By straight-forward computations it follows that the product $\#_{\mathfrak{W}}$ is given by

$$a_1 \#_{\mathfrak{W}} a_2(z, w) = \pi^{-d} \int_{\mathbf{C}^d} a_1(z, u) a_2(u, w) e^{-(z-u, w-u)} d\lambda(u), \quad z, w \in \mathbf{C}^d, \quad (1.62)$$

when the integrand belongs to $L^1(\mathbf{C}^d)$ for every $z, w \in \mathbf{C}^d$ (see e. g. [6, 32]).

1.5. Toeplitz operators. Let $\phi_1, \phi_2 \in \mathcal{S}_{1/2}(\mathbf{R}^d) \setminus 0$. Then the Toeplitz operator $\text{Tp}_{\phi_1, \phi_2}(\mathbf{a})$ with symbol $\mathbf{a} \in \mathcal{S}'_{1/2}(\mathbf{R}^{2d})$ is the linear and continuous operator from $\mathcal{S}_{1/2}(\mathbf{R}^d)$ to $\mathcal{S}'_{1/2}(\mathbf{R}^d)$, given by the formula

$$(\text{Tp}_{\phi_1, \phi_2}(\mathbf{a})f, g)_{L^2(\mathbf{R}^d)} = (\mathbf{a} \cdot V_{\phi_1} f, V_{\phi_2} g)_{L^2(\mathbf{R}^{2d})}. \quad (1.63)$$

It is proved in [35] that $\text{Tp}_{\phi_1, \phi_2}(\mathbf{a})$ is continuous on $\mathcal{S}_{1/2}(\mathbf{R}^d)$ and uniquely extendable to a continuous operator on $\mathcal{S}'_{1/2}(\mathbf{R}^d)$.

In our situation we have $\phi_1 = \phi_2$ is equal to ϕ in (1.48), and for conveniency we put $\text{Tp}(\mathbf{a}) = \text{Tp}_{\phi, \phi}(\mathbf{a})$. In fact, for suitable $a \in \widehat{A}(\mathbf{C}^{2d})$ we shall mainly consider modified Toeplitz operators $\text{Tp}_{\mathfrak{W}}(a)$ given by

$$\text{Tp}_{\mathfrak{W}}(a) = \text{Tp}(\mathbf{a}) \quad \text{when} \quad a(z, z) = \mathbf{a}(\sqrt{2}x, -\sqrt{2}\xi), \quad z = x + i\xi. \quad (1.63)'$$

By straight-forward applications of (1.50) and (1.51) it follows

$$\text{Op}_{\mathfrak{W}}^{\text{aw}}(a) = \mathfrak{W}_d \circ \text{Tp}_{\mathfrak{W}}(a) \circ \mathfrak{W}_d^{-1}. \quad (1.64)$$

(See also Section 6 in [33].)

We recall that Toeplitz operators can be formulated as pseudo-differential operators, due to the formula

$$\text{Tp}(\mathbf{a}) = \left(\frac{2}{\pi}\right)^{\frac{d}{2}} \text{Op}^w(\mathbf{a} * e^{-|\cdot|^2}), \quad (1.65)$$

which is equivalent to (1.59) and (1.60). (See e. g. [13, 28, 33] and the references therein.)

2. OPERATOR KERNELS AND MULTIPLICATIONS FOR SPACES OF POWER SERIES EXPANSIONS

In this section we deduce kernel theorems for linear operators acting between (different) \mathcal{A}_s spaces and their duals. In [31] such kernel results were explained for linear operators which map $\mathcal{A}_s(\mathbf{C}^{d_1})$ into $\mathcal{A}'_s(\mathbf{C}^{d_2})$ or which map $\mathcal{A}_{0,s}(\mathbf{C}^{d_1})$ into $\mathcal{A}'_{0,s}(\mathbf{C}^{d_2})$. The latter results can also be obtained by classical kernel results for linear operators acting on nuclear spaces (see e. g. [36]).

2.1. Kernels to linear operators acting on spaces of power series expansions.

The continuity results for kernel operators rely on duality properties between the spaces through the $\widehat{\mathcal{A}}^2$ form in (1.52) and (1.53). In fact, we have the following proposition which describes important topological properties of the spaces in Definitions 1.13 and 1.15, and which is a straight-forward consequence of Remarks 1.9 and 1.8, and the definitions. The details are left for the reader.

Proposition 2.1. *Let $s_0, s_1, s_2 \in \overline{\mathbf{R}}_b$, $s = (s_2, s_1)$ and $W = \mathbf{C}^{d_2} \times \mathbf{C}^{d_1}$. Then the following is true:*

- (1) *the spaces in (1.36) are complete Hausdorff topological vector spaces, and $\widehat{\mathcal{A}}_{0,s}(W)$ and $\widehat{\mathcal{A}}_s^*(W)$ are Fréchet spaces. It holds $\widehat{\mathcal{A}}_{(s_0, s_0)} = \widehat{\mathcal{A}}_{s_0}$ and $\widehat{\mathcal{A}}_{(s_0, s_0)}^* = \widehat{\mathcal{A}}_{s_0}^*$;*
- (2) *$\widehat{\mathcal{A}}_0(W)$ is dense in the spaces in (1.36), (1.38) and (1.39);*
- (3) *the form $(\cdot, \cdot)_{\widehat{\mathcal{A}}^2(W)}$ on $\widehat{\mathcal{A}}_0(W)$ is uniquely extendable to separate continuous non-degenerate mappings from*

$$\widehat{\mathcal{A}}_s^*(W) \times \widehat{\mathcal{A}}_s(W), \quad \widehat{\mathcal{C}}_s^*(W) \times \widehat{\mathcal{B}}_s(W) \quad \text{and} \quad \widehat{\mathcal{B}}_s^*(W) \times \widehat{\mathcal{C}}_s(W) \quad (2.1)$$

to \mathbf{C} . Furthermore, $\widehat{\mathcal{A}}_s^(W)$ and $\widehat{\mathcal{A}}_s(W)$ are (strong) duals to each others, through the form $(\cdot, \cdot)_{\widehat{\mathcal{A}}^2(W)}$.*

If instead $s_1, s_2 \in \mathbf{R}_{b, \infty}$, then the same holds true with $\widehat{\mathcal{A}}_{0,s}$, $\widehat{\mathcal{B}}_{0,s}$ and $\widehat{\mathcal{C}}_{0,s}$ in place of $\widehat{\mathcal{A}}_s$, $\widehat{\mathcal{B}}_s$ and $\widehat{\mathcal{C}}_s$, respectively, at each occurrence.

Evidently, it follows from (3) in Proposition 2.1 that

$$\widehat{\mathcal{A}}'_{(s_2, s_1)}(W) = \widehat{\mathcal{A}}^*_{(s_2, s_1)}(W), \quad (\widehat{\mathcal{A}}^*_{(s_2, s_1)})'(W) = \widehat{\mathcal{A}}_{(s_2, s_1)}(W), \quad (2.2)$$

through the form $(\cdot, \cdot)_{\widehat{\mathcal{A}}^2(W)}$.

The following kernel results characterize linear operators which map (different) \mathcal{A}_s or $\mathcal{A}_{0,s}$ spaces and their duals into each others. Here and in what follows we let $\mathcal{L}(V_1; V_2)$ be the set of linear and continuous mappings from the topological vector space V_1 into the topological vector space V_2 . We also set $\mathcal{L}(V) = \mathcal{L}(V; V)$.

Proposition 2.2. *Let $s_1, s_2 \in \overline{\mathbf{R}}_{b, \infty}$, $s = (s_2, s_1)$, $W = \mathbf{C}^{d_2} \times \mathbf{C}^{d_1}$, and let $T \in \mathcal{L}(\mathcal{A}_0(\mathbf{C}^{d_1}); \mathcal{A}'_0(\mathbf{C}^{d_2}))$. Then the following is true:*

- (1) *if $T \in \mathcal{L}(\mathcal{A}'_{s_1}(\mathbf{C}^{d_1}); \mathcal{A}_{s_2}(\mathbf{C}^{d_2}))$, then there is a unique $K \in \widehat{\mathcal{A}}_s(W)$ such that*

$$(TF)(z_2) = (F, \overline{K(z_2, \cdot)})_{\mathcal{A}^2(\mathbf{C}^{d_1})}, \quad z_2 \in \mathbf{C}^{d_2}, \quad (2.3)$$

holds true for every $F \in \mathcal{A}_{s_1}(\mathbf{C}^{d_1})$;

- (2) *if $T \in \mathcal{L}(\mathcal{A}_{s_1}(\mathbf{C}^{d_1}); \mathcal{A}'_{s_2}(\mathbf{C}^{d_2}))$, then there is a unique $K \in \widehat{\mathcal{A}}'_s(W)$ such that (2.3) holds true for every $F \in \mathcal{A}'_{s_1}(\mathbf{C}^{d_1})$.*

The same holds true with \mathcal{A}_{0,s_j} and $\widehat{\mathcal{A}}_{0,s}$ in place of \mathcal{A}_{s_j} and $\widehat{\mathcal{A}}_s$, respectively, $j = 1, 2$, at each occurrence.

Here (2.3) should be interpreted as

$$(TF, G)_{\mathcal{A}^2(\mathbf{C}^{d_2})} = (K, G \otimes \overline{F})_{\widehat{\mathcal{A}}^2(\mathbf{C}^{d_2} \times \mathbf{C}^{d_1})}, \quad (2.3)'$$

when $F \in \mathcal{A}_0(\mathbf{C}^{d_1})$ and $G \in \mathcal{A}_0(\mathbf{C}^{d_2})$.

Proposition 2.3. *Let $s_1, s_2 \in \overline{\mathbf{R}}_{b, \infty}$, $s = (s_2, s_1)$, $W = \mathbf{C}^{d_2} \times \mathbf{C}^{d_1}$, $K \in \widehat{\mathcal{A}}'_0(W)$, and let $T = T_K \in \mathcal{L}(\mathcal{A}_0(\mathbf{C}^{d_1}); \mathcal{A}'_0(\mathbf{C}^{d_2}))$ be given by*

$$F \mapsto T_K F = (z_2 \mapsto (F, \overline{K(z_2, \cdot)})_{\mathcal{A}^2(\mathbf{C}^{d_1})}). \quad (2.4)$$

Then the following is true:

- (1) if $K \in \widehat{\mathcal{A}}_s(W)$, then T_K extends uniquely to a linear and continuous map from $\mathcal{A}'_{s_1}(\mathbf{C}^{d_1})$ to $\mathcal{A}_{s_2}(\mathbf{C}^{d_2})$;
- (2) if $K \in \widehat{\mathcal{A}}'_s(W)$, then T_K extends uniquely to a linear and continuous map from $\mathcal{A}_{s_1}(\mathbf{C}^{d_1})$ to $\mathcal{A}'_{s_2}(\mathbf{C}^{d_2})$.

The same holds true with \mathcal{A}_{0,s_j} and $\widehat{\mathcal{A}}_{0,s}$ in place of \mathcal{A}_{s_j} and $\widehat{\mathcal{A}}_s$, respectively, $j = 1, 2$, at each occurrence.

We observe that if K and F are the same as in (2.4) and $z_1 \mapsto K(z_2, z_1)F(z_1)$ is integrable with respect to $d\mu(z_1)$, then (2.4) is the same as

$$(T_K F)(z_2) = \int_{\mathbf{C}^{d_1}} K(z_2, z_1)F(z_1) d\mu(z_1). \quad (2.4)'$$

Propositions 2.2 and 2.3 follow essentially from abstract kernel results for linear operators acting between topological vector spaces. (See e.g. [36]. See also [31] for more explicit approaches.) On the other hand, the following result might be more cumbersome to deduce from such abstract kernel results.

Proposition 2.4. *Let $s_1, s_2 \in \overline{\mathbf{R}}_{b,\infty}$, $s = (s_2, s_1)$, $W = \mathbf{C}^{d_2} \times \mathbf{C}^{d_1}$, and let $T \in \mathcal{L}(\mathcal{A}_0(\mathbf{C}^{d_1}); \mathcal{A}'_0(\mathbf{C}^{d_2}))$. Then the following is true:*

- (1) if $T \in \mathcal{L}(\mathcal{A}_{s_1}(\mathbf{C}^{d_1}); \mathcal{A}_{s_2}(\mathbf{C}^{d_2}))$, then there is a unique $K \in \widehat{\mathcal{B}}_s(W)$ such that (2.3) holds true for every $F \in \mathcal{A}_{s_1}(\mathbf{C}^{d_1})$;
- (2) if $T \in \mathcal{L}(\mathcal{A}'_{s_1}(\mathbf{C}^{d_1}); \mathcal{A}'_{s_2}(\mathbf{C}^{d_2}))$, then there is a unique $K \in \widehat{\mathcal{B}}_s^*(W)$ such that (2.3) holds true for every $F \in \mathcal{A}'_{s_1}(\mathbf{C}^{d_1})$.

The same holds true with \mathcal{A}_{0,s_j} and $\widehat{\mathcal{B}}_{0,s}$ in place of \mathcal{A}_{s_j} and $\widehat{\mathcal{B}}_s$, respectively, $j = 1, 2$, at each occurrence.

We also have the following converse of the preceding proposition.

Proposition 2.5. *Let $s_1, s_2 \in \overline{\mathbf{R}}_{b,\infty}$, $s = (s_2, s_1)$, $W = \mathbf{C}^{d_2} \times \mathbf{C}^{d_1}$, $K \in \widehat{\mathcal{A}}'_0(W)$, and let $T = T_K \in \mathcal{L}(\mathcal{A}_0(\mathbf{C}^{d_1}); \mathcal{A}'_0(\mathbf{C}^{d_2}))$ be given by (2.4). Then the following is true:*

- (1) if $K \in \widehat{\mathcal{B}}_s(W)$, then T_K extends uniquely to a linear and continuous map from $\mathcal{A}_{s_1}(\mathbf{C}^{d_1})$ to $\mathcal{A}_{s_2}(\mathbf{C}^{d_2})$;
- (2) if $K \in \widehat{\mathcal{B}}_s^*(W)$, then T_K extends uniquely to a linear and continuous map from $\mathcal{A}'_{s_1}(\mathbf{C}^{d_1})$ to $\mathcal{A}'_{s_2}(\mathbf{C}^{d_2})$.

The same holds true with \mathcal{A}_{0,s_j} and $\widehat{\mathcal{B}}_{0,s}$ in place of \mathcal{A}_{s_j} and $\widehat{\mathcal{B}}_s$, respectively, $j = 1, 2$, at each occurrence.

By combining Propositions 2.2–2.5, we get the following.

Corollary 2.6. *Let $s_1, s_2 \in \overline{\mathbf{R}}_b$, $s = (s_1, s_2)$, $W = \mathbf{C}^{d_2} \times \mathbf{C}^{d_1}$ and let T_K be the map in (2.4) when $K \in \widehat{\mathcal{A}}'_0(W)$. Then the mappings*

$$\widehat{\mathcal{A}}_s(W) \ni K \mapsto T_K \in \mathcal{L}(\mathcal{A}'_{s_1}(\mathbf{C}^{d_1}), \mathcal{A}_{s_2}(\mathbf{C}^{d_2})),$$

$$\widehat{\mathcal{A}}'_s(W) \ni K \mapsto T_K \in \mathcal{L}(\mathcal{A}_{s_2}(\mathbf{C}^{d_1}), \mathcal{A}'_{s_1}(\mathbf{C}^{d_2})),$$

$$\widehat{\mathcal{B}}_s(W) \ni K \mapsto T_K \in \mathcal{L}(\mathcal{A}_{s_1}(\mathbf{C}^{d_1}), \mathcal{A}_{s_2}(\mathbf{C}^{d_2}))$$

and

$$\widehat{\mathcal{B}}_s^*(W) \ni K \mapsto T_K \in \mathcal{L}(\mathcal{A}'_{s_1}(\mathbf{C}^{d_1}), \mathcal{A}'_{s_2}(\mathbf{C}^{d_2}))$$

are isomorphisms. The same holds true with $\widehat{\mathcal{A}}_{0,s}$, $\widehat{\mathcal{B}}_{0,s}$, \mathcal{A}_{0,s_1} and \mathcal{A}_{0,s_2} in place of $\widehat{\mathcal{A}}_s$, $\widehat{\mathcal{B}}_s$, \mathcal{A}_{s_1} and \mathcal{A}_{s_2} , respectively, at each occurrence.

Proposition 2.5 follows by straight-forward computations and is left for the reader.

Proof of Proposition 2.4. We only consider the case when T is continuous from $\mathcal{A}_{s_1}(\mathbf{C}^{d_1})$ to $\mathcal{A}_{s_2}(\mathbf{C}^{d_2})$. The other cases follow by similar arguments and are left for the reader. Let $\vartheta_{r,s}$ be as in Definition 1.3. By Propositions 2.2 and 2.3, there is a unique

$$K(z_2, z_1) = \sum_{\alpha_j \in \mathbf{N}^{d_j}} c(K; \alpha_2, \alpha_1) e_{\alpha_2}(z_2) e_{\alpha_1}(\bar{z}_1) \in \widehat{\mathcal{A}}'_0(W), \quad z_j \in \mathbf{C}^{d_j}, \quad (2.5)$$

such that (2.3) holds true for every $F \in \mathcal{A}_0(\mathbf{C}^{d_1})$. We need to show that $K \in \widehat{\mathcal{B}}_s(W)$.

Let $r > 0$ and set $F_\alpha(z_1) = \vartheta_{r,s_1}^{-1}(\alpha) e_\alpha(z_1)$ for every $\alpha \in \mathbf{N}^{d_1}$ and $z_1 \in \mathbf{Z}^{d_1}$. Then $\{F_\alpha\}_{\alpha \in \mathbf{N}^{d_1}}$ is a bounded subset of $\mathcal{A}_{s_1}(\mathbf{C}^{d_1})$. By the continuity of T it follows that $\{TF_\alpha\}_{\alpha \in \mathbf{N}^{d_1}}$ is a bounded subset of $\mathcal{A}_{s_2}(\mathbf{C}^{d_2})$. Hence, if

$$(TF_\alpha)(z_2) = \sum_{\alpha_2 \in \mathbf{N}^{d_2}} c(TF_\alpha; \alpha_2) e_{\alpha_2}(z_2), \quad z_2 \in \mathbf{C}^{d_2},$$

then

$$c(TF_\alpha; \alpha_2) = c(K; \alpha_2, \alpha) \vartheta_{r,s_1}^{-1}(\alpha)$$

satisfies

$$|c(TF_\alpha; \alpha_2)| \leq C \vartheta_{r_0,s_2}^{-1}(\alpha_2),$$

for some constants $C, r_0 > 0$ which are independent of α_2 and α . This gives,

$$|c(K; \alpha_2, \alpha_1)| \lesssim \vartheta_{r,s_1}(\alpha_1) \vartheta_{r_0,s_2}^{-1}(\alpha_2), \quad \alpha_j \in \mathbf{C}^{d_j}.$$

Since $r > 0$ is arbitrary, the latter estimate implies that $K \in \widehat{\mathcal{B}}_s(\mathbf{C}^{d_2} \times \mathbf{C}^{d_1})$. \square

In Section 4 we deduce related results compared to Propositions 2.2–2.5 and Corollary 2.6 with Wick and anti-Wick operators in place of kernel operators.

Example 2.7. Let $\omega \in L_{\text{loc}}^\infty(\mathbf{C}^d) \simeq L_{\text{loc}}^\infty(\mathbf{R}^{2d})$ be such that

$$\omega(z) > 0 \quad \text{and} \quad \omega(z+w) \lesssim \omega(z) \langle w \rangle^r, \quad z, w \in \mathbf{C}^d, \quad (2.6)$$

for some $r > 0$. Then we recall that the Shubin class $\text{Sh}_\rho^{(\omega)}(\mathbf{R}^{2d})$ (with respect to ρ and ω) is the set of all $\mathbf{a} \in C^\infty(\mathbf{R}^{2d})$ such that

$$|\partial^\alpha \mathbf{a}(x, \xi)| \lesssim \omega(x + i\xi) \langle (x, \xi) \rangle^{-\rho|\alpha|}.$$

(See [28].)

As in [32], we let $\widehat{\mathcal{A}}_{\text{Sh},\rho}^{(\omega)}(\mathbf{C}^{2d})$ be the class of Wick symbols which consists of all $a \in \widehat{\mathcal{A}}(\mathbf{C}^{2d})$ such that

$$|\partial_z^\alpha \bar{\partial}_w^\beta a(z, w)| \lesssim e^{\frac{1}{2}|z-w|^2} \omega(\sqrt{2}\bar{z}) \langle z+w \rangle^{-\rho|\alpha+\beta|} \langle z-w \rangle^{-N} \quad (2.7)$$

for every $\alpha, \beta \in \mathbf{N}^d$ and $N \geq 0$. Let A be areal $d \times d$ matrix. Then recall that for $\mathbf{a} \in \mathcal{S}'(\mathbf{R}^{2d})$, we have $\mathbf{a} \in \text{Sh}_\rho^{(\omega)}(\mathbf{R}^{2d})$, if and only if $\text{Op}_{\mathfrak{W}}(\mathbf{a}) = \mathfrak{W}_d \circ \text{Op}_A(\mathbf{a}) \circ \mathfrak{W}_d^{-1}$ for some $a \in \widehat{\mathcal{A}}_{\text{Sh},\rho}^{(\omega)}(\mathbf{C}^{2d})$ (cf. [32]).

Let $a \in \widehat{\mathcal{A}}_{\text{Sh},\rho}^{(\omega)}(\mathbf{C}^{2d})$ and let K be the kernel of $\text{Op}_{\mathfrak{W}}(a)$. We claim that

$$K \in \widehat{\mathcal{B}}_{0,\infty}(\mathbf{C}^{2d}) \cap \widehat{\mathcal{B}}_{0,\infty}^*(\mathbf{C}^{2d}). \quad (2.8)$$

In fact, by (2.7) and that $K(z, w) = e^{(z,w)} a(z, w)$ we get

$$|K(z, w)| \lesssim e^{\frac{1}{2}(|z|^2+|w|^2)} \omega(\sqrt{2}\bar{z}) \langle z-w \rangle^{-N} \quad \text{for every } N \geq 0. \quad (2.9)$$

Since ω satisfies (2.6) we get

$$\omega(\sqrt{2}\bar{z}) \lesssim \langle z \rangle^{N_0} \quad \text{and} \quad \omega(\sqrt{2}\bar{z}) \lesssim \omega(\sqrt{2}\bar{w}) \langle z-w \rangle^{N_0} \lesssim \langle w \rangle^{N_0} \langle z-w \rangle^{N_0},$$

for some $N_0 \geq 0$. We also have

$$\langle z - w \rangle^{-N} \lesssim \langle z \rangle^{-N} \langle w \rangle^N \quad \text{and} \quad \langle z - w \rangle^{-N} \lesssim \langle z \rangle^N \langle w \rangle^{-N}.$$

By playing with $N \geq 0$, a combination of these estimates with (2.9) shows that

$$|K(z, w)| \lesssim e^{\frac{1}{2}(|z|^2 + |w|^2)} \langle z \rangle^{N_0 + N} \langle w \rangle^{-N}$$

and

$$|K(z, w)| \lesssim e^{\frac{1}{2}(|z|^2 + |w|^2)} \langle z \rangle^{-N} \langle w \rangle^{N_0 + N},$$

which implies that (2.8) holds.

By Proposition 2.4 it now follows that $\text{Op}_{\mathfrak{M}}(a)$ is continuous on both $\mathcal{A}_{0,\infty}(\mathbf{C}^d)$ and on $\mathcal{A}'_{0,\infty}(\mathbf{C}^d)$. This shows that any Shubin operator is continuous on $\mathcal{S}(\mathbf{R}^d)$ and on $\mathcal{S}'(\mathbf{R}^d)$ (see e. g. [19, Theorem 18.6.2]).

2.2. Compositions of analytic Kernel operators. .

In what follows we identify any linear and continuous operator T from $\mathcal{A}_0(\mathbf{C}^{d_1})$ to $\mathcal{A}'_0(\mathbf{C}^{d_2})$ by its kernel $K \in \widehat{\mathcal{A}}'_0(\mathbf{C}^{d_2} \times \mathbf{C}^{d_1})$ in (2.3) and (2.3)'. By general continuity properties we get the following theorem concerning the composition map

$$(K_2, K_1) \mapsto K_2 \circ K_1. \quad (2.10)$$

Theorem 2.8. *Let $W_j = \mathbf{C}^{d_{j+1}} \times \mathbf{C}^{d_j}$, $j = 1, 2$, $W_3 = \mathbf{C}^{d_3} \times \mathbf{C}^{d_1}$, $s_1, s_2, s_3 \in \overline{\mathbf{R}}_b$ and let T be the map from $\widehat{\mathcal{A}}_0(W_2) \times \widehat{\mathcal{A}}_0(W_1)$ to $\widehat{\mathcal{A}}_0(W_3)$, given by (2.10). Then the following is true:*

- (1) *T is uniquely extendable to continuous mappings from $\widehat{\mathcal{A}}_{(s_3, s_2)}(W_2) \times \widehat{\mathcal{B}}_{(s_2, s_1)}^*(W_1)$ or from $\widehat{\mathcal{B}}_{(s_3, s_2)}(W_2) \times \widehat{\mathcal{A}}_{(s_2, s_1)}(W_1)$ to $\widehat{\mathcal{A}}_{(s_3, s_1)}(W_3)$;*
- (2) *T is uniquely extendable to continuous mappings from $\widehat{\mathcal{B}}_{(s_3, s_2)}^*(W_2) \times \widehat{\mathcal{A}}'_{(s_2, s_1)}(W_1)$ or from $\widehat{\mathcal{A}}'_{(s_3, s_2)}(W_2) \times \widehat{\mathcal{B}}_{(s_2, s_1)}(W_1)$ to $\widehat{\mathcal{A}}'_{(s_3, s_1)}(W_3)$;*

If instead $s_1, s_2, s_3 \in \mathbf{R}_{b,\infty}$, then the same hold true with $\widehat{\mathcal{A}}_{0,(s_j, s_k)}$ and $\widehat{\mathcal{B}}_{0,(s_j, s_k)}$ in place of $\widehat{\mathcal{A}}_{(s_j, s_k)}$ and $\widehat{\mathcal{B}}_{(s_j, s_k)}$, respectively, at each occurrence.

Theorem 2.9. *Let $W_j = \mathbf{C}^{d_{j+1}} \times \mathbf{C}^{d_j}$, $j = 1, 2$, $W_3 = \mathbf{C}^{d_3} \times \mathbf{C}^{d_1}$, $s_1, s_2, s_3 \in \overline{\mathbf{R}}_b$ and let T be the map from $\widehat{\mathcal{A}}_0(W_2) \times \widehat{\mathcal{A}}_0(W_1)$ to $\widehat{\mathcal{A}}_0(W_3)$, given by (2.10). Then the following is true:*

- (1) *T is uniquely extendable to continuous mappings from $\widehat{\mathcal{B}}_{(s_3, s_2)}(W_2) \times \widehat{\mathcal{C}}_{(s_2, s_1)}(W_1)$ or from $\widehat{\mathcal{C}}_{(s_3, s_2)}(W_2) \times \widehat{\mathcal{B}}_{(s_2, s_1)}(W_1)$ to $\widehat{\mathcal{C}}_{(s_3, s_1)}(W_3)$. The same holds true with $\widehat{\mathcal{B}}_{(s_j, s_k)}$ in place of $\widehat{\mathcal{C}}_{(s_j, s_k)}$ at each occurrence;*
- (2) *T is uniquely extendable to continuous mappings from $\widehat{\mathcal{B}}_{(s_3, s_2)}^*(W_2) \times \widehat{\mathcal{C}}_{(s_2, s_1)}^*(W_1)$ or from $\widehat{\mathcal{C}}_{(s_3, s_2)}^*(W_2) \times \widehat{\mathcal{B}}_{(s_2, s_1)}^*(W_1)$ to $\widehat{\mathcal{C}}_{(s_3, s_1)}^*(W_3)$. The same holds true with $\widehat{\mathcal{B}}_{(s_j, s_k)}$ in place of $\widehat{\mathcal{C}}_{(s_j, s_k)}$ at each occurrence.*

If instead $s_1, s_2, s_3 \in \mathbf{R}_{b,\infty}$, then the same hold true with $\widehat{\mathcal{A}}_{0,(s_j, s_k)}$, $\widehat{\mathcal{B}}_{0,(s_j, s_k)}$ and $\widehat{\mathcal{C}}_{0,(s_j, s_k)}$ in place of $\widehat{\mathcal{A}}_{(s_j, s_k)}$, $\widehat{\mathcal{B}}_{(s_j, s_k)}$ and $\widehat{\mathcal{C}}_{(s_j, s_k)}$, respectively, at each occurrence.

Proof of Theorems 2.8 and 2.9. The assertion, except for those parts which involve $\widehat{\mathcal{C}}_{(s_j, s_k)}$ and $\widehat{\mathcal{C}}_{(s_j, s_k)}^*$ spaces, follows from the kernel results Propositions 2.2 to 2.5 and the composition properties of the form

$$\mathcal{L}(\mathcal{A}_{s_2}(\mathbf{C}^{d_2}), \mathcal{A}_{s_3}(\mathbf{C}^{d_3})) \circ \mathcal{L}(\mathcal{A}_{s_1}(\mathbf{C}^{d_1}), \mathcal{A}_{s_2}(\mathbf{C}^{d_2})) \subseteq \mathcal{L}(\mathcal{A}_{s_1}(\mathbf{C}^{d_1}), \mathcal{A}_{s_3}(\mathbf{C}^{d_3}))$$

and

$$\mathcal{L}(\mathcal{A}_{s_2}(\mathbf{C}^{d_2}), \mathcal{A}'_{s_3}(\mathbf{C}^{d_3})) \circ \mathcal{L}(\mathcal{A}_{s_1}(\mathbf{C}^{d_1}), \mathcal{A}_{s_2}(\mathbf{C}^{d_2})) \subseteq \mathcal{L}(\mathcal{A}_{s_1}(\mathbf{C}^{d_1}), \mathcal{A}'_{s_3}(\mathbf{C}^{d_3})).$$

Next we prove $\widehat{\mathcal{B}}_{(s_3, s_2)} \circ \widehat{\mathcal{C}}_{(s_2, s_1)} \subseteq \widehat{\mathcal{C}}_{(s_3, s_1)}$. Suppose $K_1 \in \widehat{\mathcal{C}}_{(s_2, s_1)}(W_1)$ and $K_2 \in \widehat{\mathcal{B}}_{(s_3, s_2)}(W_2)$, and let $K_3 = K_2 \circ K_1$, with coefficients $c(K_1; \alpha_2, \alpha_1)$, $c(K_2; \alpha_3, \alpha_2)$ and $c(K_3; \alpha_3, \alpha_1)$, $\alpha_j \in \mathbf{N}^{d_j}$, in their power series expansions. Also let $\vartheta_{r,s}$ be the same as in Definition 1.3. Then

$$c(K_3; \alpha_3, \alpha_1) = \sum_{\beta \in \mathbf{N}^{d_2}} c(K_2; \alpha_3, \beta) c(K_1; \beta, \alpha_1).$$

By the definitions, there is an $r_{0,1} > 0$ such that

$$|c(K_1; \alpha_2, \alpha_1)| \lesssim \frac{\vartheta_{r_1, s_1}(\alpha_1)}{\vartheta_{r_{0,1}, s_2}(\alpha_2)}$$

holds for every $r_1 > 0$, and for every $r_2 > 0$ there is an $r_{0,2} > 0$ such that

$$|c(K_2; \alpha_3, \alpha_2)| \lesssim \frac{\vartheta_{r_2, s_2}(\alpha_2)}{\vartheta_{r_{0,2}, s_3}(\alpha_3)} \quad (2.11)$$

holds. Hence, if we let $r_2 < r_{0,1}$ and choose $r_{0,2} > 0$ such that (2.11) holds, then

$$|c(K_3; \alpha_3, \alpha_1)| \lesssim \sum_{\beta \in \mathbf{N}^{d_2}} \left(\frac{\vartheta_{r_2, s_2}(\beta)}{\vartheta_{r_{0,1}, s_2}(\beta)} \right) \frac{\vartheta_{r_1, s_1}(\alpha_1)}{\vartheta_{r_{0,2}, s_3}(\alpha_3)} \asymp \frac{\vartheta_{r_1, s_1}(\alpha_1)}{\vartheta_{r_{0,2}, s_3}(\alpha_3)}$$

for every $r_1 > 0$. This is the same as $K_3 \in \widehat{\mathcal{C}}_{(s_3, s_1)}(W_3)$, and the assertion follows. In the same way, we obtain the other assertions. The details are left for the reader. \square

In Section 4 we present an analogy of Theorem 2.9 for Wick operators (see Theorem 4.17).

3. BINOMIAL OPERATORS AND THEIR CONTINUITY PROPERTIES

In this section we consider linear and bilinear operators of binomial types which act on subspaces of $\ell'_0(\mathbf{N}^{2d})$ and which contain binomial expressions. In the interplay between coefficients of Wick symbols and the kernels to Wick operators, the linear binomial operators are the actions on the coefficients in power series expansions, when passing between operator kernels, Wick symbols and anti-Wick symbols of linear operators. The bilinear binomial operators are in similar ways the actions on the coefficients in power series expansions, which correspond to pure multiplications, their adjoint actions, and compositions of Wick operators.

3.1. Binomial type operators. Let $t \in \mathbf{C}$. Essential parts of our investigations concerns continuity properties of the linear binomial operators

$$(\mathcal{T}_{0,t}c)(\alpha, \beta) = \sum_{\gamma \leq \alpha, \beta} \left(\binom{\alpha}{\gamma} \binom{\beta}{\gamma} \right)^{1/2} t^{|\gamma|} c(\alpha - \gamma, \beta - \gamma), \quad t \in \mathbf{C}, \quad (3.1)$$

when $c \in \ell'_0(\mathbf{N}^{2d})$, and their formal ℓ^2 adjoint, which are given by $\mathcal{T}_{0,t}^*c$, where

$$(\mathcal{T}_{0,t}^*c)(\alpha, \beta) = \sum_{\gamma \in \mathbf{N}^d} \left(\binom{\alpha + \gamma}{\gamma} \binom{\beta + \gamma}{\gamma} \right)^{1/2} t^{|\gamma|} c(\alpha + \gamma, \beta + \gamma), \quad t \in \mathbf{C}, \quad (3.2)$$

when $c \in \ell_0(\mathbf{N}^{2d})$. In terms of the operator

$$(S_0c)(\alpha, \beta) = i^{|\alpha + \beta|} c(\alpha, \beta), \quad \alpha, \beta \in \mathbf{N}^d, \quad c \in \ell'_0(\mathbf{N}^{2d}), \quad (3.3)$$

we observe that

$$\mathcal{T}_{0,-t} = S_0^{-1} \circ \mathcal{T}_{0,t} \circ S_0 \quad \text{and} \quad \mathcal{T}_{0,-t}^* = S_0^{-1} \circ \mathcal{T}_{0,t}^* \circ S_0 \quad (3.4)$$

in the domains of $\mathcal{T}_{0,-t}$ and $\mathcal{T}_{0,-t}^*$.

The operators $\mathcal{T}_{0,1}$ and $\mathcal{T}_{0,1}^*$ are linked to transitions between kernel operators and Wick operators, and to transitions between Wick and anti-Wick operators. In fact, for any fixed $t \in \mathbf{C}$, let \mathcal{T}_t be the map on $\widehat{A}(\mathbf{C}^{2d})$, given by

$$(\mathcal{T}_t a)(z, w) = e^{t(z,w)} a(z, w), \quad a \in \widehat{A}(\mathbf{C}^{2d}) = \widehat{\mathcal{A}}_{b_1}(\mathbf{C}^{2d}), \quad a \in \widehat{A}(\mathbf{C}^{2d}). \quad (3.5)$$

It follows in particular from (1.55), (2.4) and (2.4)' that

$$T_K = \text{Op}_{\mathfrak{W}}(a) \quad \Leftrightarrow \quad K = \mathcal{T}_1 a, \quad (3.6)$$

when $a \in \widehat{A}(\mathbf{C}^{2d})$, because

$$T_K = \text{Op}_{\mathfrak{W}}(a) \quad \text{with} \quad K(z, w) = e^{(z,w)} a(z, w) = (\mathcal{T}_1 a)(z, w), \quad (3.6)'$$

As a consequence of the following proposition we have

$$\mathcal{T}_{0,t} c(a; \cdot) = c(\mathcal{T}_t a; \cdot) \quad (3.7)$$

when $a \in \widehat{\mathcal{A}}_0(\mathbf{C}^{2d})$, which gives the link between the operator $\mathcal{T}_{0,1}$ and transitions between kernel and Wick operators. We omit the proof because the result is essentially a restatement of [31, Theorem 2.6].

Proposition 3.1. *Let $s, s_0 \in \overline{\mathbf{R}}_b$ be such that $s < \frac{1}{2}$ and $s_0 \leq \frac{1}{2}$, $t \in \mathbf{C}$ and let T_A be the map in (1.26). Then \mathcal{T}_t from $\widehat{\mathcal{A}}_0(\mathbf{C}^{2d})$ to $\widehat{A}(\mathbf{C}^{2d})$ extends uniquely to continuous mappings on $\widehat{\mathcal{A}}'_s(\mathbf{C}^{2d})$ and on $\widehat{\mathcal{A}}'_{0,s_0}(\mathbf{C}^{2d})$, and the diagrams*

$$\begin{array}{ccc} \ell'_s(\mathbf{N}^{2d}) & \xrightarrow{\mathcal{T}_{0,t}} & \ell'_s(\mathbf{N}^{2d}) & & \ell'_{0,s_0}(\mathbf{N}^{2d}) & \xrightarrow{\mathcal{T}_{0,t}} & \ell'_{0,s_0}(\mathbf{N}^{2d}) \\ T_A \downarrow & & \downarrow T_A & \text{and} & T_A \downarrow & & \downarrow T_A \\ \widehat{\mathcal{A}}'_s(\mathbf{C}^{2d}) & \xrightarrow{\mathcal{T}_t} & \widehat{\mathcal{A}}'_s(\mathbf{C}^{2d}) & & \widehat{\mathcal{A}}'_{0,s_0}(\mathbf{C}^{2d}) & \xrightarrow{\mathcal{T}_t} & \widehat{\mathcal{A}}'_{0,s_0}(\mathbf{C}^{2d}) \end{array} \quad (3.8)$$

commute.

Here we observe the misprints in the commutative diagram (2.5) in [31], where $T_{\mathcal{H}}$ and $\widehat{\mathcal{A}}_{b_1}(\mathbf{C}^{2d})$ should be replaced by T_A and $\widehat{\mathcal{A}}'_{b_1}(\mathbf{C}^{2d})$, respectively, at each occurrence.

By (3.5), the commutative diagrams (3.8) and that $\mathcal{T}_{0,\bar{t}}^*$ is the ℓ^2 adjoint of $\mathcal{T}_{0,t}$ it follows that

$$\begin{aligned} \mathcal{T}_{0,t_1} \circ \mathcal{T}_{0,t_2} &= \mathcal{T}_{0,t_1+t_2}, & \mathcal{T}_{0,t}^{-1} &= \mathcal{T}_{0,-t}, \\ \mathcal{T}_{0,t_1}^* \circ \mathcal{T}_{0,t_2}^* &= \mathcal{T}_{0,t_1+t_2}^*, & (\mathcal{T}_{0,t}^*)^{-1} &= \mathcal{T}_{0,-t}^*, \quad t, t_1, t_2 \in \mathbf{C}, \end{aligned} \quad (3.9)$$

since similar facts hold true for the operator \mathcal{T}_t .

The operator $\mathcal{T}_{0,t}^*$ is linked to the operator \mathcal{T}_t^* , given by

$$(\mathcal{T}_t^* a)(t_0 z, \bar{t}_0 w) = \pi^{-d} \int_{\mathbf{C}^d} a(t_0 w_1, \bar{t}_0 w_1) e^{-(z-w_1, w-w_1)} d\lambda(w_1), \quad t_0^2 = t, \quad (3.10)$$

when $a \in \widehat{\mathcal{A}}_0(\mathbf{C}^{2d})$. We observe that due to the definitions it follows that $a^{\text{aw}}(z, w)$ in (1.58) is equal to $\mathcal{T}_1^* a(z, w)$. Hence,

$$\text{Op}_{\mathfrak{W}}(\mathcal{T}_1^* a) = \text{Op}_{\mathfrak{W}}^{\text{aw}}(a) \quad \text{and} \quad \text{Op}_{\mathfrak{W}}^{\text{aw}}(\mathcal{T}_{-1}^* a) = \text{Op}_{\mathfrak{W}}(a) \quad (3.11)$$

when $a \in \widehat{\mathcal{A}}_0(\mathbf{C}^{2d})$.

By a straight-forward consequence of (B.1) in Appendix B it follows that the diagram

$$\begin{array}{ccc} \ell_0(\mathbf{N}^{2d}) & \xrightarrow{\mathcal{T}_{0,t}^*} & \ell_0(\mathbf{N}^{2d}) \\ T_A \downarrow & & \downarrow T_A \\ \widehat{\mathcal{A}}_0(\mathbf{C}^{2d}) & \xrightarrow{\mathcal{T}_t^*} & \widehat{\mathcal{A}}_0(\mathbf{C}^{2d}) \end{array} \quad (3.12)$$

commute. A combination of (3.11) and (3.12) then gives

$$c(a^{\text{aw}}; \cdot) = \mathcal{T}_{0,1}^* c(a; \cdot), \quad (3.13)$$

when (1.58) holds and $a \in \widehat{\mathcal{A}}_0(\mathbf{C}^{2d})$, which in turn is equivalent to

$$\text{Op}_{\mathfrak{W}}(b) = \text{Op}_{\mathfrak{W}}^{\text{aw}}(a) \quad \Leftrightarrow \quad c(b; \cdot) = \mathcal{T}_{0,1}^* c(a; \cdot), \quad (3.14)$$

when $a, b \in \widehat{\mathcal{A}}_0(\mathbf{C}^{2d})$. This gives the link between $\mathcal{T}_{0,\pm 1}^*$ and transitions between Wick and anti-Wick operators.

Next we introduce certain types of bilinear binomial operators.

Definition 3.2. Let $t \in \mathbf{C}$.

- (1) The bilinear operator $(c_1, c_2) \mapsto (c_1 \bullet_t c_2)$ from $\ell_0(\mathbf{N}^d) \times \ell_0(\mathbf{N}^d)$ to $\ell_0(\mathbf{N}^d)$ is given by

$$(c_1 \bullet_t c_2)(\alpha) \equiv \sum_{\gamma \leq \alpha} \binom{\alpha}{\gamma}^{\frac{1}{2}} t^{|\gamma|} c_1(\alpha - \gamma) c_2(\gamma). \quad (3.15)$$

- (2) The bilinear operator $(c_1, c_2) \mapsto (c_1 \bullet_{(t,\diamond)} c_2)$ from $\ell_0(\mathbf{N}^d) \times \ell_0(\mathbf{N}^d)$ to $\ell_0(\mathbf{N}^d)$ is given by

$$(c_1 \bullet_{(t,\diamond)} c_2)(\alpha) \equiv t^{|\alpha|} \sum_{\gamma \in \mathbf{N}^d} \binom{\alpha + \gamma}{\gamma}^{\frac{1}{2}} c_1(\gamma) c_2(\alpha + \gamma). \quad (3.16)$$

- (3) The bilinear operator $(c_1, c_2) \mapsto (c_1 \bullet_{(t,\#)} c_2)$ from $\ell_0(\mathbf{N}^{2d}) \times \ell_0(\mathbf{N}^{2d})$ to $\ell_0(\mathbf{N}^{2d})$ is given by

$$\begin{aligned} & (c_1 \bullet_{(t,\#)} c_2)(\alpha, \beta) \\ & \equiv \sum \left(\binom{\alpha}{\alpha_0} \binom{\beta}{\beta_0} \binom{\alpha - \alpha_0 + \gamma}{\gamma} \binom{\beta - \beta_0 + \gamma}{\gamma} \right)^{\frac{1}{2}} t^{|\gamma|} c_1(\alpha_0, \beta - \beta_0 + \gamma) c_2(\alpha - \alpha_0 + \gamma, \beta_0), \end{aligned} \quad (3.17)$$

where the sum is taken over all $\alpha_0, \beta_0, \gamma \in \mathbf{N}^d$ such that $\alpha_0 \leq \alpha$, $\beta_0 \leq \beta$.

It follows by straight-forward computations that

$$(c_1 \bullet_t c_2)(\alpha) = t^{|\alpha|} (c_2 \bullet_{1/t} c_1)(\alpha) \quad (3.18)$$

and

$$(c_1 \bullet_t c_2, c_3)_{\ell^2(\mathbf{N}^d)} = (c_2, \overline{c_1} \bullet_{\overline{t}, \diamond} c_3)_{\ell^2(\mathbf{N}^d)}, \quad (3.19)$$

$c_j \in \ell_0(\Lambda)$, $j = 1, 2, 3$.

The different products in Definition 3.2 are linked into different bilinear mappings for spaces of power series expansions. Let $W = \mathbf{C}^{d_2} \times \mathbf{C}^{d_1}$ and $K_1, K_2 \in \widehat{\mathcal{A}}_0(W)$. Then

$$\begin{aligned} K_1(z_2, z_1)K_2(z_2, z_1) &= \sum_{\alpha_j \beta_j} c(K_1; \alpha_2, \alpha_1)c(K_2; \beta_2, \beta_1)e_{\alpha_2}(z_2)e_{\beta_2}(z_2)e_{\alpha_1}(z_1)e_{\beta_1}(z_1) \\ &= \sum_{\alpha_1 \alpha_2} \left(\sum_{\gamma_j \leq \alpha_j} \left(\binom{\alpha_1}{\gamma_1} \binom{\alpha_2}{\gamma_2} \right)^{\frac{1}{2}} c(K_1; \alpha_2 - \gamma_2, \alpha_1 - \gamma_1)c(K_2; \gamma_2, \gamma_1) \right) e_{\alpha_2}(z_2)e_{\alpha_1}(z_1), \end{aligned}$$

and it follows that

$$c(K_1 \cdot K_2; \cdot) = c(K_1; \cdot) \bullet_1 c(K_2; \cdot). \quad (3.20)$$

By (3.19) it follows that $\bullet_{(1, \diamond)}$ is the adjoint operation of \bullet_1 on elements in $\ell_0(\Lambda)$. In order to find corresponding relationship for elements in $\widehat{\mathcal{A}}_0(W)$, we observe that the adjoint operation of z_j is ∂_{z_j} . In fact, if $F, G \in \mathcal{A}_0(\mathbf{C}^d)$, then it follows by integrating by parts that

$$(z_j F, G)_{A^2} = (F, \partial_j G)_{A^2} \quad \text{and} \quad (\partial_j F, G)_{A^2} = (F, z_j G)_{A^2}. \quad (3.21)$$

Hence, by letting \diamond be the multiplication on $\mathcal{A}_0(\mathbf{C}^d)$, given by the formula

$$(F_1 \diamond F_2, F)_{A^2} = (F_2, F_0 \cdot F)_{A^2}, \quad F_0(z) = \overline{F_1(\bar{z})}, \quad (3.22)$$

when $F, F_1, F_2 \in \mathcal{A}_0(\mathbf{C}^d)$, it follows that

$$(F_1 \diamond F_2)(z) = \sum_{\alpha \in \mathbf{N}^d} c(F_1, \alpha) \frac{\partial_z^\alpha F_2(z)}{\alpha!^{\frac{1}{2}}}. \quad (3.23)$$

Since $\text{Op}_{\mathfrak{W}}^{\text{aw}}(\bar{w}_j) = \partial_j = \partial_{z_j}$, it also follows that

$$F_1 \diamond F_2 = \text{Op}_{\mathfrak{W}}^{\text{aw}}(F_1)F_2 = F_1(\nabla_z)F_2.$$

In similar ways, if $W = \mathbf{C}^{d_2} \times \mathbf{C}^{d_1}$ and $K_1, K_2 \in \widehat{\mathcal{A}}_0(W)$, then $K_1 \diamond K_2$ is defined by

$$(K_1 \diamond K_2)(z_2, z_1) = \sum_{\alpha_1 \in \mathbf{N}^{d_1}} \sum_{\alpha_2 \in \mathbf{N}^{d_2}} c(K_1; \alpha_2, \alpha_1) \frac{\partial_{z_2}^{\alpha_2} \partial_{z_1}^{\alpha_1} K_2(z_2, z_1)}{(\alpha_1! \alpha_2!)^{\frac{1}{2}}}. \quad (3.24)$$

It follows that

$$(K_1 \diamond K_2, K)_{\widehat{A}^2} = (K_2, K_0 \cdot K)_{\widehat{A}^2}, \quad K_0(z_2, z_1) = \overline{K_1(\bar{z}_2, \bar{z}_1)}, \quad (3.25)$$

when $K, K_1, K_2 \in \widehat{\mathcal{A}}_0(W)$. A combination of (3.19), (3.20) and (3.25), now gives

$$c(K_1 \diamond K_2; \cdot) = c(K_1; \cdot) \bullet_{(1, \diamond)} c(K_2; \cdot). \quad (3.26)$$

Example 3.3. We observe that (3.23) and straight-forward computations give

$$e_\alpha \diamond e_\beta = \begin{cases} \binom{\beta}{\alpha}^{\frac{1}{2}} e_{\beta-\alpha}, & \text{when } \beta \geq \alpha, \\ 0, & \text{otherwise,} \end{cases} \quad (3.27)$$

The following lemma shows that $\bullet_{(1, \#)}$ in Definition 3.2 is linked to compositions of Wick operators on the symbol side.

Lemma 3.4. *Let $W = \mathbf{C}^d \times \mathbf{C}^d$, $\bullet_{(1, \#)}$ be the multiplication on $\ell_0(\mathbf{N}^d \times \mathbf{N}^d)$ given by (3.17), and let*

$$u_{\alpha, \beta}(z, w) = e_\alpha(z)e_\beta(\bar{w}) \in \widehat{\mathcal{A}}_0(W).$$

Then the following is true:

(1) if $\alpha_j, \beta_j \in \mathbf{N}^d$, $j = 1, 2$, then

$$\begin{aligned} & u_{\alpha_1, \beta_1} \#_{\mathfrak{Y}} u_{\alpha_2, \beta_2} \\ &= \sum_{\gamma \leq \alpha_2, \beta_1} \left(\binom{\alpha_2}{\gamma} \binom{\beta_1}{\gamma} \binom{\alpha_1 + \alpha_2 - \gamma}{\alpha_1} \binom{\beta_1 + \beta_2 - \gamma}{\beta_2} \right)^{\frac{1}{2}} u_{\alpha_1 + \alpha_2 - \gamma, \beta_1 + \beta_2 - \gamma}; \end{aligned} \quad (3.28)$$

(2) if $a_1, a_2 \in \widehat{\mathcal{A}}_0(W)$, then

$$c(a_1 \#_{\mathfrak{Y}} a_2; \cdot) = c(a_1; \cdot) \bullet_{(1, \#)} c(a_2; \cdot). \quad (3.29)$$

Proof. By using that

$$I_{\gamma, \delta} = \pi^{-d} \int_{\mathbf{C}^d} w^\gamma \bar{w}^\delta e^{-|w|^2} d\lambda(w) = \begin{cases} \gamma!, & \gamma = \delta \\ 0, & \gamma \neq \delta, \end{cases}$$

we get

$$\begin{aligned} \pi^{-d} \int_{\mathbf{C}^d} e_\alpha(w_1) e_\beta(\bar{w}_1) e^{-|z-w_1|^2} d\lambda(w_1) &= \frac{1}{\pi^d (\alpha! \beta!)^{\frac{1}{2}}} \sum_{\gamma \leq \alpha} \sum_{\delta \leq \beta} \binom{\alpha}{\gamma} \binom{\beta}{\delta} z^{\alpha-\gamma} \bar{z}^{\beta-\delta} I_{\gamma, \delta} \\ &= \sum_{\gamma \leq \alpha, \beta} \left(\binom{\alpha}{\gamma} \binom{\beta}{\gamma} \right)^{\frac{1}{2}} e_{\alpha-\gamma}(z) e_{\beta-\delta}(\bar{z}). \end{aligned}$$

This gives

$$\begin{aligned} \pi^{-d} \int_{\mathbf{C}^d} e_\alpha(w_1) e_\beta(\bar{w}_1) e^{-(z-w_1, w-w_1)} d\lambda(w_1) \\ = \sum_{\gamma \leq \alpha, \beta} \left(\binom{\alpha}{\gamma} \binom{\beta}{\gamma} \right)^{\frac{1}{2}} e_{\alpha-\gamma}(z) e_{\beta-\delta}(\bar{w}). \end{aligned} \quad (3.30)$$

Hence, if $F_{\alpha, \beta}(z, w)$ is the left-hand side of (3.30), then we obtain

$$\begin{aligned} (u_{\alpha_1, \beta_1} \#_{\mathfrak{Y}} u_{\alpha_2, \beta_2})(z, w) &= e_{\alpha_1}(z) e_{\beta_2}(\bar{w}) F_{\alpha_2, \beta_1}(z, w) \\ &= \sum_{\gamma \leq \alpha_2, \beta_1} \left(\binom{\alpha_2}{\gamma} \binom{\beta_1}{\gamma} \right)^{\frac{1}{2}} (e_{\alpha_2-\gamma}(z) e_{\alpha_1}(z)) (e_{\beta_1-\delta}(\bar{w}) e_{\beta_2}(\bar{w})), \end{aligned}$$

which after some straight-forward computations lead to the right-hand side of (3.28). This gives (1).

Let

$$C_1(\alpha, \beta, \gamma) = \left(\binom{\alpha_2}{\gamma} \binom{\beta_1}{\gamma} \binom{\alpha_1 + \alpha_2 - \gamma}{\alpha_1} \binom{\beta_1 + \beta_2 - \gamma}{\beta_2} \right)^{\frac{1}{2}}$$

and

$$C_2(\alpha, \beta, \gamma) = \left(\binom{\alpha_2}{\alpha_1} \binom{\beta_1}{\beta_2} \binom{\alpha_2 - \alpha_1 + \gamma}{\gamma} \binom{\beta_1 - \beta_2 + \gamma}{\gamma} \right)^{\frac{1}{2}},$$

for admissible $\alpha = (\alpha_1, \alpha_2) \in \mathbf{N}^{2d}$, $\beta = (\beta_1, \beta_2) \in \mathbf{N}^{2d}$ and $\gamma \in \mathbf{N}^d$. By (1) we have

$$\begin{aligned} & (a_1 \#_{\mathfrak{B}} a_2)(z, w) \\ &= \sum_{\alpha_j, \beta_j} \sum_{\gamma \leq \alpha_2, \beta_1} C_1(\alpha, \beta, \gamma) c(a_1; \alpha_1, \beta_1) c(a_2; \alpha_2, \beta_2) e_{\alpha_1 + \alpha_2 - \gamma}(z) e_{\beta_1 + \beta_2 - \gamma}(\bar{w}) \\ &= \sum_{\alpha_1, \beta_2} \sum_{\gamma} \sum_{\alpha_2, \beta_1 \geq \gamma} C_1(\alpha, \beta, \gamma) c(a_1; \alpha_1, \beta_1) c(a_2; \alpha_2, \beta_2) e_{\alpha_1 + \alpha_2 - \gamma}(z) e_{\beta_1 + \beta_2 - \gamma}(\bar{w}). \end{aligned}$$

By taking

$$\alpha_1, \quad \alpha_1 + \alpha_2 - \gamma, \quad \beta_1 + \beta_2 - \gamma, \quad \beta_2 \quad \text{and} \quad \gamma$$

as new variables of summations, we get

$$\begin{aligned} & (a_1 \#_{\mathfrak{B}} a_2)(z, w) \\ &= \sum_{\alpha_1, \beta_2} \left(\sum_{\gamma} \sum_{\substack{\alpha_1 \leq \alpha_2 \\ \beta_1 \leq \beta_1}} C_2(\alpha, \beta, \gamma) c(a_1; \alpha_1, \beta_1 - \beta_2 + \gamma) c(a_2; \alpha_2 - \alpha_1 + \gamma, \beta_2) \right) e_{\alpha_2}(z) e_{\beta_1}(\bar{w}). \end{aligned}$$

This is the same as (3.29), and the result follows. \square

3.2. Continuity for binomial operators. The assertion (1) in the following proposition was deduced in [31], and (2) follows from (1) and duality. The details are left for the reader.

Proposition 3.5. *Let $t \in \mathbf{C}$, $s, s_0 \in \overline{\mathbf{R}_b}$ be such that $s < \frac{1}{2}$ and $0 < s_0 \leq \frac{1}{2}$, $\mathcal{T}_{0,t}, S_0 \in \mathcal{L}(\ell'_0(\mathbf{N}^{2d}))$ be given by (3.1) and (3.3), and $\mathcal{T}_{0,t}^* \in \mathcal{L}(\ell_0(\mathbf{N}^{2d}))$ be given by (3.2). Then the following is true:*

- (1) $\mathcal{T}_{0,t}$ is a homeomorphism on $\ell'_0(\mathbf{N}^{2d})$ with the inverse $\mathcal{T}_{0,-t}$. Furthermore, $\mathcal{T}_{0,t}$ restricts to homeomorphisms on $\ell'_s(\mathbf{N}^{2d})$ and on $\ell'_{0,s_0}(\mathbf{N}^{2d})$, and $\mathcal{T}_{0,-t}$ can be obtained from (3.4);
- (2) $\mathcal{T}_{0,t}^*$ is a homeomorphism on $\ell_0(\mathbf{N}^{2d})$ with the inverse $\mathcal{T}_{0,-t}^*$. Furthermore, $\mathcal{T}_{0,t}^*$ is uniquely extendable to homeomorphisms on $\ell_s(\mathbf{N}^{2d})$ and on $\ell_{0,s_0}(\mathbf{N}^{2d})$, and $\mathcal{T}_{0,-t}^*$ can be obtained from (3.4).

We have now the following complementary result to Proposition 3.5.

Proposition 3.6. *Let $t \in \mathbf{C}$, $s, s_0 \in \overline{\mathbf{R}_b}$ be such that $s < \frac{1}{2}$ and $0 < s_0 \leq \frac{1}{2}$, $\mathcal{T}_{0,t}, S_0 \in \mathcal{L}(\ell'_0(\mathbf{N}^{2d}))$ given by (3.1) and (3.3), and $\mathcal{T}_{0,t}^* \in \mathcal{L}(\ell_0(\mathbf{N}^{2d}))$ given by (3.2). Then the following is true:*

- (1) the map $\mathcal{T}_{0,t}$ restricts to homeomorphisms on

$$\ell_{\mathcal{B},s}(\mathbf{N}^{2d}), \quad \ell_{\mathcal{B},s}^*(\mathbf{N}^{2d}), \quad \ell_{\mathcal{B},0,s_0}(\mathbf{N}^{2d}), \quad \ell_{\mathcal{B},0,s_0}^*(\mathbf{N}^{2d}), \quad (3.31)$$

with inverse $\mathcal{T}_{0,-t}$, and (3.4) holds;

- (2) the map $\mathcal{T}_{0,t}^*$ on $\ell_0(\mathbf{N}^{2d})$ extends uniquely to homeomorphisms on

$$\ell_{\mathcal{C},s}(\mathbf{N}^{2d}), \quad \ell_{\mathcal{C},s}^*(\mathbf{N}^{2d}), \quad \ell_{\mathcal{C},0,s_0}(\mathbf{N}^{2d}), \quad \ell_{\mathcal{C},0,s_0}^*(\mathbf{N}^{2d}), \quad (3.32)$$

with inverse $\mathcal{T}_{0,-t}^*$, and (3.4) holds.

Proof. We only prove (1) for the spaces $\ell_{\mathcal{B},s}(\mathbf{N}^{2d})$, as well as for $\ell_{\mathcal{B},0,s_0}(\mathbf{N}^{2d})$ when $s_0 = \frac{1}{2}$, and we only prove (2) for $\ell_{\mathcal{C},s}(\mathbf{N}^{2d})$. The other assertions follow by similar arguments and are left for the reader.

First suppose that $s < \frac{1}{2}$, $r_{0,j}, r_j, \theta > 0$, $j = 1, 2$, and that

$$|c(\alpha, \beta)| \lesssim \frac{\vartheta_{r_2,s}(\beta)}{\vartheta_{r_{0,2},s}(\alpha)}. \quad (3.33)$$

Then it follows from standard estimates of binomial coefficients that

$$\begin{aligned} |(\mathcal{T}_{0,t}c)(\alpha, \beta)\vartheta_{r_{0,1},s}(\alpha)/\vartheta_{r_1,s}(\beta)| &\lesssim \sum_{\gamma \leq \alpha, \beta} \binom{\alpha}{\gamma}^{\frac{1}{2}} \binom{\beta}{\gamma}^{\frac{1}{2}} |t|^{|\gamma|} \frac{\vartheta_{r_2,s}(\beta - \gamma)\vartheta_{r_{0,1},s}(\alpha)}{\vartheta_{r_{0,2},s}(\alpha - \gamma)\vartheta_{r_1,s}(\beta)} \\ &\leq (2 + 2|t|)^{|\alpha+\beta|} \sum_{\gamma \leq \alpha, \beta} \frac{\vartheta_{r_2,s}(\beta - \gamma)\vartheta_{r_{0,1},s}(\alpha)}{\vartheta_{r_{0,2},s}(\alpha - \gamma)\vartheta_{r_1,s}(\beta)}. \end{aligned} \quad (3.34)$$

Suppose that $c \in \ell_{\mathcal{B},s}(\mathbf{N}^{2d})$ and let $r_1 > 0$ be arbitrary but fixed. Let $c_0 \in (0, 1)$ be a small constant which shall be chosen later on. By Lemma 1.2, we may choose $r_2 < r_1$ and then $r_{0,2} \in (0, r_2)$ such that

$$\vartheta_{r_2,s}(\beta - \gamma)\vartheta_{r_2,s}(\gamma) \lesssim \vartheta_{c_0 r_1,s}(\beta)$$

and that (3.33) holds. Then (3.34) gives

$$\begin{aligned} |(\mathcal{T}_{0,t}c)(\alpha, \beta)\vartheta_{r_{0,1},s}(\alpha)/\vartheta_{r_1,s}(\beta)| &\lesssim (2 + 2|t|)^{|\alpha+\beta|} \frac{\vartheta_{c_0 r_1,s}(\beta)}{\vartheta_{r_1,s}(\beta)} \sum_{\gamma \leq \alpha, \beta} \frac{\vartheta_{r_{0,1},s}(\alpha)}{\vartheta_{r_{0,2},s}(\alpha - \gamma)\vartheta_{r_2,s}(\gamma)} \\ &\lesssim (2 + 2|t|)^{|\alpha+\beta|} \frac{\vartheta_{r_{0,1},s}(\alpha)\vartheta_{c_0 r_1,s}(\beta)}{\vartheta_{r_{0,2}/C,s}(\alpha)\vartheta_{r_1,s}(\beta)} \left(\sum_{\gamma \leq \alpha, \beta} 1 \right) \\ &\lesssim (3 + 2|t|)^{|\alpha+\beta|} \frac{\vartheta_{r_{0,1},s}(\alpha)\vartheta_{c_0 r_1,s}(\beta)}{\vartheta_{r_{0,2}/C,s}(\alpha)\vartheta_{r_1,s}(\beta)} \end{aligned}$$

Since $s < \frac{1}{2}$, it follows that

$$(3 + 2|t|)^{|\alpha|} \frac{\vartheta_{r_{0,1},s}(\alpha)}{\vartheta_{r_{0,2}/C,s}(\alpha)} \leq C_0 \quad \text{and} \quad (3 + 2|t|)^{|\beta|} \frac{\vartheta_{c_0 r_1,s}(\beta)}{\vartheta_{r_1,s}(\beta)} \leq C_0,$$

for some constant $C_0 > 0$ which is independent of α and β , provided c_0 and $r_{0,1} > 0$ are chosen small enough. A combination of these estimates shows that for every $r > 0$, there is an $r_0 > 0$ such that

$$|(\mathcal{T}_{0,t}c)(\alpha, \beta)| \lesssim \frac{\vartheta_{r,s}(\beta)}{\vartheta_{r_0,s}(\alpha)}, \quad (3.35)$$

which implies that $\mathcal{T}_{0,t}c \in \ell_{\mathcal{B},s}(\mathbf{N}^{2d})$. Since it is clear that the choice of r_0 depends continuously of the interplay between r_2 and $r_{0,2}$ in (3.33), it also follows that $\mathcal{T}_{0,t}$ is continuous on $\ell_{\mathcal{B},s}(\mathbf{N}^{2d})$, which proves the asserted continuity for $\mathcal{T}_{0,t}$ on $\ell_{\mathcal{B},s}(\mathbf{N}^{2d})$.

Next suppose that $c \in \ell_{\mathcal{B},0,s_0}(\mathbf{N}^{2d})$ with $s_0 = \frac{1}{2}$. That is, for every $r > 0$, there is an $r_0 > 0$ such that

$$|c(\alpha, \beta)| \lesssim e^{-r|\alpha|+r_0|\beta|}. \quad (3.36)$$

We shall prove that for every $r > 0$, there is an $r_0 > 0$ such that

$$|(\mathcal{T}_{0,t}c)(\alpha, \beta)| \lesssim e^{-r|\alpha|+r_0|\beta|}, \quad (3.37)$$

and then it suffices to prove this when $r_0 \geq r$. By Cauchy-Schwartz inequality we get

$$\begin{aligned}
|(\mathcal{T}_{0,t}c)(\alpha, \beta)| &\lesssim \sum_{\gamma \leq \alpha, \beta} \binom{\alpha}{\gamma}^{\frac{1}{2}} \binom{\beta}{\gamma}^{\frac{1}{2}} |t|^{|\gamma|} e^{-r|\alpha-\gamma|+r_0|\beta-\gamma|} \\
&\leq e^{-r|\alpha|+r_0|\beta|} \left(\sum_{\gamma \leq \alpha} \binom{\alpha}{\gamma} |t|^{|\gamma|} \right)^{\frac{1}{2}} \left(\sum_{\gamma \leq \beta} \binom{\beta}{\gamma} |t|^{|\gamma|} \right)^{\frac{1}{2}} \\
&= e^{-(r-r_0)|\alpha|+(r_0+r_t)|\beta|},
\end{aligned}$$

where

$$r_t = \frac{1}{2} \ln(1 + |t|).$$

Since r can be chosen arbitrarily large, it follows that for every $r > 0$, there is an $r_0 > 0$ such that (3.37) holds. Consequently, $\mathcal{T}_{0,t}$ is continuous on $\ell_{\mathcal{B},0,s_0}(\mathbf{N}^{2d})$.

Next we prove (2) when $\mathcal{T}_{0,t}^*$ acts on $\ell_{\mathcal{C},s}(\mathbf{N}^{2d})$. Suppose that $c \in \ell_{\mathcal{C},s}(\mathbf{N}^{2d})$. Then there is an $r_0 = r_{0,2} > 0$ such that (3.33) holds for every $r = r_2 > 0$. We may assume that $r < c_0 r_0$ for some constant $c_0 \in (0, 1)$ which shall be determined later on.

By Lemma 1.2 we obtain

$$\begin{aligned}
|(\mathcal{T}_{0,t}^*c)(\alpha, \beta)| &\leq \sum_{\gamma} \left(\binom{\alpha + \gamma}{\gamma} \binom{\beta + \gamma}{\gamma} \right)^{\frac{1}{2}} |t|^{|\gamma|} |c(\alpha + \gamma, \beta + \gamma)| \\
&\lesssim 2^{\frac{1}{2}|\alpha+\beta|} \sum_{\gamma} |2t|^{|\gamma|} \frac{\vartheta_{r,s}(\beta + \gamma)}{\vartheta_{r_0,s}(\alpha + \gamma)} \\
&\lesssim \frac{2^{\frac{1}{2}|\alpha+\beta|} \vartheta_{C_1 r, s}(\beta)}{\vartheta_{r_0/C_1, s}(\alpha)} \cdot J \\
&\lesssim \frac{\vartheta_{C_2 r, s}(\beta)}{\vartheta_{r_0/C_2, s}(\alpha)} \cdot J,
\end{aligned}$$

for some constants $C_2 > C_1 \geq 1$ which are independent of c_0 , r and r_0 , where

$$J = \sum_{\gamma} |2t|^{|\gamma|} \frac{\vartheta_{C_1 c_0 r_0, s}(\gamma)}{\vartheta_{r_0/C_1, s}(\gamma)}.$$

In the last inequality we have used the fact that $s < \frac{1}{2}$.

By choosing c_0 small enough it follows that J is convergent. This shows that for some $r_0 > 0$, (3.35) holds for every $r > 0$. That is, $\mathcal{T}_{0,t}c \in \ell_{\mathcal{C},s}(\mathbf{N}^{2d})$ when $c \in \ell_{\mathcal{C},s}(\mathbf{N}^{2d})$. The continuity assertions of $\mathcal{T}_{0,t}$ on $\ell_{\mathcal{C},s}(\mathbf{N}^{2d})$ now follows from this fact and that r_0 in (3.35) depends continuously on $r_{0,2}$ in (3.33). This gives the result. \square

In the following three propositions we show that the bilinear mappings in Definition 3.2 can be extended in suitable ways.

Proposition 3.7. *Let $t \in \mathbf{C}$, $s_1, s_2 \in \overline{\mathbf{R}}_b$ be such that $s_1, s_2 < \frac{1}{2}$, $s = (s_1, s_2)$, $\Lambda = \mathbf{N}^{d_2} \times \mathbf{N}^{d_1}$ and let U_t be the map from $\ell_0(\Lambda) \times \ell_0(\Lambda)$ to $\ell_0(\Lambda)$, given by $U_t(c_1, c_2) = c_1 \bullet_t c_2$, where $c_1 \bullet_t c_2$ is as in Definition 3.2. Then U_t is uniquely extendable to continuous*

mappings

$$\begin{aligned}
U_t &: \ell_{\mathcal{A},s}(\Lambda) \times \ell_{\mathcal{A},s}(\Lambda) \rightarrow \ell_{\mathcal{A},s}(\Lambda), & U_t &: \ell_{\mathcal{A},s}^*(\Lambda) \times \ell_{\mathcal{A},s}^*(\Lambda) \rightarrow \ell_{\mathcal{A},s}^*(\Lambda), \\
U_t &: \ell_{\mathcal{B},s}(\Lambda) \times \ell_{\mathcal{B},s}(\Lambda) \rightarrow \ell_{\mathcal{B},s}(\Lambda), & U_t &: \ell_{\mathcal{B},s}^*(\Lambda) \times \ell_{\mathcal{B},s}^*(\Lambda) \rightarrow \ell_{\mathcal{B},s}^*(\Lambda), \\
U_t &: \ell_{\mathcal{C},s}(\Lambda) \times \ell_{\mathcal{C},s}(\Lambda) \rightarrow \ell_{\mathcal{C},s}(\Lambda) & \text{and} & U_t &: \ell_{\mathcal{C},s}^*(\Lambda) \times \ell_{\mathcal{C},s}^*(\Lambda) \rightarrow \ell_{\mathcal{C},s}^*(\Lambda).
\end{aligned} \tag{3.38}$$

If instead $0 < s_1, s_2 \leq \frac{1}{2}$, then the same holds true with $\ell_{\mathcal{A},0,s}$, $\ell_{\mathcal{B},0,s}$ and $\ell_{\mathcal{C},0,s}$ in place of $\ell_{\mathcal{A},s}$, $\ell_{\mathcal{B},s}$ and $\ell_{\mathcal{C},s}$, respectively, at each occurrence.

Proposition 3.8. Let $t \in \mathbf{C}$, $s_1, s_2 \in \overline{\mathbf{R}}_b$ be such that $s_1, s_2 < \frac{1}{2}$, $s = (s_1, s_2)$, $\Lambda = \mathbf{N}^{d_2} \times \mathbf{N}^{d_1}$ and let $U_{t,\diamond}$ be the map from $\ell_0(\Lambda) \times \ell_0(\Lambda)$ to $\ell_0(\Lambda)$, given by $U_{t,\diamond}(c_1, c_2) = c_1 \bullet_{(t,\diamond)} c_2$, where $c_1 \bullet_{(t,\diamond)} c_2$ is as in Definition 3.2. Then $U_{t,\diamond}$ is uniquely extendable to continuous mappings

$$\begin{aligned}
U_{t,\diamond} &: \ell_{\mathcal{A},s}(\Lambda) \times \ell_{\mathcal{A},s}^*(\Lambda) \rightarrow \ell_{\mathcal{A},s}^*(\Lambda), & U_{t,\diamond} &: \ell_{\mathcal{A},s}^*(\Lambda) \times \ell_{\mathcal{A},s}(\Lambda) \rightarrow \ell_{\mathcal{A},s}(\Lambda), \\
U_{t,\diamond} &: \ell_{\mathcal{B},s}(\Lambda) \times \ell_{\mathcal{C},s}^*(\Lambda) \rightarrow \ell_{\mathcal{C},s}^*(\Lambda), & U_{t,\diamond} &: \ell_{\mathcal{B},s}^*(\Lambda) \times \ell_{\mathcal{C},s}(\Lambda) \rightarrow \ell_{\mathcal{C},s}(\Lambda), \\
U_{t,\diamond} &: \ell_{\mathcal{C},s}(\Lambda) \times \ell_{\mathcal{B},s}^*(\Lambda) \rightarrow \ell_{\mathcal{B},s}^*(\Lambda) & \text{and} & U_{t,\diamond} &: \ell_{\mathcal{C},s}^*(\Lambda) \times \ell_{\mathcal{B},s}(\Lambda) \rightarrow \ell_{\mathcal{B},s}(\Lambda).
\end{aligned} \tag{3.39}$$

If instead $0 < s_1, s_2 \leq \frac{1}{2}$, then the same holds true with $\ell_{\mathcal{A},0,s}$, $\ell_{\mathcal{B},0,s}$ and $\ell_{\mathcal{C},0,s}$ in place of $\ell_{\mathcal{A},s}$, $\ell_{\mathcal{B},s}$ and $\ell_{\mathcal{C},s}$, respectively, at each occurrence.

Proposition 3.9. Let $t \in \mathbf{C}$, $s \in \overline{\mathbf{R}}_b$ be such that $s < \frac{1}{2}$, $\Lambda = \mathbf{N}^{2d}$, and let $U_{t,\#}$ be the map from $\ell_0(\Lambda) \times \ell_0(\Lambda)$ to $\ell_0(\Lambda)$, given by $U_{t,\#}(c_1, c_2) = c_1 \bullet_{(t,\#)} c_2$, where $c_1 \bullet_{(t,\#)} c_2$ is as in Definition 3.2. Then $U_{t,\#}$ is uniquely extendable to continuous mappings

$$\begin{aligned}
U_{t,\#} &: \ell_{\mathcal{A},s}(\Lambda) \times \ell_{\mathcal{A},s}(\Lambda) \rightarrow \ell_{\mathcal{A},s}(\Lambda), \\
U_{t,\#} &: \ell_{\mathcal{A},s}^*(\Lambda) \times \ell_{\mathcal{B},s}(\Lambda) \rightarrow \ell_{\mathcal{A},s}^*(\Lambda), & U_{t,\#} &: \ell_{\mathcal{B},s}^*(\Lambda) \times \ell_{\mathcal{A},s}^*(\Lambda) \rightarrow \ell_{\mathcal{A},s}^*(\Lambda), \\
U_{t,\#} &: \ell_{\mathcal{B},s}(\Lambda) \times \ell_{\mathcal{B},s}(\Lambda) \rightarrow \ell_{\mathcal{B},s}(\Lambda), & U_{t,\#} &: \ell_{\mathcal{B},s}^*(\Lambda) \times \ell_{\mathcal{B},s}^*(\Lambda) \rightarrow \ell_{\mathcal{B},s}^*(\Lambda), \\
U_{t,\#} &: \ell_{\mathcal{C},s}(\Lambda) \times \ell_{\mathcal{C},s}(\Lambda) \rightarrow \ell_{\mathcal{C},s}(\Lambda), & U_{t,\#} &: \ell_{\mathcal{C},s}^*(\Lambda) \times \ell_{\mathcal{C},s}^*(\Lambda) \rightarrow \ell_{\mathcal{C},s}^*(\Lambda),
\end{aligned} \tag{3.40}$$

If instead $0 < s \leq \frac{1}{2}$, then the same holds true with $\ell_{\mathcal{A},0,s}$, $\ell_{\mathcal{B},0,s}$ and $\ell_{\mathcal{C},0,s}$ in place of $\ell_{\mathcal{A},s}$, $\ell_{\mathcal{B},s}$ and $\ell_{\mathcal{C},s}$, respectively, at each occurrence.

In each one of the propositions above, the different statements are proved by similar arguments. We only prove selections of these statements and leave the rest of the verifications for the reader.

Proof of Proposition 3.7. We only prove that U_t is uniquely extendable to a continuous map from $\ell_{\mathcal{B},s}(\Lambda) \times \ell_{\mathcal{B},s}(\Lambda)$ to $\ell_{\mathcal{B},s}(\Lambda)$ and from $\ell_{\mathcal{C},s}(\Lambda) \times \ell_{\mathcal{C},s}(\Lambda)$ to $\ell_{\mathcal{C},s}(\Lambda)$. The other cases follow by similar arguments and are left for the reader.

Let $\vartheta_{r,s}$ be as in Definition 1.3, $\alpha = (\alpha_2, \alpha_1) \in \Lambda$, and suppose that $c_j \in \ell'_0(\Lambda)$ satisfy

$$|c_j(\alpha)| = |c_j(\alpha_2, \alpha_1)| \lesssim \frac{\vartheta_{r_j, s_1}(\alpha_1)}{\vartheta_{r_{0,j}, s_2}(\alpha_2)}, \tag{3.41}$$

for some $r_j, r_{0,j} > 0$, $j = 1, 2$. By choosing $r = \max(r_1, r_2) > 0$ and $r_0 = \min(r_{0,1}, r_{0,2}) > 0$, it is clear that (3.41) holds with r and r_0 in place of r_j and $r_{0,j}$. By Lemma 1.2 we

get

$$\begin{aligned}
|(c_1 \bullet_t c_2)(\alpha_2, \alpha_1)| &\leq \sum_{\gamma \leq \alpha} \binom{\alpha}{\gamma}^{\frac{1}{2}} |t|^{|\gamma|} |c_1(\alpha - \gamma)c_2(\gamma)| \\
&\lesssim (1 + |t|)^{|\alpha|} \sum_{\gamma \leq \alpha} 2^{\frac{1}{2}|\alpha|} \frac{\vartheta_{r,s_1}(\alpha_1 - \gamma_1)\vartheta_{r,s_1}(\gamma_1)}{\vartheta_{r_0,s_2}(\alpha_2 - \gamma_2)\vartheta_{r_0,s_2}(\gamma_2)} \\
&\lesssim (2 + 2|t|)^{|\alpha|} \sum_{\gamma \leq \alpha} \frac{\vartheta_{C_0 r, s_1}(\alpha_1)}{\vartheta_{r_0/C_0, s_2}(\alpha_2)} \\
&\lesssim (3 + 2|t|)^{|\alpha|} \frac{\vartheta_{C_0 r, s_1}(\alpha_1)}{\vartheta_{r_0/C_0, s_2}(\alpha_2)} \lesssim \frac{\vartheta_{C r, s_1}(\alpha_1)}{\vartheta_{r_0/C, s_2}(\alpha_2)},
\end{aligned} \tag{3.42}$$

for some constants $C > C_0 > 1$ which are independent of r and r_0 . In the last step we have used $s_1, s_2 < \frac{1}{2}$.

If $c_j \in \ell_{\mathcal{B},s}(\Lambda)$, $j = 1, 2$, then for every $r = r_j > 0$, there is an $r_0 = r_{0,j} > 0$ such that (3.41) holds. By (3.42) it follows that for every $r > 0$, there is an $r_0 > 0$ such that (3.41) holds with $c_1 \bullet_t c_2$ in place of c_j . This implies that U_t is extendable to a continuous map from $\ell_{\mathcal{B},s}(\Lambda) \times \ell_{\mathcal{B},s}(\Lambda)$ to $\ell_{\mathcal{B},s}(\Lambda)$. Since $\ell_0(\Lambda)$ is dense in $\ell_{\mathcal{B},s}(\Lambda)$, it also follows that the continuous extension of U_t is unique.

If $c_j \in \ell_{\mathcal{C},s}(\Lambda)$, $j = 1, 2$, then there is an $r_0 = r_{0,j} > 0$ such that (3.41) holds for every $r = r_j > 0$. By (3.42) it follows that there is an $r_0 > 0$ such that (3.41) holds for every $r > 0$, with $c_1 \bullet_t c_2$ in place of c_j . Hence U_t extends to a continuous map from $\ell_{\mathcal{C},s}(\Lambda) \times \ell_{\mathcal{C},s}(\Lambda)$ to $\ell_{\mathcal{C},s}(\Lambda)$. Since $\ell_0(\Lambda)$ is dense in $\ell_{\mathcal{C},s}(\Lambda)$, it also follows that U_t on $\ell_{\mathcal{C},s}(\Lambda)$ is uniquely defined, which gives the assertion. \square

Proof of Proposition 3.8. We only prove that $U = U_{t,\diamond}$ extends uniquely to a continuous map from $\ell_{\mathcal{C},s}^*(\Lambda) \times \ell_{\mathcal{B},s}(\Lambda)$ to $\ell_{\mathcal{B},s}(\Lambda)$. The other assertions follow by similar arguments and are left for the reader.

Since $\ell_{\mathcal{C},s}^*(\Lambda)$ is invariant under the map

$$c(\alpha_2, \alpha_1) \mapsto t^{|\alpha_1|+|\alpha_2|} c(\alpha_2, \alpha_1),$$

it suffices to prove the result for $t = 1$.

First we prove the continuity extension of U . Let $c_1 \in \ell_{\mathcal{C},s}^*(\Lambda)$ and $c_2 \in \ell_{\mathcal{B},s}(\Lambda)$. Then it follows that for some $r_0 > 0$ one has

$$|c_1(\alpha_2, \alpha_1)| \lesssim \frac{\vartheta_{r,s_2}(\alpha_2)}{\vartheta_{r_0,s_1}(\alpha_1)}, \quad (\alpha_2, \alpha_1) \in \Lambda, \tag{3.43}$$

for every $r > 0$. It also follows that for every $r > 0$, there is an $r_0 > 0$ such that

$$|c_2(\alpha_2, \alpha_1)| \lesssim \frac{\vartheta_{r,s_1}(\alpha_1)}{\vartheta_{r_0,s_2}(\alpha_2)}, \quad (\alpha_2, \alpha_1) \in \Lambda. \tag{3.44}$$

Let $r_{0,1} > 0$ be fixed such that (3.43) holds for every $r > 0$, with $r_{0,1}$ in place of r_0 . Also let $C > 1$ be a constant which only dependent on s_1, s_2 , let $r_2 \in (0, r_{0,1}/C)$ be arbitrary and choose $r_{0,2} > 0$ such that (3.44) holds with r_2 and $r_{0,2}$ in place of r and r_0 . Finally let $r_1 \in (0, r_{0,2}/C)$ be arbitrary.

If C is chosen large enough, and using the same notations as in Lemma 3.4, then it follows from Lemma 1.2 that

$$\begin{aligned} \binom{\alpha + \gamma}{\gamma}^{\frac{1}{2}} |c_1(\gamma)c_2(\alpha + \gamma)| &\lesssim 2^{|\alpha + \gamma|} \frac{\vartheta_{r_1, s_2}(\gamma_2)\vartheta_{r_2, s_1}(\alpha_1 + \gamma_1)}{\vartheta_{r_0, 1, s_1}(\gamma_1)\vartheta_{r_0, 2, s_2}(\alpha_2 + \gamma_2)} \\ &\lesssim 2^{-|\alpha + \gamma|} \frac{\vartheta_{Cr_2, s_1}(\alpha_1)}{\vartheta_{r_0, 2/C, s_2}(\alpha_2)}. \end{aligned}$$

Hence for the series on the right-hand side of (3.16) we have

$$\begin{aligned} \sum_{\gamma \in \Lambda} \binom{\alpha + \gamma}{\gamma}^{\frac{1}{2}} |c_1(\gamma_2, \gamma_1)c_2(\alpha_2 + \gamma_2, \alpha_1 + \gamma_1)| &\lesssim \sum_{\gamma \in \Lambda} 2^{-|\alpha + \gamma|} \frac{\vartheta_{Cr_2, s_1}(\alpha_1)}{\vartheta_{r_0, 2/C, s_2}(\alpha_2)} \\ &= \frac{\vartheta_{Cr_2, s_1}(\alpha_1)}{\vartheta_{r_0, 2/C, s_2}(\alpha_2)}. \end{aligned}$$

It follows that the right-hand side of (3.16) is convergent, and by defining $c_1 \bullet_{(1, \diamond)} c_2$ by (3.16), it follows that for every $r > 0$, there is an $r_0 > 0$ such that

$$|(c_1 \bullet_{(1, \diamond)} c_2)(\alpha)| \lesssim \frac{\vartheta_{r, s_1}(\alpha_1)}{\vartheta_{r_0, s_2}(\alpha_2)}. \quad (3.45)$$

This implies that $c_1 \bullet_{(1, \diamond)} c_2 \in \ell_{\mathcal{B}, s}(\Lambda)$. Furthermore, the involved parameters of the right-hand side of (3.45) depend continuously on the involved parameters of the right-hand sides of (3.43) and (3.44), which proves the continuity assertion.

Finally, the uniqueness of the extension follows from the fact that $\ell_0(\Lambda)$ is dense in $\ell_{\mathcal{B}, s}(\Lambda)$ and $\ell_{\mathcal{C}, s}^*(\Lambda)$, and the result follows. \square

Proof of Proposition 3.9. We only prove that $U = U_{t, \#}$ extends uniquely to continuous products on $\ell_{\mathcal{B}, s}(\mathbf{N}^{2d})$ and on $\ell_{\mathcal{C}, s}(\mathbf{N}^{2d})$. The other assertions follows by similar arguments and are left for the reader.

Let

$$C_0(\alpha, \alpha_0, \beta, \beta_0, \gamma) = \left(\binom{\alpha}{\alpha_0} \binom{\beta}{\beta_0} \binom{\alpha - \alpha_0 + \gamma}{\gamma} \binom{\beta - \beta_0 + \gamma}{\gamma} \right)^{\frac{1}{2}},$$

when $\alpha, \alpha_0, \beta, \beta_0, \gamma \in \mathbf{N}^d$ satisfy $\alpha_0 \leq \alpha$ and $\beta_0 \leq \beta$ (cf. $C_2(\alpha, \beta, \gamma)$ in the proof of Lemma 3.4). We observe

$$|C_0(\alpha, \alpha_0, \beta, \beta_0, \gamma)| \leq 2^{|\alpha + \beta + \gamma|}. \quad (3.46)$$

Suppose that $c_j \in \ell_{\mathcal{B}, s}(\mathbf{N}^{2d})$, $j = 1, 2$. Then it follows from Lemma 1.2 and (3.41) for $s_j = s$ that for every $r_j > 0$, there is an $r_{0, j} > 0$ such that

$$|c_1(\alpha_0, \beta - \beta_0 + \gamma)| \lesssim \frac{\vartheta_{r_1, s}(\beta - \beta_0)\vartheta_{r_1, s}(\gamma)}{\vartheta_{r_0, 1}(\alpha_0)} \quad (3.47)$$

and

$$|c_2(\alpha - \alpha_0 + \gamma, \beta_0)| \lesssim \frac{\vartheta_{r_2, s}(\beta_0)}{\vartheta_{r_0, 2, s}(\alpha - \alpha_0)\vartheta_{r_0, 2, s}(\gamma)}. \quad (3.48)$$

Let $r > 0$ be arbitrary, $C > 1$ be a constant which is independent of r which is specified later and let $r_2 > 0$ be such that $Cr_2 < r$. Choose $r_{0, 2} > 0$ such that (3.48) holds. Now choose $r_1 > 0$ such that $r_1 < r_2$ and $r_1 < r_{0, 2}/2$ and then choose $r_{0, 1} > 0$ such that $r_{0, 1} < r_{0, 2}$ and (3.47) holds.

If C is chosen large enough, then it follows from Lemma 1.2 again that

$$\begin{aligned} |c_1(\alpha_0, \beta - \beta_0 + \gamma)c_2(\alpha - \alpha_0 + \gamma, \beta_0)| &\lesssim \frac{\vartheta_{r_1,s}(\beta - \beta_0)\vartheta_{r_1,s}(\gamma)\vartheta_{r_2,s}(\beta_0)}{\vartheta_{r_0,1,s}(\alpha_0)\vartheta_{r_0,2,s}(\alpha - \alpha_0)\vartheta_{r_0,2,s}(\gamma)} \\ &\lesssim \frac{2^{-|\alpha+\beta|}2^{-|\alpha_0+\beta_0|}\vartheta_{Cr_2,s}(\beta)}{\vartheta_{r_0,1/C,s}(\alpha)\vartheta_{r_1,s}(\gamma)} \lesssim \frac{2^{-|\alpha+\beta|}2^{-|\alpha_0+\beta_0|}\vartheta_{r,s}(\beta)}{\vartheta_{r_0,s}(\alpha)\vartheta_{r_1,s}(\gamma)}, \end{aligned} \quad (3.49)$$

when $r_0 = r_{0,1}/C > 0$. By combining these estimates we obtain

$$|(c_1 \bullet_{(t,\#)} c_2)(\alpha, \beta)| \lesssim \left(\sum_{\gamma} \frac{(2 + 2|t|)^{|\gamma|}}{\vartheta_{r_1,s}(\gamma)} \right) \left(\sum_{\alpha_0, \beta_0} 2^{-|\alpha_0+\beta_0|} \frac{\vartheta_{r,s}(\beta)}{\vartheta_{r_0,s}(\alpha)} \right) = \frac{\vartheta_{r,s}(\beta)}{\vartheta_{r_0,s}(\alpha)}, \quad (3.50)$$

and it follows that U on $\ell_0(\mathbf{N}^{2d})$ extends to a continuous products on $\ell_{\mathcal{B},s}(\mathbf{N}^{2d})$.

Suppose instead that $c_1 \in \ell_{\mathcal{C},s}(\mathbf{N}^{2d})$, $c_2 \in \ell_{\mathcal{C},s}(\mathbf{N}^{2d})$. Then for some $r_{0,1}, r_{0,2} > 0$ such that $r_{0,1} < r_{0,2}$ one has that (3.47) and (3.48) hold for every $r_1, r_2 > 0$ such that $r_1 < r_2$. Hence, if C and r_0 are chosen as in the first part of the proof, then (3.49) and (3.50) show that for some $r_0 > 0$ one has that

$$|(c_1 \bullet_{(t,\#)} c_2)(\alpha, \beta)| \lesssim \frac{\vartheta_{r,s}(\beta)}{\vartheta_{r_0,s}(\alpha)},$$

for every $r > 0$. This implies that U on $\ell_0(\mathbf{N}^{2d})$ extends to a continuous products on $\ell_{\mathcal{C},s}(\mathbf{N}^{2d})$. The uniqueness assertions follow from the fact that $\ell_0(\mathbf{N}^{2d})$ is dense in $\ell_{\mathcal{B},s}(\mathbf{N}^{2d})$ and in $\ell_{\mathcal{C},s}(\mathbf{N}^{2d})$. \square

4. CONTINUITY, COMPOSITION AND TRANSITIONS OF WICK AND ANTI-WICK OPERATORS

In this section we apply the results from the previous sections to deduce estimates on kernels, Wick and anti-Wick symbols of operators. Especially we show that in several situations, any linear and continuous operator on $\mathcal{A}_{0,s}(\mathbf{C}^d)$, $\mathcal{A}_s(\mathbf{C}^d)$, $\mathcal{A}'_s(\mathbf{C}^d)$ or on $\mathcal{A}'_{0,s}(\mathbf{C}^d)$ can be expressed as Wick or anti-Wick operators (see Theorems 4.8 and 4.9). In the last part of the section we complete some analysis in Section 3 in [32]. Here we deduce some estimates of Wick symbols of anti-Wick operators with Wick or anti-Wick symbols belonging to Hilbert spaces, related to $\widehat{\mathcal{A}}'_{0,1/2}(\mathbf{C}^{2d})$.

4.1. Continuity of the operators $\mathcal{T}_t, \mathcal{T}_t^*$. We start by discussing the operator \mathcal{T}_t . A combination of Proposition 3.6 (1) and (3.8) gives the following result, related to Proposition 3.1. The details are left for the reader.

Theorem 4.1. *Let $t \in \mathbf{C}$, $s, s_0 \in \overline{\mathbf{R}}_b$ be such that $s < \frac{1}{2}$ and $s_0 \leq \frac{1}{2}$, $\mathcal{T}_{0,t}$ be the map on $\ell'_0(\mathbf{N}^{2d})$ given by (3.1), and let \mathcal{T}_t be the map on $\widehat{\mathcal{A}}'_0(\mathbf{C}^{2d})$ given by (3.5). Then the following is true:*

- (1) \mathcal{T}_t restricts to homeomorphisms on each of the spaces

$$\widehat{\mathcal{B}}_{0,s_0}(\mathbf{C}^{2d}), \widehat{\mathcal{B}}_s(\mathbf{C}^{2d}), \widehat{\mathcal{B}}_{0,s_0}^*(\mathbf{C}^{2d}) \quad \text{and} \quad \widehat{\mathcal{B}}_s^*(\mathbf{C}^{2d}); \quad (4.1)$$

- (2) the diagram (3.8) commutes, after ℓ'_s and $\widehat{\mathcal{A}}'_s$ are replaced by $\ell_{\mathcal{B},s}$ and $\widehat{\mathcal{B}}_s$, by $\ell_{\mathcal{B},0,s_0}$ and $\widehat{\mathcal{B}}_{0,s_0}$, by $\ell_{\mathcal{B},s}^*$ and $\widehat{\mathcal{B}}_s^*$, or by $\ell_{\mathcal{B},0,s_0}^*$ and $\widehat{\mathcal{B}}_{0,s_0}^*$, respectively, at each occurrence.

Next we deduce extensions of the operator \mathcal{T}_t^* , and begin with the following, which follows from (2) in Proposition 3.5 and the commutative diagram (3.12). The details are left for the reader.

Theorem 4.2. *Let $t \in \mathbf{C}$, $s, s_0 \in \overline{\mathbf{R}}_b$ be such that $s < \frac{1}{2}$ and $s_0 \leq \frac{1}{2}$, and let $\mathcal{T}_{0,t}^*$ be the map on $\ell_0(\mathbf{N}^{2d})$ given by (3.2) and let \mathcal{T}_t^* be the map on $\widehat{\mathcal{A}}_0(\mathbf{C}^{2d})$ given by (3.10). Then \mathcal{T}_t^* is uniquely extendable to homeomorphisms on $\widehat{\mathcal{A}}_s(\mathbf{C}^{2d})$ and on $\widehat{\mathcal{A}}_{0,s_0}(\mathbf{C}^{2d})$. The diagram (3.12) commutes after ℓ_0 and $\widehat{\mathcal{A}}_0$ have been replaced by ℓ_s and $\widehat{\mathcal{A}}_s$, respectively, or by ℓ_{0,s_0} and $\widehat{\mathcal{A}}_{0,s_0}$, respectively, at each occurrence.*

In the same way we get the following by combining Proposition 3.6 (2) and (3.12). Again the details are left for the reader.

Theorem 4.3. *Let $t \in \mathbf{C}$, $s, s_0 \in \overline{\mathbf{R}}_b$ be such that $s < \frac{1}{2}$ and $s_0 \leq \frac{1}{2}$, $\mathcal{T}_{0,t}^*$ be the map on $\ell_0(\mathbf{N}^{2d})$ given by (3.2), and let \mathcal{T}_t^* be the map on $\widehat{\mathcal{A}}_0(\mathbf{C}^{2d})$ given by (3.10). Then the following is true:*

- (1) \mathcal{T}_t^* is uniquely extendable to homeomorphisms on each of the spaces

$$\widehat{\mathcal{C}}_{0,s_0}(\mathbf{C}^{2d}), \widehat{\mathcal{C}}_s(\mathbf{C}^{2d}), \widehat{\mathcal{C}}_{0,s_0}^*(\mathbf{C}^{2d}) \quad \text{and} \quad \widehat{\mathcal{C}}_s^*(\mathbf{C}^{2d}); \quad (4.2)$$

- (2) the diagram (3.12) commutes, after ℓ_0 and $\widehat{\mathcal{A}}_0$ are replaced by $\ell_{\mathcal{C},s}$ and $\widehat{\mathcal{C}}_s$, by $\ell_{\mathcal{C},0,s_0}$ and $\widehat{\mathcal{C}}_{0,s_0}$, by $\ell_{\mathcal{C},s}^*$ and $\widehat{\mathcal{C}}_s^*$, or by $\ell_{\mathcal{C},0,s_0}^*$ and $\widehat{\mathcal{C}}_{0,s_0}^*$, respectively, at each occurrence.

4.2. Continuity and relationships between kernel, Wick and anti-Wick operators. By (1.58)–(1.60), Proposition 3.6, Theorem 4.3, and the fact that (3.11) holds for $a \in \widehat{\mathcal{A}}_0(\mathbf{C}^{2d})$, we may now extend the domain of anti-Wick operators as in the following definition.

Definition 4.4. Let $s_1, s_2 \in \overline{\mathbf{R}}_b$ be such that $s_1 < \frac{1}{2}$ and $s_2 \leq \frac{1}{2}$.

- (1) If $a \in \widehat{\mathcal{C}}_{s_1}(\mathbf{C}^{2d})$ ($a \in \widehat{\mathcal{C}}_{s_1}^*(\mathbf{C}^{2d})$), then the anti-Wick operator $\text{Op}_{\mathfrak{W}}^{\text{aw}}(a)$ is the linear and continuous operator on $\mathcal{A}_{s_1}(\mathbf{C}^d)$ (on $\mathcal{A}'_{s_1}(\mathbf{C}^d)$), given by (3.11).
- (2) If $a \in \widehat{\mathcal{C}}_{0,s_2}(\mathbf{C}^{2d})$ ($a \in \widehat{\mathcal{C}}_{0,s_2}^*(\mathbf{C}^{2d})$), then the anti-Wick operator $\text{Op}_{\mathfrak{W}}^{\text{aw}}(a)$ is the linear and continuous operator on $\mathcal{A}_{0,s_2}(\mathbf{C}^d)$ (on $\mathcal{A}'_{0,s_2}(\mathbf{C}^d)$), given by (3.11).

We can now combine the results in previous sections with the kernel results in Section 2 to find equalities between classes of Wick, anti-Wick operators and kernel operators. As a first taste we have the following result which follows by a combination of (3.6), and Propositions 2.2, 2.3 and 3.1. The details are left for the reader. The result is also an immediate consequence of Theorems 2.7 and 2.8 in [31].

Proposition 4.5. *Let $j \in \{1, 2\}$, $s_1, s_2 \in \overline{\mathbf{R}}_b$ be such that $s_1 < \frac{1}{2}$ and $s_2 \leq \frac{1}{2}$. Then the mappings*

$$\widehat{\mathcal{A}}'_{s_1}(\mathbf{C}^{2d}) \ni a \mapsto \text{Op}_{\mathfrak{W}}(a) \in \mathcal{L}(\mathcal{A}_{s_1}(\mathbf{C}^d), \mathcal{A}'_{s_1}(\mathbf{C}^d))$$

and

$$\widehat{\mathcal{A}}'_{0,s_2}(\mathbf{C}^{2d}) \ni a \mapsto \text{Op}_{\mathfrak{W}}(a) \in \mathcal{L}(\mathcal{A}_{0,s_2}(\mathbf{C}^d), \mathcal{A}'_{0,s_2}(\mathbf{C}^d))$$

are isomorphisms.

The corresponding result for anti-Wick operators is the following. The result follows by combining (3.11) and Theorem 4.2. The details are left for the reader.

Theorem 4.6. *Let $j \in \{1, 2\}$, $s_1, s_2 \in \overline{\mathbf{R}}_b$ be such that $s_1 < \frac{1}{2}$ and $s_2 \leq \frac{1}{2}$. Then the mappings*

$$\widehat{\mathcal{A}}_{s_1}(\mathbf{C}^{2d}) \ni a \mapsto \text{Op}_{\mathfrak{W}}^{\text{aw}}(a) \in \text{Op}_{\mathfrak{W}}(\widehat{\mathcal{A}}_{s_1}(\mathbf{C}^{2d})) \quad (4.3)$$

and

$$\widehat{\mathcal{A}}_{0,s_2}(\mathbf{C}^{2d}) \ni a \mapsto \text{Op}_{\mathfrak{Y}}^{\text{aw}}(a) \in \text{Op}_{\mathfrak{Y}}(\widehat{\mathcal{A}}_{0,s_2}(\mathbf{C}^{2d})) \quad (4.4)$$

are isomorphisms.

We notice that if $s_1 = 0$, then $\widehat{\mathcal{A}}_{s_1}(\mathbf{C}^{2d}) = \widehat{\mathcal{A}}_0(\mathbf{C}^{2d})$ is the set of all polynomials $a(z, w)$ which are analytic in z and conjugated analytic in w . Hence (4.3) in this case means that the sets of Wick operators with polynomial symbols agree with the set of anti-Wick operators with polynomial symbols, which is well-known (see e.g. [6]). On the other hand, the other cases in Theorem 4.6, seems not to be available in the literature.

Remark 4.7. Note that $\widehat{\mathcal{A}}_{s_1}(\mathbf{C}^{2d})$ and $\widehat{\mathcal{A}}_{s_2}(\mathbf{C}^{2d})$ in Theorem 4.6 are images of those Pilipović spaces, which are not Gelfand-Shilov spaces, under the map $\Theta \circ \mathfrak{Y}_{2d}$. In particular, Theorem 4.6 gives further motivations for detailed studies of Pilipović spaces.

The next two results follows from (3.6), (3.11), Propositions 2.4 and 2.5, and Theorems 4.1 and 4.3. The details are left for the reader.

Theorem 4.8. *Let $s_1, s_2 \in \overline{\mathbf{R}}_b$ be such that $s_1 < \frac{1}{2}$ and $s_2 \leq \frac{1}{2}$. Then the mappings*

$$\begin{aligned} \widehat{\mathcal{B}}_{s_1}(\mathbf{C}^{2d}) \ni a &\mapsto \text{Op}_{\mathfrak{Y}}(a) \in \mathcal{L}(\mathcal{A}_{s_1}(\mathbf{C}^d)), \\ \widehat{\mathcal{B}}_{s_1}^*(\mathbf{C}^{2d}) \ni a &\mapsto \text{Op}_{\mathfrak{Y}}(a) \in \mathcal{L}(\mathcal{A}'_{s_1}(\mathbf{C}^d)), \\ \widehat{\mathcal{B}}_{0,s_2}(\mathbf{C}^{2d}) \ni a &\mapsto \text{Op}_{\mathfrak{Y}}(a) \in \mathcal{L}(\mathcal{A}_{0,s_2}(\mathbf{C}^d)) \end{aligned}$$

and

$$\widehat{\mathcal{B}}_{0,s_2}^*(\mathbf{C}^{2d}) \ni a \mapsto \text{Op}_{\mathfrak{Y}}(a) \in \mathcal{L}(\mathcal{A}'_{0,s_2}(\mathbf{C}^d)),$$

are isomorphisms.

Theorem 4.9. *Let $s_1, s_2 \in \overline{\mathbf{R}}_b$ be such that $s_1 < \frac{1}{2}$ and $s_2 \leq \frac{1}{2}$. Then the mappings*

$$\begin{aligned} \widehat{\mathcal{C}}_{s_1}(\mathbf{C}^{2d}) \ni a &\mapsto \text{Op}_{\mathfrak{Y}}^{\text{aw}}(a) \in \text{Op}_{\mathfrak{Y}}(\widehat{\mathcal{C}}_{s_1}(\mathbf{C}^{2d})), \\ \widehat{\mathcal{C}}_{s_1}^*(\mathbf{C}^{2d}) \ni a &\mapsto \text{Op}_{\mathfrak{Y}}^{\text{aw}}(a) \in \text{Op}_{\mathfrak{Y}}(\widehat{\mathcal{C}}_{s_1}^*(\mathbf{C}^{2d})), \\ \widehat{\mathcal{C}}_{0,s_2}(\mathbf{C}^{2d}) \ni a &\mapsto \text{Op}_{\mathfrak{Y}}^{\text{aw}}(a) \in \text{Op}_{\mathfrak{Y}}(\widehat{\mathcal{C}}_{0,s_2}(\mathbf{C}^{2d})) \end{aligned}$$

and

$$\widehat{\mathcal{C}}_{0,s_2}^*(\mathbf{C}^{2d}) \ni a \mapsto \text{Op}_{\mathfrak{Y}}^{\text{aw}}(a) \in \text{Op}_{\mathfrak{Y}}(\widehat{\mathcal{C}}_{0,s_2}^*(\mathbf{C}^{2d}))$$

are isomorphisms.

Remark 4.10. Let $s \in \overline{\mathbf{R}}_b$ be such that $s_1 < \frac{1}{2}$ and $s_2 \leq \frac{1}{2}$. By the preparing results which lead to Definition 4.4, and Theorem 4.9, it follows that

$$\begin{aligned} \{ \text{Op}_{\mathfrak{Y}}^{\text{aw}}(a) ; a \in \widehat{\mathcal{C}}_{s_1}(\mathbf{C}^{2d}) \} &= \{ \text{Op}_{\mathfrak{Y}}(a) ; a \in \widehat{\mathcal{C}}_{s_1}(\mathbf{C}^{2d}) \}, \\ \{ \text{Op}_{\mathfrak{Y}}^{\text{aw}}(a) ; a \in \widehat{\mathcal{C}}_{s_1}^*(\mathbf{C}^{2d}) \} &= \{ \text{Op}_{\mathfrak{Y}}(a) ; a \in \widehat{\mathcal{C}}_{s_1}^*(\mathbf{C}^{2d}) \}, \\ \{ \text{Op}_{\mathfrak{Y}}^{\text{aw}}(a) ; a \in \widehat{\mathcal{C}}_{0,s_2}(\mathbf{C}^{2d}) \} &= \{ \text{Op}_{\mathfrak{Y}}(a) ; a \in \widehat{\mathcal{C}}_{0,s_2}(\mathbf{C}^{2d}) \}, \\ \{ \text{Op}_{\mathfrak{Y}}^{\text{aw}}(a) ; a \in \widehat{\mathcal{C}}_{0,s_2}^*(\mathbf{C}^{2d}) \} &= \{ \text{Op}_{\mathfrak{Y}}(a) ; a \in \widehat{\mathcal{C}}_{0,s_2}^*(\mathbf{C}^{2d}) \}, \end{aligned} \quad (4.5)$$

and that

$$\text{Op}_{\mathfrak{Y}}^{\text{aw}}(a_1) \neq \text{Op}_{\mathfrak{Y}}^{\text{aw}}(a_2) \Leftrightarrow \text{Op}_{\mathfrak{Y}}(a_1) \neq \text{Op}_{\mathfrak{Y}}(a_2) \Leftrightarrow a_1 \neq a_2, \quad (4.6)$$

when a_1 and a_2 belong to any of the spaces in Theorem 4.9.

In Appendix C we have listed some identities of spaces of kernels, Wick and anti-Wick operators which are immediate consequences of Corollary 2.6, Proposition 4.5 and Theorems 4.6–4.9.

For other choices of s_1 and s_2 it seems that the equalities on the left-hand sides in Theorems 4.8 and 4.9 between sets of Wick and anti-Wick operators are violated. On the other hand, the following result shows that we still may identify the operator classes on the right-hand sides in (4.8) with suitable classes of Wick operators, with Wick symbols bounded by

$$\vartheta_{1,s,r_1,r_2}(z,w) = \begin{cases} e^{\frac{1}{2}|z-w|^2 - r_1|z|^{\frac{1}{s}} + r_2|w|^{\frac{1}{s}}}, & s < \infty, \\ e^{\frac{1}{2}|z-w|^2} \langle z \rangle^{-r_1} \langle w \rangle^{r_2}, & s = \infty, \end{cases} \quad (4.7)$$

and

$$\vartheta_{2,s,r_1,r_2}(z,w) = \begin{cases} e^{\frac{1}{2}|z-w|^2 + r_2|z|^{\frac{1}{s}} - r_1|w|^{\frac{1}{s}}}, & s < \infty, \\ e^{\frac{1}{2}|z-w|^2} \langle z \rangle^{r_2} \langle w \rangle^{-r_1}, & s = \infty. \end{cases} \quad (4.8)$$

Theorem 4.11. *Let $s \in [\frac{1}{2}, \infty]$ and ϑ_{k,s,r_1,r_2} be given by (4.7) and (4.8), $k = 1, 2$. Then the following is true:*

- (1) *if $s < \infty$, then $\mathcal{L}(\mathcal{A}_s(\mathbf{C}^d))$ ($\mathcal{L}(\mathcal{A}'_s(\mathbf{C}^d))$) consists of all $\text{Op}_{\mathfrak{W}}(a)$ such that $a \in \widehat{A}(\mathbf{C}^{2d})$ and for every $r_2 > 0$, there is an $r_1 > 0$ such that*

$$|a(z,w)| \lesssim \vartheta_{1,s,r_1,r_2}(z,w) \quad (|a(z,w)| \lesssim \vartheta_{2,s,r_1,r_2}(z,w)) \quad (4.9)$$

holds;

- (2) *if $s > \frac{1}{2}$, then $\mathcal{L}(\mathcal{A}_{0,s}(\mathbf{C}^d))$ ($\mathcal{L}(\mathcal{A}'_{0,s}(\mathbf{C}^d))$) consists of all $\text{Op}_{\mathfrak{W}}(a)$ such that $a \in \widehat{A}(\mathbf{C}^{2d})$ and for every $r_1 > 0$, there is an $r_2 > 0$ such that (4.9) holds.*

Proof. We only prove (1). The assertion (2) follows by similar arguments and is left for the reader.

Suppose that T is a linear and continuous map from $\mathcal{A}_{b_1}(\mathbf{C}^d)$ to $\mathcal{A}'_{b_1}(\mathbf{C}^d) = A(\mathbf{C}^d)$ with kernel K and Wick symbol a . Then $a, K \in \widehat{\mathcal{A}}'_{b_1}(\mathbf{C}^{2d})$. By Propositions 2.4 and 2.5 it follows that $T \in \mathcal{L}(\mathcal{A}_s(\mathbf{C}^d))$ ($T \in \mathcal{L}(\mathcal{A}'_s(\mathbf{C}^d))$), if and only if for every $r_2 > 0$, there is an $r_1 > 0$ such that

$$|K(z,w)| \lesssim e^{\frac{1}{2}(|z|^2+|w|^2) - r_1|z|^{\frac{1}{s}} + r_2|w|^{\frac{1}{s}}} \quad (|K(z,w)| \lesssim e^{\frac{1}{2}(|z|^2+|w|^2) + r_2|z|^{\frac{1}{s}} - r_1|w|^{\frac{1}{s}}}).$$

The asserted equivalence now follows from the previous equivalence and the fact that $a(z,w) = K(z,w)e^{-(z,w)}$, which implies that

$$|a(z,w)e^{-\frac{1}{2}|z-w|^2}| = |K(z,w)e^{-\frac{1}{2}(|z|^2+|w|^2)}|. \quad \square$$

4.3. Multiplications, differentiations for power series expansions, and compositions of Wick operators. First we combine (3.20) and Proposition 3.7 to deduce ring structures of the spaces in (1.35), (1.36) and in Definition 1.15. The details are left for the reader.

Theorem 4.12. *Let $s_1, s_2 \in \overline{\mathbf{R}}_b$, $s = (s_1, s_2)$ and $W = \mathbf{C}^{d_2} \times \mathbf{C}^{d_1}$. Then the following is true:*

- (1) *if $s_1, s_2 < \frac{1}{2}$, then the ring $\widehat{\mathcal{A}}_0(W)$ under addition and multiplication extends uniquely to topological rings*

$$\widehat{\mathcal{A}}_s(W), \quad \widehat{\mathcal{A}}_s^*(W), \quad \widehat{\mathcal{B}}_s(W), \quad \widehat{\mathcal{B}}_s^*(W), \quad \widehat{\mathcal{C}}_s(W), \quad \text{and} \quad \widehat{\mathcal{C}}_s^*(W), \quad (4.10)$$

under addition and multiplication;

- (2) if $0 < s_1, s_2 \leq \frac{1}{2}$, then the ring $\widehat{\mathcal{A}}_0(W)$ under addition and multiplication extends uniquely to topological rings

$$\widehat{\mathcal{A}}_{0,s}(W), \widehat{\mathcal{A}}_{0,s}^*(W), \widehat{\mathcal{B}}_{0,s}(W), \widehat{\mathcal{B}}_{0,s}^*(W), \widehat{\mathcal{C}}_{0,s}(W) \quad \text{and} \quad \widehat{\mathcal{C}}_{0,s}^*(W), \quad (4.11)$$

under addition and multiplication.

Remark 4.13. In Section 6 we present trace results and results on linear pullbacks for the spaces in (4.10) and (4.11). The ring assertions in Theorem 4.12 also follows by combining such trace and pullback properties.

From now on we let the spaces in (4.10) and in (4.11) be equipped with the ring structure guaranteed by Theorem 4.12.

Next we discuss extensions of the product \diamond on $\widehat{\mathcal{A}}_0(W)$, given in (3.24). The following result is a straight-forward consequence of (3.26) and Proposition 3.8. The details are left for the reader.

Theorem 4.14. *Let $s_1, s_2 \in \overline{\mathbf{R}}_b$, $s = (s_1, s_2)$, $W = \mathbf{C}^{d_2} \times \mathbf{C}^{d_1}$ and let T_\diamond be the map from $\mathcal{A}_0(\mathbf{C}^d) \times \mathcal{A}_0(\mathbf{C}^d)$ to $\mathcal{A}_0(\mathbf{C}^d)$, given by $T_\diamond(K_1, K_2) = K_1 \diamond K_2$, where $K_1 \diamond K_2$ is given by (3.24). If $s_1, s_2 < \frac{1}{2}$, then T_\diamond is uniquely extendable to continuous mappings*

$$\begin{aligned} T_\diamond : \widehat{\mathcal{A}}_s(W) \times \widehat{\mathcal{A}}_s^*(W) &\rightarrow \widehat{\mathcal{A}}_s^*(W) & T_\diamond : \widehat{\mathcal{A}}_s^*(W) \times \widehat{\mathcal{A}}_s(W) &\rightarrow \widehat{\mathcal{A}}_s(W) \\ T_\diamond : \widehat{\mathcal{B}}_s(W) \times \widehat{\mathcal{C}}_s^*(W) &\rightarrow \widehat{\mathcal{C}}_s^*(W) & T_\diamond : \widehat{\mathcal{B}}_s^*(W) \times \widehat{\mathcal{C}}_s^*(W) &\rightarrow \widehat{\mathcal{C}}_s^*(W) \\ T_\diamond : \widehat{\mathcal{C}}_s(W) \times \widehat{\mathcal{B}}_s^*(W) &\rightarrow \widehat{\mathcal{B}}_s^*(W) & \text{and} & T_\diamond : \widehat{\mathcal{C}}_s^*(W) \times \widehat{\mathcal{B}}_s^*(W) &\rightarrow \widehat{\mathcal{B}}_s^*(W). \end{aligned}$$

If instead $0 < s_1, s_2 \leq \frac{1}{2}$, then the same holds true with $\widehat{\mathcal{A}}_{0,s}$, $\widehat{\mathcal{B}}_{0,s}$ and $\widehat{\mathcal{C}}_{0,s}$ in place of $\widehat{\mathcal{A}}_s$, $\widehat{\mathcal{B}}_s$ and $\widehat{\mathcal{C}}_s$, respectively, at each occurrence.

Remark 4.15. Let $d_2 = d$, $d_1 = 0$ and $s \in \overline{\mathbf{R}}_b$ satisfies $s < \frac{1}{2}$. Then it follows from Theorem 4.14 that

$$T_\diamond : \mathcal{A}_s(\mathbf{C}^d) \times \mathcal{A}'_s(\mathbf{C}^d) \rightarrow \mathcal{A}'_s(\mathbf{C}^d) \quad \text{and} \quad T_\diamond : \mathcal{A}'_s(\mathbf{C}^d) \times \mathcal{A}_s(\mathbf{C}^d) \rightarrow \widehat{\mathcal{A}}_s(\mathbf{C}^d) \quad (4.12)$$

are continuous. If instead $0 < s \leq \frac{1}{2}$, then the same holds true with $\mathcal{A}_{0,s}$ in place of \mathcal{A}_s at each occurrence.

Remark 4.16. It is tempting to proclaim that Theorem 4.14 is the dual result to Theorem 4.12, where the former should follow from the latter by using the adjoint relation (3.22), and that the spaces in (1.12) are close to the duals of (1.11) and vice versa. A problem here is that the literature seems not to support completeness and thereby ensure useful duality properties for some of the involved spaces (see e.g. [27, 37, 38]). It therefore seems not so straight-forward to apply duality arguments here.

Next we discuss extensions of the twisted product $\#_{\mathfrak{Y}}$ in (1.61) as a map from $\widehat{\mathcal{A}}_0(\mathbf{C}^{2d}) \times \widehat{\mathcal{A}}_0(\mathbf{C}^{2d})$ to $\widehat{\mathcal{A}}(\mathbf{C}^{2d})$. The following result is a straight-forward consequence of (3.29) and Proposition 3.9. The details are left for the reader.

Theorem 4.17. *Let $s \in \overline{\mathbf{R}}_b$ be such that $s < \frac{1}{2}$, $W = \mathbf{C}^d \times \mathbf{C}^d$, and let $T_\#$ be the map from $\widehat{\mathcal{A}}_0(W) \times \widehat{\mathcal{A}}_0(W)$ to $\widehat{\mathcal{A}}(W)$, given by $T_\#(a_1, a_2) = a_1 \#_{\mathfrak{Y}} a_2$. Then $T_\#$ is uniquely*

extendable to continuous mappings

$$\begin{aligned}
T_{\#} &: \widehat{\mathcal{A}}_s(W) \times \widehat{\mathcal{A}}_s(W) \rightarrow \widehat{\mathcal{A}}_s(W), \\
T_{\#} &: \mathcal{A}'_s(W) \times \mathcal{B}_s(W) \rightarrow \mathcal{A}'_s(W), \quad T_{\#} : \widehat{\mathcal{B}}_s^*(W) \times \widehat{\mathcal{A}}'_s(W) \rightarrow \widehat{\mathcal{A}}'_s(W), \\
T_{\#} &: \widehat{\mathcal{B}}_s(W) \times \widehat{\mathcal{B}}_s(W) \rightarrow \mathcal{B}_s(W), \quad T_{\#} : \widehat{\mathcal{B}}_s^*(W) \times \widehat{\mathcal{B}}_s^*(W) \rightarrow \widehat{\mathcal{B}}_s^*(W), \\
T_{\#} &: \widehat{\mathcal{C}}_s(W) \times \widehat{\mathcal{C}}_s(W) \rightarrow \widehat{\mathcal{C}}_s(W), \quad T_{\#} : \widehat{\mathcal{C}}_s^*(W) \times \widehat{\mathcal{C}}_s^*(W) \rightarrow \widehat{\mathcal{C}}_s^*(W),
\end{aligned} \tag{4.13}$$

If instead $0 < s \leq \frac{1}{2}$, then the same holds true with $\widehat{\mathcal{A}}_{0,s}$, $\widehat{\mathcal{B}}_{0,s}$ and $\widehat{\mathcal{C}}_{0,s}$ in place of $\widehat{\mathcal{A}}_s$, $\widehat{\mathcal{B}}_s$ and $\widehat{\mathcal{C}}_s$, respectively, at each occurrence.

By Theorems 4.9 and 4.17 the twisted anti-Wick product $\#_{\mathfrak{W}}^{\text{aw}}$, defined by

$$\text{Op}_{\mathfrak{W}}^{\text{aw}}(a_1) \circ \text{Op}_{\mathfrak{W}}^{\text{aw}}(a_2) = \text{Op}_{\mathfrak{W}}^{\text{aw}}(a_1 \#_{\mathfrak{W}}^{\text{aw}} a_2), \tag{4.14}$$

makes sense, when a_1 and a_2 belong to the symbol classes in Theorem 4.9. The following result now follows by combining Theorems 4.9 and 4.17. The details are left for the reader.

Theorem 4.18. *Let $s \in \overline{\mathbf{R}}_b$ be such that $s < \frac{1}{2}$, $W = \mathbf{C}^d \times \mathbf{C}^d$, and let $T_{\#}^{\text{aw}}$ be the map from $\widehat{\mathcal{A}}_0(W) \times \widehat{\mathcal{A}}_0(W)$ to $\widehat{\mathcal{A}}(W)$, given by $T_{\#}^{\text{aw}}(a_1, a_2) = a_1 \#_{\mathfrak{W}}^{\text{aw}} a_2$. Then $T_{\#}^{\text{aw}}$ is uniquely extendable to continuous mappings*

$$\begin{aligned}
T_{\#}^{\text{aw}} &: \widehat{\mathcal{A}}_s(W) \times \widehat{\mathcal{A}}_s(W) \rightarrow \widehat{\mathcal{A}}_s(W), \\
T_{\#}^{\text{aw}} &: \widehat{\mathcal{C}}_s(W) \times \widehat{\mathcal{C}}_s(W) \rightarrow \widehat{\mathcal{C}}_s(W), \quad T_{\#}^{\text{aw}} : \widehat{\mathcal{C}}_s^*(W) \times \widehat{\mathcal{C}}_s^*(W) \rightarrow \widehat{\mathcal{C}}_s^*(W),
\end{aligned} \tag{4.15}$$

If instead $0 < s \leq \frac{1}{2}$, then the same holds true with $\widehat{\mathcal{A}}_{0,s}$ and $\widehat{\mathcal{C}}_{0,s}$ in place of $\widehat{\mathcal{A}}_s$ and $\widehat{\mathcal{C}}_s$, respectively, at each occurrence.

4.4. Some further relationships between operator kernels, Wick and anti-Wick symbols. Next we show some transition properties between Wick, anti-Wick and kernel operators with symbols and kernels belonging to refined classes of $\widehat{\mathcal{A}}_{0,1/2}(\mathbf{C}^{2d})$ and $\widehat{\mathcal{A}}'_{0,1/2}(\mathbf{C}^{2d})$. For any $r > 0$, let $\widehat{\mathcal{A}}_{[r]}^2(\mathbf{C}^{2d})$ be the Banach space which consists of all $a \in \widehat{\mathcal{A}}(\mathbf{C}^{2d})$ such that

$$\|a\|_{\widehat{\mathcal{A}}_{[r]}^2} \equiv \left(\iint_{\mathbf{C}^{2d}} |a(z, w)|^2 e^{-r(|z|^2 + |w|^2)} d\lambda(z) d\lambda(w) \right)^{\frac{1}{2}}$$

is finite. We observe that $\widehat{\mathcal{A}}_{0,1/2}(\mathbf{C}^{2d})$ ($\widehat{\mathcal{A}}'_{0,1/2}(\mathbf{C}^{2d})$) is the projective (inductive) limit of $\widehat{\mathcal{A}}_{[r]}^2(\mathbf{C}^{2d})$ with respect to $r > 0$. First we have the following two propositions.

Proposition 4.19. *Let $r_1 \in (0, 1)$ and $r_2 > r_1/(1 - r_1)$. Then the following is true:*

- (1) *if $a \in \widehat{\mathcal{A}}_{[r_1]}^2(\mathbf{C}^{2d})$, then there is a unique $a^{\text{aw}} \in \widehat{\mathcal{A}}_{[r_2]}^2(\mathbf{C}^{2d})$ such that (1.58) holds. Furthermore,*

$$\|a^{\text{aw}}\|_{\widehat{\mathcal{A}}_{[r_2]}^2} \lesssim \|a\|_{\widehat{\mathcal{A}}_{[r_1]}^2}; \tag{4.16}$$

- (2) *if $a^{\text{aw}} \in \widehat{\mathcal{A}}_{[r_1]}^2(\mathbf{C}^{2d})$, then there is a unique $a \in \widehat{\mathcal{A}}_{[r_2]}^2(\mathbf{C}^{2d})$ such that (1.58) holds. Furthermore,*

$$\|a\|_{\widehat{\mathcal{A}}_{[r_2]}^2} \lesssim \|a^{\text{aw}}\|_{\widehat{\mathcal{A}}_{[r_1]}^2}. \tag{4.17}$$

Proposition 4.20. *Let $r_1 > 0$ and $r_2 > r_1 + 1$. Then the following is true:*

- (1) if $a \in \widehat{\mathcal{A}}_{[r_1]}^2(\mathbf{C}^{2d})$, then there is a unique $K \in \widehat{\mathcal{A}}_{[r_2]}^2(\mathbf{C}^{2d})$ such that (3.6)' holds. Furthermore,

$$\|K\|_{\widehat{\mathcal{A}}_{[r_2]}^2} \lesssim \|a\|_{\widehat{\mathcal{A}}_{[r_1]}^2}; \quad (4.18)$$

- (2) if $K \in \widehat{\mathcal{A}}_{[r_2]}^2(\mathbf{C}^{2d})$, then there is a unique $a \in \widehat{\mathcal{A}}_{[r_1]}^2(\mathbf{C}^{2d})$ such that (3.6)' holds. Furthermore,

$$\|K\|_{\widehat{\mathcal{A}}_{[r_2]}^2} \lesssim \|a\|_{\widehat{\mathcal{A}}_{[r_1]}^2}. \quad (4.19)$$

For the proofs we need the following continuity result for the operators $\mathcal{T}_{0,t}$ and $\mathcal{T}_{0,t}^*$. Here let

$$\ell_r^p(\mathbf{N}^d) = \ell_{[\vartheta_r]}^p(\mathbf{N}^d), \quad \vartheta_r(\alpha) = r^{-\frac{1}{2}|\alpha|}, \quad r > 0, \quad \alpha \in \mathbf{N}^d$$

(cf. Definition 1.3).

Lemma 4.21. *Let $p \in [1, \infty]$, $r_1 > 0$, $t \in \mathbf{C}$ be such that $|t| \leq 1$, and let $\mathcal{T}_{0,t}$ and $\mathcal{T}_{0,t}^*$ be given by (3.1) and (3.2), $j = 1, 2$. Then the following is true:*

- (1) if $r_2 \geq 1 + r_1$ with strict inequality when $p < \infty$, then $\mathcal{T}_{0,t}$ restricts to a continuous map from $\ell_{r_1}^p(\mathbf{N}^{2d})$ to $\ell_{r_2}^p(\mathbf{N}^{2d})$;
- (2) if in addition $r_1 < 1$ and $r_2 \geq r_1/(1 - r_1)$ with strict inequality when $p > 1$, then $\mathcal{T}_{0,t}^*$ restricts to a continuous map from $\ell_{r_1}^p(\mathbf{N}^{2d})$ to $\ell_{r_2}^p(\mathbf{N}^{2d})$.

For the proof we observe that

$$|b(\alpha, \beta)| \leq r^{\frac{1}{2}|\alpha+\beta|} \|b\|_{\ell_r^p}, \quad b \in \ell_r^p(\mathbf{N}^{2d}). \quad (4.20)$$

Proof. By duality it suffices to prove (1). We only prove the result for $p < \infty$. The other cases follow by similar arguments and are left for the reader.

Let $r = r_1$ and $b \in \ell_r^p(\mathbf{N}^{2d})$. Then it follows from (4.20), Cauchy-Schwartz inequality and straight-forward computations that

$$\begin{aligned} \|\mathcal{T}_{0,t} b\|_{\ell_{r_2}^p}^p &\leq \sum_{\alpha, \beta} \left(\sum_{\gamma \leq \alpha, \beta} \binom{\alpha}{\gamma}^{\frac{1}{2}} \binom{\beta}{\gamma}^{\frac{1}{2}} |b(\alpha - \gamma, \beta - \gamma)| \right)^p r_2^{-\frac{p}{2}|\alpha+\beta|} \\ &\leq \|b\|_{\ell_r^p}^p \sum_{\alpha, \beta} \left(\sum_{\gamma \leq \alpha, \beta} \binom{\alpha}{\gamma}^{\frac{1}{2}} \binom{\beta}{\gamma}^{\frac{1}{2}} r^{-|\gamma|} \right)^p r^{\frac{p}{2}|\alpha+\beta|} r_2^{-\frac{p}{2}|\alpha+\beta|} \\ &\leq \|b\|_{\ell_r^p}^p \sum_{\alpha, \beta} \left(\sum_{\gamma \leq \alpha} \binom{\alpha}{\gamma} r^{-|\gamma|} \right)^{\frac{p}{2}} \left(\sum_{\gamma \leq \beta} \binom{\beta}{\gamma} r^{-|\gamma|} \right)^{\frac{p}{2}} \left(\frac{r}{r_2} \right)^{\frac{p}{2}|\alpha+\beta|} \\ &= \|b\|_{\ell_r^p}^p \sum_{\alpha, \beta} \left(\frac{1+r}{r_2} \right)^{\frac{p}{2}|\alpha+\beta|}. \end{aligned}$$

Since $1 + r < r_2$, it follows that the last series converges, which gives the asserted continuity. \square

Proof of Propositions 4.19 and 4.20. If $r > 0$ and $a(z, w) \in \widehat{A}(\mathbf{C}^{2d})$ has the expansion

$$a(z, w) = \sum c(a; \alpha, \beta) e_\alpha(z) e_\beta(\bar{w}), \quad z, w \in \mathbf{C}^d, \quad j = 1, 2, \quad (4.21)$$

then [35, Corollary 3.8] shows that

$$\|a\|_{\widehat{\mathcal{A}}_{[r]}^2} = \sum_{\alpha, \beta \in \mathbf{N}^d} |c(a; \alpha, \beta)|^2 r^{-|\alpha|}.$$

The result now follows from the latter relation, Lemma 4.21, the fact that $\ell_0(\mathbf{N}^{2d})$ is dense in $\ell_r^2(\mathbf{N}^{2d})$, leading to that $\widehat{\mathcal{A}}_0(\mathbf{C}^{2d})$ is dense in $\widehat{\mathcal{A}}_{[r]}^2(\mathbf{C}^{2d})$, and that (1.58) and (3.6)' hold true when $a, a^{\text{aw}} \in \widehat{\mathcal{A}}_0(\mathbf{C}^{2d})$. \square

In [32, Section 3] some results concerning the Wick symbols of anti-Wick operators are presented. For example we have the following result. We omit the proof since the result is a special case of Theorems 3.1, 3.3 and Theorem 3.7 in [32]. Here $\mathcal{P}_E(\mathbf{C}^d)$ is the set of all $\omega \in L_{loc}^\infty(\mathbf{C}^d; \mathbf{R}_+)$ such that

$$\omega(z+w) \lesssim \omega(z)v(w),$$

for some $v \in L_{loc}^\infty(\mathbf{C}^d; \mathbf{R}_+)$.

Theorem 4.22. *Let $s \geq \frac{1}{2}$ ($s > \frac{1}{2}$), $a, a^{\text{aw}} \in \widehat{A}(\mathbf{C}^{2d})$ be such that (1.59)' holds, and let $\omega \in \mathcal{P}_E(\mathbf{C}^d)$. Then the following is true:*

- (1) if $|a(w, w)| \lesssim e^{-r_0|w|^{\frac{1}{s}}}$ holds for some $r_0 > 0$ (for every $r_0 > 0$), then

$$|a^{\text{aw}}(z, w)| \lesssim e^{\frac{1}{4}|z-w|^2 - r|z+w|^{\frac{1}{s}}} \quad (4.22)$$

for some $r > 0$ (for every $r > 0$);

- (2) if $|a(w, w)| \lesssim e^{r_0|w|^{\frac{1}{s}}}$ holds for every $r_0 > 0$ (for some $r_0 > 0$), then

$$|a^{\text{aw}}(z, w)| \lesssim e^{\frac{1}{4}|z-w|^2 + r|z+w|^{\frac{1}{s}}} \quad (4.23)$$

for every $r > 0$ (for some $r > 0$);

- (3) if $|a(w, w)| \lesssim e^{r|w|^2}$ for some $r < 1$, then

$$|a^{\text{aw}}(z, w)| \lesssim e^{r_0|z+w|^2 - \text{Re}(z, w)}, \quad r_0 = 4^{-1}(1-r)^{-1}; \quad (4.24)$$

- (4) if $|a(w, w)| \lesssim \omega(2w)$, then

$$|a^{\text{aw}}(z, w)| \lesssim e^{\frac{1}{4}|z-w|^2} \omega(z+w), \quad z, w \in \mathbf{C}^d. \quad (4.25)$$

By using (1.60) instead of (1.59), we get the following result in the other direction compared to previous result. The details are left for the reader.

Theorem 4.23. *Let $s \geq \frac{1}{2}$ ($s > \frac{1}{2}$), $a, a^{\text{aw}} \in \widehat{A}(\mathbf{C}^{2d})$ be such that (1.59)' holds, and let $\omega \in \mathcal{P}_E(\mathbf{C}^d)$. Then the following is true:*

- (1) if $|a^{\text{aw}}(w, -w)| \lesssim e^{-r_0|w|^{\frac{1}{s}}}$ holds for some $r_0 > 0$ (for every $r_0 > 0$), then

$$|a(z, w)| \lesssim e^{\frac{1}{4}|z+w|^2 - r|z-w|^{\frac{1}{s}}} \quad (4.26)$$

for some $r > 0$ (for every $r > 0$);

- (2) if $|a^{\text{aw}}(w, -w)| \lesssim e^{r_0|w|^{\frac{1}{s}}}$ holds for every $r_0 > 0$ (for some $r_0 > 0$), then

$$|a(z, w)| \lesssim e^{\frac{1}{4}|z+w|^2 + r|z-w|^{\frac{1}{s}}} \quad (4.27)$$

for every $r > 0$ (for some $r > 0$);

- (3) if $|a^{\text{aw}}(w, -w)| \lesssim e^{r|w|^2}$ for some $r < 1$, then

$$|a(z, w)| \lesssim e^{r_0|z-w|^2 + \text{Re}(z, w)}, \quad r_0 = 4^{-1}(1-r)^{-1}; \quad (4.28)$$

- (4) if $|a^{\text{aw}}(w, -w)| \lesssim \omega(2w)$, then

$$|a(z, w)| \lesssim e^{\frac{1}{4}|z+w|^2} \omega(z-w), \quad z, w \in \mathbf{C}^d. \quad (4.29)$$

Remark 4.24. The relationships between the Wick and anti-Wick symbols in Theorems 4.22 and 4.23 are similar, and one might believe that transitions of continuity properties for Wick operators to anti-Wick operators are similar as for transitions in reversed directions. Here we notice that this is not the case.

For example, suppose that a^{aw} satisfies (4.23) for every $r > 0$, $0 < t_0 < \frac{3}{4}$ and that $F \in A(\mathbf{C}^d)$ satisfies

$$|F(z)| \lesssim e^{t_0|z|^2+r|z|^{\frac{1}{s}}} \quad (4.30)$$

for every $r > 0$. (The case when a^{aw} satisfies conditions of the form (4.22) is treated in similar ways and leads to even stronger continuity properties.) If

$$t_1 = \frac{1 - t_0}{3 - 4t_0}$$

(which is bounded from below by t_0), then the size of the integrand in

$$|(\text{Op}_{\mathfrak{W}}(a^{\text{aw}})F)(z)| \leq \pi^{-d} \int_{\mathbf{C}^d} |a^{\text{aw}}(z, w)F(w)| e^{\text{Re}(z, w) - |w|^2} d\lambda(w)$$

can be estimated as

$$\begin{aligned} |a^{\text{aw}}(z, w)F(w)| e^{\text{Re}(z, w) - |w|^2} &\lesssim e^{\frac{1}{4}|z-w|^2+r|z+w|^{\frac{1}{s}}} e^{\text{Re}(z, w) - |w|^2} e^{t_0|w|^2+r|w|^{\frac{1}{s}}} \\ &= e^{t_1|z|^2 - \frac{3-4t_0}{4}|w - \frac{z}{3-4t_0}|^2 + r(|z+w|^{\frac{1}{s}} + |w|^{\frac{1}{s}})} \\ &\lesssim e^{t_1|z|^2 + cr|z|^{\frac{1}{s}}} e^{-\frac{3-4t_0}{4}|w - \frac{z}{3-4t_0}|^2 + cr|w - \frac{z}{3-4t_0}|^{\frac{1}{s}}}, \end{aligned} \quad (4.31)$$

for some $c \geq 1$ which is independent of $z, w \in \mathbf{C}^d$ and $r > 0$. By integrating the last estimate with respect to w , it follows that (4.30) holds true for every $r > 0$ with $\text{Op}_{\mathfrak{W}}(a^{\text{aw}})F$ and t_1 in place of F and t_0 .

By choosing $t_0 = \frac{1}{2}$, then $t_1 = \frac{1}{2}$, and we have proved that $\text{Op}_{\mathfrak{W}}(a^{\text{aw}})$ is continuous on $\mathcal{A}'_s(\mathbf{C}^d)$ for such t_0 .

As side remark we observe that stronger continuity properties for anti-Wick operators with symbols satisfying estimates of the form $a(w, w)e^{\pm r|w|^{\frac{1}{s}}}$ are obtained with more direct computational methods, without rewriting $\text{Op}_{\mathfrak{W}}^{\text{aw}}(a)$ as a Wick operator. (See Proposition 3.6 to Corollary 3.10 in [32].)

The corresponding estimates (4.26) and (4.27) when passing from Wick to anti-Wick operators lead to strongly different conclusions. In fact, let $F \in A(\mathbf{C}^d)$. For the integrand in

$$|(\text{Op}_{\mathfrak{W}}^{\text{aw}}(a)F)(z)| \leq \pi^{-d} \int_{\mathbf{C}^d} |a(w, w)F(w)| e^{\text{Re}(z, w) - |w|^2} d\lambda(w)$$

a similar type of estimate gives

$$\begin{aligned} |a(w, w)F(w)| e^{\text{Re}(z, w) - |w|^2} &\lesssim e^{\frac{1}{4}|w+w|^2 \pm r|w-w|^{\frac{1}{s}}} e^{\text{Re}(z, w) - |w|^2} |F(w)| \\ &= e^{\text{Re}(z, w)} |F(w)|. \end{aligned} \quad (4.32)$$

Since F should be entire, the right-hand side is integrable with respect to w , only when F is identically zero. Consequently, (4.26) and (4.27) might be useful only when F is the trivial zero function.

In this context we observe that continuity properties for $\text{Op}_{\mathfrak{W}}(a^{\text{aw}})$ are reachable with more direct computational methods, without rewriting it as an anti-Wick operator. (See [31, 32].)

5. APPLICATIONS TO LINEAR OPERATORS ON FUNCTIONS AND DISTRIBUTIONS
DEFINED ON \mathbf{R}^d

In this section we use the results in previous sections to extend the definition of Toeplitz operators. In the end we find that if $s \in \overline{\mathbf{R}_b}$ is suitable, then any linear and continuous operators on $\mathcal{A}_s(\mathbf{C}^d)$, $\mathcal{A}_{0,s}(\mathbf{C}^d)$ and their duals, can be expressed as Toeplitz operators.

If $a \in \widehat{\mathcal{A}}_0(\mathbf{C}^{2d})$, then $\text{Tp}_{\mathfrak{W}}(a)$ is the linear and continuous operator on $\mathcal{S}_{1/2}(\mathbf{R}^d)$, given by (1.63)". For such operators we have the following extensions.

Theorem 5.1. *Let $s_1, s_2 \in \overline{\mathbf{R}_b}$ be such that $s_1 < \frac{1}{2}$ and $0 < s_2 \leq \frac{1}{2}$. Then the following is true:*

- (1) *the map $(a, f) \mapsto \text{Tp}_{\mathfrak{W}}(a)f$ from $\widehat{\mathcal{A}}_0(\mathbf{C}^{2d}) \times \mathcal{H}_0(\mathbf{R}^d)$ to $\mathcal{H}'_0(\mathbf{R}^d)$ is uniquely extendable to a continuous map $\widehat{\mathcal{C}}_{s_1}(\mathbf{C}^{2d}) \times \mathcal{H}_{s_1}(\mathbf{R}^d)$ to $\mathcal{H}_{s_1}(\mathbf{R}^d)$, and from $\widehat{\mathcal{C}}_{s_1}^*(\mathbf{C}^{2d}) \times \mathcal{H}'_{s_1}(\mathbf{R}^d)$ to $\mathcal{H}'_{s_1}(\mathbf{R}^d)$;*
- (2) *the map $(a, f) \mapsto \text{Tp}_{\mathfrak{W}}(a)f$ from $\widehat{\mathcal{A}}_0(\mathbf{C}^{2d}) \times \mathcal{H}_0(\mathbf{R}^d)$ to $\mathcal{H}'_0(\mathbf{R}^d)$ is uniquely extendable to a continuous map $\widehat{\mathcal{C}}_{0,s_2}(\mathbf{C}^{2d}) \times \mathcal{H}_{0,s_2}(\mathbf{R}^d)$ to $\mathcal{H}_{0,s_2}(\mathbf{R}^d)$, and from $\widehat{\mathcal{C}}_{0,s_2}^*(\mathbf{C}^{2d}) \times \mathcal{H}'_{0,s_2}(\mathbf{R}^d)$ to $\mathcal{H}'_{0,s_2}(\mathbf{R}^d)$.*

Proof. The asserted continuity extensions follow from (1.63)', (1.64), Theorem 4.8 and the facts that the Bargmann transform is homeomorphic from the spaces in (1.27) to the spaces in (1.28). We need to prove the uniqueness.

By playing with r_1 and r_2 in Definition 1.5, it follows that $\ell_0(\Lambda)$ is dense in the spaces in (1.11), which implies that $\widehat{\mathcal{A}}_0(W)$ is dense in the spaces in (1.38). The uniqueness in Theorem 5.1 now follows by combining these density properties with the fact that $\mathcal{H}_0(\mathbf{R}^d)$ is dense in the spaces in (1.27). \square

Theorem 5.2. *Let $s_1, s_2 \in \overline{\mathbf{R}_b}$ be such that $s_1 < \frac{1}{2}$ and $0 < s_2 \leq \frac{1}{2}$. If $a_1, a_2 \in \widehat{\mathcal{C}}_{s_1}(\mathbf{C}^{2d})$ and $\text{Tp}_{\mathfrak{W}}(a_1)f = \text{Tp}_{\mathfrak{W}}(a_2)f$ for every $f \in \mathcal{H}_0(\mathbf{R}^d)$, then $a_1 = a_2$. The same holds true for $\widehat{\mathcal{C}}_{s_1}^*$, $\widehat{\mathcal{C}}_{0,s_2}$ and $\widehat{\mathcal{C}}_{0,s_2}^*$ in place of $\widehat{\mathcal{C}}_{s_1}$.*

Proof. Suppose that $a_1, a_2 \in \widehat{\mathcal{C}}_{s_1}(\mathbf{C}^{2d})$ satisfy $\text{Tp}_{\mathfrak{W}}(a_1)f = \text{Tp}_{\mathfrak{W}}(a_2)f$ for every $f \in \mathcal{H}_0(\mathbf{R}^d)$. Then Theorem 4.3 gives

$$\begin{aligned} \text{Tp}_{\mathfrak{W}}(a_1) = \text{Tp}_{\mathfrak{W}}(a_2) &\Leftrightarrow \text{Op}_{\mathfrak{W}}^{\text{aw}}(a_1) = \text{Op}_{\mathfrak{W}}^{\text{aw}}(a_2) \Leftrightarrow \text{Op}_{\mathfrak{W}}(\mathcal{T}_1^* a_1) = \text{Op}_{\mathfrak{W}}(\mathcal{T}_1^* a_2) \\ &\Leftrightarrow \mathcal{T}_1^* a_1 = \mathcal{T}_1^* a_2 \Leftrightarrow a_1 = a_2. \end{aligned} \quad \square$$

6. LINEAR PULLBACKS AND TRACE MAPPINGS FOR SPACES OF POWER SERIES
EXPANSIONS

In this section we show that for $0 \leq s < \frac{1}{2}$ (for $0 < s \leq \frac{1}{2}$), then linear pullbacks and trace mappings are continuous mappings on the spaces in (4.10) (in (4.11)).

For $B_j \in \mathbf{M}(d_j, \mathbf{C})$, $j = 1, 2$, we consider linear pullbacks of the form

$$K(z_2, z_1) \mapsto K(z_2, B_1 z_1) \tag{6.1}$$

$$K(z_2, z_1) \mapsto K(B_2 z_2, z_1) \tag{6.2}$$

and

$$K(z_2, z_1) \mapsto K(B_2 z_2, B_1 z_1). \tag{6.3}$$

Here $\mathbf{M}(d, \mathbf{C})$ is the set of all $d \times d$ matrices with entries in \mathbf{C} . For such pullbacks we have the following.

Theorem 6.1. *Let $s_1, s_2 \in \overline{\mathbf{R}_{b,\infty}}$, $s = (s_2, s_1)$, $W = \mathbf{C}^{d_2} \times \mathbf{C}^{d_1}$ and $B_j \in \mathbf{M}(d_j, \mathbf{C})$. Then the following is true:*

- (1) *the map (6.3) on $\hat{A}(W)$ extends uniquely to a continuous map on $\hat{\mathcal{A}}'_0(W)$;*
- (2) *if in addition $s_1 < \frac{1}{2}$ ($0 < s_1 \leq \frac{1}{2}$), then (6.1) on $\hat{\mathcal{A}}'_0(W)$ restricts to continuous mappings on the spaces in (4.10) (in (4.11));*
- (3) *if in addition $s_2 < \frac{1}{2}$ ($0 < s_2 \leq \frac{1}{2}$), then (6.2) on $\hat{\mathcal{A}}'_0(W)$ restricts to continuous mappings on the spaces in (4.10) (in (4.11));*
- (4) *if in addition $s_1, s_2 < \frac{1}{2}$ ($0 < s_1, s_2 \leq \frac{1}{2}$), then (6.3) on $\hat{\mathcal{A}}'_0(W)$ restricts to continuous mappings on the spaces in (4.10) (in (4.11)).*

Proof. We only prove (1) and (3). The assertion (2) follows from similar arguments and then (4) follows by combining (2) and (3). The details are left for the reader.

Let $B = B_2$, $d = d_2$ and $z = z_2 \in \mathbf{C}^d$. For any integer $N \geq 0$ we let

$$\Omega_N = \Omega_{N,d} = \{ \alpha \in \mathbf{N}^d; |\alpha| = N \}.$$

Then

$$e_\alpha(Bz) = \sum_{\beta \in \Omega_N} C_B(\beta, \alpha) e_\beta(z), \quad \alpha \in \Omega_N, \quad z \in \mathbf{C}^d, \quad (6.4)$$

for some constants $C_B(\beta, \alpha)$ which are uniquely defined and only depend on α , β and B . If $K \in \hat{A}(W)$ is given by (2.5), then

$$K_B(z_2, z_1) \equiv K(Bz_2, z_1) = \sum_{\alpha_j \in \mathbf{N}^{d_j}} c(K_B; \alpha_2, \alpha_1) e_{\alpha_2}(z_2) e_{\alpha_1}(\bar{z}_1), \quad (6.5)$$

where

$$c(K_B; \alpha_2, \alpha_1) = \sum_{\beta \in \Omega_{|\alpha_2|}} C_B(\alpha_2, \beta) c(K; \beta, \alpha_1). \quad (6.6)$$

Hence, if, more generally $K \in \hat{\mathcal{A}}'_0(W)$, then the only possibility to define $K(B_2 z_2, z_1)$ is given by (6.5) and (6.6). Since Ω_N is a finite set for every N , it follows that (6.6) is well-defined and that the map

$$\{c(K; \alpha_2, \alpha_1)\}_{(\alpha_2, \alpha_1) \in \mathbf{N}^{d_2} \times \mathbf{N}^{d_1}} \mapsto \{c(K_B; \alpha_2, \alpha_1)\}_{(\alpha_2, \alpha_1) \in \mathbf{N}^{d_2} \times \mathbf{N}^{d_1}}$$

is continuous

$$\begin{aligned} \ell_{\mathcal{A}_0}(\mathbf{N}^{d_2} \times \mathbf{N}^{d_1}), \quad \ell_{\mathcal{A}'_0}(\mathbf{N}^{d_2} \times \mathbf{N}^{d_1}), \quad \ell_{\mathcal{B}(0,0)}(\mathbf{N}^{d_2} \times \mathbf{N}^{d_1}), \\ \ell_{\mathcal{B}^*(0,0)}(\mathbf{N}^{d_2} \times \mathbf{N}^{d_1}), \quad \ell_{\mathcal{C}(0,0)}(\mathbf{N}^{d_2} \times \mathbf{N}^{d_1}) \quad \text{and on} \quad \ell_{\mathcal{C}^*(0,0)}(\mathbf{N}^{d_2} \times \mathbf{N}^{d_1}). \end{aligned}$$

Hence the map (6.3) on $\hat{A}(W)$ is uniquely extendable to a continuous map on $\hat{\mathcal{A}}'_0(W)$, which in turn restricts to continuous mappings on the spaces in (4.10) when $s_1 = s_2 = 0$. This gives (1) and (3) for $s_1 = s_2 = 0$.

We only prove (3) in the case when $s_1, s_2 > 0$. The case when $s_1 = s_2 = 0$ is already treated and the case when one of s_1 and s_2 is zero follows by similar arguments and is left for the reader.

Let $\{b(j, k)\}_{k=1}^d$ be the entries in the row j of B and let

$$\binom{N}{\alpha} = \frac{N!}{\alpha_1! \cdots \alpha_d!}, \quad \alpha = (\alpha_1, \dots, \alpha_d) \in \Omega_N$$

Then

$$\begin{aligned}
e_\alpha(Bz) &= \frac{1}{\alpha!^{\frac{1}{2}}} \prod_{j=1}^d (b(j, 1)z_1 + \cdots + b(j, d))^\alpha \\
&= \frac{1}{\alpha!^{\frac{1}{2}}} \prod_{j=1}^d \left(\sum_{\gamma_j \in \Omega_{\alpha_j}} \binom{\alpha_j}{\gamma_j} \prod_{m=1}^d (b(j, m)^{\gamma_{j,m}} z_m^{\gamma_{j,m}}) \right) \\
&= \frac{1}{\alpha!^{\frac{1}{2}}} \sum_{\gamma_1 \in \Omega_{\alpha_1}} \cdots \sum_{\gamma_d \in \Omega_{\alpha_d}} \left(\prod_{j=1}^d \binom{\alpha_j}{\gamma_j} \prod_{m=1}^d (b(j, m)^{\gamma_{j,m}} z_m^{\gamma_{j,m}}) \right)
\end{aligned}$$

Here $\gamma_j = (\gamma_{j,1}, \dots, \gamma_{j,d})$ in the expressions above. If we let

$$c(\gamma) = \prod_{j,m=1}^d b(j, m)^{\gamma_{j,m}}, \quad (6.7)$$

and using (1.25), then it follows by straight-forward computations that

$$e_\alpha(Bz) = \sum_{\gamma_1 \in \Omega_{\alpha_1}} \cdots \sum_{\gamma_d \in \Omega_{\alpha_d}} \left(\binom{\beta_1}{\delta_1} \cdots \binom{\beta_d}{\delta_d} \right)^{\frac{1}{2}} \left(\frac{\alpha!}{\beta!} \right)^{\frac{1}{2}} c(\gamma) e_\beta(z), \quad (6.8)$$

where

$$\delta_j = (\gamma_{1,j}, \dots, \gamma_{d,j}) \quad \text{and} \quad \beta_j = \gamma_{1,j} + \cdots + \gamma_{d,j}. \quad (6.9)$$

This in turn implies that

$$K_B(z, z_1) = \sum \left(\binom{\beta_1}{\delta_1} \cdots \binom{\beta_d}{\delta_d} \right)^{\frac{1}{2}} \left(\frac{\alpha!}{\beta!} \right)^{\frac{1}{2}} c(K; \alpha, \varrho) c(\gamma) e_\beta(z) e_\varrho(\bar{z}_1),$$

where the sum is taken over all

$$\alpha \in \mathbf{N}^d, \quad \varrho \in \mathbf{N}^{d_1} \quad \text{and} \quad \gamma_j \in \Omega_{\alpha_j}, \quad j = 1, \dots, d. \quad (6.10)$$

Hence

$$c(K_B; \beta, \varrho) = \sum \left(\binom{\beta_1}{\delta_1} \cdots \binom{\beta_d}{\delta_d} \right)^{\frac{1}{2}} \left(\frac{\alpha!}{\beta!} \right)^{\frac{1}{2}} c(K; \alpha, \varrho) c(\gamma), \quad (6.11)$$

where the sum is taken over all α and γ_j in (6.10) such that (6.9) holds. Here we observe that

$$|\alpha| = |\beta| = \sum_{j,k=1}^d \gamma_{j,k}. \quad (6.12)$$

This implies that

$$\binom{\beta_1}{\delta_1} \cdots \binom{\beta_d}{\delta_d} + \frac{\alpha!}{\beta!} \leq C^{|\alpha|} \quad \text{and} \quad |c(\gamma)| \leq M^{|\alpha|}, \quad M = \max_{1 \leq j, m \leq d} |b(j, m)|,$$

for some constant $C \geq 1$. Here the last estimate follows from (6.7). Since it is clear that the number of elements in the sum in (6.11) is bounded by

$$(1 + \alpha_1)^d \cdots (1 + \alpha_d)^d \lesssim C^{|\alpha|},$$

for some constant $C \geq 1$, it follows by combining these estimates and (6.12) that

$$\max_{|\alpha|=N} |c(K_B; \alpha, \varrho)| \leq C^N \max_{|\alpha|=N} |c(K; \alpha, \varrho)|$$

for some C which is independent of N and ϱ .

If $\vartheta_{r,(s_2,s_1)}$ and $\omega_{s_2,s_1;r_2,r_1}$ are the same as in Definitions 1.4 and 1.5, it follows by the assumptions on s_2 that

$$\begin{aligned} \sup_{\alpha \in \mathbf{N}^d} |c(K_B; \alpha, \varrho) \vartheta_{r,(s_2,s_1)}(\alpha, \varrho)| &\leq C \sup_{\alpha \in \mathbf{N}^d} |c(K; \alpha, \varrho) \vartheta_{c_1 r,(s_2,s_1)}(\alpha, \varrho)|, \\ \sup_{\alpha \in \mathbf{N}^d} |c(K_B; \alpha, \varrho) / \vartheta_{r,(s_2,s_1)}(\alpha, \varrho)| &\leq C \sup_{\alpha \in \mathbf{N}^d} |c(K; \alpha, \varrho) / \vartheta_{c_2 r,(s_2,s_1)}(\alpha, \varrho)|, \\ \sup_{\alpha \in \mathbf{N}^d} |c(K_B; \alpha, \varrho) \omega_{r_2,r_1;s_2,s_1}(\alpha, \varrho)| &\leq C \sup_{\alpha \in \mathbf{N}^d} |c(K; \alpha, \varrho) \omega_{c_1 r_2,r_1;s_2,s_1}(\alpha, \varrho)|, \end{aligned}$$

and

$$\sup_{\alpha \in \mathbf{N}^d} |c(K_B; \alpha, \varrho) / \omega_{r_2,r_1;s_2,s_1}(\alpha, \varrho)| \leq C \sup_{\alpha \in \mathbf{N}^d} |c(K; \alpha, \varrho) / \omega_{c_2 r_2,r_1;s_2,s_1}(\alpha, \varrho)|,$$

for some constants $c_1, c_2, C > 0$ which only depend on s_2 and d . The asserted continuity properties in (3) now follows from these estimates. \square

Next we consider generalized and twisted forms of trace mappings on the the spaces in (4.10) and (4.11), given in the following definition.

Definition 6.2. Let $d, d_j, n, n_j \in \mathbf{N}$, $\{e_{j,d}\}_{j=1}^d$ be the canonical basis on \mathbf{C}^d and let S_d be the permutation group on $\{1, \dots, d\}$.

(1) For any $\tau \in S_{d+n}$, $\iota_{\tau,d}$ is the linear map from \mathbf{C}^d to \mathbf{C}^{d+n} such that

$$\iota_{\tau,d} e_{j,d} = e_{\tau(j),d+n}, \quad j = 1, \dots, d.$$

(2) The *twisted trace map* $\text{Tr}_{\tau,d}$ with respect to $\tau \in S_{d+n}$, d and n is the continuous (pullback) map from $\mathcal{A}'_0(\mathbf{C}^{d+n})$ to $\mathcal{A}'_0(\mathbf{C}^d)$ given by

$$(\text{Tr}_{\tau,d} F)(z) \equiv F(\iota_{\tau,d}(z)), \quad F \in \mathcal{A}'_0(\mathbf{C}^{d+n}), \quad z \in \mathbf{C}^d.$$

(3) The *twisted trace map* $\text{Tr}_{\tau,d}$ with respect to $\tau = (\tau_2, \tau_1)$ and $d = (d_2, d_1)$, $\tau_j \in S_{d_j+n_j}$ and d_j , $j = 1, 2$, is the continuous (pullback) map from $\hat{\mathcal{A}}'_0(\mathbf{C}^{d_2+n_2} \times \mathbf{C}^{d_1+n_1})$ to $\hat{\mathcal{A}}'_0(\mathbf{C}^{d_2} \times \mathbf{C}^{d_1})$ given by

$$(\text{Tr}_{\tau,d} K)(z_2, z_1) \equiv K(\iota_{\tau_2,d_2}(z_2), \iota_{\tau_1,d_1}(z_1)) \quad K \in \hat{\mathcal{A}}'_0(\mathbf{C}^{d_2+n_2} \times \mathbf{C}^{d_1+n_1}), \quad z_j \in \mathbf{C}^{d_j}.$$

If $\tau \in S_{d+n}$ and $\iota_{\tau,d}^*$ is the map from \mathbf{N}^d to \mathbf{N}^{d+n} , given by

$$\iota_{\tau,d}^*(\alpha) = (\beta_1, \dots, \beta_{d+n}), \quad \text{where} \quad \beta_j = \begin{cases} \alpha_{\tau^{-1}(j)}, & 1 \leq \tau^{-1}(j) \leq d, \\ 0, & d < \tau^{-1}(j) \leq d+n, \end{cases} \quad (6.13)$$

then it follows that

$$(\text{Tr}_{\tau,d} F)(z) = \sum_{\alpha \in \mathbf{N}^d} c(F; \iota_{\tau,d}^*(\alpha)) e_\alpha(z), \quad z \in \mathbf{C}^d, \quad (6.14)$$

when

$$F(z) = \sum_{\alpha \in \mathbf{N}^{d+n}} c(F; \alpha) e_\alpha(z), \quad z \in \mathbf{C}^{d+n}.$$

Now let $\vartheta_{r,(s_2,s_1)}$ and $\omega_{s_2,s_1;r_2,r_1}$ be the same as in Definitions 1.4 and 1.5. Then it follows that

$$\begin{aligned} \sup_{\alpha_j \in \mathbf{N}^{d_j}} |c(K; \iota_{\tau,d_2}^*(\alpha_2), \iota_{\tau,d_1}^*(\alpha_1)) \vartheta_{r,(s_2,s_1)}(\iota_{\tau,d_2}^*(\alpha_2), \iota_{\tau,d_1}^*(\alpha_1))| \\ \leq \sup_{\alpha_j \in \mathbf{N}^{d_j+n_j}} |c(K; \alpha_2, \alpha_1) \vartheta_{r,(s_2,s_1)}(\alpha_2, \alpha_1)|, \end{aligned}$$

when $K \in \widehat{\mathcal{A}}_0(\mathbf{C}^{d_2+n_2} \times \mathbf{C}^{d_1+n_1})$, and similarly with

$$1/\vartheta_{r,(s_2,s_1)}, \quad \omega_{s_2,s_1;r_2,r_1} \quad \text{or} \quad 1/\omega_{s_2,s_1;r_2,r_1}$$

in place of $\vartheta_{r,(s_2,s_1)}$. The following twisted trace result is now a straight-forward consequence of these observations. The details are left for the reader.

Proposition 6.3. *Let $s_1, s_2 \in \overline{\mathbf{R}}_{\flat}$, $s = (s_1, s_2)$, $\text{Tr}_{\tau,d}$ be as in (3) in Definition 6.2, $W_1 = \mathbf{C}^{d_2} \times \mathbf{C}^{d_1}$ and $W_2 = \mathbf{C}^{d_2+n_2} \times \mathbf{C}^{d_1+n_1}$. Then $\text{Tr}_{\tau,d}$ restricts to continuous mappings from*

$$\widehat{\mathcal{A}}_s(W_2), \quad \widehat{\mathcal{A}}_s^*(W_2), \quad \widehat{\mathcal{B}}_s(W_2), \quad \widehat{\mathcal{B}}_s^*(W_2), \quad \widehat{\mathcal{C}}_s(W_2), \quad \text{and} \quad \widehat{\mathcal{C}}_s^*(W_2)$$

to

$$\widehat{\mathcal{A}}_s(W_1), \quad \widehat{\mathcal{A}}_s^*(W_1), \quad \widehat{\mathcal{B}}_s(W_1), \quad \widehat{\mathcal{B}}_s^*(W_1), \quad \widehat{\mathcal{C}}_s(W_1), \quad \text{and} \quad \widehat{\mathcal{C}}_s^*(W_1).$$

If instead $s_1, s_2 \in \mathbf{R}_{\flat,\infty}$, then the same holds true with $\widehat{\mathcal{A}}_{0,s}$, $\widehat{\mathcal{B}}_{0,s}$ and $\widehat{\mathcal{C}}_{0,s}$ in place of $\widehat{\mathcal{A}}_s$, $\widehat{\mathcal{B}}_s$ and $\widehat{\mathcal{C}}_s$, respectively, at each occurrence.

We observe that we may combine Proposition 6.3 and Theorem 6.1 to get an alternative proof of Theorem 4.12.

APPENDIX A. IDENTIFICATIONS OF SPACES OF POWER SERIES EXPANSIONS IN TERMS OF SPACES OF ANALYTIC FUNCTIONS

In this appendix we identify the spaces in (1.38) and (1.39) by convenient subspaces of $\widehat{\mathcal{A}}(W)$, for suitable $s \in \overline{\mathbf{R}}_{\flat}$.

We start with the following. We omit the proof since the result is an immediate consequence of Theorems 4.1, 4.2, 5.2 and 5.3 in [35] and Definition 1.13. Here let

$$\kappa_{1,r,s}(z) = \begin{cases} e^{r(\log\langle z \rangle)^{\frac{1}{1-2s}}}, & s < \frac{1}{2} \\ e^{r|z|^{\frac{2\sigma}{\sigma+1}}}, & s = \flat_{\sigma}, \sigma > 0, \\ e^{\frac{|z|^2}{2} - r|z|^{\frac{1}{s}}}, & \frac{1}{2} \leq s < \infty, \\ e^{\frac{|z|^2}{2} \langle z \rangle^{-r}}, & s = \infty, \end{cases} \quad (\text{A.1})$$

$$\kappa_{2,r,s}(z) = \begin{cases} e^{r|z|^{\frac{2\sigma}{\sigma-1}}}, & s = \flat_{\sigma}, \sigma > 1, \\ e^{\frac{|z|^2}{2} + r|z|^{\frac{1}{s}}}, & \frac{1}{2} \leq s < \infty, \\ e^{\frac{|z|^2}{2} \langle z \rangle^r}, & s = \infty, \end{cases} \quad (\text{A.2})$$

and

$$\kappa_{j,r,s}^0(z) = \begin{cases} \kappa_{j,r,s}(z), & s \neq \frac{1}{2}, \\ e^{r|z|^2}, & s = \frac{1}{2}, \end{cases} \quad j = 1, 2. \quad (\text{A.3})$$

Theorem A.1. *Let $s_j, t_j \in \overline{\mathbf{R}_{b,\infty}}$ be such that $t_j > b_1$, $j = 1, 2$, $W = \mathbf{C}^{d_2} \times \mathbf{C}^{d_1}$ and let $\kappa_{j,r,s}$ and $\kappa_{j,r,s}^0$ be given by (A.1)–(A.3), $j = 1, 2$, when $r > 0$. Then the following is true:*

- (1) *if $s_1, s_2 < \infty$, then $\widehat{\mathcal{A}}_{(s_2, s_1)}(W)$ consists of all $K \in \widehat{A}(W)$ such that $|K(z_2, z_1)| \lesssim \kappa_{1,r,s_2}(z_2)\kappa_{1,r,s_1}(z_1)$ for some $r > 0$;*
- (2) *if $s_1, s_2 > 0$, then $\widehat{\mathcal{A}}_{0,(s_2, s_1)}(W)$ consists of all $K \in \widehat{A}(W)$ such that $|K(z_2, z_1)| \lesssim \kappa_{1,r,s_2}^0(z_2)\kappa_{1,r,s_1}^0(z_1)$ for every $r > 0$;*
- (3) *if $t_1, t_2 < \infty$, then $\widehat{\mathcal{A}}'_{(t_2, t_1)}(W)$ consists of all $K \in \widehat{A}(W)$ such that $|K(z_2, z_1)| \lesssim \kappa_{2,r,t_2}(z_2)\kappa_{2,r,t_1}(z_1)$ for every $r > 0$;*
- (4) *$\widehat{\mathcal{A}}'_{0,(t_2, t_1)}(W)$ consists of all $K \in \widehat{A}(W)$ such that $|K(z_2, z_1)| \lesssim \kappa_{2,r,t_2}^0(z_2)\kappa_{2,r,t_1}^0(z_1)$ for some $r > 0$.*

By Remark 1.14 it follows that Theorem A.1 remains true after the spaces in (1.35) are replaced by corresponding spaces in (1.28).

By similar arguments as in the proofs of Theorems 4.1, 4.2, 5.2 and 5.3 in [35] we get the following two theorems. The details are left for the reader.

Theorem A.2. *Let $s_1, s_2 \in \overline{\mathbf{R}_{b,\infty}}$, $W = \mathbf{C}^{d_2} \times \mathbf{C}^{d_1}$, and let $\kappa_{j,r,s}$ and $\kappa_{j,r,s}^0$ be given by (A.1)–(A.3), $j = 1, 2$, when $r > 0$. Then the following is true:*

- (1) *if $s_1 > b_1$ and $s_1, s_2 < \infty$, then $\widehat{\mathcal{B}}_{(s_2, s_1)}(W)$ consists of all $K \in \widehat{A}(W)$ such that for every $r_1 > 0$ there is an $r_2 > 0$ such that $|K(z_2, z_1)| \lesssim \kappa_{1,r_2,s_2}(z_2)\kappa_{2,r_1,s_1}(z_1)$;*
- (2) *if $s_1 > b_1$ and $s_2 > 0$, then $\widehat{\mathcal{B}}_{0,(s_2, s_1)}(W)$ consists of all $K \in \widehat{A}(W)$ such that for every $r_2 > 0$ there is an $r_1 > 0$ such that $|K(z_2, z_1)| \lesssim \kappa_{1,r_2,s_2}^0(z_2)\kappa_{2,r_1,s_1}^0(z_1)$;*
- (3) *if $s_2 > b_1$ and $s_1, s_2 < \infty$, then $\widehat{\mathcal{B}}_{(s_2, s_1)}^*(W)$ consists of all $K \in \widehat{A}(W)$ such that for every $r_2 > 0$ there is an $r_1 > 0$ such that $|K(z_2, z_1)| \lesssim \kappa_{2,r_2,s_2}^0(z_2)\kappa_{1,r_1,s_1}^0(z_1)$;*
- (4) *if $s_1 > 0$ and $s_2 > b_1$, then $\widehat{\mathcal{B}}_{0,(s_2, s_1)}^*(W)$ consists of all $K \in \widehat{A}(W)$ such that for every $r_1 > 0$ there is an $r_2 > 0$ such that $|K(z_2, z_1)| \lesssim \kappa_{2,r_2,s_2}^0(z_2)\kappa_{1,r_1,s_1}^0(z_1)$.*

Theorem A.3. *Let $s_1, s_2 \in \overline{\mathbf{R}_{b,\infty}}$, $W = \mathbf{C}^{d_2} \times \mathbf{C}^{d_1}$, and let $\kappa_{j,r,s}$ and $\kappa_{j,r,s}^0$ be given by (A.1)–(A.3), $j = 1, 2$, when $r > 0$. Then the following is true:*

- (1) *if $s_1 > b_1$ and $s_1, s_2 < \infty$, then $\widehat{\mathcal{C}}_{(s_2, s_1)}(\mathbf{C}^{d_2} \times \mathbf{C}^{d_1})$ consists of all $K \in \widehat{A}(\mathbf{C}^{d_2} \times \mathbf{C}^{d_1})$ such that for some $r_2 > 0$ it holds $|K(z_2, z_1)| \lesssim \kappa_{1,r_2,s_2}(z_2)\kappa_{2,r_1,s_1}(z_1)$ for every $r_1 > 0$;*
- (2) *if $s_1 > b_1$ and $s_2 > 0$, then $\widehat{\mathcal{C}}_{0,(s_2, s_1)}(\mathbf{C}^{d_2} \times \mathbf{C}^{d_1})$ consists of all $K \in \widehat{A}(\mathbf{C}^{d_2} \times \mathbf{C}^{d_1})$ such that for some $r_1 > 0$ it holds $|K(z_2, z_1)| \lesssim \kappa_{1,r_2,s_2}^0(z_2)\kappa_{2,r_1,s_1}^0(z_1)$ for every $r_2 > 0$;*
- (3) *if $s_2 > b_1$ and $s_1, s_2 < \infty$, then $\widehat{\mathcal{C}}_{(s_2, s_1)}^*(\mathbf{C}^{d_2} \times \mathbf{C}^{d_1})$ consists of all $K \in \widehat{A}(\mathbf{C}^{d_2} \times \mathbf{C}^{d_1})$ such that for some $r_1 > 0$ it holds $|K(z_2, z_1)| \lesssim \kappa_{2,r_2,s_2}^0(z_2)\kappa_{1,r_1,s_1}^0(z_1)$ for every $r_2 > 0$;*
- (4) *if $s_1 > 0$ and $s_2 > b_1$, then $\widehat{\mathcal{C}}_{0,(s_2, s_1)}^*(\mathbf{C}^{d_2} \times \mathbf{C}^{d_1})$ consists of all $K \in \widehat{A}(\mathbf{C}^{d_2} \times \mathbf{C}^{d_1})$ such that for some $r_2 > 0$ it holds $|K(z_2, z_1)| \lesssim \kappa_{2,r_2,s_2}^0(z_2)\kappa_{1,r_1,s_1}^0(z_1)$ for every $r_1 > 0$.*

APPENDIX B. THE LINK BETWEEN WICK AND ANTI-WICK SYMBOLS OF RANK ONE

In this appendix we show that if

$$b_{z,w}(\alpha, \beta) = e_\alpha(z)e_\beta(\bar{w}),$$

$t, t_0 \in \mathbf{C} \setminus 0$ are such that $t_0^2 = t$ and

$$\begin{aligned} b_{t_0 z, \bar{t}_0 w}^t(\alpha, \beta) &\equiv \pi^{-d} \int_{\mathbf{C}^d} b_{t_0 w_1, \bar{t}_0 w_1}(\alpha, \beta) e^{-(z-w_1, w-w_1)} d\lambda(w_1) \\ &= \pi^{-d} \int_{\mathbf{C}^d} e_\alpha(t_0 w_1) e_\beta(t_0 \bar{w}_1) e^{-(z-w_1, w-w_1)} d\lambda(w_1) \end{aligned}$$

(cf. (1.59)), then

$$b_{z,w}^t(\alpha, \beta) = (\mathcal{T}_{0,t} b_{z,w})(\alpha, \beta), \quad \alpha, \beta \in \mathbf{N}^d. \quad (\text{B.1})$$

In fact, since all integrations can be done coordinate wise, we may assume that $d = 1$.

First we prove (B.1) for $t = 1$ and consider first the case when $\alpha = \beta = 0$, which is the same as

$$\pi^{-1} \int_{\mathbf{C}} e^{(z-w_1)(\bar{w}-\bar{w}_1)} d\lambda(w_1) = 1. \quad (\text{B.2})$$

In fact, if z and w are real, then the latter integral is equal to

$$\pi^{-1} e^{-zw} \iint_{\mathbf{R}^2} e^{(z+w)x} e^{-i(z-w)y} e^{-x^2-y^2} dx dy = e^{-zw} e^{\frac{1}{4}(z+w)^2} e^{-\frac{1}{4}(z-w)^2} = 1.$$

The searched identity now follows by analytic continuation.

For general α and β , (1.59) and integrations by parts give

$$\begin{aligned} \pi(\alpha! \beta!)^{\frac{1}{2}} b_{z,w}^1(\alpha, \beta) &= \int_{\mathbf{C}} w_1^\alpha \bar{w}_1^\beta e^{-(z-w_1)(\bar{w}-\bar{w}_1)} d\lambda(w_1) \\ &= (-1)^\beta e^{-z\bar{w}} \int_{\mathbf{C}} w_1^\alpha e^{z\bar{w}} e^{\bar{w}w_1} (\partial_{w_1}^\beta (e^{-|w_1|^2})) d\lambda(w_1) \\ &= e^{z\bar{w}} \sum_{\gamma \leq \beta} \binom{\beta}{\gamma} \int_{\mathbf{C}} (\partial^\gamma (w_1^\alpha)) e^{z\bar{w}} (\partial_{\bar{w}_1}^{\beta-\gamma} (e^{\bar{w}w_1})) e^{-|w_1|^2} d\lambda(w_1) \\ &= e^{z\bar{w}} \sum_{\gamma \leq \alpha, \beta} \binom{\alpha}{\gamma} \binom{\beta}{\gamma} \gamma! \bar{w}^{\beta-\gamma} \int_{\mathbf{C}} w_1^{\alpha-\gamma} e^{z\bar{w}} e^{\bar{w}w_1} e^{-|w_1|^2} d\lambda(w_1) \\ &= e^{z\bar{w}} \sum_{\gamma \leq \alpha, \beta} \binom{\alpha}{\gamma} \binom{\beta}{\gamma} \gamma! \bar{w}^{\beta-\gamma} (-1)^{\alpha-\gamma} \int_{\mathbf{C}} e^{z\bar{w}} e^{\bar{w}w_1} (\partial_{\bar{w}_1}^{\alpha-\gamma} (e^{-|w_1|^2})) d\lambda(w_1) \\ &= \sum_{\gamma \leq \alpha, \beta} \binom{\alpha}{\gamma} \binom{\beta}{\gamma} \gamma! z^{\alpha-\gamma} \bar{w}^{\beta-\gamma} e^{z\bar{w}} \int_{\mathbf{C}} e^{z\bar{w}} e^{\bar{w}w_1} e^{-|w_1|^2} d\lambda(w_1) \\ &= \pi(\alpha! \beta!)^{\frac{1}{2}} \sum_{\gamma \leq \alpha, \beta} \binom{\alpha}{\gamma}^{\frac{1}{2}} \binom{\beta}{\gamma}^{\frac{1}{2}} e_{\alpha-\gamma}(z) e_{\beta-\gamma}(\bar{w}), \end{aligned}$$

and (B.1) follows for general α and β . Here the last equality follows from (B.2) and the identity

$$\alpha!^{-\frac{1}{2}} \gamma!^{\frac{1}{2}} z^{\alpha-\gamma} = \binom{\alpha}{\gamma}^{-\frac{1}{2}} e_{\alpha-\gamma}(z),$$

and (B.1) follows in the case when $t = 1$.

Next suppose that $t \in \mathbf{C} \setminus 0$ and recall that $t_0^2 = t$. By the definitions it follows that

$$(\mathcal{T}_{0,t} b_{z,w})(\alpha, \beta) = t_0^{|\alpha+\beta|} (\mathcal{T}_{0,1} b_{z/t_0, w/\bar{t}_0})(\alpha, \beta),$$

and the case $t = 1$ shows that

$$\begin{aligned} (\mathcal{T}_{0,t} b_{t_0 z, \bar{t}_0 w})(\alpha, \beta) &= \pi^{-d} t_0^{|\alpha+\beta|} \int_{\mathbf{C}^d} b_{w_1, w_1}(\alpha, \beta) e^{-(z-w_1, w-w_1)} d\lambda(w_1) \\ &= \pi^{-d} \int_{\mathbf{C}^d} b_{t_0 w_1, \bar{t}_0 w_1}(\alpha, \beta) e^{-(z-w_1, w-w_1)} d\lambda(w_1), \end{aligned}$$

which gives the assertion.

APPENDIX C. IDENTITIES OF SPACES OF KERNEL OPERATORS, WICK OPERATORS AND ANTI-WICK OPERATORS

Let $s_1, s_2 \in \overline{\mathbf{R}}_b$ be such that $s_1 < \frac{1}{2}$ and $s_2 \leq \frac{1}{2}$. By Corollary 2.6, Proposition 4.5 and Theorems 4.6–4.9, it follows that

$$\text{Op}_{\mathfrak{W}}(\widehat{\mathcal{A}}'_{s_1}(\mathbf{C}^{2d})) = \{T_K; K \in \widehat{\mathcal{A}}'_{s_1}(\mathbf{C}^{2d})\} = \mathcal{L}(\mathcal{A}_{s_1}(\mathbf{C}^d), \mathcal{A}'_{s_1}(\mathbf{C}^d)), \quad (\text{C.1})$$

$$\text{Op}_{\mathfrak{W}}(\widehat{\mathcal{A}}'_{0,s_2}(\mathbf{C}^{2d})) = \{T_K; K \in \widehat{\mathcal{A}}'_{0,s_2}(\mathbf{C}^{2d})\} = \mathcal{L}(\mathcal{A}_{0,s_2}(\mathbf{C}^d), \mathcal{A}'_{0,s_2}(\mathbf{C}^d)), \quad (\text{C.2})$$

$$\text{Op}_{\mathfrak{W}}^{\text{aw}}(\widehat{\mathcal{A}}_{s_1}(\mathbf{C}^{2d})) = \text{Op}_{\mathfrak{W}}(\widehat{\mathcal{A}}_{s_1}(\mathbf{C}^{2d})), \quad \text{Op}_{\mathfrak{W}}^{\text{aw}}(\widehat{\mathcal{A}}_{0,s_2}(\mathbf{C}^{2d})) = \text{Op}_{\mathfrak{W}}(\widehat{\mathcal{A}}_{0,s_2}(\mathbf{C}^{2d})), \quad (\text{C.3})$$

$$\begin{aligned} \text{Op}_{\mathfrak{W}}^{\text{aw}}(\widehat{\mathcal{C}}_{s_1}(\mathbf{C}^{2d})) &= \text{Op}_{\mathfrak{W}}(\widehat{\mathcal{C}}_{s_1}(\mathbf{C}^{2d})) \subseteq \text{Op}_{\mathfrak{W}}(\widehat{\mathcal{B}}_{s_1}(\mathbf{C}^{2d})) \\ &= \{T_K; K \in \widehat{\mathcal{B}}_{s_1}(\mathbf{C}^{2d})\} = \mathcal{L}(\mathcal{A}_{s_1}(\mathbf{C}^d)), \quad (\text{C.4}) \end{aligned}$$

$$\begin{aligned} \text{Op}_{\mathfrak{W}}^{\text{aw}}(\widehat{\mathcal{C}}_{s_1}^*(\mathbf{C}^{2d})) &= \text{Op}_{\mathfrak{W}}(\widehat{\mathcal{C}}_{s_1}^*(\mathbf{C}^{2d})) \subseteq \text{Op}_{\mathfrak{W}}(\widehat{\mathcal{B}}_{s_1}^*(\mathbf{C}^{2d})) \\ &= \{T_K; K \in \widehat{\mathcal{B}}_{s_1}^*(\mathbf{C}^{2d})\} = \mathcal{L}(\mathcal{A}'_{s_1}(\mathbf{C}^d)), \quad (\text{C.5}) \end{aligned}$$

$$\begin{aligned} \text{Op}_{\mathfrak{W}}^{\text{aw}}(\widehat{\mathcal{C}}_{0,s_2}(\mathbf{C}^{2d})) &= \text{Op}_{\mathfrak{W}}(\widehat{\mathcal{C}}_{0,s_2}(\mathbf{C}^{2d})) \subseteq \text{Op}_{\mathfrak{W}}(\widehat{\mathcal{B}}_{0,s_2}(\mathbf{C}^{2d})) \\ &= \{T_K; K \in \widehat{\mathcal{B}}_{0,s_2}(\mathbf{C}^{2d})\} = \mathcal{L}(\mathcal{A}_{0,s_2}(\mathbf{C}^d)), \quad (\text{C.6}) \end{aligned}$$

and

$$\begin{aligned} \text{Op}_{\mathfrak{W}}^{\text{aw}}(\widehat{\mathcal{C}}_{0,s_2}^*(\mathbf{C}^{2d})) &= \text{Op}_{\mathfrak{W}}(\widehat{\mathcal{C}}_{0,s_2}^*(\mathbf{C}^{2d})) \subseteq \text{Op}_{\mathfrak{W}}(\widehat{\mathcal{B}}_{0,s_2}^*(\mathbf{C}^{2d})) \\ &= \{T_K; K \in \widehat{\mathcal{B}}_{0,s_2}^*(\mathbf{C}^{2d})\} = \mathcal{L}(\mathcal{A}'_{0,s_2}(\mathbf{C}^d)). \quad (\text{C.7}) \end{aligned}$$

REFERENCES

- [1] A. Abdeljawad, M. Capiello, J. Toft, *Pseudo-differential calculus in anisotropic Gelfand-Shilov setting*, Integr. Equ. Oper. Theory **91** (2019), 91–26.
- [2] A. Abdeljawad, S. Coriasco, N. Teofanov, *Bilinear Pseudo-differential Operators with Gevrey–Hörmander Symbols*, Mediterr. J. Math. **17** 120 (2020).
- [3] A. Abdeljawad, C. Fernandez, A. Galbis, J. Toft, R. Üster, *Characterizations of Pilipović spaces by powers of harmonic oscillator*, RACSAM **114** (2020), 131.
- [4] V. Bargmann, *On a Hilbert space of analytic functions and an associated integral transform*, Comm. Pure Appl. Math. **14** (1961), 187–214.
- [5] V. Bargmann, *On a Hilbert space of analytic functions and an associated integral transform. Part II. A family of related function spaces. Application to distribution theory*, Comm. Pure Appl. Math. **20** (1967), 1–101.
- [6] F. A. Berezin, *Wick and anti-Wick symbols of operators*, Mat. Sb. (N.S.) **86** (1971), 578–610.
- [7] F. A. Berezin, *Covariant and contravariant symbols of operators*, Izv. Akad. Nauk SSSR **36** (1972), 1134–1167.

- [8] J. M. Bony, J. Y. Chemin *Espaces Fonctionnels Associés au Calcul de Weyl-Hörmander*, Bull. Soc. math. France **122** (1994), 77–118.
- [9] M. Cappiello, R. Schulz, P. Wahlberg, *Conormal distributions in the Shubin calculus of pseudodifferential operators*, J. Math. Phys. **59** (2018), 021502.
- [10] M. Cappiello, J. Toft, *Pseudo-differential operators in a Gelfand-Shilov setting*, Math. Nachr. **290** (5-6) (2017), 738–755.
- [11] E. Cordero, S. Pilipović, L. Rodino, N. Teofanov, *Quasianalytic Gelfand-Shilov spaces with applications to localization operators*, Rocky Mountain. J. Math. **40** (2010), 1123-1147.
- [12] P. Boggiatto, E. Buzano, L. Rodino *Global Hypoellipticity and Spectral Theory*, Akademie Verlag, Berlin, 1996.
- [13] G. B. Folland, *Harmonic analysis in phase space*, Princeton University Press, Princeton, 1989.
- [14] I. M. Gelfand, G. E. Shilov, *Generalized functions, II-III*, Academic Press, New York London, 1968.
- [15] K. Gröchenig, *Foundations of time-frequency analysis*, Birkhäuser, Boston, 2001.
- [16] K. Gröchenig, *Weight functions in time-frequency analysis*, in: L. Rodino, M. W. Wong (Eds.) *Pseudodifferential Operators: Partial Differential Equations and Time-Frequency Analysis*, Fields Institute Comm., **52** (2007), pp. 343–366.
- [17] K. Gröchenig, G. Zimmermann, *Spaces of test functions via the STFT* J. Funct. Spaces Appl. **2** (2004), 25–53.
- [18] B. C. Hall, *Quantum Theory for Mathematicians*, Springer, New York, 2013.
- [19] L. Hörmander, *The analysis of linear partial differential operators*, vol I–III, Springer-Verlag, Berlin Heidelberg New York Tokyo, 1983, 1985.
- [20] E. H. Lieb, *Integral bounds for radar ambiguity functions and Wigner distributions*, J. Math. Phys. **31** (1990), 594–599.
- [21] E. H. Lieb, *The stability of matter from atoms to stars*, Bull. Amer. Math. Soc. **22** (1990), 1–49.
- [22] E. H. Lieb, J. P. Solovej, *Quantum coherent operators: a generalization of coherent states*, Lett. Math. Phys. **22** (1991), 145–154.
- [23] Z. Lozanov Crvenković, D. Perišić, *Hermite expansions of elements of Gelfand Shilov spaces in quasianalytic and non quasianalytic case*, Novi Sad J. Math. **37** (2007), 129–147.
- [24] S. Pilipović *Tempered ultradistributions*, Boll. U.M.I. **7** (1988), 235–251.
- [25] B. Prangoski, *Pseudodifferential operators of infinite order in spaces of tempered ultradistributions*, J. Pseudo-Differ. Oper. Appl. **4** (2013), 495–549.
- [26] M. Reed and B. Simon, *Methods of modern mathematical physics*, Academic Press, London New York, 1979.
- [27] H. H. Schaefer, M. P. Wolff *Topological vector spaces, 2nd. Ed.*, Graduate Texts in Mathematics, **3**, Springer-Verlag, New York, 1999.
- [28] M. A. Shubin, *Pseudodifferential operators and spectral theory*, Springer, 2001.
- [29] N. Teofanov, *Ultradistributions and time-frequency analysis*. In “Pseudo-differential operators and related topics”, 173–192, Oper. Theory Adv. Appl., **164**, Birkhäuser, Basel, 2006.
- [30] N. Teofanov, *Gelfand-Shilov spaces and localization operators*, Funct. Anal. Approx. Comput. **7** (2015), 135–158.
- [31] N. Teofanov, J. Toft, *Pseudo-differential calculus in a Bargmann setting*, Ann. Acad. Sci. Fenn. Math. **45** (2020), 227–257.
- [32] N. Teofanov, J. Toft, P. Wahlberg, *Pseudo-differential operators with isotropic symbols, Wick and anti-Wick operators, and hypoellipticity*, preprint, arXiv:2011.00313.
- [33] J. Toft, *The Bargmann transform on modulation and Gelfand-Shilov spaces, with applications to Toeplitz and pseudo-differential operators*, J. Pseudo-Differ. Oper. Appl. **3** (2012), 145–227.
- [34] *Matrix parameterized pseudo-differential calculi on modulation spaces* in: M. Oberguggenberger, J. Toft, J. Vindas, P. Wahlberg (eds), *Generalized functions and Fourier analysis*, Operator Theory: Advances and Applications **260** Birkhäuser, 2017, pp. 215–235.
- [35] J. Toft, *Images of function and distribution spaces under the Bargmann transform*, J. Pseudo-Differ. Oper. Appl. **8** (2017), 83–139.
- [36] F. Trèves, *Topological vector spaces, distributions and kernels*, Academic Press, New York London, 1967.
- [37] D. Vogt *Regularity properties of (LF)-spaces*, *Progress in functional analysis (Peñíscola, 1990)*, North-Holland Math. Stud. **170**, North-Holland, Amsterdam, 1992, pp. 57–84.
- [38] J. Wengenroth *Acyclic inductive spectra of Fréchet spaces*, Stud. Math. **120** (1996), 247–258.
- [39] G. C. Wick, *The evaluation of the collision matrix*, Phys. Rev. **80** (1950), 268–272.

DEPARTMENT OF MATHEMATICS, LINNÆUS UNIVERSITY, VÄXJÖ, SWEDEN
Email address: joachim.toft@lnu.se