

SOME MATRIX INEQUALITIES OF LOG-MAJORIZATION TYPE

BO-YAN XI AND FUZHEN ZHANG

ABSTRACT. The purpose of this paper is two-fold: we present some matrix inequalities of log-majorization type for eigenvalues indexed by a sequence; we then apply our main theorem to generalize and improve the Hua-Marcus' inequalities. Our results are stronger and more general than the existing ones.

1. INTRODUCTION

Let $\mathbb{C}^{m \times n}$ be the space of $m \times n$ complex matrices. For $A \in \mathbb{C}^{n \times n}$, we denote the eigenvalues of A by $\lambda_1(A), \lambda_2(A), \dots, \lambda_n(A)$ and the singular values of A by $\sigma_1(A), \sigma_2(A), \dots, \sigma_n(A)$. If $\lambda_1(A), \lambda_2(A), \dots, \lambda_n(A)$ are all real, we arrange them in decreasing order: $\lambda_1(A) \geq \lambda_2(A) \geq \dots \geq \lambda_n(A)$. The singular values are always ordered decreasingly: $\sigma_1(A) \geq \sigma_2(A) \geq \dots \geq \sigma_n(A)$.

Horn conjecture ...

There is a large family of matrix inequalities concerning the eigenvalues and singular values of the product (including Schur or Hadamard product) and sum of matrices. These inequalities may be sorted in four types: $\sum \zeta_{\mathcal{I}}(A+B)$, $\prod \zeta_{\mathcal{I}}(A+B)$, $\sum \zeta_{\mathcal{I}}(AB)$, $\prod \zeta_{\mathcal{I}}(AB)$, where A and B are generic matrices, and $\zeta_{\mathcal{I}}(\cdot)$ represents selected eigenvalues or singular values indexed by a sequence \mathcal{I} . The inequalities in sum \sum are usually called majorization type ([7, p. 45], [13]), while the ones in product \prod are referred to as logarithmic (log-) majorization type. See [13, p. 16], [3, 9]). For example, the following inequalities [6] (or [13, G.2.a, p. 333]) are log-majorization type with $\mathcal{I} = \{k, \dots, n\}$: for each $k = 1, \dots, n$,

$$\prod_{t=k}^n \lambda_t(A+B) \geq \prod_{t=k}^n [\lambda_t(A) + \lambda_t(B)] \geq \prod_{t=k}^n \lambda_t(A) + \prod_{t=k}^n \lambda_t(B), \quad (1.1)$$

where A and B are $n \times n$ positive semidefinite matrices.

Closely related inequalities of the same type, due to Oppenheim [14] (or [13, F.2, p. 685]), are, for $n \times n$ positive semidefinite A and B , and for each $k = 1, \dots, n$,

$$\left[\prod_{t=k}^n \lambda_t(A+B) \right]^{\frac{1}{n-k+1}} \geq \left[\prod_{t=k}^n \lambda_t(A) \right]^{\frac{1}{n-k+1}} + \left[\prod_{t=k}^n \lambda_t(B) \right]^{\frac{1}{n-k+1}}, \quad (1.2)$$

which give the Minkowski inequality by setting $k = 1$ ([13, p. 685] or [23, p. 215])

$$[\det(A+B)]^{\frac{1}{n}} \geq [\det(A)]^{\frac{1}{n}} + [\det(B)]^{\frac{1}{n}}. \quad (1.3)$$

In [20], we showed majorization inequalities of $\sum \zeta_{\mathcal{I}}(AB)$ for Hermitian matrices A and B with an arbitrary index set \mathcal{I} . In this paper, we present some

2020 *Mathematics Subject Classification.* 15A42, 15A45, 47A63.

Key words and phrases. Eigenvalue; Hua's determinant inequality; majorization inequality.

Xi's work was partially supported by NNSF of China grant No. 11361038.

log-majorization inequalities of $\prod \zeta_{\mathcal{I}}(A+B)$ with any index set \mathcal{I} for positive semidefinite matrices A and B , extending (1.1) and (1.2). Our theorems are more general and stronger than some existing results. We apply main results to improve the Hua-Marcus inequalities for contractive matrices.

2. SOME LEMMAS

Let X^* denote the conjugate transpose of matrix or vector X . For a square matrix A , we write $A \geq 0$ if A is positive semidefinite and $A > 0$ if A is positive definite. For Hermitian matrices $A, B \in \mathbb{C}^{n \times n}$, we write $A \geq B$ if $A - B \geq 0$.

Through the paper, let n be a positive integer. Let $\mathcal{I} = \{i_1, \dots, i_k\}$, where $1 \leq i_1 < \dots < i_k \leq n$ is any subsequence of $1, \dots, n$, $k = 1, \dots, n$. If $A \in \mathbb{C}^{n \times n}$ is positive semidefinite, then $\lambda_{i_1}(A) \geq \dots \geq \lambda_{i_k}(A)$ are k eigenvalues of A indexed by $\mathcal{I} = \{i_1, \dots, i_k\}$. For $t = 1, \dots, k$, setting $i_t = t$ gives the first k largest eigenvalues of A ; putting $i_t = n - k + t$ gives the last k smallest eigenvalues of A .

Lemma 2.1 (Hoffman [1, Cor. 2.5]). *Let $A \in \mathbb{C}^{n \times n}$ be positive semidefinite. Then*

$$\prod_{t=1}^k \lambda_{i_t}(A) = \max_{\substack{\mathbb{S}_1 \subset \dots \subset \mathbb{S}_k \subset \mathbb{C}^n \\ \dim \mathbb{S}_t = i_t}} \min_{\substack{x_t \in \mathbb{S}_t, (x_r, x_s) = \delta_{rs} \\ U_k = (x_1, \dots, x_k)}} \det(U_k^* A U_k),$$

where δ_{rs} is the Kronecker delta, i.e., $\delta_{rs} = 1$ if $r = s$, or 0 otherwise.

Lemma 2.2 (Lidskiĭ [11]). *Let $A, B \in \mathbb{C}^{n \times n}$ be positive semidefinite. Then*

$$\prod_{t=1}^k \lambda_{i_t}(A) \lambda_{n-t+1}(B) \leq \prod_{t=1}^k \lambda_{i_t}(AB) \leq \prod_{t=1}^k \lambda_{i_t}(A) \lambda_t(B). \quad (2.1)$$

More general inequalities of (2.1) for singular values are due to Gel'fand and Naimark (see, e.g., [13, p. 340]). The inequalities on the left-hand side of (2.1) can be found explicitly in [17].

Lemma 2.3. *Let a_1, \dots, a_n and b_1, \dots, b_n be nonnegative real numbers. Then*

$$\prod_{t=1}^n (a_t + b_t) \geq \prod_{t=1}^n a_t + \prod_{t=1}^n b_t + (2^n - 2) \left[\prod_{t=1}^n (a_t b_t) \right]^{\frac{1}{2}}. \quad (2.2)$$

In particular, for $x, y \geq 0$,

$$(x^{\frac{1}{n}} + y^{\frac{1}{n}})^n \geq x + y + (2^n - 2)(xy)^{\frac{1}{2}}. \quad (2.3)$$

Proof. Use induction on n and apply the arithmetic-geometric mean inequality. \square

We remark that Lemma 2.3 implies immediately a result of Hartfiel [8]:

$$\det(A + B) \geq \det A + \det B + (2^n - 2)[\det A \det B]^{\frac{1}{2}} \quad (2.4)$$

because of the fact that any two positive semidefinite matrices of the same size are simultaneously $*$ -congruent to diagonal matrices (see, e.g., [23, p. 209]).

3. MAIN RESULTS

We begin with a result that is important to the proof of our main theorem.

Lemma 3.1. *Let $D = \text{diag}(\lambda_1, \lambda_2, \dots, \lambda_n)$, where $\lambda_1 \geq \lambda_2 \geq \dots \geq \lambda_n \geq 0$, and let $U_k = (u_1, \dots, u_k) = (u_{ij})$ be a partial isometry, i.e., U_k is an $n \times k$ matrix such that $U_k^* U_k = I_k$. If $m \geq 1$ and $u_{i1} = 0$ for all $i > m$, then*

$$\det(U_k^* D U_k) \geq \lambda_m \det[(u_2, \dots, u_k)^* D_m (u_2, \dots, u_k)], \quad (3.1)$$

where $D_m = \text{diag}(\underbrace{\lambda_m, \dots, \lambda_m}_m, \lambda_{m+1}, \dots, \lambda_n)$.

Proof. Let $V = (u_2, \dots, u_k)$. Since $D \geq D_m$ and $u_{i1} = 0$ for $i > m$, we have

$$\begin{aligned} \det(U_k^* D U_k) &\geq \det(U_k^* D_m U_k) = \det \begin{pmatrix} \lambda_m u_1^* u_1 & \lambda_m u_1^* V \\ \lambda_m V^* u_1 & V^* D_m V \end{pmatrix} \\ &= \det \begin{pmatrix} \lambda_m & 0 \\ 0 & V^* D_m V \end{pmatrix} = \lambda_m \det(V^* D_m V) \\ &= \lambda_m \det[(u_2, \dots, u_k)^* D_m (u_2, \dots, u_k)]. \end{aligned}$$

□

We are ready to present our main result.

Theorem 3.1. *Let A and B be $n \times n$ positive semidefnite matrices. Then*

$$\prod_{t=1}^k \lambda_{i_t}^{\frac{1}{k}}(A+B) \geq \prod_{t=1}^k \lambda_{i_t}^{\frac{1}{k}}(A) + \prod_{t=1}^k \lambda_{n-t+1}^{\frac{1}{k}}(B) \quad (3.2)$$

and

$$\prod_{t=1}^k \lambda_{i_t}(A+B) \geq \prod_{t=1}^k \lambda_{i_t}(A) + \prod_{t=1}^k \lambda_{n-t+1}(B) + (2^k - 2) \prod_{t=1}^k [\lambda_{i_t}(A) \lambda_{n-t+1}(B)]^{\frac{1}{2}}. \quad (3.3)$$

Proof. If $n = 1$, the inequalities become equalities and hold trivially. Let $n \geq 2$. By spectral decomposition, there exists a unitary matrix $U = (u_1, u_2, \dots, u_n) \in \mathbb{C}^{n \times n}$, where u_1, u_2, \dots, u_n are orthonormal eigenvectors associated with $\lambda_1(A), \lambda_2(A), \dots, \lambda_n(A)$, respectively, such that

$$A = U \text{diag}(\lambda_1(A), \lambda_2(A), \dots, \lambda_n(A)) U^*.$$

For each i_t in the sequence $i_1 < i_2 < \dots < i_k$, let $\mathbb{S}_t^{(0)} = \text{Span}(u_1, u_2, \dots, u_{i_t})$, $t = 1, 2, \dots, k$. Then $\mathbb{S}_1^{(0)} \subset \mathbb{S}_2^{(0)} \subset \dots \subset \mathbb{S}_k^{(0)}$ and $\dim(\mathbb{S}_t^{(0)}) = i_t$, $t = 1, 2, \dots, k$.

Let $\{x_1, \dots, x_k\}$ be any set of orthonormal vectors, where $x_t \in \mathbb{S}_t^{(0)}$, $t = 1, 2, \dots, k$. Let $x_t = a_{1t}u_1 + \dots + a_{i_t t}u_{i_t} + 0u_{i_t+1} + \dots + 0u_{i_k} = (u_1, \dots, u_{i_k})\alpha_t$, where $\alpha_t = (a_{1t}, \dots, a_{i_t t}, 0, \dots, 0)^T$ (here T is for transpose), $t = 1, 2, \dots, k$. Then

$$U_k = (x_1, \dots, x_k) = (u_1, \dots, u_{i_k})(\alpha_1, \dots, \alpha_k).$$

Let $V_k = (\alpha_1, \dots, \alpha_k) \in \mathbb{C}^{i_k \times k}$. Since $\{x_1, \dots, x_k\}$ and $\{u_1, \dots, u_{i_k}\}$ are orthonormal sets, we see that $V_k^* V_k = I_k$. So V_k is a partial isometry, and the components of the first column of V_k are zeros except the first i_1 components (i.e., the last $i_k - i_1$ components of α_1 are all equal to zero).

In Lemma 3.1, setting $n = i_k$, $m = i_1$, $U_k = V_k$, and applying (3.1) to the $i_k \times i_k$ matrix $D_{i_1} = \text{diag}(\underbrace{\lambda_{i_1}(A), \dots, \lambda_{i_1}(A)}_{i_1}, \lambda_{i_1+1}(A), \dots, \lambda_{i_k}(A))$, we obtain

$$\begin{aligned} \det(U_k^* A U_k) &= \det [(\alpha_1, \dots, \alpha_k)^* (u_1, \dots, u_{i_k})^* A (u_1, \dots, u_{i_k}) (\alpha_1, \dots, \alpha_k)] \\ &= \det [(\alpha_1, \dots, \alpha_k)^* \text{diag}(\lambda_1(A), \dots, \lambda_{i_k}(A)) (\alpha_1, \dots, \alpha_k)] \\ &= \det [V_k^* \text{diag}(\lambda_1(A), \dots, \lambda_{i_k}(A)) V_k] \\ &\geq \lambda_{i_1}(A) \det [(\alpha_2, \dots, \alpha_k)^* D_{i_1} (\alpha_2, \dots, \alpha_k)]. \end{aligned}$$

Repeatedly using Lemma 3.1, we get

$$\det(U_k^* A U_k) \geq \prod_{t=1}^k \lambda_{i_t}(A). \quad (3.4)$$

A minimax eigenvalue result of Fan (see [4, 5] or [13, A.3.a, p. 787]) ensures

$$\det(U_k^* B U_k) \geq \min_{U^* U = I_k} \det(U^* B U) = \prod_{t=1}^k \lambda_{n-t+1}(B).$$

An application of the Minkowski inequality (1.3) together with (3.4) reveals

$$\begin{aligned} \det [U_k^* (A + B) U_k] &= \det(U_k^* A U_k + U_k^* B U_k) \\ &\geq \left[(\det(U_k^* A U_k))^{\frac{1}{k}} + (\det(U_k^* B U_k))^{\frac{1}{k}} \right]^k \\ &\geq \left[\prod_{t=1}^k \lambda_{i_t}^{\frac{1}{k}}(A) + \prod_{t=1}^k \lambda_{n-t+1}^{\frac{1}{k}}(B) \right]^k. \end{aligned}$$

Thus, by Lemma 2.1, we derive (3.2) and (3.3) as follows:

$$\begin{aligned} \prod_{t=1}^k \lambda_{i_t}(A + B) &= \max_{\substack{\mathbb{S}_1 \subset \dots \subset \mathbb{S}_k \subset \mathbb{C}^n \\ \dim \mathbb{S}_t = i_t}} \min_{\substack{y_t \in \mathbb{S}_t, (y_r, y_s) = \delta_{rs} \\ W_k = (y_1, \dots, y_k)}} \det [W_k^* (A + B) W_k] \\ &\geq \min_{\substack{x_t \in \mathbb{S}_t^{(0)} (x_r, x_s) = \delta_{rs} \\ U_k = (x_1, \dots, x_k)}} \det [U_k^* (A + B) U_k] \\ &\geq \left[\prod_{t=1}^k \lambda_{i_t}^{\frac{1}{k}}(A) + \prod_{t=1}^k \lambda_{n-t+1}^{\frac{1}{k}}(B) \right]^k \\ &\geq \prod_{t=1}^k \lambda_{i_t}(A) + \prod_{t=1}^k \lambda_{n-t+1}(B) \\ &\quad + (2^k - 2) \prod_{t=1}^k [\lambda_{i_t}(A) \lambda_{n-t+1}(B)]^{\frac{1}{2}}. \end{aligned}$$

The last inequality is by (2.3) with $x = \prod_{t=1}^k \lambda_{i_t}(A)$ and $y = \prod_{t=1}^k \lambda_{n-t+1}(B)$. \square

In (3.2), letting $i_t = n - k + t$, $t = 1, 2, \dots, k$, we arrive at the inequalities (1.2) of Oppenheim for the product of k smallest eigenvalues. Setting $i_t = t$, $t = 1, 2, \dots, k$, we obtain analogous inequalities of 1.2 for the product of k largest eigenvalues.

Corollary 3.1. *Let A and B be $n \times n$ positive semidefinite matrices. Then*

$$\prod_{t=1}^k \lambda_t^{\frac{1}{k}}(A+B) \geq \prod_{t=1}^k \lambda_t^{\frac{1}{k}}(A) + \prod_{t=1}^k \lambda_{n-t+1}^{\frac{1}{k}}(B). \quad (3.5)$$

What follows is a lower bound for the product of any two eigenvalues of the sum in terms of the eigenvalues of individual matrices.

Corollary 3.2. *Let A and B be $n \times n$ positive semidefinite matrices. Then*

$$\begin{aligned} \lambda_i(A+B)\lambda_j(A+B) &\geq \lambda_i(A)\lambda_j(A) + \lambda_{n-1}(B)\lambda_n(B) \\ &\quad + 2[\lambda_i(A)\lambda_j(A)\lambda_{n-1}(B)\lambda_n(B)]^{\frac{1}{2}}. \end{aligned}$$

Remark: In view of Fiedler's (1.1) and inequalities (3.3), it is tempting to have

$$\prod_{t=1}^k \lambda_{i_t}(A+B) \geq \prod_{t=1}^k [\lambda_{i_t}(A) + \lambda_{n-t+1}(B)]. \quad (3.6)$$

However, this need not be true. Take $A = B = \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}$, $k = 2$, $i_1 = 1$, $i_2 = 2$. Then the left-hand side of (3.6) is 0, while the right-hand side is 1. (Note: it is always true that $\lambda_i(A+B) \geq \lambda_i(A) + \lambda_n(B)$. See, e.g., [23, p. 274].)

The following inequality in (3.7) is Fiedler's 1st inequality in (1.1). Inequality in (3.8) is stronger than Fiedler's 2nd inequality. (3.8) is proved by Lemma 2.3. The inequalities in (3.9) are immediate from the inequalities (3.3) in the theorem.

Corollary 3.3. *Let $A, B \in \mathbb{C}^{n \times n}$ be positive semidefinite matrices. Then*

$$\prod_{t=n-k+1}^n \lambda_t(A+B) \geq \prod_{t=n-k+1}^n [\lambda_t(A) + \lambda_t(B)] \quad (3.7)$$

$$\begin{aligned} &\geq \prod_{t=n-k+1}^n \lambda_t(A) + \prod_{t=n-k+1}^n \lambda_t(B) \quad (3.8) \\ &\quad + (2^k - 2) \prod_{t=n-k+1}^n [\lambda_t(A)\lambda_t(B)]^{\frac{1}{2}} \end{aligned}$$

and

$$\begin{aligned} \prod_{t=1}^k \lambda_t(A+B) &\geq \prod_{t=1}^k \lambda_t(A) + \prod_{t=1}^k \lambda_{n-t+1}(B) \quad (3.9) \\ &\quad + (2^k - 2) \prod_{t=1}^k [\lambda_t(A)\lambda_{n-t+1}(B)]^{\frac{1}{2}}. \end{aligned}$$

Remark: In view of (3.7), it is appealing in the display (3.9) to have

$$\prod_{t=1}^k \lambda_t(A+B) \geq \prod_{t=1}^k [\lambda_t(A) + \lambda_t(B)] \quad \text{or} \quad \prod_{t=1}^k [\lambda_t(A) + \lambda_{n-t+1}(B)].$$

But neither one is true. Let $A = \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}$, $B = \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix}$, $k = 1$. Then $\lambda_1(A+B) = 1 < 2 = \lambda_1(A) + \lambda_1(B)$. See the counterexample below (3.6) for the second case.

4. HUA-MARCUS INEQUALITIES FOR CONTRACTIONS

A matrix $A \in \mathbb{C}^{m \times n}$ is said to be (strictly) *contractive* if $I_n - A^*A \geq 0$ (> 0), equivalently, the largest singular value (i.e., the spectral norm) of A is less than or equal to (resp. $<$) 1. We use $\mathcal{C}_{m \times n}$ to denote the set of $m \times n$ contractive matrices and $\mathcal{SC}_{m \times n}$ to denote the set of $m \times n$ strictly contractive matrices.

In this section we apply our main theorem to derive some inequalities of Hua-Marcus type for contractive matrices. We begin by citing Hua's results in [10].

Theorem 4.1 (Hua [10, Theorems 1 and 2]). *Let $A, B \in \mathcal{SC}_{n \times n}$. Then*

$$\det(I - A^*A) \det(I - B^*B) + |\det(A - B)|^2 \leq |\det(I - A^*B)|^2. \quad (4.1)$$

Consequently,

$$\det(I - A^*A) \det(I - B^*B) \leq |\det(I - A^*B)|^2. \quad (4.2)$$

Equality in (4.1) holds if and only if $A = B$.

The following [23, Theorem 7.18] is a reversal of the Hua inequality 4.1 for general $n \times n$ matrices A and B that need not be contractive:

$$|\det(I - A^*B)|^2 \leq \det(I + A^*A) \det(I + B^*B) - |\det(A + B)|^2. \quad (4.3)$$

Marcus [12] extended the Hua inequality (4.2) to the inequalities of eigenvalues.

Theorem 4.2 (Marcus [12, Theorem]). *Let $A, B \in \mathcal{C}_{n \times n}$. Then*

$$\prod_{t=1}^k |\lambda_{n-t+1}(I - A^*B)|^2 \geq \prod_{t=1}^k [1 - \lambda_t(A^*A)] [1 - \lambda_t(B^*B)]. \quad (4.4)$$

Since $|\det(I - A^*B)| = \prod_{t=1}^n |\lambda_{n-t+1}(I - A^*B)| = \prod_{t=1}^n \sigma_{n-t+1}(I - A^*B)$, by Weyl's log-majorization inequality ([18] or [13, p. 317]), the eigenvalues in absolute values are log-majorized by the singular values, we get the stronger inequality than (4.4):

$$\prod_{t=1}^k \sigma_{n-t+1}^2(I - A^*B) \geq \prod_{t=1}^k [1 - \lambda_t(A^*A)] [1 - \lambda_t(B^*B)]. \quad (4.5)$$

Related inequalities of Hua-Marcus type are seen in [2, 14, 15, 16, 21, 22].

The following result is stronger and more general than 4.5.

Theorem 4.3. *Let $A, B \in \mathcal{SC}_{n \times n}$. Then*

$$\prod_{t=1}^k \sigma_{i_t}^2(I - A^*B) \geq \prod_{t=1}^k [1 - \lambda_t(A^*A)] [1 - \lambda_{n-i_t+1}(B^*B)] \quad (4.6)$$

$$+ \prod_{t=1}^k \frac{[1 - \lambda_t(A^*A)] \sigma_{n-t+1}^2(A - B)}{1 - \lambda_{n-t+1}(A^*A)} \\ + (2^k - 2) \prod_{t=1}^k [1 - \lambda_t(A^*A)] [\lambda_{i_t}(F) \lambda_{n-t+1}(H)]^{\frac{1}{2}}, \quad (4.7)$$

where $F = I - B^*B$ and $H = (A - B)^*(I - AA^*)^{-1}(A - B)$.

Proof. For $A, B \in \mathcal{SC}_{n \times n}$, let

$$F = I - B^*B \quad \text{and} \quad H = (A - B)^*(I - AA^*)^{-1}(A - B).$$

Then (see [10, Theorem 1] or [23, pp. 230-231])

$$F + H = (I - B^*A)(I - A^*A)^{-1}(I - A^*B). \quad (4.8)$$

Applying the left inequality of (2.1) with $i_t = n - t + 1$ to H reveals

$$\begin{aligned} \prod_{t=1}^k \lambda_{n-t+1}(H) &= \prod_{t=1}^k \lambda_{n-t+1}[(I - AA^*)^{-1}(A - B)(A - B)^*] \\ &\geq \prod_{t=1}^k \lambda_{n-t+1}[(I - AA^*)^{-1}] \sigma_{n-t+1}^2(A - B) \\ &= \prod_{t=1}^k [1 - \lambda_{n-t+1}(A^*A)]^{-1} \sigma_{n-t+1}^2(A - B). \end{aligned} \quad (4.9)$$

Applying the right inequality of (2.1) to $F + H$ in the product form in (4.8) yields

$$\prod_{t=1}^k \lambda_{i_t}(F + H) \leq \prod_{t=1}^k [1 - \lambda_t(A^*A)]^{-1} \sigma_{i_t}^2(I - A^*B). \quad (4.10)$$

Thus

$$\prod_{t=1}^k \sigma_{i_t}^2(I - A^*B) \geq \prod_{t=1}^k [1 - \lambda_t(A^*A)] \lambda_{i_t}(F + H). \quad (4.11)$$

By inequalities (3.3) and (4.9), we obtain

$$\begin{aligned} &\prod_{t=1}^k \lambda_{i_t}(F + H) \\ &\geq \prod_{t=1}^k \lambda_{i_t}(F) + \prod_{t=1}^k \lambda_{n-t+1}(H) + (2^k - 2) \prod_{t=1}^k [\lambda_{i_t}(F) \lambda_{n-t+1}(H)]^{\frac{1}{2}} \\ &\geq \prod_{t=1}^k [1 - \lambda_{n-i_t+1}(B^*B)] + \prod_{t=1}^k [1 - \lambda_{n-t+1}(A^*A)]^{-1} \sigma_{n-t+1}^2(A - B) \\ &\quad + (2^k - 2) \prod_{t=1}^k [\lambda_{i_t}(F) \lambda_{n-t+1}(H)]^{\frac{1}{2}}. \end{aligned} \quad (4.12)$$

Combining (4.11) and (4.12), we obtain the desired inequalities. \square

Setting $k = n$ in Theorem 4.3 gives a stronger version of the Hua's inequality.

Corollary 4.1. *Let $A, B \in \mathcal{SC}_{n \times n}$. Then*

$$\begin{aligned} |\det(I - A^*B)|^2 &\geq \det(I - A^*A) \det(I - B^*B) + |\det(A - B)|^2 \\ &\quad + (2^n - 2) [\det(I - A^*A) \det(I - B^*B)]^{\frac{1}{2}} |\det(A - B)|. \end{aligned}$$

Below is a reversal inequality of the previous theorem.

Theorem 4.4. *Let $A, B \in \mathbb{C}^{n \times n}$. Then*

$$\begin{aligned} \prod_{t=1}^k \sigma_{i_t}^2(I - A^*B) &\leq \prod_{t=1}^k [1 + \lambda_{i_t}(A^*A)][1 + \lambda_t(B^*B)] - \prod_{t=1}^k \sigma_{n-t+1}^2(A + B) \\ &\quad - (2^k - 2) \prod_{t=1}^k \sigma_{i_t}(I - A^*B) \sigma_{n-t+1}(A + B). \end{aligned} \quad (4.13)$$

Proof. For $A, B \in \mathbb{C}^{n \times n}$, the following matrix identity holds (see, e.g., [23, pp. 228]):

$$I + A^*A = P + Q$$

where

$$\begin{aligned} P &= (A + B)^*(I + BB^*)^{-1}(A + B), \\ Q &= (I - A^*B)(I + B^*B)^{-1}(I - A^*B)^*. \end{aligned}$$

By Lemma 2.2, we have

$$\begin{aligned} \prod_{t=1}^k \lambda_{n-t+1}(P) &\geq \prod_{t=1}^k [1 + \lambda_t(BB^*)]^{-1} \sigma_{n-t+1}^2(A + B), \\ \prod_{t=1}^k \lambda_{i_t}(Q) &\geq \prod_{t=1}^k [1 + \lambda_t(B^*B)]^{-1} \sigma_{i_t}^2(I - A^*B). \end{aligned} \quad (4.14)$$

Using (3.3) in Theorem 3.1, we derive

$$\begin{aligned} \prod_{t=1}^k [1 + \lambda_{i_t}(A^*A)] &= \prod_{t=1}^k \lambda_{i_t}(P + Q) \\ &\geq \prod_{t=1}^k \lambda_{n-t+1}(P) + \prod_{t=1}^k \lambda_{i_t}(Q) + (2^k - 2) \prod_{t=1}^k [\lambda_{n-t+1}(P) \lambda_{i_t}(Q)]^{\frac{1}{2}} \\ &\geq \prod_{t=1}^k [1 + \lambda_t(B^*B)]^{-1} \left[\prod_{t=1}^k \sigma_{i_t}^2(I - A^*B) + \prod_{t=1}^k \sigma_{n-t+1}^2(A + B) \right. \\ &\quad \left. + (2^k - 2) \prod_{t=1}^k \sigma_{i_t}(I - A^*B) \sigma_{n-t+1}(A + B) \right]. \end{aligned}$$

Multiplying by $\prod_{t=1}^k [1 + \lambda_t(B^*B)]$, we obtain the desired inequalities (4.13). \square

As (4.3) is a reversal of Hua's (4.1), the following result, as a special case of the Theorem 4.4 by setting $k = n$, may be viewed as a counterpart of Corollary 4.1.

Corollary 4.2. *Let $A, B \in \mathbb{C}^{n \times n}$. Then*

$$\begin{aligned} |\det(I - A^*B)|^2 &\leq \det(I + A^*A) \det(I + B^*B) - |\det(A + B)|^2 \\ &\quad - (2^n - 2) |\det(I - A^*B)| |\det(A + B)|. \end{aligned}$$

Consequently,

$$|\det(I - A^*B)|^2 \leq \det(I + A^*A) \det(I + B^*B) - |\det(A + B)|^2.$$

REFERENCES

- [1] A.R. Amir-Moéz, *Extreme properties of eigenvalues of a Hermitian transformation and singular values of the sum and product of linear transformations*. Duke Math. J. **23** (1956), no. 3, 463–476. 2
- [2] T. Ando, *Positivity of operator-matrices of Hua-type*. Banach J. Math. Anal. **2** (2008), no. 2, 1–8. 6
- [3] T. Ando, *Log majorization and complementary Golden-Thompson type inequalities*. Linear Algebra Appl. 197/198 (1994), 113–131. 1
- [4] K. Fan, *On a Theorem of Weyl Concerning Eigenvalues of Linear Transformations I*. PNAS November 1, 1949, 35 (11) 652-655. 4
- [5] K. Fan, *On a Theorem of Weyl Concerning Eigenvalues of Linear Transformations II*. PNAS January 1, 1950, 36 (1) 31-35. 4
- [6] M. Fiedler, *Bounds for the determinant of the sum of hermitian matrices*. Proc. Amer. Math. Soc., **30** (1971), no. 1, 27–31. 1
- [7] G.H. Hardy, J.E. Littlewood and G. Pólya, *Inequalities*, Cambridge University Press, New York, 1934, 1st Ed.; 1952, 2nd Ed.; Reprint, 1994. 1
- [8] D.J. Hartfiel, *An extension of Haynsworths determinant inequality*. Proc. Amer. Math. Soc., **41** (1973), 41:463–465. 2
- [9] F. Hiai and Y. Lim, *Log-majorization and Lie-Trotter formula for the Cartan barycenter on probability measure spaces*. J. Math. Anal. Appl. 453 (2017), no. 1, 195–211. 1
- [10] L.-K. Hua, *Inequalities involving determinants*. Acta Math. Sinica 5 (1955), 463–470 (in Chinese). See also Transl. Amer. Math. Soc. Ser. II **32** (1963), 265–272. 6, 7
- [11] V.B. Lidskiĭ, *The proper values of the sum and product of symmetric matrices* (in Russian). Doklady Akad. Nauk SSSR vol. **75** (1950) 769–772. 2
- [12] M. Marcus, *On a determinantal inequality*. Amer. Math. Monthly **65** (1958), no. 4, 266–268. 6
- [13] A.W. Marshall, I. Olkin, and B.C. Arnold, *Inequalities: Theory of Majorization and Its Application* (Second Edition). Springer, New York, 2011. 1, 2, 4, 6
- [14] A. Oppenheim, *Inequalities connected with definite Hermitian forms II*. Amer. Math. Monthly **61** (1954), 463–466. 1, 6
- [15] C.C. Paige, G.P.H. Styan, B.-Y. Wang, and F. Zhang, *Huas matrix equality and Schur complements*. Internat. J. Inform. Syst. Sci. **4** (2008), no.1, 124–135. 6
- [16] R.C. Thompson and L.J. Freede, *On the eigenvalues of sums of Hermitian matrices*. Linear Algebra Appl. **4** (1971) 369–376. 6
- [17] B.-Y. Wang and F. Zhang, *Some inequalities for the Eigenvalues of the Product of Positive Semidefinite Hermitian Matrices*. Linear Algebra Appl. **160** (1992) 113–118. 2
- [18] H. Weyl, *Inequalities between two kinds of eigenvalues of a linear transformation*. Proc. Nat. Acad. Sci. U.S.A. 35 (1949), 408–411. 6
- [19] H. Wielandt, *An extremum property of sums of eigenvalues*. Proc. Amer. Math. Soc. **6** (1955), no. 1, 106–110.
- [20] B.Y. Xi and F. Zhang, *Inequalities for selected eigenvalues of the product of matrices*. Proc. Amer. Math. Soc. **147** (2019), no. 9, 3705–3713. <https://doi.org/10.1090/proc/14529>. 1
- [21] C.-Q. Xu, Z.-D. Xu, F. Zhang, *Revisiting Hua-Marcus-Bellman-Ando inequalities on contractive matrices*. Linear Algebra Appl. **430** (2009), no. 5-6, 1499–1508. 6
- [22] G.-H. Xu, C.-Q. Xu, and F. Zhang, *Contractive matrices of Hua type*. Linear Multilinear Algebra **59** (2011), no. 2, 159–172. 6
- [23] F. Zhang, *Matrix Theory: Basic Results and Techniques* (Second edition). Springer, New York, 2011. 1, 2, 5, 6, 7, 8

(B.-Y. Xi) COLLEGE OF MATHEMATICS, INNER MONGOLIA UNIVERSITY FOR NATIONALITIES, TONGLIAO CITY, INNER MONGOLIA AUTONOMOUS REGION, 028043, CHINA
Email address: baoyintu78@qq.com, baoyintu78@imn.edu.cn

(F. Zhang) DEPARTMENT OF MATHEMATICS, NOVA SOUTHEASTERN UNIVERSITY, 3301 COLLEGE AVE., FORT LAUDERDALE, FL 33314, USA
Email address: zhang@nova.edu