

Two-moment characterization of spectral measures on the real line

Paweł Pietrzycki and Jan Stochel

ABSTRACT. In [30], Kiukas, Lahti and Ylinen asked the following general question. *When is a positive operator measure projection valued?* A version of this question formulated in terms of operator moments was posed in [43]. *Let T be a selfadjoint operator and F be a Borel semispectral measure on the real line with compact support. For which positive integers $p < q$ do the equalities $T^k = \int_{\mathbb{R}} x^k F(dx)$, $k = p, q$, imply that F is a spectral measure?* In the present paper, we completely solve the second problem. The answer is affirmative if p is odd and q is even, and negative otherwise. The case $(p, q) = (1, 2)$ closely related to intrinsic noise operator was solved by several authors including Kruszyński and de Muynck as well as Kiukas, Lahti and Ylinen. The counterpart of the second problem concerning the multiplicativity of unital positive linear maps on C^* -algebras is also solved.

1. Introduction

One of the most important concepts in mathematics and physics is the notion of a normalized positive operator valued measure also known as a probability operator valued measure or a generalized observable, or else semispectral measure. This concept was introduced in the 1940s by Naimark (see [38, 37, 39]). Positive operator valued measures play a significant role in operator theory [7, 1, 25, 43] and are a standard tool in quantum information theory and quantum optics [12, 53, 23, 13]. Recall that a map $F: \mathcal{A} \rightarrow \mathcal{B}(\mathcal{H})$ defined on a σ -algebra \mathcal{A} of subsets of a set X is said to be:

- a *positive operator valued measure (POV measure)* if $\langle F(\cdot)h, h \rangle$ is a positive measure for every $h \in \mathcal{H}$,
- a *semispectral measure* if F is a POV measure such that $F(X) = I$,
- a *spectral measure* if F is a semispectral measure such that $F(\Delta)$ is an orthogonal projection for every $\Delta \in \mathcal{A}$,

where $\mathcal{B}(\mathcal{H})$ is the collection of all bounded linear operators on a Hilbert space \mathcal{H} and I is the identity operator on \mathcal{H} . The celebrated Naimark's dilation theorem (see [39] and [34, Theorem 6.4]) states that a POV measure $F: \mathcal{A} \rightarrow \mathcal{B}(\mathcal{H})$ can always be represented as the R -compression $R^*E(\cdot)R$ of a spectral measure $E: \mathcal{A} \rightarrow \mathcal{B}(\mathcal{K})$,

2020 *Mathematics Subject Classification.* Primary 47B15, 44A60; Secondary 47A63, 46G10, 46L05.

Key words and phrases. Semispectral measure, spectral measure, operator moment, positive linear map, multiplicativity, operator monotone function.

where \mathcal{K} is a Hilbert space and R is a bounded linear operator from \mathcal{H} to \mathcal{K} . By [34, p. 14], \mathcal{K} can be made minimal in the sense that $\mathcal{K} = \bigvee \{E(\Delta)R(\mathcal{H}) : \Delta \in \mathcal{A}\}$. If F is semispectral, then \mathcal{H} is a subspace of \mathcal{K} and R is the (isometric) embedding of \mathcal{H} into \mathcal{K} , and so the minimality condition takes the form $\mathcal{K} = \bigvee \{E(\Delta)\mathcal{H} : \Delta \in \mathcal{A}\}$.

It turns out that, from a mathematical and physical point of view, it is important to investigate the relationship between semispectral and spectral measures. In the classical von Neumann description of quantum mechanics selfadjoint operators or, equivalently, Borel spectral measures on the real line represent observables. This approach is insufficient in describing many natural properties of measurements, such as measurement inaccuracy. Therefore, in standard modern quantum theory, the generalization to semispectral measures is widely used. In particular, this is the case in quantum information theory and in quantum optics (to represent measurement statistics). Among the papers undertaking this line of research, the following are noteworthy [37, 22, 31, 14, 3, 4, 30, 26, 5, 6].

By a *Borel POV measure on \mathbb{R}* we mean a POV measure $F : \mathfrak{B}(\mathbb{R}) \rightarrow \mathcal{B}(\mathcal{H})$, where $\mathfrak{B}(\mathbb{R})$ stands for the σ -algebra of all Borel subsets of the real line \mathbb{R} (below, the algebra $\mathcal{B}(\mathcal{H})$ will not be explicitly mentioned unless necessary). For an integer $n \geq 1$ and a Borel POV measure F on \mathbb{R} with compact support¹, the integral

$$\int_{\mathbb{R}} x^n F(dx)$$

is a (bounded) self-adjoint operator, which is called the *n th operator moment* of F . A straightforward application of the Weierstrass approximation theorem shows that a Borel POV measure on \mathbb{R} with compact support is uniquely determined by its operator moments. One of the features of a Borel spectral measures on \mathbb{R} is the multiplicativity of the corresponding Stone-von Neumann functional calculus. In particular, if E is a Borel spectral measure on \mathbb{R} with compact support, then the following identities hold²

$$\left(\int_{\mathbb{R}} x E(dx) \right)^n = \int_{\mathbb{R}} x^n E(dx), \quad n = 1, 2, \dots \quad (1.1)$$

Hence, all operator moments of E are determined by the first one, and according to the spectral theorem there is a one-to-one correspondence between Borel spectral measures on \mathbb{R} and their first operator moments. This is no longer true for general Borel semispectral measures on \mathbb{R} . It turns out, however, that the single equality in (1.1) with $n = 2$ guarantees spectrality.

THEOREM 1.1 ([31, Proposition 1], [30, Theorem 5] and [43, Remark 5.3]). *A Borel semispectral measure F on \mathbb{R} with compact support³ is spectral if and only if*

$$\left(\int_{\mathbb{R}} x F(dx) \right)^2 = \int_{\mathbb{R}} x^2 F(dx).$$

¹For more information on closed supports of spectral and semispectral measures, see [48, p. 69] and [25, p. 1799].

²The identity (1.1) holds even if the closed support of E is not compact. Since we only deal with bounded operators in this paper, the POV measures considered have compact supports (see Section 7 for more explanation).

³The first two references contain versions of this result for semispectral measures with non-compact supports.

It is worth mentioning that if F is a Borel semispectral measure on \mathbb{R} with compact support, then the operator $\text{Var}(F)$, called *intrinsic noise operator* (see [13, p. 177]), defined by

$$\text{Var}(F) = \int_{\mathbb{R}} x^2 F(dx) - \left(\int_{\mathbb{R}} x F(dx) \right)^2 \quad (1.2)$$

is always positive (see Corollary 3.4; this can also be deduced from the Kadison inequality (2.4)). Thus, according to Theorem 1.1, equality holds in $\text{Var}(F) \geq 0$ only for spectral measures.

In this connection, it is worth emphasizing that Theorem 1.1 was developed for the needs of quantum physics. Namely, the main purpose of the quantization proposed in [29, 30] was to construct observables that are not spectral measures, and this was done by using the operator moments of these observables. To achieve this goal, it was important to be able to use these moments to determine whether a given observable is or is not a spectral measure. This led Kiukas, Lahti and Ylinen to the following question (see [30, Sec. VI], see also [28, Sec. 5]):

QUESTION 1.2. *When is a positive operator measure projection valued?*

In a recent paper [43], we gave a solution to [16, Problem 1.1] concerning subnormal square roots of quasinormal operators. In fact, the paper [43] provides two solutions to this problem that use two different approaches. The first one appeals to the theory of operator monotone functions, in particular Hansen's inequality. The second is based on the technique that utilizes operator moments of semispectral measures. A detailed analysis of both solutions led us to a new criterion for the spectrality of a Borel semispectral measure on \mathbb{R} compactly supported in $[0, \infty)$, written in terms of its two operator moments. This criterion was used to solve a generalization of [16, Problem 1.1] (see [43, Theorem 4.1]).

THEOREM 1.3 ([43, Theorem 4.2], [44]). *Let $T \in \mathcal{B}(\mathcal{H})$ be a positive operator and α, β be two distinct positive real numbers. Assume that $F: \mathfrak{B}(\mathbb{R}) \rightarrow \mathcal{B}(\mathcal{H})$ is a semispectral measure compactly supported in $[0, \infty)$. Then the following conditions are equivalent:*

- (i) F is the spectral measure of T ,
- (ii) $T^n = \int_{[0, \infty)} x^n F(dx)$ for all integers $n \geq 0$,
- (iii) $T^r = \int_{[0, \infty)} x^r F(dx)$ for all $r \in [0, \infty)$,
- (iv) $T^r = \int_{[0, \infty)} x^r F(dx)$ for $r = \alpha, \beta$.

As shown in the proof of [43, Theorem 4.2], the implication (iv) \Rightarrow (i) is equivalent to the fact that a semispectral measure $F: \mathfrak{B}(\mathbb{R}) \rightarrow \mathcal{B}(\mathcal{H})$ compactly supported in $[0, \infty)$ for which there exists $s \in (0, \infty) \setminus \{1\}$ such that

$$\left(\int_{[0, \infty)} x F(dx) \right)^s = \int_{[0, \infty)} x^s F(dx)$$

is spectral (see [43, Lemma 4.3]).

In view of Question 1.2 and Theorems 1.1 and 1.3, it seems natural to pose the following general problem in which Ξ is a fixed nonempty set of positive integers (if Ξ is finite, then we always order its elements in a non-decreasing manner). Problem 1.4 below can be regarded as a generalization of [43, Problem 5.2] which deals with two-element sets Ξ .

PROBLEM 1.4. Let $T \in \mathbf{B}(\mathcal{H})$ be a selfadjoint operator and $F: \mathfrak{B}(\mathbb{R}) \rightarrow \mathbf{B}(\mathcal{H})$ be a semispectral measure with compact support. Does the system of equations

$$T^k = \int_{\mathbb{R}} x^k F(dx), \quad k \in \Xi, \quad (1.3)$$

imply that F is spectral?

This problem can be rephrased equivalently in terms of dilation theory as follows (use Naimark's dilation theorem and Lemma 3.2):

PROBLEM 1.5. Let $T \in \mathbf{B}(\mathcal{H})$ be a selfadjoint operator, $F: \mathfrak{B}(\mathbb{R}) \rightarrow \mathbf{B}(\mathcal{H})$ be a semispectral measure with compact support, $E: \mathfrak{B}(\mathbb{R}) \rightarrow \mathbf{B}(\mathcal{K})$ be a minimal spectral dilation of F (i.e., E is a spectral measure satisfying (3.2) and (3.4)) and S be the first operator moment of E (i.e., $S = \int_{\mathbb{R}} x E(dx)$). Does the system of equations

$$T^k = PS^k|_{\mathcal{H}}, \quad k \in \Xi,$$

imply that P and S commutes?

It turns out that Problem 1.4 is closely related to the question of multiplicativity of unital positive linear maps on C^* -algebras (see Remark 4.2). In fact, the two problems are logically equivalent regardless of the cardinality of the set Ξ (see Remark 4.3). The C^* -algebra counterpart of Problem 1.4 takes the following form.

PROBLEM 1.6. Let \mathcal{A} and \mathcal{B} be unital C^* -algebras, $\Phi: \mathcal{A} \rightarrow \mathcal{B}$ be a unital positive linear map and $a \in \mathcal{A}$ and $b \in \mathcal{B}$ be selfadjoint. Does the system of equations

$$b^k = \Phi(a^k), \quad k \in \Xi,$$

imply that Φ restricted to the unital subalgebra generated by $\{a\}$ is multiplicative?

The correspondence between Problems 1.4 and 1.5 allows us to use the theory of operator monotone functions and related operator inequalities to prove the main results of this paper, which provide complete solutions to Problems 1.4, 1.5 and 1.6 for two-element sets Ξ . We begin with the affirmative solutions.

THEOREM 1.7. Let $T \in \mathbf{B}(\mathcal{H})$ be a selfadjoint operator, $F: \mathfrak{B}(\mathbb{R}) \rightarrow \mathbf{B}(\mathcal{H})$ be a semispectral measure with compact support and p, q be positive integers such that $p < q$, p is odd and q is even. Then the following conditions are equivalent:

- (i) F is the spectral measure of T ,
- (ii) $T^k = \int_{\mathbb{R}} x^k F(dx)$ for all integers $k \geq 0$,
- (iii) $T^k = \int_{\mathbb{R}} x^k F(dx)$ for $k = p, q$.

The affirmative solution to Problem 1.6 takes the form.

THEOREM 1.8. Let \mathcal{A} and \mathcal{B} be unital C^* -algebras, $\Phi: \mathcal{A} \rightarrow \mathcal{B}$ be a unital positive linear map, $a \in \mathcal{A}$ be selfadjoint and p, q be positive integers such that $p < q$, p is odd and q is even. Then the following conditions are equivalent:

- (i) Φ restricted to the unital subalgebra generated by $\{a\}$ is multiplicative,
- (ii) there exists a selfadjoint element $b \in \mathcal{B}$ such that $b^k = \Phi(a^k)$ for $k = p, q$.

Moreover, if (ii) holds, then $b = \Phi(a)$.

In the complementary result, we show that the set (with $\mathbb{N} = \{1, 2, 3, \dots\}$)

$$\Omega := \{(p, q) \in \mathbb{N}^2 : p < q, p \text{ odd and } q \text{ even}\} \quad (1.4)$$

is the largest possible subset of $\{(p, q) \in \mathbb{N}^2 : p \leq q\}$ for which Problem 1.4 has an affirmative solution for $\Xi = \{p, q\}$. Surprisingly, suitable counterexamples can be constructed even when the underlying Hilbert space \mathcal{H} is one-dimensional (see Theorem 5.2 for more details).

THEOREM 1.9. *Let $(p, q) \in \mathbb{N}^2 \setminus \Omega$ be such that $p \leq q$. Then there exist a Hilbert space \mathcal{H} , a selfadjoint operator $T \in \mathbf{B}(\mathcal{H})$ and a semispectral measure $F: \mathfrak{B}(\mathbb{R}) \rightarrow \mathbf{B}(\mathcal{H})$ with compact support which is not spectral and such that*

$$T^k = \int_{\mathbb{R}} x^k F(dx), \quad k = p, q.$$

The proofs of Theorems 1.7, 1.8 and 1.9 will be given in Sections 3, 4 and 5, respectively. In Section 2 we provide the basic facts on operator monotone functions and the related operator inequalities needed in this paper. Section 6 contains additional counterexamples (including the case of infinite dimensional spaces) related to the Fibonacci sequence. Finally, in Section 7 we discuss the possibility of adapting the two-moment characterizations of spectral measures given in Theorems 1.3 and 1.7 to the case of semispectral measures whose closed supports are not compact.

2. Prerequisites

In this paper, we use the following notation. The fields of real and complex numbers are denoted by \mathbb{R} and \mathbb{C} , respectively. The symbols \mathbb{Z}_+ , \mathbb{N} and \mathbb{R}_+ stand for the sets of nonnegative integers, positive integers and nonnegative real numbers, respectively. We write $\mathfrak{B}(X)$ for the σ -algebra of all Borel subsets of a topological Hausdorff space X . The C^* -algebra of all continuous complex functions on a compact Hausdorff space K equipped with supremum norm is denoted by $C(K)$. For $\lambda \in \mathbb{R}$, δ_λ stands for the Borel probability measure on \mathbb{R} concentrated on $\{\lambda\}$.

Let \mathcal{H} and \mathcal{K} be (complex) Hilbert spaces. Denote by $\mathbf{B}(\mathcal{H}, \mathcal{K})$ the Banach space of all bounded linear operators from \mathcal{H} to \mathcal{K} . If $A \in \mathbf{B}(\mathcal{H}, \mathcal{K})$, then A^* , $\mathcal{N}(A)$ and $\mathcal{R}(A)$ stand for the adjoint, the kernel and the range of A , respectively. It is well known that $\mathbf{B}(\mathcal{H}) := \mathbf{B}(\mathcal{H}, \mathcal{H})$ is a C^* -algebra with unit I , where $I = I_{\mathcal{H}}$ denotes the identity operator on \mathcal{H} . We say that $A \in \mathbf{B}(\mathcal{H})$ is *selfadjoint* if $A = A^*$, *positive* if $\langle Ah, h \rangle \geq 0$ for all $h \in \mathcal{H}$ and an *orthogonal projection* if $A = A^*$ and $A = A^2$.

Let \mathcal{A} be a σ -algebra of subsets of a set X and let $F: \mathcal{A} \rightarrow \mathbf{B}(\mathcal{H})$ be a semispectral measure. Denote by $L^1(F)$ the vector space of all \mathcal{A} -measurable functions $f: X \rightarrow \mathbb{C}$ such that $\int_X |f(x)| \langle F(dx)h, h \rangle < \infty$ for all $h \in \mathcal{H}$. Then for every $f \in L^1(F)$, there exists a unique operator $\int_X f dF \in \mathbf{B}(\mathcal{H})$ such that (see e.g., [51, Appendix])

$$\left\langle \int_X f dF h, h \right\rangle = \int_X f(x) \langle F(dx)h, h \rangle, \quad h \in \mathcal{H}. \quad (2.1)$$

If F is a spectral measure, then $\int_X f dF$ coincides with the usual spectral integral. In particular, if F is the spectral measure of a selfadjoint operator $A \in \mathbf{B}(\mathcal{H})$, then we write $f(A) = \int_{\mathbb{R}} f dF$ for any F -essentially bounded Borel function $f: \mathbb{R} \rightarrow \mathbb{R}$; the map $f \mapsto f(A)$ is called the Stone-von Neumann functional calculus. For more information needed in this article on spectral integrals, including the spectral theorem for selfadjoint operators and the Stone-von Neumann functional calculus, we refer the reader to [46, 54, 48].

Let $J \subseteq \mathbb{R}$ be an interval (which may be open, half-open, or closed; bounded or unbounded). A continuous function $f: J \rightarrow \mathbb{R}$ is said to be *operator monotone* if

$f(A) \leq f(B)$ for any two selfadjoint operators $A, B \in \mathbf{B}(\mathcal{H})$ such that $A \leq B$ and the spectra of A and B are contained in J . In [33], Löwner proved that a continuous function defined on an open interval is operator monotone if and only if it has an analytic continuation to the complex upper half-plane which is a Pick function (see also [19, 21]). Operator monotone functions have integral representations with respect to suitable positive Borel measures. In particular, a continuous function $f: (0, \infty) \rightarrow \mathbb{R}$ is operator monotone if and only if there exists a positive Borel measure ν on $[0, \infty)$ such that $\int_0^\infty \frac{1}{1+\lambda^2} d\nu(\lambda) < \infty$ and

$$f(t) = \alpha + \beta t + \int_0^\infty \left(\frac{\lambda}{1+\lambda^2} - \frac{1}{t+\lambda} \right) d\nu(\lambda), \quad t \in (0, \infty),$$

where $\alpha \in \mathbb{R}$ and $\beta \in \mathbb{R}_+$ (see [21, Theorem 5.2] or [9, p. 144]). The most important example of an operator monotone function is $f: [0, \infty) \ni t \rightarrow t^p \in \mathbb{R}$ for $p \in (0, 1)$. This function has the following integral representation (see [9, Exercise V.1.10(iii)] or [9, Exercise V.4.20])

$$t^p = \frac{\sin p\pi}{\pi} \int_0^\infty \frac{t\lambda^{p-1}}{t+\lambda} d\lambda, \quad t \in [0, \infty). \quad (2.2)$$

Operator monotone functions are related to the Hansen inequality [20]. In [52, Lemma 2.2], Uchiyama gave a necessary and sufficient condition for equality to hold in the Hansen inequality when the external factor is a nontrivial orthogonal projection (see the “moreover” part of Theorem 2.1 below; see also the paragraph before [43, Theorem 2.4] showing why the separability of \mathcal{H} can be dropped).

THEOREM 2.1 ([20, 52]). *Let $A \in \mathbf{B}(\mathcal{H})$ be a positive operator, $T \in \mathbf{B}(\mathcal{H})$ be a contraction and $f: [0, \infty) \rightarrow \mathbb{R}$ be a continuous operator monotone function such that $f(0) \geq 0$. Then*

$$T^* f(A) T \leq f(T^* A T). \quad (2.3)$$

Moreover, if f is not an affine function and T is an orthogonal projection such that $T \neq I$, then equality holds in (2.3) if and only if $TA = AT$ and $f(0) = 0$.

The reader is referred to [33, 19, 20, 9, 21, 49] for the fundamentals of the theory of operator monotone functions.

A linear map $\Phi: \mathcal{A} \rightarrow \mathcal{B}$ between unital C^* -algebras is said to be *positive* if $\Phi(a) \geq 0$ for every $a \in \mathcal{A}$ such that $a \geq 0$. The map Φ is called *unital* if it preserves the units. If Φ is positive and unital, then the following inequality, called Kadison’s inequality (see [27]), holds:

$$\Phi(a^2) \geq \Phi(a)^2 \text{ for all } a \in \mathcal{A} \text{ such that } a = a^*. \quad (2.4)$$

In this paper we will need the following generalization of Kadison’s inequality.

THEOREM 2.2 ([32, Theorem 2]). *Let $R \in \mathbf{B}(\mathcal{H}, \mathcal{K})$ and let $\Phi: \mathbf{B}(\mathcal{K}) \rightarrow \mathbf{B}(\mathcal{H})$ be the positive linear map defined by*

$$\Phi(X) = R^* X R, \quad X \in \mathbf{B}(\mathcal{K}).$$

Then for all $A, B \in \mathbf{B}(\mathcal{K})$, the net $\{\Phi(A^ B)(\Phi(B^* B) + \varepsilon I)^{-1} \Phi(B^* A)\}_{\varepsilon > 0}$ is convergent in the strong operator topology as $\varepsilon \downarrow 0$ and*

$$\Phi(A^* A) \geq (\text{SOT}) \lim_{\varepsilon \downarrow 0} \Phi(A^* B)(\Phi(B^* B) + \varepsilon I)^{-1} \Phi(B^* A).$$

3. Proof of Theorem 1.7

We begin with the following lemma which gives a necessary and sufficient condition for equality to hold in a Kadison type inequality (cf. (2.4)). Although this is a known fact even for unbounded operators (see [31, Lemmas 1 and 2]), we will provide a brief algebraic proof for the reader's convenience. Note also that part (iii) of Lemma 3.1 below is [18, Lemma in Sec. 6].

LEMMA 3.1. *Let $T \in \mathbf{B}(\mathcal{H})$ be a selfadjoint operator and $P \in \mathbf{B}(\mathcal{H})$ be an orthogonal projection. Then the following statements are valid:*

- (i) $(PTP)^2 \leq PT^2P$,
- (ii) *equality holds in (i) if and only if $PT = TP$,*
- (iii) *if T is an orthogonal projection, then PTP is an orthogonal projection if and only if $PT = TP$.*

PROOF. (i) This is a direct consequence of the following algebraic identities:

$$\begin{aligned} PT^2P - (PTP)^2 &= PT^2P - PTPTP \\ &= PT(I - P)TP \\ &= (TP)^*(I - P)TP \geq 0. \end{aligned} \quad (3.1)$$

(ii) It follows from (3.1) that equality holds in (i) if and only if

$$(TP)^*(I - P)TP = 0,$$

or equivalently if and only if

$$\mathcal{R}(TP) \subseteq \mathcal{N}((I - P)^{\frac{1}{2}}) = \mathcal{N}(I - P),$$

which in turn is equivalent to $(I - P)TP = 0$. The last equality holds if and only if $TP = PTP$, which by $(PTP)^* = PTP$ is equivalent to $PT = TP$.

(iii) This is a direct consequence of (ii) because PTP is an orthogonal projection if and only if $(PTP)^2 = PT^2P$. \square

For our further considerations, the following fact is fundamental. In particular, in view of Naimark's dilation theorem (see Introduction), it shows that Problems 1.4 and 1.5 are logically equivalent.

LEMMA 3.2. *Let \mathcal{H}, \mathcal{K} be Hilbert spaces such that $\mathcal{H} \subseteq \mathcal{K}$ and $P \in \mathbf{B}(\mathcal{K})$ be the orthogonal projection of \mathcal{K} onto \mathcal{H} . Suppose that $F: \mathfrak{B}(\mathbb{R}) \rightarrow \mathbf{B}(\mathcal{H})$ is a semispectral measure and $E: \mathfrak{B}(\mathbb{R}) \rightarrow \mathbf{B}(\mathcal{K})$ is a spectral measure such that*

$$F(\Delta) = PE(\Delta)|_{\mathcal{H}}, \quad \Delta \in \mathfrak{B}(\mathbb{R}). \quad (3.2)$$

Set⁴ $S := \int_{\mathbb{R}} xE(dx)$. Then the following statements are valid:

- (i) *F is spectral if and only if P commutes with E (equivalently, \mathcal{H} reduces E),*
- (ii) *if $S \in \mathbf{B}(\mathcal{K})$, then F has compact support and*

$$PS^k|_{\mathcal{H}} = \int_{\mathbb{R}} x^k F(dx), \quad k \in \mathbb{Z}_+, \quad (3.3)$$

- (iii) *if F has compact support and \mathcal{K} is minimal, that is,*

$$\mathcal{K} = \bigvee \{E(\Delta)\mathcal{H} : \Delta \in \mathcal{A}\}, \quad (3.4)$$

then E has compact support, $S \in \mathbf{B}(\mathcal{K})$ and $S = S^$.*

⁴Note that a priori the operator S may be unbounded (see [48, Theorem 5.9] for more details).

PROOF. (i) Set $\hat{F}(\Delta) = F(\Delta) \oplus 0$ for $\Delta \in \mathfrak{B}(\mathbb{R})$, where 0 stands for the zero operator on $\mathcal{K} \ominus \mathcal{H}$. Then, by (3.2), $\hat{F}(\Delta) = PE(\Delta)P$. Hence, observing that $F(\Delta)$ is an orthogonal projection if and only if $\hat{F}(\Delta)$ is an orthogonal projection and using Lemma 3.1(iii), we obtain (i).

(ii) It follows from (3.2) that the closed support of F is contained in the closed support of E . Since E has compact support (because $S \in \mathbf{B}(\mathcal{K})$, see [48, Theorem 5.9]), so does F . Applying the Stone-von Neumann functional calculus, we get

$$\begin{aligned} \langle PS^k|_{\mathcal{H}}h, h \rangle &= \langle S^k h, h \rangle = \int_{\mathbb{R}} x^k \langle E(dx)h, h \rangle \\ &\stackrel{(3.2)}{=} \int_{\mathbb{R}} x^k \langle F(dx)h, h \rangle \\ &\stackrel{(2.1)}{=} \left\langle \int_{\mathbb{R}} x^k F(dx)h, h \right\rangle, \quad h \in \mathcal{H}, k \in \mathbb{Z}_+, \end{aligned}$$

which implies (3.3).

(iii) By (3.2) and (3.4), the closed supports of the POV measures E and F coincide (see the proofs of [24, Theorem 4.4] and [25, Proposition 4(iii)]). Hence, the closed support of E is compact. As a consequence, the operator $\int_{\mathbb{R}} xE(dx)$ is bounded and selfadjoint (see [48, Theorem 5.9]). This completes the proof. \square

We are now in a position to prove the main result of this paper, which provides a two-moment characterization of spectral measures.

PROOF OF THEOREM 1.7. (i) \Rightarrow (ii) This is immediate from the Stone-von Neumann functional calculus.

(ii) \Rightarrow (iii) Obvious.

(iii) \Rightarrow (i) It follows from Naimark's dilation theorem (see Introduction) that there exist a Hilbert space \mathcal{K} containing \mathcal{H} and a spectral measure $E: \mathfrak{B}(\mathbb{R}) \rightarrow \mathbf{B}(\mathcal{K})$ such that (3.2) and (3.4) hold, where $P \in \mathbf{B}(\mathcal{K})$ is the orthogonal projection of \mathcal{K} onto \mathcal{H} . By Lemma 3.2, E has compact support, the operator $S := \int_{\mathbb{R}} xE(dx)$ is bounded and selfadjoint, and the following equalities are satisfied:

$$T^k = PS^k|_{\mathcal{H}}, \quad k = p, q. \quad (3.5)$$

First, we prove that F is a spectral measure. In view of Lemma 3.2(i), it suffices to show that P commutes with E . For this, we consider two cases.

CASE 1. $p \leq \frac{q}{2}$.

Let $\hat{T} \in \mathbf{B}(\mathcal{K})$ be defined by $\hat{T} = T \oplus 0$, where 0 stands for the zero operator on $\mathcal{K} \ominus \mathcal{H}$. Set $q' = \frac{q}{2}$. Using Lemma 3.1(i) and then applying Theorem 2.1 to the positive operator $S^{2q'}$ and the operator monotone function $f(t) = t^{\frac{p}{q'}}$ (see (2.2)), we deduce that

$$\begin{aligned} \hat{T}^{2p} &= (\hat{T}^p)^2 \stackrel{(3.5)}{=} (PS^pP)^2 \\ &\leq PS^{2p}P \\ &\leq (PS^{2q'}P)^{\frac{p}{q'}} = (PS^qP)^{\frac{2p}{q}} \stackrel{(3.5)}{=} (\hat{T}^q)^{\frac{2p}{q}} \stackrel{(*)}{=} \hat{T}^{2p}, \end{aligned}$$

where $(*)$ can be inferred from the hypothesis that q is even. This implies that

$$(PS^pP)^2 = PS^{2p}P.$$

It follows from Lemma 3.1(ii) that

$$PS^p = S^pP.$$

Hence, by [48, Theorem 5.1], P commutes with E_p , the spectral measure of S^p . By [10, Theorem 6.6.4], E_p is of the form

$$E_p(\Delta) = E(\varphi_p^{-1}(\Delta)), \quad \Delta \in \mathfrak{B}(\mathbb{R}), \quad (3.6)$$

where $\varphi_p: \mathbb{R} \rightarrow \mathbb{R}$ is a function given by

$$\varphi_p(x) = x^p, \quad x \in \mathbb{R}. \quad (3.7)$$

Since the map $\mathfrak{B}(\mathbb{R}) \ni \Delta \mapsto \varphi_p^{-1}(\Delta) \in \mathfrak{B}(\mathbb{R})$ is surjective (because p is odd), we deduce from (3.6) that P commutes with E .

CASE 2. $p > \frac{q}{2}$.

Suppose, to the contrary, that P does not commute with E . This implies that $P \neq I_{\mathcal{K}}$. Set $q' = \frac{q}{2}$ and $r = p - q'$. Since $p < q$ and q is even, we see that $r, q' \in \mathbb{N}$ and $0 < \frac{r}{q'} < 1$. By Theorem 2.1 applied to the positive operator $S^{2q'}$ and the operator monotone function $f(t) = t^{\frac{r}{q'}}$, we get

$$\hat{T}^{2r} = (\hat{T}^{2q'})^{\frac{r}{q'}} \stackrel{(3.5)}{=} (PS^{2q'}P)^{\frac{r}{q'}} \geq PS^{2r}P.$$

This implies that

$$T^{2r} = (PS^{2q'}|_{\mathcal{H}})^{\frac{r}{q'}} \geq PS^{2r}|_{\mathcal{H}}. \quad (3.8)$$

Let $\Phi: \mathbf{B}(\mathcal{K}) \rightarrow \mathbf{B}(\mathcal{H})$ be the positive unital linear map defined by

$$\Phi(X) = PX|_{\mathcal{H}}, \quad X \in \mathbf{B}(\mathcal{K}).$$

Applying Theorem 2.2 to $A = S^r$, $B = S^{q'}$ and $R \in \mathbf{B}(\mathcal{H}, \mathcal{K})$ defined by $Rh = h$ for $h \in \mathcal{H}$ leads to

$$PS^{2r}|_{\mathcal{H}} = \Phi(S^{2r}) \geq (\text{sot})\lim_{\varepsilon \downarrow 0} \Phi(S^p)(\Phi(S^q) + \varepsilon I)^{-1} \Phi(S^p). \quad (3.9)$$

Let $G: \mathfrak{B}(\mathbb{R}) \rightarrow \mathbf{B}(\mathcal{H})$ be the spectral measure of T . Using the Stone-von Neumann functional calculus, we obtain

$$\Phi(S^p)(\Phi(S^q) + \varepsilon I)^{-1} \Phi(S^p) \stackrel{(3.5)}{=} T^p(T^q + \varepsilon)^{-1}T^p = \int_{\mathbb{R}} \frac{x^{2p}}{x^q + \varepsilon} G(dx). \quad (3.10)$$

Applying Lebesgue's monotone convergence theorem and the hypothesis that q is even and $2p - q \in \mathbb{N}$, we deduce that

$$\begin{aligned} \lim_{\varepsilon \downarrow 0} \langle \Phi(S^p)(\Phi(S^q) + \varepsilon I)^{-1} \Phi(S^p)h, h \rangle &\stackrel{(3.10)}{=} \lim_{\varepsilon \downarrow 0} \left\langle \int_{\mathbb{R}} \frac{x^{2p}}{x^q + \varepsilon} G(dx)h, h \right\rangle \\ &= \lim_{\varepsilon \downarrow 0} \int_{\mathbb{R}} \frac{x^{2p}}{x^q + \varepsilon} \langle G(dx)h, h \rangle \\ &= \int_{\mathbb{R}} x^{2p-q} \langle G(dx)h, h \rangle \\ &= \langle T^{2p-q}h, h \rangle = \langle T^{2r}h, h \rangle, \quad h \in \mathcal{H}. \end{aligned}$$

Therefore, the net $\{\Phi(S^p)(\Phi(S^q) + \varepsilon I)^{-1} \Phi(S^p)\}_{\varepsilon > 0}$ is convergent to T^{2r} in the weak operator topology. Combined with (3.9), this implies that

$$PS^{2r}|_{\mathcal{H}} \geq T^{2r}. \quad (3.11)$$

Using (3.8) and (3.11), we get

$$T^{2r} = (PS^{2q'}|_{\mathcal{H}})^{\frac{r}{q'}} \geq PS^{2r}|_{\mathcal{H}} \geq T^{2r}.$$

This yields

$$(PS^{2q'}|_{\mathcal{H}})^{\frac{r}{q'}} = PS^{2r}|_{\mathcal{H}},$$

or equivalently

$$(PS^{2q'}P)^{\frac{r}{q'}} = PS^{2r}P,$$

so equality holds in the Hansen inequality. Thus, by the moreover part of Theorem 2.1, $PS^q = S^qP$ (recall that $q = 2q'$). Hence

$$\hat{T}^{qn} \stackrel{(3.5)}{=} (PS^qP)^n = (PS^q)^n = PS^{qn}P, \quad n \in \mathbb{N}.$$

Therefore, $T^{qn} = PS^{qn}|_{\mathcal{H}}$ for all $n \in \mathbb{N}$. Take any $n_0 \in \mathbb{N}$ such that $p \leq \frac{qn_0}{2}$. Then by (3.5), we have

$$T^k = PS^k|_{\mathcal{H}}, \quad k = p, qn_0.$$

Since $p \leq \frac{qn_0}{2}$, we can apply Case 1 to the pair (p, qn_0) in place of (p, q) . We then obtain that P commutes with E , which is a contradiction.

Summarizing, we have proved that in both cases F is a spectral measure. Therefore, to complete the proof it remains to show that F is the spectral measure of T . Since $T^p = \int_{\mathbb{R}} x^p F(dx)$ (by (iii)) and $T^p = \int_{\mathbb{R}} x^p G(dx)$ (by Stone-von Neumann functional calculus), an application of [10, Theorem 6.6.4] shows that $F \circ \varphi_p^{-1}$ and $G \circ \varphi_p^{-1}$ are spectral measures of T^p , where G is the spectral measure of T , φ_p is as in (3.7) and

$$(F \circ \varphi_p^{-1})(\Delta) = F(\varphi_p^{-1}(\Delta)) \text{ and } (G \circ \varphi_p^{-1})(\Delta) = G(\varphi_p^{-1}(\Delta)) \text{ for } \Delta \in \mathfrak{B}(\mathbb{R}).$$

By the uniqueness in [10, Theorem 6.1.1], $F \circ \varphi_p^{-1} = G \circ \varphi_p^{-1}$. Since the map $\mathfrak{B}(\mathbb{R}) \ni \Delta \mapsto \varphi_p^{-1}(\Delta) \in \mathfrak{B}(\mathbb{R})$ is surjective (because p is odd), we deduce that $F = G$, so F is the spectral measure of T . This completes the proof. \square

We conclude this section by providing some inequalities for moments of a semispectral measure on the real line. Though it is a well-known fact (see [11] and references therein), we outline its short proof for the reader's convenience.

PROPOSITION 3.3. *Let $F: \mathfrak{B}(\mathbb{R}) \rightarrow \mathcal{B}(\mathcal{H})$ be a semispectral measure with compact support. Then*

$$\begin{bmatrix} I & \int_{\mathbb{R}} x F(dx) & \cdots & \int_{\mathbb{R}} x^n F(dx) \\ \int_{\mathbb{R}} x F(dx) & \int_{\mathbb{R}} x^2 F(dx) & \cdots & \int_{\mathbb{R}} x^{n+1} F(dx) \\ \vdots & \vdots & \ddots & \vdots \\ \int_{\mathbb{R}} x^n F(dx) & \int_{\mathbb{R}} x^{n+1} F(dx) & \cdots & \int_{\mathbb{R}} x^{2n} F(dx) \end{bmatrix} \geq 0, \quad n \in \mathbb{Z}_+.$$

PROOF. By Naimark's dilation theorem (see Introduction), there exists a Hilbert space \mathcal{K} containing \mathcal{H} and a spectral measure $E: \mathfrak{B}(\mathbb{R}) \rightarrow \mathcal{B}(\mathcal{K})$ which satisfies

(3.2) and (3.4). By Lemma 3.2(iii), E has compact support. Applying the Stone-von Neumann functional calculus, we obtain

$$\begin{aligned} \sum_{j,k=0}^n \left\langle \int_{\mathbb{R}} x^{j+k} F(dx) h_k, h_j \right\rangle &= \sum_{j,k=0}^n \left\langle \int_{\mathbb{R}} x^{j+k} E(dx) h_k, h_j \right\rangle \\ &= \left\| \sum_{k=0}^n \int_{\mathbb{R}} x^k E(dx) h_k \right\|^2 \geq 0, \end{aligned}$$

for all finite sequences $\{h_k\}_{k=0}^n \subseteq \mathcal{H}$. \square

COROLLARY 3.4. *Let $F: \mathfrak{B}(\mathbb{R}) \rightarrow \mathcal{B}(\mathcal{H})$ be a semispectral measure with compact support. Then $\text{Var}(F) \geq 0$, where $\text{Var}(F)$ is as in (1.2).*

PROOF. Apply Proposition 3.3 with $n = 2$ and use the following well-known fact (see [17, Lemma 1]; see also [35, Theorem 5.1]): if $A, B \in \mathcal{B}(\mathcal{H})$ are selfadjoint, A is invertible in $\mathcal{B}(\mathcal{H})$ and $X \in \mathcal{B}(\mathcal{H})$, then $\begin{bmatrix} A & X \\ X^* & B \end{bmatrix} \geq 0$ if and only if $B \geq X^* A^{-1} X$. \square

4. Proof of Theorem 1.8

Before proving the main result of this section, we state the crucial lemma which seems to be of some independent interest. We provide two proofs of this lemma.

LEMMA 4.1. *Let \mathcal{A} be a unital C^* -algebra, $\Phi: \mathcal{A} \rightarrow \mathcal{B}(\mathcal{H})$ be a unital positive linear map and a be a selfadjoint element of \mathcal{A} . Then there exists a unique semispectral measure $F: \mathfrak{B}(\mathbb{R}) \rightarrow \mathcal{B}(\mathcal{H})$ such that $x^n \in L^1(F)$ for all $n \in \mathbb{Z}_+$ and*

$$\Phi(a^n) = \int_{\mathbb{R}} x^n F(dx), \quad n \in \mathbb{Z}_+.$$

Moreover, F possesses the following properties:

- (i) F has compact support,
- (ii) the closed support of F is contained in \mathbb{R}_+ whenever $a \geq 0$.

FIRST PROOF OF LEMMA 4.1. Replacing \mathcal{A} by the unital C^* -algebra generated by $\{a\}$, we may assume without loss of generality that \mathcal{A} is commutative. Let e denote the unit of \mathcal{A} . According to [41, Corollary 2.9], Φ is contractive and therefore

$$\|\Phi(a^n)\| \leq \|a\|^n, \quad n \in \mathbb{Z}_+. \quad (4.1)$$

Since a is selfadjoint, we see that

$$\begin{bmatrix} e & a^1 & \cdots & a^n \\ a^1 & a^2 & \cdots & a^{n+1} \\ \vdots & \vdots & \ddots & \vdots \\ a^n & a^{n+1} & \cdots & a^{2n} \end{bmatrix} = \begin{bmatrix} e & a^1 & \cdots & a^n \\ 0 & 0 & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & 0 \end{bmatrix}^* \begin{bmatrix} e & a^1 & \cdots & a^n \\ 0 & 0 & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & 0 \end{bmatrix} \geq 0.$$

By the Stinespring theorem (see [50, Theorem 4]), Φ is completely positive, so $[\Phi(a^{j+k})]_{j,k=0}^n \geq 0$. In particular, we have

$$\sum_{j,k=0}^n \bar{\lambda}_j \lambda_k \Phi(a^{j+k}) \geq 0, \quad \{\lambda_j\}_{j=0}^n \subseteq \mathbb{C}, \quad n \in \mathbb{Z}_+. \quad (4.2)$$

Using (4.1) and (4.2), we deduce from [11, Theorem 2] that there exists a semispectral measure $F: \mathfrak{B}(\mathbb{R}) \rightarrow \mathcal{B}(\mathcal{H})$ such that

$$\langle \Phi(a^n)h, h \rangle = \int_{\mathbb{R}} x^n \langle F(dx)h, h \rangle, \quad n \in \mathbb{Z}_+, h \in \mathcal{H}. \quad (4.3)$$

(that $F(\mathbb{R}) = I$ follows from the assumption that Φ is unital). Since $\lim_{r \rightarrow \infty} \|f\|_r = \|f\|_{\infty}$ whenever $\|f\|_r < \infty$ for some $r < \infty$ (see [47, Exercise 4, p. 71]) and

$$\lim_{n \rightarrow \infty} \left(\int_{\mathbb{R}} x^{2n} \langle F(dx)h, h \rangle \right)^{\frac{1}{2n}} \stackrel{(4.3)}{=} \lim_{n \rightarrow \infty} \langle \Phi(a^{2n})h, h \rangle^{\frac{1}{2n}} \stackrel{(4.1)}{\leq} \|a\|, \quad h \in \mathcal{H},$$

we deduce that

$$\langle F(\{x \in \mathbb{R}: |x| > \|a\|\})h, h \rangle = 0, \quad h \in \mathcal{H}.$$

Thus, the closed support of F is contained in $[-\|a\|, \|a\|]$. Combined with (2.1) and (4.3), this implies that $x^n \in L^1(F)$ for all $n \in \mathbb{Z}_+$ and

$$\Phi(a^n) = \int_{\mathbb{R}} x^n F(dx), \quad n \in \mathbb{Z}_+.$$

Using (2.1) and the well-known fact that a Hamburger moment sequence having a representing measure with compact support is determinate (see [18]), we get the uniqueness of F .

It remains to show that if $a \geq 0$, then the closed support of F is contained in \mathbb{R}_+ . Using the square root theorem (see [36, Theorem 2.2.1]), we deduce that

$$\begin{bmatrix} a^1 & a^2 & \cdots & a^{n+1} \\ a^2 & a^3 & \cdots & a^{n+2} \\ \vdots & \vdots & \ddots & \vdots \\ a^{n+1} & a^{n+2} & \cdots & a^{2n+1} \end{bmatrix} = \begin{bmatrix} a^{\frac{1}{2}} & a^{\frac{3}{2}} & \cdots & a^{\frac{2n+1}{2}} \\ 0 & 0 & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & 0 \end{bmatrix}^* \begin{bmatrix} a^{\frac{1}{2}} & a^{\frac{3}{2}} & \cdots & a^{\frac{2n+1}{2}} \\ 0 & 0 & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & 0 \end{bmatrix} \geq 0.$$

Hence, by [50, Theorem 4], $[\Phi(a^{j+k+1})]_{j,k=0}^n \geq 0$, which implies that

$$\sum_{j,k=0}^n \bar{\lambda}_j \lambda_k \Phi(a^{j+k+1}) \geq 0, \quad \{\lambda_j\}_{j=0}^n \subseteq \mathbb{C}, \quad n \in \mathbb{Z}_+. \quad (4.4)$$

Combining (4.2), (4.4) and the Stieltjes theorem (see [8, Theorem 6.2.5]) with the uniqueness of F , we conclude that the closed support of F is contained in \mathbb{R}_+ . \square

SECOND PROOF OF LEMMA 4.1. As in the first proof of Lemma 4.1, there is no loss of generality in assuming that \mathcal{A} is commutative. By the Stinespring dilation theorem (see [50, Theorems 1 and 4]), there exist a Hilbert space \mathcal{K} containing \mathcal{H} and a $*$ -representation $\pi: \mathcal{A} \rightarrow \mathcal{B}(\mathcal{K})$ such that

$$\Phi(u) = P\pi(u)|_{\mathcal{H}}, \quad u \in \mathcal{A}, \quad (4.5)$$

where $P \in \mathcal{B}(\mathcal{K})$ is the orthogonal projection of \mathcal{K} onto \mathcal{H} . Applying [46, Theorem 12.22], we deduce that there exists a spectral measure $E: \mathfrak{B}(\mathfrak{M}) \rightarrow \mathcal{B}(\mathcal{K})$ such that

$$\pi(u) = \int_{\mathfrak{M}} \widehat{\pi(u)} dE, \quad u \in \mathcal{A}, \quad (4.6)$$

where \mathfrak{M} is the maximal ideal space of the unital commutative C^* -algebra $\overline{\pi(\mathcal{A})}$, the (operator norm) closure of $\pi(\mathcal{A})$ in $\mathcal{B}(\mathcal{K})$, and $\widehat{\pi(u)}: \mathfrak{M} \rightarrow \mathbb{C}$ is the Gelfand transform of $\pi(u)$. Set $M = PE|_{\mathcal{H}}$. It follows from (4.5) and (4.6) that

$$\Phi(u) = \int_{\mathfrak{M}} \widehat{\pi(u)} dM, \quad u \in \mathcal{A}. \quad (4.7)$$

Define the semispectral measure $F: \mathfrak{B}(\mathbb{R}) \rightarrow \mathcal{B}(\mathcal{H})$ by

$$F(\Delta) = M(\widehat{\pi(a)^{-1}(\Delta)}), \quad \Delta \in \mathfrak{B}(\mathbb{R}).$$

By [46, Theorem 11.18] and the assumption that $a = a^*$, we see that $\widehat{\pi(a)}: \mathfrak{M} \rightarrow \mathbb{R}$. Since $\widehat{\pi(a)}$ is continuous and \mathfrak{M} is a compact Hausdorff space, we deduce that $\widehat{\pi(a)}(\mathfrak{M})$ is a compact subset of \mathbb{R} such that $F(\mathbb{R} \setminus \widehat{\pi(a)}(\mathfrak{M})) = 0$, which implies that the semispectral measure F has compact support. Applying (2.1) and the measure transport theorem (cf. [2, Theorem 1.6.12]), we conclude that

$$\Phi(a^n) \stackrel{(4.7)}{=} \int_{\mathfrak{M}} \widehat{\pi(a)}^n dM = \int_{\mathbb{R}} x^n F(dx), \quad n \in \mathbb{Z}_+.$$

The proof of the uniqueness of F proceeds as before.

Finally, if $a \geq 0$, then by the square root theorem and [46, Theorem 11.18], we deduce that

$$\widehat{\pi(a)} = \widehat{\pi(a^{\frac{1}{2}})}^2 \geq 0,$$

which implies that the closed support of F is contained in \mathbb{R}_+ . \square

At this point we are ready to prove the main result of this section.

PROOF OF THEOREM 1.8. (i) \Rightarrow (ii) Since the map Φ preserves selfadjointness, $b := \Phi(a)$ does the job.

(ii) \Rightarrow (i) In view of the Gelfand-Naimark theorem (see [46, Theorem 12.41]), there is no loss of generality in assuming that $\mathcal{B} = \mathcal{B}(\mathcal{H})$. By Lemma 4.1, there exists a semispectral measure $F: \mathfrak{B}(\mathbb{R}) \rightarrow \mathcal{B}(\mathcal{H})$ with compact support such that

$$\Phi(a^n) = \int_{\mathbb{R}} x^n F(dx), \quad n \in \mathbb{Z}_+. \quad (4.8)$$

Therefore, by (ii), we have

$$b^k = \Phi(a^k) \stackrel{(4.8)}{=} \int_{\mathbb{R}} x^k F(dx), \quad k = p, q.$$

Applying Theorem 1.7 to $T = b$, we conclude that F is the spectral measure of b . Using the Stone-von Neumann functional calculus, we get

$$\Phi(a^n) \stackrel{(4.8)}{=} \left(\int_{\mathbb{R}} x F(dx) \right)^n = b^n, \quad n \in \mathbb{Z}_+,$$

so $b = \Phi(a)$, which yields $\Phi(a^n) = \Phi(a)^n$ for all $n \in \mathbb{Z}_+$. This implies (i). \square

REMARK 4.2. Theorem 1.8 is somewhat related to a result of D. Petz (see [42, Theorem]) which shows that equality holds in Jensen's inequality $f(\Phi(a)) \leq \Phi(f(a))$ if and only if Φ restricted to the unital subalgebra generated by a is multiplicative, where Φ is a unital positive linear map between unital C^* -algebras, f is a non-affine operator convex function on an open subinterval J of \mathbb{R} and a is a selfadjoint element with spectrum in J . The main difference between Petz's result and Theorem 1.8 is

that the monomial x^n with $n \in \mathbb{Z}_+$ is a non-affine operator convex function on J if and only if $n = 2$ (apply rescaling and [9, Exercise V.2.11], see also [45]). \diamond

REMARK 4.3. We have deduced Theorem 1.8 from Theorem 1.7. It turns out that these two results are logically equivalent. Indeed, under the assumptions and notation of Theorem 1.7, it suffices to show that (iii) implies (i). For, define the unital positive linear map $\Phi: C(K) \rightarrow \mathcal{B}(\mathcal{H})$ by

$$\Phi(f) = \int_K f(x)F(dx), \quad f \in C(K),$$

where K stands for the closed support of F . Let $a \in C(K)$ be the function defined by $a(x) = x$ for $x \in K$ and let $b = T$. Using Theorem 1.8, we deduce that $T = \int_K xF(dx)$ and

$$\int_K x^n F(dx) = T^n = \int_K x^n G(dx), \quad n \in \mathbb{Z}_+,$$

where G is the spectral measure of T . Using (2.1) and the well-known fact that a Hamburger moment sequence having a representing measure with compact support is determinate, we conclude that $F = G$, which completes the proof.

A careful inspection of the proof of Theorem 1.8 in conjunction with the above discussion shows that in fact Problems 1.4 and 1.6 are logically equivalent regardless of the cardinality of the set Ξ . \diamond

In case where the elements a and b are positive, we get the following version of Theorem 1.8.

THEOREM 4.4. *Suppose that \mathcal{A} and \mathcal{B} are unital C^* -algebras, $\Phi: \mathcal{A} \rightarrow \mathcal{B}$ is a unital positive linear map, $a \in \mathcal{A}$ is positive and p, q are distinct positive integers. Then the following conditions are equivalent:*

- (i) Φ restricted to the unital subalgebra generated by $\{a\}$ is multiplicative,
- (ii) there exists a positive element $b \in \mathcal{B}$ such that $b^k = \Phi(a^k)$ for $k = p, q$.

Moreover, if (ii) holds, then $b = \Phi(a)$.

PROOF. It suffices to show the implication (ii) \Rightarrow (i). We present two proofs. The first one is the same as the proof of Theorem 1.8, with the only difference that we use Theorem 1.3 instead of Theorem 1.7. We leave the details to the reader.

The second proof relies upon Petz's result. Without loss of generality, we may assume that $0 < p < q$. Let $f: [0, \infty) \rightarrow \mathbb{R}$ be the function given by $f(x) = -x^{\frac{p}{q}}$ for $x \in [0, \infty)$. It follows from [9, Theorems V.1.9 and V.2.5] that f is an operator convex function. Using (ii) and the Stone-von Neumann functional calculus, we get

$$f(\Phi(a^q)) = -\Phi(a^q)^{\frac{p}{q}} = -(b^q)^{\frac{p}{q}} = -b^p,$$

and

$$\Phi(f(a^q)) = -\Phi((a^q)^{\frac{p}{q}}) = -\Phi(a^p) = -b^p.$$

Consequently,

$$f(\Phi(a^q)) = \Phi(f(a^q)).$$

Combined with [42, Theorem] and the fact that Φ is continuous (see [41, Corollary 2.9]), this implies that Φ restricted to the unital C^* -algebra generated by $\{a^q\}$ is multiplicative. Applying the Stone-von Neumann functional calculus and the

Weierstrass approximation theorem, one can show that the unital C^* -algebras generated by $\{a\}$ and $\{a^q\}$ coincide (this is a very special case of the Müntz-Szász theorem, see [47, Theorem 15.26]). Hence, (i) holds. \square

5. Proof of Theorem 1.9

We begin with a simple observation related to Problems 1.4 and 1.5. Namely, if T and F satisfy (1.3), then by (2.1) and the measure transport theorem for every $\tau \in \mathbb{R} \setminus \{0\}$, τT and F_τ satisfy (1.3), where $F_\tau: \mathfrak{B}(\mathbb{R}) \rightarrow \mathcal{B}(\mathcal{H})$ is the semispectral measure with compact support given by

$$F_\tau(\Delta) = F(\tau^{-1}\Delta), \quad \Delta \in \mathfrak{B}(\mathbb{R}).$$

Moreover, F is spectral if and only if F_τ is spectral. A similar observation applies to Problem 1.5. In other words, rescaling preserves the affirmative or negative solutions to Problems 1.4 and 1.5.

Next we prove a lemma that is central to the proof of Theorem 5.2.

LEMMA 5.1. *Suppose that $(p, q) \in \mathbb{N}^2 \setminus \Omega$ and $p \leq q$, where Ω is as in (1.4). Then for every $\tau \in \mathbb{R} \setminus \{0\}$, there exist $\alpha, \beta \in (0, 1)$ and distinct $\lambda_1, \lambda_2 \in \mathbb{R}$ such that*

$$\begin{cases} \alpha + \beta = 1, \\ \alpha \lambda_1^p + \beta \lambda_2^p = \tau^p, \\ \alpha \lambda_1^q + \beta \lambda_2^q = \tau^q. \end{cases} \quad (5.1)$$

PROOF. We may assume, without loss of generality, that $\tau = 1$. Using the substitution

$$\alpha = \frac{a}{a+b} \quad \text{and} \quad \beta = \frac{b}{a+b}$$

with $a, b \in (0, \infty)$, we obtain an equivalent system of equations:

$$\begin{cases} \frac{a}{a+b} \lambda_1^p + \frac{b}{a+b} \lambda_2^p = 1, \\ \frac{a}{a+b} \lambda_1^q + \frac{b}{a+b} \lambda_2^q = 1. \end{cases}$$

Multiplying both sides of the above equalities by $a+b$ and rearranging gives

$$\begin{cases} a(\lambda_1^p - 1) + b(\lambda_2^p - 1) = 0, \\ a(\lambda_1^q - 1) + b(\lambda_2^q - 1) = 0. \end{cases} \quad (5.2)$$

The determinant of the above system of equations (with unknowns a, b) is

$$D(\lambda_1, \lambda_2) = \det \begin{bmatrix} \lambda_1^p - 1 & \lambda_2^p - 1 \\ \lambda_1^q - 1 & \lambda_2^q - 1 \end{bmatrix} = (\lambda_1^p - 1)(\lambda_2^q - 1) - (\lambda_2^p - 1)(\lambda_1^q - 1).$$

Observe that if $D(\lambda_1, \lambda_2) \neq 0$, then the system of equations (5.2) has only one solution $a = b = 0$. Thus, the only chance to find nonzero solutions a, b of the system (5.2) is when $D(\lambda_1, \lambda_2) = 0$. Note that

$$\text{if } \lambda_1, \lambda_2 \in \mathbb{R} \setminus \{-1, 1\}, \text{ then } D(\lambda_1, \lambda_2) = 0 \text{ if and only if } \frac{\lambda_1^p - 1}{\lambda_1^q - 1} = \frac{\lambda_2^p - 1}{\lambda_2^q - 1}. \quad (5.3)$$

We will consider four cases.

CASE 1. $p = q$.

It is easily seen that for any $\alpha, \beta \in (0, 1)$ such that $\alpha + \beta = 1$, there are plenty of two-element subsets $\{\lambda_1, \lambda_2\}$ of $(0, \infty)$ solving (5.1).

CASE 2. Both p and q are even.

Set $\lambda_1 = -1$, $\lambda_2 = 1$ and $\beta = 1 - \alpha$, where $\alpha \in (0, 1)$. Then it is easily seen that (5.1) is satisfied.

CASE 3. $p < q$, p even and q odd.

Consider the function $\phi: [0, 1] \rightarrow [0, 1]$ defined by

$$\phi(x) = \frac{1 - x^p}{1 + x^q}, \quad x \in [0, 1].$$

Then ϕ is continuous, $\phi(0) = 1$, $\phi(1) = 0$ and $\phi((0, 1)) \subseteq (0, 1)$. By the Darboux property of continuous functions, $\phi((0, 1)) = (0, 1)$. Take $\lambda_1 > 1$. Then

$$\frac{1 - \lambda_1^p}{1 - \lambda_1^q} \in (0, 1).$$

Therefore, there exists $x \in (0, 1)$ such that

$$\frac{1 - \lambda_1^p}{1 - \lambda_1^q} = \phi(x). \quad (5.4)$$

Set $\lambda_2 = -x$. Then $\lambda_2 < 0$, $|\lambda_2| = x < 1$ and

$$\frac{1 - \lambda_1^p}{1 - \lambda_1^q} \stackrel{(5.4)}{=} \frac{1 - |\lambda_2|^p}{1 + |\lambda_2|^q} = \frac{1 - \lambda_2^p}{1 - \lambda_2^q},$$

which by (5.3) means that $D(\lambda_1, \lambda_2) = 0$, so the system of equations (5.2) is linearly dependent. Take any $a \in (0, \infty)$ and set

$$b = a \frac{\lambda_1^p - 1}{1 - \lambda_2^p} > 0.$$

Then the pair (a, b) is a solution of the system of equations (5.2).

CASE 4. $p < q$ and both p and q are odd.

Consider the function $\psi: [1, \infty) \rightarrow (0, 1]$ defined by

$$\psi(x) = \frac{1 + x^p}{1 + x^q}, \quad x \in [1, \infty).$$

Then ψ is continuous, $\psi(1) = 1$, $\lim_{x \rightarrow \infty} \psi(x) = 0$ and $\psi((1, \infty)) \subseteq (0, 1)$. As a consequence of the Darboux property of continuous functions, $\psi((1, \infty)) = (0, 1)$. Take $\lambda_1 > 1$. Observe that

$$\frac{1 - \lambda_1^p}{1 - \lambda_1^q} \in (0, 1).$$

Hence, there exists $x \in (1, \infty)$ such that

$$\frac{1 - \lambda_1^p}{1 - \lambda_1^q} = \psi(x). \quad (5.5)$$

Set $\lambda_2 := -x$. Then $\lambda_2 < 0$, $|\lambda_2| = x > 1$ and

$$\frac{1 - \lambda_1^p}{1 - \lambda_1^q} \stackrel{(5.5)}{=} \psi(|\lambda_2|) = \frac{1 + |\lambda_2|^p}{1 + |\lambda_2|^q} = \frac{1 - \lambda_2^p}{1 - \lambda_2^q}.$$

As in Case 3, taking any $a \in (0, \infty)$ and setting

$$b = a \frac{\lambda_1^p - 1}{1 - \lambda_2^p} = a \frac{\lambda_1^p - 1}{1 + |\lambda_2|^p} > 0,$$

we see that the pair (a, b) is a solution of the system of equations (5.2). This completes the proof. \square

We are now ready to prove the main result of this section, which provides the counter-examples mentioned earlier in Introduction. In fact, this is a stronger version of Theorem 1.9.

THEOREM 5.2. *Suppose that $(p, q) \in \mathbb{N}^2 \setminus \Omega$ and $p \leq q$, where Ω is as in (1.4). Let $\tau \in \mathbb{R} \setminus \{0\}$. Set $\mathcal{H} = \mathbb{C}$ and $T = \tau I$. Then there exist $\alpha, \beta \in (0, 1)$ and $\lambda_1, \lambda_2 \in \mathbb{R}$ such that $\alpha + \beta = 1$, $\lambda_1 \neq \lambda_2$ and*

(i) *the semispectral measure $F: \mathfrak{B}(\mathbb{R}) \rightarrow \mathbf{B}(\mathcal{H})$ defined by*

$$F(\Delta) = \alpha \delta_{\lambda_1}(\Delta)I + \beta \delta_{\lambda_2}(\Delta)I, \quad \Delta \in \mathfrak{B}(\mathbb{R}), \quad (5.6)$$

is not spectral and

$$T^k = \int_{\mathbb{R}} x^k F(dx), \quad k = p, q, \quad (5.7)$$

(ii) *the selfadjoint operator $S \in \mathbf{B}(\mathcal{H} \oplus \mathcal{H})$ defined by*

$$S = \begin{bmatrix} \alpha\lambda_1 + \beta\lambda_2 & \sqrt{\alpha\beta}(\lambda_1 - \lambda_2) \\ \sqrt{\alpha\beta}(\lambda_1 - \lambda_2) & \beta\lambda_1 + \alpha\lambda_2 \end{bmatrix}$$

does not commute with $P := \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}$ and

$$T^k = PS^k|_{\mathcal{H}}, \quad k = p, q. \quad (5.8)$$

(iii) *the unital positive linear map $\Phi: C(K) \rightarrow \mathbf{B}(\mathcal{H})$ defined by*

$$\Phi(f) = \int_K f dF, \quad f \in C(K), \quad (5.9)$$

is not multiplicative, $C(K)$ is the unital algebra generated by a and

$$b^k = \Phi(a^k), \quad k = p, q, \quad (5.10)$$

where F is as in (5.6), $K = \{\lambda_1, \lambda_2\}$, $a(x) = x$ for $x \in K$ and $b = T$.

PROOF. Let $\alpha, \beta, \lambda_1, \lambda_2$ be as in Lemma 5.1.

(i) By (5.1) and (5.6), F is a semispectral measure satisfying (5.7). However, F is not a spectral measure because $F(\{\lambda_1\}) = \alpha \in (0, 1)$.

(ii) Clearly, the operator S is selfadjoint. It follows from the first equality in (5.1) that the matrix

$$\begin{bmatrix} \sqrt{\alpha} & \sqrt{\beta} \\ -\sqrt{\beta} & \sqrt{\alpha} \end{bmatrix}$$

is unitary and consequently

$$\begin{bmatrix} \sqrt{\alpha} & \sqrt{\beta} \\ -\sqrt{\beta} & \sqrt{\alpha} \end{bmatrix}^{-1} = \begin{bmatrix} \sqrt{\alpha} & -\sqrt{\beta} \\ \sqrt{\beta} & \sqrt{\alpha} \end{bmatrix}. \quad (5.11)$$

Now it is easily seen that the Jordan decomposition of S takes the form

$$S = \begin{bmatrix} \sqrt{\alpha} & -\sqrt{\beta} \\ \sqrt{\beta} & \sqrt{\alpha} \end{bmatrix} \begin{bmatrix} \lambda_1 & 0 \\ 0 & \lambda_2 \end{bmatrix} \begin{bmatrix} \sqrt{\alpha} & \sqrt{\beta} \\ -\sqrt{\beta} & \sqrt{\alpha} \end{bmatrix}.$$

Combined with (5.11), this implies that

$$S^n = \begin{bmatrix} \alpha\lambda_1^n + \beta\lambda_2^n & \sqrt{\alpha\beta}(\lambda_1^n - \lambda_2^n) \\ \sqrt{\alpha\beta}(\lambda_1^n - \lambda_2^n) & \beta\lambda_1^n + \alpha\lambda_2^n \end{bmatrix}, \quad n \in \mathbb{Z}_+. \quad (5.12)$$

By (5.1) and (5.12), the condition (5.8) is satisfied. Since $\lambda_1 \neq \lambda_2$, the operator S does not commute with P .

(iii) It is immediate from (5.9) and (i) that Φ is the unital positive linear map which satisfies (5.10). Clearly, $C(K)$ is the unital algebra generated by a . To show that Φ is not multiplicative, consider two polynomials $u(x) = x - \lambda_1$ and $v(x) = x - \lambda_2$ and note that $\Phi(u(a)v(a)) = 0$ while $\Phi(u(a))\Phi(v(a)) = -\alpha\beta(\lambda_1 - \lambda_2)^2 \neq 0$. This completes the proof. \square

REMARK 5.3. A careful inspection of the proofs of Lemma 5.1 and Theorem 5.2 reveals that there is a lot of freedom in choosing the parameters λ_1 and λ_2 . Note that if $p < q$ and $\tau > 0$, then in view of [43, Theorem 4.2], at least one of the parameters λ_1 or λ_2 must be negative. Note also that if $p < q$, p is odd and q is even, then according to Theorem 1.7 there are no $\alpha, \beta, \lambda_1, \lambda_2$ satisfying the conclusion of Theorem 5.2. However, in this particular case, we can justify it in an elementary way. Namely, by the Hölder inequality, we infer from (5.1) that

$$\begin{aligned} |\tau|^p &= |\alpha\lambda_1^p + \beta\lambda_2^p| \leq \alpha|\lambda_1|^p + \beta|\lambda_2|^p \\ &\leq \sqrt[r]{\alpha + \beta} \sqrt[q]{\alpha|\lambda_1|^q + \beta|\lambda_2|^q} \\ &= \sqrt[r]{\alpha + \beta} \sqrt[q]{\alpha\lambda_1^q + \beta\lambda_2^q} = |\tau|^p, \end{aligned} \quad (5.13)$$

where $r \in (1, \infty)$ is such that $\frac{1}{r} + \frac{p}{q} = 1$. This means that equality in the Hölder inequality holds. As a consequence, we deduce that $|\lambda_1| = |\lambda_2|$. Combined with the first inequality in (5.13) and the assumption that p is odd, this implies that $\lambda_1 = \lambda_2$, which is a contradiction. \diamond

6. More examples

In this section we illustrate Theorem 1.9 (cf. Theorem 5.2) by considering two interesting examples for the case where $p = 2$ and $q = 3$. Now we use the dilation approach as stated in Lemma 3.2 (cf. Problems 1.4 and 1.5).

EXAMPLE 6.1. Let $\{f_n\}_{n=0}^\infty$ be the Fibonacci sequence, that is, $f_0 = 0$, $f_1 = 1$ and $f_{n+1} = f_n + f_{n-1}$ for $n \in \mathbb{N}$. It is well known and easy to prove that

$$\begin{bmatrix} 0 & 1 \\ 1 & 1 \end{bmatrix}^n = \begin{bmatrix} f_{n-1} & f_n \\ f_n & f_{n+1} \end{bmatrix}, \quad n \in \mathbb{N}. \quad (6.1)$$

Set $\mathcal{H} = \mathbb{C}$, $T = I$ and

$$S = \begin{bmatrix} 0 & 1 \\ 1 & 1 \end{bmatrix}.$$

Then S is a selfadjoint operator and the spectral measure E of S is given by

$$E(\Delta) = \frac{1}{1 + \phi^2} \delta_{1-\phi}(\Delta) \begin{bmatrix} \phi^2 & -\phi \\ -\phi & 1 \end{bmatrix} + \frac{1}{1 + \phi^2} \delta_\phi(\Delta) \begin{bmatrix} 1 & \phi \\ \phi & \phi^2 \end{bmatrix}, \quad \Delta \in \mathfrak{B}(\mathbb{R}),$$

where $\phi = \frac{1+\sqrt{5}}{2}$ is the golden ratio. It follows from (6.1) that

$$T^k = PS^k|_{\mathcal{H}} \text{ for } k = 2, 3 \text{ and } T^k \neq PS^k|_{\mathcal{H}} \text{ for } k \in \mathbb{N} \setminus \{2, 3\},$$

where $P = \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}$ is the orthogonal projection of $\mathcal{H} \oplus \mathcal{H}$ onto \mathcal{H} . The semispectral measure $F := PE|_{\mathcal{H}}$ (see (3.2) for the definition of $PE|_{\mathcal{H}}$) takes the form

$$F(\Delta) = \frac{\phi^2}{1 + \phi^2} \delta_{1-\phi}(\Delta) I + \frac{1}{1 + \phi^2} \delta_\phi(\Delta) I, \quad \Delta \in \mathfrak{B}(\mathbb{R}). \quad (6.2)$$

By Lemma 3.2(ii), we have

$$T^k = \int_{\mathbb{R}} x^k F(dx) \text{ for } k = 2, 3 \text{ and } T^k \neq \int_{\mathbb{R}} x^k F(dx) \text{ for } k \in \mathbb{N} \setminus \{2, 3\}.$$

Clearly, by (6.2), F is not a spectral measure (also because P does not commute with S , see Lemma 3.2). \diamond

The next example is a modification of the previous one.

EXAMPLE 6.2. Let $T \in \mathbf{B}(\mathcal{H})$ be a nonzero selfadjoint operator and let $S \in \mathbf{B}(\mathcal{H} \oplus \mathcal{H})$ be the selfadjoint operator given by the 2×2 block matrix

$$S = \begin{bmatrix} 0 & T \\ T & T \end{bmatrix}. \quad (6.3)$$

One can verify that

$$S^2 = \begin{bmatrix} T^2 & T^2 \\ T^2 & 2T^2 \end{bmatrix} \quad \text{and} \quad S^3 = \begin{bmatrix} T^3 & 2T^3 \\ 2T^3 & 3T^3 \end{bmatrix}.$$

Thus the operators T and S satisfy the following two identities:

$$T^k = PS^k|_{\mathcal{H}}, \quad k = 2, 3, \quad (6.4)$$

where $P = \begin{bmatrix} I_{\mathcal{H}} & 0 \\ 0 & 0 \end{bmatrix}$ is the orthogonal projection of $\mathcal{H} \oplus \mathcal{H}$ onto \mathcal{H} . Let E be the spectral measure of S and let $F := PE|_{\mathcal{H}}$ be the corresponding semispectral measure. Applying Lemma 3.2(ii) and using (6.4), we get

$$T^k = \int_{\mathbb{R}} x^k F(dx), \quad k = 2, 3.$$

Since $T \neq 0$, the operator P does not commute with S , so by Lemma 3.2, F is not a spectral measure. In contrast to Example 6.1, here it is much easier to use Lemma 3.2 to see that the semispectral measure F is not spectral. \diamond

REMARK 6.3. Regarding Example 6.2, note that the operator S given by (6.3) is unitarily equivalent to the tensor product

$$S = \begin{bmatrix} 0 & 1 \\ 1 & 1 \end{bmatrix} \otimes T.$$

Combined with (6.1), this implies that

$$S^n = \begin{bmatrix} 0 & 1 \\ 1 & 1 \end{bmatrix}^n \otimes T^n = \begin{bmatrix} f_{n-1}T^n & f_nT^n \\ f_nT^n & f_{n+1}T^n \end{bmatrix}, \quad n \in \mathbb{N}.$$

This means that tensoring and orthogonal summation enrich the class of counterexamples by allowing semispectral measures to have operator values on Hilbert spaces of arbitrary dimension. \diamond

7. Semispectral measures with non-compact supports

In this section we extend the two-moment characterizations of spectral measures given in Theorems 1.3 and 1.7 to the case of semispectral measures with non-compact supports. When considering a Borel semispectral measure F on the real line with non-compact support, it may happen that the coordinate function $\mathbb{R} \ni x \mapsto x \in \mathbb{R}$ is not in $L^1(F)$ (cf. (2.1)), which means that the expression $\int_{\mathbb{R}} x^k F(dx)$, where $k \in \mathbb{N}$, may not yield a bounded operator. On the other hand, if Problem 1.4 has an affirmative solution under the formally weaker assumption that the functions $\mathbb{R} \ni x \mapsto x^k \in \mathbb{R}$, $k \in \mathbb{N}$, are in $L^1(F)$, then F being *a posteriori* a spectral

measure must have a compact support. Indeed, by the measure transport theorem $T^k = \int_{\mathbb{R}} x(F \circ \varphi_k^{-1})(dx)$, where φ_k is as in (3.7) and k is any element of Ξ , and thus $F \circ \varphi_k^{-1}$ is the spectral measure of the (bounded) selfadjoint operator T^k . Hence, $F \circ \varphi_k^{-1}$ must have compact support (see [48, Theorem 5.9]). Consequently, F itself must have compact support. A similar argument applied in the case where a semispectral measure $F: \mathcal{A} \rightarrow \mathcal{B}(\mathcal{H})$ is considered on an abstract measurable space (X, \mathcal{A}) , and the coordinate function is replaced by a measurable real-valued function ω on X , leads to the conclusion that ω is F -essentially bounded, that is, $F(\{x \in X: |\omega(x)| > r\}) = 0$ for some $r \in \mathbb{R}_+$ (equivalently, $\omega \in L^\infty(F)$). However, to get the spectrality of F , it is not enough to assume that ω is F -essentially bounded. It turns out that the “missing” property of ω is σ -surjectivity. We say that a measurable map $f: X \rightarrow Y$ between measurable spaces (X, \mathcal{A}) and (Y, \mathcal{B}) (i.e., a map such that $f^{-1}(\Delta) \in \mathcal{A}$ for all $\Delta \in \mathcal{B}$) is σ -surjective if the corresponding map $\mathcal{B} \ni \Delta \mapsto f^{-1}(\Delta) \in \mathcal{A}$ is surjective. If Y is a topological Hausdorff space, σ -surjectivity refers to $\mathcal{B} = \mathfrak{B}(Y)$. In case $\mathcal{A} = \mathfrak{B}(X)$ and $\mathcal{B} = \mathfrak{B}(Y)$, where X and Y are topological Hausdorff spaces, σ -surjectivity is called in [15] *Borel injectivity*. It is worth emphasizing here that the property of being σ -surjective was used in the proof of Theorem 1.7.

Applying the measure transport theorem together with Theorems 1.3 and 1.7, we get the following.

THEOREM 7.1. *Assume that (X, \mathcal{A}) is a measurable space, $F: \mathcal{A} \rightarrow \mathcal{B}(\mathcal{H})$ is a semispectral measure, $T \in \mathcal{B}(\mathcal{H})$ is a selfadjoint operator and p, q are positive integers such that $p < q$. Let $\omega: X \rightarrow \mathbb{R}$ be an F -essentially bounded σ -surjective function such that*

$$T^k = \int_X \omega(x)^k F(dx), \quad k = p, q.$$

If p is odd and q is even, or if $\omega(X) \subseteq \mathbb{R}_+$, then F is a spectral measure.

It is worth pointing out that there is a wide class of measurable spaces admitting functions ω with the properties mentioned in Theorem 7.1. Namely, if X is a Borel subset of a complete separable metric space and K is a bounded Borel subset of \mathbb{R} such that X and K have the same cardinality, then by [40, Theorem 2.12] there exists a bijection $\omega_0: X \rightarrow K$ such that ω_0 and ω_0^{-1} are Borel measurable. This implies that the function $\omega: X \rightarrow \mathbb{R}$ defined by $\omega(x) = \omega_0(x)$ for $x \in X$ is bounded and σ -surjective. It turns out that the notions of injectivity and σ -surjectivity coincide for continuous maps $f: X \rightarrow Y$ between topological Hausdorff spaces whenever X is σ -compact (see [15, Proposition 16]). Coming back to the case of $X = Y = \mathbb{R}$, let us recall the well-known example of a bounded continuous and injective (consequently, σ -surjective) function $\omega: \mathbb{R} \rightarrow \mathbb{R}$ given by $\omega(x) = \frac{x}{1+|x|}$ for $x \in \mathbb{R}$.

References

- [1] N. I. Akhiezer, I. M. Glazman, *Theory of Linear Operators in Hilbert Space*, Dover Publications, Inc. New York, 1993, Vol. II.
- [2] R. B. Ash, *Probability and measure theory*, Harcourt/Academic Press, Burlington, 2000.
- [3] R. Beneduci, G. Nisticò, Sharp reconstruction of unsharp quantum observables, *J. Math. Phys.* **44** (2003), 5461.
- [4] R. Beneduci, A geometrical characterization of commutative positive operator valued measures, *J. Math. Phys.* **47** (2006), 062104.

- [5] R. Beneduci, Infinite sequences of linear functionals, positive operator valued measures and Naimark extension theorem, *B. Lond. Math. Soc.* **42** (2010), 441–451.
- [6] R. Beneduci, Positive operator valued measures and Feller Markov kernels, *J. Math. Anal. Appl.* **442** (2016), 50–71.
- [7] S. K. Berberian, *Notes on spectral theory*, Van Nostrand Mathematical Studies, No. 5, D. Van Nostrand Co., Inc., Princeton, N.J.-Toronto, Ont.-London 1966.
- [8] C. Berg, J. P. R. Christensen, P. Ressel, *Harmonic analysis on semigroups*, Springer-Verlag, Berlin, 1984.
- [9] R. Bhatia, *Matrix analysis*, Graduate Texts in Mathematics, 169, Springer-Verlag, New York, 1997.
- [10] M. Sh. Birman, M. Z. Solomjak, *Spectral Theory of selfadjoint operators in Hilbert space*, D. Reidel Publishing Co., Dordrecht, 1987.
- [11] T. M. Bisgaard, Positive definite operator sequences, *Proc. Amer. Math. Soc.* **121** (1994), 1185–1191.
- [12] H. E. Brandt, Positive operator valued measure in quantum information processing, *Am. J. Phys.* **67** (1999), 434–439.
- [13] P. Busch, P. Lahti, J.-P. Pellonpää, K. Ylinen, *Quantum measurement*, Theoretical and Mathematical Physics, Springer, [Cham], 2016.
- [14] G. Cattaneo, G. Nisticò, From unsharp to sharp quantum observables, *J. Math. Phys.* **41** (2000), 4365.
- [15] D. Cichoń, J. Stochel, F. H. Szafraniec, The complex moment problem: determinacy and extendibility, *Math. Scand.* **124** (2019), 263–288.
- [16] R. E. Curto, S. H. Lee, J. Yoon, Quasiness of powers of commuting pairs of bounded operators, *J. Funct. Anal.* **278** (2020), 108342.
- [17] C. Davis, A device for studying Hausdorff moments, *Trans. Amer. Math. Soc.* **87** (1958), 144–158.
- [18] B. Fuglede, The multidimensional moment problem, *Expo. Math.* **1** (1983), 47–65.
- [19] W. F. Donoghue, *Monotone matrix functions and analytic continuation*, Springer-Verlag, Berlin, 1974.
- [20] F. Hansen, An operator inequality, *Math. Ann.* **246** (1980), 249–250.
- [21] F. Hansen, The fast track to Löwner’s theorem, *Linear Algebra Appl.* **438** (2013), 4557–4571.
- [22] A. S. Holevo, An analog of the statistical decision theory and noncommutative probability theory, *Trans. Moscow Math. Soc.* **26** (1972), 133–149.
- [23] A. S. Holevo, *Probabilistic and statistical aspects of quantum theory*. Vol. 1. Springer Science & Business Media, 2011.
- [24] Z. J. Jabłoński, Complete hyperexpansivity, subnormality and inverted boundedness conditions, *Integr. Equ. Oper. Theory* **44** (2002), 316–336.
- [25] I. B. Jung, J. Stochel, Subnormal operators whose adjoints have rich point spectrum, *J. Funct. Anal.* **255** (2008), 1797–1816.
- [26] A. Jenčová, S. Pulmannová, How sharp are PV measures?, *Rep. Math. Phys.* **59** (2007), 257–266.
- [27] R. V. Kadison, A generalized Schwarz inequality and algebraic invariants for operator algebras, *Ann. Math.* **56** (1952), 494–503.
- [28] J. Kiukas, Phase space quantization as a moment problem, *Opt. Spectrosc.* **103** (2007), 429–433.
- [29] J. Kiukas, P. Lahti, K. Ylinen, Normal covariant quantization maps, *J. Math. Anal. Appl.* **319** (2006), 783–801.
- [30] J. Kiukas, P. Lahti, K. Ylinen, Phase space quantisation and the operator moment problem, *J. Math. Phys.* **47** (2006), 072104.
- [31] P. Kruszyński, W. M. de Muynck, Compatibility of observables represented by positive operator-valued measures, *J. Math. Phys.* **28** (1987), 1761–1763.
- [32] E. H. Lieb, M. B. Ruskai, Some operator inequalities of the Schwarz type, *Adv. Math.* **12** (1974), 269–273.
- [33] K. Löwner, Über monotone Matrixfunktionen, *Math. Z.* **38** (1934), 177–216.
- [34] W. Mlak, *Dilations of Hilbert space operators (general theory)*, Dissertationes Math. **153** (1978), 61 pp.
- [35] M. S. Moslehian, M. Kian, Q. Xu, Positivity of 2×2 block matrices of operators, *Banach J. Math. Anal.* **13** (2019), 726–743.

- [36] G. J. Murphy, *C*-algebras and operator theory*, Academic Press, Inc., Boston, MA, 1990.
- [37] M. A. Naimark, On selfadjoint extensions of second kind of a symmetric operator, *Izv. Akad. Nauk SSSR Ser. Mat.* **4** (1940), 53–104.
- [38] M. A. Naimark, Spectral Functions of Symmetric Operators, *Izv. Akad. Nauk SSSR Ser. Mat.* **4**, (1940), 277–318.
- [39] M. A. Naimark, On a Representation of Additive Operator Set Functions, *C.R. Acad. Sci. URSS* **41** (1943), 359–361.
- [40] K. R. Parthasarathy, *Probability measures on metric spaces*, Probability and Mathematical Statistics, No. 3, Academic Press, Inc., New York-London 1967.
- [41] V. I. Paulsen, *Completely bounded maps and dilations*, Pitman Research Notes in Mathematics Series, 146. Longman Scientific & Technical, Harlow; John Wiley & Sons, Inc., New York, 1986.
- [42] D. Petz, On the equality in Jensen’s inequality for operator convex functions, *Integr. Equ. Oper. Theory* **9** (1986), 744–747.
- [43] P. Pietrzycki, J. Stochel, Subnormal n th roots of quasinormal operators are quasinormal, *J. Funct. Anal.* **280** (2021), 109001.
- [44] P. Pietrzycki, J. Stochel, Corrigendum to “Subnormal n th roots of quasinormal operators are quasinormal” [*J. Funct. Anal.* **280** (2021), 109001].
- [45] M. H. Rizvi, R. W. Shorrock, A note on matrix-convexity, *Canad. J. Statist.*, **7** (1979), 39–41.
- [46] W. Rudin, *Functional analysis*, McGraw-Hill Series in Higher Math., McGraw-Hill Book Co., New York, 1973.
- [47] W. Rudin, *Real and complex analysis*, McGraw-Hill, New York, 1987.
- [48] K. Schmüdgen, *Unbounded self-adjoint operators on Hilbert space*, Graduate Texts in Mathematics, 265, Springer, Dordrecht, 2012.
- [49] B. Simon, *Loewner’s theorem on monotone matrix functions*, Grundlehren Math. Wissen. 354, Springer, 2019.
- [50] W. F. Stinespring, Positive functions on C^* -algebras, *Proc. Amer. Math. Soc.* **6** (1955), 211–216.
- [51] J. Stochel, Decomposition and disintegration of positive definite kernels on convex $*$ -semigroups, *Ann. Polon. Math.* **56** (1992), 243–294.
- [52] M. Uchiyama, Operators which have commutative polar decompositions, *Oper. Theory Adv. Appl.* **62** (1993), 197–208.
- [53] V. Vedral, *Introduction to Quantum Information Science*, Oxford University Press, USA 2007.
- [54] J. Weidmann, *Linear operators in Hilbert spaces*, Springer-Verlag, Berlin, Heidelberg, New York, 1980.

WYDZIAŁ MATEMATYKI I INFORMATYKI, UNIwersYTET Jagielloński, ul. Łojasiewicza 6,
PL-30348 KRAKÓW

Email address: `pawel.pietrzycki@im.uj.edu.pl`

WYDZIAŁ MATEMATYKI I INFORMATYKI, UNIwersYTET Jagielloński, ul. Łojasiewicza 6,
PL-30348 KRAKÓW

Email address: `jan.stochel@im.uj.edu.pl`