

Maximum Likelihood Recursive State Estimation using the Expectation Maximization Algorithm

Mohammad S. Ramadan^a, Robert R. Bitmead^b

^a*Department of Mechanical & Aerospace Engineering, University of California, San Diego, La Jolla CA 92093-0411, USA, and Electrical & Computer Engineering Department, San Diego State University, San Diego CA 92182, USA*

^b*Department of Mechanical & Aerospace Engineering, University of California, San Diego, La Jolla CA 92093-0411, USA.*

Abstract

A Maximum Likelihood recursive state estimator is derived for non-linear and non-Gaussian state-space models. The estimator combines a particle filter to generate the conditional density and the Expectation Maximization algorithm to compute the maximum likelihood state estimate iteratively. Algorithms for maximum likelihood state filtering, prediction and smoothing are presented. The convergence properties of these algorithms, which are inherited from the Expectation Maximization algorithm, are proven and examined in two examples. It is shown that, with randomized reinitialization, which is feasible because of the algorithm simplicity, these methods are able to converge to the Maximum Likelihood Estimate (MLE) of multimodal, truncated and skewed densities, as well as those of disjoint support.

Key words: Maximum Likelihood, State Estimation, Expectation Maximization, Particle Filter .

1 Introduction

We derive a maximum likelihood recursive state estimator for nonlinear systems based on: knowledge of the system equations and process noise and measurement noise densities; the operation of a particle filter to propagate conditional densities; and the Expectation Maximization (EM) algorithm [2,14]. If the state at time k is denoted ζ_k , and given the sequence of outputs up to time k , namely $\mathcal{Y}_{0:k} = \{y_0, y_1, \dots, y_k\}$, the algorithm provides an estimate of the maximizer of the posterior state density $p(\zeta_k | \mathcal{Y}_{0:k})$. This contrasts with Least-Squares estimators, which produce the conditional mean of this density. The contribution of the paper is to provide an easily implementable algorithm which is recursive and which yields this Maximum Likelihood Estimate (MLE). We propagate the conditional densities, both prior and posterior, using a particle filter, and the EM algorithm is used to compute the MLE of the posterior density from the particles of the prior density, the current measurement and the known noise densities. The observation is that knowledge of the noise densities permits computation of the gradient of successive functions whose maximizers are non-decreasing in their likelihood with respect to the posterior density, which follows from the desirable convergence prop-

erties of the EM algorithm. If the system under consideration has Gaussian additive noises and linear measurement equation, these successive functions are quadratic and the algorithm simplifies into a fixed-point iteration. The paper clarifies these details in relation to the particle filter, the EM algorithm and MLE.

In many scenarios, the conditional state density can: be multi-modal as when tracking a group of objects [13,19]; have skewness, limited support, one-sidedness as in some Stochastic Volatility Models [20,9]; or be governed by inequality state constraints [8], leading to truncated or disjoint support such as might arise in stochastic model Predictive Control. The justification for the choice of the conditional mean as the point estimate of a filter may be weakened or render an estimate which is infeasible under the density. In such scenarios, the MLE may have more reasonable grounds for use. Since finding the MLE requires optimization over the posterior density, the recursive procedure involves two stages: the propagation of the posterior density, for which we rely on a particle filter; and, an iterative optimization to find the maximizer. However, the density provided by the particle filter is a collection of point masses, usually with the same singleton mass, and so is not suited to Newton-like optimization. A novelty of this paper and its algorithms is that the previous time's density plus the current measurement are combined with the known continuous densities of the process and measurement noises to yield a gradient based

Email addresses: msramada@eng.ucsd.edu (Mohammad S. Ramadan), rbitmead@eng.ucsd.edu (Robert R. Bitmead).

maximizer. Effectively, this procedure yields a smooth interpolation of the particle filter’s posterior density and of its gradient.

In [11], a piecewise continuous cumulative distribution function is constructed from the weighted particles approximating the filtered state density $p(\zeta_k|\mathcal{Y}_{0:k})$, limited to the case when the state-space is one-dimensional. In [1], this solution is extended to a multi-dimensional state-space. However, it has a computational complexity of $\mathcal{O}(N^2)$, where N is the number of particles, and a nonstandard particle filtering scheme, as reported in [4]. Earlier, in the construction of the auxiliary particle filter [15], it was shown that it is possible to construct a smooth approximation of $p(\zeta_k|\mathcal{Y}_{0:k})$ under the name of “the empirical filtered density” using the weighted particles approximating $p(\zeta_{k-1}|\mathcal{Y}_{0:k-1})$ and the current measurement. We extend this construction with the computation of an approximate gradient, as shall be explained shortly. This constructed smooth density could be directly maximized using various search methods and the new calculation permits gradient search for the MLE. We introduce the EM algorithm as an iterative procedure to achieve this, perhaps with multiple initial starting values in the multimodal case. This yields a sequence of estimates with non-decreasing likelihood which converges to a stationary point [22,2]. This will be supported by computational examples in Sections 3.4 and 3.5. In effect, the EM algorithm smooths and recenters the empirical filtered density at the E-step before maximizing at the M-step. The iterations refine and localize the smoothing to approach the MLE. Ultimately, this is limited by the quality of the empirical density generated by the particle filter. Without such smoothing, it is not possible to find the maximizer of the particle filter’s point-mass density.

The EM algorithm [2] is versatile with many variations [12,21,5] and applications primarily in parameter estimation. It consists of two major steps: the expectation (E) step, in which an approximant to the log-likelihood function is constructed, followed by the maximization (M) step, which seeks to find the maximizer of this function. Hence, EM iterations deliver a sequence of functions for which their maximizers are non-decreasing with respect to the likelihood function under consideration [22]. Equipped with Monte-Carlo particle methods, the EM algorithm is widely used in parameter estimation for nonlinear non-Gaussian systems [18,16,10,6]. We examine the use of EM in the context of state estimation; to our knowledge this is a novel application and different in principle, since, unlike parameters, the components of the state are dynamically changing according to the tailored system model.

The particle filter [3], itself, is a common feature of nonlinear estimation and operates by propagating the conditional densities as characterized by a collection of samples, called particles, distributed with this density. For a basic particle filter, the time update pass, i.e. posterior-to-prior densities, uses the state equation and samples from the process noise density. This is followed by the measurement update, prior-

to-posterior densities, relying on weighting and possibly resampling these particles based on the output equation and the output noise density.

In this paper, we use the weighted particles’ approximation of $p(\zeta_{k-1}|\mathcal{Y}_{0:k-1})$ and the current observation y_k with the EM algorithm to construct a sequence of functions of ζ_k , whose maximizers are non-decreasing in their likelihood with respect to $p(\zeta_k|\mathcal{Y}_{0:k})$. In Section 2, the underlying assumptions on the state-space model under consideration are elucidated. In Section 3, the EM state filtering (EMSF) algorithm is introduced, and implemented in two examples 3.4 and 3.5. First, it is applied to state filtering of a linear Gaussian state-space model, and since the conditional mean is equivalent to the conditional mode of such systems [17], it is compared to the Kalman filter solution, and shown to be convergent to the conditional mode (the global maximizer) from a random initial guess. This establishes the efficacy of the algorithm and permits statements concerning its complexity. In the second example, it is used to track the MLE of a multi-modal state distribution of a nonlinear state-space model. It is also shown, in Section 3.6 that this new EM state filter (EMSF) algorithm reduces to a convergent fixed-point iteration method under mild assumption. The extensions of this algorithm to the contexts of state prediction and smoothing are provided in Sections 4 and 5.

2 Problem Statement

Consider the following nonlinear dynamical system affine in the noises ¹:

$$\zeta_{k+1} = f(\zeta_k) + w_k, \zeta_k \in \mathbb{R}^{r_\zeta}, \quad (1a)$$

$$y_k = g(\zeta_k) + v_k, y_k \in \mathbb{R}^{r_y}, \quad (1b)$$

$$w_k \sim W_k(\cdot), v_k \sim V_k(\cdot),$$

$$\zeta_0 \sim p_0(\cdot)$$

where,

- the stochastic processes $\{w_k\}$ and $\{v_k\}$ are white and independent, and their probability density functions, $W_k(\cdot)$ and $V_k(\cdot)$ are known and continuously differentiable for all k ,
- ζ_k is the state vector, with initial value, ζ_0 , having a known initial distribution which is independent of $\{w_k\}$ and $\{v_k\}$,
- y_k is the measurement vector,
- the functions f and g are known continuously differentiable functions.

Given the observation sequence, $\mathcal{Y}_{0:k} = \{y_0, y_1, \dots, y_k\}$, the problem is to find the MLE of the state-vector ζ_k using the log-likelihood function $\log p(\zeta_k|\mathcal{Y}_{0:m})$, for $m = k$ (filtering), $m > k$ (smoothing), and $m < k$ (prediction).

¹ The notation \sim here means “is distributed as the density on the right”.

3 ML state filtering by the expectation maximization algorithm

Given, at time k :

- the problem statement and assumptions as in Section 2,
- the measurement y_k ,
- the filtered density at time $k-1$, i.e. $p(\zeta_{k-1}|\mathcal{Y}_{0:k-1})$,
- initial time- k state estimate ζ_k^0 ,

we set iteration counter $i = 0$ and seek to find ζ_k^{i+1} such that $\log p(\zeta_k^{i+1}|\mathcal{Y}_{0:k}) \geq \log p(\zeta_k^i|\mathcal{Y}_{0:k})$ [22]. This is achieved by two major steps: the expectation step (E-step), followed by the maximization step (M-step). We continue with these iterations in i until we satisfy a convergence criterion, which is guaranteed by the non-decreasing property. At this stage, we either restart with a different initial condition, ζ_k^0 , or move to time $k+1$.

3.1 E-step

The E-step is performed by evaluating the following expectation,

$$\begin{aligned} Q_\zeta(\zeta_k, \zeta_k^i) &= \mathbb{E}_{\zeta_{k-1}|\zeta_k^i, \mathcal{Y}_{0:k}} \{ \log p(\zeta_k|\zeta_{k-1}, \mathcal{Y}_{0:k}) | \zeta_k^i, \mathcal{Y}_{0:k} \} \\ &= \int \log p(\zeta_k|\zeta_{k-1}, \mathcal{Y}_{0:k}) p(\zeta_{k-1}|\zeta_k^i, \mathcal{Y}_{0:k}) d\zeta_{k-1} \end{aligned} \quad (2)$$

whose calculation in terms of available densities will be presented shortly in the following theorem.

Before presenting the theorem, we first derive a lemma detailing a conditional independence result, which will be used repeatedly in the proofs and derivations.

Lemma 1 *For system (1) at time $n \geq k$, y_n conditioned on ζ_k is conditionally independent from both $\zeta_{0:k-1} = \{\zeta_0, \zeta_1, \dots, \zeta_{k-1}\}$ and $\mathcal{Y}_{0:k-1}$, that is*

$$p(y_n|\zeta_k, \zeta_{0:k-1}, \mathcal{Y}_{0:k-1}) = p(y_n|\zeta_k)$$

Proof. From (1) and for $n \geq k$, y_n is a function of ζ_k , $w_{k:n-1}$ and v_n . Since $w_{k:n-1}$ and v_n are independent from $\zeta_{0:k-1}$ and $\mathcal{Y}_{0:k-1}$, we have the result. \square

Theorem 1 *Given the dynamical system (1), the filtered density $p(\zeta_{k-1}|\mathcal{Y}_{0:k-1})$, the current measurement y_k , and ζ_k^i , the integral in (2) is, apart from a positive multiplicative constant and an additive constant both independent from ζ_k , equal to*

$$\begin{aligned} Q_\zeta(\zeta_k, \zeta_k^i) &= \\ &\int \left[\log V_k(y_k - g(\zeta_k)) + \log W_{k-1}(\zeta_k - f(\zeta_{k-1})) \right] \times \\ &W_{k-1}(\zeta_k^i - f(\zeta_{k-1})) p(\zeta_{k-1}|\mathcal{Y}_{0:k-1}) d\zeta_{k-1}, \end{aligned} \quad (3)$$

and its gradient is given by

$$\begin{aligned} \nabla_{\zeta_k} Q_\zeta(\zeta_k, \zeta_k^i) &= \int \left[\nabla_{\zeta_k} \log V_k(y_k - g(\zeta_k)) + \right. \\ &\left. \nabla_{\zeta_k} \log W_{k-1}(\zeta_k - f(\zeta_{k-1})) \right] \times \\ &W_{k-1}(\zeta_k^i - f(\zeta_{k-1})) p(\zeta_{k-1}|\mathcal{Y}_{0:k-1}) d\zeta_{k-1}. \end{aligned} \quad (4)$$

Proof. We use Bayes rule and Lemma 1, to yield

$$\begin{aligned} p(\zeta_k|\zeta_{k-1}, \mathcal{Y}_{0:k}) &= \frac{p(y_k|\zeta_k, \zeta_{k-1}, \mathcal{Y}_{0:k-1}) p(\zeta_k|\zeta_{k-1}, \mathcal{Y}_{0:k-1})}{p(y_k|\zeta_{k-1}, \mathcal{Y}_{0:k-1})} \\ &= \frac{p(y_k|\zeta_k) p(\zeta_k|\zeta_{k-1})}{p(y_k|\zeta_{k-1})}. \end{aligned}$$

Thus,

$$\begin{aligned} \log p(\zeta_k|\zeta_{k-1}, \mathcal{Y}_{0:k}) &= \log p(y_k|\zeta_k) + \log p(\zeta_k|\zeta_{k-1}) \\ &- \log p(y_k|\zeta_{k-1}) \\ &= \log V_k(y_k - g(\zeta_k)) \\ &+ \log W_{k-1}(\zeta_k - f(\zeta_{k-1})) - \tilde{C} \end{aligned} \quad (5)$$

where $\tilde{C} = \log p(y_k|\zeta_{k-1})$ is independent from ζ_k and so integrates to an additive constant in the expectation (2).

Similarly, by Bayes rule and Lemma 1,

$$\begin{aligned} p(\zeta_{k-1}|\zeta_k^i, \mathcal{Y}_{0:k}) &= p(\zeta_{k-1}|\zeta_k^i, \mathcal{Y}_{0:k-1}, y_k), \\ &= \frac{p(y_k|\zeta_k^i, \zeta_{k-1}, \mathcal{Y}_{0:k-1}) p(\zeta_{k-1}|\zeta_k^i, \mathcal{Y}_{0:k-1})}{p(y_k|\zeta_k^i, \mathcal{Y}_{0:k-1})}, \\ &= \frac{p(y_k|\zeta_k^i) p(\zeta_{k-1}|\zeta_k^i, \mathcal{Y}_{0:k-1})}{p(y_k|\zeta_k^i)}, \\ &= p(\zeta_{k-1}|\zeta_k^i, \mathcal{Y}_{0:k-1}), \end{aligned}$$

and, applying Bayes rule again,

$$p(\zeta_{k-1}|\zeta_k^i, \mathcal{Y}_{0:k}) = \frac{p(\zeta_k^i|\zeta_{k-1}) p(\zeta_{k-1}|\mathcal{Y}_{0:k-1})}{p(\zeta_k^i|\mathcal{Y}_{0:k-1})}.$$

Hence,

$$\begin{aligned} p(\zeta_{k-1}|\zeta_k^i, \mathcal{Y}_{0:k}) &= \\ &\frac{W_{k-1}(\zeta_k^i - f(\zeta_{k-1})) p(\zeta_{k-1}|\mathcal{Y}_{0:k-1})}{\hat{C}} \end{aligned} \quad (6)$$

where $\hat{C} = p(\zeta_k^i|\mathcal{Y}_{0:k-1})$ is independent from ζ_{k-1} and leads to a positive multiplicative constant in the expectation (2).

Substituting (5) and (6) into (2) and neglecting the additive

and positive multiplicative constants above yields

$$\begin{aligned} Q_\zeta(\zeta_k, \zeta_k^i) &= \int \left\{ [\log p(y_k|\zeta_k) + \log p(\zeta_k|\zeta_{k-1})] \times \right. \\ &\quad \left. p(\zeta_k^i|\zeta_{k-1})p(\zeta_{k-1}|\mathcal{Y}_{0:k-1}) \right\} d\zeta_{k-1} \\ &= \int \left\{ [\log V_k(y_k - g(\zeta_k)) + \log W_{k-1}(\zeta_k - f(\zeta_{k-1}))] \times \right. \\ &\quad \left. W_{k-1}(\zeta_k^i - f(\zeta_{k-1}))p(\zeta_{k-1}|\mathcal{Y}_{0:k-1}) \right\} d\zeta_{k-1}. \end{aligned}$$

The smoothness assumptions in Section 2 permit exchanging differentiation and integration to arrive at the gradient formula.

$$\begin{aligned} \nabla_{\zeta_k} Q_\zeta(\zeta_k, \zeta_k^i) &= \nabla_{\zeta_k} \int \left\{ [\log V_k(y_k - g(\zeta_k)) + \right. \\ &\quad \left. \log W_{k-1}(\zeta_k - f(\zeta_{k-1}))] \times \right. \\ &\quad \left. W_{k-1}(\zeta_k^i - f(\zeta_{k-1}))p(\zeta_{k-1}|\mathcal{Y}_{0:k-1}) \right\} d\zeta_{k-1} \\ &= \int \left\{ [\nabla_{\zeta_k} \log V_k(y_k - g(\zeta_k)) \right. \\ &\quad \left. + \nabla_{\zeta_k} \log W_{k-1}(\zeta_k - f(\zeta_{k-1}))] \times \right. \\ &\quad \left. W_{k-1}(\zeta_k^i - f(\zeta_{k-1}))p(\zeta_{k-1}|\mathcal{Y}_{0:k-1}) \right\} d\zeta_{k-1}. \end{aligned}$$

□

The gradient $\nabla_{\zeta_k} Q_\zeta(\zeta_k, \zeta_k^i)$ can be used within a gradient search algorithm to evaluate the M-step, which follows immediately.

3.2 M-step

The M-step maximizes the expectation in the E-step w.r.t ζ_k , choosing ζ_k^{i+1} to be the maximizer,

$$\zeta_k^{i+1} = \arg \max_{\zeta_k \in \mathbb{R}^r} Q_\zeta(\zeta_k, \zeta_k^i) \quad (7)$$

This choice of ζ_k^{i+1} will be shown in Theorem 2 to be non-decreasing in its likelihood compared to ζ_k^i . A proof of the general statement of EM algorithm can be found in [22]. We present the proof in the context of state filtering.

Theorem 2 *The sequence $\{\zeta_k^i\}_{i \geq 0}$, commencing at $\zeta_k^0 \in \mathbb{R}^r$ and satisfying the recursion (7), has the property that $\{\log p(\zeta_k^i|\mathcal{Y}_{0:k})\}_{i \geq 0}$ is monotonically non-decreasing.*

Proof. The log-likelihood function $\log p(\zeta_k|\mathcal{Y}_{0:k})$ can be expressed as

$$\begin{aligned} \log p(\zeta_k|\mathcal{Y}_{0:k}) &= \log p(\zeta_k, \zeta_{k-1}|\mathcal{Y}_{0:k}) - \log p(\zeta_{k-1}|\mathcal{Y}_{0:k}, \zeta_k) \\ &= \log p(\zeta_k|\zeta_{k-1}, \mathcal{Y}_{0:k}) + \log p(\zeta_{k-1}|\mathcal{Y}_{0:k}) \\ &\quad - \log p(\zeta_{k-1}|\mathcal{Y}_{0:k}, \zeta_k). \end{aligned}$$

Taking the expectation with $p(\zeta_{k-1}|\zeta_k^i, \mathcal{Y}_{0:k})$ as in (2),

$$\begin{aligned} \log p(\zeta_k|\mathcal{Y}_{0:k}) &= Q_\zeta(\zeta_k, \zeta_k^i) + \mathbb{E}_{\zeta_{k-1}|\zeta_k^i, \mathcal{Y}_{0:k}} \{\log p(\zeta_{k-1}|\mathcal{Y}_{0:k})\} \\ &\quad - \mathbb{E}_{\zeta_{k-1}|\zeta_k^i, \mathcal{Y}_{0:k}} \{\log p(\zeta_{k-1}|\mathcal{Y}_{0:k}, \zeta_k)\}. \end{aligned}$$

Hence, the log-likelihood of ζ_k^i ,

$$\begin{aligned} \log p(\zeta_k^i|\mathcal{Y}_{0:k}) &= Q_\zeta(\zeta_k^i, \zeta_k^i) + \mathbb{E}_{\zeta_{k-1}|\zeta_k^i, \mathcal{Y}_{0:k}} \{\log p(\zeta_{k-1}|\mathcal{Y}_{0:k})\} \\ &\quad - \mathbb{E}_{\zeta_{k-1}|\zeta_k^i, \mathcal{Y}_{0:k}} \{\log p(\zeta_{k-1}|\mathcal{Y}_{0:k}, \zeta_k^i)\}. \end{aligned}$$

Subtracting these two equations,

$$\begin{aligned} \log p(\zeta_k|\mathcal{Y}_{0:k}) - \log p(\zeta_k^i|\mathcal{Y}_{0:k}) &= Q_\zeta(\zeta_k, \zeta_k^i) - Q_\zeta(\zeta_k^i, \zeta_k^i) \\ &\quad + \mathbb{E}_{\zeta_{k-1}|\zeta_k^i, \mathcal{Y}_{0:k}} \{\log p(\zeta_{k-1}|\mathcal{Y}_{0:k}, \zeta_k^i) - \log p(\zeta_{k-1}|\mathcal{Y}_{0:k}, \zeta_k)\} \end{aligned}$$

where the term,

$$\begin{aligned} &\mathbb{E}_{\zeta_{k-1}|\zeta_k^i, \mathcal{Y}_{0:k}} \{\log p(\zeta_{k-1}|\mathcal{Y}_{0:k}, \zeta_k^i) - \log p(\zeta_{k-1}|\mathcal{Y}_{0:k}, \zeta_k)\} \\ &= \int_{\mathbb{R}^r} \frac{\log p(\zeta_{k-1}|\mathcal{Y}_{0:k}, \zeta_k^i)}{\log p(\zeta_{k-1}|\mathcal{Y}_{0:k}, \zeta_k)} p(\zeta_{k-1}|\zeta_k^i, \mathcal{Y}_{0:k}) d\zeta_{k-1} \\ &\geq 0. \end{aligned}$$

Since the right-hand side is the Kullback-Leibler divergence metric between these two densities, this quantity is non-negative. (See Example 7.7 in [7].) Thus,

$$\log p(\zeta_k|\mathcal{Y}_{0:k}) - \log p(\zeta_k^i|\mathcal{Y}_{0:k}) \geq Q_\zeta(\zeta_k, \zeta_k^i) - Q_\zeta(\zeta_k^i, \zeta_k^i).$$

It follows from the definition of ζ_k^{i+1} as the maximizer of $Q_\zeta(\zeta_k, \zeta_k^i)$ that

$$Q_\zeta(\zeta_k^{i+1}, \zeta_k^i) - Q_\zeta(\zeta_k^i, \zeta_k^i) \geq 0.$$

Therefore,

$$\log p(\zeta_k^{i+1}|\mathcal{Y}_{0:k}) - \log p(\zeta_k^i|\mathcal{Y}_{0:k}) \geq 0,$$

which completes the proof. □

Corollary 1 *If the log-likelihood function $\log p(\zeta_k|\mathcal{Y}_{0:k})$ is unimodal with the MLE being the only stationary point, then $\{\zeta_k^i\}_{i \geq 0}$ converges to this MLE.*

This parallels Corollary 1 in [22].

The EM algorithm for ML state filtering at time-step k follows.

Algorithm 1

- 1: Given the current measurement y_k , the prior filtered density $p(\zeta_{k-1}|\mathcal{Y}_{0:k-1})$ and prior MLE ζ_{k-1}^* , set $i \leftarrow 0$ and let $\zeta_k^0 = \zeta_k^i = f(\zeta_{k-1}^*) + \mathbb{E}w_k$;
- 2: Evaluate the integral:

$$Q_\zeta(\zeta_k, \zeta_k^i) = \int \left\{ \left[\log V_k(y_k - g(\zeta_k)) + \log W_{k-1}(\zeta_k - f(\zeta_{k-1})) \right] \times W_{k-1}(\zeta_k^i - f(\zeta_{k-1})) p(\zeta_{k-1}|\mathcal{Y}_{0:k-1}) \right\} d\zeta_{k-1} \quad (8)$$

- 3: Evaluate ζ_k^{i+1} :

$$\zeta_k^{i+1} = \arg \max_{\zeta_k \in \mathbb{R}^r} Q_\zeta(\zeta_k, \zeta_k^i) \quad (9)$$

- 4: If $\{\zeta_k^i\}_{i \geq 0}$ satisfies a convergence criterion: set $\zeta_k^* = \zeta_k^i$ and terminate. Else: $i \leftarrow i + 1$ and go to Step 2.
-

As guaranteed by Theorem 2, Algorithm 1 generates a sequence of estimates $\{\zeta_k^i\}_{i \geq 0}$ which is non-decreasing in its likelihoods.

Although the densities V_k and W_k are assumed to be known, the integral (8) in Algorithm 1 requires the knowledge of $p(\zeta_{k-1}|\mathcal{Y}_{0:k-1})$ as well. In general, outside the linear Gaussian case, such an integral cannot be analytically evaluated. Hence, a numerical approximation has to be considered. In this paper, we turn to using a particle filter.

3.3 EM with a particle filter

The EM algorithm, so far, relies on the provision of the filtered density $p(\zeta_{k-1}|\mathcal{Y}_{0:k-1})$ for the calculation at the E-step. A particle filter [3] provides a version of this information through a set of particles, i.e. a collection of values of $\{\tilde{\zeta}_{k-1}^j\}_{j=1}^N$, and possibly their corresponding importance weights $\{\omega_{k-1|k-1}^j\}_{j=1}^N$. The density is approximated by

$$p(\zeta_{k-1}|\mathcal{Y}_{0:k-1}) \approx \sum_{j=1}^N \omega_{k-1|k-1}^j \delta(\zeta_{k-1} - \tilde{\zeta}_{k-1}^j), \quad (10)$$

where $\delta(\cdot)$ is the delta function. If the particles are produced by resampling then each of the importance weights $\omega_{k-1|k-1}^j = 1/N$. Using these particles in (10), the expectation $Q_\zeta(\zeta_k, \zeta_k^i)$, in (8), can be replaced by its particle approximation $\hat{Q}_\zeta^N(\zeta_k, \zeta_k^i)$. Algorithm 1 can then be recast using the particles for the E-step as follows.

Algorithm 2 EM State Filter (EMSF) Algorithm

- 1: Given y_k , the particles $\{\tilde{\zeta}_{k-1}^j\}_{j=1}^N$, their normalized importance weights $\{\omega_{k-1|k-1}^j\}_{j=1}^N$ and ζ_{k-1}^* , set $i \leftarrow 0$ and let $\zeta_k^i = f(\zeta_{k-1}^*) + \mathbb{E}w_k$;
- 2: Evaluate the summation:

$$\hat{Q}_\zeta^N(\zeta_k, \zeta_k^i) = \sum_{j=1}^N \left[\log V_k(y_k - g(\zeta_k)) + \log W_{k-1}(\zeta_k - f(\tilde{\zeta}_{k-1}^j)) \right] \times W_{k-1}(\zeta_k^i - f(\tilde{\zeta}_{k-1}^j)) \omega_{k-1|k-1}^j \quad (11)$$

- 3: Evaluate ζ_k^{i+1} :

$$\zeta_k^{i+1} = \arg \max_{\zeta_k \in \mathbb{R}^r} \hat{Q}_\zeta^N(\zeta_k, \zeta_k^i) \quad (12)$$

- 4: If $\{\zeta_k^i\}_{i \geq 0}$ satisfies a convergence criterion: set $\zeta_k^* = \zeta_k^i$ and terminate. Else: $i \leftarrow i + 1$ and go to Step 2.
-

We apply EMSF in the following computed examples.

3.4 Example 1: Linear Gaussian state-space model

In the linear Gaussian state-space model case, all random variables under consideration are Gaussian. Hence, the conditional mean is equivalent to the conditional mode [17]. In the following example, the Kalman filter, which is designed to track the propagation of the conditional mean, is used in comparison with the EMSF. It will be shown that the EMSF, starting from a random initialization ζ_k^0 , converges to the Kalman Filter solution, which is guaranteed by Corollary 1.

Consider the following model,

$$\begin{aligned} \zeta_{k+1} &= F\zeta_k + w_k, \\ y_k &= H\zeta_k + v_k, \\ v_k &\sim \mathcal{N}(0, R), \quad w_k \sim \mathcal{N}(0, S) \\ \zeta_0 &\sim \mathcal{N}(\zeta_{0|-1}, \Sigma_{0|-1}) \end{aligned} \quad (13)$$

where the sequences $\{w_k\}$ and $\{v_k\}$ and the random variable ζ_0 follow the independence and whiteness assumptions as in Section 2. The term $\mathcal{N}(\mu, \Sigma)$ denotes the multivariate normal density of mean μ and covariance Σ , and $\mathcal{N}(x|\mu, \Sigma)$ denotes the value of this density when evaluated at x .

We use the following particle approximation,

$$p(\zeta_{k-1}|\mathcal{Y}_{0:k-1}) \approx \sum_{j=1}^N \omega_{k-1|k-1}^j \delta(\zeta_{k-1} - \tilde{\zeta}_{k-1}^j). \quad (14)$$

The summation in Algorithm 2, ignoring constant terms superfluous to the maximization, can be simplified to,

$$\begin{aligned} \hat{Q}_\zeta^N(\zeta_k, \zeta_k^i) = & \sum_{j=1}^N \left[- (y_k - H\zeta_k)^T R^{-1} (y_k - H\zeta_k) \right. \\ & \left. - (\zeta_k - F\tilde{\zeta}_{k-1}^j)^T S^{-1} (\zeta_k - F\tilde{\zeta}_{k-1}^j) \right] \\ & \times \lambda^j \omega_{k-1|k-1}^j \end{aligned} \quad (15)$$

where $\lambda^j = W_{k-1}(\zeta_k^i - F\tilde{\zeta}_{k-1}^j) = \mathcal{N}(\zeta_k^i - F\tilde{\zeta}_{k-1}^j | 0, S)$. Since (15) is quadratic in ζ_k , the maximizer ζ_k^{i+1} can be found analytically,

$$\zeta_k^{i+1} \text{ is such that } \nabla_{\zeta_k} Q_\zeta^N(\zeta_k^{i+1}, \zeta_k^i) = 0.$$

Hence,

$$\begin{aligned} \zeta_k^{i+1} = & \left[H^T R^{-1} H + S^{-1} \right]^{-1} \frac{1}{\sum_{l=1}^N \lambda^l \omega_{k-1|k-1}^l} \times \\ & \left\{ H^T R^{-1} y_k + S^{-1} F \sum_{j=1}^N \tilde{\zeta}_{k-1}^j \lambda^j \omega_{k-1|k-1}^j \right\}, \end{aligned} \quad (16)$$

or, using the matrix inversion lemma,

$$\begin{aligned} \zeta_k^{i+1} = & \left[F \sum_{j=1}^N \tilde{\zeta}_{k-1}^j \lambda^j \omega_{k-1|k-1}^j + \right. \\ & \left. \mathcal{B}(y_k - HF \sum_{j=1}^N \tilde{\zeta}_{k-1}^j \lambda^j \omega_{k-1|k-1}^j) \right] \frac{1}{\sum_{l=1}^N \lambda^l \omega_{k-1|k-1}^l} \end{aligned} \quad (17)$$

with

$$\mathcal{B} = SH^T (HSH^T + R)^{-1}.$$

Note that, while (17) bears a passing resemblance to the Kalman filter recursion as (16) does to the information filter, the λ^j terms and $\hat{Q}_\zeta^N(\zeta_k, \zeta_k^i)$ in (15) are dependent on the previous iterate ζ_k^i and the update follows the maximizer.

Now take

$$\begin{aligned} \zeta_k = & \begin{bmatrix} \zeta_{k,1} \\ \zeta_{k,2} \\ \zeta_{k,3} \end{bmatrix}, \quad F = \begin{bmatrix} 0.66 & -1.31 & -1.11 \\ 0.07 & 0.73 & -0.06 \\ 0.00 & 0.08 & 0.80 \end{bmatrix}, \\ H = & [0 \ 1 \ 1], \quad S = \begin{bmatrix} 0.2 & 0 & 0 \\ 0 & 0.3 & 0 \\ 0 & 0 & 0.5 \end{bmatrix}, \quad R = 0.1, \\ \zeta_{0|-1} = & \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}, \quad \Sigma_{0|-1} = \begin{bmatrix} 0.3 & 0 & 0 \\ 0 & 0.3 & 0 \\ 0 & 0 & 0.3 \end{bmatrix}. \end{aligned}$$

A simulation over 100 time-steps, with $N = 2000$ particles, is conducted with the results plotted in Figure 1. The EMSF recursion at each time-step k , as described by (17), is initialized by $\zeta_k^0 \sim \mathcal{N}(0, I_{3 \times 3})$. The results show that the EMSF converges to the Kalman Filter solution, which property follows from Corollary 1. Figure 2, depicts the convergence of




Fig. 1. The Kalman Filter solution in black, and the EMSF solution in orange dots.

the iterates at three individual times of $\zeta_{k,1}$, where the algorithm stops whenever the maximum absolute relative error is below 0.005, i.e. $\max_{q \in \{1,2,3\}} |(\zeta_{k,q}^{i+1} - \zeta_{k,q}^i) / \zeta_{k,q}^{i+1}| \leq 0.5\%$.

The EMSF algorithm is of $\mathcal{O}(N)$ computational complexity, per iteration, with N the number of particles. Ten simulations were conducted using MATLAB, an uncompiled interpretive program, on a computer with a 2.5 GHz Intel Core i5-7200U processor and 8.00 GB of RAM. The running times for the one hundred time points averaged over the ten runs were: 3 milliseconds for the Kalman filter, 3.4 seconds for the bootstrap particle filter, and 26.4 seconds for EMSF including the bootstrap particle filter.




Fig. 2. The convergence of the sequences $\{\zeta_{k,1}^i : i \geq 0\}$ for $k = 15, 55, 90$ in green, orange and purple, respectively.

3.5 Example 2: Nonlinear state-space model with multimodal/skewed state distribution

In this example, the EMSF is tested with a nonlinear state-space model with highly skewed and/or multimodal state densities. The EMSF solution is compared to the conditional mean of the bootstrap particle filter. This example also serves as an introduction to the next section, in which the EMSF is shown to reduce to a convergent fixed-point iteration for an important class of nonlinear systems.

Consider the following nonlinear state-space model,

$$\begin{aligned} \zeta_{k+1} &= \alpha_k \tanh(\pi \zeta_k) + w_k \\ y_k &= \frac{1}{2} \zeta_k + v_k \\ w_k &\sim \mathcal{N}(0, \frac{1}{5}), v_k \sim \mathcal{N}(0, 1) \\ \zeta_0 &\sim \mathcal{N}(0, 1) \end{aligned} \quad (18)$$

where $\alpha_k = (1 + 0.5 \sin(2\pi k/20))$. The summation (8) in Algorithm 2, after ignoring constant terms, can be simplified to,

$$\hat{Q}_\zeta^N(\zeta_k, \zeta_k^i) = \sum_{j=1}^N \left[- (y_k - \frac{1}{2} \zeta_k)^2 - 5(\zeta_k - \alpha_{k-1} \tanh(\pi \tilde{\zeta}_{k-1}^j))^2 \right] \times \lambda^j \omega_{k-1|k-1}^j. \quad (19)$$

where $\lambda^j = \mathcal{N}(\zeta_k^i - \alpha_{k-1} \tanh(\pi \tilde{\zeta}_{k-1}^j) | 0, 1/5)$. Since (19)

is quadratic in ζ_k , the unique maximizer is given by

$$\zeta_k^{i+1} = \frac{1}{10.5} \frac{1}{\sum_{l=1}^N \lambda^l \omega_{k-1|k-1}^l} \times \sum_{l=1}^N [y_k + 10\alpha_k \tanh(\pi \tilde{\zeta}_{k-1}^l) \lambda^l \omega_{k-1|k-1}^l]. \quad (20)$$

The simulation is carried out over 40 time-steps, and $N = 500$ particles. The EMSF, at each time k , is defined by the recursion (20) and is initialized by 5 independent initial guesses $\zeta_k^0 \sim \mathcal{U}[-2, 2]$. The number of iterations is set to be ≤ 10 , and the point of highest final likelihood is taken over the five initial conditions. Notice that

$$\begin{aligned} p(\zeta_k | \mathcal{Y}_{0:k}) &= p(\zeta_k | y_k, \mathcal{Y}_{0:k-1}) \\ &\propto p(y_k | \zeta_k) \cdot p(\zeta_k | \mathcal{Y}_{0:k-1}) \\ &= p(y_k | \zeta_k) \int p(\zeta_k | \zeta_{k-1}) p(\zeta_{k-1} | \mathcal{Y}_{0:k-1}) d\zeta_{k-1}. \end{aligned}$$

By using the particle approximation of the filtered density $p(\zeta_{k-1} | \mathcal{Y}_{0:k-1})$, we have

$$p(\zeta_k | \mathcal{Y}_{0:k}) \approx p(y_k | \zeta_k) \sum_{j=1}^N p(\zeta_k | \tilde{\zeta}_{k-1}^j) \omega_{k-1|k-1}^j$$

which is the empirical filtered density, and can be used to compare the likelihoods of the convergence points resulting from multiple initial guesses. For this example,

$$\begin{aligned} p(\zeta_k | \mathcal{Y}_{0:k}) &\approx \mathcal{N}(y_k - \frac{1}{2} \zeta_k | 0, 1) \times \\ &\sum_{j=1}^N \mathcal{N}(\zeta_k - \alpha_{k-1} \tanh(\pi \tilde{\zeta}_{k-1}^j) | 0, \frac{1}{5}) \omega_{k-1|k-1}^j \end{aligned}$$

where the particles $\{\tilde{\zeta}_{k-1}^j\}$ and their weights $\{\omega_{k-1|k-1}^j\}$ follow the definition as in the EMSF algorithm. The results are shown in Figures 3 to 5.

In this example, the EMSF reduces to a fixed-point iteration. This leads to the following result for state-space models with Gaussian noises and linear measurement equations, in which this simplification of EMSF is possible.

3.6 Gaussian state-space models with linear measurement equations

In the case of a nonlinear state-space model with linear measurement equation and affine Gaussian noises, Algorithm 2 reduces to a fixed-point iteration method which is convergent to a stationary point of the empirical filtered density.

Theorem 3 For the nonlinear state-space system (1) defined in Section 2 with $g(\zeta_k) = H\zeta_k$ for $H \in \mathbb{R}^{r_y \times r_\zeta}$ and $w_k \sim$




Fig. 3. The shading represents the empirical density function (up to a positive multiplicative constant). In black is the output of the EMSF algorithm, and in blue is the conditional mean of the particle filter.




Fig. 4. The bi-modal empirical density function of ζ_k at $k = 10$ (apart from a multiplicative constant). x_{EMSF} is the solution of the EMSF algorithm and x_{CM} is the conditional mean of the bootstrap particle filter.

$\mathcal{N}(0, S_k)$, $v_k \sim \mathcal{N}(0, R_k)$, Steps 2 and 3 of Algorithm 2 reduce to the following fixed-point iteration.

$$\zeta_k^{i+1} = \frac{1}{\sum_{l=1}^N \lambda^l \omega_{k-1|k-1}^l} \times \left[\sum_{j=1}^N f(\tilde{\zeta}_{k-1}^j) \lambda^j \omega_{k-1|k-1}^j + \mathcal{B}_k(y_k - H \sum_{j=1}^N f(\tilde{\zeta}_{k-1}^j) \lambda^j \omega_{k-1|k-1}^j) \right],$$

where

$$\mathcal{B}_k = S_{k-1} H^T (H S_{k-1} H^T + R_k)^{-1},$$

and

$$\lambda^j = \mathcal{N}(\zeta_k^i - f(\tilde{\zeta}_{k-1}^j) | 0, S_{k-1}), \quad j = 1, 2, \dots, N.$$




Fig. 5. The highly skewed empirical density function of ζ_k at $k = 25$. x_{EMSF} is the solution of the EMSF algorithm and x_{CM} is the conditional mean of the bootstrap particle filter.

Proof. The summation (8) in Algorithm 2 reduces to

$$\hat{Q}_\zeta^N(\zeta_k, \zeta_k^i) = \sum_{j=1}^N \left[- (y_k - g(\zeta_k))^T R_k^{-1} (y_k - g(\zeta_k)) - (\zeta_k - f(\tilde{\zeta}_{k-1}^j))^T S_{k-1}^{-1} (\zeta_k - f(\tilde{\zeta}_{k-1}^j)) \right] \times \lambda^j \omega_{k-1|k-1}^j$$

which is quadratic in ζ_k . Hence, the unique maximizer of $Q_\zeta^N(\zeta_k, \zeta_k^i)$ can be found by setting $\nabla_{\zeta_k} Q_\zeta^N(\zeta_k, \zeta_k^i)$ to zero. Thus,

$$\zeta_k^{i+1} = \left[H^T R_k^{-1} H + S_{k-1}^{-1} \right]^{-1} \frac{1}{\sum_{l=1}^N \lambda^l \omega_{k-1|k-1}^l} \times \left\{ H^T R_k^{-1} y_k + S_{k-1}^{-1} \sum_{j=1}^N f(\tilde{\zeta}_{k-1}^j) \lambda^j \omega_{k-1|k-1}^j \right\}.$$

Applying the matrix inversion lemma, as in Section 3.4, completes the proof. \square

We have the following ML state filtering algorithm.

Algorithm 3 Fixed-Point State Filter (FPSF) Algorithm for linear measurements and affine Gaussian noises

- 1: Subject to the assumptions of Theorem 3 and given: y_k , the particles $\{\tilde{\zeta}_{k-1}^j\}_{j=1}^N$, their normalized importance weights $\{\omega_{k-1|k-1}^j\}_{j=1}^N$ and the initial guess ζ_k^0 , set $i \leftarrow 0$;
- 2: Do:

$$\zeta_k^{i+1} = \frac{1}{\sum_{l=1}^N \lambda^l \omega_{k-1|k-1}^l} \times \left[\sum_{j=1}^N f(\tilde{\zeta}_{k-1}^j) \lambda^j \omega_{k-1|k-1}^j + \mathcal{B}_k(y_k - H \sum_{j=1}^N f(\tilde{\zeta}_{k-1}^j) \lambda^j \omega_{k-1|k-1}^j) \right]$$

where

$$\mathcal{B}_k = S_{k-1} H^T (H S_{k-1} H^T + R_k)^{-1}$$

and

$$\lambda^j = \mathcal{N}(\zeta_k^i - f(\tilde{\zeta}_{k-1}^j) | 0, S_{k-1}), \quad j = 1, 2, \dots, N$$

- 3: If $\{\zeta_k^i\}_{i \geq 0}$ satisfies a convergence criterion, terminate. Else: $i \leftarrow i + 1$ and go to step 2.
-

4 ML state prediction by the expectation maximization algorithm

Given,

- the problem statement and assumptions as in Section 2,
- the particle approximation of the time- m filtered density $p(\zeta_m | \mathcal{Y}_{0:m})$, for $m \in \{1, 2, \dots, k-1\}$, i.e.

$$p(\zeta_m | \mathcal{Y}_{0:m}) \approx \sum_{j=1}^N \omega_{m|m}^j \delta(\zeta_m - \tilde{\zeta}_{m|m}^j)$$

- initial state estimate ζ_k^0 ,

the time-update of the bootstrap particle filter can be used to propagate the particles and achieve the particle approximation

$$p(\zeta_{k-1} | \mathcal{Y}_{0:m}) \approx \sum_{j=1}^N \omega_{m|m}^j \delta(\zeta_{k-1} - \tilde{\zeta}_{k-1|m}^j). \quad (21)$$

The importance weights are unchanged because no extra measurements are available after time- m . The EM algorithm can then be used to evaluate the MLE of ζ_k of the predicted density $p(\zeta_k | \mathcal{Y}_{0:m})$, using the particle approximation (21). The E-step is performed by evaluating the following expectation

tation

$$\begin{aligned} Q_{\zeta}^p(\zeta_k, \zeta_k^i) &= \mathbb{E}_{\zeta_{k-1} | \zeta_k^i, \mathcal{Y}_{0:m}} \{ \log p(\zeta_k | \zeta_{k-1}, \mathcal{Y}_{0:m}) | \zeta_k^i, \mathcal{Y}_{0:m} \}, \\ &= \int \left\{ \log p(\zeta_k | \zeta_{k-1}, \mathcal{Y}_{0:m}) \right. \\ &\quad \left. p(\zeta_{k-1} | \zeta_k^i, \mathcal{Y}_{0:m}) \right\} d\zeta_{k-1}, \end{aligned}$$

which, analogous to the proof of Theorem 1, is equivalent to

$$Q_{\zeta}^p(\zeta_k, \zeta_k^i) = \int \left\{ [\log W_{k-1}(\zeta_k - f(\zeta_{k-1}))] \times W_{k-1}(\zeta_k^i - f(\zeta_{k-1})) p(\zeta_{k-1} | \mathcal{Y}_{0:m}) \right\} d\zeta_{k-1}. \quad (22)$$

The particle approximation (21) can be used to construct $Q_{\zeta}^{p,N}$, an approximate of (22) up to additive and multiplicative constants which are independent from ζ_k ,

$$\begin{aligned} \hat{Q}_{\zeta}^{p,N}(\zeta_k, \zeta_k^i) &= \sum_{j=1}^N \left[\log W_{k-1}(\zeta_k - f(\tilde{\zeta}_{k-1|m}^j)) \right] \\ &\quad \times W_{k-1|m}(\zeta_k^i - f(\tilde{\zeta}_{k-1|m}^j)) \omega_{m|m}^j. \end{aligned}$$

And the M-step is performed by maximizing this approximate over ζ_k . Hence, the EM algorithm for ML state prediction,

Algorithm 4 EM State Predictor (EMSP) Algorithm

- 1: Given the particles $\{\tilde{\zeta}_{k-1|m}^j\}_{j=1}^N$, their normalized importance weights $\{\omega_{m|m}^j\}_{j=1}^N$ and ζ_{k-1}^* , set $i \leftarrow 0$ and let $\zeta_k^i = f(\zeta_{k-1}^*) + \mathbb{E}w_k$;
- 2: Evaluate the summation:

$$\begin{aligned} \hat{Q}_{\zeta}^{p,N}(\zeta_k, \zeta_k^i) &= \sum_{j=1}^N \left[\log W_{k-1}(\zeta_k - f(\tilde{\zeta}_{k-1|m}^j)) \right] \\ &\quad \times W_{k-1}(\zeta_k^i - f(\tilde{\zeta}_{k-1|m}^j)) \omega_{m|m}^j \quad (23) \end{aligned}$$

- 3: Evaluate ζ_k^{i+1} :

$$\zeta_k^{i+1} = \arg \max_{\zeta_k \in \mathbb{R}^r} \hat{Q}_{\zeta}^{p,N}(\zeta_k, \zeta_k^i) \quad (24)$$

- 4: If $\{\zeta_k^i\}_{i \geq 0}$ satisfies a convergence criterion: set $\zeta_k^* = \zeta_k^i$ and terminate. Else: $i \leftarrow i + 1$ and go to step 2.
-

Similar versions of Theorem 2 and Corollary 1 hold for the sequence $\{\zeta_k^i\}_{i \geq 0}$ generated by Algorithm 4, i.e. the sequence $\{\log p(\zeta_k^i | \mathcal{Y}_m)\}_{i \geq 0}$ is non-decreasing, and if $\log p(\zeta_k | \mathcal{Y}_m)$ is unimodal and its MLE is the only stationary point, then $\{\zeta_k^i\}_{i \geq 0}$ of Algorithm 4 converges to the MLE.

5 ML state smoothing by the expectation maximization algorithm

In contrast to prediction and filtering, smoothed state estimation involves considerably greater complexity and, in the MLE environment, greater demands of the particle filter. This leads to an EM algorithm which has complexity $\mathcal{O}(N^2)$. The algorithm EMSS, however, follows directly, as we now show.

Given,

- the problem statement and assumptions as in Section 2,
- the measurements sequence $\mathcal{Y}_{0:n}$, for $n > k + 1$,
- The particle approximations of the following densities:

$$p(\zeta_{k-1}|\mathcal{Y}_{0:k-1}) \approx \sum_{j=1}^N \omega_{k-1|k-1}^j \delta(\zeta_{k-1} - \tilde{\zeta}_{k-1|k-1}^j), \quad (25)$$

$$p(\zeta_k|\mathcal{Y}_{0:k}) \approx \sum_{j=1}^N \omega_{k|k}^j \delta(\zeta_k - \tilde{\zeta}_{k|k}^j), \quad (26)$$

$$p(\zeta_{k+1}|\mathcal{Y}_{0:n}) \approx \sum_{j=1}^N \omega_{k+1|n}^j \delta(\zeta_{k+1} - \tilde{\zeta}_{k+1|n}^j), \quad (27)$$

which can be supplied by a forward–backward particle smoother [3],

- initial state estimate ζ_k^0 ,

The EM algorithm can be used to evaluate the MLE of ζ_k of the smoothed density $p(\zeta_k|\mathcal{Y}_{0:n})$. The E-step is performed by evaluating the expectation

$$\begin{aligned} Q_{\zeta}^S(\zeta_k, \zeta_k^i) &= \mathbb{E}_{\zeta_{k+1}|\zeta_k^i, \mathcal{Y}_{0:n}} \{ \log p(\zeta_k|\zeta_{k+1}, \mathcal{Y}_{0:n}) | \zeta_k^i, \mathcal{Y}_{0:n} \}, \\ &= \int \log p(\zeta_k|\zeta_{k+1}, \mathcal{Y}_{0:n}) p(\zeta_{k+1}|\zeta_k^i, \mathcal{Y}_{0:n}) d\zeta_{k+1}. \end{aligned} \quad (28)$$

Where in (28), using Bayes rule 3 times and Lemma 1, the density

$$\begin{aligned} p(\zeta_k|\zeta_{k+1}, \mathcal{Y}_{0:n}) &= \frac{p(\mathcal{Y}_{k+1:n}|\zeta_{k+1}, \zeta_k, \mathcal{Y}_{0:k}) p(\zeta_k|\zeta_{k+1}, \mathcal{Y}_{0:k})}{p(\mathcal{Y}_{k+1:n}|\zeta_{k+1}, \mathcal{Y}_{0:k})}, \\ &= \frac{p(\mathcal{Y}_{k+1:n}|\zeta_{k+1}) p(\zeta_k|\zeta_{k+1}, \mathcal{Y}_{0:k})}{p(\mathcal{Y}_{k+1:n}|\zeta_{k+1})}, \\ &= p(\zeta_k|\zeta_{k+1}, \mathcal{Y}_{0:k}), \\ &= \frac{p(\zeta_{k+1}|\zeta_k) p(\zeta_k|\mathcal{Y}_{0:k})}{p(\zeta_{k+1}|\mathcal{Y}_{0:k})}, \\ &= \frac{p(\zeta_{k+1}|\zeta_k) p(y_k|\zeta_k) p(\zeta_k|\mathcal{Y}_{0:k-1})}{p(\zeta_{k+1}|\mathcal{Y}_{0:k}) p(y_k|\mathcal{Y}_{0:k-1})}. \end{aligned}$$

The term $p(y_k|\mathcal{Y}_{0:k-1})$ is independent from ζ_k and leads to

an additive constant in the expectation (28). Thus,

$$\begin{aligned} \log p(\zeta_k|\zeta_{k+1}, \mathcal{Y}_{0:n}) &\propto \log p(\zeta_{k+1}|\zeta_k) + \log p(y_k|\zeta_k) \\ &\quad + \log p(\zeta_k|\mathcal{Y}_{0:k-1}), \\ &= \log W_k(\zeta_{k+1} - f(\zeta_k)) \\ &\quad + \log V_k(y_k - g(\zeta_k)) + \log p(\zeta_k|\mathcal{Y}_{0:k-1}). \end{aligned} \quad (29)$$

The other density in (28), using the same conditional independence result and Bayes rule,

$$\begin{aligned} p(\zeta_{k+1}|\zeta_k^i, \mathcal{Y}_{0:n}) &= \frac{p(\zeta_k^i|\zeta_{k+1}, \mathcal{Y}_{0:n}) p(\zeta_{k+1}|\mathcal{Y}_{0:n})}{p(\zeta_k^i|\mathcal{Y}_{0:n})}, \\ &= \frac{p(\zeta_k^i|\zeta_{k+1}, \mathcal{Y}_{0:k}) p(\zeta_{k+1}|\mathcal{Y}_{0:n})}{p(\zeta_k^i|\mathcal{Y}_{0:n})}, \\ &= \frac{p(\zeta_{k+1}|\zeta_k^i) p(\zeta_k^i|\mathcal{Y}_{0:k}) p(\zeta_{k+1}|\mathcal{Y}_{0:n})}{p(\zeta_{k+1}|\mathcal{Y}_{0:k}) p(\zeta_k^i|\mathcal{Y}_{0:n})}, \end{aligned} \quad (30)$$

where the terms $p(\zeta_k^i|\mathcal{Y}_{0:k})$ and $p(\zeta_k^i|\mathcal{Y}_{0:n})$ are independent from ζ_{k+1} and yield a positive multiplicative constant in the expectation (28). Hence,

$$\begin{aligned} p(\zeta_{k+1}|\zeta_k^i, \mathcal{Y}_{0:n}) &\propto \frac{p(\zeta_{k+1}|\zeta_k^i) p(\zeta_{k+1}|\mathcal{Y}_{0:n})}{p(\zeta_{k+1}|\mathcal{Y}_{0:k})}, \\ &= \frac{W_k(\zeta_{k+1} - f(\zeta_k^i)) p(\zeta_{k+1}|\mathcal{Y}_{0:n})}{p(\zeta_{k+1}|\mathcal{Y}_{0:k})}. \end{aligned} \quad (31)$$

Substituting (29) and (31) in (28), after dropping the additive and the multiplicative constants, results in

$$\begin{aligned} Q_{\zeta}^S(\zeta_k, \zeta_k^i) &= \log V_k(y_k - g(\zeta_k)) + \log p(\zeta_k|\mathcal{Y}_{0:k-1}) + \\ &\quad \int \left\{ \left[\log W_k(\zeta_{k+1} - f(\zeta_k)) \right] \times \right. \\ &\quad \left. \frac{W_k(\zeta_{k+1} - f(\zeta_k^i)) p(\zeta_{k+1}|\mathcal{Y}_{0:n})}{p(\zeta_{k+1}|\mathcal{Y}_{0:k})} \right\} d\zeta_{k+1}. \end{aligned} \quad (32)$$

Notice that for the construction of (32), the densities $p(\zeta_k|\mathcal{Y}_{0:k-1})$, $p(\zeta_{k+1}|\mathcal{Y}_{0:n})$ and $p(\zeta_{k+1}|\mathcal{Y}_{0:k})$ are required. The particle approximation (26) can be used to approximate $p(\zeta_{k+1}|\mathcal{Y}_{0:k})$,

$$\begin{aligned} p(\zeta_{k+1}|\mathcal{Y}_{0:k}) &= \int p(\zeta_{k+1}|\zeta_k) p(\zeta_k|\mathcal{Y}_{0:k}) d\zeta_k \\ &\approx \sum_{j=1}^N \omega_{k|k}^j W_k(\zeta_{k+1} - f(\tilde{\zeta}_{k|k}^j)). \end{aligned} \quad (33)$$

Similarly, the density $p(\zeta_k|\mathcal{Y}_{0:k-1})$ can be approximated using the particle approximation (25),

$$\log p(\zeta_k|\mathcal{Y}_{0:k-1}) \approx \log \sum_{j=1}^N \omega_{k-1|k-1}^j W_{k-1}(\zeta_k - f(\tilde{\zeta}_{k-1|k-1}^j)). \quad (34)$$

Finally, by using the particle approximations (34) and (33) in (32), and (27) to approximate the integral term in (32), after neglecting additive and multiplicative constants, we get $\hat{Q}_\zeta^{S,N}$, the particle approximation of Q_ζ^S ,

$$\begin{aligned} \hat{Q}_\zeta^{S,N}(\zeta_k, \zeta_k^i) &= \log V_k(y_k - g(\zeta_k)) + \\ &\log \sum_{j=1}^N \omega_{k-1|k-1}^j W_{k-1}(\zeta_k - f(\tilde{\zeta}_{k-1|k-1}^j)) + \\ &\sum_{t=1}^N \left[\log W_k(\tilde{\zeta}_{k+1|n}^t - f(\zeta_k)) \right] \times \\ &\frac{W_k(\tilde{\zeta}_{k+1|n}^t - f(\zeta_k^i))}{\sum_{d=1}^N \omega_{k|k}^d W_k(\tilde{\zeta}_{k+1|n}^t - f(\tilde{\zeta}_{k|k}^d))} \omega_{k+1|n}^t. \end{aligned} \quad (35)$$

The EM algorithm for ML state smoothing,

Algorithm 5 EM State Smoother (EMSS) Algorithm

- 1: Given the sets of particles $\{\tilde{\zeta}_{k-1|k-1}^j\}_{j=1}^N$, $\{\tilde{\zeta}_{k|k}^j\}_{j=1}^N$, $\{\tilde{\zeta}_{k+1|n}^j\}_{j=1}^N$, and their corresponding normalized importance weights, set $i \leftarrow 0$ and let the initial guess be ζ_k^i ;
- 2: Evaluate:

$$\begin{aligned} \hat{Q}_\zeta^{S,N}(\zeta_k, \zeta_k^i) &= \log V_k(y_k - g(\zeta_k)) + \\ &\log \sum_{j=1}^N \omega_{k-1|k-1}^j W_{k-1}(\zeta_k - f(\tilde{\zeta}_{k-1|k-1}^j)) + \\ &\sum_{t=1}^N \left[\log W_k(\tilde{\zeta}_{k+1|n}^t - f(\zeta_k)) \right] \times \\ &\frac{W_k(\tilde{\zeta}_{k+1|n}^t - f(\zeta_k^i))}{\sum_{d=1}^N \omega_{k|k}^d W_k(\tilde{\zeta}_{k+1|n}^t - f(\tilde{\zeta}_{k|k}^d))} \omega_{k+1|n}^t. \end{aligned} \quad (36)$$

- 3: Evaluate ζ_k^{i+1} :

$$\zeta_k^{i+1} = \arg \max_{\zeta_k \in \mathbb{R}^{\gamma_\zeta}} \hat{Q}_\zeta^{S,N}(\zeta_k, \zeta_k^i) \quad (37)$$

- 4: If $\{\zeta_k^i\}_{i \geq 0}$ satisfies a convergence criterion: set $\zeta_k^* = \zeta_k^i$ and terminate. Else: $i \leftarrow i + 1$ and go to step 2.
-

6 Conclusion

This paper contributes a novel approach to incorporating the EM algorithm, along with a particle filter, into the problem of maximum likelihood state estimation. Algorithms for maximum likelihood state filtering, prediction, and smoothing are presented, together with the desirable convergence properties they inherit from the EM algorithm. Additionally, for the wide range of nonlinear state-space systems possessing linear measurement equations and affine Gaussian

noises, these algorithms reduce to derivative-free optimization through fixed-point iteration. The EMSF algorithms is tested on two examples: one using a linear third-order Gaussian system for comparison with the Kalman filter and to demonstrate; the concordance of the estimates, the computational simplicity of the Kalman filter and the computational feasibility of the EMSF combined with its attendant particle filter; the other to estimate the MLE of a state density which exhibits multi-modality and strong skewness. Further work is being pursued to understand the role of MLE state estimates in constrained control, such as Model Predictive Control where linearity and Gaussian assumptions can contravene the problem formulation.

References

- [1] David N DeJong, Roman Liesenfeld, Guilherme V Moura, Jean-François Richard, and Hariharan Dharmarajan. Efficient likelihood evaluation of state-space representations. *Review of Economic Studies*, 80(2):538–567, 2013.
- [2] Arthur P Dempster, Nan M Laird, and Donald B Rubin. Maximum likelihood from incomplete data via the EM algorithm. *Journal of the Royal Statistical Society: Series B (Methodological)*, 39(1):1–22, 1977.
- [3] Arnaud Doucet, Simon Godsill, and Christophe Andrieu. On sequential Monte Carlo sampling methods for Bayesian filtering. *Statistics and Computing*, 10(3):197–208, 2000.
- [4] Nikolas Kantas, Arnaud Doucet, Sumeetpal S Singh, Jan Maciejowski, Nicolas Chopin, et al. On particle methods for parameter estimation in state-space models. *Statistical Science*, 30(3):328–351, 2015.
- [5] Kenneth Lange. A gradient algorithm locally equivalent to the EM algorithm. *Journal of the Royal Statistical Society: Series B (Methodological)*, 57(2):425–437, 1995.
- [6] Sylvain Le Corff and Gersende Fort. Convergence of a particle-based approximation of the block online expectation maximization algorithm. *ACM Transactions on Modeling and Computer Simulation (TOMACS)*, 23(1):1–22, 2013.
- [7] Erich L Lehmann and George Casella. *Theory of Point Estimation*. Springer Science & Business Media, 2006.
- [8] Baibing Li, Cunjia Liu, and Wen-Hua Chen. An auxiliary particle filtering algorithm with inequality constraints. *IEEE Transactions on Automatic Control*, 62(9):4639–4646, 2016.
- [9] Mengheng Li and Marcel Scharth. Leverage, Asymmetry, and Heavy Tails in the High-Dimensional Factor Stochastic Volatility Model. *Journal of Business & Economic Statistics*, pages 1–17, 2020.
- [10] Andreas Lindholm and Fredrik Lindsten. Learning dynamical systems with particle stochastic approximation EM. *arXiv preprint arXiv:1806.09548*, 2018.
- [11] Sheheryar Malik and Michael K Pitt. Particle filters for continuous likelihood evaluation and maximisation. *Journal of Econometrics*, 165(2):190–209, 2011.
- [12] Xiao-Li Meng and Donald B Rubin. Maximum likelihood estimation via the ECM algorithm: A general framework. *Biometrika*, 80(2):267–278, 1993.
- [13] Lyudmila Mihaylova, Avishy Y Carmi, François Septier, Amadou Gning, Sze Kim Pang, and Simon Godsill. Overview of Bayesian sequential Monte Carlo methods for group and extended object tracking. *Digital Signal Processing*, 25:1–16, 2014.
- [14] Todd K Moon. The expectation-maximization algorithm. *IEEE Signal Processing Magazine*, 13(6):47–60, 1996.

- [15] Michael K Pitt and Neil Shephard. Filtering via simulation: Auxiliary particle filters. *Journal of the American Statistical Association*, 94(446):590–599, 1999.
- [16] George Poyiadjis, Arnaud Doucet, and Sumeetpal S Singh. Particle approximations of the score and observed information matrix in state space models with application to parameter estimation. *Biometrika*, 98(1):65–80, 2011.
- [17] Herbert E Rauch, F Tung, and Charlotte T Striebel. Maximum likelihood estimates of linear dynamic systems. *AIAA Journal*, 3(8):1445–1450, 1965.
- [18] Thomas B Schön, Adrian Wills, and Brett Ninness. System identification of nonlinear state-space models. *Automatica*, 47(1):39–49, 2011.
- [19] François Septier, Sze Kim Pang, Simon Godsill, and Avishy Carmi. Tracking of coordinated groups using marginalised MCMC-based particle algorithm. In *2009 IEEE Aerospace Conference*, pages 1–11. IEEE, 2009.
- [20] Emilian R Vankov, Michele Guindani, Katherine B Ensor, et al. Filtering and estimation for a class of stochastic volatility models with intractable likelihoods. *Bayesian Analysis*, 14(1):29–52, 2019.
- [21] Greg CG Wei and Martin A Tanner. A Monte Carlo implementation of the EM algorithm and the poor man’s data augmentation algorithms. *Journal of the American Statistical Association*, 85(411):699–704, 1990.
- [22] CF Jeff Wu et al. On the convergence properties of the EM algorithm. *The Annals of Statistics*, 11(1):95–103, 1983.