arXiv:2103.11378v1 [math.FA] 21 Mar 2021

On the mutipliers of the Figà-Talamanca Herz algebra

Antoine Derighetti

January 6, 2022

Abstract. Let G be a locally compact group and $p, q \in \mathbb{R}$ with $p > 1$, $p \neq 2$ and q between 2 and p (if $p < 2$ then $p < q < 2$, if $p > 2$ then $2 < q < p$.) The main result of the paper is that $A_q(G)$ multiplies $A_p(G)$, more precisely we show that the Banach algebra $A_p(G)$ is a Banach module on $A_q(G)$.

1 Introduction

Let G be a locally compact group, in [1] an entirely selfcontained proof of the following result was given (section 8.3 Theorem 8 p. 158): for every $1 < p < \infty$ $A_p(G)$ is a Banach module over $A_2(G)$: for every $u \in A_2(G)$ and for every $v \in A_p(G)$ we have $uv \in A_p(G)$ and $\|uv\|_{A_p} \leq \|u\|_{A_2}\|v\|_{A_p}$. Let now $p, q \in \mathbb{R}$ with $p > 1$, $p \neq 2$ and q between 2 and p. We show (Theorem 8) that for every $u \in A_q(G)$ and for every $v \in A_p(G)$ we have $uv \in A_p(G)$ and $\|uv\|_{A_p} \leq \|u\|_{A_q}\|v\|_{A_p}$.

We recall some definitions. For φ a map of G into $\mathbb{C}$ we set $\check{\varphi}(x) = \varphi(x^{-1})$, ${}_a\varphi(x) = \varphi(ax)$ and $\varphi_a(x) = \varphi(xa)$ for $a, x \in G$. For $1 < p < \infty$ $L^p(G)$ is the L^p-space with respect to a left Haar measure of G. Let $\mathcal{L}(L^p(G))$ be the Banach algebra of all linear bounded operators of $L^p(G)$, the operator norm of an operator T is denoted $|\!|\!|T|\!|\!|_p$. An element T of $\mathcal{L}(L^p(G))$ is said to be a p-convolution operator (written $T \in CV_p(G)$) if $T({}_a\varphi) = {}_aT(\varphi)$ for every $a \in G$ and for every $\varphi \in L^p(G)$.

We denote by $C_{00}(G)$ the set of all continuous complex valued functions on G with compact support. Let μ be a Radon measure on G we put $\check{\mu}(\varphi) = \mu(\check{\varphi})$. Suppose that μ is a bounded Radon measure on G ($\mu \in M^1(G)$), there is a unique $S \in \mathcal{L}(L^p(G))$ such that

$$S[\varphi] = [\varphi * (\Delta_G^{1/p'}\check{\mu})]$$

for every $\varphi \in C_{00}(G)$. The operator S is denoted $\lambda_G^p(\mu)$. We have $\lambda_G^p(\mu) \in CV_p(G)$ and $|\!|\!|\lambda_G^p(\mu)|\!|\!|_p \leq \|\mu\|$. We also have $\lambda_G^p(\delta_a)\varphi = \varphi_a\Delta_G(a)^{1/p}$. In [1] we proved that every $1 < p < \infty$ and for every amenable locally compact

group G the following deep inequality holds: $\|\lambda_G^2(\mu)\|_2 \leq \|\lambda_G^p(\mu)\|_p$ for every $\mu \in M^1(G)$. Even for finite abelian groups this result is not trivial.

In this paper we show (Corollary 10) that for every amenable locally compact group G, for $p, q \in \mathbb{R}$ with $p > 1$, $p \neq 2$ and q between 2 and p we have $\|\lambda_G^q(\mu)\|_q \leq \|\lambda_G^p(\mu)\|_p$ for every $\mu \in M^1(G)$.

Let G be a locally compact group and $1 < p < \infty$. We denote by $\mathcal{A}_p(G)$ the set of all pairs $((k_n)_{n=1}^\infty, (l_n)_{n=1}^\infty)$ where $(k_n)_{n=1}^\infty$ is a sequence of $\mathcal{L}^p(G)$ and $(l_n)_{n=1}^\infty$ is a sequence of $\mathcal{L}^{p'}(G)$ with

$$\sum_{n=1}^{\infty} N_p(k_n) N_{p'}(l_n) < \infty.$$

The topology on $\mathcal{L}(L^p(G))$, associated to the family of seminorms

$$T \mapsto \left| \sum_{n=1}^{\infty} < T[k_n], [l_n] > \right|$$

with $((k_n)_{n=1}^\infty, (l_n)_{n=1}^\infty) \in \mathcal{A}_p(G)$, is called the ultraweak topology. The ultraweak closure of $\lambda_G^p(M^1(G))$ in $\mathcal{L}(L^p(G))$ is denoted $PM_p(G)$. Every element of $PM_p(G)$ is called a p-pseudomeasure. The space $PM_p(G)$ can be identified as the dual of a Banach algebra $A_p(G)$ of continuous continuous functions on G.

We denote by $A_p(G)$ the set:

$$\Big\{ u : G \to \mathbb{C} : \text{there is } ((k_n)_{n=1}^\infty, (l_n)_{n=1}^\infty)) \in \mathcal{A}_p(G) \text{ such that}$$

$$u(x) = \sum_{n=1}^{\infty} (\overline{k_n} * \check{l_n})(x) \text{ for every } x \in G \Big\}.$$

For $u \in A_p(G)$ we put

$$\|u\|_{A_p} := \inf \Bigg\{ \sum_{n=1}^{\infty} N_p(k_n) N_{p'}(l_n) : ((k_n)_{n=1}^\infty, (k_n)_{n=1}^\infty) \in \mathcal{A}_p(G)$$

$$\text{such that } u = \sum_{n=1}^{\infty} \overline{k_n} * \check{l_n} \Bigg\}.$$

For the pointwise product $A_p(G)$ is a Banach algebra, called the Figà-Talamanca Herz of G. We have $A_p(G) \subset C_0(G)$ and for every $u \in A_p(G)$ the following inequality holds: $\|u\|_\infty \leq \|u\|_{A_p}$. Moreover the space $A_p(G) \cap C_{00}(G)$ is dense in $A_p(G)$.

For every $T \in PM_p(G)$, we put

$$\Psi_G^p(T)(u) = \sum_{n=1}^{\infty} \overline{< T[\tau_p k_n], [\tau_{p'} l_n] >}$$

for every $u \in A_p(G)$ and for every $((k_n)_{n=1}^{\infty}, (k_n)_{n=1}^{\infty}) \in \mathcal{A}_p(G)$ such that

$$u = \sum_{n=1}^{\infty} \overline{k_n} * \check{l_n}$$

where $\tau_p\varphi(x) = \varphi(x^{-1})\Delta_G(x^{-1})^{1/p}$. Then Ψ_G^p is a conjugate linear isometry of $PM_p(G)$ onto $A_p(G)'$ with the following properties:

1. $\Psi_G^p(\lambda_G^p(\tilde{\mu})) = \mu$ for every $\mu \in M^1G$, where $\tilde{\varphi}(x) = \overline{\varphi(x^{-1})}$ and $\tilde{\mu}(\varphi) = \overline{\mu(\tilde{\varphi})}$.
2. Ψ_G^p is an homeomorphism of $PM_p(G)$, with the ultraweak topology, onto $A_p(G)'$, with the weak topology $\sigma({A_p}', A_p)$.

We will use the fact (see [1] Chapter 5) that $CV_p(G)$ carries a natural structure of left normed $A_p(G)$-module $(u, T) \mapsto uT$. We shortly recall the definition of uT: for $k \in \mathcal{L}^p(G)$, $l \in \mathcal{L}^{p'}(G)$ and $T \in CV_p(G)$ there is a unique linear bounded operator of $L^p(G)$, denoted $(\overline{k} * \check{l})T$, such that

$$\langle((\overline{k} * \check{l})T)[\varphi], [\psi]\rangle = \int_G \langle T[{}_{t^{-1}}(\check{k})\varphi], [{}_{t^{-1}}(\check{l})\psi]\rangle dt$$

for every $\varphi, \psi \in C_{00}(G)$. Then, for $u \in A_p(G)$, we put $uT = \sum_{n=1}^{\infty} (\overline{k_n} * \check{l_n})T$ for every $((k_n)_{n=1}^{\infty}, (l_n)_{n=1}^{\infty}) \in \mathcal{A}_p(G)$ such that $u = \sum_{n=1}^{\infty} \overline{k_n} * \check{l_n}$.

2 The general case

In this paragraph G is a general locally compact group, G is not assumed to be amenable.

Theorem 1. Let G be a locally compact group, $1 < p < \infty$, $u \in A_p(G), T \in CV_p(G)$ and $\varphi \in L^p(G) \cap L^2(G)$.

1. $\tau_p(uT)\tau_p\varphi \in L^2(G)$,
2. $\|\tau_p(uT)\tau_p\varphi\|_2 \leq \|u\|_{A_p} \|T\|_p \|\varphi\|_2$.

Proof. See [1] section 8.3 Theorem 1 p. 156. □

Proposition 2. Let G be a locally compact group, $p, q \in \mathbb{R}$ with $p > 1, p \neq 2$ and q between p and 2. We denote by $\mathcal{S}$ the set $\{[r] : r \in \mathcal{L}^1(G), r \text{ step function}\}$. Then for every $u \in A_p(G)$ for every $T \in CV_p(G)$ and every $\varphi \in \mathcal{S}$ we have: for every $1 < p < \infty$

1. $\tau_p(uT)\tau_p\varphi \in L^q(G)$,
2. $\|\tau_p(uT)\tau_p\varphi\|_q \leq \|u\|_{A_p} \|T\|_p \|\varphi\|_q$.

Proof. (I) We prove Proposition 2 if $p < 2$.

We put

$$t = \frac{2(q-p)}{q(2-p)}.$$

We have $0 < t < 1$ and

$$\frac{1}{q} = \frac{1-t}{p} + \frac{t}{2}.$$

According to Riesz-Thorin and to Theorem 1 $\tau_p(uT)\tau_p\varphi \in L^q(G)$ and

$$\|\tau_p(uT)\tau_p\varphi\|_q \leq \|uT\|_p^{1-t} \left(\|u\|_{A_p} \|T\|_p\right)^t \|\varphi\|_q$$

but

$$\|uT\|_p^{1-t} \left(\|u\|_{A_p} \|T\|_p\right)^t \|\varphi\|_q \leq \|u\|_{A_p}^{1-t} \|T\|_p^{1-t} \|u\|_{A_p}^t \|T\|_p^t \|\varphi\|_q$$

$$= \|u\|_{A_p} \|T\|_p \|\varphi\|_q .$$

(II) We prove Proposition 2 if $p > 2$.

We put

$$t = \frac{(2-q)p}{(2-p)q}.$$

We have $0 < t < 1$ and

$$\frac{1}{q} = \frac{1-t}{2} + \frac{t}{p}.$$

As in (I) $\tau_p(uT)\tau_p\varphi \in L^q(G)$ and

$$\|\tau_p(uT)\tau_p\varphi\|_q \leq \left(\|u\|_{A_p} \|T\|_p\right)^{1-t} \|uT\|_p^t \|\varphi\|_q$$

and consequently $\|\tau_p(uT)\tau_p\varphi\|_q \leq \|u\|_{A_p} \|T\|_p \|\varphi\|_q$. □

Theorem 3. Let G be a locally compact group, $p, q \in \mathbb{R}$ with $p > 1$, $p \neq 2$ and q between p and 2. Then for every $u \in A_p(G)$, for every $T \in CV_p(G)$ and every $\varphi \in L^p(G) \cap L^q(G)$ we have:

1. $\tau_p(uT)\tau_p\varphi \in L^q(G)$,
2. $\|\tau_p(uT)\tau_p\varphi\|_q \leq \|u\|_{A_p} \|T\|_p \|\varphi\|_q$.

Definition 1. Let G be a locally compact group, $p, q \in \mathbb{R}$ with $p > 1$, $p \neq 2$ and q between p and 2. We denote by $E_{p,q}$ the set of all $T \in CV_p(G)$ such that :

1. $\tau_p T \tau_p\varphi \in L^q(G)$ for every $\varphi \in L^p(G) \cap L^q(G)$,
2. There is $C > 0$ such that $\|\tau_p T \tau_p\varphi\|_q \leq C\|\varphi\|_q$ for every $\varphi \in L^p(G) \cap L^q(G)$.

Proposition 4. Let G be a locally compact group, $p, q \in \mathbb{R}$ with $p > 1$, $p \neq 2$ and q between p and 2. The set $E_{p,q}$ is a subalgebra of $CV_p(G)$.

Proof. Let $S, T \in E_{p,q}$, C, C' the constants of Definition 1 and $\varphi \in L^p(G) \cap L^q(G$. From $\tau_p T \tau_p \varphi \in L^p(G) \cap L^q(G)$ it follows $(\tau_p S \tau_p)(\tau_p T \tau_p \varphi) \in L^p(G) \cap L^q(G)$ and consequently $\tau_p ST \tau_p \varphi \in L^p(G) \cap L^q(G)$. Moreover

$$\|\tau_p ST \tau_p \varphi\|_q = \|\tau_p S \tau_p (\tau_p T \tau_p \varphi)\|_q \leq C \|\tau_p T \tau_p \varphi\|_q \leq CC' \|\varphi\|_q$$

this implies that $ST \in E_{p,q}$. □

Proposition 5. Let G be a locally compact group, $p, q \in \mathbb{R}$ with $p > 1$, $p \neq 2$ and q between p and 2. For every $\mu \in M^1(G)$ we have $\lambda^p_G(\mu) \in E_{p,q}$ and for every $\varphi \in L^p(G) \cap L^q(G)$ $\tau_p \lambda^p_G(\mu) \tau_p \varphi = \rho^q_G(\mu)\varphi$.

Proof. For every $\varphi \in C_{00}(G)$ we have $\tau_p(\tau_p \varphi * \Delta_G^{1/p'} \check{\mu}) = \varphi * \mu$. Let $\varphi \in L^p(G) \cap L^q(G)$ we obtain $\tau_p \lambda^p_G(\mu) \tau_p \varphi = \rho^q_G(\mu)\varphi$. But $\rho^q_G(\mu)\varphi$ belongs to $L^q(G)$ and therefore $\tau_p \lambda^p_G(\mu) \tau_p \varphi \in L^q(G)$. Finally the inequality $\|\rho^q_G(\mu)\varphi\|_q \leq \|\mu\| \|\varphi\|_q$ implies $\|\tau_p \lambda^p_G(\mu) \tau_p \varphi\|_q \leq \|\mu\| \|\varphi\|_q$.

□

Proposition 6. Let G be a locally compact group, $p, q \in \mathbb{R}$ with $p > 1$, $p \neq 2$ and q between p and 2. For every $T \in E_{p,q}$ there is a unique $S \in \mathcal{L}(L^q(G))$ such that $S\varphi = \tau_p T \tau_p \varphi$ for every $\varphi \in L^p(G) \cap L^q(G)$. We have $\tau_q S \tau_q \in CV_q(G)$.

Definition 2. Let G be a locally compact group, $p, q \in \mathbb{R}$ with $p > 1$, $p \neq 2$ and q between p and 2, $T \in E_{p,q}$ and S as in Proposition 6. We put $\alpha_{p,q}(T) = \tau_q S \tau_q$.

Theorem 7. Let G be a locally compact group, $p, q \in \mathbb{R}$ with $p > 1$, $p \neq 2$ and q between p and 2. Then:

1. $\alpha_{p,q}$ is a monomorphism of the algebra $E_{p,q}$ into $CV_q(G)$,
2. $\alpha_{p,q}(\lambda^p_G(\mu)) = \lambda^q_G(\mu)$ for every $\mu \in M^1(G)$,
3. for every $u \in A_p(G)$ and for every $T \in CV_p(G)$ we have $uT \in E_{p,q}$ and $\|\!|\alpha_{p,q}(uT)|\!\|_q \leq \|u\|_{A_p} \|\!|T|\!\|_p$,
4. for every $T \in PM_p(G) \cap E_{p,q}$ we have $\alpha_{p,q}(T) \in PM_q(G)$.

Proof. (I) We prove the assertion 1.

Let $T, T' \in E_{p,q}$ $C, C' > 0$ $S, S' \in \mathcal{L}(L^q(G))$ such that $\|\tau_p T \tau_p \varphi\|_q \leq C\|\varphi\|_q$, $\|\tau_p T' \tau_p \varphi\|_q \leq C'\|\varphi\|_q$, $S\varphi = \tau_p T \tau_p \varphi$, and $S'\varphi = \tau_p T' \tau_p \varphi$ for every $\varphi \in L^p(G) \cap L^q(G)$. We have $\alpha_{p,q}(T) = \tau_q S \tau_q$ and $\alpha_{p,q}(T') = \tau_q S' \tau_q$.

$(\mathrm{I})_1$ $\alpha_{p,q}(T + T') = \alpha_{p,q}(T) + \alpha_{p,q}(T')$.

There is $S'' \in \mathcal{L}(L^q(G))$ such that $\tau_p(T + T')\tau_p \varphi = S''\varphi$ for every $\varphi \in L^p(G) \cap L^q(G)$. We have $\alpha_{p,q}(T+T') = \tau_q S'' \tau_q$. For $\varphi \in L^p(G) \cap L^q(G)$ we get $\tau_p(T + T')\tau_p \varphi = \tau_p(T\tau_p \varphi + T'\tau_p \varphi) = \tau_p T \tau_p \varphi + \tau_p T' \tau_p \varphi = S\varphi + S'\varphi = S''\varphi$. Consequently $S + S' = S''$.Thus $\alpha_{p,q}(T) + \alpha_{p,q}(T') = \alpha_{p,q}(T + T')$.

$(\mathrm{I})_2$ $\alpha_{p,q}(\gamma T) = \gamma\, \alpha_{p,q}(T)$ for $\gamma \in \mathbb{C}$ and $\alpha_{p,q}(TT') = \alpha_{p,q}(T)\alpha_{p,q}(T')$.

The proof of $(\mathrm{I})_2$ is similar to the one of $(\mathrm{I})_1$.

(I)$_3$ If $\alpha_{p,q}(T)=0$ then $T=0$.

We have $\tau_q S\tau_q=0$, this implies $S=0$. For every $\varphi\in L^p(G)\cap L^q(G)$ $S\varphi=\tau_p T\tau_p\varphi$, consequently $\tau_p T\tau_p\varphi=0$, hence $T\tau_p\varphi=0$. Consider $r\in C_{00}(G)$, we have $T[r]=T\tau_p\tau_p[r]$, taking into account that $\tau_p[r]\in L^p(G)\cap L^q(G)$ we get $T\tau_p\tau_p[r]=0$ i.e $T[r]=0$ and finally $T=0$.

(II) The property 2. is verified.

According to Proposition 5 $\lambda_G^p(\mu)\in E_{p,q}$. Let $S\in\mathcal{L}(L^q(G))$ such that $S\varphi=\tau_p\lambda_G^p(\mu)\tau_p\varphi$ for every $\varphi\in L^p(G)\cap L^q(G)$, we have $\alpha_{p,q}(\lambda_G^p(\mu))=\tau_q S\tau_q$. By Proposition 3 for every $\varphi\in L^p(G)\cap L^q(G)$ we have $\tau_p\lambda_G^p(\mu)\tau_p\varphi=\rho_G^q(\mu)\varphi$. Consequently $S\varphi=\rho_G^q(\mu)\varphi$ every $\varphi\in L^p(G)\cap L^q(G)$, this implies $S=\rho_G^q(\mu)$. We obtain $\tau_q S\tau_q=\tau_q\rho_G^q(\mu)\tau_q$. It is straightforward to verify that $\tau_q\rho_G^q(\mu)\tau_q=\lambda_G^q(\mu)$, we finally conclude that $\alpha_{p,q}(\lambda_G^p(\mu))=\lambda_G^q(\mu)$.

(III) The property 3. is verified.

According to Theorem 3 $uT\in E_{p,q}$. Let $S\in\mathcal{L}(L^q(G))$ such that $S\varphi=\tau_p\lambda_G^p(\mu)\tau_p\varphi$ for every $\varphi\in L^p(G)\cap L^q(G)$. By Theorem 3 again for every $\varphi\in L^p(G)\cap L^q(G)$ $\|S\varphi\|_q\leq\|u\|_{A_p}\|T\|_p\|\varphi\|_q$. Consequently for every $\varphi\in L^q(G)$ we have $\|S\varphi\|_q\leq\|u\|_{A_p}\|T\|_p\|\varphi\|_q$. It follows that for every $\varphi\in L^q(G)$ we have $\|\tau_q S\tau_q\varphi\|_q=\|S\tau_q\varphi\|_q\leq u\|_{A_p}\|T\|_p\|\tau_q\varphi\|_q=\|u\|_{A_p}\|T\|_p\|\varphi\|_q$, thus $\|\tau_q S\tau_q\|_q\leq\|T\|_p\|u\|_{A_p}$. We finally have $\|\alpha_{p,q}(T)\|_q=\|\tau_q S\tau_q\|_q\leq\|T\|_p\|u\|_{A_p}$.

(IV) Proof of 4.

Let $((k_n)_{n=1}^\infty,(l_n)_{n=1}^\infty)\in\mathcal{A}_q(G)$ such that $k_n,l_n\in C_{00}(G)$ for every $n\geq1$ and such that

$$\sum_{n=1}^\infty\overline{k_n}*\check{l_n}=0.$$

We have $\langle\alpha_{p,q}(T)[\tau_q k_n],[\tau_{q'}l_n]\rangle_{L^q,L^{q'}}=\langle T[\tau_p k_n],[\tau_{p'}l_n]\rangle_{L^p,L^{p'}}$ for every $n\geq1$. By [1] Lemma 5 section 4.1 page 48

$$\sum_{n=1}^\infty\langle T[\tau_p k_n],[\tau_{p'}l_n]\rangle_{L^p,L^{p'}}=0$$

and therefore

$$\sum_{n=1}^\infty\langle\alpha_{p,q}(T)[\tau_q k_n],[\tau_{q'}l_n]\rangle_{L^q,L^{q'}}=0.$$

The Corollary 8 of [1] section 4.1 page 52 implies that $\alpha_{p,q}(T)\in PM_q(G)$. □

The following theorem is the main result of the paper.

Theorem 8. Let G be a locally compact group, $p,q\in\mathbb{R}$ with $p>1,p\neq2$ and q between p and 2. Let u be a function of $A_q(G)$ and $v\in A_p(G)$. Then:

1. $uv\in A_p(G)$ and $\|uv\|_{A_p}\leq\|u\|_{A_q}\|u\|_{A_p}$, i.e $A_p(G)$ is a Banach module on the Banach algebra $A_q(G)$,

2. for every $T \in PM_p(G)$ we have

$$\langle uv, T\rangle_{A_p,PM_p} = \langle u, \alpha_{p,q}(vT)\rangle_{A_q,PM_q}$$

(according to Theorem 7 $\alpha_{p,q}(vT) \in PM_q(G)$).

Proof. (I) Definition of a linear map ω (depending of u) of $A_p(G)'$ into $A_p(G)''$.

There is $((k_n)_{n=1}^{\infty}, (l_n)_{n=1}^{\infty}) \in \mathcal{A}_q(G)$ such that

$$u = \sum_{n=1}^{\infty} \overline{k_n} * \check{l_n}.$$

Let F be an arbitrary element of $A_p(G)'$. We put

$$\omega(F) = \sum_{n=1}^{\infty} \overline{\langle \alpha_{p,q}(v(\Psi_G^p)^{-1}(F))[\tau_q k_n], [\tau_{q'} l_n]\rangle}_{L^q,L^{q'}}.$$

It is easy to verify that for $\gamma \in \mathbb{C}$ $\omega(\gamma F) = \gamma\omega(F)$ and that $\omega(F+F') = \omega(F) + \omega(F')$ for every $F, F' \in A_p(G)'$. Moreover

$$|\omega(F)| \leq \sum_{n=1}^{\infty} \left\| \alpha_{p,q}(v(\Psi_G^p)^{-1}(F)) \right\|_q N_q(k_n) N_{q'}(l_n)$$

according to Theorem 7 point 3. the last expression is not larger than

$$\sum_{n=1}^{\infty} \|v\|_{A_p} \left\| (\Phi_G^p)^{-1}(F) \right\|_p N_q(k_n) N_{q'}(l_n)$$

$$= \|v\|_{A_p} \|F\|_{A_p'} \sum_{n=1}^{\infty} N_q(k_n) N_{q'}(l_n)$$

this implies that $\omega(F) \in A_p(G)''$.

(II) Let F be an element of $A_p(G)'$ and $(F_i)_{i\in I}$ a net of $A_p(G)'$ such that $\lim F_i = F$ for the topology $\sigma(A_p', A_p)$. We assume the existence of $C > 0$ with $\|F_i\|_{A_p'} \leq C$ for every i. Then $\lim \omega(F_i) = \omega(F)$.

By [1] Theorem 6 section 4.1 page 49 $\lim (\Psi_G^p)^{-1}(F_i) = (\Psi_G^p)^{-1}(F)$ for the ultraweak topology on $\mathcal{L}(L^p(G))$. We also have $\left\| (\Psi_G^p)^{-1}(F_i) \right\|_p \leq C$ for every $i \in I$.

(II)$_1$ Let $(S_i)_{i\in I}$ be a net of $PM_p(G)$ and $S \in PM_p(G)$. Suppose that $\lim S_i = S$ for the ultraweak topology on $\mathcal{L}(L^p(G))$. Then for every $v \in A_p(G)$ we have $\lim vS_i = vS$ for the ultraweak topology on $\mathcal{L}(L^p(G))$.

For every $a \in A_p(G)$ we have $\lim < a, S_i >_{A_p,PM_p} = < a, S >_{A_p,PM_p}$. In particular $\lim < av, S_i >_{A_p,PM_p} = < av, S >_{A_p,PM_p}$ hence $\lim < a, vS_i >_{A_p,PM_p} = < a, vS >_{A_p,PM_p}$ and consequently $\lim vS_i = vS$ for the ultraweak topology on $\mathcal{L}(L^p(G))$.

$(\mathrm{II})_2$ End of the proof of (II).

By $(\mathrm{II})_1$ $\lim v(\Psi_G^p)^{-1}(F_i) = v(\Psi_G^p)^{-1}(F)$ for the ultraweak topology on $\mathcal{L}(L^p(G))$. Moreover for every $i \in I$

$$\left\|v(\Psi_G^p)^{-1}(F_i)\right\|_p \leq \|v\|_{A_p} \left\|(\Psi_G^p)^{-1}(F_i)\right\|_p \leq \|v\|_{A_p} C.$$

Let $r, s \in C_{00}(G)$. We have

$$\lim \left\langle v(\Psi_G^p)^{-1}(F_i)[\tau_p r], [\tau_{p'} s]\right\rangle_{L^p, L^{p'}} = \left\langle v(\Psi_G^p)^{-1}(F)[\tau_p r], [\tau_{p'} s]\right\rangle_{L^p, L^{p'}}.$$

Taking in account that $[r] \in L^p(G) \cap L^q(G)$ and Theorem 3 we have

$$\tau_q \alpha_{p,q}\big(v(\Psi_G^p)^{-1}(F)\big)\tau_q[r] = \tau_p v(\Psi_G^p)^{-1}(F)\tau_p[r]$$

and

$$\tau_q \alpha_{p,q}\big(v(\Psi_G^p)^{-1}(F_i)\big)\tau_q[r] = \tau_p v(\Psi_G^p)^{-1}(F_i)\tau_p[r].$$

Consequently

$$\left\langle v(\Psi_G^p)^{-1}(F)[\tau_p r], [\tau_{p'} s]\right\rangle_{L^p, L^{p'}} = \left\langle \alpha_{p,q}\big(v(\Psi_G^p)^{-1}(F)\big)[\tau_q r], [\tau_{q'} s]\right\rangle_{L^q, L^{q'}}$$

and

$$\left\langle v(\Psi_G^p)^{-1}(F_i)[\tau_p r], [\tau_{p'} s]\right\rangle_{L^p, L^{p'}} = \left\langle \alpha_{p,q}\big(v(\Psi_G^p)^{-1}(F_i)\big)[\tau_q r], [\tau_{q'} s]\right\rangle_{L^q, L^{q'}}.$$

But for every $i \in I$

$$\left\|\alpha_{p,q}\big(v(\Psi_G^p)^{-1}(F_i)\big)\right\|_q \leq \|v\|_{A_p} \left\|(\Psi_G^p)^{-1}(F_i)\right\|_p \leq \|v\|_{A_p} C.$$

It follows that

$$\lim \alpha_{p,q}\big(v(\Psi_G^p)^{-1}(F_i)\big) = \alpha_{p,q}\big(v(\Psi_G^p)^{-1}(F)\big)$$

ultraweakly on $\mathcal{L}(L^q(G))$. In particular

$$\lim \sum_{n=1}^{\infty} \left\langle \alpha_{p,q}\big(v(\Psi_G^p)^{-1}(F_i)\big)[\tau_q k_n], [\tau_{q'} l_n]\right\rangle_{L^q, L^{q'}}$$

$$= \sum_{n=1}^{\infty} \left\langle \alpha_{p,q}\big(v(\Psi_G^p)^{-1}(F)\big)[\tau_q k_n], [\tau_{q'} l_n]\right\rangle_{L^q, L^{q'}}$$

i.e $\lim \omega(F_i) = \omega(F)$.

(III) There is $w \in A_p(G)$ such that $\omega(\Psi_G^p(T)) =< w, T >_{A_p, PM_p}$ for every $T \in PM_p(G)$.

According to [1] Theorem 6 section 4.2 page 54 there is $w \in A_p(G)$ with $\omega(F) = F(w)$ for every $F \in A_p(G)'$. For every $T \in PM_p(G)$ we get $\omega(\Psi_G^p(T)) = \Psi_G^p(T)(w) =< w, T >_{A_p, PM_p}$.

(IV) For every $T \in PM_p(G)$ we have

$$< w, T >_{A_p, PM_p} = < u, \alpha_{p,q}(vT) >_{A_q, PM_q} .$$

We have

$$< w, T >_{A_p, PM_p} = \omega(\Psi_G^p(T))$$

$$= \sum_{n=1}^{\infty} \overline{\langle \alpha_{p,q}(v(\Psi_G^p)^{-1}(\Psi_G^p(T)))[\tau_q k_n], [\tau_{q'} l_n] \rangle}_{L^q, L^{q'}}$$

$$= \sum_{n=1}^{\infty} \overline{\langle \alpha_{p,q}(vT)[\tau_q k_n], [\tau_{q'} l_n] \rangle}_{L^q, L^{q'}} = < u, \alpha_{p,q}(vT) >_{A_q, PM_q} .$$

(V) End of the proof of Theorem 8.
For every $\mu \in M^1(G)$ we have

$$< w, \lambda_G^p(\mu) >_{A_p, PM_p} = < u, \alpha_{p,q}(v\lambda_G^p(\mu)) >_{A_q, PM_q}$$

but $v\lambda_G^p(\mu) = \lambda_G^p(\tilde{v}\mu)$ and $\alpha_{p,q}(\lambda_G^p(\tilde{v}\mu)) = \lambda_G^q(\tilde{v}\mu)$. From $< w, \lambda_G^p(\mu) >_{A_p, PM_p} = \Psi_G^p(\lambda_G^p(\mu))(w) = \tilde{\mu}(w)$ and

$$\langle u, \lambda_G^q(\tilde{v}\mu) \rangle_{A_q, PM_q} = \tilde{\mu}(uv),$$

it follows $\tilde{\mu}(uv) = \tilde{\mu}(w)$. This implies $w = uv$ and therefore $uv \in A_p(G)$. We get moreover

$$\langle uv, T \rangle_{A_p, PM_p} = \langle u, \alpha_{p,q}(vT) \rangle_{A_q, PM_q}$$

for every $T \in PM_p(G)$. We obtain the following estimate

$$|\langle uv, T \rangle_{A_p, PM_p}| = |\langle u, \alpha_{p,q}(vT) \rangle_{A_q, PM_q}| \leq \|u\|_{A_q} \|\alpha_{p,q}(vT)\|_q$$

$$\leq \|u\|_{A_q} \|v\|_{A_p} \|T\|_p .$$

We finally conclude that $\|uv\|_{A_p} \leq \|u\|_{A_q} \|v\|_{A_p}$. □

Remark 1. For a related result see Theorem B of [3]. Our approach is based on properties of $A_p(G)$ contained in [1].

3 Amenable groups

We apply the results of paragraph 2 to the case of amenable groups.

Theorem 9. Let G be an amenable locally compact group, $p, q \in \mathbb{R}$ with $p > 1, p \neq 2$ and q between 2 and p. Then for every $T \in CV_p(G)$ we have $T \in E_{p,q}$ and $\|\alpha_{p,q}(T)\|_q \leq \|T\|_p$.

Proof. (I) For every $\varphi \in L^p(G) \cap L^q(G)$ we have $\tau_p T \tau_p \varphi \in L^q(G)$ and $\|\tau_p T \tau_p \varphi\|_q \leq \|T\|_p \|\varphi\|_q$.

It suffices to verify (I) for $[\varphi]$ with $\varphi \in C_{00}(G)$. Let $\psi \in C_{00}(G)$ with $N_{q'}(\psi) \leq 1$ and $\varepsilon > 0$. By [1] Lemma 1 section 5.4 page 80 there is $k, l \in C_{00}(G)$ with $k \geq 0,\ l \geq 0,\ N_p(k) = N_{p'}(l) = \int_G k(t)l(t)dt = 1$ and such that

$$\left|\left\langle ((k * \check{l})T)[\tau_p\varphi], [\tau_{p'}\psi]\right\rangle_{L^p,L^{p'}} - \left\langle T[\tau_p\varphi], [\tau_{p'}\psi]\right\rangle_{L^p,L^{p'}}\right| < \varepsilon.$$

But

$$\left\langle ((k * \check{l})T)[\tau_p\varphi], [\tau_{p'}\psi]\right\rangle_{L^p,L^{p'}} = \left\langle \tau_p((k * \check{l})T)\tau_p[\varphi], [\psi]\right\rangle_{L^p,L^{p'}}$$

and according to Theorem 3 $\tau_p((k * \check{l})T)\tau_p[\varphi]$ belongs to $L^q(G)$ this implies

$$\left\langle \tau_p((k * \check{l})T)\tau_p[\varphi], [\psi]\right\rangle_{L^p,L^{p'}} = \left\langle \tau_p((k * \check{l})T)\tau_p[\varphi], [\psi]\right\rangle_{L^q,L^{q'}}$$

and therefore

$$\left|\left\langle ((k * \check{l})T)[\tau_p\varphi], [\tau_{p'}\psi]\right\rangle_{L^p,L^{p'}}\right| \leq \|\tau_p((k * \check{l})T)\tau_p[\varphi]\|_q \|[\psi]\|_{q'}.$$

Using again Theorem 3 we have

$$\|\tau_p((k * \check{l})T)\tau_p[\varphi]\|_q \leq \|k * \check{l}\|_{A_p} \|T\|_p \|[\varphi]\|_q \leq \|T\|_p \|[\varphi]\|_q,$$

and obtain

$$\left|\left\langle ((k * \check{l})T)[\tau_p\varphi], [\tau_{p'}\psi]\right\rangle_{L^p,L^{p'}}\right| \leq \|T\|_p \|[\varphi]\|_q \|\psi\|_{q'} \leq \|T\|_p \|[\varphi]\|_q$$

and therefore

$$\left|\left\langle T[\tau_p\varphi].[\tau_{p'}\psi]\right\rangle_{L^p,L^{p'}}\right| < \varepsilon + \|T\|_p \|[\varphi]\|_q.$$

Thus

$$\left|\left\langle T[\tau_p\varphi].[\tau_{p'}\psi]\right\rangle_{L^p,L^{p'}}\right| \leq \|T\|_p \|[\varphi]\|_q,$$

$$\left|\left\langle \tau_p T \tau_p[\varphi].[\psi]\right\rangle_{L^p,L^{p'}}\right| \leq \|T\|_p \|[\varphi]\|_q$$

and finally $\tau_p T \tau_p[\varphi] \in L^q(G)$ with $\|\tau_p T \tau_p[\varphi]\|_q \leq \|T\|_p \|[\varphi]\|_q$.

(II) $\|\alpha_{p,q}(T)\|_q \leq \|T\|_p$.

There is a unique $S \in \mathcal{L}(L^q(G))$ such that $S\varphi = \tau_p T \tau_p \varphi$ for every $\varphi \in L^p(G) \cap L^q(G)$. By (I) $\|S\|_q \leq \|T\|_p$, consequently $\|\alpha_{p,q}(T)\|_q = \|\tau_q S \tau_q\|_q = \|S\|_q \leq \|T\|_p$. □

Corollary 10. Let G be an amenable locally compact group, $p, q \in \mathbb{R}$ with $p > 1, p \neq 2$ and q between 2 and p. Then:

1. $\alpha_{p,q}$ is a contractive Banach algebra monomorphism of $CV_p(G)$ into $CV_q(G)$,

2. for every $\mu \in M^1(G)$ we have $\|\lambda_G^q(\mu)\|_q \leq \|\lambda_G^p(\mu)\|_p$.

Proof. It suffices to verify 2. . By Theorem 9

$$\|\alpha_{p,q}(\lambda_G^p(\mu))\|_q \leq \|\lambda_G^p(\mu)\|_p$$

but by Theorem 7 $\alpha_{p,q}(\lambda_G^p(\mu)) = \lambda_G^q(\mu)$. □

Remark 2. See [4] Corollary p. 512. For another approach for the unimodular case see [2].

We obtain other properties of $\alpha_{p,q}$ in the next theorem.

Theorem 11. Let G be an amenable locally compact group, $p, q \in \mathbb{R}$ with $p > 1, p \neq 2$ and q between 2 and p. Then:

1. for every $u \in A_q(G)$ we have $u \in A_p(G)$ and $\|u\|_{A_p} \leq \|u\|_{A_q}$,
2. for every $u \in A_q(G)$ and for every $T \in PM_p(G)$ we have

$$\langle u, T\rangle_{A_p, PM_p} = \langle u, \alpha_{p,q}(T)\rangle_{A_q, PM_q},$$

3. for every $u \in A_q(G)$ and for every $T \in CV_p(G)$ we have

$$\alpha_{p,q}(uT) = u\,\alpha_{p,q}(T).$$

Proof. Let u be a function of $A_q(G)$.

(I) Definition of a linear map ω (depending of u) of $A_p(G)'$ into $A_p(G)''$. There is $((k_n)_{n=1}^\infty, (l_n)_{n=1}^\infty) \in \mathcal{A}_q(G)$ such that

$$u = \sum_{n=1}^\infty \overline{k_n} * \check{l_n}.$$

For every $F \in A_p(G)'$ we put

$$\omega(F) = \sum_{n=1}^\infty \overline{\langle \alpha_{p,q}((\Psi_G^p)^{-1}(F))[\tau_q k_n], [\tau_{q'} l_n]\rangle}_{L^q, L^{q'}}.$$

For every $\gamma \in \mathbb{C}$ and $F, F' \in A_p(G)'$ we have $\omega(\gamma F) = \gamma\omega(F)$ and $\omega(F + F') = \omega(F) + \omega(F')$. We now show that $\omega(F) \in A_p(G)''$. Observe at first that

$$|\omega(F)| \leq \sum_{n=1}^\infty \left|\langle \alpha_{p,q}((\Psi_G^p)^{-1}(F))[\tau_q k_n], [\tau_{q'} l_n]\rangle_{L^q, L^{q'}}\right|,$$

taking in account Theorem 9 we have for every $n \geq 1$

$$\left|\langle \alpha_{p,q}((\Psi_G^p)^{-1}(F))[\tau_q k_n], [\tau_{q'} l_n]\rangle_{L^q, L^{q'}}\right|$$

$$\leq \left\|(\Psi_G^p)^{-1}(F)\right\|_p N_q(k_n) N_{q'}(l_n) = \|F\|_{A_p'} N_q(k_n) N_{q'}(l_n)$$

hence

$$|\omega(F)| \leq \|F\|_{A'_p} \sum_{n=1}^{\infty} N_q(k_n) N_{q'}(l_n).$$

This implies that $\omega(F) \in A_p(G)''$.

(II) Let F be an element of $A_p(G)'$ and $(F_i)_{i\in I}$ a net of $A_p(G)'$ such that $\lim F_i = F$ for the topology $\sigma(A'_p, A''_p)$. We assume the existence of $C > 0$ such that $\|F_i\|_{A'_p} \leq C$ for every $i \in I$. Then $\lim \omega(F_i) = \omega(F)$.

We have $\lim (\Psi^p_G)^{-1}(F_i) = (\Psi^p_G)^{-1}(F)$ ultraweakly in $\mathcal{L}(L^p(G))$. We also have $\|\!|(\Psi^p_G)^{-1}(F_i)|\!\|_p \leq C$ for every $i \in I$.

Let $r, s \in C_{00}(G)$. We have

$$\lim \langle (\Psi^p_G)^{-1}(F_i)[\tau_p r], [\tau_{p'} s]\rangle_{L^p, L^{p'}} = \langle (\Psi^p_G)^{-1}(F)[\tau_p r], [\tau_{p'} s]\rangle_{L^p, L^{p'}}.$$

Taking in account that $[r] \in L^p(G) \cap L^q(G)$ and that $(\Psi^p_G)^{-1}(F_i), (\Psi^p_G)^{-1}(F_i) \in E_{p,q}$ we get

$$\tau_q \alpha_{p,q}\big((\Psi^p_G)^{-1}(F_i)\big)\tau_q[r] = \tau_p (\Psi^p_G)^{-1}(F_i)\tau_p[r]$$

and

$$\tau_q \alpha_{p,q}\big((\Psi^p_G)^{-1}(F)\big)\tau_q[r] = \tau_p (\Psi^p_G)^{-1}(F)\tau_p[r].$$

This implies

$$\langle (\Psi^p_G)^{-1}(F)[\tau_p r], [\tau_{p'} s]\rangle_{L^p, L^{p'}} = \langle \tau_q \alpha_{p,q}\big((\Psi^p_G)^{-1}(F)\big)\tau_q[r], [s]\rangle_{L^q, L^{q'}}$$

and for every $i \in I$

$$\langle (\Psi^p_G)^{-1}(F_i)[\tau_p r], [\tau_{p'} s]\rangle_{L^p, L^{p'}} = \langle \tau_q \alpha_{p,q}\big((\Psi^p_G)^{-1}(F_i)\big)\tau_q[r], [s]\rangle_{L^q, L^{q'}}.$$

We have therefore

$$\lim \langle \alpha_{p,q}\big((\Psi^p_G)^{-1}(F_i)\big)[\tau_q r], [\tau_{q'} s]\rangle_{L^q, L^{q'}}$$

$$= \langle \alpha_{p,q}\big((\Psi^p_G)^{-1}(F)\big)[\tau_q r], [\tau_{q'} s]\rangle_{L^q, L^{q'}}.$$

But for every $i \in I$

$$\|\!|\alpha_{p,q}\big((\Psi^p_G)^{-1}(F_i)\big)|\!\|_q \leq \|\!|(\Psi^p_G)^{-1}(F_i)|\!\|_p \leq C,$$

we obtain that

$$\lim \alpha_{p,q}\big((\Psi^p_G)^{-1}(F_i)\big) = \alpha_{p,q}\big((\Psi^p_G)^{-1}(F_i)\big)$$

ultraweakly on $\mathcal{L}(L^q(G))$. This finally implies that

$$\lim \sum_{n=1}^{\infty} \langle \alpha_{p,q}\big((\Psi^p_G)^{-1}(F_i)\big)[\tau_q k_n], [\tau_{q'} l_n]\rangle_{L^q, L^{q'}}$$

$$= \sum_{n=1}^{\infty} \langle \alpha_{p,q}((\Psi_G^p)^{-1}(F))[\tau_q k_n], [\tau_{q'} l_n] \rangle_{L^q, L^{q'}}$$

i.e. $\lim \omega(F_i) = \omega(F)$.

(III) End of the proof of Theorem 11.

As in the proof of Theorem 8 (see step (III)) there is $v \in A_p(G)$ such that $\omega(F) = F(v)$ for every $F \in A_p(G)'$. For every $T \in CV_p(G)$ we have

$$\omega(\Psi_G^p(T)) = \Psi_G^p(T)(v) = \langle v, T \rangle_{A_p, PM_p}$$

$$= \sum_{n=1}^{\infty} \overline{\langle \alpha_{p,q}((\Psi_G^p)^{-1}(\Psi_G^p(T)))[\tau_q k_n], [\tau_{q'} l_n] \rangle}_{L^q, L^{q'}}$$

$$= \sum_{n=1}^{\infty} \overline{\langle \alpha_{p,q}(T)[\tau_q k_n], [\tau_{q'} l_n] \rangle}_{L^q, L^{q'}} = \langle u, \alpha_{p,q}(T) \rangle_{A_q, PM_q}.$$

In particular for every $\mu \in M^1(G)$ we have

$$\langle v, \lambda_G^p(\mu) \rangle_{A_p, PM_p} = \langle u, \alpha_{p,q}(\lambda_G^p(\mu)) \rangle_{A_q, PM_q} = \langle u, \lambda_G^q(\mu) \rangle_{A_q, PM_q}.$$

This implies $\tilde{\mu}(v) = \tilde{\mu}(u)$ and therefore $u = v$, we obtain that $u \in A_p(G)$. We also obtain for every $u \in A_q(G)$ and for every $T \in CV_p(G)$

$$\langle u, T \rangle_{A_p, PM_p} = \langle u, \alpha_{p,q}(T) \rangle_{A_q, PM_q}.$$

This implies the following estimate:

$$|\langle u, T \rangle_{A_p, PM_p}| = |\langle u, \alpha_{p,q}(T) \rangle_{A_q, PM_q}| \leq \|u\|_{A_q} \, \|\alpha_{p,q}(T)\|_q$$

$$\leq \|u\|_{A_q} \, \|T\|_p$$

thus $\|u\|_{A_p} \leq \|u\|_{A_q}$.

It remains to verify that for every $u \in A_q(G)$ and for every $T \in CV_p(G)$ we have

$$\alpha_{p,q}(uT) = u\,\alpha_{p,q}(T).$$

Consider an arbitrary $v \in A_q(G)$. We have

$$\langle v, \alpha_{p,q}(uT) \rangle_{A_q, PM_q} = \langle v, uT \rangle_{A_p, PM_p}.$$

But

$$\langle v, uT \rangle_{A_p, PM_p} = \langle vu, T \rangle_{A_p, PM_p} = \langle vu, \alpha_{p,q}(T) \rangle_{A_q, PM_q}$$

$$= \langle v, u\,\alpha_{p,q}(T) \rangle_{A_q, PM_q}.$$

We get

$$\langle v, \alpha_{p,q}(uT) \rangle_{A_q, PM_q} = \langle v, u\,\alpha_{p,q}(T) \rangle_{A_q, PM_q}$$

and finally

$$\alpha_{p,q}(uT) = u\,\alpha_{p,q}(T).$$

□

References

[1] A. Derighetti, Convolution Operators on Groups, Lecture Notes of the Unione Matematica Italiana 11, Springer-Verlag Berlin Heidelberg 2011.

[2] A. Derighetti, Survey on the Figà-Talamanca Herz algebra, Σ Mathematics **2019**, 7, 660; doi:10.3390/math7080660.

[3] C. Herz, The theory of p-spaces with an application to convolution operators, Trans. Am. Math. **154** (1971), 69-82.

[4] C. Herz and N. Rivière, Estimates for translation-invariant operators on spaces with mixed norms, Studia Mathematica **XLIV** (1972), 511-515.