arXiv:2103.12552v2 [math.FA] 9 Jan 2022

[1]

MAPS PRESERVING TRACE OF PRODUCTS OF MATRICES

HUAJUN HUANG AND MING-CHENG TSAI

Abstract. We prove the linearity and injectivity of two maps ϕ_1 and ϕ_2 on certain subsets of M_n that satisfy $\mathrm{tr}(\phi_1(A)\phi_2(B)) = \mathrm{tr}(AB)$. We apply it to characterize maps $\phi_i : \mathcal{S} \to \mathcal{S}$ $(i = 1, \dots, m)$ satisfying

$$\mathrm{tr}(\phi_1(A_1)\cdots\phi_m(A_m)) = \mathrm{tr}(A_1\cdots A_m)$$

in which $\mathcal{S}$ is the set of n-by-n general, Hermitian, or symmetric matrices for $m \geq 3$, or positive definite or diagonal matrices for $m \geq 2$. The real versions are also given.

1. Introduction

Preserver problem is one of the most active research area in matrix theory (e.g. [9, 11, 15, 16]). Researchers would like to characterize the maps on a given space of matrices preserving certain subsets, functions or relations. In this paper, we investigate maps $\phi_1, \dots, \phi_m$ on several types of matrix spaces that have the trace-preserving property:

$$\mathrm{tr}(\phi_1(A_1)\cdots\phi_m(A_m)) = \mathrm{tr}(A_1\cdots A_m). \tag{1.1}$$

Let $M_{m,n}$ denote the set of $m \times n$ complex matrices. Let M_n, H_n, S_n, P_n, $\overline{P_n}$, and D_n denote the set of $n \times n$ complex general, Hermitian, symmetric, positive definite, positive semi-definite, and diagonal matrices respectively. We add $(\mathbb{R})$ after the above sets to denote their counterparts in real field. Let $B(X)$ denote the space of all bounded linear operators on a Banach space X.

The classical Wigner's unitary-antiunitary theorem [15, p.12] says that if a bijective map ϕ defined on the set of all rank one projections acting on a Hilbert space H with the property

$$\mathrm{tr}(\phi(P)\phi(Q)) = \mathrm{tr}(PQ),$$

that is, ϕ preserves the transition probability $\langle\phi(P), \phi(Q)\rangle = \langle P, Q\rangle$ between the pure states P and Q (or the angle between the ranges of P and Q), then there is an either unitary or antiunitary operator U on H such that $\phi(P) = U^*PU$ or $\phi(P) = U^*P^tU$ for all rank one projections P. There is a nice generalization of Wigner's theorem due to Uhlhorn. He shows that the same conclusion holds if the property $\mathrm{tr}(\phi(P)\phi(Q)) = \mathrm{tr}(PQ)$ is replaced by $\mathrm{tr}(\phi(P)\phi(Q)) = 0 \Leftrightarrow \mathrm{tr}(PQ) = 0$ (see [17]).

[1]Mathematics Subject Classification 2020: Primary 15A86, Secondary 47B49, 15A15, 15A04.

Key words and phrases. trace of products, preserver, linear operators, matrix space.

In 2002, linearity of maps ϕ on $B(X)$ satisfying

$$\operatorname{tr}(\phi(A)\phi(B)) = \operatorname{tr}(AB), \tag{1.2}$$

for $A \in B(X)$, rank one operator $B \in B(X)$ are explored by Molnár in the proof of [13, Theorem 1] to characterize surjective maps preserving the point spectrum of product of operators: $\sigma_p(\phi(A)\phi(B)) = \sigma_p(AB)$ for $A, B \in B(X)$. Later, Molnár removes the surjectivity assumption and proves the bijectivity of ϕ on the finite dimensional case M_n in the proof of [14, Theorem 1].

In 2012, Li, Plevnik, and Šemrl [10] characterize bijective maps $\phi : S \to S$ such that for a given real number c,

$$\operatorname{tr}(\phi(A)\phi(B)) = c \;\Leftrightarrow\; \operatorname{tr}(AB) = c,$$

where S is H_n, the set $S_n(\mathbb{R})$ of real symmetric matrices, or the set of rank one projections. In [6, Lemma 3.6], Huang et al show that the following statements are equivalent for a unital operator ϕ on P_n:

(1) $\operatorname{tr}(\phi(A)\phi(B)) = \operatorname{tr}(AB)$ for $A, B \in P_n$;
(2) $\operatorname{tr}(\phi(A)\phi(B)^{-1}) = \operatorname{tr}(AB^{-1})$ for $A, B \in P_n$;
(3) $\det(\phi(A) + \phi(B)) = \det(A + B)$ for $A, B \in P_n$;
(4) $\phi(A) = U^*AU$ or U^*A^tU for a unitary matrix U.

The authors also consider the cases if ϕ is not assuming unital, the set P_n is replaced by another set like M_n, S_n , or the set of upper triangular matrices in M_n. In [8, Theorem 3.8], Leung, Ng, and Wong consider the relation (1.2) in terms of Raggio transition probability between the spaces of all normal states on two von Neumann algebras.

Recently, some researchers consider similar relations of two maps preserving certain spectra on some spaces. See Miura in [12] between two standard operator algebras. See Bourhim and Lee in [2] between $B(X)$ and $B(Y)$ for some Banach spaces X and Y. See Abdelali and Aharmim in [1] between two rectangular matrices.

Let I_n (or simply I) denote the identity matrix in M_n. Given a subset S of a vector space over a field $\mathbb{F}$, let $\langle S\rangle$ denote the $\mathbb{F}$-subspace spanned by elements of S.

In Section 2, we prove the linearity and injectivity of two maps ϕ_1 and ϕ_2 on some matrix sets $\mathcal{S}$ that satisfy (1.1) for $m = 2$ (Theorems 2.1 and 2.3):

$$\operatorname{tr}(\phi_1(A)\phi_2(B)) = \operatorname{tr}(AB), \quad \forall A, B \in \mathcal{S}. \tag{1.3}$$

These preliminary results are widely used in our characterizations of maps preserving trace of products. The matrix sets discussed in this paper are closed under Hermitian conjugate. The relation (1.3) in these sets can also be interpreted as inner product preserver:

$$\langle \phi_1(A), \phi_2(B)\rangle = \langle A, B\rangle, \quad \forall A, B \in \mathcal{S}. \tag{1.4}$$

Our discussions and results on trace preserving maps will be helpful to characterize inner product preservers in related matrix spaces.

The maps $\phi_1, \dots, \phi_m : \mathcal{S} \to \mathcal{S}$ that satisfy (1.1) are characterized as follows:

$m \geq 3$:

- $\mathcal{S} = M_n$: $\phi_i(A) = N_i A N_{i+1}^{-1}$ where $N_{m+1} := N_1$ (Theorem 3.1).
- $\mathcal{S} = H_n$ (resp., P_n or $\overline{P_n}$):
  (1) when m is odd, $\phi_i(A) = c_i U^* A U$ where U is unitary;
  (2) when m is even, $\phi_i(A) = c_i M^* A M$ if i is odd, and $\phi_i(A) = c_i M^{-1} A M^{-*}$ if i is even.

  Here $c_i \in \mathbb{R}$ (resp., $c_i \in \mathbb{R}^+$) with $c_1 \cdots c_m = 1$, as stated in Theorem 4.1 (resp. Theorem 4.4).
- $\mathcal{S} = S_n$:
  (1) when m is odd, $\phi_i(A) = c_i O^t A O$ where O is orthogonal;
  (2) when m is even, $\phi_i(A) = c_i M^t A M$ if i is odd, and $\phi_i(A) = c_i M^{-1} A M^{-t}$ if i is even.

  Here $c_i \in \mathbb{C}$ with $c_1 \cdots c_m = 1$, as stated in Theorem 5.1.
- $\mathcal{S} = D_n$: $\phi_i(A) = C_i P^t A P$ where P is a permutation matrix, and $C_1, \dots, C_m$ are diagonal matrices with $C_1 \cdots C_m = I$ (Theorem 6.2).

$m = 2$:

- $\mathcal{S} = P_n$ or $\overline{P_n}$: $\phi_1(A) = M^* A M$, $\phi_2(A) = M^{-1} A M^{-*}$; or $\phi_1(A) = M^* A^t M$, $\phi_2(A) = M^{-1} A^t M^{-*}$ (Theorem 4.3).
- $\mathcal{S} = D_n$: $\phi_1(A) = \mathrm{diag}(N \mathrm{diag}^{-1}(A))$, $\phi_2(A) = \mathrm{diag}(N^{-t} \mathrm{diag}^{-1}(A))$ (Theorem 6.1).

Corollary 4.5 describes maps $\phi_i : P_n \to \overline{P_n}$ $(i = 1, \dots, m,\ m \geq 2)$ that satisfy

$$\mathrm{tr}\left(\phi_1(A_1)^{\alpha_1} \cdots \phi_m(A_m)^{\alpha_m}\right) = \mathrm{tr}\left(A_1^{\beta_1} \cdots A_m^{\beta_m}\right), \quad \forall A_1, \dots, A_m \in P_n, \tag{1.5}$$

for given nonzero real numbers $\alpha_1, \dots, \alpha_m, \beta_1, \dots, \beta_m$.

All above results have the corresponding real versions. We discuss them after the complex versions in each section.

2. Preliminary

2.1. **Two maps preserving trace of product.** We first give a general result about two maps preserving trace of product. The theorem is a powerful tool to prove linear bijectivity of various maps preserving trace of product in our later discussions. Recall that if S is a subset of a vector space, then $\langle S \rangle$ denotes the subspace spanned by S.

Theorem 2.1. *Let $\phi : V_1 \to W_1$ and $\psi : V_2 \to W_2$ be two maps between subsets of matrix spaces over a field $\mathbb{F}$ such that:*

(1) $\dim\langle V_1 \rangle = \dim\langle V_2 \rangle \geq \max\{\dim\langle W_1 \rangle, \dim\langle W_2 \rangle\}$.
(2) *AB are well-defined square matrices for $(A, B) \in (V_1 \times V_2) \cup (W_1 \times W_2)$.*

(3) *If* $A \in \langle V_1 \rangle$ *satisfies that* $\operatorname{tr}(AB) = 0$ *for all* $B \in \langle V_2 \rangle$*, then* $A = 0$.
(4) ϕ *and* ψ *satisfy that*

$$\operatorname{tr}(\phi(A)\psi(B)) = \operatorname{tr}(AB), \quad A \in V_1,\ B \in V_2. \tag{2.1}$$

Then $\dim\langle V_1 \rangle = \dim\langle V_2 \rangle = \dim\langle W_1 \rangle = \dim\langle W_2 \rangle$ *and* ϕ *and* ψ *can be extended to bijective linear map* $\widetilde{\phi} : \langle V_1 \rangle \to \langle W_1 \rangle$ *and* $\widetilde{\psi} : \langle V_2 \rangle \to \langle W_2 \rangle$*, respectively, such that*

$$\operatorname{tr}(\widetilde{\phi}(A)\widetilde{\psi}(B)) = \operatorname{tr}(AB), \quad A \in \langle V_1 \rangle,\ B \in \langle V_2 \rangle. \tag{2.2}$$

Proof. Let $\{A_1, \ldots, A_m\} \subseteq V_1$ be a basis of $\langle V_1 \rangle$, and $\{B_1, \ldots, B_m\} \subseteq V_2$ a basis of $\langle V_2 \rangle$, respectively. By assumptions (3) and (4), the matrix

$$[\operatorname{tr}(\phi(A_i)\psi(B_j))]_{m\times m} = [\operatorname{tr}(A_iB_j)]_{m\times m} \tag{2.3}$$

is invertible. So both sets $\{\phi(A_1), \ldots, \phi(A_m)\} \subseteq W_1$ and $\{\psi(B_1), \ldots, \psi(B_m)\} \subseteq W_2$ are linearly independent. By assumption (1), $\dim\langle V_1 \rangle = \dim\langle V_2 \rangle = \dim\langle W_1 \rangle = \dim\langle W_2 \rangle$, and $\{\phi(A_1), \ldots, \phi(A_m)\}$ (resp. $\{\psi(B_1), \ldots, \psi(B_m)\}$) is a basis of W_1 (resp. W_2.)

Let $\widetilde{\phi} : \langle V_1 \rangle \to \langle W_1 \rangle$ and $\widetilde{\psi} : \langle V_2 \rangle \to \langle W_2 \rangle$ be the linear maps defined in the way that for all $c_1, \ldots, c_m \in \mathbb{F}$:

$$\widetilde{\phi}\left(\sum_{i=1}^m c_iA_i\right) := \sum_{i=1}^m c_i\phi(A_i), \qquad \widetilde{\psi}\left(\sum_{i=1}^m c_iB_i\right) := \sum_{i=1}^m c_i\psi(B_i). \tag{2.4}$$

If a linear combination $\sum_{i=1}^m c_iA_i \in V_1$, then for all $j = 1, \ldots, m$,

$$\begin{aligned}
\operatorname{tr}\left[\widetilde{\phi}\left(\sum_{i=1}^m c_iA_i\right)\psi(B_j)\right] &= \sum_{i=1}^m c_i \operatorname{tr}\left(\phi(A_i)\psi(B_j)\right) \\
&= \sum_{i=1}^m c_i \operatorname{tr}(A_iB_j) = \operatorname{tr}\left[\left(\sum_{i=1}^m c_iA_i\right)B_j\right] \\
&= \operatorname{tr}\left[\phi\left(\sum_{i=1}^m c_iA_i\right)\psi(B_j)\right].
\end{aligned}$$

So by assumption (3) and (2.3), $\widetilde{\phi}\left(\sum_{i=1}^m c_iA_i\right) = \phi\left(\sum_{i=1}^m c_iA_i\right)$. Therefore, $\widetilde{\phi} : \langle V_1 \rangle \to \langle W_1 \rangle$ is a linear bijection and an extension of $\phi : V_1 \to W_1$. Similarly, $\widetilde{\psi} : \langle V_2 \rangle \to \langle W_2 \rangle$ is a linear bijection and an extension of $\psi : V_2 \to W_2$. □

Theorem 2.1 can be applied to [1, Lemma 3.5] which says that: if U and V are nonempty open subsets of $M_{m,n}$ and $M_{n,m}$ respectively, $\phi : U \to M_{p,q}$ and $\psi : V \to M_{q,p}$ satisfy that $\operatorname{tr}(\phi(A)\psi(B)) = \operatorname{tr}(AB)$ for all $A \in U$ and $B \in V$, and $mn \geq pq$, then ϕ and ψ can be extended to linear bijections $M_{m,n} \to M_{p,q}$ and $M_{n,m} \to M_{q,p}$ respectively. Theorem 2.1 on maps $M_n \to M_n$ also generalizes the result: if a map $\phi : M_n \to M_n$ satisfies $\operatorname{tr}(\phi(A)\phi(B)) = \operatorname{tr}(AB)$ for all $A, B \in M_n$, then ϕ must be a linear bijection in Molnár [14, proof of Theorem 1] and in J. T. Chan, C. K. Li, and N. S. Sze [4, Prop. 1.1].

A subset V of M_n is closed under Hermitian conjugate if $\{A^* : A \in V\} \subseteq V$. The following are some examples:

(1) Each of M_n, H_n, P_n, $\overline{P_n}$, D_n, and the sets of real/complex symmetric/skew-symmetric matrices, is closed under Hermitian conjugate.
(2) A combination of intersections, unions, sums, direct sums, Kronecker products and compounds of matrices in some sets closed under Hermitian conjugate will generate a set closed under Hermitian conjugate.

A real or complex matrix space V is closed under Hermitian conjugate if and only if V equals the direct sum of subspace of Hermitian matrices and subspace of skew-Hermitian matrices.

Corollary 2.2. *Let V be a subset of M_n closed under Hermitian conjugate. Suppose two maps $\phi, \psi : V \to V$ satisfy that*

$$\operatorname{tr}(\phi(A)\psi(B)) = \operatorname{tr}(AB), \quad A, B \in V. \tag{2.5}$$

Then ϕ and ψ can be extended to linear bijections on $\langle V\rangle$. Moreover, when V is a vector space, every linear bijection $\phi : V \to V$ corresponds to a unique linear bijection $\psi : V \to V$ such that (2.5) holds. Explicitly, given an orthonormal basis $\{A_1, \ldots, A_\ell\}$ of V with respect to the inner product $\langle A, B\rangle = \operatorname{tr}(A^*B)$, ψ is defined by $\psi(A_i) = B_i$ in which $\{B_1, \ldots, B_\ell\}$ is a basis of V with $\operatorname{tr}(\phi(A_i^*)B_j) = \delta_{i,j}$ for all $i, j \in \{1, \ldots, \ell\}$.

A similar result of Theorem 2.1 is obtained by assuming linearity of ϕ and ψ but dropping the dimension restriction on their codomains, in which ϕ and ψ can be extended to injective linear maps preserving trace of product. A map $\phi : V \to W$ between subsets of vector spaces over $\mathbb{F}$ is called linear if $v_1, \ldots, v_k, c_1v_1 + \cdots + c_kv_k \in V$ in which $c_1, \ldots, c_k \in \mathbb{F}$ implies that

$$\phi(c_1v_1 + \cdots + c_kv_k) = c_1\phi(v_1) + \cdots + c_k\phi(v_k). \tag{2.6}$$

Theorem 2.3. *Let $\phi : V_1 \to W_1$ and $\psi : V_2 \to W_2$ be two linear maps between subsets of matrix spaces over a field $\mathbb{F}$ such that:*

(1) $\dim\langle V_1\rangle = \dim\langle V_2\rangle$.
(2) *AB are well-defined square matrices for $(A, B) \in (V_1 \times V_2) \cup (W_1 \times W_2)$.*
(3) *If $A \in \langle V_1\rangle$ satisfies that $\operatorname{tr}(AB) = 0$ for all $B \in \langle V_2\rangle$, then $A = 0$.*
(4) *ϕ and ψ satisfy that*

$$\operatorname{tr}(\phi(A)\psi(B)) = \operatorname{tr}(AB), \quad A \in V_1,\ B \in V_2. \tag{2.7}$$

Then $\dim\langle V_1\rangle = \dim\langle V_2\rangle \le \min\{\dim\langle W_1\rangle, \dim\langle W_2\rangle\}$ and ϕ and ψ can be extended to two injective linear maps $\widetilde{\phi} : \langle V_1\rangle \to \langle W_1\rangle$ and $\widetilde{\psi} : \langle V_2\rangle \to \langle W_2\rangle$, respectively, such that

$$\operatorname{tr}(\widetilde{\phi}(A)\widetilde{\psi}(B)) = \operatorname{tr}(AB), \quad A \in \langle V_1\rangle,\ B \in \langle V_2\rangle. \tag{2.8}$$

Proof. We have $\dim\langle V_1\rangle = \dim\langle V_2\rangle \ge \max\{\dim\langle \operatorname{Im}\phi\rangle, \dim\langle \operatorname{Im}\psi\rangle\}$. View ϕ and ψ as $\phi : V_1 \to \operatorname{Im}\phi$ and $\psi : V_2 \to \operatorname{Im}\psi$ and apply Theorem 2.1. □

Corollary 2.4. *Let V and W be subsets of matrix spaces in which V is closed under Hermitian conjugate. Suppose two linear maps $\phi, \psi : V \to W$ satisfy that*

$$\operatorname{tr}(\phi(A)\psi(B)) = \operatorname{tr}(AB), \quad A, B \in V. \tag{2.9}$$

Then ϕ and ψ can be extended to injective linear maps $\widetilde{\phi}, \widetilde{\psi} : \langle V\rangle \to \langle W\rangle$ preserving trace of product. In particular, $\dim\langle V\rangle \leq \dim\langle W\rangle$.

3. Maps on M_n

Corollary 2.2 shows that every linear bijection $\phi : M_n \to M_n$ corresponds to another linear bijection $\psi : M_n \to M_n$ such that $\operatorname{tr}(\phi(A)\psi(B)) = \operatorname{tr}(AB)$ for all $A, B \in M_n$. If there are three or more maps $M_n \to M_n$ preserve trace of products, they can be explicitly described as follow.

Theorem 3.1. *Let $m, n \in \mathbb{Z}_+$ and $m \geq 3$. Maps $\phi_1, \ldots, \phi_m : M_n \to M_n$ satisfy that*

$$\operatorname{tr}(\phi_1(A_1)\cdots\phi_m(A_m)) = \operatorname{tr}(A_1\cdots A_m), \qquad \forall A_1, \ldots, A_m \in M_n, \tag{3.1}$$

if and only if there are invertible matrices $N_1, \ldots, N_m \in M_n$ and $N_{m+1} := N_1$ such that

$$\phi_i(A) = N_i A N_{i+1}^{-1}, \quad \forall A \in M_n. \tag{3.2}$$

Proof. It suffices to prove that (3.1) implies (3.2). Let $\psi_1(B) := \phi_2(B)\phi_3(I)\cdots\phi_m(I)$. Then $\operatorname{tr}(\phi_1(A)\psi_1(B)) = \operatorname{tr}(AB)$ for $A, B \in M_n$. By Corollary 2.2, ϕ_1 and ψ_1 are linear bijections. So $\phi_3(I), \cdots, \phi_m(I)$ are invertible. Similarly, all ϕ_i's are linear bijections and all $\phi_i(I)$'s are invertible.

For all $A, B_2, \ldots, B_m \in M_n$ we have

$$\operatorname{tr}(\phi_1(A)\phi_2(B_2)\cdots\phi_m(B_m)) = \operatorname{tr}(AB_2\cdots B_m) = \operatorname{tr}(\phi_1(A)\psi_1(B_2\cdots B_m)).$$

Since $\operatorname{Im}\phi_1 = M_n$, we get

$$\phi_2(B_2)\cdots\phi_m(B_m) = \psi_1(B_2\cdots B_m) = \phi_2(B_2\cdots B_m)\phi_3(I)\cdots\phi_m(I). \tag{3.3}$$

Let $B_4 = \cdots = B_m = I$ then

$$\phi_2(B_2)\phi_3(B_3) = \phi_2(B_2B_3)\phi_3(I). \tag{3.4}$$

Let $B_2 = I$ in (3.4), then $\phi_2(I)\phi_3(B_3) = \phi_2(B_3)\phi_3(I)$ so that $\phi_3(B_3) = \phi_2(I)^{-1}\phi_2(B_3)\phi_3(I)$. So (3.4) becomes $\phi_2(B_2)\phi_2(I)^{-1}\phi_2(B_3) = \phi_2(B_2B_3)$. Let $\widehat{\phi_2}(A) := \phi_2(A)\phi_2(I)^{-1}$ then $\widehat{\phi_2}(I) = I$ and

$$\widehat{\phi_2}(A)\widehat{\phi_2}(B) = \widehat{\phi_2}(AB), \quad \forall A, B \in M_n. \tag{3.5}$$

So $\widehat{\phi_2}$ is an algebra automorphism of M_n, which must be of the form $\widehat{\phi_2}(A) = N_2AN_2^{-1}$ for an invertible $N_2 \in M_n$. Let $N_3 := \phi_2(I)^{-1}N_2$ and $N_4 := \phi_3(I)^{-1}N_3$. Then

$$\phi_2(A) = \widehat{\phi_2}(A)\phi_2(I) = N_2AN_3^{-1}, \tag{3.6}$$

$$\phi_3(A) = \phi_2(I)^{-1}\phi_2(A)\phi_3(I) = \phi_2(I)^{-1}N_2AN_3^{-1}\phi_3(I) = N_3AN_4^{-1}. \tag{3.7}$$

Similarly, there are invertible $N_5, \dots, N_m, N_{m+1}$ and $N_1 := N_{m+1}$ such that $\phi_i(A) = N_i A N_{i+1}^{-1}$ for all $i = 1, \dots, m$. □

Remark 3.2. *Theorem 3.1 also holds if all M_n are replaced by $M_n(\mathbb{R})$, and the proof is analogous.*

The following two examples show that two maps $M_n \to M_n$ preserving trace of product may not be of the form (3.2).

Example 3.3. *Suppose that ϕ and ψ are two maps on M_n which are defined by $\phi(A) = A \circ C, \quad \psi(A) = A \circ \hat{C}$, where $C = (c_{ij}) \in S_n(\mathbb{R})$ and $\hat{C} = (\frac{1}{c_{ij}})$ with $c_{ij} \neq 0$ for all i, j. Here, $A \circ C$ denotes the Hadamard product of A and C. Then the maps ϕ and ψ send from H_n to H_n and S_n to S_n. Moreover, one can show that ϕ and ψ satisfy the equality*

$$\operatorname{tr} \phi(A)\psi(B) = \operatorname{tr} AB, \quad \forall A, B \in M_n.$$

Example 3.4. *Let $A_1, \dots, A_n, B_1, \dots, B_n \in M_n$ be invertible matrices with $A_i B_i = I_n$ for all $i = 1, \dots, n$. Let $\{P_{ij} : 1 \leq i, j \leq n\}$ and $\{Q_{ij} : 1 \leq i, j \leq n\}$ be the subsets of M_n, where $P_{ij} = E_{ij} A_i$ and $Q_{ij} = B_j E_{ij}$. Define two maps ϕ and ψ on M_n by*

$$\phi(\sum_{i,j=1}^{n} c_{ij} E_{ij}) = \sum_{i,j=1}^{n} c_{ij} P_{ij}, \quad \psi(\sum_{i,j=1}^{n} d_{ij} E_{ij}) = \sum_{i,j=1}^{n} d_{ij} Q_{ij}.$$

Then one can show that

$$\operatorname{tr} \phi(A)\psi(B) = \operatorname{tr} AB, \quad \forall A, B \in M_n.$$

3.1. **Linear Maps $M_n \to M_k$.** Now let $m \geq 2$ and we further consider linear maps $\phi_1, \dots, \phi_m : M_n \to M_k$ preserving trace of products:

$$\operatorname{tr}(\phi_1(A_1)\phi_2(A_2) \cdots \phi_m(A_m)) = \operatorname{tr}(A_1 A_2 \cdots A_m), \quad A_1, A_2, \dots, A_m \in M_n. \tag{3.8}$$

Theorem 3.5. *When $n > k$ and $m \geq 2$, there are no linear maps $\phi_1, \dots, \phi_m : M_n \to M_k$ that satisfy (3.8).*

Proof. Suppose on the contrary, there exist linear maps $\phi_1, \dots, \phi_m : M_n \to M_k$ that satisfy $\operatorname{tr}(\phi_1(A_1) \cdots \phi_m(A_m)) = \operatorname{tr}(A_1 \cdots A_m)$ for all $A_1, \dots, A_m \in M_n$. Let

$$\psi(B) := \phi_2(B)\phi_3(I_n) \cdots \phi_m(I_n).$$

Then both ϕ_1 and ψ are linear maps and $\operatorname{tr}(\phi_1(A)\psi(B)) = \operatorname{tr}(AB)$ for $A, B \in M_n$. Let $\{B_1, B_2, \dots, B_{n^2}\}$ be a basis of M_n. The matrix $\Big(\operatorname{tr}(B_i B_j)\Big)_{n^2 \times n^2}$ has full rank n^2, but the matrix $\Big(\operatorname{tr}(\phi_1(B_i)\psi(B_j))\Big)_{n^2 \times n^2}$ has rank no more than $k^2 < n^2$. This is a contradiction. □

Let 0_n denote the zero matrix of size $n \times n$.

Theorem 3.6. *Suppose $n \le k$ and linear maps $\phi_1, \phi_2 : M_n \to M_k$ satisfy (3.8) for $m = 2$:*

$$\operatorname{tr}(\phi_1(A)\phi_2(B)) = \operatorname{tr}(AB), \quad A, B \in M_n. \tag{3.9}$$

Then there are linear bijections $\widetilde{\phi}_1, \widetilde{\phi}_2 : M_k \to M_k$ such that

$$\operatorname{tr}(\widetilde{\phi}_1(\widetilde{A})\widetilde{\phi}_2(\widetilde{B})) = \operatorname{tr}(\widetilde{A}\widetilde{B}), \quad \widetilde{A}, \widetilde{B} \in M_k, \tag{3.10}$$

$$\widetilde{\phi}_i(A \oplus 0_{k-n}) = \phi_i(A), \quad A \in M_n,\ i = 1, 2. \tag{3.11}$$

Moreover, if ϕ_1 and ϕ_2 both preserve Hermitian matrices, then $\widetilde{\phi}_1$ and $\widetilde{\phi}_2$ can be chosen to preserve Hermitian matrices.

Proof. Denote

$$V := \{A \oplus 0_{k-n} : A \in M_n\}, \tag{3.12}$$

$$W := \left\{ \begin{pmatrix} 0_n & B \\ C & D \end{pmatrix}_{k\times k} : B \in M_{n,k-n}, C \in M_{k-n,n}, D \in M_{k-n} \right\}. \tag{3.13}$$

Then $M_k = V \oplus W$ and $\operatorname{tr}(\widetilde{A}\widetilde{B}) = 0$ for all $\widetilde{A} \in V$ and $\widetilde{B} \in W$. Let $\{\widetilde{C_1}, \dots, \widetilde{C_{k^2-n^2}}\}$ be a basis of W. The matrix $\left(\operatorname{tr}(\widetilde{C_p}\widetilde{C_q})\right)_{(k^2-n^2)\times(k^2-n^2)}$ is invertible.

For $i = 1, 2$, let

$$X_i := \left\{ \widetilde{B} \in M_k : \operatorname{tr}(\phi_i(A)\widetilde{B}) = 0 \text{ for all } A \in M_n \right\}. \tag{3.14}$$

The (3.9) implies that $\operatorname{Im}(\phi_1) \cap X_2 = \operatorname{Im}(\phi_2) \cap X_1 = \{0_k\}$. Corollary 2.4 shows that ϕ_1 and ϕ_2 are injections. Hence

$$M_k = \operatorname{Im}(\phi_1) \oplus X_2 = \operatorname{Im}(\phi_2) \oplus X_1. \tag{3.15}$$

Fix a basis $\{\widetilde{E_1}, \dots, \widetilde{E_{k^2-n^2}}\}$ of X_1. There is a unique basis $\{\widetilde{F_1}, \dots, \widetilde{F_{k^2-n^2}}\}$ of X_2 such that

$$\left(\operatorname{tr}(\widetilde{E_i}\widetilde{F_j})\right)_{(k^2-n^2)\times(k^2-n^2)} = \left(\operatorname{tr}(\widetilde{C_i}\widetilde{C_j})\right)_{(k^2-n^2)\times(k^2-n^2)}. \tag{3.16}$$

Let $\widetilde{\phi}_1, \widetilde{\phi}_2 : M_k \to M_k$ be the linear maps defined by

- $\widetilde{\phi}_i(A \oplus 0_{k-n}) = \phi_i(A)$ for $A \in M_n$ and $i = 1, 2$;
- $\widetilde{\phi}_1(\widetilde{C_j}) = \widetilde{E_j}$ and $\widetilde{\phi}_2(\widetilde{C_j}) = \widetilde{F_j}$ for $j = 1, \dots, k^2 - n^2$.

Then $\widetilde{\phi}_1$ and $\widetilde{\phi}_2$ are linear bijections, $\widetilde{\phi}_i(A \oplus 0_{k-n}) = \phi_i(A)$ for $A \in M_n$ and $i = 1, 2$, and (3.10) holds.

Finally, suppose ϕ_1 and ϕ_2 both preserve Hermitian matrices. Since M_n has a basis consisting of Hermitian matrices, so do $\operatorname{Im}(\phi_1)$ and $\operatorname{Im}(\phi_2)$. We can let the sets $\{\widetilde{C_1}, \dots, \widetilde{C_{k^2-n^2}}\}$ and $\{\widetilde{E_1}, \dots, \widetilde{E_{k^2-n^2}}\}$ consist entirely of Hermitian matrices. Then each $\widetilde{F_j}$ in the set $\{\widetilde{F_1}, \dots, \widetilde{F_{k^2-n^2}}\}$ satisfies that

$$\operatorname{tr}(\phi_2(A)\widetilde{F_j}) = 0 \in \mathbb{R}, \quad \operatorname{tr}(\widetilde{E_i}\widetilde{F_j}) = \operatorname{tr}(\widetilde{C_i}\widetilde{C_j}) \in \mathbb{R},$$

which is possible only when each $\widetilde{F_j}$ is a Hermitian matrix. Therefore, the maps $\widetilde{\phi}_1$ and $\widetilde{\phi}_2$ constructed above preserve Hermitian matrices. □

Example 3.7. *Analogous linear map extension result of Theorem 3.6 for $m > 2$ does not hold. Consider $k = 2n$ and $m = 3$. Let $X \in M_n$ be an arbitrary nonscalar matrix. Let $\phi_1, \phi_2, \phi_3 : M_n \to M_{2n}$ such that*

$$\phi_1(A) = A \oplus 0_n, \quad \phi_2(B) = B \oplus B, \quad \phi_3(C) = C \oplus XCX^*, \quad A, B, C \in M_n.$$

Obviously,

$$\operatorname{tr}(\phi_1(A)\phi_2(B)\phi_3(C)) = \operatorname{tr}(ABC), \quad A, B, C \in M_n.$$

Suppose there are linear bijections $\widetilde{\phi}_1, \widetilde{\phi}_2, \widetilde{\phi}_3 : M_{2n} \to M_{2n}$ such that $\widetilde{\phi}_i(A \oplus 0_n) = \phi_i(A)$ for $A \in M_n$ and $i = 1, 2, 3$, and

$$\operatorname{tr}(\widetilde{\phi}_1(\widetilde{A})\widetilde{\phi}_2(\widetilde{B})\widetilde{\phi}_3(\widetilde{C})) = \operatorname{tr}(\widetilde{A}\widetilde{B}\widetilde{C}), \quad \widetilde{A}, \widetilde{B}, \widetilde{C} \in M_{2n}.$$

Then by (3.4), it is necessary that for all $B, C \in M_n$,

$$\begin{aligned} \widetilde{\phi}_2(B \oplus 0_n)\widetilde{\phi}_3(C \oplus 0_n) &= \widetilde{\phi}_2(BC \oplus 0_n)\widetilde{\phi}_3(I_{2n}), \quad \textit{that is,} \\ (B \oplus B)(C \oplus XCX^*) = BC \oplus BXCX^* &= (BC \oplus BC)\widetilde{\phi}_3(I_{2n}). \end{aligned} \tag{3.17}$$

However, since X is selected from nonscalar matrices, no matrix $\widetilde{\phi}_3(I_{2n}) \in M_{2n}$ can satisfy (3.17) for all $B, C \in M_n$. So the statement in Theorem 3.6 is not true if $m > 2$. This example also indicates that analogous statement on maps $H_n \to H_k$ is not true for $m > 2$ either.

4. Maps on H_n and P_n

In this section, we investigate maps $H_n \to H_n$ and $P_n \to \overline{P_n}$ that preserve a trace of products property. These two types of maps lead to different results, especially in the trace of products of two maps.

4.1. Maps $H_n \to H_n$ preserving trace of products.

By Corollary 2.2, every linear bijection $\phi : H_n \to H_n$ corresponds to another linear bijection $\psi : H_n \to H_n$ such that $\operatorname{tr}(\phi(A)\psi(B)) = \operatorname{tr}(AB)$ for all $A, B \in H_n$. Some examples can be analogously developed from Example 3.3.

We investigate the problem of three or more maps $H_n \to H_n$ preserving trace of products.

Theorem 4.1. *Suppose $m \geq 3$. Maps $\phi_1, \ldots, \phi_m : H_n \to H_n$ satisfy that*

$$\operatorname{tr}(\phi_1(A_1) \cdots \phi_m(A_m)) = \operatorname{tr}(A_1 \cdots A_m), \quad \forall A_1, \ldots, A_m \in H_n, \tag{4.1}$$

if and only if they can be described as follow:

(1) *When m is odd, there exist a unitary matrix $U \in M_n$ and nonzero real scalars $c_1, \ldots, c_m$ such that $c_1 \cdots c_m = 1$ and*

$$\phi_i(A) = c_i U^* A U, \quad \forall A \in H_n, \ i = 1, \ldots, m. \tag{4.2}$$

(2) *When m is even, there exist an invertible $M \in M_n$ and nonzero real scalars $c_1, \dots, c_m$ such that $c_1 \cdots c_m = 1$ and*

$$\phi_i(A) = \begin{cases} c_i M^* A M, & i \text{ is odd}, \\ c_i M^{-1} A M^{-*}, & i \text{ is even}, \end{cases} \quad \forall A \in H_n. \tag{4.3}$$

Proof. It suffices to prove that (4.1) implies (4.2) (when m is odd) or (4.3) (when m is even).

Let $\psi_1 : H_n \to M_n$ be defined by

$$\psi_1(B) = \phi_2(B)\phi_3(I) \cdots \phi_m(I), \qquad B \in H_n.$$

The (4.1) implies that $\operatorname{tr}(\phi_1(A)\psi_1(B)) = \operatorname{tr}(AB)$ for $A, B \in H_n$. By Theorem 2.1 and the fact that the complex span of H_n is M_n, ϕ_1 can be extended to a linear bijection $\widetilde{\phi}_1$ on M_n. Similarly, each ϕ_i $(1 \le i \le m)$ can be extended to a linear bijection $\widetilde{\phi}_i$ on M_n.

Now both sides of (4.1) can be viewed as multilinear functions on $(A_1, \dots, A_m) \in {H_n}^{\times m}$. Therefore, $\widetilde{\phi}_1, \dots, \widetilde{\phi}_m$ satisfy equality (3.1) after replacing each ϕ_i with $\widetilde{\phi}_i$. Theorem 3.1 says that there are invertible $N_1, \dots, N_m \in M_n$ and $N_{m+1} := N_1$ such that each $\widetilde{\phi}_i(A) = N_i A N_{i+1}^{-1}$ for $A \in M_n$. So for all $A \in H_n$, $\phi_i(A) = N_i A N_{i+1}^{-1} \in H_n$. This could happen if and only if $N_{i+1}^{-1} = c_i' N_i^*$ for a nonzero real scalar c_i', in which $\phi_i(A) = c_i' N_i A N_i^*$. Let $M := N_1^*$. Then $\phi_1(A) = c_1 M^* A M$ for a nonzero real number c_1, and $\phi_2(A) = c_2 M^{-1} A M^{-*}$, $\phi_3(A) = c_3 M^* A M$, and so on.

(1) When m is odd, $\phi_m(A) = c_m M^* A M = N_m A N_{m+1}^{-1} = N_m A N_1^{-1} = N_m A M^{-*}$. So $M = cM^{-*}$ or $MM^* = cI$ for a scalar c. By adjusting the scalar, we may assume that $U := M$ is unitary, and ϕ_i are given by (4.2). Then necessarily $c_1 \cdots c_m = 1$.
(2) When m is even, we get the form (4.3). Clearly $c_1 \cdots c_m = 1$. □

Remark 4.2. *Real Hermitian matrices are real symmetric matrices. The real version of Theorem 4.1 also holds. See Theorem 5.2 on $S_n(\mathbb{R})$.*

4.2. **Maps $P_n \to \overline{P_n}$ preserving trace of products.** Now consider the maps $P_n \to \overline{P_n}$ that preserve trace of products. Many properties of these maps can be generalized to their counterparts for the maps $P_n \to P_n$ and $\overline{P_n} \to \overline{P_n}$. Note that linear maps $\overline{P_n} \to \overline{P_n}$ are positive maps, which are popular and useful in quantum information theory.

Unlike the situations in $M_n \to M_n$ and $H_n \to H_n$, two maps $\phi, \psi : P_n \to \overline{P_n}$ (similarly for $P_n \to P_n$ and $\overline{P_n} \to \overline{P_n}$) preserving trace of products have the following special forms.

Theorem 4.3. *Two maps $\phi : P_n \to \overline{P_n}$ and $\psi : P_n \to \overline{P_n}$ satisfy*

$$\operatorname{tr}(\phi(A)\psi(B)) = \operatorname{tr}(AB), \quad \forall A, B \in P_n, \tag{4.4}$$

if and only if the following claims hold:

(1) *If $\phi(I) = I$, then there exists a unitary matrix $U \in M_n$ such that*

$$\phi(A) = \psi(A) = U^* A U \quad \text{or} \quad \phi(A) = \psi(A) = U^* A^t U, \quad \forall A \in P_n. \tag{4.5}$$

(2) *In general, there exists an invertible $M \in M_n$ such that for all $A \in P_n$:*

$$\phi(A) = M^*AM,\ \psi(A) = M^{-1}AM^{-*}; \quad \textit{or} \tag{4.6a}$$

$$\phi(A) = M^*A^tM,\ \psi(A) = M^{-1}A^tM^{-*}. \tag{4.6b}$$

Proof. The maps defined by (4.5) and (4.6) obviously satisfy (4.4). We shall prove the converse. By Theorem 2.1, ϕ and ψ can be extended to linear bijections $H_n \to H_n$, still denoted as ϕ and ψ here, that satisfy (4.4).

(1) If $\phi(I) = I$, then (4.4) implies that $\operatorname{tr}\psi(A) = \operatorname{tr} A$ for $A \in P_n$. Given a rank one projection P, we have $\phi(P), \psi(P) \in \overline{P_n}$ so that

$$\begin{aligned} 1 &= \operatorname{tr} P = \operatorname{tr}(P^2) = \operatorname{tr}(\phi(P)\psi(P)) && (4.7)\\ &\le [\operatorname{tr}(\phi(P)^2)\operatorname{tr}(\psi(P)^2)]^{1/2} && (4.8)\\ &\le \operatorname{tr}\phi(P)\operatorname{tr}\psi(P) = \operatorname{tr}\phi(P)\operatorname{tr}P = \operatorname{tr}\phi(P). && (4.9) \end{aligned}$$

Let $E_{ij} \in M_n$ be the matrix with a sole nonzero value 1 in the (i,j) entry. Write $P = WE_{11}W^*$ for a unitary W. Then I is a sum of n rank one projections $I = P + P^{(2)} + \cdots + P^{(n)}$ in which $P^{(i)} := WE_{ii}W^*$ for $2 \le i \le n$.

$$n = \operatorname{tr}\phi(I) = \operatorname{tr}\phi(P) + \operatorname{tr}\phi(P^{(2)}) + \cdots + \operatorname{tr}\phi(P^{(n)}) \ge n. \tag{4.10}$$

Therefore, $\operatorname{tr}\phi(P) = 1$. The Cauchy-Schwartz inequality (4.8) becomes equality only if $\psi(P) = c\phi(P)$ for some $c \ge 0$; the inequality in (4.9) becomes equality only if both $\phi(P)$ and $\psi(P)$ are rank one. Then $\operatorname{tr}\phi(P) = 1$ and $\operatorname{tr}\psi(P) = \operatorname{tr}P = 1$ imply that $\psi(P) = \phi(P)$ are rank one projections.

Since all rank one projections span H_n, we have $\phi(A) = \psi(A)$ for all $A \in P_n$. Then (4.4) becomes

$$\operatorname{tr}(\phi(A)\phi(B)) = \operatorname{tr}(AB), \qquad \forall A, B \in P_n. \tag{4.11}$$

By [6, Lemma 3.6], there is a unitary U such that $\phi = \psi$ are given by (4.5):

$$\phi(A) = \psi(A) = U^*AU \qquad \text{or} \qquad \phi(A) = \psi(A) = U^*A^tU.$$

(2) In general, we first show that $\phi(I)$ is invertible. Suppose on the contrary, $\phi(I)$ is not invertible. Then $\phi(I)v = 0$ for certain nonzero $v \in \mathbb{C}^n$. By the linear bijectivity of ψ on H_n, there exists a nonzero $B \in H_n$ such that $\psi(B) = vv^*$. If $B \in \overline{P_n}$ then

$$0 = \operatorname{tr}(\phi(I)vv^*) = \operatorname{tr}(\phi(I)\psi(B)) = \operatorname{tr}(B), \tag{4.12}$$

which contradicts $B \ne 0$. So $B \notin \overline{P_n}$ and B has a negative eigenvalue. Using spectral decomposition, there exists $A \in P_n$ such that $\operatorname{tr}(AB) < 0$. Then $\phi(A) \in \overline{P_n}$ and

$$0 > \operatorname{tr}(AB) = \operatorname{tr}(\phi(A)\psi(B)) \ge 0, \tag{4.13}$$

which is also a contradiction. Therefore, $\phi(I) \in \overline{P_n}$ must be invertible.

Define $\phi_1(A) := \phi(I)^{-1/2}\phi(A)\phi(I)^{-1/2}$ and $\psi_1(B) := \phi(I)^{1/2}\psi(B)\phi(I)^{1/2}$. Then $\phi_1 : P_n \to \overline{P_n}$ and $\psi_1 : P_n \to \overline{P_n}$ satisfy (4.4) and $\phi_1(I) = I$. By preceding argument,

there is a unitary $U \in M_n$ such that $\phi_1(A) = \psi_1(A) = U^*AU$ or $\phi_1(A) = \psi_1(A) = U^*A^tU$. Therefore, let $M := U\phi(I)^{1/2}$, then either $\phi(A) = M^*AM$ and $\psi(A) = M^{-1}AM^{-*}$, or $\phi(A) = M^*A^tM$ and $\psi(A) = M^{-1}A^tM^{-*}$. □

Next consider $m \geq 3$ maps $\phi_1, \dots, \phi_m$ from P_n to $\overline{P_n}$ that preserve trace of products. The following result can be proved by Theorem 4.1. We will give another proof by Theorem 4.3.

Theorem 4.4. *Let $m \geq 3$. Maps $\phi_i : P_n \to \overline{P_n}$ $(i = 1, \dots, m)$ satisfy that*

$$\operatorname{tr}(\phi_1(A_1) \cdots \phi_m(A_m)) = \operatorname{tr}(A_1 \cdots A_m), \quad \forall A_1, \dots, A_m \in P_n, \tag{4.14}$$

if and only if they are described as follow:

(1) *When m is odd, there exist a unitary matrix $U \in M_n$ and scalars $c_1, \dots, c_m \in \mathbb{R}_+$ such that $c_1 \cdots c_m = 1$ and*

$$\phi_i(A) = c_iU^*AU, \quad \forall A \in P_n,\ i = 1, \dots, m. \tag{4.15}$$

(2) *When m is even, there exist an invertible $M \in M_n$ and scalars $c_1, \dots, c_m \in \mathbb{R}_+$ such that $c_1 \cdots c_m = 1$ and*

$$\phi_i(A) = \begin{cases} c_iM^*AM, & i \text{ is odd}, \\ c_iM^{-1}AM^{-*}, & i \text{ is even}, \end{cases} \quad \forall A \in P_n. \tag{4.16}$$

Proof. It suffices to prove that (4.14) implies (4.15) when m is odd and (4.16) when m is even. Let $C := \phi_3(I) \cdots \phi_m(I)$. In Theorem 2.1, let $\phi : P_n \to \overline{P_n}$ such that $\phi(A) := \phi_1(A)$, and $\psi : P_n \to \overline{P_n}C$ such that $\psi(A) := \phi_2(A)C$. The complex spanning space $\langle P_n \rangle = M_n$. So ϕ and ψ satisfy the assumptions of Theorem 2.1. It implies that ϕ_1 and ϕ_2 can be extended to linear bijections $\widetilde{\phi_1} : M_n \to M_n$ and $\widetilde{\phi_2} : M_n \to M_n$ respectively, which necessarily map H_n to H_n bijectively.

Let $A_1, A_2 \in P_n$ and $B_1 \in H_n$. There exists $t \in \mathbb{R}_+$ such that $A_1 + tB_1 \in P_n$. (4.14) implies that

$$\begin{aligned} & \operatorname{tr}(A_1A_2) + t \operatorname{tr}(\widetilde{\phi_1}(B_1)\phi_2(A_2)C) \\ = \ & \operatorname{tr}(\phi_1(A_1)\phi_2(A_2)C) + t \operatorname{tr}(\widetilde{\phi_1}(B_1)\phi_2(A_2)C) \\ = \ & \operatorname{tr}(\phi_1(A_1 + tB_1)\phi_2(A_2)C) \\ = \ & \operatorname{tr}[(A_1 + tB_1)A_2] \in \mathbb{R}. \end{aligned}$$

Therefore, $\operatorname{tr}(\widetilde{\phi_1}(B_1)\phi_2(A_2)C) \in \mathbb{R}$ for any Hermitian matrix B_1. The bijectivity of $\widetilde{\phi_1}|_{H_n} : H_n \to H_n$ implies that $\operatorname{tr}(B\phi_2(A_2)C) \in \mathbb{R}$ for all $B \in H_n$.

Write $\phi_2(A_2)C = A' + \mathbf{i}A''$ for $A', A'' \in H_n$. Then

$$\operatorname{tr}(A''A') + \mathbf{i} \operatorname{tr}(A''A'') = \operatorname{tr}(A''\phi_2(A_2)C) \in \mathbb{R}.$$

So $\operatorname{tr}(A''A'') = 0$ and $A'' = 0$. Hence $\phi_2(A_2)C \in H_n$ for all $A_2 \in P_n$.

Every Hermitian matrix is the difference of two positive definite matrices. Thus for all $B \in H_n$ we have $\widetilde{\phi_2}(B)C \in H_n$. The bijectivity of $\widetilde{\phi_2}|_{H_n} : H_n \to H_n$ implies that $AC \in H_n$ for all $A \in H_n$. This is true only when $\phi_3(I)\cdots\phi_m(I) = C = cI$ for some $c \in \mathbb{R}$; and (4.14) implies that $c \in \mathbb{R}_+$. Let $c_{1,2} := \frac{1}{c}$. Then $c_{1,2} \in \mathbb{R}_+$ and (4.14) becomes

$$\operatorname{tr}(\phi_1(A_1)\phi_2(A_2)) = c_{1,2}\operatorname{tr}(A_1A_2), \quad \forall A_1, A_2 \in P_n. \tag{4.17}$$

For each $i \in \{1, 2, \dots, m\}$, the identity (4.14) is equivalent to

$$\operatorname{tr}(\phi_i(A_i)\cdots\phi_m(A_m)\phi_1(A_1)\cdots\phi_{i-1}(A_{i-1})) = \operatorname{tr}(A_i\cdots A_mA_1\cdots A_{i-1}) \tag{4.18}$$

for all $A_1, \dots, A_m \in P_n$. Similarly, we can prove that

$$\phi_{i+2}(I)\cdots\phi_m(I)\phi_1(I)\cdots\phi_{i-1}(I) = \frac{1}{c_{i,i+1}}I \tag{4.19}$$

for some $c_{i,i+1} \in \mathbb{R}_+$. So (4.18) implies that

$$\operatorname{tr}(\phi_i(A_i)\phi_{i+1}(A_{i+1})) = c_{i,i+1}\operatorname{tr}(A_iA_{i+1}), \quad \forall A_i, A_{i+1} \in P_n. \tag{4.20}$$

Theorem 4.3 and (4.17) indicate that there exist an invertible $M \in M_n$ and $c_1, c_2 \in \mathbb{R}_+$ such that

$$\phi_1(A) = c_1M^*AM, \ \phi_2(A) = c_2M^{-1}AM^{-*}; \quad \text{or} \tag{4.21}$$

$$\phi_1(A) = c_1M^*A^tM, \ \phi_2(A) = c_2M^{-1}A^tM^{-*}. \tag{4.22}$$

(1) If (4.21) holds, then (4.20) for $i = 2$ implies that

$$\begin{aligned}\operatorname{tr}(A_2M^{-*}\phi_3(A_3)M^{-1}) &= \operatorname{tr}(M^{-1}A_2M^{-*}\phi_3(A_3))\\ &= \frac{1}{c_2}\operatorname{tr}(\phi_2(A_2)\phi_3(A_3)) = \frac{c_{2,3}}{c_2}\operatorname{tr}(A_2A_3)\end{aligned}$$

for all $A_2, A_3 \in P_n$. So $\phi_3(A) = c_3M^*AM$ for some $c_3 \in \mathbb{R}_+$. Similarly,

$$\phi_i(A) = \begin{cases} c_iM^*AM & \text{if } \ 2 \nmid i,\\ c_iM^{-1}AM^{-*} & \text{if } \ 2 \mid i,\end{cases} \quad \text{for } \ i \in \{1, \dots, m\}. \tag{4.23}$$

If m is even, then (4.14) implies that $c_1c_2\cdots c_m = 1$; so (4.16) holds. If m is odd, then (4.20) for $i = 1$ and $i = m$ imply that there are $c_1, c_{m+1} \in \mathbb{R}_+$ such that $c_1M^*AM = c_{m+1}M^{-1}AM^{-*}$ and thus

$$c_1MM^*AMM^* = c_{m+1}A, \quad \forall A \in P_n.$$

Hence MM^* is a scalar matrix. We may adjust the matrix M and the scalars c_i's such that $U := M$ is unitary, and $c_1\cdots c_m = 1$. So (4.15) holds.

(2) If (4.22) holds, then similarly we have

$$\phi_i(A) = \begin{cases} c_iM^*A^tM & \text{if } \ 2 \nmid i,\\ c_iM^{-1}A^tM^{-*} & \text{if } \ 2 \mid i,\end{cases} \quad \text{for } \ i \in \{1, \dots, m\}. \tag{4.24}$$

Moreover, $c_1 \cdots c_m = 1$, and M is unitary when m is odd. However, when $m \geq 3$, none of the expressions (4.24) satisfies (4.14). □

Theorem 4.3 and Theorem 4.4 can be generalized to a result involving powers.

Corollary 4.5. *Let an integer $m \geq 2$ and nonzero real numbers $\alpha_1, \ldots, \alpha_m, \beta_1, \ldots, \beta_m$ be given. Maps $\phi_i : P_n \to \overline{P_n}$ $(i = 1, \ldots, m)$ satisfy that*

$$\operatorname{tr}\left(\phi_1(A_1)^{\alpha_1} \cdots \phi_m(A_m)^{\alpha_m}\right) = \operatorname{tr}\left(A_1^{\beta_1} \cdots A_m^{\beta_m}\right), \quad \forall A_1, \ldots, A_m \in P_n, \tag{4.25}$$

if and only if they are described as follow:

(1) *When m is odd, there exists a unitary matrix $U \in M_n$ and scalars $c_1, \ldots, c_m \in \mathbb{R}_+$ such that $c_1 \cdots c_m = 1$ and*

$$\phi_i(A) = c_i^{1/\alpha_i} U^* A^{\beta_i/\alpha_i} U, \quad \forall A \in P_n,\ i = 1, \ldots, m. \tag{4.26}$$

(2) *When m is even, there exists an invertible $M \in M_n$ and scalars $c_1, \ldots, c_m \in \mathbb{R}_+$ such that $c_1 \cdots c_m = 1$ and*

$$\phi_i(A) = \begin{cases} c_i^{1/\alpha_i} \left(M^* A^{\beta_i} M\right)^{1/\alpha_i}, & i \text{ is odd}, \\ c_i^{1/\alpha_i} \left(M^{-1} A^{\beta_i} M^{-*}\right)^{1/\alpha_i}, & i \text{ is even}, \end{cases} \quad \forall A \in P_n. \tag{4.27}$$

Proof. It suffices to prove that (4.25) implies (4.26) (when m is odd) or (4.27) (when m is even). Define $\psi_i : P_n \to \overline{P_n}$ such that $\psi_i(A) := \phi_i(A^{1/\beta_i})^{\alpha_i}$ for $i = 1, \ldots, m$. Then

$$\operatorname{tr}(\psi_1(A_1) \cdots \psi_m(A_m)) = \operatorname{tr}(A_1 \cdots A_m), \quad \forall A_1, \ldots, A_m \in P_n.$$

Apply Theorem 4.3 for $m = 2$ and Theorem 4.4 for $m \geq 3$ to complete the proof. □

All theorems in this subsection have counterparts in the maps $P_n(\mathbb{R}) \to \overline{P_n(\mathbb{R})}$. The real versions of Theorem 4.3 and Theorem 4.4 are as follow.

Theorem 4.6. *Let $m \geq 2$. Maps $\phi_i : P_n(\mathbb{R}) \to \overline{P_n(\mathbb{R})}$ $(i = 1, \ldots, m)$ satisfy that*

$$\operatorname{tr}(\phi_1(A_1) \cdots \phi_m(A_m)) = \operatorname{tr}(A_1 \cdots A_m) \tag{4.28}$$

for all $A_1, \ldots, A_m \in P_n(\mathbb{R})$, if and only if they are described as follow:

(1) *When m is odd, there exist an orthogonal matrix $O \in M_n(\mathbb{R})$ and scalars $c_1, \ldots, c_m \in \mathbb{R}_+$ such that $c_1 \cdots c_m = 1$ and*

$$\phi_i(A) = c_i O^t A O, \quad \forall A \in P_n(\mathbb{R}),\ i = 1, \ldots, m. \tag{4.29}$$

(2) *When m is even, there exist an invertible $M \in M_n(\mathbb{R})$ and scalars $c_1, \ldots, c_m \in \mathbb{R}_+$ such that $c_1 \cdots c_m = 1$ and*

$$\phi_i(A) = \begin{cases} c_i M^t A M, & i \text{ is odd}, \\ c_i M^{-1} A M^{-t}, & i \text{ is even}, \end{cases} \quad \forall A \in P_n(\mathbb{R}). \tag{4.30}$$

Proof. We extend each ϕ_i to a map $S_n(\mathbb{R}) \to S_n(\mathbb{R})$ and still denote it ϕ_i by abuse of terminologies. For each $A \in S_n(\mathbb{R})$, choose arbitrary $c \in \mathbb{R}$ such that $A + cI_n \in P_n(\mathbb{R})$. Define

$$\phi_i(A) = \phi_i(A + cI_n) - c\phi_i(I_n). \tag{4.31}$$

The equality (4.28) guarantees that each $\phi_i : S_n(\mathbb{R}) \to S_n(\mathbb{R})$ is well-defined, and they satisfy (4.28) for all $A_1, \ldots, A_m \in S_n(\mathbb{R})$.

When $m \geq 3$, the statement can be proved by Theorem 5.2 in the next section.

When $m = 2$, we have $\operatorname{tr}(\phi_1(A_1)\phi_2(A_2)) = \operatorname{tr}(A_1A_2)$ for all $A_1, A_2 \in P_n(\mathbb{R})$. Similar to the proof of Theorem 4.3 (2), we see that $\phi_1(I)$ is invertible. Replacing $(\phi_1(A_1), \phi_2(A_2))$ by $(\phi_1(I)^{-1/2}\phi_1(A_1)\phi_1(I)^{-1/2}, \phi_1(I)^{1/2}\phi_2(A_2)\phi_1(I)^{1/2})$ if necessary, we may assume that $\phi_1(I) = I$. Similar to the proof of Theorem 4.3 (1), we get $\phi_1 = \phi_2$ and $\operatorname{tr}(\phi_1(A)) = \operatorname{tr}(A)$ for $A \in P_n(\mathbb{R})$. Therefore,

$$\operatorname{tr}(\phi_1(A)^2) = \operatorname{tr}(\phi_1(A)\phi_2(A)) = \operatorname{tr}(A^2) = \operatorname{tr}(\phi_1(A^2)), \quad \forall A \in P_n(\mathbb{R}). \tag{4.32}$$

By Kadison's inequality [7, p. 495], $\phi_1(A)^2 \leq \phi_1(A^2)$. The equality (4.32) forces $\phi_1(A)^2 = \phi_1(A^2)$ for all $A \in P_n(\mathbb{R})$ and thus for all $A \in S_n(\mathbb{R})$. Now extend ϕ_1 to a linear map $\widetilde{\phi_1} : S_n \to S_n$ such that for all $A, B \in S_n(\mathbb{R})$:

$$\widetilde{\phi_1}(A + \mathbf{i}B) = \phi_1(A) + \mathbf{i}\phi_1(B). \tag{4.33}$$

Then

$$\begin{aligned}
\widetilde{\phi_1}((A + \mathbf{i}B)^2) &= \phi_1(A^2 - B^2) + \mathbf{i}\phi_1((A + B)^2 - A^2 - B^2) && (4.34)\\
&= \phi_1(A)^2 - \phi_1(B)^2 + \mathbf{i}\left(\phi_1(A + B)^2 - \phi_1(A)^2 - \phi_1(B)^2\right) && (4.35)\\
&= \phi_1(A)^2 - \phi_1(B)^2 + \mathbf{i}\left(\phi_1(A)\phi_1(B) + \phi_1(B)\phi_1(A)\right) && (4.36)\\
&= \widetilde{\phi_1}(A + \mathbf{i}B)^2. && (4.37)
\end{aligned}$$

By [3, Theorem 2], there is an orthogonal matrix $O = [o_{ij}] \in M_n$ such that $\phi_1(A) = \phi_2(A) = O^tAO$ for $A \in S_n(\mathbb{R})$. Note that every $B \in M_n(\mathbb{R})$ is a product of two real symmetric matrices, say $B = A_1A_2$ for $A_1, A_2 \in S_n(\mathbb{R})$, so that

$$O^tBO = O^tA_1OO^tA_2O = \phi_1(A_1)\phi_2(A_2) \in M_n(\mathbb{R}).$$

The (p, q) entry of $O^tE_{ij}O$ is $o_{ip}o_{jq} \in \mathbb{R}$ for all $i, j, p, q \in \{1, \ldots, n\}$. Hence the entries of O are either all real or all pure imaginary. Since O is orthogonal, it must be real orthogonal. The statement for $m = 2$ has been proved. □

5. Maps on S_n

By Corollary 2.2, every linear bijection $\phi : S_n \to S_n$ corresponds to another linear bijection $\psi : S_n \to S_n$ such that $\operatorname{tr}(\phi(A)\psi(B)) = \operatorname{tr}(AB)$ for all $A, B \in S_n$. Some examples can be analogously developed from Example 3.3.

We now study the problem of three or more maps $S_n \to S_n$ preserving trace of products.

Theorem 5.1. *Suppose $m \geq 3$. Maps $\phi_1, \dots, \phi_m : S_n \to S_n$ satisfy that*

$$\operatorname{tr}(\phi_1(A_1) \cdots \phi_m(A_m)) = \operatorname{tr}(A_1 \cdots A_m), \quad \forall A_1, \dots, A_m \in S_n, \tag{5.1}$$

if and only if they are described as follow:

(1) *When m is odd, there exist an orthogonal matrix $O \in M_n$ and nonzero complex scalars $c_1, \dots, c_m$ such that $c_1 \cdots c_m = 1$ and*

$$\phi_i(A) = c_i O^t A O, \quad \forall A \in S_n, \ i = 1, \dots, m. \tag{5.2}$$

(2) *When m is even, there exist an invertible $M \in M_n$ and nonzero complex scalars $c_1, \dots, c_m$ such that $c_1 \cdots c_m = 1$ and*

$$\phi_i(A) = \begin{cases} c_i M^t A M, & i \text{ is odd}, \\ c_i M^{-1} A M^{-t}, & i \text{ is even}, \end{cases} \quad \forall A \in S_n. \tag{5.3}$$

Proof. It suffices to prove that (5.1) implies (5.2) (when m is odd) or (5.3) (when m is even). Let $C_1 := \phi_3(I) \cdots \phi_m(I)$, and $\psi_1 : S_n \to S_n C_1$ such that $\psi_1(B) := \phi_2(B) C_1$ for $B \in S_n$. Then $\operatorname{tr}(\phi_1(A)\psi_1(B)) = \operatorname{tr}(AB)$ for all $A, B \in S_n$. Theorem 2.1 implies that ϕ_1 and ψ_1 are linear bijections. Similarly, each ϕ_i's for $i = 1, \dots, m$ is a linear bijection. If C_1 is singular, then $v^t C_1 = 0$ for some $v \in \mathbb{C}^n \setminus \{0\}$, so that $vv^t \in S_n \setminus \{0\}$ but

$$\psi_1(\phi_2^{-1}(vv^t)) = vv^t C_1 = 0.$$

This contradicts the bijectivity of ψ_1. Hence C_1 is invertible. So each of $\phi_3(I), \dots, \phi_m(I)$, as well as $\phi_1(I)$ and $\phi_2(I)$, is invertible.

First we prove the theorem for the case $m = 3$.

For all $A, B, C \in S_n$ we have

$$\begin{aligned} & \operatorname{tr}\left[\phi_1(A)\left(\phi_2(B)\phi_3(C) - \phi_3(B)\phi_2(C)\right)\right] \\ = \ & \operatorname{tr}(\phi_1(A)\phi_2(B)\phi_3(C)) - \operatorname{tr}(\phi_2(C)\phi_3(B)\phi_1(A))^t \\ = \ & \operatorname{tr}(ABC) - \operatorname{tr}(ACB) = \operatorname{tr}(ABC) - \operatorname{tr}(ABC)^t = 0. \end{aligned}$$

Since $\phi_1 : S_n \to S_n$ is bijective, we get

$$\phi_2(B)\phi_3(C) - \phi_3(B)\phi_2(C) \in K_n. \tag{5.4}$$

Set $B = \phi_2^{-1}(I)$ and denote $J := \phi_3 \circ \phi_2^{-1}(I) \in S_n$ in (5.4). Then

$$\phi_3(C) - J\phi_2(C) \in K_n, \qquad \forall C \in S_n. \tag{5.5}$$

Therefore,

$$\phi_3(C) = \frac{1}{2}(J\phi_2(C) + \phi_2(C)J), \qquad \forall C \in S_n. \tag{5.6}$$

Apply the expression of ϕ_3 in (5.6) to (5.4):

$$\begin{aligned} & \frac{1}{2}\phi_2(B)(J\phi_2(C)+\phi_2(C)J)-\frac{1}{2}(J\phi_2(B)+\phi_2(B)J)\phi_2(C) \\ = \ & \frac{1}{2}\phi_2(B)\phi_2(C)J-\frac{1}{2}J\phi_2(B)\phi_2(C)\in K_n. \end{aligned} \tag{5.7}$$

By surjectivity of ϕ_2, the product $\frac{\phi_2(B)}{2}\phi_2(C)$ of two symmetric matrices can be any matrix in M_n (cf. [5, Theorem 3.2.3.2]). So $JA-AJ\in K_n$ for all $A\in M_n$, which is equivalent to

$$J(A-A^t)\in K_n, \qquad \forall A\in M_n. \tag{5.8}$$

The matrix $A-A^t$ in (5.8) goes through all matrices in K_n. So (5.8) holds if and only if $J=cI$ for some $c\in\mathbb{C}\setminus\{0\}$. Then (5.6) implies that

$$\phi_3(C)=c\phi_2(C), \qquad \forall C\in S_n. \tag{5.9}$$

Similarly, ϕ_1 is a nonzero constant multiple of ϕ_2. So there exists a linear bijection $\phi: S_n\to S_n$ such that $\phi_i=c_i'\phi$ for $i=1,2,3$, $c_1'c_2'c_3'=1$, and

$$\operatorname{tr}(\phi(A)\phi(B)\phi(C))=\operatorname{tr}(ABC), \qquad \forall A,B,C\in S_n. \tag{5.10}$$

If $A\in S_n$ satisfies $A^2=A$, then $\operatorname{tr}(\phi(A)^2\phi(B))=\operatorname{tr}(\phi(A)\phi(I)\phi(B))$ for all $B\in S_n$. So

$$\phi(A)^2-\phi(A)\phi(I) \in K_n, \quad \text{which implies that} \tag{5.11}$$

$$2\phi(A)^2 = \phi(A)\phi(I)+\phi(I)\phi(A), \tag{5.12}$$

$$\phi(A)^2\phi(I)+\phi(A)\phi(I)\phi(A) = 2\phi(A)^3=\phi(A)\phi(I)\phi(A)+\phi(I)\phi(A)^2, \tag{5.13}$$

$$\phi(A)^2\phi(I) = \phi(I)\phi(A)^2. \tag{5.14}$$

Fix a basis $\{A_i : i=1,\dots,\binom{n+1}{2},\ A_i^2=A_i\}$ of S_n. We claim that $\{\phi(A_i)^2 : i=1,\dots,\binom{n+1}{2}\}$ is also a basis of S_n. Suppose $a_1,\dots,a_{\binom{n+1}{2}}\in\mathbb{C}$ and

$$a_1\phi(A_1)^2+\cdots+a_{\binom{n+1}{2}}\phi(A_{\binom{n+1}{2}})^2=0. \tag{5.15}$$

Then for all $C\in S_n$:

$$\begin{aligned} 0 &= \operatorname{tr}\left(a_1\phi(A_1)^2+\cdots+a_{\binom{n+1}{2}}\phi(A_{\binom{n+1}{2}})^2\right)\phi(C) \\ &= \operatorname{tr}\left(a_1A_1^2+\cdots+a_{\binom{n+1}{2}}A_{\binom{n+1}{2}}^2\right)C \\ &= \operatorname{tr}\left(a_1A_1+\cdots+a_{\binom{n+1}{2}}A_{\binom{n+1}{2}}\right)C. \end{aligned} \tag{5.16}$$

Therefore, $a_1A_1+\cdots+a_{\binom{n+1}{2}}A_{\binom{n+1}{2}}=0$, so that $a_1=\cdots=a_{\binom{n+1}{2}}=0$.

By (5.14),

$$\phi(A_i)^2\phi(I)=\phi(I)\phi(A_i)^2, \qquad i=1,\dots,\binom{n+1}{2}. \tag{5.17}$$

Hence $\phi(I)$ commutes with all matrices in S_n. Therefore, $\phi(I)=c_0I$ for some $c_0\in\mathbb{C}$. By

$$c_0^3n=\operatorname{tr}(\phi(I)^3)=\operatorname{tr}(I^3)=n \tag{5.18}$$

we get $c_0^3 = 1$.

Denote the linear bijection $\widetilde{\phi} = \frac{1}{c_0}\phi$. Then $\widetilde{\phi}(I) = I$ and

$$\operatorname{tr}(\widetilde{\phi}(A)\widetilde{\phi}(B)\widetilde{\phi}(C)) = \operatorname{tr}(ABC), \qquad \forall A, B, C \in S_n. \tag{5.19}$$

Given $B \in S_n$, for all $A \in S_n$,

$$\operatorname{tr}(\widetilde{\phi}(A)\widetilde{\phi}(B^2)) = \operatorname{tr}(\widetilde{\phi}(A)\widetilde{\phi}(B^2)\widetilde{\phi}(I)) = \operatorname{tr}(AB^2) = \operatorname{tr}(\widetilde{\phi}(A)\widetilde{\phi}(B)^2). \tag{5.20}$$

So the symmetric matrices $\widetilde{\phi}(B^2) = \widetilde{\phi}(B)^2$. By [3, Theorem 2], there exists an orthogonal matrix $O \in M_n$ such that $\widetilde{\phi}(A) = O^t AO$ for $A \in S_n$. Let $c_i = c_0 c_i'$ for $i = 1, 2, 3$. Then we get (5.2) for $m = 3$.

From now on, we assume that $m \geq 4$.

The matrix $\phi_3(A_3)\cdots\phi_m(A_m)$ for $A_3, \ldots, A_m \in S_n$ can be any matrix in M_n (cf. [5, Theorem 3.2.3.2]). For all $A, A_3, \ldots, A_m \in S_n$ we have

$$\operatorname{tr}\left[(\phi_1(A)\phi_2(I) - \phi_1(I)\phi_2(A))\,\phi_3(A_3)\cdots\phi_m(A_m)\right] = 0. \tag{5.21}$$

So $\phi_1(A)\phi_2(I) = \phi_1(I)\phi_2(A)$ for all $A \in S_n$. Hence

$$\phi_2(A) = \phi_1(I)^{-1}\phi_1(A)\phi_2(I). \tag{5.22}$$

Since $\phi_1(A)$ and $\phi_2(A)$ are symmetric and ϕ_1 is surjective, there exists $c_2' \in \mathbb{C}\setminus\{0\}$ such that

$$\phi_2(I) = c_2'\phi_1(I)^{-t} = c_2'\phi_1(I)^{-1}, \quad \text{so that by (5.22)}, \tag{5.23}$$

$$\phi_2(A) = c_2'\phi_1(I)^{-1}\phi_1(A)\phi_1(I)^{-1}, \qquad \forall A \in S_n. \tag{5.24}$$

Similarly, there is $c_3' \in \mathbb{C}\setminus\{0\}$ such that

$$\phi_3(A) = c_3'\phi_2(I)^{-1}\phi_2(A)\phi_2(I)^{-1} = \frac{c_3'}{c_2'}\phi_1(A), \qquad \forall A \in S_n. \tag{5.25}$$

Therefore, the maps $\phi_1, \phi_3, \phi_5, \ldots$ (resp. $\phi_2, \phi_4, \phi_6, \ldots$) differ only by nonzero scalar multiples.

(1) If m is odd, then $\phi_1 = a_m\phi_m$ for some $a_m \in \mathbb{C}\setminus\{0\}$ and $\phi_m = a_2\phi_2$ for some $a_2 \in \mathbb{C}\setminus\{0\}$. Therefore, $\phi_1, \phi_2, \phi_3, \ldots, \phi_m$ differ only by nonzero scalar multiples. (5.22) implies that $\phi_1(A) = \phi_1(I)^{-1}\phi_1(A)\phi_1(I)$ for all $A \in S_n$, so that $\phi_1(I) = c_1 I$ for some $c_1 \in \mathbb{C}\setminus\{0\}$. Let $\phi(A) := \frac{1}{c_1}\phi_1(A)$. Then $\phi(I) = I$ and $\phi_1 = c_1\phi$. There exist $c_2, \ldots, c_m \in \mathbb{C}\setminus\{0\}$ such that $\phi_i = c_i\phi$ for $i = 2, \ldots, m$. We have

$$c_1c_2\cdots c_m \operatorname{tr}(I^m) = c_1c_2\cdots c_m \operatorname{tr}(\phi(I)^m) = \operatorname{tr}(\phi_1(I)\cdots\phi_m(I)) = \operatorname{tr}(I^m).$$

Therefore, $c_1\cdots c_m = 1$, and

$$\operatorname{tr}(\phi(A_1)\cdots\phi(A_m)) = \operatorname{tr}(A_1\cdots A_m), \qquad \forall A_1, \ldots, A_m \in S_n. \tag{5.26}$$

(2) If m is even, write $\phi_1(I) = M^tM$ for an invertible $M \in M_n$, then (5.24) implies that

$$M\phi_2(A)M^t = c_2'M^{-t}\phi_1(A)M^{-1}, \qquad \forall A \in S_n. \tag{5.27}$$

There exist a map $\phi(A) = M^{-t}\phi_1(A)M^{-1}$ and scalars $c_1 = 1, c_2, \ldots, c_m \in \mathbb{C}\setminus\{0\}$ such that

(a) If $i \in \{1, \dots, m\}$ is odd, then $M^{-t}\phi_i(A)M^{-1} = c_i\phi(A)$ for $A \in S_n$.
(b) If $i \in \{1, \dots, m\}$ is even, then $M\phi_i(A)M^t = c_i\phi(A)$ for $A \in S_n$.
(c) $c_1 \cdots c_m = 1$, $\phi(I) = I$, and (5.26) holds:

$$\operatorname{tr}(\phi(A_1)\cdots\phi(A_m)) = \operatorname{tr}(A_1\cdots A_m), \qquad \forall A_1, \dots, A_m \in S_n.$$

In both m even and m odd cases, we successfully find a linear bijection $\phi : S_n \to S_n$ that satisfies $\phi(I) = I$ and (5.26). For all $A, A_3, \dots, A_m \in S_n$ and $k \in \mathbb{Z}_+$ we have

$$\operatorname{tr}\left[\left(\phi(A)\phi(A^{k-1}) - \phi(I)\phi(A^k)\right)\phi(A_3)\cdots\phi(A_m)\right] = 0. \tag{5.28}$$

Hence $\phi(A)\phi(A^{k-1}) = \phi(I)\phi(A^k) = \phi(A^k)$ and $\phi(A)^k = \phi(A^k)$ for $k \in \mathbb{Z}_+$. Therefore, the following identities hold for every $A, B \in S_n$:

$$\operatorname{tr}(\phi(A)\phi(B)^k) = \operatorname{tr}(\phi(A)\phi(B^k)\phi(I)\cdots\phi(I)) = \operatorname{tr}(AB^k), \quad \forall k \in \mathbb{Z}_+. \tag{5.29}$$

By the proof of (3) $\Rightarrow$ (1) in [6, Lemma 4.2] or by [3, Theorem 2], there exists an orthogonal matrix $O \in M_n$ such that $\phi(A) = O^tAO$ for $A \in S_n$. When m is odd, we get (5.2). When m is even, we get (5.3). □

The real version of Theorem 5.1 holds. We state and prove it below.

Theorem 5.2. *Suppose $m \geq 3$. Maps $\phi_1, \dots, \phi_m : S_n(\mathbb{R}) \to S_n(\mathbb{R})$ satisfy that*

$$\operatorname{tr}(\phi_1(A_1)\cdots\phi_m(A_m)) = \operatorname{tr}(A_1\cdots A_m), \quad \forall A_1, \dots, A_m \in S_n(\mathbb{R}), \tag{5.30}$$

if and only if the following are true:

(1) *When m is odd, there exist a real orthogonal matrix $O \in M_n(\mathbb{R})$ and nonzero real scalars $c_1, \dots, c_m$ such that $c_1 \cdots c_m = 1$ and*

$$\phi_i(A) = c_iO^tAO, \quad \forall A \in S_n(\mathbb{R}),\ i = 1, \dots, m. \tag{5.31}$$

(2) *When m is even, there exist an invertible $M \in M_n(\mathbb{R})$ and nonzero real scalars $c_1, \dots, c_m$ such that $c_1 \cdots c_m = 1$ and*

$$\phi_i(A) = \begin{cases} c_iM^tAM, & i \text{ is odd}, \\ c_iM^{-1}AM^{-t}, & i \text{ is even}, \end{cases} \quad \forall A \in S_n(\mathbb{R}). \tag{5.32}$$

Proof. For every $A \in S_n$, denote $\operatorname{Re} A, \operatorname{Im} A \in S_n(\mathbb{R})$ such that $A = \operatorname{Re} A + \mathbf{i}\operatorname{Im} A$. For each $i = 1, \dots, m$, define $\widetilde{\phi_i} : S_n \to S_n$ in the way that

$$\widetilde{\phi_i}(A) = \phi_i(\operatorname{Re} A) + \mathbf{i}\phi_i(\operatorname{Im} A), \quad \forall A \in S_n. \tag{5.33}$$

Then

$$\operatorname{tr}(\widetilde{\phi_1}(A_1)\cdots\widetilde{\phi_m}(A_m)) = \operatorname{tr}(A_1\cdots A_m), \quad \forall A_1, \dots, A_m \in S_n. \tag{5.34}$$

By Theorem 5.1:

(1) When m is odd, there exist an orthogonal matrix $O \in M_n$ and $c_1, \dots, c_m \in \mathbb{C}$ such that $c_1 \cdots c_m = 1$ and

$$\widetilde{\phi}_i(A) = c_i O^t A O, \quad \forall A \in S_n,\ i = 1, \dots, m. \tag{5.35}$$

Then $\widetilde{\phi}_i(I) = c_i I \in S_n(\mathbb{R})$ implies that each $c_i \in \mathbb{R}$. Write $O = [o_{ij}]_{n\times n}$. The (p,q) entry of $\widetilde{\phi_1}(E_{ij})$ is $c_1 o_{ip} o_{jq}$, which must be real for all $i, j, p, q \in \{1, \dots, n\}$. Hence the entries of O are either all real or all pure imaginary. Since O is orthogonal, it has to be real orthogonal.

(2) When m is even, there exist an invertible $M \in M_n$ and $c_1, \dots, c_m \in \mathbb{C}$ such that $c_1 \cdots c_m = 1$ and

$$\widetilde{\phi}_i(A) = \begin{cases} c_i M^t A M, & i \text{ is odd}, \\ c_i M^{-1} A M^{-t}, & i \text{ is even}, \end{cases} \quad \forall A \in S_n. \tag{5.36}$$

We may assume that $c_1 \in \mathbb{R}$ by adjusting M with a scalar factor if necessary. Then $\widetilde{\phi_1}(I)\widetilde{\phi_2}(I) = c_1 c_2 I$ is real, which implies $c_2 \in \mathbb{R}$. Similarly, $c_i \in \mathbb{R}$ for all $i = 1, \dots, n$. Using the fact that $\widetilde{\phi_1}(E_{jj})$ is real for all $j = 1, \dots, n$, we see that the entries of M are either all real or all pure imaginary. We can assure that $M \in M_n(\mathbb{R})$ by adjusting the signs of all c_i's if necessary.

Overall, the theorem is proved. □

6. Maps on D_n

The results about maps $D_n \to D_n$ in this section can be applied to maps between the spaces of upper (resp. lower) triangular matrices with respective properties. We view

$$\mathrm{diag} : \mathbb{C}^n \to D_n \tag{6.1}$$

as the linear bijection that sends $(c_1, \cdots, c_n)^t$ to the diagonal matrix $\mathrm{diag}(c_1, \dots, c_n)$, and

$$\mathrm{diag}^{-1} : D_n \to \mathbb{C}^n \tag{6.2}$$

the inverse map of diag.

Let $\phi_1, \phi_2 : D_n \to D_n$ satisfy that

$$\mathrm{tr}(\phi_1(A_1)\phi_2(A_2)) = \mathrm{tr}(A_1 A_2), \quad \forall A_1, A_2 \in D_n. \tag{6.3}$$

Then Theorem 2.1 and Corollary 2.2 imply that ϕ_1 (resp. ϕ_2) can be any linear automorphism, and ϕ_1 and ϕ_2 uniquely determine each other. The following theorem gives the explicit description of these (ϕ_1, ϕ_2) pairs.

Theorem 6.1. *Two maps $\phi_1, \phi_2 : D_n \to D_n$ satisfy (6.3) if and only if there is an invertible matrix $N \in M_n$ such that*

$$\phi_1(A) = \mathrm{diag}(N\mathrm{diag}^{-1}(A)), \quad \phi_2(A) = \mathrm{diag}(N^{-t}\mathrm{diag}^{-1}(A)), \quad \forall A \in D_n. \tag{6.4}$$

Proof. For two vector $u, v \in \mathbb{C}^n$,

$$\mathrm{tr}(\mathrm{diag}(u)\mathrm{diag}(v)) = u^t v. \tag{6.5}$$

Define the linear automorphisms $\widetilde{\phi}_i : \mathbb{C}^n \to \mathbb{C}^n$ by

$$\widetilde{\phi}_i := \mathrm{diag}^{-1} \circ \phi_i \circ \mathrm{diag}, \qquad i = 1, 2. \tag{6.6}$$

There exist invertible matrices $N, M \in M_n$ such that $\widetilde{\phi}_1(v) = Nv$ and $\widetilde{\phi}_2(v) = Mv$ for $v \in \mathbb{C}^n$. Express elements of D_n as $\mathrm{diag}(v)$ for $v \in \mathbb{C}^n$. Then (6.3) becomes

$$\widetilde{\phi}_1(u)^t \widetilde{\phi}_2(v) = u^t v, \quad \forall u, v \in \mathbb{C}^n. \tag{6.7}$$

It implies that $M = N^{-t}$. Therefore, for $A \in D_n$,

$$\phi_1(A) = \mathrm{diag} \circ \widetilde{\phi}_1 \circ \mathrm{diag}^{-1}(A) = \mathrm{diag}(N\mathrm{diag}^{-1}(A)) \tag{6.8}$$

and similarly $\phi_2(A) = \mathrm{diag}(N^{-t}\mathrm{diag}^{-1}(A))$. □

Theorem 6.2. *Suppose $m \geq 3$ and $\phi_1, \ldots, \phi_m : D_n \to D_n$ satisfy that*

$$\mathrm{tr}(\phi_1(A_1) \cdots \phi_m(A_m)) = \mathrm{tr}(A_1 \cdots A_m), \quad \forall A_1, \ldots, A_m \in D_n. \tag{6.9}$$

Then there exist a permutation matrix $P \in M_n$ and invertible diagonal matrices $C_1, \ldots, C_m \in D_n$ with $C_1 \cdots C_m = I$ such that

$$\phi_i(A) = C_i P^t A P, \qquad \forall A \in D_n, \ i = 1, \ldots, m. \tag{6.10}$$

Proof. Let $C = \phi_3(I) \cdots \phi_m(I)$ and $\psi_1 : D_n \to D_n$ such that $\psi_1(A) = \phi_2(A)C$. Then $\mathrm{tr}(\phi_1(A)\psi_1(B)) = \mathrm{tr}(AB)$ for all $A, B \in D_n$, which implies that both ϕ_1 and ψ_1 are linear bijections. So each of $\phi_3(I), \ldots, \phi_m(I)$ is invertible. Similarly, each of $\phi_1(I)$ and $\phi_2(I)$ is invertible, and each of $\phi_2, \ldots, \phi_m$ is a linear bijection.

If $AB = A'B'$ for $A, B, A', B' \in D_n$, then for all $A_3, \ldots, A_m \in D_n$,

$$\begin{aligned} & \mathrm{tr}(\phi_1(A)\phi_2(B)\phi_3(A_3) \cdots \phi_m(A_m)) \\ = \ & \mathrm{tr}(ABA_3 \cdots A_m) = \mathrm{tr}(A'B'A_3 \cdots A_m) \qquad (6.11) \\ = \ & \mathrm{tr}(\phi_1(A')\phi_2(B')\phi_3(A_3) \cdots \phi_m(A_m)). \qquad (6.12) \end{aligned}$$

The bijectivities of $\phi_3, \ldots, \phi_m$ imply that $\phi_1(A)\phi_2(B) = \phi_1(A')\phi_2(B')$. In particular,

$$\phi_2(B) = \phi_1(I)^{-1}\phi_1(B)\phi_2(I), \qquad \forall B \in D_n. \tag{6.13}$$

For all $A, B \in D_n$ we also have

$$\phi_1(AB)\phi_2(I) = \phi_1(A)\phi_2(B) = \phi_1(A)\phi_1(I)^{-1}\phi_1(B)\phi_2(I). \tag{6.14}$$

Let $\widetilde{\phi}_1(A) := \phi_1(A)\phi_1(I)^{-1}$ then

$$\widetilde{\phi}_1(A)\widetilde{\phi}_1(B) = \widetilde{\phi}_1(AB), \quad \forall A, B \in D_n. \tag{6.15}$$

So $\widetilde{\phi}_1$ is an algebra automorphism of D_n, and the restriction of $\widetilde{\phi}_1$ on the group of invertible diagonal matrices in D_n gives a group automorphism. Hence there is a permutation matrix $P \in M_n$ such that

$$\widetilde{\phi}_1(A) = P^t AP, \quad \forall A \in D_n. \tag{6.16}$$

Let $C_i := \phi_i(I)$ for $i = 1, \ldots, m$. Then $\phi_1(A) = C_1 P^t AP$, and by (6.13),

$$\phi_2(A) = (P^t AP)C_2 = C_2 P^t AP. \tag{6.17}$$

Similarly,

$$\phi_i(A) = C_i P^t AP, \qquad \forall A \in D_n, \quad i = 1, \ldots, m, \tag{6.18}$$

in which $C_1 \cdots C_m = I$ by (6.9). □

Both Theorem 6.1 and Theorem 6.2 have counterparts in maps on the set of real diagonal matrices and the proofs are analogeous.

References

[1] Z.E.A. Abdelali, B. Aharmim, *Multiplicatively spectrum preserving maps on rectangular matrices,* Linear Multilinear Algebra (2020). doi = 10.1080/03081087.2019.1708237

[2] A. Bourhim, J.E. LEE, *Multiplicatively local spectrum-preserving maps,* Linear Algebra Appl. **549** (2018) 291–308.

[3] G. H. Chan, M. H. Lim, *Linear preservers on powers of matrices,* Linear Algebra Appl. **162–164** (1992) 615–626.

[4] J. T. Chan, C. K. Li, N. S. Sze, *Mappings preserving spectra of product of matrices,* Proc. Amer. Math. Soc., **135** (2007) 977–986.

[5] R. A. Horn, C. R. Johnson, Matrix Analysis, 2nd ed, Cambridge University Press.

[6] H. Huang, C.N. Liu, M.C. Tsai, P. Szokol, J. Zhang, *Trace and determinant preserving maps of matrices,* Linear Alg. Appl., **507** (2016) 373–388.

[7] R.V. Kadison, *A generalized Schwarz inequality and algebraic invariants for operator algebras*, Ann. of Math., **56** (1952) 494–503.

[8] C.W. Leung, C.K. Ng, N.C. Wong, *Transition probabilities of normal states determine the Jordan structure of a quantum system,* J. Math. Phys., **57** (2016) 015212.

[9] C. K. Li, S. Pierce, *Linear preserver problems,* Amer. Math. Monthly **108** (2001) 591–605.

[10] C. K. Li, L. Plevnik, P. Šemrl, *Preservers of matrix pairs with a fixed inner product value,* Oper. Matrices **6** (2012) 433–464.

[11] C.K. Li, N.K. Tsing, *Linear preserver problems: a brief introduction and some special techniques,* Linear Alg. Appl., **162–164** (1992) 217–235.

[12] T. Miura, D. Honma, *A generalization of peripherally-multiplicative surjections between standard operator algebras,* Cent. Eur. J. Math., **7(3)** (2009) 479–486.

[13] L. Molnár, Some characterizations of the automorphisms of $B(H)$ and $C(X)$, Proc. Amer. Math. Soc., **130(1)** (2002) 111–120.

[14] L. Molnár, *Local automorphisms of operator algebras on Banach spaces*, Proc. Amer. Math. Soc., **131(6)** (2003) 1867–1874.

[15] L. Molnár, Selected preserver problems on algebraic structures of linear operators and on function spaces, Lecture Notes in Mathematics 1895, Springer-Verlag, Berlin, 2007.

[16] S. Pierce et. al., *A survey of linear preserver problems*, Linear Multilinear Algebra, **33** (1992) 1–129.

[17] U. Uhlhorn, *Representation of symmetry transformations in quantum mechanics,* Ark. Fysik **23** (1963) 307–340.

Department of Mathematics and Statistics, Auburn University, AL 36849–5310, USA

Email address: huanghu@auburn.edu

General Education Center, Taipei University of Technology, Taipei 10608, Taiwan

Email address: mctsai2@mail.ntut.edu.tw