

Extending the Mann-Kendall test to allow for measurement uncertainty

Stavros Nikolakopoulos^{*1,2}, Eric Cator³, and Mart P. Janssen⁴

¹Department of Statistics, Athens University of Economics and Business

²Department of Biostatistics, University Medical Center Utrecht

³Department of Mathematics, Radboud University Nijmegen

⁴Transfusion Technology Assessment Group, Department of Donor Medicine Research, Sanquin Research, Amsterdam

Abstract

The Mann-Kendall test for trend has gained a lot of attention in a range of disciplines, especially in the environmental sciences. One of the drawbacks of the Mann-Kendall test when applied to real data is that no distinction can be made between meaningful and non-meaningful differences in subsequent observations. We introduce the concept of partial ties, which allows inferences while accounting for (non)meaningful difference. We introduce the modified statistic that accounts for such a concept and derive its variance estimator. We also present analytical results for the behavior of the test in a class of contiguous alternatives. Simulation results which illustrate the added value of the test are presented. We apply our extended version of the test to some real data concerning blood donation in Europe.

Keywords: Mann-Kendall test; Measurement error; Regional Kendall test; Relevant difference; Ties

1 Introduction

1.1 Trend analysis and measurement uncertainty

The report "Trends and Observations on the Collection, Testing and Use of Blood and Blood Components in Europe, 2001-2011" (Janssen et al., 2015) presents and analyzes data provided annually to the Council of Europe by its Member States. Since 2001, data concerning blood donation in the European Union (EU) has been collected by means of a, newly developed, standardized questionnaire which was distributed to EU's Member States. Naturally, during the first years of the questionnaire administration, improvements and amendments took place, following recommendations from expert committees (Janssen et al., 2015). Given this was the first time this kind of data were annually collected and analyzed, the assessment of a trend over time was an important component of the data analysis (Van der Poel et al., 2011; Janssen et al., 2015). However, given the newly developed tool and overall discrepancies among the Member States, there were several possible sources of *measurement uncertainty*. These include over- or under- reporting, different ways of registering per country (e.g. electronic or paper archives) or

*corresponding author, email: sknikolak@aueb.gr

lack of uniform definitions among the Member States. Furthermore, data were first analyzed in 2005 (Van der Poel et al., 2011), with only (max) five datapoints per country on top of the aforementioned difficulties. Thus non-parametric methods were the methods of choice for trend analysis.

1.2 The Mann-Kendall test

Kendall's tau coefficient (t) (Kendall, 1938) was originally derived to test the degree of agreement, or correlation, between two sets of observations. It is a non-parametric statistic, estimating population's τ , also referred to as rank correlation coefficient. The t statistic is a standardized version of the S statistic, which in turn is calculated directly from the ranks of the observations within each of the sets of observations.

Mann (1945), suggested the use of t for testing against the existence of a trend over time. The Mann-Kendall test has gained a lot of attention in a range of disciplines, especially in the environmental sciences where its use has been extended in combining several sets of observations of comparable times of the year (Seasonal Kendall test for trend, Hirsch et al. (1982)) or data obtained from different regions (Regional Kendall test for trend, Helsel and Frans (2006)). The Mann-Kendall test is still popular for trend detection (Dawood et al., 2017; Nourani et al., 2018)

The Man-Kendall test is a straightforward application of Kendall's τ where one of the sets of observations is timepoints, accompanied with measurements obtained at these time points. Ranks of the time variable are always in the natural order $(1, 2, \dots, n)$ and the S statistic is given by

$$S = \sum_{i < j} \text{sign}(X_j - X_i) \quad (1.1)$$

where X_i refer to the variable of interest. Values of X_i could be either the original values measured or their ranks within the sample obtained.

Kendall (Kendall and Dickinson Gibbons, 1990; Kendall, 1938) discusses normality of S , and also proves that:

$$E(S) = 0$$

and

$$\text{Var}(S) = \frac{1}{18}n(n-1)(2n+5).$$

1.3 Presence of ties

When ties are present in the data, the formula of $\text{Var}(S)$ requires some modification. Given that there are ties of extent w , (e.g., if there are 3 tied groups, one duplet and 2 triplets, $w=\{2,3,3\}$) and w_i is the number of tied values per group, the variance is calculated by:

$$\text{Var}(S) = \frac{1}{18} \left\{ n(n-1)(2n+5) - \sum_i w_i(w_i-1)(2w_i+5) \right\}.$$

Behavior of t and S in rankings containing ties has been discussed by Kendall (Kendall and Dickinson Gibbons, 1990; Kendall, 1945) and others (Sillitto, 1947; Robillard, 1972; Panneton and Robillard, 1972).

Based on the above, inference can be made, in the sense of statistical significance and calculating p-values, on the existence or not of a positive or negative trend over time in a given dataset.

Table 1: Example dataset on consecutive measurements of systolic blood pressure of an individual

Time point	1	2	3	4	5	6	7	8	9	10
<i>DBP</i>	90.9	95.2	98.6	95.8	100.7	94.9	92.8	101.5	99	98.7
Rank(<i>DBP</i>)	1	4	7	5	9	3	2	10	8	6
ordered <i>DBP</i>	90.9	92.8	94.9	95.2	95.8	98.6	98.7	99	100.7	101.5
Rank(<i>DBP</i>)	1	2	3	4	5	6	7	8	9	10

1.4 Introducing partially tied observations

One of the characteristics (and strengths) of the Mann-Kendall test, due to its non-parametric nature, is that it can provide information concerning the existence of a trend over time without any further assumptions. One of the limitations of this approach is that any increase or decrease between two observations is interpreted as such, irrespective of the absolute difference between the values actually compared. We found in practice (considering the blood donation data described in Section 1.1) that when applying the Mann-Kendall test, this in some cases lead to a rejection of clearly visible trends. This made us explore the possibility of extending the traditional test to only value a positive or negative contribution to the test statistic in those cases where a difference in observed values is actually considered meaningful in the light of the context or measurement accuracy applicable.

From the calculation of S from Equation 1.1, it is obvious that pairs of values compared contribute to S either +1 or -1 irrespectively of the magnitude of the difference between each pair's elements. One could argue though, that two observations without the exact same numerical value but with a *relatively* small difference, are not reasonable to be accounted for as different and therefore should not contribute to the value of S . For demonstration, let us consider an example of observations on diastolic blood pressure of an individual (*DBP*) measured over 10 consecutive time points like the ones in Table 1.

As Manios et al. (2007) report, diastolic blood pressure is overestimated when recorded by non-invasive methods while systolic blood pressure is underestimated when measured intravenously. One could then argue that 98.6 is not really different from 98.7. Such small differences could be attributed to measurement error, or could be considered clinically irrelevant. The example could be extended to other kinds of variables with alternative sources of uncertainty. Uncertainty could also be assumed to have one direction only, for example known rates of under or over reporting of financial or demographic measurements. To our knowledge, this problem has not been dealt with in the literature, with the only exception being a simulation study by Soderberg and Hennet (2007).

In light of the above, it might be required to apply a *level of relevant difference (LRD)* on observations, that is, define d such that a pair (X_i, X_j) will be considered a tie if $|X_i - X_j| \leq d$. The value of d could be defined on the basis of known or assumed measurement error, given the observations nature and/or the way in which they were obtained. Such an approach could have awkward implications for the application of the traditional Mann-Kendall test. In the above example, by applying a *LRD* of $d = 0.6$ we get the seemingly paradoxical situation where in the ordered sample (lower part of Table 1 and denoting by $DBP_{(i)}$ the i -ith ranked observation) $DBP_{(3)} = DBP_{(4)}, DBP_{(4)} = DBP_{(5)}$ but $DBP_{(3)} \neq DBP_{(5)}$. One can imagine that applying a *LRD* rule can result in highly complex tie structures, which we will refer to as *partially tied observations*. The S statistic cannot account for such complications as presented in Equation

1.1.

The paper is organized as follows: In the next section, the Mann-Kendall test is extended to be able to account for partially tied observation with the key extension being the generalization of the variance formula of S . In section 3 a simulation study is presented to illustrate the performance of the extended test compared to the classical test. In section 4 the extended version is applied to a real data set where it is implemented within the Regional test for trend. The final section provides a discussion on the extended test.

2 Properties of S and t in the presence of partially tied observations

2.1 Properties of S

Suppose we have a sample X_1, \dots, X_n of measurements taken at subsequent times $1, 2, \dots, n$. Our test wishes to distinguish a trend in data from the null hypothesis H_0 that all X_i 's are iid. We introduce the *extended* Mann-Kendall test where we allow for the above introduced *LRD* which we denote by $d > 0$ and thus we define

$$S_{ex} = \sum_{i < j} \text{sign}(X_j - X_i) \mathbb{1}_{\{|X_i - X_j| > d\}} \quad (2.1)$$

where $\mathbb{1}_{\{\cdot\}}$ denotes the indicator function. The conditions of this criterion could vary as discussed above, for example the absolute could be replaced if we were interested in uncertainty towards one direction of the measurements or $>$ could be replaced by \geq in Equation 2.1, depending on the measurement units or accuracy of the measurement instrument employed.

The statistic S_{ex} will be strongly positive if there is a positive trend in the data, and it will be strongly negative if there is a negative trend in the data. We will base our test on a normalized version of S_{ex} (t_{ex}) and for this we need to estimate its variance under H_0 . It is easy to see that under the null hypothesis,

$$\mathbb{E}(S_{ex}|H_0) = 0$$

For the variance of S_{ex} , since $\mathbb{E}(S_{ex}|H_0) = 0$, we need to calculate

$$\text{Var}(S_{ex}|H_0) = \mathbb{E}(S_{ex}^2|H_0) = \sum_{i < k} \sum_{j < l} \mathbb{E}(\text{sign}(X_k - X_i) \mathbb{1}_{\{|X_i - X_k| > d\}} \text{sign}(X_l - X_j) \mathbb{1}_{\{|X_j - X_l| > d\}}).$$

Define the index set

$$\Omega_0 = \{(i, j, k, l) \in \mathbb{N}^4 : 1 \leq i < k \leq n \text{ and } 1 \leq j < l \leq n\}.$$

We split up this set into six disjoint parts:

$$\begin{aligned}
\Omega_1 &= \{(i, j, k, l) \in \Omega_0 : i \neq j, k \neq l, i \neq l, j \neq k\} \\
\Omega_2 &= \{(i, j, k, l) \in \Omega_0 : i = j, k \neq l\} \\
\Omega_3 &= \{(i, j, k, l) \in \Omega_0 : k = l, i \neq j\} \\
\Omega_4 &= \{(i, j, k, l) \in \Omega_0 : i = l\} \\
\Omega_5 &= \{(i, j, k, l) \in \Omega_0 : j = k\} \\
\Omega_6 &= \{(i, j, k, l) \in \Omega_0 : i = j, k = l\}
\end{aligned}$$

Now define $N_i = \#\Omega_i$. Then

$$\begin{aligned}
N_0 &= \frac{1}{4}n^4 - \frac{1}{2}n^3 + \frac{1}{4}n^2 \\
N_1 &= \frac{1}{4}n^4 - \frac{3}{2}n^3 + \frac{11}{4}n^2 - \frac{3}{2}n \\
N_2 = N_3 &= \frac{1}{3}n^3 - n^2 + \frac{2}{3}n \\
N_4 = N_5 &= \frac{1}{6}n^3 - \frac{1}{2}n^2 + \frac{1}{3}n = \frac{1}{2}N_2 \\
N_6 &= \frac{1}{2}n^2 - \frac{1}{2}n.
\end{aligned}$$

Furthermore we define

$$\alpha_+ = \mathbb{E}(\mathbb{1}_{\{X_1 > X_2 + d\}} \mathbb{1}_{\{X_1 > X_3 + d\}}) \quad \text{and} \quad \alpha_- = \mathbb{E}(\mathbb{1}_{\{X_1 < X_2 - d\}} \mathbb{1}_{\{X_1 < X_3 - d\}})$$

and

$$\beta = \mathbb{E}(\mathbb{1}_{\{X_1 > X_2 + d\}} \mathbb{1}_{\{X_1 < X_3 - d\}}) \quad \text{and} \quad \gamma = \mathbb{E}(\mathbb{1}_{\{X_1 > X_2 + d\}}) = \mathbb{E}(\mathbb{1}_{\{X_1 < X_2 - d\}}).$$

Note that $\mathbb{1}_{\{X_1 > X_2 + d\}}$ and $\mathbb{1}_{\{X_1 > X_3 + d\}}$ are positively correlated, $\mathbb{1}_{\{X_1 < X_2 - d\}}$ and $\mathbb{1}_{\{X_1 < X_3 - d\}}$ are also positively correlated, but $\mathbb{1}_{\{X_1 > X_2 + d\}}$ and $\mathbb{1}_{\{X_1 > X_3 - d\}}$ are negatively correlated (of course, this also holds if we use any triplet X_i, X_j, X_k , for different values of i, j and k). This implies

$$\min(\alpha_-, \alpha_+) \geq \gamma^2 \geq \beta.$$

We then easily check the following implications under H_0 :

$$\begin{aligned}
(i, j, k, l) \in \Omega_1 &\implies \mathbb{E}(\text{sign}(X_k - X_i) \mathbb{1}_{\{|X_i - X_k| > d\}} \text{sign}(X_l - X_j) \mathbb{1}_{\{|X_j - X_l| > d\}}) = 0 \\
(i, j, k, l) \in \Omega_2 \cup \Omega_3 &\implies \mathbb{E}(\text{sign}(X_k - X_i) \mathbb{1}_{\{|X_i - X_k| > d\}} \text{sign}(X_l - X_j) \mathbb{1}_{\{|X_j - X_l| > d\}}) = \alpha_- + \alpha_+ - 2\beta \\
(i, j, k, l) \in \Omega_4 \cup \Omega_5 &\implies \mathbb{E}(\text{sign}(X_k - X_i) \mathbb{1}_{\{|X_i - X_k| > d\}} \text{sign}(X_l - X_j) \mathbb{1}_{\{|X_j - X_l| > d\}}) = 2\beta - \alpha_- - \alpha_+ \\
(i, j, k, l) \in \Omega_6 &\implies \mathbb{E}(\text{sign}(X_k - X_i) \mathbb{1}_{\{|X_i - X_k| > d\}} \text{sign}(X_l - X_j) \mathbb{1}_{\{|X_j - X_l| > d\}}) = 2\gamma
\end{aligned}$$

Our conclusion is that under H_0 ,

$$\text{Var}(S_{ex} | H_0) = \left(\frac{1}{3}n^3 - n^2 + \frac{2}{3}n\right)(\alpha_- + \alpha_+ - 2\beta) + (n^2 - n)\gamma.$$

2.2 Estimating the variance of S_{ex}

A natural way to estimate $\text{Var}(S_{ex}|H_0)$ would be to use unbiased estimates of α_- , α_+ , β and γ . Since these constants are defined as expectations, we will estimate them by averages. Also, since under H_0 all variables are iid, we will use a symmetric average:

$$\begin{aligned}
\hat{\alpha}_+ &= \frac{1}{n(n-1)(n-2)} \sum_{\{i,j,k : i \neq j \neq k\}} \mathbb{1}_{\{X_i > X_j + d\}} \mathbb{1}_{\{X_i > X_k + d\}} \\
&= \frac{1}{n(n-1)(n-2)} \sum_{i=1}^n \sum_{j \neq k} \mathbb{1}_{\{X_i > X_j + d\}} \mathbb{1}_{\{X_i > X_k + d\}} \\
&= \frac{1}{n(n-1)(n-2)} \sum_{i=1}^n \left(\sum_{j=1}^n \mathbb{1}_{\{X_i > X_j + d\}} \cdot \sum_{k=1}^n \mathbb{1}_{\{X_i > X_k + d\}} - \sum_{j=1}^n \mathbb{1}_{\{X_i > X_j + d\}} \right) \\
&= \frac{1}{n(n-1)(n-2)} \sum_{i=1}^n u_i^2 - u_i.
\end{aligned}$$

Here we define

$$u_i = \#\{j \in \{1, \dots, n\} : X_i > X_j + d\} \quad \text{and} \quad v_i = \#\{j \in \{1, \dots, n\} : X_i < X_j - d\}.$$

Similar calculations show

$$\begin{aligned}
\hat{\alpha}_- &= \frac{1}{n(n-1)(n-2)} \sum_{i=1}^n v_i^2 - v_i \\
\hat{\beta} &= \frac{1}{n(n-1)(n-2)} \sum_{i=1}^n u_i v_i \\
\hat{\gamma} &= \frac{1}{n(n-1)} \sum_{i=1}^n u_i = \frac{1}{n(n-1)} \sum_{i=1}^n v_i.
\end{aligned}$$

Finally we get as an estimator for the variance of $S_{ex}|H_0$:

$$\widehat{\text{Var}}(S_{ex}|H_0) = \frac{1}{3} \sum_{i=1}^n (u_i - v_i)^2 + \frac{1}{3} \sum_{i=1}^n u_i.$$

Our adapted Mann-Kendall test statistic T therefore becomes

$$T = \frac{S_{ex}}{\sqrt{\widehat{\text{Var}}(S_{ex})}} = \frac{\sum_{i < j} \text{sign}(X_j - X_i) \mathbb{1}_{\{|X_i - X_j| > d\}}}{\sqrt{\frac{1}{3} \sum_{i=1}^n (u_i - v_i)^2 + \frac{1}{3} \sum_{i=1}^n u_i}}.$$

2.3 Contiguous alternatives

We wish to study the power of our test for different choices of d in a theoretical manner. A good way to do this, is by using what is known as *contiguous alternatives* to the null hypothesis that there is no trend, by which we mean the following: suppose Z_1, \dots, Z_n are iid random variables, choose $\lambda \in \mathbb{R}$ and define

$$X_i = \lambda \frac{i}{n^{3/2}} + Z_i.$$

This means that the X_i 's have a trend (and so we are not in the null hypothesis), but as the sample size n grows, the trend becomes smaller: in the end-point the difference decreases like $n^{-1/2}$, precisely enough to still be noticeable by the classic Mann-Kendall test. The advantage of such alternatives is that we are able to write down the asymptotic power of our test in a more or less closed form, so that we can compare this power for different choices of d in our test. We need some notation: suppose $f(\cdot)$ is the density of Z_1 , and define $g(\cdot)$ as the density of $Z_1 - Z_2$. It is not hard to show that g has a maximum at 0 and it is symmetric, so $g(z) = g(-z)$ for all $z \in \mathbb{R}$. In fact we have

$$g(z) = \int_{\mathbb{R}} f(x+z)f(x) dx.$$

Now note that

$$\begin{aligned} |\mathbb{1}_{\{X_i > X_j + d\}} \mathbb{1}_{\{X_i > X_k + d\}} - \mathbb{1}_{\{Z_i > Z_j + d\}} \mathbb{1}_{\{Z_i > Z_k + d\}}| &\leq \mathbb{1}_{\{Z_i - Z_j \in [d - |\lambda(j-i)|n^{-3/2}, d + |\lambda(j-i)|n^{-3/2}]\}} + \\ &\quad \mathbb{1}_{\{Z_i - Z_k \in [d - |\lambda(k-i)|n^{-3/2}, d + |\lambda(k-i)|n^{-3/2}]\}} \\ &\leq \mathbb{1}_{\{Z_i - Z_j \in [d - |\lambda|n^{-1/2}, d + |\lambda|n^{-1/2}]\}} + \\ &\quad \mathbb{1}_{\{Z_i - Z_k \in [d - |\lambda|n^{-1/2}, d + |\lambda|n^{-1/2}]\}} \end{aligned}$$

This means that

$$\hat{\alpha}_{+,X} = \hat{\alpha}_{-,Z} + O_p(n^{-1/2}). \quad (2.2)$$

Here, the subscript X or Z means that we are either using the sample X_1, \dots, X_n , or the coupled sample Z_1, \dots, Z_n which satisfies the null hypothesis. The same conclusion holds for $\hat{\alpha}_-$ and $\hat{\beta}$, which means that

$$\frac{\widehat{\text{Var}}(S_X)}{(\frac{1}{3}n^3 - n^2 + \frac{2}{3}n)} \rightarrow \alpha_- + \alpha_+ - 2\beta \quad (\text{as } n \rightarrow \infty),$$

where the constants are defined for the distribution of Z_1 . In other words, the contiguous alternative does not significantly alter the estimation of the variance of S_{ex} ; in fact

$$\text{Var}(S_X) = \text{Var}(S_Z)(1 + O(n^{-1/2})).$$

However, the alternative does significantly change the expected value of S_{ex} . Under the null hypothesis, we know that $S_Z = O_p(n^{3/2})$. Now let us calculate $\mathbb{E}(S_X)$:

$$\begin{aligned} \mathbb{E}(S_X) &= \sum_{i < j} \mathbb{P}(X_i < X_j - d) - \mathbb{P}(X_i > X_j + d) \\ &= \sum_{i < j} \mathbb{P}(Z_i - Z_j < -d + \lambda(j-i)n^{-3/2}) - \mathbb{P}(Z_i - Z_j > d + \lambda(j-i)n^{-3/2}) \\ &= \left(\sum_{i < j} 2g(d)\lambda(j-i)n^{-3/2} \right) (1 + o(1)) \\ &= \frac{1}{3} \lambda g(d) n^{3/2} (1 + o(1)). \end{aligned}$$

Here we used that $\mathbb{P}(Z_1 - Z_2 < -d) = \mathbb{P}(Z_1 - Z_2 > d)$ and the fact that $g(-d) = g(d)$. As you can see, the difference is of the order square root of the variance. Furthermore, from (2.2) we conclude that

$$T_X \sim N \left(\frac{\lambda g(d)}{\sqrt{3(\alpha_- + \alpha_+ - 2\beta)}}, 1 \right).$$

To investigate the power of our test against this contiguous alternative as a function of d (remember, we reject the null hypothesis if $|T_X| \geq z_{\alpha/2}$, the normal critical value), it is therefore enough that we investigate $\mathbb{E}_d(T_X)$, the expected value of T_X as a function of d , given by

$$\mathbb{E}_d(T_X) = \frac{\lambda g(d)}{\sqrt{3(\alpha_- + \alpha_+ - 2\beta)}}.$$

Here, we have left out the lower order terms as n goes to infinity. The power of our test is clearly an increasing function of $\mathbb{E}_d(T_X)$: if $\mathbb{E}_d(T_X) = 0$, the power equals α , and if $\mathbb{E}_d(T_X) \rightarrow \infty$, then the power increases to 1. We start by investigating the behaviour around $d = 0$. It is clear that $\mathbb{E}_d(T_X)$ is a symmetric function of d . After some tedious calculus, it turns out that this function has a positive second derivative at 0, if and only if

$$\int f(z)^2 dz \cdot \int f(z)^3 dz > \frac{1}{6} \int f'(z)^2 dz. \quad (2.3)$$

If this is the case, it means that we can gain power even asymptotically by choosing d slightly larger than 0. There exist distributions f of Z_i that do not satisfy Condition (2.3), but we will see soon that it is satisfied if the Z_i 's are normally distributed. For large values of d , things are not as clear. We will see that if the Z_i 's are normally distributed, $\mathbb{E}(T_X)$ will go to 0 when $d \rightarrow \infty$ (this means that the power of our test would go to α). We expect this behaviour in similar distributions, mainly because we know that $\alpha_+, \alpha_- > \gamma^2 > \beta$. If the first inequality is substantial, for example if we would have that for d large enough,

$$\alpha_+ > C\gamma^{2\eta},$$

for some $\frac{1}{2} \leq \eta < 1$ and some $C > 0$ (we always have that $\alpha_+ \leq \gamma$), then we conclude that

$$\mathbb{E}_d(T_X) < -\tilde{C}[(1 - G)^{1-\eta}(d)]'$$

where $\tilde{C} > 0$ and

$$G(d) = \mathbb{P}(Z_1 - Z_2 \leq d) = \int_{-\infty}^d g(s) ds = 1 - \gamma.$$

Since $(1 - G)(d) \rightarrow 0$ as $d \rightarrow \infty$, we would have that under these conditions and if g is eventually decreasing, $\mathbb{E}(T_X)$ goes to zero as $d \rightarrow \infty$ and the power of our test to α . A notable exception is when X_1 has a uniform distribution on, for example, $[0, 1]$. In that case, it would be good to choose d slightly larger than 1, and the power with respect to the contiguous alternative would go to 1. This is of course due to the fact that if we would see a difference between two X_i 's larger than 1, we could always reject the null hypothesis. However, also when for example g has local maxima around the integers, we expect it might be possible to increase the power of our test for these contiguous alternatives all the way to 1.

As mentioned before, an interesting case is if all Z_i 's have a normal distribution with standard deviation σ . Figure 1 shows $\mathbb{E}_d(T_X)$ as a function of d for $\sigma = 1$ and $\lambda = 1$.

The expected value for T is maximal around $d = 1.34\sigma$, but there is only a very slight increase in power. Note that if you pick d too large, the power quickly drops, and furthermore, the increase of power for small d is only quadratic in d (this holds for any differentiable f , and if Condition (2.3) is violated, there would be a quadratic *decrease* of power).




Figure 1: $\mathbb{E}_d(T_X)$ (vertical axis) against d .

2.4 Properties of t

For the standard S statistic (no *LRD* applied) and when no ties are present in the data, then $\max(S) = \frac{1}{2}n(n-1)$ and t is defined as :

$$t = \frac{2S}{n(n-1)}$$

In case of tied ranks, t_a or t_b can be used. If by c_{ij} we denote the score allotted to any pair of measurements subject to the conditions that $c_{ij} = -c_{ji}$ and $c_{ij} = 0$ if $i = j$ (so that, for S and given the natural order ($i < j$), $c_{ij} = 1$). For the variable representing time, we denote the score e_{ij} , subject to the same conditions. Then, t_a and t_b are defined as follows: t_a is still S divided by $\frac{1}{2}n(n-1)$ while in t_b the denominator is replaced by $\frac{1}{2}\sqrt{\sum c^2_{ij} \sum e^2_{ij}}$. For our extended Mann-Kendall test, it is straightforward to see that $\sum c^2_{ij} = \sum_i u_i + \sum_i v_i = 2 \sum_i u_i = 2 \sum_i v_i$. It then follows that in the case of the Mann-Kendall test where $\sum e^2_{ij} = n(n-1)$ since $e_{ij} = 1, \forall i \neq j$ for the time variable rankings, t_b will take the form:

$$t_{b_{ex}} = \frac{S_{ex}}{\frac{1}{2}\sqrt{\{2 \sum_i u_i\} \{n(n-1)\}}}$$

We refer the interested reader to the discussion provided by Kendall and Dickinson (1990, chapter 3) on whether to use t_a or t_b .

2.5 Distribution of S and T

Kendall and Dickinson Gibbons (1990) note that "the distribution of T for any fixed number of ties tends to normality as n increases, and there is probably little important error involved in using the normal approximation for $n \geq 10$, unless the ties are very extensive or very numerous in which case a special investigation may be necessary". This conjecture was verified by Burr (1960) who examined tied ranks in both rankings and the normal approximation was found not to be adequate only in cases where only a few of the extreme values of S are significant, e.g. the distribution of S is sparsely represented in the critical region. It is suggested that these should be taken under consideration when applying a *LRD* to a sample and ties structure should be monitored.

The distribution of S_{ex} (and T_{ex} since it is a multiple of S_{ex}) is always symmetric and

centered around 0 (under H_0), with the only modification with respect to its variance, when we are dealing with partially tied observations. Therefore, the continuity correction suggested by Kendall and Dickinson Gibbons (1990) is still applicable. This results in the Z statistic to be used for testing against trend where:

$$Z = \begin{cases} \frac{S_{ex}+1}{\sigma_{S_{ex}}} & \text{if } S_{ex} < 0 \\ 0 & \text{if } S_{ex} = 0 \\ \frac{S_{ex}-1}{\sigma_{S_{ex}}} & \text{if } S_{ex} > 0 \end{cases} \quad (2.4)$$

and $\sigma_{S_{ex}} = \sqrt{\text{Var}(S_{ex})}$ so that $Z|H_0 \sim N(0, 1)$

3 Simulation study

A simulation study is conducted to assess the operational characteristics of the proposed extended version of the Mann-Kendall test and its added value compared to the standard one. The study aims at evaluating the performance of the test in the settings of linear and quadratic trends over time. The general data-generating model is:

$$Y = \theta X^p + \epsilon$$

Where Y denotes the variable of interest, $X = \{1, \dots, n\}$ is the time variable and ϵ is the measurement error associated with this process. The simulation parameters include $\theta \in \{0, 1\}$, $p \in \{1, 2\}$, $n \in \{20, 30\}$ and ϵ is generated from either Normal or Uniform distributions with mean 0 and standard deviations $\sigma^{1/p} \in \{10, 15, 20\}$. The parameters of the Uniform distributions $U(a, b)$ are thus calculated as $a = -\sigma\sqrt{3}$ and $b = a$. This corresponds to σ ranging from 10 to 400 (for $p = 2$ and $\sigma^{1/p} = 20$) and a ranging from -17 to -690. For each simulation scenario, the extended Mann-Kendall test is applied, with LRD 's $d/\sigma \in \{0, 0.5, 1, 1.5, 2\}$. Thus, d is supposed to be determined relative to the error of the measurements, where $d=0$ refers to the standard Mann-Kendall test with which the comparison is the main goal. Testing is conducted at the 5% significance level, two-sided. The choice of the sample sizes and effect sizes was based on the pursuit to explore the validity of the normality assumption of the extended test in borderline small sample sizes in combination with the concept of partial ties. All simulations were conducted with the R language for statistical computing. The code is publicly available on the first author's github repository (

The simulations results are presented in Tables 2 and 3. Table 2 presents empirical power values (and thus type I error for $\theta = 0$) for the explored scenaria while Table 3 presents the average proportion of ties observed in each scenario, $\pi_t = \mathbb{E} \left(\frac{maxS - \sum_i u_i}{maxS} \right)$, where $maxS = \frac{1}{2}n(n-1)$. In order to evaluate the performance of the extended test we jointly refer at each respective cell from the two tables, so that validity of the normality assumption is assessed together with the resulting operational characteristics.

In general, normality assumption seems to hold for $\pi_t < 0.6$ and might be judged adequate for even larger values when the errors distribution is assumed to be Uniform (type I errors close to nominal levels). The latter however seems to be a matter of both the sample size and the extent of tied values, therefore attention should be paid, as with the standard version of the test. The extended version of the Mann-Kendall test is doing what is intended, namely increasing the

power of the Mann-Kendall test in the presence of measurement error. For normally distributed errors and a sample size of 30, (Table 2a) applying a *LRD* increases power for values of d/σ up to 1.5 and a slight drop is observed for $d/\sigma = 2$. The drop could however be attributed to the inadequacy of the normal approximation due to an extensive number of resulting ties when applying such a (relatively) large d , as the type I error is shown to deviate from the nominal level as well.

The gains in power seem to be even more substantial in the case of uniformly distributed errors (2b) in combinations of sample sizes and π_t 's where the normal approximation seems to hold (type I error levels close to nominal). In such cases there is shown to be a substantial increase in power for large measurement error, when this is taken into account through the application of a *LRD*. It is evident from the simulation results that both the magnitude and the distribution of the measurement error should be taken into account so that an optimal d to be chosen for the analysis.

4 Application

We demonstrate the extended Mann-Kendall test by applying it to the blood donation data introduced in Section 1.1. We will limit the application to the first five years of the study (2001-2005) during which measurement uncertainty was of greater concern, given that the questionnaire was newly developed and amendments and uniform concepts definitions took place during those years (Janssen et al., 2015).

A dataset included in this report (number of donated platelet therapeutic units per 1000 inhabitants) was chosen for demonstrating the extended Mann-Kendall test in a combined version of it, namely using the Regional Kendall test for trend (Helsel and Frans, 2006). The Regional Kendall test for trend, in the same way as the Seasonal Kendall test for trend (Hirsch et al., 1982), combines datasets across different regions in an overall test to decide upon the existence of a trend across these regions. If data are available over m regions for the same time periods, the overall statistic, S_r can be computed by :

$$S_r = \sum_{L=1}^m S_L, \quad (4.1)$$

where S_L denotes the S statistic calculated from the annual (or any period) data of the L 'th region. Its variance is the sum of the regional variances and the corrected score, Z_r is computed as above (Equation 2.4). When the product of the number of regions and number of periods is more than 25 (Hirsch et al., 1982), the distribution of S_r can be approximated by a normal distribution, therefore Z_r can be used for testing against trend.

Table 4 presents the data, considering countries without missing values only.

The extended Mann-Kendall test can provide a mean for applying an overall *LRD*, or a *LRD* that differs per country. The results presented are just to illustrate the use of the extended Mann-Kendall test and not to contradict or doubt the results presented in the report, which were drawn with different methods and based on the complete dataset. Let us demonstrate some possible ways of applying the extended Mann-Kendall test and the consequences for conclusions drawn.

First, if a simple Regional Kendall test is conducted, so $LRD=0$. This yields a value for $S_r=41$ and a two-sided p -value of approximately 0.024. Testing on the 0.05 significance level, one would conclude that there is a significant upward trend in platelets donated within the European Union in the period of 2001-2005. When one considers a difference of 0.4 platelets donation per 1000 inhabitants irrelevant, the p -value would be 0.07 and this would result in a

Table 2: Probability of rejection per simulation scenario. Data are generated from $\theta X^p + \epsilon$ where ϵ follows the respective distribution. See text for details.

(a) Normal distribution						(b) Uniform dist.					
			θ			θ					
			0	1		0	1				
			p			p					
n	$\sigma^{\frac{1}{p}}$	d/σ		1	2		1	2			
20	10	0.0	0.047	0.632	0.996	0.047	0.592	0.998			
		0.5	0.044	0.643	0.997	0.044	0.591	0.999			
		1.0	0.042	0.643	0.997	0.043	0.586	0.999			
		1.5	0.038	0.619	0.997	0.040	0.595	0.999			
		2.0	0.027	0.553	0.996	0.034	0.621	0.998			
	15	0.0	0.047	0.333	0.564	0.047	0.309	0.524			
		0.5	0.044	0.337	0.574	0.045	0.298	0.522			
		1.0	0.042	0.334	0.573	0.043	0.285	0.516			
		1.5	0.038	0.313	0.552	0.040	0.287	0.524			
		2.0	0.027	0.259	0.488	0.034	0.300	0.545			
20	0.0	0.047	0.207	0.222	0.047	0.195	0.208				
	0.5	0.045	0.208	0.223	0.045	0.184	0.197				
	1.0	0.043	0.205	0.220	0.043	0.174	0.187				
	1.5	0.038	0.189	0.204	0.040	0.173	0.187				
	2.0	0.027	0.151	0.164	0.034	0.176	0.191				
30	10	0.0	0.049	0.992	1.000	0.049	0.992	1.000			
		0.5	0.047	0.993	1.000	0.047	0.993	1.000			
		1.0	0.046	0.994	1.000	0.046	0.994	1.000			
		1.5	0.044	0.993	1.000	0.045	0.996	1.000			
		2.0	0.038	0.990	1.000	0.042	0.997	1.000			
	15	0.0	0.049	0.833	1.000	0.049	0.803	1.000			
		0.5	0.047	0.840	1.000	0.047	0.800	1.000			
		1.0	0.046	0.843	1.000	0.046	0.800	1.000			
		1.5	0.044	0.835	1.000	0.045	0.824	1.000			
		2.0	0.038	0.801	1.000	0.042	0.874	1.000			
20	0.0	0.049	0.593	0.918	0.049	0.559	0.901				
	0.5	0.047	0.599	0.924	0.047	0.546	0.903				
	1.0	0.046	0.602	0.928	0.046	0.533	0.908				
	1.5	0.044	0.589	0.924	0.044	0.557	0.924				
	2.0	0.038	0.545	0.904	0.041	0.627	0.946				

Table 3: Average proportion of ties per simulation scenario

(a) Normal distribution						(b) Uniform dist.		
n	$\sigma^{\frac{1}{p}}$	d/σ	θ		p	θ		p
			0	1		0	1	
						1	2	
		0.0	0.00	0.00	0.00	0.00	0.00	0.00
		0.5	0.28	0.21	0.10	0.27	0.20	0.10
		1.0	0.52	0.4	0.19	0.49	0.39	0.19
		1.5	0.71	0.57	0.28	0.68	0.56	0.28
		2.0	0.84	0.71	0.36	0.82	0.70	0.36
30	15	0.0	0.00	0.00	0.00	0.00	0.00	0.00
		0.5	0.28	0.24	0.17	0.27	0.23	0.17
		1.0	0.52	0.46	0.33	0.49	0.44	0.33
		1.5	0.71	0.64	0.48	0.68	0.62	0.48
		2.0	0.84	0.78	0.61	0.82	0.76	0.61
	20	0.0	0.00	0.00	0.00	0.00	0.00	0.00
		0.5	0.28	0.25	0.23	0.27	0.24	0.22
		1.0	0.52	0.48	0.43	0.49	0.46	0.42
		1.5	0.71	0.67	0.61	0.68	0.65	0.60
		2.0	0.84	0.80	0.75	0.82	0.79	0.74
	10	0.0	0.00	0.00	0.00	0.00	0.00	0.00
		0.5	0.28	0.24	0.17	0.27	0.23	0.17
		1.0	0.52	0.46	0.33	0.50	0.44	0.33
		1.5	0.71	0.64	0.48	0.68	0.62	0.47
		2.0	0.84	0.78	0.60	0.82	0.76	0.60
20	15	0.0	0.00	0.00	0.00	0.00	0.00	0.00
		0.5	0.28	0.26	0.24	0.27	0.24	0.23
		1.0	0.52	0.49	0.46	0.49	0.47	0.44
		1.5	0.71	0.67	0.64	0.68	0.65	0.62
		2.0	0.84	0.81	0.78	0.82	0.8	0.77
	20	0.0	0.00	0.00	0.00	0.00	0.00	0.00
		0.5	0.28	0.27	0.26	0.27	0.25	0.25
		1.0	0.52	0.50	0.50	0.49	0.48	0.48
		1.5	0.71	0.69	0.69	0.68	0.66	0.66
		2.0	0.84	0.82	0.69	0.82	0.81	0.80

Table 4: Platelets donated per 1000 inhabitants for countries with data available for all 5-years.

Country	Year				
	2001	2002	2003	2004	2005
Belgium	4.65	4.57	4.87	5.82	6.98
Bulgaria	1.38	2.02	2.16	0.71	0.70
Czech Republic	2.11	2.29	2.19	2.37	2.10
Finland	6.67	6.66	6.07	6.17	6.42
France	3.26	3.22	3.32	3.35	3.50
Germany	3.27	3.27	4.04	4.53	4.45
Greece	11.67	12.88	12.63	15.85	14.95
Iceland	3.21	2.42	3.38	3.17	3.66
Italy	8.26	12.84	3.37	2.16	2.75
Latvia	1.06	1.70	1.80	1.66	1.74
Netherlands	9.39	3.45	2.92	3.23	3.19
Norway	3.43	3.12	2.99	3.48	3.39
Poland	0.94	1.49	1.04	1.30	1.62
Romania	1.57	1.84	1.84	2.72	3.13
Slovak Republic	1.36	1.72	1.60	1.60	1.87
Slovenia	14.00	11.92	11.07	13.08	13.96
Sweden	3.52	3.79	3.63	3.90	3.67
Switzerland	2.43	2.02	3.12	2.51	2.69
United Kingdom	4.50	4.42	4.53	4.44	4.39

different conclusion. Another way of applying an LRD could be by making different assumptions per country. For example one could argue that it is not reasonable to apply the same LRD on the data from Greece, where the average platelets per 1000 inhabitants are 13.6 and on the data from Bulgaria where the average platelets per 1000 inhabitants are 1.39. So, a LRD as a percentage of the average reported platelets per 1000 inhabitants could be chosen. Results of the Regional Kendall test for trend when a 0.05, 0.10 and 0.20 of the average reported platelets per 1000 inhabitants is used as an LRD alongside with the results discussed above are presented in Table5.

We see that the choice of LRD is crucial for the results obtained. A different LRD will lead to different results as far as the significance of the trend is concerned. Therefore it is recommended that prior knowledge on uncertainty in the observed values or consensus on a meaningful difference in subsequent observations is obtained before applying the extended Mann-Kendall test.

5 Discussion

In this paper, we extend the Mann-Kendall test for trend to allow for measurement error and partial ties as defined here. That is, we propose that two values do not have to be exactly equal to be accounted as such but rather a LRD is applied so that two values are considered equal if their difference is smaller than some value d . This provides flexibility for the Mann-Kendall test to allow for measurement error and/or incorporation of a meaningful interpretation of observed outcomes. The idea of partial ties could essentially be extended to any rank-based test, even though this is beyond the scope of the present manuscript.

Table 5: Results of Regional Kendall test applying extended Mann-Kendall test per country.

LRD	Results		
	S_r	$\text{Var}(S_r)$	2-sided p -value
0	41	315.67	0.0244
0.05	45	295.67	0.0105
0.20	41	223.67	0.0075
5% of c.m.*	49	239.67	0.0019
10% of c.m.	41	175	0.0025

* countries' mean
platelet use

We derive a formula for the variance of the extended statistic and provide analytical results for a class of contiguous alternatives. We also demonstrated the behavior of the test in the presence of a linear or quadratic trend with normally or uniformly distributed errors via a simulation study. The operational characteristics of the extended Mann-Kendall test are appealing compared to the conventional Mann-Kendall test, as the extended test increases power while retaining the type I error to nominal levels.

We demonstrate that for small d , there is (asymptotically) a marginal gain in power for the class of contiguous alternatives studied here. This gain is shown to be a function of the error variance when the errors are normally distributed. Via simulation we explore linear and quadratic alternatives for normal and uniform errors, and we show that there can be substantial increase in power for the case of uniformly distributed errors. Our main conclusion is that the extended test will be useful once a user has a substantial understanding of the data generating mechanism and the measurement error.

Attention should be paid when using a LRD with respect to the resulting number of ties. Applying a relatively large LRD can lead to a large number of tied values in the data. In extreme occasions, this could lead to implications concerning the normal approximation of the distribution of S_{ex} . In such cases a permutation test will be more appropriate in order to acquire exact p -values which would always be valid for such a model (Ernst et al., 2004).

As shown in the examples presented, the approach can also be used when combining several Mann-Kendall tests over time (Seasonal Kendall test for trend) or regions (Regional Kendall test for trend). In this case extra attention should be paid so the LRD applied is realistic for a specific dataset. Different conclusions can result from different values of LRD , therefore d should be meaningful and case-specific.

The extension can be useful for applied researchers in many disciplines where non-parametric testing for trend is applied. It requires knowledge of the measurement error in the observed data, but it will contribute to more robust conclusions about the existence of a trend over time.

References

- Burr, E. J. (1960). The distribution of kendall's score s for a pair of tied rankings. *Biometrika* 47(1/2), 151–171.
- Dawood, M. et al. (2017). Spatio-statistical analysis of temperature fluctuation using mann-kendall and sen's slope approach. *Climate dynamics* 48(3-4), 783–797.

- Ernst, M. D. et al. (2004). Permutation methods: a basis for exact inference. *Statistical Science* 19(4), 676–685.
- Helsel, D. R. and L. M. Frans (2006). Regional kendall test for trend. *Environmental science and technology* 40, 4066–4073.
- Hirsch, R. M., J. R. Slack, and R. A. Smith (1982). Techniques of trend analysis for monthly water quality data. *Water resources Research* 18, 10–121.
- Janssen, M., L. van Hoeven, and G. Rautmann (2015). Trends and observations on the collection, testing and use of blood and blood components in europe 2001–2011 report. *Directorate for the Quality of Medicines and Health-Care of the Council of Europe (EDQM)*.
- Kendall, M. G. (1938). A new measure of rank correlation. *Biometrika* 30, 81–93.
- Kendall, M. G. (1945). The treatment of ties in ranking problems. *Biometrika* 33, 239–245.
- Kendall, M. G. and J. Dickinson Gibbons (1990). *Rank correlation methods*. Oxford University Press.
- Manios, E., K. Vemmos, G. Tsivgoulis, G. Barlas, K. Eleni, K. Spengos, and N. Zakopoulos (2007). Comparison of noninvasive oscillometric and intra-arterial blood pressure measurements in hyperacute stroke. *Blood pressure monitoring* 12(3), 149–156.
- Mann, H. B. (1945). Nonparametric tests against trend. *Econometrica* 13, 245–259.
- Nourani, V., A. D. Mehr, and N. Azad (2018). Trend analysis of hydroclimatological variables in urmia lake basin using hybrid wavelet mann–kendall and şen tests. *Environmental earth sciences* 77(5), 207.
- Panneton, M. and P. Robillard (1972). The exact distribution of kendall’s s with ties in one ranking. *Journal of the Royal statistical society series c - applied statistics* 21, 321–323.
- Robillard, P. (1972). Kendall’s s distribution with ties in one ranking. *Journal of the American statistical association* 67, 36–40.
- Sillitto, G. (1947). The distribution of kendall’s t coefficient of rank correlation in rankings containing ties. *Biometrika* 34, 36–40.
- Soderberg, K. and R. J. Hennet (2007). Uncertainty and trend analysis - radium in ground water and drinking water. *Ground Water Monitoring and Remediation* 27, 122–129.
- Van der Poel, C., M. Janssen, and M. Behr-Gross (2011). Trends and observations on the collection, testing and use of blood and blood components in europe. 2001-2005 report. *Directorate for the Quality of Medicines and Health-Care of the Council of Europe (EDQM)*.