Maximum pairwise-rank-likelihood-based inference for the semiparametric transformation model

By Tao Yu

Department of Statistics & Applied Probability, National University of Singapore, Singapore, 117546

Email: stayt@nus.edu.sg

Pengfei Li

Department of Statistics and Actuarial Science, University of Waterloo, Waterloo, Ontario, Canada N2L 3G1 Email: pengfei.li@uwaterloo.ca

Baojiang Chen

Department of Biostatistics and Data Science, University of Texas Health Science Center at Houston, School of Public Health in Austin, Austin, Texas 78701, USA

Ao Yuan

Department of Biostatistics, Bioinformatics and Biomathematics, Georgetown University, Washington DC, 20057, USA Email: ay312@georgetown.edu

Jing Qin National Institute of Allergy and Infectious Diseases, National Institutes of Health, MD 20892, USA Email: jingqin@niaid.nih.gov

Abstract

In this paper, we study the linear transformation model in the most general setup. This model includes many important and popular models in statistics and econometrics as special cases. Although it has been studied for many years, the methods in the literature are based on kernel-smoothing techniques or make use of only the ranks of the responses in the estimation of the parametric components. The former approach needs a tuning parameter, which is not easily optimally specified in practice; and the latter is computationally expensive and may not make full use of the information in the data. In this paper, we propose two methods: a pairwise rank likelihood method and a score-function-based method based on this pairwise rank likelihood. We also explore the theoretical properties of the proposed estimators. Via extensive numerical studies, we demonstrate that our methods are appealing in that the estimators are not only robust to the distribution of the random errors but also lead to mean square errors that are in many cases comparable to or smaller than those of existing methods.

Keywords: Linear transformation model, M-estimation, profile likelihood, pairwise rank likelihood, pseudo-likelihood, semiparametric inference

1 Introduction

In this paper, we consider the linear transformation model in the most general setup. Let $\{X_i, Y_i\}, i = 1, ..., n$ be independent and identically distributed (i.i.d.) copies of $\{X, Y\}$, where Y denotes the response and X is a $p \times 1$ random vector of covariates. The transformation model assumes that

$$H(Y_i) = X_i^T \boldsymbol{\beta} + \epsilon_i, \tag{1}$$

where ϵ_i is the random error with unknown cumulative distribution function (c.d.f.) $F_{\epsilon}(\cdot)$, and we define $F(\cdot)$ to be the c.d.f. of $\epsilon_i - \epsilon_j$, $i \neq j$. We assume that $H(\cdot)$, $F(\cdot)$, and β are all unknown parameters, where $H(\cdot)$ is a strictly increasing function, and $E(\epsilon_i) = 0$. Furthermore, we need to impose some conditions so that this model is identifiable. One such condition is that $\|\beta\|_2 = 1$, $F_0(\cdot)$ is strictly increasing, and the support of X_i , denoted by \mathcal{X} , contains at least one interior point in \mathbb{R}^p . Here, $F_0(\cdot)$ denotes the true value of $F(\cdot)$; see Remark 10 in the Appendix for more discussion of the identifiability of the model.

The class of linear transformation models is of great practical value; it includes many important and popular models in statistics and econometrics as special cases. A comprehensive review of these special cases is as follows. The popular Box–Cox model (Box and Cox 1964) can be viewed as a special case of the linear transformation model with both $H(\cdot)$ and the distribution of the random errors modeled parametrically; in particular, the distribution of the random errors is typically assumed to follow a normal distribution (see Carroll and Ruppert 1988). The log-linear regression and accelerated failure time models model the distribution of the random errors nonparametrically but assume that the transformation function $H(\cdot)$ satisfies a specific parametric form (see Bickel and Doksum 1981 and the references therein). The parametric and semiparametric proportional hazard models (Cox 1972) can also be formulated as special cases of linear transformation models. Specifically, they can be formulated as the linear transformation model with the transformation function model as the linear transformation of the random errors modeled parametrically (Zeng and Lin 2007); see Horowitz (1996) for a more detailed discussion.

Developing appropriate methodology and fast algorithms that accurately estimate the parameters in transformation models is a challenging task because of the nonparametric unknown components. To overcome this difficulty, there are two general strategies. The first is kernel-smoothing techniques (see for example Song et al. 2007; Lin and Peng 2013; and Zhang et al. 2018). These methods rely on a tuning parameter, which may not be specified optimally. The second strategy is rank-based methods: the main idea is to use $\text{Rank}(Y_i)$ to replace $H(Y_i)$ in the loss function in the estimation of the parametric parameters. This is motivated by the fact that $\text{Rank}(Y_i) = \text{Rank}(H(Y_i))$ because of the monotonicity of $H(\cdot)$. Examples include the maximum rank correlation (MRC) method (Han 1987), the monotone rank (MR) method (Cavanagh and Sherman 1998), and the pairwise-difference rank (PDR) method (Abrevaya 1999a,b). These methods are fast and do not need a tuning parameter. However, they use the rank information (rather than the full information) of the responses and may be less efficient. Furthermore, estimating the transformation function $H(\cdot)$ and the c.d.f. $F(\cdot)$ may be of particular interest in some applications. For example, the estimate of $H(\cdot)$, together with that of β , can be used to predict the response Y for every given X. The estimates of $H(\cdot)$ and $F(\cdot)$ can be

used to check the model assumptions of some parametric and semiparametric models, e.g., the multiple linear regression model and the Cox PH model reviewed in Section 2.2. Estimation methods for $H(\cdot)$ are available in the literature (e.g., Horowitz 1996; Chen 2002; Zhang 2013), but those for $F(\cdot)$ are not.

We consider the transformation model (1) by treating $H(\cdot), F(\cdot)$, and β as unknown parameters. We propose two methods for estimating β and $F(\cdot)$. First, in the spirit of the pseudolikelihood method (Besag 1975), we propose a pairwise rank likelihood method; then, using a strategy similar to that of Groeneboom and Hendrickx (2018), we propose a score-functionbased method based on this pairwise rank likelihood. Numerical studies show that both methods lead to desirable β and $F(\cdot)$ estimates. Furthermore, we explore the asymptotic properties of the proposed estimators. We expect that the methodology, theoretical results, and technical tools of this paper will benefit the study of similar models (e.g., the Box–Cox model). Via extensive simulation studies, we show that our methods are robust to the distribution function $F_{\epsilon}(\cdot)$ and lead to mean square errors (MSEs) that are in many cases comparable to or smaller than those of existing methods.

The rest of the paper is organized as follows. Section 2 reviews two popular methods; they are the main competitors of our methods in the numerical studies. Section 3 presents our pairwise likelihood and score-function-based methods for the estimation of the unknown parameters in the linear transformation model (1). Section 4 investigates the asymptotic properties of these estimates, and Section 5 discusses the simulation studies. Section 6 applies our methods to a real-data example, and Section 7 ends the paper with a discussion. The technical conditions are relegated to the Appendix, and the technical details are given in the supplementary material.

2 Existing Methods

We now briefly review several existing methods that are relevant to our work. Our numerical studies will use them for comparison purposes.

2.1 Pairwise-difference rank estimators

Abrevaya (2003) gave a thorough review of rank estimators for the linear transformation model (1). These include the MRC estimator (Han 1987), the MR estimator (Cavanagh and Sherman 1998), and the PDR estimators (Abrevaya 1999a, 1999b). There are several versions of the PDR estimators. PDR3 is based on observation triples, and PDR4 is based on observation quadruples. As demonstrated by Abrevaya (2003), PDR4 has the best performance of all the above estimators.

PDR4 considers the following objective function:

$$S_4(\beta) = \sum_{(i,j,k,l) \in \mathcal{C}_4} I\{(X_i - X_j)^T \beta > (X_k - X_l)^T \beta\} \cdot \{I(Y_i > Y_j) - I(Y_k > Y_l)\},\$$

with

$$C_4 = \{(i, j, k, l) : i, j, k, l \in \{1, \dots, n\}, i \neq j, i \neq k, i \neq l, j \neq k, j \neq l, k \neq l\}$$

being the quadruples of the index set $\{1, ..., n\}$. The estimator of the unknown parameter β is defined to be

$$\widehat{\boldsymbol{\beta}}_{PDR4} = \operatorname{argmax}_{\boldsymbol{\beta}} S_4(\boldsymbol{\beta}).$$

Although PDR4 performs well, it does not make full use of the data information, and it does not result in an estimator for the nonparametric function $F(\cdot)$. We propose a pairwise rank likelihood method for estimating both $F(\cdot)$ and the parametric parameter β .

2.2 Proportional hazard model

The proportional hazard model (Cox 1972, 1975), called the Cox PH model hereafter, has been frequently studied and widely applied to analyze censored survival data from the medical sciences. It is a special case of the linear transformation model (Doksum 1987). Let Y_i be a survival time with c.d.f. $F_i(y)$ and probability density function (p.d.f.) $f_i(y)$, and let X_i be the $p \times 1$ clinical covariate. Let $r_i(y) = f_i(y)/\{1 - F_i(y)\}$ be the hazard rate. The Cox PH model assumes

$$r_i(y) = r(y) \exp(X_i^T \boldsymbol{\beta}),$$

which has an equivalent form (Lehmann 1953):

$$F_i(y) = 1 - \{1 - h(y)\}^{\exp(X_i^T / \beta)},$$

where $h(y) = 1 - \exp \left\{ -\int_0^y r(t) dt \right\}$. This immediately leads to the model

$$H(Y_i) = -X_i^T \boldsymbol{\beta} + \epsilon_i,$$

where $H(y) = \log \left[-\log\{1 - h(y)\}\right]$ and the ϵ_i 's are i.i.d. with the extreme value distribution. The partial likelihood method (Cox 1972, 1975) is commonly used to estimate β . Since the error is specified as the extreme value distribution, the Cox PH model may not be robust if this distribution is misspecified. For example, another popular model in survival analysis is the proportional odds ratio model where the error distribution is the logistic distribution (Bennett, 1983a, 1983b). If this is the underlying model, then a method based on the Cox model may not be robust and may lead to larger errors in the β estimation.

3 Proposed Estimation Methods

Existing methods for semiparametric regression models include the profile likelihood method (Breslow, 1972), the partial likelihood method (Cox 1972, 1975), and the rank likelihood method (Kalbfleisch and Prentice, 1973). Because of the existence of sets of infinitely many

nuisance parameters $F_{\epsilon}(\cdot)$ and $H(\cdot)$, none of these methods can be applied directly. In this section, we propose two methods to estimate both $F(\cdot)$ and β in the transformation model (1) effectively. First, in the spirit of the pseudo-likelihood method (Besag 1975), we propose a pairwise rank likelihood method. Second, using this pairwise rank likelihood and a strategy similar to that of Groeneboom and Hendrickx (2018), we propose a score-function-based method.

3.1 Pairwise rank likelihood estimation

We propose the pairwise rank likelihood and establish a fast algorithm for the estimation of β and $F(\cdot)$, where we recall that $F(\cdot)$ is the c.d.f. of $\epsilon_i - \epsilon_j$, $i \neq j$.

Given the monotonicity of $H(\cdot)$, we immediately have

$$P(Y_i > Y_j | X_i, X_j) = P(H(Y_i) > H(Y_j) | X_i, X_j)$$

= $P(X_i^T \beta + \epsilon_i > X_j^T \beta + \epsilon_j | X_i, X_j)$
= $P(\epsilon_j - \epsilon_i < (X_i - X_j)^T \beta | X_i, X_j)$
= $F((X_i - X_j)^T \beta).$ (2)

This motivates the pairwise rank log-likelihood, given by

$$\ell(\boldsymbol{\beta}, F) = \sum_{i \neq j} \left[I(Y_i > Y_j) \log\{P(Y_i > Y_j | X_i, X_j)\} + I(Y_i \leqslant Y_j) \log\{P(Y_i \leqslant Y_j | X_i, X_j)\} \right]$$

$$= \sum_{i \neq j} \left[I(Y_i > Y_j) \log\{F((X_i - X_j)^T \boldsymbol{\beta})\} + I(Y_i \leqslant Y_j) \log\{1 - F((X_i - X_j)^T \boldsymbol{\beta})\} \right].$$
(3)

Note that $\ell(\beta, F)$ is essentially the log-likelihood function of the generalized linear model with $I(Y_i > Y_j), i, j = 1, ..., n$ and $i \neq j$ being the responses and $F(\cdot)$ being the link function. Consequently

$$(\widehat{\boldsymbol{\beta}}, \widehat{F}) = \operatorname{argmax}_{\boldsymbol{\beta} \in \mathcal{B}, F \in \mathcal{F}} \ell(\boldsymbol{\beta}, F),$$
(4)

where \mathcal{B} is a compact subspace of \mathbb{R}^p and

 $\mathcal{F} = \{F(\cdot) : F(x) \in [0, 1] \text{ and is monotonically increasing}\}.$

We propose the following two-stage algorithm that solves the optimization problem (4) numerically:

Stage 1. For a given β , profile F to obtain the profile likelihood of β through the following steps:

(a) Let $(v_1(\beta), \ldots, v_K(\beta))$ be a vector composed of $\{(X_i - X_j)^T \beta : i \neq j\}$, and (I_1, \ldots, I_K) be the corresponding vector composed of $\{I(Y_i > Y_j) : i \neq j\}$. Sort $v_k(\beta)$ in increasing order:

$$v_{(1)}(\boldsymbol{\beta}) \leq \ldots \leq v_{(K)}(\boldsymbol{\beta}).$$

The corresponding I_j 's are denoted by $I_1(\beta), \ldots, I_K(\beta)$. Substituting $v_{(j)}(\beta)$ and the corresponding $I_j(\beta)$ into (3), we have

$$\ell(\boldsymbol{\beta}, F) = \sum_{j=1}^{K} [I_j(\boldsymbol{\beta}) \log\{F(v_{(j)}(\boldsymbol{\beta}))\} + \{1 - I_j(\boldsymbol{\beta})\} \log\{1 - F(v_{(j)}(\boldsymbol{\beta}))\}]$$

(b) For any β and the $\ell(\beta, F)$ given in (a), let

$$\widehat{F}_{\boldsymbol{\beta}} = \operatorname{argmax}_{F \in \mathcal{F}} \ell(\boldsymbol{\beta}, F).$$
(5)

(c) The profile pairwise rank log-likelihood is given by

$$\ell(\boldsymbol{\beta}, \widehat{F}_{\boldsymbol{\beta}}) = \sum_{i \neq j} \Big[I(Y_i > Y_j) \log\{\widehat{F}_{\boldsymbol{\beta}}((X_i - X_j)^T \boldsymbol{\beta})\} + I(Y_i \leqslant Y_j) \log\{1 - \widehat{F}_{\boldsymbol{\beta}}((X_i - X_j)^T \boldsymbol{\beta})\} \Big].$$
(6)

Stage 2. Maximize (6) with respect to β to obtain $\hat{\beta}$. More details of the numerical implementations for $\hat{\beta}$ are given in Section 9 of the supplementary material.

Note that the optimization problem (5) in Stage 1(b) can be readily solved by the classical pool-adjacent-violation-algorithm (PAVA; Ayer et al. 1955) and active set methods (de Leeuw et al. 2009). This is because, based on Ayer et al. (1955), we immediately conclude that

$$\widehat{F}_{\boldsymbol{\beta}}(v_{(j)}(\boldsymbol{\beta})) = \max_{s \leqslant j} \min_{t \geqslant j} \frac{\sum_{h=s}^{t} I_h(\boldsymbol{\beta})}{t-s+1}, j = 1, \dots, K,$$

which is equivalent to the solution from the classical isotonic regression problem (Robertson et al. 1988),

$$\widehat{F}_{\boldsymbol{\beta}} = \operatorname{argmin}_{F \in \mathcal{F}} \sum_{j=1}^{K} \left\{ I_j(\boldsymbol{\beta}) - F(v_{(j)}(\boldsymbol{\beta})) \right\}^2$$

subject to F being a nondecreasing function.

Remark 1. The linear transformation model (1) is essentially a semiparametric model; it contains two nonparametric parameters, $F_{\epsilon}(\cdot)$ and $H(\cdot)$, and one $(p \times 1)$ -dimensional parametric parameter β . It may be difficult to construct and directly maximize a single objective function that includes all the unknown parameters (e.g., the full semiparametric likelihood). Our method solves this problem by first omitting the nuisance nonparametric component $H(\cdot)$ using the pseudo-likelihood technique; a profile-type method is then applied to further overcome the potential difficulty that may be caused by $F(\cdot)$. Thus, the objective function carries only the parametric parameter β , and it can be readily estimated by existing algorithms.

Remark 2. Our pairwise rank likelihood approach takes all the orders of $(X_i - X_j)^T \beta$ into account, while PDR4 considers only the paired order of $(X_i - X_j)^T \beta$ and $(X_k - X_l)^T \beta$. Therefore, our method has used the information from the data more effectively. Moreover, our method can be considered a pairwise-rank-based approach in which the transformation function $H(\cdot)$ is eliminated.

Remark 3. For presentational convenience, in all our analyses we have assumed that the responses Y_i have no ties. However, in practice, ties may occasionally occur in the outcome data, e.g., if the collected responses are subject to some rounding mechanism. In Section 10 of the supplementary material, we suggest a solution for this problem in the framework of our pairwise rank log-likelihood.

3.2 Score-function-based method based on the pairwise rank likelihood

As observed in Section 4 and discussed in Remark 6, it is difficult to establish the root *n* consistency and asymptotic distribution for the maximum pairwise rank log-likelihood estimator $\hat{\beta}$. In this section, using a strategy similar to that of Groeneboom and Hendrickx (2018), we consider an alternative estimator $\hat{\beta}$ whose asymptotic distribution can be established; see Section 4.2.

We need the following notation. For any $\beta \in \mathcal{B}$, define

$$F_{\boldsymbol{\beta}}(t) = E\left\{I(Y_1 > Y_2) \middle| (X_1 - X_2)^T \boldsymbol{\beta} = t\right\}$$
$$= \int F_0(t + \boldsymbol{v}^T(\boldsymbol{\beta}_0 - \boldsymbol{\beta})) f_{(X_1 - X_2)|(X_1 - X_2)^T} \boldsymbol{\beta}_{=t}(\boldsymbol{v}) d\boldsymbol{v}$$

and

$$\varphi_{\boldsymbol{\beta}}(t) = E(X_1 - X_2 | (X_1 - X_2)^T \boldsymbol{\beta} = t), \quad \varphi_0(t) = \varphi_{\boldsymbol{\beta}_0}(t).$$
(7)

We observe that for any β , \hat{F}_{β} defined by (5) is essentially an estimator for F_{β} , and

$$\widehat{\boldsymbol{\beta}} = \arg \max_{\boldsymbol{\beta} \in \boldsymbol{\mathcal{B}}} \ell\left(\boldsymbol{\beta}, \widehat{F}_{\boldsymbol{\beta}}\right),$$

where the form of $\ell\left(\boldsymbol{\beta}, \hat{F}_{\boldsymbol{\beta}}\right)$ is given in (6).

The score function for $\ell\left(\boldsymbol{\beta}, \hat{F}_{\boldsymbol{\beta}}\right)$ (if it exists) has the form

$$\overline{\psi}_n(\boldsymbol{\beta}) = \sum_{i \neq j} \xi(X_i, X_j; \widehat{F}_{\boldsymbol{\beta}}, \boldsymbol{\beta}) \left\{ I(Y_i > Y_j) - \widehat{F}_{\boldsymbol{\beta}} \left((X_i - X_j)^T \boldsymbol{\beta} \right) \right\},\$$

with $\xi(\cdot)$ being a function depending on the data and the unknown parameter β . Then $\hat{\beta}$ can

be viewed as the root of this score function. However, there are two aspects that complicate the development of the optimal convergence rate and asymptotic distribution for $\hat{\beta}$: (1) this score function may not exist, since $\hat{F}_{\beta}((X_i - X_j)^T\beta)$ may not be differentiable with respect to β ; (2) the function $\xi(\cdot)$ has a complicated form and may depend on β . To bypass these two difficulties, we first replace $\xi(X_i, X_j; \hat{F}_{\beta}, \beta)$ by $X_i - X_j$ in this score function. Second, we would like to define $\tilde{\beta}$ to be the root of $\psi_n(\beta) = 0$, where

$$\psi_n(\boldsymbol{\beta}) = \frac{1}{n^2} \sum_{i \neq j} (X_i - X_j) \left\{ I(Y_i > Y_j) - \hat{F}_{\boldsymbol{\beta}} \left((X_i - X_j)^T \boldsymbol{\beta} \right) \right\}.$$
(8)

However, since $\psi_n(\beta)$ may not be continuous, this root may not exist, so, similarly to Groeneboom and Hendrickx (2018), we define:

 $\widetilde{\boldsymbol{\beta}}$ is the *zero-crossing* of $\psi_n(\boldsymbol{\beta})$,

where the zero-crossing of a function (or a mapping) is defined below.

Definition 1. For a function $\psi: \mathcal{X} \to \mathbb{R}$, x is called the zero-crossing of ψ if every open neighborhood of x contains x_1, x_2 such that $\psi(x_1)\psi(x_2) \leq 0$. For a mapping $\widetilde{\psi}: \mathcal{X} \to \mathbb{R}^d$, xis called the zero-crossing of $\widetilde{\psi}$ if x is the zero-crossing of each component of $\widetilde{\psi}$.

Details of the numerical implementations of $\tilde{\beta}$ are given in Section 9 of the supplementary material. With $\tilde{\beta}$ and the corresponding techniques given in Section 3.1, we estimate $F(\cdot)$ by $\tilde{F}(\cdot) = \hat{F}_{\tilde{\beta}}(\cdot)$.

Remark 4. With the estimators of β and $F(\cdot)$, estimation methods for $H(\cdot)$ are available in the literature (see, e.g., Horowitz 1996; Chen 2002; Zhang 2013). Chen (2002) proposed a rank-based estimator for $H(\cdot)$, assuming that the β estimate is available, and Zhang (2013) proposed a self-induced smoothing method based on Chen (2002)'s estimator. If the estimate for β is root n consistent, both estimates can achieve root n consistency and converge weakly to Gaussian processes. In Section 4.2, we show that $\tilde{\beta}$ is root n consistent; therefore, with $\tilde{\beta}$, the corresponding $H(\cdot)$ estimates from both papers have desirable asymptotic properties.

Remark 5. In this paper, we have focused on the case where the Y_i 's are complete data. Our methods, however, can be extended to analyze the transformation model when some Y_i 's are right censored. See the supplementary material for this extension.

4 Asymptotic Properties

In this section, we explore the asymptotic properties of our estimators. Similarly to Section 3, we organize this section into two subsections, respectively establishing the asymptotic properties of the estimators given in Sections 3.1 and 3.2. For space limitations and presentational continuity, we have relegated the technical conditions to the Appendix, and the technical details are in the supplementary material.

4.1 Asymptotic properties for the pairwise rank likelihood estimators

We first investigate the asymptotic properties of the pairwise rank likelihood estimators proposed in Section 3.1. Let

$$D(\boldsymbol{\beta}_{2}, F_{2}; \boldsymbol{\beta}_{1}, F_{1}) = \left[\int \int \left\{ F_{1}((\boldsymbol{x}_{1} - \boldsymbol{x}_{2})^{T} \boldsymbol{\beta}_{1}) -F_{2}((\boldsymbol{x}_{1} - \boldsymbol{x}_{2})^{T} \boldsymbol{\beta}_{2}) \right\}^{2} dF_{X}(\boldsymbol{x}_{1}) dF_{X}(\boldsymbol{x}_{2}) \right]^{1/2}, \quad (9)$$

where $F_X(\cdot)$ is the c.d.f. of X_i . Theorem 1 below establishes the convergence rates of $\hat{\beta}$ and $\hat{F}(\cdot)$.

Theorem 1. Assume Conditions 0–6 in the Appendix. We have

(a)
$$D(\hat{\beta}, \hat{F}; \beta_0, F_0) = O_p(n^{-1/3}),$$

(b) $\hat{\beta} - \beta_0 = O_p(n^{-1/3}),$

where β_0 and F_0 are the true values of β and F.

Remark 6. The development of the asymptotic properties given in the above theorem is a challenging task for two reasons. First, the structure of the pairwise rank likelihood function (3) is complicated. It is not a sum of i.i.d. components, and the M-estimator theory from empirical processes is not directly applicable. Second, the monotonic nonparametric components in the transformation model complicate the development of the convergence rate for the estimator of the parametric component. Such problems are long-standing. For example, Huang and Wellner (1993) encountered a similar challenge. They studied current status data under the accelerated failure time model assumption, and they proved only the $o_p(1)$ convergence of their estimators. In other words, they proved consistency but did not prove a convergence rate. In the theorem above, we show that the convergence rate for $\hat{\beta}$ is at least $O_p(n^{-1/3})$. However, we conjecture that this rate is not optimal; the best rate may be $O_p(n^{-1/2})$. We leave this important and interesting problem for future research.

4.2 Asymptotic distribution for the score-function-based estimators

In this section, we establish the asymptotic distributions for $\tilde{\beta}$ and $\tilde{F}(\cdot)$.

Recall the definition of $\psi_n(\cdot)$ in (8). Its population version $\psi_0(\beta)$ can be defined to be:

$$\psi_0(\boldsymbol{\beta}) = \int \int (\boldsymbol{x}_1 - \boldsymbol{x}_2) \left\{ I(y_1 > y_2) - F_{\boldsymbol{\beta}}((\boldsymbol{x}_1 - \boldsymbol{x}_2)^T \boldsymbol{\beta}) \right\} \\ \times dF_{X,Y}(\boldsymbol{x}_1, y_1) dF_{X,Y}(\boldsymbol{x}_2, y_2),$$
(10)

where $F_{X,Y}(\boldsymbol{x}, y)$ is the joint c.d.f. of (X, Y).

We have the following lemma for $\psi_0(\cdot)$.

Lemma 1. Assume Conditions 6, A1, and A2 in the Appendix. We have the following:

- (1) $\psi_0(\boldsymbol{\beta}_0) = 0;$ (2) $\psi'_0(\boldsymbol{\beta}_0) = \frac{\partial \psi_0(\boldsymbol{\beta})}{\partial \boldsymbol{\beta}} \Big|_{\boldsymbol{\beta} = \boldsymbol{\beta}_0}$ exists with rank p - 1.
- (3) Since $\|\beta_0\|_2 = 1$, there exists an *i* such that the *i*th component of β_0 is nonzero. Define $A = \psi'_0(\beta_0) + \overline{A}$, where \overline{A} is a $p \times p$ matrix with *i*th row β_0^T and all other entries 0. Then A is of full rank, and

$$\psi_0(\boldsymbol{\beta}) = A \cdot (\boldsymbol{\beta} - \boldsymbol{\beta}_0) + o(\boldsymbol{\beta} - \boldsymbol{\beta}_0).$$

We have the following theorem, which establishes the asymptotic properties of $\tilde{\beta}$.

Theorem 2. Assume Conditions 0–2, 6, and A1–A3 in the Appendix. Denote by $\tilde{\beta}$ the zerocrossing of $\psi_n(\beta)$ (if it exists). We have the following:

- (1) When $n \to \infty$, a zero-crossing $\widetilde{\beta}$ of $\psi_n(\beta)$ exists with probability tending to 1.
- (2) $\widetilde{\boldsymbol{\beta}} \to \boldsymbol{\beta}_0$ in probability.
- (3) Recalling the matrix A defined in Lemma 1 Part (3), we have

$$\widetilde{\boldsymbol{\beta}} - \boldsymbol{\beta}_{0} = \frac{A^{-1}}{n^{2}} \sum_{i \neq j} \left[\left\{ X_{i} - X_{j} - \varphi_{0} \left((X_{i} - X_{j})^{T} \boldsymbol{\beta}_{0} \right) \right\} \times \left\{ I(Y_{i} > Y_{j}) - F_{0} \left((X_{i} - X_{j})^{T} \boldsymbol{\beta}_{0} \right) \right\} \right] + o_{p} (\widetilde{\boldsymbol{\beta}} - \boldsymbol{\beta}_{0}) + o_{p} (n^{-1/2}).$$
(11)

To derive the asymptotic distribution for $\tilde{\beta}$, we need to work on a U-statistic with the kernel:

$$h(\boldsymbol{x}_1, y_1; \boldsymbol{x}_2, y_2) = (\boldsymbol{x}_1 - \boldsymbol{x}_2) \left\{ I(y_1 > y_2) - F_0 \left((\boldsymbol{x}_1 - \boldsymbol{x}_2)^T \boldsymbol{\beta}_0 \right) \right\}.$$

Denote

$$\hbar(\boldsymbol{x}_1, y_1) = E\{h(\boldsymbol{x}_1, y_1; X_2, Y_2)\}.$$
(12)

The theorem above leads to the asymptotic distribution of $\tilde{\beta}$.

Corollary 1. Assume the conditions of Theorem 2. Furthermore, assume that both $\Sigma_X = var(X)$ and $\Sigma_{\hbar} = var \{\hbar(X, Y)\}$ are of full rank, and

$$\boldsymbol{\alpha}^{T} E\left(X_{2} - X_{1} | (X_{2} - X_{1})^{T} \boldsymbol{\beta}_{0}\right) = 0,$$

for any $\boldsymbol{\alpha} \in \mathbb{R}^{p}$, and $\boldsymbol{\alpha}^{T} \Sigma_{X} \boldsymbol{\beta}_{0} = 0.$ (13)

Denote $\widetilde{A} = \psi'_0(\beta_0)\Sigma_X^{-1} + \overline{A}$, where \overline{A} is defined in Lemma 1 Part (3). We have

$$\sqrt{n}(\widetilde{\boldsymbol{\beta}} - \boldsymbol{\beta}_0) \to 2A^{-1}\widetilde{A}^{-1}\psi_0'(\boldsymbol{\beta}_0)\Sigma_X^{-1}N(\mathbf{0}, \Sigma_\hbar)$$
(14)

in distribution, as $n \to \infty$.

Remark 7. Write $Z_1 = \alpha^T (X_2 - X_1)$, $Z_2 = \beta_0^T (X_2 - X_1)$. Then the condition given by (S.7) essentially requires that $E(Z_1|Z_2) = E(Z_1)$ is implied by $cov(Z_1, Z_2) = 0$, since the latter is straightforwardly verified by $\alpha^T \Sigma_X \beta_0 = 0$. This condition is satisfied, for example, when X_i follows a multivariate normal distribution.

Remark 8. In the above lemma, we have derived the explicit formula for the asymptotic variance of $\tilde{\beta}$; therefore, plug-in methods can be employed to estimate this variance when the data are available. However, this formula has a complicated form, and plug-in methods may not lead to the desired accuracy. We suggest that one can incorporate bootstrap methods to estimate this variance and perform statistical inference for β ; in Section 5, we suggest a bootstrap percentile confidence interval (BPCI) method that can be used for this inference. From our numerical studies, we observe that the coverage probability of the BPCI based on $\tilde{\beta}$ is reasonably close to the nominal level.

The proof of this corollary uses standard results from the literature for the asymptotic distribution of the U-statistic; the details are given in the supplementary material.

Next, we establish the asymptotic distribution of $\widetilde{F}(\cdot) = \widehat{F}_{\widetilde{A}}(\cdot)$.

Theorem 3. Assume that all the conditions of Corollary 1 are satisfied, and also assume the following conditions:

<u>Condition F1</u>: Recall that $F_{\epsilon}(s)$ is the c.d.f. for ϵ_i . Assume that it is continuous for $s \in \mathbb{R}$ and is continuously differentiable and strictly monotone for s in its support.

<u>Condition F2</u>: Denote by $F_{X^T\beta}(s)$ the c.d.f. for $X^T\beta$. Assume that it is continuously differentiable for s in its support. Let $f_{X^T\beta}(s) \equiv \frac{\partial F_{X^T\beta}(s)}{\partial s}$; assume that for any s, it is continuous for β in the neighborhood of β_0 ; and $f_{X^T\beta_0}(s)$ is continuous for s in its support.

Then, we have for every t,

$$n^{1/2}\left\{\widehat{F}_{\widetilde{\boldsymbol{\beta}}}(t) - F_0(t)\right\} \to N\left(0, \sigma^2(t)/g_0^2(t)\right),$$

in distribution, where $g_0(t)$ denotes the p.d.f. for $(X_2 - X_1)^T \boldsymbol{\beta}_0$, $\sigma^2(t) = var \{\zeta_t(X, Y)\}$,

$$\begin{aligned} \zeta_t(\boldsymbol{x}, y) &= \left\{ 1 - F_{\epsilon}(H_0(y) - \boldsymbol{x}^T \boldsymbol{\beta}_0 - t) - F_0(t) \right\} f_{X^T \boldsymbol{\beta}_0}(\boldsymbol{x}^T \boldsymbol{\beta}_0 + t) \\ &+ \left\{ F_{\epsilon}(H_0(y) - \boldsymbol{x}^T \boldsymbol{\beta}_0 + t) - F_0(t) \right\} f_{X^T \boldsymbol{\beta}_0}(\boldsymbol{x}^T \boldsymbol{\beta}_0 - t) \\ &+ 2g_0(t) \dot{F}_0^T(t) A^{-1} \tilde{A}^{-1} \psi_0'(\boldsymbol{\beta}_0) \Sigma_X^{-1} \hbar(\boldsymbol{x}, y), \end{aligned}$$

with $\dot{F}_0(t) = \partial F_{\beta}(t) / \partial \beta|_{\beta=\beta_0}$ and $\hbar(\cdot, \cdot)$ defined by (S.6).

With developments similar to but simpler than those of Theorem 3, we are able to establish the asymptotic distribution for $\hat{F}_{\beta}(t)$ for every given $\beta \in \mathcal{B}$ and $t \in \mathbb{R}$; we summarize this result in the following corollary.

Corollary 2. Assume Conditions 0–2 and A1 in the Appendix and Conditions F1' and F2' given below.

<u>Condition F1'</u>: $F_{\epsilon,\beta}(y,s)$ defined by (S.10) is continuous for $s \in \mathbb{R}$ and is continuously differentiable for s in the support and $\beta \in \mathcal{B}$.

<u>Condition F2'</u>: Denote by $F_{X^T\beta}(s)$ the c.d.f. for $X^T\beta$. Assume that it is continuously differentiable for s in the support. Recall $f_{X^T\beta}(s)$ defined by Condition F2 in Theorem 3; assume that it is continuous for s in the support and $\beta \in \beta$.

For every $t \in \mathbb{R}$ *and* $\beta \in \mathcal{B}$ *, we have*

$$n^{1/2}\left\{\widehat{F}_{\boldsymbol{\beta}}(t) - F_{\boldsymbol{\beta}}(t)\right\} \to N\left(0, \sigma_{\boldsymbol{\beta}}^2(t)/g_{\boldsymbol{\beta}}^2(t)\right),$$

in distribution, where $g_{\beta}(\cdot)$ denotes the p.d.f. of $(X_2 - X_1)^T \beta$, and

$$\sigma_{\boldsymbol{\beta}}(t) = E \Big[var \Big\{ f_{X^T \boldsymbol{\beta}}(X_1^T \boldsymbol{\beta} + t) F_{\epsilon, \boldsymbol{\beta}}(Y_1, X_1^T \boldsymbol{\beta} + t) - f_{X^T \boldsymbol{\beta}}(X_1^T \boldsymbol{\beta} - t) F_{\epsilon, \boldsymbol{\beta}}(Y_1, X_1^T \boldsymbol{\beta} - t) \Big| X_1 \Big\} \Big]$$

$$F_{\epsilon, \boldsymbol{\beta}}(y, s) = E \Big\{ F_{\epsilon}(H(y) - X^T \boldsymbol{\beta}_0) \Big| X^T \boldsymbol{\beta} = s \Big\}.$$
(15)
(15)
(15)
(15)
(15)
(16)

Interestingly, we observe from Theorem 3 and Corollary 2 that with any fixed β or our β estimate, our $F(\cdot)$ estimate is able to achieve root n consistency and is asymptotically normally distributed. As far as we are aware, we are the first to establish a root n consistent estimate for this $F(\cdot)$. A good estimate may be valuable in both theory and practice. First, we have illustrated in the real-data example that this $F(\cdot)$ estimate can be used to perform diagnostics for the distribution of ϵ_i . Second, this estimate with its desirable theoretical properties may benefit other research problems. For example, in the Harrells C-index measurement in medical applications, $F(\cdot)$ plays an important role because

$$E\left\{I(Y_i < Y_j)I(X_i^T \boldsymbol{\beta} < X_j^T \boldsymbol{\beta}) \middle| X_i, X_j\right\}$$

= $F\left((X_j - X_i)^T \boldsymbol{\beta}\right)I\left((X_i - X_j)^T \boldsymbol{\beta} < 0\right).$

5 Simulation Studies

In this section, we investigate the performance of our methods. We compare our pairwise rank likelihood estimators from Section 3.1 (referred to as "Our-PRL") and the score-function-based estimators from Section 3.2 (referred to as "Our-score") with: (1) the PDR4 method

reviewed in Section 2.1; (2) the partial likelihood method based on the Cox PH model reviewed in Section 2.2; (3) the smoothed rank correlation estimator (Ma and Huang 2005, Lin and Peng 2013) using the normal cumulative density function to approximate the indicator function (denoted "Smooth-Normal"); (4) the smoothed rank correlation estimator using the sigmoid function to approximate the indicator function (denoted "Smooth-Sigmoid"; Song et al. 2007); and (5) the smoothed maximum rank correlation estimator (denoted "SMRCE"; Zhang et al. 2018). Clearly, all the methods are built on semiparametric models. However, the Cox PH method requires a stronger model assumption than the others, i.e., it requires that the random error follows the extreme value distribution.

As discussed in Remark 8, the variance formula for β has a complicated form; estimating this variance by a plug-in method may not be practical. On the other hand, in many applications it may be of particular interest to estimate the standard error of β_k for k = 1, ..., p and to carry out statistical inference for this parameter. For example, it may be of interest to test the hypothesis H_0 : $\beta_k = \beta_{k0}$ versus H_1 : $\beta_k \neq \beta_{k0}$ for some given $\beta_{k0} \in \mathbb{R}$. We propose to estimate the standard error of β_k and perform this test via a nonparametric bootstrap method (Efron 1979). The procedure is summarized as follows:

- Generate B bootstrap samples, {(X^{*}_{i,b}, Y^{*}_{i,b}), i = 1,..., n}, b = 1,..., B, with B being a large number. For each b = 1,..., B, {(X^{*}_{i,b}, Y^{*}_{i,b}), i = 1,..., n} is a random sample from {(X_i, Y_i), i = 1,..., n} with replacement.
- For each b = 1, ..., B, based on Our-score, we can estimate $\tilde{\beta}_{k,b}^*$ from the bootstrap sample $\{(X_{i,b}^*, Y_{i,b}^*), i = 1, ..., n\}$. The standard error of $\tilde{\beta}_k$ can be estimated by the sample standard deviation of these $\{\tilde{\beta}_{k,b}^*\}_{b=1}^B$. The level 1α (BPCI) for β_k can be constructed as $[L_k^*, U_k^*]$, where L_k^* and U_k^* are respectively the $(\alpha/2)$ th and $(1 \alpha/2)$ th quantiles of $\{\tilde{\beta}_{k,b}^*\}_{b=1}^B$; this can be used to perform the aforementioned test.

For the other methods given at the beginning of this section, we can similarly establish the corresponding standard errors of the β_k estimates and the level $1 - \alpha$ BPCIs. In this section, we will also compare the performance of $\hat{\beta}$ with that of the other methods based on their BPCIs. Since we have not established the asymptotic normality of $\hat{\beta}$, we do not include the simulation results of the BPCI based on $\hat{\beta}$.

We have not established the theoretical justification for the BPCI based on the proposed β . The main obstacle is the nonparametric monotone component $\hat{F}_{\beta}(\cdot)$ in $\psi_n(\beta)$ (Shao and Tu, 1995; Kosorok, 2008). We use simulation studies to gain insight into the BPCI based on $\tilde{\beta}$, and we observe that the coverage proportions of the BPCIs based on $\tilde{\beta}$ are reasonably close to the nominal level. We leave the theoretical justification for future research.

5.1 Data simulation

We simulate the responses Y_i from the following model:

$$H(Y_i) = \beta_1 X_{i1} + \beta_2 X_{i2} + \epsilon_i.$$

The covariates X_{i1} and X_{i2} are simulated by $X_{i1} \sim \chi_1^2$ and $X_{i2} \sim N(X_{i1}, 1)$. We set $\beta_1 = \beta_2 = 1$ as the true value in the data simulation. To evaluate the performance of the methods for different true values of the nonparametric components, we experiment with different combinations of the transformation function $H(\cdot)$ and the distribution of the random errors ϵ_i . In particular,

- we consider (1) H(y) = y and (2) $H(y) = \log(y)$;
- for the distribution of the random errors ε_i, we use (1) the extreme value distribution, i.e., *F*_ε(t) = 1 − exp{−exp(t)}; (2) the N(0, π²/6) distribution; (3) the logistic distribution with mean 0 and scale 1/√2; so that all these distributions have the same variance π²/6.

For each combination, we consider the sample sizes n = 100 and 200. We repeat each simulation 1000 times.

5.2 Results

We apply the seven methods to the simulated data in Section 5.1, and in the results we normalize $\|\beta\|_2 = 1$ for comparison purposes. Tables 1–3 report the relative bias (RB), variance (Var), and MSE for β_1 and β_2 , over 1000 repetitions; here RB is defined as (estimate – true value)/(true value). All the values reported in these tables are $100 \times \text{computed values}$. Table 4 gives the proportions of the level 95% BPCIs that cover the true values of β_1 and β_2 , called the coverage proportion (CP), based on the Our-score, Cox PH, Smooth-Normal, Smooth-Sigmoid, and SMRCE methods over 1000 repetitions. Here we use B = 200 to compute the BPCI for each method. We do not include the BPCIs for Our-PRL because the asymptotic distribution of $\hat{\beta}$ is not available; we also omit those for PDR4 because they are too computationally intensive. Furthermore, since all the methods under comparison are established only on the ranks of the Y_i in the observed sample, they are invariant to the specific function form of $H(\cdot)$ in the estimation of β . We observe the same β estimates when we simulate the data by H(y) = y or $H(y) = \log(y)$. Therefore, we have reported only the results for data simulated from H(y) = y.

Table 1 presents the results for the case where the ϵ_i 's follow the extreme value distribution. We observe in this table, as well as in Tables 2 and 3, that the kernel-based methods (Smooth-Normal, Smooth-Sigmoid, and SMRCE) usually lead to larger MSE values than those of our methods (Our-PRL and Our-score); hereafter, we focus our discussion on the non-kernel methods (Our-PRL, Our-score, PDR4, and Cox PH). In this setup, the assumptions for all the methods are satisfied. The Cox PH method has the best performance; this is not surprising, since the assumption that the random error follows the extreme value distribution is satisfied, whereas the other methods do not need this assumption. Our-score, Our-PRL, and PDR4 have comparable MSE values for both β_1 and β_2 in this example.

Tables 2 and 3 present the results for the cases where the ϵ_i 's follow the normal and logistic distributions respectively. In both cases, especially the latter, the model assumption for the Cox PH method is violated, and this method has the worst performance among the non-kernel methods. In both tables, Our-score has the smallest MSE values. The MSEs of Our-PRL and PDR4 are comparable, but PDR4 is much more computationally intensive.

Table 1: Comparison of β_1 and β_2 estimation; the error follows the extreme value distribution with c.d.f. $F(t) = 1 - \exp\{-\exp(t)\}$

			β_1			β_2	
n	Method	RB	Var	MSE	RB	Var	MSE
100	Our-PRL	-1.47	1.64	1.65	-1.61	1.42	1.43
	Our-score	-1.98	1.48	1.50	-0.80	1.28	1.28
	PDR4	-2.62	1.52	1.55	-0.23	1.30	1.30
	Cox PH	-1.85	0.87	0.89	0.17	0.79	0.79
	Smooth-Normal	-4.11	1.95	2.03	0.47	1.61	1.61
	Smooth-Sigmoid	-5.52	1.70	1.85	2.33	1.31	1.34
	SMRCE	-11.37	2.75	3.40	6.07	1.72	1.90
200	Our-PRL	0.07	0.76	0.76	-1.58	0.74	0.76
	Our-score	-0.53	0.73	0.73	-0.91	0.70	0.70
	PDR4	-0.84	0.73	0.73	-0.58	0.69	0.69
	Cox PH	-0.03	0.42	0.42	-0.80	0.41	0.41
	Smooth-Normal	-1.83	0.96	0.97	0.00	0.86	0.86
	Smooth-Sigmoid	-2.60	0.86	0.89	0.95	0.76	0.76
	SMRCE	-6.22	1.19	1.39	3.86	0.90	0.98

Table 2: Comparison of β_1 and β_2 estimation; the error follows the normal distribution

			β_1			β_2	
n	Method	RB	Var	MSE	RB	Var	MSE
100	Our-PRL	-2.38	1.66	1.68	-0.76	1.46	1.46
	Our-score	-2.09	1.45	1.47	-0.68	1.30	1.30
	PDR4	-2.77	1.62	1.66	-0.31	1.42	1.42
	Cox PH	-4.99	1.69	1.81	1.73	1.43	1.45
	Smooth-Normal	-4.64	2.51	2.61	-0.01	2.04	2.04
	Smooth-Sigmoid	-6.01	2.20	2.37	1.93	1.69	1.71
	SMRCE	-12.73	3.50	4.30	6.08	2.16	2.34
200	Our-PRL	-0.91	0.72	0.72	-0.52	0.71	0.71
	Our-score	-0.81	0.69	0.69	-0.55	0.67	0.67
	PDR4	-1.11	0.74	0.75	-0.36	0.73	0.73
	Cox PH	-3.62	0.83	0.90	1.99	0.72	0.74
	Smooth-Normal	-1.87	1.07	1.09	-0.24	1.02	1.02
	Smooth-Sigmoid	-2.85	1.00	1.04	0.91	0.91	0.91
	SMRCE	-7.83	1.50	1.80	4.82	1.09	1.21

Table 4 shows that the CPs from Our-score are reasonably close to the nominal level (i.e., 0.95), indicating that the suggested bootstrap method controls the type I error well if we apply it to the testing problems $H_0: \beta_j = \beta_{j,0}$ versus $\beta_j \neq \beta_{j,0}; j = 1$ or 2.

			β_1			β_2	
n	Method	RB	Var	MSE	RB	Var	MSE
100	Our-PRL	-1.33	1.58	1.58	-1.72	1.45	1.47
	Our-score	-1.27	1.40	1.40	-1.43	1.28	1.29
	PDR4	-1.97	1.53	1.54	-0.94	1.36	1.37
	Cox PH	-5.38	2.06	2.21	1.53	1.63	1.64
	Smooth-Normal	-3.81	2.47	2.54	-0.67	1.95	1.95
	Smooth-Sigmoid	-5.22	2.10	2.24	1.40	1.58	1.59
	SMRCE	-11.44	3.24	3.89	5.36	2.05	2.19
200	Our-PRL	-0.55	0.72	0.73	-0.88	0.70	0.71
	Our-score	-0.42	0.67	0.67	-0.91	0.66	0.66
	PDR4	-0.71	0.69	0.70	-0.67	0.68	0.69
	Cox PH	-5.73	1.13	1.30	3.46	0.91	0.97
	Smooth-Normal	-1.59	1.01	1.02	-0.42	0.99	0.98
	Smooth-Sigmoid	-2.59	0.91	0.95	0.79	0.86	0.86
	SMRCE	-6.59	1.30	1.52	3.97	1.02	1.10

Table 3: Comparison of β_1 and β_2 estimation; the error follows the logistic distribution

We now examine the numerical performance of our $\tilde{F}(\cdot)$ in the estimation of $F(\cdot)$. We evaluate the mean and the sample standard deviation (SD) of the integrated square errors (ISEs) of $\tilde{F}(\cdot)$ over 1000 replicates, for all the simulated data in Section 5.1. The results are given in Table 5. We observe that as the sample size n increases, both the mean and SD of the ISEs decrease; this agrees with the asymptotic results we derived in Section 4. To further illustrate the performance of $\tilde{F}(\cdot)$ in the estimation of $F(\cdot)$, we construct the mean and the percentile bands. In Figure 1, we summarize the estimates of $\tilde{F}(\cdot)$ from our method for the case where n = 200 and $F(t) = 1 - \exp\{-\exp(t)\}$, i.e., the error ϵ_i follows the extreme value distribution. We have included the true curve, the 2.5% percentile, the 97.5% percentile, and the mean $\tilde{F}(\cdot)$ curves over 1000 repetitions. We observe that the mean curve of $\tilde{F}(\cdot)$ almost overlaps with the true curve; the 95% pointwise confidence bands contain the true curve; and the widths of the bands are quite small. This reinforces the asymptotic results derived for $\tilde{F}(\cdot)$ in Section 4.

In summary, our methods perform well for all the simulation setups. Among the PDR4, Our-PRL, and Our-score methods, Our-score leads to the smallest MSE values in most of the simulation examples. Our-PRL and PDR4 have similar performance in the β estimation, but PDR4 is much more computationally intensive. The Cox PH method performs the best when its model assumptions are satisfied. It fails when these assumptions are (partially) invalid, so these assumptions are clearly restrictive in practice. We therefore recommend Our-score unless there is strong prior scientific evidence that the assumptions of the Cox PH model are satisfied.

		Extreme Value		Normal		Logistic	
n	Method	β_1	β_2	β_1	β_2	β_1	β_2
100	Our-score	95.8	95.8	95.3	95.3	94.3	94.3
	Cox PH	93.8	93.8	93.2	93.2	91.8	91.8
	Smooth-Normal	96.1	96.1	96.3	96.3	96.5	96.5
	Smooth-Sigmoid	95.7	95.7	95.4	95.4	95.2	95.2
	SMRCE	97.2	98.1	97.8	99.1	96.8	97.7
200	Our-score	94.2	94.2	93.7	93.7	94.1	94.1
	Cox PH	93.8	93.8	93.7	93.7	92.2	92.2
	Smooth-Normal	97.5	97.5	96.2	96.2	96.8	96.8
	Smooth-Sigmoid	96.2	96.2	95.4	95.4	95.9	95.9
	SMRCE	98.1	98.1	97.4	97.4	98.0	98.0

Table 4: CP (%) of BPCI for β_1 and β_2

Table 5: Mean (×1000 and SD (×1000) of ISE for $\widetilde{F}(\cdot)$

	n = 100		n = 1	200
Distribution	MISE	SD	MISE	SD
Extreme Value	1.20	0.97	0.52	0.41
Normal	1.12	0.94	0.50	0.42
Logistic	1.15	0.91	0.53	0.44




Figure 1: Summarized $F(\cdot)$ estimation based on 1000 simulations; n = 200 and $F(t) = 1 - \exp\{-\exp(t)\}$.

6 Application to an Alzheimer's Disease Study

In this section, we analyze the National Alzheimer's Coordinating Center (NACC) uniform data set (UDS) (Beekly et al. 2007). Approximately five million people in the United States and more than thirty-seven million people worldwide suffer from Alzheimer's disease. This disease gradually destroys the patient's memory and ability to learn and to carry out daily activities such as talking and eating. As the disease progresses, there may also be changes in personality and behavior. Unfortunately, to date there is no cure and no effective way to predict how quickly an individual will progress through the stages of the disease. However, early diagnosis and appropriate treatment can slow its progression.

A popular test for Alzheimer's disease is the mini-mental state examination (MMSE) test (Folstein et al. 1975). This is a 30-point test that gives a numerical measurement of the cognitive impairment. We use the MMSE score as our response (Y) and study the association between this score and the covariates of interest, including the age (in years; X_1) and education level (in years; X_2) of the corresponding individual. In particular, we consider the linear transformation model, given by

$$H(Y) = \beta_1 X_1 + \beta_2 X_2 + \epsilon.$$

A sample of 100 subjects enrolled in 2005 was used to study the relationship between the covariates and the responses.

	β_1			β_2			
Method	Estimate	95% BPCI	Length	Estimate	95% BPCI	Length	
Our-Score	0.931	[0.796, 0.977]	0.181	-0.365	[-0.605, -0.212]	0.393	
PDR4	0.934	[0.736, 0.979]	0.243	-0.359	[-0.677, -0.205]	0.472	
Smooth-Normal	0.922	[0.669, 0.984]	0.315	-0.388	[-0.743, -0.180]	0.563	
Smooth-Sigmoid	0.924	[0.672, 0.984]	0.312	-0.383	[-0.740, -0.179]	0.561	
SMRCE	0.905	[0.627, 0.981]	0.354	-0.426	[-0.779, -0.194]	0.585	
Cox PH	0.957	[0.812, 0.988]	0.176	-0.289	[-0.584, -0.157]	0.427	
LR	0.934	[0.648, 0.983]	0.335	-0.358	[-0.761, -0.185]	0.576	

Table 6: Estimates and 95% BPCI of β estimates for the NACC UDS data; the Length columns report the length of the corresponding BPCIs

As we have observed in the simulation studies, Our-score has better performance than Our-PRL; in this real-data example, we do not include the results from Our-PRL. We apply the following methods: (1) Our-score; (2) PDR4; (3) Smooth-Normal; (4) Smooth-Sigmoid; (5) SMRCE; (6) Cox PH; and (7) classical linear regression (LR). We summarize the β estimates in Table 6, where we regularize $\|\beta\|_2 = 1$ for comparison purposes; the 95% BPCIs are based on the method given in Section 5 with B = 200. We observe that the β estimates from PDR4, Smooth-Normal, Smooth-Sigmoid, and LR are close to those from our methods; whereas those from Cox PH and SMRCE are slightly different. The lengths of the BPCIs from Our-score are comparable to those from Cox PH and much shorter than those from the other methods.

The estimates of $F(\cdot)$ from our methods can be used to check the model assumptions of LR and Cox PH. Specifically, note that $\widetilde{F}(\cdot)$ is the estimate for $F(\cdot)$, which is the c.d.f. of $\epsilon_i - \epsilon_j$. This c.d.f. can also be estimated under the LR model with normal errors or the Cox PH model; the estimated $F(\cdot)$'s are respectively N(0, 153.979) and Logistic(0, 6.807), the logistic distribution with mean 0 and scale 6.807. Figure 2 shows quantile-quantile plots of the $\widetilde{F}(\cdot)$ from Our-Score versus those of N(0, 153.979) (left panel) and Logistic(0, 6.807) (right panel). This figure indicates that the distribution of the errors may deviate from the normal and extreme value distributions.

Combining the observations above with those of Section 5.2, we expect that in this example, the bias from Cox PH could be larger than that from the other methods; our methods might have produced good estimates of the unknown parameters. However, we are unable to compare the estimates with their true values.




Figure 2: Quantile-quantile plots of the estimated $\tilde{F}(\cdot)$ from Our-Score versus those from the LR method (left panel) and the Cox PH method (right panel) for the NACC UDS data.

7 Discussion

The linear transformation model has been widely studied in statistics and econometrics. It includes as special cases many important and popular models that are commonly applied in practice. The development of appropriate methodology, fast algorithms, and solid statistical theory is an important but challenging task. In the literature, there are two general strategies: kernel-smoothing techniques and rank-based methods. However, the former approach needs a tuning parameter, and the latter may not make full use of the data information or may be computationally expensive.

We have developed two methods for analyzing the linear transformation model: (1) a pairwise rank likelihood method; and (2) a score-function-based method. Our methods do not need a tuning parameter and make effective use of the information carried in the original data. For the pairwise rank likelihood estimator, we have established a theoretical upper bound on the asymptotic convergence rate; for the score-function-based estimator, we have established asymptotic normality. Furthermore, via extensive numerical studies, we have demonstrated that our methods are more appealing than existing methods because they are not only robust to the distribution of the random errors but also in many cases lead to comparable or smaller MSEs in the estimation of the model parameters.

We expect that the methodology, theoretical results, and technical tools of this paper will benefit the study of similar models, e.g., the popular Box–Cox model (Box and Cox 1964) has a similar structure. We have assumed that the effect of the covariates on the transformed response is linear; future work will include relaxing and performing hypothesis testing on this assumption by considering more complicated models. The pairwise rank likelihood method could therefore be extended to these models. Some of the technical tools and theoretical results in this paper could facilitate these studies. Furthermore, the development of the asymptotic properties of the estimators for the pairwise rank likelihood (3) is challenging; see the more detailed discussion in Remark 6. Thus far, we have proved only $O_p(n^{-1/3})$ convergence. We conjecture that this rate may not be sharp for the estimator $\hat{\beta}$; there may be room for improvement. In addition, we have developed the theory only for the random design, i.e., where the X_i 's are random variables. We conjecture that for the fixed design, these theoretical results are still valid under appropriate regularity conditions. We leave these important and interesting topics for future research.

Acknowlegement

Dr. Yu's research is supported in part by Singapore Ministry Education Academic Research Fund Tier 1: R-155-000-157-112, R-155-000-202-114, and Ministry of Education of Singapore: MOE2014-T2-1-072. Dr. Li's research is supported in part by NSERC Grant RGPIN-2020-04964. Dr. Chen was supported in part by National Institute on Aging grant U01AG016976. The NACC database is funded by NIA/NIH Grant U01 AG016976. The NACC data are contributed by the NIA-funded ADCs: P30 AG019610 (PI Eric Reiman, MD), P30 AG013846 (PI Neil Kowall, MD), P50 AG008702 (PI Scott Small, MD), P50 AG025688 (PI Allan Levey, MD, PhD), P30 AG010133 (PI Andrew Saykin, PsyD), P50 AG005146 (PI Marilyn Albert, PhD), P50 AG005134 (PI Bradley Hyman, MD, PhD), P50 AG016574 (PI Ronald Petersen, MD, PhD), P50 AG005138 (PI Mary Sano, PhD), P30 AG008051 (PI Steven Ferris, PhD), P30 AG013854 (PI M. Marsel Mesulam, MD), P30 AG008017 (PI Jeffrey Kaye, MD), P30 AG010161 (PI David Bennett, MD), P30 AG010129 (PI Charles DeCarli, MD), P50 AG016573 (PI Frank LaFerla, PhD), P50 AG016570 (PI David Teplow, PhD), P50 AG005131 (PI Douglas Galasko, MD), P50 AG023501 (PI Bruce Miller, MD), P30 AG035982 (PI Russell Swerdlow, MD), P30 AG028383 (PI Linda Van Eldik, PhD), P30 AG010124 (PI John Trojanowski, MD, PhD), P50 AG005133 (PI Oscar Lopez, MD), P50 AG005142 (PI Helena Chui, MD), P30 AG012300 (PI Roger Rosenberg, MD), P50 AG005136 (PI Thomas Montine, MD, PhD), P50 AG033514 (PI Sanjay Asthana, MD, FRCP), and P50 AG005681 (PI John Morris, MD).

Appendix: Technical Conditions

We impose the following regularity conditions to establish the theoretical results in Section 4.1. They are not necessarily the weakest possible.

<u>Condition 0</u>: H(y) is a strictly increasing function on the support \mathcal{Y} of Y, and ϵ_i and X_i are independent.

<u>Condition 1</u>: $\beta \in \mathcal{B}$ and $F_X(x)$ is supported on \mathcal{X} , where both \mathcal{B} and \mathcal{X} are compact subspaces of \mathbb{R}^p .

<u>Condition 2</u>: For any $x_1, x_2 \in \mathbb{R}$, we have

$$\sup_{\boldsymbol{\beta}\in\mathcal{B}} \left| F_{\boldsymbol{\beta}^{T}_{X}}(x_{1}) - F_{\boldsymbol{\beta}^{T}_{X}}(x_{2}) \right| \lesssim |x_{1} - x_{2}|.$$

<u>Condition 3</u>: $\inf_{\boldsymbol{x}_1, \boldsymbol{x}_2 \in \mathcal{X}, \boldsymbol{\beta} \in \mathcal{B}} F_0((\boldsymbol{x}_1 - \boldsymbol{x}_2)^T \boldsymbol{\beta}) > 0.$

<u>Condition 4</u>: If F(x) is a c.d.f. such that $F((\boldsymbol{x}_1 - \boldsymbol{x}_2)^T \boldsymbol{\beta}) = F_0((\boldsymbol{x}_1 - \boldsymbol{x}_2)^T \boldsymbol{\beta}_0)$ for all $\boldsymbol{x}_1, \boldsymbol{x}_2 \in \mathcal{X}$, then $\boldsymbol{\beta} = \boldsymbol{\beta}_0$ and $F(\cdot) = F_0(\cdot)$.

<u>Condition 5</u>: $F_0(u)$ is second-order differentiable with bounded second-order derivatives and $\inf_{\boldsymbol{v}\in\mathcal{X}-\mathcal{X},\boldsymbol{\beta}\in\mathcal{B}}F'_0(\boldsymbol{v}^T\boldsymbol{\beta}) > 0.$

<u>Condition 6</u>: There exists an $\eta_0 > 0$ such that

$$\inf_{\boldsymbol{\beta}\in\mathcal{O}_{\eta_0}}\lambda_2\Big(\left[\operatorname{var}\{(X_1-X_2)-E((X_1-X_2)|(X_1-X_2)^T\boldsymbol{\beta})\}\right]\Big)>0,$$

where $\mathcal{O}_{\eta_0} = \{ \boldsymbol{\beta} : \| \boldsymbol{\beta} - \boldsymbol{\beta}_0 \|_2 \leq \eta_0, \| \boldsymbol{\beta} \|_2 = 1 \}; \lambda_2(B)$ denotes the second smallest eigenvalue of matrix B.

Furthermore, we need the following additional conditions to establish the theoretical results for $\tilde{\beta}$ in Section 4.2.

<u>Condition A1</u>: $F_{\beta}(t)$ is continuously differentiable in β and t, and

$$\inf_{t \in \{\boldsymbol{v}^T \boldsymbol{\beta}: \boldsymbol{v} \in \boldsymbol{\mathcal{X}} - \boldsymbol{\mathcal{X}}, \boldsymbol{\beta} \in \boldsymbol{\mathcal{B}}\}, \boldsymbol{\beta} \in \boldsymbol{\mathcal{B}}} \frac{\partial F_{\boldsymbol{\beta}}(t)}{\partial t} > 0.$$

Condition A2: $f_{X_1-X_2|(X_1-X_2)^T} \boldsymbol{\beta}_{=t}(\boldsymbol{v})$ is continuously differentiable in $\boldsymbol{\beta} \in \boldsymbol{\mathcal{B}}$ and $t \in \{\boldsymbol{x}^T \boldsymbol{\beta} : \boldsymbol{x} \in \boldsymbol{\mathcal{X}}, \boldsymbol{\beta} \in \boldsymbol{\mathcal{B}}\}.$

<u>Condition A3</u>: We have $\beta = \beta_0$ if

$$\operatorname{cov}\left\{\boldsymbol{\beta}_{0}^{T}(X_{1}-X_{2}), F_{0}(\boldsymbol{\beta}_{0}^{T}(X_{1}-X_{2})) \middle| \boldsymbol{\beta}^{T}(X_{1}-X_{2})\right\} = 0$$

almost surely.

Remark 9. Some of the conditions above are not intuitive and not easily checked in practice. In the supplementary document, we provide more discussion and give stronger but more intuitive conditions.

Remark 10. Condition 4 is required to ensure the identifiability of the model; this is because based on this condition and (2), we can conclude that β and $F(\cdot)$ are uniquely determined by the transformation model, and so is $H(\cdot)$. Based on our derivation in Section 3.2 of the supplementary document, this condition can be replaced with a more intuitive condition: $\|\beta\|_2 = 1$, $F_0(\cdot)$ is strictly increasing, and χ contains at least one interior point in \mathbb{R}^p . Furthermore, we observe that Condition A3 replaces Condition 4 when we ensure the identifiability of β in the proof of Theorem 2.

References

- Abrevaya, J. (1999a). Computation of the maximum rank correlation estimator. *Economics Letters*, 62, 279-285.
- Abrevaya, J. (1999b). Leapfrog estimation of a fixed-effects model with unknown transformation of the dependent variable. *Journal of Econometrics*, 93, 203-228.
- Abrevaya, J. (2003). Pairwise-difference rank estimation of the transformation model. *Journal* of Business & Economic Statistics, 21, 437-447.
- Ayer, M., Brunk, H. D., Ewing, G. M., Reid, W. T., Silverman, E. (1955). An empirical distribution function for sampling with incomplete information. *The Annals of Mathematical Statistics*, 26, 641-647.
- Beekly, D. L., Ramos, E. M., Lee, W. W., Deitrich, W. D., Jacka, M. E., Wu, J., et al. (2007). The National Alzheimers Coordinating Center (NACC) database: The uniform data set. *Alzheimer Disease and Associated Disorders*, 21(3), 249-258.
- Bennett, S. (1983a). Analysis of survival data by the proportional odds model. *Statistics in Medicine*, 2, 273-277.
- Bennett, S. (1983b). Log-logistic regression models for survival data. *Applied Statistics*, 32, 165-171.
- Besag, J. (1975). Statistical analysis of non-lattice data. The Statistician, 24, 179-195.
- Bickel, P. J. and Doksum, K. A. (1981). An analysis of transformations revisited. *Journal of the American Statistical Association*, 76, 296-311.
- Box, G. E. P. and Cox, D. R. (1964). An analysis of transformations. *Journal of the Royal Statistical Society, Series B*, 26, 211-252.

- Breslow, N. E. (1972). Contribution to the discussion of paper by D.R. Cox. *Journal of the Royal Statistical Society, Series B*, 34, 216-217.
- Carroll, R. J. and Ruppert, D. (1988). *Transformation and Weighting in Regression*. Chapman and Hall: London and New York.
- Cavanagh, C. and Sherman, R. P. (1998). Rank estimators for monotonic index models. *Journal* of *Econometrics*, 84, 351-381.
- Chen, S. (2002). Rank estimation of transformation models. *Econometrica*, 70, 1683-1697.
- Cox, D. R. (1972). Regression models and life tables. *Journal of the Royal Statistical Society, Series B*, 34, 187-220.
- Cox, D. R. (1975). Partial likelihood. *Biometrika*, 62, 269-276.
- de Leeuw, J., Hornik, K., and Patrick, M. (2009). Isotone optimization in R: Pool-adjacentviolators-algorithm (PAVA) and active set methods. *Journal of Statistical Software*, 32, 1-24.
- Doksum, K. A. (1987). An extension of partial likelihood methods for proportional hazard models to general transformation models. *Annals of Statistics*, 15, 325-345.
- Efron, B. (1979). Bootstrap methods: Another look at the jackknife. *The Annals of Statistics*, 7, 1–26.
- Folstein, M. F., Folstein, S. E., and McHugh, P. R. (1975). "Mini-mental state". A practical method for grading the cognitive state of patients for the clinician. *Journal of Psychiatric Research*, 12, 189-198.
- Groeneboom, P. and Hendrickx, K. (2018). Current status linear regression. *The Annals of Statistics*, 46, 1415-1444.
- Han, A. K. (1987). Non-parametric analysis of a generalized regression model: The maximum rank correlation estimator. *Journal of Econometrics*, 35, 303-316.
- Horowitz, J. L. (1996). Semiparametric estimation of a regression model with an unknown transformation of the dependent variable. *Econometrica*, 64, 103-137.
- Huang, J. and Wellner, J. A. (1993). Regression models with interval censoring. *Technical Report*, No. 261, University of Washington.
- Kalbfleisch, J. D. and Prentice, R. L. (1973). Marginal likelihoods based on Cox's regression and life model. *Biometrika*, 60, 267-278.
- Kosorok, M. R. (2008). Introduction to Empirical Processes and Semiparametric Inference. New York: Springer.
- Lehmann, E. L. (1953). The power of rank tests. Annals of Mathematical Statistics, 24, 23-43.

- Lin, H., and Peng, H. (2013). Smoothed rank correlation of the linear transformation regression model. *Computational Statistics & Data Analysis*, 57, 615-630.
- Ma, S., and Huang, J. (2005). Regularized ROC method for disease classification and biomarker selection with microarray data. *Bioinformatics*, 21, 4356-4362.
- Robertson, T., Wright, F. T., and Dykstra, R. L. (1988). *Order Restricted Statistical Inference*. Chichester, U.K.: John Wiley.
- Shao, J. and Tu, D. (1995). The Jackknife and Bootstrap. New York: Springer.
- Song, X., Ma, S., Huang, J., and Zhou, X. H. (2007). A semiparametric approach for the nonparametric transformation survival model with multiple covariates. *Biostatistics*, 8, 197-211.
- Zeng, D., and Lin, D. Y. (2007). Maximum likelihood estimation in semiparametric regression models with censored data (with discussion). *Journal of the Royal Statistical Society, Series B*, 69, 507-564.
- Zhang, J. (2013). Estimation and testing methods for monotone transformation models. PhD thesis, Columbia University.
- Zhang, J., Jin, Z., Shao, Y., and Ying, Z. (2018). Statistical inference on transformation models: A self-induced smoothing approach. *Journal of Nonparametric Statistics*, 30, 308-331.

Supplementary Material for "Maximum pairwise-rank-likelihood-based inference for the semiparametric transformation model"

This supplementary document contains technical details for the theoretical results in Section 4 of the main article (Sections 1–7), some details for the numerical algorithms of our $\hat{\beta}$ and $\hat{\beta}$ estimates (Section 8), an extension of our methods to data with ties in the responses (Section 9), and an extension of our methods to right-censored data (Section 10).

1 Review of Technical Conditions and Theorems in the Main Article

In this section, we review the technical conditions in the Appendix and theorems in Section 4 of the main article.

1.1 Technical conditions

We have imposed the following technical conditions in the Appendix of the main article.

<u>Condition 0</u>: H(y) is a strictly increasing function on the support \mathcal{Y} of Y, and ϵ_i and X_i are independent.

<u>Condition 1</u>: $\beta \in \mathcal{B}$ and $F_X(x)$ is supported on \mathcal{X} , where both \mathcal{B} and \mathcal{X} are compact subspaces of \mathbb{R}^p .

<u>Condition 2</u>: For any $x_1, x_2 \in \mathbb{R}$, we have

$$\sup_{\boldsymbol{\beta}\in\boldsymbol{\mathcal{B}}} \left| F_{\boldsymbol{\beta}^T X}(x_1) - F_{\boldsymbol{\beta}^T X}(x_2) \right| \lesssim |x_1 - x_2|.$$

<u>Condition 3</u>: $\inf_{\boldsymbol{x_1}, \boldsymbol{x_2} \in \mathcal{X}, \boldsymbol{\beta} \in \mathcal{B}} F_0((\boldsymbol{x_1} - \boldsymbol{x_2})^T \boldsymbol{\beta}) > 0.$

<u>Condition 4</u>: If F(x) is a c.d.f. such that $F((\boldsymbol{x}_1 - \boldsymbol{x}_2)^T \boldsymbol{\beta}) = F_0((\boldsymbol{x}_1 - \boldsymbol{x}_2)^T \boldsymbol{\beta}_0)$ for all $\boldsymbol{x}_1, \boldsymbol{x}_2 \in \mathcal{X}$, then $\boldsymbol{\beta} = \boldsymbol{\beta}_0$ and $F(\cdot) = F_0(\cdot)$.

<u>Condition 5</u>: $F_0(u)$ is second-order differentiable with bounded second-order derivatives and $\inf_{\boldsymbol{v}\in\mathcal{X}-\mathcal{X},\boldsymbol{\beta}\in\mathcal{B}}F'_0(\boldsymbol{v}^T\boldsymbol{\beta}) > 0.$

<u>Condition 6</u>: There exists an $\eta_0 > 0$ such that

$$\inf_{\boldsymbol{\beta}\in\mathcal{O}_{\eta_0}}\lambda_2\Big(\left[\operatorname{var}\{(X_1-X_2)-E((X_1-X_2)|(X_1-X_2)^T\boldsymbol{\beta})\}\right]\Big)>0,$$

where $\mathcal{O}_{\eta_0} = \{ \boldsymbol{\beta} : \| \boldsymbol{\beta} - \boldsymbol{\beta}_0 \|_2 \leq \eta_0, \| \boldsymbol{\beta} \|_2 = 1 \}; \lambda_2(B)$ denotes the second smallest eigenvalue of matrix B.

Furthermore, we need the following additional conditions to establish the asymptotic distribution for $\tilde{\beta}$.

<u>Condition A1</u>: $F_{\beta}(t)$ is continuously differentiable in β and t, and

$$\inf_{t \in \{\boldsymbol{v}^{T}\boldsymbol{\beta}: \boldsymbol{v} \in \boldsymbol{\mathcal{X}} - \boldsymbol{\mathcal{X}}, \boldsymbol{\beta} \in \boldsymbol{\mathcal{B}}\}, \boldsymbol{\beta} \in \boldsymbol{\mathcal{B}}} \frac{\partial F_{\boldsymbol{\beta}}(t)}{\partial t} > 0$$

<u>Condition A2</u>: $f_{X_1-X_2|(X_1-X_2)^T} \beta_{=t}(v)$ is continuously differentiable in $\beta \in \beta$ and $t \in \{x^T \beta : x \in \mathcal{X}, \beta \in \beta\}$.

<u>Condition A3</u>: We have $\beta = \beta_0$ if

$$\operatorname{cov}\left\{\boldsymbol{\beta}_{0}^{T}(X_{1}-X_{2}), F_{0}(\boldsymbol{\beta}_{0}^{T}(X_{1}-X_{2})) \middle| \boldsymbol{\beta}^{T}(X_{1}-X_{2})\right\} = 0$$

almost surely.

1.2 Theorem in Section 4.1 of the main article

In Section 4.1 of the main article, we have presented the following theorem, which establish the asymptotic properties of $\hat{\beta}$ and $\hat{F}(\cdot)$. Let

$$D(\boldsymbol{\beta}_{2}, F_{2}; \boldsymbol{\beta}_{1}, F_{1})$$

$$= \left[\int \int \left\{ F_{1}((\boldsymbol{x}_{1} - \boldsymbol{x}_{2})^{T} \boldsymbol{\beta}_{1}) - F_{2}((\boldsymbol{x}_{1} - \boldsymbol{x}_{2})^{T} \boldsymbol{\beta}_{2}) \right\}^{2} dF_{X}(\boldsymbol{x}_{1}) dF_{X}(\boldsymbol{x}_{2}) \right]^{1/2}.$$
(S.1)

The following is Theorem 1 in the main article.

Theorem 1. Assume Conditions 0–6. We have

(a)
$$D(\hat{\beta}, \hat{F}; \beta_0, F_0) = O_p(n^{-1/3}),$$

(b) $\hat{\beta} - \beta_0 = O_p(n^{-1/3}),$

where β_0 and F_0 are the true values of β and F.

1.3 Theorems in Section 4.2 of the main article

In Section 4.2 of the main article, we have presented the following theorems, which establish the asymptotic distributions for $\hat{\beta}$, \hat{F}_{β} , and $\hat{F}_{\tilde{\beta}}$.

Recall that we have the following notation. For any $\beta \in \mathcal{B}$, define

$$F_{\boldsymbol{\beta}}(t) = E\left\{I(Y_1 > Y_2) \middle| (X_1 - X_2)^T \boldsymbol{\beta} = t\right\}$$
$$= \int F_0(t + \boldsymbol{v}^T(\boldsymbol{\beta}_0 - \boldsymbol{\beta})) f_{(X_1 - X_2)|(X_1 - X_2)^T} \boldsymbol{\beta}_{=t}(\boldsymbol{v}) d\boldsymbol{v}$$
(S.2)

and

$$\varphi_{\boldsymbol{\beta}}(t) = E(X_1 - X_2 | (X_1 - X_2)^T \boldsymbol{\beta} = t), \quad \varphi_0(t) = \varphi_{\boldsymbol{\beta}_0}(t).$$
(S.3)

 $\widetilde{\boldsymbol{\beta}}$ is defined to be the zero-crossing of $\psi_n(\boldsymbol{\beta})$, where

$$\psi_n(\boldsymbol{\beta}) = \frac{1}{n^2} \sum_{i \neq j} (X_i - X_j) \left\{ I(Y_i > Y_j) - \hat{F}_{\boldsymbol{\beta}} \left((X_i - X_j)^T \boldsymbol{\beta} \right) \right\}$$
$$= \int \int (\boldsymbol{x}_1 - \boldsymbol{x}_2) \left\{ I(y_1 > y_2) - \hat{F}_{\boldsymbol{\beta}} ((\boldsymbol{x}_1 - \boldsymbol{x}_2)^T \boldsymbol{\beta}) \right\} d\mathbb{F}_{X,Y}(\boldsymbol{x}_1, y_1) d\mathbb{F}_{X,Y}(\boldsymbol{x}_2, y_2),$$

and the zero-crossing of a function (or a mapping) is defined below.

Definition 1. For a function $\psi: \mathcal{X} \to \mathbb{R}$, x is called the zero-crossing of ψ if every open neighborhood of x contains x_1, x_2 such that $\psi(x_1)\psi(x_2) \leq 0$. For a mapping $\tilde{\psi}: \mathcal{X} \to \mathbb{R}^d$, x is called the zero-crossing of $\tilde{\psi}$ if x is the zero-crossing of each component of $\tilde{\psi}$.

The population version of $\psi_n(\cdot)$ is given by

$$\psi_{0}(\boldsymbol{\beta}) = \iint (\boldsymbol{x}_{1} - \boldsymbol{x}_{2}) \left\{ I(y_{1} > y_{2}) - F_{\boldsymbol{\beta}}((\boldsymbol{x}_{1} - \boldsymbol{x}_{2})^{T} \boldsymbol{\beta}) \right\}$$
(S.4)

$$\times dF_{X,Y}(\boldsymbol{x}_{1}, y_{1}) dF_{X,Y}(\boldsymbol{x}_{2}, y_{2}).$$

We have the following lemma for $\psi_0(\cdot)$; it is Lemma 1 in the main article. The proof is given in Section 5.1.

Lemma 1. Assume Conditions 6, A1, and A2. We have the following:

- (1) $\psi_0(\beta_0) = 0;$
- (2) $\psi'_0(\boldsymbol{\beta}_0) = \frac{\partial \psi_0(\boldsymbol{\beta})}{\partial \boldsymbol{\beta}} \Big|_{\boldsymbol{\beta} = \boldsymbol{\beta}_0}$ exists with rank p 1.
- (3) Since $\|\beta_0\|_2 = 1$, there exists an *i* such that the *i*th component of β_0 is nonzero. Define $A = \psi'_0(\beta_0) + \overline{A}$, where \overline{A} is a $p \times p$ matrix with *i*th row β_0^T and all other entries 0. Then *A* is of full rank, and

$$\psi_0(\boldsymbol{\beta}) = A \cdot (\boldsymbol{\beta} - \boldsymbol{\beta}_0) + o(\boldsymbol{\beta} - \boldsymbol{\beta}_0).$$

We have the following theorem, which is Theorem 2 in the main article; it establishes the asymptotic properties of β , and the proof is given in Sections 5.2–5.4.

Theorem 2. Assume Conditions 0–2, 6, and A1–A3. Denote by $\tilde{\beta}$ the zero-crossing of $\psi_n(\beta)$ (if it exists). We have the following:

(1) When $n \to \infty$, a zero-crossing $\widetilde{\beta}$ of $\psi_n(\beta)$ exists with probability tending to 1.

(2) $\widetilde{\boldsymbol{\beta}} \to \boldsymbol{\beta}_0$ in probability.

(3) Recalling the matrix A defined in Lemma 1 Part (3), we have

$$\widetilde{\boldsymbol{\beta}} - \boldsymbol{\beta}_{0} = \frac{A^{-1}}{n^{2}} \sum_{i \neq j} \left[\left\{ X_{i} - X_{j} - \varphi_{0} \left((X_{i} - X_{j})^{T} \boldsymbol{\beta}_{0} \right) \right\} \\ \times \left\{ I(Y_{i} > Y_{j}) - F_{0} \left((X_{i} - X_{j})^{T} \boldsymbol{\beta}_{0} \right) \right\} \right] \\ + o_{p} (\widetilde{\boldsymbol{\beta}} - \boldsymbol{\beta}_{0}) + o_{p} (n^{-1/2}).$$
(S.5)

To derive the asymptotic distribution for $\tilde{\beta}$, we need to work on a U-statistic with the kernel:

$$h(\boldsymbol{x}_1, y_1; \boldsymbol{x}_2, y_2) = (\boldsymbol{x}_1 - \boldsymbol{x}_2) \left\{ I(y_1 > y_2) - F_0 \left((\boldsymbol{x}_1 - \boldsymbol{x}_2)^T \boldsymbol{\beta}_0 \right) \right\}.$$

Denote

$$\hbar(\boldsymbol{x}_1, y_1) = E\{h(\boldsymbol{x}_1, y_1; X_2, Y_2)\}.$$
(S.6)

The theorem above leads to the asymptotic distribution of β .

Corollary 1. Assume the conditions of Theorem 2. Furthermore, assume that both $\Sigma_X = var(X)$ and $\Sigma_{\hbar} = var \{\hbar(X, Y)\}$ are of full rank, and

$$\boldsymbol{\alpha}^{T} E\left(X_{2} - X_{1} | (X_{2} - X_{1})^{T} \boldsymbol{\beta}_{0}\right) = 0,$$

for any $\boldsymbol{\alpha} \in \mathbb{R}^{p}$, and $\boldsymbol{\alpha}^{T} \Sigma_{X} \boldsymbol{\beta}_{0} = 0.$ (S.7)

Denote $\widetilde{A} = \psi'_0(\beta_0)\Sigma_X^{-1} + \overline{A}$, where \overline{A} is defined in Lemma 1. We have

$$\sqrt{n}(\widetilde{\boldsymbol{\beta}} - \boldsymbol{\beta}_0) \to 2A^{-1}\widetilde{A}^{-1}\psi_0'(\boldsymbol{\beta}_0)\Sigma_X^{-1}N(\mathbf{0}, \Sigma_\hbar)$$
(S.8)

in distribution, as $n \rightarrow \infty$ *.*

Theorem 3 below is the Theorem 3 in the main article; it establishes the asymptotic distribution of $\widetilde{F}(\cdot) = \widehat{F}_{\alpha}(\cdot)$; the proof is given in Section 6.

Theorem 3. Assume that all the conditions of Corollary 1 are satisfied, and also assume the following conditions:

<u>Condition F1</u>: Recall that $F_{\epsilon}(s)$ is the c.d.f. for ϵ_i . Assume that it is continuous for $s \in \mathbb{R}$ and is continuously differentiable and strictly monotone for s in its support.

<u>Condition F2</u>: Denote by $F_{X^T\beta}(s)$ the c.d.f. for $X^T\beta$. Assume that it is continuously differentiable for s in its support. Let $f_{X^T\beta}(s) \equiv \frac{\partial^F_{X^T\beta}(s)}{\partial s}$; assume that for any s, it is continuous for β in the neighborhood of β_0 ; and $f_{X^T\beta_0}(s)$ is continuous for s in its support.

Then, we have for every t,

$$n^{1/2}\left\{\widehat{F}_{\widetilde{\boldsymbol{\beta}}}(t) - F_0(t)\right\} \rightarrow N\left(0, \sigma^2(t)/g_0^2(t)\right),$$

in distribution, where $g_0(t)$ denotes the p.d.f. for $(X_2 - X_1)^T \beta_0$, $\sigma^2(t) = var \{\zeta_t(X, Y)\}$,

$$\begin{aligned} \zeta_t(\boldsymbol{x}, y) &= \left\{ 1 - F_{\epsilon}(H_0(y) - \boldsymbol{x}^T \boldsymbol{\beta}_0 - t) - F_0(t) \right\} f_{X^T \boldsymbol{\beta}_0}(\boldsymbol{x}^T \boldsymbol{\beta}_0 + t) \\ &+ \left\{ F_{\epsilon}(H_0(y) - \boldsymbol{x}^T \boldsymbol{\beta}_0 + t) - F_0(t) \right\} f_{X^T \boldsymbol{\beta}_0}(\boldsymbol{x}^T \boldsymbol{\beta}_0 - t) \\ &+ 2g_0(t) \dot{F}_0^T(t) A^{-1} \widetilde{A}^{-1} \psi_0'(\boldsymbol{\beta}_0) \Sigma_X^{-1} \hbar(\boldsymbol{x}, y), \end{aligned}$$

with $\dot{F}_0(t) = \partial F_{\beta}(t) / \partial \beta |_{\beta=\beta_0}$ and $\hbar(\cdot, \cdot)$ defined by (S.6).

Remark 1. Based on the definition

$$F_0(s) = P(\epsilon_1 - \epsilon_2 \leqslant s) = E\{F_\epsilon(\epsilon_2 + s)\},\$$

the dominant convergence theorem, and Condition F1, we can verify that $F'_0(s) \equiv \partial F_0(s)/\partial s$ exists and continuous, and $F'_0(t) > 0$ for any t in the support of $F_0(\cdot)$. Similarly, based on the definition

$$\begin{aligned} G_{\boldsymbol{\beta}}(s) &= P((X_1 - X_2)^T \boldsymbol{\beta} < s) \\ &= E\left\{F_{X^T \boldsymbol{\beta}}(X_2^T \boldsymbol{\beta} + s)\right\}, \end{aligned}$$

Condition F2 implies that $G'_{\beta}(s) \equiv \partial G_{\beta}(s)/\partial s$ exists and continuous for (s,β) in a neighbourhood of (t,β_0) , and $G'_0(t) > 0$. We denote $g_{\beta}(s) = G'_{\beta}(s)$ and $g_0(s) = g_{\beta_0}(s)$.

With developments similar to but simpler than those of Theorem 3, we are able to establish the asymptotic distribution for $\hat{F}_{\beta}(t)$ for every given $\beta \in \mathcal{B}$ and $t \in \mathbb{R}$; we summarise this result in the following corollary; a sketched proof is given in Section 7.

Corollary 2. Assume Conditions 0–2 and A1, and Conditions F1' and F2' given below.

<u>Condition F1'</u>: $F_{\epsilon,\beta}(y,s)$ defined by (S.10) is continuous for $s \in \mathbb{R}$ and is continuously differentiable for s in the support and $\beta \in \mathcal{B}$.

<u>Condition F2'</u>: Denote by $F_{X^T\beta}(s)$ the c.d.f. for $X^T\beta$. Assume that it is continuously differentiable for s in the support. Recall $f_{X^T\beta}(s)$ defined by Condition F2 in Theorem 3; assume that it is continuous for s in the support and $\beta \in \beta$.

For every $t \in \mathbb{R}$ *and* $\beta \in \mathcal{B}$ *, we have*

$$n^{1/2}\left\{\widehat{F}_{\boldsymbol{\beta}}(t) - F_{\boldsymbol{\beta}}(t)\right\} \to N\left(0, \sigma_{\boldsymbol{\beta}}^2(t)/g_{\boldsymbol{\beta}}^2(t)\right),$$

in distribution, where $g_{\beta}(\cdot)$ denotes the p.d.f. of $(X_2 - X_1)^T \beta$, and

$$\sigma_{\boldsymbol{\beta}}(t) = E \Big[var \Big\{ f_{X^T \boldsymbol{\beta}}(X_1^T \boldsymbol{\beta} + t) F_{\epsilon, \boldsymbol{\beta}}(Y_1, X_1^T \boldsymbol{\beta} + t) - f_{X^T \boldsymbol{\beta}}(X_1^T \boldsymbol{\beta} - t) F_{\epsilon, \boldsymbol{\beta}}(Y_1, X_1^T \boldsymbol{\beta} - t) \Big| X_1 \Big\} \Big]$$

$$F_{\epsilon, \boldsymbol{\beta}}(y, s) = E \Big\{ F_{\epsilon}(H(y) - X^T \boldsymbol{\beta}_0) \Big| X^T \boldsymbol{\beta} = s \Big\}.$$
(S.9)
(S.10)

2 Some Discussion and Special Cases for Conditions 2, 4, and 6

In this section, we discuss some special cases under which Conditions 2, 4, and 6 are satisfied.

2.1 Special cases for Condition 2

Recall that Condition 2 requires that for any $x_1, x_2 \in \mathbb{R}$, we have

$$\sup_{\boldsymbol{\beta}\in\mathcal{B}} \left| F_{\boldsymbol{\beta}^T X}(x_1) - F_{\boldsymbol{\beta}^T X}(x_2) \right| \lesssim |x_1 - x_2|.$$

Clearly, a sufficient condition for this is the following Condition 2':

<u>Condition 2'</u>: The p.d.f. of $\beta^T X$, denoted by $f_{\beta^T X}(x)$, exists and

$$\sup_{\boldsymbol{\beta}\in\mathcal{B},x\in\mathbb{R}}f_{\boldsymbol{\beta}^{T}X}(x)\lesssim1$$

Moreover, Condition 2' is easily satisfied for many popular distributions of X; for example,

• if $X \sim N(\mu, \Sigma_X)$ with $\lambda_{\min}(\Sigma_X) > 0$, then Condition 2' is satisfied, since $\beta^T X \sim N(\beta^T \mu, \beta^T \Sigma_X \beta)$, and the variance

$$\inf_{\boldsymbol{\beta}\in\boldsymbol{\mathcal{B}}}\boldsymbol{\beta}^{T}\Sigma_{X}\boldsymbol{\beta} \geq \lambda_{\min}(\Sigma_{X})\inf_{\boldsymbol{\beta}\in\boldsymbol{\mathcal{B}}}\|\boldsymbol{\beta}\|_{2}^{2} = \lambda_{\min}(\Sigma_{X}) > 0,$$

where we have used the fact that $\|\beta\|_2 = 1$;

- if $X \sim \text{Uniform}(\Omega)$, where Ω is a bounded subset of \mathbb{R}^p , then Condition 2' is satisfied;
- more generally, it is satisfied if the distribution of X belongs to a location-scale family, whose density has the structure f_X(x) = det(Σ_X^{-1/2})f(Σ_X^{-1/2}(x μ_X)), where det(·) denotes the determinant of a matrix, Σ_X and μ_X are unknown parameters satisfying λ_{min}(Σ_X) > 0, and f(·) is some given density function such that sup_{x∈ℝ} f(x) ≤ 1.

2.2 Special cases for Condition 4

Condition 4 is to ensure that $F(\cdot)$ and β are identifiable. The following lemma gives a sufficient, but more intuitive condition under which Condition 4 is satisfied. We need the following Condition 4':

• <u>Condition 4'</u>: There exists an $\tilde{x} \in \mathcal{X}$ and $\eta > 0$ such that $\{x : \|x - \tilde{x}\|_2 \leq \eta\} \subset \mathcal{X}$.

Remark 2. Condition 4' is satisfied if we assume that $\lambda_{\min} \{ var(X) \} > 0$ and $f_X(x)$, the p.d.f. of X, is continuous.

Lemma 2. Assume $\|\beta\|_2 = 1$, $F_0(\cdot)$ is strictly increasing, and Condition 4' given above. Then Condition 4 is satisfied.

Proof: We need to show only that if $F(v^T \beta) = F_0(v^T \beta_0)$ for all $v \in \mathcal{X} - \mathcal{X}$, then $\beta = \beta_0$. We prove this by contradiction. Suppose otherwise, then there exists a $\beta_1 \neq \beta_0$ such that $F(v^T \beta_1) = F_0(v^T \beta_0)$ for all $v \in \mathcal{X} - \mathcal{X}$. Since $F \in \mathcal{F}$ is nondecreasing and $F_0(\cdot)$ is strictly increasing, we must have for any $v_1, v_2 \in \mathcal{X} - \mathcal{X}$,

• if $v_1^T \beta_0 > v_2^T \beta_0$, then $v_1^T \beta_1 \ge v_2^T \beta_1$; otherwise, since $F(\cdot)$ is nondecreasing,

$$F_0(\boldsymbol{v}_1^T\boldsymbol{\beta}_0) = F(\boldsymbol{v}_1^T\boldsymbol{\beta}_1) \leqslant F(\boldsymbol{v}_2^T\boldsymbol{\beta}_1) = F_0(\boldsymbol{v}_2^T\boldsymbol{\beta}_0)$$

which contradicts $\boldsymbol{v}_1^T \boldsymbol{\beta}_0 > \boldsymbol{v}_2^T \boldsymbol{\beta}_0$ and $F_0(\cdot)$ is strictly increasing;

• likewise if $v_1^T \beta_0 < v_2^T \beta_0$, then $v_1^T \beta_1 \leq v_2^T \beta_1$.

In summary, we must have

$$(\boldsymbol{v}_1 - \boldsymbol{v}_2)^T \boldsymbol{\beta}_1 \boldsymbol{\beta}_0^T (\boldsymbol{v}_1 - \boldsymbol{v}_2) \ge 0.$$
(S.11)

On the other hand, since $\beta_1 \neq \beta_0$ and $\|\beta_1\|_2 = \|\beta_0\|_2 = 1$, we have $|\beta_1^T \beta_0| < 1$. Because of Condition 4', we can verify that there exists an $\tilde{\eta} > 0$ such that $0 \neq \tilde{v} \in \{v_1 - v_2 : v_1 \in \mathcal{X} - \mathcal{X}, v_2 \in \mathcal{X} - \mathcal{X}\}$, where $\tilde{v} = \tilde{\eta} (\beta_1 - \beta_0)$. As a consequence, noting that $\beta_1^T \beta_1 = \beta_0^T \beta_0 = 1$, we have

$$\widetilde{\boldsymbol{v}}^{T}\boldsymbol{\beta}_{1}\boldsymbol{\beta}_{0}^{T}\widetilde{\boldsymbol{v}}=-\widetilde{\eta}^{2}\left(1-\boldsymbol{\beta}_{0}^{T}\boldsymbol{\beta}_{1}\right)^{2}<0,$$

which contradicts (S.11). This completes the proof of this lemma.

2.3 Special cases for Condition 6

In this section, we consider some stronger but more intuitive conditions than Condition 6.

Let $\beta \in \mathcal{O}_{\eta_0}$ and $\Sigma(\beta) = \operatorname{var} (V - E(V|V^T\beta))$. Then $\Sigma(\beta)$ is positive semidefinite; there exists an orthonormal matrix $P(\beta)$ and a diagonal matrix $\Lambda(\beta) = \operatorname{diag}(\lambda_1(\beta), \ldots, \lambda_p(\beta))$ with $0 \leq \lambda_1(\beta) \leq \ldots \leq \lambda_p(\beta)$ such that $\Sigma(\beta) = P(\beta)\Lambda(\beta)P(\beta)^T$. Furthermore, since

$$\boldsymbol{\beta}^{T} \Sigma(\boldsymbol{\beta}) \boldsymbol{\beta} = \boldsymbol{\beta}^{T} \operatorname{var} \left(V - E(V|V^{T}\boldsymbol{\beta}) \right) \boldsymbol{\beta}$$

= $\operatorname{var} \left(\boldsymbol{\beta}^{T} V - E(\boldsymbol{\beta}^{T} V|V^{T}\boldsymbol{\beta}) \right) = \operatorname{var}(0) = 0,$ (S.12)

meaning that $\Sigma(\beta)$ is singular, we have $\lambda_1(\beta) = 0$. As a consequence,

$$\Lambda(\boldsymbol{\beta}) = \operatorname{diag}(0, \lambda_2(\boldsymbol{\beta}), \dots, \lambda_p(\boldsymbol{\beta})),$$

and Condition 6 is equivalent to

$$\inf_{\boldsymbol{\beta}\in\mathcal{O}_{\eta_0}}\lambda_2(\boldsymbol{\beta})>0.$$

We shall show that the following Condition 6' is a sufficient condition for Condition 6.

<u>Condition 6'</u>: There exists an $\eta_1 > 0$ such that all entries of $\Sigma(\beta)$ are continuous functions for $\beta \in \mathcal{O}_{\eta_1}$; and for any vector $\mathbf{0} \neq \boldsymbol{\alpha} \in \mathbb{R}^p$ such that $\boldsymbol{\alpha}^T \boldsymbol{\beta}_0 = 0$, we have

$$\operatorname{var}\left\{\boldsymbol{\alpha}^{T}V - E(\boldsymbol{\alpha}^{T}V|\boldsymbol{\beta}_{0}^{T}V)\right\} > 0.$$

Lemma 3. Condition 6 holds if Condition 6' holds.

Proof: We need the following notational convention for matrices: for any matrix B, B_i denotes the *i*th row, and $B_{i:j}$ denotes the *i*th to *j*th rows of B; $B_{i,j}$ denotes the (i, j)th entry of B.

We first show that $\lambda_2(\boldsymbol{\beta}_0) > 0$. In fact, there exists an orthonormal matrix $U(\boldsymbol{\beta}_0)$ whose first row is given by $U_1(\boldsymbol{\beta}_0) = \boldsymbol{\beta}_0^T / \|\boldsymbol{\beta}_0\|_2$, and since $U_{2:p}(\boldsymbol{\beta}_0) \Sigma(\boldsymbol{\beta}_0) U_{2:p}(\boldsymbol{\beta}_0)$ is positive semidefinite, there exists a $(p-1) \times (p-1)$ orthonormal matrix $R(\boldsymbol{\beta}_0)$ such that

$$R^{T}(\boldsymbol{\beta}_{0})U_{2:p}(\boldsymbol{\beta}_{0})\Sigma(\boldsymbol{\beta}_{0})U_{2:p}^{T}(\boldsymbol{\beta}_{0})R(\boldsymbol{\beta}_{0}) = \operatorname{diag}(\tau_{2}(\boldsymbol{\beta}_{0}),\ldots,\tau_{p}(\boldsymbol{\beta}_{0})),$$

where $0 \leq \tau_2(\boldsymbol{\beta}_0) \leq \ldots \leq \tau_p(\boldsymbol{\beta}_0)$. It is straightforward to verify that $\widetilde{R}(\boldsymbol{\beta}_0) = (U_1^T(\boldsymbol{\beta}_0), U_{2:p}^T(\boldsymbol{\beta}_0)R(\boldsymbol{\beta}_0))^T$ is a $p \times p$ orthonormal matrix, and

$$\widetilde{R}(\beta_{0})\Sigma(\beta_{0})\widetilde{R}^{T}(\beta_{0})$$

$$= \widetilde{R}(\beta_{0})\operatorname{var}\left(V - E(V|V^{T}\beta_{0})\right)\widetilde{R}^{T}(\beta_{0})(\beta_{0})$$

$$= \operatorname{var}\left(\begin{pmatrix}\beta_{0}^{T}/\|\beta_{0}\|_{2}\\R^{T}(\beta_{0})U_{2:p}(\beta_{0})\end{pmatrix}V - E\left(\begin{pmatrix}\beta_{0}^{T}/\|\beta_{0}\|_{2}\\R^{T}(\beta_{0})U_{2:p}(\beta_{0})\end{pmatrix}V^{T}\beta_{0}\end{pmatrix}\right)$$

$$= \operatorname{var}\left(\begin{pmatrix}0\\R^{T}(\beta_{0})U_{2:p}(\beta_{0})(V - E(V|V^{T}\beta_{0}))\end{pmatrix}\right)$$

$$= \begin{pmatrix}0\\0\\R^{T}(\beta_{0})U_{2:p}(\beta_{0})\Sigma(\beta_{0})U_{2:p}^{T}(\beta_{0})R(\beta_{0})\end{pmatrix}$$

$$= \begin{pmatrix}0\\0\\\mathrm{diag}(\tau_{2}(\beta_{0}), \dots, \tau_{p}(\beta_{0}))\end{pmatrix},$$
(S.13)

indicating that $0 = \tau_1(\boldsymbol{\beta}_0) \leq \tau_2(\boldsymbol{\beta}_0) \leq \ldots \leq \tau_p(\boldsymbol{\beta}_0)$ are the eigenvalues of $\Sigma(\boldsymbol{\beta}_0)$. Based on the uniqueness of eigenvalues, $\lambda_i(\boldsymbol{\beta}_0) = \tau_i(\boldsymbol{\beta}_0)$ for $i = 2, \ldots, p$. If $\lambda_2(\boldsymbol{\beta}_0) = 0$, then $\tau_2(\boldsymbol{\beta}_0) = \lambda_2(\boldsymbol{\beta}_0) = 0$. Consider $\widetilde{R}_2(\boldsymbol{\beta}_0)$, the second row of the orthonormal matrix $\widetilde{R}(\boldsymbol{\beta}_0)$; we have

- (i) $\|\widetilde{R}_2^T(\boldsymbol{\beta}_0)\|_2 = 1 \neq 0$, since $\widetilde{R}(\boldsymbol{\beta}_0)$ is orthonormal;
- (ii) $\widetilde{R}_2(\boldsymbol{\beta}_0)\boldsymbol{\beta}_0 = 0$, since $\boldsymbol{\beta}_0^T/\|\boldsymbol{\beta}_0\|_2$ is the first row of $\widetilde{R}(\boldsymbol{\beta}_0)$;
- (iii) based on (S.13),

$$\operatorname{var}\left(\widetilde{R}_{2}(\boldsymbol{\beta}_{0})V - E(\widetilde{R}_{2}(\boldsymbol{\beta}_{0})V|\boldsymbol{\beta}_{0}^{T}V)\right) = \widetilde{R}_{2}(\boldsymbol{\beta}_{0})\Sigma(\boldsymbol{\beta}_{0})\widetilde{R}_{2}^{T}(\boldsymbol{\beta}_{0}) = \tau_{2}(\boldsymbol{\beta}_{0}) = 0.$$

We observe that (i)–(iii) introduce a contradiction to Condition 6'. Therefore, we must have $\lambda_2(\beta_0) > 0$. Furthermore, as assumed by Condition 6', we must have that $\Sigma(\beta)$ is continuous for $\beta \in \mathcal{O}_{\eta_1}$; and its eigenvalues are the solutions of the polynomial det $(\lambda I_p - \Sigma(\beta))$. Therefore, $\lambda_2(\beta)$ is a continuous function for $\beta \in \mathcal{O}_{\eta_1}$; see Zedek (1965). As a consequence, there exists $0 < \eta_0 \leq \eta_1$ such that $\inf_{\beta \in \mathcal{O}_{\eta_0}} \lambda_2(\beta) > 0$. Similarly, $\lambda_{\min}(\Sigma_{-1,-1}(\beta))$ is a continuous function for $\beta \in \mathcal{O}_{\eta_0}$ such that

$$\lambda_{\min}(\Sigma_{-1,-1}(\boldsymbol{\beta}')) = \inf_{\boldsymbol{\beta} \in \mathcal{O}_{\eta_0}} \lambda_{\min}(\Sigma_{-1,-1}(\boldsymbol{\beta})).$$

Clearly,

$$\lambda_2(\boldsymbol{\beta}') \ge \inf_{\boldsymbol{\beta} \in \mathcal{O}_{\eta_0}} \lambda_2(\boldsymbol{\beta}) > 0.$$

We complete the proof of this lemma.

Finally, we give one special example under which Condition 6' is satisfied.

Example 1. Assume X_i , i = 1, ..., n, are *i.i.d.* and follow the multivariate normal distribution with a variance matrix Σ_X that is of full rank. Then Condition 6' is satisfied.

Proof: Clearly, $\Sigma_V = \operatorname{var}(X_i - X_j) = 2\Sigma_X$, which is strictly positive definite; thus, $\Sigma_V = P_V^T \Lambda_V P_V$ with P_V being an orthonormal matrix and $\Lambda_V = \operatorname{diag}(\nu_1, \ldots, \nu_p), 0 < \nu_1 \leq \nu_2 \leq \ldots \leq \nu_p$. For any $\beta \in \mathcal{O}_{\eta_1}$, let $U_V(\beta)$ be an orthonormal matrix whose first row is given by

$$U_{V,1} = \frac{\boldsymbol{\beta}^T P_V^T \Lambda_V^{1/2}}{\sqrt{\boldsymbol{\beta}^T \Sigma_V \boldsymbol{\beta}}}$$

Let $Q_V = U_V(\boldsymbol{\beta})\Lambda_V^{-1/2}P_V$. Clearly, the first row of Q_V is $Q_{V,1} = \frac{\boldsymbol{\beta}^T}{\sqrt{\boldsymbol{\beta}^T \Sigma_V \boldsymbol{\beta}}}$, and

$$\operatorname{var}(Q_V V) = Q_V \Sigma_V Q_V^T = U_V(\boldsymbol{\beta}) \Lambda_V^{-1/2} P_V(P_V^T \Lambda_V P_V) P_V^T \Lambda_V^{-1/2} U_V^T(\boldsymbol{\beta}) = I_p,$$

which together with the normality assumption indicates that $Q_{V,2:p}V$ is independent of $\beta^T V$. Furthermore, note that $Q_V^{-1} = P_V^T \Lambda_V^{1/2} U_V^T(\beta)$. Therefore,

$$E(V|\boldsymbol{\beta}^{T}V) = Q_{V}^{-1}E(Q_{V}V|\boldsymbol{\beta}^{T}V) = P_{V}^{T}\Lambda_{V}^{1/2}U_{V}^{T}(\boldsymbol{\beta})\begin{pmatrix}\boldsymbol{\beta}^{T}V/\sqrt{\boldsymbol{\beta}^{T}\Sigma_{V}\boldsymbol{\beta}}\\\mathbf{0}\end{pmatrix}$$
$$= P_{V}^{T}\Lambda_{V}P_{V}\boldsymbol{\beta}\boldsymbol{\beta}^{T}V/(\boldsymbol{\beta}^{T}\Sigma_{V}\boldsymbol{\beta}) = \Sigma_{V}\boldsymbol{\beta}\boldsymbol{\beta}^{T}V/(\boldsymbol{\beta}^{T}\Sigma_{V}\boldsymbol{\beta}),$$

and thus

$$\Sigma(\boldsymbol{\beta}) = \operatorname{var}\left(V - E(V|V^{T}\boldsymbol{\beta})\right)$$
$$= \left(I_{p} - \frac{\Sigma_{V}\boldsymbol{\beta}\boldsymbol{\beta}^{T}}{\boldsymbol{\beta}^{T}\Sigma_{V}\boldsymbol{\beta}}\right)\Sigma_{V}\left(I_{p} - \frac{\boldsymbol{\beta}\boldsymbol{\beta}^{T}\Sigma_{V}}{\boldsymbol{\beta}^{T}\Sigma_{V}\boldsymbol{\beta}}\right),$$
(S.14)

which is clearly a continuous function of β .

It remains to show that for any vector $\mathbf{0} \neq \boldsymbol{\alpha} \in \mathbb{R}^p$ that satisfies $\boldsymbol{\alpha}^T \boldsymbol{\beta}_0 = 0$, we must have

$$\operatorname{var}\left(\boldsymbol{\alpha}^{T}V - E(\boldsymbol{\alpha}^{T}V|\boldsymbol{\beta}_{0}^{T}V)\right) > 0.$$
(S.15)

In fact, noting (S.14), we have

$$\operatorname{var}\left(\boldsymbol{\alpha}^{T}V - E(\boldsymbol{\alpha}^{T}V|\boldsymbol{\beta}_{0}^{T}V)\right) = \boldsymbol{\alpha}^{T}\Sigma(\boldsymbol{\beta}_{0})\boldsymbol{\alpha}$$
$$= \boldsymbol{\alpha}^{T}\left(I_{p} - \frac{\Sigma_{V}\boldsymbol{\beta}_{0}\boldsymbol{\beta}_{0}^{T}}{\boldsymbol{\beta}_{0}^{T}\Sigma_{V}\boldsymbol{\beta}_{0}}\right)\Sigma_{V}\left(I_{p} - \frac{\boldsymbol{\beta}_{0}\boldsymbol{\beta}_{0}^{T}\Sigma_{V}}{\boldsymbol{\beta}_{0}^{T}\Sigma_{V}\boldsymbol{\beta}_{0}}\right)\boldsymbol{\alpha}.$$

Therefore, because Σ_V is of full rank, (S.15) is verified if

$$\boldsymbol{\alpha}^{T}\left(I_{p}-\frac{\Sigma_{V}\boldsymbol{\beta}_{0}\boldsymbol{\beta}_{0}^{T}}{\boldsymbol{\beta}_{0}^{T}\Sigma_{V}\boldsymbol{\beta}_{0}}\right)\neq\mathbf{0}.$$

Suppose otherwise; then

$$0 = \boldsymbol{\alpha}^{T} \left(I_{p} - \frac{\Sigma_{V} \boldsymbol{\beta}_{0} \boldsymbol{\beta}_{0}^{T}}{\boldsymbol{\beta}_{0}^{T} \Sigma_{V} \boldsymbol{\beta}_{0}} \right) \boldsymbol{\alpha}$$
$$= \|\boldsymbol{\alpha}\|_{2}^{2} - \boldsymbol{\alpha}^{T} \frac{\Sigma_{V} \boldsymbol{\beta}_{0} (\boldsymbol{\beta}_{0}^{T} \boldsymbol{\alpha})}{\boldsymbol{\beta}_{0}^{T} \Sigma_{V} \boldsymbol{\beta}_{0}} = \|\boldsymbol{\alpha}\|_{2}^{2},$$

which contradicts $\alpha \neq 0$. This completes the proof of this example.

3 Notations and Some Preliminary Results

We first introduce some notation used throughout the technical development. Let " \leq " (" \geq ") denote smaller (greater) than, up to a universal constant. If not otherwise stated, for any arbitrary random variable (vector) Z, denote by $F_Z(\cdot)$ the c.d.f. of Z; likewise for Z_1, \ldots, Z_n , denote by $\mathbb{F}_Z(\cdot)$ the empirical c.d.f. We use $\|\cdot\|_q$ to denote the l_q norm in Euclidean space; for any probability measure P, we use $\|f(\cdot)\|_{q,P}$ to denote the $L_q(P)$ norm of f. We need the following definitions of covering number, bracketing number, and entropy for a class of functions; these concepts play key roles in modern empirical process theory. These definitions are adapted from Definitions 2.1 and 2.2 in van de Geer (2000).

Definition 2. Let \Im be a class of functions. For any $\delta > 0$ and q > 0, let $N_q(\delta, \Im, P)$ be the smallest value of N for which there exists a collection of functions $\{g_1, \ldots, g_N\}$ such that for

any $g \in \mathfrak{S}$, there exists a $j = j(g) \in \{1, \ldots, N\}$, such that $||g - g_j||_{q,P} \leq \delta$. $N_q(\delta, \mathfrak{S}, P)$ is called the δ -covering number of \mathfrak{S} , and

$$H_q(\delta, \mathfrak{F}, P) = \log N_q(\delta, \mathfrak{F}, P)$$

is called the δ -entropy of \Im (for the $L_q(P)$ -metric).

Definition 3. Let \Im be a class of functions. For any $\delta > 0$ and q > 0, let $N_{q,B}(\delta, \Im, P)$ be the smallest value of N for which there exists a set of pairs of functions $\{(g_j^L, g_j^U)\}_{j=1}^N$ such that (i) $\|g_j^U - g_j^L\|_{q,P} \leq \delta$, and (ii) for any $g \in \Im$, there exists a j = j(g) such that

$$g_j^L \leqslant g \leqslant g_j^U$$

 $N_{a,B}(\delta, \Im, P)$ is called the δ -bracketing number of \Im , and

$$H_{q,B}(\delta,\mathfrak{F},P) = \log N_{q,B}(\delta,\mathfrak{F},P)$$

is called the δ -entropy with bracketing of \Im .

We reiterate the following notational convention for matrices: for any matrix B, B_i denotes the *i*th row, and $B_{i:j}$ denotes the *i*th to *j*th rows of B; $B_{i,j}$ denotes the (i, j)th entry of B; $B_{-i,-j}$ denotes the matrix by removing the *i*th row and *j*th column B.

We first introduce two lemmas. Lemma 4 is from Lemma 5.13 in van de Geer (2000).

Lemma 4. Let \Im be a class of functions and Z_1, \ldots, Z_n be an i.i.d. sample. Assume that

$$\sup_{g \in \Im} |g - g_0|_{\infty} \leq 1,$$
$$H_{2,B}(\delta, \Im, F_Z) \leq A\delta^{-\alpha},$$

for every $\delta > 0$ and some $0 < \alpha < 2$ and some constant A. Then, for some constants c and n_0 depending on α and A, we have for all $T \ge c$ and $n \ge n_0$,

$$P\left(\sup_{g\in\Im, \|g-g_0\|_{2,F_Z} \leqslant n^{-1/(2+\alpha)}} \left| \int (g-g_0) d(\mathbb{F}_Z - F_Z) \right| \ge T n^{-2/(2+\alpha)} \right)$$

$$\leqslant c \exp\left\{ -\frac{T n^{\alpha/(2+\alpha)}}{c^2} \right\},$$

$$P\left(\sup_{g\in\Im, \|g-g_0\|_{2,F_Z} > n^{-1/(2+\alpha)}} \frac{\sqrt{n} \left| \int (g-g_0) d(\mathbb{F}_Z - F_Z) \right|}{\|g-g_0\|_{2,F_Z}^{1-\alpha/2}} \ge T \right)$$

$$\leqslant c \exp\left(-\frac{T}{c^2}\right).$$

To facilitate our subsequent development, we define the following function classes and

derive their δ -entropies:

$$\mathcal{F} = \{F(\cdot) : F(\boldsymbol{x}) \in [0, 1] \text{ and is monotonically increasing} \}, \mathcal{F}_1 = \{F((\boldsymbol{x}_1 - \boldsymbol{x}_2)^T \boldsymbol{\beta}) : F \in \mathcal{F}, \boldsymbol{\beta} \in \boldsymbol{\mathcal{B}} \}, \\ \mathcal{F}_2 = \{F((\boldsymbol{x}_1 - \boldsymbol{x}_2)^T \boldsymbol{\beta}) : F \in \mathcal{F}, \boldsymbol{\beta} \in \boldsymbol{\mathcal{B}}, \boldsymbol{x}_2 \in \boldsymbol{\mathcal{X}} \}, \\ \mathcal{D} = \{d_{\boldsymbol{\beta}, F}(\boldsymbol{x}_2) = \left[\int \{F_0((\boldsymbol{x}_1 - \boldsymbol{x}_2)^T \boldsymbol{\beta}_0) - F((\boldsymbol{x}_1 - \boldsymbol{x}_2)^T \boldsymbol{\beta})\}^2 dF_X(\boldsymbol{x}_1)\right]^{1/2} : \\ F \in \mathcal{F}, \boldsymbol{\beta} \in \boldsymbol{\mathcal{B}} \}.$$

The following lemma establishes the δ -entropy of $\mathcal{F}_1, \mathcal{F}_2$, and \mathcal{D} .

Lemma 5. Assume Conditions 1 and 2. For any arbitrary $1 \le q < \infty$, we have

$$H_{q,B}(\delta, \mathcal{F}_1, F_{X_1, X_2}) \lesssim 1/\delta,$$
 (S.16)

$$H_{q,B}(\delta, \mathcal{F}_2, F_X) \lesssim 1/\delta,$$
 (S.17)

$$H_{q,B}(\delta, \mathcal{D}, F_X) \leq 1/\delta,$$
 (S.18)

where " \leq " is up to a universal constant depending only on q.

Proof. We first prove a preliminary result, which is helpful in the proof of Lemma 5.

Lemma 6. Let \Im be an arbitrary class of function such that $H_{q,B}(\delta, \Im, P) < \infty$. Then

$$H_{q,B}(2\delta, |\mathfrak{S}|, P) \leq H_{q,B}(\delta, \mathfrak{S}, P)$$

Proof. By the definition of $H_{q,B}(\delta, \Im, P)$, there exists a set of brackets $\{[l_i, u_i]\}_{i=1}^N$ that covers \Im , where $N = \exp\{H_{q,B}(\delta, \Im, P)\}$. Let |g| be an arbitrary function in $|\Im|$ and hence $g \in \Im$. Let $[l_i, u_i]$ be the bracket such that $l_i \leq g \leq u_i$. We immediately have $l_i^+ + u_i^- \leq g^+ + g^- \leq l_i^- + u_i^+$, that is $l_i^+ + u_i^- \leq |g| \leq l_i^- + u_i^+$. Furthermore,

$$|l_i^- + u_i^+ - l_i^+ - u_i^-| \le |l_i^- - u_i^-| + |u_i^+ - l_i^+| \le 2|u_i - l_i|,$$

and hence

$$||l_i^- + u_i^+ - l_i^+ - u_i^-||_{q,P} \le 2||u_i - l_i||_{q,P}.$$

This indicates, every δ -bracket under $L_q(P)$ in \Im leads to a 2δ -bracket under $L_q(P)$ in $|\Im|$. This completes our proof of this lemma.

We now move back to the proof of Lemma 5. First, we show (S.17); (S.16) can be obtained with very similar but simpler arguments. For any $\delta > 0$, let $\delta_1 = \delta^q$. Because of Condition 1, we have the following:

(i) There exist $\beta_1, \ldots, \beta_{N_1} \in \mathbb{R}^p$, such that $\mathcal{B} = \bigcup_{j_1=1}^{N_1} \{ \boldsymbol{\beta} : \| \boldsymbol{\beta} - \boldsymbol{\beta}_{j_1} \|_1 < \delta_1, \boldsymbol{\beta} \in \mathcal{B} \}$, where $N_1 \leq 1/\delta_1^p$.

(ii) There exist $\{[a_{j_2}, b_{j_2}] : a_{j_2} \in \mathbb{R}, b_{j_2} \in \mathbb{R}\}_{j_2=1}^{N_2}$ with $N_2 \leq 1/\delta_1^{pq}$ and $|b_{j_2} - a_{j_2}| < \delta_1$, such that for every $\boldsymbol{\beta} \in \boldsymbol{\mathcal{B}}$ and $\boldsymbol{x}_2 \in \boldsymbol{\mathcal{X}}$, we have $\boldsymbol{x}_2^T \boldsymbol{\beta} \in [a_{j_2}, b_{j_2}]$ for some $j_2 \in \{1, \dots, N_2\}$.

Next, we define a set of brackets $\{[l_j(\boldsymbol{x}_1), u_j(\boldsymbol{x}_1)] : j = 1..., N\}$ that covers $\{(\boldsymbol{x}_1 - \boldsymbol{x}_2)^T \boldsymbol{\beta} : \boldsymbol{\beta} \in \boldsymbol{\mathcal{B}}, \boldsymbol{x}_2 \in \boldsymbol{\mathcal{X}}\}$. Here $N = N_1 N_2 \leq 1/\delta_1^{p(1+q)}$. To this end, for $j_1 = 1..., N_1$ let

$$\boldsymbol{\beta}_{j_{1}+} = \boldsymbol{\beta}_{j_{1}} + (\delta_{1}, \dots, \delta_{1})^{T}; \qquad \boldsymbol{\beta}_{j_{1}-} = \boldsymbol{\beta}_{j_{1}} - (\delta_{1}, \dots, \delta_{1})^{T}.$$

Now, for every $j_1 = 1, \ldots, N_1$ and $j_2 = 1, \ldots, N_2$, we define

$$l_{j_1,j_2}(\boldsymbol{x}_1) = \boldsymbol{\beta}_{j_1-}^T \boldsymbol{x}_1^+ - \boldsymbol{\beta}_{j_1+}^T \boldsymbol{x}_1^- - b_{j_2}; \quad u_{j_1,j_2}(\boldsymbol{x}_1) = \boldsymbol{\beta}_{j_1+}^T \boldsymbol{x}_1^+ - \boldsymbol{\beta}_{j_1-}^T \boldsymbol{x}_1^- - a_{j_2},$$

where \boldsymbol{x}_1^+ and \boldsymbol{x}_1^- respectively denote the positive and negative parts of \boldsymbol{x}_1 . Based on (i) and (ii) above, we immediately observe that for every $\boldsymbol{\beta} \in \boldsymbol{\beta}$ and $\boldsymbol{x}_2 \in \boldsymbol{\mathcal{X}}$, there exist $j_1 \in \{1, \ldots, N_1\}$ and $j_2 \in \{1, \ldots, N_2\}$ such that $l_{j_1, j_2}(\boldsymbol{x}_1) \leq (\boldsymbol{x}_1 - \boldsymbol{x}_2)^T \boldsymbol{\beta} \leq u_{j_1, j_2}(\boldsymbol{x}_1)$. This indicates $\{[l_{j_1, j_2}(\boldsymbol{x}_1), u_{j_1, j_2}(\boldsymbol{x}_1)] : j_1 = 1 \ldots, N_1, j_2 = 1, \ldots, N_2\}$ covers $\{(\boldsymbol{x}_1 - \boldsymbol{x}_2)^T \boldsymbol{\beta} : \boldsymbol{\beta} \in \boldsymbol{\mathcal{B}}, \boldsymbol{x}_2 \in \boldsymbol{\mathcal{X}}\}$. For notational convenience, with a little abuse of notation, we denote this set to be $\{[l_j(\boldsymbol{x}_1), u_j(\boldsymbol{x}_1)] : j = 1 \ldots, N\}$, where $N = N_1 N_2 \leq 1/\delta_1^{p(q+1)}, l_j(\boldsymbol{x}_1) = \boldsymbol{\beta}_{j-}^T \boldsymbol{x}_1^+ - \boldsymbol{\beta}_{j+}^T \boldsymbol{x}_1^- - b_j$, and $u_j(\boldsymbol{x}_1) = \boldsymbol{\beta}_{j+}^T \boldsymbol{x}_1^+ - \boldsymbol{\beta}_{j-}^T \boldsymbol{x}_1^- - a_j$.

Next, we establish a set of brackets that cover \mathcal{F}_2 . Since \mathcal{F} is the subset of the class of monotonically increasing functions. Based on the well known results in the empirical process literature (see for example Theorem 9.24 in Kosorok 2008), there exists a set $\{[L_{i,j}(u), U_{i,j}(u)]\}_{i=1}^{M_j}$ of brackets that cover \mathcal{F} , $\log M_j \leq 1/\delta$, and for every $i = 1, \ldots, M_j$,

$$\|L_{i,j} - U_{i,j}\|_{q,F_j} \le \delta, \tag{S.19}$$

where F_j is the c.d.f. of $u_j(X)$. Without loss of generality, we assume that $L_{i,j}$ and $U_{i,j}$ are monotonically increasing functions, $L_{i,j}(u) \in [0, 1]$, and $U_{i,j}(u) \in [0, 1]$. We consider the set of brackets

$$\{[L_{i,j}(l_j(\boldsymbol{x}_1)), U_{i,j}(u_j(\boldsymbol{x}_1))], i = 1, \dots, M_j; \quad j = 1, \dots, N\},$$
(S.20)

which contains the number of brackets

$$\sum_{j=1}^{N} M_j; \quad \text{ and } \log \sum_{j=1}^{N} M_j \lesssim \log \left(N e^{1/\delta} \right) \lesssim 1/\delta.$$

Finally, (S.17) follows if we can verify (S.21) and (S.22) given below:

$$L_{i,j}(l_j(\boldsymbol{x}_1)) \leqslant F((\boldsymbol{x}_1 - \boldsymbol{x}_2)^T \boldsymbol{\beta}) \leqslant U_{i,j}(u_j(\boldsymbol{x}_2)),$$

for every $F((\boldsymbol{x}_1 - \boldsymbol{x}_2)^T \boldsymbol{\beta}) \in \mathcal{F}_2;$ (S.21)

and uniformly in $i \in \{1, \ldots, M\}$ and $j \in \{1, \ldots, N\}$,

$$\|U_{ij}(u_j) - L_{i,j}(l_j)\|_{q,F_X} \lesssim \delta.$$
(S.22)

We first verify (S.21). Based on the definition of $\{[l_j(\boldsymbol{x}_1), u_j(\boldsymbol{x}_2)] : j = 1..., N\}$, for every $\boldsymbol{\beta} \in \boldsymbol{\mathcal{B}}$ and $\boldsymbol{x}_2 \in \boldsymbol{\mathcal{X}}$, there exists $j \in \{1, ..., N\}$ such that $l_j(\boldsymbol{x}_1) \leq (\boldsymbol{x}_1 - \boldsymbol{x}_2)^T \boldsymbol{\beta} \leq u_j(\boldsymbol{x}_1)$ for all $\boldsymbol{x}_1 \in \boldsymbol{\mathcal{X}}$. For this j, based on (S.19), for every $F \in \boldsymbol{\mathcal{F}}$ there exists an $i \in \{1, ..., M\}$ such that $L_{i,j}(u) \leq F(u) \leq U_{i,j}(u)$. Therefore, we have

$$L_{i,j}(l_j(\boldsymbol{x}_1)) \leqslant F(l_j(\boldsymbol{x}_1)) \leqslant F((\boldsymbol{x}_1 - \boldsymbol{x}_2)^T \boldsymbol{\beta}) \leqslant F(u_j(\boldsymbol{x}_1)) \leqslant U_{i,j}(u_j(\boldsymbol{x}_1)),$$

which proves (S.21). We proceed to show (S.22). Consider

$$\begin{aligned} \|U_{ij}(u_j) - L_{i,j}(l_j)\|_{q,F_X} \\ \leqslant \|U_{ij}(u_j) - L_{i,j}(u_j)\|_{q,F_X} + \|L_{i,j}(u_j) - L_{i,j}(l_j)\|_{q,F_X} \\ = \|U_{i,j} - L_{i,j}\|_{q,F_j} + \|L_{i,j}(u_j) - L_{i,j}(l_j)\|_{q,F_X} \\ \leqslant \delta + \|L_{i,j}(u_j) - L_{i,j}(l_j)\|_{q,F_X}. \end{aligned}$$

$$(S.23)$$

Therefore, it is left to bound $||L_{i,j}(u_j) - L_{i,j}(l_j)||_{q,F_X}$. Since $L_{i,j}(u) \in [0, 1]$ is a monotonically increasing function, there exists a monotonically increasing function $S(u) = \sum_l c_l I(d_l < u)$ such that $0 \leq c_l \leq 1$, $\sum_l c_l \leq 1$, and

$$\sup_{u} |S(u) - L_{i,j}(u)| \lesssim \delta.$$

Now that

$$\begin{split} \|L_{i,j}(u_{j}) - L_{i,j}(l_{j})\|_{q,F_{X}} &\leq 2\delta + \|S(u_{j}) - S(l_{j})\|_{q,F_{X}} \\ &= 2\delta + \left[\int \left\{\sum_{l} c_{l} I(l_{j}(\boldsymbol{x}_{1}) \leq d_{l} < u_{j}(\boldsymbol{x}_{1}))\right\}^{q} dF_{X}(\boldsymbol{x}_{1})\right]^{1/q} \\ &\leq 2\delta + \sum_{l} c_{l} \left[\int I(l_{j}(\boldsymbol{x}_{1}) \leq d_{l} < u_{j}(\boldsymbol{x}_{1})) dF_{X}(\boldsymbol{x}_{1})\right]^{1/q} \\ &= 2\delta + \sum_{l} c_{l} \left[P(l_{j}(\boldsymbol{x}_{1}) \leq d_{l}) - P(u_{j}(\boldsymbol{x}_{1}) \leq d_{l})\right]^{1/q} \\ &= 2\delta + \sum_{l} c_{l} \left[P\left(\beta_{j}^{T}X \leq \delta_{1} \|X\|_{1} + b_{j} + d_{l}\right) - P\left(\beta_{j}^{T}X \leq -\delta_{1} \|X\|_{1} + a_{j} + d_{l}\right)\right]^{1/q} \\ &\leq 2\delta + \sum_{l} c_{l} \left[P\left(\beta_{j}^{T}X \leq \delta_{1} \sup_{\boldsymbol{x}_{1} \in \mathcal{X}} \|\boldsymbol{x}_{1}\|_{1} + b_{j} + d_{l}\right) \\ &- P\left(\beta_{j}^{T}X \leq -\delta_{1} \sup_{\boldsymbol{x}_{1} \in \mathcal{X}} \|\boldsymbol{x}_{1}\|_{1} + a_{j} + d_{l}\right)\right]^{1/q} \\ &= 2\delta + \sum_{l} c_{l} \left[F_{\beta_{j}^{T}X}(\delta_{1} \sup_{\boldsymbol{x}_{1} \in \mathcal{X}} \|\boldsymbol{x}_{1}\|_{1} + b_{j} + d_{l}) - F_{\beta_{j}^{T}X}(-\delta_{1} \sup_{\boldsymbol{x}_{1} \in \mathcal{X}} \|\boldsymbol{x}_{1}\|_{1} + a_{j} + d_{l})\right]^{1/q} \end{split}$$

By Condition 2, we further have

$$\|U_{ij}(u_j) - L_{i,j}(l_j)\|_{q,F_X} \lesssim 2\delta + \sum_l c_l \left(2\delta_1 \sup_{\boldsymbol{x}_1 \in \mathcal{X}} \|\boldsymbol{x}_1\|_1 + |a_j - b_j| \right)^{1/q}$$

$$\lesssim 2\delta + \delta_1^{1/q} \lesssim \delta. \tag{S.24}$$

Combining (S.24) with (S.23), (S.22) follows, and we finish the proof of (S.17).

We proceed to show (S.18). Based on (S.16) and Lemma 6, we immediately have

$$H_{q_1,B}(\delta, |F_0((\boldsymbol{x}_1 - \boldsymbol{x}_2)^T \boldsymbol{\beta}_0 - \mathcal{F}_1|, F_{X_1,X_2}) \lesssim 1/\delta,$$

for any arbitrary $1 \le q_1 < \infty$. Now for any arbitrary q, we set $q_1 = 2$ when $1 \le q \le 2$ and $q_1 = q$ when $1 \le q \le 2$ in above. Then, there exist a set of δ -brackets $\{[l_i(\boldsymbol{x}_1, \boldsymbol{x}_2), u_i(\boldsymbol{x}_1, \boldsymbol{x}_2)]\}_{i=1}^{N_3}$ to cover $|F_0((\boldsymbol{x}_1 - \boldsymbol{x})^T \boldsymbol{\beta}_0 - \mathcal{F}_1|$ under the $L_{q_1}(F_{X_1,X_2})$ norm, where $\log N_3 \le 1/\delta$. Without loss of generality, we assume $l_i(\boldsymbol{x}_1, \boldsymbol{x}_2) \ge 0$ and $u_i(\boldsymbol{x}_1, \boldsymbol{x}_2) \ge 0$. For $i = 1, \ldots, N_3$, define

$$L_{i}(\boldsymbol{x}_{2}) = \left[\int l_{i}^{2}(\boldsymbol{x}_{1}, \boldsymbol{x}_{2}) dF_{X}(\boldsymbol{x}_{1}) \right]^{1/2}$$

$$U_{i}(\boldsymbol{x}_{2}) = \left[\int u_{i}^{2}(\boldsymbol{x}_{1}, \boldsymbol{x}_{2}) dF_{X}(\boldsymbol{x}_{1}) \right]^{1/2}.$$
 (S.25)

Now for any $d_{\boldsymbol{\beta},F}(\boldsymbol{x}_2) = \left[\int \{F_0((\boldsymbol{x}_1 - \boldsymbol{x}_2)^T \boldsymbol{\beta}_0) - F((\boldsymbol{x}_1 - \boldsymbol{x}_2)^T \boldsymbol{\beta})\}^2 dF_X(\boldsymbol{x}_1)\right]^{1/2} \in \mathcal{D},$ there exists $1 \leq i \leq N_3$, such that

$$0 \leq l_i(\boldsymbol{x}_1, \boldsymbol{x}_2) \leq \left| F_0((\boldsymbol{x}_1 - \boldsymbol{x}_2)^T \boldsymbol{\beta}_0) - F((\boldsymbol{x}_1 - \boldsymbol{x}_2)^T \boldsymbol{\beta}) \right| \leq u_i(\boldsymbol{x}_1, \boldsymbol{x}_2),$$

which immediately implies

$$L_i(\boldsymbol{x}_2) \leq d_{\boldsymbol{\beta},F}(\boldsymbol{x}_2) \leq U_i(\boldsymbol{x}_2).$$

This indicates that the set of brackets $\{[L_i, U_i]\}_{i=1}^{N_3}$ with L_i, U_i defined by (S.25) covers \mathcal{D} . Furthermore, for every $i = 1, \ldots, N_3$, we have

$$\begin{array}{ll} 0 &\leqslant & U_{i}(\boldsymbol{x}_{2}) - L_{i}(\boldsymbol{x}_{2}) = \left[\int u_{i}^{2}(\boldsymbol{x}_{1}, \boldsymbol{x}_{2}) dF_{X}(\boldsymbol{x}_{1}) \right]^{1/2} - \left[\int l_{i}^{2}(\boldsymbol{x}_{1}, \boldsymbol{x}_{2}) dF_{X}(\boldsymbol{x}_{1}) \right]^{1/2} \\ &= & \frac{\int \{u_{i}(\boldsymbol{x}_{1}, \boldsymbol{x}_{2}) - l_{i}(\boldsymbol{x}_{1}, \boldsymbol{x}_{2})\} \{u_{i}(\boldsymbol{x}_{1}, \boldsymbol{x}_{2}) + l_{i}(\boldsymbol{x}_{1}, \boldsymbol{x}_{2})\} dF_{X}(\boldsymbol{x}_{1})}{\left[\int u_{i}^{2}(\boldsymbol{x}_{1}, \boldsymbol{x}_{2}) dF_{X}(\boldsymbol{x}_{1}) \right]^{1/2} + \left[\int l_{i}^{2}(\boldsymbol{x}_{1}, \boldsymbol{x}_{2}) dF_{X}(\boldsymbol{x}_{1}) \right]^{1/2}} \\ &\leqslant & \frac{\left[\int \{u_{i}(\boldsymbol{x}_{1}, \boldsymbol{x}_{2}) - l_{i}(\boldsymbol{x}_{1}, \boldsymbol{x}_{2}) \}^{2} dF_{X}(\boldsymbol{x}_{1}) \right]^{1/2} + \left[\int l_{i}^{2}(\boldsymbol{x}_{1}, \boldsymbol{x}_{2}) + l_{i}(\boldsymbol{x}_{1}, \boldsymbol{x}_{2}) \}^{2} dF_{X}(\boldsymbol{x}_{1}) \right]^{1/2}}{\left[\int u_{i}^{2}(\boldsymbol{x}_{1}, \boldsymbol{x}_{2}) dF_{X}(\boldsymbol{x}_{1}) \right]^{1/2}} \\ &\leqslant & \left[\int \{u_{i}(\boldsymbol{x}_{1}, \boldsymbol{x}_{2}) - l_{i}(\boldsymbol{x}_{1}, \boldsymbol{x}_{2}) \}^{2} dF_{X}(\boldsymbol{x}_{1}) \right]^{1/2}} \\ \end{cases}$$

where the last " \leq " follows from the fact that

$$\int \{u_i(\boldsymbol{x}_1, \boldsymbol{x}_2) + l_i(\boldsymbol{x}_1, \boldsymbol{x}_2)\}^2 dF_X(\boldsymbol{x}_1)$$

$$-\left[\left\{\int u_i^2(\boldsymbol{x}_1, \boldsymbol{x}_2)dF_X(\boldsymbol{x}_1)\right\}^{1/2} + \left\{\int l_i^2(\boldsymbol{x}_1, \boldsymbol{x}_2)dF_X(\boldsymbol{x}_1)\right\}^{1/2}\right]^2$$

= $2\int u_i(\boldsymbol{x}_1, \boldsymbol{x}_2)l_i(\boldsymbol{x}_1, \boldsymbol{x}_2)dF_X(\boldsymbol{x}_1)$
 $-2\left\{\int u_i^2(\boldsymbol{x}_1, \boldsymbol{x}_2)dF_X(\boldsymbol{x}_1)\int l_i^2(\boldsymbol{x}_1, \boldsymbol{x}_2)dF_X(\boldsymbol{x}_1)\right\}^{1/2} \leq 0.$

Therefore

$$\begin{split} \|U_{i} - L_{i}\|_{q,F_{X}} \\ \leqslant & \left[\int \left[\int \left\{ u_{i}(\boldsymbol{x}_{1}, \boldsymbol{x}_{2}) - l_{i}(\boldsymbol{x}_{1}, \boldsymbol{x}_{2}) \right\}^{2} dF_{X}(\boldsymbol{x}_{1}) \right]^{q/2} dF_{X}(\boldsymbol{x}_{2}) \right]^{1/q} \\ \leqslant & \left\{ \begin{bmatrix} \int \int \left\{ u_{i}(\boldsymbol{x}_{1}, \boldsymbol{x}_{2}) - l_{i}(\boldsymbol{x}_{1}, \boldsymbol{x}_{2}) \right\}^{q} dF_{X}(\boldsymbol{x}_{1}) dF_{X}(\boldsymbol{x}_{2}) \right]^{1/q} & \text{when } q > 2 \\ \begin{bmatrix} \int \int \left\{ u_{i}(\boldsymbol{x}_{1}, \boldsymbol{x}_{2}) - l_{i}(\boldsymbol{x}_{1}, \boldsymbol{x}_{2}) \right\}^{2} dF_{X}(\boldsymbol{x}_{1}) dF_{X}(\boldsymbol{x}_{2}) \end{bmatrix}^{1/2} & \text{when } 1 \leqslant q \leqslant 2 \\ = & \|u_{i} - l_{i}\|_{q_{1}, F_{X_{1}, X_{2}}} \leqslant \delta, \end{split} \end{split}$$

based on the fact that $[l_i(\boldsymbol{x}_1, \boldsymbol{x}_2), u_i(\boldsymbol{x}_1, \boldsymbol{x}_2)], i = 1, ..., N_3$ are δ -brackets that cover \mathcal{F}_1 . This completes our proof of (S.18). The proof of Lemma 5 is completed.

4 Proof of Theorem 1

We first show Part (a) of Theorem 1. It is structured as Lemmas 7 and 8 below.

Lemma 7 establishes a basic inequality that plays a key role in developing the asymptotic property for $D(\hat{\beta}, \hat{F}; \beta_0, F_0)$. We need the following notation:

$$\gamma_1(\boldsymbol{v}; \hat{F}, \hat{\boldsymbol{\beta}}) = 4 \left\{ \sqrt{\frac{\hat{F}(\boldsymbol{v}^T \hat{\boldsymbol{\beta}})}{F_0(\boldsymbol{v}^T \boldsymbol{\beta}_0)}} - 1 \right\},$$

$$\gamma_2(\boldsymbol{v}; \hat{F}, \hat{\boldsymbol{\beta}}) = 4 \left\{ \sqrt{\frac{1 - \hat{F}(\boldsymbol{v}^T \hat{\boldsymbol{\beta}})}{1 - F_0(\boldsymbol{v}^T \boldsymbol{\beta}_0)}} - 1 \right\}.$$

Lemma 7. Recall the definition of $D(\hat{\beta}, \hat{F}; \beta_0, F_0)$ in (S.1):

$$D(\hat{\boldsymbol{\beta}}, \hat{F}; \boldsymbol{\beta}_0, F_0) = \left[\int \int \left\{ F_0((\boldsymbol{x}_1 - \boldsymbol{x}_2)^T \boldsymbol{\beta}) - \hat{F}((\boldsymbol{x}_1 - \boldsymbol{x}_2)^T \hat{\boldsymbol{\beta}}) \right\}^2 dF_X(\boldsymbol{x}_1) dF_X(\boldsymbol{x}_2) \right]^{1/2}.$$

We have

$$D^{2}\left(\widehat{\boldsymbol{\beta}},\widehat{F};\boldsymbol{\beta}_{0},F_{0}\right) \leq 2\int\left\{I(u>0)\gamma_{1}(\boldsymbol{v};\widehat{F},\widehat{\boldsymbol{\beta}}) + I(u<0)\gamma_{2}(\boldsymbol{v};\widehat{F},\widehat{\boldsymbol{\beta}})\right\} \quad (S.26)$$

$$\times d \left\{ \mathbb{F}_{U,V}(u, \boldsymbol{v}) - F_{U,V}(u, \boldsymbol{v}) \right\},$$

where $F_{U,V}$ is the joint c.d.f. of $(U_{i,j}, V_{i,j})$ and $\mathbb{F}_{U,V}(u, v)$ is the empirical c.d.f. of $\{(U_{i,j}, V_{i,j})\}_{i,j=1}^{n}$, $U_{i,j} = Y_i - Y_j$, $V_{i,j} = X_i - X_j$.

Proof. Since $P(Y_i = Y_j) = 0$ for $i \neq j$ by Conditions 0 and 5, without loss of generality, we assume $Y_i \neq Y_j$ when $i \neq j$. Then, we can write

$$\ell(\boldsymbol{\beta}, F) = \sum_{i,j} [I(Y_i > Y_j) \log\{F((X_i - X_j)^T \boldsymbol{\beta})\} + I(Y_i < Y_j) \log\{1 - F((X_i - X_j)^T \boldsymbol{\beta})\}].$$

We have

$$\begin{array}{ll}
0 & \geq & \ell(\beta_{0}, F_{0}) - \ell(\widehat{\beta}, \widehat{F}) \\
& = & -\sum_{i,j} I(U_{i,j} > 0) \log \frac{\widehat{F}(V_{i,j}^{T} \widehat{\beta})}{F_{0}(V_{i,j}^{T} \beta_{0})} - \sum_{i,j} I(U_{i,j} < 0) \log \frac{1 - \widehat{F}(V_{i,j}^{T} \widehat{\beta})}{1 - F_{0}(V_{i,j}^{T} \beta_{0})} \\
& = & -n^{2} \int I(u > 0) \log \frac{\widehat{F}(\boldsymbol{v}^{T} \widehat{\beta})}{F_{0}(\boldsymbol{v}^{T} \beta_{0})} d\mathbb{F}_{U,V}(u, \boldsymbol{v}) \\
& & -n^{2} \int I(u < 0) \log \frac{1 - \widehat{F}(\boldsymbol{v}^{T} \widehat{\beta})}{1 - F_{0}(\boldsymbol{v}^{T} \beta_{0})} d\mathbb{F}_{U,V}(u, \boldsymbol{v}).
\end{array}$$
(S.27)

On the other hand, using the inequality $0.5 \log x \leq \sqrt{x} - 1$ for any x > 0, we have

$$-\log\frac{\widehat{F}(\boldsymbol{v}^{T}\widehat{\boldsymbol{\beta}})}{F_{0}(\boldsymbol{v}^{T}\boldsymbol{\beta}_{0})} \ge 2\left\{1-\sqrt{\frac{\widehat{F}(\boldsymbol{v}^{T}\widehat{\boldsymbol{\beta}})}{F_{0}(\boldsymbol{v}^{T}\boldsymbol{\beta}_{0})}}\right\} = -0.5\gamma_{1}(\boldsymbol{v};\widehat{F},\widehat{\boldsymbol{\beta}}), \quad (S.28)$$

and likewise

$$-\log\frac{1-\hat{F}(\boldsymbol{v}^{T}\hat{\boldsymbol{\beta}})}{1-F_{0}(\boldsymbol{v}^{T}\boldsymbol{\beta}_{0})} \ge 2\left\{1-\sqrt{\frac{1-\hat{F}(\boldsymbol{v}^{T}\hat{\boldsymbol{\beta}})}{1-F_{0}(\boldsymbol{v}^{T}\boldsymbol{\beta}_{0})}}\right\} = -0.5\gamma_{2}(\boldsymbol{v};\hat{F},\hat{\boldsymbol{\beta}}).$$
(S.29)

Combining (S.27)–(S.29), we have

$$0 \geq -\int \left\{ I(u>0)\gamma_{1}(\boldsymbol{v};\hat{F},\hat{\boldsymbol{\beta}}) + I(u<0)\gamma_{2}(\boldsymbol{v};\hat{F},\hat{\boldsymbol{\beta}}) \right\} d\mathbb{F}_{U,V}(u,\boldsymbol{v})$$

$$= -\int \left\{ I(u>0)\gamma_{1}(\boldsymbol{v};\hat{F},\hat{\boldsymbol{\beta}}) + I(u<0)\gamma_{2}(\boldsymbol{v};\hat{F},\hat{\boldsymbol{\beta}}) \right\} \times d\left\{ \mathbb{F}_{U,V}(u,\boldsymbol{v}) - F_{U,V}(u,\boldsymbol{v}) \right\}$$

$$-\int \left\{ I(u>0)\gamma_{1}(\boldsymbol{v};\hat{F},\hat{\boldsymbol{\beta}}) + I(u<0)\gamma_{2}(\boldsymbol{v};\hat{F},\hat{\boldsymbol{\beta}}) \right\} dF_{U,V}(u,\boldsymbol{v}).$$
(S.30)

Therefore, to show (S.26), we need to show only that

$$-\int \left\{ I(u>0)\gamma_1(\boldsymbol{v};\hat{F},\hat{\boldsymbol{\beta}}) + I(u<0)\gamma_2(\boldsymbol{v};\hat{F},\hat{\boldsymbol{\beta}}) \right\} dF_{U,V}(u,\boldsymbol{v})$$

$$\geq 0.5D^2(\hat{\boldsymbol{\beta}},\hat{F};\boldsymbol{\beta}_0,F_0).$$
(S.31)

To this end, note that

$$-\int \left\{ I(u>0)\gamma_{1}(\boldsymbol{v};\hat{F},\hat{\boldsymbol{\beta}}) + I(u<0)\gamma_{2}(\boldsymbol{v};\hat{F},\hat{\boldsymbol{\beta}}) \right\} dF_{U,V}(u,\boldsymbol{v})$$

$$= -\int \left\{ \gamma_{1}(\boldsymbol{v};\hat{F},\hat{\boldsymbol{\beta}}) \int I(u>0) dF_{U|\boldsymbol{v}}(u) + \gamma_{2}(\boldsymbol{v};\hat{F},\hat{\boldsymbol{\beta}}) \int I(u<0) dF_{U|\boldsymbol{v}}(u) \right\} dF_{V}(\boldsymbol{v}), \qquad (S.32)$$

where $F_{U|\boldsymbol{v}}(u)$ denotes the conditional c.d.f. of $U_{i,j}|V_{i,j} = \boldsymbol{v}$ and $F_V(\boldsymbol{v})$ is the marginal c.d.f. of $V_{i,j}$. Note that by Condition 0,

$$\int I(u > 0) dF_{U|\boldsymbol{v}}(u) = E \left\{ I(U_{i,j} > 0) \middle| V_{i,j} = \boldsymbol{v} \right\}$$

= $E \left\{ I(Y_i - Y_j > 0) \middle| X_i - X_j = \boldsymbol{v} \right\}$
= $P(Y_i > Y_j | X_i - X_j = \boldsymbol{v}) = F_0(\boldsymbol{v}^T \boldsymbol{\beta}_0),$ (S.33)

and likewise

$$\int I(u < 0)dF_{U|\boldsymbol{v}}(u) = 1 - F_0(\boldsymbol{v}^T \boldsymbol{\beta}_0).$$
(S.34)

Combining (S.28), (S.29), (S.32), (S.33), and (S.34), we have

$$-\int \left\{ I(u > 0)\gamma_{1}(\boldsymbol{v}; \hat{F}, \hat{\boldsymbol{\beta}}) + I(u < 0)\gamma_{2}(\boldsymbol{v}; \hat{F}, \hat{\boldsymbol{\beta}}) \right\} dF_{U,V}(u, \boldsymbol{v})$$

$$= 4\int \left\{ F_{0}(\boldsymbol{v}^{T}\boldsymbol{\beta}_{0}) - \sqrt{F_{0}(\boldsymbol{v}^{T}\boldsymbol{\beta}_{0})}\sqrt{\hat{F}(\boldsymbol{v}^{T}\hat{\boldsymbol{\beta}})} + 1 - F_{0}(\boldsymbol{v}^{T}\boldsymbol{\beta}_{0}) - \sqrt{1 - F_{0}(\boldsymbol{v}^{T}\boldsymbol{\beta}_{0})}\sqrt{1 - \hat{F}(\boldsymbol{v}^{T}\hat{\boldsymbol{\beta}})} \right\} dF_{V}(\boldsymbol{v})$$

$$= 4\int \left\{ 1 - \sqrt{F_{0}(\boldsymbol{v}^{T}\boldsymbol{\beta}_{0})}\sqrt{\hat{F}(\boldsymbol{v}^{T}\hat{\boldsymbol{\beta}})} - \sqrt{1 - F_{0}(\boldsymbol{v}^{T}\boldsymbol{\beta}_{0})}\sqrt{1 - \hat{F}(\boldsymbol{v}^{T}\hat{\boldsymbol{\beta}})} \right\} dF_{V}(\boldsymbol{v})$$

$$= 2\int \left\{ \sqrt{F_{0}(\boldsymbol{v}^{T}\boldsymbol{\beta}_{0})} - \sqrt{\hat{F}(\boldsymbol{v}^{T}\hat{\boldsymbol{\beta}})} \right\}^{2} dF_{V}(\boldsymbol{v})$$

$$+ \int \left\{ \sqrt{1 - F_0(\boldsymbol{v}^T \boldsymbol{\beta}_0)} - \sqrt{1 - \hat{F}(\boldsymbol{v}^T \hat{\boldsymbol{\beta}})} \right\}^2 dF_V(\boldsymbol{v})$$

$$\geq 0.5 \int \left\{ F_0(\boldsymbol{v}^T \boldsymbol{\beta}_0) - \hat{F}(\boldsymbol{v}^T \hat{\boldsymbol{\beta}}) \right\}^2 dF_V(\boldsymbol{v}), \qquad (S.35)$$

where to derive the last " \geq " we used the fact that $F_0(\cdot) \in [0,1]$ and $\hat{F}(\cdot) \in [0,1]$.

On the other hand,

$$F_V(\boldsymbol{v}) = P(X_i - X_j \leq \boldsymbol{v}) = E \{ P(X_i \leq X_j + \boldsymbol{v} | X_j) \}$$
$$= E \{ F_X(X_j + \boldsymbol{v}) \} = \int F_X(\boldsymbol{x}_2 + \boldsymbol{v}) dF_X(\boldsymbol{x}_2).$$

Therefore,

$$\int \left\{ F_0(\boldsymbol{v}^T \boldsymbol{\beta}_0) - \hat{F}(\boldsymbol{v}^T \hat{\boldsymbol{\beta}}) \right\}^2 dF_V(\boldsymbol{v})$$

=
$$\int \int \left\{ F_0(\boldsymbol{v}^T \boldsymbol{\beta}_0) - \hat{F}(\boldsymbol{v}^T \hat{\boldsymbol{\beta}}) \right\}^2 dF_X(\boldsymbol{x}_2 + \boldsymbol{v}) dF_X(\boldsymbol{x}_2).$$
(S.36)

Setting $x_1 = x_2 + v$, (S.36) immediately implies

$$\begin{split} &\int \left\{ F_0(\boldsymbol{v}^T \boldsymbol{\beta}_0) - \hat{F}(\boldsymbol{v}^T \hat{\boldsymbol{\beta}}) \right\}^2 dF_V(\boldsymbol{v}) \\ &= \int \int \left\{ F_0((\boldsymbol{x}_1 - \boldsymbol{x}_2)^T \boldsymbol{\beta}_0) - \hat{F}((\boldsymbol{x}_1 - \boldsymbol{x}_2)^T \hat{\boldsymbol{\beta}}) \right\}^2 dF_X(\boldsymbol{x}_1) dF_X(\boldsymbol{x}_2) \\ &= D^2 \left(\hat{\boldsymbol{\beta}}, \hat{F}; \boldsymbol{\beta}_0, F_0 \right), \end{split}$$

which together with (S.35) leads to (S.31), and therefore proves (S.26).

Lemma 8 below establishes the asymptotic order of the right-hand side of (S.26).

Lemma 8. Assume Conditions 1–3. We have

$$\int \left\{ I(u>0)\gamma_1(\boldsymbol{v}; \hat{F}, \hat{\boldsymbol{\beta}}) + I(u<0)\gamma_2(\boldsymbol{v}; \hat{F}, \hat{\boldsymbol{\beta}}) \right\} \\
\times d \left\{ \mathbb{F}_{U,V}(u, \boldsymbol{v}) - F_{U,V}(u, \boldsymbol{v}) \right\} \\
= O_p(n^{-2/3}) \vee \left\{ O_p(n^{-1/2})D^{1/2}(\hat{\boldsymbol{\beta}}, \hat{F}; \boldsymbol{\beta}_0, F_0) \right\}.$$
(S.37)

Proof. Recall that we have defined the following notations in Lemma 7. $F_{U,V}$ denotes the joint c.d.f. of $(U_{i,j}, V_{i,j})$ and likewise, $\mathbb{F}_{U,V}(u, v)$ denotes the empirical c.d.f. of $\{(U_{i,j}, V_{i,j})\}_{i,j=1}^n$, where $U_{i,j} = Y_i - Y_j$ and $V_{i,j} = X_i - X_j$.

In the subsequent proof of this lemma, we need Lemma 3.4.2 in van der Vaart (1996), which is reviewed as follows.

Lemma 9. Let \Im be a class of functions such that for every $g \in \Im$, $\int g^2 dP < \widetilde{\delta}^2$ and $\|g\|_{\infty} < M$

for some M > 0, and Z_1, \ldots, Z_n be i.i.d. sample. Then

$$\sup_{g\in\mathfrak{S}} \left| \int gd(\mathbb{F}_Z - F_Z) \right| \lesssim n^{-1/2} \widetilde{J}_{[]}(\widetilde{\delta}, \mathfrak{S}, L_2(F_Z)) \left\{ 1 + \frac{\widetilde{J}_{[]}(\widetilde{\delta}, \mathfrak{S}, L_2(F_Z))}{\widetilde{\delta}^2 \sqrt{n}} M \right\},$$

where

$$\widetilde{J}_{[]}(\widetilde{\delta}, \mathfrak{S}, L_2(F_Z)) = \int_0^{\widetilde{\delta}} \sqrt{1 + H_{2,B}(\delta, \mathfrak{S}, F_Z)}.$$

We now move back to the proof of Lemma 8. Note that we only need to show

$$\int I(u>0)\gamma_1(\boldsymbol{v};\hat{F},\hat{\boldsymbol{\beta}})d\left\{\mathbb{F}_{U,V}(u,\boldsymbol{v})-F_{U,V}(u,\boldsymbol{v})\right\}$$
$$= O_p(n^{-2/3}) \vee \left\{O_p(n^{-1/2})D^{1/2}(\hat{\boldsymbol{\beta}},\hat{F};\boldsymbol{\beta}_0,F_0)\right\}.$$
(S.38)

The same arguments can be applied to show

$$\int I(u < 0)\gamma_2(\boldsymbol{v}; \hat{F}, \hat{\boldsymbol{\beta}})d\left\{\mathbb{F}_{U,V}(u, \boldsymbol{v}) - F_{U,V}(u, \boldsymbol{v})\right)\right\}$$

= $O_p(n^{-2/3}) \lor \left\{O_p(n^{-1/2})D^{1/2}(\hat{\boldsymbol{\beta}}, \hat{F}; \boldsymbol{\beta}_0, F_0)\right\}.$ (S.39)

Then, combining (S.38) and (S.39) leads to (S.37).

To this end, note that by the definition of $\mathbb{F}_{U,V}(u, v)$,

$$\int I(u > 0)\gamma_1(\boldsymbol{v}; \hat{F}, \hat{\boldsymbol{\beta}}) d\mathbb{F}_{U,V}(u, \boldsymbol{v})$$

$$= \frac{1}{n} \sum_{j=1}^n \frac{1}{n} \sum_{i=1}^n I(Y_i > Y_j)\gamma_1(X_i - X_j; \hat{F}, \hat{\boldsymbol{\beta}})$$

$$= \int \int I(y_1 > y_2)\gamma_1(\boldsymbol{x}_1 - \boldsymbol{x}_2; \hat{F}, \hat{\boldsymbol{\beta}}) d\mathbb{F}_{X,Y}(\boldsymbol{x}_1, y_1) d\mathbb{F}_{X,Y}(\boldsymbol{x}_2, y_2), \quad (S.40)$$

where $\mathbb{F}_{X,Y}(\boldsymbol{x}, y)$ denotes the empirical c.d.f. of $\{(X_i, Y_i)\}_{i=1}^n$. Furthermore by the definition of $F_{U,V}(u, \boldsymbol{v})$, we have

$$F_{U,V}(u, \boldsymbol{v}) = P(Y_1 - Y_2 \leq u, X_1 - X_2 \leq \boldsymbol{v})$$

= $E \{ P(Y_1 - Y_2 \leq u, X_1 - X_2 \leq \boldsymbol{v} | Y_2, X_2) \}$
= $E \{ F_{X,Y}(X_2 + \boldsymbol{v}, Y_2 + u) \}$
= $\int F_{X,Y}(\boldsymbol{x}_2 + \boldsymbol{v}, y_2 + u) dF_{X,Y}(\boldsymbol{x}_2, y_2),$

which together with the transformation $u = y_1 - y_2$, $v = x_1 - x_2$ immediately leads to

$$\int I(u > 0)\gamma_1(\boldsymbol{v}; \hat{F}, \hat{\boldsymbol{\beta}}) dF_{U,V}(u, \boldsymbol{v})$$

$$= \int I(u > 0)\gamma_1(\boldsymbol{v}; \hat{F}, \hat{\boldsymbol{\beta}}) \int dF_{X,Y}(\boldsymbol{x}_2 + \boldsymbol{v}, y_2 + u) dF_{X,Y}(\boldsymbol{x}_2, y_2)$$

$$= \int \int I(y_1 - y_2 > 0)\gamma_1(\boldsymbol{x}_1 - \boldsymbol{x}_2; \hat{F}, \hat{\boldsymbol{\beta}}) dF_{X,Y}(\boldsymbol{x}_1, y_1) dF_{X,Y}(\boldsymbol{x}_2, y_2). \quad (S.41)$$

Combining (S.40) with (S.41), to show (S.38), we only need to show

$$\int \int I(y_1 > y_2) \gamma_1(\boldsymbol{x}_1 - \boldsymbol{x}_2; \hat{F}, \hat{\boldsymbol{\beta}}) d\mathbb{F}_{X,Y}(\boldsymbol{x}_1, y_1) d\mathbb{F}_{X,Y}(\boldsymbol{x}_2, y_2)
- \int \int I(y_1 - y_2 > 0) \gamma_1(\boldsymbol{x}_1 - \boldsymbol{x}_2; \hat{F}, \hat{\boldsymbol{\beta}})
\times dF_{X,Y}(\boldsymbol{x}_1, y_1) dF_{X,Y}(\boldsymbol{x}_2, y_2)
= O_p(n^{-2/3}) \vee \left\{ O_p(n^{-1/2}) D^{1/2}(\hat{\boldsymbol{\beta}}, \hat{F}; \boldsymbol{\beta}_0, F_0) \right\}.$$
(S.42)

We will use the following two steps to establish (S.42).

• In step 1, we show

$$\int \left| \int I(y_1 > y_2) \gamma_1(\boldsymbol{x}_1 - \boldsymbol{x}_2; \hat{F}, \hat{\boldsymbol{\beta}}) d\left\{ \mathbb{F}_{X,Y}(\boldsymbol{x}_1, y_1) - F_{X,Y}(\boldsymbol{x}_1, y_1) \right\} \right| \quad (S.43)$$

$$\times d\mathbb{F}_{X,Y}(\boldsymbol{x}_2, y_2)$$

$$= O_p(n^{-2/3}) \vee \left[O_p(n^{-1/2}) \left\{ D\left(\hat{\boldsymbol{\beta}}, \hat{F}; \boldsymbol{\beta}_0, F_0\right) + O_p(n^{-1/2}) \right\}^{1/2} \right].$$

• In *step 2*, we show

$$\int \left\{ \int I(y_1 - y_2 > 0) \gamma_1(\boldsymbol{x}_1 - \boldsymbol{x}_2; \hat{F}, \hat{\boldsymbol{\beta}}) dF_{X,Y}(\boldsymbol{x}_1, y_1) \right\}$$

$$\times d(\mathbb{F}_{X,Y}(\boldsymbol{x}_1, y_1) - F_{X,Y}(\boldsymbol{x}_2, y_2))$$

$$= O_p(n^{-2/3}) \vee \left\{ O_p(n^{-1/2}) D^{1/2} \left(\hat{\boldsymbol{\beta}}, \hat{F}; \boldsymbol{\beta}_0, F_0 \right) \right\}.$$
(S.44)

We start with step 1. Consider the class of functions

$$\widetilde{\mathcal{G}} = \left\{ I(y_1 > y_2) \gamma_1(\boldsymbol{x}_1 - \boldsymbol{x}_2; F, \boldsymbol{\beta}) : \\ F \in \mathcal{F}, \boldsymbol{\beta} \in \boldsymbol{\mathcal{B}}, y_2 \in \boldsymbol{\mathcal{Y}}, \boldsymbol{x}_2 \in \boldsymbol{\mathcal{X}} \right\} = \mathcal{G}_1 \cdot \mathcal{G}_2,$$
(S.45)

where

$$\mathcal{G}_1 = \{ I(y_1 > y_2) : y_2 \in \mathcal{Y} \}; \quad \mathcal{G}_2 = \{ \gamma_1(\boldsymbol{x}_1 - \boldsymbol{x}_2; F, \boldsymbol{\beta}) : F \in \mathcal{F}, \boldsymbol{\beta} \in \mathcal{B}, \boldsymbol{x}_2 \in \mathcal{X} \}.$$

It is straightforward to check that there exists a universal constant M > 0, such that

$$\sup_{g\in\tilde{\mathcal{G}}}|g|_{\infty}\leqslant M.$$
(S.46)

We can also check

$$H_{q,B}(\delta, \mathcal{G}_1, F_{X,Y}) \lesssim -\log \delta.$$
 (S.47)

Furthermore, applying Lemma 5 and Condition 3, we can easily check that

$$H_{q,B}(\delta, \mathcal{G}_2, F_{X,Y}) \lesssim 1/\delta. \tag{S.48}$$

Combining (S.45), (S.47), and (S.48), and applying Lemma 9.25 in Kosorok (2008), we can conclude that

$$H_{q,B}(\delta, \widetilde{\mathcal{G}}, F_{X,Y}) \lesssim 1/\delta.$$
 (S.49)

Now, let $\mathcal{G} = \frac{1}{M} \widetilde{\mathcal{G}}$. Then, we have

$$\begin{split} \sup_{g \in \mathcal{G}} |g|_{\infty} &\leq 1 \\ H_{q,B}(\delta, \mathcal{G}, F_{X,Y}) \lesssim 1/\delta, \end{split}$$

which together with Lemma 4 lead to

$$P\left(\sup_{g\in\mathcal{G}(n^{-1/3})}\left|\int g(\boldsymbol{x}_{1},y_{1})d\left\{\mathbb{F}_{X,Y}(\boldsymbol{x}_{1},y_{1})-F_{X,Y}(\boldsymbol{x}_{1},y_{1})\right\}\right| \ge Tn^{-2/3}\right)$$

$$\leqslant c\exp\left(-Tn^{1/3}/c^{2}\right),$$
(S.50)

and

$$P\left(\sup_{g\in\mathcal{G},\|g\|_{2,F_{X,Y}}>n^{-1/3}}\frac{\left|\int g(\boldsymbol{x}_{1},y_{1})d\left\{\mathbb{F}_{X,Y}(\boldsymbol{x}_{1},y_{1})-F_{X,Y}(\boldsymbol{x}_{1},y_{1})\right\}\right|}{\|g\|_{2,F_{X,Y}}^{1/2}} \geqslant Tn^{-1/2}\right)$$

$$\leqslant c\exp\left(-T/c^{2}\right),\qquad(S.51)$$

for all $T \ge c$ and $n \ge n_0$, where c and n_0 are constants.

Note that (S.51) leads to

$$(\mathbf{S.52})$$

$$1 - c \exp\left(-T/c^2\right)$$

$$< P\left(\sup_{g \in \mathcal{G}, \|g\|_{2, F_{X,Y}} > n^{-1/3}} \frac{\left| \int g(\boldsymbol{x}_{1}, y_{1}) d\left\{ \mathbb{F}_{X,Y}(\boldsymbol{x}_{1}, y_{1}) - F_{X,Y}(\boldsymbol{x}_{1}, y_{1}) \right\} \right|}{\|g\|_{2, F_{X,Y}}^{1/2}} < Tn^{-1/2} \right)$$

$$\leq P\left(\bigcap_{i=1}^{n} \left\{ \sup_{g \in \mathcal{G}_{X_{i}, Y_{i}}, \|g\|_{2, F_{X,Y}} > n^{-1/3}} \frac{\left| \int g(\boldsymbol{x}_{1}, y_{1}) d\left\{ \mathbb{F}_{X,Y}(\boldsymbol{x}_{1}, y_{1}) - F_{X,Y}(\boldsymbol{x}_{1}, y_{1}) \right\} \right|}{\|g\|_{2, F_{X,Y}}^{1/2}} < Tn^{-1/2} \right\} \right)$$

$$\leq P\left(\bigcap_{i=1}^{n} \left\{ \frac{\left| \int \hat{g}_{i}(\boldsymbol{x}_{1}, y_{1}) d\left\{ \mathbb{F}_{X,Y}(\boldsymbol{x}_{1}, y_{1}) - F_{X,Y}(\boldsymbol{x}_{1}, y_{1}) \right\} \right| I\left(\|\hat{g}_{i}\|_{2, F_{X,Y}} > Mn^{-1/3}\right)}{\|\hat{g}_{i}\|_{2, F_{X,Y}}^{1/2}} < Tn^{-1/2} \right\} \right)$$

$$\leq P\left(\frac{\frac{1}{n} \sum_{i=1}^{n} \left| \int \hat{g}_{i}(\boldsymbol{x}_{1}, y_{1}) d\left\{ \mathbb{F}_{X,Y}(\boldsymbol{x}_{1}, y_{1}) - F_{X,Y}(\boldsymbol{x}_{1}, y_{1}) \right\} \right| I\left(\|\hat{g}_{i}\|_{2, F_{X,Y}} > Mn^{-1/3}\right)}{\|\hat{g}_{i}\|_{2, F_{X,Y}}^{1/2}} < Tn^{-1/2} \right\} \right),$$

where

$$\mathcal{G}_{X_i,Y_i} = \frac{1}{M} \Big\{ I(y_1 > Y_i) \gamma_1(\boldsymbol{x}_1 - X_i; F, \boldsymbol{\beta}) : F \in \mathcal{F}, \boldsymbol{\beta} \in \boldsymbol{\beta} \Big\}$$
$$\hat{g}_i(\boldsymbol{x}, y) = \frac{1}{M} I(y > Y_i) \gamma_1(\boldsymbol{x} - X_i; \hat{F}, \boldsymbol{\beta}).$$
(S.53)

Applying the similar arguments on (S.50) leads to

$$1 - c \exp\left(-Tn^{1/3}/c^{2}\right)$$

$$< P\left(\frac{1}{n}\sum_{i=1}^{n}\left|\int \hat{g}_{i}(\boldsymbol{x}_{1}, y_{1})d\left\{\mathbb{F}_{X,Y}(\boldsymbol{x}_{1}, y_{1}) - F_{X,Y}(\boldsymbol{x}_{1}, y_{1})\right\}\right|$$

$$\times I\left(\|\hat{g}_{i}\|_{2, F_{X,Y}} \leq Mn^{-1/3}\right) < Tn^{-2/3}\right).$$
(S.54)

Combining (S.52) and (S.54), we have

$$P\left(\frac{1}{n}\sum_{i=1}^{n}\left|\int \widehat{g}_{i}(\boldsymbol{x}_{1}, y_{1})d\left\{\mathbb{F}_{X,Y}(\boldsymbol{x}_{1}, y_{1}) - F_{X,Y}(\boldsymbol{x}_{1}, y_{1})\right\}\right|$$

$$< Tn^{-2/3} + Tn^{-1/2}\frac{1}{n}\sum_{i=1}^{n}\|\widehat{g}_{i}\|_{2,F_{X,Y}}^{1/2}\right)$$

$$> 1 - c\exp\left(-Tn^{1/3}/c^{2}\right) - c\exp\left(-T/c^{2}\right),$$

which together the definition of \hat{g}_i in (S.53) indicates

$$\int \left| \int I(y_1 > y_2) \gamma_1(\boldsymbol{x}_1 - \boldsymbol{x}_2; \hat{F}, \hat{\boldsymbol{\beta}}) d\left\{ \mathbb{F}_{X,Y}(\boldsymbol{x}_1, y_1) - F_{X,Y}(\boldsymbol{x}_1, y_1) \right\} \right| \quad (S.55) \\
\times d\mathbb{F}_{X,Y}(\boldsymbol{x}_2, y_2) \\
= O_p(n^{-2/3}) \lor O_p(n^{-1/2}) \frac{1}{n} \sum_{i=1}^n \| I(y_1 > Y_i) \gamma_1(\boldsymbol{x}_1 - X_i; \hat{F}, \hat{\boldsymbol{\beta}}) \|_{2, F_{X,Y}}^{1/2}.$$

Next, we shall study the convergence of

$$J_0 = \frac{1}{n} \sum_{i=1}^n \|I(y_1 > Y_i)\gamma_1(\boldsymbol{x}_1 - X_i; \hat{F}, \hat{\boldsymbol{\beta}})\|_{2, F_{X, Y}}^{1/2}.$$
 (S.56)

Applying Condition 3, we have

$$\begin{split} \|I(y_{1} > Y_{i})\gamma_{1}(\boldsymbol{x}_{1} - X_{i}; \hat{F}, \hat{\boldsymbol{\beta}})\|_{2, F_{X,Y}}^{2} \\ &= \int I(y_{1} > Y_{i})\gamma_{1}^{2}(\boldsymbol{x}_{1} - X_{i}; \hat{F}, \hat{\boldsymbol{\beta}})dF_{X,Y}(\boldsymbol{x}_{1}, y_{1}) \\ &\leq \int \gamma_{1}^{2}(\boldsymbol{x}_{1} - X_{i}; \hat{F}, \hat{\boldsymbol{\beta}})dF_{X}(\boldsymbol{x}_{1}) = 16\int \left\{1 - \sqrt{\frac{\hat{F}((\boldsymbol{x}_{1} - X_{i})^{T} \hat{\boldsymbol{\beta}})}{F_{0}((\boldsymbol{x}_{1} - X_{i})^{T} \boldsymbol{\beta}_{0})}}\right\}^{2} dF_{X}(\boldsymbol{x}_{1}) \\ &\leq \int \left\{\sqrt{F_{0}((\boldsymbol{x}_{1} - X_{i})^{T} \boldsymbol{\beta}_{0})} - \sqrt{\hat{F}((\boldsymbol{x}_{1} - X_{i})^{T} \hat{\boldsymbol{\beta}})}\right\}^{2} dF_{X}(\boldsymbol{x}_{1}) \\ &= \int \left\{\frac{F_{0}((\boldsymbol{x}_{1} - X_{i})^{T} \boldsymbol{\beta}_{0}) - \hat{F}((\boldsymbol{x}_{1} - X_{i})^{T} \hat{\boldsymbol{\beta}})}{\sqrt{F_{0}((\boldsymbol{x}_{1} - X_{i})^{T} \boldsymbol{\beta}_{0})} + \sqrt{\hat{F}((\boldsymbol{x}_{1} - X_{i})^{T} \hat{\boldsymbol{\beta}})}\right\}^{2} dF_{X}(\boldsymbol{x}_{1}) \\ &\leq \int \left\{F_{0}((\boldsymbol{x}_{1} - X_{i})^{T} \boldsymbol{\beta}_{0}) - \hat{F}((\boldsymbol{x}_{1} - X_{i})^{T} \hat{\boldsymbol{\beta}})\right\}^{2} dF_{X}(\boldsymbol{x}_{1}) \\ &\leq \int \left\{F_{0}((\boldsymbol{x}_{1} - X_{i})^{T} \boldsymbol{\beta}_{0}) - \hat{F}((\boldsymbol{x}_{1} - X_{i})^{T} \hat{\boldsymbol{\beta}})\right\}^{2} dF_{X}(\boldsymbol{x}_{1}) \\ &= d_{\hat{\boldsymbol{\beta}}, \hat{F}}^{2}(X_{i}), \end{split}$$

which together with the fact that $x^{1/2}$ for any x > 0 is a strictly concave function leads to

$$0 \leq J_{0} \leq \left\{ \frac{1}{n} \sum_{i=1}^{n} \|I(y_{1} > Y_{i})\gamma_{1}(\boldsymbol{x}_{1} - X_{i}; \widehat{F}, \widehat{\boldsymbol{\beta}})\|_{2, F_{X, Y}} \right\}^{1/2} \\ \leq \left\{ \frac{1}{n} \sum_{i=1}^{n} d_{\widehat{\boldsymbol{\beta}}, \widehat{F}}(X_{i}) \right\}^{1/2}.$$
(S.57)

Note that $d_{\widehat{\boldsymbol{\beta}},\widehat{F}}(\boldsymbol{x}) \in \mathcal{D}$. Incorporating (S.18) with Lemma 9 leads to

$$\frac{1}{n} \sum_{i=1}^{n} d_{\widehat{\beta},\widehat{F}}(X_{i})$$

$$= \int d_{\widehat{\beta},\widehat{F}}(\boldsymbol{x}_{2}) dF_{X}(\boldsymbol{x}_{2}) + O_{p}(n^{-1/2})$$

$$= \int \left[\int \left\{ F_{0}((\boldsymbol{x}_{1} - \boldsymbol{x}_{2})^{T} \boldsymbol{\beta}_{0}) - \widehat{F}((\boldsymbol{x}_{1} - \boldsymbol{x}_{2})^{T} \widehat{\boldsymbol{\beta}}) \right\}^{2} dF_{X}(\boldsymbol{x}_{1}) \right]^{1/2} dF_{X}(\boldsymbol{x}_{2})$$

$$+ O_{p}(n^{-1/2})$$
(S.58)

$$\leq \left[\int \int \left\{ F_0((\boldsymbol{x}_1 - \boldsymbol{x}_2)^T \boldsymbol{\beta}_0) - \widehat{F}((\boldsymbol{x}_1 - \boldsymbol{x}_2)^T \widehat{\boldsymbol{\beta}}) \right\}^2 dF_X(\boldsymbol{x}_1) dF_X(\boldsymbol{x}_2) \right]^{1/2} \\ + O_p(n^{-1/2}) \\ = D(\widehat{\boldsymbol{\beta}}, \widehat{F}; \boldsymbol{\beta}_0, F_0) + O_p(n^{-1/2}).$$

Combining (S.55)-(S.58) leads to

$$\int \left| \int I(y_1 > y_2) \gamma_1(\boldsymbol{x}_1 - \boldsymbol{x}_2; \hat{F}, \hat{\boldsymbol{\beta}}) d\left\{ \mathbb{F}_{X,Y}(\boldsymbol{x}_1, y_1) - F_{X,Y}(\boldsymbol{x}_1, y_1) \right\} \right| d\mathbb{F}_{X,Y}(\boldsymbol{x}_2, y_2)$$

= $O_p(n^{-2/3}) \lor \left[O_p(n^{-1/2}) \left\{ D\left(\hat{\boldsymbol{\beta}}, \hat{F}; \boldsymbol{\beta}_0, F_0\right) + O_p(n^{-1/2}) \right\}^{1/2} \right],$

which proves (S.43) and therefore completes our proof in step 1.

We proceed to show (S.44) claimed in step 2. Let

$$\widetilde{\mathcal{G}}_{I} = \left\{ \int g_{\beta,F}(\boldsymbol{x}_{1}, \boldsymbol{x}_{2}, y_{1}, y_{2}) dF_{X,Y}(\boldsymbol{x}_{1}, y_{1}) : g_{\beta,F}(\boldsymbol{x}_{1}, \boldsymbol{x}_{2}, y_{1}, y_{2}) \in \mathcal{G}_{I,1} \right\}$$
(S.59)

where

$$\begin{aligned} \mathcal{G}_{I,1} &= \left\{ g_{\beta,F}(\boldsymbol{x}_{1}, \boldsymbol{x}_{2}, y_{1}, y_{2}) = I(y_{1} > y_{2})\gamma_{1}(\boldsymbol{x}_{1} - \boldsymbol{x}_{2}; F, \beta) : F \in \mathcal{F}, \beta \in \mathcal{B} \right\} \\ &= I(y_{1} > y_{2}) \cdot \mathcal{G}_{I,2}, \\ \mathcal{G}_{I,2} &= \{\gamma_{1}(\boldsymbol{x}_{1} - \boldsymbol{x}_{2}; F, \beta) : F \in \mathcal{F}, \beta \in \mathcal{B} \}. \end{aligned}$$

Based on Condition 3, it is easy to check that there exists a universal constant M > 0, such that

$$\sup_{g \in \widetilde{\mathcal{G}}_I} |g|_{\infty} < M. \tag{S.60}$$

Define $G_I = \frac{1}{M} \widetilde{G}_I$. With (S.16) in Lemma 5 and Condition 3, it is straightforward to show that

$$H_{q,B}\left(\delta, \frac{1}{M}\mathcal{G}_{I,1}, F_{X_1, X_2, Y_1, Y_2}\right) \lesssim 1/\delta,$$

where $F_{X_1,X_2,Y_1,Y_2}(\boldsymbol{x_1},\boldsymbol{x_2},y_1,y_2)$ denotes the joint c.d.f. of $(X_1^T,X_2^T,Y_1,Y_2)^T$. Therefore there exists a set of δ -brackets { $[l_i(\boldsymbol{x}_1,\boldsymbol{x}_2,y_1,y_2), u_i(\boldsymbol{x}_1,\boldsymbol{x}_2,y_1,y_2)]_{i=1}^{N_4}$ } under $L_q(F_{X_1,X_2,Y_1,Y_2})$ that cover $\frac{1}{M}\mathcal{G}_{I,1}$, where $N_4 \leq 1/\delta$. Consider the set of brackets

$$\left\{ [L_i(\boldsymbol{x}_2, y_2), U_i(\boldsymbol{x}_2, y_2)] :$$
(S.61)

$$L_i(\boldsymbol{x}_2, y_2) = \int l_i(\boldsymbol{x}_1, \boldsymbol{x}_2, y_1, y_2) dF_{X,Y}(\boldsymbol{x}_1, y_1); \\ U_i(\boldsymbol{x}_2, y_2) = \int u_i(\boldsymbol{x}_1, \boldsymbol{x}_2, y_1, y_2) dF_{X,Y}(\boldsymbol{x}_1, y_1); \quad i = 1, \dots, N_4 \bigg\}.$$

Clearly, for every $\frac{1}{M} \int g_{\beta,F}(\boldsymbol{x}_1, \boldsymbol{x}_2, y_1, y_2) dF_{X,Y}(\boldsymbol{x}_1, y_1) \in \mathcal{G}_I$, there exists an $i \in \{1, \ldots, N_4\}$ such that

$$L_{i}(\boldsymbol{x}_{2}, y_{2}) \leq \frac{1}{M} \int g_{\boldsymbol{\beta}, F}(\boldsymbol{x}_{1}, \boldsymbol{x}_{2}, y_{1}, y_{2}) dF_{X, Y}(\boldsymbol{x}_{1}, y_{1}) \leq U_{i}(\boldsymbol{x}_{2}, y_{2}),$$
(S.62)

which indicates the set of brackets given in (S.61) covers G_I , with bracket length

$$\left[\int \{U_i(\boldsymbol{x}_2, y_2) - L_i(\boldsymbol{x}_2, y_2)\}^q dF_{X,Y}(\boldsymbol{x}_2, y_2) \right]^{1/q}$$

$$= \left[\int \left\{ \int (u_i(\boldsymbol{x}_1, \boldsymbol{x}_2, y_1, y_2) - l_i(\boldsymbol{x}_1, \boldsymbol{x}_2, y_1, y_2)) dF_{X,Y}(\boldsymbol{x}_1, y_1) \right\}^q dF_{X,Y}(\boldsymbol{x}_2, y_2) \right]^{1/q}$$

$$\leq \left[\int \int \{u_i(\boldsymbol{x}_1, \boldsymbol{x}_2, y_1, y_2) - l_i(\boldsymbol{x}_1, \boldsymbol{x}_2, y_1, y_2)\}^q dF_{X,Y}(\boldsymbol{x}_1, y_1) dF_{X,Y}(\boldsymbol{x}_2, y_2) \right]^{1/q}$$

$$= \|u_i - l_i\|_{q, F_{X_1, X_2, Y_1, Y_2}} \leq \delta.$$

$$(S.63)$$

(S.62) and (S.63) imply that the set of brackets given in (S.61) are δ -brackets under $L_q(F_{X,Y})$ that cover \mathcal{G}_I . Therefore

$$H_{q,B}(\delta, \mathcal{G}_I, F_{X,Y}) \lesssim 1/\delta,$$

which together with the fact in (S.60) and Lemma 4 leads to

$$P\left(\sup_{g\in\mathcal{G}_{I}(n^{-1/3})}\left|\int g(\boldsymbol{x}_{2},y_{2})d\left\{\mathbb{F}_{X,Y}(\boldsymbol{x}_{2},y_{2})-F_{X,Y}(\boldsymbol{x}_{2},y_{2})\right\}\right| \ge Tn^{-2/3}\right)$$
$$\leqslant c\exp\left(-Tn^{1/3}/c^{2}\right),$$

and

$$P\left(\sup_{g\in\mathcal{G}_{I},\|g\|_{2,F_{X,Y}}>n^{-1/3}}\frac{\left|\int g(\boldsymbol{x}_{2},y_{2})d\left\{\mathbb{F}_{X,Y}(\boldsymbol{x}_{2},y_{2})-F_{X,Y}(\boldsymbol{x}_{2},y_{2})\right\}\right|}{\|g\|_{2,F_{X,Y}}^{1/2}} \geqslant Tn^{-1/2}\right)$$
$$\leqslant c\exp\left(-T/c^{2}\right),$$

for all $T \ge c$ and $n \ge n_0$, where c and n_0 are constants. This together with the fact that

$$\int I(y_1 > y_2)\gamma_1(\boldsymbol{x}_1 - \boldsymbol{x}_2; \hat{F}, \hat{\boldsymbol{\beta}})dF_{X,Y}(\boldsymbol{x}_1, y_1) \in \mathcal{G}_I$$

immediately implies

$$\int \left\{ \int I(y_1 > y_2) \gamma_1(\boldsymbol{x}_1 - \boldsymbol{x}_2; \hat{F}, \hat{\boldsymbol{\beta}}) dF_{X,Y}(\boldsymbol{x}_1, y_1) \right\}$$
(S.64)
$$\times d \left\{ \mathbb{F}_{X,Y}(\boldsymbol{x}_2, y_2) - F_{X,Y}(\boldsymbol{x}_2, y_2) \right\}$$
$$= O_p(n^{-2/3}) \lor O_p(n^{-1/2})$$
$$\times \left\| \int I(y_1 > y_2) \gamma_1(\boldsymbol{x}_1 - \boldsymbol{x}_2; \hat{F}, \hat{\boldsymbol{\beta}}) dF_{X,Y}(\boldsymbol{x}_1, y_1) \right\|_{2, F_{X,Y}}^{1/2}.$$

On the other hand, with Condition 3,

$$\begin{split} \left\| \int I(y_{1} > y_{2})\gamma_{1}(\boldsymbol{x}_{1} - \boldsymbol{x}_{2}; \hat{F}, \hat{\boldsymbol{\beta}})dF_{X,Y}(\boldsymbol{x}_{1}, y_{1}) \right\|_{2,F_{X,Y}}^{2} \tag{S.65} \\ &= \int \left\{ \int I(y_{1} > y_{2})\gamma_{1}(\boldsymbol{x}_{1} - \boldsymbol{x}_{2}; \hat{F}, \hat{\boldsymbol{\beta}})dF_{X,Y}(\boldsymbol{x}_{1}, y_{1}) \right\}^{2} dF_{X,Y}(\boldsymbol{x}_{2}, y_{2}) \\ &\leq \int \int I(y_{1} > y_{2})\gamma_{1}^{2}(\boldsymbol{x}_{1} - \boldsymbol{x}_{2}; \hat{F}, \hat{\boldsymbol{\beta}})dF_{X,Y}(\boldsymbol{x}_{1}, y_{1})dF_{X,Y}(\boldsymbol{x}_{2}, y_{2}) \\ &\leq \int \int \gamma_{1}^{2}(\boldsymbol{x}_{1} - \boldsymbol{x}_{2}; \hat{F}, \hat{\boldsymbol{\beta}})dF_{X}(\boldsymbol{x}_{1})dF_{X}(\boldsymbol{x}_{2}) \\ &= 16 \int \int \left\{ 1 - \sqrt{\frac{\hat{F}((\boldsymbol{x}_{1} - \boldsymbol{x}_{2})^{T}\hat{\boldsymbol{\beta}})}{F_{0}((\boldsymbol{x}_{1} - \boldsymbol{x}_{2})^{T}\beta_{0}} \right\}^{2} dF_{X}(\boldsymbol{x}_{1})dF_{X}(\boldsymbol{x}_{2}) \\ &\leq \int \int \left\{ \sqrt{F_{0}((\boldsymbol{x}_{1} - \boldsymbol{x}_{2})^{T}\beta_{0}} - \sqrt{\hat{F}((\boldsymbol{x}_{1} - \boldsymbol{x}_{2})^{T}\hat{\boldsymbol{\beta}})} \right\}^{2} dF_{X}(\boldsymbol{x}_{1})dF_{X}(\boldsymbol{x}_{2}) \\ &= \int \int \left\{ \frac{F_{0}((\boldsymbol{x}_{1} - \boldsymbol{x}_{2})^{T}\beta_{0}) - \hat{F}((\boldsymbol{x}_{1} - \boldsymbol{x}_{2})^{T}\hat{\boldsymbol{\beta}})}{\sqrt{F_{0}((\boldsymbol{x}_{1} - \boldsymbol{x}_{2})^{T}\beta_{0})} + \sqrt{\hat{F}((\boldsymbol{x}_{1} - \boldsymbol{x}_{2})^{T}\hat{\boldsymbol{\beta}})} \right\}^{2} dF_{X}(\boldsymbol{x}_{1})dF_{X}(\boldsymbol{x}_{2}) \\ &\leq \int \int \left\{ F_{0}((\boldsymbol{x}_{1} - \boldsymbol{x}_{2})^{T}\beta_{0}) - \hat{F}((\boldsymbol{x}_{1} - \boldsymbol{x}_{2})^{T}\hat{\boldsymbol{\beta}}) \right\}^{2} dF_{X}(\boldsymbol{x}_{1})dF_{X}(\boldsymbol{x}_{2}) \\ &= \int \int \left\{ F_{0}((\boldsymbol{x}_{1} - \boldsymbol{x}_{2})^{T}\beta_{0}) - \hat{F}((\boldsymbol{x}_{1} - \boldsymbol{x}_{2})^{T}\hat{\boldsymbol{\beta}}) \right\}^{2} dF_{X}(\boldsymbol{x}_{1})dF_{X}(\boldsymbol{x}_{2}) \\ &\leq \int \int \left\{ F_{0}((\boldsymbol{x}_{1} - \boldsymbol{x}_{2})^{T}\beta_{0}) - \hat{F}((\boldsymbol{x}_{1} - \boldsymbol{x}_{2})^{T}\hat{\boldsymbol{\beta}}) \right\}^{2} dF_{X}(\boldsymbol{x}_{1})dF_{X}(\boldsymbol{x}_{2}) \\ &= D^{2} \left(\hat{\boldsymbol{\beta}}, \hat{F}; \beta_{0}, F_{0} \right), \end{split}$$

which together with (S.64) leads to (S.44), and therefore completes our proof of <u>step 2</u>. The proof of Lemma 8 is completed.

If we combine Lemmas 7 and 8, the result claimed in Part (a) follows immediately. We proceed to show Part (b). We first establish the consistency of $\hat{\beta}$.

Lemma 10. Assume Conditions 1–4. We have

$$\widehat{\boldsymbol{\beta}} - \boldsymbol{\beta}_0 = o_p(1). \tag{S.66}$$

Proof. Let

$$M^*(\boldsymbol{\beta}) = \inf_{F \in \mathcal{F}} D(\boldsymbol{\beta}; F, \boldsymbol{\beta}_0, F_0).$$

Then, from the arguments in Wald (1949), to show (S.66), we need to show only that

- (i) $M^*(\hat{\beta}) = o_p(1);$
- (ii) $M^*(\boldsymbol{\beta}) = 0$ implies that $\boldsymbol{\beta} = \boldsymbol{\beta}_0$;
- (iii) $M^*(\beta)$ is continuous in \mathcal{B} .

Part (i) is easily obtained from Part (a) of this theorem, since

$$0 \leq M^*(\widehat{\boldsymbol{\beta}}) \leq D(\widehat{\boldsymbol{\beta}}, \widehat{F}; \boldsymbol{\beta}_0, F_0) = o_p(1).$$

Part (ii) holds because of Condition 4. We now show Part (iii). Similarly to the proof of (S.17) in Lemma 5, for an arbitrary $\delta_1 > 0$ and β^* there exists a set of brackets $[L_i(\boldsymbol{x}_1 - \boldsymbol{x}_2), U_i(\boldsymbol{x}_1 - \boldsymbol{x}_2)]_{i=1}^{N^*}$ satisfying

$$\left[\int \int \left\{ U_i(\boldsymbol{x}_1 - \boldsymbol{x}_2) - L_i(\boldsymbol{x}_1 - \boldsymbol{x}_2) \right\}^2 dF_X(\boldsymbol{x}_1) dF_X(\boldsymbol{x}_2) \right]^{1/2} \leqslant C^* \delta_1, \quad (S.67)$$

where C^* is a universal constant. Moreover, for any $F \in \mathcal{F}$, there exists an $i \in \{1, \ldots, N^*\}$ such that

$$0 \leq L_i(\boldsymbol{x}_1 - \boldsymbol{x}_2) \leq F((\boldsymbol{x}_1 - \boldsymbol{x}_2)^T \boldsymbol{\beta}) \leq U_i(\boldsymbol{x}_1 - \boldsymbol{x}_2) \leq 1, \text{ and} 0 \leq L_i(\boldsymbol{x}_1 - \boldsymbol{x}_2) \leq F((\boldsymbol{x}_1 - \boldsymbol{x}_2)^T \boldsymbol{\beta}^*) \leq U_i(\boldsymbol{x}_1 - \boldsymbol{x}_2) \leq 1,$$
(S.68)

for any $\|\beta - \beta^*\|_1 \leq \delta_1$. Now for any $\epsilon > 0$, we set $\delta_1 = \epsilon/C^*$. Then when $\|\beta - \beta^*\|_1 \leq \delta_1$, we have

$$D(\boldsymbol{\beta}, F; \boldsymbol{\beta}_0, F_0) \ge D(\boldsymbol{\beta}^*, F; \boldsymbol{\beta}_0, F_0) - D(\boldsymbol{\beta}^*, F; \boldsymbol{\beta}, F),$$
(S.69)

from the triangle inequality. On the other hand, from (S.67), (S.68), and the definition of $D(\beta^*, F; \beta, F)$, we have

$$D(\boldsymbol{\beta}^{*}, F; \boldsymbol{\beta}, F)$$

$$= \left[\int \int \left\{ F((\boldsymbol{x}_{1} - \boldsymbol{x}_{2})^{T} \boldsymbol{\beta}) - F((\boldsymbol{x}_{1} - \boldsymbol{x}_{2})^{T} \boldsymbol{\beta}^{*}) \right\}^{2} dF_{X}(\boldsymbol{x}_{1}) dF_{X}(\boldsymbol{x}_{2}) \right]^{1/2}$$

$$\leq \left[\int \int \left\{ U_{i}(\boldsymbol{x}_{1} - \boldsymbol{x}_{2}) - L_{i}(\boldsymbol{x}_{1} - \boldsymbol{x}_{2}) \right\}^{2} dF_{X}(\boldsymbol{x}_{1}) dF_{X}(\boldsymbol{x}_{2}) \right]^{1/2}$$

$$\leq C^{*} \delta_{1} = \epsilon,$$

which together with (S.69) implies

$$D(\boldsymbol{\beta}, F; \boldsymbol{\beta}_0, F_0) \ge D(\boldsymbol{\beta}^*, F; \boldsymbol{\beta}_0, F_0) - \epsilon.$$

Therefore,

$$M^{*}(\boldsymbol{\beta}) = \inf_{F \in \mathcal{F}} D(\boldsymbol{\beta}; F, \boldsymbol{\beta}_{0}, F_{0})$$

$$\geq \inf_{F \in \mathcal{F}} D(\boldsymbol{\beta}^{*}, F; \boldsymbol{\beta}_{0}, F_{0}) - \epsilon = M^{*}(\boldsymbol{\beta}^{*}) - \epsilon.$$
(S.70)

Using similar arguments, we can show that

$$M^*(\boldsymbol{\beta}) \leqslant M^*(\boldsymbol{\beta}^*) + \epsilon. \tag{S.71}$$

Combining (S.70) and (S.71), we prove Part (iii). This completes the proof of this lemma. \blacksquare

Recall that in the proof of Lemma 7, we have shown that

$$D^{2}(\boldsymbol{\beta}; F, \boldsymbol{\beta}_{0}, F_{0}) = \int \left\{ F_{0}(\boldsymbol{v}^{T}\boldsymbol{\beta}_{0}) - \widehat{F}(\boldsymbol{v}^{T}\widehat{\boldsymbol{\beta}}) \right\}^{2} dF_{V}(\boldsymbol{v}), \qquad (S.72)$$

where $F_V(\boldsymbol{v})$ is the c.d.f. of $X_i - X_j$ for $i \neq j$. Let V be a random variable independent of but sharing the same distribution with $X_i - X_j$, and let $\psi(\boldsymbol{v}^T\boldsymbol{\beta}) = E(F_0(V^T\boldsymbol{\beta}_0)|V^T\boldsymbol{\beta} = \boldsymbol{v}\boldsymbol{\beta})$. Combining (S.72) with the results in Part (a) leads to

$$O_{p}(n^{-2/3}) = \int \left\{ F_{0}(\boldsymbol{v}^{T}\boldsymbol{\beta}_{0}) - \hat{F}(\boldsymbol{v}^{T}\hat{\boldsymbol{\beta}}) \right\}^{2} dF_{V}(\boldsymbol{v})$$

$$= \int \left\{ F_{0}(\boldsymbol{v}^{T}\boldsymbol{\beta}_{0}) - \psi(\boldsymbol{v}^{T}\hat{\boldsymbol{\beta}}) \right\}^{2} dF_{V}(\boldsymbol{v}) + \int \left\{ \psi(\boldsymbol{v}^{T}\hat{\boldsymbol{\beta}}) - \hat{F}(\boldsymbol{v}^{T}\hat{\boldsymbol{\beta}}) \right\}^{2} dF_{V}(\boldsymbol{v})$$

$$+ \int \left\{ F_{0}(\boldsymbol{v}^{T}\boldsymbol{\beta}_{0}) - \psi(\boldsymbol{v}^{T}\hat{\boldsymbol{\beta}}) \right\} \left\{ \psi(\boldsymbol{v}^{T}\hat{\boldsymbol{\beta}}) - \hat{F}(\boldsymbol{v}^{T}\hat{\boldsymbol{\beta}}) \right\} dF_{V}(\boldsymbol{v})$$

$$\geq \int \left\{ F_{0}(\boldsymbol{v}^{T}\boldsymbol{\beta}_{0}) - \psi(\boldsymbol{v}^{T}\hat{\boldsymbol{\beta}}) \right\}^{2} dF_{V}(\boldsymbol{v})$$

$$= E \left[\left\{ F_{0}(V^{T}\boldsymbol{\beta}_{0}) - \psi(V^{T}\hat{\boldsymbol{\beta}}) \right\}^{2} \left| \hat{\boldsymbol{\beta}} \right],$$
(S.73)

where the " \geq " is because

$$\int \left\{ F_0(\boldsymbol{v}^T\boldsymbol{\beta}_0) - \psi(\boldsymbol{v}^T\boldsymbol{\hat{\beta}}) \right\} \left\{ \psi(\boldsymbol{v}^T\boldsymbol{\hat{\beta}}) - \hat{F}(\boldsymbol{v}^T\boldsymbol{\hat{\beta}}) \right\} dF_V(\boldsymbol{v})$$

$$= E \left[\left\{ F_0(V^T\boldsymbol{\beta}_0) - \psi(V^T\boldsymbol{\hat{\beta}}) \right\} \left\{ \psi(V^T\boldsymbol{\hat{\beta}}) - \hat{F}(V^T\boldsymbol{\hat{\beta}}) \right\} \left| \{(X_i, Y_i)\}_{i=1}^n \right]$$

$$= E \left[E \left[\left\{ F_0(V^T\boldsymbol{\beta}_0) - \psi(V^T\boldsymbol{\hat{\beta}}) \right\} \left\{ \psi(V^T\boldsymbol{\hat{\beta}}) - \hat{F}(V^T\boldsymbol{\hat{\beta}}) \right\} \left| \{(X_i, Y_i)\}_{i=1}^n, V^T\boldsymbol{\hat{\beta}} \right] \left| \{(X_i, Y_i)\}_{i=1}^n \right]$$

$$= E \left[\left\{ \psi(V^T\boldsymbol{\hat{\beta}}) - \hat{F}(V^T\boldsymbol{\hat{\beta}}) \right\} E \left[\left\{ F_0(V^T\boldsymbol{\beta}_0) - \psi(V^T\boldsymbol{\hat{\beta}}) \right\} \left| \{(X_i, Y_i)\}_{i=1}^n, V^T\boldsymbol{\hat{\beta}} \right] \left| \{(X_i, Y_i)\}_{i=1}^n \right]$$

$$= E \left[\left\{ \psi(V^T\boldsymbol{\hat{\beta}}) - \hat{F}(V^T\boldsymbol{\hat{\beta}}) \right\} E \left[\left\{ F_0(V^T\boldsymbol{\beta}_0) - \psi(V^T\boldsymbol{\hat{\beta}}) \right\} \left| V^T\boldsymbol{\hat{\beta}} \right] \left| \{(X_i, Y_i)\}_{i=1}^n \right]$$

$$= E \left[\left\{ \psi(V^T\boldsymbol{\hat{\beta}}) - \hat{F}(V^T\boldsymbol{\hat{\beta}}) \right\} E \left[\left\{ F_0(V^T\boldsymbol{\beta}_0) - \psi(V^T\boldsymbol{\hat{\beta}}) \right\} \left| V^T\boldsymbol{\hat{\beta}} \right] \right| \left\{ (X_i, Y_i)\}_{i=1}^n \right]$$

Furthermore, let

$$R(V,\widehat{\boldsymbol{\beta}}) = F_0(V^T\boldsymbol{\beta}_0) - F_0(V^T\widehat{\boldsymbol{\beta}}) - F_0'(V^T\widehat{\boldsymbol{\beta}})V^T(\boldsymbol{\beta}_0 - \widehat{\boldsymbol{\beta}}).$$

Then

$$|R(V,\widehat{\boldsymbol{\beta}})| \leq 0.5 \sup_{u \in \mathbb{R}} |F_0''(u)| \sup_{\boldsymbol{v} \in \mathcal{X} - \mathcal{X}} \|\boldsymbol{v}\|_2^2 \|\widehat{\boldsymbol{\beta}} - \boldsymbol{\beta}_0\|^2 \lesssim \|\widehat{\boldsymbol{\beta}} - \boldsymbol{\beta}_0\|^2.$$

Hence, there exists a constant $C_R > 0$ such that

$$|R(V,\hat{\boldsymbol{\beta}})| \leq C_R \|\hat{\boldsymbol{\beta}} - \boldsymbol{\beta}_0\|^2.$$
(S.74)

Noting (S.74) and Condition 5, we have

$$E\left[\left\{F_{0}(V^{T}\beta_{0})-\psi(V^{T}\widehat{\beta})\right\}^{2}\left|\widehat{\beta}\right]\right]$$

$$=E\left[\left\{F_{0}'(V^{T}\widehat{\beta})(\beta_{0}-\widehat{\beta})^{T}\left(V-E(V|V^{T}\widehat{\beta},\widehat{\beta})\right)\right.$$

$$\left.+R(V,\widehat{\beta})-E(R(V,\widehat{\beta})|V^{T}\widehat{\beta},\widehat{\beta})\right\}^{2}\left|\widehat{\beta}\right]\right]$$

$$\geq 0.5E\left[\left\{F_{0}'(V^{T}\widehat{\beta})(\beta_{0}-\widehat{\beta})^{T}\left(V-E(V|V^{T}\widehat{\beta},\widehat{\beta})\right)\right\}^{2}\left|\widehat{\beta}\right]$$

$$\left.-E\left[\left\{R(V,\widehat{\beta})-E(R(V,\widehat{\beta})|V^{T}\widehat{\beta},\widehat{\beta})\right\}^{2}\left|\widehat{\beta}\right]\right]$$

$$\gtrsim (\beta_{0}-\widehat{\beta})^{T}E\left[\left(V-E(V|V^{T}\widehat{\beta},\widehat{\beta})\right)\left(V-E(V|V^{T}\widehat{\beta},\widehat{\beta})\right)^{T}\left|\widehat{\beta}\right]$$

$$\times(\beta_{0}-\widehat{\beta})-C_{R}\|\widehat{\beta}-\beta_{0}\|^{4}$$

$$= (\beta_{0}-\widehat{\beta})^{T}\operatorname{var}\left[\left(V-E(V|V^{T}\widehat{\beta})\right)\left|\widehat{\beta}\right](\beta_{0}-\widehat{\beta})-C_{R}\|\widehat{\beta}-\beta_{0}\|^{4}.$$
(S.75)

Therefore, it is left to verify that with probability arbitrarily large,

$$(\boldsymbol{\beta}_0 - \hat{\boldsymbol{\beta}})^T \operatorname{var}\left[\left(V - E(V|V^T \hat{\boldsymbol{\beta}})\right) | \hat{\boldsymbol{\beta}}\right] (\boldsymbol{\beta}_0 - \hat{\boldsymbol{\beta}}) \gtrsim \| \hat{\boldsymbol{\beta}} - \boldsymbol{\beta}_0 \|^2.$$

To this end, note that based on Condition 6, there exists an orthonormal matrix P (may depend on $\hat{\beta}$), and eigenvalues $0 < \lambda_2(\hat{\beta}) \leq \ldots \leq \lambda_p(\hat{\beta})$ such that

$$\operatorname{var}\left[\left(V - E(V|V^{T}\widehat{\boldsymbol{\beta}})\right) |\widehat{\boldsymbol{\beta}}\right]$$

= $P^{T}\operatorname{diag}(0, \lambda_{2}(\widehat{\boldsymbol{\beta}}), \dots, \lambda_{p}(\widehat{\boldsymbol{\beta}}))P$
= $\left(\widehat{\boldsymbol{\beta}}, P_{2:p}^{T}\right)\operatorname{diag}\left(0, \lambda_{2}(\widehat{\boldsymbol{\beta}}), \dots, \lambda_{p}(\widehat{\boldsymbol{\beta}})\right) \left(\widehat{\boldsymbol{\beta}}, P_{2:p}^{T}\right)^{T}$
= $\bar{P}^{T}\operatorname{diag}\left(0, \lambda_{2}(\widehat{\boldsymbol{\beta}}), \dots, \lambda_{p}(\widehat{\boldsymbol{\beta}})\right)\bar{P},$ (S.76)

where $P_{2:p}$ denotes the matrix formed by the 2nd to pth row of P, $\overline{P} = (\hat{\beta}, P_{2:p}^T)^T$. We can verify that \overline{P} is still orthonormal; this is because that the first row of \overline{P} is $\hat{\beta}^T$, which satisfies $\|\hat{\beta}\|_2 = 1$, and

$$0 = \widehat{\boldsymbol{\beta}}^{T} \operatorname{var} \left[\left(V - E(V|V^{T}\widehat{\boldsymbol{\beta}}) \right) |\widehat{\boldsymbol{\beta}} \right] \widehat{\boldsymbol{\beta}} = \widehat{\boldsymbol{\beta}}^{T} \overline{P}^{T} \operatorname{diag} \left(0, \lambda_{2}(\widehat{\boldsymbol{\beta}}), \dots, \lambda_{p}(\widehat{\boldsymbol{\beta}}) \right) \overline{P} \widehat{\boldsymbol{\beta}} \ge \lambda_{2}(\widehat{\boldsymbol{\beta}}) \left\| \overline{P}_{2:p} \widehat{\boldsymbol{\beta}} \right\|_{2}^{2},$$

which concludes $\bar{P}_{2:p}\hat{\boldsymbol{\beta}} = 0$. Denote by $a_i, b_i, i = 1, \dots, p$, the entries of $\bar{P}\boldsymbol{\beta}_0$ and $\bar{P}\hat{\boldsymbol{\beta}}$ respectively; then we have $a_1 = \hat{\boldsymbol{\beta}}^T \boldsymbol{\beta}_0 > 0.5$ with probability arbitrarily large for sufficiently large n, and $b_1 = 1 > 0.5$; and $\sum_i a_i^2 = 1$, $\sum_i b_i^2 = 1$. Then, denoting by \bar{P}_1 the first row of \bar{P} , we have

$$\left\{ \bar{P}_{1} \left(\boldsymbol{\beta}_{0} - \hat{\boldsymbol{\beta}} \right) \right\}^{2}$$

$$= \left(a_{1} - b_{1} \right)^{2} = \left(\frac{a_{1}^{2} - b_{1}^{2}}{a_{1} + b_{1}} \right)^{2}$$

$$\leq \left\{ (a_{2}^{2} - b_{2}^{2}) + \ldots + (a_{p}^{2} - b_{p}^{2}) \right\}^{2}$$

$$= \left\{ (a_{2} - b_{2}, \ldots, a_{p} - b_{p}) \cdot (a_{2} + b_{2}, \ldots, a_{p} + b_{p})^{T} \right\}^{2}$$

$$\leq \left\{ (a_{2} - b_{2})^{2} + \ldots + (a_{p} - b_{p})^{2} \right\} \cdot \left\{ (a_{2} + b_{2})^{2} + (a_{p} + b_{p})^{2} \right\}$$

$$\leq \left(a_{2} - b_{2})^{2} + \ldots + (a_{p} - b_{p})^{2} \right\} \cdot \left\{ (a_{2} + b_{2})^{2} + (a_{p} + b_{p})^{2} \right\}$$

$$= \left\| \bar{P}_{2:p} \left(\boldsymbol{\beta}_{0} - \hat{\boldsymbol{\beta}} \right) \right\|_{2}^{2}.$$

$$(S.77)$$

Combining (S.76) with (S.77), and noting that \overline{P} is orthonormal, we have

$$(\boldsymbol{\beta}_{0} - \boldsymbol{\hat{\beta}})^{T} \operatorname{var} \left[\left(V - E(V|V^{T} \boldsymbol{\hat{\beta}}) \right) \left| \boldsymbol{\hat{\beta}} \right] (\boldsymbol{\beta}_{0} - \boldsymbol{\hat{\beta}}) \right]$$

$$\geq \lambda_{2}(\boldsymbol{\hat{\beta}}) \left\| \bar{P}_{2:p}(\boldsymbol{\beta}_{0} - \boldsymbol{\hat{\beta}}) \right\|_{2}^{2} \geq \left\| \bar{P}(\boldsymbol{\beta}_{0} - \boldsymbol{\hat{\beta}}) \right\|_{2}^{2} = \left\| \boldsymbol{\beta}_{0} - \boldsymbol{\hat{\beta}} \right\|_{2}^{2}$$

where we have used Condition 6. This together with (S.75) and (S.73) completes the proof of Part (b). We complete the proof of Theorem 1.

5 Technical Details for Lemma 1, Theorem 2, and Corollary 1

This section is composed of five subsections. Section 5.1 gives the proof of Lemma 1; Sections 5.2-5.4 presents proof of Theorem 2 Parts (1)–(3); Section 5.5 proves Corollary 1.

5.1 Proof of Lemma 1

Proof. The proof for Part (1), i.e., $\psi(\beta_0) = 0$, is straightforward.

We proceed to show Parts (2) and (3). For Part (2), let $V = X_1 - X_2$ and $U = Y_1 - Y_2$. We then have

$$\psi_0(\boldsymbol{\beta}) = E\left[V\left\{I(U>0) - F_{\boldsymbol{\beta}}(V^T\boldsymbol{\beta})\right\}\right],$$

and therefore based on Conditions A1 and A2, $\psi_0'(\beta)$ exists and

$$\frac{\partial \psi_{0}(\boldsymbol{\beta})}{\partial \boldsymbol{\beta}}\Big|_{\boldsymbol{\beta}=\boldsymbol{\beta}_{0}} = -E\left\{V\frac{\partial F_{\boldsymbol{\beta}}(V^{T}\boldsymbol{\beta})}{\partial \boldsymbol{\beta}}\right\}\Big|_{\boldsymbol{\beta}=\boldsymbol{\beta}_{0}}$$

$$= -E\left\{V\int\frac{\partial F_{0}(V^{T}\boldsymbol{\beta}+\boldsymbol{v}^{T}(\boldsymbol{\beta}_{0}-\boldsymbol{\beta}))}{\partial \boldsymbol{\beta}}f_{V|V^{T}}\boldsymbol{\beta}(\boldsymbol{v})d\boldsymbol{v}\right\}\Big|_{\boldsymbol{\beta}=\boldsymbol{\beta}_{0}}$$

$$-E\left\{V\int F_{0}(V^{T}\boldsymbol{\beta}+\boldsymbol{v}^{T}(\boldsymbol{\beta}_{0}-\boldsymbol{\beta}))\frac{\partial f_{V|V^{T}}\boldsymbol{\beta}(\boldsymbol{v})}{\partial \boldsymbol{\beta}}d\boldsymbol{v}\right\}\Big|_{\boldsymbol{\beta}=\boldsymbol{\beta}_{0}}$$

$$= -E\left\{V\int(V^{T}-\boldsymbol{v}^{T})f_{0}(V^{T}\boldsymbol{\beta}_{0})f_{V|V^{T}}\boldsymbol{\beta}_{0}(\boldsymbol{v})d\boldsymbol{v}\right\} + 0$$

$$= -E[V\{V^{T}-E(V^{T}|V^{T}\boldsymbol{\beta}_{0})\}f_{0}(V^{T}\boldsymbol{\beta}_{0})]$$

$$= -E[\{V-E(V|V^{T}\boldsymbol{\beta}_{0})\}\{V^{T}-E(V^{T}|V^{T}\boldsymbol{\beta}_{0})\}f_{0}(V^{T}\boldsymbol{\beta}_{0})]. \quad (S.78)$$

As a consequence, based on Conditions 6 and A1, we immediately conclude that $\psi'_0(\beta_0)$ has rank p-1. This completes our proof of Part (2).

Last, we show Part (3). Without loss of generality, assume that $\beta_{0,1} \neq 0$, i.e, the first element of β_0 is nonzero. Let

$$\check{\psi}(\boldsymbol{\beta}) = \psi_0(\boldsymbol{\beta}) + \begin{pmatrix} 0.5\boldsymbol{\beta}^T\boldsymbol{\beta} \\ \mathbf{0} \end{pmatrix}.$$

Then by the fact that $\|\beta\|_2^2 = 1$, we have $\psi_0(\beta) = \check{\psi}(\beta) - \check{\psi}(\beta_0)$; by Lemma 11 given below and the fact that $\psi_0(\beta_0) \cdot \beta_0 = 0$, we immediately conclude that

$$\check{\psi}'(oldsymbol{eta}_0) = \psi'(oldsymbol{eta}_0) + egin{pmatrix}oldsymbol{eta}_0^T\oldsymbol{0}\end{pmatrix}$$

is of full rank. Define $A(\beta_0) = \check{\psi}'(\beta_0)$; by Taylor expansion, we conclude

$$\psi_0(\boldsymbol{\beta}) = \check{\psi}(\boldsymbol{\beta}) - \check{\psi}(\boldsymbol{\beta}_0) = A(\boldsymbol{\beta}_0)(\boldsymbol{\beta} - \boldsymbol{\beta}_0) + o(\boldsymbol{\beta} - \boldsymbol{\beta}_0).$$

This completes the proof of this lemma.

Lemma 11. Assume Condition 6 and $\beta_{0,1} \neq 0$, i.e., the first component of β_0 is non-zero, then $\psi'_0(\beta_0)_{2:p}$ is of full rank.

Proof:Based on Conditions 6 and A1, and (S.78), there exist eigenvalues of $\psi'_0(\beta_0)$: 0 <

 $\lambda_2 \leq \ldots \leq \lambda_p$ and orthonormal matrix P, such that

$$\Sigma \equiv \psi'_0(\boldsymbol{\beta}_0) = P diag(0, \lambda_2, \dots, \lambda_p) P^T.$$

To show that $\psi'_0(\beta_0)_{2:p}$ is of full rank, it suffices to show that $\psi'_0(\beta_0)_{-1,-1} = \sum_{-1,-1} is$ of full rank.

Denote $\widetilde{\Lambda} = diag(\lambda_2, \lambda_2, \dots, \lambda_p)$ and $\widetilde{I}_p = diag(0, 1, \dots, 1)$. Then, we have

$$\widetilde{\Lambda}^{-1/2} P^T \Sigma P \widetilde{\Lambda}^{-1/2} = \widetilde{I}_p.$$
(S.79)

On the other hand, let U be an orthonormal matrix whose first row is given by

$$U_1 = \frac{\boldsymbol{\beta}_0^T P \widetilde{\Lambda}^{1/2}}{\sqrt{\boldsymbol{\beta}_0^T \widetilde{\Sigma} \boldsymbol{\beta}_0}},\tag{S.80}$$

where $\tilde{\Sigma} = P\tilde{\Lambda}P^T$. Let $Q = U\tilde{\Lambda}^{-1/2}P^T$; then its first row is $Q_1 = \frac{\beta_0^T}{\sqrt{\beta_0^T\tilde{\Sigma}\beta_0}}$ and $Q^{-1} = P\tilde{\Lambda}^{1/2}U^T$. We consider

$$\begin{split} \Sigma &= \operatorname{var}\left(V - E(V|V^{T}\boldsymbol{\beta}_{0})\right) = Q^{-1}\operatorname{var}\left(QV - E(QV|V^{T}\boldsymbol{\beta}_{0})\right)(Q^{-1})^{T} \\ &= Q^{-1}\operatorname{var}\left(\left(\frac{\boldsymbol{\beta}_{0}^{T}\boldsymbol{\Sigma}}{\sqrt{\boldsymbol{\beta}_{0}^{T}}\tilde{\boldsymbol{\Sigma}}\boldsymbol{\beta}_{0}}\right) - E\left(\left(\frac{\boldsymbol{\beta}_{0}^{T}\boldsymbol{V}}{\sqrt{\boldsymbol{\beta}_{0}^{T}}\tilde{\boldsymbol{\Sigma}}\boldsymbol{\beta}_{0}}\right)\right)|V^{T}\boldsymbol{\beta}_{0}\right)\right)(Q^{-1})^{T} \\ &= Q^{-1}\operatorname{var}\left(\begin{pmatrix}0\\Q_{2:p}V\\Q_{2$$

and therefore

$$\begin{split} \Sigma_{-1,-1} &= (Q^{-1})_{-1,-1} U_{2:p} \widetilde{I}_p U_{2:p}^T (Q^{-1})_{-1,-1}^T \\ &= P_{2:p} \widetilde{\Lambda}^{1/2} U_{2:p}^T U_{2:p} \widetilde{I}_p U_{2:p}^T U_{2:p} \widetilde{\Lambda}^{1/2} P_{2:p}^T \end{split}$$

Our goal is to show that $\Sigma_{-1,-1}$ is of full rank, and we prove this by contradiction. Suppose otherwise, then there exists $\mathbf{0} \neq \mathbf{x} \in \mathbb{R}^{p-1}$ such that $\mathbf{x}^T \Sigma_{-1,-1} \mathbf{x} = 0$, which is equivalent to

$$0 = \boldsymbol{x}^{T} P_{2:p} \widetilde{\Lambda}^{1/2} U_{2:p}^{T} U_{2:p} \widetilde{I}_{p} U_{2:p}^{T} U_{2:p} \widetilde{\Lambda}^{1/2} P_{2:p}^{T} \boldsymbol{x} = \| \widetilde{I}_{p} U_{2:p}^{T} U_{2:p} \widetilde{\Lambda}^{1/2} P_{2:p}^{T} \boldsymbol{x} \|_{2}^{2},$$

which is further equivalent to

$$\mathbf{0} = \widetilde{I}_p U_{2:p}^T U_{2:p} \widetilde{\Lambda}^{1/2} P_{2:p}^T \boldsymbol{x}$$
$$= \begin{pmatrix} 0\\ U_{-1,-1}^T U_{2:p} \widetilde{\Lambda}^{1/2} P_{2:p}^T \boldsymbol{x} \end{pmatrix}$$

and therefore

$$U_{-1,-1}^{T}U_{2:p}\widetilde{\Lambda}^{1/2}P_{2:p}^{T}\boldsymbol{x} = \boldsymbol{0}.$$
 (S.81)

Note that U is orthonormal, its determinant det(U) = 1 or -1, and since $U^T = U^{-1} = \frac{1}{det(U)}U^*$, where U^* denotes the adjoint matrix of U, its (1,1)th entry is $det(U_{-1,-1})$. As a consequence,

$$det(U_{-1,-1}) = U_{1,1} \cdot det(U) = \frac{\boldsymbol{\beta}_0^T P(\sqrt{\lambda_2}, 0, \dots, 0)^T}{\sqrt{\boldsymbol{\beta}_0^T \widetilde{\Sigma} \boldsymbol{\beta}_0}} \cdot det(U) \neq 0$$
(S.82)

if

$$\boldsymbol{\beta}_0^T P(\sqrt{\lambda_2}, 0, \dots, 0)^T \neq 0.$$
(S.83)

For presentational continuity, we postpone the proof of (S.83) to the end of the proof for this lemma. (S.82) indicates that $U_{-1,-1}$ is invertible, and this together with (S.81) gives

$$U_{2:p}\widetilde{\Lambda}^{1/2}P_{2:p}^T\boldsymbol{x}=\boldsymbol{0}.$$

This together with the fact that U is orthonormal and the definition of U_1 in (S.80) gives

$$0 = \boldsymbol{x}^{T} P_{2:p} \widetilde{\Lambda}^{1/2} U_{2:p}^{T} U_{2:p} \widetilde{\Lambda}^{1/2} P_{2:p}^{T} \boldsymbol{x}$$

$$= \boldsymbol{x}^{T} P_{2:p} \widetilde{\Lambda}^{1/2} (I_{p} - U_{1}^{T} U_{1}) \widetilde{\Lambda}^{1/2} P_{2:p}^{T} \boldsymbol{x}$$

$$= \boldsymbol{x}^{T} \widetilde{\Sigma}_{-1,-1} \boldsymbol{x} - \boldsymbol{x}^{T} P_{2:p} \widetilde{\Lambda}^{1/2} \frac{\widetilde{\Lambda}^{1/2} P^{T} \boldsymbol{\beta}_{0}}{\sqrt{\boldsymbol{\beta}_{0}^{T} \widetilde{\Sigma} \boldsymbol{\beta}_{0}}} \frac{\boldsymbol{\beta}_{0}^{T} P \widetilde{\Lambda}^{1/2}}{\sqrt{\boldsymbol{\beta}_{0}^{T} \widetilde{\Sigma} \boldsymbol{\beta}_{0}}} \widetilde{\Lambda}^{1/2} P_{2:p}^{T} \boldsymbol{x},$$

which leads to

$$\boldsymbol{x}^{T} \widetilde{\Sigma}_{-1,-1} \boldsymbol{x} = \boldsymbol{x}^{T} P_{2:p} \widetilde{\Lambda}^{1/2} \frac{\widetilde{\Lambda}^{1/2} P^{T} \boldsymbol{\beta}_{0}}{\sqrt{\boldsymbol{\beta}_{0}^{T} \widetilde{\Sigma} \boldsymbol{\beta}_{0}}} \frac{\boldsymbol{\beta}_{0}^{T} P \widetilde{\Lambda}^{1/2}}{\sqrt{\boldsymbol{\beta}_{0}^{T} \widetilde{\Sigma} \boldsymbol{\beta}_{0}}} \widetilde{\Lambda}^{1/2} P_{2:p}^{T} \boldsymbol{x}$$

$$= \frac{\left(\boldsymbol{x}^{T} P_{2:p} \widetilde{\Lambda}^{1/2} \widetilde{\Lambda}^{1/2} P^{T} \boldsymbol{\beta}_{0}\right)^{2}}{\boldsymbol{\beta}_{0}^{T} \widetilde{\Sigma} \boldsymbol{\beta}_{0}}$$

$$\leqslant \frac{\|\widetilde{\Lambda}^{1/2} P_{2:p}^{T} \boldsymbol{x}\|_{2}^{2} \cdot \|\widetilde{\Lambda}^{1/2} P^{T} \boldsymbol{\beta}_{0}\|_{2}^{2}}{\boldsymbol{\beta}_{0}^{T} \widetilde{\Sigma} \boldsymbol{\beta}_{0}} = \boldsymbol{x}^{T} \widetilde{\Sigma}_{-1,-1} \boldsymbol{x}, \quad (S.84)$$

where " \leq " is from Cauchy's inequality, and "=" holds if and only if

$$\widetilde{\Lambda}^{1/2} P_{2:p}^T \boldsymbol{x} = a \cdot \widetilde{\Lambda}^{1/2} P^T \boldsymbol{\beta}_0$$
(S.85)

for some $a \in \mathbb{R}$. Therefore, (S.84) introduces a contradiction, if we can show that (S.85) cannot hold. In fact, if

$$\widetilde{\Lambda}^{1/2} P_{2:p}^T \boldsymbol{x} = a \cdot \widetilde{\Lambda}^{1/2} P^T \boldsymbol{\beta}_0$$

then

$$\widetilde{\Lambda}^{1/2} P^T \left\{ (0, \boldsymbol{x}^T)^T - a \cdot \boldsymbol{\beta}_0 \right\} = 0,$$

which leads to $(0, \mathbf{x}^T)^T = a \cdot \boldsymbol{\beta}_0$, since $\tilde{\Lambda}^{1/2} P^T$ is invertible. Recall that the first element of $\boldsymbol{\beta}_0$ is nonzero by assumption, so we must have a = 0, which further implies that $\mathbf{x} = \mathbf{0}$; and this contradicts $\mathbf{x} \neq 0$.

Finally, we verify (S.82). Consider

$$\boldsymbol{\beta}_0^T P \Lambda P^T \boldsymbol{\beta}_0 = \boldsymbol{\beta}_0^T \Sigma \boldsymbol{\beta}_0 = 0.$$
(S.86)

On the other hand, since $\tilde{\Sigma}$ is of full rank,

$$\boldsymbol{\beta}_0^T P \tilde{\boldsymbol{\Lambda}} P^T \boldsymbol{\beta}_0 = \boldsymbol{\beta}_0^T \tilde{\boldsymbol{\Sigma}} \boldsymbol{\beta}_0 > 0.$$
(S.87)

Combining (S.86) with (S.87) leads to

$$(\boldsymbol{\beta}_0^T P(\sqrt{\lambda_2}, 0, \dots, 0)^T)^2 = \boldsymbol{\beta}_0^T P \cdot diag(\lambda_2, 0, \dots, 0) \cdot P^T \boldsymbol{\beta}_0$$
$$= \boldsymbol{\beta}_0^T P(\widetilde{\Lambda} - \Lambda) P^T \boldsymbol{\beta}_0 > 0,$$

giving (S.82). We have completed the proof of this lemma.

5.2 **Proof of Theorem 2 Part (1)**

In this section, we show the existence of the zero-crossing for $\psi_n(\beta)$. We need the following lemma, which establishes the L_2 consistency of \hat{F}_{β} . Let

$$D_{\beta}(F_2, F_1) = \left[\int \int \left\{ F_1((\boldsymbol{x}_1 - \boldsymbol{x}_2)^T \boldsymbol{\beta}) - F_2((\boldsymbol{x}_1 - \boldsymbol{x}_2)^T \boldsymbol{\beta}) \right\}^2 dF_X(\boldsymbol{x}_1) dF_X(\boldsymbol{x}_2) \right]^{1/2}$$

Lemma 12. Assume Conditions 0–2 and Condition A1. We have

$$\sup_{\boldsymbol{\beta}\in\boldsymbol{\mathcal{B}}} D_{\boldsymbol{\beta}}(\hat{F}_{\boldsymbol{\beta}}, F_{\boldsymbol{\beta}}) = O_p(n^{-1/3}),$$

where we recall that \hat{F}_{β} is defined by (3.4) in the main article.

Proof. The proof follows the same lines as that of Theorem 1 Part (a); the details are omitted. ■

We return to the proof of Theorem 2 Part (1). Consider

$$\psi_{n}(\boldsymbol{\beta}) = \int \int (\boldsymbol{x}_{1} - \boldsymbol{x}_{2}) \left\{ I(y_{1} > y_{2}) - \widehat{F}_{\boldsymbol{\beta}}((\boldsymbol{x}_{1} - \boldsymbol{x}_{2})^{T}\boldsymbol{\beta}) \right\} \\ \times d\mathbb{F}_{X,Y}(\boldsymbol{x}_{1}, y_{1})d\mathbb{F}_{X,Y}(\boldsymbol{x}_{2}, y_{2}) \\ = \int \int (\boldsymbol{x}_{1} - \boldsymbol{x}_{2}) \left\{ I(y_{1} > y_{2}) - F_{\boldsymbol{\beta}}((\boldsymbol{x}_{1} - \boldsymbol{x}_{2})^{T}\boldsymbol{\beta}) \right\} \\ \times d\mathbb{F}_{X,Y}(\boldsymbol{x}_{1}, y_{1})d\mathbb{F}_{X,Y}(\boldsymbol{x}_{2}, y_{2}) \\ + \int \int (\boldsymbol{x}_{1} - \boldsymbol{x}_{2}) \left\{ F_{\boldsymbol{\beta}}((\boldsymbol{x}_{1} - \boldsymbol{x}_{2})^{T}\boldsymbol{\beta}) - \widehat{F}_{\boldsymbol{\beta}}((\boldsymbol{x}_{1} - \boldsymbol{x}_{2})^{T}\boldsymbol{\beta}) \right\} \\ \times d\mathbb{F}_{X,Y}(\boldsymbol{x}_{1}, y_{1})d\mathbb{F}_{X,Y}(\boldsymbol{x}_{2}, y_{2}) \\ = I_{1} + I_{2}. \tag{S.88}$$

We consider I_1 first. Note that we can further decompose $I_1 = I_{1,1} + I_{1,2} + I_{1,3}$, where

$$I_{1,1} = \iint (\boldsymbol{x}_1 - \boldsymbol{x}_2) \left\{ I(y_1 > y_2) - F_{\boldsymbol{\beta}}((\boldsymbol{x}_1 - \boldsymbol{x}_2)^T \boldsymbol{\beta}) \right\} \\ \times d \left\{ \mathbb{F}_{X,Y}(\boldsymbol{x}_1, y_1) - F_{X,Y}(\boldsymbol{x}_1, y_1) \right\} d\mathbb{F}_{X,Y}(\boldsymbol{x}_2, y_2) \\ I_{1,2} = \iint (\boldsymbol{x}_1 - \boldsymbol{x}_2) \left\{ I(y_1 > y_2) - F_{\boldsymbol{\beta}}((\boldsymbol{x}_1 - \boldsymbol{x}_2)^T \boldsymbol{\beta}) \right\} \\ \times dF_{X,Y}(\boldsymbol{x}_1, y_1) d \left\{ \mathbb{F}_{X,Y}(\boldsymbol{x}_2, y_2) - F_{X,Y}(\boldsymbol{x}_2, y_2) \right\} \\ I_{1,3} = \iint (\boldsymbol{x}_1 - \boldsymbol{x}_2) \left\{ I(y_1 > y_2) - F_{\boldsymbol{\beta}}((\boldsymbol{x}_1 - \boldsymbol{x}_2)^T \boldsymbol{\beta}) \right\} \\ \times dF_{X,Y}(\boldsymbol{x}_1, y_1) dF_{X,Y}(\boldsymbol{x}_2, y_2).$$

Without loss of generality, we assume that $I_{1,1}$, $I_{1,2}$, and $I_{1,3}$ are one-dimensional. If not, the following developments can be applied entry-wise. Furthermore, this argument is applicable to the similar developments in Sections 5.3 and 5.4.

For $I_{1,1}$, we consider the function class

$$\mathcal{J}_1 = \left\{ (\boldsymbol{x}_1 - \boldsymbol{x}_2) \left\{ I(y_1 > y_2) - F_{\boldsymbol{\beta}} ((\boldsymbol{x}_1 - \boldsymbol{x}_2)^T \boldsymbol{\beta}) \right\} : \boldsymbol{x}_2 \in \mathcal{X}, y_2 \in \mathcal{Y}, \boldsymbol{\beta} \in \boldsymbol{\beta} \right\},\$$

and with Lemma 5 and Condition A1 that $F_{\beta}(t)$ is monotone in t, it is straightforward to verify that

$$\begin{split} \sup_{g \in \mathcal{J}_1} |g| &\lesssim 1, \\ H_{q,B}(\delta, \mathcal{J}_1, F_{X,Y}) &\lesssim 1/\delta. \end{split}$$

Applying Lemma 4 leads to

$$\sup_{\boldsymbol{x}_{2}\in\mathcal{X}, y_{2}\in\mathcal{Y}, \boldsymbol{\beta}\in\mathcal{B}} \left| \int (\boldsymbol{x}_{1} - \boldsymbol{x}_{2}) \left\{ I(y_{1} > y_{2}) - F_{\boldsymbol{\beta}}((\boldsymbol{x}_{1} - \boldsymbol{x}_{2})^{T}\boldsymbol{\beta}) \right\} \\ \times d \left\{ \mathbb{F}_{X,Y}(\boldsymbol{x}_{1}, y_{1}) - F_{X,Y}(\boldsymbol{x}_{1}, y_{1}) \right\} \right|$$
$$= O_{p}(n^{-1/2}),$$

and therefore

$$\sup_{\boldsymbol{\beta}\in\mathcal{B}}|I_{1,1}|=O_p(n^{-1/2}).$$

For $I_{1,2}$, we consider the function class

$$\mathcal{J}_2 = \left\{ \int (\boldsymbol{x}_1 - \boldsymbol{x}_2) \left\{ I(y_1 > y_2) - F_{\boldsymbol{\beta}} ((\boldsymbol{x}_1 - \boldsymbol{x}_2)^T \boldsymbol{\beta}) \right\} dF_{X,Y}(\boldsymbol{x}_1, y_1) : \boldsymbol{\beta} \in \boldsymbol{\mathcal{B}} \right\}.$$

Clearly, $\sup_{g \in \mathcal{J}_2} |g| \leq 1$. With the same strategy used to compute the bracketing number for \mathcal{G}_I in Step 2 of the proof of Lemma 8, we have

$$H_{q,B}(\delta, \mathcal{J}_2, F_{X,Y}) \lesssim 1/\delta.$$

Applying Lemma 4, we have

$$\sup_{\boldsymbol{\beta}\in\mathcal{B}}I_{1,2}=O_p(n^{-1/2}).$$

Furthermore, we have $I_{1,3} = \psi_0(\beta)$. As a consequence, we have established

$$\sup_{\boldsymbol{\beta} \in \mathcal{B}} |I_1 - \psi_0(\boldsymbol{\beta})| = O_p(n^{-1/2}).$$
(S.89)

For I_2 , we can use the same strategy to decompose $I_2 = I_{2,1} + I_{2,2} + I_{2,3}$ and verify $\sup_{\beta \in \mathcal{B}} I_{2,1} = O_p(n^{-1/2})$, $\sup_{\beta \in \mathcal{B}} I_{2,2} = O_p(n^{-1/2})$. Applying the Cauchy–Schwarz inequality and Lemma 12, we have

$$\begin{aligned} \sup_{\boldsymbol{\beta} \in \mathcal{B}} |I_{2,3}| \\ &= \sup_{\boldsymbol{\beta} \in \mathcal{B}} \left| \int \int (\boldsymbol{x}_1 - \boldsymbol{x}_2) \left\{ F_{\boldsymbol{\beta}} ((\boldsymbol{x}_1 - \boldsymbol{x}_2)^T \boldsymbol{\beta}) - \hat{F}_{\boldsymbol{\beta}} ((\boldsymbol{x}_1 - \boldsymbol{x}_2)^T \boldsymbol{\beta}) \right\} \\ &\times dF_{X,Y}(\boldsymbol{x}_1, y_1) dF_{X,Y}(\boldsymbol{x}_2, y_2) \right| \\ &\lesssim \sup_{\boldsymbol{\beta} \in \mathcal{B}} D_{\boldsymbol{\beta}} (\hat{F}_{\boldsymbol{\beta}}, F_{\boldsymbol{\beta}}) = O_p(n^{-1/3}). \end{aligned}$$

As a consequence,

$$\sup_{\boldsymbol{\beta}\in\mathcal{B}}|I_2| = O_p(n^{-1/3}).$$
(S.90)

Combining (S.88)–(S.90) leads to

$$\psi_n(\boldsymbol{\beta}) = \psi_0(\boldsymbol{\beta}) + O_p(n^{-1/3}),$$
(S.91)

uniformly in $\beta \in \mathcal{B}$.

With Lemma 1, the rest of the proof for Part (1) of Theorem 2 follows the same lines as those of Part 1 of Theorem 4.1 in Groeneboom and Hendrickx (2018); the details are omitted.

5.3 **Proof of Theorem 2 Part (2)**

Similarly to the proof of Lemma 10, we establish Part (2) by showing the following:

- (i) $\psi_0(\widetilde{\boldsymbol{\beta}}) = o_p(1);$
- (ii) $\psi_0(\boldsymbol{\beta}) = 0$ implies that $\boldsymbol{\beta} = \boldsymbol{\beta}_0$;
- (iii) $\psi_0(\boldsymbol{\beta})$ is continuous in $\boldsymbol{\mathcal{B}}$.

and

(iii) holds because of Condition A1. We proceed to show (i). Since $\tilde{\beta}$ is the zero-crossing of $\psi_n(\beta)$, there exist $\tilde{\beta}^{(U)}$ and $\tilde{\beta}^{(L)}$ such that

$$\widetilde{\boldsymbol{\beta}}^{(U)} = \widetilde{\boldsymbol{\beta}} + O(1/n), \qquad \widetilde{\boldsymbol{\beta}}^{(L)} = \widetilde{\boldsymbol{\beta}} + O(1/n), \psi_n(\widetilde{\boldsymbol{\beta}}^{(U)}) \ge 0, \qquad \psi_n(\widetilde{\boldsymbol{\beta}}^{(L)}) \le 0.$$
(S.92)

As a consequence, there exists $\alpha_n = (\alpha_{n,1}, \ldots, \alpha_{n,p})^T$, $0 \leq \alpha_{n,j} \leq 1$ for $j = 1, \ldots, p$, such that

$$\boldsymbol{\alpha}_{n}^{T}\psi_{n}(\widetilde{\boldsymbol{\beta}}^{(U)}) + (1-\boldsymbol{\alpha}_{n})^{T}\psi_{n}(\widetilde{\boldsymbol{\beta}}^{(L)}) = 0.$$
(S.93)

On the other hand, applying (S.91) and based on the continuity of $\psi_0(\cdot)$, we have

$$\boldsymbol{\alpha}_{n}^{T}\psi_{n}(\widetilde{\boldsymbol{\beta}}^{(U)}) + (1-\boldsymbol{\alpha}_{n})^{T}\psi_{n}(\widetilde{\boldsymbol{\beta}}^{(L)})$$

= $\boldsymbol{\alpha}_{n}^{T}\psi_{0}(\widetilde{\boldsymbol{\beta}}^{(U)}) + (1-\boldsymbol{\alpha}_{n})^{T}\psi_{0}(\widetilde{\boldsymbol{\beta}}^{(L)}) + O_{p}(n^{-1/3})$
= $\psi_{0}(\widetilde{\boldsymbol{\beta}}) + O(n^{-1}) + O_{p}(n^{-1/3}),$

which together with (S.93) proves (i). We now show (ii). Based on the definition of $\psi_0(\beta)$,

$$\begin{split} \boldsymbol{\beta}_{0}^{T}\psi_{0}(\boldsymbol{\beta}) &= E\left[\boldsymbol{\beta}_{0}^{T}V\left\{I(U>0) - E(I(U>0)|V^{T}\boldsymbol{\beta})\right\}\right] \\ &= E\left[\boldsymbol{\beta}_{0}^{T}V\left\{F_{0}(\boldsymbol{\beta}_{0}^{T}V) - E(F_{0}(\boldsymbol{\beta}_{0}^{T}V)|V^{T}\boldsymbol{\beta})\right\}\right] \\ &= E\left[\operatorname{cov}\left\{\boldsymbol{\beta}_{0}^{T}V, F_{0}(\boldsymbol{\beta}_{0}^{T}V)\Big|V^{T}\boldsymbol{\beta}\right\}\right] \\ &= E\left[\operatorname{cov}\left\{Z_{1}, F_{0}(Z_{1})\Big|Z_{2}\right\}\right], \end{split}$$

where we denote $Z_1 = \beta_0^T V$ and $Z_2 = V^T \beta$, with $Z_1 \ge C$, for some $C > -\infty$. Using the same arguments as in the proof of Part (ii) of Lemma 4.1 in Groeneboom and Hendrickx (2018), we conclude that

$$\operatorname{cov}\left\{Z_1, F_0(Z_1) \middle| Z_2\right\} \ge 0,$$

almost surely. As a consequence, if there exists some $\beta \neq \beta_0$ such that $\psi_0(\beta) = 0$, we must have

$$\operatorname{cov}\left\{\boldsymbol{\beta}_{0}^{T}V, F_{0}(\boldsymbol{\beta}_{0}^{T}V)\middle|V^{T}\boldsymbol{\beta}\right\} = 0$$

almost surely, which contradicts Condition A3. This completes the proof of (ii).

5.4 **Proof of Theorem 2 Part (3)**

Recall the definition of φ_{β} in (S.3). For any piecewise constant distribution function F with finitely many jumps $\tau_1 < \tau_2 < \ldots$, we define

$$\overline{\varphi}_{\boldsymbol{\beta},F}(t) = \begin{cases} \varphi_{\boldsymbol{\beta}}(\tau_i) & \text{if } F_{\boldsymbol{\beta}}(t) > F(\tau_i), \ t \in [\tau_i, \tau_{i+1}) \\ \varphi_{\boldsymbol{\beta}}(s) & \text{if } F_{\boldsymbol{\beta}}(t) = F(s), \ \text{for some } s \in [\tau_i, \tau_{i+1}) \\ \varphi_{\boldsymbol{\beta}}(\tau_{i+1}) & \text{if } F_{\boldsymbol{\beta}}(t) < F(\tau_i), \ t \in [\tau_i, \tau_{i+1}) \end{cases}.$$

Then, we have the following lemma.

Lemma 13. Assume Conditions 1, A1, and A2. For $\beta \in \beta$ and any piecewise constant distribution function F with finitely many jumps $\tau_1 < \tau_2 < \ldots$, there exists a constant C > 0 not depending on F and β such that

$$\left|\varphi_{\boldsymbol{\beta}}(t) - \overline{\varphi}_{\boldsymbol{\beta},F}(t)\right| \leq C|F(t) - F_{\boldsymbol{\beta}}(t)|,$$

where if $\varphi_{\beta}(\cdot)$ and $\overline{\varphi}_{\beta,F}(\cdot)$ are multi-dimensional, this inequality holds for $\varphi_{\beta}(t) - \overline{\varphi}_{\beta,F}(t)$ entry-wise.

Proof. Without loss of generality, assume that $\varphi_{\beta}(\cdot)$ and $\overline{\varphi}_{\beta,F}(\cdot)$ are one-dimensional. For

any $t \in [\tau_i, \tau_{i+1})$, we have

$$\varphi_{\boldsymbol{\beta}}(t) - \overline{\varphi}_{\boldsymbol{\beta},F}(t) = \begin{cases} \varphi_{\boldsymbol{\beta}}(t) - \varphi_{\boldsymbol{\beta}}(\tau_{i}) & \text{if } F_{\boldsymbol{\beta}}(t) > F(\tau_{i}), \ t \in [\tau_{i}, \tau_{i+1}) \\ \varphi_{\boldsymbol{\beta}}(t) - \varphi_{\boldsymbol{\beta}}(s) & \text{if } F_{\boldsymbol{\beta}}(t) = F(s), \ \text{for some } s \in [\tau_{i}, \tau_{i+1}) \\ \varphi_{\boldsymbol{\beta}}(t) - \varphi_{\boldsymbol{\beta}}(\tau_{i+1}) & \text{if } F_{\boldsymbol{\beta}}(t) < F(\tau_{i}), \ t \in [\tau_{i}, \tau_{i+1}) \end{cases}.$$

We need to show only that if $F_{\beta}(t) > F(\tau_i)$ for $t \in [\tau_i, \tau_{i+1})$,

$$\left|\varphi_{\boldsymbol{\beta}}(t) - \overline{\varphi}_{\boldsymbol{\beta},F}(t)\right| \lesssim |F_{\boldsymbol{\beta}}(t) - F(t)|$$

up to a constant not depending on F and β . The other two cases are similar. In fact, based on Conditions 1, A1, and A2,

$$\begin{aligned} \left|\varphi_{\boldsymbol{\beta}}(t) - \overline{\varphi}_{\boldsymbol{\beta},F}(t)\right| &= \left|\varphi_{\boldsymbol{\beta}}(t) - \varphi_{\boldsymbol{\beta}}(\tau_{i})\right| \lesssim |t - \tau_{i}| \\ &\leqslant \frac{1}{\inf_{\boldsymbol{\beta} \in \mathcal{B}, t \in [\tau_{i}, \tau_{i+1})} f_{\boldsymbol{\beta}}(t)} |F_{\boldsymbol{\beta}}(t) - F_{\boldsymbol{\beta}}(\tau_{i})| \lesssim |F_{\boldsymbol{\beta}}(t) - F_{\boldsymbol{\beta}}(\tau_{i})| \\ &\leqslant |F_{\boldsymbol{\beta}}(t) - F(\tau_{i})| = |F_{\boldsymbol{\beta}}(t) - F(t)|, \end{aligned}$$

where both " \leq " are up to constants not depending on β .

The next lemma establishes the bracketing numbers of the function classes

$$\mathcal{J}_{3,1} = \{ \overline{\varphi}_{\boldsymbol{\beta},F}((\boldsymbol{x}_1 - \boldsymbol{x}_2)^T \boldsymbol{\beta}) : \boldsymbol{\beta} \in \boldsymbol{\mathcal{B}}, F \in \widetilde{\boldsymbol{\mathcal{F}}} \}, \\ \mathcal{J}_{3,2} = \{ \overline{\varphi}_{\boldsymbol{\beta},F}((\boldsymbol{x}_1 - \boldsymbol{x}_2)^T \boldsymbol{\beta}) : \boldsymbol{x}_2 \in \boldsymbol{\mathcal{X}}, \boldsymbol{\beta} \in \boldsymbol{\mathcal{B}}, F \in \widetilde{\boldsymbol{\mathcal{F}}} \},$$

where $\widetilde{\mathcal{F}}$ is the class of piecewise constant distribution functions with finitely many jumps, and all jumps are uniformly bounded, i.e., there exists a constant M > 0 such that if τ is a jump for some $F \in \widetilde{F}$, then $|\tau| \leq M$.

Lemma 14. Assume Conditions 1, 2, and A2. For any arbitrary $1 \le q \le \infty$, we have

$$\begin{aligned} H_{q,B}(\delta, \mathcal{J}_{3,1}, F_{X_1, X_2}) &\lesssim 1/\delta, \\ H_{q,B}(\delta, \mathcal{J}_{3,2}, F_{X_1, X_2}) &\lesssim 1/\delta. \end{aligned}$$

Proof. Based on the definition of $\overline{\varphi}_{\beta,F}(\cdot)$, for any $\beta \in \mathcal{B}$ and $F \in \widetilde{\mathcal{F}}$ with jumps $\tau_0 \leq \ldots \leq \tau_m$, there exist $\xi_j \in [\tau_{j-1}, \tau_j], j = 1, \ldots, m$, which may depend on β and F, such that

$$\overline{\varphi}_{\boldsymbol{\beta},F}(t) = \sum_{j=1}^{m} \varphi_{\boldsymbol{\beta}}(\xi_j) I\left(t \in [\tau_{j-1}, \tau_j)\right)$$
$$= \varphi_{\boldsymbol{\beta}}(\xi_1) + \sum_{\{j:j \ge 2, t \ge \tau_j\}} \left\{\varphi_{\boldsymbol{\beta}}(\xi_j) - \varphi_{\boldsymbol{\beta}}(\xi_{j-1})\right\}$$

$$= \varphi_{\boldsymbol{\beta}}(\xi_{1}) + \sum_{\{j:j \ge 2, t \ge \tau_{j}\}} \left\{ \varphi_{\boldsymbol{\beta}}(\xi_{j}) - \varphi_{\boldsymbol{\beta}}(\xi_{j-1}) \right\}_{+} - \sum_{\{j:j \ge 2, t \ge \tau_{j}\}} \left\{ \varphi_{\boldsymbol{\beta}}(\xi_{j}) - \varphi_{\boldsymbol{\beta}}(\xi_{j-1}) \right\}_{-}$$
$$= \overline{\varphi}_{\boldsymbol{\beta},F,+}(t) - \overline{\varphi}_{\boldsymbol{\beta},F,-}(t),$$

where

$$\overline{\varphi}_{\boldsymbol{\beta},F,+}(t) = \varphi_{\boldsymbol{\beta}}(\xi_{1}) + \sum_{\{j:j\geq 2,t\geq \tau_{j}\}} \left\{ \varphi_{\boldsymbol{\beta}}(\xi_{j}) - \varphi_{\boldsymbol{\beta}}(\xi_{j-1}) \right\}_{+}$$
$$\overline{\varphi}_{\boldsymbol{\beta},F,-}(t) = \sum_{\{j:j\geq 2,t\geq \tau_{j}\}} \left\{ \varphi_{\boldsymbol{\beta}}(\xi_{j}) - \varphi_{\boldsymbol{\beta}}(\xi_{j-1}) \right\}_{-}$$

are both monotone in t. Here for any $a \in \mathbb{R}$, $a_+ = \max\{a, 0\}$, $a_- = \min\{a, 0\}$. Based on Conditions 1 and A2, and the fact that by definition all functions in $\tilde{\mathcal{F}}$ have uniformly bounded jumps, we have

$$\begin{aligned} \left| \overline{\varphi}_{\boldsymbol{\beta},F,+}(t) \right| &= \left| \varphi_{\boldsymbol{\beta}}(\xi_{1}) + \sum_{\{j:j \ge 2,t \ge \tau_{j}\}} \left\{ \varphi_{\boldsymbol{\beta}}(\xi_{j}) - \varphi_{\boldsymbol{\beta}}(\xi_{j-1}) \right\}_{+} \right| \\ &\leqslant \left| \varphi_{\boldsymbol{\beta}}(\xi_{1}) \right| + \sum_{j=1}^{m} \left| \varphi_{\boldsymbol{\beta}}(\xi_{j}) - \varphi_{\boldsymbol{\beta}}(\xi_{j-1}) \right| \\ &\lesssim \left| \varphi_{\boldsymbol{\beta}}(\xi_{1}) \right| + \sum_{j=1}^{m} \left| \xi_{j} - \xi_{j-1} \right| = \left| \varphi_{\boldsymbol{\beta}}(\xi_{1}) \right| + \xi_{m} - \xi_{1} \lesssim 1 \end{aligned}$$

uniformly in β and F, and likewise

$$\overline{\varphi}_{\boldsymbol{\beta},F,-}(t) \leq 1$$

uniformly in β and F. With Lemma 5, we can immediately conclude that

$$\mathcal{J}_{3,1,+} = \{ \overline{\varphi}_{\boldsymbol{\beta},F,+} ((\boldsymbol{x}_1 - \boldsymbol{x}_2)^T \boldsymbol{\beta}) : \boldsymbol{\beta} \in \boldsymbol{\mathcal{B}}, F \in \widetilde{\boldsymbol{\mathcal{F}}} \}$$
$$\mathcal{J}_{3,1,-} = \{ \overline{\varphi}_{\boldsymbol{\beta},F,-} ((\boldsymbol{x}_1 - \boldsymbol{x}_2)^T \boldsymbol{\beta}) : \boldsymbol{\beta} \in \boldsymbol{\mathcal{B}}, F \in \widetilde{\boldsymbol{\mathcal{F}}} \}$$

are uniformly bounded function classes and satisfy

$$H_{2,B}(\delta, \mathcal{J}_{3,1,+}, F_{X_1,X_2}) \lesssim 1/\delta$$
 and $H_{2,B}(\delta, \mathcal{J}_{3,1,-}, F_{X_1,X_2}) \lesssim 1/\delta$.

Now it is straightforward to verify that $\mathcal{J}_{3,1} \subset \mathcal{J}_{3,1,+} - \mathcal{J}_{3,1,-}$ satisfies

$$H_{q,B}(\delta, \mathcal{J}_{3,1}, F_{X_1,X_2}) \lesssim 1/\delta.$$

With the same strategy, we can verify that

$$H_{q,B}(\delta, \mathcal{J}_{3,2}, F_{X_1,X_2}) \leq 1/\delta.$$

Consider an estimator $\check{\beta}$ of β that satisfies

$$\dot{\boldsymbol{\beta}} \to \boldsymbol{\beta}_0 \tag{S.94}$$

in probability as $n \to \infty$. We have

$$\begin{split} \psi_{n}(\breve{\boldsymbol{\beta}}) &= \int \int (\boldsymbol{x}_{1} - \boldsymbol{x}_{2}) \left\{ I(y_{1} > y_{2}) - \widehat{F}_{\breve{\boldsymbol{\beta}}}((\boldsymbol{x}_{1} - \boldsymbol{x}_{2})^{T}\breve{\boldsymbol{\beta}}) \right\} \\ &\times d\mathbb{F}_{X,Y}(\boldsymbol{x}_{1}, y_{1})d\mathbb{F}_{X,Y}(\boldsymbol{x}_{2}, y_{2}) \\ &= \int \int \left\{ \boldsymbol{x}_{1} - \boldsymbol{x}_{2} - \varphi_{\breve{\boldsymbol{\beta}}}\left((\boldsymbol{x}_{1} - \boldsymbol{x}_{2})^{T}\breve{\boldsymbol{\beta}}\right) \right\} \\ &\times \left\{ I(y_{1} > y_{2}) - \widehat{F}_{\breve{\boldsymbol{\beta}}}\left((\boldsymbol{x}_{1} - \boldsymbol{x}_{2})^{T}\breve{\boldsymbol{\beta}}\right) \right\} d\mathbb{F}_{X,Y}(\boldsymbol{x}_{1}, y_{1})d\mathbb{F}_{X,Y}(\boldsymbol{x}_{2}, y_{2}) \\ &+ \int \int \left\{ \varphi_{\breve{\boldsymbol{\beta}}}\left((\boldsymbol{x}_{1} - \boldsymbol{x}_{2})^{T}\breve{\boldsymbol{\beta}}\right) - \bar{\varphi}_{\breve{\boldsymbol{\beta}},\tilde{F}_{\breve{\boldsymbol{\beta}}}}\left((\boldsymbol{x}_{1} - \boldsymbol{x}_{2})^{T}\breve{\boldsymbol{\beta}}\right) \right\} \\ &\times \left\{ I(y_{1} > y_{2}) - \widehat{F}_{\breve{\boldsymbol{\beta}}}\left((\boldsymbol{x}_{1} - \boldsymbol{x}_{2})^{T}\breve{\boldsymbol{\beta}}\right) \right\} d\mathbb{F}_{X,Y}(\boldsymbol{x}_{1}, y_{1})d\mathbb{F}_{X,Y}(\boldsymbol{x}_{2}, y_{2}) \\ &= I_{3} + I_{4}, \end{split}$$
(S.95)

where we have used the fact that

$$\int \int \bar{\varphi}_{\breve{\boldsymbol{\beta}},\widehat{F}_{\breve{\boldsymbol{\beta}}}} \left((\boldsymbol{x}_1 - \boldsymbol{x}_2)^T \breve{\boldsymbol{\beta}} \right) \left\{ I(y_1 > y_2) - \widehat{F}_{\breve{\boldsymbol{\beta}}} \left((\boldsymbol{x}_1 - \boldsymbol{x}_2)^T \breve{\boldsymbol{\beta}} \right) \right\} \\ \times d\mathbb{F}_{X,Y}(\boldsymbol{x}_1, y_1) d\mathbb{F}_{X,Y}(\boldsymbol{x}_2, y_2) = 0.$$

This is because $\bar{\varphi}_{\check{\beta},\hat{F}_{\check{\beta}}}(t)$ is a piecewise constant function with the same jumps as $\hat{F}_{\check{\beta}}(t)$; on the other hand, recall the definition of $\hat{F}_{\check{\beta}}(t)$: it is the slope of the greatest convex minorant of the corresponding cusum diagram, based on the values of $I(Y_i > Y_j)$ in the order of $(X_i - X_j)^T \check{\beta}$. Therefore, for every constant piece $[\tau_j, \tau_{j+1})$ of $\hat{F}_{\check{\beta}}(t)$,

$$\int_{(\boldsymbol{x}_1-\boldsymbol{x}_2)^T \widecheck{\boldsymbol{\beta}} \in [\tau_j,\tau_{j+1})} \left\{ I(y_1 > y_2) - \widehat{F}_{\widecheck{\boldsymbol{\beta}}} \left((\boldsymbol{x}_1 - \boldsymbol{x}_2)^T \widecheck{\boldsymbol{\beta}} \right) \right\} \\ \times d\mathbb{F}_{X,Y}(\boldsymbol{x}_1,y_1) d\mathbb{F}_{X,Y}(\boldsymbol{x}_2,y_2) = 0.$$

Without loss of generality, we assume that $\psi_n(\check{\beta})$ is one-dimensional; if not, the rest of the proof in this subsection can be applied entry-wise.

We derive the asymptotic properties of I_3 and I_4 given in (S.95) separately. We consider I_4 first. We will show that

$$I_4 = O_p(n^{-2/3}), (S.96)$$

using a strategy similar to that in the proof of Lemma 8. Denote

$$g_{4,\boldsymbol{\beta},F}(\boldsymbol{x}_{1}-\boldsymbol{x}_{2},y_{1}-y_{2})$$

$$=\left\{\varphi_{\boldsymbol{\beta}}\left((\boldsymbol{x}_{1}-\boldsymbol{x}_{2})^{T}\boldsymbol{\beta}\right)-\overline{\varphi}_{\boldsymbol{\beta},F}\left((\boldsymbol{x}_{1}-\boldsymbol{x}_{2})^{T}\boldsymbol{\beta}\right)\right\}$$

$$\times\left\{I(y_{1}>y_{2})-F\left((\boldsymbol{x}_{1}-\boldsymbol{x}_{2})^{T}\boldsymbol{\beta}\right)\right\}.$$
(S.97)

Then

$$I_4 = \int \int g_{4,\widetilde{\boldsymbol{\beta}},\widehat{F}_{\widetilde{\boldsymbol{\beta}}}}(\boldsymbol{x}_1 - \boldsymbol{x}_2, y_1 - y_2) d\mathbb{F}_{X,Y}(\boldsymbol{x}_1, y_1) d\mathbb{F}_{X,Y}(\boldsymbol{x}_2, y_2).$$
(S.98)

Consider the function class

$$\mathcal{J}_4 = \left\{ g_{4,\boldsymbol{\beta},F}(\boldsymbol{x}_1 - \boldsymbol{x}_2, y_1 - y_2) : \boldsymbol{x}_2 \in \mathcal{X}, y_2 \in \mathcal{Y}, \boldsymbol{\beta} \in \mathcal{B}, F \in \widetilde{\mathcal{F}} \right\}.$$

With Conditions 1 and A2, we can verify that for any $\beta_1, \beta_2 \in \mathcal{B}$, we have

$$\left| arphi_{oldsymbol{eta}_1}((oldsymbol{x}_1 - oldsymbol{x}_2)oldsymbol{eta}_1) - arphi_{oldsymbol{eta}_2}((oldsymbol{x}_1 - oldsymbol{x}_2)oldsymbol{eta}_2)
ight| \lesssim \|oldsymbol{eta}_1 - oldsymbol{eta}_2\|_2.$$

Based on Theorem 9.23 in Kosorok (2008), we immediately conclude that the funciton class

$$\mathcal{J}_{4,1} = \left\{ \varphi_{\boldsymbol{\beta}} ((\boldsymbol{x}_1 - \boldsymbol{x}_2)^T \boldsymbol{\beta}) : \boldsymbol{\beta} \in \boldsymbol{\beta} \right\}$$

satisfies for any $1 \leq q \leq \infty$,

$$H_{q,B}(\delta, \mathcal{J}_{4,1}, F_{X_1,X_2}) \lesssim -\log \delta_{q}$$

up to a constant depending only on q. This, together with Lemma 5 and Lemma 14, leads to

$$H_{q,B}(\delta, \mathcal{J}_4, F_{X_1,X_2}) \lesssim 1/\delta.$$

With the same strategy as in the derivation of (S.55), we can show that

$$\int \left| \int g_{4,\widetilde{\boldsymbol{\beta}},\widehat{F}_{\widetilde{\boldsymbol{\beta}}}}(\boldsymbol{x}_{1} - \boldsymbol{x}_{2}, y_{1} - y_{2}) d \left\{ \mathbb{F}_{X,Y}(\boldsymbol{x}_{1}, y_{1}) - F_{X,Y}(\boldsymbol{x}_{1}, y_{1}) \right\} \right|$$
(S.99)
$$\times d\mathbb{F}_{X,Y}(\boldsymbol{x}_{2}, y_{2})$$
$$= O_{p}(n^{-2/3}) \lor O_{p}(n^{-1/2}) \frac{1}{n} \sum_{i=1}^{n} \left\| g_{4,\widetilde{\boldsymbol{\beta}},\widehat{F}_{\widetilde{\boldsymbol{\beta}}}}(\boldsymbol{x}_{1} - X_{i}, y_{1} - Y_{i}) \right\|_{2,F_{X,Y}}^{1/2}$$

and

$$\frac{1}{n}\sum_{i=1}^{n} \left\| g_{4,\widetilde{\boldsymbol{\beta}},\widehat{\boldsymbol{F}}_{\widetilde{\boldsymbol{\beta}}}}(\boldsymbol{x}_{1}-X_{i},y_{1}-Y_{i}) \right\|_{2,F_{X,Y}}^{1/2} \leq \left\{ \frac{1}{n}\sum_{i=1}^{n} \left\| g_{4,\widetilde{\boldsymbol{\beta}},\widehat{\boldsymbol{F}}_{\widetilde{\boldsymbol{\beta}}}}(\boldsymbol{x}_{1}-X_{i},y_{1}-Y_{i}) \right\|_{2,F_{X,Y}} \right\}^{1/2} \leq \left\{ \frac{1}{n}\sum_{i=1}^{n} \left\| \widehat{\boldsymbol{F}}_{\widetilde{\boldsymbol{\beta}}}((\boldsymbol{x}_{1}-X_{i})^{T}\widetilde{\boldsymbol{\beta}}) - F_{\widetilde{\boldsymbol{\beta}}}((\boldsymbol{x}_{1}-X_{i})^{T}\widetilde{\boldsymbol{\beta}}) \right\|_{2,F_{X,Y}} \right\}^{1/2} = \left\{ \frac{1}{n}\sum_{i=1}^{n} g_{5,\widetilde{\boldsymbol{\beta}},\widehat{\boldsymbol{F}}_{\widetilde{\boldsymbol{\beta}}}}(X_{i}) \right\}^{1/2}, \qquad (S.100)$$

where

$$g_{5,\boldsymbol{\beta},F}(\boldsymbol{x}_{2}) = \left\| F((\boldsymbol{x}_{1} - \boldsymbol{x}_{2})^{T}\boldsymbol{\beta}) - F_{\boldsymbol{\beta}}((\boldsymbol{x}_{1} - \boldsymbol{x}_{2})^{T}\boldsymbol{\beta}) \right\|_{2,F_{X,Y}} \\ = \left[\int \left\{ F((\boldsymbol{x}_{1} - \boldsymbol{x}_{2})^{T}\boldsymbol{\beta}) - F_{\boldsymbol{\beta}}((\boldsymbol{x}_{1} - \boldsymbol{x}_{2})^{T}\boldsymbol{\beta}) \right\}^{2} dF_{X}(\boldsymbol{x}_{1}) \right]^{1/2};$$

and to derive " \leq " we have applied Lemma 13. With the same development as Part (3) of Lemma 5, we can show that the function class

$$\mathcal{J}_{5} = \left\{ g_{5,\boldsymbol{\beta},F}(\boldsymbol{x}_{2}) : \boldsymbol{\beta} \in \boldsymbol{\mathcal{B}}, F \in \widetilde{F} \right\}$$

satisfies for any arbitrary $1\leqslant q\leqslant\infty$,

$$H_{q,B}(\delta, \mathcal{J}_5, F_X) \leq 1/\delta,$$

with " \leq " being up to a universal constant depending only on p and q. Therefore, applying Lemma 9 given at the beginning of the proof of Lemma 8, we have

$$\frac{1}{n}\sum_{i=1}^{n}g_{5,\breve{\beta},\widetilde{F}_{\breve{\beta}}}(X_{i}) = \int g_{5,\breve{\beta},\widetilde{F}_{\breve{\beta}}}(\boldsymbol{x}_{2})dF_{X}(\boldsymbol{x}_{2}) + O_{p}(n^{-1/2})$$

$$= \int \left[\int \left\{\widehat{F}_{\breve{\beta}}((\boldsymbol{x}_{1}-\boldsymbol{x}_{2})^{T}\breve{\beta}) - F_{\breve{\beta}}((\boldsymbol{x}_{1}-\boldsymbol{x}_{2})^{T}\breve{\beta})\right\}^{2}dF_{X}(\boldsymbol{x}_{1})\right]^{1/2}dF_{X}(\boldsymbol{x}_{2})$$

$$+ O_{p}(n^{-1/2})$$

$$\leq \left[\int \int \left\{\widehat{F}_{\breve{\beta}}((\boldsymbol{x}_{1}-\boldsymbol{x}_{2})^{T}\breve{\beta}) - F_{\breve{\beta}}((\boldsymbol{x}_{1}-\boldsymbol{x}_{2})^{T}\breve{\beta})\right\}^{2}dF_{X}(\boldsymbol{x}_{1})dF_{X}(\boldsymbol{x}_{2})\right]^{1/2}$$

$$+ O_{p}(n^{-1/2})$$

$$= D_{\widetilde{\boldsymbol{\beta}}}(\widehat{F}_{\widetilde{\boldsymbol{\beta}}}, F_{\widetilde{\boldsymbol{\beta}}}) + O_p(n^{-1/2})$$

$$= O_p(n^{-1/3}),$$

by Lemma 12. This, together with (S.99) and (S.100), leads to

$$\int \left| \int g_{4,\widecheck{\boldsymbol{\beta}},\widehat{F}_{\widecheck{\boldsymbol{\beta}}}}(\boldsymbol{x}_{1}-\boldsymbol{x}_{2},y_{1}-y_{2})d\left\{ \mathbb{F}_{X,Y}(\boldsymbol{x}_{1},y_{1})-F_{X,Y}(\boldsymbol{x}_{1},y_{1})\right\} \right| \times d\mathbb{F}_{X,Y}(\boldsymbol{x}_{2},y_{2}) = O_{p}(n^{-2/3}).$$
(S.101)

Furthermore, using a development similar to the proof of (S.44), we can derive

$$\int \int g_{4,\breve{\beta},\widetilde{F}_{\breve{\beta}}}(\boldsymbol{x}_{1}-\boldsymbol{x}_{2},y_{1}-y_{2})dF_{X,Y}(\boldsymbol{x}_{1},y_{1})d\left\{\mathbb{F}_{X,Y}(\boldsymbol{x}_{2},y_{2})-F_{X,Y}(\boldsymbol{x}_{2},y_{2})\right\} \\ = O_{p}(n^{-2/3}),$$

which together with (S.98), (S.101), and the definition of $g_{4,\beta,F}(\cdot,\cdot)$ given in (S.97) leads to

$$\begin{split} |I_4| &= \left| \int \int g_{4,\widetilde{\boldsymbol{\beta}},\widehat{\boldsymbol{F}}_{\widetilde{\boldsymbol{\beta}}}}(\boldsymbol{x}_1 - \boldsymbol{x}_2, y_1 - y_2) dF_{X,Y}(\boldsymbol{x}_1, y_1) dF_{X,Y}(\boldsymbol{x}_2, y_2) + O_p(n^{-2/3}) \right| \\ &\leqslant \left| \int \int \left\{ \varphi_{\widetilde{\boldsymbol{\beta}}}\left((\boldsymbol{x}_1 - \boldsymbol{x}_2)^T \widetilde{\boldsymbol{\beta}} \right) - \bar{\varphi}_{\widetilde{\boldsymbol{\beta}},\widehat{\boldsymbol{F}}_{\widetilde{\boldsymbol{\beta}}}}\left((\boldsymbol{x}_1 - \boldsymbol{x}_2)^T \widetilde{\boldsymbol{\beta}} \right) \right\} \\ &\times \left\{ I(y_1 > y_2) - \widehat{\boldsymbol{F}}_{\widetilde{\boldsymbol{\beta}}}\left((\boldsymbol{x}_1 - \boldsymbol{x}_2)^T \widetilde{\boldsymbol{\beta}} \right) \right\} dF_{X,Y}(\boldsymbol{x}_1, y_1) dF_{X,Y}(\boldsymbol{x}_2, y_2) \right| \\ &\quad + O_p(n^{-2/3}) \\ &= \left| \int \int \left\{ \varphi_{\widetilde{\boldsymbol{\beta}}}\left((\boldsymbol{x}_1 - \boldsymbol{x}_2)^T \widetilde{\boldsymbol{\beta}} \right) - \bar{\varphi}_{\widetilde{\boldsymbol{\beta}},\widehat{\boldsymbol{F}}_{\widetilde{\boldsymbol{\beta}}}}\left((\boldsymbol{x}_1 - \boldsymbol{x}_2)^T \widetilde{\boldsymbol{\beta}} \right) \right\} \\ &\times \left\{ F_{\widetilde{\boldsymbol{\beta}}}\left((\boldsymbol{x}_1 - \boldsymbol{x}_2)^T \widetilde{\boldsymbol{\beta}} \right) - \widehat{\boldsymbol{F}}_{\widetilde{\boldsymbol{\beta}}}\left((\boldsymbol{x}_1 - \boldsymbol{x}_2)^T \widetilde{\boldsymbol{\beta}} \right) \right\} dF_X(\boldsymbol{x}_1) dF_X(\boldsymbol{x}_2) \right| \\ &\quad + O_p(n^{-2/3}) \\ &\leqslant \left\| \varphi_{\widetilde{\boldsymbol{\beta}}}\left((\boldsymbol{x}_1 - \boldsymbol{x}_2)^T \widetilde{\boldsymbol{\beta}} \right) - \bar{\varphi}_{\widetilde{\boldsymbol{\beta}},\widehat{\boldsymbol{F}}_{\widetilde{\boldsymbol{\beta}}}}\left((\boldsymbol{x}_1 - \boldsymbol{x}_2)^T \widetilde{\boldsymbol{\beta}} \right) \right\|_{2,F_{X_1,X_2}} \\ &\times \left\| F_{\widetilde{\boldsymbol{\beta}}}\left((\boldsymbol{x}_1 - \boldsymbol{x}_2)^T \widetilde{\boldsymbol{\beta}} \right) - \widehat{\boldsymbol{F}}_{\widetilde{\boldsymbol{\beta}}}\left((\boldsymbol{x}_1 - \boldsymbol{x}_2)^T \widetilde{\boldsymbol{\beta}} \right) \right\|_{2,F_{X_1,X_2}} + O_p(n^{-2/3}) \\ &\lesssim \left\| F_{\widetilde{\boldsymbol{\beta}}}\left((\boldsymbol{x}_1 - \boldsymbol{x}_2)^T \widetilde{\boldsymbol{\beta}} \right) - \widehat{\boldsymbol{F}}_{\widetilde{\boldsymbol{\beta}}}\left((\boldsymbol{x}_1 - \boldsymbol{x}_2)^T \widetilde{\boldsymbol{\beta}} \right) \right\|_{2,F_{X_1,X_2}}^2 + O_p(n^{-2/3}) \\ &\lesssim \left\| F_{\widetilde{\boldsymbol{\beta}}}\left((\boldsymbol{x}_1 - \boldsymbol{x}_2)^T \widetilde{\boldsymbol{\beta}} \right) - \widehat{\boldsymbol{F}}_{\widetilde{\boldsymbol{\beta}}}\left((\boldsymbol{x}_1 - \boldsymbol{x}_2)^T \widetilde{\boldsymbol{\beta}} \right) \right\|_{2,F_{X_1,X_2}}^2 + O_p(n^{-2/3}) \\ &\leq \left\| F_{\widetilde{\boldsymbol{\beta}}}\left((\boldsymbol{x}_1 - \boldsymbol{x}_2)^T \widetilde{\boldsymbol{\beta}} \right) - \widehat{\boldsymbol{F}}_{\widetilde{\boldsymbol{\beta}}}\left((\boldsymbol{x}_1 - \boldsymbol{x}_2)^T \widetilde{\boldsymbol{\beta}} \right) \right\|_{2,F_{X_1,X_2}}^2 + O_p(n^{-2/3}) \\ &\leq \left\| F_{\widetilde{\boldsymbol{\beta}}}\left((\boldsymbol{x}_1 - \boldsymbol{x}_2)^T \widetilde{\boldsymbol{\beta}} \right) - \widehat{\boldsymbol{F}}_{\widetilde{\boldsymbol{\beta}}}\left((\boldsymbol{x}_1 - \boldsymbol{x}_2)^T \widetilde{\boldsymbol{\beta}} \right) \right\|_{2,F_{X_1,X_2}}^2 + O_p(n^{-2/3}) \\ &= O_p(n^{-2/3}), \end{aligned}$$

where " \leq " is because of Lemma 13, and the last "=" is because of Lemma 12. Therefore, we have completed the proof of (S.96).

Next, we consider I_3 . Denote

$$\widetilde{g}_{6,\boldsymbol{\beta},F}(\boldsymbol{x}_1 - \boldsymbol{x}_2, y_1 - y_2) = \left\{ \boldsymbol{x}_1 - \boldsymbol{x}_2 - \varphi_{\boldsymbol{\beta}} \left((\boldsymbol{x}_1 - \boldsymbol{x}_2)^T \boldsymbol{\beta} \right) \right\} \\ \times \left\{ I(y_1 > y_2) - F \left((\boldsymbol{x}_1 - \boldsymbol{x}_2)^T \boldsymbol{\beta} \right) \right\}$$
(S.102)

and

$$g_{6,\boldsymbol{\beta},F}(\boldsymbol{x}_{1}-\boldsymbol{x}_{2},y_{1}-y_{2}) = \widetilde{g}_{6,\boldsymbol{\beta},F}(\boldsymbol{x}_{1}-\boldsymbol{x}_{2},y_{1}-y_{2}) \\ -\widetilde{g}_{6,\boldsymbol{\beta}_{0},F_{0}}(\boldsymbol{x}_{1}-\boldsymbol{x}_{2},y_{1}-y_{2}).$$

Then

$$I_3 = \int \int \widetilde{g}_{6,\widetilde{\boldsymbol{\beta}},\widehat{F}_{\widetilde{\boldsymbol{\beta}}}}(\boldsymbol{x}_1 - \boldsymbol{x}_2, y_1 - y_2) d\mathbb{F}_{X,Y}(\boldsymbol{x}_1, y_1) d\mathbb{F}_{X,Y}(\boldsymbol{x}_2, y_2).$$

Based on Conditions 1 and A2, we have

$$\begin{split} \left| g_{6,\widecheck{\boldsymbol{\beta}},\widehat{F}_{\widecheck{\boldsymbol{\beta}}}}(\boldsymbol{v},\boldsymbol{u}) \right| \lesssim \left| \widecheck{\boldsymbol{\beta}} - \boldsymbol{\beta}_{0} \right| + \left| \widehat{F}_{\widecheck{\boldsymbol{\beta}}}\left(\boldsymbol{v}^{T} \widecheck{\boldsymbol{\beta}} \right) - F_{\widecheck{\boldsymbol{\beta}}}\left(\boldsymbol{v}^{T} \widecheck{\boldsymbol{\beta}} \right) \right| \\ + \left| F_{\widecheck{\boldsymbol{\beta}}}\left(\boldsymbol{v}^{T} \widecheck{\boldsymbol{\beta}} \right) - F_{0}(\boldsymbol{v}^{T} \boldsymbol{\beta}_{0}) \right|, \end{split}$$

up to a constant not depending on $v \in \mathcal{X} - \mathcal{X}$ and $u \in \mathcal{Y} - \mathcal{Y}$. With this inequality, working on function classes formed by $g_{6,\beta,F}(x_1 - x_2, y_1 - y_2)$ and using the same arguments as in the proof of Lemma 8, we can derive

$$\int \int g_{6,\breve{\beta},\widehat{F}_{\breve{\beta}}}(\bm{x}_{1}-\bm{x}_{2},y_{1}-y_{2})d\mathbb{F}_{X,Y}(\bm{x}_{1},y_{1})d\mathbb{F}_{X,Y}(\bm{x}_{2},y_{2})$$

$$= \int \int g_{6,\breve{\beta},\widehat{F}_{\breve{\beta}}}(\bm{x}_{1}-\bm{x}_{2},y_{1}-y_{2})dF_{X,Y}(\bm{x}_{1},y_{1})dF_{X,Y}(\bm{x}_{2},y_{2}) + o_{p}(n^{-1/2}).$$

As a consequence,

$$I_{3} = \int \int \widetilde{g}_{6,\beta_{0},F_{0}}(\boldsymbol{x}_{1} - \boldsymbol{x}_{2}, y_{1} - y_{2}) d\mathbb{F}_{X,Y}(\boldsymbol{x}_{1}, y_{1}) d\mathbb{F}_{X,Y}(\boldsymbol{x}_{2}, y_{2}) + \int \int g_{6,\beta,\hat{F}_{\beta}}(\boldsymbol{x}_{1} - \boldsymbol{x}_{2}, y_{1} - y_{2}) dF_{X,Y}(\boldsymbol{x}_{1}, y_{1}) dF_{X,Y}(\boldsymbol{x}_{2}, y_{2}) + o_{p}(n^{-1/2}).$$
(S.103)

We consider the second term on the right-hand side of the equality above. Note that it is

straightforward to verify that

$$\int \int \widetilde{g}_{6,\boldsymbol{\beta}_{0},F_{0}}(\boldsymbol{x}_{1}-\boldsymbol{x}_{2},y_{1}-y_{2})dF_{X,Y}(\boldsymbol{x}_{1},y_{1})dF_{X,Y}(\boldsymbol{x}_{2},y_{2})=0.$$

Therefore, using the notation $\boldsymbol{u} = y_1 - y_2$ and $\boldsymbol{v} = \boldsymbol{x}_1 - \boldsymbol{x}_2$, we have

$$\int \int g_{6,\widetilde{\boldsymbol{\beta}},\widehat{F}_{\widetilde{\boldsymbol{\beta}}}}(\boldsymbol{x}_{1}-\boldsymbol{x}_{2},y_{1}-y_{2})dF_{X,Y}(\boldsymbol{x}_{1},y_{1})dF_{X,Y}(\boldsymbol{x}_{2},y_{2})$$

$$= \int \int \widetilde{g}_{6,\widetilde{\boldsymbol{\beta}},\widehat{F}_{\widetilde{\boldsymbol{\beta}}}}(\boldsymbol{x}_{1}-\boldsymbol{x}_{2},y_{1}-y_{2})dF_{X,Y}(\boldsymbol{x}_{1},y_{1})dF_{X,Y}(\boldsymbol{x}_{2},y_{2})$$

$$= \int \int \left\{ \boldsymbol{v}-\varphi_{\widetilde{\boldsymbol{\beta}}}\left(\boldsymbol{v}^{T}\widetilde{\boldsymbol{\beta}}\right)\right\} \left\{ I(u>0)-\widehat{F}_{\widetilde{\boldsymbol{\beta}}}\left(\boldsymbol{v}^{T}\widetilde{\boldsymbol{\beta}}\right)\right\} dF_{V,U}(\boldsymbol{v},u)$$

$$= \int \int \left\{ \boldsymbol{v}-\varphi_{\widetilde{\boldsymbol{\beta}}}\left(\boldsymbol{v}^{T}\widetilde{\boldsymbol{\beta}}\right)\right\} \left\{ I(u>0)-F_{\widetilde{\boldsymbol{\beta}}}\left(\boldsymbol{v}^{T}\widetilde{\boldsymbol{\beta}}\right)\right\} dF_{V,U}(\boldsymbol{v},u)$$

$$+ \int \int \left\{ \boldsymbol{v}-\varphi_{\widetilde{\boldsymbol{\beta}}}\left(\boldsymbol{v}^{T}\widetilde{\boldsymbol{\beta}}\right)\right\} \left\{ F_{\widetilde{\boldsymbol{\beta}}}\left(\boldsymbol{v}^{T}\widetilde{\boldsymbol{\beta}}\right)-\widehat{F}_{\widetilde{\boldsymbol{\beta}}}\left(\boldsymbol{v}^{T}\widetilde{\boldsymbol{\beta}}\right)\right\} dF_{V}(\boldsymbol{v}) \quad (S.104)$$

$$= \iint \left\{ \boldsymbol{v} - \varphi_{\boldsymbol{\breve{\beta}}} \left(\boldsymbol{v}^T \boldsymbol{\breve{\beta}} \right) \right\} \left\{ I(u > 0) - F_{\boldsymbol{\breve{\beta}}} \left(\boldsymbol{v}^T \boldsymbol{\breve{\beta}} \right) \right\} dF_{V,U}(\boldsymbol{v}, u) \qquad (S.105)$$

$$= \iint_{\sim} \boldsymbol{v} \left\{ I(u > 0) - F_{\boldsymbol{\beta}} \left(\boldsymbol{v}^{T} \boldsymbol{\beta} \right) \right\} dF_{V,U}(\boldsymbol{v}, u)$$
(S.106)

$$= \psi_0(\check{\boldsymbol{\beta}}). \tag{S.107}$$

Some of the above steps need explanation. From (S.104) to (S.105), we have used the fact that $\varphi_{\beta}(V^T \check{\beta}), F_{\beta}(V^T \check{\beta}), \text{ and } \hat{F}_{\beta}(V^T \check{\beta})$ are all measurable functions of $V^T \check{\beta}, \check{\beta}, (X_1, Y_1) \dots, (X_n, Y_n)$, so by considering V as a random vector independent of $\check{\beta}, (X_1, Y_1) \dots, (X_n, Y_n)$, we have

$$\int \left\{ \boldsymbol{v} - \varphi_{\check{\boldsymbol{\beta}}} \left(\boldsymbol{v}^{T} \check{\boldsymbol{\beta}} \right) \right\} \left\{ F_{\check{\boldsymbol{\beta}}} \left(\boldsymbol{v}^{T} \check{\boldsymbol{\beta}} \right) - \widehat{F}_{\check{\boldsymbol{\beta}}} \left(\boldsymbol{v}^{T} \check{\boldsymbol{\beta}} \right) \right\} dF_{V}(\boldsymbol{v}) \\
= E \left[\left\{ V - \varphi_{\check{\boldsymbol{\beta}}} \left(V^{T} \check{\boldsymbol{\beta}} \right) \right\} \left\{ F_{\check{\boldsymbol{\beta}}} \left(V^{T} \check{\boldsymbol{\beta}} \right) - \widehat{F}_{\check{\boldsymbol{\beta}}} \left(V^{T} \check{\boldsymbol{\beta}} \right) \right\} \left| \{(X_{i}, Y_{i})\}_{i=1}^{n}, \check{\boldsymbol{\beta}} \right] \\
= E \left[\left\{ E \left(V \middle| V^{T} \check{\boldsymbol{\beta}} \right) - \varphi_{\check{\boldsymbol{\beta}}} \left(V^{T} \check{\boldsymbol{\beta}} \right) \right\} \left\{ F_{\check{\boldsymbol{\beta}}} \left(V^{T} \check{\boldsymbol{\beta}} \right) - \widehat{F}_{\check{\boldsymbol{\beta}}} \left(V^{T} \check{\boldsymbol{\beta}} \right) \right\} \left| \{(X_{i}, Y_{i})\}_{i=1}^{n}, \check{\boldsymbol{\beta}} \right] \\
= E \left[\left\{ \varphi_{\check{\boldsymbol{\beta}}} \left(V^{T} \check{\boldsymbol{\beta}} \right) - \varphi_{\check{\boldsymbol{\beta}}} \left(V^{T} \check{\boldsymbol{\beta}} \right) \right\} \left\{ F_{\check{\boldsymbol{\beta}}} \left(V^{T} \check{\boldsymbol{\beta}} \right) - \widehat{F}_{\check{\boldsymbol{\beta}}} \left(V^{T} \check{\boldsymbol{\beta}} \right) \right\} \left| \{(X_{i}, Y_{i})\}_{i=1}^{n}, \check{\boldsymbol{\beta}} \right] \\
= 0.$$

From (S.105) to (S.106), we have used the fact that $\varphi_{\check{\beta}}\left(V^T\check{\beta}\right)$ and $F_{\check{\beta}}\left(V^T\check{\beta}\right)$ are measurable functions of $V^T\check{\beta},\check{\beta},(X_1,Y_1),\ldots,(X_n,Y_n)$, so by considering (U,V) as independent of

 $\check{\boldsymbol{\beta}}, (X_1, Y_1) \dots, (X_n, Y_n)$, we have

$$\begin{split} &\int \int \varphi_{\widetilde{\boldsymbol{\beta}}} \left(\boldsymbol{v}^{T} \check{\boldsymbol{\beta}} \right) \left\{ I(u > 0) - F_{\widetilde{\boldsymbol{\beta}}} \left(\boldsymbol{v}^{T} \check{\boldsymbol{\beta}} \right) \right\} dF_{V,U}(\boldsymbol{v}, u) \\ &= E \left[\varphi_{\widetilde{\boldsymbol{\beta}}} \left(V^{T} \check{\boldsymbol{\beta}} \right) \left\{ I(U > 0) - F_{\widetilde{\boldsymbol{\beta}}} \left(V^{T} \check{\boldsymbol{\beta}} \right) \right\} \left| \check{\boldsymbol{\beta}}, \{ (X_{1}, Y_{1}) \}_{i=1}^{n} \right] \\ &= E \left[\varphi_{\widetilde{\boldsymbol{\beta}}} \left(V^{T} \check{\boldsymbol{\beta}} \right) \left\{ E \left\{ I(U > 0) \middle| V^{T} \check{\boldsymbol{\beta}} \right\} - F_{\widetilde{\boldsymbol{\beta}}} \left(V^{T} \check{\boldsymbol{\beta}} \right) \right\} \middle| \check{\boldsymbol{\beta}}, \{ (X_{1}, Y_{1}) \}_{i=1}^{n} \right] \\ &= E \left[\varphi_{\widetilde{\boldsymbol{\beta}}} \left(V^{T} \check{\boldsymbol{\beta}} \right) \left\{ F_{\widetilde{\boldsymbol{\beta}}} \left(V^{T} \check{\boldsymbol{\beta}} \right) - F_{\widetilde{\boldsymbol{\beta}}} \left(V^{T} \check{\boldsymbol{\beta}} \right) \right\} \left| \check{\boldsymbol{\beta}}, \{ (X_{1}, Y_{1}) \}_{i=1}^{n} \right] = 0. \end{split}$$

From (S.106) to (S.107), we have used the definition of $\psi_0(\cdot)$ given by (S.4).

Applying Lemma 1, and combining (S.94), (S.102), (S.103), and (S.107), we have

$$I_{3} = \iint \left\{ \boldsymbol{x}_{1} - \boldsymbol{x}_{2} - \varphi_{0} \left((\boldsymbol{x}_{1} - \boldsymbol{x}_{2})^{T} \boldsymbol{\beta}_{0} \right) \right\} \\ \times \left\{ I(y_{1} > y_{2}) - F_{0} \left((\boldsymbol{x}_{1} - \boldsymbol{x}_{2})^{T} \boldsymbol{\beta}_{0} \right) \right\} d\mathbb{F}_{X,Y}(\boldsymbol{x}_{1}, y_{1}) d\mathbb{F}_{X,Y}(\boldsymbol{x}_{2}, y_{2}) \\ + A(\boldsymbol{\check{\beta}} - \boldsymbol{\beta}_{0}) + o_{p}(\boldsymbol{\check{\beta}} - \boldsymbol{\beta}_{0}) + o_{p}(n^{-1/2}).$$
(S.108)

Combining (S.95), (S.96), and (S.108), we conclude that for any estimator $\check{\beta}$ of β_0 such that $\check{\beta} - \beta_0 = o_p(1)$, we have

$$\psi_{n}(\check{\boldsymbol{\beta}}) = \iint \left\{ \boldsymbol{x}_{1} - \boldsymbol{x}_{2} - \varphi_{0} \left((\boldsymbol{x}_{1} - \boldsymbol{x}_{2})^{T} \boldsymbol{\beta}_{0} \right) \right\} \\ \times \left\{ I(y_{1} > y_{2}) - F_{0} \left((\boldsymbol{x}_{1} - \boldsymbol{x}_{2})^{T} \boldsymbol{\beta}_{0} \right) \right\} d\mathbb{F}_{X,Y}(\boldsymbol{x}_{1}, y_{1}) d\mathbb{F}_{X,Y}(\boldsymbol{x}_{2}, y_{2}) \\ + A(\check{\boldsymbol{\beta}} - \boldsymbol{\beta}_{0}) + o_{p}(\check{\boldsymbol{\beta}} - \boldsymbol{\beta}_{0}) + o_{p}(n^{-1/2}).$$
(S.109)

Furthermore, the conclusion in Part (b) of this theorem implies that $\tilde{\beta}^{(U)}$ and $\tilde{\beta}^{(L)}$ given by (S.92) are both consistent estimators for β_0 . Combining (S.93) with (S.109) leads to

$$0 = \boldsymbol{\alpha}_{n}^{T} \psi_{n}(\widetilde{\boldsymbol{\beta}}^{(U)}) + (1 - \boldsymbol{\alpha}_{n})^{T} \psi_{n}(\widetilde{\boldsymbol{\beta}}^{(L)}) = \iint \{ \boldsymbol{x}_{1} - \boldsymbol{x}_{2} - \varphi_{0} ((\boldsymbol{x}_{1} - \boldsymbol{x}_{2})^{T} \boldsymbol{\beta}_{0}) \} \times \{ I(y_{1} > y_{2}) - F_{0} ((\boldsymbol{x}_{1} - \boldsymbol{x}_{2})^{T} \boldsymbol{\beta}_{0}) \} d\mathbb{F}_{X,Y}(\boldsymbol{x}_{1}, y_{1}) d\mathbb{F}_{X,Y}(\boldsymbol{x}_{2}, y_{2}) + A(\widetilde{\boldsymbol{\beta}} - \boldsymbol{\beta}_{0}) + o_{p}(\widetilde{\boldsymbol{\beta}} - \boldsymbol{\beta}_{0}) + o_{p}(n^{-1/2}), \end{cases}$$

which completes our proof of Theorem 2.

5.5 **Proof of Corollary 1**

Based on Theorem 2, we need to work with the asymptotics of the statistic

$$\mathbb{U}_{n} = \sum_{i \neq j} \left\{ X_{i} - X_{j} - \varphi_{0} \left((X_{i} - X_{j})^{T} \beta_{0} \right) \right\} \left\{ I(Y_{i} > Y_{j}) - F_{0} \left((X_{i} - X_{j})^{T} \beta_{0} \right) \right\}.$$
 (S.110)

Without loss of generality, assume $\beta_{0,1} \neq 0$, i.e., the first element of β_0 is nonzero. Then we have $\widetilde{A} = \psi'_0(\beta_0)\Sigma_X^{-1} + \begin{pmatrix} \beta_0^T \\ \mathbf{0} \end{pmatrix}$; and it is invertible. If otherwise, there exists $\mathbf{0} \neq \boldsymbol{\alpha} \in \mathbb{R}^p$ such that

$$\boldsymbol{\alpha}^{T}\psi_{0}^{\prime}(\boldsymbol{\beta}_{0})\boldsymbol{\Sigma}_{X}^{-1} + \boldsymbol{\alpha}_{1}\boldsymbol{\beta}_{0}^{T} = 0.$$
(S.111)

• If $\alpha_1 \neq 0$, (S.111) leads to

$$\boldsymbol{\beta}_0^T \Sigma_X \boldsymbol{\beta}_0 = -\frac{\boldsymbol{\alpha}^T}{\boldsymbol{\alpha}_1} \psi_0'(\boldsymbol{\beta}_0) \boldsymbol{\beta}_0 = 0,$$

which contradicts that $\beta_0 \neq 0$ and Σ_X is of full rank.

• If $\alpha_1 = 0$, (S.111) leads to

$$\boldsymbol{\alpha}_{2:p}^T \psi_0'(\boldsymbol{\beta}_0)_{2:p} \Sigma_X^{-1} = 0,$$

which together with our assumption that Σ_X is of full rank contradicts the conclusion given by Lemma 11.

For any $i \neq j$, noting that $\{\psi'_0(\beta_0)\Sigma_X^{-1}\}\Sigma_X\beta_0 = 0$ and the assumption given in (S.7), we have

$$\psi_{0}'(\boldsymbol{\beta}_{0})\Sigma_{X}^{-1}\left[X_{i}-X_{j}-E\left\{X_{i}-X_{j}|(X_{i}-X_{j})^{T}\boldsymbol{\beta}_{0}\right\}\right]$$

= $\psi_{0}'(\boldsymbol{\beta}_{0})\Sigma_{X}^{-1}(X_{i}-X_{j})-E\left\{\psi_{0}'(\boldsymbol{\beta}_{0})\Sigma_{X}^{-1}(X_{i}-X_{j})|(X_{i}-X_{j})^{T}\boldsymbol{\beta}_{0}\right\}$
= $\psi_{0}'(\boldsymbol{\beta}_{0})\Sigma_{X}^{-1}(X_{i}-X_{j}),$

and

$$\boldsymbol{\beta}_0^T \Big[X_i - X_j - E \left\{ X_i - X_j | (X_i - X_j)^T \boldsymbol{\beta}_0 \right\} \Big] = 0.$$

As a consequence, for any $i \neq j$, recalling that

$$\varphi_0\left((X_i - X_j)^T \boldsymbol{\beta}_0\right) = E\left\{X_i - X_j | (X_i - X_j)^T \boldsymbol{\beta}_0\right\},\,$$

we have

$$\widetilde{A}^{-1}\widetilde{A}\left\{X_i - X_j - \varphi_0\left((X_i - X_j)^T \boldsymbol{\beta}_0\right)\right\} = \widetilde{A}^{-1}\psi_0'(\boldsymbol{\beta}_0)\Sigma_X^{-1}(X_i - X_j).$$

Therefore

$$\mathbb{U}_{n} = \widetilde{A}^{-1}\psi_{0}'(\beta_{0})\Sigma_{X}^{-1}\sum_{i\neq j}(X_{i}-X_{j})\left\{I(Y_{i}>Y_{j})-F_{0}\left((X_{i}-X_{j})^{T}\beta_{0}\right)\right\} \\
= \widetilde{A}^{-1}\psi_{0}'(\beta_{0})\Sigma_{X}^{-1}\sum_{i\neq j}h(X_{i},Y_{i};X_{j},Y_{j}),$$

where

$$h(\boldsymbol{x}_1, y_1; \boldsymbol{x}_2, y_2) = (\boldsymbol{x}_1 - \boldsymbol{x}_2) \left\{ I(y_1 > y_2) - F_0 \left((\boldsymbol{x}_1 - \boldsymbol{x}_2)^T \boldsymbol{\beta}_0 \right) \right\}.$$

Because of Conditions 0 and A1, with probability equal to 1, $Y_i \neq Y_j$ for $i \neq j$. Using the fact that $F_0(\cdot)$ is the c.d.f. of a symmetric distribution about 0, we have

$$I(y_1 > y_2) - F_0 \left((\boldsymbol{x}_1 - \boldsymbol{x}_2)^T \boldsymbol{\beta}_0 \right) \\ = 1 - I(y_1 < y_2) - \left\{ 1 - F_0 \left(-(\boldsymbol{x}_1 - \boldsymbol{x}_2)^T \boldsymbol{\beta}_0 \right) \right\} \\ = - \left\{ I(y_2 > y_1) - F_0 \left((\boldsymbol{x}_2 - \boldsymbol{x}_1)^T \boldsymbol{\beta}_0 \right) \right\},$$

and

$$m{x}_1 - m{x}_2 = - \left(m{x}_2 - m{x}_1
ight).$$

Therefore,

$$h(\boldsymbol{x}_1, y_1; \boldsymbol{x}_2, y_2) = h(\boldsymbol{x}_2, y_2; \boldsymbol{x}_1, y_1),$$

i.e., h is a symmetric kernel for the U-statistic contained in \mathbb{U}_n . Furthermore, it is straightforward to verify that $E\{h(X_1, Y_1; X_2, Y_2)\} = 0$; we have

$$\overline{\mathbb{U}}_n = \sum_{i=1}^n E(\mathbb{U}_n | X_i, Y_i)$$
$$= \frac{2(n-1)}{n^2} \widetilde{A}^{-1} \psi_0'(\boldsymbol{\beta}_0) \Sigma_X^{-1} \sum_{i=1}^n \hbar(X_i, Y_i).$$

where we recall that

$$\hbar(\boldsymbol{x}_1, y_1) = E\{h(\boldsymbol{x}_1, y_1; X_2, Y_2)\}.$$

With the condition that $\lambda_{\max}(\Sigma_{\hbar}) < \infty$, and applying Theorem 12.3 in van der Vaart (1998), we have that

$$\mathbb{U}_n - \overline{\mathbb{U}}_n = o_p(n^{-1/2}) \tag{S.112}$$

holds entry-wise. By the condition $0 < \lambda_{\min}(\Sigma_{\hbar}) \leq \lambda_{\max}(\Sigma_{\hbar}) < \infty$ and the Central Limit

Theorem, we have

$$\sqrt{n\overline{\mathbb{U}}}_n \to 2\widetilde{A}^{-1}\psi_0'(\boldsymbol{\beta}_0)\Sigma_X^{-1}N(0,\Sigma_\hbar), \qquad (S.113)$$

where we have used the fact that $E \{\hbar(X_1, Y_1)\} = 0$. Combining (S.110), (S.112), (S.113) with (S.5) leads to (S.8). This completes the proof of this corollary.

6 Proof of Theorem 3

In this section, we shall adopt the following notations. For any function $f(x_1, y_1; x_2, y_2)$, we denote

$$\mathbb{P}^{2}f = \int f(\boldsymbol{x}_{1}, y_{1}; \boldsymbol{x}_{2}, y_{2}) dF_{X,Y}(\boldsymbol{x}_{1}, y_{1}) dF_{X,Y}(\boldsymbol{x}_{2}, y_{2}),$$

and $\mathbb{U}_n^2 f$ to be the second order U-statistic with f being the kernel. For any function f(x, y), we denote

$$\mathbb{P}f = \mathbb{P}f(\cdot, \cdot) = \int f(\boldsymbol{x}, y) dF_{X,Y}(\boldsymbol{x}, y)$$
$$\mathbb{P}_n f = \mathbb{P}_n f(\cdot, \cdot) = \frac{1}{n} \sum_{i=1}^n f(X_i, Y_i)$$
$$\mathbb{G}_n f = \sqrt{n} \left(\mathbb{P}_n f - \mathbb{P}f\right).$$

We generally use " \overline{F} " to denote the envelope function of a function class. Denote $T_{i,j} = (X_i - X_j)^T \beta$, $\widetilde{T}_{i,j} = (X_i - X_j)^T \widetilde{\beta}$, $T_{i,j,0} = (X_i - X_j)^T \beta_0$, and $\Delta_{i,j} = I(Y_i > Y_j)$; then the expression for $\ell(\beta, F)$ can be rewritten to be

$$\ell(\boldsymbol{\beta}, F) = \sum_{i,j} \left[\Delta_{i,j} \log\{F(T_{i,j})\} + (1 - \Delta_{i,j}) \log\{1 - F(T_{i,j})\} \right].$$
(S.114)

For any β , let $G_{\beta}(\cdot)$ and $G_n(\cdot)$ be the c.d.f. and empirical c.d.f. of $T_{i,j}$ respectively. Denote $G_0 = G_{\beta_0}, g_{\beta}(t) = \frac{\partial G_{\beta}(t)}{\partial t}$, and $g_0(t) = \frac{\partial G_0(t)}{\partial t}$. Let

$$V_n(t) = \frac{1}{n^2} \sum_{i,j} \Delta_{i,j} I(T_{i,j} \leqslant t).$$

In addition, we add tilde to a notation to denote the corresponding expression whose β is replaced by $\tilde{\beta}$; for example,

$$\widetilde{V}_n(t) = \frac{1}{n^2} \sum_{i,j} \Delta_{i,j} I(\widetilde{T}_{i,j} \leqslant t).$$

We sketch a blueprint of the proof for this theorem first; the technical details are then

organised in the lemmas afterwards. Note that for every β , \hat{F}_{β} is the maximiser of $\ell(\beta, F)$ given by (S.114); it is the left derivative of the greatest convex minorant of the cumulative sum diagram defined by $(G_n(t), V_n(t)), t \in \mathbb{R}$. Furthermore, let

$$U_n(s) = \operatorname{argmin}_{t \in \mathbb{R}} \{ V_n(t) - sG_n(t) \}.$$
(S.115)

Following the discussion given at the beginning of the proof for Theorem 3.7 of Groeneboom and Jongbloed (2014), we have the switching relation:

$$\widehat{F}_{\boldsymbol{\beta}}(t) \ge s \Leftrightarrow G_n(t) \ge G_n(U_n(s)) \Leftrightarrow t \ge U_n(s).$$

Set $s_0 \equiv F_0(t)$; we have

$$P\left(n^{1/2}\left\{\widehat{F}_{\widetilde{\boldsymbol{\beta}}}(t) - F_{0}(t)\right\} \ge x\right) = P\left(\widehat{F}_{\widetilde{\boldsymbol{\beta}}}(t) \ge s_{0} + n^{-1/2}x\right)$$
$$= P\left(t \ge \widetilde{U}_{n}(s_{0} + n^{-1/2}x)\right)$$
$$= P\left(n^{1/2}\left\{\widetilde{U}_{n}(s_{0} + n^{-1/2}x) - t\right\} \le 0\right).$$
(S.116)

Therefore, to derive the asymptotic distribution for $n^{1/2} \left\{ \widehat{F}_{\widetilde{\beta}}(t) - F_0(t) \right\}$, we can work on the asymptotic distribution of $n^{1/2} \left\{ \widetilde{U}_n(s_0 + n^{-1/2}x) - t \right\}$ instead. This is achieved by three lemmas.

- We first establish that for any M > 0, $n^{1/3} \sup_{x \in [-M,M]} \left| \widetilde{U}_n(s_0 + n^{-1/2}x) t \right| = O_p(1)$ in Lemma 15.
- We then show that $n^{1/2} \sup_{x \in [-M,M]} \left| \widetilde{U}_n(s_0 + n^{-1/2}x) t \right| = O_p(1)$ in Lemma 17.
- Last with the argmax theorem reviewed by Lemma 18, we establish the asymptotic distribution for $n^{1/2} \left\{ \widetilde{U}_n(s_0 + n^{-1/2}x) t \right\}$ in Lemma 19:

$$n^{1/2} \left\{ \widetilde{U}_n(s_0 + n^{-1/2}x) - t \right\} \leadsto \frac{g_0(t)x - \sigma(t)Z}{f_0(t)g_0(t)}.$$
(S.117)

Combining (S.117) with (S.116) leads to

$$P\left(n^{1/2}\left\{\widehat{F}_{\widetilde{\boldsymbol{\beta}}}(t) - F_{0}(t)\right\} \ge x\right) \rightarrow P\left(\frac{g_{0}(t)x - \sigma(t)Z}{g_{0}(t)f_{0}(t)} \le 0\right)$$
$$= P\left(\frac{\sigma(t)}{g_{0}(t)}Z \ge x\right),$$

where $Z \sim N(0,1)$, which implies $n^{1/2} \left\{ \widehat{F}_{\widetilde{\beta}}(t) - F_0(t) \right\} \rightsquigarrow N(0, \sigma^2(t)/g_0^2(t))$. This com-

pletes the proof of this theorem.

We first establish the tightness of $n^{1/3} \sup_{x \in [-M,M]} \left| \widetilde{U}_n(s_0 + n^{-1/2}x) - t \right| = O_p(1)$ in the following lemma.

Lemma 15. Assume that all conditions for Theorem 3 are effective. For every M > 0, we have

$$n^{1/3} \sup_{x \in [-M,M]} \left| \widetilde{U}_n(s_0 + n^{-1/2}x) - t \right| = O_p(1).$$
(S.118)

Proof:Based on Corollary 1, $\tilde{\boldsymbol{\beta}} - \boldsymbol{\beta}_0 = O_p(n^{-1/2})$; therefore for any $\epsilon > 0$, there exists a universal constant C > 0, such that

$$P\left(\widetilde{\boldsymbol{\beta}}\in \mathcal{B}_{n,C}\right) > 1-\epsilon,$$

where $\mathcal{B}_{n,C} = \{ \boldsymbol{\beta} : \| \boldsymbol{\beta} - \boldsymbol{\beta}_0 \|_2 \leq n^{-1/2}C \}$. Hereafter, if appropriate we treat $\boldsymbol{\beta} \in \mathcal{B}_{n,C}$ to avoid some presentational complication; for example, in some occasions, we are to show that the probability of some event, which involves $\boldsymbol{\beta}$, can be arbitrarily small; we can treat $\boldsymbol{\beta} \in \mathcal{B}_{n,C}$, since otherwise we can use $\{ \boldsymbol{\beta} \in \mathcal{B}_{n,C} \}$ to intersect the event of interest.

Note that $U_n(s)$ is nondecreasing in s; this is because for any $s_1 < s_2$, let $t_1 = U_n(s_1)$ and $t_2 = U_n(s_2)$, then we must have $t_1 \leq t_2$; since if otherwise, recalling the definition of $U_n(\cdot)$ given by (S.115), we have

$$V_{n}(t_{2}) - s_{2}G_{n}(t_{2})$$

$$= \frac{1}{n^{2}}\sum_{i,j}(\Delta_{i,j} - s_{2})I(T_{i,j} \leq t_{2})$$

$$= \frac{1}{n^{2}}\sum_{i,j}(\Delta_{i,j} - s_{1})I(T_{i,j} \leq t_{2}) + \frac{1}{n^{2}}\sum_{i,j}(s_{1} - s_{2})I(T_{i,j} \leq t_{2})$$

$$\geq \frac{1}{n^{2}}\sum_{i,j}(\Delta_{i,j} - s_{1})I(T_{i,j} \leq t_{1}) + \frac{1}{n^{2}}\sum_{i,j}(s_{1} - s_{2})I(T_{i,j} \leq t_{1})$$

$$= \frac{1}{n^{2}}\sum_{i,j}(\Delta_{i,j} - s_{2})I(T_{i,j} \leq t_{1})$$

$$= V_{n}(t_{1}) - s_{2}G_{n}(t_{1}),$$

which contradicts the definition of t_2 . Therefore, to show (S.118), we need to show that for any $\epsilon > 0$, there exists an M' > 0, such that

$$P\left(n^{1/3}\left\{\widetilde{U}_{n}(s_{0}+n^{-1/2}M)-t\right\}>M'\right)<\epsilon$$
(S.119)

$$P\left(n^{1/3}\left\{\widetilde{U}_{n}(s_{0}-n^{-1/2}M)-t\right\}<-M'\right)<\epsilon.$$
(S.120)

We need to show only (S.119); (S.120) can be shown using exactly the same arguments. Note

that

$$P\left(n^{1/3}\left\{\widetilde{U}_{n}(s_{0}+n^{-1/2}M)-t\right\} > M'\right)$$

$$\leq P\left(\exists u > M': \widetilde{V}_{n}(t+n^{-1/3}u) - (s_{0}+n^{-1/2}M)\widetilde{G}_{n}(t+n^{-1/3}u)\right)$$

$$\leq \widetilde{V}_{n}(t) - (s_{0}+n^{-1/2}M)\widetilde{G}_{n}(t)\right)$$

$$= P\left(\exists u > M': \widetilde{W}_{n,1}(u) + \widetilde{W}_{n,2}(u) - \widetilde{W}_{n,3}(u) + \widetilde{W}_{n,4}(u) \leq 0\right),$$

where we recall that $s_0 = F_0(t)$, and

$$W_{n,1}(u) = n^{2/3} \left\{ V_n(t + n^{-1/3}u) - V_n(t) \right\} - \frac{n^{2/3}}{n^2} \sum_{i,j} F_0(T_{i,j,0}) I(T_{i,j} \in [t, t + n^{-1/3}u]) W_{n,2}(u) = n^{2/3} \int_{w \in [t, t + n^{-1/3}u]} \{F_0(w) - F_0(t)\} dG_n(w) W_{n,3}(u) = n^{1/6} M \left\{ G_n(t + n^{-1/3}u) - G_n(t) \right\} W_{n,4}(u) = \frac{n^{2/3}}{n^2} \sum_{i,j} \left[\left\{ F_0(T_{i,j,0}) - F_0(T_{i,j}) \right\} \times I(T_{i,j} \in [t, t + n^{-1/3}u]) \right].$$
(S.121)

We derive the asymptotic properties of $\widetilde{W}_{n,i}(u)$ for i = 1, 2, 3, 4 separately. In particular, we show that

(a) for any $\eta > 0, \epsilon > 0$, and M' > 0, when M_1 is sufficiently large, we have

$$P\left(\exists u > M' : \left|\widetilde{W}_{n,1}(u)\right| \ge \eta u^2 + M_1\right) < \epsilon;$$
(S.122)

(b) there exists a universal constant c > 0, such that for any $\epsilon > 0$, when M' is sufficiently large, we have

$$P\left(\exists u > M' : \widetilde{W}_{n,2}(u) \ge cu^2\right) > 1 - \epsilon;$$
(S.123)

(c) for any $\epsilon > 0$ and M' > 0, when M_1 is sufficiently large, we have

$$P\left(\exists u > M' : \left|\widetilde{W}_{n,3}(u)\right| \ge M_1\right) < \epsilon;$$

(d) for any $\epsilon > 0$ and M' > 0, when M_1 is sufficiently large, we have

$$P\left(\exists u > M' : \left|\widetilde{W}_{n,4}(u)\right| \ge M_1\right) < \epsilon$$

We show (a) first. Note that

$$W_{n,1}(u) = \frac{n^{2/3}}{n^2} \sum_{i,j} \{\Delta_{i,j} - F_0(T_{i,j,0})\} I(T_{i,j} \in [t, t + n^{-1/3}u])$$
$$\equiv \frac{n^{2/3}}{n^2} \sum_{i,j} f_{n,u}(X_i, Y_i, X_j, Y_j),$$

where

$$f_{n,u}(X_i, Y_i; X_j, Y_j) = \{\Delta_{i,j} - F_0(T_{i,j,0})\} I(T_{i,j} \in [t, t + n^{-1/3}u]).$$
(S.124)

We observe that the main part of $W_{n,1}(u)$ is a second order U statistic, and that $\mathbb{P}^2 f_{n,u}(\cdot, \cdot; \cdot, \cdot) = 0$; we have the decomposition (see Serfling, 1980, pages 177–178):

$$W_{n,1}(u) = n^{2/3} \left\{ \mathbb{U}_n^2 \bar{f}_{n,u} + \mathbb{P}_n f_{n,u,1} + \mathbb{P}_n f_{n,u,2} \right\},$$
(S.125)

where $\bar{f}_{n,u} = f_{n,u} - f_{n,u,1} - f_{n,u,2}$; $\mathbb{U}_n^2 \bar{f}_{n,u}$ is a degenerate U-statistic. For $\mathbb{U}_n^2 \bar{f}_{n,u}$, based on the discussion in van der Vaart and Wellner (1996) (page 98), we observe that the concept "covering number" in (S.136) can be replaced with "packing number" given by Definition 1 in Sherman (1994). Then, we immediately have that the function class

$$\mathcal{F}_n = \{ f_{n,u}(\cdot, \cdot, \cdot, \cdot) : u \in \mathbb{R}, \boldsymbol{\beta} \in \boldsymbol{\beta} \}$$
(S.126)

is Euclidean(C, 8p + 12) for envelope function F = 2 with C being a universal constant not depending on n and t; see Definition 3 in Sherman (1994) for the definition of Euclidean. With an application of Lemma 6 and then Corollary 4 in Sherman (1994), we conclude that

$$n \sup_{u \in \mathbb{R}, \boldsymbol{\beta} \in \mathcal{B}} \left| \mathbb{U}_n^2 \bar{f}_{n,u} \right| = O_p(1).$$
(S.127)

For $\mathbb{P}_n f_{n,u,1}$, we consider $[M', \infty) \subset \bigcup_{j=\lfloor M' \rfloor}^{\infty} A_j \equiv \bigcup_{j=\lfloor M' \rfloor}^{\infty} [j, (j+1)]$, where $\lfloor M' \rfloor$ denotes the largest integer that is less than or equal to M'. Then we have

$$P\left(\exists u > M': n^{2/3} \sup_{\boldsymbol{\beta} \in \mathcal{B}_{n,C}} |\mathbb{P}_n f_{n,u,1}| \ge \eta u^2 + M_1\right)$$

$$\leq \sum_{j=\lfloor M' \rfloor}^{\infty} P\left(\exists u \in A_j: n^{2/3} \sup_{\boldsymbol{\beta} \in \mathcal{B}_{n,C}} |\mathbb{P}_n f_{n,u,1}| \ge \eta u^2 + M_1\right)$$

$$\leq \sum_{j=\lfloor M' \rfloor}^{\infty} P\left(\exists u \in A_j : n^{2/3} \sup_{\boldsymbol{\beta} \in \mathcal{B}_{n,C}} |\mathbb{P}_n f_{n,u,1}| \geq \eta j^2 + M_1\right)$$

$$\leq n^{4/3} \sum_{j=1}^{\infty} \frac{E\left\{\sup_{u \in A_j, \boldsymbol{\beta} \in \mathcal{B}_{n,C}} |\mathbb{P}_n f_{n,u,1}|^2\right\}}{(\eta j^2 + M_1)^2}.$$
(S.128)

Note that $\mathcal{F}_{n,1,j} = \{f_{n,u,1} : u \in A_j, \beta \in \mathcal{B}_{n,C}\}$ *is a subset of the function class*

$$\mathcal{F}_{n,1} = \{ f_{n,u,1}(\boldsymbol{x}, y) = \mathbb{P}f_{n,u}(\cdot, \cdot, \boldsymbol{x}, y) : u \in \mathcal{R}, \boldsymbol{\beta} \in \mathcal{B} \},\$$

considered in Lemma 16. Based on the definition of $f_{n,u,1}$, $\mathcal{B}_{n,C}$, we can find a universal constant $C_1 > 0$, such that

$$\sup_{u \in A_{j}, \boldsymbol{\beta} \in \mathcal{B}_{n,C}} |f_{n,u,1}(\boldsymbol{x}, \boldsymbol{y})|$$

$$\leq C_{1} \int_{\boldsymbol{w} \in \mathcal{X}} I\left\{(\boldsymbol{w} - \boldsymbol{x})^{T} \boldsymbol{\beta}_{0} \in [t - C_{1}n^{-1/2}, t + n^{-1/3}(j+1) + C_{1}n^{-1/2}]\right\}$$

$$\times dF_{X}(\boldsymbol{w})$$

$$= C_{1} \left[F_{X^{T}} \boldsymbol{\beta}_{0} \left(\boldsymbol{x}^{T} \boldsymbol{\beta}_{0} + t + n^{-1/3}(j+1) + C_{1}n^{-1/2}\right) -F_{X^{T}} \boldsymbol{\beta}_{0} \left(\boldsymbol{x}^{T} \boldsymbol{\beta}_{0} + t - C_{1}n^{-1/2}\right)\right]$$

$$= \bar{F}(\boldsymbol{x}),$$

which serves as an envelope function of $\mathcal{F}_{n,1,j}$. Based on Condition 2, we have $\|\bar{F}\|_{2,P} \leq (j+1)n^{-1/3}$. Then, applying Theorem 11.1 in Kosorok (2008), we have

$$nE \left\{ \sup_{u \in A_{j}, \boldsymbol{\beta} \in \mathcal{B}_{n,C}} |\mathbb{P}_{n} f_{n,u,1}|^{2} \right\}$$

$$\lesssim \{J^{*}(1, \mathcal{F}_{n,1})\}^{2} \|\bar{F}\|_{2,P}^{2} \lesssim (j+1)^{2} n^{-2/3},$$
(S.129)

where we have used the fact that

$$J^{*}(1, \mathcal{F}_{n,1}) \equiv \sup_{Q} \int_{0}^{1} \sqrt{1 + H_{2}(\epsilon \|\bar{F}\|_{2,Q}, \mathcal{F}_{n,1,j}, Q)} d\epsilon \lesssim 1.$$

Combining (S.128) with (S.129) leads to

$$P\left(\exists u > M' : n^{2/3} \sup_{\beta \in \mathcal{B}_{n,C}} |\mathbb{P}_n f_{n,u,1}| \ge \eta u^2 + M_1\right)$$

$$\lesssim n^{-1/3} \sum_{j=\lfloor M' \rfloor}^{\infty} \frac{(j+1)^2}{(\eta j^2 + M_1)^2} < \epsilon/3,$$
(S.130)

with M_1 chosen to be sufficiently large. Using exactly the same strategy as the derivation for (S.130), we can obtain

$$P\left(\exists u > M' : n^{2/3} \sup_{\beta \in \mathcal{B}_{n,C}} |\mathbb{P}_n f_{n,u,2}| \ge \eta u^2 + M_1\right)$$

$$\lesssim n^{-1/3} \sum_{j=\lfloor M' \rfloor}^{\infty} \frac{(j+1)^2}{(\eta j^2 + M_1)^2} < \epsilon/3.$$
(S.131)

Combining (S.127), (S.130), and (S.131) leads to (S.122); we complete the proof for (a). We proceed to show (b). We can write

$$W_{n,2} = \frac{n^{2/3}}{n^2} \sum_{i,j} \{F_0(T_{i,j}) - F_0(t)\} I\left(T_{i,j} \in [t, t + n^{-1/3}u)\right)$$

$$\equiv \frac{n^{2/3}}{n^2} \sum_{i,j} g_{n,u}(X_i, X_j)$$

$$= \frac{n^{2/3}}{n^2} \sum_{i,j} \{g_{n,u}(X_i, X_j) - \mathbb{P}^2 g_{n,u}(\cdot, \cdot)\} + n^{2/3} \mathbb{P}^2 g_{n,u}(\cdot, \cdot).$$
(S.132)

Using exactly the same procedure as the proof for (a), we are able to show that for any $\eta > 0$, $\epsilon > 0$, and M' > 0, when M_1 is sufficiently large, we have

$$P\left(\exists u > M' : \left|\frac{n^{2/3}}{n^2} \sum_{i,j} \left\{ \widetilde{g}_{n,u}(X_i, X_j) - \mathbb{P}^2 \widetilde{g}_{n,u}(\cdot, \cdot) \right\} \right| \ge \eta u^2 + M_1 \right)$$

< $\epsilon.$ (S.133)

Furthermore, recall that $G_{\beta}(w)$ is the c.d.f. of $T_{i,j}$; based on Conditions F1 and F2 and the discussion in Remark 1, we can verify that there exists a universal constant c > 0, such that

$$n^{2/3} \inf_{\beta \in \mathcal{B}_{n,C}} \mathbb{P}^2 g_{n,u}(\cdot, \cdot)$$

= $n^{2/3} \inf_{\beta \in \mathcal{B}_{n,C}} \int_{w \in [t,t+n^{-1/3}u)} \{F_0(w) - F_0(t)\} dG_{\beta}(w) \ge cu^2.$ (S.134)

Combining (S.132)–(S.134) leads to (S.123); we complete the proof for (b).

Last, the proofs for (c) and (d) are straightforward, and are omitted.

Now, combining (a), (b), (c), and (d), we have that for any $\epsilon > 0$, when M' is sufficiently large,

$$P\left(\exists u > M' : \widetilde{W}_{n,1}(u) + \widetilde{W}_{n,2}(u) - \widetilde{W}_{n,3}(u) + \widetilde{W}_{n,4}(u) \leq 0\right) \qquad (S.135)$$

$$\leq P\left(\exists u > M' : \widetilde{W}_{n,2}(u) < cu^2\right)$$

$$+P\left(\exists u > M': cu^{2} \leqslant \widetilde{W}_{n,2}(u) \\ \leqslant -\widetilde{W}_{n,1}(u) + \widetilde{W}_{n,3}(u) - \widetilde{W}_{n,4}(u)\right)$$

$$\leqslant \epsilon + P\left(\exists u > M': cu^{2} \leqslant \widetilde{W}_{n,2}(u) \\ \leqslant |\widetilde{W}_{n,1}(u)| + |\widetilde{W}_{n,3}(u)| + |\widetilde{W}_{n,4}(u)|\right)$$

$$\leqslant \epsilon + P\left(\exists u > M': |\widetilde{W}_{n,1}(u)| \ge cu^{2}/3\right) \\ + P\left(\exists u > M': |\widetilde{W}_{n,3}(u)| \ge cu^{2}/3\right) \\ + P\left(\exists u > M': |\widetilde{W}_{n,4}(u)| \ge cu^{2}/3\right)$$

$$\leqslant \epsilon + \epsilon + \epsilon + \epsilon.$$

This together with (S.121) and that ϵ is arbitrary leads to (S.119). We complete the proof of this lemma.

In the proof of Lemma 15, we have used the entropy conditions given in the following lemma.

Lemma 16. Recall $f_{n,u}$ defined by (S.124). Consider the function classes

$$\begin{aligned} \mathcal{F}_n &= \{f_{n,u}(\cdot, \cdot; \cdot, \cdot) : u \in \mathbb{R}; \boldsymbol{\beta} \in \boldsymbol{\beta} \} \\ \mathcal{F}_{n,1} &= \{f_{n,u,1}(\boldsymbol{x}, y) = \mathbb{P}f_{n,u}(\cdot, \cdot, \boldsymbol{x}, y) : u \in \mathbb{R}, \boldsymbol{\beta} \in \boldsymbol{\beta} \} \\ \mathcal{F}_{n,1} &= \{f_{n,u,2}(\boldsymbol{x}, y) = \mathbb{P}f_{n,u}(\boldsymbol{x}, y, \cdot, \cdot) : u \in \mathbb{R}, \boldsymbol{\beta} \in \boldsymbol{\beta} \}. \end{aligned}$$

Then

(a) for any probability measure Q defined on the σ -algebra $\sigma(X_1, Y_1) \times \sigma(X_2, Y_2)$, $r \ge 1$ and any envelope function \overline{F} with $\|\overline{F}\|_{r,Q} > 0$; and any $0 < \epsilon < 1$, we have

$$N_r(\epsilon \|\bar{F}\|_{r,Q}, \mathcal{F}_n, Q) \lesssim (1/\epsilon)^{4pr+6r}, \tag{S.136}$$

where recall that p is the dimension of β ;

(b) for any probability measure Q defined on the σ -algebra $\sigma(X, Y)$, $r \ge 1$ and any envelope function \overline{F} with $\|\overline{F}\|_{r,Q} > 0$, we have

$$N_r(\epsilon \|\bar{F}\|_{r,Q}, \mathcal{F}_{n,1}, Q) \lesssim (1/\epsilon)^{4pr+6r}$$
 (S.137)

$$N_r(\epsilon \|\bar{F}\|_{r,Q}, \mathcal{F}_{n,2}, Q) \lesssim (1/\epsilon)^{4pr+6r}.$$
 (S.138)

Proof:Note that for any given t and n, based on Lemma 9.6 in Kosorok (2008), we have

that the function classes

$$\left\{ \boldsymbol{x}^{T}\boldsymbol{\beta} - t - n^{-1/2}\boldsymbol{u} : \boldsymbol{\beta} \in \boldsymbol{\mathcal{B}}, \boldsymbol{u} \in \mathbb{R} \right\} \\ \left\{ \boldsymbol{x}^{T}\boldsymbol{\beta} - t : \boldsymbol{\beta} \in \boldsymbol{\mathcal{B}} \right\}$$

are both VC-subgraph classes with VC indexes p + 3 and p + 2 respectively. Then, based on Lemma 9.9 (iii) and (iv) in Kosorok (2008), the sets

$$\mathcal{C}_1 = \{ \{ \boldsymbol{x}^T \boldsymbol{\beta} - t - n^{-1/2} u \leq 0 \} : \boldsymbol{\beta} \in \boldsymbol{\mathcal{B}}, u \in \mathbb{R} \}$$

$$\mathcal{C}_2 = \{ \{ \boldsymbol{x}^T \boldsymbol{\beta} - t \geq 0 \} : \boldsymbol{\beta} \in \boldsymbol{\mathcal{B}}, u \in \mathbb{R} \}$$

are both VC-classes of sets with $V_{C_1} = p + 3$ and $V_{C_2} = p + 2$. Applying Lemma 9.7 (ii) in Kosorok (2008), we conclude that

$$\left\{\left\{\boldsymbol{x}^{T}\boldsymbol{\beta}\in\left[t,t+n^{-1/2}u\right]\right\}:\boldsymbol{\beta}\in\boldsymbol{\mathcal{B}},u\in\mathbb{R}\right\}\subset\mathcal{C}_{1}\sqcap\mathcal{C}_{2}$$

is VC with index $\leq 2p + 5 - 1$. By Lemma 9.8, and Lemma 9.9 (vi) and (vii) in Kosorok (2008), we have that the function class \mathcal{F}_n is VC with index $\leq 2(2p + 4) - 1 = 4p + 7$.

Consequently, applying Theorem 9.3 in Kosorok (2008) leads to (S.136), based on which and applying Jensen inequality leads to (S.137) and (S.138). We complete the proof of this lemma. \blacksquare

Lemma 17. Assume all conditions for Theorem 3 are effective. For every M > 0, we have

$$n^{1/2} \sup_{x \in [-M,M]} \left| \widetilde{U}_n(s_0 + n^{-1/2}x) - t \right| = O_p(1).$$
(S.139)

Proof: Similar in strategy to the proof of Lemma 15, because of the monotonicity of $U_n(\cdot)$, we need to show only that for any $\epsilon > 0$, there exists M' > 0 such that

$$P\left(n^{1/2}\left\{\widetilde{U}_{n}(s_{0}+n^{-1/2}M)-t\right\} > M'\right) < \epsilon$$
(S.140)

$$P\left(n^{1/2}\left\{\widetilde{U}_{n}(s_{0}-n^{-1/2}M)-t\right\}<-M'\right)<\epsilon.$$
(S.141)

We need to show only (*S.140*); (*S.141*) *can be shown using exactly the same arguments. Note that*

$$P\left(n^{1/2}\left\{\widetilde{U}_{n}(s_{0}+n^{-1/2}M)-t\right\} > M'\right)$$

$$\leq P\left(t+n^{-1/2}M' < \widetilde{U}_{n}(s_{0}+n^{-1/2}M) \leq t+M_{2}n^{-1/3}\right)$$

$$+P\left(\widetilde{U}_{n}(s_{0}+n^{-1/2}M) \geq t+M_{2}n^{-1/3}\right).$$
(S.142)

Consider the right hand side of the above expression; based on Lemma 15, the second term can be arbitrarily small for sufficiently large M_2 ; therefore, we only need to consider the first

term. Recall that $s_0 = F_0(t)$ *, we have*

$$P\left(t+n^{-1/2}M' < \widetilde{U}_{n}(s_{0}+n^{-1/2}M) \leq t+M_{2}n^{-1/3}\right)$$

$$= P\left(\exists u \in [M', M_{2}n^{1/6}] : \widetilde{V}_{n}(t+n^{-1/2}u) - (s_{0}+n^{-1/2}M)\widetilde{G}_{n}(t+n^{-1/2}u) \\ \leq \widetilde{V}_{n}(t) - (s_{0}+n^{-1/2}M)\widetilde{G}_{n}(t)\right)$$

$$= P\left(\exists u \in [M', M_{2}n^{1/6}] : \widetilde{W}_{n,1}(u) + \widetilde{W}_{n,2}(u) - \widetilde{W}_{n,3}(u) + \widetilde{W}_{n,4} \leq 0\right),$$
(S.143)

where

$$\mathcal{W}_{n,1}(u) = n \left\{ V_n(t + n^{-1/2}u) - V_n(t) \right\}$$

$$-\frac{n}{n^2} \sum_{i,j} F_0(T_{i,j,0}) I(T_{i,j} \in [t, t + n^{-1/2}u])$$
(S.144)

$$\mathcal{W}_{n,2}(u) = n \int_{w \in [t,t+n^{-1/2}u]} \{F_0(w) - F_0(t)\} dG_n(w)$$
(S.145)

$$\mathcal{W}_{n,3}(u) = n^{1/2} M \left\{ G_n(t + n^{-1/2}u) - G_n(t) \right\}$$
(S.146)

$$\mathcal{W}_{n,4}(u) = \frac{n}{n^2} \sum_{i,j} \left[\{F_0(T_{i,j,0}) - F_0(T_{i,j})\} \\ \times I(T_{i,j} \in [t, t + n^{-1/2}u]) \right].$$
(S.147)

In analogous to "(a), (b), (c), (d)" in Lemma 15, we proceed to verify

(a') for any $\eta > 0, \epsilon > 0$, $M_2 > 0$, and M' > 0, when M_1 is sufficiently large, we have

$$P\left(\exists u \in [M', M_2 n^{1/6}] : \left|\widetilde{\mathcal{W}}_{n,1}(u)\right| \ge \eta u^2 + M_1\right) < \epsilon;$$
(S.148)

(b') there exists a universal constant c > 0, such that for any $\epsilon > 0$ and $M_2 > 0$, when M' is sufficiently large, we have

$$P\left(\exists u \in [M', M_2 n^{1/6}] : \widetilde{\mathcal{W}}_{n,2}(u) \ge c u^2\right) > 1 - \epsilon;$$
(S.149)

(c') there exists a universal constant C > 0, such that for any $\epsilon > 0$, $M_2 > 0$, and M > 0, when M' and M_1 are sufficiently large, we have

$$P\left(\exists u \in [M', M_2 n^{1/6}] : \widetilde{\mathcal{W}}_{n,3}(u) \ge Cu + M_1\right) < \epsilon;$$
(S.150)

(d') there exists a universal constant C > 0, such that for any $\epsilon > 0$, $M_2 > 0$, and M > 0, when M' and M_1 are sufficiently large, we have

$$P\left(\exists u \in [M', M_2 n^{1/6}] : \left|\widetilde{\mathcal{W}}_{n,4}(u)\right| \ge Cu + M_1\right) < \epsilon.$$

We show (a') first; note that

$$\mathcal{W}_{n,1}(u) = \frac{1}{n} \sum_{i,j} \{\Delta_{i,j} - F_0(T_{i,j,0})\} I(T_{i,j} \in [t, t + n^{-1/2}u])$$

$$\equiv \frac{1}{n} \sum_{i,j} \mathcal{L}_{n,u}(X_i, Y_i, X_j, Y_j),$$

where

$$\pounds_{n,u}(X_i, Y_i, X_j, Y_j) = \{\Delta_{i,j} - F_0(T_{i,j,0})\} I(T_{i,j} \in [t, t + n^{-1/2}u]).$$

Using similar developments as those from (S.125) to (S.127), we have the U-statistic decomposition

$$\mathcal{W}_{n,1}(u) = n \left\{ \mathbb{U}_n^2 \overline{\mathcal{L}}_{n,u} + \mathbb{P}_n \mathcal{L}_{n,u,1} + \mathbb{P}_n \mathcal{L}_{n,u,2} \right\},$$
(S.151)

where $\overline{\pounds}_{n,u} = \pounds_{n,u} - \pounds_{n,u,1} - \pounds_{n,u,2}$, such that $\mathbb{U}_n^2 \overline{\pounds}_{n,u}$ is a degenerate second order U-statistic, and that the function class

$$\mathfrak{S}_n = \left\{ n^{1/6} \mathcal{L}_{n,u} : u \in [M', M_2 n^{1/6}]; \boldsymbol{\beta} \in \mathcal{B}_{n,C} \right\}$$

is Euclidean(C, 8p + 12) with an envelope function

$$\bar{F}(X_i, Y_i, X_j, Y_j) = 2n^{1/6} I\left(T_{i,j} \in \left[t - C_1 n^{-1/2}, t + n^{-1/2} M_2 n^{1/6} + C_1 n^{-1/2}\right]\right)$$

for some universal constant $C_1 > 0$; based on Condition F2 and the discussion in Remark 1, it satisfies $\|\bar{F}\|_{2,P} \leq 1$. With an application of Lemma 6 and then Corollary 4 in Sherman (1994), we conclude that

$$n^{7/6} \sup_{u \in [M', M_2 n^{1/6}], \boldsymbol{\beta} \in \mathcal{B}_{n,C}} \left| \mathbb{U}_n^2 \overline{\mathcal{L}}_{n,u} \right| = O_p(1).$$
(S.152)

Furthermore, using exactly the same arguments as the proof for (S.130) and (S.131), we can establish

$$P\left(\exists u > M': n \sup_{\boldsymbol{\beta} \in \mathcal{B}_{n,C}} |\mathbb{P}_n \mathcal{L}_{n,u,1}| \ge \eta u^2 + M_1\right)$$

$$\lesssim \sum_{j=\lfloor M' \rfloor}^{\infty} \frac{(j+1)^2}{(\eta j^2 + M_1)^2} < \epsilon/3 \qquad (S.153)$$

$$P\left(\exists u > M': n \sup_{\boldsymbol{\beta} \in \mathcal{B}_{n,C}} |\mathbb{P}_n \mathcal{L}_{n,u,2}| \ge \eta u^2 + M_1\right)$$

$$\lesssim \sum_{j=\lfloor M' \rfloor}^{\infty} \frac{(j+1)^2}{(\eta j^2 + M_1)^2} < \epsilon/3, \qquad (S.154)$$

when M_1 is sufficiently large.

Combining (S.152)–(S.154) leads to (S.148); we complete the proof for (a'). We consider (b') next. We can write

$$\mathcal{W}_{n,2}(u) = n \int_{w \in [t,t+n^{-1/2}u]} \{F_0(w) - F_0(t)\} d\{G_n(w) - G_{\beta}(w)\}$$

+ $n \int_{w \in [t,t+n^{-1/2}u]} \{F_0(w) - F_0(t)\} dG_{\beta}(w)$
= $\mathcal{W}_{n,2,1}(u) + \mathcal{W}_{n,2,2}(u).$ (S.155)

Using exactly the same procedure as the proof for (a'), we have the decomposition

$$\mathcal{W}_{n,2,1} = n \left[\mathbb{U}_n^2 \overline{\varrho}_{n,u} + \left(\mathbb{P}_n \varrho_{n,u,1} - \mathbb{P}^2 \varrho_{n,u} \right) + \left\{ \mathbb{P}_n \varrho_{n,u,2} - \mathbb{P}^2 \varrho_{n,u} \right\} \right],$$
(S.156)

where $\overline{\varrho}_{n,u} = \varrho_{n,u} - \varrho_{n,u,1} - \varrho_{n,u,2} + \mathbb{P}^2 \varrho_{n,u}$, $\varrho_{n,u,1}(X_j, Y_j) = \mathbb{P} \varrho_{n,u}(\cdot, \cdot; X_j, Y_j)$, $\varrho_{n,u,2}(X_i, Y_i) = \mathbb{P} \varrho_{n,u}(X_i, Y_i; \cdot, \cdot)$ with

$$\varrho_{n,u}(X_i, Y_i; X_j, Y_j) = \{F_0(T_{i,j}) - F_0(t)\} I \left(T_{i,j} \in [t, t + n^{-1/2}u)\right)$$

$$\mathbb{P}^2 \varrho_{n,u} = \mathcal{W}_{n,2,2}(u) = \int_{w \in [t, t+n^{-1/2}u)} \{F_0(w) - F_0(t)\} dG_{\boldsymbol{\beta}}(w).(S.157)$$

Note that $\mathbb{U}_n^2 \overline{\varrho}_{n,u}$ is a degenerate U-statistic. Using exactly the same developments as those for (S.152), we have

$$n^{7/6} \sup_{u \in [M', M_2 n^{1/6}], \boldsymbol{\beta} \in \mathcal{B}_{n,C}} \left| \mathbb{U}_n^2 \overline{\varrho}_{n,u} \right| = O_p(1)$$
(S.158)

Then, using exactly the same arguments as from (S.151) to (S.154), we are able to verify that for any $\eta > 0$, $\epsilon > 0$, M_2 , and M' > 0, when M_1 is sufficiently large,

$$P\left(\exists u \in [M', M_2 n^{1/6}] : \widetilde{\mathcal{W}}_{n,2,1}(u) \ge \eta u^2 + M_1\right) < \epsilon.$$
(S.159)

Furthermore, based on Conditions F1 and F2 and the discussion given in Remark 1, we can find a universal constant c > 0, such that

$$\mathcal{W}_{n,2,2} = n \int_{w \in [t,t+n^{-1/2}u)} \{F_0(w) - F_0(t)\} dG_{\beta}(w) \ge cu^2.$$
(S.160)

Combining (S.155)–(S.160) leads to (S.149); we complete the proof for (b').

We proceed to consider (c'). Using a similar strategy as the proof for (a'), we can show that

$$\sup_{u \in [M', M_2 n^{1/6}], \boldsymbol{\beta} \in \mathcal{B}_{n, C}} n^{1/2} M \left| \left\{ G_n(t + n^{-1/2} u) - G_n(t) \right\} \right.$$

$$-\left\{G_{\beta}(t+n^{-1/2}u) - G_{\beta}(t)\right\} = o_p(1). \quad (S.161)$$

Furthermore, based on Condition F2 and the discussion given in Remark 1, there exists a universal constant $C_1 > 0$, such that

$$n^{1/2} \sup_{\boldsymbol{\beta} \in \mathcal{B}_{n,C}} \left| G_{\boldsymbol{\beta}}(t+n^{-1/2}u) - G_{\boldsymbol{\beta}}(t) \right| \leq C_1 u.$$
(S.162)

Combining (S.161) with (S.162) leads to (S.150), and we complete the proof for (c').

Last, we consider (d'). Note that $\tilde{\boldsymbol{\beta}} - \boldsymbol{\beta}_0 = O_p(n^{-1/2})$ and based on Condition F1 and the discussion given in Remark 1, with probability arbitrarily large,

$$\begin{aligned} \mathcal{W}_{n,4}(u) &| &\lesssim \quad \frac{n^{1/2}}{n^2} \sum_{i,j} I(T_{i,j} \in [t, t + n^{-1/2}u]) \\ &= \quad n^{1/2} \left\{ G_n(t + n^{-1/2}u) - G_n(t) \right\} \\ &= \quad \mathcal{W}_{n,3}(u) / M. \end{aligned}$$

Therefore, the conclusion of (d') follows by using the result of (c').

Finally, combining (a')–(d'), and using a similar development to (S.135), and incorporating (S.142) and (S.143) leads to (S.140). We complete the proof of this lemma.

We need to apply the following Lemma 18, which is the argmax theorem adapted from Theorem 14.1 in Kosorok (2008) (see also Theorem 3.2.2 in van der Vaart and Wellner, 1996) to establish the asymptotic normality of $n^{1/2} \left\{ \widetilde{U}_n(s + n^{-1/2}x) - t \right\}$.

Lemma 18. Let W_n , W be stochastic processes indexed by a metric space \mathcal{H} , such that $W_n \rightsquigarrow W$ in $L^{\infty}(H)$ for every compact $H \subset \mathcal{H}$. Suppose also that almost all sample paths $h \mapsto W(h)$ are upper semicontinuous and possess a unique maximum at a (random) point \hat{h} , which as a random map in \mathcal{H} is tight. If the sequence \hat{h}_n is uniformly tight and satisfies $W_n(\hat{h}_n) \ge \sup_{h \in H} W_n(h) - o_p(1)$, then $\hat{h}_n \rightsquigarrow \hat{h}$ in \mathcal{H} .

Lemma 19. Assume all conditions for Theorem 3 are effective. We have

$$n^{1/2} \left\{ \widetilde{U}_n(s_0 + n^{-1/2}x) - t \right\} \leadsto \frac{g_0(t)x - \sigma(t)Z}{f_0(t)g_0(t)}.$$
(S.163)

Proof: We shall apply Lemma 18 to establish the asymptotic result (S.163). Note that in Lemma 17, we have verified that $n^{1/2} \left\{ \tilde{U}_n(s_0 + n^{-1/2}x) - t \right\}$ is uniformly tight. Therefore, to apply the argmax theorem, i.e., Lemma 18, we need to establish the asymptotic convergence of the corresponding stochastic process that it minimises.

Based on the definition of U_n given in (S.115) and note that $s_0 = F_0(t)$, we have

$$n^{1/2} \left\{ \widetilde{U}_{n}(s_{0} + n^{-1/2}x) - t \right\}$$

= $\arg \min_{u} \left[n \left\{ \widetilde{V}_{n}(t + n^{-1/2}u) + (s_{0} + n^{-1/2}x)\widetilde{G}_{n}(t + n^{-1/2}u) \right\} \right]$
= $\arg \min_{u} \left\{ \widetilde{\mathcal{W}}_{n,1}(u) + \widetilde{\mathcal{W}}_{n,2}(u) - \widetilde{\mathcal{W}}_{n,5}(u) + \widetilde{\mathcal{W}}_{n,4}(u) \right\}$ (S.164)

where $\widetilde{W}_{n,1}(u)$, $\widetilde{W}_{n,2}(u)$, and $\widetilde{W}_{n,4}(u)$ are respectively defined by (S.144), (S.145), and (S.147); and

$$\mathcal{W}_{n,5}(u) = n^{1/2} x \left\{ G_n(t + n^{-1/2}u) - G_n(t) \right\}.$$

Consider $W_{n,4}(u)$. Recall

=

$$\mathcal{W}_{n,4}(u) = \frac{n}{n^2} \sum_{i,j} \left\{ F_0(T_{i,j,0}) - F_0(T_{i,j}) \right\} I(T_{i,j} \in [t, t + n^{-1/2}u])$$

$$= \frac{n}{n^2} \sum_{i,j} h_{n,u}(X_i, X_j)$$

$$= \frac{n}{n^2} \sum_{i,j} \left\{ h_{n,u}(X_i, X_j) - \mathbb{P}^2 h_{n,u} \right\} + n \mathbb{P}^2 h_{n,u}.$$
 (S.165)

Then, applying exactly the same arguments as (S.151) and (S.152), we have uniformly in $u \in [-M, M]$ and $\beta \in \mathcal{B}_{n,C}$,

$$\frac{n}{n^2} \sum_{i,j} \left\{ h_{n,u}(X_i, X_j) - \mathbb{P}^2 h_{n,u} \right\}$$

= $n \left\{ \mathbb{P}_n h_{n,u,1} - \mathbb{P}^2 h_{n,u} \right\} + n \left\{ \mathbb{P}_n h_{n,u,2} - \mathbb{P}^2 h_{n,u} \right\} + o_p(1),$ (S.166)

where $h_{n,u,1}(\boldsymbol{x}) = \mathbb{P}h_{n,u}(\cdot, \boldsymbol{x})$ and $h_{n,u,2}(\boldsymbol{x}) = \mathbb{P}h_{n,u}(\boldsymbol{x}, \cdot)$. Furthermore, recall the definition $F_{\boldsymbol{\beta}}(s) = E\left\{I(Y_1 > Y_2) \middle| (X_1 - X_2)^T \boldsymbol{\beta} = s\right\}$ and denote $\dot{F}_0(s) = \frac{\partial^F \boldsymbol{\beta}^{(s)}}{\partial \boldsymbol{\beta}} \middle|_{\boldsymbol{\beta} = \boldsymbol{\beta}_0}$. Then based on Condition A1, $\dot{F}_0(s)$ is continuous in s; and based on Condition F2, $g_{\boldsymbol{\beta}}(s) = G'_{\boldsymbol{\beta}}(s)$ is continuous in the neighbourhood of s = t and $\boldsymbol{\beta} = \boldsymbol{\beta}_0$. We have that uniformly in $u \in [-M, M]$ and $\boldsymbol{\beta} \in \mathcal{B}_{n,C}$,

$$n\mathbb{P}^{2}h_{n,u} = nE\left[\left\{\Delta_{i,j} - F_{0}(T_{i,j})\right\}I(T_{i,j} \in [t, t + n^{-1/2}u])\right]$$

$$= nE\left(E\left[\left\{\Delta_{i,j} - F_{0}(T_{i,j})\right\}I(T_{i,j} \in [t, t + n^{-1/2}u])\Big|T_{i,j}\right]\right)$$

$$= nE\left[\left\{F_{\beta}(T_{i,j}) - F_{0}(T_{i,j})\right\}I(T_{i,j} \in [t, t + n^{-1/2}u])\right]$$

$$= n(\beta - \beta_{0})^{T}E\left\{\dot{F}_{0}(T_{i,j})I(T_{i,j} \in [t, t + n^{-1/2}u])\right\}\{1 + o(1)\}$$

$$= n(\boldsymbol{\beta} - \boldsymbol{\beta}_0)^T \{1 + o(1)\} \int_t^{t+n^{-1/2}u} \dot{F}_0(w) dG_{\boldsymbol{\beta}}(w)$$

$$= \sqrt{n}(\boldsymbol{\beta} - \boldsymbol{\beta}_0)^T \dot{F}_0(t) ug_0(t) \{1 + o(1)\}.$$
(S.167)

Combining (S.151), (S.152), (S.155), (S.156), (S.158), (S.161), (S.165), and (S.166), we have uniformly in $u \in [-M, M]$ and $\beta \in \mathcal{B}_{n,C}$,

$$\mathcal{W}_{n,1}(u) + \mathcal{W}_{n,2}(u) - \mathcal{W}_{n,5}(u) + \mathcal{W}_{n,4}(u)
= n\mathbb{P}_n(\mathcal{L}_{n,u,1} + \mathcal{L}_{n,u,2} + \varrho_{n,u,1} + \varrho_{n,u,2} + h_{n,u,1} + h_{n,u,2})
-2n\mathbb{P}^2(\varrho_{n,u} + h_{n,u}) + n\mathbb{P}^2\varrho_{n,u} + n\mathbb{P}^2h_{n,u}
-n^{1/2}x\left\{G_{\beta}(t + n^{-1/2}u) - G_{\beta}(t)\right\} + o_p(1)
= \mathbb{G}_n(p_{n,u,1} + p_{n,u,2}) + n\mathbb{P}^2\varrho_{n,u} + n\mathbb{P}^2h_{n,u}
-n^{1/2}x\left\{G_{\beta}(t + n^{-1/2}u) - G_{\beta}(t)\right\} + o_p(1),$$
(S.168)

where $p_{n,u,1}(X_j, Y_j) = \mathbb{P}p_{n,u}(\cdot, \cdot; X_j, Y_j), p_{n,u,2}(X_i, Y_i) = \mathbb{P}p_{n,u}(X_i, Y_i; \cdot, \cdot)$ with

$$p_{n,u}(X_i, Y_i; X_j, Y_j) = \sqrt{n} \{ \Delta_{i,j} - F_0(t) \} I(T_{i,j} \in [t, t + n^{-1/2}u]),$$

and therefore

$$p_{n,u,1}(\boldsymbol{x}, y) = \sqrt{n} \int_{\boldsymbol{w}^T \boldsymbol{\beta} \in [\boldsymbol{x}^T \boldsymbol{\beta} + t, \boldsymbol{x}^T \boldsymbol{\beta} + t + n^{-1/2}u]} \left\{ 1 - F_{\epsilon}(H(y) - \boldsymbol{w}^T \boldsymbol{\beta}_0) - F_0(t) \right\} dF_X(\boldsymbol{w})$$

$$p_{n,u,2}(\boldsymbol{x}, y) = \sqrt{n} \int_{\boldsymbol{w}^T \boldsymbol{\beta} \in [\boldsymbol{x}^T \boldsymbol{\beta} - t - n^{-1/2}u, \boldsymbol{x}^T \boldsymbol{\beta} - t]} \left\{ F_{\epsilon}(H(y) - \boldsymbol{w}^T \boldsymbol{\beta}_0) - F_0(t) \right\} dF_X(\boldsymbol{w}).$$

Denote

$$\begin{split} & \breve{p}_{n,u,1}(\boldsymbol{x}, y) \\ &= \sqrt{n} \int_{\boldsymbol{w}^T \boldsymbol{\beta} \in [\boldsymbol{x}^T \boldsymbol{\beta} + t, \boldsymbol{x}^T \boldsymbol{\beta} + t + n^{-1/2} u]} \left\{ 1 - F_{\epsilon}(H(y) - \boldsymbol{w}^T \boldsymbol{\beta}) - F_{0}(t) \right\} dF_X(\boldsymbol{w}) \\ &= \sqrt{n} \int_{\boldsymbol{x}^T \boldsymbol{\beta} + t}^{\boldsymbol{x}^T \boldsymbol{\beta} + t + n^{-1/2} u} \left\{ 1 - F_{\epsilon}(H(y) - w) - F_{0}(t) \right\} dF_{X^T \boldsymbol{\beta}}(w) \\ & \breve{p}_{n,u,2}(\boldsymbol{x}, y) \\ &= \sqrt{n} \int_{\boldsymbol{w}^T \boldsymbol{\beta} \in [\boldsymbol{x}^T \boldsymbol{\beta} - t - n^{-1/2} u, \boldsymbol{x}^T \boldsymbol{\beta} - t]} \left\{ F_{\epsilon}(H(y) - \boldsymbol{w}^T \boldsymbol{\beta}) - F_{0}(t) \right\} dF_X(\boldsymbol{w}) \\ &= \sqrt{n} \int_{\boldsymbol{x}^T \boldsymbol{\beta} - t}^{\boldsymbol{x}^T \boldsymbol{\beta} - t} \left\{ F_{\epsilon}(H(y) - w) - F_{0}(t) \right\} dF_{X^T \boldsymbol{\beta}}(w). \end{split}$$

Then, based on Condition F1, considering the function class

$$\{p_{n,u,1} - \breve{p}_{n,u,1} : u \in [-M, M], \boldsymbol{\beta} \in \mathcal{B}_{n,C}\},\$$

and applying Theorem 11.1 in Kosorok (2008), we can establish

$$\sup_{u \in [-M,M], \boldsymbol{\beta} \in \mathcal{B}_{n,C}} |\mathbb{G}_n(p_{n,u,1} - \breve{p}_{n,u,1})| \lesssim n^{-1/2}.$$
(S.169)

Similarly,

$$\sup_{u\in[-M,M],\boldsymbol{\beta}\in\mathcal{B}_{n,C}} |\mathbb{G}_n(p_{n,u,2}-\breve{p}_{n,u,2})| \lesssim n^{-1/2}.$$
(S.170)

We further consider

$$\bar{p}_{n,u,1,\boldsymbol{\beta}}(\boldsymbol{x},y) = u \left\{ 1 - F_{\epsilon}(H(y) - \boldsymbol{x}^{T}\boldsymbol{\beta} - t) - F_{0}(t) \right\} f_{X^{T}\boldsymbol{\beta}}(\boldsymbol{x}^{T}\boldsymbol{\beta} + t)$$

$$\bar{p}_{n,u,2,\boldsymbol{\beta}}(\boldsymbol{x},y) = u \left\{ F_{\epsilon}(H(y) - \boldsymbol{x}^{T}\boldsymbol{\beta} + t) - F_{0}(t) \right\} f_{X^{T}\boldsymbol{\beta}}(\boldsymbol{x}^{T}\boldsymbol{\beta} - t).$$

Based on Conditions F1 and F2, considering the function classes

$$\left\{ \breve{p}_{n,u,1} - \bar{p}_{n,u,1,\boldsymbol{\beta}} : u \in [-M, M], \boldsymbol{\beta} \in \mathcal{B}_{n,C} \right\}$$
$$\left\{ \bar{p}_{n,u,1,\boldsymbol{\beta}} : u \in [-M, M], \boldsymbol{\beta} \in \mathcal{B}_{n,C} \right\},$$

and applying Theorem 11.1 in Kosorok (2008), we can obtain

$$\sup_{u \in [-M,M], \boldsymbol{\beta} \in \mathcal{B}_{n,C}} \left| \mathbb{G}_n(\breve{p}_{n,u,1} - \bar{p}_{n,u,1,\boldsymbol{\beta}}) \right| = O_p(n^{-1/2})$$
(S.171)

$$\sup_{u \in [-M,M], \boldsymbol{\beta} \in \mathcal{B}_{n,C}} \left| \mathbb{G}_n(\bar{p}_{n,u,1,\boldsymbol{\beta}} - \bar{p}_{n,u,1,\boldsymbol{\beta}_0}) \right| = O_p(n^{-1/2}).$$
(S.172)

Likewise, we are able to derive

$$\sup_{u \in [-M,M], \boldsymbol{\beta} \in \mathcal{B}_{n,C}} \left| \mathbb{G}_n(\breve{p}_{n,u,2} - \bar{p}_{n,u,2,\boldsymbol{\beta}}) \right| = O_p(n^{-1/2})$$
(S.173)

$$\sup_{u \in [-M,M], \boldsymbol{\beta} \in \mathcal{B}_{n,C}} \left| \mathbb{G}_n(\bar{p}_{n,u,2,\boldsymbol{\beta}} - \bar{p}_{n,u,2,\boldsymbol{\beta}_0}) \right| = O_p(n^{-1/2}).$$
(S.174)

Combining (S.168)–(S.174) and noting (S.157) and (S.167), we have uniformly in $u \in [-M, M]$ and $\beta \in \mathcal{B}_{n,C}$,

$$\mathcal{W}_{n,1}(u) + \mathcal{W}_{n,2}(u) - \mathcal{W}_{n,5}(u) + \mathcal{W}_{n,4}(u)$$

= $\mathbb{G}_n(\bar{p}_{n,u,1,\beta_0} + \bar{p}_{n,u,2,\beta_0}) + n \int_{w \in [t,t+n^{-1/2}u]} \{F_0(w) - F_0(t)\} \, dG_{\beta}(w)$

$$\begin{aligned} &+\sqrt{n}(\boldsymbol{\beta}-\boldsymbol{\beta}_{0})^{T}\dot{F}_{0}(t)ug_{0}(t)\left\{1+o(1)\right\}-n^{1/2}x\left\{G_{\boldsymbol{\beta}}(t+n^{-1/2}u)-G_{\boldsymbol{\beta}}(t)\right\}+o_{p}(1)\\ &= \mathbb{G}_{n}(\bar{p}_{n,u,1,\boldsymbol{\beta}_{0}}+\bar{p}_{n,u,2,\boldsymbol{\beta}_{0}})+n\int_{w\in[t,t+n^{-1/2}u]}\left\{F_{0}(w)-F_{0}(t)\right\}dG_{0}(w)\\ &+\sqrt{n}(\boldsymbol{\beta}-\boldsymbol{\beta}_{0})^{T}\dot{F}_{0}(t)ug_{0}(t)\left\{1+o(1)\right\}-xug_{0}(t)+o_{p}(1)\\ &= \mathbb{G}_{n}(\bar{p}_{n,u,1,\boldsymbol{\beta}_{0}}+\bar{p}_{n,u,2,\boldsymbol{\beta}_{0}})+\sqrt{n}(\boldsymbol{\beta}-\boldsymbol{\beta}_{0})^{T}\dot{F}_{0}(t)ug_{0}(t)+\frac{1}{2}g_{0}(t)f_{0}(t)u^{2}-g_{0}(t)xu+o_{p}(1).\end{aligned}$$

Replacing β with $\tilde{\beta}$ in the expression above and based on Theorem 2 and the proof of Corollary 1 that

$$\sqrt{n}\left(\widetilde{\boldsymbol{\beta}}-\boldsymbol{\beta}_{0}\right)=2A^{-1}\widetilde{A}^{-1}\psi_{0}'(\boldsymbol{\beta}_{0})\Sigma_{X}^{-1}\mathbb{G}_{n}\hbar+o_{p}(1)$$

leads to

$$\widetilde{\mathcal{W}}_{n,1}(u) + \widetilde{\mathcal{W}}_{n,2}(u) - \widetilde{\mathcal{W}}_{n,5}(u) + \widetilde{\mathcal{W}}_{n,4}(u)$$

= $u \mathbb{G}_n \zeta_t + \frac{1}{2} g_0(t) f_0(t) u^2 - g_0(t) x u + o_p(1),$

where

$$\begin{aligned} \zeta_t(\boldsymbol{x}, y) &= \left\{ 1 - F_{\epsilon}(H(y) - \boldsymbol{x}^T \boldsymbol{\beta}_0 - t) - F_0(t) \right\} f_{X^T \boldsymbol{\beta}_0}(\boldsymbol{x}^T \boldsymbol{\beta}_0 + t) \\ &+ \left\{ F_{\epsilon}(H(y) - \boldsymbol{x}^T \boldsymbol{\beta}_0 + t) - F_0(t) \right\} f_{X^T \boldsymbol{\beta}_0}(\boldsymbol{x}^T \boldsymbol{\beta}_0 - t) \\ &+ 2g_0(t) \dot{F}_0^T(t) A^{-1} \widetilde{A}^{-1} \psi_0'(\boldsymbol{\beta}_0) \Sigma_X^{-1} \hbar(\boldsymbol{x}, y). \end{aligned}$$

Denoting $\sigma(t) = var(\zeta_t(X_1, Y_1))$, and applying central limit theorem, we have

$$\widetilde{\mathcal{W}}_{n,1}(u) + \widetilde{\mathcal{W}}_{n,2}(u) - \widetilde{\mathcal{W}}_{n,5}(u) + \widetilde{\mathcal{W}}_{n,4}(u)$$

$$\longleftrightarrow u\sigma(t)Z + \frac{1}{2}g_0(t)f_0(t)u^2 - g_0(t)xu,$$

where Z is a standard normal random variable. Applying Lemma 18, we have

$$n^{1/2} \left\{ \widetilde{U}_n(s+n^{-1/2}x) - t \right\} \leadsto \frac{g_0(t)x - \sigma(t)Z}{f_0(t)g_0(t)}$$

which completes our proof of this lemma.

7 Proof of Corollary 2

We give a sketched proof of Corollary 2; we omit details that are similar to those in the proof of Theorem 3 to avoid lengthy repetition. With the switching relation and using similar devel-

opments to (S.116), we can obtain

$$P\left(n^{1/2}\left\{\widehat{F}_{\boldsymbol{\beta}}(t) - F_{\boldsymbol{\beta}}(t)\right\} \ge x\right)$$

$$= P\left(n^{1/2}\left\{U_n\left(s_{\boldsymbol{\beta}} + n^{-1/2}x\right) - t\right\} \le 0\right),$$
(S.175)

where $s_{\beta} = F_{\beta}(t)$. Therefore, we only need to consider the asymptotics of the process $n^{1/2} \left\{ U_n \left(s_{\beta} + n^{-1/2} x \right) - t \right\}$. For any $t \in \mathbb{R}$ and $\beta \in \beta$, using similar arguments as Lemmas 15 and 17, assuming that

For any $t \in \mathbb{R}$ and $\beta \in \beta$, using similar arguments as Lemmas 15 and 17, assuming that all the technical conditions are effective, we are able to verify that for any M > 0,

$$n^{1/2} \sup_{x \in [-M,M]} \left| U_n \left(s_{\beta} + n^{-1/2} x \right) - t \right| = O_p(1).$$

Then, similar to the developments in (S.164) and (S.168), we can verify

$$n^{1/2} \left[U_n \left(s_{\beta} + n^{-1/2} x \right) - t \right]$$

= $\arg \min_u \left[\mathbb{G}_n(q_{n,u,1} + q_{n,u,2}) + n \int_{w \in [t,t+n^{-1/2}u]} \left\{ F_{\beta}(w) - F_{\beta}(t) \right\} dG_{\beta}(w) + n^{1/2} x \left\{ G_{\beta}(t+n^{-1/2}u) - G_{\beta}(t) \right\} + o_p(1) \right],$

where $o_p(1)$ is uniform in $u \in [-M, M]$ and $\boldsymbol{\beta} \in \boldsymbol{\mathcal{B}}$; $q_{n,u,1}(X_j, Y_j) = \mathbb{P}q_{n,u}(\cdot, \cdot; X_j, Y_j)$, $q_{n,u,2}(X_i, Y_i) = \mathbb{P}q_{n,u}(X_i, Y_i; \cdot, \cdot)$ with

$$\begin{split} q_{n,u} &= \sqrt{n} \left\{ \Delta_{i,j} - F_{\pmb{\beta}}(t) \right\} I(T_{i,j} \in [t, t + n^{-1/2}u]). \\ q_{n,u,1}(\pmb{x}, y) &= \sqrt{n} \int_{\pmb{x}^T \pmb{\beta} + t}^{\pmb{x}^T \pmb{\beta} + t + n^{-1/2}u} \left\{ 1 - F_{\epsilon, \pmb{\beta}}(y, w) - F_{\pmb{\beta}}(t) \right\} dF_{X^T \pmb{\beta}}(w) \\ q_{n,u,2}(\pmb{x}, y) &= \sqrt{n} \int_{\pmb{x}^T \pmb{\beta} - t}^{\pmb{x}^T \pmb{\beta} - t} \left\{ F_{\epsilon, \pmb{\beta}}(y, w) - F_{\pmb{\beta}}(t) \right\} dF_{X^T \pmb{\beta}}(w) \\ F_{\epsilon, \pmb{\beta}}(y, w) &= E \left\{ F_{\epsilon}(H(y) - X^T \pmb{\beta}_0) \middle| X^T \pmb{\beta} = w \right\}. \end{split}$$

Consider

$$\begin{split} \bar{q}_{n,u,1}(\boldsymbol{x},y) &= u \left\{ 1 - F_{\boldsymbol{\epsilon},\boldsymbol{\beta}}(y,\boldsymbol{x}^{T}\boldsymbol{\beta} + t) - F_{\boldsymbol{\beta}}(t) \right\} f_{X^{T}\boldsymbol{\beta}}(\boldsymbol{x}^{T}\boldsymbol{\beta} + t) \\ \bar{q}_{n,u,2}(\boldsymbol{x},y) &= u \left\{ F_{\boldsymbol{\epsilon},\boldsymbol{\beta}}(y,\boldsymbol{x}^{T}\boldsymbol{\beta} - t) - F_{\boldsymbol{\beta}}(t) \right\} f_{X^{T}\boldsymbol{\beta}}(\boldsymbol{x}^{T}\boldsymbol{\beta} - t). \end{split}$$

Then based on Conditions F1' and F2', considering the function classes

$$\left\{q_{n,u,1} - \bar{q}_{n,u,1} : u \in [-M, M], \boldsymbol{\beta} \in \boldsymbol{\mathcal{B}}\right\}$$

and applying Theorem 11.1 in Kosorok (2008), we can obtain

$$\sup_{u \in [-M,M], \boldsymbol{\beta} \in \mathcal{B}} |\mathbb{G}_n(q_{n,u,1} - \bar{q}_{n,u,1})| = O_p(n^{-1/2}), \qquad (S.176)$$

for any M > 0. Similarly, we have

$$\sup_{u \in [-M,M], \boldsymbol{\beta} \in \mathcal{B}} |\mathbb{G}_n(q_{n,u,2} - \bar{q}_{n,u,2})| = O_p(n^{-1/2}).$$
(S.177)

,

Therefore, we have

$$\begin{aligned} \mathbb{G}_{n}(q_{n,u,1} + q_{n,u,2}) + n \int_{w \in [t,t+n^{-1/2}u]} \left\{ F_{\beta}(w) - F_{\beta}(t) \right\} dG_{\beta}(w) \\ &+ n^{1/2}x \left\{ G_{\beta}(t+n^{-1/2}u) - G_{\beta}(t) \right\} \\ = & \mathbb{G}_{n}(\bar{q}_{n,u,1} + \bar{q}_{n,u,2}) + \frac{u^{2}}{2} f_{\beta}(t) g_{\beta}(t) + uxg_{\beta}(t) + o_{p}(1) \\ & \longleftrightarrow u\sigma_{\beta}(t)Z + \frac{u^{2}}{2} f_{\beta}(t)g_{\beta}(t) + uxg_{\beta}(t), \end{aligned}$$

where $o_p(1)$ is uniform in $u \in [-M, M]$ and $\beta \in \beta$; the weak convergence is in $L^{\infty}[-M, M]$ for any given $\beta \in \beta$; Z is a standard normal random variable; and

$$\sigma_{\boldsymbol{\beta}}(t) = \operatorname{var}\left\{\bar{q}_{n,u,1}(X_{1}, Y_{1}) + \bar{q}_{n,u,2}(X_{1}, Y_{1})\right\}$$
$$= E\left[\operatorname{var}\left\{f_{X^{T}\boldsymbol{\beta}}(X_{1}^{T}\boldsymbol{\beta} + t)F_{\epsilon,\boldsymbol{\beta}}(Y_{1}, X_{1}^{T}\boldsymbol{\beta} + t) - f_{X^{T}\boldsymbol{\beta}}(X_{1}^{T}\boldsymbol{\beta} - t)F_{\epsilon,\boldsymbol{\beta}}(Y_{1}, X_{1}^{T}\boldsymbol{\beta} - t)\right|X_{1}\right\}$$

by noting that $E\{\bar{q}_{n,u,1}(X_1,Y_1)\} = E\{\bar{q}_{n,u,2}(X_1,Y_1)\} = 0$. Applying Lemma 18, we have

$$n^{1/2}\left\{U_n\left(s_{\boldsymbol{\beta}}+n^{-1/2}x\right)-t\right\} \leadsto \frac{g_{\boldsymbol{\beta}}(t)x-\sigma_{\boldsymbol{\beta}}(t)Z}{f_{\boldsymbol{\beta}}(t)g_{\boldsymbol{\beta}}(t)},$$

which together with (S.175) completes the proof of this corollary.

8 Some Details for the Numerical Algorithms of our Estimates

In this section, we provide some details of the numerical algorithms for computing our $\hat{\beta}$ and $\tilde{\beta}$ estimates.

8.1 Some details of the numeral algorithm for $\hat{\beta}$

Recall that our $\hat{\beta}$ estimate is obtained by maximizing the profile pairwise rank log-likelihood:

$$\ell(\boldsymbol{\beta}, \widehat{F}_{\boldsymbol{\beta}}) = \sum_{i \neq j} \left[I(Y_i > Y_j) \log\{\widehat{F}_{\boldsymbol{\beta}}((X_i - X_j)^T \boldsymbol{\beta})\} + I(Y_i \leqslant Y_j) \log\{1 - \widehat{F}_{\boldsymbol{\beta}}((X_i - X_j)^T \boldsymbol{\beta})\} \right],$$

subject to the constraint $\|\beta\|_2 = 1$.

To incorporate the constraint $\|\boldsymbol{\beta}\|_2 = 1$, we consider the following polar transformation for $\boldsymbol{\beta} = (\beta_1, \dots, \beta_p)^T$:

$$\begin{array}{rcl} \beta_1 &=& \cos \theta_1, \\ \beta_2 &=& \sin \theta_1 \cos \theta_2, \\ \cdots & \cdots & \cdots \\ \beta_{p-1} &=& \sin \theta_1 \cdots \sin \theta_{p-2} \cos \theta_{p-1}, \\ \beta_p &=& \sin \theta_1 \cdots \sin \theta_{p-2} \sin \theta_{p-1}, \end{array}$$

where $\theta_1, \ldots, \theta_{p-1} \in [-\pi, \pi]$. Denote $\boldsymbol{\theta} = (\theta_1, \ldots, \theta_{p-1})^T$, and let $pl(\boldsymbol{\theta}) = \ell(\boldsymbol{\beta}, \hat{F}_{\boldsymbol{\beta}})$ with $\boldsymbol{\beta}$ being replaced by the polar transformation. We have

$$\hat{\boldsymbol{\theta}} = \arg \max_{\boldsymbol{\theta}} pl(\boldsymbol{\theta}). \tag{S.178}$$

In our implementation, we have used the R function optim() with the default method "Nelder-Mead". We observe that the likelihood function (S.178) is neither concave nor smooth, so the algorithm can be trapped in a local maximum. To tackle this problem, we used multiple initial values in our numerical studies. Specifically, we used 25 randomly generated initial values and chose the one that led to the greatest value of $pl(\theta)$. This strategy worked well. We tested the method using different sets of initial values in a number of simulation examples. In all the examples, the different sets of initial values gave the same $\hat{\beta}$ estimates to the third decimal.

8.2 Some details of the numerical algorithm for $\tilde{\beta}$

Recall that $\widetilde{\boldsymbol{\beta}}$ is the zero-crossing of $\psi_n(\boldsymbol{\beta})$, where

$$\psi_n(\boldsymbol{\beta}) = \frac{1}{n^2} \sum_{i \neq j} (X_i - X_j) \left\{ I(Y_i > Y_j) - \widehat{F}_{\boldsymbol{\beta}} \left((X_i - X_j)^T \boldsymbol{\beta} \right) \right\}.$$

Let $S_n(\theta) = \psi_n(\beta)$ with β being replaced by the polar transformation given in Section 9.1. We define $\tilde{\theta}$ to be the zero-crossing of $S_n(\theta)$. Clearly, once $\tilde{\theta}$ is obtained, $\tilde{\beta}$ can be obtained accordingly based on the polar transformation.

Therefore, we focus on how to calculate $\tilde{\theta}$ numerically. Note that the dimension of $S_n(\theta)$ is

p, while the dimension of $\boldsymbol{\theta}$ is p-1. Therefore, the zero-crossing of $S_n(\boldsymbol{\theta})$ may not exist when n is small, although we have theoretically established its asymptotic existence in Theorem 2. To bypass this difficulty, we define $S_n^{(-k)}(\boldsymbol{\theta})$ to be the vector obtained by removing the kth element of $S_n(\boldsymbol{\theta})$, so that the dimensions of $S_n^{(-k)}(\boldsymbol{\theta})$ and $\boldsymbol{\theta}$ are both p-1. If we can obtain $\widetilde{\boldsymbol{\theta}}^{(-k)}$ the zero-crossing of $S_n^{(-k)}(\boldsymbol{\theta})$, we can then combine $\widetilde{\boldsymbol{\theta}}^{(-1)}, \dots, \widetilde{\boldsymbol{\theta}}^{(-p)}$ to get

$$\widetilde{\boldsymbol{\theta}} = \frac{1}{p} \sum_{k=1}^{p} \widetilde{\boldsymbol{\theta}}^{(-k)}$$

Next, we discuss how to obtain $\tilde{\theta}^{(-k)}$ from $S_n^{(-k)}(\theta)$. When p = 2, the dimension of $S_n^{(-k)}(\theta)$ is 1. We can apply the R function *uniroot.all()* in the rootSolve package to find the zero-crossings of $S_n^{(-k)}(\theta)$; if the solution is not unique, we select the one that leads to the largest value of $pl(\theta)$. When p > 2, the dimension of $S_n^{(-k)}(\theta)$ is greater than 1. The existing R functions for finding the roots of multivariate equations may not lead to stable results, since it is necessary to evaluate the Jacobian, which may not exist for $S_n^{(-k)}(\theta)$. Following a similar strategy used by Balabdaoui et al. (2019), we suggest minimizing $\|S_n^{(-k)}(\theta)\|_2$ to obtain $\tilde{\theta}^{(-k)}$. This can be done by using R function optim() with the default method "Nelder–Mead" and multiple initial values, say 25. If multiple zero-crossings of $S_n^{(-k)}(\theta)$ exist, we select the one that leads to the largest value of $pl(\theta)$.

9 Extension to Data with Ties in Responses

In the main article, we have assumed that the responses Y_i have continuous distributions, and there are no ties. However, our pairwise rank likelihood method can readily be extended to accommodate the case where the observed responses Y_i contain ties. In particular, noting the derivation of (3.1) in the main article, we have

$$P(Y_i \leq Y_j | X_i, X_j) = P(\epsilon_i - \epsilon_j \leq (X_j - X_i)^T \boldsymbol{\beta} | X_i, X_j) = F((X_j - X_i)^T \boldsymbol{\beta}),$$

where we have used only the fact that $F(\cdot)$ is the c.d.f. of $\epsilon_i - \epsilon_j$, and therefore

$$P(Y_i > Y_j | X_i, X_j) = 1 - P(Y_i \le Y_j | X_i, X_j) = 1 - F((X_j - X_i)^T \beta).$$

In the spirit of (3.2) in the main article, the pairwise rank log-likelihood can be defined to be

$$\widetilde{\ell}(\boldsymbol{\beta}, F) = \sum_{i \neq j} \left[I(Y_i > Y_j) \log\{P(Y_i > Y_j | X_i, X_j)\} + I(Y_i \leqslant Y_j) \log\{P(Y_i \leqslant Y_j | X_i, X_j)\} \right]$$
$$= \sum_{i \neq j} \left[I(Y_i > Y_j) \log\{1 - F((X_j - X_i)^T \boldsymbol{\beta})\} + I(Y_i \leqslant Y_j) \log\{F((X_j - X_i)^T \boldsymbol{\beta})\} \right].$$

As a consequence, the algorithm for estimating β and $F(\cdot)$ is based on the procedures given in Section 3.1 of the main article. We observe that $\tilde{\ell}(\beta, F)$ accounts for the potential effects of ties in the responses; the reason is as follows. When there are ties, the effects of the event $Y_i = Y_j$ and its probability $P(Y_i = Y_j | X_i, X_j)$ have been respectively accounted for in $I(Y_i \leq Y_j)$ and the corresponding probability $P(Y_i \leq Y_j | X_i, X_j)$. For a more intuitive view of this, note that for every given $i \neq j$, the only terms in $\tilde{\ell}(\beta, F)$ that contain data from both the *i*th and *j*th subjects are

$$I(Y_i > Y_j) \log\{1 - F((X_j - X_i)^T \boldsymbol{\beta})\} + I(Y_i \leq Y_j) \log\{F((X_j - X_i)^T \boldsymbol{\beta})\}$$

and

$$I(Y_{i} < Y_{j}) \log\{1 - F((X_{i} - X_{j})^{T} \boldsymbol{\beta})\} + I(Y_{i} \ge Y_{j}) \log\{F((X_{i} - X_{j})^{T} \boldsymbol{\beta})\}.$$

As a consequence, from the table below, we observe that for different relationships of Y_i and Y_j (i.e., $Y_i = Y_j$ or $Y_i > Y_j$ or $Y_i < Y_j$), the contribution to $\tilde{\ell}(\boldsymbol{\beta}, F)$ is different.

	Contribution of observations i and j to $\widetilde{\ell}(\boldsymbol{\beta},F)$
$Y_i = Y_j$	
$Y_i > Y_j$	$\log\{1 - F((X_j - X_i)^T \boldsymbol{\beta})\} + \log\{F((X_i - X_j)^T \boldsymbol{\beta})\} \\ \log\{F((X_j - X_i)^T \boldsymbol{\beta})\} + \log\{1 - F((X_i - X_j)^T \boldsymbol{\beta})\}$
$Y_i < Y_j$	$\log\{F((X_j - X_i)^T \boldsymbol{\beta})\} + \log\{1 - F((X_i - X_j)^T \boldsymbol{\beta})\}$

10 Extension to Right-Censored Data

In this section, we extend our methods to accommodate the right-censored data. Let T_i be the survival time of a subject *i*, and let C_i be the corresponding censoring time, i = 1, ..., n. Denote by $\{Y_i, \Delta_i, X_i\}_{i=1}^n$ the observed i.i.d. data, where $Y_i = \min\{T_i, C_i\}$ is the observed survival time of the *i*th subject with censoring and Δ_i is the censoring indicator; namely $\Delta_i = 1$ if $Y_i = T_i$, and $\Delta_i = 0$ otherwise.

10.1 The pairwise rank likelihood for the right-censored data

Assuming that T_i , i = 1, ..., n, are observed, the pairwise rank likelihood, based on (T_i, X_i) , i = 1, ..., n, is given by

$$\ell(\boldsymbol{\beta}, F) = \sum_{i < j} \left[I(T_i > T_j) \log\{F((X_i - X_j)^T \boldsymbol{\beta})\} + I(T_i \leqslant T_j) \log\{1 - F((X_i - X_j)^T \boldsymbol{\beta})\}\right].$$

When the data are subject to censoring, we can incorporate the idea of Cheng et al. (1995), i.e., the inverse probability estimation method, in this objective function. In particular, we replace $I(T_i > T_j)$ by $\Delta_j I(Y_i > Y_j) \{G^{-2}(Y_j)\}$, where $G(y) = P(C_i > y)$ is the unknown survival function for the censoring time. As a consequence, the pairwise rank likelihood based

on $(Y_i, \Delta_i, X_i), i = 1, \ldots, n$, is given by

$$\ell(\boldsymbol{\beta}, F, G) = \sum_{i < j} \left[\frac{\Delta_j I(Y_i > Y_j)}{G^2(Y_j)} \log\{F((X_i - X_j)^T \boldsymbol{\beta})\} + \frac{\Delta_i I(Y_i \le Y_j)}{G^2(Y_i)} \log\{1 - F((X_i - X_j)^T \boldsymbol{\beta})\} \right].$$
 (S.179)

We observe that $\ell(\beta, F, G)$ is an unbiased estimator of $\ell(\beta, F)$ if we assume that T_i and C_i are independent and G(y) is continuous, since

$$E\left\{\frac{\Delta_{j}I(Y_{i} > Y_{j})}{G^{2}(Y_{j})} \middle| X_{i}, X_{j}\right\}$$

$$= E\left\{\frac{I(T_{j} \leq C_{j})I(\min(T_{i}, C_{i}) > \min(T_{j}, C_{j}))}{G^{2}(Y_{j})} \middle| X_{i}, X_{j}\right\}$$

$$= E\left\{\frac{I(T_{j} \leq C_{j})I(\min(T_{i}, C_{i}) > T_{j})}{G^{2}(T_{j})} \middle| X_{i}, X_{j}\right\}$$

$$= E\left[E\left\{\frac{I(T_{j} \leq C_{j})I(T_{i} > T_{j})I(C_{i} > T_{j})}{G^{2}(T_{j})} \middle| T_{j}, X_{i}, X_{j}\right\} \middle| X_{i}, X_{j}\right]$$

$$= E\left\{I(T_{i} > T_{j})|X_{i}, X_{j}\right\}.$$
(S.180)

Note that the survival function $G(\cdot)$ can be estimated by the Kaplan–Meier estimator, denoted by $\hat{G}(\cdot)$. As a consequence, β and F are estimated by

$$(\widehat{\boldsymbol{\beta}}, \widehat{F}) = \operatorname{argmax}_{\boldsymbol{\beta} \in \mathcal{B}, F \in \mathcal{F}} \ell(\boldsymbol{\beta}, F, \widehat{G}),$$

where \mathcal{B} and \mathcal{F} share the same definitions as those without censoring in this article. This optimization problem can be solved using similar techniques to Section 3.1 in the main article and Section 8.1.

10.2 Score-function-based method for right-censored data

Similar in strategy to the developments of Section 3.2 in the main article, we can also derive a score-function-based method to estimate β for right-censored data. The procedure is sketched as follows.

Based on (S.179), we can define

$$\widehat{F}_{\boldsymbol{\beta},G}(\cdot) = \operatorname{argmax}_{F \in \mathcal{F}} \ell(\boldsymbol{\beta}, F, G)$$

for each $G(\cdot)$ and $\beta \in \mathcal{B}$. This leads to the profile likelihood for β : $\ell(\beta, \hat{F}_{\beta}, G)$, whose score function (if it exists) has the form:

$$\overline{\psi}_n(\boldsymbol{\beta}, G) = \sum_{i \neq j} \xi(X_i, X_j; \widehat{F}_{\boldsymbol{\beta}, G}, \boldsymbol{\beta}) \left[\frac{\Delta_j I(Y_i > Y_j)}{G^2(Y_j)} \right]$$

$$-\left\{\frac{\Delta_j I(Y_i > Y_j)}{G^2(Y_j)} + \frac{\Delta_i I(Y_i \leqslant Y_j)}{G^2(Y_i)}\right\} \widehat{F}_{\boldsymbol{\beta},G}\left((X_i - X_j)^T \boldsymbol{\beta}\right) \bigg].$$

By replacing $\xi(X_i, X_j; \hat{F}_{\beta,G}, \beta)$ with $X_i - X_j$, this score function is reduced to

$$\psi_n(\boldsymbol{\beta}, G) = \sum_{i \neq j} (X_i - X_j) \left[\frac{\Delta_j I(Y_i > Y_j)}{G^2(Y_j)} - \left\{ \frac{\Delta_j I(Y_i > Y_j)}{G^2(Y_j)} + \frac{\Delta_i I(Y_i \leqslant Y_j)}{G^2(Y_i)} \right\} \hat{F}_{\boldsymbol{\beta}, G} \left((X_i - X_j)^T \boldsymbol{\beta} \right) \right]$$

Consequently, we can define $\hat{\beta}$, the score-function-based estimate for β , to be the zero crossing of $\psi_n(\beta, \hat{G})$, where similar to Section 10.1, $\hat{G}(\cdot)$ is the Kaplan–Meier estimator for the survival function $G(\cdot)$.

References

- Balabdaoui, F, Groeneboom, P, and Hendrickx, K. (2019) Score estimation in the monotone single-index model. *Scandinavian Journal of Statistics*, 46, 517–544.
- Cheng, S. C., Wei, L. J., and Ying, Z. (1995). Analysis of transformation models with censored data. *Biometrika*, 82, 835-845.
- Groeneboom, P. and Hendrickx, K. (2018). Current status linear regression. *The Annals of Statistics*, 46, 1415-1444.
- Groeneboom, P. and Jongbloed (2014). *Nonparametric Estimation under Shape Constraints*. New York: Cambridge University Press.
- Kosorok, M. R. (2008). Introduction to Empirical Processes and Semiparametric Inference. New York: Springer.
- Serfling, R.J. (1980). Approximation Theorems of Mathematical Statistics. Wiley: New York.
- Sherman, R.P. (1994). Maximal inequalities for degenerate U-processes with applications to optimization estimators. *The Annals of Probability*, 22, 439-459.
- van der Vaart, A. W. (1998). Asymptotic Statistics. New York: Cambridge University Press.
- van der Vaart, A. W. and Wellner, J. A. (1996). *Weak Convergence and Empirical Processes: With Applications to Statistics*. New York: Springer.
- van de Geer, S. A. (2000). *Empirical Processes in M-Estimation*. Cambridge University Press, New York.
- Wald, A. (1949). Note on the consistency of the maximum likelihood estimate. *Annals of Mathematical Statistics*, 20, 595-601.