CHARACTERIZATIONS OF MATRIX VALUED ASYMMETRIC TRUNCATED TOEPLITZ OPERATORS

REWAYAT KHAN, YAGOUB AMEUR, AND JAMROZ KHAN

ABSTRACT. Matrix valued asymmetric truncated Toeplitz operators are compressions of multiplication operators acting between two possibly different model spaces. In this paper, we characterize matrix valued asymmetric truncated Toeplitz operators by using compressed shifts.

1. Introduction

Let H^2 be the classical Hardy space in the unit disk $\mathbb{D} = \{\lambda \in \mathbb{C} : |\lambda| < 1\}$. Truncated Toeplitz operators (TTO's) and asymmetric truncated Toeplitz operators (ATTO's) are compressions of multiplication operator to the backward shift invariant subspaces of H^2 (with two possibly different underlying subspaces in the asymmetric case). Each of these subspaces is of the form $K_{\theta} = (\theta H^2)^{\perp} = H^2 \ominus \theta H^2$, where θ a complex-valued inner function: $\theta \in H^{\infty}$ and $|\theta(z)| = 1$ a.e. on the unit circle $\mathbb{T} = \partial \mathbb{D} = \{z \in \mathbb{C} : |z| = 1\}$. Since D. Sarasons paper [26] TTO's, and later on ATTO's [3,6,7], have been intensly studied (see [1,8,10,12,14,27] and [5,15–18,22,23]).

It is natural to consider TTO's and ATTO's defined on subspaces of vector valued Hardy space $H^2(\mathcal{H})$ with \mathcal{H} a separable finite dimensional complex Hilbert space (see Sections 2 and 3 for detailed definitions). A vector valued model space $K_{\Theta} \subset H^2(\mathcal{H})$ is the orthogonal complement of $\Theta H^2(\mathcal{H})$, that is, $K_{\Theta} = H^2(\mathcal{H}) \ominus \Theta H^2(\mathcal{H})$. Here Θ is an operator valued inner function: a function with values in $\mathcal{L}(\mathcal{H})$ (the algebra of all bounded linear operators on \mathcal{H}), analytic in \mathbb{D} , bounded and such that the boundary values $\Theta(z)$ are unitary operators a.e. on \mathbb{T} . These spaces appear in connection with model theory of Hilbert space contractions (see [24]). Let P_{Θ} be the orthogonal projection from $L^2(\mathcal{H})$ onto K_{Θ} .

²⁰¹⁰ Mathematics Subject Classification. Primary 47B35, Secondary 47B32, 30D20.

Key words and phrases. model space, model for a contraction, truncated Toeplitz operator, asymmetric truncated Toeplitz operator, matrix valued truncated Toeplitz operator, matrix valued asymmetric truncated Toeplitz operator.

For two operator valued inner functions $\Theta_1, \Theta_2 \in H^{\infty}(\mathcal{L}(\mathcal{H}))$ and $\Phi \in L^2(\mathcal{L}(\mathcal{H}))$ (again, see Sections 2 and 3 for definitions) let

$$(1.1) A_{\Phi}^{\Theta_1,\Theta_2} f = P_{\Theta_2}(\Phi f), f \in K_{\Theta_1} \cap H^{\infty}(\mathcal{L}(\mathcal{H})).$$

The operator $A_{\Phi}^{\Theta_1,\Theta_2}$ is called a matrix valued asymmetric truncated Toeplitz operator (MATTO), while $A_{\Phi}^{\Theta_1} = A_{\Phi}^{\Theta_1,\Theta_1}$ is called a matrix valued truncated Toeplitz operator (MTTO, see [21]). Both are densely defined. Let $\mathcal{MT}(\Theta_1, \Theta_2)$ be the set of all MATTO's of the form (1.1) which can be extended boundedly to the whole space K_{Θ_1} and for $\Theta_1 = \Theta_2 = \Theta$ let $\mathcal{MT}(\Theta) = \mathcal{MT}(\Theta, \Theta)$.

Two important examples of operators from $\mathcal{MT}(\Theta)$ are the model operators

(1.2)
$$S_{\Theta} = A_z^{\Theta} = A_{zI_{\mathcal{U}}}^{\Theta} \quad \text{and} \quad S_{\Theta}^* = A_{\bar{z}}^{\Theta} = A_{\bar{z}I_{\mathcal{U}}}^{\Theta}.$$

It is known that each C_0 contraction with finite defect indices is unitarily equivalent to S_{Θ} for some operator valued inner function Θ (see [24, Chapter IV]). On the other hand, operators from $\mathcal{MT}(\Theta_1, \Theta_2)$ with certain bounded analytic symbols appear as the operators intertwining S_{Θ_1} and S_{Θ_2} (see [2, p. 238]).

Some algebraic properties of MTTO's were studied in [21], while the asymmetric case was investigated in [20]. Here we continue the investigation started in [20].

Sections 2 and 3 contain preliminary material on spaces of vector valued functions (Section 2), model spaces and MATTO's (Section 3). In Section 4 we consider some model space operators and their action on $\mathcal{MT}(\Theta_1, \Theta_2)$. Section 5 is devoted to characterizations of MATTO's in terms of S_{Θ_1} , S_{Θ_2} and their adjoints.

2. Spaces of vector valued functions and their operators

Let \mathcal{H} be a complex separable Hilbert space. In what follows $\|\cdot\|_{\mathcal{H}}$ and $\langle\cdot,\cdot\rangle_{\mathcal{H}}$ will denote the norm and the inner product in \mathcal{H} , respectively. Moreover, we will assume that dim $\mathcal{H} < \infty$. The space $L^2(\mathcal{H})$ can be defined as

$$L^2(\mathcal{H}) = \{ f \colon \mathbb{T} \to \mathcal{H} : f \text{ is measurable and } \int_{\mathbb{T}} \|f(z)\|_{\mathcal{H}}^2 \, dm(z) < \infty \}$$

(*m* being the normalized Lebesgue measure on \mathbb{T}). As usual, each $f \in L^2(\mathcal{H})$ is interpreted as a class of functions equal to the representing f a.e. on \mathbb{T} with respect to m. The space $L^2(\mathcal{H})$ is a (separable) Hilbert space with the inner product given by

$$\langle f, g \rangle_{L^2(\mathcal{H})} = \int_{\mathbb{T}} \langle f(z), g(z) \rangle_{\mathcal{H}} dm(z), \quad f, g \in L^2(\mathcal{H}).$$

Equivalently, $L^2(\mathcal{H})$ consists of elements $f: \mathbb{T} \to \mathcal{H}$ of the form

(2.1)
$$f(z) = \sum_{n=-\infty}^{\infty} a_n z^n \quad \text{(a.e. on } \mathbb{T})$$
 with $\{a_n\} \subset \mathcal{H} \text{ such that } \sum_{n=-\infty}^{\infty} \|a_n\|_{\mathcal{H}}^2 < \infty.$

The *n*-th Fourier coefficient a_n of $f \in L^2(\mathcal{H})$ is determined by

(2.2)
$$\langle a_n, x \rangle_{\mathcal{H}} = \int_{\mathbb{T}} \overline{z}^n \langle f(z), x \rangle_{\mathcal{H}} dm(z) \text{ for all } x \in \mathcal{H}.$$

If $f \in L^2(\mathcal{H})$ is given by (2.1), then its Fourier series converges in the $L^2(\mathcal{H})$ norm and

$$||f||_{L^2(\mathcal{H})}^2 = \int_{\mathbb{T}} ||f(z)||_{\mathcal{H}}^2 dm(z) = \sum_{n=-\infty}^{\infty} ||a_n||_{\mathcal{H}}^2.$$

Moreover, for $g(z) = \sum_{n=-\infty}^{\infty} b_n z^n \in L^2(\mathcal{H})$ we have

$$\langle f, g \rangle_{L^2(\mathcal{H})} = \sum_{n=-\infty}^{\infty} \langle a_n, b_n \rangle_{\mathcal{H}}.$$

For $\mathcal{H} = \mathbb{C}$ we denote $L^2 = L^2(\mathbb{C})$.

The vector valued Hardy space $H^2(\mathcal{H})$ is defined as the set of all the elements of $L^2(\mathcal{H})$ whose Fourier coefficients with negative indices vanish, that is,

$$H^2(\mathcal{H}) = \left\{ f \in L^2(\mathcal{H}) : f(z) = \sum_{n=0}^{\infty} a_n z^n \right\}.$$

Each $f \in H^2(\mathcal{H})$, $f(z) = \sum_{n=0}^{\infty} a_n z^n$, can also be identified with a function

$$f(\lambda) = \sum_{n=0}^{\infty} a_n \lambda^n, \quad \lambda \in \mathbb{D},$$

analytic in the unit disk \mathbb{D} (the boundary values f(z) can be obtained from the radial limits, which converge to the boundary function in the $L^2(\mathcal{H})$ norm). Denote by P_+ the orthogonal projection $P_+: L^2(\mathcal{H}) \to H^2(\mathcal{H})$,

$$P_{+}\left(\sum_{n=-\infty}^{\infty}a_{n}z^{n}\right) = \sum_{n=0}^{\infty}a_{n}z^{n},$$

and let $H^2 = H^2(\mathbb{C})$.

We can also consider the spaces

$$L^{\infty}(\mathcal{H}) = \left\{ f \colon \mathbb{T} \to \mathcal{H} : \begin{array}{l} f \text{ is measurable and} \\ \|f\|_{\infty} = \text{ess sup}_{z \in \mathbb{T}} \|f(z)\|_{\mathcal{H}} < \infty \end{array} \right\}$$

(clearly, $L^{\infty}(\mathcal{H}) \subset L^{2}(\mathcal{H})$) and

$$H^{\infty}(\mathcal{H}) = L^{\infty}(\mathcal{H}) \cap H^{2}(\mathcal{H}),$$

the latter seen also as a space of all bounded \mathcal{H} -valued functions which are analytic in \mathbb{D} .

Now let $\mathcal{L}(\mathcal{H})$ be the algebra of all bounded linear operators on \mathcal{H} equipped with the operator norm $\|\cdot\|_{\mathcal{L}(\mathcal{H})}$. In the case dim $\mathcal{H} = d < \infty$ each element of $\mathcal{L}(\mathcal{H})$ can be identified with a $d \times d$ matrix. Denote

$$L^{\infty}(\mathcal{L}(\mathcal{H})) = \left\{ \mathbf{F} \colon \mathbb{T} \to \mathcal{L}(\mathcal{H}) : \begin{array}{l} \mathbf{F} \text{ is measurable and} \\ \|\mathbf{F}\|_{\infty} = \text{ess sup}_{z \in \mathbb{T}} \|\mathbf{F}(z)\|_{\mathcal{L}(\mathcal{H})} < \infty \end{array} \right\}$$

(a function $\mathbf{F} \colon \mathbb{T} \to \mathcal{L}(\mathcal{H})$ is measurable if $\mathbf{F}(\cdot)x \colon \mathbb{T} \to \mathcal{H}$ is measurable for every $x \in \mathcal{H}$). Each $\mathbf{F} \in L^{\infty}(\mathcal{L}(\mathcal{H}))$ admits a formal Fourier expansion (a.e. on \mathbb{T})

(2.3)
$$\mathbf{F}(z) = \sum_{n=-\infty}^{\infty} F_n z^n \quad \text{with } \{F_n\} \subset \mathcal{L}(\mathcal{H})$$

defined by

(2.4)
$$F_n x = \int_{\mathbb{T}} \overline{z}^n \mathbf{F}(z) x \, dm(z) \quad \text{for } x \in \mathcal{H}$$

(integrated in the strong sense). Let

$$H^{\infty}(\mathcal{L}(\mathcal{H})) = \left\{ \mathbf{F} \in L^{2}(\mathcal{L}(\mathcal{H})) : \mathbf{F}(z) = \sum_{n=0}^{\infty} F_{n} z^{n} \right\}.$$

The space $H^{\infty}(\mathcal{L}(\mathcal{H}))$ can equivalently be defined as the space of all analytic functions $\mathbf{F}: \mathbb{D} \to \mathcal{L}(\mathcal{H})$ such that

$$\|\mathbf{F}\|_{\infty} = \sup_{\lambda \in \mathbb{D}} \|\mathbf{F}(\lambda)\|_{\mathcal{L}(\mathcal{H})} < \infty.$$

Each such bounded analytic \mathbf{F} is of the form

(2.5)
$$\mathbf{F}(\lambda) = \sum_{n=0}^{\infty} F_n \lambda^n, \quad z \in \mathbb{D},$$

and can be identified with the boundary function

(2.6)
$$\mathbf{F}(z) = \sum_{n=0}^{\infty} F_n z^n \in L^{\infty}(\mathcal{L}(\mathcal{H})).$$

Conversely, each $\mathbf{F} \in L^{\infty}(\mathcal{L}(\mathcal{H}))$ given by (2.6) can be extended by (2.5) to a function bounded and analytic in \mathbb{D} . In each case the coefficients $\{F_n\}$ can be obtained by (2.4) and the norms $\|\cdot\|_{\infty}$ of the boundary function and its extension coincide (see [2, p. 232]).

Note that for each $\lambda \in \mathbb{D}$ the function $\mathbf{k}_{\lambda}(z) = (1 - \bar{\lambda}z)^{-1}I_{\mathcal{H}}$ belongs to $H^{\infty}(\mathcal{L}(\mathcal{H}))$. Moreover, for every $x \in \mathcal{H}$ the function $\mathbf{k}_{\lambda}x : z \mapsto \mathbf{k}_{\lambda}(z)x$ belongs to $H^{\infty}(\mathcal{H})$ and has the following reproducing property

$$\langle f, \mathbf{k}_{\lambda} x \rangle_{L^2(\mathcal{H})} = \langle f(\lambda), x \rangle_{\mathcal{H}}, \quad f \in H^2(\mathcal{H}).$$

To each $\mathbf{F} \in L^{\infty}(\mathcal{L}(\mathcal{H}))$ there corresponds a multiplication operator $M_{\mathbf{F}}: L^2(\mathcal{H}) \to L^2(\mathcal{H})$: for $f \in L^2(\mathcal{H})$,

$$(M_{\mathbf{F}}f)(z) = \mathbf{F}(z)f(z)$$
 a.e. on \mathbb{T} .

We will write $\mathbf{F}f$ instead of $M_{\mathbf{F}}f$. For a constant \mathbf{F} , that is, $\mathbf{F}(z) = F$ a.e. on \mathbb{T} for some $F \in \mathcal{L}(\mathcal{H})$, we will also write Ff instead of $\mathbf{F}f$. By $T_{\mathbf{F}}$ we will denote the compression of $M_{\mathbf{F}}$ to the Hardy space: $T_{\mathbf{F}}: H^2(\mathcal{H}) \to H^2(\mathcal{H})$,

$$T_{\mathbf{F}}f = P_{+}M_{\mathbf{F}}f$$
 for $f \in H^{2}(\mathcal{H})$.

It is clear that $(M_{\mathbf{F}})^* = M_{\mathbf{F}^*}$ and $(T_{\mathbf{F}})^* = T_{\mathbf{F}^*}$, where $\mathbf{F}^*(z) = \mathbf{F}(z)^*$ a.e. on \mathbb{T} . It is also not difficult to verify that for $\mathbf{F} \in L^{\infty}(\mathcal{L}(\mathcal{H}))$ we have that $\mathbf{F} \in H^{\infty}(\mathcal{L}(\mathcal{H}))$ if and only if $M_{\mathbf{F}}(H^2(\mathcal{H})) \subset H^2(\mathcal{H})$. In particular, for $M_z = M_{zI_{\mathcal{H}}}$ we have $M_z^* = M_{\bar{z}} = M_{\bar{z}I_{\mathcal{H}}}$ and $M_z(H^2(\mathcal{H})) \subset H^2(\mathcal{H})$. The operator $S = T_z = M_{z|H^2(\mathcal{H})}$ is called the (forward) shift operator. Its adjoint, the backward shift operator $S^* = T_{\bar{z}}$, is given by the formula

$$S^*f(z) = \bar{z}\big(f(z) - f(0)\big).$$

Considering $\mathcal{L}(\mathcal{H})$ as a Hilbert space with the Hilbert–Schmidt norm we may also define as above the spaces $L^2(\mathcal{L}(\mathcal{H}))$ and $H^2(\mathcal{L}(\mathcal{H}))$. We then have

$$L^2(\mathcal{L}(\mathcal{H})) = \left[zH^2(\mathcal{L}(\mathcal{H}))\right]^* \oplus H^2(\mathcal{L}(\mathcal{H})).$$

Here we assume that $\dim \mathcal{H} < \infty$. Hence the Hilbert–Schmidt norm and the operator norm are equivalent, and

$$L^{\infty}(\mathcal{L}(\mathcal{H})) \subset L^{2}(\mathcal{L}(\mathcal{H})), \quad H^{\infty}(\mathcal{L}(\mathcal{H})) \subset H^{2}(\mathcal{L}(\mathcal{H})).$$

For $\mathbf{F} \in L^2(\mathcal{L}(\mathcal{H}))$ the operators $M_{\mathbf{F}}$ and $T_{\mathbf{F}}$ can be defined densely, on $L^{\infty}(\mathcal{H})$ and $H^{\infty}(\mathcal{H})$, respectively. For more details on spaces of vector valued functions we refer the reader to [2,24,25].

3. Model spaces and MATTO's

An element $\Theta \in H^{\infty}(\mathcal{L}(\mathcal{H}))$ is called an (operator valued) inner function if its boundary values $\Theta(z)$ are unitary operators a.e. on \mathbb{T} (in general it is assumed that the boundary values are isometries, see e.g. [2, p. 113], but here dim $\mathcal{H} < \infty$). We will consider only pure inner functions, that is Θ such that $\|\Theta(0)\|_{\mathcal{L}(\mathcal{H})} < 1$.

The model space

$$K_{\Theta} = H^2(\mathcal{H}) \ominus \Theta H^2(\mathcal{H})$$

corresponding to an inner function Θ is invariant under the backward shift S^* . Moreover, by the vector valued version of Beurling's invariant subspace theorem, each closed (nontrivial) S^* -invariant subspace of $H^2(\mathcal{H})$ is a model space ([2, Chapter 5, Theorem 1.10]). Let P_{Θ} be the orthogonal projection from $L^2(\mathcal{H})$ onto K_{Θ} . Then

$$P_{\Theta} = P_{+} - M_{\Theta} P_{+} M_{\Theta^*}.$$

For each $\lambda \in \mathbb{D}$ we can consider

$$\mathbf{k}_{\lambda}^{\Theta}(z) = \frac{1}{1-\lambda z} (I_{\mathcal{H}} - \Theta(z)\Theta(\lambda)^*) \in H^{\infty}(\mathcal{L}(\mathcal{H})).$$

For $x \in \mathcal{H}$ we will denote the function $z \mapsto \mathbf{k}_{\lambda}^{\Theta}(z)x$ simply by $\mathbf{k}_{\lambda}^{\Theta}x$. Then, for each $x \in \mathcal{H}$ and $\lambda \in \mathbb{D}$, the function $\mathbf{k}_{\lambda}^{\Theta}x = P_{\Theta}(\mathbf{k}_{\lambda}x)$ belongs to $K_{\Theta}^{\infty} = K_{\Theta} \cap H^{\infty}(\mathcal{L}(\mathcal{H}))$ and has the following reproducing property

$$\langle f, \mathbf{k}_{\lambda}^{\Theta} x \rangle_{L^{2}(\mathcal{H})} = \langle f(\lambda), x \rangle_{\mathcal{H}} \text{ for every } f \in K_{\Theta}.$$

It follows in particular that $K_{\Theta}^{\infty} = K_{\Theta} \cap H^{\infty}(\mathcal{H})$ is a dense subset of K_{Θ} .

Now let $\Theta_1, \Theta_2 \in H^{\infty}(\mathcal{L}(\mathcal{H}))$ be two inner functions. For any $\Phi \in L^2(\mathcal{L}(\mathcal{H}))$ define

$$A_{\Phi}^{\Theta_1,\Theta_2}f = P_{\Theta_2}M_{\Phi}f = P_{\Theta_2}(\Phi f), \quad f \in K_{\Theta_1}^{\infty}.$$

The operator $A_{\Phi}^{\Theta_1,\Theta_2}$ is called a matrix valued asymmetric truncated Toeplitz operator (MATTO) with symbol $\Phi \in L^2(\mathcal{L}(\mathcal{H}))$. It is densely defined and if bounded, it can be extended to a bounded linear operator $A_{\Phi}^{\Theta_1,\Theta_2}: K_{\Theta_1} \to K_{\Theta_2}$ (in which case we simply say that $A_{\Phi}^{\Theta_1,\Theta_2}$ is bounded). Let us denote

$$\mathcal{MT}(\Theta_1, \Theta_2) = \{A_{\Phi}^{\Theta_1, \Theta_2} : \Phi \in L^2(\mathcal{L}(\mathcal{H})) \text{ and } A_{\Phi}^{\Theta_1, \Theta_2} \text{ is bounded}\}.$$

For $\Theta_1 = \Theta_2 = \Theta$ we put $A_{\Phi}^{\Theta} = A_{\Phi}^{\Theta,\Theta}$ (a matrix valued truncated Toeplitz operator, MTTO) and $\mathcal{MT}(\Theta) = \mathcal{MT}(\Theta,\Theta)$. Observe that $(A_{\Phi}^{\Theta_1,\Theta_2})^* = A_{\Phi^*}^{\Theta_2,\Theta_1}$, so $A \in \mathcal{MT}(\Theta_1,\Theta_2)$ if and only if $A^* \in \mathcal{MT}(\Theta_2,\Theta_1)$.

Two important examples of MTTO's are the model operators: the compressed shift S_{Θ} and its adjoint S_{Θ}^* , defined by (1.2). Clearly,

 $S_{\Theta}f = P_{\Theta}Sf = P_{\Theta}(M_zf)$ and since K_{Θ} is S^* -invariant, we have $S_{\Theta}^*f = S^*f = P_+(M_{\bar{z}}f)$.

These operators are models for a class of Hilbert space contractions. For example, each C_0 contraction with finite defect indices is unitarily equivalent to S_{Θ} for some operator valued inner function Θ (see [24, Chapter IV]).

Let

$$\mathcal{D}_{\Theta} = \{ (I_{\mathcal{H}} - \Theta\Theta(0)^*)x : x \in \mathcal{H} \} = \{ \mathbf{k}_0^{\Theta} x : x \in \mathcal{H} \} \subset K_{\Theta}.$$

Then for $f \in K_{\Theta}$ we have $f \perp \mathcal{D}_{\Theta}$ if and only if f(0) = 0. It follows that

$$(S_{\Theta}^* f)(z) = \begin{cases} \bar{z} f(z) & \text{for } f \perp \mathcal{D}_{\Theta}, \\ -\bar{z} (\Theta(z) - \Theta(0)) \Theta(0)^* x & \text{for } f = \mathbf{k}_0^{\Theta} x \in \mathcal{D}_{\Theta}. \end{cases}$$

Now denote (the defect operator)

$$D_{\Theta} = I_{K_{\Theta}} - S_{\Theta} S_{\Theta}^*.$$

Since for each $f \in H^2(\mathcal{H})$ we have $(I_{H^2(\mathcal{H})} - SS^*)f = f(0)$ (a constant function in $H^2(\mathcal{H})$), it follows that for $f \in K_{\Theta}$,

(3.1)
$$D_{\Theta}f = (I_{K_{\Theta}} - S_{\Theta}S_{\Theta}^{*})f = P_{\Theta}(I_{H^{2}(\mathcal{H})} - SS^{*})f$$
$$= (I_{\mathcal{H}} - \Theta\Theta(0)^{*})f(0) = \mathbf{k}_{0}^{\Theta}f(0) \in \mathcal{D}_{\Theta}.$$

More precisely,

$$D_{\Theta}f = \begin{cases} 0 & \text{for } f \perp \mathcal{D}_{\Theta}, \\ \mathbf{k}_{0}^{\Theta}(I_{\mathcal{H}} - \Theta(0)\Theta(0)^{*})x & \text{for } f = \mathbf{k}_{0}^{\Theta}x \in \mathcal{D}_{\Theta}. \end{cases}$$

Since \mathbf{k}_0^{Θ} is invertible in $H^{\infty}(\mathcal{L}(\mathcal{H}))$, the formula

$$\Omega_{\Theta}(\mathbf{k}_0^{\Theta}x) = x, \quad x \in \mathcal{H},$$

gives a well defined operator $\Omega_{\Theta}: \mathcal{D}_{\Theta} \to \mathcal{H}$. Clearly, Ω_{Θ} is bounded (here for example as an operator acting between two finite dimensional Hilbert spaces). Since \mathcal{H} can be identified with a subspace of $H^2(\mathcal{H})$ (the space of all constant \mathcal{H} -valued functions), Ω_{Θ} can be seen as an operator from \mathcal{D}_{Θ} into $H^2(\mathcal{H})$. For each $f \in K_{\Theta}$ we then have

(3.2)
$$\Omega_{\Theta} D_{\Theta} f = \Omega_{\Theta} (\mathbf{k}_{0}^{\Theta} f(0)) = f(0) = (I_{H^{2}(\mathcal{H})} - SS^{*}) f.$$

4. MATTO'S AND SOME MODEL SPACE OPERATORS

In this section we consider the action of some model space operators on $\mathcal{MT}(\Theta_1, \Theta_2)$.

In [19] the author considers the generalized Crofoot transform. For a pure inner function $\Theta \in H^{\infty}(\mathcal{L}(\mathcal{H}))$ and $W \in \mathcal{L}(\mathcal{H})$ such that

 $||W||_{\mathcal{L}(\mathcal{H})} < 1$ define the generalized Crofoot transform $J_W^{\Theta}: L^2(\mathcal{H}) \to L^2(\mathcal{H})$ by

$$J_W^{\Theta} f = D_{W^*} (I_{L^2(\mathcal{H})} - \Theta W^*)^{-1} f, \quad f \in L^2(\mathcal{H}).$$

Then J_W^{Θ} is unitary and maps K_{Θ} onto K_{Θ^W} , where

$$\Theta^W(z) = -W + D_{W^*} (I_{L^2(\mathcal{H})} - \Theta(z)W^*)^{-1} \Theta(z) D_W.$$

The following theorem describes the action of the Crofoot transform on $\mathcal{MT}(\Theta_1, \Theta_2)$ (see [19] for the case $\Theta_1 = \Theta_2$ and [16] for the scalar case):

Theorem 4.1. Let $\Theta_1, \Theta_2 \in H^{\infty}(\mathcal{L}(\mathcal{H}))$ be two pure inner functions and let $W_1, W_2 \in \mathcal{L}(\mathcal{H})$ be such that $\|W_1\|_{\mathcal{L}(\mathcal{H})} < 1$ and $\|W_2\|_{\mathcal{L}(\mathcal{H})} < 1$. A bounded linear operator $A: K_{\Theta_1} \to K_{\Theta_2}$ belongs to $\mathcal{MT}(\Theta_1, \Theta_2)$ if and only if $J_{W_2}^{\Theta_2} A(J_{W_1}^{\Theta_1})^*$ belongs to $\mathcal{MT}(\Theta_1^{W_1}, \Theta_2^{W_2})$. More precisely, $A = A_{\Phi}^{\Theta_1,\Theta_2} \in \mathcal{MT}(\Theta_1, \Theta_2)$ if and only if $J_{W_2}^{\Theta_2} A(J_{W_1}^{\Theta_1})^* = A_{\Psi}^{\Theta_1^{W_1},\Theta_2^{W_2}} \in \mathcal{MT}(\Theta_1^{W_1}, \Theta_2^{W_2})$ with

$$\Psi = D_{W_2^*} (I_{\mathcal{L}(\mathcal{H})} - \Theta_2 W_2^*)^{-1} \Phi D_{W_1^*} (I_{\mathcal{L}(\mathcal{H})} + \Theta_1^{W_1} W_1^*)^{-1}.$$

Recall that if $\Theta \in H^{\infty}(\mathcal{L}(\mathcal{H}))$ is an inner function, then so is

$$\widetilde{\Theta}(z) = \Theta(\bar{z})^*$$
.

Let us now consider the map $\tau_{\Theta}: L^2(\mathcal{H}) \to L^2(\mathcal{H})$ defined for $f \in L^2(\mathcal{H})$ by

(4.1)
$$(\tau_{\Theta} f)(z) = \bar{z} \Theta(\bar{z})^* f(\bar{z}) = \bar{z} \widetilde{\Theta}(z) f(\bar{z}) \quad \text{a.e. on } \mathbb{T}.$$

The map τ_{Θ} is an isometry and its adjoint $\tau_{\Theta}^* = \tau_{\widetilde{\Theta}}$ is also its inverse. Hence τ_{Θ} is unitary. Moreover, it is easy to verify that

$$\tau_{\Theta}(\Theta H^2(\mathcal{H})) \subset H^2(\mathcal{H})^{\perp}$$
 and $\tau_{\Theta}(H^2(\mathcal{H})^{\perp}) \subset \widetilde{\Theta} H^2(\mathcal{H})$,

which implies that

$$\tau_{\Theta}(K_{\Theta}) = K_{\widetilde{\Theta}}.$$

Theorem 4.2. Let $\Theta_1, \Theta_2 \in H^{\infty}(\mathcal{L}(\mathcal{H}))$ be two pure inner functions. A bounded linear operator $A: K_{\Theta_1} \to K_{\Theta_2}$ belongs to $\mathcal{MT}(\Theta_1, \Theta_2)$ if and only if $\tau_{\Theta_2} A \tau_{\Theta_1}^*$ belongs to $\mathcal{MT}(\widetilde{\Theta}_1, \widetilde{\Theta}_2)$. More precisely, $A = A_{\Phi}^{\Theta_1,\Theta_2} \in \mathcal{MT}(\Theta_1, \Theta_2)$ if and only if $\tau_{\Theta_2} A \tau_{\Theta_1}^* = A_{\Psi}^{\widetilde{\Theta}_1,\widetilde{\Theta}_2} \in \mathcal{MT}(\widetilde{\Theta}_1, \widetilde{\Theta}_2)$ with

(4.2)
$$\begin{aligned} \Psi(z) &= \Theta_2(\bar{z})^* \Phi(\bar{z}) \Theta_1(\bar{z}) \\ &= \widetilde{\Theta}_2(z) \Phi(\bar{z}) \widetilde{\Theta}_1(z)^* \quad a.e. \ on \ \mathbb{T}. \end{aligned}$$

Proof. Let $A: K_{\Theta_1} \to K_{\Theta_2}$ be a bounded linear operator. Assume that $A = A_{\Phi}^{\Theta_1,\Theta_2} \in \mathcal{MT}(\Theta_1,\Theta_2)$ with some $\Phi \in L^2(\mathcal{L}(\mathcal{H}))$, and take $f \in K_{\widetilde{\Theta}_1}^{\infty}$ and $g \in K_{\widetilde{\Theta}_2}^{\infty}$. Note that $\tau_{\widetilde{\Theta}_1} f \in K_{\Theta_1}^{\infty}$ and $\tau_{\widetilde{\Theta}_2} g \in K_{\Theta_2}^{\infty}$. Therefore

$$\begin{split} \langle \tau_{\Theta_2} A \, \tau_{\Theta_1}^* f, g \rangle_{L^2(\mathcal{H})} &= \langle A_\Phi^{\Theta_1,\Theta_2} \tau_{\widetilde{\Theta}_1} f, \tau_{\Theta_2}^* g \rangle_{L^2(\mathcal{H})} \\ &= \langle \Phi \, \tau_{\widetilde{\Theta}_1} f, \tau_{\Theta_2}^* g \rangle_{L^2(\mathcal{H})} = \langle \tau_{\Theta_2} (\Phi \, \tau_{\widetilde{\Theta}_1} f), g \rangle_{L^2(\mathcal{H})} \\ &= \int_{\mathbb{T}} \langle \bar{z} \widetilde{\Theta}_2(z) (\Phi \, \tau_{\widetilde{\Theta}_1} f)(\bar{z}), g(z) \rangle_{\mathcal{H}} \, dm(z) \\ &= \int_{\mathbb{T}} \langle \bar{z} \widetilde{\Theta}_2(z) \Phi(\bar{z}) z \Theta_1(\bar{z}) f(z), g(z) \rangle_{\mathcal{H}} \, dm(z) \\ &= \int_{\mathbb{T}} \langle \Psi(z) f(z), g(z) \rangle_{\mathcal{H}} \, dm(z) = \langle A_\Psi^{\widetilde{\Theta}_1, \widetilde{\Theta}_2} f, g \rangle_{L^2(\mathcal{H})} \end{split}$$

with $\Psi \in L^2(\mathcal{L}(\mathcal{H}))$ given by (4.2).

Now, if $\tau_{\Theta_2} A \tau_{\Theta_1}^* = A_{\Psi}^{\widetilde{\Theta}_1, \widetilde{\Theta}_2} \in \mathcal{MT}(\widetilde{\Theta}_1, \widetilde{\Theta}_2)$ for some $\Psi \in L^2(\mathcal{L}(\mathcal{H}))$, then $A = \tau_{\widetilde{\Theta}_2} A_{\Psi}^{\widetilde{\Theta}_1, \widetilde{\Theta}_2} \tau_{\widetilde{\Theta}_1}^*$ and by the first part of the proof $A = A_{\Phi}^{\Theta_1, \Theta_2} \in \mathcal{MT}(\Theta_1, \Theta_2)$ with

(4.3)
$$\Phi(z) = \widetilde{\Theta}_2(\bar{z})^* \Psi(\bar{z}) \widetilde{\Theta}_1(\bar{z}) = \Theta_2(z) \Psi(\bar{z}) \Theta_1(z)^* \quad \text{a.e. on } \mathbb{T}.$$
Hence $\Psi(z) = \Theta_2(\bar{z})^* \Phi(\bar{z}) \Theta_1(\bar{z})$ and (4.2) is satisfied. \square

Denote

$$\widetilde{D}_{\Theta} = I - S_{\Theta}^* S_{\Theta}.$$

Applying Theorem 4.2 to the model operator S_{Θ} ($\Theta_1 = \Theta_2 = \Theta$) we obtain

$$\tau_{\Theta} S_{\Theta} \tau_{\Theta}^* = \tau_{\Theta} S_{\Theta} \tau_{\widetilde{\Theta}} = S_{\widetilde{\Theta}}^*$$

(see [21, p. 1001]). It follows that

$$(4.5) \widetilde{D}_{\Theta} = \tau_{\widetilde{\Theta}} D_{\widetilde{\Theta}} \tau_{\Theta} = \tau_{\widetilde{\Theta}} D_{\widetilde{\Theta}} \tau_{\widetilde{\Theta}}^*$$

and by (3.1),

$$\widetilde{D}_{\Theta}f = \tau_{\widetilde{\Theta}}(\mathbf{k}_0^{\widetilde{\Theta}}(\tau_{\Theta}f)(0)) \text{ for all } f \in K_{\Theta}.$$

For $\lambda \in \mathbb{D}$ let $\widetilde{\mathbf{k}}_{\lambda}^{\Theta} x = \tau_{\widetilde{\Theta}}(\mathbf{k}_{\overline{\lambda}}^{\widetilde{\Theta}} x), x \in \mathcal{H}$. Then (a.e. on \mathbb{T})

$$\begin{split} \widetilde{\mathbf{k}}_{\lambda}^{\Theta}(z)x &= \tau_{\widetilde{\Theta}}(\mathbf{k}_{\widetilde{\lambda}}^{\widetilde{\Theta}}(z)x) = \bar{z}\Theta(z)\mathbf{k}_{\widetilde{\lambda}}^{\widetilde{\Theta}}(\bar{z})x \\ &= \frac{\bar{z}}{1-\lambda\bar{z}}\Theta(z)(I_{\mathcal{H}} - \widetilde{\Theta}(\bar{z})\widetilde{\Theta}(\bar{\lambda})^*)x \\ &= \frac{1}{z-\lambda}\Theta(z)(I_{\mathcal{H}} - \Theta(z)^*\Theta(\lambda))x \\ &= \frac{1}{z-\lambda}(\Theta(z) - \Theta(\lambda))x \in K_{\Theta} \quad \text{for each } x \in \mathcal{H}. \end{split}$$

In paticular,

$$\widetilde{\mathbf{k}}_0^{\Theta}(z)x = \bar{z}(\Theta(z) - \Theta(0))x$$

and

$$\widetilde{D}_{\Theta}f = \widetilde{\mathbf{k}}_{0}^{\Theta}(\tau_{\Theta}f)(0) \in \widetilde{\mathcal{D}}_{\Theta},$$

where

$$\widetilde{\mathcal{D}}_{\Theta} = \tau_{\widetilde{\Theta}} \mathcal{D}_{\widetilde{\Theta}} = \{ \widetilde{\mathbf{k}}_0^{\Theta} x : x \in \mathcal{H} \} = \{ \overline{z} (\Theta(z) - \Theta(0)) x : x \in \mathcal{H} \}.$$

It can be verified that

$$(S_{\Theta}f)(z) = \begin{cases} zf(z) & \text{for } f \perp \widetilde{\mathcal{D}}_{\Theta}, \\ -(I_{\mathcal{H}} - \Theta(z)\Theta(0)^*)\Theta(0)x & \text{for } f = \widetilde{\mathbf{k}}_{0}^{\Theta}x \in \widetilde{\mathcal{D}}_{\Theta}. \end{cases}$$

Hence

$$\widetilde{D}_{\Theta} f = \begin{cases} 0 & \text{for } f \perp \widetilde{\mathcal{D}}_{\Theta}, \\ \widetilde{\mathbf{k}}_{0}^{\Theta} (I_{\mathcal{H}} - \Theta(0)\Theta(0)^{*}) x & \text{for } f = \widetilde{\mathbf{k}}_{0}^{\Theta} x \in \widetilde{\mathcal{D}}_{\Theta}. \end{cases}$$

Finally observe that for $f \in K_{\Theta}$, $x \in \mathcal{H}$,

$$\langle f, \widetilde{\mathbf{k}}_{\lambda}^{\Theta} x \rangle_{L^{2}(\mathcal{H})} = \langle f, \tau_{\widetilde{\Theta}}(\mathbf{k}_{\overline{\lambda}}^{\widetilde{\Theta}} x) \rangle_{L^{2}(\mathcal{H})}$$
$$= \langle \tau_{\Theta} f, \mathbf{k}_{\overline{\lambda}}^{\widetilde{\Theta}} x \rangle_{L^{2}(\mathcal{H})} = \langle (\tau_{\Theta} f)(\overline{\lambda}), x \rangle_{\mathcal{H}}.$$

5. Characterizations with compressed shift operators

In [21](see Theorem 5.2 and Remark 5.4) characterizations of matrix valued truncated Toeplitz operators in $\mathcal{MT}(\Theta)$ were given by using the model operators S_{Θ} , S_{Θ}^* and the defect operators D_{Θ} , \widetilde{D}_{Θ} . These characterizations generalized D. Sarasons results for the scalar case [26]. Here we obtain analogous results for matrix valued asymmetric truncated Toeplitz operators from $\mathcal{MT}(\Theta_1, \Theta_2)$. We use reasoning analogous to that from [21] (see [15] for the scalar case).

Lemma 5.1. If $\Phi \in H^2((\mathcal{L}(E)))$, then

$$A_{\Phi}^{\Theta_1,\Theta_2} - S_{\Theta_2} A_{\Phi}^{\Theta_1,\Theta_2} S_{\Theta_1}^* = P_{\Theta_2} M_{\Phi} (I_{H^2(\mathcal{H})} - SS^*) \quad on \ K_{\Theta_1}^{\infty}.$$

Proof. Recall that $S_{\Theta} = P_{\Theta} M_{z|K_{\Theta}}$ and $S_{\Theta}^* = P_{+} M_{\bar{z}|K_{\Theta}}$. Hence, for $f \in K_{\Theta_1}^{\infty}$,

$$A_{\Phi}^{\Theta_1,\Theta_2}f - S_{\Theta_2}A_{\Phi}^{\Theta_1,\Theta_2}S_{\Theta_1}^*f = P_{\Theta_2}M_{\Phi}f - P_{\Theta_2}M_zP_{\Theta_2}M_{\Phi}P_{\Theta_1}M_{\bar{z}}f$$

(note that $S_{\Theta_1}^* f \in K_{\Theta_1}^{\infty}$). Since $P_{\Theta_2} M_z P_{\Theta_2} = P_{\Theta_2} M_z$ on $H^2(\mathcal{H})$ (as $M_z(\Theta_2 H^2(\mathcal{H})) \subset \Theta_2 H^2(\mathcal{H})$), we have

$$\begin{split} A_{\Phi}^{\Theta_{1},\Theta_{2}}f - S_{\Theta_{2}}A_{\Phi}^{\Theta_{1},\Theta_{2}}S_{\Theta_{1}}^{*}f &= P_{\Theta_{2}}M_{\Phi}f - P_{\Theta_{2}}M_{z}M_{\Phi}P_{+}M_{\bar{z}}f \\ &= P_{\Theta_{2}}(M_{\Phi} - M_{z}M_{\Phi}P_{+}M_{\bar{z}})f \\ &= P_{\Theta_{2}}(M_{\Phi} - M_{\Phi}M_{z}P_{+}M_{\bar{z}})f \\ &= P_{\Theta_{2}}M_{\Phi}(I_{H^{2}(\mathcal{H})} - SS^{*})f. \end{split}$$

Recall that

$$D_{\Theta} = I_{K_{\Theta}} - S_{\Theta} S_{\Theta}^*, \quad \widetilde{D}_{\Theta} = I_{K_{\Theta}} - S_{\Theta}^* S_{\Theta}$$

and

 $\mathcal{D}_{\Theta} = \{ (I_{\mathcal{H}} - \Theta(z)\Theta(0)^*)x : x \in \mathcal{H} \}, \quad \widetilde{\mathcal{D}}_{\Theta} = \{ \overline{z}(\Theta(z) - \Theta(0))x : x \in \mathcal{H} \},$ while the operator $\Omega_{\Theta} : \mathcal{D}_{\Theta} \to \mathcal{H} \subset H^2(\mathcal{H})$ is defined by

$$\Omega_{\Theta}(\mathbf{k}_0^{\Theta}x) = x.$$

Theorem 5.2. Let $\Theta_1, \Theta_2 \in H^{\infty}(\mathcal{L}(\mathcal{H}))$ be two pure inner functions and let $A: K_{\Theta_1} \to K_{\Theta_2}$ be a bounded linear operator. Then A belongs to $\mathcal{MT}(\Theta_1, \Theta_2)$ if and only if there exist bounded linear operators $B_1: \mathcal{D}_{\Theta_1} \to K_{\Theta_2}$ and $B_2: \mathcal{D}_{\Theta_2} \to K_{\Theta_1}$, such that

$$(5.1) A - S_{\Theta_2} A S_{\Theta_1}^* = B_1 D_{\Theta_1} + D_{\Theta_2} B_2^*$$

Proof. Assume first that $A \in \mathcal{MT}(\Theta_1, \Theta_2)$, $A = A_{\Psi + \Xi^*}^{\Theta_1, \Theta_2}$ with $\Psi, \Xi \in H^2(\mathcal{L}(\mathcal{H}))$. Then for each $f \in K_{\Theta_1}^{\infty}$ (note that $S_{\Theta_1}^* f \in K_{\Theta_1}^{\infty}$) we have

$$\begin{split} (A - S_{\Theta_2} A S_{\Theta_1}^*) f &= \\ &= (A_{\Psi}^{\Theta_1, \Theta_2} - S_{\Theta_2} A_{\Psi}^{\Theta_1, \Theta_2} S_{\Theta_1}^*) f + (A_{\Xi^*}^{\Theta_1, \Theta_2} - S_{\Theta_2} A_{\Xi^*}^{\Theta_1, \Theta_2} S_{\Theta_1}^*) f. \end{split}$$

Since $\Psi, \Xi \in H^2(\mathcal{L}(\mathcal{H}))$, it follows from Lemma 5.1 and (3.2) that

$$(A_{\Psi}^{\Theta_{1},\Theta_{2}} - S_{\Theta_{2}}A_{\Psi}^{\Theta_{1},\Theta_{2}}S_{\Theta_{1}}^{*})f = P_{\Theta_{2}}M_{\Psi}(I_{H^{2}(\mathcal{H})} - SS^{*})f$$

$$= P_{\Theta_{2}}M_{\Psi}\Omega_{\Theta_{1}}(I_{K_{\Theta_{1}}} - S_{\Theta_{1}}S_{\Theta_{1}}^{*})f$$

$$= B_{1}D_{\Theta_{1}}f,$$

where $B_1 = P_{\Theta_2} M_{\Psi} \Omega_{\Theta_1} : \mathcal{D}_{\Theta_1} \to K_{\Theta_2}$. Similarly, for each $g \in K_{\Theta_2}^{\infty}$.

$$(A_{\Xi}^{\Theta_{2},\Theta_{1}} - S_{\Theta_{1}} A_{\Xi}^{\Theta_{2},\Theta_{1}} S_{\Theta_{2}}^{*}) g = P_{\Theta_{1}} M_{\Xi} (I_{H^{2}(\mathcal{H})} - SS^{*}) g$$

= $B_{2} D_{\Theta_{2}} g$,

where $B_2 = P_{\Theta_1} M_{\Xi} \Omega_{\Theta_2} : \mathcal{D}_{\Theta_2} \to K_{\Theta_1}$. Note that both B_1 and B_2 are bounded since Ω_{Θ_1} and Ω_{Θ_2} are bounded. Thus

$$(A_{\Xi^*}^{\Theta_1,\Theta_2} - S_{\Theta_2} A_{\Xi^*}^{\Theta_1,\Theta_2} S_{\Theta_1}^*) f = D_{\Theta_2} B_2^* f.$$

It follows that A satisfies (5.1).

Assume now that a bounded linear operator $A: K_{\Theta_1} \to K_{\Theta_2}$ satisfies (5.1). Then

$$S_{\Theta_2}^n A S_{\Theta_1}^{*n} - S_{\Theta_2}^{n+1} A S_{\Theta_1}^{*(n+1)} = S_{\Theta_2}^n B_1 D_{\Theta_1} S_{\Theta_1}^{*n} + S_{\Theta_2}^n D_{\Theta_2} B_2^* S_{\Theta_1}^{*n}$$

for n = 0, 1, ... We thus see that for each integer $N \ge 0$,

$$A = \sum_{n=0}^{N} \left(S_{\Theta_2}^n B_1 D_{\Theta_1} S_{\Theta_1}^{*n} + S_{\Theta_2}^n D_{\Theta_2} B_2^* S_{\Theta_1}^{*n} \right) + S_{\Theta_2}^{N+1} A S_{\Theta_1}^{*(N+1)}.$$

Hence, for for all $f \in K_{\Theta_1}$, and $g \in K_{\Theta_2}$ we have

$$\langle Af, g \rangle_{L^{2}(\mathcal{H})} = \sum_{n=0}^{N} \left(\langle S_{\Theta_{2}}^{n} B_{1} D_{\Theta_{1}} S_{\Theta_{1}}^{*n} f, g \rangle_{L^{2}(\mathcal{H})} + \langle S_{\Theta_{2}}^{n} D_{\Theta_{2}} B_{2}^{*} S_{\Theta_{1}}^{*n} f, g \rangle_{L^{2}(\mathcal{H})} \right) + \langle AS_{\Theta_{1}}^{*(N+1)} f, S_{\Theta_{2}}^{*(N+1)} g \rangle_{L^{2}(\mathcal{H})}.$$

Since $S_{\Theta_1}^{*N}f \to 0$ and $S_{\Theta_2}^{*N}g \to 0$ as $N \to \infty$, we obtain

$$\langle Af, g \rangle_{L^{2}(\mathcal{H})} =$$

$$= \sum_{n=0}^{\infty} \left(\langle S_{\Theta_{2}}^{n} B_{1} D_{\Theta_{1}} S_{\Theta_{1}}^{*n} f, g \rangle_{L^{2}(\mathcal{H})} + \langle f, S_{\Theta_{1}}^{n} B_{2} D_{\Theta_{2}} S_{\Theta_{2}}^{*n} g \rangle_{L^{2}(\mathcal{H})} \right).$$

Let us now define $\Psi, \Xi \in H^2(\mathcal{L}(\mathcal{H}))$ a.e. on \mathbb{T} by

$$\Psi(z)x = (B_1\mathbf{k}_0^{\Theta_1}x)(z)$$
 and $\Xi(z)x = (B_2\mathbf{k}_0^{\Theta_2}x)(z)$, $x \in \mathcal{H}$.

Take
$$f(z) = \sum_{k=0}^{\infty} a_n z^n \in K_{\Theta_1}^{\infty}$$
 and $g(z) = \sum_{k=0}^{\infty} b_n z^n \in K_{\Theta_2}^{\infty}$. We then have

$$D_{\Theta_1} S_{\Theta_1}^{*n} f = \mathbf{k}_0^{\Theta_1} (S_{\Theta_1}^{*n} f)(0) = (I_{\mathcal{H}} - \Theta_1 \Theta_1(0)^*) a_n = \mathbf{k}_0^{\Theta_1} a_n$$

and

$$D_{\Theta_2} S_{\Theta_2}^{*n} g = \mathbf{k}_0^{\Theta_2} (S_{\Theta_2}^{*n} g)(0) = (I_{\mathcal{H}} - \Theta_2 \Theta_2(0)^*) b_n = \mathbf{k}_0^{\Theta_2} b_n.$$

It follows from (5.2) that

$$\langle A_{\Psi+\Xi^*}^{\Theta_1,\Theta_2} f, g \rangle_{L^2(\mathcal{H})} =$$

$$= \langle \Psi f, g \rangle_{L^2(\mathcal{H})} + \langle f, \Xi g \rangle_{L^2(\mathcal{H})}$$

$$= \sum_{n=0}^{\infty} \left(\langle z^n \Psi a_n, g \rangle_{L^2(\mathcal{H})} + \langle f, z^n \Xi b_n \rangle_{L^2(\mathcal{H})} \right)$$

$$= \sum_{n=0}^{\infty} \left(\langle z^n B_1 \mathbf{k}_0^{\Theta_1} a_n, g \rangle_{L^2(\mathcal{H})} + \langle f, z^n B_2 \mathbf{k}_0^{\Theta_2} b_n \rangle_{L^2(\mathcal{H})} \right)$$

$$= \sum_{n=0}^{\infty} \left(\langle S_{\Theta_2}^n B_1 D_{\Theta_1} S_{\Theta_1}^{*n} f, g \rangle_{L^2(\mathcal{H})} + \langle f, S_{\Theta_1}^n B_2 D_{\Theta_2} S_{\Theta_2}^{*n} g \rangle_{L^2(\mathcal{H})} \right)$$

$$= \langle Af, g \rangle_{L^2(\mathcal{H})}$$

and so
$$A = A_{\Psi + \Xi^*}^{\Theta_1, \Theta_2} \in \mathcal{MT}(\Theta_1, \Theta_2)$$
.

Corollary 5.3. Let $\Theta_1, \Theta_2 \in H^{\infty}(\mathcal{L}(\mathcal{H}))$ be two pure inner functions and let $A: K_{\Theta_1} \to K_{\Theta_2}$ be a bounded linear operator.

(a) If A satisfies (5.1), then
$$A = A_{\Psi + \Xi^*}^{\Theta_1, \Theta_2} \in \mathcal{MT}(\Theta_1, \Theta_2)$$
 with

(5.3)
$$\Psi(z)x = (B_1\mathbf{k}_0^{\Theta_1}x)(z)$$
 and $\Xi(z)x = (B_2\mathbf{k}_0^{\Theta_2}x)(z)$, $x \in \mathcal{H}$.

(b) If
$$A = A_{\Psi+\Xi^*}^{\Theta_1,\Theta_2} \in \mathcal{MT}(\Theta_1,\Theta_2)$$
, then A satisfies (5.1) with

(5.4)
$$B_1 = P_{\Theta_2} M_{\Psi} \Omega_{\Theta_1} \quad and \quad B_2 = P_{\Theta_1} M_{\Xi} \Omega_{\Theta_2}.$$

As in [21] we can use the unitary operator τ_{Θ} defined by (4.1) and obtain the following theorem.

Theorem 5.4. Let $\Theta_1, \Theta_2 \in H^{\infty}(\mathcal{L}(\mathcal{H}))$ be two pure inner functions and let $A: K_{\Theta_1} \to K_{\Theta_2}$ be a bounded linear operator. Then A belongs to $\mathcal{MT}(\Theta_1, \Theta_2)$ if and only if there exist bounded linear operators $\widetilde{B}_1: \widetilde{\mathcal{D}}_{\Theta_1} \to K_{\Theta_2}$ and $\widetilde{B}_2: \widetilde{\mathcal{D}}_{\Theta_2} \to K_{\Theta_1}$, such that

$$(5.5) A - S_{\Theta_2}^* A S_{\Theta_1} = \widetilde{B}_1 \widetilde{D}_{\Theta_1} + \widetilde{D}_{\Theta_2} \widetilde{B}_2^*.$$

Proof. Let $A: K_{\Theta_1} \to K_{\Theta_2}$ be a bounded linear operator. By Theorem 4.2, A belongs to $\mathcal{MT}(\Theta_1, \Theta_2)$ if and only if $\widetilde{A} = \tau_{\Theta_2} A \tau_{\Theta_1}^*$ belongs to $\mathcal{MT}(\widetilde{\Theta}_1, \widetilde{\Theta}_2)$. By Theorem 5.2 the latter happens if and only if there exist bounded linear operators $B_1: \mathcal{D}_{\widetilde{\Theta}_1} \to K_{\widetilde{\Theta}_2}$ and $B_2: \mathcal{D}_{\widetilde{\Theta}_2} \to K_{\widetilde{\Theta}_1}$, such that

$$(5.6) \ \widetilde{A} - S_{\widetilde{\Theta}_2} \widetilde{A} S_{\widetilde{\Theta}_1}^* = \tau_{\Theta_2} A \tau_{\Theta_1}^* - S_{\widetilde{\Theta}_2} \tau_{\Theta_2} A \tau_{\Theta_1}^* S_{\widetilde{\Theta}_1}^* = B_1 D_{\widetilde{\Theta}_1} + D_{\widetilde{\Theta}_2} B_2^*.$$
In other words,

$$A - \tau_{\Theta_2}^* S_{\widetilde{\Theta}_2} \tau_{\Theta_2} A \tau_{\Theta_1}^* S_{\widetilde{\Theta}_1}^* \tau_{\Theta_1} = \tau_{\Theta_2}^* B_1 D_{\widetilde{\Theta}_1} \tau_{\Theta_1} + \tau_{\Theta_2}^* D_{\widetilde{\Theta}_2} B_2^* \tau_{\Theta_1}.$$

By (4.4) we have

$$\tau_{\Theta_2}^* S_{\widetilde{\Theta}_2} \tau_{\Theta_2} = \tau_{\widetilde{\Theta}_2} S_{\widetilde{\Theta}_2} \tau_{\widetilde{\Theta}_2}^* = S_{\Theta_2}^* \text{ and } \tau_{\Theta_1}^* S_{\widetilde{\Theta}_1}^* \tau_{\Theta_1} = \tau_{\widetilde{\Theta}_1} S_{\widetilde{\Theta}_1} \tau_{\widetilde{\Theta}_1}^* = S_{\Theta_1},$$

while from (4.5) it follows that

$$D_{\widetilde{\Theta}_1} \tau_{\Theta_1} = \tau_{\Theta_1} \widetilde{D}_{\Theta_1}$$
 and $\tau_{\Theta_2}^* D_{\widetilde{\Theta}_2} = \widetilde{D}_{\Theta_2} \tau_{\Theta_2}^*$.

Thus (5.6) is equivalent to

$$A - S_{\Theta_2}^* A S_{\Theta_1} = \tau_{\Theta_2}^* B_1 \tau_{\Theta_1} \widetilde{D}_{\Theta_1} + \widetilde{D}_{\Theta_2} \tau_{\Theta_2}^* B_2^* \tau_{\Theta_1} = \widetilde{B}_1 \widetilde{D}_{\Theta_1} + \widetilde{D}_{\Theta_2} \widetilde{B}_2^*$$
 with

$$\widetilde{B}_1 = \tau_{\Theta_2}^* B_1 \, \tau_{\Theta_1 | \widetilde{\mathcal{D}}_{\Theta_1}}, \quad \widetilde{B}_1 : \widetilde{\mathcal{D}}_{\Theta_1} \to K_{\Theta_2}$$

and

$$\widetilde{B}_2 = \left(\tau_{\Theta_2}^* B_2^* \tau_{\Theta_1}\right)^* = \tau_{\Theta_1}^* B_2 \tau_{\Theta_2 \mid \widetilde{\mathcal{D}}_{\Theta_2}}, \quad \widetilde{B}_2 : \widetilde{\mathcal{D}}_{\Theta_2} \to K_{\Theta_1}.$$

Note that $\tau_{\Theta_i}^* \mathcal{D}_{\widetilde{\Theta_i}} = \widetilde{\mathcal{D}}_{\Theta_i}$, i = 1, 2. This allows us to treat $\tau_{\Theta_2}^* B_2^* \tau_{\Theta_1}$ as an operator from K_{Θ_1} to $\widetilde{\mathcal{D}}_{\Theta_2}$. Moreover, we have

(5.7)
$$B_1 = \tau_{\Theta_2} \widetilde{B}_1 \tau_{\Theta_1 | \mathcal{D}_{\widetilde{\Theta}_1}}^* \quad \text{and} \quad B_2 = \tau_{\Theta_1} \widetilde{B}_2 \tau_{\Theta_2 | \mathcal{D}_{\widetilde{\Theta}_2}}^*.$$

References

- A. Baranov, I. Chalendar, E. Fricain, J. Mashreghi and A. Timotin, Bounded symbols and reproducing kernel thesis for truncated Toeplitz operators, J. Funct. Anal. 259, 2010, 2673–2701.
- [2] H. Bercovici, Operator theory and aritmetic in H^{∞} , Mathematical Surveys and Monographs No. 26, Amer. Math. Soc., Procidence, Rhode Island 1988.
- [3] C. Câmara, J. Jurasik, K. Kliś-Garlicka and M. Ptak, Characterizations of asymmetric truncated Toeplitz operators, Banach J. Math. Anal. 11 (2017), 899–922.
- [4] M. C. Câmara, K. Kliś-Garlicka, B. Lanucha, M. Ptak, Conjugations in $L^2(\mathcal{H})$, Integr. Equ. Oper. Theory **92** (2020), 48.
- [5] C. Câmara, K. Kliś-Garlicka and M. Ptak, Asymmetric truncated Toeplitz operators and conjugations, Filomat, 33 (2019), 3697–3710.
- [6] M. C. Câmara, J. R. Partington, Spectral properties of truncated Toeplitz operators by equivalence after extension, J. Math. Anal. Appl. 433 (2016), no. 2, 762–784.
- [7] M. C. Câmara, J. R. Partington, Asymmetric truncated Toeplitz operators and Toeplitz operators with matrix symbol, J. Oper. Theory 77 (2017), no. 2, 455– 479.
- [8] I. Chalendar, E. Fricain, D. Timotin, A survey of some recent results on truncated Toeplitz operators, in Recent progress on operator theory and approximation in spaces of analytic functions, 59–77, Contemp. Math., 679, Amer. Math. Soc., Providence, RI, 2016.

- [9] N. Chevrot, E. Fricain, and D. Timotin, *The characteristic function of a complex symmetric contraction*, Proc. Amer. Math. Soc. **135** (2007), 2877–2886.
- [10] J. A. Cima, S. R. Garcia, W. T. Ross, W. R. Wogen, Truncated Toeplitz operators: spatial isomorphism, unitary equivalence, and similarity, Indiana Univ. Math. J. 59 (2010), no. 2, 595–620.
- [11] J. B. Conway, A course in operator theory, Graduate Studies in Mathematics, 21. Amer. Math. Soc., Providence, RI, 2000.
- [12] S. R. Garcia, J. E. Mashreghi, W. Ross, Introduction to Model Spaces and their Operators, Cambridge Studies in Advanced Mathematics, 148, Cambridge University Press, 2016.
- [13] S. R. Garcia and M. Putinar, Complex symmetric operators and applications, Trans. Amer. Math. Soc., 358 (2006), 1285–1315.
- [14] S. R. Garcia, W. T. Ross, Recent progress on truncated Toeplitz operators, in: J. Mashreghi, E. Fricain (Eds.), Blaschke products and their applications, Fields Inst. Commun., 65, Springer, New York, 2013, 275–319.
- [15] C. Gu, B. Lanucha, M. Michalska, Characterizations of asymmetric truncated Toeplitz and Hankel operators, Complex Anal. Oper. Theory, 13, 673–684 (2019).
- [16] J. Jurasik, B. Łanucha, Asymmetric truncated Toeplitz operators equal to the zero operator, Ann. Univ. Mariae Curie-Skłodowska, Sect. A 70 (2016), no. 2, 51–62.
- [17] J. Jurasik, B. Lanucha, Asymmetric truncated Toeplitz operators on finite-dimensional spaces, Operators and Matrices 11 (2017), no. 1, 245–262.
- [18] J. Jurasik, B. Lanucha, Matrix Representations of Asymmetric Truncated Toeplitz Operators, Bull. Malays. Math. Sci. Soc. (2020), https://doi.org/10.1007/s40840-020-01001-x
- [19] R. Khan, *The Generalized Crofoot Transform*, accepted for publication in Oper. Matrices (2021), https://arxiv.org/abs/2002.02337.
- [20] R. Khan, The Generalized Crofoot transform and matrix valued asymmetric truncated Toeplitz operators, preprint, https://arxiv.org/abs/2008.04670.
- [21] R. Khan and D. Timotin, *Matrix valued transated Toeplitz operators: basic properties*, Complex Anal. Oper. Theory **12** (2018), 997–1014.
- [22] B. Lanucha, Asymmetric truncated Toeplitz operators of rank one, Comput. Methods Funct. Theory 18 (2018), no. 2, 259–267.
- [23] B. Lanucha, On rank one asymmetric truncated Toeplitz operators on finite-dimensional spaces, J. Math. Anal. Appl. 454 (2017), no. 2, 961–980.
- [24] B. Sz.-Nagy, C. F. Foias, H. Bercovici, L. Kérchy, *Harmonic Analysis of Operators on a Hilbert Space*, second edition, Springer, London 2010.
- [25] H. Radjavi, P. Rosenthal, *Invariant Subspaces*, Springer, New York 1973.
- [26] D. Sarason Algebraic properties of truncated Toeplitz operators, Oper. Matrices 1 (2007), 491–526.
- [27] N. A. Sedlock, Algebras of truncated Toeplitz operators, Oper. Matrices 5 (2011), 309–326.

R. Khan, Abbottabad University of Science and Technology, Pakistan

 $Email\ address: {\tt rewayat.khan@gmail.com}$

Y. Ameur, Laboratoire de mathematiques pures et appliques Universite de Amar telidji Laghouat Algerie 03000

 $Email\ address: \verb"a.yagoub@lagh-univ.dz"$

J. Khan, Government College of Management Science Hangu, Pakistan

Email address: jamroz.khan73@gmail.com