

Superdiffusion of Aerosols Emitted After Sneezing - Nonequilibrium Statistical Mechanics Approach -

Satoshi SUGIMOTO*

*Department of Applied Mathematics and Physics,
Graduate School of Informatics, Kyoto University*

Ken UMENO†

*Department of Applied Mathematics and Physics,
Graduate School of Informatics, Kyoto University*

(Dated: March 30, 2024)

We study a stochastic behavior of aerosols by the non-equilibrium statistical mechanics approach using the analytical approach of the Langevin equation. We firstly show that superdiffusion can possibly occur right after the emission, which may be attributed to the physical mechanism of the outbreak of the COVID-19 pandemic. We also provide clear evidence of the least required distance to prevent infections occurred by aerosols. In particular, the required distance to prevent aerosol infections is derived to be about 42 m when we assume Cauchy distribution as an initial velocity distribution. This fact implies that due to superdiffusion the aerosol infection can occur even far away from a long distance as compared to the previous considerations.

I. INTRODUCTION

Diffusion is one of the important issues in non-equilibrium statistical mechanics, such as the behavior of a drop of ink spreading into water. This phenomenon is observed not only in physical phenomena but also in biological phenomena. Okubo and Umeno studied the phenomenon of diffusion in dynamical systems [1]. They showed that in systems with a Hamiltonian with time-reversibility, an irreversible behavior occurs in the course of time. Alexander analyzed the invasion of cancer cells in tissues using the diffusion equation with real derivatives, and observed *superdiffusion* [2]. P.Siegle *et.al* numerically studied a minimal non-Markovian model of superdiffusion which originates from long-range velocity correlations within the generalized Langevin equation approach [3].

In this Letter, we investigate the diffusion phenomenon of aerosols. Aerosols are many particles dispersed in gas, discussed in different fields of variant researches, such as environmental problems, air purification, medical applications, and so on [4]. These studies are mostly based on experimental or theoretical approaches and simulation approaches of fluid dynamics. One of the most general issues on aerosol dynamics is droplets generated by sneezes. Silwal *et.al* determine how many aerosols are dispersed by each type of pronunciation [5]. Bourouiba *et.al* photographed the behavior of the aerosols generated by a sneeze [6]. Fluid dynamics simulations are used to investigate the difference in the degree of aerosol diffusion with and without a mask [7]. However, aerosol infection cannot be explained without simulation, which cannot avoid simulation errors due to its complex fluid phenom-

ena such as turbulent behavior. We discuss the behavior of aerosols from the viewpoint of stochastic processes, which is different from the previous studies. Our findings provide an evidence that the degree of diffusion of aerosols well describes the actual phenomena, and show that an approach based on stochastic processes is also effective for analyzing aerosol dynamics. Furthermore, if the initial distribution of velocity is a Cauchy distribution, it is theoretically expected that *superdiffusion* occurs right after a sneeze. Note that *superdiffusion* is not subdiffusion due to turbulent fields of fluid dynamics. In this paper, we suppose that the behavior of an aerosol follows the Langevin equation. We ignore the effect of turbulent field and flows of air. We can obtain the distribution of their positions from the Langevin equation for their velocity [10].

II. CALCULATING THE DISTRIBUTION OF POSITION USING LANGEVIN EQUATION OF VELOCITY

Langevin equation for velocity of a particle of mass m is

$$\dot{u}(t) = -fu + R(t)/m$$

where f is a friction coefficient of a particle and $R(t)$ is a white noise, $\langle R(t_1)R(t_2) \rangle = 2mfkT\delta(t_1 - t_2)$. Here, we consider a white noise $R(t)$ to describe the source of random fluctuations of aerosols. When the initial distribution of a velocity is a normal distribution with its average U and standard deviation $V_0 = \frac{\tan 10^\circ}{1 - \tan 10^\circ}U$ [15], the result is

$$P(x, t) = \mathcal{N}_x \left(U \frac{1 - e^{-ft}}{f}, 2\sigma^2 + V_0^2 \left(\frac{1 - e^{-ft}}{f} \right)^2 \right), \quad (1)$$

* alohngoabnp@gmail.com

† umeno.ken.8z@kyoto-u.ac.jp

where $\mathcal{N}_x(\mu, \sigma^2)$ represents a normal distribution with a mean μ and a variance σ^2 with respect to x , and σ^2 is defined as $\frac{kT}{mf} \left(t - \frac{2}{f}(1 - e^{-ft}) + \frac{1}{2f}(1 - e^{-2ft}) \right)$ [11]. To examine the distribution of positions, we calculate the characteristic functional of $x(t)$

$$\langle e^{i\xi x(t)} \rangle = \left\langle \exp \left(i\xi u_0 \frac{1 - e^{-ft}}{f} \right) \right\rangle \left\langle \exp \left(i\xi \int_0^t d\tau \frac{1 - e^{-f(t-\tau)}}{f} \frac{R(\tau)}{m} \right) \right\rangle,$$

where $\langle A \rangle$ means an expected value of A over all the possible stochastic processes $x(t)$ computed by a stochastic integral. The calculation of the first factor is as follows:

$$\begin{aligned} & \left\langle \exp \left(i\xi u_0 \frac{1 - e^{-ft}}{f} \right) \right\rangle \\ &= \int \exp \left(i\xi u_0 \frac{1 - e^{-ft}}{f} \right) p(u_0) du_0 \\ &= \mathcal{F}[p](\tilde{\xi}) \end{aligned}$$

where $\tilde{\xi} = \frac{1 - e^{-ft}}{f} \xi$ and $\mathcal{F}[p](\xi) = \int_{-\infty}^{\infty} p(x) e^{i\xi x} dx$.

Then $\langle e^{i\xi x(t)} \rangle$ can be calculated as:

$$\langle e^{i\xi x(t)} \rangle = \exp(-\xi^2 \sigma^2) \mathcal{F}[p](\tilde{\xi}),$$

$$\text{where } \sigma^2 = \frac{kT}{mf} \left(t - \frac{2}{f}(1 - e^{-ft}) + \frac{1}{2f}(1 - e^{-2ft}) \right).$$

By an inverse Fourier transform of this, we get $P(x, t)$:

$$P(x, t) = \frac{1}{2\pi} \int_{-\infty}^{\infty} \exp(-\sigma^2 \xi^2 - i\xi x) \mathcal{F}[p](\tilde{\xi}) d\xi.$$

We assume that value of σ^2 is small. We analyze this integral by using the Taylor expansion [12]:

$$\begin{aligned} &= \sum_{n=0}^{\infty} \frac{1}{2\pi} \int_{-\infty}^{\infty} (-1)^n \frac{\sigma^{2n} \xi^{2n}}{n!} \exp(-i\xi x) \mathcal{F}[p](\tilde{\xi}) d\xi \\ &= \sum_{n=0}^{\infty} \frac{1}{2\pi} \frac{\partial^{2n}}{\partial x^{2n}} \int_{-\infty}^{\infty} \frac{\sigma^{2n}}{n!} \exp(-i\xi x) \mathcal{F}[p](\tilde{\xi}) d\xi \\ &= \sum_{n=0}^{\infty} \frac{f}{1 - e^{-ft}} \frac{\sigma^{2n}}{2\pi n!} \frac{\partial^{2n}}{\partial x^{2n}} \int_{-\infty}^{\infty} \exp \left(-i\tilde{\xi} \frac{fx}{1 - e^{-ft}} \right) \mathcal{F}[p](\tilde{\xi}) d\tilde{\xi} \\ &= \frac{f}{1 - e^{-ft}} \sum_{n=0}^{\infty} \frac{\sigma^{2n}}{n!} \frac{\partial^{2n}}{\partial x^{2n}} p \left(\frac{fx}{1 - e^{-ft}} \right) \end{aligned} \quad (2)$$

In particular, when σ^2 is evaluated as zero, $P(x, t)$ is wrtitten as

$$P(x, t) = \frac{f}{1 - e^{-ft}} p \left(\frac{fx}{1 - e^{-ft}} \right).$$

Now we assume that the initial velocity $u_0 = u(0)$ follows Cauchy distribution $\frac{1}{\pi} \frac{\gamma}{(u_0 - U)^2 + \gamma^2}$, where γ is the scaling parameter and satisfies $\gamma = V_0$. The reason why we consider the Cauchy distribution is based on the following fact: when we launch particles uniformly and randomly at the plate with respect to the angle, the number of particles on the plate follows Cauchy distribution. Firstly we consider the distribution of velocity of y-axis v_y . Suppose that aerosols are exhaled for x-axis direction. Then we have

$$p(\alpha \leq \theta \leq \alpha + \delta\alpha) = \frac{\delta\alpha}{\pi},$$

where $\theta = \tan^{-1} \frac{v_y}{v}$, α is a real number which satisfies $|\alpha| < \frac{\pi}{2}$, and $0 < \delta\alpha \ll 1$. Secondly, we rewrite this equation without using θ ,

$$p(v \tan \alpha \leq v_y \leq v \tan(\alpha + \delta\alpha)) = \frac{\delta\alpha}{\pi}.$$

By using the following expansion about $\delta\alpha$, the relation $v \tan(\alpha + \delta\alpha) = v \tan \alpha + \frac{v \delta\alpha}{\cos^2 \alpha} = v_y + \frac{v^2 + v_y^2}{v} \delta\alpha$ holds. Thus we have:

$$p(v_y) \delta v_y = \frac{1}{\pi} \frac{v}{v_y^2 + v^2} \delta v_y.$$

This means that the vertical distribution of aerosol follows the Cauchy distribution. Thus, by symmetry, we can suppose that the aerosols follows the same Cauchy distribution in terms of v_x [8]. This means that the aerosol distribution follows the Cauchy distribution on v_x . As a result, we get the distribution of aerosol $P(x, t)$:

$$P(x, t) \approx \frac{1}{\pi} \frac{\tilde{\gamma}}{\tilde{x}^2 + \tilde{\gamma}^2} \left(1 - 2\sigma^2 \frac{\tilde{\gamma}^2 - 3\tilde{x}^2}{(\tilde{x}^2 + \tilde{\gamma}^2)^2} \right), \quad (3)$$

where $\tilde{\gamma} = \frac{1 - e^{-ft}}{f} \gamma$ and $\tilde{x} = x - \frac{1 - e^{-ft}}{f} U$. Note that $\int_{-\infty}^{\infty} \frac{1}{\pi} \frac{\tilde{\gamma}}{\tilde{x}^2 + \tilde{\gamma}^2} \left(-2\sigma^2 \frac{\tilde{\gamma}^2 - 3\tilde{x}^2}{(\tilde{x}^2 + \tilde{\gamma}^2)^2} \right) = 0$. In particular, when $\sigma^2 \ll 1$ such that we can ignore the second term, we can thus derive the Cauchy distribution. This formula implies the deceleration against the time. The properties of the distribution are also preserved even after this approximation with higher-order expansions as long as $\frac{\sigma^2}{\gamma^2} \ll 1$. In particular, when we assume $t \ll \frac{1}{f}$,

$$P(x, t) \approx \frac{1}{\pi} \frac{\gamma t}{(\gamma t)^2 + (x - Ut)^2}$$

holds, which show that $x(t)$ obeys a Cauchy distribution centered at $x = Ut$ with scale number γt . This result indicates the occurrence of *superdiffusion* and the mean-square displacement (MSD) obeys to the order of $O(t^2)$, where the order of diffusion speed is much faster than the normal diffusion having the speed as $\text{MSD} = O(t)$.

III. NUMERICAL ESTIMATION

The properties in numerical estimation are as follows:

1. The density of the aerosol is assumed to be the same as that of water.
2. Temperature is set to the typical temperature as 291K [14].
3. U is assumed 50 m/s [14].
4. The Friction Coefficient of Aerosols f

The Stokes formula[4] is valid in this case, so the friction coefficient of aerosols f is $\frac{3\pi\mu d_p}{m}$. Note that μ is the viscosity coefficient of gas and d_p is the diameter of the aerosol. m is calculated as $\frac{1000}{6}\pi d_p^3$ kg at this time. We consider aerosol particle size and the corresponding f in the TABLE I.

TABLE I. Aerosol particle size and f

Droplet diameter μm	f s ⁻¹
60	90.0
80	50.6
100	32.4
120	22.5
140	16.5
200	8.1

5. Aerosol Residence Time t

TABLE II shows aerosol residence time at different relative humidity [14].

TABLE II. Aerosol residence time t s at different relative humidity

d_p μm	0%	30%	50%	70%	90%
60	3	4	12	7	35
80	5	7	18	12	13
100	8	12	12	17	9
120	10	8	6	7	6
140	7	6	6	6	6
200	5	5	5	5	5

6. The mean free path of the aerosol l_p

The mean free path corresponds to the expected distance that a particle travels between two collisions [16]. Collisions between aerosol particles are negligible. We assume that the volume of exhaled air is 0.001 m³ [17]. Since the mean free path cannot be uniquely determined for a non-equilibrium system, we calculate the velocity and particle size distribution of aerosols at the moment of sneezing. The result of the calculation is shown in TABLE III.

TABLE III. Aerosol particle diameter d_p μm and the mean free path l_p m due to collisions of air constituent molecules. The notation of the numbers is in significant figures, e.g. 6.3e-3 represents 6.3×10^{-3} .

d_p μm	l_p m
60	2.9×10^{-18}
80	1.6×10^{-18}
100	1.0×10^{-18}
120	7.3×10^{-19}
140	5.4×10^{-19}
200	2.6×10^{-19}

7. When an initial velocity $u_0 = u(0)$ follows Cauchy distribution, we can evaluate the second term of (3) as follows:

$$\begin{aligned}
 \left| 2\sigma^2 \frac{\tilde{\gamma}^2 - 3\tilde{x}^2}{(\tilde{x}^2 + \tilde{\gamma}^2)^2} \right| &\leq 6 \frac{\sigma^2}{\tilde{x}^2 + \tilde{\gamma}^2} \\
 &\leq \frac{6\sigma^2}{\tilde{\gamma}^2} \\
 &= \frac{6fkT}{m\gamma^2} \frac{\int_0^t \left(\frac{1-e^{-f\tau}}{f} \right)^2 d\tau}{\left(\frac{1-e^{-ft}}{f} \right)^2} \\
 &\leq \frac{6fkTt}{m\gamma^2}.
 \end{aligned}$$

When the diameter of an particle d_p is 200 μm , we have $\frac{fkTt}{m\gamma^2} = 6.8 \times 10^{-14}t$. This means that the second term of Eq. (3) is negligible in this numerical estimation.

A. Calculation Results

TABLE IV shows the numerical results in the condition that the distribution for an initial velocity u_0 is the normal distribution with its average U and the standard deviation V_0 . The time t is calculated as the mean and the variance of the aerosols at the time listed in the previous TABLE II.

TABLE IV. The mean and the standard deviation in meters of the distribution of aerosols at each particle size d_p . The effect of relative humidity is negligible.

d_p μm	Mean m	Deviation m
60	0.56	0.119
80	0.99	0.211
100	1.54	0.330
120	2.22	0.476
140	3.03	0.648
200	6.17	1.321

TABLE V shows the numerical results in the condition that the distribution for an initial velocity u_0 is the Cauchy distribution. The time t is calculated as the mean and the variance of the aerosols at the time listed in the previous TABLE II.

TABLE V. $U \frac{1-e^{-ft}}{f}$ and the scale number $\tilde{\gamma}$ at each particle size d_p when the initial velocity u_0 follows the Cauchy distribution. The effect of relative humidity is negligible.

$d_p \text{ } \mu\text{m}$	$U \frac{1-e^{-ft}}{f} \text{ m}$	$\tilde{\gamma} \text{ m}$
60	0.56	0.119
80	0.99	0.211
100	1.54	0.330
120	2.22	0.476
140	3.03	0.648
200	6.17	1.321

FIG.1 shows how droplets diffuse calculated from Eqs. (1) and (3). The result indicates that the maximum point moves forward as the amount of time increases.




FIG. 1. The results of how droplets diffuse in one sneeze ($\gamma = \frac{50 \tan 10^\circ}{1 - \tan 10^\circ}$ m/s). $d_p = 200 \text{ } \mu\text{m}$ is used.

We evaluate a required distance α to prevent aerosol infection. α is defined as the distance for that 1% of aerosols diffuse: $\int_{\alpha}^{\infty} P(x,t) dx = 0.01$. When $P(x,t)$ follows a normal distribution $P(x,t) = \mathcal{N}_x(\mu, \sigma^2)$, we have

$$\begin{aligned} \int_{\alpha}^{\infty} \mathcal{N}_x(\mu, \sigma^2) &= \frac{1}{\sqrt{\pi}} \int_{\frac{\alpha-\mu}{\sqrt{2}\sigma}}^{\infty} e^{-x^2} dx \\ &= \frac{1}{2} \left\{ 1 - \operatorname{erf} \left(\frac{\alpha - \mu}{\sqrt{2}\sigma} \right) \right\}, \end{aligned}$$

where $\operatorname{erf}(x) = \frac{2}{\sqrt{\pi}} \int_0^x e^{-t^2} dt$. Then we can calculate

α by using this formula: $\alpha = \sqrt{2}\sigma \operatorname{erf}^{-1}(0.98) + \mu = \frac{\sqrt{2kT}}{mf} \left(t - \frac{2}{f}(1 - e^{-ft}) + \frac{1}{2f}(1 - e^{-2ft}) \right) \operatorname{erf}^{-1}(0.98) + U \frac{1 - e^{-ft}}{f}$.

When $P(x,t)$ is a Cauchy distribution $P(x,t) = \frac{1}{\pi} \frac{\Gamma}{(x-x_0)^2 + \Gamma^2}$, we have

$$\begin{aligned} \int_{\alpha}^{\infty} \frac{1}{\pi} \frac{\Gamma}{(x-x_0)^2 + \Gamma^2} dx &= \frac{1}{\pi} \int_{\frac{\alpha-x_0}{\Gamma}}^{\infty} \frac{1}{x^2 + 1} dx \\ &= \frac{1}{2} - \frac{1}{\pi} \tan^{-1} \left(\frac{\alpha - x_0}{\Gamma} \right), \end{aligned}$$

where Γ is given by the formula: $\Gamma = \tilde{\gamma} = \frac{1-e^{-ft}}{f} \gamma$. Then we can calculate α by using this formula: $\alpha = \gamma \frac{1 - e^{-ft}}{f} \tan(0.49\pi) + U \frac{1 - e^{-ft}}{f}$, where $\gamma = U \frac{\tan 10^\circ}{1 - \tan 10^\circ}$.

TABLE VI shows the distance α for that 1% of aerosols diffuse: $\int_{\alpha}^{\infty} P(x,t) dx = 0.01$. The result indicates that aerosols can diffuse farther when their initial distribution of velocity is Cauchy.

TABLE VI. The distance α m for that 1% of aerosols diffuse

$t \text{ s}$	(Gaussian) m	(Cauchy) m
0.05	5.22	16.1
0.1	8.71	22.8
0.15	11.0	23.9
0.2	12.6	38.7
0.25	13.6	41.9

IV. CONCLUSION AND FUTURE WORKS

In this paper, we investigate the kinetics of aerosols generated by sneezes from the viewpoint of stochastic processes and then compare with the mean free path. In deriving the probability distribution for position from the Langevin equation for velocity, we show that when the initial distribution is Cauchy, superdiffusion occurs immediately after emission. It means that aerosols can spread very widely in a short time, which is consistent with the fact that COVID-19 pandemic was spread to all over the world in a short period in 2020. By using the results from the calculation using Cauchy distribution as an initial velocity distribution, we find that the required distance to prevent aerosol infection is about 42 m. Alternatively, using Gaussian distribution as an initial velocity distribution, our analytical estimation based on non-equilibrium statistical mechanics shows that the required distance is about 14 m, which is even greater than the several meter-scale (5 m) that has been previously reported in the world[14]. The results also indicate that the required distance is sufficiently larger than that

of the mean free path and that the larger aerosol particles cause the farther dispersion. As a future study, the results obtained by fluid dynamics approach [6] need to be derived even from the approach of this study.

ACKNOWLEDGEMENT

The authors thank Shinji Kakinaka, Assistant Professor Atsushi Iwasaki and Minghui Kao for their comments on the manuscript. The authors acknowledge the project “Study of Wireless Power Transmission Coloring for 5G Electricity” of *MinnaDenryoku, Inc.* for financial support.

-
- [1] Okubo, K. I., & Umeno, K. (2017). Proof of Irreversibility for Systems with Time Reversal Symmetry. arXiv preprint arXiv:1703.10888.
 - [2] Iomin, A. (2005). Superdiffusion of cancer on a comb structure. In *Journal of Physics: conference series* (Vol. 7, No. 1, p. 005). IOP Publishing.
 - [3] Siegle, P., Goychuk, I., & Hänggi, P. (2010). Origin of hyperdiffusion in generalized Brownian motion. *Physical review letters*, 105(10), 100602.
 - [4] Friedlander, S. K. (1977). *Smoke, dust and haze: Fundamentals of aerosol behavior*. New York.
 - [5] Silwal, L., Bhatt, S. P., & Raghav, V. (2021). Experimental characterization of speech aerosol dispersion dynamics. *Scientific reports*, 11(1), 1-12.
 - [6] Bourouiba, L., Dehandschoewercker, E., & Bush, J. W. (2014). Violent expiratory events: on coughing and sneezing. *Journal of Fluid Mechanics*, 745, 537-563.
 - [7] Khosronejad, A., Santoni, C., Flora, K., Zhang, Z., Kang, S., Payabvash, S., & Sotiropoulos, F. (2020). Fluid dynamics simulations show that facial masks can suppress the spread of COVID-19 in indoor environments. *AIP Advances*, 10(12), 125109.
 - [8] Khalid, M., Amin, O., Ahmed, S., Shihada, B., & Alouini, M. S. (2020). Modeling of viral aerosol transmission and detection. *IEEE Transactions on Communications*, 68(8), 4859-4873.
 - [9] Epstein, P. S. (1924). On the resistance experienced by spheres in their motion through gases. *Physical Review*, 23(6), 710.
 - [10] Toda, M., Kubo, R., Saitō, N., & Hashitsume, N. (1991). *Statistical physics II: nonequilibrium statistical mechanics* (Vol. 2). Springer Science & Business Media.
 - [11] Van Kampen, N. G. (1992). *Stochastic processes in physics and chemistry* (Vol. 1). Elsevier.
 - [12] L.D.Landau & E.M.Lifshitz (1980). *Statistical Physics 3rd Edition Part 1*. Pergamon Press.
 - [13] Löhner, R., Antil, H., Idelsohn, S., & Oñate, E. (2020). Detailed simulation of viral propagation in the built environment. *Computational Mechanics*, 66(5), 1093-1107.
 - [14] Xie, X., Li, Y., Chwang, A. T. Y., Ho, P. L., & Seto, W. H. (2007). How far droplets can move in indoor environments-revisiting the Wells evaporation-falling curve. *Indoor air*, 17(3), 211-225.
 - [15] Li, H., Leong, F. Y., Xu, G., Kang, C. W., Lim, K. H., Tan, B. H., & Loo, C. M. (2020). Airborne dispersion of droplets during coughing: a physical model of viral transmission. arXiv preprint arXiv:2008.01912.
 - [16] Atkins, P. W., & De Paula, J. (1998). *Physical chemistry*.
 - [17] Koeppen, B. M., & Stanton, B. A. (2009). *Berne & Levy Physiology, Updated Edition E-Book*. Elsevier Health Sciences.