

Kolmogorov widths of a finite-dimensional balls intersection

A.A. Vasil'eva

1 Introduction

When studying the problem of estimating the widths of Sobolev weight classes with restrictions on derivatives in various L_p -metrics [1], the question arose about the widths of intersections of two finite-dimensional balls. Their order estimates were obtained in [2]. Here we generalize this result to the intersection of an arbitrary family of balls.

Let X be a normed space, $M \subset X$, $n \in \mathbb{Z}_+$. We denote by $\mathcal{L}_n(X)$ the family of subspaces in X of dimension at most n . The Kolmogorov n -width of the set M in the space X is defined as follows:

$$d_n(M, X) = \inf_{L \in \mathcal{L}_n(X)} \sup_{x \in M} \inf_{y \in L} \|x - y\|.$$

Given $N \in \mathbb{N}$, $1 \leq p \leq \infty$, we denote by l_p^N the space \mathbb{R}^N with the norm

$$\|(x_1, \dots, x_N)\|_p = \begin{cases} \left(\sum_{i=1}^N |x_i|^p \right)^{1/p}, & \text{if } 1 \leq p < \infty, \\ \max_{1 \leq i \leq N} |x_i|, & \text{if } p = \infty. \end{cases}$$

By B_p^N we denote the unit ball in the space l_p^N .

Let A be a non-empty set, $1 \leq p_\alpha \leq \infty$, $\nu_\alpha > 0$ for every $\alpha \in A$ and $p_\alpha \neq p_\beta$ for $\alpha \neq \beta$, and let

$$M = \cap_{\alpha \in A} \nu_\alpha B_{p_\alpha}^N. \quad (1)$$

In this paper we obtain order estimates of the widths $d_n(M, l_q^N)$ for $n \leq N/2$.

The explicit values of $d_n(B_p^N, l_q^N)$ were obtained by Pietsch and Stesin [3, 4] for $p \geq q$, and by Kolmogorov, Petrov, Smirnov [5] and Stechkin [6] for $p = 1$, $q = 2$. For $p < q < \infty$, Gluskin [9, 10] obtained order estimates of $d_n(B_p^N, l_q^N)$. The case $q = \infty$ was investigated by Kashin [7, 8], Gluskin and Garnaev [11]; for $p \geq 2$, order estimates were obtained, and for $1 \leq p < 2$, the upper and the lower estimates differ by a logarithmic multiplier. For details, see [12].

Galeev [13] obtained order estimates for $d_n(M, l_q^{2n})$, where M is defined by (1), $A \subset \mathbb{R}$, $\nu_\alpha = (2n)^{-\alpha}$. This result was applied for estimating the widths of intersections of some function classes [13, 14].

Given $\alpha, \beta \in A$, we set

$$\varkappa_{\alpha, \beta} = \begin{cases} \left(\frac{\nu_\beta}{\nu_\alpha} \right)^{\frac{p_\alpha p_\beta}{p_\alpha - p_\beta}}, & \text{if } \alpha \neq \beta, \\ 1, & \text{if } \alpha = \beta. \end{cases} \quad (2)$$

Then

$$\nu_\alpha \varkappa_{\alpha, \beta}^{-1/p_\alpha} = \nu_\beta \varkappa_{\alpha, \beta}^{-1/p_\beta}. \quad (3)$$

Notice that $\varkappa_{\alpha, \beta} = \varkappa_{\beta, \alpha}$.

Suppose that

$$1 \leq \varkappa_{\alpha, \beta} \leq N, \quad \alpha, \beta \in A \quad (4)$$

(in §3 it will be shown how to reduce the general case to this case). Then $\{\nu_\alpha : \alpha \in A\}$ is bounded and separated from zero.

Now we formulate the main result of this paper. First we give notation for order equalities and inequalities.

Let X, Y be sets, and let $f_1, f_2 : X \times Y \rightarrow \mathbb{R}_+$. We write $f_1(x, y) \underset{y}{\lesssim} f_2(x, y)$ (or $f_2(x, y) \underset{y}{\gtrsim} f_1(x, y)$) if for any $y \in Y$ there exists $c(y) > 0$ such that $f_1(x, y) \leq c(y)f_2(x, y)$ for every $x \in X$; $f_1(x, y) \underset{y}{\asymp} f_2(x, y)$ if $f_1(x, y) \underset{y}{\lesssim} f_2(x, y)$ and $f_2(x, y) \underset{y}{\lesssim} f_1(x, y)$.

We denote

$$K = \{(\nu_\alpha, 1/p_\alpha)\}_{\alpha \in A}. \quad (5)$$

Theorem 1. *Let $n, N \in \mathbb{N}$, $n \leq N/2$, and let M be defined by (1). Suppose that (4) holds.*

1. *Let $p_\alpha \geq q$ for every $\alpha \in A$. Then*

$$d_n(M, l_q^N) \asymp \inf_{\alpha \in A} \nu_\alpha N^{1/q-1/p_\alpha}. \quad (6)$$

2. *Let $q \leq 2$, $p_\alpha \leq q$ for any $\alpha \in A$. Then*

$$d_n(M, l_q^N) \asymp \inf_{\alpha \in A} \nu_\alpha. \quad (7)$$

3. *Let $q \leq 2$, $A_1 = \{\alpha \in A : p_\alpha > q\} \neq \emptyset$, $A_2 = \{\alpha \in A : p_\alpha < q\} \neq \emptyset$, $A'_1 = \{\alpha \in A : p_\alpha \geq q\}$, $A'_2 = \{\alpha \in A : p_\alpha \leq q\}$. Then*

$$d_n(M, l_q^N) \asymp \inf\{\nu_\alpha \varkappa_{\alpha, \beta}^{1/q-1/p_\alpha} : \alpha \in A'_1, \beta \in A'_2\}. \quad (8)$$

4. Let $2 < q < \infty$, and let $p_\alpha \leq 2$ for every $\alpha \in A$. Then

$$d_n(M, l_q^N) \asymp_q \inf_{\alpha \in A} \nu_\alpha \min\{1, n^{-1/2} N^{1/q}\}. \quad (9)$$

5. Let $2 < q < \infty$, $A'_1 = \{\alpha \in A : p_\alpha \geq q\}$, $A'_2 = \{\alpha \in A : 2 \leq p_\alpha \leq q\}$, $A'_3 = \{\alpha \in A : p_\alpha \leq 2\}$. Suppose that $A \not\subset A'_1$, $A \not\subset A'_3$. Let K be defined by (5). Then

$$\begin{aligned} d_n(M, l_q^N) &\asymp_q \Phi(n, N, q, K) := \\ &= \min\{\Phi_1(n, N, q, K), \Phi_2(n, N, q, K), \Phi_3(n, N, q, K)\}, \end{aligned} \quad (10)$$

where

$$\Phi_1(n, N, q, K) = \inf\{\nu_\alpha \mathfrak{K}_{\alpha,\beta}^{1/q-1/p_\alpha} : \alpha \in A'_1, \beta \in A'_2 \cup A'_3\}, \quad (11)$$

$$\Phi_2(n, N, q, K) = \inf\left\{\nu_\alpha (\min\{1, n^{-1/2} N^{1/q}\})^{\frac{1/p_\alpha-1/q}{1/2-1/q}}, \alpha \in A'_2\right\}, \quad (12)$$

$$\Phi_3(n, N, q, K) = \min\{\nu_\alpha \mathfrak{K}_{\alpha,\beta}^{1/2-1/p_\alpha} \min\{1, n^{-1/2} N^{1/q}\} : \alpha \in A'_1 \cup A'_2, \beta \in A'_3\} \quad (13)$$

(the infimum of the empty set is $+\infty$).

Remark 1. If $p_\alpha = q$, we have

$$\nu_\alpha \mathfrak{K}_{\alpha,\beta}^{1/q-1/p_\alpha} = \nu_\alpha = \nu_\alpha (\min\{1, n^{-1/2} N^{1/q}\})^{\frac{1/p_\alpha-1/q}{1/2-1/q}};$$

if $p_\beta = q$, then

$$\nu_\alpha \mathfrak{K}_{\alpha,\beta}^{1/q-1/p_\alpha} \stackrel{(3)}{=} \nu_\beta \mathfrak{K}_{\alpha,\beta}^{1/q-1/p_\beta} = \nu_\beta = \nu_\beta (\min\{1, n^{-1/2} N^{1/q}\})^{\frac{1/p_\beta-1/q}{1/2-1/q}}.$$

If $p_\alpha = 2$, we have

$$\begin{aligned} &\nu_\alpha (\min\{1, n^{-1/2} N^{1/q}\})^{\frac{1/p_\alpha-1/q}{1/2-1/q}} = \\ &= \nu_\alpha \min\{1, n^{-1/2} N^{1/q}\} = \nu_\alpha \mathfrak{K}_{\alpha,\beta}^{1/2-1/p_\alpha} \min\{1, n^{-1/2} N^{1/q}\}; \end{aligned}$$

if $p_\beta = 2$, then

$$\begin{aligned} &\nu_\alpha \mathfrak{K}_{\alpha,\beta}^{1/2-1/p_\alpha} \min\{1, n^{-1/2} N^{1/q}\} \stackrel{(3)}{=} \nu_\beta \mathfrak{K}_{\alpha,\beta}^{1/2-1/p_\beta} \min\{1, n^{-1/2} N^{1/q}\} = \\ &= \nu_\beta \min\{1, n^{-1/2} N^{1/q}\} = \nu_\beta (\min\{1, n^{-1/2} N^{1/q}\})^{\frac{1/p_\beta-1/q}{1/2-1/q}}. \end{aligned}$$

Theorem 2. Let $n, N \in \mathbb{N}$, $n \leq N-1$, let M be defined by (1), $p_\alpha \geq 2$ for every $\alpha \in A$. Suppose that (4) holds. Then

$$d_n(M, l_\infty^N) \asymp \inf\left\{\nu_\alpha \min\left(1, n^{-1} \log \frac{2N}{n}\right)^{1/p_\alpha} : \alpha \in A\right\}. \quad (14)$$

2 Proof of the main results

First we formulate some well-known results.

Theorem A. [10] *Let $1 \leq p \leq q < \infty$, $0 \leq n \leq N/2$.*

1. *Let $1 \leq q \leq 2$. Then $d_n(B_p^N, l_q^N) \underset{p,q}{\asymp} 1$.*
2. *Let $2 < q < \infty$, $\lambda_{pq} = \min \left\{ 1, \frac{1/p-1/q}{1/2-1/q} \right\}$. Then*

$$d_n(B_p^N, l_q^N) \underset{p,q}{\asymp} \min\{1, n^{-1/2} N^{1/q}\}^{\lambda_{pq}}.$$

Theorem B. [3,4] *Let $1 \leq q \leq p \leq \infty$, $0 \leq n \leq N$. Then*

$$d_n(B_p^N, l_q^N) = (N - n)^{1/q-1/p}. \quad (15)$$

Theorem C. [11] *Let $p \geq 2$, $n \leq N - 1$. Then*

$$d_n(B_p^N, l_\infty^N) \asymp \min\{1, n^{-1/p} \log^{1/p}(1 + N/n)\}. \quad (16)$$

Given $k \in \{1, \dots, N\}$, we define the set $V_k \subset \mathbb{R}^N$ by

$$V_k = \text{conv} \{(\varepsilon_1 \hat{x}_{\sigma(1)}, \dots, \varepsilon_N \hat{x}_{\sigma(N)}) : \varepsilon_j = \pm 1, 1 \leq j \leq N, \sigma \in S_N\},$$

where $\hat{x}_j = 1$ for $1 \leq j \leq k$, $\hat{x}_j = 0$ for $k+1 \leq j \leq N$, S_N is the group of permutations of N elements. Notice that $V_1 = B_1^N$, $V_N = B_\infty^N$.

For $2 \leq q < \infty$, the lower estimates for $d_n(V_k, l_q^N)$ were obtained by Gluskin [9].

Theorem D. [9] *Let $2 \leq q < \infty$, $1 \leq k \leq N$, $n \leq \min\{N^{\frac{2}{q}} k^{1-\frac{2}{q}}, N/2\}$. Then*

$$d_n(V_k, l_q^N) \underset{q}{\gtrsim} k^{1/q}. \quad (17)$$

Let $2 \leq q < \infty$, $1 \leq k \leq N$, $N^{\frac{2}{q}} k^{1-\frac{2}{q}} \leq n \leq N/2$. Then

$$d_n(V_k, l_q^N) \underset{q}{\gtrsim} k^{1/2} n^{-1/2} N^{1/q}. \quad (18)$$

The following result was obtained by Gluskin [15] (the constant in the order inequality depends on q), and by Malykhin and Rjutin [16] (the constant does not depend on q). In [15, p. 39] it was noticed that Galeev obtained the equality $d_n(V_k, l_1^N) = \min\{k, N - n\}$.

Theorem E. [15, 16]. *Let $1 \leq q \leq 2$, $n \leq N/2$. Then*

$$d_n(V_k, l_q^N) \gtrsim k^{1/q}. \quad (19)$$

The following result was obtained by Konyagin, Malykhin and Rjutin [17].

Theorem F. [17]. *Let $n \leq N - 1$, $1 \leq k \leq N$. Then there is an absolute constant $C \geq 1$ such that for $k \geq Cn \log^{-1} \frac{2N}{n}$*

$$d_n(V_k, l_\infty^N) \geq 1/2.$$

Proof of Theorem 1. Let K be defined by (5). From (4) it follows that K is bounded and $\inf_{\alpha \in A} \nu_\alpha > 0$. We show that it suffices to consider the case of a compact K . Indeed, let $\overline{K} = \{(\nu_\alpha, 1/p_\alpha) : \alpha \in A'\}$ be the closure of K . For $\alpha, \beta \in A'$ we define the numbers $\varkappa_{\alpha,\beta}$ by (2). We prove that

1. $M = \cap_{\alpha \in A'} \nu_\alpha B_{p_\alpha}^N$,
2. on the right-hand sides of (6)–(13) the set A can be replaced by A' , and the minimum is attained.

Let $\nu_{\alpha_k} \xrightarrow{k \rightarrow \infty} \nu_\alpha$, $1/p_{\alpha_k} \xrightarrow{k \rightarrow \infty} 1/p_\alpha$. Then $\cap_{k \in \mathbb{N}} \nu_{\alpha_k} B_{p_{\alpha_k}}^N \subset \nu_\alpha B_{p_\alpha}^N$. This implies the first assertion.

Let $(\nu_{\alpha_k}, 1/p_{\alpha_k}) \xrightarrow{k \rightarrow \infty} (\nu_\alpha, 1/p_\alpha)$, $(\nu_{\beta_k}, 1/p_{\beta_k}) \xrightarrow{k \rightarrow \infty} (\nu_\beta, 1/p_\beta)$, $\alpha \neq \beta$. Then

$$\varkappa_{\alpha_k, \beta_k} \xrightarrow{k \rightarrow \infty} \varkappa_{\alpha, \beta}. \quad (20)$$

In particular, $1 \leq \varkappa_{\alpha, \beta} \leq N$ for all $\alpha, \beta \in A'$.

It is clear that if $(\nu_{\alpha_k}, 1/p_{\alpha_k}) \xrightarrow{k \rightarrow \infty} (\nu_\alpha, 1/p_\alpha)$, then $\nu_{\alpha_k} \xrightarrow{k \rightarrow \infty} \nu_\alpha$, $\nu_{\alpha_k} N^{1/q-1/p_{\alpha_k}} \xrightarrow{k \rightarrow \infty} \nu_\alpha N^{1/q-1/p_\alpha}$; if, in addition, $q > 2$, then

$$\nu_{\alpha_k} (\min\{1, n^{-1/2} N^{1/q}\})^{\frac{1/p_{\alpha_k}-1/q}{1/2-1/q}} \xrightarrow{k \rightarrow \infty} \nu_\alpha (\min\{1, n^{-1/2} N^{1/q}\})^{\frac{1/p_\alpha-1/q}{1/2-1/q}}.$$

Let $(\nu_{\alpha_k}, 1/p_{\alpha_k}) \xrightarrow{k \rightarrow \infty} (\nu_\alpha, 1/p_\alpha)$, $(\nu_{\beta_k}, 1/p_{\beta_k}) \xrightarrow{k \rightarrow \infty} (\nu_\beta, 1/p_\beta)$, $p_{\alpha_k} \geq s$, $p_{\beta_k} \leq s$ for every k . We show that $\nu_{\alpha_k} \varkappa_{\alpha_k, \beta_k}^{1/s-1/p_{\alpha_k}} \xrightarrow{k \rightarrow \infty} \nu_\alpha \varkappa_{\alpha, \beta}^{1/s-1/p_\alpha}$. If $\alpha \neq \beta$, it follows from (20). If $\alpha = \beta$, then $p_\alpha = p_\beta = s$, $\varkappa_{\alpha, \beta} = 1$, $1 \leq \varkappa_{\alpha_k, \beta_k}^{1/s-1/p_{\alpha_k}} \leq N^{1/s-1/p_{\alpha_k}} \xrightarrow{k \rightarrow \infty} 1$.

This completes the proof of the second assertion.

In what follows we suppose that K is compact.

Let $p_\beta \leq s \leq p_\alpha$. Then $\frac{1}{s} = \frac{1-\lambda}{p_\beta} + \frac{\lambda}{p_\alpha}$ for some $\lambda \in [0, 1]$. From Hölder's inequality or from [13, Theorem 2] it follows that

$$B_{p_\alpha}^N \cap \varkappa_{\alpha, \beta}^{1/p_\beta-1/p_\alpha} B_{p_\beta}^N \subset \varkappa_{\alpha, \beta}^{(1-\lambda)(1/p_\beta-1/p_\alpha)} B_s^N = \varkappa_{\alpha, \beta}^{1/s-1/p_\alpha} B_s^N. \quad (21)$$

Now we prove (6)–(10).

Case 1. The upper estimate holds, since for every $\alpha \in A$

$$d_n(M, l_q^N) \leq \nu_\alpha d_n(B_{p_\alpha}^N, l_q^N) \stackrel{(15)}{=} \nu_\alpha (N-n)^{1/q-1/p_\alpha} \leq \nu_\alpha N^{1/q-1/p_\alpha}.$$

Let us prove the lower estimate. Let

$$\nu_{\alpha_*} N^{1/q-1/p_{\alpha_*}} = \min_{\alpha \in A} \nu_{\alpha} N^{1/q-1/p_{\alpha}}. \quad (22)$$

We prove that

$$\nu_{\alpha_*} N^{-1/p_{\alpha_*}} B_{\infty}^N \subset M. \quad (23)$$

It suffices to check that $\nu_{\alpha_*} N^{1/p_{\alpha}-1/p_{\alpha_*}} \leq \nu_{\alpha}$ for all $\alpha \in A$. Indeed, this inequality holds, since $\nu_{\alpha_*} N^{1/q-1/p_{\alpha_*}} \leq \nu_{\alpha} N^{1/q-1/p_{\alpha}}$ by (22).

Hence,

$$d_n(M, l_q^N) \stackrel{(23)}{\geq} \nu_{\alpha_*} N^{-1/p_{\alpha_*}} d_n(B_{\infty}^N, l_q^N) \stackrel{(15)}{=} \nu_{\alpha_*} N^{-1/p_{\alpha_*}} (N-n)^{1/q} \gtrsim \nu_{\alpha_*} N^{1/q-1/p_{\alpha_*}}$$

(recall that $n \leq N/2$).

Cases 2 and 4. Let $\nu_{\alpha_*} = \min_{\alpha \in A} \nu_{\alpha}$. Then $\nu_{\alpha_*} B_1^N \subset M \subset \nu_{\alpha_*} B_{\min\{q, 2\}}^N$. It remains to apply Theorem A.

Case 3. Let us prove the upper estimate. Let $\alpha \in A'_1$, $\beta \in A'_2$. Then

$$\nu_{\alpha} B_{p_{\alpha}}^N \cap \nu_{\beta} B_{p_{\beta}}^N \stackrel{(3)}{=} \nu_{\alpha} (B_{p_{\alpha}}^N \cap \varkappa_{\alpha, \beta}^{1/p_{\beta}-1/p_{\alpha}} B_{p_{\beta}}^N) \stackrel{(21)}{\subset} \nu_{\alpha} \varkappa_{\alpha, \beta}^{1/q-1/p_{\alpha}} B_q^N.$$

Hence,

$$d_n(M, l_q^N) \leq \nu_{\alpha} \varkappa_{\alpha, \beta}^{1/q-1/p_{\alpha}} d_n(B_q^N, l_q^N) = \nu_{\alpha} \varkappa_{\alpha, \beta}^{1/q-1/p_{\alpha}}.$$

It remains to take the minimum over $\alpha \in A'_1$, $\beta \in A'_2$.

Now we obtain the lower estimate. Let

$$\nu_{\alpha_*} \varkappa_{\alpha_*, \beta_*}^{1/q-1/p_{\alpha_*}} = \min_{\alpha \in A'_1, \beta \in A'_2} \nu_{\alpha} \varkappa_{\alpha, \beta}^{1/q-1/p_{\alpha}}. \quad (24)$$

First we consider the case $p_{\alpha_*} > q$, $p_{\beta_*} < q$.

Let $k = [\varkappa_{\alpha_*, \beta_*}]$. By (4), we have $1 \leq k \leq N$. We prove that

$$\nu_{\alpha_*} k^{-1/p_{\alpha_*}} V_k \subset 2M. \quad (25)$$

It suffices to check that

$$\nu_{\alpha_*} \varkappa_{\alpha_*, \beta_*}^{1/p_{\gamma}-1/p_{\alpha_*}} \leq \nu_{\gamma}, \quad \gamma \in A \quad (26)$$

(then $\nu_{\alpha_*} k^{1/p_{\gamma}-1/p_{\alpha_*}} \leq 2\nu_{\gamma}$ for all $\gamma \in A$, and (25) holds).

Let $\gamma \in A'_2$. We have

$$\nu_{\alpha_*} \varkappa_{\alpha_*, \beta_*}^{1/q-1/p_{\alpha_*}} \stackrel{(24)}{\leq} \nu_{\alpha_*} \varkappa_{\alpha_*, \gamma}^{1/q-1/p_{\alpha_*}};$$

since $p_{\alpha_*} > q$, the inequality $\varkappa_{\alpha_*, \beta_*} \leq \varkappa_{\alpha_*, \gamma}$ holds. By the conditions $p_{\alpha_*} > q \geq p_{\gamma}$,

we get $\nu_{\alpha_*} \varkappa_{\alpha_*, \beta_*}^{1/p_{\gamma}-1/p_{\alpha_*}} \leq \nu_{\alpha_*} \varkappa_{\alpha_*, \gamma}^{1/p_{\gamma}-1/p_{\alpha_*}} \stackrel{(3)}{=} \nu_{\gamma}$.

Let $\gamma \in A'_1$. We have

$$\nu_{\alpha_*} \varkappa_{\alpha_*, \beta_*}^{1/q-1/p_{\alpha_*}} \stackrel{(24)}{\leq} \nu_{\gamma} \varkappa_{\gamma, \beta_*}^{1/q-1/p_{\gamma}}.$$

By (3), it is equivalent to

$$\nu_{\beta_*} \varkappa_{\alpha_*, \beta_*}^{1/q-1/p_{\beta_*}} \leq \nu_{\beta_*} \varkappa_{\gamma, \beta_*}^{1/q-1/p_{\beta_*}}.$$

Since $p_{\beta_*} < q$, this implies that $\varkappa_{\alpha_*, \beta_*} \geq \varkappa_{\gamma, \beta_*}$. By the conditions $p_{\beta_*} < q \leq p_{\gamma}$, we get

$$\nu_{\alpha_*} \varkappa_{\alpha_*, \beta_*}^{1/p_{\gamma}-1/p_{\alpha_*}} \stackrel{(3)}{=} \nu_{\beta_*} \varkappa_{\alpha_*, \beta_*}^{1/p_{\gamma}-1/p_{\beta_*}} \leq \nu_{\beta_*} \varkappa_{\gamma, \beta_*}^{1/p_{\gamma}-1/p_{\beta_*}} \stackrel{(3)}{=} \nu_{\gamma}.$$

This concludes the proof of (26). Hence,

$$d_n(M, l_q^N) \stackrel{(25)}{\gtrsim} d_n(\nu_{\alpha_*} k^{-1/p_{\alpha_*}} V_k, l_q^N) \stackrel{(19)}{\gtrsim} \nu_{\alpha_*} k^{1/q-1/p_{\alpha_*}} \asymp \nu_{\alpha_*} \varkappa_{\alpha_*, \beta_*}^{1/q-1/p_{\alpha_*}}.$$

Let now $p_{\alpha_*} = q$. Then $\nu_{\alpha_*} \varkappa_{\alpha_*, \beta_*}^{1/q-1/p_{\alpha_*}} = \nu_{\alpha_*}$.

First we consider the case $\inf\{p_{\alpha} : \alpha \in A_1\} > q$. We take $\alpha_0 \in A_1$ such that

$$\varkappa_{\alpha_0, \alpha_*} = \max_{\gamma \in A_1} \varkappa_{\gamma, \alpha_*} \tag{27}$$

(the maximum is attained, since $\{(\nu_{\gamma}, 1/p_{\gamma}) : \gamma \in A_1\}$ is compact and we can apply (20)).

We have

$$p_{\alpha_0} > q. \tag{28}$$

Let $k = \lfloor \varkappa_{\alpha_0, \alpha_*} \rfloor$. By (4), we have $1 \leq k \leq N$. We check that

$$\nu_{\alpha_0} k^{-1/p_{\alpha_0}} V_k \subset 2M. \tag{29}$$

It suffices to prove that

$$\nu_{\alpha_0} \varkappa_{\alpha_0, \alpha_*}^{1/p_{\gamma}-1/p_{\alpha_0}} \leq \nu_{\gamma}, \quad \gamma \in A. \tag{30}$$

Let $\gamma \in A'_2$. Then

$$\nu_{\alpha_0} \varkappa_{\alpha_0, \alpha_*}^{1/q-1/p_{\alpha_0}} = \nu_{\alpha_0} \varkappa_{\alpha_0, \alpha_*}^{1/p_{\alpha_*}-1/p_{\alpha_0}} \stackrel{(3)}{=} \nu_{\alpha_*} \stackrel{(24)}{\leq} \nu_{\alpha_0} \varkappa_{\alpha_0, \gamma}^{1/q-1/p_{\alpha_0}},$$

this together with (28) yields that $\varkappa_{\alpha_0, \alpha_*} \leq \varkappa_{\alpha_0, \gamma}$. Since $p_{\gamma} \leq q < p_{\alpha_0}$, we have

$$\nu_{\alpha_0} \varkappa_{\alpha_0, \alpha_*}^{1/p_{\gamma}-1/p_{\alpha_0}} \leq \nu_{\alpha_0} \varkappa_{\alpha_0, \gamma}^{1/p_{\gamma}-1/p_{\alpha_0}} \stackrel{(3)}{=} \nu_{\gamma}.$$

Let $\gamma \in A_1$. By (3), the inequality (30) is equivalent to

$$\nu_{\alpha_*} \varkappa_{\alpha_0, \alpha_*}^{1/p_{\gamma}-1/p_{\alpha_*}} \leq \nu_{\alpha_*} \varkappa_{\gamma, \alpha_*}^{1/p_{\gamma}-1/p_{\alpha_*}}.$$

Since $p_\gamma > q = p_{\alpha_*}$, it is equivalent to $\varkappa_{\alpha_0, \alpha_*} \geq \varkappa_{\gamma, \alpha_*}$. The last inequality follows from (27).

This completes the proof of (30). Hence,

$$d_n(M, l_q^N) \stackrel{(29)}{\gtrsim} \nu_{\alpha_0} k^{-1/p_{\alpha_0}} d_n(V_k, l_q^N) \stackrel{(19)}{\gtrsim} \nu_{\alpha_0} \varkappa_{\alpha_0, \alpha_*}^{1/q-1/p_{\alpha_0}} \stackrel{(3)}{=} \nu_{\alpha_*} \varkappa_{\alpha_0, \alpha_*}^{1/q-1/p_{\alpha_*}} = \nu_{\alpha_*}. \quad (31)$$

Now let $\inf\{p_\alpha : \alpha \in A_1\} = q$. For $\delta > 0$ we set

$$A^\delta = A'_2 \cup \{\alpha \in A_1 : 1/q - 1/p_\alpha \geq \delta\}, \quad (32)$$

$M_\delta = \cap_{\alpha \in A^\delta} \nu_\alpha B_{p_\alpha}^N$. By (31), we have

$$d_n(M_\delta, l_q^N) \gtrsim \nu_{\alpha_*}.$$

We prove that for small $\delta > 0$ the inclusion $M_\delta \subset 2M$ holds. It suffices to take

$$0 < \delta \leq \frac{\ln 2}{\ln N} \quad (33)$$

and to check that

$$2 \cap_{\alpha \in A \setminus A^\delta} \nu_\alpha B_{p_\alpha}^N \supset \nu_{\alpha_*} B_q^N. \quad (34)$$

Notice that $p_\alpha > q$ for $\alpha \in A \setminus A^\delta$. Hence (34) holds if $2\nu_\alpha \geq \nu_{\alpha_*}$ for all $\alpha \in A \setminus A^\delta$. We have

$$\nu_{\alpha_*}/\nu_\alpha \stackrel{(3)}{=} \varkappa_{\alpha, \alpha_*}^{1/p_{\alpha_*}-1/p_\alpha} \stackrel{(4)}{\leq} N^{1/q-1/p_\alpha} \stackrel{(32)}{\leq} N^\delta \stackrel{(33)}{\leq} 2;$$

this implies (34) and completes the proof of the lower estimate of $d_n(M, l_q^N)$ for $p_{\alpha_*} = q$.

Let now $p_{\beta_*} = q$. Then $\nu_{\alpha_*} \varkappa_{\alpha_*, \beta_*}^{1/q-1/p_{\alpha_*}} \stackrel{(3)}{=} \nu_{\beta_*} \varkappa_{\alpha_*, \beta_*}^{1/q-1/p_{\beta_*}} = \nu_{\beta_*}$. Hence, $d_n(M, l_q^N) \gtrsim \nu_{\beta_*}$.

Case 5. We prove the upper estimate. Let $\alpha \in A'_2$. By Theorem A,

$$d_n(M, l_q^N) \leq d_n(\nu_\alpha B_{p_\alpha}^N, l_q^N) \lesssim_q \nu_\alpha (\min\{1, n^{-1/2} N^{1/q}\})^{\frac{1/p_\alpha-1/q}{1/2-1/q}}.$$

Let $\alpha \in A'_1$, $\beta \in A'_2 \cup A'_3$. Then

$$M \subset \nu_\alpha B_{p_\alpha}^N \cap \nu_\beta B_{p_\beta}^N \stackrel{(3),(21)}{\subset} \nu_\alpha \varkappa_{\alpha, \beta}^{1/q-1/p_\alpha} B_q^N;$$

hence,

$$d_n(M, l_q^N) \leq \nu_\alpha \varkappa_{\alpha, \beta}^{1/q-1/p_\alpha} d_n(B_q^N, l_q^N) = \nu_\alpha \varkappa_{\alpha, \beta}^{1/q-1/p_\alpha}.$$

Let $\alpha \in A'_1 \cup A'_2$, $\beta \in A'_3$. Then

$$M \subset \nu_\alpha B_{p_\alpha}^N \cap \nu_\beta B_{p_\beta}^N \stackrel{(3),(21)}{\subset} \nu_\alpha \varkappa_{\alpha, \beta}^{1/2-1/p_\alpha} B_2^N;$$

by Theorem A,

$$d_n(M, l_q^N) \leq d_n(\nu_\alpha \varkappa_{\alpha,\beta}^{1/2-1/p_\alpha} B_2^N, l_q^N) \lesssim_q \nu_\alpha \varkappa_{\alpha,\beta}^{1/2-1/p_\alpha} \min\{1, n^{-1/2} N^{1/q}\}.$$

Now we prove the lower estimate. We denote $A_1 = \{\alpha \in A : p_\alpha > q\}$, $A_2 = \{\alpha \in A : 2 < p_\alpha < q\}$, $A_3 = \{\alpha \in A : p_\alpha < 2\}$.

1. Let

$$\Phi(n, N, q, K) = \nu_{\alpha_*}(\min\{1, n^{-1/2} N^{1/q}\})^{\frac{1/p_{\alpha_*}-1/q}{1/2-1/q}}, \quad \alpha_* \in A'_2; \quad (35)$$

in addition, we suppose that $p_{\alpha_*} \neq q$ if $A_1 \neq \emptyset$, and $p_{\alpha_*} \neq 2$ if $A_3 \neq \emptyset$.

We set

$$l = (\max\{1, n^{1/2} N^{-1/q}\})^{\frac{1}{1/2-1/q}}, \quad k = \lceil l \rceil. \quad (36)$$

Notice that $1 \leq l \leq N$ and $1 \leq k \leq N$. We prove that

$$\nu_{\alpha_*} k^{-1/p_{\alpha_*}} V_k \subset 2M. \quad (37)$$

It suffices to check that

$$\nu_{\alpha_*} l^{1/p_\gamma-1/p_{\alpha_*}} \leq \nu_\gamma, \quad \gamma \in A.$$

If $\gamma \in A'_2$, it follows from the inequality

$$\nu_{\alpha_*}(\min\{1, n^{-1/2} N^{1/q}\})^{\frac{1/p_{\alpha_*}-1/q}{1/2-1/q}} \stackrel{(10),(12),(35)}{\leq} \nu_\gamma(\min\{1, n^{-1/2} N^{1/q}\})^{\frac{1/p_\gamma-1/q}{1/2-1/q}}.$$

Let $\gamma \in A_1$ (hence, $A_1 \neq \emptyset$; by our assumption, we have $p_{\alpha_*} < q$). Then

$$\nu_{\alpha_*}(\min\{1, n^{-1/2} N^{1/q}\})^{\frac{1/p_{\alpha_*}-1/q}{1/2-1/q}} \stackrel{(10),(11),(35)}{\leq} \nu_\gamma \varkappa_{\gamma,\alpha_*}^{1/q-1/p_\gamma} \stackrel{(3)}{=} \nu_{\alpha_*} \varkappa_{\gamma,\alpha_*}^{1/q-1/p_{\alpha_*}}.$$

Hence, $l^{1/q-1/p_{\alpha_*}} \leq \varkappa_{\gamma,\alpha_*}^{1/q-1/p_{\alpha_*}}$, and $l \geq \varkappa_{\gamma,\alpha_*}$. Since $p_\gamma > q > p_{\alpha_*}$, we have

$$\nu_{\alpha_*} l^{1/p_\gamma-1/p_{\alpha_*}} \leq \nu_{\alpha_*} \varkappa_{\gamma,\alpha_*}^{1/p_\gamma-1/p_{\alpha_*}} \stackrel{(3)}{=} \nu_\gamma.$$

Let $\gamma \in A_3$ (hence, $A_3 \neq \emptyset$; by our assumption, we have $p_{\alpha_*} > 2$). Then

$$\nu_{\alpha_*}(\min\{1, n^{-1/2} N^{1/q}\})^{\frac{1/p_{\alpha_*}-1/q}{1/2-1/q}} \stackrel{(10),(13),(35)}{\leq} \nu_{\alpha_*} \varkappa_{\alpha_*,\gamma}^{1/2-1/p_{\alpha_*}} \min\{1, n^{-1/2} N^{1/q}\}.$$

Therefore, $l^{1/2-1/p_{\alpha_*}} \leq \varkappa_{\alpha_*,\gamma}^{1/2-1/p_{\alpha_*}}$. Hence, $l \leq \varkappa_{\alpha_*,\gamma}$; since $p_\gamma < 2 < p_{\alpha_*}$, we have

$$\nu_{\alpha_*} l^{1/p_\gamma-1/p_{\alpha_*}} \leq \nu_{\alpha_*} \varkappa_{\alpha_*,\gamma}^{1/p_\gamma-1/p_{\alpha_*}} \stackrel{(3)}{=} \nu_\gamma.$$

This concludes the proof of (37). Notice that $n \stackrel{(36)}{\leq} N^{2/q} k^{1-2/q}$. Hence,

$$\begin{aligned} d_n(M, l_q^N) &\stackrel{(37)}{\gtrsim} d_n(\nu_{\alpha_*} k^{-1/p_{\alpha_*}} V_k, l_q^N) \stackrel{(17)}{\gtrsim}_q \nu_{\alpha_*} k^{1/q-1/p_{\alpha_*}} \stackrel{(36)}{\gtrsim} \\ &\gtrsim \nu_{\alpha_*}(\min\{1, n^{-1/2} N^{1/q}\})^{\frac{1/p_{\alpha_*}-1/q}{1/2-1/q}}. \end{aligned}$$

2. Let

$$\Phi(n, N, q, K) = \nu_{\alpha_*}, \quad p_{\alpha_*} = q, \quad A_1 \neq \emptyset. \quad (38)$$

First we suppose that

$$\sup\{p_\alpha : \alpha \in A_2 \cup A'_3\} < q. \quad (39)$$

We take $\alpha_0 \in A_2 \cup A'_3$ such that

$$\varkappa_{\alpha_*, \alpha_0} = \min_{\gamma \in A_2 \cup A'_3} \varkappa_{\alpha_*, \gamma} \quad (40)$$

(by (39), the minimum is attained and $p_{\alpha_0} < q$). Let $k = \lceil \varkappa_{\alpha_*, \alpha_0} \rceil$. From (4) it follows that $1 \leq k \leq N$. We prove that

$$\nu_{\alpha_0} k^{-1/p_{\alpha_0}} V_k \subset 2M. \quad (41)$$

It suffices to check the inequality

$$\nu_{\alpha_0} \varkappa_{\alpha_*, \alpha_0}^{1/p_\gamma - 1/p_{\alpha_0}} \leq \nu_\gamma, \quad \gamma \in A. \quad (42)$$

Let $\gamma \in A_2 \cup A'_3$. From (40) it follows that $\varkappa_{\alpha_*, \alpha_0} \leq \varkappa_{\alpha_*, \gamma}$. Since $p_{\alpha_*} = q > p_\gamma$, we get $\nu_{\alpha_*} \varkappa_{\alpha_*, \alpha_0}^{1/p_\gamma - 1/p_{\alpha_*}} \leq \nu_{\alpha_*} \varkappa_{\alpha_*, \gamma}^{1/p_\gamma - 1/p_{\alpha_*}}$. By (3), this inequality is equivalent to (42).

Let $\gamma \in A'_1$. By (3), the inequality (42) is equivalent to $\nu_{\alpha_0} \varkappa_{\alpha_*, \alpha_0}^{1/p_\gamma - 1/p_{\alpha_0}} \leq \nu_{\alpha_0} \varkappa_{\gamma, \alpha_0}^{1/p_\gamma - 1/p_{\alpha_0}}$. Since $p_{\alpha_0} < p_\gamma$, it is equivalent to

$$\varkappa_{\alpha_*, \alpha_0} \geq \varkappa_{\gamma, \alpha_0}. \quad (43)$$

Let us prove this inequality. If $p_\gamma = q$, then $\gamma = \alpha_*$. Let $p_\gamma > q$. We have

$$\nu_{\alpha_*} \stackrel{(10),(11),(38)}{\leq} \nu_\gamma \varkappa_{\gamma, \alpha_0}^{1/q - 1/p_\gamma};$$

by (3), it is equivalent to $\nu_{\alpha_0} \varkappa_{\alpha_*, \alpha_0}^{1/p_{\alpha_*} - 1/p_{\alpha_0}} \leq \nu_{\alpha_0} \varkappa_{\gamma, \alpha_0}^{1/q - 1/p_{\alpha_0}} = \nu_{\alpha_0} \varkappa_{\gamma, \alpha_0}^{1/p_{\alpha_*} - 1/p_{\alpha_0}}$. Since $p_{\alpha_0} < q = p_{\alpha_*}$, we get (43).

This completes the proof of (41).

Now we check that

$$n \leq N^{2/q} \varkappa_{\alpha_*, \alpha_0}^{1-2/q}. \quad (44)$$

It suffices to consider $n \geq N^{2/q}$.

If $\alpha_0 \in A_2$, (44) follows from

$$\nu_{\alpha_*} \stackrel{(10),(12),(38)}{\leq} \nu_{\alpha_0} (n^{-1/2} N^{1/q})^{\frac{1/p_{\alpha_0} - 1/q}{1/2 - 1/q}} \stackrel{(3)}{=} \nu_{\alpha_*} \varkappa_{\alpha_*, \alpha_0}^{1/p_{\alpha_0} - 1/p_{\alpha_*}} (n^{-1/2} N^{1/q})^{\frac{1/p_{\alpha_0} - 1/q}{1/2 - 1/q}}.$$

If $\alpha_0 \in A'_3$, (44) follows from

$$\nu_{\alpha_*} \stackrel{(10),(13),(38)}{\leq} \nu_{\alpha_*} \mathfrak{N}_{\alpha_*, \alpha_0}^{1/2-1/p_{\alpha_*}} n^{-1/2} N^{1/q}.$$

Hence,

$$d_n(M, l_q^N) \stackrel{(17),(41),(44)}{\underset{q}{\gtrsim}} \nu_{\alpha_0} k^{1/q-1/p_{\alpha_0}} \asymp \nu_{\alpha_0} \mathfrak{N}_{\alpha_*, \alpha_0}^{1/p_{\alpha_*}-1/p_{\alpha_0}} \stackrel{(3)}{=} \nu_{\alpha_*}. \quad (45)$$

Let now

$$\sup\{p_\alpha : \alpha \in A_2 \cup A'_3\} = q.$$

We set

$$A^\delta = A'_1 \cup \{\alpha \in A_2 \cup A'_3 : 1/p_\alpha - 1/q \geq \delta\}, \quad (46)$$

$M_\delta = \cap_{\alpha \in A^\delta} \nu_\alpha B_{p_\alpha}^N$. By (45),

$$d_n(M_\delta, l_q^N) \underset{q}{\gtrsim} \nu_{\alpha_*}.$$

We prove that if

$$0 < \delta \leq \frac{\ln 2}{\ln N}, \quad (47)$$

then $M_\delta \subset 2M$. It suffices to check that

$$2 \cap_{\alpha \in A \setminus A^\delta} \nu_\alpha B_{p_\alpha}^N \supset \nu_{\alpha_*} B_q^N. \quad (48)$$

Since $p_\alpha < q$ for all $\alpha \in A \setminus A^\delta$, (48) holds if

$$\nu_{\alpha_*} \leq 2\nu_\alpha \cdot N^{1/q-1/p_\alpha}, \quad \alpha \in A \setminus A^\delta. \quad (49)$$

We have $\nu_\alpha \stackrel{(3)}{=} \nu_{\alpha_*} \mathfrak{N}_{\alpha_*, \alpha}^{1/p_\alpha-1/p_{\alpha_*}} \stackrel{(4),(46)}{\geq} \nu_{\alpha_*}$. This together with (46) and (47) yields (49).

3. Let

$$\Phi(n, N, q, K) = \nu_{\alpha_*} n^{-1/2} N^{1/q}, \quad p_{\alpha_*} = 2, \quad A_3 \neq \emptyset, \quad (50)$$

and let $n \geq N^{2/q}$.

First we consider the case

$$\inf\{p_\alpha : \alpha \in A'_1 \cup A_2\} > 2. \quad (51)$$

We take $\alpha_0 \in A'_1 \cup A_2$ such that

$$\varkappa_{\alpha_0, \alpha_*} = \max_{\gamma \in A'_1 \cup A_2} \varkappa_{\gamma, \alpha_*} \quad (52)$$

(by (51), the maximum is attained and $p_{\alpha_0} > 2$). Let $k = \lfloor \varkappa_{\alpha_0, \alpha_*} \rfloor$.

We check that

$$\nu_{\alpha_0} k^{-1/p_{\alpha_0}} V_k \subset 2M. \quad (53)$$

It suffices to prove the inequality

$$\nu_{\alpha_0} \varkappa_{\alpha_0, \alpha_*}^{1/p_{\gamma}-1/p_{\alpha_0}} \leq \nu_{\gamma}, \quad \gamma \in A. \quad (54)$$

If $\gamma \in A'_3$, then

$$\begin{aligned} \nu_{\alpha_0} \varkappa_{\alpha_0, \gamma}^{1/2-1/p_{\alpha_0}} n^{-1/2} N^{1/q} &\stackrel{(10),(13),(50)}{\geq} \nu_{\alpha_*} n^{-1/2} N^{1/q} \stackrel{(3)}{=} \\ &= \nu_{\alpha_0} \varkappa_{\alpha_0, \alpha_*}^{1/p_{\alpha_*}-1/p_{\alpha_0}} n^{-1/2} N^{1/q} = \nu_{\alpha_0} \varkappa_{\alpha_0, \alpha_*}^{1/2-1/p_{\alpha_0}} n^{-1/2} N^{1/q}; \end{aligned}$$

this together with the condition $p_{\alpha_0} > 2$ yields that $\varkappa_{\alpha_0, \gamma} \geq \varkappa_{\alpha_0, \alpha_*}$. Since $p_{\gamma} \leq 2 < p_{\alpha_0}$, we have

$$\nu_{\alpha_0} \varkappa_{\alpha_0, \alpha_*}^{1/p_{\gamma}-1/p_{\alpha_0}} \leq \nu_{\alpha_0} \varkappa_{\alpha_0, \gamma}^{1/p_{\gamma}-1/p_{\alpha_0}} \stackrel{(3)}{=} \nu_{\gamma}.$$

If $\gamma \in A'_1 \cup A_2$, then by (3) inequality (54) is equivalent to $\nu_{\alpha_*} \varkappa_{\alpha_0, \alpha_*}^{1/p_{\gamma}-1/p_{\alpha_*}} \leq \nu_{\alpha_*} \varkappa_{\gamma, \alpha_*}^{1/p_{\gamma}-1/p_{\alpha_*}}$; since $p_{\gamma} > p_{\alpha_*}$, it is equivalent to $\varkappa_{\alpha_0, \alpha_*} \geq \varkappa_{\gamma, \alpha_*}$; the last inequality holds by (52).

This completes the proof of (53).

Let us prove that

$$n \geq N^{2/q} \varkappa_{\alpha_0, \alpha_*}^{1-2/q}. \quad (55)$$

For $\alpha_0 \in A'_1$, it follows from

$$\nu_{\alpha_*} n^{-1/2} N^{1/q} \stackrel{(10),(11),(50)}{\leq} \nu_{\alpha_0} \varkappa_{\alpha_0, \alpha_*}^{1/q-1/p_{\alpha_0}} \stackrel{(3)}{=} \nu_{\alpha_*} \varkappa_{\alpha_0, \alpha_*}^{1/q-1/p_{\alpha_*}} = \nu_{\alpha_*} \varkappa_{\alpha_0, \alpha_*}^{1/q-1/2},$$

for $\alpha_0 \in A_2$, it follows from

$$\nu_{\alpha_*} n^{-1/2} N^{1/q} \stackrel{(10),(12),(50)}{\leq} \nu_{\alpha_0} (n^{-1/2} N^{1/q})^{\frac{1/p_{\alpha_0}-1/q}{1/2-1/q}} \stackrel{(3)}{=} \nu_{\alpha_*} \varkappa_{\alpha_0, \alpha_*}^{1/p_{\alpha_0}-1/p_{\alpha_*}} (n^{-1/2} N^{1/q})^{\frac{1/p_{\alpha_0}-1/q}{1/2-1/q}}.$$

Hence, we get

$$\begin{aligned} d_n(M, l_q^N) &\stackrel{(53)}{\gtrsim} d_n(\nu_{\alpha_0} k^{-1/p_{\alpha_0}} V_k, l_q^N) \stackrel{(18),(55)}{\gtrsim}_q \\ &\gtrsim \nu_{\alpha_0} \varkappa_{\alpha_0, \alpha_*}^{1/2-1/p_{\alpha_0}} n^{-1/2} N^{1/q} \stackrel{(3)}{=} \nu_{\alpha_*} n^{-1/2} N^{1/q}. \end{aligned}$$

If $\inf\{p_{\alpha} : \alpha \in A'_1 \cup A_2\} = 2$, we argue as in the previous cases and obtain the desired estimate.

4. Let

$$\Phi(n, N, q, K) = \nu_{\alpha_*} \varkappa_{\alpha_*, \beta_*}^{1/q-1/p_{\alpha_*}}, \quad \alpha_* \in A'_1, \quad \beta_* \in A'_2 \cup A'_3. \quad (56)$$

Notice that if $p_{\alpha_*} = q$, the right-hand side of (56) is equal to ν_{α_*} ; if $p_{\beta_*} = q$, we get by (3) $\nu_{\alpha_*} \varkappa_{\alpha_*, \beta_*}^{1/q-1/p_{\alpha_*}} = \nu_{\beta_*}$; thus, we have arrived at the case already considered. Further we suppose that $p_{\alpha_*} > q$, $p_{\beta_*} < q$.

Let $k = \lceil \varkappa_{\alpha_*, \beta_*} \rceil$.

We prove that

$$\nu_{\alpha_*} k^{-1/p_{\alpha_*}} V_k \subset 2M. \quad (57)$$

It suffices to check that

$$\nu_{\alpha_*} \varkappa_{\alpha_*, \beta_*}^{1/p_{\gamma}-1/p_{\alpha_*}} \leq \nu_{\gamma}, \quad \gamma \in A. \quad (58)$$

Let $\gamma \in A'_1$. We have $\nu_{\alpha_*} \varkappa_{\alpha_*, \beta_*}^{1/q-1/p_{\alpha_*}} \stackrel{(10),(11),(56)}{\leq} \nu_{\gamma} \varkappa_{\gamma, \beta_*}^{1/q-1/p_{\gamma}}$; by (3), it is equivalent to $\nu_{\beta_*} \varkappa_{\alpha_*, \beta_*}^{1/q-1/p_{\beta_*}} \leq \nu_{\beta_*} \varkappa_{\gamma, \beta_*}^{1/q-1/p_{\beta_*}}$; since $p_{\beta_*} < q$, we have $\varkappa_{\alpha_*, \beta_*} \geq \varkappa_{\gamma, \beta_*}$. This together with $p_{\gamma} \geq q > p_{\beta_*}$ implies that

$$\nu_{\alpha_*} \varkappa_{\alpha_*, \beta_*}^{1/p_{\gamma}-1/p_{\alpha_*}} \stackrel{(3)}{=} \nu_{\beta_*} \varkappa_{\alpha_*, \beta_*}^{1/p_{\gamma}-1/p_{\beta_*}} \leq \nu_{\beta_*} \varkappa_{\gamma, \beta_*}^{1/p_{\gamma}-1/p_{\beta_*}} \stackrel{(3)}{=} \nu_{\gamma}.$$

Let $\gamma \in A'_2 \cup A'_3$. We have

$$\nu_{\alpha_*} \varkappa_{\alpha_*, \beta_*}^{1/q-1/p_{\alpha_*}} \stackrel{(10),(11),(56)}{\leq} \nu_{\alpha_*} \varkappa_{\alpha_*, \gamma}^{1/q-1/p_{\alpha_*}};$$

hence, $\varkappa_{\alpha_*, \beta_*} \leq \varkappa_{\alpha_*, \gamma}$ by the condition $p_{\alpha_*} > q$. Since $p_{\gamma} \leq q < p_{\alpha_*}$, we have

$$\nu_{\alpha_*} \varkappa_{\alpha_*, \beta_*}^{1/p_{\gamma}-1/p_{\alpha_*}} \leq \nu_{\alpha_*} \varkappa_{\alpha_*, \gamma}^{1/p_{\gamma}-1/p_{\alpha_*}} \stackrel{(3)}{=} \nu_{\gamma}.$$

This concludes the proof of (58).

Now we check the inequality

$$n \leq N^{2/q} \varkappa_{\alpha_*, \beta_*}^{1-2/q}. \quad (59)$$

It suffices to consider $n \geq N^{2/q}$.

If $\beta_* \in A_2$, then (59) follows from

$$\nu_{\beta_*} \varkappa_{\alpha_*, \beta_*}^{1/q-1/p_{\beta_*}} \stackrel{(3)}{=} \nu_{\alpha_*} \varkappa_{\alpha_*, \beta_*}^{1/q-1/p_{\alpha_*}} \stackrel{(10),(12),(56)}{\leq} \nu_{\beta_*} (n^{-1/2} N^{1/q})^{\frac{1/p_{\beta_*}-1/q}{1/2-1/q}}.$$

If $\beta_* \in A'_3$, then (59) follows from

$$\nu_{\alpha_*} \varkappa_{\alpha_*, \beta_*}^{1/q-1/p_{\alpha_*}} \stackrel{(10),(13),(56)}{\leq} \nu_{\alpha_*} \varkappa_{\alpha_*, \beta_*}^{1/2-1/p_{\alpha_*}} n^{-1/2} N^{1/q}.$$

Hence,

$$\begin{aligned} d_n(M, l_q^N) &\stackrel{(57)}{\gtrsim} d_n(\nu_{\alpha_*} k^{-1/p_{\alpha_*}} V_k, l_q^N) \stackrel{(17),(59)}{\gtrsim_q} \\ &\gtrsim \nu_{\alpha_*} k^{1/q-1/p_{\alpha_*}} \gtrsim \nu_{\alpha_*} \varkappa_{\alpha_*, \beta_*}^{1/q-1/p_{\alpha_*}}. \end{aligned}$$

5. Let

$$\Phi(n, N, q, K) = \nu_{\alpha_*} \varkappa_{\alpha_*, \beta_*}^{1/2-1/p_{\alpha_*}} n^{-1/2} N^{1/q}, \quad \alpha_* \in A'_1 \cup A'_2, \quad \beta_* \in A'_3, \quad (60)$$

and let $n \geq N^{2/q}$. Notice that if $p_{\alpha_*} = 2$, the right-hand side of (60) equals to $\nu_{\alpha_*} n^{-1/2} N^{1/q}$; if $p_{\beta_*} = 2$, we have $\nu_{\alpha_*} \varkappa_{\alpha_*, \beta_*}^{1/2-1/p_{\alpha_*}} n^{-1/2} N^{1/q} \stackrel{(3)}{=} \nu_{\beta_*} n^{-1/2} N^{1/q}$. Thus, we have arrived at the case already considered.

Further we suppose that $p_{\alpha_*} > 2$, $p_{\beta_*} < 2$.

Let $k = \lfloor \varkappa_{\alpha_*, \beta_*} \rfloor$.

We show that

$$\nu_{\alpha_*} k^{-1/p_{\alpha_*}} V_k \subset 2M. \quad (61)$$

It suffices to check that

$$\nu_{\alpha_*} \varkappa_{\alpha_*, \beta_*}^{1/p_\gamma-1/p_{\alpha_*}} \leq \nu_\gamma, \quad \gamma \in A. \quad (62)$$

If $\gamma \in A'_1 \cup A'_2$, we have

$$\nu_{\alpha_*} \varkappa_{\alpha_*, \beta_*}^{1/2-1/p_{\alpha_*}} n^{-1/2} N^{1/q} \stackrel{(10),(13),(60)}{\leq} \nu_\gamma \varkappa_{\gamma, \beta_*}^{1/2-1/p_\gamma} n^{-1/2} N^{1/q};$$

this together with (3) yields that $\nu_{\beta_*} \varkappa_{\alpha_*, \beta_*}^{1/2-1/p_{\beta_*}} \leq \nu_{\beta_*} \varkappa_{\gamma, \beta_*}^{1/2-1/p_{\beta_*}}$. Since $p_{\beta_*} < 2$, it implies that $\varkappa_{\alpha_*, \beta_*} \geq \varkappa_{\gamma, \beta_*}$. Hence, by $p_\gamma \geq 2 > p_{\beta_*}$ we get

$$\nu_{\alpha_*} \varkappa_{\alpha_*, \beta_*}^{1/p_\gamma-1/p_{\alpha_*}} \stackrel{(3)}{=} \nu_{\beta_*} \varkappa_{\alpha_*, \beta_*}^{1/p_\gamma-1/p_{\beta_*}} \leq \nu_{\beta_*} \varkappa_{\gamma, \beta_*}^{1/p_\gamma-1/p_{\beta_*}} \stackrel{(3)}{=} \nu_\gamma.$$

If $\gamma \in A'_3$, we have

$$\nu_{\alpha_*} \varkappa_{\alpha_*, \beta_*}^{1/2-1/p_{\alpha_*}} n^{-1/2} N^{1/q} \stackrel{(10),(13),(60)}{\leq} \nu_{\alpha_*} \varkappa_{\alpha_*, \gamma}^{1/2-1/p_{\alpha_*}} n^{-1/2} N^{1/q}.$$

Since $p_{\alpha_*} > 2$, it yields that $\varkappa_{\alpha_*, \beta_*} \leq \varkappa_{\alpha_*, \gamma}$. Recall that $p_{\alpha_*} > 2 \geq p_\gamma$. Hence,

$$\nu_{\alpha_*} \varkappa_{\alpha_*, \beta_*}^{1/p_\gamma-1/p_{\alpha_*}} \leq \nu_{\alpha_*} \varkappa_{\alpha_*, \gamma}^{1/p_\gamma-1/p_{\alpha_*}} \stackrel{(3)}{=} \nu_\gamma.$$

This completes the proof of (62).

Now we show that

$$n \geq \varkappa_{\alpha_*, \beta_*}^{1-2/q} N^{2/q}. \quad (63)$$

If $\alpha_* \in A'_1$, then (63) follows from

$$\nu_{\alpha_*} \varkappa_{\alpha_*, \beta_*}^{1/2-1/p_{\alpha_*}} n^{-1/2} N^{1/q} \stackrel{(10),(11),(60)}{\leq} \nu_{\alpha_*} \varkappa_{\alpha_*, \beta_*}^{1/q-1/p_{\alpha_*}}.$$

If $\alpha_* \in A_2$, then (63) follows from

$$\nu_{\alpha_*} \varkappa_{\alpha_*, \beta_*}^{1/2-1/p_{\alpha_*}} n^{-1/2} N^{1/q} \stackrel{(10),(12),(60)}{\leq} \nu_{\alpha_*} (n^{-1/2} N^{1/q})^{\frac{1/p_{\alpha_*}-1/q}{1/2-1/q}}.$$

Hence,

$$\begin{aligned} d_n(M, l_q^N) &\stackrel{(61)}{\gtrsim} d_n(\nu_{\alpha_*} k^{-1/p_{\alpha_*}} V_k, l_q^N) \stackrel{(18),(63)}{\gtrsim} \\ &\gtrsim \nu_{\alpha_*} k^{1/2-1/p_{\alpha_*}} n^{-1/2} N^{1/q} \asymp \nu_{\alpha_*} \varkappa_{\alpha_*, \beta_*}^{1/2-1/p_{\alpha_*}} n^{-1/2} N^{1/q}. \end{aligned}$$

6. Let $n \leq N^{2/q}$, $\Phi(n, N, q, K) = \nu_{\alpha_*} \varkappa_{\alpha_*, \beta_*}^{1/2-1/p_{\alpha_*}}$, $\alpha_* \in A'_1 \cup A_2$, $\beta_* \in A_3$. If $\alpha_* \in A'_1$, it follows from (10), (11) that $\nu_{\alpha_*} \varkappa_{\alpha_*, \beta_*}^{1/2-1/p_{\alpha_*}} \leq \nu_{\alpha_*} \varkappa_{\alpha_*, \beta_*}^{1/q-1/p_{\alpha_*}}$; therefore, $\varkappa_{\alpha_*, \beta_*} \leq 1$. By (4), we get $\varkappa_{\alpha_*, \beta_*} = 1$ and $\Phi(n, N, q, K) = \nu_{\alpha_*} \varkappa_{\alpha_*, \beta_*}^{1/q-1/p_{\alpha_*}}$. We have arrived at the case already considered. If $\alpha_* \in A_2$, it follows from (10), (12) that $\nu_{\alpha_*} \varkappa_{\alpha_*, \beta_*}^{1/2-1/p_{\alpha_*}} \leq \nu_{\alpha_*}$. Hence, $\varkappa_{\alpha_*, \beta_*} \stackrel{(4)}{=} 1$ and $\Phi(n, N, q, K) = \nu_{\alpha_*}$. We have arrived at the case already considered.

7. Let $n \leq N^{2/q}$, $p_{\alpha_*} = 2$, $\Phi(n, N, q, K) = \nu_{\alpha_*}$. We prove that $\nu_{\alpha_*} \leq \nu_{\alpha}$ for all $\alpha \in A$. If $\alpha \in A_2$, it follows from (10), (12). If $\alpha \in A'_3$, then $\nu_{\alpha} \stackrel{(3)}{=} \nu_{\alpha_*} \varkappa_{\alpha_*, \alpha}^{1/p_{\alpha}-1/p_{\alpha_*}} \stackrel{(4)}{\geq} \nu_{\alpha_*}$. If $\alpha \in A'_1$, then $\nu_{\alpha_*} \leq \nu_{\alpha} \varkappa_{\alpha, \alpha_*}^{1/q-1/p_{\alpha}}$ by (10), (11). Applying (3), we get $\nu_{\alpha_*} \leq \nu_{\alpha_*} \varkappa_{\alpha, \alpha_*}^{1/q-1/p_{\alpha_*}}$, or $\varkappa_{\alpha, \alpha_*} \leq 1$. By (4), we have $\varkappa_{\alpha, \alpha_*} = 1$ and $\nu_{\alpha_*} = \nu_{\alpha}$. Hence, $M \supset \nu_{\alpha_*} B_1^N$, and by Theorem A

$$d_n(M, l_q^N) \geq \nu_{\alpha_*} d_n(B_1^N, l_q^N) \gtrsim_q \nu_{\alpha_*}.$$

This completes the proof. \square

Proof of Theorem 2. The upper estimate follows from Theorem C.

Let us prove the lower estimate. As in Theorem 1, we may assume that the set K is compact and the minimum in the right-hand side of (14) is attained at some $\alpha_* \in A$. Then

$$\nu_{\alpha_*} \left[n \log^{-1} \frac{2N}{n} \right]^{-1/p_{\alpha_*}} \leq \nu_{\alpha} \left[n \log^{-1} \frac{2N}{n} \right]^{-1/p_{\alpha}}, \quad \alpha \in A. \quad (64)$$

Let $C > 0$ be the constant from Theorem F, $k_* = \min \left\{ \lceil Cn \log^{-1} \frac{2N}{n} \rceil, N \right\}$. Notice that if $k_* = N$, then $n \asymp N$, $d_n(V_{k_*}, l_\infty^N) = d_n(B_\infty^N, l_\infty^N) = 1$.

We have

$$d_n(\nu_{\alpha_*} k_*^{-1/p_{\alpha_*}} V_{k_*}, l_\infty^N) \geq \nu_{\alpha_*} k_*^{-1/p_{\alpha_*}} / 2 \gtrsim \nu_{\alpha_*} \min \left\{ 1, n^{-1} \log \frac{2N}{n} \right\}^{1/p_{\alpha_*}}.$$

By (64), there is an absolute constant $C_1 > 0$ such that $\nu_{\alpha_*} k_*^{-1/p_{\alpha_*}} V_{k_*} \subset C_1 M$. This concludes the proof. \square

3 The case of arbitrary ν_α .

Notice that if $\inf_{\alpha \in A} \nu_\alpha = 0$, then $\cap_{\alpha \in A} \nu_\alpha B_{p_\alpha}^N = \{0\}$. Therefore, further we assume that $\inf_{\alpha \in A} \nu_\alpha > 0$.

We set $Z = \{1/p_\alpha : \alpha \in A\} \subset [0, 1]$; for $z = 1/p_\alpha$ we set $\nu(z) = \nu_\alpha$. Then $M = \cap_{z \in Z} \nu(z) B_{1/z}^N$, and (4) is equivalent to

$$1 \leq \frac{\nu(z)}{\nu(w)} \leq N^{z-w}, \quad z, w \in Z, \quad z \geq w.$$

Let $\nu : Z \rightarrow (0, \infty)$ be an arbitrary function, and let $\inf_{z \in Z} \nu(z) > 0$. We construct the function $\nu_{**} : Z \rightarrow (0, \infty)$ such that

$$1 \leq \frac{\nu_{**}(z)}{\nu_{**}(w)} \leq N^{z-w}, \quad z, w \in Z, \quad z \geq w, \quad (65)$$

$$M = \cap_{z \in Z} \nu_{**}(z) B_{1/z}^N. \quad (66)$$

Let

$$\nu_*(z) = \inf \{ \nu(w) : w \geq z \}, \quad z \in Z.$$

Then $\nu_*(z) \leq \nu(z)$ and the function ν_* is non-decreasing. We show that

$$M = \cap_{z \in Z} \nu_*(z) B_{1/z}^N. \quad (67)$$

Since $\nu_*(z) \leq \nu(z)$ for all $z \in Z$, it suffices to check that

$$\nu_*(w) B_{1/w}^N \supset \cap_{z \in Z} \nu(z) B_{1/z}^N, \quad w \in Z. \quad (68)$$

Let $w_n \geq w$, $\nu(w_n) \xrightarrow{n \rightarrow \infty} \nu_*(w)$. Then

$$\nu_*(w) B_{1/w}^N \supset \cap_{n \in \mathbb{N}} \nu(w_n) B_{1/w}^N \supset \cap_{n \in \mathbb{N}} \nu(w_n) B_{1/w_n}^N,$$

which implies (68).

Now we set

$$\nu_{**}(z) = \inf\{\nu_*(w) \cdot N^{z-w} : w \leq z\}.$$

Notice that

$$\nu_{**}(z) \leq \nu_*(z), \quad z \in Z. \quad (69)$$

We show that ν_{**} is a non-decreasing function. Indeed, let $z_1 \leq z_2$. If $w \leq z_1$, then $\nu_*(w) \cdot N^{z_1-w} \leq \nu_*(w) \cdot N^{z_2-w}$; hence, $\nu_{**}(z_1) \leq \inf\{\nu_*(w) \cdot N^{z_2-w} : w \leq z_1\}$. If $z_1 \leq w \leq z_2$, by the monotonicity of the function ν_* we get

$$\nu_*(w) \cdot N^{z_2-w} \geq \nu_*(z_1) \geq \nu_{**}(z_1).$$

Therefore, $\nu_{**}(z_2) \geq \nu_{**}(z_1)$.

Now we prove that

$$\frac{\nu_{**}(z_2)}{\nu_{**}(z_1)} \leq N^{z_2-z_1}, \quad z_1, z_2 \in Z, \quad z_1 \leq z_2. \quad (70)$$

This yields (65). For every $w \leq z_1$ we have

$$\nu_{**}(z_2) \leq \nu_*(w) \cdot N^{z_2-w} = \nu_*(w) \cdot N^{z_1-w} \cdot N^{z_2-z_1};$$

taking infimum over $w \leq z_1$, we get (70).

Now we show that

$$\cap_{z \in Z} \nu_{**}(z) B_{1/z}^N = \cap_{z \in Z} \nu_*(z) B_{1/z}^N. \quad (71)$$

By (69), it suffices to check that for every $w \in Z$

$$\nu_{**}(w) B_{1/w}^N \supset \cap_{z \in Z} \nu_*(z) B_{1/z}^N. \quad (72)$$

Let $w_n \leq w$, $\nu_*(w_n) \cdot N^{w-w_n} \xrightarrow{n \rightarrow \infty} \nu_{**}(w)$. Then

$$\nu_{**}(w) B_{1/w}^N \supset \cap_{n \in \mathbb{N}} \nu_*(w_n) N^{w-w_n} B_{1/w_n}^N \supset \cap_{n \in \mathbb{N}} \nu_*(w_n) B_{1/w_n}^N.$$

Hence, we obtain (72).

Now, (67) and (71) imply (66).

REFERENCES

- [1] A.A. Vasil'eva, "Kolmogorov widths of weighted Sobolev classes on a multi-dimensional domain with conditions on the derivatives of order r and zero", arXiv:2004.06013.
- [2] A.A. Vasil'eva, "Kolmogorov widths of the intersection of two finite-dimensional balls", arXiv:2007.04894v1.

- [3] A. Pietsch, “ s -numbers of operators in Banach space”, *Studia Math.*, **51** (1974), 201–223.
- [4] M.I. Stesin, “Aleksandrov diameters of finite-dimensional sets and of classes of smooth functions”, *Dokl. Akad. Nauk SSSR*, **220**:6 (1975), 1278–1281 [Soviet Math. Dokl.].
- [5] A.N. Kolmogorov, A. A. Petrov, Yu. M. Smirnov, “A formula of Gauss in the theory of the method of least squares”, *Izvestiya Akad. Nauk SSSR. Ser. Mat.* **11** (1947), 561–566 (in Russian).
- [6] S. B. Stechkin, “On the best approximations of given classes of functions by arbitrary polynomials”, *Uspekhi Mat. Nauk*, **9**:1(59) (1954) 133–134 (in Russian).
- [7] B.S. Kashin, “The diameters of octahedra”, *Usp. Mat. Nauk* **30**:4 (1975), 251–252 (in Russian).
- [8] B.S. Kashin, “The widths of certain finite-dimensional sets and classes of smooth functions”, *Math. USSR-Izv.*, **11**:2 (1977), 317–333.
- [9] E.D. Gluskin, “On some finite-dimensional problems of the theory of diameters”, *Vestn. Leningr. Univ.*, **13**:3 (1981), 5–10 (in Russian).
- [10] E.D. Gluskin, “Norms of random matrices and diameters of finite-dimensional sets”, *Math. USSR-Sb.*, **48**:1 (1984), 173–182.
- [11] A.Yu. Garnaev and E.D. Gluskin, “On widths of the Euclidean ball”, *Dokl. Akad. Nauk SSSR*, **277**:5 (1984), 1048–1052 [Sov. Math. Dokl. **30** (1984), 200–204]
- [12] V.M. Tikhomirov, “Theory of approximations”. In: *Current problems in mathematics. Fundamental directions*. vol. 14. (*Itogi Nauki i Tekhniki*) (Akad. Nauk SSSR, Vsesoyuz. Inst. Nauchn. i Tekhn. Inform., Moscow, 1987), pp. 103–260 [Encycl. Math. Sci. vol. 14, 1990, pp. 93–243].
- [13] E.M. Galeev, “The Kolmogorov diameter of the intersection of classes of periodic functions and of finite-dimensional sets”, *Math. Notes*, **29**:5 (1981), 382–388.
- [14] E.M. Galeev, “Widths of functional classes and finite-dimensional sets”, *Vladikavkaz. Mat. Zh.*, **13**:2 (2011), 3–14.
- [15] E. D. Gluskin, “Intersections of a cube with an octahedron are poorly approximated by subspaces of small dimension”, *Approximation of functions by special classes of operators*, Interuniv. Collect. Sci. Works, Vologda, 1987, pp. 35–41 (in Russian).
- [16] Yu. V. Malykhin, K. S. Ryutin, “The Product of Octahedra is Badly Approximated in the $l_{2,1}$ -Metric”, *Math. Notes*, **101**:1 (2017), 94–99.
- [17] S. V. Konyagin, Yu. V. Malykhin, C. S. Rjutin, “On Exact Recovery of Sparse Vectors from Linear Measurements”, *Math. Notes*, **94**:1 (2013), 107–114.