# Blow-up phenomena in a parabolic-elliptic-elliptic attraction-repulsion chemotaxis system with superlinear logistic degradation

Yutaro Chiyo\*<sup>†</sup>, Monica Marras<sup>‡</sup>, Yuya Tanaka\*, Tomomi Yokota\*

\*Department of Mathematics, Tokyo University of Science 1-3, Kagurazaka, Shinjuku-ku, Tokyo 162-8601, Japan

<sup>‡</sup>Department of Mathematics and Computer Sciences, University of Cagliari via Ospedale 72, 09123 Cagliari, Italy

November 4, 2021

**Abstract.** This paper is concerned with the attraction-repulsion chemotaxis system with superlinear logistic degradation,

$$\begin{cases} u_t = \Delta u - \chi \nabla \cdot (u \nabla v) + \xi \nabla \cdot (u \nabla w) + \lambda u - \mu u^k, & x \in \Omega, \ t > 0, \\ 0 = \Delta v + \alpha u - \beta v, & x \in \Omega, \ t > 0, \\ 0 = \Delta w + \gamma u - \delta w, & x \in \Omega, \ t > 0, \end{cases}$$

under homogeneous Neumann boundary conditions, in a ball  $\Omega \subset \mathbb{R}^n$   $(n \geq 3)$ , with constant parameters  $\lambda \in \mathbb{R}, \ k > 1, \ \mu, \chi, \xi, \alpha, \beta, \gamma, \delta > 0$ . Blow-up phenomena in the system have been well investigated in the case  $\lambda = \mu = 0$ , whereas the attraction-repulsion chemotaxis system with logistic degradation has been not studied. Under the condition that k > 1 is close to 1, this paper ensures a solution which blows up in  $L^{\infty}$ -norm and  $L^{\sigma}$ -norm with some  $\sigma > 1$  for some nonnegative initial data. Moreover, a lower bound of blow-up time is derived.

<sup>2010</sup> Mathematics Subject Classification. Primary: 35B44; Secondary: 35Q92, 92C17.

Key words and phrases: finite-time blow-up; chemotaxis; attraction-repulsion; logistic source.

<sup>&</sup>lt;sup>†</sup>Corresponding author.

E-mail: ycnewssz@gmail.com, mmarras@unica.it, yuya.tns.6308@gmail.com, yokota@rs.tus.ac.jp

#### 1. Introduction

Chemotaxis is a property of cells to move in response to the concentration gradient of a chemical substance produced by the cells. More precisely, it accounts for a process in which cells exhibit in response to chemoattractant and chemorepellent which are produced by themselves, that is, moving towards higher concentrations of an attractive signal and keeping away from a repulsive signal. A fully parabolic attraction-repulsion chemotaxis system was proposed by Painter and Hillen [18] to show the quorum effect in the chemotactic process and Luca et al. [10] to describe the aggregation of microglia observed in Alzheimer's disease, and can be approximated by a parabolic-elliptic-elliptic system.

In this paper we consider the parabolic–elliptic–elliptic attraction-repulsion chemotaxis system with superlinear logistic degradation,

m with superlinear logistic degradation,
$$\begin{cases}
u_{t} = \Delta u - \chi \nabla \cdot (u \nabla v) + \xi \nabla \cdot (u \nabla w) + \lambda u - \mu u^{k}, & x \in \Omega, \ t > 0, \\
0 = \Delta v + \alpha u - \beta v, & x \in \Omega, \ t > 0, \\
0 = \Delta w + \gamma u - \delta w, & x \in \Omega, \ t > 0, \\
\frac{\partial u}{\partial \nu} = \frac{\partial v}{\partial \nu} = \frac{\partial w}{\partial \nu} = 0, & x \in \partial\Omega, \ t > 0, \\
u(x,0) = u_{0}(x), & x \in \Omega,
\end{cases}$$
(1.1)

where  $\Omega := B_R(0) \subset \mathbb{R}^n$   $(n \geq 3)$  is an open ball centered at the origin with radius R > 0;  $\lambda \in \mathbb{R}$ , k > 1 and  $\mu, \chi, \xi, \alpha, \beta, \gamma, \delta$  are positive constants;  $\frac{\partial}{\partial \nu}$  is the outward normal derivative on  $\partial \Omega$ . Moreover, the initial data  $u_0$  is supposed to satisfy

$$u_0 \in C^0(\overline{\Omega})$$
 is radially symmetric and nonnegative. (1.2)

The functions u, v and w represent the cell density, the concentration of attractive and repulsive chemical substances, respectively.

Blow-up phenomena correspond to the concentration of organisms on chemical substances. Hence it is important to investigate whether a solution of system (1.1) blows up or not. In this paper we show finite-time blow-up in  $L^{\infty}$ -norm and  $L^{\sigma}$ -norm with some  $\sigma > 1$ , and derive a lower bound of blow-up time. Still more, not only blow-up phenomena but also global existence and boundedness have been studied in many literatures on chemotaxis systems (see [7], [2] and [1]). Before presenting the main results, we give an overview of known results about some problems related to (1.1).

We first focus on the chemotaxis system

$$\begin{cases} u_t = \Delta u - \chi \nabla \cdot (u \nabla v) + g(u), \\ \tau v_t = \Delta v + \alpha u - \beta v \end{cases}$$
 (1.3)

under homogeneous Neumann boundary conditions, where  $\chi, \alpha, \beta$  are positive constants and g is a function of logistic type,  $\tau \in \{0, 1\}$ . The system with  $g(u) \equiv 0$  was proposed by Keller and Segel [8]. Since then, system (1.3) was extensively investigated as listed below.

- If  $\tau = 1$ ,  $g(u) \equiv 0$  and  $\alpha = \beta = 1$ , global existence and boundedness as well as finite-time blow-up were investigated as follows. In the one-dimensional setting, Osaki and Yagi [17] showed that all solutions are global in time and bounded. In the two-dimensional setting, Nagai et al. [15] established global existence and boundedness under the condition  $\int_{\Omega} u_0(x) dx < \frac{4\pi}{\chi}$ . On the other hand, Herrero and Velázquez [6] presented existence of radially symmetric solutions which blow up in finite time. Winkler in [24] with  $\chi = 1$  and  $n \geq 3$ , derived that if  $\|u_0\|_{L^{\frac{n}{2}+\varepsilon}(\Omega)}$  and  $\|\nabla v_0\|_{L^{n+\varepsilon}(\Omega)}$  are small for sufficiently small  $\varepsilon > 0$ , then a solution is global and bounded. Also, Winkler in [25] proved finite-time blow-up under some conditions for initial data  $(u_0, v_0)$ .
- If  $\tau=1$  and  $g(u)=\lambda u-\mu u^k$  with  $\lambda,\mu>0$ , global existence for any k>1 and stabilization for  $k\geq 2-\frac{2}{n}$  were achieved in a generalized solution concept by Winkler [27]. Also, for certain choices of  $\lambda,\mu$ , Yan and Fuest in [28], derived global existence of weak solutions under the condition  $k>\min\{2-\frac{2}{n},\ 2-\frac{4}{n+4}\},\ n\geq 2$  and  $\alpha=\beta=1$ . In particular for n=2, they showed that taking any k>1 suffices to exclude the possibility of collapse into a persistent Dirac distribution.
- If  $\tau=0$ ,  $g(u)\equiv 0$  and  $\beta=1$ , Nagai in [13] proved global existence and boundedness when n=1, or n=2 and  $\int_{\Omega}u_0(x)\,dx<\frac{4\pi}{\chi\alpha}$ , and finite-time blow-up under some condition for the energy function and the moment of u when  $n\geq 2$ . Also, in the two-dimensional setting, Nagai in [14] obtained global existence and boundedness under the condition  $\int_{\Omega}u_0(x)\,dx<\frac{8\pi}{\chi\alpha}$ , and finite-time blow-up under the conditions that  $\alpha=1$ ,  $\int_{\Omega}u_0(x)\,dx>\frac{8\pi}{\chi}$  and that

$$\int_{\Omega} u_0(x)|x-x_0|^2 dx \text{ is sufficiently small for some } x_0 \in \Omega.$$
 (1.4)

- If  $\tau = 0$ ,  $g(u) \le a \mu u^2$  with a > 0,  $\mu > 0$   $(n \le 2)$ ,  $\mu > \frac{n-2}{n}\chi$   $(n \ge 3)$  and  $\alpha = \beta = 1$ , Tello and Winkler in [22] showed global existence and boundedness.
- If  $\tau = 0$  and  $\chi = \alpha = \beta = 1$ , when  $g(u) = \lambda u \mu u^k$  with  $\lambda \in \mathbb{R}$ ,  $\mu > 0$  and k > 1, Winkler in [26] established finite-time blow-up in  $L^{\infty}$ -norm under suitable conditions on data; more precisely, the author asserted that if  $\Omega = B_R(0) \subset \mathbb{R}^n$  with  $n \geq 3$ , R > 0 and  $1 < k < \frac{7}{6}$   $(n \in \{3,4\})$ ,  $1 < k < 1 + \frac{1}{2(n-1)}$   $(n \geq 5)$ , then system (1.3) admits a solution which blows up in  $L^{\infty}$ -norm at finite time. In [12], Marras and Vernier derived finite-time blow-up in  $L^{\sigma}$ -norm with  $\sigma > \frac{n}{2}$  and finally obtained a lower bound of blow-up time. Moreover, as to system (1.3) with nonlinear diffusion, finite-time blow-up in  $L^{\infty}$ -norm was obtained by Black et al. in [3] (see also [19], [20] for weak chemotactic sensitivity and [11] for finite-time blow-up in  $L^{p}$ -norm to more general chemotaxis system).

We now shift our attention to the attraction-repulsion chemotaxis system

$$\begin{cases} u_t = \Delta u - \chi \nabla \cdot (u \nabla v) + \xi \nabla \cdot (u \nabla w) + g(u), \\ \tau v_t = \Delta v + \alpha u - \beta v, \\ \tau w_t = \Delta w + \gamma u - \delta w \end{cases}$$
(1.5)

under homogeneous Neumann boundary conditions, where  $\chi, \xi, \alpha, \beta, \gamma, \delta > 0$  are constants and  $\tau \in \{0, 1\}$ . The system with  $\tau = 0$  and  $g(u) = \lambda u - \mu u^k$  coincides with (1.1), whereas the previous works on this system are collected as follows.

• If  $\tau = 0$  and  $g(u) \equiv 0$ , existence of solutions which blow up in  $L^{\infty}$ -norm at finite time was studied in [21] and [9]. More precisely, in the two-dimensional setting, Tao and Wang [21] derived finite-time blow-up under the conditions (1.4) and

(i) 
$$\chi \alpha - \xi \gamma > 0$$
,  $\delta = \beta$  and  $\int_{\Omega} u_0(x) dx > \frac{8\pi}{\chi \alpha - \xi \gamma}$ .

Also, in the two-dimensional setting, Li and Li [9] extended the above (i) to the following two conditions:

(ii) 
$$\chi \alpha - \xi \gamma > 0$$
,  $\delta \ge \beta$  and  $\int_{\Omega} u_0(x) \, dx > \frac{8\pi}{\chi \alpha - \xi \gamma}$ ;  
(iii)  $\chi \alpha \delta - \xi \gamma \beta > 0$ ,  $\delta < \beta$  and  $\int_{\Omega} u_0(x) \, dx > \frac{8\pi}{\chi \alpha \delta - \xi \gamma \beta}$ .

- If  $\tau = 0$  and  $g(u) \equiv 0$ , Yu et al. [29] replaced  $\chi \alpha \delta \xi \gamma \beta$  with  $\chi \alpha \xi \gamma$  in (iii) and filled the gap between the above (ii) and (iii). In [21], [9] and [29], blow-up phenomena were analyzed by introducing the linear combination of the solution components v, w such that  $z := \chi v \xi w$  (as to the fully parabolic case  $\tau = 1$ , see [5]).
- If  $\tau = 0$  and  $g(u) \equiv 0$ , explicit lower bound of blow-up time for system (1.5) was provided under the condition  $\chi \alpha \xi \gamma > 0$  in the two-dimensional setting (see [23]).

In summary, blow-up phenomena have been well studied in both a parabolic–elliptic Keller–Segel system and an attraction-repulsion one when logistic sources are missing. However, blow-up with effect of logistic degradation in a Keller–Segel system has been investigated, while for an attraction-repulsion system it is still an open problem.

The purpose of this paper is to solve the above open problem. Namely, we examine finite-time blow-up in the attraction-repulsion system (1.1) and we achieve a lower bound of the blow-up time.

We now state main theorems. The first one asserts finite-time blow-up in  $L^{\infty}$ -norm. The statement reads as follows.

**Theorem 1.1** (Finite-time blow-up in  $L^{\infty}$ -norm). Let  $\Omega = B_R(0) \subset \mathbb{R}^n$ ,  $n \geq 3$  and R > 0, and let  $\lambda \in \mathbb{R}$ ,  $\mu > 0$ ,  $\chi, \xi, \alpha, \beta, \gamma, \delta > 0$ . Assume that k > 1 satisfies

$$k < \begin{cases} \frac{7}{6} & \text{if } n \in \{3, 4\}, \\ 1 + \frac{1}{2(n-1)} & \text{if } n \ge 5, \end{cases}$$
 (1.6)

and  $\chi, \xi, \alpha, \gamma > 0$  fulfill  $\chi \alpha - \xi \gamma > 0$ . Then, for all L > 0, m > 0 and  $m_0 \in (0, m)$  one can find  $r_0 = r_0(R, \lambda, \mu, k, L, m, m_0) \in (0, R)$  with the property that whenever  $u_0$  satisfies (1.2) and is such that

$$u_0(x) \le L|x|^{-n(n-1)}$$
 for all  $x \in \Omega$  (1.7)

as well as

$$\int_{\Omega} u_0(x) dx \le m \quad but \quad \int_{B_{r_0}(0)} u_0(x) dx \ge m_0,$$

there exist  $T_{\text{max}} \in (0, \infty)$  and a classical solution (u, v, w) of system (1.1), uniquely determined by

$$u \in C^{0}(\overline{\Omega} \times [0, T_{\max})) \cap C^{2,1}(\overline{\Omega} \times (0, T_{\max})),$$

$$v, w \in \bigcap_{\vartheta > n} C^{0}([0, T_{\max}); W^{1,\vartheta}(\Omega)) \cap C^{2,1}(\overline{\Omega} \times (0, T_{\max})),$$

which blows up at  $t = T_{\text{max}}$  in the sense that

$$\lim_{t \nearrow T_{\text{max}}} \|u(\cdot, t)\|_{L^{\infty}(\Omega)} = \infty.$$
(1.8)

We next state a result which guarantees a solution blows up in  $L^{\sigma}$ -norm at the blow-up time in  $L^{\infty}$ -norm. The theorem is the following.

**Theorem 1.2** (Finite-time blow-up in  $L^{\sigma}$ -norm). Let  $\Omega = B_R(0) \subset \mathbb{R}^n$ ,  $n \geq 3$  and R > 0. Then, a classical solution (u, v, w) for  $t \in (0, T_{\text{max}})$ , provided by Theorem 1.1, is such that for all  $\sigma > \frac{n}{2}$ ,

$$\limsup_{t \nearrow T_{\max}} \|u(\cdot, t)\|_{L^{\sigma}(\Omega)} = \infty.$$

Define for all  $\sigma > 1$  the energy function

$$\Psi(t) := \frac{1}{\sigma} \|u(\cdot, t)\|_{L^{\sigma}(\Omega)}^{\sigma} \quad \text{with} \quad \Psi_0 := \Psi(0) = \frac{1}{\sigma} \|u_0\|_{L^{\sigma}(\Omega)}^{\sigma}. \tag{1.9}$$

The third theorem provides a lower bound of blow-up time. The result reads as follows.

**Theorem 1.3** (Lower bound of blow-up time). Let  $\Omega = B_R(0) \subset \mathbb{R}^n$ ,  $n \geq 3$  and R > 0. Then, for all  $\sigma > \frac{n}{2}$  and some  $B_1 \geq 0$ ,  $B_2, B_3 > 0$  depending on  $\lambda$ ,  $\mu$ ,  $\sigma$ , and n, the blow-up time  $T_{\text{max}}$ , provided by Theorem 1.1, satisfies the estimate

$$T_{\text{max}} \ge \int_{\Psi_0}^{\infty} \frac{d\eta}{B_1 \eta + B_2 \eta^{\gamma_1} + B_3 \eta^{\gamma_2}},$$
 (1.10)

with  $\gamma_1 := \frac{\sigma+1}{\sigma}$ ,  $\gamma_2 := \frac{2(\sigma+1)-n}{2\sigma-n}$ .

One of the difficulties in the proofs of the above theorems is that the transformation  $z := \chi v - \xi w$  does not work to reduce (1.1) to the Keller–Segel system in the case  $\beta \neq \delta$ , in contrast to the case  $\beta = \delta$  which ensures the simplification of (1.1) as

$$\begin{cases} u_t = \Delta u - \nabla \cdot (u\nabla z) + \lambda u - \mu u^k, & x \in \Omega, \ t > 0, \\ 0 = \Delta z + (\chi \alpha - \xi \gamma)u - \beta z, & x \in \Omega, \ t > 0, \end{cases}$$

which has already been studied in [12, 26]. To overcome the difficulty, we carry out the arguments in the literatures without using the above transformation z. In particular, we need to handle the effect caused by the repulsion term  $\xi \nabla \cdot (u \nabla w)$ .

This paper is organized as follows. In Section 2 we give preliminary results on local existence of classical solutions to (1.1) and some basic and useful facts. In Sections 3 and 4 we prove finite-time blow-up in  $L^{\infty}$ -norm and  $L^{\sigma}$ -norm (Theorems 1.1 and 1.2), respectively. Section 5 is devoted to the derivation of a lower bound of blow-up time (Theorem 1.3).

#### 2. Preliminaries

We start with the following lemma on local existence of classical solutions to (1.1). This lemma can be proved by a standard fixed point argument (see e.g., [22]).

**Lemma 2.1.** Let  $\Omega = B_R(0) \subset \mathbb{R}^n$ ,  $n \geq 3$  and R > 0, and let  $\lambda \in \mathbb{R}$ ,  $\mu > 0$ , k > 1,  $\chi, \xi, \alpha, \beta, \gamma, \delta > 0$ . Then for all nonnegative  $u_0 \in C^0(\overline{\Omega})$  there exists  $T_{\text{max}} \in (0, \infty]$  such that (1.1) possesses a unique classical solution (u, v, w) such that

$$u \in C^{0}(\overline{\Omega} \times [0, T_{\max})) \cap C^{2,1}(\overline{\Omega} \times (0, T_{\max})),$$
  
$$v, w \in \bigcap_{\vartheta > n} C^{0}([0, T_{\max}); W^{1,\vartheta}(\Omega)) \cap C^{2,1}(\overline{\Omega} \times (0, T_{\max})),$$

and

$$u \ge 0$$
,  $v \ge 0$ ,  $w \ge 0$  for all  $t \in (0, T_{\text{max}})$ .

Moreover,

if 
$$T_{\text{max}} < \infty$$
, then  $\limsup_{t \nearrow T_{\text{max}}} \|u(\cdot, t)\|_{L^{\infty}(\Omega)} = \infty$ . (2.1)

Remark 2.1. We can use  $\lim_{t\nearrow T_{\max}} \|u(\cdot,t)\|_{L^{\infty}(\Omega)}$  instead of  $\limsup_{t\nearrow T_{\max}} \|u(\cdot,t)\|_{L^{\infty}(\Omega)}$  in the blow-up criterion (2.1), because we can construct a classical solution on [0,T] with some positive time T depending only on  $\|u_0\|_{L^{\infty}(\Omega)}$  and discuss the extension of the classical solution in a neighborhood of its maximal existence time  $T_{\max}$ , if  $T_{\max} < \infty$ .

We next give some properties of the Neumann heat semigroup which will be used later. For the proof, see [4, Lemma 2.1] and [24, Lemma 1.3].

**Lemma 2.2.** Suppose  $(e^{t\Delta})_{t\geq 0}$  is the Neumann heat semigroup in  $\Omega$ , and let  $\mu_1 > 0$  denote the first non zero eigenvalue of  $-\Delta$  in  $\Omega$  under Neumann boundary conditions. Then there exist  $k_1, k_2 > 0$  which only depend on  $\Omega$  and have the following properties:

(i) if  $1 \le q \le p \le \infty$ , then

$$||e^{t\Delta}z||_{L^p(\Omega)} \le k_1 t^{-\frac{n}{2}(\frac{1}{q}-\frac{1}{p})} ||z||_{L^q(\Omega)}, \quad \forall \ t > 0$$
 (2.2)

holds for all  $z \in L^q(\Omega)$ .

(ii) If  $1 < q \le p \le \infty$ , then

$$\|e^{t\Delta}\nabla \cdot \boldsymbol{z}\|_{L^{p}(\Omega)} \le k_{2}\left(1 + t^{-\frac{1}{2} - \frac{n}{2}(\frac{1}{q} - \frac{1}{p})}\right)e^{-\mu_{1}t}\|\boldsymbol{z}\|_{L^{q}(\Omega)}, \quad \forall \ t > 0$$
(2.3)

is valid for any  $\mathbf{z} \in (L^q(\Omega))^n$ , where  $e^{t\Delta} \nabla \cdot$  is the extension of the operator  $e^{t\Delta} \nabla \cdot$  on  $(C_0^{\infty}(\Omega))^n$  to  $(L^q(\Omega))^n$ .

In Section 5 we will use the following lemma which is obtained by a minor adjustment of the power of the Gagliardo-Nirenberg inequality.

**Lemma 2.3.** Let  $\Omega$  be a bounded and smooth domain of  $\mathbb{R}^n$  with  $n \geq 1$ . Let  $r \geq 1$ ,  $0 < q \leq p \leq \infty$ , s > 0. Then there exists a constant  $C_{GN} > 0$  such that

$$||f||_{L^{\mathsf{p}}(\Omega)}^{\mathsf{p}} \le C_{\mathsf{GN}} \Big( ||\nabla f||_{L^{\mathsf{r}}(\Omega)}^{\mathsf{p}a} ||f||_{L^{\mathsf{q}}(\Omega)}^{\mathsf{p}(1-a)} + ||f||_{L^{\mathsf{s}}(\Omega)}^{\mathsf{p}} \Big)$$
(2.4)

for all  $f \in L^{q}(\Omega)$  with  $\nabla f \in (L^{r}(\Omega))^{n}$ , and  $a := \frac{\frac{1}{q} - \frac{1}{p}}{\frac{1}{q} + \frac{1}{n} - \frac{1}{r}} \in [0, 1]$ .

*Proof.* Following from the Gagliardo-Nirenberg inequality (see [16] for more details):

$$||f||_{L^{\mathsf{p}}(\Omega)}^{\mathsf{p}} \le \left[ c_{\mathsf{GN}} \Big( ||\nabla f||_{L^{\mathsf{p}}(\Omega)}^{a} ||f||_{L^{\mathsf{q}}(\Omega)}^{1-a} + ||f||_{L^{\mathsf{s}}(\Omega)} \Big) \right]^{\mathsf{p}},$$

with some  $c_{\rm GN} > 0$ , and then from the inequality

$$(a + b)^{\alpha} \le 2^{\alpha} (a^{\alpha} + b^{\alpha})$$
 for any  $a, b \ge 0, \alpha > 0$ ,

we arrive to (2.4) with  $C_{\text{GN}} = 2^{\mathsf{p}} c_{\text{GN}}^{\mathsf{p}}$ .

# 3. Finite-time blow-up in $L^{\infty}$ -norm

Throughout the sequel, we suppose that  $\Omega = B_R(0) \subset \mathbb{R}^n$   $(n \geq 3)$  with R > 0 and  $u_0$  satisfies condition (1.2) as well as  $\lambda \in \mathbb{R}$ ,  $\mu > 0$ , k > 1,  $\chi, \xi, \alpha, \beta, \gamma, \delta > 0$ . Then we denote by (u, v, w) = (u(r, t), v(r, t), w(r, t)) the local classical solution of (1.1) given in Lemma 2.1 and by  $T_{\text{max}} \in (0, \infty)$  its maximal existence time.

The goal of this section is to prove finite-time blow-up in  $L^{\infty}$ -norm. To this end, noting that  $u_0$  is radially symmetric and so are u, v, w, we first define the functions

$$U(s,t) := \int_0^{s^{\frac{1}{n}}} \rho^{n-1} u(\rho,t) \, d\rho, \quad s \in [0,R^n], \ t \in [0,T_{\text{max}}),$$

$$V(s,t) := \int_0^{s^{\frac{1}{n}}} \rho^{n-1} v(\rho,t) \, d\rho, \quad s \in [0,R^n], \ t \in [0,T_{\text{max}}),$$

$$W(s,t) := \int_0^{s^{\frac{1}{n}}} \rho^{n-1} w(\rho,t) \, d\rho, \quad s \in [0,R^n], \ t \in [0,T_{\text{max}}).$$

Then we prove the following lemma.

**Lemma 3.1.** Under the above notation, we have

$$U_{t}(s,t) = n^{2} s^{2-\frac{2}{n}} U_{ss}(s,t) + n\chi \alpha U(s,t) U_{s}(s,t) - n\chi \beta V(s,t) U_{s}(s,t) - n\xi \gamma U(s,t) U_{s}(s,t) + n\xi \delta W(s,t) U_{s}(s,t) + \lambda U(s,t) - n^{k-1} \mu \int_{0}^{s} U_{s}^{k}(\sigma,t) d\sigma$$
(3.1)

for all  $s \in (0, R^n)$ ,  $t \in (0, T_{\text{max}})$ .

*Proof.* By the definitions of U, V, W, we obtain

$$U_{s}(s,t) = \frac{1}{n}u(s^{\frac{1}{n}},t), \quad U_{ss}(s,t) = \frac{1}{n^{2}}s^{\frac{1}{n}-1}u_{r}(s^{\frac{1}{n}},t),$$

$$V_{s}(s,t) = \frac{1}{n}v(s^{\frac{1}{n}},t), \quad V_{ss}(s,t) = \frac{1}{n^{2}}s^{\frac{1}{n}-1}v_{r}(s^{\frac{1}{n}},t),$$

$$W_{s}(s,t) = \frac{1}{n}w(s^{\frac{1}{n}},t), \quad W_{ss}(s,t) = \frac{1}{n^{2}}s^{\frac{1}{n}-1}w_{r}(s^{\frac{1}{n}},t),$$

for all  $s \in (0, \mathbb{R}^n)$ ,  $t \in (0, T_{\text{max}})$ . Since u, v, w are radially symmetric functions, we see from the second and third equations in (1.1) that

$$\frac{1}{r^{n-1}}(r^{n-1}v_r(r,t))_r = -\alpha u(r,t) + \beta v(r,t),$$

$$\frac{1}{r^{n-1}}(r^{n-1}w_r(r,t))_r = -\gamma u(r,t) + \delta w(r,t),$$

from which we obtain

$$r^{n-1}v_r(r,t) = -\alpha U(r^n,t) + \beta V(r^n,t), \tag{3.2}$$

$$r^{n-1}w_r(r,t) = -\gamma U(r^n,t) + \delta W(r^n,t)$$
(3.3)

for all  $r \in (0, R)$ ,  $t \in (0, T_{\text{max}})$ . Moreover, rewriting the first equation in (1.1) in the radial coordinates as

$$u_t(r,t) = \frac{1}{r^{n-1}} (r^{n-1}u_r(r,t))_r - \chi \frac{1}{r^{n-1}} (u(r,t)r^{n-1}v_r(r,t))_r$$
$$+ \xi \frac{1}{r^{n-1}} (u(r,t)r^{n-1}w_r(r,t))_r$$
$$+ \lambda u(r,t) - \mu u^k(r,t)$$

and integrating it with respect to r over  $[0, s^{\frac{1}{n}}]$ , we have

$$U_t(s,t) = n^2 s^{2-\frac{2}{n}} U_{ss}(s,t) - n\chi U_s(s,t) s^{1-\frac{1}{n}} v_r(s^{\frac{1}{n}},t)$$

$$+ n\xi U_s(s,t) s^{1-\frac{1}{n}} w_r(s^{\frac{1}{n}},t)$$

$$+ \lambda U(s,t) - n^{k-1} \mu \int_0^s U_s^k(\sigma,t) d\sigma$$

for all  $s \in (0, \mathbb{R}^n)$ ,  $t \in (0, T_{\text{max}})$ . Thanks to (3.2) and (3.3), we arrive at (3.1).

Given  $p \in (0,1)$ ,  $s_0 \in (0,R^n)$ , we next derive a differential inequality for the moment-type function  $\Phi$  defined as

$$\Phi(t) := \int_0^{s_0} s^{-p}(s_0 - s)U(s, t) \, ds, \quad t \in [0, T_{\text{max}}).$$

**Lemma 3.2.** Let  $\lambda \in \mathbb{R}$ ,  $\mu > 0$ ,  $\chi, \xi, \alpha, \beta, \gamma, \delta > 0$  and let  $\chi \alpha - \xi \gamma > 0$ . Assume that k > 1 satisfies (1.6). Then there is  $p \in (1 - \frac{2}{n}, 1)$  with the following property: For all m > 0 and L > 0 there exist  $s_* \in (0, R^n)$  and  $C_1 > 0$  such that whenever  $u_0$  fulfills (1.2), (1.7) and  $\int_{\Omega} u_0(x) dx \leq m$ , for any  $s_0 \in (0, s_*)$  the function  $\Phi$  satisfies

$$\Phi'(t) \ge \frac{1}{C_1} s_0^{p-3} \Phi^2(t) - C_1 s_0^{\frac{2}{n} + 1 - p}$$
(3.4)

for all  $t \in (0, \widehat{T}_{\max})$ , where  $\widehat{T}_{\max} := \min\{1, T_{\max}\}$ . Moreover, for all  $m_0 \in (0, m)$  one can find  $s_0 \in (0, s_*)$  and  $r_0 = r_0(R, \lambda, \mu, k, L, m, m_0) \in (0, R)$  such that if  $\int_{B_{r_0}(0)} u_0(x) dx \ge m_0$  and  $\widehat{T}_{\max} > \frac{1}{2}$ , then for all  $t \in (0, \frac{1}{2})$ ,

$$\Phi'(t) \ge C_2 s_0^{p-3} \Phi^2(t), \tag{3.5}$$

where  $C_2$  is a positive constant.

*Proof.* By the definition of  $\Phi$  and equation (3.1), we have

$$\begin{split} \Phi'(t) &= \int_0^{s_0} s^{-p}(s_0 - s) U_t(s, t) \, ds \\ &= n^2 \int_0^{s_0} s^{2 - \frac{2}{n} - p}(s_0 - s) U_{ss}(s, t) \, ds \\ &+ n(\chi \alpha - \xi \gamma) \int_0^{s_0} s^{-p}(s_0 - s) U(s, t) U_s(s, t) \, ds \\ &- n \chi \beta \int_0^{s_0} s^{-p}(s_0 - s) V(s, t) U_s(s, t) \, ds \\ &+ n \xi \delta \int_0^{s_0} s^{-p}(s_0 - s) W(s, t) U_s(s, t) \, ds \\ &+ \lambda \int_0^{s_0} s^{-p}(s_0 - s) U(s, t) \, ds - n^{k-1} \mu \int_0^{s_0} s^{-p}(s_0 - s) \left[ \int_0^s U_s^k(\sigma, t) \, d\sigma \right] ds. \end{split}$$
(3.6)

Since  $U_s(s,t) = \frac{1}{n}u(s^{\frac{1}{n}},t) \ge 0$  and hence the fourth term on the right-hand side of (3.6) is nonnegative, we obtain

$$\Phi'(t) \ge n^2 \int_0^{s_0} s^{2-\frac{2}{n}-p} (s_0 - s) U_{ss}(s, t) \, ds$$

$$+ n(\chi \alpha - \xi \gamma) \int_0^{s_0} s^{-p} (s_0 - s) U(s, t) U_s(s, t) \, ds$$

$$- n\chi \beta \int_0^{s_0} s^{-p} (s_0 - s) V(s, t) U_s(s, t) \, ds$$

$$+ \lambda \int_0^{s_0} s^{-p} (s_0 - s) U(s, t) \, ds - n^{k-1} \mu \int_0^{s_0} s^{-p} (s_0 - s) \left[ \int_0^s U_s^k(\sigma, t) \, d\sigma \right] ds$$

for all  $t \in (0, T_{\text{max}})$ . Since  $\chi \alpha - \xi \gamma > 0$  by assumption, following the steps in [26, (4.3)], we can derive the differential inequalities (3.4) and (3.5); note that, in the assumption  $\widehat{T}_{\text{max}} > \frac{1}{2}$  for (3.5) the value  $\frac{1}{2}$  can be replaced with other positive values less than 1.  $\square$ 

Now, we can prove Theorem 1.1.

**Proof of Theorem 1.1.** Thanks to Lemma 3.2, in particular, from (3.5), we can see that  $T_{\text{max}} < \infty$ . Therefore, from blow-up criterion (2.1), we conclude that the finite-time blow-up in  $L^{\infty}$ -norm occurs. Namely, (1.8) is proved.

# 4. Finite-time blow-up in $L^{\sigma}$ -norm

In these next sections we will assume the conditions contained in Theorem 1.1. In order to prove Theorem 1.2, first we state the following lemmas.

**Lemma 4.1.** Let  $\Omega \subset \mathbb{R}^n$ ,  $n \geq 3$  be a bounded and smooth domain, and  $\lambda \in \mathbb{R}$ ,  $\mu > 0$ , k > 1. Then for a classical solution (u, v, w) of (1.1) we have

$$\int_{\Omega} u \, dx \le m_* \quad \text{for all } t \in (0, T_{\text{max}}), \tag{4.1}$$

with

$$m_* := \max \left\{ \int_{\Omega} u_0 \, dx, \, \left( \frac{\lambda_+}{\mu} |\Omega|^{k-1} \right)^{\frac{1}{k-1}} \right\},$$
 (4.2)

where  $\lambda_+ := \max\{0, \lambda\}.$ 

*Proof.* Integrating the first equation in (1.1) and applying the divergence theorem and boundary conditions of (1.1), we obtain

$$\frac{d}{dt} \int_{\Omega} u \, dx = \lambda \int_{\Omega} u \, dx - \mu \int_{\Omega} u^k \, dx \le \lambda_+ \int_{\Omega} u \, dx - \mu |\Omega|^{1-k} \left( \int_{\Omega} u \, dx \right)^k, \tag{4.3}$$

where in the last term we used Hölder's inequality:  $\int_{\Omega} u \leq |\Omega|^{\frac{k-1}{k}} (\int_{\Omega} u^k)^{\frac{1}{k}}$ . From (4.3) we deduce that  $y := \int_{\Omega} u \, dx$  fulfills

$$\begin{cases} y'(t) \le \lambda_+ y(t) - \bar{\mu} y^k(t), & \bar{\mu} := \mu |\Omega|^{1-k} & \text{for all } t \in (0, T_{\text{max}}), \\ y(0) = y_0, & y_0 := \int_{\Omega} u_0 \, dx. \end{cases}$$

Upon an ODE comparison argument this implies that

$$y(t) \le m_*$$
 for all  $t \in (0, T_{\text{max}})$ .

The lemma is proved.

We next prove the following lemma which plays an important role in the proof of Theorem 1.2.

**Lemma 4.2.** Let  $\Omega \subset \mathbb{R}^n$ ,  $n \geq 3$  be a bounded and smooth domain. Let (u, v, w) be a classical solution of system (1.1). If for some  $\sigma_0 > \frac{n}{2}$  there exists C > 0 such that

$$||u(\cdot,t)||_{L^{\sigma_0}(\Omega)} \le C$$
 for all  $t \in (0,T_{\max})$ ,

then, for some  $\hat{C} > 0$ ,

$$||u(\cdot,t)||_{L^{\infty}(\Omega)} \le \hat{C} \quad \text{for all } t \in (0,T_{\text{max}}). \tag{4.4}$$

*Proof.* For any  $x \in \Omega$ ,  $t \in (0, T_{\text{max}})$ , we set  $t_0 := \max\{0, t - 1\}$  and we consider the representation formula for u:

$$u(\cdot,t) = e^{(t-t_0)\Delta}u(\cdot,t_0) - \chi \int_{t_0}^t e^{(t-s)\Delta}\nabla \cdot (u(\cdot,s)\nabla v(\cdot,s)) ds$$
$$+ \xi \int_{t_0}^t e^{(t-s)\Delta}\nabla \cdot (u(\cdot,s)\nabla w(\cdot,s)) ds + \int_{t_0}^t e^{(t-s)\Delta} \left[\lambda u(\cdot,s) - \mu u^k(\cdot,s)\right] ds$$
$$=: u_1(\cdot,t) + u_2(\cdot,t) + u_3(\cdot,t) + u_4(\cdot,t)$$

and

$$0 \le u(\cdot, t) \le \|u_1(\cdot, t)\|_{L^{\infty}(\Omega)} + \|u_2(\cdot, t)\|_{L^{\infty}(\Omega)} + \|u_3(\cdot, t)\|_{L^{\infty}(\Omega)} + u_4(\cdot, t). \tag{4.5}$$

We have

$$||u_1(\cdot,t)||_{L^{\infty}(\Omega)} \le \max\{||u_0||_{L^{\infty}(\Omega)}, m_*k_1\} =: C_5, \tag{4.6}$$

with  $k_1 > 0$  and  $m_*$  defined in (4.2). In fact, if  $t \le 1$ , then  $t_0 = 0$  and hence the maximum principle yields  $u_1(\cdot,t) \le ||u_0||_{L^{\infty}(\Omega)}$ . If t > 1, then  $t - t_0 = 1$  and from (2.2) with  $p = \infty$  and q = 1, we deduce from (4.1) that  $||u_1(\cdot,t)||_{L^{\infty}(\Omega)} \le k_1(t-t_0)^{-\frac{n}{2}}||u(\cdot,t_0)||_{L^1(\Omega)} \le m_*k_1$ . We next use (2.3) with  $p = \infty$ , which leads to

$$||u_2(\cdot,t)||_{L^{\infty}(\Omega)} \le k_2 \chi \int_{t_0}^t (1 + (t-s)^{-\frac{1}{2} - \frac{n}{2q}}) e^{-\mu_1(t-s)} ||u(\cdot,s)\nabla v(\cdot,s)||_{L^q(\Omega)} ds. \tag{4.7}$$

Here, we may assume that  $\frac{n}{2} < \sigma_0 < n$ , and then we can fix q > n such that  $1 - \frac{(n - \sigma_0)q}{n\sigma_0} > 0$ , which enables us to pick  $\theta \in (1, \infty)$  fulfilling  $\frac{1}{\theta} < 1 - \frac{(n - \sigma_0)q}{n\sigma_0}$ , that is,  $\frac{q\theta}{\theta - 1} < \frac{n\sigma_0}{n - \sigma_0}$ . Then by Hölder's inequality, we can estimate

$$\begin{aligned} \|u(\cdot,s)\nabla v(\cdot,s)\|_{L^{q}(\Omega)} &\leq \|u(\cdot,s)\|_{L^{q\theta}(\Omega)} \|\nabla v(\cdot,s)\|_{L^{\frac{q\theta}{\theta-1}}(\Omega)} \\ &\leq C_6 \|u(\cdot,s)\|_{L^{q\theta}(\Omega)} \|\nabla v(\cdot,s)\|_{L^{\frac{n\sigma_0}{n-\sigma_0}}(\Omega)} \quad \text{for all } s \in (0,T_{\text{max}}), \end{aligned}$$

with some  $C_6 > 0$ . The Sobolev embedding theorem and elliptic regularity theory applied to the second equation in (1.1) tell us that  $\|v(\cdot,s)\|_{W^{1,\frac{n\sigma_0}{n-\sigma_0}}(\Omega)} \leq C_7\|v(\cdot,s)\|_{W^{2,\sigma_0}(\Omega)} \leq C_8$  with some  $C_7, C_8 > 0$ . Thus again by Hölder's inequality and (4.1), we obtain

$$||u(\cdot,s)\nabla v(\cdot,s)||_{L^{q}(\Omega)} \leq C_9||u(\cdot,s)||_{L^{q\theta}(\Omega)}$$
  
$$\leq C_{10}||u(\cdot,s)||_{L^{\infty}(\Omega)}^{\bar{\theta}} \quad \text{for all } s \in (0,T_{\text{max}}),$$

with some  $\bar{\theta} \in (0,1)$ ,  $C_9 := C_6 C_8$  and  $C_{10} := C_9 m_*^{1-\bar{\theta}}$ . Hence, combining this estimate and (4.7), we infer

$$||u_2(\cdot,t)||_{L^{\infty}(\Omega)} \le C_{10}k_2\chi \int_{t_0}^t (1+(t-s)^{-\frac{1}{2}-\frac{n}{2q}})e^{-\mu_1(t-s)}||u(\cdot,s)||_{L^{\infty}(\Omega)}^{\bar{\theta}} ds.$$

Now fix any  $T \in (0, T_{\text{max}})$ . Then, since  $t - t_0 \le 1$ , we have

$$||u_{2}(\cdot,t)||_{L^{\infty}(\Omega)} \leq C_{10}k_{2}\chi \int_{t_{0}}^{t} (1+(t-s)^{-\frac{1}{2}-\frac{n}{2q}}e^{-\mu_{1}(t-s)}) ds \cdot \sup_{t \in [0,T]} ||u(\cdot,t)||_{L^{\infty}(\Omega)}^{\bar{\theta}}$$

$$\leq C_{11}\chi \sup_{t \in [0,T]} ||u(\cdot,t)||_{L^{\infty}(\Omega)}^{\bar{\theta}}, \tag{4.8}$$

where  $C_{11} := C_{10}k_2(1 + \mu_1^{\frac{n}{2q} - \frac{1}{2}} \int_0^\infty r^{-\frac{1}{2} - \frac{n}{2q}} e^{-r} dr) > 0$  is finite, because  $-\frac{1}{2} - \frac{n}{2q} > -1$ . Analogously we can conclude

$$||u_3(\cdot,t)||_{L^{\infty}(\Omega)} \le C_{11}\xi \sup_{t \in [0,T]} ||u(\cdot,t)||_{L^{\infty}(\Omega)}^{\bar{\theta}}.$$
(4.9)

We next prove that there exists a constant  $C_{12} \ge 0$  such that  $u_4(\cdot, t) \le C_{12}$ . To this end, we firstly observe that

$$h(u) := \lambda u - \mu u^k \le h(u_*) =: C_{12},$$

with  $u_* := \left(\frac{\lambda_+}{\mu k}\right)^{\frac{1}{k-1}}$ . We have

$$u_4(\cdot,t) = \int_{t_0}^t e^{(t-s)\Delta} \left[ \lambda u(\cdot,s) - \mu u^k(\cdot,s) \right] ds \le C_{12} \int_{t_0}^t ds \le C_{12}.$$
 (4.10)

Plugging (4.6), (4.8), (4.9) and (4.10) into (4.5), we see that

$$0 \le u(x,t) \le C_5 + C_{12} + C_{11}(\chi + \xi) \sup_{t \in [0,T]} \|u(\cdot,t)\|_{L^{\infty}(\Omega)}^{\bar{\theta}},$$

which implies

$$\sup_{t \in [0,T]} \|u(\cdot,t)\|_{L^{\infty}(\Omega)} \le C_{13} + C_{14} \left( \sup_{t \in [0,T]} \|u(\cdot,t)\|_{L^{\infty}(\Omega)} \right)^{\bar{\theta}} \quad \text{for all } T \in (0,T_{\max}),$$

with  $C_{13} := C_5 + C_{12}$  and  $C_{14} := C_{11}(\chi + \xi)$ . From this inequality with  $\bar{\theta} \in (0, 1)$ , we arrive at (4.4).

**Proof of Theorem 1.2.** Since Theorem 1.1 holds, the unique local classical solution of (1.1) blows up at  $t = T_{\text{max}}$  in the sense  $\limsup_{t \nearrow T_{\text{max}}} \|u(\cdot,t)\|_{L^{\infty}(\Omega)} = \infty$  (i.e., (1.8)). By contradiction, we prove that it blows up also in  $L^{\sigma}$ -norm. In fact, if there exist  $\sigma_0 > \frac{n}{2}$  and C > 0 such that

$$||u(\cdot,t)||_{L^{\sigma_0}(\Omega)} \le C$$
 for all  $t \in (0,T_{\text{max}})$ ,

then, from Lemma 4.2, there exists  $\hat{C} > 0$  such that

$$||u(\cdot,t)||_{L^{\infty}(\Omega)} \leq \hat{C}$$
 for all  $t \in (0,T_{\max})$ ,

which is in contradiction to (1.8), so that, if u blows up in  $L^{\infty}$ -norm, then u blows up also in  $L^{\sigma}$ -norm for all  $\sigma > \frac{n}{2}$ .

# 5. A lower bound for $T_{\text{max}}$ , the proof of Theorem 1.3

Let us consider  $\Psi(t) = \frac{1}{\sigma} \int_{\Omega} u^{\sigma}(x,t) dx$ , u(x,t) the first component of solutions to (1.1) and we prove that  $\Psi$  satisfies a first order differential inequality.

In the proof of Theorem 1.3 we need an estimate for  $\int_{\Omega} u^{\sigma+1} dx$ . To this end, we use the Gagliardo-Nirenberg inequality (2.4) with  $f = u^{\frac{\sigma}{2}}$ ,  $p = \frac{2(\sigma+1)}{\sigma}$ , r = 2, q = 2, s = 2. Since  $\sigma > \frac{n}{2}$ , we have

$$\int_{\Omega} u^{\sigma+1} dx = \|u^{\frac{\sigma}{2}}\|_{L^{2(\sigma+1)}\sigma}^{\frac{2(\sigma+1)}{\sigma}}(\Omega)$$

$$\leq C_{\text{GN}} \|\nabla u^{\frac{\sigma}{2}}\|_{L^{2}(\Omega)}^{\frac{2(\sigma+1)}{\sigma}}(0) \|u^{\frac{\sigma}{2}}\|_{L^{2}(\Omega)}^{\frac{2(\sigma+1)}{\sigma}}(1-\theta_{0}) + C_{\text{GN}} \|u^{\frac{\sigma}{2}}\|_{L^{2}(\Omega)}^{\frac{2(\sigma+1)}{\sigma}}$$

$$= C_{\text{GN}} \left(\int_{\Omega} |\nabla u^{\frac{\sigma}{2}}|^{2} dx\right)^{\frac{\sigma+1}{\sigma}\theta_{0}} \left(\int_{\Omega} u^{\sigma} dx\right)^{\frac{\sigma+1}{\sigma}(1-\theta_{0})} + C_{\text{GN}} \left(\int_{\Omega} u^{\sigma} dx\right)^{\frac{\sigma+1}{\sigma}}$$

$$\leq C_{\text{GN}}\varepsilon_{1}\beta_{0} \int_{\Omega} |\nabla u^{\frac{\sigma}{2}}|^{2} dx + C_{\text{GN}}\varepsilon_{1}^{-\frac{\beta_{0}}{1-\beta_{0}}} (1-\beta_{0}) \left(\int_{\Omega} u^{\sigma} dx\right)^{\frac{(\sigma+1)(1-\theta_{0})}{\sigma(1-\beta_{0})}}$$

$$+ C_{\text{GN}} \left(\int_{\Omega} u^{\sigma} dx\right)^{\frac{\sigma+1}{\sigma}}$$

$$= c_{1}(\varepsilon_{1}) \int_{\Omega} |\nabla u^{\frac{\sigma}{2}}|^{2} dx + c_{2}(\varepsilon_{1}) \left(\int_{\Omega} u^{\sigma} dx\right)^{\frac{2(\sigma+1)-n}{2\sigma-n}} + c_{3} \left(\int_{\Omega} u^{\sigma} dx\right)^{\frac{\sigma+1}{\sigma}}, \quad (5.1)$$

with  $\varepsilon_1 > 0$ ,  $\theta_0 := \frac{n}{2(\sigma+1)} \in (0,1)$  and  $\beta_0 := \frac{\sigma+1}{\sigma}\theta_0 = \frac{n}{2\sigma} \in (0,1)$ . Now, we derive a differential inequality of the first order for  $\Psi(t)$ .

$$\Psi'(t) = \int_{\Omega} u^{\sigma-1} \Delta u \, dx - \chi \int_{\Omega} u^{\sigma-1} \nabla \cdot (u \nabla v) \, dx + \xi \int_{\Omega} u^{\sigma-1} \nabla \cdot (u \nabla w) \, dx$$
$$+ \lambda \int_{\Omega} u^{\sigma} \, dx - \mu \int_{\Omega} u^{\sigma+k-1} \, dx$$
$$=: \mathcal{I}_{1} + \mathcal{I}_{2} + \mathcal{I}_{3} + \mathcal{I}_{4} + \mathcal{I}_{5}. \tag{5.2}$$

We have:

$$\mathcal{I}_{1} = \int_{\Omega} u^{\sigma-1} \Delta u \, dx = -(\sigma - 1) \int_{\Omega} u^{\sigma-2} |\nabla u|^{2} \, dx$$

$$= -\frac{4(\sigma - 1)}{\sigma^{2}} \int_{\Omega} |\nabla u^{\frac{\sigma}{2}}|^{2} \, dx, \tag{5.3}$$

and

$$\mathcal{I}_{2} = -\chi \int_{\Omega} u^{\sigma-1} \nabla \cdot (u \nabla v) \, dx = \chi \frac{\sigma - 1}{\sigma} \int_{\Omega} \nabla u^{\sigma} \cdot \nabla v \, dx \\
= -\chi \frac{\sigma - 1}{\sigma} \int_{\Omega} u^{\sigma} \Delta v \, dx \\
= -\chi \beta \frac{\sigma - 1}{\sigma} \int_{\Omega} u^{\sigma} v \, dx + \chi \alpha \frac{\sigma - 1}{\sigma} \int_{\Omega} u^{\sigma+1} \, dx \\
\leq \chi \alpha \frac{\sigma - 1}{\sigma} \int_{\Omega} u^{\sigma+1} \, dx \tag{5.4}$$

as well as

$$\mathcal{I}_{3} = \xi \int_{\Omega} u^{\sigma-1} \nabla \cdot (u \nabla w) \, dx 
= \xi \delta \frac{\sigma - 1}{\sigma} \int_{\Omega} u^{\sigma} w \, dx - \xi \gamma \frac{\sigma - 1}{\sigma} \int_{\Omega} u^{\sigma+1} \, dx 
\leq \xi \delta \frac{\sigma - 1}{\sigma} \Big( \int_{\Omega} u^{\sigma+1} \, dx \Big)^{\frac{\sigma}{\sigma+1}} \Big( \int_{\Omega} w^{\sigma+1} \, dx \Big)^{\frac{1}{\sigma+1}} - \xi \gamma \frac{\sigma - 1}{\sigma} \int_{\Omega} u^{\sigma+1} \, dx 
\leq \xi \gamma \frac{\sigma - 1}{\sigma} \int_{\Omega} u^{\sigma+1} \, dx - \xi \gamma \frac{\sigma - 1}{\sigma} \int_{\Omega} u^{\sigma+1} \, dx 
= 0,$$
(5.5)

where the last inequality holds from  $(\int_{\Omega} w^{\sigma+1})^{\frac{1}{\sigma+1}} \leq \frac{\gamma}{\delta} (\int_{\Omega} u^{\sigma+1})^{\frac{1}{\sigma+1}}$  established by standard testing procedures in the equation for w. We now use (5.1) in (5.4) to obtain

$$\mathcal{I}_{2} \leq \tilde{c}_{1}(\varepsilon_{1}) \int_{\Omega} |\nabla u^{\frac{\sigma}{2}}|^{2} dx + \tilde{c}_{2}(\varepsilon_{1}) \left( \int_{\Omega} u^{\sigma} dx \right)^{\frac{2(\sigma+1)-n}{2\sigma-n}} + \tilde{c}_{3} \left( \int_{\Omega} u^{\sigma} dx \right)^{\frac{\sigma+1}{\sigma}}, \tag{5.6}$$

with  $\tilde{c}_1(\varepsilon_1) := \chi \alpha \frac{\sigma - 1}{\sigma} c_1(\varepsilon_1)$ ,  $\tilde{c}_2(\varepsilon_1) := \chi \alpha \frac{\sigma - 1}{\sigma} c_2(\varepsilon_1)$ ,  $\tilde{c}_3 := \chi \alpha \frac{\sigma - 1}{\sigma} c_3$ . Also, using Hölder's inequality, we see that

$$\mathcal{I}_{4} + \mathcal{I}_{5} = \lambda \int_{\Omega} u^{\sigma} dx - \mu \int_{\Omega} u^{\sigma+k-1} dx$$

$$\leq \lambda_{+} \int_{\Omega} u^{\sigma} dx - \mu |\Omega|^{\frac{1-k}{\sigma}} \left( \int_{\Omega} u^{\sigma} dx \right)^{\frac{\sigma+k-1}{\sigma}}.$$
(5.7)

Substituting (5.3), (5.5), (5.6) and (5.7) in (5.2) we get

$$\Psi' \leq B_1 \Psi + B_2 \Psi^{\frac{\sigma+1}{\sigma}} + B_3 \Psi^{\frac{2(\sigma+1)-n}{2\sigma-n}} - B_4 \Psi^{\frac{\sigma+k-1}{\sigma}} + \left(\tilde{c}_1(\varepsilon_1) - \frac{4(\sigma-1)}{\sigma^2}\right) \int_{\Omega} |\nabla u^{\frac{\sigma}{2}}|^2 dx, \tag{5.8}$$

with  $B_1 := \lambda_+ \sigma$ ,  $B_2 := \tilde{c}_3 \sigma^{\frac{\sigma+1}{\sigma}}$ ,  $B_3 := \tilde{c}_2(\varepsilon_1) \sigma^{\frac{2(\sigma+1)-n}{2\sigma-n}}$ ,  $B_4 := \mu |\Omega|^{\frac{1-k}{\sigma}} \sigma^{\frac{\sigma+k-1}{\sigma}}$ . In (5.8) we choose  $\varepsilon_1$  such that  $\tilde{c}_1(\varepsilon_1) - \frac{4(\sigma-1)}{\sigma^2} \leq 0$  and neglecting the negative terms, we obtain

$$\Psi' \le B_1 \Psi + B_2 \Psi^{\frac{\sigma+1}{\sigma}} + B_3 \Psi^{\frac{2(\sigma+1)-n}{2\sigma-n}}. \tag{5.9}$$

Integrating (5.9) from 0 to  $T_{\text{max}}$ , we arrive to (1.10).

Remark 5.1. Since u blows up in  $L^{\sigma}(\Omega)$ -norm at finite time  $T_{\text{max}}$ , then there exists a time  $t_1 \in [0, T_{\text{max}})$ , where  $\Psi(t_1) = \Psi_0$ . As a consequence,  $\Psi(t) \geq \Psi_0$ ,  $t \in [t_1, T_{\text{max}})$  so that  $\Psi^{\rho} \leq \Psi^{\gamma_2} \Psi_0^{\rho - \gamma_2}$  for some  $\rho \leq \gamma_2$ . Moreover, taking into account that  $1 < \frac{\sigma+1}{\sigma} \leq \frac{2(\sigma+1)-n}{2\sigma-n} = \gamma_2$ , it follows that

$$\Psi' \le A\Psi^{\gamma_2} \quad \text{in } (t_1, T_{\text{max}}), \tag{5.10}$$

with  $A := B_1 \Psi_0^{-\frac{2}{2\sigma-n}} + B_2 \Psi_0^{-\frac{n}{\sigma(2\sigma-n)}} + B_3$ . Integrating (5.10) from  $t_1$  to  $T_{\text{max}}$ , we derive the following explicit lower bound of the blow-up time  $T_{\text{max}}$ :

$$T_{\text{max}} \ge \frac{1}{A(\gamma_2 - 1)\Psi_0^{\gamma_2 - 1}}.$$

### Acknowledgments

The authors would like to express their gratitude to Professor Stella Vernier-Piro for giving them the opportunity of a joint study and her encouragement. YC, YT and TY are partially supported by Tokyo University of Science Grant for International Joint Research. MM is a member of the Gruppo Nazionale per l'Analisi Matematica, la Probabilità e le loro Applicazioni (GNAMPA) of the Istituto Nazionale di Alta Matematica (INdAM) and is partially supported by the research project Evolutive and stationary Partial Differential Equations with a focus on biomathematics (Fondazione di Sardegna 2019).

## References

- [1] G. Arumugam and J. Tyagi. Keller–Segel chemotaxis models: A review. *Acta Appl. Math.*, 171(6):82pp., 2021.
- [2] N. Bellomo, A. Bellouquid, Y. Tao, and M. Winkler. Toward a mathematical theory of Keller–Segel models of pattern formation in biological tissues. *Math. Models Methods Appl. Sci.*, 25(9):1663–1763, 2015.
- [3] T. Black, M. Fuest, and J. Lankeit. Relaxed parameter conditions for chemotactic collapse in logistic-type parabolic-elliptic Keller-Segel systems. arXiv: 2005.12089.
- [4] X. Cao. Global bounded solutions of the higher-dimensional Keller–Segel system under smallness conditions in optimal spaces. *Discrete Contin. Dyn. Syst.*, 35(5):1891–1904, 2015.
- [5] Y. Chiyo and T. Yokota. Remarks on finite-time blow-up in a fully parabolic attraction-repulsion chemotaxis system via reduction to the Keller–Segel system. arXiv:2103.02241.
- [6] M. A. Herrero and J. J. L. Velázquez. A blow-up mechanism for a chemotaxis model. *Ann. Scuola Norm. Sup. Pisa Cl. Sci.* (4), 24(4):633–683 (1998), 1997.
- [7] T. Hillen and K. J. Painter. A user's guide to PDE models for chemotaxis. *J. Math. Biol.*, 58(1–2):183–217, 2009.
- [8] E. F. Keller and L. A. Segel. Initiation of slime mold aggregation viewed as an instability. *J. Theoret. Biol.*, 26(3):399–415, 1970.
- [9] Y. Li and Y. Li. Blow-up of nonradial solutions to attraction-repulsion chemotaxis system in two dimensions. *Nonlinear Anal. Real World Appl.*, 30:170–183, 2016.
- [10] M. Luca, A. Chavez-Ross, L. Edelstein-Keshet, and A. Mogliner. Chemotactic signalling, microglia, and Alzheimer's disease senile plague: Is there a connection? *Bull. Math. Biol.*, 65:673–730, 2003.
- [11] M. Marras, T. Nishino, and G. Viglialoro. A refined criterion and lower bounds for the blow-up time in a parabolic–elliptic chemotaxis system with nonlinear diffusion. *Nonlinear Anal.*, 195:111725, 15, 2020.

- [12] M. Marras and S. Vernier-Piro. Finite time collapse in chemotaxis systems with logistic-type superlinear source. *Math. Methods Appl. Sci.*, 43(17):10027–10040, 2020.
- [13] T. Nagai. Blow-up of radially symmetric solutions to a chemotaxis system. Adv. Math. Sci. Appl., 5(2):581–601, 1995.
- [14] T. Nagai. Blowup of nonradial solutions to parabolic–elliptic systems modeling chemotaxis in two-dimensional domains. *J. Inequal. Appl.*, 6(1):37–55, 2001.
- [15] T. Nagai, T. Senba, and K. Yoshida. Application of the Trudinger–Moser inequality to a parabolic system of chemotaxis. *Funkcial. Ekvac.*, 40(3):411–433, 1997.
- [16] L. Nirenberg. On elliptic partial differential equations. Ann. Scuola Norm. Sup. Pisa Cl. Sci. (3), 13:115–162, 1959.
- [17] K. Osaki and A. Yagi. Finite dimensional attractor for one-dimensional Keller–Segel equations. *Funkcial. Ekvac.*, 44(3):441–469, 2001.
- [18] K. J. Painter and T. Hillen. Volume-filling and quorum-sensing in models for chemosensitive movement. *Can. Appl. Math. Q.*, 10(4):501–543, 2002.
- [19] Y. Tanaka. Blow-up in a quasilinear parabolic–elliptic Keller–Segel system with logistic source. arXiv: 2103.00159.
- [20] Y. Tanaka and T. Yokota. Blow-up in a parabolic–elliptic Keller–Segel system with density-dependent sublinear sensitivity and logistic source. *Math. Methods Appl. Sci.*, 43(12):7372–7396, 2020.
- [21] Y. Tao and Z-A. Wang. Competing effects of attraction vs. repulsion in chemotaxis. *Math. Models Methods Appl. Sci.*, 23(1):1–36, 2013.
- [22] J. I. Tello and M. Winkler. A chemotaxis system with logistic source. *Comm. Partial Differential Equations*, 32(4-6):849–877, 2007.
- [23] G. Viglialoro. Explicit lower bound of blow-up time for an attraction-repulsion chemotaxis system. J. Math. Anal. Appl., 479(1):1069–1077, 2019.
- [24] M. Winkler. Aggregation vs. global diffusive behavior in the higher-dimensional Keller–Segel model. *J. Differential Equations*, 248(12):2889–2905, 2010.
- [25] M. Winkler. Finite-time blow-up in the higher-dimensional parabolic-parabolic Keller-Segel system. J. Math. Pures Appl. (9), 100(5):748-767, 2013.
- [26] M. Winkler. Finite-time blow-up in low-dimensional Keller–Segel systems with logistic-type superlinear degradation. *Z. Angew. Math. Phys.*, 69(2):Paper No. 69, 40, 2018.
- [27] M. Winkler. Attractiveness of constant states in logistic-type Keller–Segel systems involving subquadratic growth restrictions. *Adv. Nonlinear Stud.*, 20(4):795–817, 2020.

- [28] J. Yan and M. Fuest. When do Keller–Segel systems with heterogeneous logistic sources admit generalized solutions? *Discrete Contin. Dyn. Syst. Ser. B*, to appear.
- [29] H. Yu, Q. Guo, and S. Zheng. Finite time blow-up of nonradial solutions in an attraction-repulsion chemotaxis system. *Nonlinear Anal. Real World Appl.*, 34:335–342, 2017.