

# Formulas and properties\*

S.V. Sudoplatov

## Abstract

We study links between first-order formulas and arbitrary properties for families of theories, classes of structures and their isomorphism types. Possibilities for ranks and degrees for formulas and theories with respect to given properties are described. Characteristics for generic sentences and generic theories with respect to properties are described and characterized.

**Key words:** formula, property, rank, degree, generic sentence, generic theory.

First-order formulas are used to express semantic and syntactic definable properties. Since in general there are more natural properties than definable ones, these formulas can express them in a partial way. In the present paper we study links between formulas and arbitrary properties, consider characteristics reflecting measures of their correspondence.

The paper is organized as follows. Preliminary notions, notations and results are represented in Section 1. In Section 2 we consider links between formulas and properties for semantic and syntactic families. We study and characterize these links with respect to constructions of formulas, set-theoretic operations and closures. In Section 3 we study, characterize and describe rank values and degree values for formulas with respect to given properties. In Section 4, we describe spectra for cardinalities of definable properties. In Section 5, generic sentences and theories with respect to properties are introduced, and their links and ranks are described.

Throughout we use the standard terminology in Mathematical Logic [1], notions, notations and results of [2, 3, 4].

## 1 Preliminaries

Let  $\Sigma$  be a language. If  $\Sigma$  is relational we denote by  $\mathcal{T}_\Sigma$  the family of all theories of the language  $\Sigma$ . If  $\Sigma$  contains functional symbols  $f$  then  $\mathcal{T}_\Sigma$  is the family of all theories of the language  $\Sigma'$ , which is obtained by replacements of all  $n$ -ary symbols  $f$  with  $(n + 1)$ -ary predicate symbols  $R_f$  interpreted by  $R_f = \{(\bar{a}, b) \mid f(\bar{a}) = b\}$ .

Following [2] we define the *rank*  $RS(\cdot)$  for families  $\mathcal{T} \subseteq \mathcal{T}_\Sigma$ , similar to Morley rank for a fixed theory, and a hierarchy with respect to these ranks in the following way.

By  $F(\Sigma)$  we denote the set of all formulas in the language  $\Sigma$  and by  $\text{Sent}(\Sigma)$  the set of all sentences in  $F(\Sigma)$ .

For a sentence  $\varphi \in \text{Sent}(\Sigma)$  we denote by  $\mathcal{T}_\varphi$  the set of all theories  $T \in \mathcal{T}$  with  $\varphi \in T$ .

---

\*The study was carried out within the framework of the state contract of the Sobolev Institute of Mathematics (project No. 0314-2019-0002), and the Committee of Science in Education and the Science Ministry of the Republic of Kazakhstan (Grant No. AP08855544).

Any set  $\mathcal{T}_\varphi$  is called the  $\varphi$ -neighbourhood, or simply a *neighbourhood*, for  $\mathcal{T}$ , or the ( $\varphi$ -)definable subset of  $\mathcal{T}$ . The set  $\mathcal{T}_\varphi$  is also called (*formula- or sentence-*)definable (by the sentence  $\varphi$ ) with respect to  $\mathcal{T}$ , or (*sentence-*) $\mathcal{T}$ -definable, or simply *s-definable*.

**Definition** [2]. For the empty family  $\mathcal{T}$  we put the rank  $\text{RS}(\mathcal{T}) = -1$ , for finite nonempty families  $\mathcal{T}$  we put  $\text{RS}(\mathcal{T}) = 0$ , and for infinite families  $\mathcal{T} - \text{RS}(\mathcal{T}) \geq 1$ .

For a family  $\mathcal{T}$  and an ordinal  $\alpha = \beta + 1$  we put  $\text{RS}(\mathcal{T}) \geq \alpha$  if there are pairwise inconsistent  $\Sigma(\mathcal{T})$ -sentences  $\varphi_n$ ,  $n \in \omega$ , such that  $\text{RS}(\mathcal{T}_{\varphi_n}) \geq \beta$ ,  $n \in \omega$ .

If  $\alpha$  is a limit ordinal then  $\text{RS}(\mathcal{T}) \geq \alpha$  if  $\text{RS}(\mathcal{T}) \geq \beta$  for any  $\beta < \alpha$ .

We set  $\text{RS}(\mathcal{T}) = \alpha$  if  $\text{RS}(\mathcal{T}) \geq \alpha$  and  $\text{RS}(\mathcal{T}) \not\geq \alpha + 1$ .

If  $\text{RS}(\mathcal{T}) \geq \alpha$  for any  $\alpha$ , we put  $\text{RS}(\mathcal{T}) = \infty$ .

A family  $\mathcal{T}$  is called *e-totally transcendental*, or *totally transcendental*, if  $\text{RS}(\mathcal{T})$  is an ordinal.

If  $\mathcal{T}$  is *e-totally transcendental*, with  $\text{RS}(\mathcal{T}) = \alpha \geq 0$ , we define the *degree*  $\text{ds}(\mathcal{T})$  of  $\mathcal{T}$  as the maximal number of pairwise inconsistent sentences  $\varphi_i$  such that  $\text{RS}(\mathcal{T}_{\varphi_i}) = \alpha$ .

**Definition** [5]. An infinite family  $\mathcal{T}$  is called *e-minimal* if for any sentence  $\varphi \in \Sigma(\mathcal{T})$ ,  $\mathcal{T}_\varphi$  is finite or  $\mathcal{T}_{\neg\varphi}$  is finite.

By the definition a family  $\mathcal{T}$  is *e-minimal* iff  $\text{RS}(\mathcal{T}) = 1$  and  $\text{ds}(\mathcal{T}) = 1$  [2], and iff  $\mathcal{T}$  has unique accumulation point [5].

In the paper [6] the notion of *E-closure* was introduced and characterized as follows:

**Proposition 1.1.** *If  $\mathcal{T} \subseteq \mathcal{T}_\Sigma$  is an infinite set and  $T \in \mathcal{T}_\Sigma \setminus \mathcal{T}$  then  $T \in \text{Cl}_E(\mathcal{T})$  (i.e.,  $T$  is an accumulation point for  $\mathcal{T}$  with respect to *E-closure*  $\text{Cl}_E$ ) if and only if for any sentence  $\varphi \in T$  the set  $\mathcal{T}_\varphi$  is infinite.*

The following theorem characterizes the property of *e-total transcendency* for countable languages.

**Theorem 1.2** [2]. *For any family  $\mathcal{T}$  with  $|\Sigma(\mathcal{T})| \leq \omega$  the following conditions are equivalent:*

- (1)  $|\text{Cl}_E(\mathcal{T})| = 2^\omega$ ;
- (2)  $e\text{-Sp}(\mathcal{T}) = 2^\omega$ ;
- (3)  $\text{RS}(\mathcal{T}) = \infty$ .

**Theorem 1.3** [3]. *For any language  $\Sigma$  either  $\text{RS}(\mathcal{T}_\Sigma)$  is finite, if  $\Sigma$  consists of finitely many 0-ary and unary predicates, and finitely many constant symbols, or  $\text{RS}(\mathcal{T}_\Sigma) = \infty$ , otherwise.*

For a language  $\Sigma$  we denote by  $\mathcal{T}_{\Sigma,n}$  the family of all theories in  $\mathcal{T}_\Sigma$  having  $n$ -element models,  $n \in \omega$ , as well as by  $\mathcal{T}_{\Sigma,\infty}$  the family of all theories in  $\mathcal{T}_\Sigma$  having infinite models.

**Theorem 1.4** [3]. *For any language  $\Sigma$  either  $\text{RS}(\mathcal{T}_{\Sigma,n}) = 0$ , if  $\Sigma$  is finite or  $n = 1$  and  $\Sigma$  has finitely many predicate symbols, or  $\text{RS}(\mathcal{T}_{\Sigma,n}) = \infty$ , otherwise.*

**Theorem 1.5** [3]. *For any language  $\Sigma$  either  $\text{RS}(\mathcal{T}_{\Sigma,\infty})$  is finite, if  $\Sigma$  is finite and without predicate symbols of arities  $m \geq 2$  as well as without functional symbols of arities  $n \geq 1$ , or  $\text{RS}(\mathcal{T}_{\Sigma,\infty}) = \infty$ , otherwise.*

By the definition the families  $\mathcal{T}_\Sigma$ ,  $\mathcal{T}_{\Sigma,n}$ ,  $\mathcal{T}_{\Sigma,\infty}$  are *E-closed*. Thus, combining Theorem 1.2 with Theorems 1.3–1.5 we obtain the following possibilities of cardinalities for the families  $\mathcal{T}_\Sigma$ ,  $\mathcal{T}_{\Sigma,n}$ ,  $\mathcal{T}_{\Sigma,\infty}$  depending on  $\Sigma$  and  $n \in \omega$ :

**Proposition 1.6.** For any language  $\Sigma$  either either  $\mathcal{T}_\Sigma$  is countable, if  $\Sigma$  consists of finitely many 0-ary and unary predicates, and finitely many constant symbols, or  $|\mathcal{T}_\Sigma| \geq 2^\omega$ , otherwise.

**Proposition 1.7.** For any language  $\Sigma$  either  $\mathcal{T}_{\Sigma,n}$  is finite, if  $\Sigma$  is finite or  $n = 1$  and  $\Sigma$  has finitely many predicate symbols, or  $|\mathcal{T}_{\Sigma,n}| \geq 2^\omega$ , otherwise.

**Proposition 1.8.** For any language  $\Sigma$  either  $\mathcal{T}_{\Sigma,\infty}$  is at most countable, if  $\Sigma$  is finite and without predicate symbols of arities  $m \geq 2$  as well as without functional symbols of arities  $n \geq 1$ , or  $|\mathcal{T}_{\Sigma,\infty}| \geq 2^\omega$ , otherwise.

**Definition [4].** If  $\mathcal{T}$  is a family of theories and  $\Phi$  is a set of sentences, then we put  $\mathcal{T}_\Phi = \bigcap_{\varphi \in \Phi} \mathcal{T}_\varphi$  and the set  $\mathcal{T}_\Phi$  is called (*type-*) or (*diagram-*)*definable* (by the set  $\Phi$ ) with respect to  $\mathcal{T}$ , or (*diagram-*) $\mathcal{T}$ -*definable*, or simply *d-definable*.

Clearly, finite unions of *d*-definable sets are again *d*-definable. Considering infinite unions  $\mathcal{T}'$  of *d*-definable sets  $\mathcal{T}_{\Phi_i}$ ,  $i \in I$ , one can represent them by sets of sentences with infinite disjunctions  $\bigvee_{i \in I} \varphi_i$ ,  $\varphi_i \in \Phi_i$ . We call these unions  $\mathcal{T}'$  are called *d<sub>∞</sub>-definable* sets.

**Definition [4].** Let  $\mathcal{T}$  be a family of theories,  $\Phi$  be a set of sentences,  $\alpha$  be an ordinal  $\leq \text{RS}(\mathcal{T})$  or  $-1$ . The set  $\Phi$  is called  $\alpha$ -*ranking* for  $\mathcal{T}$  if  $\text{RS}(\mathcal{T}_\Phi) = \alpha$ . A sentence  $\varphi$  is called  $\alpha$ -*ranking* for  $\mathcal{T}$  if  $\{\varphi\}$  is  $\alpha$ -*ranking* for  $\mathcal{T}$ .

The set  $\Phi$  (the sentence  $\varphi$ ) is called *ranking* for  $\mathcal{T}$  if it is  $\alpha$ -*ranking* for  $\mathcal{T}$  with some  $\alpha$ .

**Proposition 1.9 [4].** For any ordinals  $\alpha \leq \beta$ , if  $\text{RS}(\mathcal{T}) = \beta$  then  $\text{RS}(\mathcal{T}_\varphi) = \alpha$  for some ( $\alpha$ -*ranking*) sentence  $\varphi$ . Moreover, there are  $\text{ds}(\mathcal{T})$  pairwise  $\mathcal{T}$ -*inconsistent*  $\beta$ -*ranking* sentences for  $\mathcal{T}$ , and if  $\alpha < \beta$  then there are infinitely many pairwise  $\mathcal{T}$ -*inconsistent*  $\alpha$ -*ranking* sentences for  $\mathcal{T}$ .

**Theorem 1.10 [4].** Let  $\mathcal{T}$  be a family of a countable language  $\Sigma$  and with  $\text{RS}(\mathcal{T}) = \infty$ ,  $\alpha$  be a countable ordinal,  $n \in \omega \setminus \{0\}$ . Then there is a *d<sub>∞</sub>-definable* subfamily  $\mathcal{T}^* \subset \mathcal{T}$  such that  $\text{RS}(\mathcal{T}^*) = \alpha$  and  $\text{ds}(\mathcal{T}^*) = n$ .

**Theorem 1.11 [7].** For any two disjoint subfamilies  $\mathcal{T}_1$  and  $\mathcal{T}_2$  of an *E-closed* family  $\mathcal{T}$  the following conditions are equivalent:

- (1)  $\mathcal{T}_1$  and  $\mathcal{T}_2$  are separated by some sentence  $\varphi$ :  $\mathcal{T}_1 \subseteq \mathcal{T}_\varphi$  and  $\mathcal{T}_2 \subseteq \mathcal{T}_{\neg\varphi}$ ;
- (2) *E-closures* of  $\mathcal{T}_1$  and  $\mathcal{T}_2$  are disjoint in  $\mathcal{T}$ :  $\text{Cl}_E(\mathcal{T}_1) \cap \text{Cl}_E(\mathcal{T}_2) \cap \mathcal{T} = \emptyset$ ;
- (3) *E-closures* of  $\mathcal{T}_1$  and  $\mathcal{T}_2$  are disjoint:  $\text{Cl}_E(\mathcal{T}_1) \cap \text{Cl}_E(\mathcal{T}_2) = \emptyset$ .

**Definition [6].** Let  $\mathcal{T}_0$  be a family of theories. A subset  $\mathcal{T}'_0 \subseteq \mathcal{T}_0$  is said to be *generating* if  $\mathcal{T}_0 = \text{Cl}_E(\mathcal{T}'_0)$ . The generating set  $\mathcal{T}'_0$  (for  $\mathcal{T}_0$ ) is *minimal* if  $\mathcal{T}'_0$  does not contain proper generating subsets. A minimal generating set  $\mathcal{T}'_0$  is *least* if  $\mathcal{T}'_0$  is contained in each generating set for  $\mathcal{T}_0$ .

**Theorem 1.12 [6].** If  $\mathcal{T}'_0$  is a generating set for a *E-closed* set  $\mathcal{T}_0$  then the following conditions are equivalent:

- (1)  $\mathcal{T}'_0$  is the least generating set for  $\mathcal{T}_0$ ;
- (2)  $\mathcal{T}'_0$  is a minimal generating set for  $\mathcal{T}_0$ ;
- (3) any theory in  $\mathcal{T}'_0$  is isolated by some set  $(\mathcal{T}'_0)_\varphi$ , i.e., for any  $T \in \mathcal{T}'_0$  there is  $\varphi \in T$  such that  $(\mathcal{T}'_0)_\varphi = \{T\}$ ;
- (4) any theory in  $\mathcal{T}'_0$  is isolated by some set  $(\mathcal{T}_0)_\varphi$ , i.e., for any  $T \in \mathcal{T}'_0$  there is  $\varphi \in T$  such that  $(\mathcal{T}_0)_\varphi = \{T\}$ .

## 2 Relations between formulas and properties

**Definition.** Let  $\Sigma$  be a language,  $\varphi \equiv \varphi(\bar{x})$  be a formula in  $F(\Sigma)$ ,  $P_s$  be a subclass of the class  $K(\Sigma)$  of all structures  $\mathcal{A}$  in the language  $\Sigma$ . We say that  $\varphi(\bar{x})$  *partially* (respectively, *totally*) *satisfies*  $P_s$ , denoted by  $\varphi \triangleright_{ps} P_s$  or  $\varphi \triangleright_s^{\exists} P_s$  ( $\varphi \triangleright_{ts} P_s$  or  $\varphi \triangleright_s^{\forall} P_s$ ), if there are  $\mathcal{A} \in P_s$  and  $\bar{a} \in A$  (for any  $\mathcal{A} \in P_s$  there is  $\bar{a} \in A$ ) such that  $\mathcal{A} \models \varphi(\bar{a})$ .

If  $P_{is}$  is a subclass of the class  $\text{ITK}(\Sigma)$  of isomorphism types for the class  $K(\Sigma)$  then we say that  $\varphi(\bar{x})$  *partially* (respectively, *totally*) *satisfies*  $P_{its}$ , denoted by  $\varphi \triangleright_{pits} P_{its}$  or  $\varphi \triangleright_{its}^{\exists} P_{its}$  ( $\varphi \triangleright_{tits} P_{its}$  or  $\varphi \triangleright_{its}^{\forall} P_{its}$ ) if  $\varphi \triangleright_{ps} P_s$  ( $\varphi \triangleright_{ts} P_s$ , where  $P_s$  consists of all structures whose isomorphism types belong to  $P_{its}$ ).

If  $P_t$  is a subset of the set  $\mathcal{T}_\Sigma$  of all complete theories in the language  $\Sigma$  then we say that  $\varphi(\bar{x})$  *partially* (respectively, *totally*) *satisfies*  $P_t$ , denoted by  $\varphi \triangleright_{pt} P_t$  or  $\varphi \triangleright_t^{\exists} P_t$  ( $\varphi \triangleright_{tt} P_t$  or  $\varphi \triangleright_t^{\forall} P_t$ ), if there are  $T \in P_t$ ,  $\mathcal{M} \models T$ , and  $\bar{a} \in M$  (for any  $T \in P_t$  there are  $\mathcal{M} \models T$  and  $\bar{a} \in M$ ) such that  $\mathcal{M} \models \varphi(\bar{a})$ .

We write  $\not\triangleright_\xi$  if a  $\triangleright_\xi$ -relation does not hold.

**Remark 2.1.** By the definition we have the following obvious properties.

1. If  $P_s \neq \emptyset$  and  $\varphi \triangleright_{ts} P_s$  then  $\varphi \triangleright_{ps} P_s$ . Similarly, if  $\varphi \triangleright_{tits} P_{its}$  for nonempty  $P_{its}$  then  $\varphi \triangleright_{pits} P_{its}$ , and if  $\varphi \triangleright_{tt} P_t$  for nonempty  $P_t$  then  $\varphi \triangleright_{pt} P_t$ .

2. For any singleton  $P_s$ ,  $\varphi \triangleright_{ps} P_s$  implies  $\varphi \triangleright_{ts} P_s$ . Similarly,  $\varphi \triangleright_{pits} P_{its}$  implies  $\varphi \triangleright_{tits} P_{its}$  for any singleton  $P_{its}$ , and  $\varphi \triangleright_{pt} P_t$  implies  $\varphi \triangleright_{tt} P_t$  for any singleton  $P_t$ .

3. If  $\varphi \triangleright_{ps} \{\mathcal{A}\}$  and  $\mathcal{A} \equiv \mathcal{B}$  then  $\varphi \triangleright_{ps} \{\mathcal{B}\}$ . It implies that the relations  $\varphi \triangleright_{pt} P_t$  and  $\varphi \triangleright_{tt} P_t$  do not depend on the choice of models  $\mathcal{M} \models T$  for  $T \in P_t$ .

4. (Reflexivity) For any sentence  $\varphi$  and a (nonempty) family  $\mathcal{T}_\varphi \subseteq \mathcal{T}_\Sigma$  we have  $\varphi \triangleright_{tt} \mathcal{T}_\varphi$  (and  $\varphi \triangleright_{pt} \mathcal{T}_\varphi$ ).

5. (Monotony) If  $\varphi \triangleright_{ps} P_s$ ,  $\varphi \vdash \psi$  and  $P_s \subseteq P'_s \subseteq K(\Sigma)$  then  $\psi \triangleright_{ps} P'_s$ . If  $\varphi \triangleright_{ts} P_s$ ,  $\varphi \vdash \psi$  and  $P_s \supseteq P'_s$  then  $\psi \triangleright_{ps} P'_s$ . If  $\varphi \triangleright_{pits} P_{its}$ ,  $\varphi \vdash \psi$  and  $P_{its} \subseteq P'_{its} \subseteq \text{ITK}(\Sigma)$  then  $\psi \triangleright_{pits} P'_{its}$ . If  $\varphi \triangleright_{tits} P_{its}$ ,  $\varphi \vdash \psi$  and  $P_{its} \supseteq P'_{its}$  then  $\psi \triangleright_{ps} P'_{its}$ . If  $\varphi \triangleright_{pt} P_t$ ,  $\varphi \vdash \psi$  and  $P_t \subseteq P'_t \subseteq \mathcal{T}_\Sigma$  then  $\psi \triangleright_{ps} P'_t$ . If  $\varphi \triangleright_{tt} P_t$ ,  $\varphi \vdash \psi$  and  $P_t \supseteq P'_t$  then  $\psi \triangleright_{tt} P'_t$ .

For a property  $P_s$  we denote by  $\text{ITK}(P_s)$  the class of isomorphism types for structures in  $P_s$ , and by  $\text{Th}(P_s)$  the set  $\{T \in \mathcal{T}_\Sigma \mid \mathcal{A} \models T \text{ for some } \mathcal{A} \in P_s\}$ .

For a property  $P_{its}$  we denote by  $K(P_{its})$  the class of all structures whose isomorphism types are represented in  $P_{its}$ , and by  $\text{Th}(P_{its})$  the set  $\text{Th}(K(P_{its}))$ .

For a property  $P_t$  we denote by  $K(P_t)$  the class of all models of theories in  $P_t$ , and by  $\text{ITK}(P_t)$  the class  $\text{ITK}(K(P_t))$ .

In terms of these notations by the definition we have the following natural links between semantic properties  $P_s$  and  $P_{its}$  and syntactic properties  $P_t$ :

**Proposition 2.2.** *For any formula  $\varphi \in F(\Sigma)$  and properties  $P_s$ ,  $P_{its}$ ,  $P_t$  the following conditions hold:*

- (1)  $\varphi \triangleright_{ps} P_s$  iff  $\varphi \triangleright_{pits} \text{ITK}(P_s)$ , and iff  $\varphi \triangleright_{pt} \text{Th}(P_s)$ ;
- (2)  $\varphi \triangleright_{ts} P_s$  iff  $\varphi \triangleright_{tits} \text{ITK}(P_s)$ , and iff  $\varphi \triangleright_{tt} \text{Th}(P_s)$ ;
- (3)  $\varphi \triangleright_{pits} P_{its}$  iff  $\varphi \triangleright_{ps} K(P_{its})$ , and iff  $\varphi \triangleright_{pt} \text{Th}(P_{its})$ ;
- (4)  $\varphi \triangleright_{tits} P_{its}$  iff  $\varphi \triangleright_{ts} K(P_{its})$ , and iff  $\varphi \triangleright_{tt} \text{Th}(P_{its})$ ;
- (5)  $\varphi \triangleright_{pt} P_t$  iff  $\varphi \triangleright_{ps} K(P_t)$ , and iff  $\varphi \triangleright_{pits} \text{ITK}(P_t)$ ;
- (6)  $\varphi \triangleright_{tt} P_t$  iff  $\varphi \triangleright_{ts} K(P_t)$ , and iff  $\varphi \triangleright_{tits} \text{ITK}(P_t)$ .

In the items (3) and (4) the class  $K(P_{\text{its}})$  can be replaced by a subclass  $K'$  such that  $\text{ITK}(K') = P_{\text{its}}$ . Similarly, in the items (5) and (6) the class  $K(P_t)$  can be replaced by a subclass  $K'$  such that  $\text{Th}(K') = P_t$ , and independently  $\text{ITK}(P_t)$  can be replaced by a subclass  $K''$  such that  $\text{Th}(K'') = P_t$ .

By Proposition 2.2 semantic properties  $P_s$  and  $P_{\text{its}}$  can be naturally transformed into syntactic ones  $P_t$ , and vice versa. It means that natural model-theoretic properties such as  $\omega$ -categoricity, stability, simplicity etc. can be formulated both for theories, for structures and for their isomorphism types.

The links between  $\triangleright$ -relations which pointed out in Proposition 2.2 allow to reduce our consideration to the relations  $\triangleright_{\text{pt}}$  and  $\triangleright_{\text{tt}}$ . Besides, for the simplicity we will principally consider sentences  $\varphi$  instead of formulas in general. Reductions of formulas  $\psi(\bar{x})$  to sentences use the operators  $\psi(\bar{x}) \mapsto \forall \bar{x} \psi(\bar{x})$  and  $\psi(\bar{x}) \mapsto \exists \bar{x} \psi(\bar{x})$ .

**Proposition 2.3.** *For any sentences  $\varphi, \psi \in \text{Sent}(\Sigma)$  and properties  $P_t, P'_t \subseteq \mathcal{T}_\Sigma$  the following conditions hold:*

- (1) *if  $(\varphi \wedge \psi) \triangleright_{\text{pt}} (P_t \cap P'_t)$  then  $\varphi \triangleright_{\text{pt}} P_t$  and  $\psi \triangleright_{\text{pt}} P'_t$ ; the converse implication does not hold: there are  $\varphi', \psi' \in \text{Sent}(\Sigma)$  and  $P''_t \in \mathcal{T}_\Sigma$  such that  $\varphi' \triangleright_{\text{pt}} P''_t$ ,  $\psi' \triangleright_{\text{pt}} P''_t$ , and  $(\varphi' \wedge \psi') \not\triangleright_{\text{pt}} P''_t$ ;*
- (2)  *$(\varphi \wedge \psi) \triangleright_{\text{tt}} P_t$  iff  $\varphi \triangleright_{\text{tt}} P_t$  and  $\psi \triangleright_{\text{tt}} P_t$ ;*
- (3) *if  $\varphi \triangleright_{\text{tt}} P_t$  and  $\psi \triangleright_{\text{tt}} P'_t$  then  $(\varphi \wedge \psi) \triangleright_{\text{tt}} (P_t \cap P'_t)$ ; the converse implication does not hold: there are  $\varphi', \psi' \in \text{Sent}(\Sigma)$  and  $P''_t, P'''_t \in \mathcal{T}_\Sigma$  such that  $(\varphi \wedge \psi) \triangleright_{\text{tt}} (P''_t \cap P'''_t)$  whereas  $\varphi' \not\triangleright_{\text{tt}} P''_t$  and  $\psi' \not\triangleright_{\text{tt}} P'''_t$ .*

Proof. (1) If  $(\varphi \wedge \psi) \triangleright_{\text{pt}} (P_t \cap P'_t)$  then there is  $T \in P_t \cap P'_t$  with  $(\varphi \wedge \psi) \in T$ . Since  $\varphi, \psi \in T$ ,  $T \in P_t$  and  $T \in P'_t$ , we obtain  $\varphi \triangleright_{\text{pt}} P_t$  and  $\psi \triangleright_{\text{pt}} P'_t$ . Therefore it suffices to notice for (1) that sentences  $\varphi', \psi'$  asserting distinct finite cardinalities  $m$  and  $n$  for universes partially satisfy a property  $P''_t$  containing a theory  $T_1$  with an  $m$ -element model and a theory  $T_2$  with an  $n$ -element model. At the same time  $(\varphi' \wedge \psi') \not\triangleright_{\text{pt}} P''_t$  since  $(\varphi' \wedge \psi')$  is inconsistent.

(2) If  $(\varphi \wedge \psi) \triangleright_{\text{tt}} P_t$  then  $(\varphi \wedge \psi)$ , and so,  $\varphi$  and  $\psi$  belong to all theories in  $P_t$  implying  $\varphi \triangleright_{\text{tt}} P_t$  and  $\psi \triangleright_{\text{tt}} P_t$ . Conversely, if  $\varphi \triangleright_{\text{tt}} P_t$  and  $\psi \triangleright_{\text{tt}} P_t$  then  $\varphi, \psi$ , and so  $(\varphi \wedge \psi)$  belong to all theories in  $P_t$  implying  $(\varphi \wedge \psi) \triangleright_{\text{tt}} P_t$ .

(3) If  $\varphi \triangleright_{\text{tt}} P_t$  and  $\psi \triangleright_{\text{tt}} P'_t$  then  $\varphi \in \bigcap P_t$  and  $\psi \in \bigcap P'_t$ , implying  $(\varphi \wedge \psi) \in \bigcap P_t \cap \bigcap P'_t$ , i.e.,  $(\varphi \wedge \psi) \triangleright_{\text{tt}} (P_t \cap P'_t)$ . Finally, if  $P''_t$  and  $P'''_t$  are nonempty with  $P''_t \cap P'''_t = \emptyset$ , and  $\varphi', \psi'$  are inconsistent sentences then  $(\varphi \wedge \psi) \triangleright_{\text{tt}} (P''_t \cap P'''_t)$  and  $\varphi' \not\triangleright_{\text{pt}} P''_t$  and  $\psi' \not\triangleright_{\text{pt}} P'''_t$  implying  $\varphi' \not\triangleright_{\text{tt}} P''_t$  and  $\psi' \not\triangleright_{\text{tt}} P'''_t$ .  $\square$

**Proposition 2.4.** *For any sentences  $\varphi, \psi \in \text{Sent}(\Sigma)$  and properties  $P_t, P'_t \subseteq \mathcal{T}_\Sigma$  the following conditions hold:*

- (1) *if  $\varphi \triangleright_{\text{pt}} P_t$  or  $\psi \triangleright_{\text{pt}} P'_t$  then  $(\varphi \vee \psi) \triangleright_{\text{pt}} (P_t \cup P'_t)$ ; the converse implication does not hold: there are  $\varphi', \psi' \in \text{Sent}(\Sigma)$  and  $P''_t \in \mathcal{T}_\Sigma$  such that  $(\varphi' \vee \psi') \triangleright_{\text{pt}} P''_t$ , and  $\varphi' \not\triangleright_{\text{pt}} P''_t$  or  $\psi' \not\triangleright_{\text{pt}} P''_t$ ;*
- (2)  *$(\varphi \vee \psi) \triangleright_{\text{pt}} P_t$  iff  $\varphi \triangleright_{\text{pt}} P_t$  or  $\psi \triangleright_{\text{pt}} P_t$ ;*
- (3) *if  $\varphi \triangleright_{\text{tt}} P_t$  and  $\psi \triangleright_{\text{tt}} P'_t$  then  $(\varphi \vee \psi) \triangleright_{\text{tt}} (P_t \cup P'_t)$ ; the converse implication does not hold: there are  $\varphi', \psi' \in \text{Sent}(\Sigma)$  and  $P''_t \in \mathcal{T}_\Sigma$  such that  $(\varphi' \vee \psi') \triangleright_{\text{tt}} (P''_t)$  whereas  $\varphi' \not\triangleright_{\text{tt}} P''_t$  and  $\psi' \not\triangleright_{\text{tt}} P''_t$ .*

Proof. (1) If  $\varphi \triangleright_{\text{pt}} P_t$  or  $\psi \triangleright_{\text{pt}} P'_t$  then  $\varphi \in T$  for some  $T \in P_t$  or  $\psi \in T'$  for some  $T' \in P'_t$ . Therefore  $T$  or  $T'$  witness that  $(\varphi \vee \psi) \triangleright_{\text{pt}} (P_t \cup P'_t)$ . If  $\varphi'$  is a tautology and  $\psi'$  is an inconsistent sentence, then for any nonempty  $P''_t \subseteq \mathcal{T}_\Sigma$  we have  $(\varphi' \vee \psi') \triangleright_{\text{pt}} P''_t$ ,  $\varphi' \triangleright_{\text{pt}} P''_t$ , and  $\psi' \not\triangleright_{\text{pt}} P''_t$ .

(2) It holds since a sentence  $(\varphi \vee \psi)$  belongs to a complete theory  $T$  if and only if  $\varphi \in T$  or  $\psi \in T$ .

(3) If  $\varphi \triangleright_{tt} P_t$  and  $\psi \triangleright_{tt} P'_t$  then  $\varphi \in \bigcap P_t$  and  $\psi \in \bigcap P'_t$ . Therefore  $(\varphi \vee \psi) \in \bigcap P_t$  and  $(\varphi \vee \psi) \in \bigcap P'_t$  implying  $(\varphi \vee \psi) \in \bigcap P_t \cap \bigcap P'_t$ , hence  $(\varphi \vee \psi) \triangleright_{tt} (P_t \cup P'_t)$ .

Now let  $P''_t = \mathcal{T}_\Sigma$ ,  $\varphi'$  be a sentence belonging to some but not all theories in  $P''_t$ . For the sentence  $\psi' = \neg\varphi'$  we have  $(\varphi' \vee \psi') \triangleright_{tt} (P''_t)$  since  $(\varphi' \vee \psi')$  is a tautology,  $\varphi' \not\triangleright_{tt} P''_t$  and  $\psi' \not\triangleright_{tt} P''_t$  by the choice of  $\varphi'$ .  $\square$

**Proposition 2.5.** *For any sentence  $\varphi \in \text{Sent}(\Sigma)$  and a property  $P_t \subseteq \mathcal{T}_\Sigma$  the following conditions hold:*

- (1)  $\varphi \triangleright_{pt} P_t$  iff  $\neg\varphi \not\triangleright_{tt} P_t$ ;
- (2)  $\varphi \triangleright_{tt} P_t$  iff  $\neg\varphi \not\triangleright_{pt} P_t$ .

Proof. (1) If  $\varphi \triangleright_{pt} P_t$  then there is  $T \in P_t$  such that  $\varphi \in T$ . Since  $T$  is complete then  $\neg\varphi \notin T$  implying  $\neg\varphi \not\triangleright_{tt} P_t$ . Conversely if  $\neg\varphi \not\triangleright_{tt} P_t$  then  $\neg\varphi$  does not belong to some theory  $T \in P_t$ . Since  $T$  is complete then  $\varphi \in T$  implying  $\varphi \triangleright_{pt} P_t$ .

(2) is immediately follows from (1).  $\square$

**Proposition 2.6.** *For any formula  $\varphi \in F(\Sigma)$  and a property  $P_t \subseteq \mathcal{T}_\Sigma$  the following conditions hold:*

- (1) if  $\varphi = \forall x\psi$  and  $\varphi \triangleright_{pt} P_t$  then  $\psi \triangleright_{pt} P_t$ ;
- (2) if  $\varphi = \forall x\psi$  and  $\varphi \triangleright_{tt} P_t$  then  $\psi \triangleright_{tt} P_t$ ;
- (3) if  $\varphi = \exists x\psi$  and  $\psi \triangleright_{pt} P_t$  then  $\varphi \triangleright_{pt} P_t$ ;
- (3) if  $\varphi = \exists x\psi$  and  $\psi \triangleright_{tt} P_t$  then  $\varphi \triangleright_{tt} P_t$ .

Proof. (1) Let  $\varphi = \forall x\psi$  and  $\varphi \triangleright_{pt} P_t$ . Then there are  $T \in P_t$ ,  $\mathcal{M} \models T$ ,  $\bar{a} \in M$  such that  $\mathcal{M} \models \varphi(\bar{a})$ . It implies  $\mathcal{M} \models \forall x\psi(x, \bar{a})$ , therefore there is  $b \in M$  with  $\mathcal{M} \models \psi(b, \bar{a})$ , i.e.,  $\psi \triangleright_{pt} \{T\}$ , hence  $\psi \triangleright_{pt} P_t$ .

(2) We repeat arguments for (1) replacing some  $T \in P_t$  by an arbitrary one.

(3) Let  $\varphi = \exists x\psi$  and  $\psi \triangleright_{pt} P_t$ . Then there are  $T \in P_t$ ,  $\mathcal{M} \models T$ ,  $\bar{a}, b \in M$  such that  $\mathcal{M} \models \psi(b, \bar{a})$ . It implies  $\mathcal{M} \models \exists x\psi(x, \bar{a})$ , i.e.,  $\varphi \triangleright_{pt} \{T\}$ , hence  $\varphi \triangleright_{pt} P_t$ .

(4) As for (2) we repeat arguments for (3) replacing some  $T \in P_t$  by an arbitrary one.  $\square$

The following two theorems assert that the relations  $\triangleright_{pt}$  and  $\triangleright_{tt}$  are preserved under  $E$ -closures.

**Theorem 2.7.** *For any sentence  $\varphi \in \text{Sent}(\Sigma)$  and a property  $P_t \subseteq \mathcal{T}_\Sigma$  the following conditions are equivalent:*

- (1)  $\varphi \triangleright_{pt} P_t$ ,
- (2)  $\varphi \triangleright_{pt} \text{Cl}_E(P_t)$ ,
- (3)  $\varphi \triangleright_{pt} P'_t$  for any/some  $P'_t$  with  $\text{Cl}_E(P'_t) = \text{Cl}_E(P_t)$ .

Proof. (1)  $\Rightarrow$  (2) holds in view of  $P_t \subseteq \text{Cl}_E(P_t)$  and the monotony of the relation  $\triangleright_{pt}$ .

(2)  $\Rightarrow$  (3). It suffices to show that  $\varphi \triangleright_{pt} P'_t$  for any  $P'_t$  with  $\text{Cl}_E(P'_t) = \text{Cl}_E(P_t)$ . Since  $\varphi \triangleright_{pt} \text{Cl}_E(P_t)$  there is a theory  $T \in \text{Cl}_E(P_t)$  with  $\varphi \in T$ . If  $T \in P'_t$  we have  $\varphi \triangleright_{pt} P'_t$ . Otherwise  $T$  is an accumulation point of  $P'_t$  implying, in view of Proposition 1.1, that  $\text{varphi}$  belongs to infinitely many theories in  $P'_t$ . Therefore  $\varphi \triangleright_{pt} P'_t$ .

(3)  $\Rightarrow$  (1) is obvious.  $\square$

**Theorem 2.8.** *For any sentence  $\varphi \in \text{Sent}(\Sigma)$  and a property  $P_t \subseteq \mathcal{T}_\Sigma$  the following conditions are equivalent:*

- (1)  $\varphi \triangleright_{tt} P_t$ ,

- (2)  $\varphi \triangleright_{\text{tt}} \text{Cl}_E(P_t)$ ,
- (3)  $\varphi \triangleright_{\text{tt}} P'_t$  for any/some  $P'_t$  with  $\text{Cl}_E(P'_t) = \text{Cl}_E(P_t)$ .

Proof. (1)  $\Rightarrow$  (2). Let  $\varphi \triangleright_{\text{tt}} P_t$ . If  $P_t$  is finite then  $\text{Cl}_E(P_t) = P_t$  and we have  $\varphi \triangleright_{\text{tt}} \text{Cl}_E(P_t)$ . If  $P_t$  is infinite then by Proposition 1.1,  $\text{Cl}_E(P_t)$  consists of theories in  $P_t$  and of theories  $T \in \mathcal{T}_\Sigma \setminus P_t$  such that for any sentence  $\psi \in T$  the set  $(P_t)_\psi$  is infinite. Since  $\varphi \triangleright_{\text{tt}} P_t$ ,  $\varphi$  belongs to each such theory  $T$ . Thus,  $\varphi \triangleright_{\text{tt}} \text{Cl}_E(P_t)$ .

(2)  $\Rightarrow$  (1) and (2)  $\Rightarrow$  (3) are obvious.

(3)  $\Rightarrow$  (2) follows assuming  $\varphi \triangleright_{\text{tt}} P'_t$  for any/some  $P'_t$  with  $\text{Cl}_E(P'_t) = \text{Cl}_E(P_t)$  repeating the arguments for (1)  $\Rightarrow$  (2).  $\square$

For a property  $P_t \subseteq \mathcal{T}_\Sigma$  we denote by  $\nabla(P_t)$  the set of all sentences  $\varphi \in \text{Sent}(\Sigma)$  with  $\varphi \triangleright_{\text{pt}} P_t$ , and by  $\Delta(P_t)$  the set of all sentences  $\varphi \in \text{Sent}(\Sigma)$  with  $\varphi \triangleright_{\text{tt}} P_t$ .

By the definition  $\nabla(\emptyset) = \emptyset$ ,  $\Delta(\emptyset) = \text{Sent}(\Sigma)$ ,  $\nabla(\mathcal{T}_\Sigma)$  consists of all consistent sentences  $\varphi \in \text{Sent}(\Sigma)$ , and  $\Delta(\mathcal{T}_\Sigma)$  consists of all tautologies  $\varphi \in \text{Sent}(\Sigma)$ .

**Proposition 2.9.** *For any property  $P_t \subseteq \mathcal{T}_\Sigma$  the following conditions hold:*

- (1)  $\nabla(P_t) = \bigcup P_t$ ;
- (2)  $\nabla(P_t)$  is consistent iff  $|P_t| \leq 1$ , and  $\nabla(P_t)$  is a complete theory iff  $P_t$  is a singleton;
- (3)  $\Delta(P_t) = \bigcap P_t$ ;
- (4)  $\Delta(P_t)$  is a consistent theory iff  $P_t \neq \emptyset$ , and  $\Delta(P_t)$  is complete iff  $P_t$  is a singleton;
- (5) for any  $P_t \neq \emptyset$ ,  $\nabla(P_t) \supseteq \Delta(P_t)$ , and  $\nabla(P_t) = \Delta(P_t)$  iff  $P_t$  is a singleton.

Proof. (1) If  $\varphi \in \nabla(P_t)$  then  $\varphi \triangleright_{\text{pt}} P_t$  and  $\varphi \in T$  for some  $T \in P_t$  implying  $\nabla(P_t) \subseteq \bigcup P_t$ . Conversely, if  $\varphi \in \bigcup P_t$  then  $\varphi \in T$  for some  $T \in P_t$  implying  $\varphi \triangleright_{\text{pt}} P_t$  and therefore  $\varphi \in \nabla(P_t)$ .

(2) Since  $\nabla(\emptyset) = \emptyset$  it is consistent. If  $P_t = \{T\}$  then  $\nabla(P_t) = T$ , i.e.,  $\nabla(P_t)$  is consistent and complete. If  $P_t$  contains two distinct theories  $T_1$  and  $T_2$  then  $T_1 \cup T_2 \subseteq \nabla(P_t)$  implying that  $\nabla(P_t)$  is inconsistent since there are sentences  $\psi$  such that  $\psi \in T_1$  and  $\neg\psi \in T_2$ .

(3) If  $\varphi \in \Delta(P_t)$  then  $\varphi \triangleright_{\text{tt}} P_t$  and  $\varphi \in T$  for any  $T \in P_t$  implying  $\Delta(P_t) \subseteq \bigcap P_t$ . Conversely, if  $\varphi \in \bigcap P_t$  then  $\varphi \in T$  for any  $T \in P_t$  implying  $\varphi \triangleright_{\text{tt}} P_t$  and therefore  $\varphi \in \Delta(P_t)$ .

(4) Since  $\Delta(\emptyset) = \text{Sent}(\Sigma)$  it is inconsistent. If  $P_t \neq \emptyset$  then  $\Delta(P_t) = \bigcap P_t$  by (3) implying that  $\Delta(P_t)$  is a consistent theory as an intersection of complete theories. If  $\Delta(P_t)$  is complete then both  $P_t$  is nonempty and does not contain two distinct theories, i.e.,  $P_t$  is a singleton. Conversely, if  $P_t = \{T\}$  then  $\Delta(P_t) = T$  which is a complete theory.

(5) If  $P_t \neq \emptyset$  then by (1) and (3) we have  $\nabla(P_t) = \bigcup P_t \supseteq \bigcap P_t = \Delta(P_t)$ . If  $P_t = \{T\}$  then  $\Delta(P_t) = T = \nabla(P_t)$ . If  $T_1, T_2 \in P_t$  for some  $T_1 \neq T_2$  then  $\nabla(P_t) = \bigcup P_t \supseteq T_1 \cup T_2 \supsetneq T_1 \cap T_2 \supseteq \bigcap P_t = \Delta(P_t)$ .  $\square$

Theorems 1.11, 2.7, 2.8 and Proposition 2.9 immediately imply the following corollary on the separability of properties with respect to the relations  $\triangleright_{\text{pt}}$  and  $\triangleright_{\text{tt}}$ .

**Corollary 2.10.** (1) *For any properties  $P_1, P_2 \subseteq \mathcal{T}_\Sigma$  the following conditions hold:*

(1) *there exists  $\varphi \in \text{Sent}(\Sigma)$  such that  $\varphi \triangleright_{\text{pt}} P_1$  and  $\neg\varphi \triangleright_{\text{pt}} P_2$  iff  $P_1$  and  $P_2$  are nonempty and  $|P_1 \cup P_2| \geq 2$ ; in particular, there exists  $\varphi \in \text{Sent}(\Sigma)$  such that  $\varphi \triangleright_{\text{pt}} P_1$  and  $\neg\varphi \triangleright_{\text{pt}} P_1$  iff  $|P_1| \geq 2$ ;*

(2) *there exists  $\varphi \in \text{Sent}(\Sigma)$  such that  $\varphi \triangleright_{\text{tt}} P_1$  and  $\neg\varphi \triangleright_{\text{tt}} P_2$  iff  $\text{Cl}_E(P_1) \cap \text{Cl}_E(P_2) = \emptyset$ .*

**Corollary 2.11.** *For any nonempty property  $P_t \subseteq \mathcal{T}_\Sigma$  the following conditions hold:*

(1) *the set  $\Delta(P_t)$  forms a filter  $\Delta(P_t)/\equiv$  on  $\{\equiv(\varphi) \mid \varphi \in \text{Sent}(\Sigma)\}$  with respect to  $\vdash$ ;*

(2) *the filter  $\Delta(P_t)/\equiv$  is principal iff  $\bigcap P_t$  is forced by some its sentence, i.e.,  $\bigcap P_t$  is a finitely axiomatizable theory, which is incomplete for  $|P_t| \geq 2$ ;*

(3) the filter  $\Delta(P_t)/\equiv$  is an ultrafilter iff  $P_t$  is a singleton.

Proof. (1) holds by Monotony and Proposition 2.3, (2).

(2) immediately follows by Proposition 2.9, (3).

(3) is satisfied in view of Proposition 2.9, (4).  $\square$

### 3 Ranks of sentences and spectra with respect to properties

In this section we introduce a measure of complexity for sentences satisfying a property using the RS-rank for families of theories [2, 3, 4].

**Definition.** For a sentence  $\varphi \in \text{Sent}(\Sigma)$  and a property  $P = P_t \subseteq \mathcal{T}_\Sigma$  we put  $\text{RS}_P(\varphi) = \text{RS}(P_\varphi)$ , and  $\text{ds}_P(\varphi) = \text{ds}(P_\varphi)$  if  $\text{ds}(P_\varphi)$  is defined.

If  $P = \mathcal{T}_\Sigma$  then we omit  $P$  and write  $\text{RS}(\varphi)$ ,  $\text{ds}(\varphi)$  instead of  $\text{RS}_P(\varphi)$  and  $\text{ds}_P(\varphi)$ , respectively.

Clearly, if  $P \subseteq P' \subseteq \mathcal{T}_\Sigma$  and  $\varphi \in \text{Sent}(\Sigma)$  then  $\text{RS}_P(\varphi) \leq \text{RS}_{P'}(\varphi)$ , and if  $\text{RS}_P(\varphi) = \text{RS}_{P'}(\varphi) \in \text{Ord}$  then  $\text{ds}_P(\varphi) \leq \text{ds}_{P'}(\varphi)$ . Besides, we have:

**Proposition 3.1.** (1)  $\varphi \triangleright_{\text{tt}} P$  iff  $\text{RS}_P(\neg\varphi) = -1$ .

(2)  $\varphi \triangleright_{\text{pt}} P$  iff  $\text{RS}_P(\varphi) \geq 0$ .

**Definition.** For a sentence  $\varphi \in \text{Sent}(\Sigma)$  and a property  $P \subseteq \mathcal{T}_\Sigma$  we say that  $\varphi$  is *P-totally transcendental* if  $\text{RS}_P(\varphi)$  is an ordinal. If  $P = \mathcal{T}_\Sigma$  then a *P-totally transcendental* sentence is said to be *totally transcendental*.

A sentence  $\varphi$  is *co-(P)-totally transcendental* if  $\neg\varphi$  is *P-totally transcendental*.

We omit  $P$  and say about totally transcendental and co-totally transcendental sentences if  $P = \mathcal{T}_\Sigma$ .

By the definition each sentence  $\varphi \in \text{Sent}(\Sigma)$  obtains the characteristics  $\text{RS}_P(\varphi)$  and  $\text{RS}_P(\neg\varphi)$  saying that  $\varphi$  is (co)-rich enough with respect to the property  $P$ , The characteristics  $\text{ds}_P(\varphi)$  and  $\text{ds}_P(\neg\varphi)$ , if they are defined, give an additional information on “*P*-richness” of  $\varphi$ .

For instance, if  $\text{RS}_P(\varphi) = 0$  and  $\text{ds}_P(\varphi) = n$  then  $\varphi$  is *P-finite* satisfying exactly  $n$  theories in  $P$ . Respectively, if  $\text{RS}_P(\neg\varphi) = 0$  and  $\text{ds}_P(\neg\varphi) = n$  then  $\varphi$  is *P-cofinite*, i.e., it does not satisfy exactly  $n$  theories in  $P$ .

Clearly,  $\varphi$  is both *P-finite* and *P-cofinite* iff  $P$  is nonempty and is finite.

**Theorem 3.2.** For a language  $\Sigma$  there is a totally transcendental sentence  $\varphi \in \text{Sent}(\Sigma)$  iff  $\Sigma$  has finitely many predicate symbols.

Proof. If  $\Sigma$  has finitely many predicate symbols we choose a sentence  $\varphi$  assigning that a universe is a singleton. Since functional symbols have unique interpretations and there are finitely many possibilities for (non)empty language predicate we obtain  $\text{RS}(\varphi) = 0$  that is  $\varphi$  is totally transcendental.

Conversely, if  $\Sigma$  has infinitely many predicate symbols then each consistent sentence  $\varphi$  obtains a 2-tree with respect to (non)empty predicates  $Q \in \Sigma \setminus \Sigma(\varphi)$ . This 2-tree witnesses that  $\text{RS}(\varphi) = \infty$ , i.e.,  $\varphi$  is not totally transcendental.  $\square$

**Remark 3.3.** If  $\Sigma$  is finite then for the proof of Theorem 3.2 it suffices to choose a sentence  $\varphi$  assigning that a universe is finite, since there are finitely many many possibilities, up to isomorphism, for interpretations of language symbols implying  $\text{RS}(\varphi) = 0$ .  $\square$

The following definition introduces values for richness of a sentence with respect to a property.

**Definition.** For a language  $\Sigma$ , a property  $P \subseteq \mathcal{T}_\Sigma$ , an ordinal  $\alpha$  and a natural number  $n \geq 1$ , a sentence  $\varphi \in \text{Sent}(\Sigma)$  is called  $(P, \alpha, n)$ -(co-)rich if  $\text{RS}_P(\varphi) = \alpha$  and  $\text{ds}_P(\varphi) = n$  (respectively,  $\text{RS}_P(\neg\varphi) = \alpha$  and  $\text{ds}_P(\neg\varphi) = n$ ).

A sentence  $\varphi \in \text{Sent}(\Sigma)$  is called  $(P, \infty)$ -(co-)rich if  $\text{RS}_P(\varphi) = \infty$  (respectively,  $\text{RS}_P(\neg\varphi) = \infty$ ).

If  $P = \mathcal{T}_\Sigma$  we write that  $\varphi$  is  $(\alpha, n)$ -(co-)rich instead of  $(P, \alpha, n)$ -(co-)rich, and  $\infty$ -(co-)rich instead of  $(P, \infty)$ -(co-)rich.

If for a property  $P$  there is a  $(P, *)$ -(co-)rich sentence  $\varphi$ , we say that  $P$  has a  $(P, *)$ -(co-)rich sentence, where  $*$  =  $\alpha, n$  or  $\alpha = \infty$ .

By the definition if a sentence  $\varphi$  is  $(P, \alpha, n)$ -rich then  $\text{RS}(P_\varphi) = \alpha$ ,  $\text{ds}(P_\varphi) = n$ .

**Theorem 3.4.** (1) *If a property  $P \subseteq \mathcal{T}_\Sigma$  has a  $(P, \alpha, m)$ -rich sentence  $\varphi$  which is  $(P, \beta, n)$ -co-rich then  $\text{RS}(P) = \max\{\alpha, \beta\}$ ,  $\text{ds}(P) = m$  for  $\alpha > \beta$ ,  $\text{ds}(P) = n$  for  $\alpha < \beta$ , and  $\text{ds}(P) = m + n$  for  $\alpha = \beta$ .*

(2) *If for a property  $P \subseteq \mathcal{T}_\Sigma$ ,  $\text{RS}(P) = \alpha$  and  $\text{ds}(P) = n$ , then for each sentence  $\varphi \in \text{Sent}(\Sigma)$  the following assertions hold:*

(i)  $\text{RS}_P(\varphi) \leq \alpha$ ,

(ii) *if  $\text{RS}_P(\varphi) = \alpha$  then  $\varphi$  is  $(P, \alpha, m)$ -rich for some  $m \leq n$ , and for  $m = n$  either  $\varphi \triangleright_{\text{tt}} P$  or  $\varphi$  is  $(P, \beta, k)$ -co-rich for some  $\beta < \alpha$  and  $k \in \omega$ , and if  $m < n$  then  $\varphi$  is  $(P, \alpha, n - m)$ -co-rich.*

*Proof.* (1) For the sentence  $\varphi$  we have  $\text{RS}_P(\varphi) = \alpha$ ,  $\text{ds}_P(\varphi) = m$ ,  $\text{RS}_P(\neg\varphi) = \beta$ ,  $\text{ds}_P(\neg\varphi) = n$ . It means that  $P$  is divided into two disjoint parts  $P_\varphi$  and  $P_{\neg\varphi}$  with given characteristics  $\text{RS}(P_\varphi) = \alpha$ ,  $\text{ds}(P_\varphi) = m$ ,  $\text{RS}(P_{\neg\varphi}) = \beta$ ,  $\text{ds}(P_{\neg\varphi}) = n$ .

If  $\text{RS}(P) = 0$  then  $|P| = \text{ds}(P)$ ,  $\text{RS}_P(\varphi) = \text{RS}_P(\neg\varphi) = 0$ ,  $\text{ds}(P) = |P_\varphi| + |P_{\neg\varphi}| = m + n$ .

If  $\text{RS}(P) > 0$  then a tree witnessing the value  $\text{RS}(P) = \gamma$  can be transformed step-by-step using theories either in  $P_\varphi$  or in  $P_{\neg\varphi}$ : in each step witnessing  $\text{RS}(P) = \gamma$  there are infinitely many branches of previous values related to  $P_\varphi$  or to  $P_{\neg\varphi}$ .

In the first case, related to  $P_\varphi$ , we have  $\gamma = \alpha$  and in the second case, related to  $P_{\neg\varphi}$ ,  $\gamma = \beta$ . If  $\alpha > \beta$  a tree for  $P_\varphi$  witnesses  $l = m$ . If  $\alpha < \beta$  a tree for  $P_{\neg\varphi}$  witnesses  $l = n$ . If  $\alpha = \beta$  then both trees for  $P_\varphi$  and for  $P_{\neg\varphi}$  witness  $\gamma = \alpha$  and  $l = m + n$ , since there are exactly  $l + m$   $s$ -definable subsets of  $P$  having the rank  $\gamma$  and the degree 1.

(2) We have (i) by the monotony of rank (if  $P_1 \subseteq P_2$  then  $\text{RS}(P_1) \leq \text{RS}(P_2)$ ) and the inclusion  $P_\varphi \subseteq P$ . (ii) holds by the monotony of degree for a fixed rank (if  $P_1 \subseteq P_2$  and  $\text{RS}(P_1) = \text{RS}(P_2) \in \text{Ord}$  then  $\text{ds}(P_1) \leq \text{ds}(P_2)$ ) and the inclusion  $P_\varphi \subseteq P$ . Besides if  $\text{ds}(P_\varphi) = m = n = \text{ds}(P)$  then  $P$  can not have a tree in  $P_{\neg\varphi} = P \setminus P_\varphi$  witnessing  $\text{RS}(P_{\neg\varphi}) = \alpha$  since otherwise  $\text{ds}(P)$  should be more than  $n$ . Therefore either  $\neg\varphi$  is  $P$ -inconsistent, i.e.,  $\varphi \triangleright_{\text{tt}} P$ , or  $\varphi$  is  $(P, \beta, k)$ -co-rich for some  $\beta < \alpha$  and  $k \in \omega$ . If  $m < n$  then  $\varphi$  is  $(P, \alpha, n - m)$ -co-rich in view of (1).  $\square$

By Theorem 3.4 for any  $e$ -totally transcendental property  $P$  and any  $\alpha \leq \text{RS}(P)$  there are  $s$ -definable subfamilies  $P_\varphi$  with  $\text{RS}(P_\varphi) = \alpha$ . Similarly all values  $m \leq \text{ds}(P)$  are also realized by appropriate  $s$ -definable subfamilies.

Thus the *spectrum*  $\text{Sp}_{\text{Rd}}(P)$  for the pairs  $(\text{RS}_P(\varphi), \text{ds}_P(\varphi))$  with nonempty  $P_\varphi$  forms the set

$$\{(\text{RS}(P), m) \mid 1 \leq m \leq \text{ds}(P)\} \cup \{(\alpha, m) \mid \alpha < \text{RS}(P), m \in \omega \setminus \{0\}\}, \quad (1)$$

which is an initial segment  $O[(\beta, n)]$  consisting of all pairs  $(\alpha, m) \in \text{Ord} \times (\omega \setminus 0)$  with  $\alpha \leq \beta$  and  $m \leq n$  for  $\alpha = \beta$ ,  $\text{RS}(P) = \beta$ ,  $\text{ds}(P) = n$ .

**Remark 3.5.** If  $\text{RS}(P) = \infty$  then  $s$ -definable subfamilies  $P_\varphi$  can have only values  $\text{RS}(P_\varphi) = \infty$  or both the value  $\text{RS}(P) = \infty$  and pairs forming some initial segment  $O[(\beta, n)]$ .

Indeed, let  $P$  be a family of theories in a language  $\Sigma$  of independent 0-ary predicates  $Q_l$ ,  $l \in \lambda$ ,  $\lambda \geq \omega$ , such that each sentence  $Q_{i_1}^{\delta_1} \wedge \dots \wedge Q_{i_k}^{\delta_k}$ ,  $i_1 < \dots < i_k < \lambda$ ,  $\delta_1, \dots, \delta_k \in \{0, 1\}$ ,  $k \in \omega$ , is  $P$ -consistent. Each  $P$ -consistent sentence  $\varphi \in \text{Sent}(\Sigma)$  is divided into 2-tree witnessing that  $\text{RS}_P(\varphi) = \infty$ . In such a case we say that the spectrum  $\text{Sp}_{\text{Rd}}(P)$  equals  $\{\infty\}$ .

The family  $P$  above can be extended by a family  $P'$  with dependent predicates  $Q_l$  producing given RS-rank and ds-degree for a subfamily with, say,  $Q_0 \leftrightarrow Q_1$ . Therefore the arguments for Theorem 3.4 produce an initial segment  $O[(\beta, n)]$  for the spectrum  $\text{Sp}_{\text{Rd}}(P')$  of  $s$ -definable family  $P'$ . Thus,  $\text{Sp}_{\text{Rd}}(P \cup P') = O[(\beta, n)] \cup \{\infty\}$ .

Since each nonempty  $s$ -definable subfamily has a spectrum of the form  $O[(\beta, n)]$ , or  $\{\infty\}$ , or  $O[(\beta, n)] \cup \{\infty\}$ , initial segments are well-ordered with respect to the relation  $\subseteq$ , and the ordinal  $\text{RS}(P)$ -ranks are bounded by  $|\mathcal{T}_\Sigma| \leq 2^{\max\{|\Sigma|, \omega\}}$ , all values  $\text{Sp}_{\text{Rd}}(P)$ , for nonempty properties  $P \subseteq \mathcal{T}_\Sigma$ , are exhausted by these three possibilities, and we obtain the following:

**Theorem 3.6.** *For any nonempty property  $P \subseteq \mathcal{T}_\Sigma$  one of the following possibilities holds for some  $\beta \in \text{Ord}$  and  $n \in \omega \setminus \{0\}$ :*

- (1)  $\text{Sp}_{\text{Rd}}(P) = O[(\beta, n)]$ ,
- (2)  $\text{Sp}_{\text{Rd}}(P) = \{\infty\}$ ,
- (3)  $\text{Sp}_{\text{Rd}}(P) = O[(\beta, n)] \cup \{\infty\}$ .

*All possibilities above are realized by appropriate languages  $\Sigma$  and properties  $P \subseteq \mathcal{T}_\Sigma$ .*

**Remark 3.7.** Any value  $\text{Sp}_{\text{Rd}}(P)$  can be naturally extended till  $\overline{\text{Sp}}_{\text{Rd}}(P) = \text{Sp}_{\text{Rd}}(P) \cup \{-1\}$  corresponding to the value  $\text{RS}(\emptyset) = -1$  of the empty subfamily of  $\mathcal{T}_\Sigma$ . It is also natural to put  $\overline{\text{Sp}}_{\text{Rd}}(P) = \{-1\}$  for empty  $P \subset \mathcal{T}_\Sigma$ . In view of Theorem 3.6 we have the following description of extended spectra  $\overline{\text{Sp}}_{\text{Rd}}(P)$ :

- i)  $\overline{\text{Sp}}_{\text{Rd}}(P) = \{-1\}$ ,
- ii)  $\overline{\text{Sp}}_{\text{Rd}}(P) = O[(\beta, n)] \cup \{-1\}$ ,
- iii)  $\overline{\text{Sp}}_{\text{Rd}}(P) = \{-1, \infty\}$ ,
- iv)  $\overline{\text{Sp}}_{\text{Rd}}(P) = O[(\beta, n)] \cup \{-1, \infty\}$ .

**Theorem 3.8** [4]. *Let  $\mathcal{T}$  be a family of a countable language  $\Sigma$  and with  $\text{RS}(\mathcal{T}) = \infty$ ,  $\alpha$  be a countable ordinal,  $n \in \omega \setminus \{0\}$ . Then there is a  $d_\infty$ -definable subfamily  $\mathcal{T}^* \subset \mathcal{T}$  such that  $\text{RS}(\mathcal{T}^*) = \alpha$  and  $\text{ds}(\mathcal{T}^*) = n$ .*

Theorems 3.6 and 3.8 immediately imply:

**Corollary 3.9.** *Let  $\mathcal{T}$  be a family of a countable language  $\Sigma$  and with  $\text{RS}(\mathcal{T}) = \infty$ ,  $\alpha$  be a countable ordinal,  $n \in \omega \setminus \{0\}$ . Then there is a  $d_\infty$ -definable property  $P \subset \mathcal{T}$  such that  $\text{Sp}_{\text{Rd}}(P) = O[(\alpha, n)]$ .*

## 4 Spectra for cardinalities of definable subproperties

In this section we study some refinements of the relation  $\triangleright_{\text{pt}}$ .

For a cardinality  $\lambda \geq 1$ , a sentence  $\varphi \in \text{Sent}(\Sigma)$  and a property  $P \subseteq \mathcal{T}_\Sigma$  we write  $\varphi \triangleright_{\text{pt}}^\lambda P$  if  $\varphi$  satisfies exactly  $\lambda$  theories in  $P$ , i.e.,  $|P_\varphi| = \lambda$ .

By the definition if  $P \neq \emptyset$  and  $\varphi \triangleright_{\text{tt}} P$  then  $\varphi \triangleright_{\text{pt}}^{|P|} P$ , and conversely  $\varphi \triangleright_{\text{pt}}^{|P|} P$  implies  $\varphi \triangleright_{\text{tt}} P$  for finite  $P$ . For infinite  $P$  the converse implication can fail. Moreover, since infinite sets can be divided into two parts of same cardinality, one can easily introduce an expansion  $P'$  of  $P$  by a 0-ary predicate  $Q$  such that  $Q \triangleright_{\text{pt}}^{|P'|} P'$  and  $\neg Q \triangleright_{\text{pt}}^{|P'|} P'$ , implying that  $Q \not\triangleright_{\text{tt}} P'$ .

For a property  $P$  we denote by  $\text{Sp}_{\text{pt}}(P)$  the set  $\{\lambda \mid \varphi \triangleright_{\text{pt}}^\lambda P \text{ for some sentence } \varphi\}$ . This set is called the *pt-spectrum* of  $P$ .

By the definition  $|P| \in \text{Sp}_{\text{pt}}(P)$  for any nonempty  $P$ , and  $\lambda \leq |P|$  for any  $\lambda \in \text{Sp}_{\text{pt}}(P)$ , i.e.,  $\text{supSp}_{\text{pt}}(P) = ||P| + 1|$ .

A natural question arises on a description of pt-spectra.

This question is easily answered for finite  $P$ , since in such a case all subsets of theories are separated as  $s$ -definable singletons from their complements, and we obtain the following:

**Proposition 4.1.** *For any finite property  $P \subseteq \mathcal{T}_\Sigma$ ,  $\text{Sp}_{\text{pt}}(P) = (|P| + 1) \setminus \{0\}$ .*

The following assertion generalizes Proposition 4.1 in terms of isolated points due Theorem 1.12:

**Proposition 4.2.** *If  $P$  has exactly  $n \in \omega \setminus \{0\}$  isolated points then  $\text{Sp}_{\text{pt}}(P) \cap \omega = (n + 1) \setminus \{0\}$ .*

Proof. Let  $T_1, \dots, T_n$  are isolated points in  $P$ . If  $k \in \text{Sp}_{\text{pt}}(P) \cap \omega$ , i.e., there is a sentence  $\varphi$  with  $\varphi \triangleright_{\text{pt}}^k P$  then  $P_\varphi$  consists of isolated points  $T_{i_1}, \dots, T_{i_k}$  since elements of the finite set  $P_\varphi$  are separated as  $s$ -definable singletons from their complements in  $P_\varphi$ . Then  $k \in (n + 1) \setminus \{0\}$ . Conversely, if  $k \in \omega \setminus \text{Sp}_{\text{pt}}(P)$  then  $k > n$  since each  $k \leq n$  equals  $|P_\varphi|$  for some sentence  $\varphi$ , implying  $k \in \text{Sp}_{\text{pt}}(P)$ .  $\square$

**Proposition 4.3.** *For any nonempty property  $P \subseteq \mathcal{T}_\Sigma$  either  $\text{Sp}_{\text{pt}}(P) \cap \omega$  equals an initial segment  $(n + 1) \setminus \{0\}$  for some  $n \in \omega \setminus \{0\}$  or  $\text{Sp}_{\text{pt}}(P) \cap \omega = \omega \setminus \{0\}$ .*

Proof. If there are sentences  $\varphi$  with finite  $P_\varphi$  then either there is  $P_\varphi$  with the greatest finite cardinality, implying  $\text{Sp}_{\text{pt}}(P) \cap \omega = (n + 1) \setminus \{0\}$  by the arguments of Proposition 4.2, or the finite cardinalities  $|P_\varphi|$  are unbounded that means  $\text{Sp}_{\text{pt}}(P) \cap \omega = \omega \setminus \{0\}$ .  $\square$

**Proposition 4.4.** *For any infinite property  $P \subseteq \mathcal{T}_\Sigma$  there is a nonempty set  $Y \subseteq |P|$  of infinite cardinalities such that either there is  $n \in \omega \setminus \{0\}$  with  $\text{Sp}_{\text{pt}}(P) = Y \cup (n + 1) \setminus \{0\}$ , or  $\text{Sp}_{\text{pt}}(P) = Y \cup \omega \setminus \{0\}$ , or  $\text{Sp}_{\text{pt}}(P) = Y$ . All values  $Y \cup (n + 1) \setminus \{0\}$ ,  $Y \cup \omega \setminus \{0\}$  and  $Y$ , for a nonempty set  $Y$  of infinite cardinalities and  $n \in \omega \setminus \{0\}$ , are realized as  $\text{Sp}_{\text{pt}}(P)$  for appropriate properties  $P$ .*

Proof. Since  $\text{supSp}_{\text{pt}}(P) \leq |P|$  and Proposition 4.3 describes all possibilities for  $\text{Sp}_{\text{pt}}(P) \cap \omega$ , it suffices, for a nonempty set  $Y$  of infinite cardinalities and  $n \in \omega \setminus \{0\}$ , to find a property  $P_1$  with  $\text{Sp}_{\text{pt}}(P_1) = Y \cup (n + 1) \setminus \{0\}$ , a property  $P_2$  with  $\text{Sp}_{\text{pt}}(P_2) = Y \cup \omega \setminus \{0\}$ , and a property  $P_3$  with  $\text{Sp}_{\text{pt}}(P_3) = Y$ . For the property  $P_1$  one can take a finite  $n$ -element family  $Y_n$ , expanded by a 0-ary predicate  $Q_0$  marking all theories in  $Y_n$ , and extend  $Y_n$  by families  $Y_\lambda$ , for each  $\lambda \in Y$ , of  $\lambda$  theories of  $\lambda$  independent 0-ary predicates  $Q_i^\lambda$  expanded by a 0-ary predicate  $Q_\lambda$  marking all theories in  $Y_\lambda$ . Any  $P_1$ -consistent sentence  $\varphi$  satisfies either  $k \leq n$  theories in  $Y_n$  or  $\lambda$  many theories in  $Z_\lambda$  and possibly  $\mu$  many theories in  $Y_\mu$  for finitely many  $\mu < \lambda$ . It means that the cardinalities  $|(P_1)_\varphi|$  witness the equality  $\text{Sp}_{\text{pt}}(P_1) = Y \cup (n + 1) \setminus \{0\}$ .

For the property  $P_2$  we repeat the process for  $P_1$  replacing the part  $Y_n$  by an  $e$ -minimal family  $Y_0^\lambda$  consisting of some  $\lambda \in Y$  theories all of which are marked by new 0-ary predicate  $Q_0$ . Realizing this process we obtain that  $s$ -definable sets are finite, cofinite, or consists of  $\lambda$  theories for  $\lambda \in Y$ . Thus,  $\text{Sp}_{\text{pt}}(P_1) = Y \cup \omega \setminus \{0\}$ .

For the property  $P_3$  we repeat the process for  $P_1$  without the part  $Y_n$  obtaining  $\text{Sp}_{\text{pt}}(P_1) = Y$ .  $\square$

Summarizing Propositions 4.1–4.4 we obtain the following theorem describing pt-spectra of properties.

**Theorem 4.5.** *For any nonempty property  $P \subseteq \mathcal{T}_\Sigma$  one of the following conditions holds:*

- (1)  $\text{Sp}_{\text{pt}}(P) = (n+1) \setminus \{0\}$  for some  $n \in \omega \setminus \{0\}$ ; it is satisfied iff  $P$  is finite with  $|P| = n$ ;
- (2)  $\text{Sp}_{\text{pt}}(P) = Y \cup (n+1) \setminus \{0\}$  for some nonempty set  $Y \subseteq |P|$  of infinite cardinalities and  $n \in \omega \setminus \{0\}$ ;

(3)  $\text{Sp}_{\text{pt}}(P) = Y \cup \omega \setminus \{0\}$  for some nonempty set  $Y \subseteq |P|$  of infinite cardinalities;

(4)  $\text{Sp}_{\text{pt}}(P) = Y$  for some nonempty set  $Y \subseteq |P|$  of infinite cardinalities.

All values  $(n+1) \setminus \{0\}$ ,  $Y \cup (n+1) \setminus \{0\}$ ,  $Y \cup \omega \setminus \{0\}$ , and  $Y$ , for a nonempty set  $Y$  of infinite cardinalities and  $n \in \omega \setminus \{0\}$ , are realized as  $\text{Sp}_{\text{pt}}(P)$  for an appropriate property  $P$ .

The following assertion shows that  $Y$  in Theorem 4.5 is finite for a property  $P$  with  $\text{RS}(P) = 1$ .

**Proposition 4.6.** *If  $\text{RS}(P) = 1$  then  $\text{Sp}_{\text{pt}}(P) = Y \cup \omega \setminus \{0\}$  for some finite nonempty set  $Y \subseteq |P|$  of infinite cardinalities with  $|Y| \leq \text{ds}(P)$ .*

*Proof.* Since  $\text{RS}(P) = 1$ ,  $P$  is divided into  $k = \text{ds}(P)$  infinite  $s$ -definable  $e$ -minimal parts  $P_1, \dots, P_k$ . Each part  $P_i$  has only finite and cofinite  $s$ -definable subset producing  $\text{Sp}_{\text{pt}}(P_i) = \{|P_i|\} \cup \omega \setminus \{0\}$ . Since each  $s$ -definable subset of  $P$  is a Boolean combination of  $s$ -definable subsets of  $P_i$  and  $|Z_i \cup Z_j| = \max\{|Z_i|, |Z_j|\}$  for infinite  $s$ -definable  $Z_i \subseteq P_i$ ,  $Z_j \subseteq P_j$ ,  $i, j \leq k$ , we obtain  $\text{Sp}_{\text{pt}}(P) = \{|P_1|, \dots, |P_k|\} \cup \omega \setminus \{0\}$ .  $\square$

**Remark 4.7.** Describing possibilities for pt-spectra  $\text{Sp}_{\text{pt}}(P)$  we admit that properties  $P$  may be not  $E$ -closed. If we assume that  $P$  is infinite and  $E$ -closed then we have two cases: either  $P$  is  $e$ -totally transcendental with the least generating set of a cardinality  $\mu_1 \leq \max\{|\Sigma|, \omega\}$  and with  $\mu_2 \leq \max\{|\Sigma|, \omega\}$  accumulation points, or  $\text{RS}(P) = \infty$  with  $|P| \geq 2^\omega$  by Theorem 1.2. In the first case values for  $\lambda \in \text{Sp}_{\text{pt}}(P)$  are exhausted by all cardinalities in  $\omega \setminus \{0\}$  and by some infinite cardinalities  $\leq \max\{|\Sigma|, \omega\}$ . In particular, for countable  $\Sigma$ , since  $\text{RS}(P)$  is a countable ordinal, we have  $\text{Sp}_{\text{pt}}(P) = (\omega + 1) \setminus \{0\}$ . In the second case values for  $\lambda \in \text{Sp}_{\text{pt}}(P)$  are exhausted by cardinalities in  $\omega \setminus \{0\}$  or by cardinalities of some its initial segment, depending on existence of the least generating set for  $P$ , and by some infinite cardinalities  $\leq \max\{|\Sigma|, \omega\}$  and cardinalities  $\geq 2^\omega$ . In particular, for countable  $\Sigma$ ,  $\text{Sp}_{\text{pt}}(P)$  includes  $2^\omega$ , and possibly  $\omega$  depending on existence of infinite totally transcendental definable part. Thus, in Theorem 4.5, for  $E$ -closed  $P$  some cases are not realized:  $\{\omega\} \cup (n+1) \setminus \{0\}$ ,  $\{\omega\}$ ,  $\{\omega_1\}$ ,  $\{\omega, \omega_1\}$ , etc.

By Remark 4.7 using Theorem 1.2 we have the following:

**Theorem 4.8.** *For any nonempty  $E$ -closed property  $P \subseteq \mathcal{T}_\Sigma$  with at most countable language  $\Sigma$  one of the following possibilities holds:*

- (1)  $\text{Sp}_{\text{pt}}(P) = (n+1) \setminus \{0\}$  for some  $n \in \omega \setminus \{0\}$ , if  $P$  is finite with  $|P| = n$ ;
- (2)  $\text{Sp}_{\text{pt}}(P) = \{2^\omega\} \cup (n+1) \setminus \{0\}$  for some  $n \in \omega$ , if  $P$  is infinite and has  $n$  isolated points;

(3)  $\text{Sp}_{\text{pt}}(P) = (\omega + 1) \setminus \{0\}$ , if  $P$  is infinite and totally transcendental;

(4)  $\text{Sp}_{\text{pt}}(P) = \{\omega, 2^\omega\} \cup \omega \setminus \{0\}$ , if  $P$  has an infinite totally transcendental definable subfamily but  $P$  itself is not totally transcendental;

(5)  $\text{Sp}_{\text{pt}}(P) = \{2^\omega\} \cup \omega \setminus \{0\}$ , if  $P$  has infinitely many isolated points but does not have infinite totally transcendental definable subfamilies.

**Remark 4.9.** Possibilities in Theorem 4.7 give low bounds for correspondent cases in uncountable languages.

## 5 $P$ -generic sentences and $P$ -generic theories

**Definition.** (Cf. [8, 9, 10]) For a property  $P \subseteq \mathcal{T}_\Sigma$  a sentence  $\varphi \in \text{Sent}(\Sigma)$  is called  $P$ -generic if  $\text{RS}_P(\varphi) = \text{RS}(P)$ , and  $\text{ds}_P(\varphi) = \text{ds}(P)$  if  $\text{ds}(P)$  is defined.

If  $P = \mathcal{T}_\Sigma$  then we omit  $P$  and a  $P$ -generic sentence is called *generic*.

By the definition we have the following:

**Proposition 5.1.** *Any  $P$ -generic sentence  $\varphi$  is  $(P, \text{RS}(P), \text{ds}(P))$ -rich if  $\text{RS}(P)$  is an ordinal, and  $(P, \infty)$ -rich if  $\text{RS}(P) = \infty$ . And vice versa, each  $(P, \text{RS}(P), \text{ds}(P))$ -rich sentence, for an ordinal  $\text{RS}(P)$ , is  $P$ -generic, and each  $(P, \infty)$ -rich sentence, for  $\text{RS}(P) = \infty$ , is  $P$ -generic.*

**Proposition 5.2.** *If  $\varphi \triangleright_{\text{tt}} P$  then  $\varphi$  is  $P$ -generic.*

In view of Proposition 5.2 any property  $P \subseteq \mathcal{T}_\Sigma$  has  $P$ -generic sentences.

**Corollary 5.3.** *If a property  $P \subseteq \mathcal{T}_\Sigma$  is finite and  $\varphi \in \text{Sent}(\Sigma)$  then  $\varphi \triangleright_{\text{tt}} P$  iff  $\varphi$  is  $P$ -generic.*

*Proof.* By Proposition 5.2 it suffices to show that if  $\varphi$  is  $P$ -generic then  $\varphi \triangleright_{\text{tt}} P$ . If  $P = \emptyset$  then both  $\text{RS}(P) = \text{RS}_P(\varphi) = -1$  and  $\varphi \triangleright_{\text{tt}} P$ . If  $P$  consists of  $n \geq 1$  theories, then  $\text{RS}(P) = 0$ ,  $\text{ds}(P) = n$ . Assuming that  $\varphi$  is  $P$ -generic we have  $\text{RS}(P_\varphi) = 0$ ,  $\text{ds}(P_\varphi) = n$  implying that  $\varphi$  belongs to all  $n$  theories in  $P$  and  $\varphi \triangleright_{\text{tt}} P$ .  $\square$

**Remark 5.4.** In view of Corollary 5.3 the converse implication for Proposition 5.2 holds iff  $P$  is finite. Indeed, if  $P = \mathcal{T}_\Sigma$  for a countable language  $\Sigma$  with  $\text{RS}(P) = \infty$ , which is characterized in Theorem 1.2, then we can construct a 2-tree of sentences  $\varphi_\delta$ ,  $\delta \in < \omega^2$  witnessing this value of  $\text{RS}(P)$ . It means that  $P$  is divided into two disjoint definable parts  $P_{\varphi_0}$  and  $P_{\varphi_1}$  with  $\text{RS}(P_{\varphi_0}) = \text{RS}(P_{\varphi_1}) = \infty$ . Thus,  $\varphi_0$  and  $\varphi_1$  are generic whereas  $\varphi_0 \not\triangleright_{\text{tt}} P$  and  $\varphi_1 \not\triangleright_{\text{tt}} P$ . Moreover, this effect works both for an arbitrary property  $P$  with  $\text{RS}(P) = \infty$  and for an arbitrary property  $P$  with  $\text{RS}(P) \in \text{Ord}$  and  $\text{RS}(P) \geq 1$ . In the latter case we can remove a branch in the tree witnessing the values  $\text{RS}(P)$  and  $\text{ds}(P)$  just considering a sentence  $\varphi_1 \wedge \neg\psi$ , where  $\varphi_1$  is a tautology and  $P_\psi$  is nonempty with  $\text{RS}(P_\psi) < \text{RS}(P)$ . In such a case  $\varphi_1 \wedge \neg\psi$  is  $P$ -generic and  $(\varphi_1 \wedge \neg\psi) \not\triangleright_{\text{tt}} P$ .  $\square$

In view of Remark 5.4 we have the following:

**Proposition 5.5.** *For a property  $P \subseteq \mathcal{T}_\Sigma$  there is a  $P$ -generic sentence  $\varphi \in \text{Sent}(\Sigma)$  with minimal/least  $P_\varphi$  iff  $P$  is finite. If that  $\varphi$  exists then  $P_\varphi = P$ .*

By Proposition 5.5 for a property  $P \subseteq \mathcal{T}_P$  with  $\text{RS}(P) \geq 1$ ,  $P$ -generic sentences  $\varphi$  produce infinite descending chains of  $s$ -definable subfamilies  $P_\varphi$ .

**Proposition 5.6.** *If for a property  $P \subseteq \mathcal{T}_\Sigma$ ,  $\text{RS}(P) = \alpha \in \text{Ord}$ ,  $\text{ds}(P) = 1$ , then for any sentence  $\varphi \in \text{Sent}(\Sigma)$  either  $\text{RS}_P(\varphi) = \alpha$  and  $\text{ds}_P(\varphi) = 1$ , or  $\text{RS}_P(\neg\varphi) = \alpha$  and  $\text{ds}_P(\neg\varphi) = 1$ .*

Proof. By the conjecture for  $P$  and the monotony for pairs of values of RS and ds we have  $\text{RS}_P(\varphi) \leq \alpha$  and  $\text{RS}_P(\neg\varphi) \leq \alpha$  for any  $\varphi \in \text{Sent}(\Sigma)$ , and if the RS-rank equals  $\alpha$  then the ds-degree equals 1. We can not have  $\text{RS}_P(\varphi) < \alpha$  and  $\text{RS}_P(\neg\varphi) < \alpha$  for  $\varphi \in \text{Sent}(\Sigma)$  by Theorem 3.4. Thus  $\text{RS}_P(\varphi) = \alpha$  and  $\text{ds}_P(\varphi) = 1$ , or  $\text{RS}_P(\neg\varphi) = \alpha$  and  $\text{ds}_P(\neg\varphi) = 1$ . The latter conditions can not be satisfied simultaneously since otherwise  $\text{ds}(P) \geq 2$ .  $\square$

Since any property  $P \subseteq \mathcal{T}_\Sigma$  with  $\text{RS}(P) = \alpha \in \text{Ord}$  is represented as a disjoint union of  $\text{ds}(P)$   $s$ -definable subfamilies  $P_1, \dots, P_{\text{ds}(P)}$  with  $\text{RS} = \alpha$  and  $\text{ds} = 1$ , Proposition 5.6 immediately implies:

**Corollary 5.7.** *For any property  $P \subseteq \mathcal{T}_\Sigma$  with  $\text{RS}(P) = \alpha \in \text{Ord}$  and any sentence  $\varphi \in \text{Sent}(\Sigma)$  either  $\varphi$  is  $P$ -generic or  $\neg\varphi$  is  $P$ -generic, or, for  $\text{ds}(P) > 1$  with non- $P$ -generic  $\varphi$  and  $\neg\varphi$ ,  $\varphi$  is represented as a disjunction of  $k$   $(P, \alpha, 1)$ -rich sentences and  $\neg\varphi$  is represented as a disjunction of  $m$   $(P, \alpha, 1)$ -rich sentences such that  $k + m = \text{ds}(P)$ ,  $k > 0$ ,  $m > 0$ .*

**Remark 5.8.** By Proposition 5.6 for any property  $P \subseteq \mathcal{T}_\Sigma$  with  $\text{RS}(P) = \alpha \in \text{Ord}$  and  $\text{ds}(P) = 1$  there is unique ultrafilter  $U_P$  consisting of  $P$ -generic sentences. By Proposition 5.5 this ultrafilter is principal if and only if  $P$  is finite, i.e., in such a case it is a singleton. Anyway  $U_P$  produces a theory  $T \in \mathcal{T}_\Sigma$  consisting of  $P$ -generic sentences only. This theory  $T$  is called  *$P$ -generic*.

If  $P$  is infinite then  $T$  belongs to the  $E$ -closure  $\text{Cl}_E(P)$  [2, 6] of  $P$  as unique element of  $\alpha$ -th Cantor–Bendixson derivative of  $\text{Cl}_E(P)$ , i.e., an element of  $\text{Cl}_E(P)$  having Cantor–Bendixson rank  $\text{CB}(\text{Cl}_E(P)) = \alpha$  [2].

If  $\text{ds}(P) > 1$  we can divide  $P$  into  $\text{ds}(P)$   $s$ -definable parts  $P_i$  with  $\text{RS}(P_i) = \text{RS}(P)$  and  $\text{ds}(P_i) = 1$ , each of which has unique  $P_i$ -generic theory  $T_i$ . The set  $\{T_1, \dots, T_{\text{ds}(P)}\}$  is called the *set of  $P$ -generic theories*.

Thus we have the following:

**Proposition 5.9.** *Each  $e$ -totally transcendental property  $P$  has finitely many, exactly  $\text{ds}(P) \geq 1$ ,  $P$ -generic theories. These theories have Cantor–Bendixson rank  $\text{CB}(\text{Cl}_E(P)) = \alpha = \text{RS}(P)$ .*

Now we extend the results above on generic sentences and theories for properties  $P$  with  $\text{RS}(P) = \infty$ .

**Proposition 5.10.** *If for a property  $P \subseteq \mathcal{T}_\Sigma$ ,  $\text{RS}(P) = \infty$ , then for any sentence  $\varphi \in \text{Sent}(\Sigma)$  either  $\text{RS}_P(\varphi) = \infty$  or  $\text{RS}_P(\neg\varphi) = \infty$ .*

Proof. Assume that for a sentence  $\varphi \in \text{Sent}(\Sigma)$  we have  $\text{RS}_P(\varphi) < \infty$  and  $\text{RS}_P(\neg\varphi) < \infty$ . We can suppose that  $\varphi$  and  $\neg\varphi$  are both  $P$ -consistent. Then  $\varphi$  is  $(P, \alpha, m)$ -rich and  $(P, \beta, n)$ -co-rich for some  $\alpha, \beta \in \text{Ord}$ ,  $m, n \in \omega$ . Applying Theorem 3.4 we obtain  $\text{RS}(P) \in \text{Ord}$  contradicting  $\text{RS}(P) = \infty$ .  $\square$

Proposition 5.10 immediately implies:

**Corollary 5.11.** *For any property  $P \subseteq \mathcal{T}_\Sigma$  with  $\text{RS}(P) = \infty$  and any sentence  $\varphi \in \text{Sent}(\Sigma)$  either  $\varphi$  is  $P$ -generic or  $\neg\varphi$  is  $P$ -generic.*

**Remark 5.12.** By Proposition 5.10 for any property  $P \subseteq \mathcal{T}_\Sigma$  with  $\text{RS}(P) = \infty$  there is ultrafilters  $U_P$  consisting of  $P$ -generic sentences. Since the condition  $\text{RS}(P) = \infty$  implies the existence of 2-tree of  $\Sigma$ -sentences there are at least continuum many these ultrafilters, and by Theorem 1.2 there are exactly continuum many ones for the language  $\Sigma$  with  $|\Sigma| \leq \omega$ .

Each  $U_P$  produces a theory  $T \in \mathcal{T}_\Sigma$  consisting of  $P$ -generic sentences only. This theory  $T$  is called  $P$ -generic.

The  $P$ -generic theories form the perfect kernel with respect to Cantor–Bendixson derivatives of  $\text{Cl}_E(P)$ , i.e., the set of element of  $\text{Cl}_E(P)$  having Cantor–Bendixson rank  $\text{CB}(\text{Cl}_E(P)) = \infty$ .

Applying Theorem 3.6 and summarizing Remarks 5.8 and 5.12 we obtain the following:

**Theorem 5.13.** (1) *For any nonempty property  $P \subseteq \mathcal{T}_\Sigma$  there are  $\text{ds}(P)$   $P$ -generic theories if  $P$  is totally transcendental, and at least continuum many if  $P$  is not totally transcendental. In the latter case either all theories in  $P$  are  $P$ -generic if  $\text{Sp}_{\text{Rd}}(P) = \{\infty\}$ , or  $P$  has at least  $\beta \cdot \omega + n$  non- $P$ -generic theories if  $\text{Sp}_{\text{Rd}}(P) = O[(\beta, n)] \cup \{\infty\}$ .*

(2) *The CB-rank of each  $P$ -generic theory equals  $\text{RS}(P)$ .*

**Definition** [5]. For a property  $P \subseteq \mathcal{T}_\Sigma$  a sentence  $\varphi \in \text{Sent}(\Sigma)$  is called  $P$ -complete if  $\varphi$  isolates a unique theory  $T$  in  $P$ , i.e.,  $P_\varphi$  is a singleton. In such a case the theory  $T \in P_\varphi$  is called  $P$ -finitely axiomatizable (by the sentence  $\varphi$ ).

Since  $P$ -finitely axiomatizable theories are isolated points we have the following:

**Proposition 5.14.** *For any nonempty property  $P \subseteq \mathcal{T}_\Sigma$  a  $P$ -finitely axiomatizable theory  $T$  is  $P$ -generic iff  $P$  is finite.*

## 6 Conclusion

We studied links between formulas and properties, considered and described characteristics for properties with respect to satisfying formulas, their ranks and degrees. Possibilities of spectra for ranks, degrees, and cardinalities of definable properties are shown. Generic formulas and theories are introduced and characterized.

Possibilities for ranks and degrees for formulas and theories with respect to given properties are described. Their top values form generic sentences and theories which are also described and characterized.

There are many natural model-theoretic and other properties that can be studied and described in this context. In this case, the relations  $\triangleright_{\text{pt}}$  and  $\triangleright_{\text{tt}}$  are preserved under natural closures. At the same time the operator of  $E$ -closure does not preserve a series of natural model-theoretic properties. For instance, there are families of strongly minimal whose accumulation points have the strict order properties. It implies that families of  $\omega$ -stable, superstable and stable theories can be not  $E$ -closed in these classes. Natural questions arise on characteristics and characterizations of families in these classes  $\mathcal{T}$  whose  $E$ -closures are contained in  $\mathcal{T}$ .

## References

- [1] *Ershov Yu. L.* Mathematical logic / Yu. L. Ershov, E. A. Palyutin. — Moscow : FIZMATLIT, 2011. — 356 p. [in Russian]
- [2] *Sudoplatov S. V.* Ranks for families of theories and their spectra / S. V. Sudoplatov // arXiv:1901.08464v1 [math.LO]. — 2019. — 17 p.

- [3] *Markhabatov N. D.* Ranks for families of all theories of given languages / N. D. Markhabatov, S. V. Sudoplatov // Eurasian Mathematical Journal. — 2021 (to appear). arXiv:1901.09903v1 [math.LO]. — 2019. — 9 p.
- [4] *Markhabatov N. D.* Definable subfamilies of theories, related calculi and ranks / N. D. Markhabatov, S. V. Sudoplatov // Siberian Electronic Mathematical Reports. — 2020. — Vol. 17. — P. 700–714.
- [5] *Sudoplatov S. V.* Approximations of theories / S. V. Sudoplatov // Siberian Electronic Mathematical Reports. — 2020. — Vol. 17. — P. 715–725.
- [6] *Sudoplatov S. V.* Closures and generating sets related to combinations of structures / S. V. Sudoplatov // Bulletin of Irkutsk State University. Series “Mathematics”. — 2016. — Vol. 16. — P. 131–144.
- [7] *Sudoplatov S. V.* Hierarchy of families of theories and their rank characteristics / S. V. Sudoplatov // Bulletin of Irkutsk State University. Series “Mathematics”. — 2020. — Vol. 33. — P. 80–95.
- [8] *Poizat B.* Groupes Stables / B. Poizat. — Villeurbanne : Nur Al-Mantiq Wal-Ma’rifah, 1987.
- [9] *Truss J. K.* Generic Automorphisms of Homogeneous Structures / J. K. Truss // Proceedings of the London Mathematical Society. — 1992. — Vol. 65, No. 3. — P. 121–141.
- [10] *Tent K.* A Course in Model Theory (Lecture Notes in Logic. No. 40) / K. Tent, M. Ziegler. — Cambridge : Cambridge University Press, 2012. — 258 p.

**Sergey Vladimirovich Sudoplatov**, Doctor of Sciences (Physics and Mathematics), docent; Leading researcher, Sobolev Institute of Mathematics, Novosibirsk; Head of Chair, Novosibirsk State Technical University; Professor, Novosibirsk State University.  
 e-mail: sudoplat@math.nsc.ru