Graph of uv-paths in 2-connected graphs *

Eduardo Rivera-Campo

Departamento de Matemáticas Universidad Autónoma Metropolitana - Iztapalapa erc@xanum.uam.mx

Abstract

For a 2-connected graph G and vertices u, v of G we define an abstract graph $\mathcal{P}(G_{uv})$ whose vertices are the paths joining u and vin G, where paths S and T are adjacent if T is obtained from S by replacing a subpath S_{xy} of S with an internally disjoint subpath T_{xy} of T. We prove that $\mathcal{P}(G_{uv})$ is always connected and give a necessary and a sufficient condition for connectedness in cases where the cycles formed by the replacing subpaths are restricted to a specific family of cycles of G.

1 Introduction

For any vertices x, y of a path L, we denote by L_{xy} the subpath of L that joins x and y. Let G be a 2-connected graph and u and v be vertices of G. The uv path graph of G is the graph $\mathcal{P}(G_{uv})$ whose vertices are the paths joining u and v in G, where two paths S and T are adjacent if T is obtained from S by replacing a subpath S_{xy} of S with an internally disjoint subpath T_{xy} of T. The uv path graph $\mathcal{P}(G_{uv})$ is closely related to the graph G(P, f)of f-monotone paths on a polytope P (see C. A. Athanasiadis *et al* [1, 2]), whose vertices are the f-monotone paths on P and where two paths S and Tare adjacent if there is a 2-dimensional face F of P such that T is obtained

^{*}Partially supported by Conacyt, México.

from S by replacing an f-monotone subpath of S contained in F with the complementary f-monotone subpath of T contained in F.

In Section 2 we show that the graphs $\mathcal{P}(G_{uv})$ are always connected as is the case for the graphs G(P, f).

If S and T are adjacent paths in a uv-path graph $\mathcal{P}(G_{uv})$, then $S \cup T$ is a subgraph of G consisting of a unique cycle σ joined to u and v by disjoint paths P_u and P_v . See Figure 1.




Figure 1: $S \cup T$

Let \mathcal{C} be a set of of cycles of G; the *uv*-path graph of G defined by \mathcal{C} is the spanning subgraph $\mathcal{P}_{\mathcal{C}}(G_{uv})$ of $\mathcal{P}(G_{uv})$ where two paths S and T are adjacent if and only if the unique cycle σ which is contained in $S \cup T$ lies in \mathcal{C} . A graph $\mathcal{P}_{\mathcal{C}}(G_{uv})$ may be disconnected.

The uv path graph $\mathcal{P}(G_{uv})$ is also related to the well-known tree graph $\mathcal{T}(G)$ of a connected graph G, studied by R. L. Cummins [3], in which the vertices are the spanning trees of G and the edges correspond to pairs of trees S and R which are obtained from each other by a single edge exchange. As in the uv path graph, if two trees S and R are adjacent in $\mathcal{T}(G)$, then $S \cup R$ is a subgraph of G containing a unique cycle. X. Li *et al* [5] define, in an analogous way, a subgraph $\mathcal{T}_{\mathcal{C}}(G)$ of $\mathcal{T}(G)$ for a set of cycles \mathcal{C} of G and give a necessary condition and a sufficient condition for $\mathcal{T}_{\mathcal{C}}(G)$ to be connected. In sections 3 and 4 we show that the same conditions apply to uv path graphs $\mathcal{P}_{\mathcal{C}}(G_{uv})$.

Similar results are obtained by A. P. Figueroa *et al* [4] with respect to the *perfect matching graph* $\mathcal{M}(G)$ of a graph G where the vertices are the perfect matchings of G and in which two matchings L and N are ajacent if their symmetric difference is a cycle of G. Again, if L and N are adjacent matchings in $\mathcal{M}(G)$, then $L \cup M$ contains a unique cycle of G.

For any subgraphs F and H of a graph G, we denote by $F\Delta H$ the subgraph of G induced by the set of edges $(E(F) \setminus E(H)) \cup (E(H) \setminus E(F))$.

2 Preliminary results

In this section we prove that the uv path graph is connected for any 2connected graph G and give an upper bound for the diameter of a graph $\mathcal{P}(G_{uv})$.

Theorem 1. Let G be a 2-connected graph. The uv-path graph $\mathcal{P}(G_{uv})$ is connected for every pair of vertices u, v of G.

Proof. For any different uv paths Q and R in G denote by n(Q, R) the number of consecutive initial edges Q and R have in common. Assume the result is false and choose two uv paths $S: u = x_0, x_1, \ldots, x_s = v$ and $T: u = y_0, y_1, \ldots, y_t = v$ in different components of $\mathcal{P}(G_{uv})$ for which $n^* = n(S, T)$ is maximum.

Since edges $x_{n^*}x_{n^*+1}$ and $y_{n^*}y_{n^*+1}$ are not equal, $x_{n^*+1} \neq y_{n^*+1}$. Let $j = min\{i : x_{n^*+i} \in V(T)\}$ and $k = min\{i : y_{n^*+i} \in V(S)\}$ and let l and m be integers such that $y_l = x_{n^*+j}, x_m = y_{n^*+k}$. Consider the path:

 $S': u = x_0, x_1, \dots, x_{n^*}, y_{n^*+1}, y_{n^*+2}, \dots, y_{n^*+k}, x_{m+1}, x_{m+2}, \dots, x_s = v$

Paths S and S' are adjacent in $\mathcal{P}(G_{uv})$ since S' is obtained from S by replacing the subpath $x_{n^*}, x_{n^*+1}, \ldots, x_m$ of S with the subpath $y_{n^*}, y_{n^*+1}, \ldots, y_{n^*+k}$ of S'. Notice that $n(S', T) \ge n(S, T) + 1$ since $x_0x_1, x_1x_2, \ldots, x_{n^*-1}x_{n^*}, x_{n^*}y_{n^*+1} \in E(S') \cap E(T)$. By the choice of S, and T, paths S' and T are connected in $\mathcal{P}(G_{uv})$. This implies that S and T are also connected in $\mathcal{P}(G_{uv})$ which is a contradiction.

For any two vertices u and v of a connected graph G we denote by $d_G(u, v)$ the distance between u and v in G, that is the length of a shortest uv path in G. The diameter of a connected graph G is the maximum distance among pairs of vertices of G. For a path P, we denote by l(P) the length of P.

Theorem 2. Let u and v be vertices of a 2-connected graph G. The diameter of the graph $\mathcal{P}(G_{uv})$ is at most $2d_G(u, v)$.

Proof. Let S and T be uv paths in G and let P be a shortest uv path in G. From the proof of Theorem 1 one can see that there are two paths Q_S and Q_T in $\mathcal{P}(G_{uv})$, each with length at most l(P), joining S to P and T to P, respectively. Clearly $Q_S \cup Q_T$ contains a path joining S and T in $\mathcal{P}(G_{uv})$ with length at most $2l(P) = 2d_G(u, v)$. In Figure 2 we show a 2-connected graph G^2 and paths S and T joining vertices u and v of G^2 such that $d_{G^2}(u,v) = 2$ and $d_{\mathcal{P}(G^2_{uv})}(S,T) = 4$. For any positive integer k > 2 the graph G^2 can be extended to a graph G^k such that $d_{G^k}(u,v) = k$ and that the diameter of $\mathcal{P}(G^k_{uv})$ is 2k. This shows that Theorem 2 is tight.




Figure 2: Graph G^2 and paths S and T.

3 Necessary condition

Let u and v be vertices of a 2-connected graph G and S and T be two uv paths adjacent in $\mathcal{P}(G_{uv})$. Since T is obtained from S by replacing a subpath S_{xy} of S with an internally disjoint subpath T_{xy} of T, the graph $S\Delta T$ is the cycle $S_{xy} \cup T_{xy}$.

Theorem 3. Let G be a 2-connected graph, u and v be vertices of G and C be a set of cycles of G. If the graph $\mathcal{P}_{\mathcal{C}}(G_{uv})$ is connected, then C spans the cycle space of G.

Proof. Let σ be a cycle of G. Since G is 2-connected, there are two disjoint paths P_u and P_v joining, respectively, u and v to σ . Denote by u' and v'the unique vertices of P_u and P_v , respectively, that lie in σ . Vertices u'and v' partition cycle σ into two internally disjoint paths Q and R. Let $S = P_u \cup Q \cup P_v$ and $T = P_u \cup R \cup P_v$. Clearly S and T are two different uvpaths in G such that $S\Delta T = \sigma$.

Since $\mathcal{P}_{\mathcal{C}}(G_{uv})$ is connected, there are uv paths $S = W_0, W_1, \ldots, W_k = T$ such that for $i = 1, 2, \ldots k$, paths W_{i-1} and W_i are adjacent in $\mathcal{P}_{\mathcal{C}}(G_{uv})$. For $i = 1, 2, \ldots k$ let $\alpha_i = W_{i-1} \Delta W_i$. Then $\alpha_1, \alpha_2, \ldots, \alpha_k$ are cycles in \mathcal{C} such that:

$$\alpha_1 \Delta \alpha_2 \Delta \cdots \Delta \alpha_k = (W_0 \Delta W_1) \Delta (W_1 \Delta W_2) \Delta \cdots \Delta (W_{k-1} \Delta W_k) = W_0 \Delta W_k = \sigma$$

Therefore \mathcal{C} spans σ .

Let G be a complete graph with four vertices u, x, y, v and let $\mathcal{C} = \{\alpha, \beta, \delta\}$, where $\alpha = uxv$, $\beta = uyv$ and $\delta = uxyv$. Set \mathcal{C} spans the cycle space of G but the graph $\mathcal{P}_{\mathcal{C}}(G_{uv})$ is not connected since the uv path uyxv is an isolated vertex of $\mathcal{P}_{\mathcal{C}}(G_{uv})$, see Fig 3. This shows that the condition in Theorem 3 is not sufficient for $\mathcal{P}_{\mathcal{C}}(G_{uv})$ to be connected.




Figure 3: Graph G, set $\mathcal{C} = \{\alpha, \beta, \delta\}$ and graph $\mathcal{P}_C(G_{uv})$.

4 Sufficient condition

A unicycle of a connected graph G is a spanning subgraph \mathcal{U} of G that contains a unique cycle. Let u and v be vertices of a 2-connected graph G. A uv-monocle of G is a subgraph of G that consists of a cycle σ and two disjoint paths P_u and P_v that join, respectively u and v to σ , see Fig. 1. Clearly for each uv-monocle \mathcal{M} of a 2-connected graph G, there is a unicycle \mathcal{U} of G that contains \mathcal{M} .

Let \mathcal{C} be a set of cycles of G. A cycle σ of G has Property Δ^* with respect to \mathcal{C} if for every unicycle \mathcal{U} containing σ there is an edge e of G, not in \mathcal{U} and two cycles $\alpha, \beta \in \mathcal{C}$, contained in $\mathcal{U} + e$, such that $\sigma = \alpha \Delta \beta$. **Lemma 1.** Let G be a 2-connected graph and u and v be vertices of G. Also let C be a set of cycles of G and σ be a cycle having Property Δ^* with respect to C. The graph $\mathcal{P}_{\mathcal{C}\cup\{\sigma\}}(G_{uv})$ is connected if and only if $\mathcal{P}_{\mathcal{C}}(G_{uv})$ is connected. Proof. If $\mathcal{P}_{\mathcal{C}}(G_{uv})$ is connected, then $\mathcal{P}_{\mathcal{C}\cup\{\sigma\}}(G_{uv})$ is connected since the former is a subgraph of the latter.

Assume now $\mathcal{P}_{\mathcal{C}\cup\{\sigma\}}(G_{uv})$ is connected and let S and T be uv paths in G which are adjacent in $\mathcal{P}_{\mathcal{C}\cup\{\sigma\}}(G_{uv})$. We show next that S and T are connected in $\mathcal{P}_{\mathcal{C}}(G_{uv})$ by a path of length at most 2.

If $\omega = S\Delta T \in \mathcal{C}$, then S and T are adjacent in $\mathcal{P}_{\mathcal{C}}(G_{uv})$. For the case $\omega = \sigma$ denote by \mathcal{M} the *uv*-monocle given by $S \cup T$.

Let \mathcal{U} be a unicycle of G containing \mathcal{M} . Since σ has Property Δ^* with respect to \mathcal{C} , there exists an edge e = xy of G, not in \mathcal{U} , and two cycles $\alpha, \beta \in \mathcal{C}$ contained in $\mathcal{U} + e$ such that $\sigma = \alpha \Delta \beta$.

Let x' and y' denote the vertices in \mathcal{M} which are closest in \mathcal{U} to x and y, respectively. Then there exists a path $R_{x'y'}$ in G, with edges in $E(\mathcal{U} + e) \setminus E(\mathcal{M})$ joining x' and y' and such that cycles α and β are contained in $\mathcal{M} \cup R_{x'y'}$. We analyze several cases according to the location of x' and y' in \mathcal{M} .

Denote by P_u and P_v the unique paths, contained in \mathcal{M} , that join u and v to σ and by u' and v' the vertices where P_u and P_v , respectively, meet σ .

Case 1.- $x' \in V(P_u), y' \in V(P_v)$. Without loss of generality we assume $\alpha = S_{x'y'} \cup R_{y'x'}$ and $\beta = T_{x'y'} \cup R_{y'x'}$, see Fig. 4.




Figure 4: Left: $\mathcal{M} \cup R_{y'x'}$. Right: Cycles α and β .

Let Q be the uv-path obtained from S by replacing $S_{x'y'}$ with $R_{x'y'}$. Notice that Q can also be obtained from T by replacing $T_{x'y'}$ with $R_{x'y'}$. Case 2.- $x' \in V(P_u), y' \in S \cap \sigma$. Without loss of generality we assume $\alpha = S_{x'y'} \cup R_{y'x'}$ and $\beta = T_{x'v'} \cup S_{v'y'} \cup R_{y'x'}$, see Fig. 5.




Figure 5: Left: $\mathcal{M} \cup R_{y'x'}$. Right: Cycles α and β .

Again let Q be the uv-path obtained from S by replacing $S_{x'y'}$ with $R_{x'y'}$. In this case, Q can also be obtained from T by replacing $T_{x'v'}$ with $R_{x'y'} \cup S_{y'v'}$.

Case 3.- $x', y' \in S \cap \sigma$. Without loss of generality we assume $\alpha = S_{x'y'} \cup R_{y'x'}$ and $\beta = S_{u'x'} \cup R_{x'y'} \cup S_{y'v'} \cup T_{v'u'}$, see Fig. 6.




Figure 6: Left: $\mathcal{M} \cup R_{y'x'}$. Right: Cycles α and β .

Let Q be the uv-path obtained from S by replacing $S_{x'y'}$ with $R_{x'y'}$. Path Q is also obtained from T by replacing $T_{u'v'}$ with $S_{u'x'} \cup R_{x'y'} \cup S_{y'v'}$.

Case 4.- $x' \in S \cap \sigma$ and $y' \in T \cap \sigma$. Without loss of generality we assume $\alpha = S_{u'x'} \cup R_{x'y'} \cup T_{y'u'}$ and $\beta = T_{y'v'} \cup S_{v'x'} \cup R_{x'y'}$. , see Fig. 7.

Let Q be the uv-path obtained from S by replacing $S_{u'x'}$ with $T_{u'y'} \cup R_{y'x'}$. Now Q can also be obtained from T by replacing $T_{y'v'}$ with $R_{y',x'} \cup S_{x'v'}$




Figure 7: Left: $\mathcal{M} \cup R_{u'x'}$. Right: Cycles α and β .

In each case $S\Delta Q = \alpha$ and $Q\Delta T = \beta$. Since $\alpha, \beta \in C$, path S is adjacent to Q and path Q is adjacent to T in $P_{\mathcal{C}}(G_{uv})$. Therefore S and T are connected in $\mathcal{P}_{\mathcal{C}}(G_{uv})$ by a path with length at most 2.

All remaining cases are analogous to either Case 2 or to Case 3.

Consider a 2-connected graph G with two specified vertices u and v and let \mathcal{C} be a set of cycles of G. Construct a sequence of sets of cycles $\mathcal{C} = \mathcal{C}_0, \mathcal{C}_1, \ldots, \mathcal{C}_k$ as follows: If there is a cycle σ_1 not in \mathcal{C}_0 that has Property Δ^* with respect to \mathcal{C}_0 add σ_1 to \mathcal{C}_0 to obtain \mathcal{C}_1 . At step t add to \mathcal{C}_t a new cycle σ_{t+1} (if it exists) that has Property Δ^* with respect to \mathcal{C}_t to obtain \mathcal{C}_{t+1} . Stop at a step k where there are no cycles, not in \mathcal{C}_k , having Property Δ^* with respect to \mathcal{C}_k . We denote by $Cl(\mathcal{C})$ the final set obtained with this process. Li *et al* [5] proved that the final set of cycles obtained is independent of which cycle σ_t is added at each step in the case of multiple possibilities.

A set of cycles of G is Δ^* -dense if $Cl(\mathcal{C})$ is the whole set of cycles of G.

Theorem 4. If \mathcal{C} is Δ^* -dense, then $\mathcal{P}_{\mathcal{C}}(G_{uv})$ is connected.

Proof. Since C is Δ^* -dense, Cl(C) is the set of cycles of G and therefore $\mathcal{P}_{Cl(C)}(G_{uv}) = \mathcal{P}(G_{uv})$ which is connected by Theorem 1.

Let $C = C_0, C_1, \ldots, C_k = Cl(C)$ be a sequence of sets of cycles obtained from C as above. By Lemma 1, all graphs $\mathcal{P}_{Cl(C)}(G_{uv}) = \mathcal{P}_{C_k}(G_{uv}), \mathcal{P}_{C_{k-1}}(G_{uv}), \ldots, \mathcal{P}_{C_0}(G_{uv}) = \mathcal{P}_{C}(G_{uv})$ are connected. \Box

Li *et al* [5] proved the following:

Theorem 5. If G is a plane 2-connected graph and C is the set of internal faces of G, then C is Δ^* -dense.

Theorem 6. If G is a 2-connected graph and C is the set of cycles that contain a given edge e of G, then C is Δ^* -dense.

We end this section with the following immediate corollaries.

Collorary 1. Let u and v be vertices of a 2-connected plane graph G. If C is the set of internal faces of G, then $\mathcal{P}_{\mathcal{C}}(G_{uv})$ is connected.

Proof. By Theorem 5, \mathcal{C} is Δ^* -dense and by Theorem 4, $\mathcal{P}_{\mathcal{C}}(G_{uv})$ is connected.

Collorary 2. Let u and v be vertices of a 2-connected graph G. If C is the set of cycles of G that contain a given edge e, then $\mathcal{P}_{\mathcal{C}}(G_{uv})$ is connected.

Proof. By Theorem 6, \mathcal{C} is Δ^* -dense and by Theorem 4, $\mathcal{P}_{\mathcal{C}}(G_{uv})$ is connected.

Collorary 3. Let u and v be vertices of a 2-connected graph G. If C_u is the set of cycles of G that contain vertex u, then $\mathcal{P}_{C_u}(G_{uv})$ is connected.

Proof. Let e be an edge of G incident with vertex u. Clearly the set $\mathcal{C}(e)$ of cycles that contain edge e is a subset of the set \mathcal{C}_u . Therefore $\mathcal{P}_{\mathcal{C}(e)}(G_{uv})$ is a subgraph of $\mathcal{P}_{\mathcal{C}_u}(G_{uv})$. By Corollary 2, the graph $\mathcal{P}_{\mathcal{C}(e)}(G_{uv})$ is connected. \Box

References

- Athanasiadis, C.A.; de Loera, J. A.; Zhang, Z.; Enumerative problems for arborescences and monotone paths on polytopes, arXiv:2002.00999v1 [math.CO].
- [2] Athanasiadis, C.A.; Edelman, P. H.; Reiner, V., Monotone paths on polytopes, Math. Z. 235 (2000), 315 – 334.
- [3] Cummins, R. L., Hamilton circuits in tree graphs, IEEE Trans. Circuit Theory, vol. CT-13 (1966), 82–90,
- [4] Figueroa, A. P.; Fresán-Figueroa, J.; Rivera-Campo, E., On the perfect matching graph defined by a set of cycles, Bol. Soc. Mat. Mex. 23 Issue 2 (2017), 549 – 556.

[5] Li, X.; Neumann-Lara, V.; Rivera-Campo, E., On a tree graph defined by a set of cycles, Discrete Math. 271 (2003), 303–310.