

ON HOLOMORPHIC CONTRACTIBILITY OF TEICHMÜLLER SPACES

SAMUEL L. KRUSHKAL

ABSTRACT. The problem of holomorphic contractibility of the Teichmüller spaces $\mathbf{T}(0, n)$ of punctured spheres ($n > 4$) arose in the 1970s in connection with solving algebraic equations in Banach algebras. Recently it was solved by the author in [12].

In the present paper we improve the statement of Lemma 3 in [12] and provide an alternate proof of holomorphic contractibility of low dimensional Teichmüller spaces.

2010 Mathematics Subject Classification: Primary: 30C55, 30F60; Secondary: 30F35, 46G20

Key words and phrases: Teichmüller spaces, Fuchsian group, quasiconformal deformations, holomorphic contractibility, univalent function, Schwarzian derivative, holomorphic sections

1. PREAMBLE

1.1. A complex Banach manifold X is contractible to its point x_0 if there exists a continuous map $F : X \times [0, 1] \rightarrow X$ with $F(x, 0) = x$ and $F(x, 1) = x_0$ for all $x \in X$. If the map F can be chosen so that for every $t \in [0, 1]$ the map $F_t : x \mapsto F(x, t)$ of X to itself is holomorphic and $F_t(x_0) = x_0$, then X is called holomorphically contractible to x_0 .

The problem of holomorphic contractibility of Teichmüller spaces $\mathbf{T}(0, n)$ of the punctured spheres ($n > 4$) arose in the 1970s in connection with solving the algebraic equations in Banach algebras. It was caused by the fact that in the space $\mathbb{C}^m$, $m > 1$, there are domains (even bounded), which are only topologically but not holomorphically contractible (see [7], [8], [15], [16]).

Recently this problem was solved positively in [12]. There is established that all space $\mathbf{T}(0, n)$, $n > 4$, are holomorphically contractible.

The proof of Lemma 3 in this paper contains a wrongly assertion (which does not be used there) that the map s_m (giving the inclusion of the space $T(\Gamma_0)$ into $T(\Gamma_0^m)$) is a section of the forgetful map $\chi_m : T(X_{a^0}^m) \rightarrow T(X_{a^0})$. Such sections do not exist if $n > 6$.

In fact, there was only used in the proof that s_m is an open holomorphic map (of a domain onto manifold), and the openness is preserved for the limit map $s = \lim_{m \rightarrow \infty} s_m$ which determines an $(n - 3)$ -dimensional complex submanifold $s(T(X_{a^0}))$ in the universal Teichmüller space $\mathbf{T}$.

In the present paper, we improve the statement of the indicated Lemma 3 (without changing other arguments in [12] concerning the contractibility of $\mathbf{T}(0, n)$). In the second part of the paper, we provide an alternate proof of holomorphic contractibility of low dimensional Teichmüller spaces (of dimensions two and three), which has an independent interest in view of importance of the problem. The underlying idea of this proof is different from [12].

1.2. There are two Teichmüller spaces of dimension two: the space $\mathbf{T}(0, 5)$ of the spheres with five punctures and the space $\mathbf{T}(1, 2)$ of twice punctured tori; these spaces are biholomorphically equivalent. Such spheres and tori are uniformized by the corresponding Fuchsian groups Γ and

Γ' so that Γ is a subgroup of index two in Γ' ; letting $\mathbf{T}(0, 5) = \mathbf{T}(\Gamma)$, $\mathbf{T}(1, 2) = \mathbf{T}(\Gamma')$, we have $\mathbf{T}(\Gamma') = \mathbf{T}(\Gamma)$.

In a similar way, the Teichmüller spaces $\mathbf{T}(0, 6)$ of spheres with six punctures and $\mathbf{T}(2, 0)$ of closed Riemann surfaces of genus 2 also are biholomorphically equivalent, and in terms of the corresponding Fuchsian groups Γ and Γ' we have the same relation $\mathbf{T}(\Gamma') = \mathbf{T}(\Gamma)$. We state:

Theorem 1. *The spaces $\mathbf{T}(0, 5)$, $\mathbf{T}(1, 2)$, $\mathbf{T}(0, 6)$, $\mathbf{T}(2, 0)$ are holomorphically contractible.*

It remains the Teichmüller space $\mathbf{T}(1, 3)$ of tori with three punctures also having dimension three; it will not be involved here.

2. UNDERLYING RESULTS

2.1. Consider the ordered n -tuples of points

$$\mathbf{a} = (0, 1, a_1, \dots, a_{n-3}, \infty), \quad n > 4, \quad (1)$$

with distinct $a_j \in \widehat{\mathbb{C}} \setminus \{1, -1, i\}$ and the corresponding punctured spheres

$$X_{\mathbf{a}} = \widehat{\mathbb{C}} \setminus \{0, 1, a_1, \dots, a_{n-3}, \infty\}, \quad \widehat{\mathbb{C}} = \mathbb{C} \cup \{\infty\},$$

regarded as the Riemann surfaces of genus zero. Fix a collection $\mathbf{a}^0 = (0, 1, a_1^0, \dots, a_{n-3}^0, 1, \infty)$ defining the base point $X_{\mathbf{a}^0}$ of Teichmüller space $\mathbf{T}(0, n) = \mathbf{T}(X_{\mathbf{a}^0})$ of n -punctured spheres. Its points are the equivalence classes $[\mu]$ of Beltrami coefficients from the ball

$$\text{Belt}(\mathbb{C})_1 = \{\mu \in L_\infty(\mathbb{C}) : \|\mu\|_\infty < 1\},$$

under the relation that $\mu_1 \sim \mu_2$ if the corresponding quasiconformal homeomorphisms $w^{\mu_1}, w^{\mu_2} : X_{\mathbf{a}^0} \rightarrow X_{\mathbf{a}}$ (the solutions of the Beltrami equation $\bar{\partial}w = \mu\partial w$ with $\mu = \mu_1, \mu_2$) are homotopic on $X_{\mathbf{a}^0}$ (and hence coincide in the points $0, 1, a_1^0, \dots, a_{n-3}^0, \infty$). This models $\mathbf{T}(0, n)$ as the quotient space

$$\mathbf{T}(0, n) = \text{Belt}(\mathbb{C})_1 / \sim$$

with complex Banach structure of dimension $n - 3$ inherited from the ball $\text{Belt}(\mathbb{C})_1$.

Another canonical model of $\mathbf{T}(0, n) = \mathbf{T}(X_{\mathbf{a}^0})$ is obtained using the uniformization of Riemann surfaces and the holomorphic Bers embedding of Teichmüller spaces. Consider the upper and lower half-planes

$$U = \{z = x + iy : y > 0\}, \quad U^* = \{z \in \widehat{\mathbb{C}} : y < 0\}$$

The holomorphic universal covering map $h : U \rightarrow X_{\mathbf{a}^0}$ provides a torsion free Fuchsian group Γ_0 of the first kind acting discontinuously on $U \cup U^*$, and the surface $X_{\mathbf{a}^0}$ is represented (up to conformal equivalence) as the quotient space U/Γ_0 . The functions $\mu \in L_\infty(X_{\mathbf{a}^0}) = L_\infty(\mathbb{C})$ are lifted to U as the Beltrami $(-1, 1)$ -measurable forms $\tilde{\mu}d\bar{z}/dz$ on U with respect to Γ_0 satisfying $(\tilde{\mu} \circ \gamma)\overline{\gamma'}/\gamma' = \tilde{\mu}$, $\gamma \in \Gamma_0$ and form the corresponding Banach space $L_\infty(U, \Gamma_0)$. We extend these $\tilde{\mu}$ by zero to U^* and consider the unit ball $\text{Belt}(U, \Gamma_0)_1$ of this space $L_\infty(U, \Gamma_0)$. The corresponding quasiconformal maps $w^{\hat{\mu}}$ are conformal on the half-plane U^* , and their *Schwarzian derivatives*

$$S_w(z) = \left(\frac{w'''(z)}{w'(z)} \right)' - \frac{1}{2} \left(\frac{w''(z)}{w'(z)} \right)^2, \quad w = w^{\hat{\mu}},$$

fill a bounded domain in the complex $(n-3)$ -dimensional space $\mathbf{B}(\Gamma_0) = \mathbf{B}(U^*, \Gamma_0)$ of hyperbolically bounded holomorphic Γ_0 -automorphic forms of degree -4 on U^* (i.e., satisfy $(\varphi \circ \gamma)(\gamma')^2 = \varphi$, $\gamma \in \Gamma_0$), with norm

$$\|\varphi\| = \sup_{U^*} 4y^2 |\varphi(z)|.$$

This domain models the *Teichmüller space* $\mathbf{T}(\Gamma_0)$ of the group Γ_0 . It is canonically isomorphic to the space $\mathbf{T}(X_{\mathbf{a}^0})$ (and to a bounded domain in the complex Euclidean space $\mathbb{C}^{n-3}$).

The indicated map $\widehat{\mu} \rightarrow S_{w\widehat{\mu}}$ determines a holomorphic map $\phi_{\mathbf{T}} : \text{Belt}(U, \Gamma_0)_1 \rightarrow \mathbf{B}(\Gamma_0)$; it has only local holomorphic sections.

Note also that $\mathbf{T}(\Gamma_0) = \mathbf{T} \cap \mathbf{B}(\Gamma_0)$, where $\mathbf{T}$ is the universal Teichmüller space (modelled as domain of the Schwarzian derivatives of all univalent functions on U^* admitting quasiconformal extension to U).

2.2. The collections (1) fill a domain $\mathcal{D}_n$ in $\mathbb{C}^{n-3}$ obtained by deleting from this space the hyperplanes $\{z = (z_1, \dots, z_{n-3}) : z_j = z_l, j \neq l\}$, and with $z_1 = 0, z_2 = 1$. This domain represents the Torelli space of the spheres $X_{\mathbf{a}}$ and is covered by $\mathbf{T}(0, n)$, which is given by the following lemma (cf. e.g., [10]; [13], Section 2.8).

Lemma 1. *The holomorphic universal covering space of $\mathcal{D}_n$ is the Teichmüller space $\mathbf{T}(0, n)$. This means that for each punctured sphere $X_{\mathbf{a}}$ there is a holomorphic universal covering*

$$\pi_{\mathbf{a}} : \mathbf{T}(0, n) = \mathbf{T}(X_{\mathbf{a}}) \rightarrow \mathcal{D}_n.$$

The covering map $\pi_{\mathbf{a}}$ is well defined by

$$\pi_{\mathbf{a}} \circ \phi_{\mathbf{a}}(\mu) = (0, 1, w^{\mu}(a_1), \dots, w^{\mu}(a_{n-3}), \infty),$$

where $\phi_{\mathbf{a}}$ denotes the canonical projection of the ball $\text{Belt}(\mathbb{C})_1$ onto the space $\mathbf{T}(X_{\mathbf{a}})$.

Lemma 1 yields also that the truncated collections $\mathbf{a}_* = (a_1, \dots, a_{n-3})$ provide the local complex coordinates on the space $\mathbf{T}(0, n)$ and define its complex structure.

These coordinates are simply connected with the Bers local complex coordinates on $\mathbf{T}(0, n)$ (related to bases of the tangent spaces to $\mathbf{T}(0, n)$ at its points, see [1]) via standard variation of quasiconformal maps of $X_{\mathbf{a}} = U/\Gamma_{\mathbf{a}}$

$$\begin{aligned} w^{\mu}(z) &= z - \frac{z(z-1)}{\pi} \iint_{\mathbb{C}} \frac{\mu(\zeta)}{\zeta(\zeta-1)(\zeta-z)} d\xi d\eta + O(\|\mu\|_{\infty}^2) \\ &= z - \frac{z(z-1)}{\pi} \sum_{\gamma \in \Gamma_{\mathbf{a}} U/\Gamma_{\mathbf{a}}} \iint \frac{\mu(\gamma\zeta) |\gamma'(\zeta)|^2}{\gamma\zeta(\gamma\zeta-1)(\gamma\zeta-z)} d\xi d\eta + O(\|\mu\|_{\infty}^2). \end{aligned}$$

with uniform estimate of the ratio $O(\|\mu\|_{\infty}^2)/\|\mu\|_{\infty}^2$ on compacts in $\mathbb{C}$ (see e.g., [11]).

It turns out that one can obtain the whole space $\mathbf{T}(X_{\mathbf{a}^0})$ using only the similar equivalence classes $[\mu]$ of the Beltrami coefficients from the ball $\mu \in \text{Belt}(U)_1$ (vanishing on U^*), requiring that the corresponding quasiconformal homeomorphisms w^{μ} are homotopic on the punctured sphere $X_{\mathbf{a}^0}$. Surjectivity of this holomorphic quotient map

$$\chi : \text{Belt}(U)_1 \rightarrow \mathbf{T}(0, n),$$

is a consequence of following interpolation result from [3].

Lemma 2. *Given two cyclically ordered collections of points $(z_1, \dots, z_m)$ and $(\zeta_1, \dots, \zeta_m)$ on the unit circle $S^1 = \{|z| = 1\}$, there exists a holomorphic univalent function f in the closure of the unit disk $\mathbb{D} = \{|z| < 1\}$ such that $|f(z)| < 1$ for $z \in \mathbb{D}$ distinct from $z_1, \dots, z_m$, and $f(z_k) = \zeta_k$ for all $k = 1, \dots, m$. Moreover, there exist univalent polynomials f with such an interpolation property.*

It follows that the function f given by Lemma 2 actually is holomorphic and univalent (hence, maps conformally) in a broader disk $\mathbb{D}_r$, $r > 1$.

First of all, $f'(z) \neq 0$ on the unit circle S^1 . Indeed, were $f'(z_0) = 0$ at some point $z_0 \in S^1$, then in its neighborhood $f(z) = c_s(z - z_0)^s + O((z - z_0)^{s+1}) = c_s \zeta^s$, where $c_s \neq 0$ for some $s > 1$, which contradicts to injectivity of $f(z)$ on S^1 . So f is univalent in some disk $\mathbb{D}_r = \{|z| < r\}$, $r > 1$.

Now, assuming, in the contrary, that f is not globally univalent in any admissible disk $\mathbb{D}_r$ with $r > 1$, one obtains the distinct sequences $\{z'_n\}, \{z''_n\} \subset \mathbb{D}_r$ with $f(z'_n) = f(z''_n)$ for any n , whose limit points z'_0, z''_0 lie on S^1 . Then also in the limit, $f(z'_0) = f(z''_0)$, which in the case $z'_0 \neq z''_0$ contradicts to univalence of f on S^1 given by Lemma 2 and in the case $z'_0 = z''_0 = z_0$ contradicts to local univalence of f in a neighborhood of z_0 .

The interpolating function f given by Lemma 2 is extended quasiconformally to the whole sphere $\widehat{\mathbb{C}}$ across any circle $\{|z| = r\}$ with $r > 1$ indicated above. Hence, given a cyclically ordered collection $(z_1, \dots, z_m)$ of points on S^1 , then for any ordered collection $(\zeta_1, \dots, \zeta_m)$ in $\widehat{\mathbb{C}}$, there exists a quasiconformal homeomorphism $\widehat{f}$ of the sphere $\widehat{\mathbb{C}}$ carrying the points z_j to ζ_j , $j = 1, \dots, m$, and such that its restriction to the closed disk $\overline{\mathbb{D}}$ is biholomorphic on $\overline{\mathbb{D}}$.

Taking the quasicircles L passing through the points $\zeta_1, \dots, \zeta_m$ and such that each ζ_j belongs to an analytic subarc of L , one obtains quasiconformal extensions of the interpolating function f , which are biholomorphic on the union of $\overline{\mathbb{D}}$ and some neighborhoods of the initial points $z_1, \dots, z_m \in S^1$. Now Lemma 1 provides quasiconformal extensions of f lying in prescribed homotopy classes of homeomorphisms $X_{\mathbf{z}} \rightarrow X_{\mathbf{w}}$.

2.3. Pick a space $\mathbf{T}(0, n) = \mathbf{T}(X_{\mathbf{a}^0})$ with $n \geq 5$ and let

$$X'_{\mathbf{a}^0} = X_{\mathbf{a}^0} \setminus \{a_{n-3}^0\} = U/\Gamma'_0.$$

Due to the Bers isomorphism theorem, *the space $T(X'_{\mathbf{a}^0})$ is biholomorphically isomorphic to the Bers fiber space*

$$\mathcal{F}(0, n) := \mathcal{F}(T(X_{\mathbf{a}^0})) = \{(\phi_{\mathbf{T}}(\mu), z) \in \mathbf{T}(X_{\mathbf{a}^0}) \times \mathbb{C} : \mu \in \text{Belt}(U, \Gamma'_0)_1, z \in w^\mu(\mathbb{D})\}$$

over $\mathbf{T}(X_{\mathbf{a}^0})$ with holomorphic projection $\pi(\varphi, z) = \varphi$ ($\varphi \in T(X_{\mathbf{a}^0})$) (see [2]).

This fiber space is a bounded hyperbolic domain in $\mathbf{B}(\Gamma_0) \times \mathbb{C}$ and represents the collection of domains $D_\mu = w^\mu(U)$ (the universal covers of the surfaces $X_{\mathbf{a}^0}$) as a holomorphic family over the space $\mathbf{T}(0, n-1) = T(X_{\mathbf{a}^0})$.

The indicated isomorphism between $\mathbf{T}(0, n+1)$ and $\mathcal{F}(0, n)$ is induced by the inclusion map $j : \mathbb{D}_* \hookrightarrow \mathbb{D}$ forgetting the puncture at a_n^0 , via

$$\mu \mapsto (S_{w^{\mu_1}}, w^{\mu_1}(\widehat{a}_{n-3}^0)) \quad \text{with} \quad \mu_1 = j_*\mu := (\mu \circ \widehat{j}_0)\widehat{j}'/\widehat{j}', \quad (2)$$

where $\widehat{j}$ is the lift of j to U and $\widehat{a}_{n-3}^0$ is one of the points from the fiber over a_n^0 under the quotient map $U \rightarrow U/\Gamma_0$.

Note also that the holomorphic universal covering maps $h : U^* \rightarrow U^*/\Gamma_0$ and $h' : U^* \rightarrow U^*/\Gamma'_0$ (and similarly the corresponding covering maps in U) are related by

$$j \circ h' = h \circ \widehat{j},$$

where $\widehat{j}$ is the lift of j . This induces a surjective homomorphism of the covering groups $\theta : \Gamma_0 \rightarrow \Gamma'_0$ by

$$\widehat{j} \circ \gamma = \theta(\gamma) \circ \widehat{j}, \quad \gamma \in \Gamma_0$$

and the norm preserving isomorphism $\widehat{j}_* : \mathbf{B}(\Gamma_0) \rightarrow \mathbf{B}(\Gamma'_0)$ by

$$\widehat{j}_*\varphi = (\varphi \circ \widehat{j})(\widehat{j}')^2, \quad (3)$$

which projects to the surfaces $X_{\mathbf{a}^0}$ and $X'_{\mathbf{a}^0}$ as the inclusion of the space $Q(X_{\mathbf{a}^0})$ of holomorphic quadratic differentials corresponding to $\mathbf{B}(\Gamma_0)$ into the space $Q(X'_{\mathbf{a}^0})$ (cf. [5]).

The Bers theorem is valid for Teichmüller space $\mathbf{T}(X_0 \setminus \{x_0\})$ of any punctured hyperbolic Riemann surface $X_0 \setminus \{x_0\}$ and implies that $\mathbf{T}(X_0 \setminus \{x_0\})$ is biholomorphically isomorphic to the Bers fiber space $\mathcal{F}(\mathbf{T}(X_0))$ over $\mathbf{T}(X_0)$.

2.4. The group Γ'_0 uniformizing the surface $X_{\mathbf{a}^0}$ acts discontinuously on the fiber space $\mathcal{F}(\Gamma_0)$ as a group of biholomorphic maps by

$$\gamma(\phi_{\mathbf{T}}(\mu), z) = (\phi_{\mathbf{T}}(\mu), \gamma^\mu z), \quad (4)$$

where $\mu \in \text{Belt}(U, \Gamma_0)$, $z \in w^\mu(U)$, $\gamma \in \Gamma_0$, and

$$\gamma^\mu \circ w^\mu = w^\mu \circ \gamma$$

(see [2]). The quotient space

$$\mathcal{V}(0, n) := \mathcal{V}(\Gamma_0) = \mathbf{T}(0, n+1)/\Gamma_0$$

is called the *n-punctured Teichmüller curve* and depends only on the analytic type of the group Γ_0 . The projection $\pi : \mathcal{F}(0, n) \rightarrow \mathbf{T}(0, n)$ induces a holomorphic projection

$$\pi_1 : \mathcal{V}(0, n) \rightarrow \mathbf{T}(0, n). \quad (5)$$

This curve also is a complex manifold with fibers $\pi^{-1}(x) = X_{\mathbf{a}}$.

Due to the deep Hubbard-Earle-Kra theorem [5], [9], the projections $\mathcal{V}(0, n) \rightarrow \mathbf{T}(0, n)$ and (4) have no holomorphic sections for any $n \geq 7$ (more generally, for all spaces $\mathbf{T}(\Gamma)$ corresponding to groups Γ without elliptic elements, excluding the spaces $\mathbf{T}(1, 2) \simeq \mathbf{T}(0, 5)$ and $\mathbf{T}(2, 0) \simeq \mathbf{T}(0, 6)$). Such sections exist for groups Γ containing elliptic elements.

In the exceptional cases of $\mathbf{T}(1, 2)$ and $\mathbf{T}(2, 0)$, there is a group Γ' which contains Γ as a subgroup of index two. Then $\mathbf{T}(\Gamma') = \mathbf{T}(\Gamma)$, $\mathcal{F}(\Gamma') = \mathcal{F}(\Gamma)$, and the elliptic elements $\gamma \in \Gamma'$ produce the indicated holomorphic sections s as the maps

$$\phi_{\mathbf{T}}(\mu) \mapsto (\phi_{\mathbf{T}}(\mu), w^\mu(z_0)), \quad (6)$$

where z_0 is a fixed point of γ in the half-plane U . These sections are called the *Weierstrass sections* (in view of their connection with the Weierstrass points of hyperelliptic surface U/Γ). We describe these sections following [5].

Consider also the *punctured fiber space* $\mathcal{F}_0(\Gamma)$ to be the largest open dense subset of $\mathcal{F}(\Gamma)$ on which Γ acts freely, and let

$$\mathcal{V}'(\Gamma) = \mathcal{F}_0(\Gamma)/\Gamma.$$

For Γ with no elliptic elements, the universal covering space for $\mathcal{V}'(g, n) = \mathcal{V}'(\Gamma)$ is $\mathbf{T}(g, n+1)$.

If Γ contains elliptic elements γ , then any holomorphic section $\mathbf{T}(\Gamma) \rightarrow \mathcal{F}(\Gamma)$ is determined by the maps (6) so that $w^\mu(z_0)$ is exactly one fixed point of the corresponding map (4) in the fiber $w^\mu(U)$. These holomorphic sections take their values in the set $\mathcal{V}(\Gamma) \setminus \mathcal{V}'(\Gamma)$ and do not provide in general sections of projection (5).

In the case of spaces $\mathbf{T}(1, 2)$ and $\mathbf{T}(2, 0)$, either of the corresponding curves $\mathcal{V}(1, 2)$ and $\mathcal{V}(2, 0)$ has a unique biholomorphic self-map γ of order two which maps each fiber into itself. The fixed-point locus of γ is a finite set of connected closed complex submanifolds of $\mathcal{V}'(g, n)$, and the restriction of the map (5) to one of these submanifolds is holomorphic map onto $\mathbf{T}(0, n)$; its inverse is a Weierstrass section. The restriction of γ to each fiber is a conformal involution of the corresponding hyperelliptic Riemann surface interchanging its sheets, and the fixed points of γ are the Weierstrass points on this surface.

In dimension one, there are three biholomorphically isomorphic Teichmüller spaces $\mathbf{T}(1, 0)$, $\mathbf{T}(1, 1)$ and $\mathbf{T}(0, 4)$ (see, e.g., [2], [14]). We shall use the last two spaces. Their fiber space $\mathcal{F}(0, 4)$ is isomorphic to $\mathbf{T}(0, 5)$.

As a special case of the Hubbard-Earle-Kra theorem [5], [9], we have:

Lemma 3. (a) The curve $\mathcal{V}(0, 4)$ has for any its point x a unique holomorphic section s with $s(\pi_1(x)) = x$.

(b) If $\dim \mathcal{V}(g, n)' > 1$, the only curves $\mathcal{V}(1, 2)'$ and $\mathcal{V}(2, 0)'$ have holomorphic sections, which are their Weierstrass sections.

In particular, the curve $\mathcal{V}(2, 0)$ has six disjoint holomorphic sections corresponding to the Weierstrass points of hyperelliptic surfaces of genus two.

3. HOLOMORPHIC MAPS OF $\mathbf{T}(0, n)$ INTO UNIVERSAL TEICHMÜLLER SPACE

3.1. The universal Teichmüller space $\mathbf{T} = \text{Teich}(U)$ is the space of quasimetric homeomorphisms of the unit circle factorized by Möbius maps; all Teichmüller spaces have their isometric copies in $\mathbf{T}$.

The canonical complex Banach structure on $\mathbf{T}$ is defined by factorization of the ball of the Beltrami coefficients

$$\text{Belt}(U)_1 = \{\mu \in L_\infty(\mathbb{C}) : \mu|_{U^*} = 0, \|\mu\|_\infty < 1\}$$

(i.e., supported in the upper-half plane), letting $\mu_1, \mu_2 \in \text{Belt}(U)_1$ be equivalent if the corresponding quasiconformal maps w^{μ_1}, w^{μ_2} coincide on $\overline{\mathbb{R}} = \mathbb{R} \cup \{\infty\} = \partial U^*$ (hence, on $\overline{U^*}$). Such μ and the corresponding maps w^μ are called **T-equivalent**. The equivalence classes $[w^\mu]_{\mathbf{T}}$ are in one-to-one correspondence with the Schwarzian derivatives S_w in U^* , which fill a bounded domain in the space $\mathbf{B} = \mathbf{B}(U^*)$ (see **2.1**).

The map $\phi_{\mathbf{T}} : \mu \rightarrow S_{w^\mu}$ is holomorphic and descends to a biholomorphic map of the space $\mathbf{T}$ onto this domain, which we will identify with $\mathbf{T}$. As was mentioned above, it contains as complex submanifolds the Teichmüller spaces of all hyperbolic Riemann surfaces and of Fuchsian groups.

We also define on this ball another equivalence relation, letting $\mu, \nu \in \text{Belt}(U)_1$ be equivalent if $w^\mu(a_j^0) = w^\nu(a_j^0)$ for all j and the homeomorphisms w^μ, w^ν are homotopic on the punctured sphere $X_{\mathbf{a}^0}$. Let us call such μ and ν *strongly n -equivalent*.

Lemma 4. *If the coefficients $\mu, \nu \in \text{Belt}(U)_1$ are **T-equivalent**, then they are also strongly n -equivalent.*

The proof of this lemma is given in [6].

In view of Lemmas 1 and 4, the above factorizations of the ball $\text{Belt}(U)_1$ generate (by descending to the equivalence classes) a holomorphic map χ of the underlying space $\mathbf{T}$ into $\mathbf{T}(0, n) = \mathbf{T}(X_{\mathbf{a}^0})$.

This map is a split immersion (has local holomorphic sections), which is a consequence, for example, of the following existence theorem from [11], which we present here as

Lemma 5. *Let D be a finitely connected domain on the Riemann sphere $\widehat{\mathbb{C}}$. Assume that there are a set E of positive two-dimensional Lebesgue measure and a finite number of points $z_1, \dots, z_m$ distinguished in D . Let $\alpha_1, \dots, \alpha_m$ be non-negative integers assigned to $z_1, \dots, z_m$, respectively, so that $\alpha_j = 0$ if $z_j \in E$.*

Then, for a sufficiently small $\varepsilon_0 > 0$ and $\varepsilon \in (0, \varepsilon_0)$, and for any given collection of numbers $w_{sj}, s = 0, 1, \dots, \alpha_j, j = 1, 2, \dots, m$, which satisfy the conditions $w_{0j} \in D$,

$$|w_{0j} - z_j| \leq \varepsilon, \quad |w_{1j} - 1| \leq \varepsilon, \quad |w_{sj}| \leq \varepsilon \quad (s = 0, 1, \dots, \alpha_j, \quad j = 1, \dots, m),$$

there exists a quasiconformal automorphism h of domain D which is conformal on $D \setminus E$ and satisfies

$$h^{(s)}(z_j) = w_{sj} \quad \text{for all } s = 0, 1, \dots, \alpha_j, \quad j = 1, \dots, m.$$

Moreover, the Beltrami coefficient μ_h of h on E satisfies $\|\mu_h\|_\infty \leq M\varepsilon$. The constants ε_0 and M depend only upon the sets D, E and the vectors $(z_1, \dots, z_m)$ and $(\alpha_1, \dots, \alpha_m)$.

3.2. In fact, we have more, given by the following theorem which corrects the assertion of Lemma 3 in [12], as was mentioned in Preamble.

Theorem 2. *The map χ is surjective and generates an open holomorphic map s of the space $\mathbf{T}(0, n) = \mathbf{T}(X_{\mathbf{a}^0})$ into the universal Teichmüller space $\mathbf{T}$ embedding $\mathbf{T}(0, n)$ into $\mathbf{T}$ as a $(n - 3)$ -dimensional submanifold.*

Proof. The surjectivity of χ is a consequence of Lemma 2. To construct s , take a dense subset

$$e = \{x_1, x_2, \dots\} \subset X_{\mathbf{a}^0} \cap \mathbb{R}$$

accumulating to all points of $\mathbb{R}$ and consider the punctured spheres $X_{\mathbf{a}^0}^m = X_{\mathbf{a}^0} \setminus \{x_1, \dots, x_m\}$ with $m > 1$. The equivalence relations on $\text{Belt}(\mathbb{C})_1$ for $X_{\mathbf{a}^0}^m$ and $X_{\mathbf{a}^0}$ generate the corresponding holomorphic map $\chi_m : \mathbf{T}(X_{\mathbf{a}^0}^m) \rightarrow \mathbf{T}(X_{\mathbf{a}^0})$.

Uniformizing the surfaces $X_{\mathbf{a}^0}$ and $X_{\mathbf{a}^0}^m$ by the corresponding torsion free Fuchsian groups Γ_0 and Γ_0^m of the first kind acting discontinuously on $U \cup U^*$ and applying to U^*/Γ_0 and U^*/Γ_0^m the construction from Section 2.3 (forgetting the additional punctures), one obtains similar to (3) the norm preserving isomorphism $\hat{j}_{m,*} : \mathbf{B}(\Gamma_0) \rightarrow \mathbf{B}(\Gamma_0^m)$ by

$$\hat{j}_{m,*}\varphi = (\varphi \circ \hat{j})(\hat{j}')^2,$$

which projects to the surfaces $X_{\mathbf{a}^0}$ and $X_{\mathbf{a}^0}^m$ as the inclusion of the space $Q(X_{\mathbf{a}^0})$ of quadratic differentials corresponding to $\mathbf{B}(\Gamma_0)$ into the space $Q(X_{\mathbf{a}^0}^m)$, and (since the projection η_m has local holomorphic sections) geometrically this relation yields a holomorphic embedding of the space $\mathbf{T}(\Gamma_0)$ into $\mathbf{T}(\Gamma_0^m)$ as an $(n - 3)$ -dimensional submanifold. Denote this embedding by s_m .

To investigate the limit function for $m \rightarrow \infty$, we compose the maps s_m with the canonical biholomorphic isomorphisms

$$\eta_m : \mathbf{T}(X_{\mathbf{a}^0}^m) \rightarrow \mathbf{T}(\Gamma_0^m) = \mathbf{T} \cap \mathbf{B}(\Gamma_0^m) \quad (m = 1, 2, \dots).$$

Then the elements of $\mathbf{T}(\Gamma_0^m)$ are given by

$$\hat{s}_m(X_{\mathbf{a}}) = \eta_m \circ s_m(X_{\mathbf{a}}),$$

and this is a collection of the Schwarzians $S_{f^m}(z)$ corresponding to the points $X_{\mathbf{a}}$ of $\mathbf{T}(X_{\mathbf{a}^0})$. So for any surface $X_{\mathbf{a}}$, we have

$$\hat{s}_m(X_{\mathbf{a}}) = S_{f^m}(z). \quad (7)$$

Each Γ_0^m is the covering group of the universal cover $h_m : U^* \rightarrow X_{\mathbf{a}^0}^m$, which can be normalized (conjugating appropriately Γ_0^m) by $h_m(-i) = -i$, $h'_m(-i) > 0$. Take its fundamental polygon P_m obtained as the union of the circular m -gon in U^* centered at $z = -i$ with the zero angles at the vertices and its reflection with respect to one of the boundary arcs. These polygons increasingly exhaust the half-plane U^* from inside; hence, by the Carathéodory kernel theorem, the maps h_m converge to the identity map locally uniformly in U^* .

Since the set of punctures e is dense on $\mathbb{R}$, it completely determines the equivalence classes $[w^\mu]$ and S_{w^μ} as the points of the universal space $\mathbf{T}$. The limit function $\hat{s} = \lim_{m \rightarrow \infty} \hat{s}_m$ maps $\mathbf{T}(X_{\mathbf{a}^0}) = \mathbf{T}(0, n)$ into the space $\mathbf{T}$ and also is canonically defined by the marked spheres $X_{\mathbf{a}}$.

Similar to (7), the function $\hat{s}$ is represented as the Schwarzian of some univalent function f^n on U^* with quasiconformal extension to $\hat{\mathbb{C}}$ determined by $X_{\mathbf{a}}$. Then, by the well-known property of elements in the functional spaces with sup norms, $\hat{s}$ is holomorphic also in $\mathbf{B}$ -norm on $\mathbf{T}$.

Lemma 5 yields that $\hat{s}$ is a locally open map from $\mathbf{T}(X_{\mathbf{a}^0})$ to $\mathbf{T}$. So the image $\hat{s}(\mathbf{T}(X_{\mathbf{a}^0}))$ is an $(n - 3)$ -dimensional complex submanifold in $\mathbf{T}$ biholomorphically equivalent to $\mathbf{T}(\Gamma_0)$. The proof of Theorem 2 is completed.

The holomorphy property indicated above is based on the following lemma of Earle [4].

Lemma 6. *Let E, T be open subsets of complex Banach spaces X, Y and $B(E)$ be a Banach space of holomorphic functions on E with sup norm. If $\varphi(x, t)$ is a bounded map $E \times T \rightarrow B(E)$ such that $t \mapsto \varphi(x, t)$ is holomorphic for each $x \in E$, then the map φ is holomorphic.*

Holomorphy of $\varphi(x, t)$ in t for fixed x implies the existence of complex directional derivatives

$$\varphi'_t(x, t) = \lim_{\zeta \rightarrow 0} \frac{\varphi(x, t + \zeta v) - \varphi(x, t)}{\zeta} = \frac{1}{2\pi i} \int_{|\xi|=1} \frac{\varphi(x, t + \xi v)}{\xi^2} d\xi,$$

while the boundedness of φ in sup norm provides the uniform estimate

$$\|\varphi(x, t + c\zeta v) - \varphi(x, t) - \varphi'_t(x, t)c\zeta v\|_{B(E)} \leq M|c|^2,$$

for sufficiently small $|c|$ and $\|v\|_Y$.

3.2. Now the desired holomorphic homotopy of $\mathbf{T}(0, n) = \mathbf{T}(X_{\mathbf{a}0})$ into its base point is constructed as follows.

Using the canonical embedding of $\mathbf{T}(0, n)$ in $\mathbf{T}$ via $\mathbf{T}(\Gamma_0)$, we define on the space $\mathbf{T}(\Gamma_0)$ a holomorphic homotopy applying the maps

$$W^\mu = \sigma^{-1} \circ w^\mu \circ \sigma, \quad \mu \in \text{Belt}(U)_1; \quad \sigma(\zeta) = i(1 + \zeta)/(1 - \zeta), \quad \zeta \in \mathbb{D},$$

and $w_t^\mu(z) := w^\mu(z, t) = \sigma \circ W_t^\mu \circ \sigma^{-1}(z)$; then

$$S_{w^\mu}(\cdot, t) = t^2 S_{w^\mu}(\cdot) = t^{-2} (S_{W^\mu} \circ \sigma^{-1})(\sigma')^{-2}.$$

This point-wise equality determines holomorphic map $\eta(\varphi, t) = S_{w_t^\mu} : \mathbf{T} \times \mathbb{D} \rightarrow \mathbf{T}$ with $\eta(\mathbf{0}, t) = \mathbf{0}$, $\eta(\varphi, 0) = \mathbf{0}$, $\eta(\varphi, 1) = \varphi$. It is not compatible with the group Γ_0 ; hence, there are images $\eta(\varphi, t) = S_{w_t^\mu}$ which are located in $\mathbf{T}$ outside of $\mathbf{T}(\Gamma_0)$.

Composition of $\eta(\varphi, t)$ with maps χ and s carries these images to the points of the space $\mathbf{T}(0, n) = \mathbf{T}(X_{\mathbf{a}0})$ and implies the desired holomorphic homotopy of $\mathbf{T}(0, n)$.

4. HOLOMORPHIC CONTRACTIBILITY OF LOW DIMENSIONAL TEICHMÜLLER SPACES (PROOF OF THEOREM 1)

(a) *Case $n = 5$ (dimension two).* It is enough to establish holomorphic contractibility of the space $\mathbf{T}(0, 5) \simeq \mathbf{F}(0, 4)$ for the spheres

$$X_{\mathbf{a}} = \widehat{\mathbb{C}} \setminus \{0, 1, a_1, a_2, \infty\}.$$

The fibers of $\mathbf{T}(0, 5)$ are the spheres with quadruples of punctures $\{0, 1, a_1, \infty\}$.

We start with construction of the needed holomorphic homotopy of the space $\mathbf{T}(0, 5)$ to its base point $X_{\mathbf{a}0}$ and first apply the assertion (a) of Lemma 3 about holomorphic sections over $\mathbf{T}(0, 4)$. It implies for any point

$$x = (S_{w^{\mu_1}}, w^{\mu_1}(\widehat{a}_{n-3}^0)) \in \mathbf{T}(0, 5)$$

a unique holomorphic section $s : \mathbf{T}(0, 5) \rightarrow \mathbf{T}(0, 4)$ with $s(\pi_1(x)) = x$. This section has a common point with each fiber $\pi^{-1}(x)$ over $\mathbf{T}(0, 4)$.

Since $\mathbf{T}(0, 4)$ is (up to a biholomorphic equivalence) a simply connected bounded Jordan domain $D \subset \mathbb{C}$ containing the origin, there is a holomorphic isotopy $h(\zeta, t) : D \times [0, 1] \rightarrow D$ with $h(\zeta, 0) = \zeta$, $h(z, 1) = 0$. Using this isotopy, we define a homotopy $h_1(\varphi, t)$ on $\mathbf{T}(0, 5)$, which carries each point $x = (S_{w^\mu}, w^\mu(\widehat{a}_2^0)) \in \mathbf{T}(0, 5)$ to its image on the section s passing from x , that is

$$h_1(\varphi, w^\mu(\widehat{a}_2^0)) = (h(\varphi), \widetilde{a}_2), \quad \varphi = S_{w^\mu}, \quad \mu \in \text{Belt}(\mathbb{C})_1, \quad (8)$$

where $\widetilde{a}_2$ is the common point of the fiber $h(\varphi)$ and the selected section s . The holomorphy of this homotopy in variables $x = (S_{w^\mu}, w^\mu(\widehat{a}_2^0))$ for any fixed $t \in [0, 1]$ follows from Lemmas 1, 2 and the Bers isomorphism theorem. The limit map

$$h_1^*(x) = \lim_{t \rightarrow 1} h_1(x, t),$$

carries each fiber $w^\mu(U)$, to the initial half-plane U .

There is a canonical holomorphic isotopy

$$h_2(\zeta, t) : U \times [0, 1] \rightarrow U \quad (9)$$

of U into its point corresponding to the origin of $\mathbf{T}(0, 5)$. Now put $\mathbf{h}(x, t)$ to be equal to $h_1(x, 2t)$ for $t \leq 1/2$ and equal to $h_2(x, 2t - 1)$ for $x = \zeta \in U$ and $1/2 \leq t \leq 1$.

This function is holomorphic in $x \in \mathbf{T}(0, 5)$ for any fixed $t \in [0, 1]$ and hence provides the desired holomorphic homotopy of the space $\mathbf{T}(0, 5)$ into its base point.

(b) *Case $n = 6$ (dimension three).* This case is more complicate.

We prescribe to each ordered sextuple $\mathbf{a} = \{0, 1, a_1, a_2, a_3, \infty\}$ of distinct points the corresponding punctured sphere

$$X_{\mathbf{a}} = \widehat{\mathbb{C}} \setminus \{0, 1, a_1, a_2, a_3, \infty\} \quad (10)$$

and the two-sheeted closed hyperelliptic surface $\widehat{X}_{\mathbf{a}}$ of genus two with the branch points $0, 1, a_1, a_2, a_3, \infty$. The corresponding Teichmüller spaces $\mathbf{T}(0, 6)$ and $\mathbf{T}(2, 0)$ coincide up to a natural biholomorphic isomorphism. Note also that the collections $\mathbf{a} = \{0, 1, a_1, a_2, a_3, \infty\}$ provide the local complex coordinates on each from the spaces $\mathbf{T}(0, 6)$ and $\mathbf{T}(2, 0)$.

In view of the symmetry of hyperelliptic surfaces, it suffices to deal with the Beltrami differentials $\mu d\bar{z}/dz$ on $\widehat{X}_{\mathbf{a}}$, which are compatible with conformal involution $J_{\mathbf{a}}$ of $\widehat{X}_{\mathbf{a}}$, hence, satisfy $\mu(J_{\mathbf{a}}z) = \mu(z)J'_{\mathbf{a}}/\overline{J_{\mathbf{a}}}$. In other words, these μ are obtained by lifting to $\widehat{X}_{\mathbf{a}}$ of the Beltrami coefficients on $X_{\mathbf{a}}$. This extends Lemma 2 and its consequences on holomorphy in the neighborhoods of the boundary interpolation points to the corresponding two-sheeted disks on hyperelliptic surfaces.

Fix a base point of $\mathbf{T}(2, 0)$ determining a Fuchsian group Γ for which $\mathbf{T}(\Gamma) = \mathbf{T}(2, 0)$. The corresponding Teichmüller curve $\mathcal{V}(2, 0)$ is a 4-dimensional complex analytic manifold with projection $\pi_1 : \mathcal{V}(2, 0) \rightarrow \mathbf{T}(2, 0)$ onto $\mathbf{T}(2, 0)$ such that for every $\varphi \in \mathbf{T}(2, 0)$ the fiber $\pi_1^{-1}(\varphi)$ is a hyperelliptic surface, determined by φ (see 2.4).

Due to assertion (b) of Lemma 3, this curve has for any point

$$\widehat{X}_{\mathbf{a}} = (S_{w^{\mu_1}}, w^{\mu_1}(\widehat{a}_{n-3}^0)) \in \mathbf{T}(2, 0)$$

six distinct holomorphic sections $\widehat{s}_1, \dots, \widehat{s}_6$, corresponding to the Weierstrass points of the surface $X_{\mathbf{a}}$, with $\widehat{s}_j(\pi_1(X_{\mathbf{a}})) = X_{\mathbf{a}}$, and either from these sections has one common point with every fiber over $\mathbf{T}(2, 0)$. We lift these sections to the Bers fiber space $\mathcal{F}(\Gamma)$ covering $\mathcal{V}(2, 0)$.

As was mentioned in 2.4, these sections are generated by the space $\mathcal{F}(\Gamma') = \mathcal{F}(\Gamma)$ corresponding to the extension Γ' of Γ , for which Γ is a subgroup of index two. Every section $\widehat{s}_j$ acts on $\mathbf{T}(\Gamma')$ via (6), where z_0 is now the corresponding Weierstrass point of hyperelliptic surface $\widehat{X}_{\mathbf{a}}$, and $\widehat{s}_j$ is compatible with the action (2) of the Bers isomorphism.

Thus each $\widehat{s}_j$ descends to a holomorphic map $s_j : \mathbf{T}(0, 6) \rightarrow \mathcal{V}(0, 6)$ of the underlying space $\mathbf{T}(0, 6)$ for the punctured spheres (10). We choose one from these maps and denote it by s .

The features of sections $\widehat{s}_j$ provide that the descended map s also determines for each point $z_0 \in X_{\mathbf{a}}$ its unique image on every fiber $w^{\mu}(X_{\mathbf{a}})$ with $\mu \in \text{Belt}(X_{\mathbf{a}})_1$, and this image is the point $w^{\mu}(z_0)$.

The next preliminary construction consists of embedding the space $\mathbf{T}(0, 5)$ into $\mathbf{T}(0, 6)$, using the forgetting map (3). Its image $j_*\mathbf{T}(0, 5)$ is a connected submanifold in $\mathbf{T}(0, 6)$, and the corresponding fibers of the curve $\mathcal{V}(0, 6)$ over the points $j_*\varphi \in j_*\mathbf{T}(0, 5)$ are the surfaces $w^{j_*\mu}(X_{\mathbf{a}})$ with $j_*\mu(z) = \mu(\widehat{j}(z))\widehat{j}'(z)/\overline{\widehat{j}'(z)}$. The covering domains $w^{j_*\mu}(U)$ over these surfaces fill a submanifold $\widehat{\mathbf{T}}(0, 7) \subset \mathbf{T}(0, 7)$, which is biholomorphically equivalent to the space $\mathbf{T}(0, 6)$.

Using the biholomorphic equivalence of the space $\mathbf{T}(0, 5)$ to its image $j_*\mathbf{T}(0, 5)$ in $\mathbf{T}(0, 6)$, we carry over to $j_*\mathbf{T}(0, 5)$ the result of the previous step (a) on holomorphic contractibility of $\mathbf{T}(0, 5)$, which provides a holomorphic homotopy

$$h(j_*\varphi, t) : j_*\mathbf{T}(0, 5) \times [0, 1] \rightarrow j_*\mathbf{T}(0, 5) \quad \text{with} \quad h(j_*\varphi, 0) = j_*\varphi, \quad h(j_*\varphi, 1) = \mathbf{0} \quad (11)$$

(here $\mathbf{0}$ stands for the origin of $j_*\mathbf{T}(0, 5)$).

Now we may construct the desired holomorphic homotopy of $\mathbf{T}(0, 6)$ contracting this space to its origin.

First, regarding $\mathbf{T}(0, 6)$ as the Bers fiber space $\mathcal{F}(0, 5)$ over $\mathbf{T}(0, 5)$ (whose fibers are the covers of surfaces $X_{\mathbf{a}'}$ with collections of five punctures $\mathbf{a}' = (0, 1, a_1, a_2, \infty)$), we apply the homotopy (11) and define for any pair $x = (j_*\varphi, z)$ with $\varphi \in \mathbf{T}(0, 5)$ and $z \in X_{\mathbf{a}}$ the map

$$\tilde{h}_1((j_*\varphi, z), t) = (h(j_*\varphi, t), w_t^{j_*\mu}(z)), \quad \varphi \in \mathbf{T}(0, 5), \quad (12)$$

noting that the image point $w_t^{j_*\mu}(z)$ is uniquely determined on the surface $w^{h(j_*\mu)}(X_{\mathbf{a}})$ by the map s , as was indicated above.

The pairs $(j_*\varphi, z)$ are located in the space $\mathcal{F}(0, 6)$ and fill there a three-dimensional submanifold $\tilde{\mathbf{T}}(0, 6)$ biholomorphically equivalent to $\mathbf{T}(0, 6)$.

The homotopy (12) is well defined on $\tilde{\mathbf{T}}(0, 6) \times [0, 1]$ and contracts the set $\tilde{\mathbf{T}}(0, 6)$ into the fiber $\tilde{U}$ over the base point. It is holomorphic with respect to the space variable $x = (j_*\varphi, z)$ for any fixed $t \in [0, 1]$ and continuous in both variables.

In view of biholomorphic equivalence of $\tilde{\mathbf{T}}(0, 6)$ to $\mathbf{T}(0, 6)$, (12) generates a holomorphic homotopy $h_1(x, t)$ of the space $\mathbf{T}(0, 6)$ onto the initial fiber (half-plane) U over the origin of $\mathbf{T}(0, 5)$.

It remains to combine this homotopy h_1 with the additional homotopy (9) of U into its point corresponding to the origin of $\mathbf{T}(0, 6)$. This provides the desired homotopy h and completes the proof of Theorem 1.

REFERENCES

- [1] L. Bers, *Spaces of Riemann surfaces*, Proc. Internat. Congress Math. 1958, Cambridge Univ. Press, NY, 1960, pp. 349-361.
- [2] L. Bers, *Fiber space over Teichmüller spaces*, Acta Math. **130** (1973), 89-126.
- [3] J.G. Clunie, D.J. Hallenbeck and T.H. MacGregor, *A peaking and interpolation problem for univalent functions*, J. Math. Anal. Appl. **111** (1985), 559-570.
- [4] C.J. Earle, *On quasiconformal extensions of the Beurling-Ahlfors type*, Contribution to Analysis, Academic Press, New York, 1974, pp. 99-105.
- [5] C.J. Earle and I. Kra, *On holomorphic mappings between Teichmüller spaces*, Contributions to Analysis, Academic Press, New York, 1974, pp. 107-124.
- [6] F.P. Gardiner, *Teichmüller Theory and Quadratic Differentials*, Wiley, New York, 1987.
- [7] E.A. Gorin, Zapiski Nauchn. Seminarov Leningrad. Otdel. Mat. Inst. Steklov. (LOMI) **81** (1978), 58-61 (Problem 9.2); Engl. transl. in J. Soviet Math. **26** (1984), no. 5.
- [8] A. Hirschowitz, *A propos de principe d'Oka*, C.R. Acad. Sci. Paris **272** (1971), A792-A794.
- [9] J. Hubbard, *Sur le non-existence de sections analytiques a la courbe universelle de Teichmüller*, C. R. Acad. Sci. Paris Ser A-B. **274** (1972), A978-A979.
- [10] I. Kaliman, *Holomorphic universal covering spaces of polynomials without multiple roots*, Funktsional Anal. i Prilozh. **71** (1975), no. 1, 71 (Russian).
- [11] S. L. Krushkal, *Quasiconformal Mappings and Riemann Surfaces*, Wiley, New York, 1979.
- [12] S.L. Krushkal, *Holomorphic contractibility of Teichmüller spaces*, Complex Anal. Oper. Theory **13(6)** (2019), 2829-2838.
<https://doi.org/10.1007/s11785-018-0848-8>
- [13] S. Nag, *The Complex Analytic Theory of Teichmüller Spaces*, Wiley, 1988.
- [14] D.B. Patterson, *The Teichmüller spaces are distinct*, Proc. Amer. Math. Soc. **35** (1972), 179-182.
- [15] M.G. Zaidenberg and V.Ya. Lin, *On bounded domains of holomorphy that are not holomorphically contractible*, Soviet Math. Dokl. **20** (1979), 1262-1266.

- [16] M.G. Zaidenberg and V.Ya. Lin, *The finiteness theorems for holomorphic maps*, Current problems in mathematics. Fundamental directions, vol. 9. Itogi Nauki i Tekniki, Vsesoyuzn. Inst. Nauchn. i Technich. Inform., 1986, pp. 127-193 (Russian).

Department of Mathematics, Bar-Ilan University, 5290002 Ramat-Gan, Israel
and Department of Mathematics, University of Virginia, Charlottesville, VA 22904-4137, USA