arXiv:2104.01621v2 [math.GR] 13 Apr 2021

PROPERTY (T) IN K-GONAL RANDOM GROUPS

MURPHYKATE MONTEE

Abstract. The k-gonal models of random groups are defined as the quotients of free groups on n generators by cyclically reduced words of length k. As k tends to infinity, this model approaches the Gromov density model. In this paper we show that for any fixed $d_0 \in (0,1)$, if positive k-gonal random groups satisfy Property (T) with overwhelming probability for densities $d > d_0$, then so do nk-gonal random groups, for any $n \in \mathbb{N}$. In particular, this shows that for densities above $1/3$, groups in $3k$-gonal models satisfy Property (T) with probability 1 as n approaches infinity.

1. Introduction

The study of random groups is one way to approach questions of the form 'Does a *typical* group have property P?' To answer this question, one needs to choose a model of random groups, and then calculate the probability that property P is satisfied. Generally speaking we are interested in the asymptotic behavior of this probability as some constraint goes to infinity.

There are many models of random groups (an excellent introduction to this topic is [Oll05]), the most studied of which is the Gromov density model, first introduced by Gromov in [Gro93]. In this model, a group presentation is chosen uniformly at random by fixing the number of generators, and choosing cyclically reduced relators of a certain length. The number of relators chosen is controlled by a fixed constant called the *density*. One then analyzes the probability of a property being satisfied as both the length and the number of relators goes to infinity. This probability generally tends to 0 or 1, and in many important cases there is a *sharp threshold* d_0, so that for densities above d_0 the probability is 1, and below d_0 the probability is 0 (or vice versa). Some examples of this phase shift behavior include properties such as hyperbolicity [Gro93; Oll04], small cancellation [Gro93], and satisfying Dehn's algorithm [Alo+91].

It is known that there is a sharp threshold for Kazhdan's Property (T); however its precise value is unknown. Property (T) was introduced by Kazhdan in [Kaž67] as a tool to study lattices in Lie groups, but turned out to be important in several areas of mathematics and computer science, especially in the study of expander graphs (for more information, see [Lub94]). In [Żuk03], Żuk stated and sketched a proof that for $d > 1/3$, random groups

Date: March 2021.




FIGURE 1. Possible values of the sharp threshold for Property (T) in the k-gonal models. Red indicates (some of the) results from this paper.

in the Gromov density model satisfy, with overwhelming probability, Property (T). The proof was completed in [KK13].

Similarly, we do not yet know a sharp threshold for *cubulation* of random groups (i.e. admitting a cocompact action on a CAT(0) cube complex without a global fixed point). In [MP15], Mackay-Przytycki built on the work of Ollivier-Wise [OW11] to show that at densities $d < 5/24$, Gromov random groups are, with probability 1, cubulated. This bound was improved to $d < 3/14$ in [Mon] using a generalization of the methods in [MP15].

For infinite hyperbolic groups, being cubulated and having Property (T) are mutually exclusive, so together these results show that the sharp threshold at which Gromov random groups fail to satisfy Property (T) must be between $1/3$ and $3/14$.

In order to prove his theorem, Żuk introduced a new model of random groups, called the *triangular model*. This was extended to the *square model* in [Odr19], and in [AR20] this was further generalized this to a class of models, here called the *k-gonal models*. Roughly speaking, as k approaches infinity the behavior of k-gonal random groups should reflect the behavior of Gromov random groups. Odrzygóźdź showed that in the 6-gonal model, $d = 1/3$ is the sharp threshold at which the probability of a group satisfying Property (T) switches from 1 to 0. Figure 1 illustrates what is currently known about values of the sharp threshold for Property (T) in the k-gonal models.

In this paper, we show that show that any upper bound on the sharp threshold for Property (T) in the positive k-gonal model gives an upper

bound in the jk-gonal model for all $j \in \mathbb{N}$. More precisely, we prove the following theorem:

Theorem 1.1. *Let d_0 be some fixed value in $(0,1)$, and suppose that with overwhelming probability a random group at density $d > d_0$ in $\mathcal{M}_k^+(n,d)$ satisfies Property (T). Then with overwhelming probability, for any $d' \in (d_0, d)$ and for any $j \in \mathbb{N}$, a random group in $\mathcal{M}_{jk}(n,d')$ or $\mathcal{M}_{jk}^+(n,d')$ satisfies Property (T).*

In particular, this improves the known upper bounds in the $3j$-gonal models from $1/2$ to $1/3$. It also shows that to find upper bounds for the sharp threshold of Property (T), it suffices to investigate the *positive* k-gonal models. Further, this provides evidence supporting the following conjecture:

Conjecture 1.2. *Let $d_{(T)}(k)$ denote the sharp threshold for Property (T) in the k-gonal model. Then $\lim_{k\to\infty} d_{(T)}(k)$ exists, and is equal to the sharp threshold of the Gromov model.*

Acknowledgements. I would like to thank Danny Calegari and Tomasz Odrzygóźdź for valuable comments and feedback. This research was partially supported by a Clare Boothe Luce Professorship in Mathematics from the Henry Luce Foundation.

2. Background

We begin by defining the k-gonal model of random groups.

Definition 2.1 ([AR20])**.** Fix a set S of cardinality n. The *k-gonal model of random groups*, denoted $\mathcal{M}_k(n,d)$, is the set of group presentations $\langle S|R\rangle$ such that R is a set of $(2n-1)^{kd}$ cyclically reduced words on $S \cup S^{-1}$ of length k, along with the uniform probability measure. We say that a k-gonal random group at density d satisfies a property P *with overwhelming probability*, abbreviated w.o.p, if

$$\lim_{n\to\infty} \mathbb{P}(\text{the group defined by a presentation in } \mathcal{M}_k(n,d) \text{ satisfies P}) = 1.$$

A related family of models of random groups restricts possible relators to *positive* words, i.e. words on S but not on S^{-1}. These are called the *positive k-gonal models of random groups* and are denoted $\mathcal{M}_k^+(n,d)$.

We now define (Kazhdan's) Property (T) for discrete, finitely generated groups. The notation is suggestive - the idea of Property (T) is that the trivial representation of the group must be isolated in some metric. More formally, we can define this property via (almost) invariant vectors: Let Γ be a finite generated group with finitely generating set S. If $\mathcal{H}$ is a Hilbert space and $\pi : \Gamma \to U(\mathcal{H})$ is a unitary representation of Γ on $\mathcal{H}$, then π has *almost invariant vectors* if, for every $\varepsilon > 0$, there exists some $u_\varepsilon \in \mathcal{H}$ such that for every $s \in S$, $\|\pi(s)u_\varepsilon - u_\varepsilon\| < \varepsilon\|u_\varepsilon\|$. A vector $u \in \mathcal{H}$ is *invariant* if $\pi(g)(u) = u$ for every $g \in \Gamma$.

Definition 2.2 ([Kaž67])**.** A finitely generated group Γ has *Property (T)* if, for every $\mathcal{H}$ and π, the following holds: if π has almost invariant vectors, then π has a nonzero invariant vector.

The first results towards demonstrating Property (T) in random groups were first stated in [Żuk03], and fully proved in [KK13]. The proof involved passing between several different models of random groups and analyzing spectral properties of random graphs.

Theorem 2.3 ([KK13] Theorem 3.14)**.** *A random group in $\mathcal{M}_3^+(n, d')$ with density $d' > 1/3$ satisfies Property (T) with overwhelming probability.*

In fact, they also showed that this is true in $\mathcal{M}_3(n, d)$ and in the Gromov model of random groups (at the same density).

Remark 2.4. Property (T) is preserved by surjections and finite extensions (see [KK13] Remark 2.10 and [BHV08] Theorem 1.7.1). In other words, if H is a group with Property (T) and H surjects onto H', then H' has Property (T); and if $H \overset{f.i.}{\leq} H''$ is a sub-group of finite index, then H'' has Property (T).

The tempting argument is to say if $G \in \mathcal{M}_{jk}(n, d)$ then $G = \langle S|R\rangle$ and any relator in R can be interpreted as a word of length k over the alphabet of words of length j on $S \cup S^{-1}$, denoted $|W_j|$. Then G is in $\mathcal{M}_k(|W_j|, d)$. However, as is pointed out in [KK13], problems arise around the requirement that relators be *reduced*. In particular, a word of length kj is a word of length k in the alphabet of words of length j; however, a reduced word of length k in the alphabet of words of length j might not be reduced as a word of length jk. So we must first consider an intermediate group, G^+, whose relators are *positive* words on S. This removes the reduction problem, and we can safely interpret this group as a random group in $\mathcal{M}_k^+(n, d)$.

The method of proof closely follows the argument used to prove Theorem 4.3 in [Odr19], which shows that groups in the 6-gonal model w.o.p. satisfy Property (T). In particular, to prove that a random group $G \in \mathcal{M}_{jk}(n, d)$ satisfies Property (T), we construct two groups G^+ and Γ so that

$$\Gamma \overset{\phi}{\twoheadrightarrow} \phi(\Gamma) \overset{f.i.}{\leq} G^+ \twoheadrightarrow G,$$

and then show that Γ satisfies Property (T).

There are two parts to this proof. The first is showing that $\phi(\Gamma)$ is finite index in G^+. We find a finite set of coset representatives to prove this explicitly, generalizing the method used in [Odr19]. The second is in showing that the group Γ can be interpreted as a typical representative of random groups in $\mathcal{M}_k^+(m, d')$ for $d' > 1/3$. For this part of the argument, we mimic the method developed in the proof of Theorem A in [KK13].

3. Property (T) in jk-Gonal Models

Fix a generating set S of cardinality n. Let W^+ be the set of positive words on S and let W_j be the words of length j on S. For any set R of words on S, let $R^+ = R \cap W^+$. Then for any group $G = \langle S|R\rangle$ in $\mathcal{M}_{jk}(n,d)$, there is a group $G^+ = \langle S|R^+\rangle$ that surjects onto G.

Since G is in the jk-gonal model, every word in R and R^+ has length jk. There is a natural injective map $\phi : W_j^+ \to F(S)$, where $F(S)$ denotes the free group on S, given by the labeling on each word in W_j^+. Define $R_k^+ = \phi^{-1}(R^+)$, and notice that every word in R_k^+ has length k over W_j^+. Let $\Gamma = \langle W_j^+|R_k^+\rangle$. Then ϕ induces a homomorphism, also denoted ϕ, mapping $\Gamma \to G^+$.

Note also that $|R_k^+| = |\phi(R_k^+)| = |R^+|$, so if $G^+ \in \mathcal{M}_{jk}^+(n,d')$, then $\Gamma \in \mathcal{M}_k^+(|W_j^+|,d')$. Furthermore, this map takes the uniform probability distribution on $\mathcal{M}_{jk}^+(n,d')$ to the uniform probability distribution on $\mathcal{M}_k^+(|W_j^+|,d')$, because ϕ induces a 1-1 map on possible sets of relators.

We will show that $\phi(\Gamma)$ is a finite index subgroup of G^+ by finding an explicit set of coset representatives of $\phi(\Gamma)$ in G^+. We note that an element of G^+ is in $\phi(\Gamma)$ if and only if it is expressible as a string of words of length k, all of which are positive or the inverse thereof. We do this by inserting trivial words that allow us to group positive and negative letters into subwords of length j.

In the following proof, for any word w, $|w|_j$ denotes the word length of w modulo j.

Lemma 3.1. *$\phi(\Gamma)$ is a finite index subgroup of G^+.*

Proof. Let w be a word on $S \cup S^{-1}$. Without loss of generality, assume that the first letter of w is in S^{-1}. Then w can be written as

$$w = A_\ell a_\ell \cdots A_2 a_2 A_1 a_1,$$

where $\{a_i\}$ are positive words on S, $\{A_i\}$ are negative words on S, and a_1 might be empty.

Let x be a positive word of length $j - |a_1|_j$ and let y be a positive word of length $j - |A_1 x^{-1}|_j$. Then we have

$$w = A_\ell a_\ell \cdots A_2 a_2 y y^{-1} A_1 x^{-1} x a_1.$$

Note that xa is a positive word with length divisible by j, and $y^{-1}A_1x^{-1}$ is a negative word with length divisible by j. Therefore $w = A_\ell a_\ell \cdots A_2 a_2' u$, where $u \in \phi(\Gamma)$. By induction, we see that $w \in A_\ell' \phi(\Gamma)$ where A_ℓ' is a positive word on S. Then $A_\ell' = az$, where j divides $|z|$ and $|a| < j$, so $z \in \phi(\Gamma)$. Therefore $w = azv$, where $zv \in \phi(\Gamma)$ and $|a| < j$.

Thus the (finite) set of words of length $< j$ contains a transversal of $\phi(\Gamma)$ in G^+, and $\phi(\Gamma)$ has finite index in G^+. □

We are now ready to prove Theorem 1.1:

Theorem 1.1. *Let d_0 be some fixed value in $(0,1)$, and suppose that with overwhelming probability a random group at density $d > d_0$ in $\mathcal{M}_k^+(n,d)$ satisfies Property (T). Then with overwhelming probability, for any $d' \in (d_0, d)$ and for any $j \in \mathbb{N}$, a random group in $\mathcal{M}_{jk}(n,d')$ or $\mathcal{M}_{jk}^+(n,d')$ satisfies Property (T).*

Proof. We first show that a random group G^+ in $\mathcal{M}_{kj}^+(n,d)$ has Property (T). Let $m = |W_j^+|$. By the construction above, given such a G^+ we can find a group Γ in $\mathcal{M}_k^+(m,d)$ and a map $\phi : \Gamma to G^+$. The distribution induced on $\mathcal{M}_k^+(m,d)$ in this way is uniform and by Lemma 3.1 $\phi(\Gamma)$ is a finite index subgroup of G^+. Since Γ satisfies Property (T) with overwhelming probability, by Remark 2.4, so does G^+.

Now let $G = \langle S|R\rangle$ be a group in $\mathcal{M}_{kj}(n,d)$. Note that $|W_j^+| > \frac{1}{2^j}|W_j|$. Therefore, the expected number of elements of R^+ is larger than $\frac{1}{2^j}|R|$. Furthermore, for any $d' < d$ there is some $N_{d'}$ so that for $n > N_{d'}$, we have

$$|R^+| > \frac{1}{2^j}|R| = \frac{1}{2^j}(2n-1)^{kjd} > (2n-1)^{kjd'}.$$

Fix $d' \in (d_0, d)$. If we enumerate the elements of W_j^+ and take R' to be the set of the first $(2n-1)^{kjd'}$ elements of W_j^+ which are in R, we can find a group $G^+\langle S|R'\rangle \in \mathcal{M}_{kj}^+(n,d')$ that surjects onto G. This gives a uniform distribution on $\mathcal{M}_{kj}^+(n,d')$, so w.o.p G^+ satisfies Property (T), and by Remark 2.4 so does G. □

By Theorem 3.14 in [KK13], with overwhelming probability random groups in $\mathcal{M}_3^+(n,d)$ satisfy Property (T) at density $d > 1/3$. By applying Theorem 1.1 we get the following result.

Corollary 3.2. *With overwhelming probability, a random group at density $d > 1/3$ in $\mathcal{M}_{3k}(n,d)$ or $\mathcal{M}_{3k}^+(n,d)$ satisfies Property (T).*

References

[Alo+91] J. M. Alonso et al. "Notes on word hyperbolic groups". In: *Group theory from a geometrical viewpoint (Trieste, 1990)*. Edited by Short. World Sci. Publ., River Edge, NJ, 1991, pp. 3–63.

[AR20] Calum J. Ashcroft and Colva M. Roney-Dougal. "On random presentations with fixed relator length". In: *Comm. Algebra* 48.5 (2020), pp. 1904–1918. ISSN: 0092-7872. DOI: 10.1080/00927872.2019.1710161. URL: https://doi.org/10.1080/00927872.2019.1710161.

[BHV08] Bachir Bekka, Pierre de la Harpe, and Alain Valette. *Kazhdan's property (T)*. Vol. 11. New Mathematical Monographs. Cambridge University Press, Cambridge, 2008, pp. xiv+472. ISBN: 978-0-521-88720-5. DOI: 10.1017/CBO9780511542749. URL: https://doi.org/10.1017/CBO9780511542749.

[Gro93] M. Gromov. "Asymptotic invariants of infinite groups". In: *Geometric group theory, Vol. 2 (Sussex, 1991)*. Vol. 182. London Math. Soc. Lecture Note Ser. Cambridge Univ. Press, Cambridge, 1993, pp. 1–295.

[Kaž67] D. A. Každan. "On the connection of the dual space of a group with the structure of its closed subgroups". In: *Funkcional. Anal. i Priložen.* 1 (1967), pp. 71–74. ISSN: 0374-1990.

[KK13] Marcin Kotowski and Michał Kotowski. "Random groups and property (T): Żuk's theorem revisited". In: *J. Lond. Math. Soc. (2)* 88.2 (2013), pp. 396–416. ISSN: 0024-6107. DOI: 10.1112/jlms/jdt024. URL: https://doi.org/10.1112/jlms/jdt024.

[Lub94] Alexander Lubotzky. *Discrete groups, expanding graphs and invariant measures*. Vol. 125. Progress in Mathematics. With an appendix by Jonathan D. Rogawski. Birkhäuser Verlag, Basel, 1994, pp. xii+195. ISBN: 3-7643-5075-X. DOI: 10.1007/978-3-0346-0332-4. URL: https://doi.org/10.1007/978-3-0346-0332-4.

[Mon] MurphyKate Montee. "Cubulating random groups at densities less than 3/14". In preparation.

[MP15] John M. Mackay and Piotr Przytycki. "Balanced walls for random groups". In: *Michigan Math. J.* 64.2 (2015), pp. 397–419. ISSN: 0026-2285. DOI: 10.1307/mmj/1434731930. URL: https://doi.org/10.1307/mmj/1434731930.

[Odr19] Tomasz Odrzygóźdź. *Bent walls for random groups in the square and hexagonal model*. 2019. arXiv: 1906.05417 [math.GR].

[Oll04] Y. Ollivier. "Sharp phase transition theorems for hyperbolicity of random groups". In: *Geom. Funct. Anal.* 14.3 (2004), pp. 595–679. ISSN: 1016-443X. DOI: 10.1007/s00039-004-0470-y. URL: https://doi.org/10.1007/s00039-004-0470-y.

[Oll05] Yann Ollivier. *A January 2005 invitation to random groups*. Vol. 10. Ensaios Matemáticos [Mathematical Surveys]. Sociedade Brasileira de Matemática, Rio de Janeiro, 2005, pp. ii+100. ISBN: 85-85818-30-1.

[OW11] Yann Ollivier and Daniel T. Wise. "Cubulating random groups at density less than 1/6". In: *Trans. Amer. Math. Soc.* 363.9 (2011), pp. 4701–4733. ISSN: 0002-9947. DOI: 10.1090/S0002-9947-2011-05197-4. URL: https://doi.org/10.1090/S0002-9947-2011-05197-4.

[Żuk03] A. Żuk. "Property (T) and Kazhdan constants for discrete groups". In: *Geom. Funct. Anal.* 13.3 (2003), pp. 643–670. ISSN: 1016-443X. DOI: 10.1007/s00039-003-0425-8. URL: https://doi.org/10.1007/s00039-003-0425-8.

Department of Mathematics and Statistics, Carleton College
Email address: murphykate.montee@gmail.com