

THE BRAUER GROUP OF A VALUATION RING

VIVEK SADHU

ABSTRACT. Let V be a valuation ring and K be its field of fraction. We show that the canonical map $\mathrm{Br}(V) \rightarrow \mathrm{Br}(K)$ is injective.

1. INTRODUCTION

In recent years, it has been observed by various authors that a valuation ring (not necessarily Noetherian) behaves almost like regular ring in many situations. One can see [8] for some K -theoretic evidence. In this article, we give some evidence for Brauer groups. The Brauer group $\mathrm{Br}(R)$ of a commutative ring R consists of Morita equivalence classes of Azumaya algebras over R (see [1], [10]). The group operation on $\mathrm{Br}(R)$ is \otimes_R . An element of $\mathrm{Br}(R)$ is represented by a class $[A]$, where A is an Azumaya algebra over R .

A classical result of Auslander and Goldman says that for a regular noetherian domain R with the field of fraction K , the canonical map $\mathrm{Br}(R) \rightarrow \mathrm{Br}(K)$ sending $[A]$ to $[A \otimes_R K]$ is injective (see Theorem 7.2 of [1]). We observe that the same is true for valuation rings (see below Theorem 1.1). A subring V of a field K is said to be valuation ring if for each nonzero element $a \in K$, either $a \in V$ or $a^{-1} \in V$. Clearly, K itself a valuation ring and we call it a trivial valuation ring of K . If $V \neq K$ then we say that V is a nontrivial valuation ring of K . Every valuation ring is a local normal domain. If \mathfrak{p} is a prime ideal of a valuation ring V then the quotient V/\mathfrak{p} and the localization $V_{\mathfrak{p}}$ both are valuation rings. The rank of a valuation ring is equal to its krull dimension. Here is our main result.

Theorem 1.1. *Let V be a valuation ring with the field of fraction K . Then the canonical map $\mathrm{Br}(V) \rightarrow \mathrm{Br}(K)$ is injective.*

Acknowledgement: The author would like to thank Prof. Jean-Louis Colliot Thélène for his valuable comments on the first draft of this article.

Date: December 23, 2021.

2020 Mathematics Subject Classification. 14F20, 14F22, 13F30.

Key words and phrases. Brauer groups, Valuation rings, Étale cohomology.

Author was supported by SERB-DST MATRICS grant MTR/2018/000283.

2. ONE DIMENSIONAL CASE

In this section, we prove Theorem 1.1 in the case when V is a one dimensional valuation ring with the field of fraction K . Throughout this section, let V be a one dimensional valuation ring with the field of fraction K and $f : \text{Spec}(K) \rightarrow \text{Spec}(V)$ be the map of schemes induced by the inclusion $V \hookrightarrow K$. Given a ring R , R^\times always denotes the group of units.

The Leray spectral sequence of the map $f : \text{Spec}(K) \rightarrow \text{Spec}(V)$ gives

$$(2.1) \quad \begin{aligned} 0 \rightarrow H_{et}^1(\text{Spec}(V), f_*\mathbb{G}_{m,K}) &\rightarrow H_{et}^1(\text{Spec}(K), \mathbb{G}_{m,K}) \rightarrow H_{et}^0(\text{Spec}(V), R^1f_*\mathbb{G}_{m,K}) \\ &\rightarrow H_{et}^2(\text{Spec}(V), f_*\mathbb{G}_{m,K}) \rightarrow H_{et}^2(\text{Spec}(K), \mathbb{G}_{m,K}), \end{aligned}$$

where \mathbb{G}_m denotes the étale unit sheaf. We know $H_{et}^1(\text{Spec}(K), \mathbb{G}_{m,K}) \cong \text{Pic}(K) = 0$ and $H_{et}^2(\text{Spec}(K), \mathbb{G}_{m,K}) \cong \text{Br}(K)$. The stalk of $R^1f_*\mathbb{G}_{m,K}$ at a geometric point \bar{v} is $H_{et}^1(\text{Spec}(K_{\bar{v}}), \mathbb{G}_{m,K_{\bar{v}}}) \cong \text{Pic}(K_{\bar{v}})$ where $K_{\bar{v}} = K \otimes_V \mathcal{O}_{V,\bar{v}}$. Since $\dim(V) = 1$, $\mathcal{O}_{V,\bar{v}}$ is either K^{sep} or V^{sh} , where K^{sep} is any separable closure of K . Moreover, V^{sh} is a normal domain of dimension one (see [12, Tag 07QL]). Thus, $K_{\bar{v}}$ is either K^{sep} or the field of fraction of V^{sh} . Therefore, $R^1f_*\mathbb{G}_{m,K} = 0$. By (2.1) we get

$$(2.2) \quad H_{et}^1(\text{Spec}(V), f_*\mathbb{G}_{m,K}) = 0 \text{ and } 0 \rightarrow H_{et}^2(\text{Spec}(V), f_*\mathbb{G}_{m,K}) \rightarrow \text{Br}(K).$$

Now, the long exact sequence associated to the exact sequence

$$1 \rightarrow \mathbb{G}_{m,V} \rightarrow f_*\mathbb{G}_{m,K} \rightarrow f_*\mathbb{G}_{m,K}/\mathbb{G}_{m,V} \rightarrow 1$$

of étale sheaves on $\text{Spec}(V)$ implies (using (2.2) and an wellknown fact that the Picard group of a local ring is trivial)

$$(2.3) \quad H_{et}^0(\text{Spec}(V), f_*\mathbb{G}_{m,K}/\mathbb{G}_{m,V}) \cong K^\times/V^\times$$

and

$$(2.4) \quad 0 \rightarrow H_{et}^1(\text{Spec}(V), f_*\mathbb{G}_{m,K}/\mathbb{G}_{m,V}) \rightarrow H_{et}^2(\text{Spec}(V), \mathbb{G}_{m,V}) \rightarrow H_{et}^2(\text{Spec}(V), f_*\mathbb{G}_{m,K}).$$

Our first goal is to prove the following result:

Theorem 2.1. *Let V be a one dimensional valuation ring with the field of fraction K . Let $f : \text{Spec}(K) \rightarrow \text{Spec}(V)$ be the map of schemes induced by the inclusion $V \hookrightarrow K$. Then $H_{et}^1(\text{Spec}(V), f_*\mathbb{G}_{m,K}/\mathbb{G}_{m,V}) = 0$.*

We need some preparations to prove Theorem 2.1. Let us begin with a definition.

Let S be any scheme. The support of an abelian sheaf \mathcal{F} on S_{et} is the set of points $s \in S$ such that $\mathcal{F}_{\bar{s}} \neq 0$ for any geometric point \bar{s} lying over s . It is denoted by $\text{Supp}(\mathcal{F})$.

Lemma 2.2. *Suppose V , K and f are as in Theorem 2.1. Let \mathfrak{m} denote the unique maximal ideal of V . Then $\text{Supp}(f_*\mathbb{G}_{m,K}/\mathbb{G}_{m,V})$ is $\text{Spec}(V/\mathfrak{m})$.*

Proof. Write \mathcal{I} for the étale sheaf $f_*\mathbb{G}_{m,K}/\mathbb{G}_{m,V}$. The stalk of \mathcal{I} at a geometric point \bar{v} is $\mathcal{I}_{\bar{v}} = (K \otimes_V \mathcal{O}_{V,\bar{v}})^\times / \mathcal{O}_{V,\bar{v}}^\times$. Since $\dim V = 1$, $\mathrm{Spec}(V) = \{\xi, \mathfrak{m}\}$ where ξ is the generic point of $\mathrm{Spec}(V)$. Then $\mathcal{I}_\xi = 0$ and $\mathcal{I}_{\bar{\mathfrak{m}}} = K(V^{sh})^\times / (V^{sh})^\times$, where $K(V^{sh})$ denotes the field of fraction of V^{sh} . Note that $K(V^{sh})^\times \neq (V^{sh})^\times$ because V^{sh} is a one dimensional local domain. Thus, the support of \mathcal{I} is $\mathrm{Spec}(V/\mathfrak{m})$. \square

Let k denote the residue field V/\mathfrak{m} . We have a closed immersion $i : \mathrm{Spec}(k) \rightarrow \mathrm{Spec}(V)$. Write \mathcal{I} for $f_*\mathbb{G}_{m,K}/\mathbb{G}_{m,V}$ as before. By Lemma 2.2, $\mathrm{Supp}(\mathcal{I}) = \mathrm{Spec}(k)$. Then Proposition 58.46.4 of [12, Tag 04E1] implies that $i_*(i^{-1}\mathcal{I}) \cong \mathcal{I}$. Here $i^{-1}\mathcal{I}$ is a sheaf on $(\mathrm{Spec}(k))_{et}$. Set $\mathcal{I}' = i^{-1}\mathcal{I}$. Since i_* is an exact functor, $R^q i_* \mathcal{I}' = 0$ for all $q > 0$. The spectral sequence

$$H_{et}^p(\mathrm{Spec}(V), R^q i_* \mathcal{I}') \Rightarrow H_{et}^{p+q}(\mathrm{Spec}(k), \mathcal{I}')$$

gives the following:

Lemma 2.3. *With the above notations, there is an isomorphism of groups*

$$H_{et}^j(\mathrm{Spec}(V), \mathcal{I}) \cong H_{et}^j(\mathrm{Spec}(k), i^{-1}\mathcal{I}) \text{ for all } j \geq 0.$$

The absolute Galois group of k is $\mathrm{Gal}(k^{sep}/k)$ and is denoted by G_k . Here k^{sep} is the separable closure of k in the algebraically closed field \bar{k} . Note that $(i^{-1}\mathcal{I})_{\bar{\mathfrak{m}}} = \mathcal{I}_{\bar{\mathfrak{m}}} = K(V^{sh})^\times / (V^{sh})^\times$. Moreover, $K(V^{sh})^\times / (V^{sh})^\times$ is a G_k -module (see [12, Tag 03QW]). So, we can consider its Galois cohomology group $H^j(G_k, K(V^{sh})^\times / (V^{sh})^\times)$ for $j \geq 0$. Some details related to Galois cohomology can be found in [6] and [11]. Given a G -module M , the invariant subgroup $M^G = \{x \in M \mid g.x = x \text{ for all } g \in G\}$. Here G is any group. Now, Lemma 58.58.2 of [12, Tag 03QQ] implies the following:

Lemma 2.4. *With the above notations, we have*

- (1) $H_{et}^0(\mathrm{Spec}(k), i^{-1}\mathcal{I}) = (K(V^{sh})^\times / (V^{sh})^\times)^{G_k}$,
- (2) $H_{et}^j(\mathrm{Spec}(k), i^{-1}\mathcal{I}) = H^j(G_k, K(V^{sh})^\times / (V^{sh})^\times)$ for $j > 0$.

By combining the above two lemmas, we obtain

$$(2.5) \quad H_{et}^0(\mathrm{Spec}(V), \mathcal{I}) \cong H_{et}^0(\mathrm{Spec}(k), i^{-1}\mathcal{I}) = (K(V^{sh})^\times / (V^{sh})^\times)^{G_k},$$

$$(2.6) \quad H_{et}^j(\mathrm{Spec}(V), \mathcal{I}) \cong H_{et}^j(\mathrm{Spec}(k), i^{-1}\mathcal{I}) = H^j(G_k, K(V^{sh})^\times / (V^{sh})^\times) \text{ for } j > 0.$$

We are now ready to prove Theorem 2.1.

Proof of Theorem 2.1: The strict henselization V^{sh} of V is also a one dimensional valuation ring (see Lemma 15.118.5 of [12, Tag 0ASF]). Recall $K(V^{sh})$ is the field of fraction of V^{sh} . Then, we have $K(V^{sh})^\times / (V^{sh})^\times \cong K^\times / V^\times \cong (K(V^{sh})^\times / (V^{sh})^\times)^{G_k}$, where the first isomorphism by Lemma 15.118.5 of [12, Tag 0ASF] and the second

isomorphism follows from (2.3) and (2.5). Hence, G_k acts trivially on $K(V^{sh})^\times / (V^{sh})^\times$. By (2.6), we also have $H_{et}^1(\text{Spec}(V), f_* \mathbb{G}_{m,K} / \mathbb{G}_{m,V}) \cong H^1(G_k, K(V^{sh})^\times / (V^{sh})^\times)$. So, it is enough to show that $H^1(G_k, K(V^{sh})^\times / (V^{sh})^\times) = 0$. Since $\dim V^{sh} = 1$, the value group $K(V^{sh})^\times / (V^{sh})^\times$ is isomorphic to a non-trivial additive subgroup of \mathbb{R} (see chapter VI, section 4.5, Proposition 8 of [3]). In fact, $K(V^{sh})^\times / (V^{sh})^\times$ is a torsion free $\mathbb{Z}[G_k]$ -module. As G_k is a profinite group (see Proposition 4.1.3 of [6]), $G_k = \varprojlim \text{Gal}(L/k)$, where the limit is over all finite Galois subextensions L/k of k^{sep}/k . Thus,

$$H^1(G_k, K(V^{sh})^\times / (V^{sh})^\times) \cong \varprojlim H^1(\text{Gal}(L/k), K(V^{sh})^\times / (V^{sh})^\times).$$

There are natural surjections $G_k \rightarrow \text{Gal}(L/k)$ for all finite Galois subextensions L/k of k^{sep}/k (see Corollary 4.1.4 of [6]). Hence, all $\text{Gal}(L/k)$ acts trivially on $K(V^{sh})^\times / (V^{sh})^\times$. In this situation, we know (see Exercise 6.1.5 of [13])

$$H^1(\text{Gal}(L/k), K(V^{sh})^\times / (V^{sh})^\times) \cong \text{Hom}_{\text{Groups}}(\text{Gal}(L/k), K(V^{sh})^\times / (V^{sh})^\times),$$

where $\text{Hom}_{\text{Groups}}(-, -)$ denotes the hom set in the category of groups. Since $\text{Gal}(L/k)$ is a finite group, $\text{Hom}_{\text{Groups}}(\text{Gal}(L/k), K(V^{sh})^\times / (V^{sh})^\times) = 0$. Hence the result. \square

Corollary 2.5. *Let V be a one dimensional valuation ring with the field of fraction K . Then the following are true:*

- (1) $H_{et}^2(\text{Spec}(V), \mathbb{G}_{m,V})$ is a torsion group.
- (2) $\text{Br}(V) \cong H_{et}^2(\text{Spec}(V), \mathbb{G}_{m,V})$.
- (3) the canonical map $\text{Br}(V) \rightarrow \text{Br}(K)$ is injective.

Proof. (1) The Brauer group of a field is always a torsion group (see Corollary 2.8.5 of [6]). Thus, $H_{et}^2(\text{Spec}(V), f_* \mathbb{G}_{m,K})$ is a torsion group by (2.2). Hence the result by Theorem 2.1 and (2.4).

(2) By a theorem of Gabber (see [4]), $\text{Br}(R) \cong (H_{et}^2(\text{Spec}(R), \mathbb{G}_{m,R}))_{tor}$ for any commutative ring R . We get the assertion by (1).

(3) This is now immediate from Theorem 2.1, (2.2) and (2.4). \square

3. GENERAL CASE

In this section, we prove the general case of Theorem 1.1. We also discuss some consequences of Theorem 1.1.

Proof of Theorem 1.1: We may assume that $V \neq K$. Note that $K = \bigcup K_i$, where K_i is a finitely generated subfield of K . Then V can be written as filtered limit of V_i , where $V_i = V \cap K_i$ is a valuation ring of K_i with finite rank (see Lemma 2.22 of [2]). Since Br commutes with filtered limit, we may also assume that V is a valuation ring of finite rank $d > 0$. We use induction on d . If $d = 1$ then the result is true by Corollary 2.5(3). Suppose that the result is true for all valuation rings of rank less than d . Let

V be a valuation ring of rank d with the field of fraction K . Let \mathfrak{p} be a nonzero, non-maximal prime ideal of V . Note that $\text{Spec}(V_{\mathfrak{p}})$ is open in $\text{Spec}(V)$. Moreover, $V_{\mathfrak{p}}$ and V/\mathfrak{p} both are valuation rings with the field of fractions K and $k(\mathfrak{p}) = V_{\mathfrak{p}}/\mathfrak{p}V_{\mathfrak{p}}$. We also have $1 \leq \dim(V_{\mathfrak{p}}) < d$ and $1 \leq \dim(V/\mathfrak{p}) < d$. Since V is a valuation ring, the following square (see Proposition 2.8 of [2])

$$\begin{array}{ccc} V & \longrightarrow & V_{\mathfrak{p}} \\ \downarrow & & \downarrow \\ V/\mathfrak{p} & \longrightarrow & k(\mathfrak{p}) \end{array}$$

is Milnor. By Corollary 2.5 of [9], we get an exact sequence

$$0 \rightarrow \text{Br}(V) \rightarrow \text{Br}(V_{\mathfrak{p}}) \times \text{Br}(V/\mathfrak{p}) \rightarrow \text{Br}(k(\mathfrak{p})).$$

Now consider the following commutative diagram

$$\begin{array}{ccccc} 0 & \longrightarrow & \text{Br}(V) & \longrightarrow & \text{Br}(V_{\mathfrak{p}}) \times \text{Br}(V/\mathfrak{p}) \\ & & \downarrow & & \downarrow \\ & & \text{Br}(K) & \longrightarrow & \text{Br}(K) \times \text{Br}(k(\mathfrak{p})), \end{array}$$

where the second vertical map is injective by the induction hypothesis. Hence the result. \square

Corollary 3.1. *Let V be a valuation ring with the field of fraction K . Let $\mathfrak{p}, \mathfrak{q} \in \text{Spec}(V)$. Then either $\text{Br}(V_{\mathfrak{p}}) \hookrightarrow \text{Br}(V_{\mathfrak{q}})$ or $\text{Br}(V_{\mathfrak{q}}) \hookrightarrow \text{Br}(V_{\mathfrak{p}})$. In particular, $\text{Br}(V) \hookrightarrow \text{Br}(V_{\mathfrak{p}})$ for all $\mathfrak{p} \in \text{Spec}(V)$.*

Proof. Since V is a valuation ring, either $\mathfrak{p} \subseteq \mathfrak{q}$ or $\mathfrak{q} \subseteq \mathfrak{p}$. Note that the both $V_{\mathfrak{p}}$ and $V_{\mathfrak{q}}$ are valuation rings with the field of fraction K . The assertion now clear from Theorem 1.1. For the second part, consider $\mathfrak{p} \subseteq \mathfrak{m}$, where \mathfrak{m} is the unique maximal ideal. \square

Remark 3.2. For a valuation ring V , we have exactly one chain of prime ideals, say

$$(0) \subseteq \mathfrak{p} \subseteq \cdots \subseteq \mathfrak{q} \subseteq \mathfrak{m}.$$

By Corollary 3.1, we get

$$\text{Br}(V) \hookrightarrow \text{Br}(V_{\mathfrak{q}}) \hookrightarrow \cdots \hookrightarrow \text{Br}(V_{\mathfrak{p}}) \hookrightarrow \text{Br}(K).$$

In particular, $\text{Br}(V) = \bigcap_{\mathfrak{p} \in \text{Spec}(V)} \text{Br}(V_{\mathfrak{p}})$.

Remark 3.3. Kestutis Česnavičius inform me in [7] that a stronger version of Theorem 1.1 is also true over valuation rings. More explicitly, for a valuation ring V with the field of fraction K , the canonical map $H_{et}^2(\text{Spec}(V), \mathbb{G}_{m,V}) \rightarrow H_{et}^2(\text{Spec}(K), \mathbb{G}_{m,K})$ is injective.

REFERENCES

- [1] M. Auslander and O. Goldman, *The Brauer group of a commutative ring*. Trans. Amer. Math. Soc. **97**, (1960) 367-409.
- [2] B. Bhatt and A. Mathew, *The arc-topology*, arXiv:1807.04725v4, to appear in Duke.
- [3] N. Bourbaki, *Commutative algebra, chapters 1-7*, Springer, 1989.
- [4] A.J. De Jong, *A result of Gabber*, Preprint, 2003.
- [5] A. Grothendieck, Le Groupe de Brauer II, Dix Exposes sur la Cohomologie des Schemas, North Holland, Amsterdam, (1969), 66-87.
- [6] P. Gille and T. Szamuely, *Central Simple Algebras and Galois Cohomology*, Second edition, Cambridge Studies in Advanced Mathematics, **165**. Cambridge University Press, Cambridge, 2017.
- [7] Kestutis Česnavičius, an email on April 9, 2021.
- [8] S. Kelly and M. Morrow, *K-theory of valuation rings*, arxiv: 1810.12203, 2018, to appear in Compositio Mathematica.
- [9] M.A. Knus and M. Ojanguren, *A Mayer-Vietoris sequence for the Brauer group*, Journal of pure and applied algebra **5**, (1974) 345-360.
- [10] J. Milne, *Étale Cohomology*. Princeton: Princeton University Press 1980.
- [11] J. P. Serre, *Galois Cohomology*, Springer, 2002.
- [12] Stack Project, <https://stacks.math.columbia.edu>.
- [13] C. Weibel, *An introduction to homological algebra*, Cambridge studies in advanced mathematics 38, 1994.

DEPARTMENT OF MATHEMATICS, INDIAN INSTITUTE OF SCIENCE EDUCATION AND RESEARCH,
 BHOPAL, BHOPAL BYPASS ROAD, BHAURI, BHOPAL-462066, MADHYA PRADESH, INDIA
Email address: vsadhu@iiserb.ac.in, viveksadhu@gmail.com