

ALGEBRAIC SPECTRAL THEORY AND SERRE MULTIPLICITY FORMULA

ANAR DOSI (DOSIEV)

ABSTRACT. The present paper is devoted to an algebraic treatment of the joint spectral theory within the framework of Noetherian modules over an algebra finite extension of an algebraically closed field. We prove the spectral mapping theorem and analyze the index of tuples in purely algebraic case. The index function over tuples from the coordinate ring of a variety is naturally extended up to a numerical Tor-polynomial. Based on Serre's multiplicity formula, we deduce that Tor-polynomial is just the Samuel polynomial of the local algebra.

1. INTRODUCTION

The joint spectral theory plays a central role in operator theory and the complex analytic geometry. Its origin goes back to Gelfand's commutative Banach algebras and their representations. The joint spectral theory was mainly developed within the context of holomorphic functional calculus problem by J. Taylor [25], [26]. An interesting link to the complex analytic geometry was found by M. Putinar [23], [13] by constructing Taylor's holomorphic functional calculus within the context of Stein spaces. Taylor's spectral theory has a strong homological background developed independently by J. L. Taylor in [26] and A. Ya. Helemskii in [17], that allows to have a further generalization of the theory in the noncommutative setting. That played a crucial role in foundations of noncommutative complex analytic geometry (see [6], [8], [9] and [21]).

It is well known that many key results from the complex analytic geometry have their analogs in schemes such as the fundamental theorem of Serre on vanishing [16, 3.3.7]. More interesting sight is to have a scheme-theoretic analog of the joint spectra found in [10] and [11] that indicates to the fundamental nature of the joint spectral theory. Furthermore, Taylor's multivariable functional calculus has a scheme version suggested in [11]. Namely, Putinar spectrum can be defined for a module over a scheme and it plays the same central role in the functional calculus problem over Noetherian schemes as it does in the complex analytic geometry. Actually, in the affine case the same spectrum occurred in [18] and [19] based on the result of A. Neeman [20] that there is a bijection between subsets of $\text{Spec}(R)$ and localizing subcategories of the derived category of complexes of R -modules. Taylor's functional calculus through derived categories was proposed in [22] by A. Yu. Pirkovskii, which is a complex analytic version of Neeman's result.

In the paper we intend to develop a spectral theory in purely algebraic case with all its key properties and their links to algebraic geometry. As a basic tool we are exploiting many results from commutative algebra. Actually, some key results of commutative algebra can be retreated from point of view the joint spectral theory. For example, associated primes of a Noetherian module play the role of eigenvalues whereas support primes are spectral values. That approach inherits Koszul homology groups of an algebraic variety and the related index. In the case of a variety the index is reduced to the multiplicity from the local theory [24], and we use the multiplicity formula of Serre for calculation of Koszul homology groups.

Date: April 8, 2021.

2000 Mathematics Subject Classification. Primary 47A60, 13D03; Secondary 13D40, 46H30.

Key words and phrases. Taylor spectrum, Noetherian modules, integral extensions, Samuel polynomial, Serre's formula.

Fix a commutative k -algebra R/k with an n -tuple x in R , and an R -module M . It turns out that x is a family of commuting linear transformations acting on the k -vector space M , and we have their *Taylor spectrum* $\sigma(x, M)$ to be the set of those $a \in \mathbb{A}^n$ such that the Koszul complex $\text{Kos}(x - a, M)$ fails to be exact. If the homology groups $H_p(x - a, M)$ of $\text{Kos}(x - a, M)$ are finite dimensional k -vector spaces we have the index $i(x - a) = \sum_{p=0}^n (-1)^p \dim_k(H_p(x - a, M))$ of the tuple $x - a$. The last homology group $H_n(x - a, M)$ responds to the submodule of all joint eigenvectors in M whenever a is a joint eigenvalue of the tuple x . The set of all eigenvalues of x is called *the point spectrum* $\sigma_p(x, M)$ of x . The central result of Section 2 is the spectral mapping formula

$$(1.1) \quad \sigma(p(x), M) = p(\sigma(x, M))$$

for all tuples x and $p(x)$ (a polynomial tuple in x) from R whenever R/k is an algebra finite extension of an algebraically closed field k and M is a Noetherian R -module. Moreover, $i(x) < \infty$ for an n -tuple x from R whenever $R' = k[x] \subseteq R$ is an integral extension. In this case, $i(x) = i(y)$ for every n -tuple y from the subalgebra R' generating the same maximal ideal $\langle x \rangle \subseteq R'$. It is an analog of the index stability result from analysis.

In Section 3, we analyze the general case of a scheme $(\mathfrak{X}, \mathcal{O}_{\mathfrak{X}})$ and a module M over the global sections $R = \Gamma(\mathfrak{X}, \mathcal{O}_{\mathfrak{X}})$. The spectrum $\sigma(\mathfrak{X}, M)$ of the R -module M over the scheme $(\mathfrak{X}, \mathcal{O}_{\mathfrak{X}})$ is defined as the complement to the set of those $x \in \mathfrak{X}$ such that $\text{Tor}_i^R(\mathcal{O}_{\mathfrak{X}}(U), M) = 0$, $i \geq 0$ for an affine neighborhood U of x . This spectrum was introduced in [11] as a scheme-theoretic analog of Putinar spectrum of analytic sheaves [23]. In the case of an affine scheme $\mathfrak{X} = \text{Spec}(R)$ the spectrum $\sigma(\mathfrak{X}, M)$ is reduced to the closure of the support $\text{Supp}_R(M)$ of the R -module M . If M is a finitely generated R -module then we come up with the support $\text{Supp}_R(M)$ rather than its closure. The point spectrum $\sigma_p(\mathfrak{X}, M)$ is defined to be the set $\text{Ass}_R(M)$ of all associated primes of M . The spectral mapping theorem for modules over schemes was proposed in [11]. Its affine version states that

$$(1.2) \quad \sigma(\mathfrak{Y}, M) = f(\sigma(\mathfrak{X}, M))^-$$

whenever $\mathfrak{X} = \text{Spec}(R)$ and $\mathfrak{Y} = \text{Spec}(R')$ are affine schemes, $f = \varphi^* : \mathfrak{X} \rightarrow \mathfrak{Y}$ is a morphism responding to a ring map $\varphi : R' \rightarrow R$, and $M \in R\text{-mod}$, which is R' -module through φ either. For a ring extension $\iota : R' \subseteq R$ and a finitely generated R -module M we obtain that $\iota^*(\text{Supp}_R(M))$ is dense in $\text{Supp}_{R'}(M)$. In the case of the point spectrum we prove more valuable equality without the related closure in (1.2) (being so weak, the closure of Zariski topology covers up too much). If $\iota : R' \subseteq R$ is a ring extension with Noetherian R , and $M \in R\text{-mod}$, then (see Theorem 3.1)

$$\sigma_p(\mathfrak{X}', M) = \iota^*(\sigma_p(\mathfrak{X}, M)).$$

Notice that similar result for the spectrum (or support) is not true (see below Remark 3.1). But that is the case if $R' \subseteq R$ is integral and we come with classics Krull-Cohen-Seidenberg Theory [1, Ch. 14], [2, 5.2].

The spectrum over a scheme and Taylor spectrum linked with each other when $R = k[x]$ is an algebra finite extension. Namely, $\mathfrak{X} = \text{Spec}(R) \subseteq \mathbb{A}_k^n$ up to a homeomorphism, $\sigma(\mathfrak{X}, M) = \sigma(\mathbb{A}_k^n, M)$, $\sigma_p(\mathfrak{X}, M) = \sigma_p(\mathbb{A}_k^n, M)$, and $\sigma_p(x, M) = \sigma_p(\mathbb{A}_k^n, M) \cap \mathbb{A}^n$. If M is a Noetherian R -module then $\sigma(x, M) = \sigma(\mathbb{A}_k^n, M) \cap \mathbb{A}^n$ and it is a nonempty closed subset of \mathbb{A}^n whose closure in the scheme \mathbb{A}_k^n is reduced to $\sigma(\mathfrak{X}, M)$. If $k[p(x)] \subseteq R$ is integral then

$$\sigma_p(p(x), M) = p(\sigma_p(x, M)).$$

Actually, (1.1) and (1.2) can be driven from the classics whenever $k[p(x)] \subseteq R$ is integral and M is Noetherian. But the spectral mapping formula (1.1) is much stronger than classics, which is true for all tuples. Thus Taylor spectrum with all its properties have an independent value.

In Section 4 we consider the case of $M = R$ of the coordinate ring R of a variety $Y \subseteq \mathbb{A}^n$. In this case, the tuple $x \subseteq R$ consists of the coordinate functions and $\sigma(x, R) = Y$ whereas $\sigma_p(x, R) = \emptyset$. In the case of an algebraic set Y the point spectrum $\sigma_p(x, R)$ is the set of all isolated points of Y . For every point $a \in \mathbb{A}^n$ we have $i(x - a) = 0$ due to Serre's multiplicity formula from the local algebra. For a singular point $a \in Y$ the dimension of $H_1(x - a, R)$ could get high integers depending on the depth of singularity (see below Lemma 4.6). The link between the index and the dimension of Y is obtained through the numerical polynomial $p : \mathbb{Z} \rightarrow \mathbb{Z}$, $p(r) = \sum_{i=1}^n (-1)^i \dim_k (\text{Tor}_i^P(P/\langle X_1, \dots, X_n \rangle^r, R))$ called the Tor-polynomial, which is reduced to Hilbert-Samuel polynomial of the localization $R_{\langle x-a \rangle}$.

Finally, I wish to thank G. G. Amosov, O. Yu. Aristov, B. Bilich and A. Yu. Pirkovskii for their interest to the paper, useful discussions and to draw my attention to the papers [18] and [19].

2. THE PROJECTION PROPERTY OF SPECTRUM

In the present section we prove the projection property of spectrum based on homology groups of the Koszul complex. The technical back up is the homology of commutative rings related to Koszul homology groups.

2.1. Lemma Bourbaki. Let R be a (unital) commutative ring, $K_i \in R\text{-mod}$, $i = 0, 1$, and let $\alpha : K_1 \rightarrow K_0$ be an R -linear map (module morphism). The chain complex $0 \leftarrow K_0 \xleftarrow{\alpha} K_1$ is denoted by \mathcal{K} . If \mathcal{C} is a chain complex $0 \leftarrow C_0 \xleftarrow{d_0} C_1 \xleftarrow{d_1} \dots$ in $R\text{-mod}$ then $\mathcal{K} \otimes_R \mathcal{C}$ is the following complex

$$\begin{array}{ccccccc} & & (\mathcal{K} \otimes_R \mathcal{C})_{p-1} & & & & (\mathcal{K} \otimes_R \mathcal{C})_p \\ & & \parallel & & & & \parallel \\ \dots & \xleftarrow{\partial_{p-2}} & (K_0 \otimes_R C_{p-1}) \oplus (K_1 \otimes_R C_{p-2}) & \xleftarrow{\partial_{p-1}} & (K_0 \otimes_R C_p) \oplus (K_1 \otimes_R C_{p-1}) & \xleftarrow{\partial_p} & \dots \end{array}$$

with the differential

$$(2.1) \quad \partial_{p-1}(z_{0,p}, z_{1,p-1}) = ((1 \otimes d_{p-1})(z_{0,p}) + (-1)^{p-1}(\alpha \otimes 1)(z_{1,p-1}), (1 \otimes d_{p-2})(z_{1,p-1})),$$

where $z_{i,j} \in K_i \otimes_R C_j$. The canonical embedding i_p and projection π_p define the exact sequence

$$0 \leftarrow K_1 \otimes_R C_{p-1} \xleftarrow{\pi_p} (K_0 \otimes_R C_p) \oplus (K_1 \otimes_R C_{p-1}) \xleftarrow{i_p} K_0 \otimes_R C_p \leftarrow 0$$

which splits. Since the diagram

$$\begin{array}{ccccccc} & & 0 & & 0 & & \\ & & \downarrow & & \downarrow & & \\ \dots & \xleftarrow{1 \otimes d_{p-2}} & K_0 \otimes_R C_{p-1} & \xleftarrow{1 \otimes d_{p-1}} & K_0 \otimes_R C_p & \xleftarrow{1 \otimes d_p} & \dots \\ & & \downarrow i_{p-1} & & \downarrow i_p & & \\ \dots & \xleftarrow{\partial_{p-2}} & (\mathcal{K} \otimes_R \mathcal{C})_{p-1} & \xleftarrow{\partial_{p-1}} & (\mathcal{K} \otimes_R \mathcal{C})_p & \xleftarrow{\partial_p} & \dots \\ & & \downarrow \pi_{p-1} & & \downarrow \pi_p & & \\ \dots & \xleftarrow{1 \otimes d_{p-3}} & K_1 \otimes_R C_{p-2} & \xleftarrow{1 \otimes d_{p-2}} & K_1 \otimes_R C_{p-1} & \xleftarrow{1 \otimes d_{p-1}} & \dots \\ & & \downarrow & & \downarrow & & \\ & & 0 & & 0 & & \end{array}$$

commutes, there is an exact sequence

$$0 \leftarrow K_1 \otimes_R \mathcal{C} \xleftarrow{\pi} \mathcal{K} \otimes_R \mathcal{C} \xleftarrow{i} K_0 \otimes_R \mathcal{C} \leftarrow 0$$

of complexes with $\deg(\pi) = -1$ and $\deg(i) = 0$. The sequence in turn generates a long exact sequence of homology groups

$$(2.2) \quad \cdots \longleftarrow H_{p-1}(K_0 \otimes_R \mathcal{C}) \xleftarrow{\delta_{p-1}} H_{p-1}(K_1 \otimes_R \mathcal{C}) \longleftarrow H_p(\mathcal{K} \otimes_R \mathcal{C}) \longleftarrow H_p(K_0 \otimes_R \mathcal{C}) \xleftarrow{\delta_p} \cdots$$

with the connecting morphisms δ_p , $p \geq 0$. If \mathcal{K} is flat (that is, K_i are flat modules) then $H_p(K_i \otimes_R \mathcal{C}) = K_i \otimes_R H_p(\mathcal{C})$ for all i and p .

Lemma 2.1. *The morphism $\delta_{p-1} : H_{p-1}(K_1 \otimes_R \mathcal{C}) \rightarrow H_{p-1}(K_0 \otimes_R \mathcal{C})$ is acting by the rule $\delta_{p-1}(\omega^\sim) = (-1)^{p-1}((\alpha \otimes 1)\omega)^\sim$ for all $\omega \in \ker(1 \otimes d_{p-2})$, that is, $\delta_{p-1} = (-1)^{p-1}(\alpha \otimes 1)^\sim$ for all p . In the case of a flat complex \mathcal{K} the morphism δ_{p-1} is reduced to the following morphism $\delta_{p-1} : K_1 \otimes_R H_{p-1}(\mathcal{C}) \rightarrow K_0 \otimes_R H_{p-1}(\mathcal{C})$, $\delta_{p-1} = (-1)^{p-1}\alpha \otimes 1$.*

Proof. Take $\omega \in \ker(1 \otimes d_{p-2})$. Then $(0, \omega) \in (K_0 \otimes_R \mathcal{C}_p) \oplus (K_1 \otimes_R \mathcal{C}_{p-1}) = (\mathcal{K} \otimes_R \mathcal{C})_p$ and

$$\partial_{p-1}(0, \omega) = ((-1)^{p-1}(\alpha \otimes 1)(\omega), 0) = i_{p-1}((-1)^{p-1}(\alpha \otimes 1)(\omega))$$

(see (2.1)). Hence $(-1)^{p-1}(\alpha \otimes 1)(\omega) \in \ker(1 \otimes d_{p-2})$ and $\delta_{p-1}(\omega^\sim) = (-1)^{p-1}((\alpha \otimes 1)\omega)^\sim$.

Finally, if \mathcal{K} is flat and $\omega = x_1 \otimes y_{p-1}$ with $x_1 \in K_1$, $y_{p-1} \in \ker(d_{p-2})$ then

$$\delta_{p-1}(x_1 \otimes y_{p-1}^\sim) = \delta_{p-1}(\omega^\sim) = (-1)^{p-1}(\alpha(x_1) \otimes y_{p-1})^\sim = (-1)^{p-1}\alpha(x_1) \otimes y_{p-1}^\sim,$$

that is, $\delta_{p-1} = (-1)^{p-1}\alpha \otimes 1$. □

The following key assertion is taken from [3, 9.5, Lemma 3].

Lemma 2.2. *(N. Bourbaki) If $\alpha : K_1 \rightarrow K_0$ is a morphism of flat R -modules and \mathcal{C} is a complex in $R\text{-mod}$ then the long homology sequence generates the following short exact sequences*

$$0 \longleftarrow \ker(\alpha) \otimes_R H_{p-1}(\mathcal{C}) \longleftarrow H_p(\mathcal{K} \otimes_R \mathcal{C}) \longleftarrow \text{coker}(\alpha) \otimes_R H_p(\mathcal{C}) \longleftarrow 0$$

of R -modules, $p \geq 1$.

Proof. Using the long homology sequence (2.2) and Lemma 2.1, we obtain the following exact sequence

$$K_0 \otimes_R H_{p-1}(\mathcal{C}) \xleftarrow{(-1)^{p-1}\alpha \otimes 1} K_1 \otimes_R H_{p-1}(\mathcal{C}) \longleftarrow H_p(\mathcal{K} \otimes_R \mathcal{C}) \longleftarrow K_0 \otimes_R H_{p-1}(\mathcal{C}) \xleftarrow{(-1)^p \alpha \otimes 1} K_1 \otimes_R H_p(\mathcal{C})$$

of R -modules. Using again the flatness, we deduce that $\ker((-1)^{p-1}\alpha \otimes 1) = \ker(\alpha) \otimes_R H_{p-1}(\mathcal{C})$ and $\text{coker}((-1)^p \alpha \otimes 1) = \text{coker}(\alpha) \otimes_R H_p(\mathcal{C})$. The rest is clear. □

As a practical use of Lemma 2.2, let us consider the case of $K_0 = K_1 = R$ and $\alpha = x$ is an element of the ring R , which is acting over all R -modules as a multiplication operator. Moreover, $K_i \otimes_R H_p(\mathcal{C}) = H_p(\mathcal{C})$ and $\alpha \otimes 1$ is the same action $x : H_p(\mathcal{C}) \rightarrow H_p(\mathcal{C})$ of x over the R -module $H_p(\mathcal{C})$ denoted by $x|H_p(\mathcal{C})$. The complex $\mathcal{K} \otimes_R \mathcal{C}$ is reduced to the cone $\text{Con}(x, \mathcal{C})$ of the morphism $x : \mathcal{C} \rightarrow \mathcal{C}$. Namely, $\text{Con}(x, \mathcal{C})$ is the following complex

$$\cdots \longleftarrow C_{p-1} \oplus C_{p-2} \xleftarrow{\partial_{p-1}} C_p \oplus C_{p-1} \longleftarrow \cdots$$

with the differential $\partial_{p-1}(c_p, c_{p-1}) = (d_{p-1}(c_p) + (-1)^{p-1}xc_{p-1}, d_{p-2}(c_{p-1}))$.

Corollary 2.1. *If $x \in R$ and \mathcal{C} is a complex of R -modules then the following canonical sequences*

$$0 \longleftarrow \ker(x|H_{p-1}(\mathcal{C})) \longleftarrow H_p(x, \mathcal{C}) \longleftarrow \text{coker}(x|H_p(\mathcal{C})) \longleftarrow 0$$

are exact, where $H_p(x, \mathcal{C})$ are homology groups of the cone $\text{Con}(x, \mathcal{C})$.

Proof. Since $\ker(\alpha) \otimes_R H_{p-1}(\mathcal{C}) = \ker(\alpha \otimes 1) = \ker(x|H_{p-1}(\mathcal{C}))$ and $\text{coker}(\alpha) \otimes_R H_p(\mathcal{C}) = \text{coker}(\alpha \otimes 1) = \text{coker}(x|H_p(\mathcal{C}))$, the result follows from Lemma 2.2. □

In particular, $H_p(x, \mathcal{C}) = 0$ iff $x : H_{p-1}(\mathcal{C}) \rightarrow H_{p-1}(\mathcal{C})$ is an injection and $x : H_p(\mathcal{C}) \rightarrow H_p(\mathcal{C})$ is a surjection.

2.2. Koszul homology groups. Fix a field k , R/k a commutative k -algebra, $x = (x_1, \dots, x_n)$ an n -tuple in R , and let M be an R -module, which in turn is a k -vector space with an n -tuple x of mutually commuting linear transformations acting on it. If it is necessary one can replace R by an algebra finite extension $k[x]$ of the field k . For every point a from the affine space \mathbb{A}^n we have the tuple $x - a = (x_1 - a_1, \dots, x_n - a_n)$ on M , which in turn defines the Koszul complex $\text{Kos}(x - a, M)$:

$$0 \leftarrow M \xleftarrow{\partial_0} M \otimes_k k^n \xleftarrow{\partial_1} \dots \xleftarrow{\partial_{p-2}} M \otimes_k \wedge^{p-1} k^n \xleftarrow{\partial_{p-1}} M \otimes_k \wedge^p k^n \xleftarrow{\partial_p} \dots \xleftarrow{\partial_{n-1}} M \leftarrow 0$$

in R -mod with the differential

$$\partial_{p-1}(m \otimes v_p) = \sum_{s=1}^p (-1)^{s+1} (x_{i_s} - a_{i_s}) m \otimes v_{p,s},$$

where $m \in M$, $v_p = e_{i_1} \wedge \dots \wedge e_{i_p}$, $v_{p,s} = e_{i_1} \wedge \dots \wedge \widehat{e_{i_s}} \wedge \dots \wedge e_{i_p}$ (the notation $\widehat{e_{i_s}}$ stands for skipping e_{i_s} from the p -vector) and (e_1, \dots, e_n) is the standard basis for k^n . The homology groups of $\text{Kos}(x - a, M)$ are denoted by $H_p(x - a, M)$, $p \geq 0$, which are R -modules. We put

$$i(x) = \sum_{p=0}^n (-1)^{p+1} \dim_k(H_p(x, M))$$

to be the index of the tuple x whenever $\dim_k(H_p(x, M)) < \infty$ for all p . In the latter case we write $i(x) < \infty$. Recall that the index $i(t)$ of a single k -linear transformation $t : M \rightarrow M$ is given by $i(t) = \dim_k \ker(t) - \dim_k \text{coker}(t)$ whenever both dimensions are finite. If $\dim_k(M) < \infty$ then $i(t) = 0$ for every t . The index of tuples of bounded linear operators acting on a Banach space is a subject of Fredholm theory from analysis [5], [14].

Lemma 2.3. *If $x' = (x_1, \dots, x_{n-1})$ then there is an exact sequence*

$$0 \leftarrow \ker(x_n | H_{p-1}(x', M)) \longleftarrow H_p(x, M) \longleftarrow \text{coker}(x_n | H_p(x', M)) \leftarrow 0$$

of R -modules. In particular, if $H_p(x, M) \neq 0$ then $H_p(x', M) \neq 0$ or $H_{p-1}(x', M) \neq 0$. Moreover, if $i(x) < \infty$ then $i(x_n | H_p(x', M)) < \infty$ for every p , and $i(x) = \sum_{p=0}^{n-1} (-1)^p i(x_n | H_p(x', M))$. If $i(x') < \infty$ then $i(x) = 0$.

Proof. Put $\mathcal{C} = \text{Kos}(x', M)$ to be a complex of R -modules. Then x_n defines an endomorphism of \mathcal{C} , which in turn generates the cone $\text{Con}(x_n, \mathcal{C})$. The fact $\text{Con}(x_n, \mathcal{C}) = \text{Kos}(x, M)$ is known even in the noncommutative case (see [15, Lemma 1.5]). One needs to use the vector space isomorphism $\wedge^p k^{n-1} \oplus \wedge^{p-1} k^{n-1} \rightarrow \wedge^p k^n$, $(v_1, v_2) \mapsto v_1 + v_2 \wedge e_n$, which generates P -module isomorphism $C_p \oplus C_{p-1} \rightarrow M \otimes_k \wedge^p k^n$ for every p . If $m_p = m \otimes v_p \in M \otimes_k \wedge^p k^n$ with $j_p \neq n$, then m_p is identified with an element $(c_p, 0)$ of $C_p \oplus C_{p-1}$ and $\partial(c_p, 0) = (\partial'(c_p), 0)$, where ∂' is the differential of \mathcal{C} . If $j_p = n$ then m_p is identified with $(0, c_{p-1})$, where $c_{p-1} = m \otimes v_{p-1} \in C_{p-1}$, $v_{p-1} \wedge e_n = v_p$. Moreover,

$$\begin{aligned} \partial(0, c_{p-1}) &= \partial(m_p) = \sum_{s=1}^{p-1} (-1)^{s+1} x_{i_s} m \otimes v_{p-1,s} \wedge e_n + (-1)^{p+1} x_n m \otimes v_{p-1} \\ &= \partial'(c_{p-1}) \wedge e_n + (-1)^{p+1} x_n c_{p-1} = ((-1)^{p+1} x_n c_{p-1}, \partial'(c_{p-1})). \end{aligned}$$

Hence $\partial(c_p, c_{p-1}) = (\partial'(c_p) + (-1)^{p-1} x_n c_{p-1}, \partial'(c_{p-1}))$ for all $(c_p, c_{p-1}) \in C_p \oplus C_{p-1}$, which means that ∂ is the morphism of $\text{Con}(x_n, \mathcal{C})$. It remains to use Corollary 2.1.

Finally, consider the case of $i(x) < \infty$. Since $\dim_k(H_p(x, M)) < \infty$ for every p , it follows that $\dim_k \ker(x_n|H_{p-1}(x', M)) < \infty$ and $\dim_k \operatorname{coker}(x_n|H_p(x', M)) < \infty$ for every p , which means that $i(x_n|H_p(x', M)) < \infty$ for every p . Then

$$\begin{aligned} i(x) &= \sum_{p=0}^n (-1)^{p+1} (\dim_k \ker(x_n|H_{p-1}(x', M)) + \dim_k \operatorname{coker}(x_n|H_p(x', M))) \\ &= \sum_{p=0}^n (-1)^{p+1} \dim_k \operatorname{coker}(x_n|H_p(x', M)) + (-1)^p \dim_k \ker(x_n|H_p(x', M)) \\ &= \sum_{p=0}^n (-1)^p i(x_n|H_p(x', M)), \end{aligned}$$

that is, $i(x) = \sum_{p=0}^{n-1} (-1)^p i(x_n|H_p(x', M))$ (for $H_n(x', M) = 0$). In particular, if $i(x') < \infty$ then $\dim_k(H_p(x', M)) < \infty$ and $i(x_n|H_p(x', M)) = 0$ for all p . Therefore $i(x) = 0$. \square

The Taylor spectrum $\sigma(x, M)$ of the operator tuple x on M is defined to be a subset of \mathbb{A}^n of those a such that $\operatorname{Kos}(x - a, M)$ is not exact. The point spectrum $\sigma_p(x, M)$ consists of those $a \in \mathbb{A}^n$ such that $H_n(x - a, M) \neq 0$. Taking into account $H_n(x - a, M) = \ker \partial_{n-1}$, we conclude that $a \in \sigma_p(x, M)$ iff there is a nonzero $m \in M$ such that $x_i m = a_i m$ for all i , that is, a turns out to be a joint eigenvalue and m is the related joint eigenvector (see [4]).

Corollary 2.2. *Let R/k be an algebra, $M \in R\text{-mod}$, x an n -tuple from R . Then $\pi(\sigma(x, M)) \subseteq \sigma(x', M)$, where $\pi : \mathbb{A}^n \rightarrow \mathbb{A}^{n-1}$ is the canonical projection onto first $n - 1$ coordinates.*

Proof. If $a \in \sigma(x, M)$ then $H_p(x - a, M) \neq 0$ for some p . By Lemma 2.3, either $H_p(x' - a', M) \neq 0$ or $H_{p-1}(x' - a', M) \neq 0$, where $a' = \pi(a)$. Whence $a' \in \sigma(x', M)$. \square

Put $M_i = M/\langle x_1, \dots, x_i \rangle M$, $i \geq 1$. Recall [3, 9.6], [1, Ch. 23] that if $M_n \neq \{0\}$ and $x_i \notin \operatorname{zdiv}(M_{i-1})$ (zero-divisors) for all i , then x is called a regular sequence. In the case of a nonzero Noetherian module M and $x \subseteq \operatorname{rad}(M)$ the condition $M_n \neq \{0\}$ is satisfied automatically. Indeed, otherwise $\langle x \rangle M = M$ implies that $(1 + a)M = \{0\}$ for some $a \in \langle x \rangle$ (Nakayama Lemma), which means that $1 + a \in \operatorname{Ann}(M) \subseteq \operatorname{rad}(M)$ or $1 \in \operatorname{rad}(M)$, a contradiction.

Corollary 2.3. *Let R/k be a k -algebra, $M \in R\text{-mod}$, and let $x = (x_1, \dots, x_n) \subseteq R$ be a regular sequence for M . Then $H_i(x, M) = 0$ for all $i > 0$.*

Proof. We proceed by induction on n . If $n = 1$ the result follows. In the general case, we put $x' = (x_1, \dots, x_{n-1})$, which is regular either. By induction hypothesis, $H_i(x', M) = 0$ for all $i > 0$. By Lemma 2.3, $H_p(x, M) = 0$ for all $p \geq 2$. Moreover, $H_1(x, M) = \ker(x_n|H_0(x', M))$ and $H_0(x', M) = M/\langle x' \rangle M$. But $x_n : M/\langle x' \rangle M \rightarrow M/\langle x' \rangle M$ is injective by assumption, therefore $H_1(x, M) = 0$. Whence $H_i(x, M) = 0$ for all $i > 0$. \square

In the case of a local ring R and a Noetherian module M the statement of Corollary 2.3 turns out to be a criteria for regularity of a tuple (see [7] for regularity in the noncommutative setting). One can easily see that the tuple $X = (X_1, \dots, X_n)$ is a regular sequence for the polynomial algebra $P = k[X]$ to be a P -module. By Corollary 2.3, $H_i(X, P) = 0$ for all $i > 0$, and $H_0(X, P) = P/\mathfrak{t} = k$, where $\mathfrak{t} = \langle X \rangle$ is the maximal ideal of P . Thus $0 \leftarrow k \leftarrow \operatorname{Kos}(X, P)$ is exact or $\operatorname{Kos}(X, P)$ provides a free P -module resolution for the P -module k . In particular, for every P -module M we obtain that $\operatorname{Kos}(X, P) \otimes_P M = \operatorname{Kos}(x, M)$ and $\operatorname{Tor}_i^P(k, M) = H_i(\operatorname{Kos}(X, P) \otimes_P M) = H_i(x, M)$, where x is the n -tuple of linear transformations on M given by X -action on M (see also below Subsection 3.4).

Let R/k be a k -algebra, $R' = k[x] \subseteq R$ a subalgebra with an n -tuple x . An m -tuple y from R' is said to be *related to* x if x and y generate the same (maximal) ideal $\langle x \rangle = \langle y \rangle$ in R' . In particular, x is always related to x itself.

The following assertion is well know [25]. Below we provide its modified (a bit) version with the detailed proof.

Lemma 2.4. *If R/k is an algebra, x an n -tuple in R and y is an m -tuple related to x , then the $k[x]$ -module structure of $H_p(y, M)$ is reduced to its k -vector space structure for every $p \geq 0$. In particular, the action of a polynomial $p(x) \in R$ on $H_p(y, M)$ is just the multiplication by $p(0)$ operator.*

Proof. For every i let us define the P -linear map

$$\gamma_i : M \otimes_k \wedge^p k^m \rightarrow M \otimes_k \wedge^{p+1} k^m, \quad \gamma_i(u_p) = u \otimes (e_i \wedge v_p),$$

where $u_p = u \otimes v_p$, $e_i \wedge v_p = (-1)^{s-1} e_{i_1} \wedge \dots \wedge e_i \wedge \dots \wedge e_{i_p} = (-1)^{s-1} v_{p+1}$ whenever $v_p = e_{i_1} \wedge \dots \wedge e_{i_p}$ and $i_1 < \dots < i_{s-1} < i < i_s < \dots < i_p$ for some s . Then

$$\begin{aligned} \partial_p \gamma_i(u_p) &= (-1)^{s-1} \sum_{k < s} (-1)^{k+1} y_{i_k} u \otimes v_{p+1,k} + x_i u \otimes v_p + (-1)^{s-1} \sum_{k \geq s} (-1)^k y_{i_k} u \otimes v_{p+1,k} \\ &= - \sum_{k < s} (-1)^{k+1} y_{i_k} u \otimes e_i \wedge v_{p,k} + \sum_{k \geq s} (-1)^k y_{i_k} u \otimes e_i \wedge v_{p,k} + x_i m \otimes v_p \\ &= -\gamma_i \partial_{p-1}(u_p) + y_i u_p. \end{aligned}$$

If $u_p \in \ker(\partial_{p-1})$ then $y_i m_p = \partial_p \gamma_i(u_p) \in \text{im}(\partial_p)$, which means that the action of y_i over $H_p(y, M)$ is trivial, that is, $y H_p(y, M) = \{0\}$. But $H_p(y, M)$ is an R -module, therefore $\langle y \rangle H_p(y, M) = \{0\}$. Taking into account that $\langle x \rangle = \langle y \rangle$ in R' , we deduce that $\langle x \rangle H_p(y, M) = \{0\}$.

Finally, take $p(x) \in R'$. Then $p(x) = p(0) + g$, $g \in \langle x \rangle$ and $p(x) H_p(y, M) = p(0) H_p(y, M)$. \square

Since $R' = k[x] = k[x - a]$ and $\langle x - a \rangle \subseteq R'$ is a maximal ideal, we conclude that $k[x]$ -module structure of $H_p(x - a, M)$ is reduced to its k -vector space structure for every $p \geq 0$. Actually, the action of a polynomial $p(x) \in R$ on $H_p(x - a, M)$ is just the multiplication by $p(a)$ operator.

Again consider $R' = k[x] \subseteq R$ a subalgebra with an n -tuple x , and an m -tuple y from R' related to x . If $p(x) \in R'$ then $z = (y, p(x))$ is an $m + 1$ -tuple in R .

Proposition 2.1. *If $b = (0, \lambda) \in \sigma(z, M)$ for some $\lambda \in k$, then $0 \in \sigma(y, M)$ and $\lambda = p(0)$. Moreover, $\dim_k(H_p(z - b, M)) = \dim_k H_{p-1}(y, M) + \dim_k H_p(y, M)$ for all p , and $i(z - b) = 0$ whenever $i(y) < \infty$.*

Proof. Suppose $b \in \sigma(z, M)$ and $H_j(z - b, M) \neq 0$ for some j . Using Corollary 2.2, we derive that $0 \in \sigma(y, M)$. By Lemma 2.3, the following sequence

$$0 \leftarrow \ker(p(x) - \lambda | H_{j-1}(y, M)) \longleftarrow H_j(z - b, M) \longleftarrow \text{coker}(p(x) - \lambda | H_j(y, M)) \leftarrow 0$$

turns out to be exact. By Lemma 2.4, $p(x) H_p(y, M) = p(0) H_p(y, M)$ for every $p(x) \in R'$. Thus the actions of $p(x) - \lambda$ over the homology groups $H_i(y, M)$ are reduced to the constant multiplication operators by $p(0) - \lambda$. If $\lambda \neq p(0)$ then $p(x) - \lambda$ defines an invertible action over homology groups. Therefore the corners of the exact sequence are vanishing, which means that $H_j(z - b, M) = 0$, a contradiction. Hence $\lambda = p(0)$. Using again Lemma 2.3, for every p we obtain the following

$$0 \leftarrow H_{p-1}(y, M) \longleftarrow H_p(z - b, M) \longleftarrow H_p(y, M) \leftarrow 0$$

exact sequence. In particular, $\dim_k (H_p(z - b, M)) = \dim_k H_{p-1}(y, M) + \dim_k H_p(y, M)$ whenever $i(y) < \infty$. Finally,

$$\begin{aligned} i(z - b) &= \sum_{p=0}^{n+1} (-1)^{p+1} \dim_k (H_p(z - b, M)) \\ &= \sum_{p=0}^{n+1} (-1)^{p+1} \dim_k H_{p-1}(y, M) + (-1)^{p+1} \dim_k H_p(y, M) = 0, \end{aligned}$$

that is, $i(z - b) = 0$. \square

Remark 2.1. *The assertion just proven is equally true for the tuple $z = (p(x), y)$ swapped. Namely, if $b = (\lambda, 0) \in \sigma(z, M)$ for some $\lambda \in k$, then $0 \in \sigma(y, M)$ and $\lambda = p(0)$.*

Corollary 2.4. *If $w = (x, p(x))$ is an $n+1$ -tuple in R and $c = (a, \lambda) \in \sigma(w, M)$ for some $a \in \mathbb{A}^n$ and $\lambda \in k$, then $a \in \sigma(x, M)$ and $\lambda = p(a)$.*

Proof. Put $y = x - a$, $R' = k[y]$, $z = (y, q(y))$ and $b = (0, \lambda)$, where $q(y) = p(y + a)$. Then $z - b = (y, q(y) - \lambda) = w - c$, which means that $b \in \sigma(z, M)$. It remains to use Proposition 2.1 for y instead of x . Then $0 \in \sigma(y, M)$ (or $a \in \sigma(x, M)$) and $\lambda = q(0) = p(a)$. \square

Now let us define the following one-to-one function

$$f_n : \mathbb{Z}_+^{n+1} \rightarrow \mathbb{Z}_+^{n+2}, \tau_n(d_0, \dots, d_n) = (d_0, d_0 + d_1, d_1 + d_2, \dots, d_{n-1} + d_n, d_n)$$

that generates Fibonacci numbers by increasing $n+1$ -tuples of nonnegative integers. If x is an n -tuple from R with $i(x) < \infty$ then $d(x) = (d_0, \dots, d_n) \in \mathbb{Z}_+^{n+1}$ is a tuple of dimensions $d_p = \dim_k H_p(x, M)$, $0 \leq p \leq n$.

Corollary 2.5. *Let R/k be an algebra, $R' = k[x] \subseteq R$ a subalgebra with an n -tuple x , and related to x an m -tuple y . If $(0, \lambda) \in \sigma((x, y), M)$ and both $i(x)$ and $i(y)$ are finite, then $\lambda = 0$, $d(x) = f_{n-m-1} \dots f_m(d(y))$ and $i(x) = \delta_{nm} i(y)$ whenever $m \leq n$.*

Proof. Put $z = (x, y)$ to be $n+m$ -tuple from R . Note that y is a tuple of polynomials from R' such that $y \subseteq \langle x \rangle \subseteq R'$, that is, $y(0) = 0$. By Corollary 2.4, $0 \in \sigma(x, M)$ and $\lambda = y(0) = 0 \in \sigma(y, M)$ (see Corollary 2.2). Thus $0 \in \sigma(z, M)$ and $d(z) = f_{n+m-1} \dots f_n(d(x))$ thanks Proposition 2.1. By symmetry (see Remark 2.1), we deduce $d(z) = f_{n+m-1} \dots f_m(d(y))$ either. If $m \leq n$ then $d(x) = f_{n-m-1} \dots f_m(d(y))$ due to the property to be a one-to-one function. If $n > m$ then $i(x) = 0$ (see Proposition 2.1). But if $m = n$ then $d(x) = d(y)$ and $i(x) = i(y)$. \square

Remark 2.2. *If $R' = k[x] = k[y]$ for an m -tuple $y \subseteq \langle x \rangle$ then y is a tuple related to x . Indeed, $x_i = q_i(y) \in k[y] = R'$ for every i . It follows that $x_i = q_i(0) + h_i(y)$, $h_i(y) \in \langle y \rangle \subseteq \langle x \rangle$. Hence $q_i(0) = 0$, that is, $x_i \in \langle y \rangle$. Thus $\langle x \rangle = \langle y \rangle$ in R' .*

2.3. Noetherian modules and triangular actions. As above we have an algebra R/k and an R -module M with an n -tuple x from R . If R is Noetherian and M is a finitely generated R -module then it turns out to be Noetherian automatically. Conversely, a Noetherian R -module has a finitely many generators. For brevity we say that M is a Noetherian R -module with an acting tuple x on M . Since $M \otimes_k \wedge^i k^n$ is a Noetherian R -module, it follows that so is $H_i(x, M)$ for every i .

Lemma 2.5. *Let $R = k[x]$ be an algebra finite extension with an n -tuple x , $\mathfrak{m} = \langle x \rangle \subseteq R$ a maximal ideal, and let M be an R -module. Then $H_i(x, M)_{\mathfrak{m}} = H_i(x, M)$ for all $i \geq 0$. If M is Noetherian and $H_p(x, M) = 0$ for some p , then $H_j(x, M) = 0$ for all $j \geq p$.*

Proof. By Lemma 2.4, the action of every $p(x) \notin \mathfrak{m}$ on $H_i(x, M)$ is reduced to the diagonal operator $p(0)$, which is invertible. Therefore $H_i(x, M)_{\mathfrak{m}} = H_i(x, M)$ (see [1, 12.1, 12.4]).

Now assume that M is a Noetherian R -module and prove that $H_p(x, M) = 0$ implies that $H_j(x, M) = 0$ for all $j \geq p$. If R is a local k -algebra with its maximal ideal \mathfrak{m} and $x \subseteq \mathfrak{m}$ then the result follows thanks to Lemma 2.3. Namely, we proceed by induction on n for a nonzero Noetherian R -module M . Since $M/x_1M \neq 0$ (Nakayama lemma), the result follows in the case of $n = 1$. Put $x' = (x_1, \dots, x_{n-1})$ and suppose $H_p(x, M) = 0$. By Lemma 2.3, we obtain that $\text{coker}(x_n | H_p(x', M)) = 0$. Since $H_p(x', M)$ is a Noetherian R -module and $x_n \in \text{rad } R$, it follows that $H_p(x', M)/x_n H_p(x', M) \neq 0$ whenever $H_p(x', M) \neq 0$ (Nakayama lemma). Hence $H_p(x', M) = 0$, and $H_j(x', M) = 0$, $j \geq p$ by induction hypothesis. Using again Lemma 2.3, we obtain the exact sequence

$$0 \leftarrow \ker(x_n | H_j(x', M)) \longleftarrow H_{j+1}(x, M) \longleftarrow \text{coker}(x_n | H_{j+1}(x', M)) \leftarrow 0$$

for every $j \geq p$. It follows that $H_{j+1}(x, M) = 0$ for all $j \geq p$. Optionally one can use [12, Theorem 17.6].

In the general case, $M_{\mathfrak{m}}$ is a Noetherian $R_{\mathfrak{m}}$ -module and $H_i(x, M) = H_i(x, M)_{\mathfrak{m}} = H_i(x/1, M_{\mathfrak{m}})$ (the localization is an exact functor) $i \geq 0$, where $x/1$ is the tuple representing $(x_1/1, \dots, x_n/1) \subseteq \mathfrak{m}R_{\mathfrak{m}} = \text{rad } R_{\mathfrak{m}}$. If $H_p(x, M) = 0$ then $H_p(x/1, M_{\mathfrak{m}}) = 0$ and we come up with the local k -algebra case $R_{\mathfrak{m}}$. Based on the fact just proven, we conclude that $H_j(x/1, M_{\mathfrak{m}}) = 0$ for all $j \geq p$. Hence $H_j(x, M) = 0$ for all $j \geq p$. \square

Now we analyze tuples y from an algebra finite extension R/k which admit finite index. Everywhere below we assume that k is an algebraically closed field. A tuple y whose extension $k[y] \subseteq k[x]$ is integral plays a key role in this manner.

Theorem 2.1. *Let $k \subseteq R' \subseteq R$ be a ring extensions of the field k such that R/k is an algebra finite extension and R/R' is integral. If M is a Noetherian R -module then $i(y) < \infty$ whenever y is a tuple in R generating a maximal ideal in R' . In this case, the R -module structure on every $H_p(y, M)$ is triangularizable (or it is a semilocal R -module) whereas its R' -module structure is diagonalizable.*

Proof. Since R/R' is algebra finite and integral, it follows that R is module finite over R' [1, 10.18]. But M is module finite over R , therefore M is module finite over R' (see [1, 10.16]). Moreover, R'/k is an algebra finite extension by Artin-Tate [1, 16.17], and it is Noetherian (Hilbert Basis). In particular, M is a Noetherian R' -module either, and so are all homology groups $H_p(y, M)$ for all tuples y in R' . Assume that y is an m -tuple generating a maximal ideal $\langle y \rangle$ in R' . But $R' = k[y']$ for a certain s -tuple $y' \subseteq R'$, and the maximal ideals of R' respond to points from \mathbb{A}^s by Zariski Nullstellensatz [1, 15.4]. In particular, $\langle y \rangle = \langle y' - a' \rangle$ for some $a' \in \mathbb{A}^s$. Thus we can assume that $R' = k[y']$ and $\langle y \rangle = \langle y' \rangle$, which means that y is a tuple related to y' . By Lemma 2.4, the R' -module structure of $H_p(y, M)$ is diagonalizable, and it is just the k -vector space structure. In particular, every ascending chain of vector subspaces in $H_p(y, M)$ turns out to be a chain of R' -submodules which has to stabilize being a Noetherian R' -module. Hence $H_p(y, M)$ is a finite dimensional k -vector space, and $i(y) < \infty$.

But $H_p(y, M)$ has also Noetherian R -module structure on. In particular, $H_p(y, M)$ has the finite length as an R -module, which means that there is a Jordan-Hölder series with the quotients R/\mathfrak{m}_j for some maximal ideals $\mathfrak{m}_j \subseteq R$. Thus $H_p(y, M)$ is a semilocal R -module, that is, $\text{Supp}_R(H_p(y, M))$ is finite. As above $R = k[x]$ for some n -tuple x , and maximal ideals respond to points from \mathbb{A}^n . Hence there is a chain $0 = V_s \subsetneq V_{s-1} \subsetneq \dots \subsetneq V_0 = H_p(y, M)$ of R -submodules such that $V_j/V_{j+1} = R/\mathfrak{m}_j = k$ with $\mathfrak{m}_j = \langle x - b^{(j)} \rangle$, $b^{(j)} \in \mathbb{A}^n$, $1 \leq j \leq s$. Thus

$s = \dim_k(H_p(x, M))$ and

$$\text{Supp}_R(H_p(y, M)) = \text{Ass}_R(H_p(y, M)) = \text{Max}_R(H_p(y, M)) = \{\mathfrak{m}_1, \dots, \mathfrak{m}_s\}$$

with their multiplicities (see [1, 19.4]). Based on the chain we can easily construct a basis $\omega = (\omega_1, \dots, \omega_s)$ for $H_p(y, M)$ such that

$$x_i |_{H_p(y, M)} = \begin{bmatrix} b_i^{(1)} & & 0 \\ & \ddots & \\ * & & b_i^{(s)} \end{bmatrix}$$

for all i , $1 \leq i \leq n$ with respect to ω . In particular,

$$p(x) |_{H_p(y, M)} = \begin{bmatrix} p(b^{(1)}) & & 0 \\ & \ddots & \\ * & & p(b^{(s)}) \end{bmatrix}$$

for every $p(x) \in R$, which means that R -action on $H_p(y, M)$ is triangularizable. \square

Corollary 2.6. *Let R/k be an algebra finite extension, and let $\mathfrak{a} \subsetneq R$ be an ideal. The following assertions are equivalent: (i) there is a tuple y generating \mathfrak{a} such that $k[y] \subseteq R$ is integral; (ii) there is a tuple y generating \mathfrak{a} such that $i(y) < \infty$ for every Noetherian R -module M and the R -module structure on every $H_p(y, M)$ is triangularizable; (iii) the radical $\sqrt{\mathfrak{a}}$ is a finite intersection of maximal ideals of R .*

Proof. The implication (i) \Rightarrow (ii) is due to Theorem 2.1. Let prove the implication (ii) \Rightarrow (iii). Suppose that $\mathfrak{a} = \langle y \rangle$ for some tuple y such that $i(y) < \infty$ for every Noetherian R -module M , and the R -module structure on every $H_p(y, M)$ is triangularizable. By Lemma 2.4, $k[y]$ -module structure on $H_p(y, M)$ is diagonalizable. In particular, $\mathfrak{a} = \langle y \rangle \subseteq \text{Ann}_R(H_p(y, M))$, where $\text{Ann}_R(H_p(y, M))$ is the annihilator of the R -module $H_p(y, M)$. Taking into account that R -module $H_p(y, M)$ is semilocal (or triangularizable), we deduce that

$$\begin{aligned} \sqrt{\mathfrak{a}} &\subseteq \sqrt{\text{Ann}_R(H_p(y, M))} = \text{nil}_R(H_p(y, M)) = \cap \text{Supp}_R(H_p(y, M)) = \cap \text{Ass}_R(H_p(y, M)) \\ &= \cap_{j=1}^s \mathfrak{m}_j, \end{aligned}$$

that is, $\sqrt{\mathfrak{a}} \subseteq \cap_{j=1}^s \mathfrak{m}_j$ (see to the proof of Theorem 2.1, and [1, 13.6]). If $p = 0$ and $M = R$, then we obtain that $H_0(y, R) = R/\langle y \rangle = R/\mathfrak{a}$ and $\cap_{j=1}^s \mathfrak{m}_j = \text{nil}_R(R/\mathfrak{a}) = \sqrt{\mathfrak{a}}$.

Finally prove that (iii) \Rightarrow (i). Suppose that $\sqrt{\mathfrak{a}} = \cap_{j=1}^s \mathfrak{m}_j$ for some maximal ideals $\{\mathfrak{m}_j\}$. It is a primary decomposition and $\text{Ass}(R/\sqrt{\mathfrak{a}}) \subseteq \{\mathfrak{m}_j\} \subseteq \text{Max}(R/\sqrt{\mathfrak{a}})$ (see [1, 18.17]). Since $R/\sqrt{\mathfrak{a}}$ is Noetherian, we deduce that $\text{Spec}(R/\sqrt{\mathfrak{a}}) = \text{Max}(R/\sqrt{\mathfrak{a}})$ (see [1, 17.14]) or the Krull dimension $\dim(R/\sqrt{\mathfrak{a}}) = 0$. By Akizuki-Hopkins Theorem [1, 19.8], we conclude that $R/\sqrt{\mathfrak{a}}$ is an Artinian ring. Using again Zariski Nullstellensatz, we deduce that $\dim_k(R/\sqrt{\mathfrak{a}}) < \infty$ (the gaps of a Jordan-Hölder series are $R/\mathfrak{m}_j = k$). But $R = k[x]$ for some n -tuple x , and every x_i defines a linear transformation on $R/\sqrt{\mathfrak{a}}$. By Cayley-Hamilton Theorem, $q_i(x_i) \in \sqrt{\mathfrak{a}}$ for a monic polynomial $q_i \in k[t]$. It follows that $y_i = p_i(x_i) \in \mathfrak{a}$ for a monic polynomial $p_i \in k[t]$, $1 \leq i \leq n$. Put $y = (y_1, \dots, y_n)$ and consider the subalgebra $R' = k[y] \subseteq R$. For every i we have $(p_i - y_i)(x_i) = 0$, that is, x_i is integral over R' and $R = R'[x]$. Hence R is module finite over R' or $R' \subseteq R$ is an integral extension [1, 10.18]. But y can easily be extended up to generators of the ideal \mathfrak{a} . Just consider $\mathfrak{b} = \langle y \rangle \subseteq \mathfrak{a}$ and pick up generators from $\mathfrak{b}/\mathfrak{a}$, or just add up another generators of \mathfrak{a} to y being a Noetherian ideal. Whence $\mathfrak{a} = \langle y \rangle$ and $k[y] \subseteq R$ is an integral extension. \square

Recall that if M is a module over a k -algebra R and $t \in R$, which defines a linear transformation $t|M$ over the k -vector space M , then its spectrum $\sigma(t|M)$ is defined to be a subset of those $\lambda \in k$ such that $t - \lambda$ is not an invertible linear transformation over M . Actually, it is just the Taylor spectrum $\sigma(t, M)$ of the single tuple t from the algebra R . Note also that it is not expected that $t^{-1} \in R$ in the case of an invertible linear transformation t over M , that is, $0 \notin \sigma(t)$ (see below Example 2.2).

Proposition 2.2. *(Non voidness) Let R/k be an algebra finite extension and M a Noetherian R -module. Then $\sigma(t|M) \neq \emptyset$ for every $t \in R$. For every chain $0 = M_0 \subsetneq M_1 \subsetneq \cdots \subsetneq M_{n-1} \subsetneq M_n = M$ of submodules with $M_i/M_{i-1} = R/\mathfrak{p}_i$ and $\text{Ass}(M) \subseteq \{\mathfrak{p}_1, \dots, \mathfrak{p}_n\} \subseteq \text{Supp}(M)$, we have $\sigma(t|M) = \cup_i \sigma(t|R/\mathfrak{p}_i)$. In particular, $\sigma(t|M)$ is either finite set or it is a dense subset of \mathbb{A}^1 .*

Proof. We can assume that $M \neq 0$ and $R = k[x]$ for an n -tuple x . Then $\text{Supp}(M) \neq \emptyset$. Actually, $\text{Supp}(M) = V(\text{Ann}(M))$ (see [1, 13.4]) to be the set of all primes containing the annihilator $\text{Ann}(M)$ of M . In particular, there is a maximal ideal $\mathfrak{m} \subseteq R$ from $\text{Supp}(M)$, that is, $M_{\mathfrak{m}} \neq 0$. By Zariski Nullstellensatz, $\mathfrak{m} = \langle x - a \rangle$ for some $a \in \mathbb{A}^n$. Take $t = t(x) \in R$. Then $t - t(a) \in \mathfrak{m}$ and $\langle (t/1) - t(a) \rangle \subseteq \mathfrak{m}R_{\mathfrak{m}} = \text{rad}(R_{\mathfrak{m}})$. By Nakayama lemma, $(M/\langle t - t(a) \rangle M)_{\mathfrak{m}} = M_{\mathfrak{m}}/\langle (t/1) - t(a) \rangle M_{\mathfrak{m}} \neq 0$. It follows that $M/\langle t - t(a) \rangle M \neq 0$ or $\text{im}(t - t(a)) \neq M$, that is, $t(a) \in \sigma(t|M)$.

Further, consider a chain $0 = M_0 \subsetneq M_1 \subsetneq \cdots \subsetneq M_{n-1} \subsetneq M_n = M$ of submodules with $M_i/M_{i-1} = R/\mathfrak{p}_i$ and $\text{Ass}(M) \subseteq \{\mathfrak{p}_1, \dots, \mathfrak{p}_n\} \subseteq \text{Supp}(M)$ (see [1, 17.16]). If $t - \lambda$ is invertible on M then $(t - \lambda)^{-1}$ is an R -linear map too. Hence $t - \lambda$ is invertible iff so are all $(t - \lambda)|M_i/M_{i-1}$, that is, $\sigma(t|M) = \cup_i \sigma(t|R/\mathfrak{p}_i)$.

Finally, consider the case of $M = R/\mathfrak{p}$ for some prime \mathfrak{p} , which is the coordinate ring of the variety $Y = V(\mathfrak{p}) \cap \mathbb{A}^n \subseteq \mathbb{A}^n$. Then $\text{zdiv}(M) = \{0\}$, and $\lambda \in \sigma(t|M)$ iff $t - \lambda \in \mathfrak{m}$ for some maximal ideal \mathfrak{m} containing \mathfrak{p} . By Zariski Nullstellensatz, $\mathfrak{m} = \langle x - a \rangle$ for some $a \in Y$, that is, $\lambda = t(a)$. Hence $\sigma(t|M) = t(Y)$ is an irreducible subset of \mathbb{A}^1 (see [2, 2.4.4]). But $\sigma(t|M)$ is irreducible iff so is its closure $\sigma(t|M)^-$ in \mathbb{A}^1 (see [2, 2.4.2]). So $\sigma(t|M)^-$ is either singleton or $\sigma(t|M)^- = \mathbb{A}^1$. But $\sigma(t|M) \neq \emptyset$, therefore $\sigma(t|M)$ is either singleton or it is a dense irreducible subset of \mathbb{A}^1 . \square

Example 2.1. *As an example of the case $\sigma(t|M) = \mathbb{A}^1$ consider the action of X on the PID $k[X]$. If $\ell_R(M) < \infty$ then $\text{Ass}(M) = \text{Max}(M)$ and M is a finite dimensional k -vector space (see [1, 19.4]). In this case, t is a triangularizable whose diagonal entries consist of its finite spectrum $\sigma(t|M)$.*

Example 2.2. *The spectrum can be empty set unless M is Noetherian. Put $R = k[X]$ and $M = k(X) = \text{Frac}(R)$. Then M is an R -module and X is acting as a multiplication operator over M . For every $\lambda \in k$ we have $X - \lambda$ is invertible and $(X - \lambda)^{-1}$ is the multiplication operator on M by $1/(X - \lambda)$. Note also that $(X - \lambda)^{-1} \notin R$.*

Example 2.3. *The spectrum $\sigma(t|M)$ can be an open subset of \mathbb{A}^1 . Let M be the coordinate ring of the hyperbola $Y = \{xy = 1\}$ in \mathbb{A}^2 , and $t = x$. Then $\sigma(t|M) = t(Y) = \mathbb{A}^1 - \{0\}$ is an open dense subset of \mathbb{A}^1 .*

Now we can prove a key result of the present section.

Theorem 2.2. *(The projection property) Let R/k be an algebra finite extension of the field k , M a Noetherian R -module, $y = (y_1, \dots, y_m)$ an m -tuple in R , and let $y' = (y_1, \dots, y_{m-1})$. Then $\sigma(y', M) = \pi(\sigma(y, M))$, where $\pi : \mathbb{A}^m \rightarrow \mathbb{A}^{m-1}$ is the canonical projection onto first $m - 1$ coordinates.*

Proof. The inclusion $\pi(\sigma(y, M)) \subseteq \sigma(y', M)$ holds in the general case thanks to Corollary 2.2. Conversely, take $a' \in \sigma(y', M)$. Then $H_p(y' - a', M) \neq 0$ for some p . But $H_p(y' - a', M)$ is a Noetherian R -module, and y_m defines a linear transformation on it. By Proposition 2.2, $\sigma(y_m|H_p(y' - a', M)) \neq \emptyset$, that is,

$$y_m - a_m : H_p(y' - a', M) \rightarrow H_p(y' - a', M)$$

is not invertible for some $a_m \in k$. If $\text{coker}(y_m - a_m|H_p(y' - a', M)) \neq 0$ then using the following exact (Lemma 2.3) sequence

$$0 \leftarrow \ker(y_m - a_m|H_{p-1}(y' - a', M)) \longleftarrow H_p(y - a, M) \longleftarrow \text{coker}(y_m - a_m|H_p(y' - a', M)) \leftarrow 0,$$

we conclude that $H_p(y - a, M) \neq 0$. If $\ker(y_m - a_m|H_p(y' - a', M)) \neq 0$ then using the exact sequence

$$0 \leftarrow \ker(y_m - a_m|H_p(y' - a', M)) \longleftarrow H_{p+1}(y - a, M) \longleftarrow \text{coker}(y_m - a_m|H_{p+1}(y' - a', M)) \leftarrow 0,$$

we conclude that $H_{p+1}(y - a, M) \neq 0$. Hence either $H_p(y - a, M) \neq 0$ or $H_{p+1}(y - a, M) \neq 0$. Anyway $a \in \sigma(y, M)$ and $\pi(a) = a'$. \square

Corollary 2.7. *Let R/k be an algebra finite extension of the field k and let M be a Noetherian R -module. Then $\sigma(y, M) \neq \emptyset$ for every tuple y in R .*

Proof. Indeed, using Theorem 2.2, we conclude that $\pi_1(\sigma(y, M)) = \sigma(y_1, M) = \sigma(y_1|M)$, where $\pi_1 : \mathbb{A}^m \rightarrow \mathbb{A}^1$ is the canonical projection onto the first coordinate. By Proposition 2.2, $\sigma(y_1|M) \neq \emptyset$. Therefore $\sigma(y, M) \neq \emptyset$. \square

Nonetheless the point spectrum $\sigma_p(y, M)$ could be an empty set, which detects only zero-divisors in the module M .

2.4. The spectral mapping property. Finally, let us prove the main result for the present section. As above we fix an algebra finite extension R/k , an n -tuple with $x = (x_1, \dots, x_n)$ from R , and an m -tuple $p(x) = (p_1(x), \dots, p_m(x))$ from the subalgebra $k[x]$. If M is an R -module then $k[p(x)] \subseteq R$ is a subalgebra and M has the natural $k[p(x)]$ -module structure. Moreover, the tuple $p(x)$ defines a morphism mapping $p : \mathbb{A}^n \rightarrow \mathbb{A}^m$, $a \mapsto p(a) = (p_1(a), \dots, p_m(a))$.

Theorem 2.3. *If R/k is an algebra finite extension of the field k , M a Noetherian R -module, then $\sigma(p(x), M) = p(\sigma(x, M))$ for all tuples x and $p(x)$ from R .*

Proof. Consider the $n + m$ -tuple $y = (x, p(x))$ in R . By Theorem 2.2, $\sigma(x, M) = \pi_n(\sigma(y, M))$, where $\pi_n : \mathbb{A}^{n+m} \rightarrow \mathbb{A}^n$ is the canonical projection onto the first n coordinates. Take $(a, b) \in \sigma(y, M)$. By Corollary 2.4, $b_j = p_j(a)$ for all j , $1 \leq j \leq m$, that is, $b = p(a)$. Thus $\sigma(y, M) = \{(a, p(a)) : a \in \sigma(x, M)\}$. Using again Theorem 2.2, we conclude that $\sigma(p(x), M) = \pi_m(\sigma(y, M)) = p(\sigma(x, M))$, where $\pi_m : \mathbb{A}^{n+m} \rightarrow \mathbb{A}^m$ is the canonical projection onto the last m coordinates. \square

Corollary 2.8. *If R/k is an algebra finite extension of the field k , M a Noetherian R -module, then $\sigma(p(x)|M) = p(\sigma(x, M))$ for every polynomial $p(x) \in R$.*

Corollary 2.9. *Let R/k be an algebra, $R' = k[x] \subseteq R$ a k -subalgebra with an n -tuple x such that R/R' is integral, and let M be a Noetherian R -module. If y is an m -tuple in R' related to x with $m \leq n$, and $0 \in \sigma(x, M)$, then $d(x) = f_{n-m-1} \dots f_m(d(y))$ and $i(x) = \delta_{nm}i(y)$.*

Proof. By Theorem 2.3, $(0, \lambda) \in \sigma((x, y), M)$ for a certain $\lambda \in \mathbb{A}^m$. Actually, $\lambda = y(0) = 0$, and $0 \in \sigma((x, y), M)$. Since $R' = k[x]$ and y generates the maximal ideal $\langle x \rangle$ in R' , we conclude that $i(x) < \infty$ and $i(y) < \infty$ thanks to Theorem 2.1. It remains to use Corollary 2.5. \square

3. SPECTRA OF A MODULE OVER A SCHEME AND EXTENSION PROPERTY

In the present section we review some key facts on spectrum of a module over a scheme from [11], and use them regarding our framework of spectra. We also introduce the point spectrum in the general setting and investigate its invariance under integral extensions. Thereafter we switch to the case of the affine scheme \mathbb{A}_k^n over a field k .

3.1. Spectra of a module over scheme. Let $(\mathfrak{X}, \mathcal{O}_{\mathfrak{X}})$ be a scheme with the ring $R = \Gamma(\mathfrak{X}, \mathcal{O}_{\mathfrak{X}})$ of global sections, and let $M \in R\text{-mod}$. We say that a point $x \in \mathfrak{X}$ belongs to a *resolvent set* $\text{res}(\mathfrak{X}, M)$ if there is an affine neighborhood U of x such that $\mathcal{O}_{\mathfrak{X}}(U) \perp_R M$, that is, $\text{Tor}_i^R(\mathcal{O}_{\mathfrak{X}}(U), M) = 0$ for all $i \geq 0$. Since U is an affine neighborhood for all points close to x , it follows that $\text{res}(\mathfrak{X}, M)$ is an open set, whose complement set $\sigma(\mathfrak{X}, M)$ is called *the spectrum of M over the scheme*. The property $\mathcal{O}_{\mathfrak{X}}(U) \perp_R M$ implies that $\mathcal{O}_{\mathfrak{X}}(V) \perp_R M$ for every open affine $V \subseteq U$. Indeed, suppose $U = \text{Spec}(B)$ and take a free resolution

$$\mathcal{P} = R \otimes G_{\circ}, \quad 0 \leftarrow R \otimes G_0 \xleftarrow{\partial_0} R \otimes G_1 \xleftarrow{\partial_1} \dots$$

of the module M , where G_j are free abelian groups. Since $\mathcal{O}_{\mathfrak{X}}$ is a quasi-coherent sheaf, it follows that $\mathcal{O}_{\mathfrak{X}}|_U = \tilde{B}$ with $\mathcal{O}_{\mathfrak{X}}(U) = B$, and $B \otimes_R \mathcal{P} = B \otimes G_{\circ}$, which is exact iff so are $B_x \otimes G_{\circ}$, $x \in \text{Spec}(B)$ (see [11, Lemma 2.1]). But $B_x = \mathcal{O}_{\mathfrak{X},x}$ for all $x \in U$. Thus $\mathcal{O}_{\mathfrak{X}}(U) \perp_R M$ is equivalent to $\mathcal{O}_{\mathfrak{X},x} \perp_R M$ for all $x \in U$. In particular, $\mathcal{O}_{\mathfrak{X},x} \perp_R M$ for all $x \in V$ imply that $\mathcal{O}_{\mathfrak{X}}(V) \perp_R M$.

If $\mathfrak{X} = \text{Spec}(R)$ is an affine scheme, then $\mathcal{O}_{\mathfrak{X},x} \perp_R M$ means that $R_x \otimes_R \mathcal{P}$ is exact. But R_x is a flat R -module, therefore the sequence $0 \leftarrow R_x \otimes_R M \leftarrow R_x \otimes_R \mathcal{P}$ remains exact. Hence $\mathcal{O}_{\mathfrak{X},x} \perp_R M$ iff $M_x = 0$. Thus $x \in \text{res}(\mathfrak{X}, M)$ iff $M_y = 0$ for all y from a neighborhood of x . Since $\sigma(\mathfrak{X}, M) = \mathfrak{X} - \text{res}(\mathfrak{X}, M)$, we derive that $x \in \sigma(\mathfrak{X}, M)$ such that property $M_y \neq 0$ holds for some y in every neighborhood of x , that is, $\sigma(\mathfrak{X}, M)$ is the closure of $\text{Supp}(M)$ in \mathfrak{X} . Thus

$$(3.1) \quad \sigma(\mathfrak{X}, M) = \text{Supp}(M)^- \subseteq V(\text{Ann}(M)).$$

If M is a finitely generated R -module then $\text{Supp}(M)$ is closed, $\text{res}(\mathfrak{X}, M) = \{x \in \mathfrak{X} : R_x \perp_R M\} = \{x \in \mathfrak{X} : M_x = 0\}$, that is,

$$(3.2) \quad \sigma(\mathfrak{X}, M) = \text{Supp}(M) = V(\text{Ann}(M)).$$

In the case of a general scheme the spectrum can be localized using an affine covering. In particular, $\sigma(\mathbb{P}_k^r, M) = \mathbb{P}_k^r$ whenever M is a nonzero vector space over a field k (see [11]). The related spectral mapping formula was mentioned in (1.2).

3.2. The point spectrum. Let $\mathfrak{X} = \text{Spec}(R)$ be an affine scheme and let M be an R -module. The set $\text{Ass}_R(M)$ of all primes associated to M is called *the point spectrum* $\sigma_p(\mathfrak{X}, M)$ of M . If R is Noetherian then $\text{Supp}(M) = \bigcup \{V(\mathfrak{p}) : \mathfrak{p} \in \sigma_p(\mathfrak{X}, M)\}$ and the minimal primes $\text{Min}(M)$ from $\text{Supp}(M)$ belong to $\sigma_p(\mathfrak{X}, M)$ [1, 17.14]. In particular, $\sigma_p(\mathfrak{X}, M)$ is dense in $\text{Supp}(M)$, which in turn implies that $\sigma_p(\mathfrak{X}, M)^- = \sigma(\mathfrak{X}, M)$ in \mathfrak{X} (see (3.1)).

Now let $R' \subseteq R$ be a ring extension, $\iota : R' \rightarrow R$ is the realted inclusion map, and $\mathfrak{X}' = \text{Spec}(R')$ with the canonical mapping $\iota^* : \mathfrak{X} \rightarrow \mathfrak{X}'$, $\iota^*(\mathfrak{p}) = \iota^{-1}(\mathfrak{p}) = \mathfrak{p} \cap R$. If $\mathfrak{p}' = \mathfrak{p} \cap R = \iota^*(\mathfrak{p})$ we say that \mathfrak{p} lies over the prime \mathfrak{p}' . Note also that M has the canonical R' -module structure along the embedding ι . If $\mathfrak{a} = \text{Ann}(m)$ is the annihilator of $m \in M$ in R then $\mathfrak{a}' = \mathfrak{a} \cap R' = \text{Ann}'(m)$ is the annihilator of m in R' . It follows that $\iota^*(\sigma_p(\mathfrak{X}, M)) \subseteq \sigma_p(\mathfrak{X}', M)$.

Notice also that $\iota^*(\text{Supp}(M)) \subseteq \text{Supp}'(M)$, where $\text{Supp}'(M) = \text{Supp}_{R'}(M)$. Indeed, if $M_{\mathfrak{p}} \neq \{0\}$ for some $\mathfrak{p} \in \mathfrak{X}$, then $sm \neq 0$ for some $m \in M$ and all $s \in R - \mathfrak{p}$. But $\mathfrak{p}' = \mathfrak{p} \cap R'$, and for

$s' \in R'$ we have $s' \in R' - \mathfrak{p}'$ iff $s' \in R - \mathfrak{p}$. Hence $s'm \neq 0$ for all $s' \in R' - \mathfrak{p}'$, which means that $m/1 \neq 0$ in $M_{\mathfrak{p}'}$ or $\mathfrak{p}' \in \text{Supp}'(M)$.

Using (3.1) and continuity of the map ι^* , we deduce that $\iota^*(\sigma(\mathfrak{X}, M)) \subseteq \sigma(\mathfrak{X}', M)$. Actually $\iota^*(\sigma(\mathfrak{X}, M))$ is dense in $\sigma(\mathfrak{X}', M)$ due to the spectral mapping property (1.2).

Theorem 3.1. *If $R' \subseteq R$ is a ring extension and $M \in R\text{-mod}$ then $\sigma_{\mathfrak{p}}(\mathfrak{X}', M) = \iota^*(\sigma_{\mathfrak{p}}(\mathfrak{X}, M))$ whenever R is Noetherian.*

Proof. First note that $M \neq \{0\}$ iff $\sigma_{\mathfrak{p}}(\mathfrak{X}, M) \neq \emptyset$ (see [1, 17.10]), and the result follows in the case of $\sigma_{\mathfrak{p}}(\mathfrak{X}, M) = \emptyset$. Thus we can assume that $M \neq \{0\}$, and take $\mathfrak{q} \in \sigma_{\mathfrak{p}}(\mathfrak{X}', M)$. Then $\iota_{\mathfrak{p}} : R'_{\mathfrak{q}} \rightarrow R_{\mathfrak{q}}$ is a ring extension and $M_{\mathfrak{q}} = R'_{\mathfrak{q}} \otimes_{R'} M = R'_{\mathfrak{q}} \otimes_{R'} R \otimes_R M = R_{\mathfrak{q}} \otimes_R M$ is an $R_{\mathfrak{q}}$ -module either, where $R_{\mathfrak{q}} = \iota(R' - \mathfrak{q})^{-1} R$. Moreover, $\mathfrak{q}R'_{\mathfrak{q}} \in \text{Ass}_{R'_{\mathfrak{q}}}(M_{\mathfrak{q}})$ [1, 17.8], say $\mathfrak{q}R'_{\mathfrak{q}} = \text{Ann}'(m/1)$ for some $m/1 \in M_{\mathfrak{q}} \setminus \{0\}$. If $\text{Ann}(m/1)$ is the annihilator of $m/1$ in $R_{\mathfrak{q}}$ then $\text{Ann}(m/1) \cap R'_{\mathfrak{q}} = \text{Ann}'(m/1) = \mathfrak{q}R'_{\mathfrak{q}}$. Pick

$$\mathfrak{S} = \{ \text{Ann}(n) : n \in M_{\mathfrak{q}}, \text{Ann}(n) \cap R'_{\mathfrak{q}} = \mathfrak{q}R'_{\mathfrak{q}} \},$$

which is a nonempty set of ideals of the ring $R_{\mathfrak{q}}$. Since R is Noetherian, so is $R_{\mathfrak{q}}$ and \mathfrak{S} has a maximal element $\mathfrak{l} = \text{Ann}(n)$ for some $n \in M_{\mathfrak{q}}$. If $x \in R_{\mathfrak{q}} - \mathfrak{l}$ then $xn \neq 0$ (or $1 \notin \text{Ann}(xn)$), $\mathfrak{l} \subseteq \text{Ann}(xn) \neq R_{\mathfrak{q}}$, which in turn implies that

$$\mathfrak{q}R'_{\mathfrak{q}} = \text{Ann}(n) \cap R'_{\mathfrak{q}} \subseteq \text{Ann}(xn) \cap R'_{\mathfrak{q}} \subsetneq R'_{\mathfrak{q}}.$$

But $R'_{\mathfrak{q}}$ is local with its unique maximal ideal $\mathfrak{q}R'_{\mathfrak{q}}$, therefore $\mathfrak{q}R'_{\mathfrak{q}} = \text{Ann}(xn) \cap R'_{\mathfrak{q}}$ or $\text{Ann}(xn) \in \mathfrak{S}$. It follows that $\mathfrak{l} = \text{Ann}(xn)$ whenever $x \in R_{\mathfrak{q}} - \mathfrak{l}$. In particular, if $xy \in \mathfrak{l}$ with $x \in R_{\mathfrak{q}} - \mathfrak{l}$ then $yx = 0$ or $y \in \text{Ann}(xn) = \mathfrak{l}$, which means that \mathfrak{l} is a prime or $\mathfrak{l} \in \text{Ass}_{R_{\mathfrak{q}}}(M_{\mathfrak{q}})$, and it is lying over $\mathfrak{q}R'_{\mathfrak{q}}$. But $\mathfrak{l} = \mathfrak{p}R_{\mathfrak{q}}$ for some $\mathfrak{p} \in \text{Ass}_R(M)$ [1, 17.8], [2, 4.1.5] with $\mathfrak{p} \cap (R' - \mathfrak{q}) = \emptyset$ (or $\mathfrak{p} \cap R' \subseteq \mathfrak{q}$). Prove that $\mathfrak{p} \cap R' = \mathfrak{q}$. Take $x' \in \mathfrak{q}$. Then $x'/1 \in \mathfrak{q}R'_{\mathfrak{q}} \subseteq \text{Ann}(n) = \mathfrak{p}R_{\mathfrak{q}}$ or $x'/1 = x/t'$ for some $x \in \mathfrak{p}$ and $t' \in R' - \mathfrak{q}$. It follows that $sx' \in \mathfrak{p}$ for some $s \in R' - \mathfrak{q}$. Taking into account that $s \in R - \mathfrak{p}$, we conclude that $x' \in \mathfrak{p}$ or $x' \in \mathfrak{p} \cap R'$. Hence $\mathfrak{q} = \mathfrak{p} \cap R' = \iota^*(\mathfrak{p})$ and $\mathfrak{p} \in \sigma_{\mathfrak{p}}(\mathfrak{X}, M)$. \square

Remark 3.1. *Note that $\text{Supp}'(M)$ can be much larger than $\iota^*(\text{Supp}(M))$. For example, put $R' = \mathbb{Z} \subset \mathbb{Q} = R$ and $M = \mathbb{Q}$. Then $\sigma(\mathfrak{X}, M) = \text{Supp}(M) = \{0\}$ whereas $\sigma(\mathfrak{X}', M) = \text{Supp}'(M)^- = \text{Spec}(\mathbb{Z}) = \mathfrak{X}'$. But $\sigma_{\mathfrak{p}}(\mathfrak{X}', M) = \text{Ass}_{\mathbb{Z}}(\mathbb{Q}) = \{0\} = \iota^*(\{0\}) = \iota^*(\text{Ass}_{\mathbb{Q}}(\mathbb{Q})) = \iota^*(\sigma_{\mathfrak{p}}(\mathfrak{X}, M))$, and it is dense in \mathfrak{X}' .*

Corollary 3.1. *Let k be a field and let $k \subseteq R' \subseteq R$ be k -algebra extensions such that R/k is algebra-finite. Then $\sigma_{\mathfrak{p}}(\mathfrak{X}', M) = \iota^*(\sigma_{\mathfrak{p}}(\mathfrak{X}, M))$ for every R -module M .*

Proof. The ring R is Noetherian by Hilbert Basis Theorem. It remains to use Theorem 3.1. \square

Corollary 3.2. *Let $R' \subseteq R$ be a ring extension with Noetherian R , $M \in R\text{-mod}$ and let $Q \subseteq M$ be an R -submodule. If Q is \mathfrak{p} -primary then Q is \mathfrak{p}' -primary R' -submodule of M , where $\mathfrak{p} \in \mathfrak{X}$ and $\mathfrak{p}' = \iota^*(\mathfrak{p})$. If M is a Noetherian R -module with its submodule Q then Q is \mathfrak{p}' -primary R' -submodule of M iff $Q = Q_1 \cap \dots \cap Q_r$ admits an irredundant primary decomposition in $R\text{-mod}$ such that every Q_j is \mathfrak{p}' -primary R' -module.*

Proof. Suppose that Q is a \mathfrak{p} -primary submodule, that is, $\text{Ass}_R(M/Q) = \{\mathfrak{p}\}$ or $\sigma_{\mathfrak{p}}(\mathfrak{X}, M/Q) = \{\mathfrak{p}\}$. Since $\mathfrak{p}' = \iota^*(\mathfrak{p}) \in \iota^*(\sigma_{\mathfrak{p}}(\mathfrak{X}, M/Q)) \subseteq \sigma_{\mathfrak{p}}(\mathfrak{X}', M/Q)$, it follows that $\mathfrak{p}' \in \text{Ass}_{R'}(M/Q)$. Conversely, if $\mathfrak{q} \in \text{Ass}_{R'}(M/Q)$ then

$$\mathfrak{q} \in \sigma_{\mathfrak{p}}(\mathfrak{X}', M/Q) = \iota^*(\sigma_{\mathfrak{p}}(\mathfrak{X}, M/Q)) = \iota^*(\text{Ass}_R(M/Q)) = \iota^*(\{\mathfrak{p}\}) = \{\mathfrak{p}'\}$$

thanks to Theorem 3.1, that is, $\mathfrak{q} = \mathfrak{p}'$. Hence $\text{Ass}_{R'}(M/Q) = \{\mathfrak{p}'\}$, which means that Q is a \mathfrak{p}' -primary R' -submodule of M .

Finally, assume that M is a finitely generated R -module. By Lasker-Noether Theorem [1, 18.19], $Q = Q_1 \cap \cdots \cap Q_r$ admits an irredundant primary decomposition in $R\text{-mod}$, where Q_j is \mathfrak{p}_j -primary. In this case, $\{\mathfrak{p}_1, \dots, \mathfrak{p}_r\}$ are uniquely defined, in fact they are all distinct primes of $\text{Ass}_R(M/Q)$ (First Uniqueness [1, 18.18]). In particular,

$$\text{Ass}_{R'}(M/Q) = \sigma_{\mathfrak{p}}(\mathfrak{X}', M/Q) = \iota^*(\text{Ass}_R(M/Q)) = \iota^*(\{\mathfrak{p}_1, \dots, \mathfrak{p}_r\})$$

by virtue of Theorem 3.1. If Q is \mathfrak{p}' -primary R' -submodule then $\text{Ass}_{R'}(M/Q) = \{\mathfrak{p}'\}$ and

$$\text{Ass}_{R'}(M/Q_j) = \sigma_{\mathfrak{p}}(\mathfrak{X}', M/Q_j) = \iota^*(\text{Ass}_R(M/Q_j)) = \iota^*(\{\mathfrak{p}_j\}) = \{\mathfrak{p}'\}$$

for every j . Thus every Q_j is \mathfrak{p}' -primary R' -module.

Conversely, suppose that $Q = Q_1 \cap \cdots \cap Q_r$ admits an irredundant primary decomposition in $R\text{-mod}$ such that every Q_j is \mathfrak{p}' -primary R' -module. Then $\iota^*(\{\mathfrak{p}_j\}) = \{\mathfrak{p}'\}$ for every j , and $\text{Ass}_{R'}(M/Q) = \iota^*(\{\mathfrak{p}_1, \dots, \mathfrak{p}_r\}) = \{\mathfrak{p}'\}$, which means that Q is a \mathfrak{p}' -primary R' -module. \square

3.3. Integral extensions. Now assume that $R' \subseteq R$ is an integral extension of rings and $M \in R\text{-mod}$. In this case, the mapping $\iota^* : \mathfrak{X} \rightarrow \mathfrak{X}'$ is surjective due to Krull-Cohen-Seidenberg Theory [1, Ch. 14], [2, 5.2].

Proposition 3.1. *If $R' \subseteq R$ is integral and $M \in R\text{-mod}$ then $\text{Supp}'(M) = \iota^*(\text{Supp}(M))$.*

Proof. The inclusion $\iota^*(\text{Supp}(M)) \subseteq \text{Supp}'(M)$ was proved above. Conversely, take $\mathfrak{p}' \in \text{Supp}'(M)$. Then $M_{\mathfrak{p}'} \neq \{0\}$, which means that $s'm \neq 0$ for all $s' \in R' - \mathfrak{p}'$ and some $m \in M$. Put $\mathfrak{a} = \text{Ann}(m) \subseteq R$ and $\mathfrak{a}' = \text{Ann}'(m) \subseteq R'$ to be ideals with $\mathfrak{a}' = \mathfrak{a} \cap R'$. Then $\mathfrak{a}' \subseteq \mathfrak{p}'$, and using the key property Going Up [1, 14.3], we deduce that $\mathfrak{p}' = \iota^*(\mathfrak{p})$ for some $\mathfrak{p} \in \mathfrak{X}$ such that $\mathfrak{a} \subseteq \mathfrak{p}$. The latter means that $sm \neq 0$ for all $s \in R - \mathfrak{p}$, that is, $m/1 \neq 0$ in $M_{\mathfrak{p}}$. Thus $\mathfrak{p} \in \text{Supp}(M)$ and $\mathfrak{p}' = \iota^*(\mathfrak{p}) \in \iota^*(\text{Supp}(M))$. \square

Remark 3.2. *In the case of an integral algebra finite extension R/R' the set $(\iota^*)^{-1}(\mathfrak{q})$ is finite for every $\mathfrak{q} \in \text{Supp}'(M)$. Indeed, since R/R' is module finite, it follows that so is $R_{\mathfrak{q}}/R'_{\mathfrak{q}}$, which in turn implies that A/k is module finite either, where $A = R_{\mathfrak{q}}/\mathfrak{q}R_{\mathfrak{q}}$ and $k = R'_{\mathfrak{q}}/\mathfrak{q}R'_{\mathfrak{q}}$ is the residue field of the local ring $R'_{\mathfrak{q}}$. In particular, $\dim_k(A) < \infty$, and A turns out to be an Artinian ring. By Akizuki-Hopkins Theorem [1, 19.8], $\dim(A) = 0$, $\text{Spec}(A)$ consists of maximal ideals and it is finite. But $(\iota^*)^{-1}(\mathfrak{q})$ is canonically identified with a subset of $\text{Spec}(A)$.*

Corollary 3.3. *If $R' \subseteq R$ is integral with Noetherian R and $M \in R\text{-mod}$ with finite length $\ell_{R'}(M) < \infty$ as an R' -module then $\ell_R(M) < \infty$.*

Proof. Since $\ell_{R'}(M) < \infty$, it follows that $\text{Supp}'(M)$ consists of maximal ideals by Jordan-Hölder Theorem [1, 19.3]. By Proposition 3.1, $\text{Supp}'(M) = \iota^*(\text{Supp}(M))$ and $\text{Supp}(M)$ consists of primes lying over the maximal ideals from $\text{Supp}'(M)$. By Maximality [1, 14.3], $\text{Supp}(M)$ consists of maximal ideals of R either. But M is a finitely generated R' -module having finite length $\ell_{R'}(M)$, therefore so is M as an R -module. It follows that $\ell_R(M) < \infty$ (see [1, 19.4 or 17.16]). \square

Remark 3.3. *Let k be a field, $k \subseteq R' \subseteq R$ algebra extensions such that R/k is algebra-finite and R/R' is integral, and let $M \in R\text{-mod}$. Then $\ell_R(M) < \infty$ iff $\ell_{R'}(M) < \infty$. By Artin-Tate lemma [1, 16.21], R'/k is algebra-finite too. In particular, R' is Noetherian by Hilbert Basis. If $\{M_i : 0 \leq i \leq n\}$ is a Jordan-Hölder chain of submodules in $R'\text{-mod}$ (or $R\text{-mod}$) then $M_i/M_{i-1} = R'/\mathfrak{m}'_i$ for some maximal ideals \mathfrak{m}'_i , $i \geq 1$. But $k \subseteq R'/\mathfrak{m}'_i$ is an algebra finite extension, which is finite by Zariski Nullstellensatz [1, 15.4]. It follows that $\dim_k(M_i/M_{i-1}) < \infty$ for every i . Hence $\dim_k(M) < \infty$. Thus $\ell_{R'}(M) < \infty$ iff $\dim_k(M) < \infty$, which in turn is equivalent to $\ell_R(M) < \infty$.*

3.4. Algebraic varieties and spectra. Now let $P = k[X]$ be the algebra of all polynomials in n variables $X = (X_1, \dots, X_n)$ over the field k . The P -bimodule $P \otimes_k P$ has the n -tuple $T = (T_1, \dots, T_n)$ of mutually commuting operators $T_i = 1 \otimes X_i - X_i \otimes 1$ acting on, and we have the Koszul complex $\text{Kos}(T, P \otimes_k P)$ augmented with the multiplication (bi)operator $\pi : P \otimes_k P \rightarrow P$ provides a free P -bimodule resolution of the algebra P . Actually, $\text{Kos}(T, P \otimes_k P)$ splits as a complex of P -bimodules. If $M \in P\text{-mod}$ with the actions $x_i(m) = X_i \cdot m$, $1 \leq i \leq n$, then derive that the complex $0 \leftarrow P \otimes_P M \xleftarrow{1 \otimes \pi} \text{Kos}(P \otimes_k P, T) \otimes_P M$ remains exact. But $P \otimes_P M = M$ and $\text{Kos}(T, P \otimes_k P) \otimes_P M = \text{Kos}(t, P \otimes_k M)$ with $t_i = 1 \otimes x_i - X_i \otimes 1$, and $1 \otimes \pi$ is reduced to the homomorphism $\pi_M : P \otimes_k M \rightarrow M$, $\pi_M(r \otimes_k m) = rm$. Thus $\mathcal{P} = \text{Kos}(t, P \otimes_k M)$ provides a finite free resolution of the module M . If R/k is an algebra finite extension with $R = k[x]$ for an n -tuple x , then R is a quotient of P and $\text{Kos}(t, P \otimes_k R)$ turns out to be a free R -module resolution of R either. Therefore it splits in $R\text{-mod}$.

Consider the affine space $\mathfrak{X} = \mathbb{A}_k^n$ over k . Then $\mathfrak{p} \in \text{res}(\mathfrak{X}, M)$ iff there is an open affine neighborhood $U = \text{Spec}(B)$ of \mathfrak{p} such that $B \perp_P M$. It means that

$$0 \leftarrow B \otimes_P M \longleftarrow B \otimes_P \mathcal{P} = \text{Kos}(t^B, B \otimes_k M)$$

with $t_i^B = 1 \otimes x_i - (X_i|_U) \otimes 1$ is exact. Put $k(\mathfrak{p}) = P_{\mathfrak{p}}/\mathfrak{p}P_{\mathfrak{p}} = (P/\mathfrak{p})_{\mathfrak{p}} = \text{Frac}(P/\mathfrak{p})$ to be residue field of \mathfrak{p} . Then $k(\mathfrak{p}) \otimes_P \mathcal{P} = \text{Kos}(t^{\mathfrak{p}}, k(\mathfrak{p}) \otimes_k M)$ with $t_i^{\mathfrak{p}} = 1 \otimes x_i - X_i(\mathfrak{p}) \otimes 1$, $1 \leq i \leq n$.

The following key result was proved in [11].

Theorem 3.2. *Let $\mathfrak{X} = \mathbb{A}_k^n$ be the affine space over k , $P = k[X_1, \dots, X_n]$ and let $M \in P\text{-mod}$. If $\mathfrak{p} \in \text{res}(\mathfrak{X}, M)$ then $k(\mathfrak{p}) \perp_P M$, that is, the complex $\text{Kos}(t^{\mathfrak{p}}, k(\mathfrak{p}) \otimes_k M)$ is exact. Thus*

$$\text{res}(\mathfrak{X}, M) \subseteq \{\mathfrak{p} \in \mathfrak{X} : k(\mathfrak{p}) \perp_P M\} \subseteq \{\mathfrak{p} \in \mathfrak{X} : M_{\mathfrak{p}} = \mathfrak{m}_{\mathfrak{p}} M_{\mathfrak{p}}\},$$

where $\mathfrak{m}_{\mathfrak{p}} = \text{rad}(P_{\mathfrak{p}})$. If M is a Noetherian P -module then $\text{res}(\mathfrak{X}, M) = \{\mathfrak{p} \in \mathfrak{X} : k(\mathfrak{p}) \perp_P M\}$.

If $\mathfrak{p} = \langle X - a \rangle$ is a maximal ideal corresponding to a closed point $a \in \mathfrak{X}$ then $k(\mathfrak{p}) = \text{Frac}(P/\mathfrak{p}) = k$ and $k(\mathfrak{p}) \otimes_P \mathcal{P} = \text{Kos}(t^{\mathfrak{p}}, M)$ with $t_i^{\mathfrak{p}} = x_i - a_i$, $1 \leq i \leq n$, that is, $t^{\mathfrak{p}} = x - a$. Thus if M is a finitely generated P -module, then $\sigma(\mathfrak{X}, M) = \text{Supp}(M)$ (see (3.2)), and $a \in \text{res}(\mathfrak{X}, M)$ iff $\text{Kos}(x - a, M)$ is exact thanks to Theorem 3.2.

Lemma 3.1. *Let R/k be an algebra finite extension of the field k , $\mathfrak{X} = \text{Spec}(R)$, M an R -module, and let $R = k[x]$ for an n -tuple x . Then M is a P -module, $\mathfrak{X} \subseteq \mathbb{A}_k^n$ up to a homeomorphism, $\text{Supp}_P(M) = \text{Supp}_R(M)$, $\sigma(\mathfrak{X}, M) = \sigma(\mathbb{A}_k^n, M)$, $\sigma_{\mathfrak{p}}(\mathfrak{X}, M) = \sigma_{\mathfrak{p}}(\mathbb{A}_k^n, M)$, and*

$$\sigma_{\mathfrak{p}}(x, M) = \sigma_{\mathfrak{p}}(\mathbb{A}_k^n, M) \cap \mathbb{A}^n.$$

If M is a Noetherian module then

$$\sigma(x, M) = \sigma(\mathbb{A}_k^n, M) \cap \mathbb{A}^n$$

and it is a nonempty closed subset of \mathbb{A}^n whose closure in the scheme \mathbb{A}_k^n is reduced to $\sigma(\mathfrak{X}, M)$.

Proof. Put $R = P/\mathfrak{a}$ for some ideal $\mathfrak{a} \subseteq P$. Then M turns out to be a P -module with the actions $X_i u = x_i u$, $u \in M$, $1 \leq i \leq n$ and $\mathfrak{a} \subseteq \text{Ann}_P(M)$. Note that $\mathfrak{X} = V(\mathfrak{a}) \subseteq \text{Spec}(P) = \mathbb{A}_k^n$ up to a homeomorphism. Moreover, $M_{\mathfrak{p}} = M_{\mathfrak{p}'}$ whenever $\mathfrak{p} \in V(\mathfrak{a})$ and $\mathfrak{p}' = \mathfrak{p}/\mathfrak{a}$. Indeed, $R_{\mathfrak{p}'} = (R - \mathfrak{p}')^{-1} R = (P - \mathfrak{p})^{-1} R = R_{\mathfrak{p}} = P_{\mathfrak{p}} \otimes_P R$ (see [1, (11.15.1)]) and

$$M_{\mathfrak{p}'} = R_{\mathfrak{p}'} \otimes_R M = P_{\mathfrak{p}} \otimes_P R \otimes_R M = P_{\mathfrak{p}} \otimes_P M = M_{\mathfrak{p}}.$$

In particular, $\text{Supp}_P(M) = \text{Supp}_R(M)$. But $R/\mathfrak{p}' = (P/\mathfrak{a})/(\mathfrak{p}/\mathfrak{a}) = P/\mathfrak{p}$, which means that $R/\mathfrak{p}' \hookrightarrow M$ iff $P/\mathfrak{p} \hookrightarrow M$. Thus $\text{Ass}_P(M) = \text{Ass}_R(M)$ (see [1, 17.3]) or $\sigma_{\mathfrak{p}}(\mathbb{A}_k^n, M) = \sigma_{\mathfrak{p}}(\mathfrak{X}, M)$. Using (3.1), we obtain that $\sigma(\mathfrak{X}, M) = \text{Supp}_R(M)^- = \text{Supp}_P(M)^- = \sigma(\mathbb{A}_k^n, M)$.

The variety \mathbb{A}^n is the set of all closed points in \mathbb{A}_k^n obtained by means of the natural homeomorphism $\beta : \mathbb{A}^n \rightarrow \mathbb{A}_k^n$, $\beta(a) = \langle X - a \rangle$ (see [16, 2.2.6]) onto the closed points. Put $\sigma_c(\mathbb{A}_k^n, M)$ to be $\sigma(\mathbb{A}_k^n, M) \cap \mathbb{A}^n$ or $\beta^{-1}(\sigma(\mathbb{A}_k^n, M))$. Since $\sigma(\mathbb{A}_k^n, M)$ is closed, the set $\sigma_c(\mathbb{A}_k^n, M)$ turns out to be a closed subset of \mathbb{A}^n . If $\mathfrak{p} \in \sigma(\mathbb{A}_k^n, M)$ then $V(\mathfrak{p}) = \overline{\{\mathfrak{p}\}} \subseteq \sigma(\mathbb{A}_k^n, M)$ and $\mathfrak{m} \in \sigma_c(\mathbb{A}_k^n, M)$ whenever $\mathfrak{m} \in V(\mathfrak{p}) \cap \mathbb{A}^n$ is a maximal ideal. By Hilbert Nullstellensatz [1, 15.7], $\sigma(\mathbb{A}_k^n, M)$ is the closure of $\sigma_c(\mathbb{A}_k^n, M)$ in \mathbb{A}_k^n .

Notice that $\mathfrak{m} \in \sigma_p(\mathbb{A}_k^n, M) \cap \mathbb{A}^n$ iff $\mathfrak{m} = \langle X - a \rangle = \text{Ann}(u)$ for some point $a \in \mathbb{A}^n$ and $u \neq 0$ from M . The latter means that $x_i u = a_i u$ for all i , which means that $a \in \sigma_p(x, M)$ is a joint eigenvalue of x with the related eigenvector u .

Finally, assume that M is a Noetherian R -module (or P -module). Based on Theorem 3.2, we conclude that $\text{res}(\mathbb{A}_k^n, M) \cap \mathbb{A}^n = \{a \in \mathbb{A}^n : k(\langle X - a \rangle) \perp_P M\}$ or

$$\sigma_c(\mathbb{A}_k^n, M) = \{a \in \mathbb{A}^n : \text{Kos}(x - a, M) \text{ is not exact}\} = \sigma(x, M)$$

is the Taylor spectrum of the tuple x on M . Hence $\sigma(x, M)$ is a closed subset of \mathbb{A}^n and $\sigma(x, M)^- = \sigma(\mathbb{A}_k^n, M) = \sigma(\mathfrak{X}, M)$. By Corollary 2.7, $\sigma(x, M) \neq \emptyset$. \square

Corollary 3.4. *Let R/k be an algebra finite extension of the field k , M an R -module with $\ell_R(M) < \infty$, and let x be an n -tuple in R such that $k[x] \subseteq R$ is integral. Then $\sigma_p(x, M) = \sigma_p(\mathbb{A}_k^n, M) = \sigma(\mathbb{A}_k^n, M) = \sigma(x, M)$ and the $k[x]$ -module structure on M is triangularizable. In this case, $i(x - a) = 0$ for all $a \in \mathbb{A}^n$.*

Proof. Based on Corollary 3.3, we can assume that $R = k[x]$ (see also Remark 3.3). In this case, $\text{Ass}(M) = \text{Supp}(M) = \text{Max}(M) \subseteq \mathbb{A}^n$ [1, 19.4]. It remains to use Lemma 3.1. Further, all the gaps of a Jordan-Hölder chain is k by Zariski Nullstellensatz. There is a k -basis $\omega = (\omega_1, \dots, \omega_s)$ for M such that $\text{Ass}(M) = \{a^{(1)}, \dots, a^{(s)}\}$ and

$$x = \begin{bmatrix} a^{(1)} & & 0 \\ & \ddots & \\ * & & a^{(s)} \end{bmatrix},$$

which means that the $k[x]$ -module structure on M is triangularizable.

Finally, for every $a \in \mathbb{A}^n$ the complex $\text{Kos}(x - a, M)$ consists of finite dimensional k -vector space. It follows that its index coincides with its Euler characteristics. Hence

$$\begin{aligned} i(x - a) &= \sum_{j=0}^n (-1)^{j+1} \dim_k(H_j(x - a, M)) = \sum_{j=0}^n (-1)^{j+1} \dim_k(M \otimes_k \wedge^j k^n) \\ &= \dim_k(M) \sum_{j=0}^n (-1)^{j+1} \binom{n}{j} = 0, \end{aligned}$$

that is, $i(x - a) = 0$ for all $a \in \mathbb{A}^n$. \square

The following reformulation of Theorem 3.1 results in spectral mapping theorem for the joint point spectrum.

Theorem 3.3. *Let $R = k[x]$ be an algebra finite extension with an n -tuple x , and let M be an R -module. Then*

$$\sigma_p(p(x), M) = p^*(\sigma_p(\mathbb{A}_k^n, M)) \cap \mathbb{A}^m$$

for every m -tuple $p(x)$ from R . If $k[p(x)] \subseteq R$ is integral then $\sigma_p(p(x), M) = p(\sigma_p(x, M))$ and $\sigma(p(x), M)$ is a closed set.

Proof. Put $\mathfrak{X} = \text{Spec}(R)$, $R' = k[p(x)]$ to be a subalgebra of R generated by an m -tuple $p(x)$ from R with the inclusion map $\iota : R' \rightarrow R$, and $\iota^* : \mathfrak{X} \rightarrow \mathfrak{X}'$ is the natural map with $\mathfrak{X}' = \text{Spec}(R')$. There are canonical surjective maps $k[X] \rightarrow R$ and $k[Y] \rightarrow R'$, where Y is an m -tuple. In particular, $\mathfrak{X} \subseteq \mathbb{A}_k^n$ and $\mathfrak{X}' \subseteq \mathbb{A}_k^m$ up to a canonical homeomorphisms. We have the polynomial ring map $p : k[Y] \rightarrow k[X]$, $Y_j \mapsto p_j(X)$, $1 \leq j \leq m$, which in turn generates the map $p^* : \mathbb{A}_k^n \rightarrow \mathbb{A}_k^m$ of affine spaces. If $\mathfrak{m}_a = \langle X - a \rangle \in \mathbb{A}^n$ then $p^*(\mathfrak{m}_a) = p^{-1}(\mathfrak{m}_a) \supseteq \langle Y - p(a) \rangle = \mathfrak{m}_{p(a)}$, for $p(\langle Y - p(a) \rangle) \subseteq \langle p(X) - p(a) \rangle \subseteq \mathfrak{m}_a$. Since $\mathfrak{m}_{p(a)} \in \mathbb{A}^m$ is maximal, we conclude that $p^*(\mathfrak{m}_a) = \mathfrak{m}_{p(a)}$. Thus $p^*(\mathbb{A}^n) \subseteq \mathbb{A}^m$ and $p^* : \mathbb{A}^n \rightarrow \mathbb{A}^m$, $a \mapsto p(a)$ is a polynomial morphism. Moreover, $p^*(\mathbb{A}_k^n) \cap \mathbb{A}^m = p^*(\mathbb{A}^n)$. Indeed, if $p^*(\mathfrak{q}) = \mathfrak{m}_b$ for some $\mathfrak{q} \in \mathbb{A}_k^n$ and $b \in \mathbb{A}^m$, then $\mathfrak{q} \subseteq \mathfrak{m}_a$ for some $a \in \mathbb{A}^n$ and $\mathfrak{m}_b \subseteq p^*(\mathfrak{m}_a) = \mathfrak{m}_{p(a)}$, which implies that $\mathfrak{m}_b = \mathfrak{m}_{p(a)}$ or $b = p(a)$.

Using the duality correspondence between affine schemes and rings, we obtain the following commutative diagrams

$$\begin{array}{ccccc} k[Y] & \xrightarrow{p} & k[X] & & \mathbb{A}_k^m & \xleftarrow{p^*} & \mathbb{A}_k^n \\ \downarrow & & \downarrow & \Leftrightarrow & \uparrow & & \uparrow \\ R' & \xrightarrow{\iota} & R & & \mathfrak{X}' & \xleftarrow{\iota^*} & \mathfrak{X} \end{array}$$

supporting each other. Using Theorem 3.1 and Lemma 3.1, we deduce that

$$\begin{aligned} \sigma_p(p(x), M) &= \sigma_p(\mathbb{A}_k^m, M) \cap \mathbb{A}^m = \sigma_p(\mathfrak{X}', M) \cap \mathbb{A}^m = \iota^*(\sigma_p(\mathfrak{X}, M)) \cap \mathbb{A}^m \\ &= p^*(\sigma_p(\mathbb{A}_k^n, M)) \cap \mathbb{A}^m. \end{aligned}$$

If $R' \subseteq R$ is integral, then $\sigma_p(\mathfrak{X}', M) \cap \mathbb{A}^m = \iota^*(\sigma_p(\mathfrak{X}, M) \cap \mathbb{A}^n)$ by Maximality [1, 14.3] and Theorem 3.1. Using again Lemma 3.1, we deduce that $\sigma_p(p(x), M) = p^*(\sigma_p(\mathbb{A}_k^n, M) \cap \mathbb{A}^n) = p(\sigma_p(x, M))$.

Finally, prove that Taylor spectrum $\sigma(p(x), M)$ is a closed set. Since R/R' is integral, M is module finite over R' . By Lemma 3.1, $\sigma(p(x), M) = \sigma(\mathbb{A}_k^m, M) \cap \mathbb{A}^m$ is a closed set. \square

Remark 3.4. *The same argument from the proof of Theorem 3.3, and Proposition 3.1, result in the spectral mapping property $\sigma(p(x), M) = p(\sigma(x, M))$ for Taylor spectrum whenever $k[p(x)] \subseteq R$ is integral and M is Noetherian. But as we have seen above in Theorem 2.3, the formula holds in the general case of all tuples $p(x)$.*

If $\mathfrak{a} \subseteq P$ is a radical ideal with the related algebraic set $Y = Z(\mathfrak{a}) = V(\mathfrak{a}) \cap \mathbb{A}^n$ (that is $I(Y) = \mathfrak{a}$), then the coordinate ring R of Y is reduced to the Noetherian ring P/\mathfrak{a} . Put $\mathfrak{Y} = \text{Spec}(R)$. By (3.2), we obtain that $\sigma(\mathbb{A}_k^n, R) = \text{Supp}(R) = V(\mathfrak{a}) = \mathfrak{Y}$ is the set of all subvarieties of Y , and by Lemma 3.1, $Y = \mathfrak{Y} \cap \mathbb{A}^n = \sigma(x, R)$ is the Taylor spectrum of the operator tuple x in R , which consists of all coordinate functions. For a Noetherian $k[x]$ -module M , we obtain that the tuple x on the module M is triangularizable whose diagonal entries are varieties (see [11, 4.4]).

Let $p = (p_1, \dots, p_m) : Y \rightarrow \mathbb{A}^m$ be a morphism given by means of a ring extension $R' = k[y] \subseteq R$ with $y = p(x)$ to be an m -tuple. In particular, we have the spectrum $\sigma(\mathbb{A}_k^m, R)$ of the R' -module R . If R/R' is integral then p^* turns out to be a finite morphism (see [16, 2.3]). Using (1.2), we deduce that $\sigma(\mathbb{A}_k^m, R) \cap \mathbb{A}^m = p(Y)^-$ (see [11] for the details).

Corollary 3.5. *If $Y \subseteq \mathbb{A}^n$ is an algebraic set over k with its coordinate ring R , then $\sigma_p(x, R)$ is the set Y_{is} of all isolated points of Y . If $p : Y \rightarrow \mathbb{A}^m$ is a finite morphism given by a ring extension $k[p(x)] \subseteq R$, then $\sigma(p(x), R) = \sigma(\mathbb{A}_k^m, R) \cap \mathbb{A}^m = p(Y)$ and $\sigma_p(p(x), R) = p(Y_{is})$.*

Proof. Note that $a \in \sigma_p(x, R)$ iff $\langle X - a \rangle g(X) \subseteq \mathfrak{a}$ for some $g(X) \notin \mathfrak{a}$. It means that $Y = Z(\mathfrak{a}) \subseteq \{a\} \cup Z(g)$ and $Y \not\subseteq Z(g)$. Since $a \in Y$, we conclude that $a \notin Z(g)$ and $\{a\} = Y \cap (\mathbb{A}^n - Z(g))$ is open in Y , that is, $\{a\}$ is an isolated point. Conversely, take $a \in Y_{is}$.

Since the complements to hypersurfaces from \mathbb{A}^n is a topology base in \mathbb{A}^n , it follows that $\{a\} = Y \cap (\mathbb{A}^n - Z(g))$ for a certain hypersurface $Z(g)$. Then $Y = Z(\mathfrak{a}) \subseteq \{a\} \cup Z(g)$, $Y \not\subseteq Z(g)$ and

$$\begin{aligned} \langle X - a \rangle g &\subseteq \langle X - a \rangle \cap \sqrt{g} = I(\{a\}) \cap I(Z(g)) = I(\{a\} \cup Z(g)) \\ &\subseteq I(Y) = \sqrt{\mathfrak{a}} = \mathfrak{a}, \end{aligned}$$

which means that $a \in \sigma_p(x, R)$. Further, if p is a finite morphism then R is a Noetherian $k[p(x)]$ -module and $\sigma(p(x), R) = \sigma(\mathbb{A}_k^m, R) \cap \mathbb{A}^m$ is closed (see Lemma 3.1). Using Lemma 3.1 and Theorem 2.3, we derive that $\sigma(\mathbb{A}_k^m, R) \cap \mathbb{A}^m = \sigma(p(x), R) = p(\sigma(x, R)) = p(Y)$, and $\sigma_p(p(x), R) = p(\sigma_p(x, R)) = p(Y_{is})$ thanks to Theorem 3.3. \square

4. KOSZUL HOMOLOGY GROUPS OF A VARIETY

In the present section we focus on the case of a module M which is the coordinate ring R of a variety Y , and investigate its Koszul homology groups. We are targeting to find out a link between the dimensions of Koszul homology groups of the standard tuple in R and the dimension $d = \dim(Y)$.

4.1. Multiplicity formula of Serre. Let $P = k[X]$ be the polynomial algebra over the field k with n -tuple $X = (X_1, \dots, X_n)$, and put $h(r) = \dim_k(\mathfrak{t}^r/\mathfrak{t}^{r+1}) = \binom{r+n-1}{n-1}$ to be the Hilbert polynomial ($\deg(h) = n-1$) of the graded P -algebra $G^\circ P$ associated with the filtration $\{\mathfrak{t}^r\}$, where $\mathfrak{t} = \langle X \rangle \subseteq P$ is the maximal ideal generated by X , and $\binom{z}{s} = \frac{1}{s!} z(z-1)\cdots(z-s+1)$ is the binomial coefficient. If $\mathfrak{s}(P, r) = \dim_k(P/\mathfrak{t}^r)$ is the Samuel polynomial of the filtration then $\mathfrak{s}(P, r) = \sum_{i=0}^{r-1} h(i) = \sum_{i=0}^{r-1} \binom{i+n-1}{n-1} = \binom{r+n-1}{n}$. If M is a P -module given by an n -tuple x of k -linear maps on M , then $P/\mathfrak{t}^r \otimes_k M = M^{\binom{r+n-1}{n}}$ and there are nilpotent (shift) operators $X_i \otimes 1$ acting on $M^{\binom{r+n-1}{n}}$. Put $x_i^{(r)} = (1 \otimes x_i) - X_i \otimes 1$ and $x^{(r)} = (x_1^{(r)}, \dots, x_n^{(r)})$ to be an n -tuple of mutually commuting operators on the inflation $M^{\binom{r+n-1}{n}}$, which defines its P -module structure. For every $a \in \mathbb{A}^n$ the P -module structure on $M^{\binom{r+n-1}{n}}$ given by $x^{(r)} - a$ is the same inflation of the P -module M given by the tuple $x - a$. Just notice that $x_i^{(r)} - a_i = 1 \otimes (x_i - a_i) - X_i \otimes 1 = (x_i - a_i)^{(r)}$ for all i , or $x^{(r)} - a = (x - a)^{(r)} = 1 \otimes (x - a) - X \otimes 1$. In this case one can also replace \mathfrak{t} by $\langle X - a \rangle$.

Lemma 4.1. *If M is a Noetherian P -module then so is $M^{\binom{r+n-1}{n}}$ and $\sigma(x, M) = \sigma(x^{(r)}, M^{\binom{r+n-1}{n}})$ for all r . Thus $i((x - a)^{(r)}) < \infty$ for all $a \in \mathbb{A}^n$ and $r \geq 1$.*

Proof. For every $r \geq 1$ consider the following canonical exact sequence

$$(4.1) \quad 0 \leftarrow P/\mathfrak{t}^r \leftarrow P/\mathfrak{t}^{r+1} \leftarrow \mathfrak{t}^r/\mathfrak{t}^{r+1} \leftarrow 0,$$

in P -mod. Notice that the P -module structure of $\mathfrak{t}^r/\mathfrak{t}^{r+1}$ is reduced to its k -vector space one with $\dim_k(\mathfrak{t}^r/\mathfrak{t}^{r+1}) = h(r)$. Using (4.1), we generate the following exact sequences

$$0 \leftarrow P/\mathfrak{t}^r \otimes_k M \leftarrow P/\mathfrak{t}^{r+1} \otimes_k M \leftarrow \mathfrak{t}^r/\mathfrak{t}^{r+1} \otimes_k M \leftarrow 0$$

such that the P -module structure of $\mathfrak{t}^r/\mathfrak{t}^{r+1} \otimes_k M$ is diagonal and therefore it is Noetherian. By induction on r we deduce that $P/\mathfrak{t}^r \otimes_k M$ is Noetherian. By Spectral Mapping Theorem 2.3, we

have

$$\begin{aligned}\sigma\left(x^{(r)}, M^{\binom{r+n-1}{n}}\right) &= \sigma\left(1 \otimes x - X \otimes 1, M^{\binom{r+n-1}{n}}\right) \\ &= \left\{a - b : (a, b) \in \sigma\left((1 \otimes x, X \otimes 1), M^{\binom{r+n-1}{n}}\right)\right\}\end{aligned}$$

and $b \in \sigma\left(X \otimes 1, M^{\binom{r+n-1}{n}}\right)$. For every $a \in \sigma\left(1 \otimes x, M^{\binom{r+n-1}{n}}\right)$ there is b such that $(a, b) \in \sigma\left((1 \otimes x, X \otimes 1), M^{\binom{r+n-1}{n}}\right)$ (see Theorem 2.2). Moreover, $b_i \in \sigma\left((X_i \otimes 1) | M^{\binom{r+n-1}{n}}\right) = \{0\}$, for $X_i \otimes 1$ is nilpotent. Hence $\sigma\left(x^{(r)}, M^{\binom{r+n-1}{n}}\right) = \sigma\left(1 \otimes x, M^{\binom{r+n-1}{n}}\right)$. Finally,

$$\text{Kos}\left(1 \otimes (x - a), M^{\binom{r+n-1}{n}}\right) = P/\mathfrak{t}^r \otimes_k \text{Kos}(x - a, M),$$

which is exact iff so is $\text{Kos}(x - a, M)$, that is, $\sigma\left(1 \otimes x, M^{\binom{r+n-1}{n}}\right) = \sigma(x, M)$. It remains to use Theorem 2.1. \square

Let A/k be a local Noetherian algebra with an n -tuple x contained in the maximal ideal of A , and let M be a Noetherian A -module such that $\ell(M/xM) < \infty$, that is, x is a system of parameters for M . The filtration $\{\langle x \rangle^r M\}$ in M defines the Samuel polynomial $\mathfrak{s}(M, r) = \ell(M/\langle x \rangle^r M)$ whose degree is at most n , and $\mathfrak{s}(M, r) = e_{x,M}(n) \frac{r^n}{n!} + q(r)$ with $e_{x,M}(n) = \Delta^n \mathfrak{s}(M, \circ)$ (n th difference operator) and $\deg(q) < n$. By Theorem 2.1, $\ell(H_p(x, M)) = \dim_k(H_p(x, M))$ for all p , and the Euler characteristic $\chi(x, M) = \sum_{p=0}^n (-1)^p \ell(H_p(x, M))$ from [24, 4.A.3] is reduced to the opposite index $-i(x)$ of the tuple x . The multiplicity formula of Serre [24, 4.A.3. Theorem 1] is expressed in the following way

$$(4.2) \quad i(x) = -e_{x,M}(n).$$

If $\dim(M) = n$ then $i(x) < 0$, whereas $i(x) = 0$ in the case of $\dim(M) < n$.

4.2. The numerical Tor-polynomial. Now let $\mathfrak{p} \subseteq P$ be a nonzero prime ideal of a variety $Y \subseteq \mathbb{A}^n$ with its coordinate ring $R = P/\mathfrak{p}$. The actions of X_i on R are denoted by x_i , and $x = (x_1, \dots, x_n)$ is an operator tuple on R . Thus $R = k[x]$ is an algebra finite extension of k which is a domain. In particular, $\sigma_{\mathfrak{p}}(x, R) = \emptyset$ or $H_n(x) = 0$. For every r we have the inflated P -module $R^{\binom{r+n-1}{n}}$ given by means of the n -tuple $x^{(r)}$. Using Lemma 3.1 and Lemma 4.1, we deduce that

$$Y = \text{Spec}(R) \cap \mathbb{A}^n = \sigma(\mathbb{A}_k^n, R) \cap \mathbb{A}^n = \sigma(x, R) = \sigma\left(x^{(r)}, R^{\binom{r+n-1}{n}}\right)$$

for all r . For every $a \in Y$ the homology groups $H_i\left(x^{(r)} - a, R^{\binom{r+n-1}{n}}\right)$ denoted by $H_i(x^{(r)} - a)$ are finite dimensional k -vector spaces (Lemma 4.1). Put $d_p(a^{(r)}) = \dim_k(H_p(x^{(r)} - a))$ and define the tuple $d(a^{(r)}) = (d_p(a^{(r)}))_p \in \mathbb{Z}_+^{n+1}$. In particular, $i(x^{(r)} - a) = \sum_{p=0}^n (-1)^{p+1} d_p(a^{(r)})$ and it defines the function $i_Y^{(r)} : Y \rightarrow \mathbb{Z}$, $a \mapsto i(x^{(r)} - a)$. We also put i_Y instead of $i_Y^{(1)}$.

Lemma 4.2. *If $Y \subseteq \mathbb{A}^n$ is a variety then $i_Y = -\delta_{nd}$, where $d = \dim(Y)$.*

Proof. If $Y = \mathbb{A}^n$ then $\text{Kos}(X - a, P)$ is a free P -module resolution for the P -module k (see Corollary 2.3), that is, $H_0(X - a, P) = k$ and $H_i(X - a, P) = 0$, $i \geq 1$. In particular, $d_0(a) = 1$, $d_i(a) = 0$, $i \geq 1$, and $i_Y(a) = -1$ for all $a \in Y$.

Now assume that $Y \subseteq \mathbb{A}^n$ is a variety of dimension $d < n$, and $a = 0 \in Y$, which responds to the maximal ideal $\langle x \rangle \subseteq R$. Then $R_{\langle x \rangle}$ is a local Noetherian algebra with its maximal ideal

$\mathfrak{m} = \langle x \rangle R_{\langle x \rangle}$. By Lemma 2.5, we have $H_p(x, R) = H_p(x, R)_{\langle x \rangle} = H_p(x/1, R_{\langle x \rangle})$ for all $p \geq 0$, and $\mathfrak{m} = \langle x/1 \rangle$ for the n -tuple $x/1$. Thus $i(x) = i(x/1)$, $x/1$ is a system of parameters for $R_{\langle x \rangle}$ and the Samuel polynomial $\mathfrak{s}(R_{\langle x \rangle}, r)$ has the degree $\dim(R_{\langle x \rangle})$ [1, Theorem 21.4]. But $\dim(R_{\langle x \rangle}) = \dim(R) = d < n$. Using Serre's formula (4.2), we conclude that $i(x) = 0$. \square

Similar result for the inflated P -module $R^{(r+n-1)}_n$ is not trivial (at least is not straightforward), for its P -module structure is not diagonal. The related result is proved below in Proposition 4.1.

Remark 4.1. *In the case of a nonsingular point $a = 0 \in Y$ of a variety $Y \subseteq \mathbb{A}^n$ with $d < n$, one can skip Serre's formula. Namely, $d = \dim_k(\mathfrak{m}/\mathfrak{m}^2)$ is the minimal number of generators of the $R_{\langle x \rangle}$ -module \mathfrak{m} [1, 10.9]. As above $\mathfrak{m} = \langle x/1 \rangle$ for the n -tuple $x/1$, and there is a new d -tuple $y \subseteq \mathfrak{m}$ generating \mathfrak{m} due to the regularity of $R_{\langle x \rangle}$, that is, y is a tuple related to $x/1$. Note that $0 \in Y = \sigma(x, R)$ or $H_p(x, R) \neq 0$ for some p , which in turn implies that $0 \in \sigma(x/1, R_{\langle x \rangle})$. Using Corollary 2.9, we deduce that $i(x) = i(x/1) = \delta_{nd}i(y) = 0$.*

Remark 4.2. *A singular point a of a variety Y is said to be integral if $x_i - a_i$ is integral over the subalgebra $k[x' - a'] \subseteq R$ with i -th term skipped tuple x' . Suppose that $a = 0$, and say x_n is integral over $R' = k[x']$ with $x' = (x_1, \dots, x_{n-1})$. Then $R' \subseteq R$ is integral and $i(x') < \infty$ thanks to Theorem 2.1. Moreover, $0 \in \sigma(x', R)$ thanks to Corollary 2.2. By Lemma 2.3, $i(x) = 0$. Again we can avoid (4.2).*

Example 4.1. *In the case of a point $Y = \{a\}$, we have $R = k$ and $i = 0$ thanks to Corollary 3.4. Actually, $\text{Kos}(x, k)$ is the following complex*

$$0 \leftarrow k \leftarrow k^n \leftarrow \dots \leftarrow \wedge^p k^n \leftarrow \dots \leftarrow k \leftarrow 0$$

with trivial morphisms, $H_j(x, k) = \wedge^j k^n$, $d_j(a) = \binom{n}{j}$, $j \geq 0$, and $i_Y = \sum_{j=0}^n (-1)^{j+1} \binom{n}{j} = 0$.

Now assume that Y is a variety and $a = 0 \in Y$, that is, $\mathfrak{p} \subseteq \langle X \rangle$. For every $r \geq 1$ consider again the sequence (4.1). Notice that $\text{Tor}_i^P(\mathfrak{t}^r/\mathfrak{t}^{r+1}, R) = \text{Tor}_i^P(k, R)^{h(r)} = H_i(x)^{h(r)}$ for all $i \geq 0$. The functor $\circ \otimes_P R$ generates the following long homology exact sequence

$$\begin{aligned} 0 \leftarrow P/\mathfrak{t}^r \otimes_P R \leftarrow P/\mathfrak{t}^{r+1} \otimes_P R \leftarrow k^{h(r)} \leftarrow \text{Tor}_1^P(P/\mathfrak{t}^r, R) \leftarrow \text{Tor}_1^P(P/\mathfrak{t}^{r+1}, R) \leftarrow H_1(x)^{h(r)} \\ \dots \leftarrow H_{i-1}(x)^{h(r)} \leftarrow \text{Tor}_i^P(P/\mathfrak{t}^r, R) \leftarrow \text{Tor}_i^P(P/\mathfrak{t}^{r+1}, R) \leftarrow H_i(x)^{h(r)} \leftarrow \dots \end{aligned}$$

But $P/\mathfrak{t}^r \otimes_P R = R/\mathfrak{t}^r R = R/\langle x \rangle^r$ and $0 \leftarrow R/\langle x \rangle^r \leftarrow R/\langle x \rangle^{r+1} \leftarrow \langle x \rangle^r / \langle x \rangle^{r+1} \leftarrow 0$ is exact. Thus we come with the exact sequence

$$0 \leftarrow \langle x \rangle^r / \langle x \rangle^{r+1} \leftarrow k^{h(r)} \leftarrow \text{Tor}_1^P(P/\mathfrak{t}^r, R) \leftarrow \text{Tor}_1^P(P/\mathfrak{t}^{r+1}, R) \leftarrow H_1(x)^{h(r)} \leftarrow \dots$$

Note that $\mathcal{P} = \text{Kos}(t, P \otimes_k R)$ is a free resolution of R , and $P/\mathfrak{t}^r \otimes_P \mathcal{P} = \text{Kos}(x^{(r)}, P/\mathfrak{t}^r \otimes_k R) = \text{Kos}(x^{(r)}, R^{(r+n-1)}_n)$ with $x^{(r)} = 1 \otimes x - X \otimes 1$ (see Lemma 4.1). In particular, $\text{Tor}_i^P(P/\mathfrak{t}^r, R) = H_i(P/\mathfrak{t}^r \otimes_P \mathcal{P}) = H_i(x^{(r)})$ for all i and r . Now we apply the localization functor $\circ \otimes_R R_{\langle x \rangle}$ (or $\circ \otimes_P P_{\langle X \rangle}$) at $\langle x \rangle$. By Lemma 2.5, we have

$$H_i\left(x^{(r)}/1, R_{\langle x \rangle}^{(r+n-1)}_n\right) = H_i\left(x^{(r)}, R^{(r+n-1)}_n\right)_{\langle x \rangle} = H_i\left(x^{(r)}, R^{(r+n-1)}_n\right) = H_i(x^{(r)})$$

for all i and r . We obtain the following exact sequence

$$\begin{aligned} (4.3) \quad 0 \leftarrow \mathfrak{m}^r / \mathfrak{m}^{r+1} \leftarrow k^{h(r)} \leftarrow H_1(x^{(r)}) \leftarrow H_1(x^{(r+1)}) \leftarrow H_1(x)^{h(r)} \leftarrow \dots \\ \leftarrow H_{n-1}(x)^{h(r)} \leftarrow H_n(x^{(r)}) \leftarrow H_n(x^{(r+1)}) \leftarrow H_n(x)^{h(r)} = 0, \end{aligned}$$

where $\mathfrak{m} = \langle x \rangle R_{\langle x \rangle}$ is the maximal ideal of the local ring $R_{\langle x \rangle}$.

Lemma 4.3. *If $H_{j+1}(x) = 0$ for some j , then $H_i(x^{(r)}) = 0$ for all $i > j$ and $r \geq 1$, and the sequence*

$$\begin{aligned} 0 \leftarrow \mathfrak{m}^r / \mathfrak{m}^{r+1} \leftarrow k^{h(r)} \leftarrow \cdots \leftarrow H_{i-1}(x)^{h(r)} \leftarrow H_i(x^{(r)}) \leftarrow H_i(x^{(r+1)}) \leftarrow H_i(x)^{h(r)} \leftarrow \cdots \\ \leftarrow H_{j-1}(x^{(r)}) \leftarrow H_j(x^{(r)}) \leftarrow H_j(x)^{h(r)} \leftarrow 0 \end{aligned}$$

remains exact. In particular, $H_n(x^{(r)}) = 0$ for all r , and the sequence

$$0 \leftarrow \mathfrak{m}^r / \mathfrak{m}^{r+1} \leftarrow k^{h(r)} \leftarrow \cdots \leftarrow H_{n-1}(x^{(r)}) \leftarrow H_{n-1}(x^{(r+1)}) \leftarrow H_{n-1}(x)^{h(r)} \leftarrow 0$$

is exact.

Proof. Suppose that $H_{j+1}(x) = 0$ for some j . Prove that $H_i(x^{(r)}) = 0$ for all $i > j$, $r \geq 1$ by induction on r . By Lemma 2.5, $H_i(x) = 0$ for all $i > j$. If $r = 1$, then we have $\text{Tor}_i^P(P/\mathfrak{t}, R) = \text{Tor}_i^P(k, R) = H_i(x)$, $i \geq 0$ and we come with the exact pieces

$$\cdots \leftarrow H_i(x) \leftarrow H_i(x^{(2)}) \leftarrow H_i(x)^{h(r)} \leftarrow \cdots$$

of the sequence (4.3) for $r = 1$. If $i > j$ then $H_i(x^{(2)}) = 0$. By induction hypothesis, $H_i(x^{(r)}) = 0$ for all $i > j$. The exact part

$$\cdots \leftarrow H_{i-1}(x)^{h(r)} \leftarrow H_i(x^{(r)}) \leftarrow H_i(x^{(r+1)}) \leftarrow H_i(x)^{h(r)} \leftarrow \cdots$$

of the sequence (4.3) for $i > j$, implies that $H_i(x^{(j+1)}) = 0$, and the long homology sequence terminates at $H_j(x)^{h(r)}$. Finally, since $H_n(x) = 0$ we conclude that $H_n(x^{(r)}) = 0$ for all $r \geq 1$. \square

Based on Lemma 4.1, we have $i(x^{(r)}) < \infty$ for all r , and we define the following function

$$p : \mathbb{Z} \longrightarrow \mathbb{Z}, \quad p(r) = \sum_{i=1}^n (-1)^{i+1} \dim_k (\text{Tor}_i^P(P/\mathfrak{t}^r, R))$$

called as of Tor-polynomial. Note that $-\dim_k(H_0(x^{(r)})) + p(r) = i(x^{(r)})$ for all r . Recall that $R_{\langle x \rangle}$ is a local Noetherian algebra, and $d = \dim(Y) = \dim(R_{\langle x \rangle}) = \deg(h_R) + 1 = \deg \mathfrak{s}(R_{\langle x \rangle}, \circ) \leq n$, where $h_R(r) = \dim_k(\mathfrak{m}^r / \mathfrak{m}^{r+1})$ is the Hilbert polynomial of $R_{\langle x \rangle}$ associated with $\{\mathfrak{m}^r\}$.

Proposition 4.1. *The equality $p(r) = (1 - \delta_{nd}) \mathfrak{s}(R_{\langle x \rangle}, r)$ holds for all r , that is, the function p is a numerical polynomial of degree at most d . In particular, $i_Y^{(r)} = \binom{r+n-1}{n} i_Y$ for all r .*

Proof. Using the exact sequence from Lemma 4.3 and its Euler characteristics, we deduce that

$$\begin{aligned} 0 = -h_R(r) + h(r) + \sum_{i=1}^{n-1} (-1)^i \dim_k(H_i(x^{(r)})) + \sum_{i=1}^{n-1} (-1)^{i+1} \dim_k(H_i(x^{(r+1)})) \\ + h(r) \sum_{i=1}^{n-1} (-1)^i \dim_k(H_i(x)). \end{aligned}$$

Note that $H_0(x) = R/\langle x \rangle = k$, $H_n(x^{(r)}) = 0$, and $\sum_{i=0}^n (-1)^i \dim_k(H_i(x)) = 1 - p(1) = -i(x)$. If $d < n$ then $i(x) = 0$ thanks to Lemma 4.2. Thus $p : \mathbb{Z} \rightarrow \mathbb{Z}$ is a function with the property

$$p(z+1) - p(z) = h_R(z),$$

that is, $p(z+1) - p(z)$ is a numerical polynomial for large z . That means $p(z) \in \mathbb{Q}[z]$ is a numerical polynomial (see [16, 1.7.3]) and $p(r) = \sum_{j=0}^{r-1} h_R(j) = \mathfrak{s}(R_{\langle x \rangle}, r)$ for all $r \geq 1$. If

$Y = \mathbb{A}^n$ then $h_R(r) = h(r)$, $H_i(x) = 0$, $i \geq 1$, and $i(x) = -1$ (see Lemma 4.2). It follows that $p(r+1) = p(r)$ for all $r \geq 1$. But $p(1) = i(x) + 1 = 0$, therefore $p = 0$.

Finally, taking into account that

$$H_0(x^{(r)}) = H_0(x^{(r)})_{\langle x \rangle} = \text{Tor}_0^P(P/\mathfrak{t}^r, R)_{\langle x \rangle} = (P/\mathfrak{t}^r \otimes_P R)_{\langle x \rangle} = (R/\langle x \rangle^r)_{\langle x \rangle} = R_{\langle x \rangle}/\mathfrak{m}^r,$$

we deduce that $\dim_k(H_0(x^{(r)})) = \dim_k(R_{\langle x \rangle}/\mathfrak{m}^r) = \mathfrak{s}(R_{\langle x \rangle}, r)$. It follows that

$$i(x^{(r)}) = -\dim_k(H_0(x^{(r)})) + p(r) = -\mathfrak{s}(R_{\langle x \rangle}, r) + (1 - \delta_{nd})\mathfrak{s}(R_{\langle x \rangle}, r) = -\delta_{nd}\mathfrak{s}(R_{\langle x \rangle}, r).$$

Thus $i_Y^{(r)} = 0$ whenever $d < n$, and $i_Y^{(r)} = -\mathfrak{s}(P_{\langle X \rangle}, r) = -\mathfrak{s}(P, r) = -\binom{r+n-1}{n}$ for $Y = \mathbb{A}^n$. \square

Example 4.2. *If $d < n$ and $H_2(x) = 0$, then $H_i(x^{(r)}) = 0$ for all $i \geq 2$, $r \geq 1$ thanks to Lemma 4.3. Then $p(r) = \dim_k(H_1(x^{(r)}))$, $i(x) = -1 + \dim_k(H_1(x)) = 0$ and $\dim_k(H_1(x^{(r)})) = \mathfrak{s}(R_{\langle x \rangle}, r)$, that is, $r \mapsto \dim_k(H_1(x^{(r)}))$ is a polynomial of degree d . If $Y \subseteq \mathbb{A}^2$ is an irreducible curve and $a \in Y$, then $d_0(a) = 1$ and $d_2(a) = 0$. By Lemma 4.2, $i_Y(a) = 0$, therefore $d_1(a) = 1$, and $r \mapsto \dim_k(H_1((x-a)^{(r)}))$ is a linear polynomial.*

Remark 4.3. *The presence of nonzero $H_1(x)$ is well known. If $H_1(x) = 0$ then $R \xleftarrow{\partial_0} R \otimes k^n \xleftarrow{\partial_1} R \otimes \wedge^2 k^n$ is exact and so is its localization $R_{\langle x \rangle} \xleftarrow{\partial_0} R_{\langle x \rangle} \otimes k^n \xleftarrow{\partial_1} R_{\langle x \rangle} \otimes \wedge^2 k^n$, that is, $x/1 \subseteq \text{rad}(R_{\langle x \rangle})$ with $H_1(x/1) = 0$. Using [3, 9.7, Theorem 1], we conclude that $x/1$ is a regular sequence in $R_{\langle x \rangle}$. In particular, $\text{depth}(R_{\langle x \rangle}) \geq n$. But $\text{depth}(R_{\langle x \rangle}) \leq \dim(R_{\langle x \rangle}) = d < n$, a contradiction.*

The evaluation map $a : P \rightarrow k$, $f \mapsto f(a)$ can be lifted to a ring homomorphism $a : R \rightarrow k$, which is the quotient mapping $R \rightarrow R/\langle x \rangle$. In particular, there are k -linear maps $\wedge^r a : R \otimes_k \wedge^r k^n \rightarrow \wedge^r k^n$, $r \geq 1$, and $(\wedge^r a)\partial_{r-1} = 0$ for all r , which in turn define k -linear maps $a^{(r)} : H_r(x) \rightarrow \wedge^r k^n$ on the homology groups. Note that $H_0(x) = R/\text{im}(\partial_0) = k$, $a^{(0)} = 1$, and $a^{(1)} : H_1(x) \rightarrow k^n$ is a k -linear map.

4.3. Upper-triangular matrices. Now let us analyze the first homology group $H_1(x)$. We assume that $\text{char}(k) \neq 2$. Every $g = \sum_{i < j} g_{ij}e_i \wedge e_j \in R \otimes_k \wedge^2 k^n$ can be represented by means of an upper triangular matrix $g = [g_{ij}]_{i,j} \in M_n(R)$ whose i -th column C_i and i -th row R_i (they are n -tuples in R) have the shapes

$$\begin{array}{ccccccc} & & & & g_{1i} & & \\ & & & & g_{2i} & & \\ & & & & \vdots & & \\ & & & & g_{i-1,i} & & \\ \cdots & 0 & g_{ii+1} & \cdots & g_{in-1} & g_{in} & \\ & & & & \vdots & & \end{array}$$

for all $i > 1$, and C_1, R_n consist of zeros. For n -tuples α and β from R^n we write $\langle \alpha, \beta \rangle$ instead of the sum $\sum_{i=1}^n \alpha_i \beta_i$ in R . Note that $\partial_1 : R \otimes_k \wedge^2 k^n \rightarrow R \otimes_k k^n$ is an R -linear map acting by the rule

$$(4.4) \quad \partial_1(g) = \sum_{i < j} x_i g_{ij} e_j - x_j g_{ij} e_i = \sum_{i=1}^n \left(\sum_{s < i} x_s g_{si} - \sum_{i < s} x_s g_{is} \right) e_i = \sum_{i=1}^n (\langle C_i, x \rangle - \langle R_i, x \rangle) e_i.$$

For example, if

$$g = \begin{bmatrix} 0 & \cdots & g_j & \cdots & g_n \\ 0 & \cdots & 0 & \cdots & 0 \\ & & \vdots & & \\ 0 & & 0 & & 0 \end{bmatrix} \quad \text{with} \quad \sum_{k=j}^n x_k g_k = 0$$

for some g_j, \dots, g_n then $C_i = 0$, $1 \leq i \leq j-1$, $R_i = 0$, $i > 1$ and

$$\begin{aligned} \partial_1(g) &= -\langle R_1, x \rangle e_1 + \langle C_j, x \rangle e_j + \cdots + \langle C_n, x \rangle e_n = -\left(\sum_{k=j}^n x_k g_k\right) e_1 + x_1 g_j e_j + \cdots + x_1 g_n e_n \\ &= x_1 \sum_{k=j}^n g_k e_k. \end{aligned}$$

The following assertion is a special case of Lemma 2.4. But for our purposes we provide an independent proof in the case of $H_1(x)$ exploiting upper triangular matrices.

Lemma 4.4. *The equality $\langle x \rangle H_1(x) = \{0\}$ holds.*

Proof. Take $\omega = \sum_{j=1}^n g_j e_j \in \ker(\partial_0)$ or $\sum_{j=1}^n x_j g_j = 0$ in R . Fix i and prove that $x_i \omega \in \text{im}(\partial_1)$. First note that $x_i g_i = -\sum_{j \neq i} x_j g_j$ and consider the matrix

$$g = \begin{bmatrix} 0 & \cdots & -g_1 & 0 & \cdots & 0 \\ & & \vdots & \vdots & & \vdots \\ & & -g_{i-1} & 0 & & 0 \\ 0 & \cdots & 0 & g_{i+1} & \cdots & g_n \\ & & \vdots & \vdots & & \vdots \\ & & 0 & 0 & & 0 \end{bmatrix}$$

with nontrivial i -th column and i -th row. Since

$$\langle C_i, x \rangle - \langle R_i, x \rangle = -\sum_{j=1}^{i-1} x_j g_j - \sum_{j=i+1}^n x_j g_j = -\sum_{j \neq i} x_j g_j = x_i g_i,$$

we deduce using (4.4) that

$$\begin{aligned} \partial_1(g) &= -x_i g_1 e_1 - \cdots - x_i g_{i-1} e_{i-1} + (\langle C_i, x \rangle - \langle R_i, x \rangle) e_i + x_i g_{i+1} e_{i+1} + \cdots + x_i g_n e_n \\ &= x_i \left(\sum_{j < i} g_j e_j + \sum_{j \geq i} g_j e_j \right) = x_i \omega, \end{aligned}$$

that is, $x_i H_1(x) = \{0\}$. □

4.4. Minimal generators. Now let $F = \{f_1, \dots, f_p\}$ be a set of generators of the ideal \mathfrak{p} . Since $\mathfrak{p} \subseteq \langle X \rangle$ in P , it follows that every $f \in \mathfrak{p}$ has a standard representation $f = X_1^{m_1} h_1 + \cdots + X_n^{m_n} h_n$ with $h_j(a) \neq 0$. In particular, $f_j = \sum_i X_i^{m_{ij}} h_{ij}$ with $h_{ij}(a) \neq 0$ for all i, j . For every i , $1 \leq i \leq n$, we put $m_i = \min \{m_{ij} : 1 \leq j \leq p\}$. A generator $f \in F$ is said to be *minimal* if $f = f_j$ and $m_i = m_{ij}$ for some i . Every $f = \sum_i X_i^{m_i} h_i \in \mathfrak{p}$ associates in turn $\omega = \sum_{i=1}^n x_i^{m_i-1} h_i e_i \in R \otimes_k k^n$ such that $\partial_0(\omega) = \sum_{i=1}^n x_i^{m_i} h_i = 0$ in $R = P/\mathfrak{p}$. In particular, $\omega^\sim \in H_1(x)$.

Lemma 4.5. *If $f \in F$ is a minimal generator then $\omega^\sim \neq 0$ in $H_1(x)$, and $a^{(1)}(\omega^\sim) \neq 0$ whenever $m_i = 1$.*

Proof. Suppose $\omega^\sim = 0$, which means that $\omega = \partial_1(g)$ for some upper triangular matrix $g \in M_n(R)$. It follows that $\langle C_i, x \rangle - \langle R_i, x \rangle = x_i^{m_i-1} h_i$, $1 \leq i \leq n$ in R or

$$\sum_{s < i} X_s g_{si} - X_i^{m_i-1} h_i - \sum_{i < s} X_s g_{is} = \sum_{q=1}^p f_q l_q = \sum_{q=1}^p \sum_{k=1}^n X_k^{m_{kq}} h_{kq} l_q$$

in P for some $l_q \in P$. But $f = f_j$ for some j , therefore

$$\sum_{s < i} X_s g_{si} - X_i^{m_i-1} h_i - \sum_{i < s} X_s g_{is} = \sum_{k=1}^n X_k^{m_k} h_k l_j + \sum_{q \neq j} \sum_{k=1}^n X_k^{m_{kq}} h_{kq} l_q.$$

By passing to the quotient algebra $P / \langle X_1, \dots, \widehat{X_i}, \dots, X_n \rangle = k[X_i]$, we obtain that $-X_i^{m_i-1} h_i^\sim = X_i^{m_i} h_i^\sim l_j^\sim + \sum_{q \neq j} X_i^{m_{iq}} h_{iq}^\sim l_q^\sim$. But $m_i = \min \{m_{iq} : 1 \leq q \leq p\}$, therefore $m_{iq} - m_i \geq 0$ and

$$h_i^\sim = -X_i h_i^\sim l_j^\sim - \sum_{q \neq j} X_i^{m_{iq}-m_i+1} h_{iq}^\sim l_q^\sim \in \langle X_i \rangle.$$

It follows that $h_i(a) = 0$, a contradiction. Finally, $a(x_i^{m_i-1} h_i) = h_i(a) \neq 0$ whenever $m_i = 1$. Therefore $a^{(1)}(\omega^\sim) = (a(x_k^{m_k-1} h_k))_k \neq 0$ in k^n . \square

Remark 4.4. In the case of an irreducible curve $Y \subseteq \mathbb{A}^2$ given by $f = X_1^n h_1 + X_2^m h_2 \in k[X_1, X_2]$, $h_j \notin \langle X_1, X_2 \rangle$, $n, m \geq 1$, we have $H_1(x) = k\omega^\sim$ with $\omega = (x_2^{m-1} h_2, x_1^{n-1} h_1) \in \ker(\partial_0)$ and $a^{(1)}(\omega^\sim) = (a(x_2^{m-1} h_2), a(x_1^{n-1} h_1))$, which is not trivial whenever $m = 1$ or $n = 1$. The latter is equivalent to the presence of a linear part of f , which means that f is nonsingular at a . Thus $a^{(1)} : H_1(x - a) \rightarrow k^2$ is a nonzero linear map iff f is nonsingular at a .

For every i we put $S_i = \{j : m_{ij} = m_i\}$ and define the vector $v_j = (v_{1j}, v_{2j}, \dots, v_{nj}) \in k^n$ with

$$v_{ij} = \begin{cases} h_{ij}(a) & \text{if } j \in S_i, \\ 0 & \text{if } j \notin S_i. \end{cases}$$

Thus $v_j = 0$ whenever $j \notin \cup_i S_i$. If $j \in S_i$ for some i , then $m_{ij} = m_i$, f_j turns out to be a minimal generator and $v_{ij} = h_{ij}(a) \neq 0$. Hence $v_j \neq 0$ in k^n iff f_j is a minimal generator. Suppose f_1, \dots, f_t are minimal generators in F . Then we have the related nonzero vectors $\{v_1, \dots, v_t\}$ from k^n , and related nonzero vectors $\{\omega_1^\sim, \dots, \omega_t^\sim\}$ (see Lemma 4.5) from $H_1(x)$.

Lemma 4.6. If $\{v_1, \dots, v_t\}$ is a linearly independent set of vectors in k^n then so is the set $\{\omega_1^\sim, \dots, \omega_t^\sim\}$ and $\dim_k(H_1(x)) \geq t$.

Proof. Suppose that $\sum_{j=1}^t \lambda_j \omega_j^\sim = 0$, that is, $\sum_{i=1}^n \left(\sum_{j=1}^t \lambda_j x_i^{m_{ij}-1} h_{ij} \right) e_i = \partial_1(g)$ for some g . As in the proof of Lemma 4.5, we derive that

$$\sum_{s < i} X_s g_{si} - \sum_{j=1}^t X_i^{m_{ij}-1} \lambda_j h_{ij} - \sum_{i < s} X_s g_{is} = \sum_{s=1}^p X_i^{m_{is}} h_{is} l_s + \sum_{s=1}^p \sum_{k \neq i} X_k^{m_{ks}} h_{ks} l_s$$

for some $l_s \in P$. Again by passing to the quotient algebra $k[X_i]$, we obtain that $\sum_{j=1}^t X_i^{m_{ij}-1} \lambda_j h_{ij}^\sim = -\sum_{s=1}^p X_i^{m_{is}} h_{is}^\sim l_s^\sim$. But

$$\sum_{j=1}^t X_i^{m_{ij}-1} \lambda_j h_{ij}^\sim = \sum_{j \in S_i} + \sum_{j \notin S_i} = X_i^{m_i-1} \sum_{j \in S_i} \lambda_j h_{ij}^\sim + \sum_{j \notin S_i} X_i^{m_{ij}-1} \lambda_j h_{ij}^\sim,$$

therefore

$$X_i^{m_i-1} \sum_{j \in S_i} \lambda_j h_{ij}^\sim = - \sum_{j \notin S_i} X_i^{m_{ij}-1} \lambda_j h_{ij}^\sim - \sum_{s=1}^p X_i^{m_{is}} h_{is}^\sim l_s^\sim.$$

Thus $\sum_{j \in S_i} \lambda_j h_{ij}^\sim \in \langle X_i \rangle$, which in turn implies that $\sum_{j \in S_i} \lambda_j h_{ij}(a) = 0$ for every i . In particular,

$$\sum_{j=1}^t \lambda_j v_{ij} = \sum_{j \in S_i} \lambda_j h_{ij}(a) = 0$$

for every i , which means that $\sum_{j=1}^t \lambda_j v_j = 0$ in k^n . Using the fact of independence, we conclude that $\lambda_j = 0$ for all j . Hence $\{\omega_1^\sim, \dots, \omega_t^\sim\}$ is a linearly independent set of vectors and the R -module structure of $H_1(x)$ is reduced to its k -vector space one (see Lemma 4.4). Whence $\dim_k(H_1(x)) \geq t$. \square

For every j we have the well defined vector $f'_j = \left(\frac{\partial f_j}{\partial X_1}(a), \dots, \frac{\partial f_j}{\partial X_n}(a) \right) \in k^n$. Notice that $v_j \neq 0$ whenever $f'_j \neq 0$, and in this case $v_j = f'_j$ and $a^{(1)}(\omega_j^\sim) = f'_j$.

Corollary 4.1. *If the Jacobian matrix $J_a = \left[\frac{\partial f_j}{\partial X_i}(a) \right]_{j,i}$ has the rank t then $\dim_k(H_1(x-a)) \geq t$. If Y is nonsingular at point a then $d_1(a) \geq n-d$.*

Proof. There are linearly independent columns $f'_{j_1}, \dots, f'_{j_t}$ of the Jacobian matrix. In particular, so are vectors v_{j_1}, \dots, v_{j_t} . By Lemma 4.6, $\{\omega_{j_1}^\sim, \dots, \omega_{j_t}^\sim\}$ is an independent set of vectors from $H_1(x)$, therefore $\dim_k(H_1(x)) \geq t$. If Y is nonsingular at point a then the rank of the matrix J_a is $n-d$, where $d = \dim(Y)$. It follows that $\dim_k(H_1(x)) \geq n-d$. \square

4.5. Examples. Let $Y \subseteq \mathbb{A}^3$ be a variety given by a prime $\mathfrak{p} \subseteq P = k[X_1, X_2, X_3]$. For the actions of X_j on $R = P/\mathfrak{p}$ we use the notations x, y and z , respectively. If $a = 0 \in Y$ then the Koszul complex $\text{Kos}((x, y, z), R)$ looks like that

$$0 \leftarrow R \xleftarrow{\partial_0} R^3 \xleftarrow{\partial_1} R^3 \xleftarrow{\partial_2} R \leftarrow 0$$

with the operators

$$\partial_0 = \begin{bmatrix} x & y & z \end{bmatrix}, \partial_1 = \begin{bmatrix} -y & -z & 0 \\ x & 0 & -z \\ 0 & x & y \end{bmatrix}, \partial_2 = \begin{bmatrix} z \\ -y \\ x \end{bmatrix}$$

whose homology groups are denoted by H_j and $d_j = \dim_k(H_j)$, $0 \leq j \leq 3$. Recall that $d_0 = 1$, $d_3 = 0$.

Proposition 4.2. *The equalities $\partial_1(\partial_1^2 + 2xz - y^2) = 0$ and $\text{im}(\partial_1^2 + 2xz - y^2) \oplus \text{im}(\partial_2|k \cdot 1) = \text{im}(\partial_2)$ hold.*

Proof. One can easily verify that $Q(t) = t(t^2 + 2xz - y^2) \in R[t]$ is the characteristic polynomial of ∂_1 . By Cayley-Hamilton Theorem, we obtain that $Q(\partial_1) = 0$. In particular, $\text{im}(\partial_1^2 + 2xz - y^2) \subseteq \ker(\partial_1)$. But

$$\partial_1^2 + 2xz - y^2 = \begin{bmatrix} xz & yz & z^2 \\ -xy & -y^2 & -yz \\ x^2 & xy & xz \end{bmatrix},$$

which in turn implies that $(\partial_1^2 + 2xz - y^2)g = \partial_2(xg_1 + yg_2 + zg_3)$ for every $g = (g_1, g_2, g_3) \in R^3$. Thus $\text{im}(\partial_1^2 + 2xz - y^2) \subseteq \text{im}(\partial_2)$. Finally, for every $h \in R$ we have $h = h(0) + xh_1 + yh_2 + zh_3$ and

$$\partial_2(h) = h(0)\partial_2(1) + \partial_2(xh_1 + yh_2 + zh_3) = (\partial_1^2 + 2xz - y^2)g + h(0)\partial_2(1),$$

where $g = (h_1, h_2, h_3) \in R^3$. It remains to use the fact $H_2 = 0$ and the k -linear projection $\partial_2(h) \mapsto h(0)\partial_2(1)$. \square

Thus $g \in \ker(\partial_1) - \text{im}(\partial_2)$ iff $xg_1 = zg_3$, $xg_2 = -yg_3$, $yg_1 = -zg_3$, and

$$g \neq \lambda \begin{bmatrix} z \\ -y \\ x \end{bmatrix} + (2xz - y^2)h$$

for all $\lambda \in k$ and $h \in R^3$, that is, the structure of H_2 is really complicated. But Lemma 4.2 turns out to be useful in this manner. Namely, consider the following hypersurface $Y = \{xy + yz + zx = 0\}$, whose singularity at $a = 0$ is not integral (see Remark 4.2). Then

$$H_1 = k\omega^\sim \text{ with } \omega = \begin{bmatrix} y + z \\ z \\ 0 \end{bmatrix}.$$

Indeed, $\partial_0(\omega) = x(y + z) + yz = 0$ means that $\omega \in \ker(\partial_0)$. Take $g \in \ker(\partial_0)$, that is, $xg_1 + yg_2 + zg_3 = 0$ in R . Then $X_1g_1 + X_2g_2 + X_3g_3 = (X_1(X_2 + X_3) + X_2X_3)h \in \mathfrak{p}$ or $X_3g_3 = X_1((X_2 + X_3)h - g_1) + X_2(X_3h - g_2) \in \langle X_1, X_2 \rangle$ in P . Then $g_3 = X_1q_1 + X_2q_2 \in \langle X_1, X_2 \rangle$ and

$$\begin{aligned} (X_2 + X_3)h - g_1 &= X_3q_1 - X_2l, \\ X_3h - g_2 &= X_3q_2 + X_1l \end{aligned}$$

for some $q_j, l \in P$. It follows that

$$\begin{aligned} g &= \begin{bmatrix} g_1 \\ g_2 \\ g_3 \end{bmatrix} = \begin{bmatrix} (y + z)h + yl - zq_1 \\ zh - xl - zq_2 \\ xq_1 + yq_2 \end{bmatrix} = h \begin{bmatrix} y + z \\ z \\ 0 \end{bmatrix} + \begin{bmatrix} (-y)(-l) + (-z)q_1 \\ x(-l) - zq_2 \\ xq_1 + yq_2 \end{bmatrix} \\ &= h\omega + \partial_1 \begin{bmatrix} -l \\ q_1 \\ q_2 \end{bmatrix}, \end{aligned}$$

that is, $g^\sim = h^\sim\omega^\sim = h(0)\omega^\sim$ (see Lemma 4.4). Using Lemma 4.5, we deduce that $\omega^\sim \neq 0$ and $H_1 = k\omega^\sim$, that is, $d_1 = 1$. By Lemma 4.2, we have $d_2 = d_0 - d_1 + d_2 - d_3 = -i_Y(a) = 0$.

Another example of a singular (at $a = 0$) variety Y is given by the following generators

$$f_1 = X_1^2 - X_2(1 - X_1X_3) + X_3^3, \quad f_2 = X_1^3X_2 + X_2 + (X_2 - 1)X_3^2, \quad f_3 = X_1^3 + X_2^2X_3 + X_3^4$$

in P . Notice that f_1 and f_2 are minimal generators, $m_1 = m_3 = 2$, $m_2 = 1$, $S_1 = \{1\}$, $S_2 = \{1, 2\}$, $S_3 = \{2\}$ and $v_1 = (1, -1, 0)$, $v_2 = (0, 1, -1)$. Note also that these vectors can also be represented by

$$v_1 = \left(\frac{\partial^2 f_1}{\partial X_1^2}(0), \frac{\partial f_1}{\partial X_2}(0), \frac{\partial^2 f_1}{\partial X_3^2}(0) \right), v_2 = \left(f_2(0), \frac{\partial f_2}{\partial X_2}(0), \frac{\partial^2 f_2}{\partial X_3^2}(0) \right)$$

up to constant multiplies. Based on Lemma 4.6, we deduce that $d_1 \geq 2$. Notice also that $x^3 - g = 0$ in $R = k[x]$, where $g = -y^2z - z^4$ (since $f_3 = 0$ in R). That means that x is integral over $k[y, z]$ or a is integral with $i_Y(a) = 0$ (see Remark 4.2).

REFERENCES

- [1] Altman A., Kleinman S., A term of commutative algebra, W.C.Math. (LLC) 2017.
- [2] Bourbaki N., Commutative algebra, Ch. 1-7, Moscow (MIR) 1971.
- [3] Bourbaki N., Homological algebra, Ch. 10, Moscow (Nauka) 1987.
- [4] Cho M., Takaguchi M., *Joint spectra of matrices*, Sci. Rep. Hirosaki Univ. 26 (1979), 15-19.
- [5] Curto R.E., *Fredholm and invertible tuples of operators. The deformation problem*, Trans. Amer. Math. Soc. 266 (1) (1981) 129-159.
- [6] Dosi A. A., *Noncommutative holomorphic functions in elements of a Lie algebra and noncommutative localizations*, Izvestiya Math. RAN, 73 (06) (2009) 77-100.
- [7] Dosiev A. A., *Local left invertibility for operator tuples and noncommutative localizations*. J. K-Theory 4 (1) (2009) 163-191.
- [8] Dosi A. A., *Noncommutative affine spaces and Lie-complete rings*, Comptes Rendus Math. 353 (2) (2015) 149-153.
- [9] Dosi A. A., *Noncommutative localizations of Lie-complete rings*, Comm. Algebra. 44 (2016) 4892-4944.
- [10] Dosi A. A., *Functional calculus on noetherian schemes*, C. R. Acad. Sci. Paris, Ser I, 353 (2015) 57-61.
- [11] Dosi A. A., *The spectrum of a module along scheme morphism and multi-operator functional calculus*, Moscow Math. J. 21 (2) (2021) 1-38
- [12] Eisenbud D., Commutative algebra with a view toward algebraic geometry, Springer New York (1995)
- [13] Eschmeier J., Putinar M., Spectral decompositions and analytic sheaves, London Math. Soc., Clarendon Press, Oxford (1996).
- [14] Fainshtein A. S., *Joint essential spectrum of a family of linear operators*, Funct. Anal. Appl. 14 (2) (1980) 152-153.
- [15] Fainshtein A. S., *Taylor joint spectrum for families of operators generating nilpotent Lie algebra*, J. Operator Theory 29 (1993), 3-27.
- [16] Hartshorne R., Algebraic geometry, Grad. Text. Math. 52 (Springer-Verlag) 1977.
- [17] Helemskii A. Ya., Homology of Banach and topological algebras, MGU, 1986.
- [18] Nakamura T., Yoshimo Y., *Localization functors and cosupport in derived categories of commutative Noetherian rings*, Pacific J. Math., 296 (2) (2018) 405-435
- [19] Nakamura T., Yoshimo Y., *A local duality principle in derived categories of commutative Noetherian rings*, J. Pure Appl. Algebra, 222 (9) (2018) 2580-2595
- [20] Neeman A., *The chromatic tower for $D(R)$* , Topology, 31 (3) (1992) 519-532
- [21] Pirkovskii A. Yu., *Holomorphically finitely generated algebras*, J. Noncom. Geometry (2015).
- [22] Pirkovskii A. Yu., *Taylor's functional calculus and derived categories*, Mathematical methods and applications. Proc. of the 19th RSSU mathematical lectures, 148-163, APKiPPRO, Moscow, 2010.(Russian)
- [23] Putinar M., *Functional calculus with sections of an analytic space*, J. Oper. Th. 4 (1980) 297-306.
- [24] Serre J. P., Local algebra, Springer (2000).
- [25] Taylor J. L., *A joint spectrum for several commuting operators*, J. Funct. Anal. 6 (1970), 172-191.
- [26] Taylor J. L., *A general framework for a multi-operator functional calculus*, Adv. Math. 9 (1972), 183-252.

MIDDLE EAST TECHNICAL UNIVERSITY NORTHERN CYPRUS CAMPUS, GUZELYURT, KKTC, MERSIN 10, TURKEY

Email address: (dosiev@yahoo.com), (dosiev@metu.edu.tr)