

# EQUIVALENT VERSION OF HUPPERT'S CONJECTURE FOR $K_3$ -GROUPS

MOHSEN GHASEMI AND SOMAYEH HEKMATARA

**ABSTRACT.** For a character  $\chi$  of a finite group  $G$  the number  $\text{cod}(\chi) = |G : \text{Ker}(\chi)|/\chi(1)$  is called the codegree of  $\chi$ . In this note, we verify the equivalent version of Huppert's conjecture for  $K_3$ -groups.

## 1. INTRODUCTION

Throughout this paper,  $G$  will be a finite group and  $\text{cd}(G)$  will be the set of the degrees of the complex irreducible characters of  $G$ . The number  $\text{cod}(\chi) = |G : \text{ker}(\chi)|/\chi(1)$  is called the codegree of  $\chi$ . Also the set of codegrees of the irreducible characters of  $G$  is denoted by  $\text{cod}(G)$ . This definition for codegrees first appeared in [16], where the authors use a graph-theoretic approach to compare the structure of a group with its set of codegrees. In [4, 5, 6, 17] some properties of the codegrees of irreducible characters of finite groups have been studied.

In 1990, Huppert proposed the following conjecture:

**Huppert's Conjecture.** Let  $H$  be any finite non-abelian simple group and  $G$  a finite group such that  $\text{cd}(G) = \text{cd}(H)$ . Then,  $G \cong H \times A$ , where  $A$  is abelian.

Many people were devoted to the study of this problem. An analogues of Huppert's conjecture can be proposed and studied for any set of integers related to a finite group. For instance, a dual version of Huppert's conjecture for the set of conjugacy class sizes is considered. For more results see [1, 2, 3]. In this paper, we are concerned with the following conjecture, inspired by Huppert's conjecture:

**Conjecture:** Let  $G$  be a finite group and  $H$  a non-abelian simple group. If  $\text{cod}(G) = \text{cod}(H)$ , then  $G \cong H$ .

This conjecture first proposed in [7], where the authors verified the above conjecture for all projective special linear groups of degree 2. In this article by using the same method in [7], we verify this conjecture for  $K_3$ -groups. Also we recall that Huppert's conjecture is true for  $K_3$ -groups (see [12]).

We refer to [13] for the notation of character theory of finite groups. If  $L \leq G$  and  $\theta \in \text{Irr}(L)$ , then we write  $\text{Irr}(G \mid \theta) = \{\chi \in \text{Irr}(G) \mid [\chi_L, \theta] \neq 0\}$ . Moreover,  $\text{Irr}(G \mid L) = \text{Irr}(G) - \text{Irr}(G/L)$ .

---

*Date:* January 7, 2022.

*2010 Mathematics Subject Classification.* 05C15, 20D05.

*Key words and phrases.* finite group; irreducible character; codegree.

## 2. PRELIMINARIES

In this section we give some results which will be used later in our proofs.

**Proposition 2.1.** [15, Lemma 2] *Let  $S$  be a non-abelian finite simple group. Then there exists  $1_s \neq \theta \in \text{Irr}(S)$  that extends to  $\text{Aut}(S)$ .*

**Proposition 2.2.** [8, Lemma 5] *Let  $N$  be a minimal normal subgroup of  $G$  such that  $N = S_1 \times S_2 \times \cdots \times S_t$ , where  $S_i \cong S$  is a non-abelian simple group. If  $\theta \in \text{Irr}(S)$  extends to  $\text{Aut}(S)$  then  $\theta \times \theta \times \cdots \times \theta \in \text{Irr}(N)$  extends to  $G$ .*

**Proposition 2.3.** [15, Theorem C] *Let  $G$  be a nonabelian finite simple group. Then either  $|\text{cd}(G)| \geq 8$  or one of the following cases holds.*

- (i):  $|\text{cd}(G)| = 4$  and  $G = \text{PSL}(2, 2^f)$ , where  $f \geq 2$ .
- (ii):  $|\text{cd}(G)| = 5$  and  $G = \text{PSL}(2, q)$ , where  $q = p^f$ ,  $p \neq 2$  and  $p^f > 5$ .
- (iii):  $|\text{cd}(G)| = 6$  and  $G = \text{PSL}(3, 4)$  or  $G/N = {}^2B_2(2^{f+1})$ , where  $f \geq 1$ .
- (iv):  $|\text{cd}(G)| = 7$  and  $G = \text{PSL}(3, 3)$ ,  $A_7$ ,  $M_{11}$  or  $J_1$ .

**Remark.** By [10] we see that  $\text{cod}(U_3(3)) = \{1, 2^4.3^2.7, 2^5.3^3, 2^4.3^3, 2^5.3^2, 2^5.7, 2^3.3^3, 3^3.7\}$ . Also  $\text{cod}(U_4(2)) = \{1, 2^6.3^4, 2^5.3^3.5, 2^5.3^4, 2^6.3^3, 2^4.3^4, 2^3.3^3.5, 2^5.3^3, 2^3.3^4, 2^6.3^2, 2^4.3^3, 3^4.5, 2^6.5\}$ .

By [16, Lemma 2.1] we have the following result.

**Proposition 2.4.** *Let  $\chi \in \text{Irr}(G)$ .*

- (i) *For any  $N \trianglelefteq G$  with  $N \leq \ker(\chi)$ ,  $\chi$  may be viewed as an irreducible character of  $G/N$ . The codegree  $\text{cod}(\chi)$  of  $\chi$  is the same whenever  $\chi$  is seen as an irreducible character of  $G$  or  $G/N$ . Furthermore,  $\text{cod}(\chi)$  is independent of the choice of such  $N$ .*
- (ii) *If  $M$  is a subnormal subgroup of  $G$  and  $\varphi$  is an irreducible constituent of  $\chi_M$ , then  $\text{cod}(\varphi)$  divides  $\text{cod}(\chi)$ .*

## 3. MAIN RESULTS

We recall that if  $G$  is a finite simple  $K_3$ -group, then  $G$  is isomorphic to one of the following groups:

$$A_5, A_6, L_2(7), L_2(8), L_2(17), L_3(3), U_3(3), U_4(2).$$

Also by [7], we see that the conjecture is true for  $A_5, A_6, L_2(7), L_2(8), L_2(17), L_3(3)$ . Thus in this section we just verify the conjecture for  $U_3(3)$  and  $U_4(2)$ .

**Theorem 3.1.** *Suppose that  $G$  is a group and  $\text{cod}(G) = \text{cod}(U_3(3))$ . If  $N$  is a maximal normal subgroup of  $G$  then  $G/N \cong U_3(3)$ .*

*Proof.* By our assumption  $G/N$  is a simple group. If  $G/N$  is abelian then  $G' \leq N$  and so  $G \neq G'$ . Now by Proposition 2.4,  $\text{cod}(G/G') \subseteq \text{cod}(G)$  and so  $\text{cod}(G)$  contains a prime power, a contradiction. Thus we may suppose that  $G/N$  is nonabelian simple group. If  $|\text{cd}(G/N)| \geq 8$  then by  $\text{cod}(G/N) \subseteq \text{cod}(G)$  and  $\text{cod}(G) = \text{cod}(U_3(3))$  we see

that  $|\text{cod}(G/N)| = |\text{cod}(U_3(3))| = 8$ . Thus  $|\text{cd}(G/N)| = |\text{cd}(U_3(3))|$  and by Huppert's conjecture  $G/N \cong U_3(3)$ . Thus we may suppose that  $|\text{cd}(G/N)| < 8$  and we find contradiction. Now by Proposition 2.3, we considering the following cases.

**Case 1.**  $|\text{cd}(G/N)| = 4$  and  $G/N \cong \text{PSL}(2, 2^f)$ , where  $f \geq 2$ .

By [19], we know that  $\text{cod}(\text{PSL}(2, 2^f)) = \{1, q(q-1), q(q+1), q^2+1\}$ , where  $q = 2^f$  and  $\text{cod}(U_3(3)) = \{1, 2^4.3^2.7, 2^5.3^3, 2^4.3^3, 2^5.3^2, 2^5.7, 2^3.3^3, 3^3.7\}$ . It is easy to see that  $\text{cod}(\text{PSL}(2, 2^f)) \not\subseteq \text{cod}(U_3(3))$ , a contradiction.

**Case 2.**  $|\text{cd}(G/N)| = 5$  and  $G/N \cong \text{PSL}(2, q)$ , where  $q = p^f$ ,  $p \neq 2$  and  $q > 5$ .

By [19], we know that  $\text{cd}(\text{PSL}(2, q)) = \{1, q(q-1)/2, q(q+1)/2, (q^2-1)/2, q(q-\epsilon(q))\}$ , where  $\epsilon(q) = (-1)^{(q-1)/2}$ . It is easy to see that  $\text{cod}(\text{PSL}(2, p^f)) \not\subseteq \text{cod}(U_3(3))$ , a contradiction.

**Case 3.**  $|\text{cd}(G/N)| = 6$  and  $G/N = \text{PSL}(3, 4)$  or  $G/N = {}^2B_2(2^{2f+1})$ , where  $f \geq 1$ .

First suppose that  $G/N = \text{PSL}(3, 4)$ . We know that  $\text{cd}(\text{PSL}(3, 4)) = \{1, 20, 35, 45, 63, 64\}$  and so  $\text{cod}(\text{PSL}(3, 4)) = \{1, 2^4.3^2.7, 2^6.3^2, 2^6.7, 2^6.5, 3^2.5.7\}$ . Now  $2^6.5 \in \text{cod}(\text{PSL}(3, 4))$  and so  $2^6.5 \notin \text{cod}(U_3(3))$ , a contradiction. Thus we may suppose  $G/N = {}^2B_2(2^{2f+1})$ , where  $q^2 = 2^{2f+1}$ ,  $r = 2^{f+1}$  and  $f \geq 1$ . We know that  $\text{cd}({}^2B_2(2^{2f+1})) = \{1, q^4, q^4+1, (q^2-r+1)(q^2-1), (q^2+r+1)(q^2-1), r(q^2-1)/2\}$  and so  $\text{cod}({}^2B_2(2^{2f+1})) = \{1, (q^4+1)(q^2-1), q^4(q^2-1), \frac{q^4(q^4+1)}{q^2-r+1}, \frac{q^4(q^4+1)}{q^2+r+1}, \frac{2q^4(q^4+1)}{r}\}$ . Now with the simple check we can see that  $\text{cod}({}^2B_2(2^{2f+1})) \not\subseteq \text{cod}(U_3(3))$ , a contradiction.

**Case 4.**  $|\text{cd}(G/N)| = 7$  and  $G/N = \text{PSL}(3, 3)$ ,  $A_7$ ,  $M_{11}$  or  $J_1$ .

We know that  $\text{cd}(\text{PSL}(3, 3)) = \{1, 12, 13, 16, 26, 27, 39\}$  and so  $\text{cod}(\text{PSL}(3, 3)) = \{1, 2^2.3^2.13, 2^4.3^3, 3^3.13, 2^3.3^3, 2^4.13, 2^4.3^2\}$ . Now it is easy to see that  $\text{cod}(\text{PSL}(3, 3)) \not\subseteq \text{cod}(U_3(3))$ . Also if  $G/N \cong A_7$  then  $\text{cd}(A_7) = \{1, 6, 10, 14, 15, 21, 35\}$  and so  $\text{cod}(A_7) = \{1, 2^2.3.5.7, 2^2.3^2.7, 2^2.3^2.5, 2^3.3.7, 2^3.3.5, 2^3.3^2\}$ . Again we see that  $\text{cod}(A_7) \not\subseteq \text{cod}(U_3(3))$ . If  $G/N \cong M_{11}$  then  $\text{cd}(M_{11}) = \{1, 10, 11, 16, 44, 45, 55\}$ . If  $\chi(1) = 16$  then  $\text{cod}(\chi(1)) = 3^2.5.11$  and so  $3^2.5.11 \in \text{cod}(U_3(3))$ , a contradiction. Finally if  $G/N \cong J_1$  then  $\text{cd}(J_1) = \{1, 56, 76, 77, 120, 133, 209\}$ . If  $\chi(1) = 56$  then  $\text{cod}(\chi(1)) = 3.5.11.19$  and so  $3.5.11.19 \in \text{cod}(U_3(3))$ , a contradiction.

□

**Lemma 3.2.** *Suppose that  $G$  is a simple group which is isomorphic to alternating group  $A_n$  ( $n \geq 5$ ) or sporadic group. Then  $4 \leq |\text{cod}(G)| \leq 12$  if and only if  $G$  is isomorphic to one of the following groups.*

$$A_5, A_6, A_7, A_8, M_{11}, M_{12}, M_{22}, M_{23}, J_1.$$

*Proof.* We use the classification of finite simple groups. For the sporadic groups by [10], with the simple check we see that  $G$  is isomorphic to  $M_{11}$  or  $M_{12}$  or  $M_{22}$  or  $M_{23}$  or  $J_1$ . For  $A_n$  where  $5 \leq n \leq 13$ , by [10] we see that  $G \cong A_5$  or  $A_6$  or  $A_7$  or  $A_8$ . Thus we may assume that  $n \geq 14$ . Now the characters corresponding to the non-self-associated partitions  $(n-1, 1)$ ,  $(n-2, 2)$ ,  $(n-2, 1, 1)$ ,  $(n-3, 3)$ ,  $(n-3, 2, 1)$ ,  $(n-3, 1, 1, 1)$ ,  $(n-4, 4)$ ,  $(n-4, 3, 1)$ ,  $(n-4, 2, 2)$ ,  $(n-4, 1, 1, 1, 1)$ ,  $(n-5, 5)$ ,  $(n-5, 4, 1)$ ,  $(n-5, 3, 2)$  have distinct degrees  $n-1$ ,  $\frac{n(n-3)}{2}$ ,  $\frac{(n-1)(n-2)}{2}$ ,

$$\frac{n(n-1)(n-5)}{(n-1)(n-2)(n-3)(n-4)}^6, \frac{n(n-2)(n-4)}{(n-1)(n-2)(n-3)(n-4)}^3, \frac{(n-1)(n-2)(n-3)}{(n-1)(n-2)(n-3)(n-4)}^6, \frac{n(n-1)(n-2)(n-7)}{(n-1)(n-2)(n-3)(n-4)}^{24}, \frac{n(n-1)(n-3)(n-6)}{(n-1)(n-2)(n-3)(n-4)}^8, \frac{n(n-1)(n-4)(n-5)}{(n-1)(n-2)(n-3)(n-4)}^{12},$$

□

**Theorem 3.3.** *Suppose that  $G$  is a group and  $\text{cod}(G) = \text{cod}(U_4(2))$ . If  $N$  is a maximal normal subgroup of  $G$  then  $G/N \cong U_4(2)$ .*

*Proof.* By our assumption  $G/N$  is a nonabelian simple group and so  $|\text{cd}(G/N)| \geq 4$ . We know that  $\text{cod}(G/N) \subseteq \text{cod}(G)$  and  $\text{cod}(G) = \text{cod}(U_4(2))$  and so  $4 \leq |\text{cod}(G/N)| \leq 13$ . if  $|\text{cod}(G/N)| = 13$  then  $|\text{cd}(G/N)| = |\text{cd}(U_4(2))|$  and by Huppert's conjecture  $G/N \cong U_4(2)$ . Thus we may suppose that  $4 \leq |\text{cd}(G/N)| \leq 12$ .

If  $G/N$  is alternating group  $A_n$  or sporadic group then by Lemma 3.2  $G/N$  is isomorphic to  $A_5$  or  $A_6$  or  $A_7$  or  $A_8$  or  $M_{11}$  or  $M_{12}$  or  $M_{22}$  or  $M_{23}$  or  $J_1$ . By [10] we see  $\text{cod}(G/N) \not\subseteq \text{cod}(U_4(2))$ , a contradiction. In what follows, we assume that  $G/N$  is a simple group of Lie type.

*Exceptional groups of Lie type.* Let  $G/N$  be one of the groups  ${}^3D_4(q)$ ,  $G_2(q)$ ,  $F_4(q)$ ,  ${}^2B_2(q^2)(q^2 = 2^{2f+1})$ ,  $E_6(q)$ ,  ${}^2E_6(q)$ ,  $E_7(q)$ ,  $E_8(q)$ ,  ${}^2G_2(q)(q = 3^{2f+1})$  and  ${}^2F_4(q)(q = 2^{2f+1})$ . According to [9, Section 13.9], we see that  $|\text{cod}(G/N)| \geq 14$ , where  $G/N$  is isomorphic to  $F_4(q)$ ,  $E_6(q)$ ,  ${}^2E_6(q)$ ,  $E_7(q)$ ,  $E_8(q)$  or  ${}^2F_4(q)$ . Thus we may suppose that  $G/N$  is isomorphic to  ${}^3D_4(q)$ ,  $G_2(q)$ ,  ${}^2B_2(q^2)$  or  ${}^2G_2(q)$ .

Also by [11] We see that  $|\text{cod}(G/N)| \geq 14$ , where  $G/N$  is isomorphic to  ${}^3D_4(q)$ . If  $G/N = {}^2B_2(2^{2f+1})$ ,  $f \geq 1$ , where  $q^2 = 2^{2f+1}$ , and  $f \geq 1$ , then  $|\text{cd}(G/N)| = 6$ . Now with the simple check we can see that  $\text{cod}({}^2B_2(2^{2f+1})) \not\subseteq \text{cod}(U_4(2))$ , a contradiction. Also if  $G/N$  is isomorphic to  $G_2(q)$  then by [9, Section 13.9] we see that  $G$  has cuspidal character of degree  $\frac{q(q+1)^2(q^2+q+1)}{6}$ , say  $\phi$ . Now it is easy to see that  $\text{cod}(\phi) = \frac{6q^5(q^4+q^2+1)(q-1)^2}{q^2+q+1}$ . Now we see that  $\text{cod}(\phi) \notin \text{cod}(U_4(2))$ , a contradiction. Finally if  $G$  is isomorphic to  ${}^2G_2(q)$ , then by [18] we see that  $G/N$  has character of degree  $(q-1)(q^2-q+1)$ , say  $\delta$ . Now it is easy to see that  $\text{cod}(\delta) = q^3(q+1)$ . No we see that  $\text{cod}(\delta) \notin \text{cod}(U_4(2))$ , a contradiction.

*Type  $A_n$ : Linear groups.* Therefore  $G/N \cong L_{n+1}(q)$ , where  $q$  is a prime power of  $p$  and  $n \geq 1$ . If  $n = 1$  then  $G/N \cong L_2(q)$ . If  $q = 2^f$  where  $f \geq 2$  then it is easy to see that  $\text{cod}(\text{PSL}(2, q)) \not\subseteq \text{cod}(U_4(2))$ . If  $q = p^f$ , where  $p \neq 2$  and  $q > 5$ , then again it is easy to see that  $\text{cod}(\text{PSL}(2, q)) \not\subseteq \text{cod}(U_4(2))$ , a contradiction. Thus we may suppose  $n \geq 2$ , then  $G/N$  has a character of degree  $\frac{q(q^n-1)}{q-1}$ , say  $\chi$ . Now  $\text{cod}(\chi) = q^{\frac{n^2+n-2}{2}} \prod_{i=1}^n \frac{(q^{i+1}-1)(q-1)}{(n+1, q-1)(q^n-1)}$  and  $\text{cod}(\chi) \notin \text{cod}(U_4(2))$ , a contradiction.

*Type  ${}^2A_n$ : Unitary groups.* Therefore  $G/N \cong U_{n+1}(q)$ , where  $n \geq 2$ . Now by [14],  $G/N$  has a character of degree  $\frac{q(q^n-(-1)^n)}{q+1}$ , say  $\chi$ . Now  $\text{cod}(\chi) = q^{\frac{n^2+n-2}{2}} \prod_{i=1}^n \frac{(q^{i+1}-(-1)^{i+1})(q+1)}{(n+1, q+1)(q^n-(-1)^n)}$  and  $\text{cod}(\chi) \notin \text{cod}(U_4(2))$ , a contradiction.

*Type  ${}^\circ B_n$  or  $C_n$ : Odd dimensional orthogonal groups and Symplectic groups.* Therefore  $G/N \cong S_{2n}(q)$  or  $O_{2n+1}(q)$ , where  $n \geq 2$ . If  $n = 2$  then by [9, Section 13.8],  $G/N$  has a unipotent character of  $\frac{q(q+1)^2}{2}$ , say  $\chi$ . Now  $\text{cod}(\chi) = \frac{2q^3(q-1)^2(q^2+1)}{(2, q-1)}$  and  $\text{cod}(\chi) \notin \text{cod}(U_4(2))$ , where  $q \neq 3$ , a contradiction. If  $q = 3$  then  $G/N \cong U_4(2)$ , as wanted. Also if  $n > 2$  then by [9, Section 13.8]  $G/N$  has a steinberg character of degree

$q^{n^2}$ , say  $\theta$ . Now  $\text{cod}(\theta) = \frac{\prod_{i=1}^n (q^{2i}-1)}{(2, q-1)}$  and  $\text{cod}(\chi) \notin \text{cod}(U_4(2))$ , a contradiction.

*Type  $D_n$ : Even dimensional orthogonal groups.* Therefore  $G/N \cong O_{2n}^+(q)$ , where  $n \geq 2$ . If  $n = 2$  then  $G/N \cong L_2(q) \times L_2(q)$ . Suppose that  $q = 2^f$ . Then  $\text{cod}(\theta) = q^2(q^2-1)$ , where  $f \geq 2$  and  $\theta \in \text{Irr}(L_2(q) \times L_2(q))$ . Now suppose that  $q = p^f$ , where  $p \neq 2$  and  $p^f > 5$ . Then  $\text{cod}(\theta) = \frac{q^2(q^2-1)}{4}$ , where  $\theta \in \text{Irr}(L_2(q) \times L_2(q))$ . In each case we see that  $\text{cod}(\theta) \notin \text{cod}(U_4(2))$ , a contradiction.

Also if  $n = 3$  then  $G/N \cong L_4(q)$  which is isomorphic to type  $A_3$ . Thus we may suppose that  $n \geq 4$ . Now by [14],  $G/N$  has an irreducible character of degree  $\frac{q(q^{n-2}+1)(q^n-1)}{q^2-1}$ ,

say  $\phi$ . Now  $\text{cod}(\phi) = \frac{q^{n^2-n-1} \prod_{i=1}^{n-1} (q^{2i}-1)(q^2-1)}{(4, q^n-1)(q^{n-2}+1)}$  and  $\text{cod}(\phi) \notin \text{cod}(U_4(2))$ , a contradiction.

*Type  ${}^2D_n$ : Even dimensional orthogonal groups.* Therefore  $G/N \cong O_{2n}^-(q)$ , where  $n \geq 2$ . If  $n = 2$  then  $G/N \cong L_2(q^2)$  which is isomorphic to type  $A_1$ . Also if  $n = 3$  then  $G/N \cong U_4(q)$  which is isomorphic to type  ${}^2A_3$ . Thus we may suppose that  $n \geq 4$ . Now by [14],  $G/N$  has an irreducible character of degree  $\frac{q(q^{n-2}-1)(q^n+1)}{q^2-1}$ , say  $\sigma$ . Now

$\text{cod}(\sigma) = \frac{q^{n^2-n-1} \prod_{i=1}^{n-1} (q^{2i}-1)(q^2-1)}{(4, q^n+1)(q^{n-2}-1)}$  and  $\text{cod}(\sigma) \notin \text{cod}(U_4(2))$ , a contradiction.

□

**Theorem 3.4.** *Suppose that  $G$  is a finite group and  $\text{cod}(G) = \text{cod}(U_3(3))$  or  $\text{cod}(G) = \text{cod}(U_4(2))$ . Then  $G \cong U_3(3)$  or  $G \cong U_4(2)$ .*

*Proof.* Suppose that  $N$  is the maximal normal subgroup of  $G$ . By Theorem 3.1 and Theorem 3.3,  $G/N \cong U_3(3)$  or  $G/N \cong U_4(2)$ . We show that  $N = 1$ . Suppose to the contrary that  $G$  is a counterexample with minimal order. Thus there is no groups with order less than  $G$ , say  $H$ , such that has a maximal normal subgroup, say  $M$ , such that  $H/M \cong U_3(3)$  or  $U_4(2)$  and  $M \neq 1$ . Assume that  $H \trianglelefteq G$ , where  $H \neq 1$  and  $H < N$ . Now  $G/N \cong (G/H)/(N/H)$  and so  $\text{cod}(G/N) \subseteq \text{cod}(G/H) \subseteq \text{cod}(G)$ . Thus  $\text{cod}(G/H) = \text{cod}(G)$  and so by minimality of  $G$  we get a contradiction. Thus  $N$  is also a minimal normal subgroup of  $G$ . If  $N$  is not abelian then  $N \cong T \times T \times \cdots \times T = T^n$ , where  $T$  is a non-abelian simple group. By Propositions 2.1, 2.2, there exists  $1_N \neq \phi \in \text{Irr}(N)$  which extends to some irreducible character of  $G$ , say  $\theta$ . We have  $\ker(\theta)N \trianglelefteq G$  and by the maximality of  $N$  we have that either  $\ker(\theta)N = N$  or  $\ker(\theta)N = G$ . If  $\ker(\theta)N = N$  then  $\ker(\theta) \subseteq N$ . Since  $N$  is a minimal normal subgroup we have either  $\ker(\theta) = 1$  or  $\ker(\theta) = N$ . We know that  $\theta_N \neq 1$  and so we  $\ker(\theta) = 1$  or  $G = \ker(\theta) \times N$ . If  $\ker(\theta) = 1$  then  $\text{cod}(\theta) = |G|/\theta(1)$ . Since  $\theta(1) \mid |N|$ , it implies that  $|N| = m\theta(1)$  for some positive integer  $m$ . Thus  $|G|/|N| = |G|/m\theta(1)$  and so  $m|G|/|N| = |G|/\theta(1)$ . Therefore  $|G|/|N| \mid |G|/\theta(1) = \text{cod}(\theta)$ . We know that  $\text{cod}(\theta) \in \text{cod}(G)$  and  $\text{cod}(G) = \text{cod}(U_3(3))$  or  $\text{cod}(G) = \text{cod}(U_4(2))$ . We know that  $|G/N| = |U_3(3)| = 2^5 \cdot 3^3 \cdot 7$  and  $|G/N| = |U_4(2)| = 2^6 \cdot 3^4 \cdot 5$ . Now we get contradiction easily. Thus we may suppose that  $G = \ker(\theta) \times N$ . Since  $N$  is the minimal normal subgroup of  $G$ , it follows that  $\ker(\theta)$  is the maximal normal subgroup of  $G$ . Now by Theorem 3.1 and Theorem 3.3 we see that  $G/\ker(\theta) \cong U_3(3)$  or  $G/\ker(\theta) \cong U_4(2)$ . Also we have  $N \cong G/\ker(\theta)$ . Therefore  $N \cong \ker(\theta) \cong G/\ker(\theta)$ . We know that

$\text{cod}(G/\ker(\theta)) \subseteq \text{cod}(G)$  and  $\theta \in \text{Irr}(G/\ker(\theta))$ . Now  $\text{cod}(\theta) = |G|/\theta(1) = |N|^2/\theta(1)$ . Since  $\theta(1) \mid |N|$  and  $\text{cod}(\theta) \in \text{cod}(G)$  we have a contradiction. Thus  $N$  is an abelian group. We know that  $N \leq C_G(N)$  and  $C_G(N) \trianglelefteq G$ . Now since  $N$  is the maximal normal subgroup of  $G$  we conclude that  $N = C_G(N)$  or  $C_G(N) = G$ . If  $C_G(N) = G$  then  $N \leq Z(G)$ . On the other hand if  $G' < G$  then by  $\text{cod}(G/G') \subseteq \text{cod}(G)$ , where  $G \cong U_3(3)$  or  $U_4(2)$ , we see that  $\text{cod}(G)$  contains a prime, a contradiction. Thus  $G$  is perfect and  $N \cap G' = N$ . Thus  $N \leq G'$  and so  $N \leq Z(G) \cap G'$ . Now since  $G/N \cong U_3(3)$  or  $U_4(2)$  we see that  $G$  is a central extension of  $U_3(3)$  or  $U_4(2)$ . Also by [10], we know that  $\text{Mult}(U_3(3)) = 1$  and  $\text{Mult}(U_4(2)) \cong \mathbb{Z}_2$ . Thus  $G \cong U_3(3)$  or  $G \cong U_4(2)$  or  $G \cong 2.U_4(2)$ . Also by [10], we know that  $\text{cd}(2.U_4(2)) = \{1, 4, 20, 36, 60, 64, 80\}$  and we have a contradiction. Thus  $G \cong U_3(3)$  or  $G \cong U_4(2)$  and the assertion is hold. Therefore we may assume that  $N = C_G(N)$ . Now suppose that  $\theta \in \text{Irr}(G|N)$ . We show that  $\theta$  is faithful. If  $\ker(\theta) \neq 1$  then we know that  $N \trianglelefteq \ker(\theta)N \trianglelefteq G$  and so by our assumption about  $N$  we see that either  $N = \ker(\theta)N$  or  $\ker(\theta)N = G$ . First suppose that  $\ker(\theta)N = G$ . Now  $\ker(\theta) \cap N \subseteq N$  and by our assumption about  $N$  we see that  $\ker(\theta) \cap N = 1$  or  $\ker(\theta) \cap N = N$ . If  $\ker(\theta) \cap N = N$  then  $N \subseteq \ker(\theta) \subseteq G$ , a contradiction. Thus  $\ker(\theta) \cap N = 1$ . Thus  $G = \ker(\theta) \times N$ . Now by considering  $G = G'$  we get a contradiction. Therefore  $\ker(\theta) \leq N$  and so by the minimality of  $N$  we have  $\ker(\theta) = N$ , a contradiction. Thus  $\theta$  is faithful for each  $\theta \in \text{Irr}(G|N)$  as asserted. Now we show that if  $1_N \neq \lambda \in \text{Irr}(N)$  then we have  $|I_G(\lambda)|/\theta(1) \in \text{cod}(G)$  for all  $\theta \in \text{Irr}(I_G(\lambda)|\lambda)$ . By [13, Theorem 6.11], we see that  $\theta^G \in \text{Irr}(G)$  for all  $\theta \in \text{Irr}(I_G(\lambda)|\lambda)$ . If  $N \leq \ker(\theta)$  then for each  $n \in N$  we have  $\theta(n) = \theta(1)$  and by [13, Theorem 6.2] we have  $\theta_N(1) = et\theta(1)$  where  $t = |G : I_G(\theta)|$  and  $e$  divides  $|G : H|$ . Now it is easy to see that for each  $n \in N$  we have  $\lambda(n) = \lambda(1)$ , a contradiction. Thus we have  $N \not\leq \ker(\theta)$  and  $\theta \in \text{Irr}(G|N)$ . Now as we showed that we see that  $\ker(\theta^G) = 1$ . Thus we have  $\text{cod}(\theta^G) \in \text{cod}(G)$ , as desired. We have  $\text{cod}(G/N) = \text{cod}(G)$  and  $|I_G(\lambda)|/\theta(1) \in \text{cod}(G)$ . Therefore  $|I_G(\lambda)||N|/|N|\theta(1) \in \text{cod}(G)$ . By [13, Theorem 6.15] we have  $\theta(1) \mid |I_G(\lambda)/N|$ . Thus  $|N|$  divides the codegree of some irreducible character of  $G/N$ , and so  $|N| \mid |G/N|$ .

We know that  $C_G(N) = N$  and so  $|G/N| \mid |\text{Aut}(N)|$ . Also we know that  $N$  is an elementary abelian  $r$ -group. If  $G/N \cong U_3(3)$  then  $r \in \{2, 3, 7\}$ . Thus  $|\text{Aut}(N)|$  divides  $|\text{GL}(5, 2)|$  or  $|\text{GL}(3, 3)|$  or  $|\mathbb{Z}_6|$ . Now since  $|N| \mid |G/N|$  we have a contradiction. Also if  $G \cong U_4(2)$  then  $r \in \{2, 3, 5\}$ . If  $|N| \mid 2^5$  or  $|N| \mid 3^3$  or  $|N| = 5$  then  $|\text{Aut}(N)|$  divides  $|\text{GL}(5, 2)|$  or  $|\text{GL}(3, 3)|$  or  $|\mathbb{Z}_4|$ , a contradiction. Thus we may suppose that  $|N| = 2^6$  or  $|N| = 3^4$ . If  $|N| = 2^6$  then  $\text{Aut}(N) \cong \text{GL}(6, 2)$ . Now it is easy to see that  $\text{cd}(U_4(2)) \not\subseteq \text{cd}(\text{GL}(6, 2))$ , a contradiction. Also if  $|N| = 3^4$  then  $\text{Aut}(N) \cong \text{GL}(4, 3)$  and again  $\text{cd}(U_4(2)) \not\subseteq \text{cd}(\text{GL}(4, 3))$ , a contradiction.  $\square$

## REFERENCES

- [1] N. Ahanjideh, Finite groups with the same conjugacy class sizes as a finite simple group, *International J. Group Theory* 8(1) (2019), 23-33.
- [2] Z. Akhlaghi, T. Le, M. Khatami, J. Moori, H.P. Tong-viet, A dual version of Huppert's conjecture on conjugacy class sizes, *J. Group Theory* 18(1) (2015), 115-131



- [3] Z. Akhlaghi, M. Khatami, Improving Thompson's Conjecture for Suzuki Groups, *Comm. Algebra* 44 (2016) 3927-3932.
- [4] F. Alizadeh, H. Behraves, M. Ghaffarzadeh, M. Ghasemi, S. Hekmatara Groups with few codegrees of irreducible characters, *Comm. Algebra* 47 (2019) 1-7.
- [5] F. Alizadeh, M. Ghasemi, M. Ghaffarzadeh, Finite groups whose codegrees are almost prime, *Comm. Algebra* 49 (2020) 1-7
- [6] R. Bahramian, N. Ahanjideh,  $p$ -divisibility of co-degrees of irreducible characters, *Bull. Austral. Math. Soc.* 103 (2020) 1-5.
- [7] A. Bahri, Z. Akhlaghi, B. Khosravi, An analogue of Huppert's conjecture for character codegrees, *accepted to Bull. Austral. Math. Soc.*
- [8] M. Bianchi, D. Chillag, M. L. Lewis, E. Pacifici, Character degree graphs that are complete graphs, *Proc. Amer. Math. Soc.* 135 (2007), 3, 671-676
- [9] R.W. Carter, Finite Groups of Lie Type: Conjugacy Classes and Complex Characters, Wilwy, New York, 1985.
- [10] J.H. Conway, R.T. Curtis, S.P. Norton, R.A. Parker, R.A. Wilson, *Atlas of Finite Groups*, Clarendon Press, Oxford, 1985.
- [11] D.I. Derizitis, G.O. Michler, Character table and blocks of finite simple triality groups  ${}^3D_4(q)$ , *Trans. Amer. Math. Soc.* 303 (1987) 39-70.
- [12] B. Huppert, Some simple groups which are determined by the set of their character degrees I, *Illinois J. Math.* 44 (2000) 828-842.
- [13] I. M. Isaacs, *Character Theory of Finite Groups*, Academic Press, San Diego, California, 1976
- [14] L. He, G. Zhu, Nonsolvable normal subgroups and irreducible character degrees, *J. Algebra* 372 (2012), 68-84.
- [15] G. Malle, A. Moretó, Nonsolvable groups with few character degrees, *J. Algebra* 294 (1) (2005), 117-126
- [16] G. Qian, Y. Wang, H. Wei, Co-degrees of irreducible characters in finite groups, *J. Algebra* 312 (2007) 946-955.
- [17] Z. Sayanjali, Z. Akhlaghi, B. Khosravi, On the codegrees of finite groups, *Comm. Algebra* (48) (2020) 1327-1332.
- [18] H.N. Ward On Ree's series of simple groups, *Trans. Amer. Math. Soc.* 121 (1966) 62-89
- [19] D. L. White, Character degrees of extensions of  $\text{PSL}(2, q)$  and  $\text{SL}(2, q)$ , *J. Group Theory* 16 (1) (2013), 1-33.

DEPARTMENT OF MATHEMATICS, URMIA UNIVERSITY, URMIA 57135, IRAN  
 Email address: m.ghasemi@urmia.ac.ir

DEPARTMENT OF MATHEMATICS, URMIA UNIVERSITY, URMIA 57135, IRAN  
 Email address: somayehhekmatara93@gmail.com