

STABILIZERS IN HIGMAN–THOMPSON GROUPS

JAMES BELK, JAMES HYDE, AND FRANCESCO MATUCCI

ABSTRACT. We investigate stabilizers of finite sets of rational points in Cantor space for the Higman–Thompson groups $V_{n,r}$. We prove that the pointwise stabilizer is an iterated ascending HNN extension of $V_{n,q}$ for any $q \geq 1$. We also prove that the commutator subgroup of the pointwise stabilizer is simple, and we compute the abelianization. Finally, for each n we classify such pointwise stabilizers up to isomorphism.

In recent work, Golan and Sapir use iterated ascending HNN extensions to describe the stabilizer of any finite set of rational points in $(0, 1)$, and they classify such stabilizers up to isomorphism [4]. Here we record analogous results for stabilizers in Thompson’s group V , and more generally for the Higman–Thompson group $V_{n,r}$.

For $n \geq 2$ and $r \geq 1$, let $\mathfrak{C}_{n,r}$ denote the Cantor space $\{1, \dots, r\} \times \{1, \dots, n\}^\omega$, and let $V_{n,r}$ be the associated Higman–Thompson group (denoted $G_{n,r}$ in [2, 5]). A point in $\mathfrak{C}_{n,r}$ is said to be **rational** if the associated sequence is eventually repeating. If S is a finite set of rational points in $\mathfrak{C}_{n,r}$, the **stabilizer** $\text{Stab}(S)$ is the group of elements of $V_{n,r}$ that fix S setwise, the **pointwise stabilizer** $\text{Fix}(S)$ is the subgroup of elements that fix S pointwise. Let $\text{Fix}_0(S)$ be the subgroup consisting of elements that are the identity in some neighborhood of S .

Our main theorem is the following:

Theorem 1. *Let $n \geq 2$ and $q, r \geq 1$, and let $S \subset \mathfrak{C}_{n,r}$ be a finite, nonempty set of rational points. Then:*

- (1) *$\text{Fix}(S)$ is an iterated ascending HNN extension of $V_{n,q}$.*
- (2) *$\text{Fix}(S)$ and $\text{Stab}(S)$ have type F_∞ .*
- (3) *The commutator subgroup of $\text{Fix}(S)$ is simple, and is the intersection of $\text{Fix}_0(S)$ with the commutator subgroup of $V_{n,r}$.*
- (4) *The abelianization of $\text{Fix}(S)$ is the direct sum of $\mathbb{Z}^{|S|}$ with the abelianization of $V_{n,r}$.*

For (1), recall that a group G is an **ascending HNN extension** of a subgroup H if G has presentation

$$G = \langle H, t \mid tht^{-1} = \varphi(h) \text{ for all } h \in H \rangle$$

The first author has been partially supported by EPSRC grant EP/R032866/1 during the creation of this paper.

The third author is a member of the Gruppo Nazionale per le Strutture Algebriche, Geometriche e le loro Applicazioni (GNSAGA) of the Istituto Nazionale di Alta Matematica (INdAM) and gratefully acknowledges the support of the Fundação para a Ciência e a Tecnologia (CEMAT-Ciências FCT projects UIDB/04621/2020 and UIDP/04621/2020) and of the Università degli Studi di Milano–Bicocca (FA project ATE-2016-0045 “Strutture Algebriche”).

where $\varphi: H \rightarrow H$ is a monomorphism. More generally, G is an **iterated ascending HNN extension** of H if there exists a chain

$$H = H_0 \leq H_1 \leq \cdots \leq H_n = G$$

of subgroups of H such that each H_i is an ascending HNN extension of H_{i-1} . We prove (1) in Section 2 below.

For (2), it is well known that an ascending HNN extension of a group of type F_∞ has type F_∞ . Since the groups $V_{n,r}$ all have type F_∞ by a result of Brown [2, Theorem 7.3.1], it follows from statement (1) above that $\text{Fix}(S)$ always has type F_∞ when S is a finite set of rational points. Since $\text{Fix}(S)$ has finite index in $\text{Stab}(S)$, the stabilizer has type F_∞ as well, and thus (2) follows from (1). The authors will use these results in [1] to prove finiteness properties for certain interesting groups, including a class of Röver–Nekrashevych groups.

For parts (3) and (4), recall that Higman proved that the commutator subgroup of $V_{n,r}$ is always simple, and that the abelianization of $V_{n,r}$ is trivial if n is even and cyclic of order two if n is odd [5, Chapter 5]. Thus the abelianization of $\text{Fix}(S)$ is $\mathbb{Z}^{|S|}$ if n is even and $\mathbb{Z}^{|S|} \oplus \mathbb{Z}_2$ if n is odd. We prove (3) and (4) in Theorems 10 and 11 below.

In addition to the above results, the following theorem classifies the isomorphism types of the groups $\text{Fix}(S)$.

Theorem 2. *Let $n \geq 2$ and $q, r \geq 1$, and let $S \subset \mathfrak{C}_{n,q}$ and $S' \subset \mathfrak{C}_{n,r}$ be finite, nonempty sets of rational points. Then $\text{Fix}(S)$ and $\text{Fix}(S')$ are isomorphic if and only if $|S| = |S'|$.*

The proof of this theorem can be found at the end of Section 3.

1. BACKGROUND

1.1. Thompson-like homeomorphisms. We view $\mathfrak{C}_{n,r}$ as a space of infinite sequences of digits, where the first digit of each sequence lies in $\{1, \dots, r\}$ and the remaining digits lie in $\{1, \dots, n\}$. If $\alpha \in \{1, \dots, r\} \times \{1, \dots, n\}^*$ is any finite prefix, the associated **cone** C_α consists of all sequences in $\mathfrak{C}_{n,r}$ that have α as a prefix. Note that the clopen sets in $\mathfrak{C}_{n,r}$ are precisely the finite unions of cones.

Given any two cones $C_\alpha \subseteq \mathfrak{C}_{n,q}$ and $C_\beta \subseteq \mathfrak{C}_{n,r}$, the corresponding **prefix-replacement homeomorphism** from C_α to C_β is the function that maps any word $\alpha\psi \in C_\alpha$ to the corresponding word $\beta\psi \in C_\beta$. If $U \subseteq \mathfrak{C}_{n,q}$ and $U' \subseteq \mathfrak{C}_{n,r}$ are open, we say that a homeomorphism $f: U \rightarrow U'$ is **Thompson-like** if each point of U is contained in a cone on which f restricts to a prefix replacement. The **Higman–Thompson group** $V_{n,r}$ is the group of all Thompson-like homeomorphisms of $\mathfrak{C}_{n,r}$.

1.2. Types of clopen sets. Every clopen set $E \subseteq \mathfrak{C}_{n,r}$ can be expressed as a finite, disjoint union $C_{\alpha_1} \sqcup \cdots \sqcup C_{\alpha_s}$ of cones. Such a decomposition is not unique, since we can replace any cone C_{α_i} by the disjoint union of the n subcones $\{C_{\alpha_i j}\}_{j=1}^n$. However, any two decompositions of the same clopen set into cones must have the same number of cones modulo $n-1$. We refer to this element of \mathbb{Z}_{n-1} as the **type** of E , denoted $\text{type}(E)$. Note that any cone has type 1, the whole Cantor space $\mathfrak{C}_{n,r}$ has type r (modulo $n-1$), and

$$\text{type}(E_1 \cup E_2) = \text{type}(E_1) + \text{type}(E_2) - \text{type}(E_1 \cap E_2)$$

for any two clopen sets $E_1, E_2 \subseteq \mathfrak{C}_{n,r}$. As an immediate consequence, we see that $\text{type}(E \setminus F) = \text{type}(E) - \text{type}(E \cap F)$, for any clopen sets $E, F \subseteq \mathfrak{C}_{n,r}$.

Proposition 3. *Let $n \geq 2$, let $q, r \geq 1$, and let $E \subseteq \mathfrak{C}_{n,q}$ and $E' \subseteq \mathfrak{C}_{n,r}$ be clopen sets. Then there exists a Thompson-like homeomorphism $f: E \rightarrow E'$ if and only if $\text{type}(E) = \text{type}(E')$*

Proof. If a Thompson-like homeomorphism $f: E \rightarrow E'$ exists, then there is a suitable subdivision of E into cones so that their images are cones too, and therefore $\text{type}(E) = \text{type}(E')$. Conversely, if $\text{type}(E) = \text{type}(E')$, there exists a positive integer m with

$$m \equiv \text{type}(E) \pmod{n-1}$$

and suitable cones $C_1, \dots, C_m \subseteq \mathfrak{C}_{n,q}$ and $C'_1, \dots, C'_m \subseteq \mathfrak{C}_{n,r}$ so that $E = \biguplus_{i=1}^m C_i$ and $E' = \biguplus_{i=1}^m C'_i$. In this case, we can define a Thompson-like homeomorphism $f: E \rightarrow E'$ by mapping each C_i to C'_i by a prefix replacement. \square

Corollary 4. *Let $E \subset \mathfrak{C}_{n,r}$ be a proper, nonempty clopen set, and let $q = \text{type}(E)$. Then $\text{Fix}(\mathfrak{C}_{n,r} \setminus E) \cong V_{n,q}$.*

Proof. By Proposition 3, there is a Thompson-like homeomorphism $h: E \rightarrow \mathfrak{C}_{n,q}$. Then $f \mapsto h(f|_E)h^{-1}$ defines an isomorphism $\text{Fix}(\mathfrak{C}_{n,r} \setminus E) \rightarrow V_{n,q}$. \square

1.3. Germs at fixed points. If $p \in \mathfrak{C}_{n,r}$ is a rational point, the **group of germs** at p is the quotient

$$(V_{n,r})_p = \text{Fix}(p)/\text{Fix}_0(p).$$

If $f \in \text{Fix}(p)$, we let $(f)_p$ denote its image in $(V_{n,r})_p$. Note that if $f, g \in \text{Fix}(p)$, then $(f)_p = (g)_p$ if and only if f and g agree on a neighborhood of p .

Proposition 5. *Let $p \in \mathfrak{C}_{n,r}$ be a rational point, and write $p = \alpha\bar{\beta}$ for some finite sequences α and β , where β is not a power of a shorter sequence. Then there exists an $f \in V_{n,r}$ that agrees with the prefix replacement $\alpha\psi \mapsto \alpha\beta\psi$ in a neighborhood of p , and $(V_{n,r})_p$ is the infinite cyclic group generated by $(f)_p$*

Proof. Note that $C_{\alpha\beta}$ has nonempty complement. Since $C_{\alpha\beta}$ and $C_{\alpha\beta^2}$ have type 1, their complements have type $r-1$, so by Proposition 3 there exists a Thompson-like homeomorphism $h: \mathfrak{C}_{n,r} \setminus C_{\alpha\beta} \rightarrow \mathfrak{C}_{n,r} \setminus C_{\alpha\beta^2}$. Let f be the element of $V_{n,r}$ which agrees with h on $\mathfrak{C}_{n,r} \setminus C_{\alpha\beta}$ and maps $C_{\alpha\beta}$ to $C_{\alpha\beta^2}$ by a prefix replacement. Then f agrees with the prefix replacement $\alpha\psi \mapsto \alpha\beta\psi$ on $C_{\alpha\beta}$.

Now observe that if $g \in \text{Fix}(p)$, then g must act as a prefix replacement of the form $\alpha\psi \mapsto \alpha\beta^k\psi$ or $\alpha\beta^k\psi \mapsto \alpha\psi$ in a neighborhood of p for some $k \geq 0$. In the first case, g agrees with f^k in a neighborhood of p , and hence $(g)_p = (f)_p^k$. In the second case, g agrees with f^{-k} in a neighborhood of p , and hence $(g)_p = (f)_p^{-k}$. Thus $(V_{n,r})_p$ is precisely the infinite cyclic group generated by $(f)_p$. \square

We refer to the germ $(f)_p$ described in Proposition 5 as the **attracting generator** for $(V_{n,r})_p$.

We will need the following generalization of Proposition 5.

Proposition 6. *Let $S \subset \mathfrak{C}_{n,r}$ be a finite set of rational points, and let $s \in S$. Then there exists an $f \in \text{Fix}_0(S \setminus \{s\})$ that fixes s such that $(f)_s$ is the attracting generator for $(V_{n,r})_s$.*

Proof. Let E be a clopen neighborhood of $S \setminus \{s\}$ that does not contain s . Write $s = \alpha\bar{\beta}$, where α and β are finite words and β is not a power of a shorter word. Replacing α with $\alpha\beta^k$ for some k , we may assume that C_α is properly contained in $\mathfrak{C}_{n,r} \setminus E$. Since $\text{type}(C_\alpha) = \text{type}(C_{\alpha\beta}) = 1$, by the argument at the end of Subsection 1.2 we see that

$$\text{type}(\mathfrak{C}_{n,r} \setminus (E \uplus C_\alpha)) = \text{type}(\mathfrak{C}_{n,r} \setminus (E \uplus C_{\alpha\beta}))$$

so by Proposition 3 there is a Thompson-like homeomorphism

$$h: \mathfrak{C}_{n,r} \setminus (E \uplus C_\alpha) \rightarrow \mathfrak{C}_{n,r} \setminus (E \uplus C_{\alpha\beta}).$$

Then the element $f \in V_{n,r}$ that is the identity on E , maps C_α to $C_{\alpha\beta}$ by a prefix replacement, and agrees with h on $\mathfrak{C}_{n,r} \setminus (E \uplus C_\alpha)$ has the desired properties. \square

Corollary 7. *If $S \subset \mathfrak{C}_{n,r}$ is a finite set of rational points, then there exists an $f \in \text{Fix}(S)$ so that $(f)_s$ is the attracting generator for $(V_{n,r})_s$ for each $s \in S$.*

Proof. Let $S = \{s_1, \dots, s_m\}$. By Proposition 6, there exists for each i an element $f_i \in \text{Fix}_0(S \setminus \{s_i\})$ that fixes s_i such that $(f_i)_{s_i}$ is the attracting generator for $(V_{n,r})_{s_i}$. Then the product $f = f_1 \cdots f_m$ has the desired property. \square

2. ASCENDING HNN EXTENSIONS

In this section we prove that pointwise stabilizers are ascending HNN extensions. It follows from Bass-Serre theory (see [6]) that a group G is an ascending HNN extension of a group H if and only if G acts by automorphisms on some directed tree Γ with the following properties:

- (1) Γ has exactly one orbit of vertices under G .
- (2) The stabilizer of each vertex of Γ is isomorphic to H .
- (3) Each vertex of Γ has exactly one outgoing edge.

The following proposition translates this geometry into algebra.

Proposition 8. *Let G be a group. Let $t \in G$ have infinite order, let $H \leq G$, and suppose that*

- (1) $\langle t \rangle \cap H = 1$,
- (2) $t^{-1}Ht \leq H$, and
- (3) $\bigcup_{i,j \in \mathbb{N}} t^i H t^{-j} = G$.

Then G is an ascending HNN extension of H by t .

Proof. Let Γ be the directed graph whose vertices are the left cosets of H in G , with a directed edge from gH to gtH for every vertex gH . Note that if $gH = g'H$ for some $g, g' \in G$, then since $t^{-1}Ht \leq H$ we have $gtH = g'tH$, and thus Γ has only one directed edge emanating from each vertex. Clearly G acts on Γ by automorphisms with one orbit of vertices and one orbit of edges, so it suffices to prove that Γ is a directed tree.

Note first that Γ is connected, for by condition (3) if gH is any vertex of Γ then $gt^jH = t^iH$ for some $i, j \in \mathbb{N}$, and hence gH lies in the same component of Γ as the vertex H . To prove that Γ has no directed cycles, let K be the ascending union of the subgroups $\{t^n H t^{-n}\}_{n \in \mathbb{N}}$. It follows from condition (1) that $K \cap \langle t \rangle = 1$, so by condition (3) the group G is the disjoint union of the cosets $\{t^j K\}_{j \in \mathbb{Z}}$, each of which is a disjoint union of cosets of H . Since $gtH \subseteq t^{j+1}K$ whenever $gH \subseteq t^j K$,

we conclude that the graph Γ has no directed cycles. Since every vertex of Γ has only one outgoing edge, it follows that Γ is a directed tree. \square

We can now give a proof of the first part of Theorem 1.

Theorem 9. *Let $n \geq 2$ and $q, r \geq 1$, and let S be a finite, nonempty set of rational points in $\mathfrak{C}_{n,r}$. Then $\text{Fix}(S)$ is an iterated ascending HNN extension of $V_{n,q}$.*

Proof. We prove the result by induction on $|S|$. Fix a point $s \in S$. By Proposition 6 there exists an element $f \in \text{Fix}(S)$ such that $(f)_s$ is the attracting generator for $(V_{n,r})_s$. Let C_α be a cone containing s on which f acts as a prefix replacement $\alpha\psi \mapsto \alpha\beta\psi$, and note that C_α is disjoint from $S \setminus \{s\}$. Let T be a clopen subset of C_α that contains $C_{\alpha\beta}$ and satisfies

$$\text{type}(T) \equiv r - q \pmod{n-1},$$

and let $H = \text{Fix}(S \cup T)$. Since

$$\text{type}(\mathfrak{C}_{n,r} \setminus T) \equiv q \pmod{n-1}$$

it follows from Proposition 3 that there exists a Thompson-like homeomorphism $k: \mathfrak{C}_{n,r} \setminus T \rightarrow \mathfrak{C}_{n,q}$, which determines an isomorphism $\text{Fix}(T) \rightarrow V_{n,q}$ as in Corollary 4. If $|S| = 1$, then $H = \text{Fix}(T)$ and hence $H \cong V_{n,q}$. If $|S| \geq 2$ then this isomorphism instead maps H to the subgroup $\text{Fix}(k(S \setminus \{s\}))$ of $V_{n,q}$, which by our induction hypothesis is an iterated ascending HNN extension of $V_{n,q}$. Thus it suffices to prove that $\text{Fix}(S)$ is an ascending HNN extension of H .

We verify the three conditions in Proposition 8. Clearly f has infinite order. Next, since $f^{-1}(T)$ contains T we have that $f^{-1}Hf = \text{Fix}(S \cup f^{-1}(T)) \leq H$. Finally, if $g \in \text{Fix}(S)$ then $(g)_s = (f)_s^i$ for some integer $i \in \mathbb{Z}$. Let U be a neighborhood of s on which g agrees with f^i , and let $j \geq |i|$ so that $C_{\alpha\beta^j} \subseteq U$ and hence $f^j(T) \subseteq U$. Since $f^{-i}g$ is the identity on U , it follows that $f^{-i-j}gf^j$ is the identity on T . Then $f^{-i-j}gf^j$ lies in H , and therefore $g \in f^{i+j}Hf^{-j}$. \square

3. THE COMMUTATOR SUBGROUP

In this section we analyze the structure of the commutator subgroup $[\text{Fix}(S), \text{Fix}(S)]$ as well as the abelianization of $\text{Fix}(S)$.

Theorem 10. *Let $n \geq 2$ and $r \geq 1$, and let $S \subset \mathfrak{C}_{n,r}$ be a finite set of rational points. Then*

$$[\text{Fix}(S), \text{Fix}(S)] = \text{Fix}_0(S) \cap [V_{n,r}, V_{n,r}].$$

and this group is simple.

Proof. Let $S = \{s_1, \dots, s_m\}$, and let $\pi: \text{Fix}(S) \rightarrow \prod_{i=1}^m (V_{n,r})_{s_i}$ be the homomorphism defined by $\pi(f) = ((f)_{s_1}, \dots, (f)_{s_m})$. Then the kernel of π is $\text{Fix}_0(S)$ and the codomain is a free abelian group, so it follows that $[\text{Fix}(S), \text{Fix}(S)] \subseteq \text{Fix}_0(S)$, and hence $[\text{Fix}(S), \text{Fix}(S)] \subseteq \text{Fix}_0(S) \cap [V_{n,r}, V_{n,r}]$.

For the opposite inclusion, let $E_1 \subseteq E_2 \subseteq \dots$ be an ascending sequence of nonempty clopen sets whose union is $\mathfrak{C}_{n,r} \setminus S$, and let G_i denote the subgroup of $V_{n,r}$ consisting of elements that are supported on E_i . Note that $\text{Fix}_0(S)$ is the ascending union of the subgroups G_i , and that each G_i is isomorphic to V_{n,t_i} , where $t_i = \text{type}(E_i)$. We claim that $[G_i, G_i] = [V_{n,r}, V_{n,r}] \cap G_i$ for each i .

The inclusion $[G_i, G_i] \subseteq [V_{n,r}, V_{n,r}] \cap G_i$ is clear. For the opposite inclusion, if n is even then $[G_i, G_i] = G_i$ and $[V_{n,r}, V_{n,r}] = V_{n,r}$ and we are done. If n is odd,

then $[V_{n,r}, V_{n,r}]$ has index two in $V_{n,r}$, and its complement contains all elements of order two (see [5, Chapter 5]). Then any element of order two in G_i is not contained in $[V_{n,r}, V_{n,r}]$, which means that $[V_{n,r}, V_{n,r}] \cap G_i$ is a proper subgroup of G_i . Since $[G_i, G_i]$ has index two in G_i and $[G_i, G_i] \subseteq [V_{n,r}, V_{n,r}] \cap G_i$, it follows that $[G_i, G_i] = [V_{n,r}, V_{n,r}] \cap G_i$.

We conclude that

$$\text{Fix}_0(S) \cap [V_{n,r}, V_{n,r}] = \bigcup_{i \in \mathbb{N}} G_i \cap [V_{n,r}, V_{n,r}] = \bigcup_{i \in \mathbb{N}} [G_i, G_i] \subseteq [\text{Fix}(S), \text{Fix}(S)]$$

and therefore $[\text{Fix}(S), \text{Fix}(S)] = \text{Fix}_0(S) \cap [V_{n,r}, V_{n,r}]$. Moreover, since each G_i is isomorphic to $V_{n,q}$ for some $q \geq 1$, each of the commutator subgroups $[G_i, G_i]$ is simple [5], so the ascending union $[\text{Fix}(S), \text{Fix}(S)]$ must be simple as well. \square

Theorem 11. *Let $S \subseteq \mathfrak{C}_{n,r}$ be a finite set of rational points. Then*

$$\text{Fix}(S)/[\text{Fix}(S), \text{Fix}(S)] \cong \begin{cases} \mathbb{Z}^{|S|} & \text{if } n \text{ is even,} \\ \mathbb{Z}^{|S|} \oplus \mathbb{Z}_2 & \text{if } n \text{ is odd.} \end{cases}$$

Proof. Let $S = \{s_1, \dots, s_m\}$ and let $\pi: \text{Fix}(S) \rightarrow \prod_{i=1}^m (V_{n,r})_{s_i}$ be the homomorphism defined by

$$\pi(f) = ((f)_{s_1}, \dots, (f)_{s_m}).$$

Let $A = V_{n,r}/[V_{n,r}, V_{n,r}]$, and let $\rho: V_{n,r} \rightarrow A$ be the quotient homomorphism. Note that $\prod_{i=1}^m (V_{n,r})_{s_i} \cong \mathbb{Z}^m$, and that A is trivial if n is even and $A \cong \mathbb{Z}_2$ if n is odd (see [5, Chapter 5]). Let $\sigma: \text{Fix}(S) \rightarrow \prod_{i=1}^m (V_{n,r})_{s_i} \times A$ be the homomorphism $\pi \times \rho$. Then

$$\ker(\sigma) = \ker(\pi) \cap \ker(\rho) = \text{Fix}_0(S) \cap [V_{n,r}, V_{n,r}] = [\text{Fix}(S), \text{Fix}(S)]$$

by Theorem 10, so it suffices to prove that σ is surjective.

By Proposition 6, there exists for each i an element $f_i \in \text{Fix}_0(S \setminus \{s_i\})$ so that $(f_i)_{s_i}$ is a generator for $(V_{n,r})_{s_i}$. Then π maps f_1, \dots, f_m to a generating set for $\prod_{i=1}^m (V_{n,r})_{s_i}$, and therefore π is surjective. Thus, all that remains is to prove that $\text{Fix}_0(S) = \ker(\pi)$ maps onto A under ρ .

If n is even then A is trivial and we are done. If n is odd, then any element of order two in $V_{n,r}$ maps to the nontrivial element of A under ρ . In particular, if we choose disjoint clopen sets E, E' of the same type that are disjoint from S and let $h: E \rightarrow E'$ be a Thompson-like homeomorphism, then the element $f \in V_{n,r}$ that agrees with h on E , agrees with h^{-1} on E' , and is the identity elsewhere lies in $\text{Fix}_0(S)$ and maps to the nontrivial element of A under ρ . \square

Finally, we are ready to prove the following

Theorem 2. *Let $n \geq 2$, let $q, r \geq 1$, and let $S \subset \mathfrak{C}_{n,q}$ and $S' \subset \mathfrak{C}_{n,r}$ be finite sets of rational points. Then $\text{Fix}(S)$ is isomorphic to $\text{Fix}(S')$ if and only if $|S| = |S'|$.*

Proof. If $\text{Fix}(S) \cong \text{Fix}(S')$ then $|S| = |S'|$ by Theorem 11. For the converse, suppose $|S| = |S'|$, and choose a bijection $\varphi: S \rightarrow S'$.

By Corollary 7, there exists an $f \in \text{Fix}(S)$ so that for each $s \in S$, the germ $(f)_s$ is the attracting generator for $(V_{n,q})_s$. For each $s \in S$, choose a cone containing s on which f acts as a prefix replacement, and let E be the complement of the union of these cones. Shrinking the cones if necessary, we may assume that E is nonempty. Note then that E is clopen, $f(E) \supset E$, and $\bigcup_{n \in \mathbb{N}} f^n(E) = \mathfrak{C}_{n,q} \setminus S$.

Repeating the last paragraph for S' , we obtain an element $f' \in \text{Fix}(S')$ and a clopen set $E' \subset \mathfrak{C}_{n,r}$ so that $(f')_s$ is the attracting generator for $(V_{n,r})_s$ for each $s \in S'$, and moreover $f'(E') \supset E'$ and $\bigcup_{n \in \mathbb{N}} (f')^n(E') = \mathfrak{C}_{n,r} \setminus S'$. Replacing E' by a larger clopen subset of $f(E')$ if necessary, we may assume that $\text{type}(E') = \text{type}(E)$.

Now, observe that $\mathfrak{C}_{n,q} \setminus S$ is the disjoint union of E and the sets

$$E_k = f^k(E) \setminus f^{k-1}(E)$$

for $k \geq 1$. Similarly, $\mathfrak{C}_{n,r} \setminus S'$ is the disjoint union of E' and analogous sets E'_k ($k \geq 1$). Note that $\text{type}(E_k) = \text{type}(E'_k) = 0$ for each $k \geq 1$. Choose Thompson-like homeomorphisms $h_0: E \rightarrow E'$ and $h_1: E_1 \rightarrow E'_1$, and let $h: \mathfrak{C}_{n,q} \rightarrow \mathfrak{C}_{n,r}$ be the homeomorphism that agrees with φ on S , agrees with h_0 on E , and agrees with $(f')^{n-1}h_1f^{1-n}$ on E_n for each $n \geq 1$. We claim that $h\text{Fix}(S)h^{-1} = \text{Fix}(S')$, and hence $\text{Fix}(S) \cong \text{Fix}(S')$.

By symmetry, it suffices to prove that $h\text{Fix}(S)h^{-1} \subseteq \text{Fix}(S')$. If $g \in \text{Fix}(S)$, then clearly hgh^{-1} fixes S' pointwise, so it suffices to prove that hgh^{-1} is an element of $V_{n,r}$. Since h restricts to a Thompson-like homeomorphism $\mathfrak{C}_{n,q} \setminus S \rightarrow \mathfrak{C}_{n,r} \setminus S'$, it suffices to prove that hgh^{-1} is Thompson-like in a neighborhood of S' .

To see this, observe first that hfh^{-1} agrees with f' on $\mathfrak{C}_{n,r} \setminus E'$. More generally, for every $k \in \mathbb{Z}$ the functions hf^kh^{-1} and $(f')^k$ agree in a neighborhood of S' . For each $s \in S$, we know that $(f)_s$ is a generator for the infinite cyclic group $(V_{n,q})_s$, so g must agree with some power of f in a neighborhood of s . It follows that hgh^{-1} agrees with some power of f' in a neighborhood of $h(s)$, and therefore hgh^{-1} is Thompson-like in a neighborhood of $h(s)$. \square

Remark 12. Note that $\text{Fix}_0(S) \cong \text{Fix}_0(S')$ for any finite sets S and S' (regardless of size), since any Thompson-like homeomorphism $\mathfrak{C}_{n,q} \setminus S \rightarrow \mathfrak{C}_{n,r} \setminus S'$ conjugates $\text{Fix}_0(S)$ to $\text{Fix}_0(S')$. (More generally, if $U \subset \mathfrak{C}_{n,q}$ and $U' \subset \mathfrak{C}_{n,r}$ are nonempty, non-compact open sets, then the subgroup of $V_{n,q}$ consisting of elements supported on a compact subset of U is isomorphic to the subgroup of $V_{n,r}$ consisting of elements supported on a compact subset of U' .) Since $[\text{Fix}(S), \text{Fix}(S)]$ is the commutator subgroup of $\text{Fix}_0(S)$, it follows that $[\text{Fix}(S), \text{Fix}(S)] \cong [\text{Fix}(S'), \text{Fix}(S')]$ for any nonempty finite sets $S \subset \mathfrak{C}_{n,q}$ and $S' \subset \mathfrak{C}_{n,r}$.

REFERENCES

- [1] J. Belk, J. Hyde, and F. Matucci, Finite germ extensions. Preprint in preparation.
- [2] K. Brown, Finiteness properties of groups. *Journal of Pure and Applied Algebra* **44.1–3** (1987): 45–75. Crossref
- [3] R. Geoghegan, *Topological Methods in Group Theory*. Graduate Texts in Mathematics **243**, Springer, 2008. Crossref.
- [4] G. Golan and M. Sapir, On the stabilizers of finite sets of numbers in the R. Thompson group F . *St. Petersburg Mathematical Journal* **29.1** (2018): 51–79. Crossref.
- [5] G. Higman, Finitely presented infinite simple groups. Notes on Pure Mathematics 8 (1974), Australian National University, Canberra.
- [6] J.P. Serre, *Trees*. Springer, 1980. Crossref.

SCHOOL OF MATHEMATICS & STATISTICS, UNIVERSITY OF ST ANDREWS, ST ANDREWS, KY16 9SS,
SCOTLAND.

Email address: `jmb42@st-andrews.ac.uk`

DEPARTMENT OF MATHEMATICS, CORNELL UNIVERSITY, ITHACA, NEW YORK 14853.

Email address: `jth263@cornell.edu`

DIPARTIMENTO DI MATEMATICA E APPLICAZIONI, UNIVERSITÀ DEGLI STUDI DI MILANO–BICOCCA,
MILAN 20125, ITALY.

Email address: `francesco.matucci@unimib.it`