

RECONSTRUCTING CURVES FROM THEIR HODGE CLASSES

MARIA GIOIA CIFANI, GIAN PIETRO PIROLA AND ENRICO SCHLESINGER

ABSTRACT. Let S be a smooth algebraic surface in $\mathbb{P}^3(\mathbb{C})$. A curve C in S has a cohomology class $\eta_C \in H^1(\Omega_S^1)$. Define $\alpha(C)$ to be the equivalence class of η_C in the quotient of $H^1(\Omega_S^1)$ modulo the subspace generated by the class η_H of a plane section of S . In the paper "Reconstructing subvarieties from their periods" the authors Movasati and Sertöz pose several interesting questions about the reconstruction of C from the annihilator $I_{\alpha(C)}$ of $\alpha(C)$ in the polynomial ring $R = H_*^0(\mathcal{O}_{\mathbb{P}^3})$. It contains the homogeneous ideal of C , but is much larger as $R/I_{\alpha(C)}$ is artinian. We give sharp numerical conditions that guarantee C is reconstructed by forms of low degree in $I_{\alpha(C)}$. We also show it is not always the case that the class $\alpha(C)$ is *perfect*, that is, that $I_{\alpha(C)}$ could be bigger than the sum of the Jacobian ideal of S and of the homogeneous ideals of curves D in S for which $I_{\alpha(D)} = I_{\alpha(C)}$.

1. INTRODUCTION

The Hodge conjecture, one of the most challenging and interesting open questions in algebraic geometry, can be regarded as a reconstruction problem. Even when the Hodge conjecture is known, as for curves on surfaces, there are a series of somewhat related problems that might shed a new light on some aspects of the cycle map.

A good example is given by [GH83, Theorem 4.b.26] where it is proven that, given a smooth surface $S \subset \mathbb{P}^3$ and an integral class γ in $H^1(\Omega_S^1)$ with the same numerical properties as the fundamental class of a curve $C \subset S$, then γ is itself the fundamental class of an effective divisor $D \subset S$ provided $\deg(S)$ is large relative to the self-intersection of γ and to $\deg(C)$.

In a similar vein, very interesting recent work by Movasati and Sertöz [MS19] concerns the reconstruction of subvarieties of \mathbb{P}^N from their periods. Our purpose is to give an answer, in the special case of curves lying on a smooth algebraic surface S in complex projective space, to two questions raised in [MS19] that we now illustrate. A curve C in S has a fundamental cohomology class $\eta_C \in H^1(\Omega_S^1)$. We denote by $\alpha(C)$ the equivalence class of η_C in the quotient of $H^1(\Omega_S^1)$ modulo the subspace generated by the class η_H of a plane section of S : the class $\alpha(C)$ depends on the embedding of S in \mathbb{P}^3 , and can be seen as a linear form on the primitive cohomology $H^1(\Omega_S^1)^{\perp_H}$. Following [MS19] we focus our analysis on the *annihilator* $I_{\alpha(C)}$ of $\alpha(C)$ in the polynomial ring $R = H_*^0(\mathcal{O}_{\mathbb{P}^3})$. Note - see Proposition 2.3 - that $I_{\alpha(C)} = I_{\alpha(D)}$ for two curves C and D in S if and only if $mC + nD + pH$ is linearly equivalent to zero for some choice of integers m, n and p with m and n non zero and relatively prime. Thus the annihilator $I_{\alpha(C)}$, which

contains the homogeneous ideal I_C of C and the Jacobian ideal J_S of S , in general it is much larger than $I_C + J_S$, as it contains the ideal I_D for any curve D for which there is a relation $mC + nD + pH \sim 0$ as above. Still, one can ask whether C can be reconstructed from $I_{\alpha(C)}$ when $\deg(S)$ is large with respect to the degree or other invariants of C , and Movasati and Sertöz in [MS19] investigate, in a more general context than ours, the following questions:

- (1) under which conditions $I_{\alpha(C)}$ reconstructs C , in the sense that forms of low degree in $I_{\alpha(C)}$ cut out the curve C scheme-theoretically? To be precise, we will say that C is *reconstructed at level m by $I_{\alpha(C)}$* if its homogeneous ideal I_C is generated over R by $I_{\alpha(C), \leq m}$ - that is, by forms of degree $\leq m$ in $I_{\alpha(C)}$.
- (2) as the example of complete intersection suggests, they define a class $\alpha \in H^1(\Omega_S^1)/\mathbb{C}\eta_H$ to be *perfect at level m* if there exist effective divisors D_1, \dots, D_q in S such that $I_{\alpha(D_i)} = I_\alpha$ for every $i = 1, \dots, q$ and

$$I_{\alpha, j} = \sum_{i=1}^q I_{D_i, j} + J_{S, j} \quad \text{for every } j \leq m$$

where J_S denotes the Jacobian ideal of S . The question is under which conditions the class $\alpha(C)$ is perfect at level m , and whether all classes $\alpha(C)$ are perfect at every level m .

In this paper we prove two theorems that give partial answers to these questions. Our first theorem extends known results on complete intersections [Dan17, MS19] to arithmetically Cohen-Macaulay curves (ACM curves for short). The tools we need for this are provided by a very nice paper by Ellingsrud and Peskine [EP93] which unfortunately seems to be little known. In [EP93] the authors were interested in the study of the Noether-Lefschetz locus, and the invariant $\alpha(C)$ plays a prominent role in their work because it vanishes if and only if the curve is a complete intersection of S and another surface. Their paper connects the class $\alpha(C)$ to the normal sequence arising from the inclusions $C \subset S \subset \mathbb{P}^3$ and gives an effective tool for computing its annihilator $I_{\alpha(C)}$ - see Lemma 2.5. To state our first theorem, given a curve C in \mathbb{P}^3 , we let $s(C)$ be the minimum degree of a surface containing C , and $e(C)$ the index of speciality of C , that is, the maximum n such that $\mathcal{O}_C(n)$ is special, that is, $h^1(\mathcal{O}_C(n)) > 0$.

Theorem 1.1. *Suppose C is an ACM curve on the smooth surface $S \subseteq \mathbb{P}^3(\mathbb{C})$. Let s denote the degree of S . Then*

- (1) *if $s \geq 2e(C) + 8 - s(C)$, the curve C is reconstructed at level $e(C) + 3$ by $I_{\alpha(C)}$;*
- (2) *the class $\alpha(C)$ is perfect at level m for every m .*

For example, let C be a twisted cubic curve: C is then ACM with invariants $s(C) = 2$ and $e(C) = -1$. By Theorem 1.1, if S is a quartic surface containing C , then C is cut out scheme-theoretically by the quadrics whose equations lie in $I_{\alpha(C, S)}$. This was suggested and verified for thousands of randomly chosen quartic surfaces containing C in [MS19, Sections 2.3 and 3.2].

Our second theorem provides a first example of a non-perfect algebraic class $\alpha(C)$, giving a negative answer to Question 2.12 in [MS19].

Theorem 1.2. *Let $C \subset \mathbb{P}^3$ be a smooth rational curve of degree 4 contained in a smooth surface S of degree $s = 4$. The class $\alpha(C)$ in S is not perfect at level 3.*

It would be very interesting to determine conditions for a class $\alpha(C)$ to be perfect, and we don't know whether ACM curves form the largest set of curves C whose classes $\alpha(C)$, in any smooth surface S containing C , are perfect.

Finally, we note that Movasati and Sertöz pose their questions of reconstruction and perfectness in a more general context, namely for classes in $H^n(\Omega_X^n)$ of varieties of dimension n in smooth hypersurfaces X in \mathbb{P}^{2n+1} . An interesting and challenging problem is trying to answer those question for every n , generalizing as far as it is possible the results of this paper to higher dimension.

The paper is structured as follows. In section 2 we collect some well known facts we need, and, for the benefit of the reader, we recall in some detail the constructions from [EP93] we will need in the sequel of the paper. In section 3 we prove Proposition 3.1 - a numerical criterion that guarantees, when the degree of S is large to respect to that of C , that the curve C is reconstructed at a certain level m by $I_{\alpha(C)}$. As an example we prove in Corollary 3.2 a reconstruction result for a general rational curve of degree d . In section 4 we prove Theorem 1.1, which is split in Theorems 4.1 and 4.8. In section 5 we prove Theorem 1.2.

The authors are members of GNSAGA of INdAM. The authors were partially supported by national MIUR funds, PRIN 2017 Moduli and Lie theory, and by MIUR: Dipartimenti di Eccellenza Program (2018-2022) - Dept. of Math. Univ. of Pavia.

2. PRELIMINARIES

We work in the projective space \mathbb{P}^3 over the field \mathbb{C} of complex numbers. Given a coherent sheaf \mathcal{F} on \mathbb{P}^3 and $i \in \mathbb{N}$, we define

$$H_*^i(\mathcal{F}) = \bigoplus_{n \in \mathbb{N}} H^i(\mathbb{P}^3, \mathcal{F}(n)).$$

These are graded module over the polynomial ring

$$R = H_*^0(\mathcal{O}_{\mathbb{P}^3}) \cong \mathbb{C}[x, y, z, w].$$

Given a subscheme X of \mathbb{P}^3 , we will denote by \mathcal{I}_X its sheaf of ideals, and by $I_X = H_*^0(\mathcal{I}_X)$ its saturated homogeneous ideal in R . We will write $I_{X,n}$ to denote its n^{th} graded piece $H^0(\mathbb{P}^3, \mathcal{I}_X(n))$.

If $M = \bigoplus_{n \in \mathbb{Z}} M_n$ a graded R -module, the *graded \mathbb{C} -dual* module M^* of M is defined by setting $(M^*)_m = \text{Hom}_{\mathbb{C}}(M_{-m}, \mathbb{C})$ with multiplication $R_n \times (M^*)_m \rightarrow (M^*)_{m+n}$ defined by

$$g\lambda(v) = \lambda(gv), \quad \forall g \in R_n, \lambda \in (M^*)_m, v \in M_{-m-n}.$$

By Serre's duality, if $X \subseteq \mathbb{P}^N$ is an equidimensional Cohen-Macaulay subscheme of dimension d , then for any locally free sheaf \mathcal{F} on X there is an isomorphism of graded R -modules

$$(H_*^i(X, \mathcal{F}))^* \cong H_*^{d-i}(X, \mathcal{F}^\vee \otimes \omega_X)$$

Let S be a *smooth algebraic surface* of degree s in \mathbb{P}^3 , and $C \subset S$ a *curve*, that is, an effective Cartier divisor in S . The curve C has a cohomology class $\eta_C \in H^1(S, \Omega_S^1)$. It can be defined as follows: the curve C defines a linear form λ_C on the set of $(1, 1)$ forms by integration; abstractly one can define this linear form as the image of the trace map $H^1(C, \Omega_C^1) \rightarrow \mathbb{C}$ under the transpose of the morphism $H^1(S, \Omega_S^1) \rightarrow H^1(C, \Omega_C^1)$ obtained by restricting differentials on S to C [Har77, Chapter III Ex. 7.4]. The cohomology class η_C is the image of λ_C under the Serre's duality isomorphism $H^1(\Omega_S^1)^* \cong H^1(\Omega_S^1)$.

If $\mathcal{O}_S(C)$ denotes the invertible sheaf on S corresponding to C , then $\eta_C = c(\mathcal{O}_S(C))$ where c denotes the first Chern class homomorphism

$$(1) \quad c : \text{Pic}(S) \rightarrow H^1(S, \Omega_S^1).$$

The perfect pairing $\langle \ , \ \rangle$ of Serre's duality is compatible with the intersection product of divisor classes [Har77, Chapter V Ex. 1.8] in the sense that for every pair of Cartier divisors D and E on S

$$\langle c(\mathcal{O}_S(D)), c(\mathcal{O}_S(E)) \rangle = D \cdot E.$$

Since S is a surface in \mathbb{P}^3 , numerically equivalent divisors on S are linearly equivalent, and the first Chern class map $\text{Pic}(S) \rightarrow H^1(S, \Omega_S^1)$ is injective.

The cotangent bundles of S and \mathbb{P}^3 are related by the exact sequence

$$(2) \quad 0 \rightarrow \mathcal{O}_S(-s) \rightarrow \Omega_{\mathbb{P}^3}^1 \otimes \mathcal{O}_S \rightarrow \Omega_S^1 \rightarrow 0.$$

It is well known (see e.g. [EGPSm85]) that $H^1(\Omega_{\mathbb{P}^3}^1 \otimes \mathcal{O}_S) \cong \mathbb{C}$ and that its image in $H^1(\Omega_S^1)$ is the class η_H of a plane section H of S . We look at a portion of the long cohomology sequence arising from (2)

$$(3) \quad H^1(\Omega_S^1) \xrightarrow{\delta} H^2(\mathcal{O}_S(-s)) \xrightarrow{\epsilon} H^2(\Omega_{\mathbb{P}^3}^1 \otimes \mathcal{O}_S)$$

Dualizing and using Serre's duality we get an exact sequence

$$(4) \quad H^0(\mathcal{T}_{\mathbb{P}^3}(s-4)) \xrightarrow{\epsilon^*} H^0(\mathcal{O}_S(2s-4)) \rightarrow \text{Im}(\delta)^* \rightarrow 0$$

We denote by J_S the Jacobian ideal of S , that is, the ideal of R generated by the partial derivatives of an equation of S . Then the above discussion is summarized by Griffith's theorem: the primitive first cohomology group of S is isomorphic to the $(2s-4)$ -graded piece of the Jacobian ring of S :

$$H^1(\Omega_S^1)^{\perp_H} \cong \text{Im}(\delta)^* \cong \frac{H^0(\mathcal{O}_{\mathbb{P}^3}(2s-4))}{J_{S, 2s-4}}$$

Definition 2.1. Given a curve C in S , we will denote by $\alpha(C) = \alpha(C, S, \mathbb{P}^3)$ the image of its cohomology class η_C under the map

$$H^1(\Omega_S^1) \xrightarrow{\delta} H^2(\mathcal{O}_S(-s)) \cong H^0(\mathcal{O}_S(2s-4))^*$$

Thus $\alpha(C)$ is a linear form on $H^0(\mathcal{O}_S(2s-4))$ that vanishes on $J_{S, 2s-4}$.

Given $\alpha \in H^0(\mathcal{O}_S(2s-4))^*$, we denote by I_α the annihilator of α in the polynomial ring R : it is the homogeneous ideal in R whose n^{th} graded piece is

$$I_{\alpha,n} = \{f \in R_n \mid \alpha(fg) = 0, \forall g \in H^0(\mathcal{O}_S(2s-4-n))\}.$$

Remark 2.2. When writing the paper, we decided to take all ideals in the polynomial ring $R = H^0(\mathcal{O}_{\mathbb{P}^3})$: thus J_S and I_α are for us ideals of R , and $J_S \subset I_{\alpha(C)}$. Our motivation is that we would like to compare I_α with the ideal of C as a curve in \mathbb{P}^3 . In [MS19] the author's denote by I_α the annihilator of α in the Jacobian ring and by \tilde{I}_α its preimage in R .

Let $T = R/I_S = H^0(\mathcal{O}_S)$. Then $\alpha \in (T_{2s-4})^*$, and the ideal I_α is determined by $\text{Ker}(\alpha) \subseteq T_{2s-4}$; conversely, one can recover $\text{Ker}(\alpha)$ as the image of $I_{\alpha,2s-4}$ via the quotient map $R_{2s-4} \rightarrow T_{2s-4}$. The perfect pairing

$$R_n/I_{\alpha,n} \times (R_{2s-4-n}/I_{\alpha,2s-4-n})^* \rightarrow \mathbb{C}$$

shows $A := R/I_\alpha = \bigoplus_{n=0}^{2s-4} A_n$ is an artinian Gorenstein ring of socle $2s-4$ [EP93, Prop 1.3].

In [EP93] the authors were interested in the study of the Noether-Lefschetz locus, and the invariant $\alpha(C)$ plays a prominent role in their work because it vanishes if and only if the curve is a complete intersection of S and another surface. More generally, a Lefschetz type theorem about the Picard group of S (see [SGA73, B78, Voi02],) implies the following fact:

Proposition 2.3. *Let C and D be effective divisors on a smooth surface $S \in \mathbb{P}^3$, and let H denote a plane section of S . Then $I_{\alpha(C)} = I_{\alpha(D)}$ if and only if there exist $m, n, p \in \mathbb{Z}$, $m, n \neq 0$ and relatively prime, such that $mC + nD + pH$ is linearly equivalent to zero.*

Proof. Suppose $mC + nD + pH$ is linearly equivalent to zero and m and n are nonzero. The cotangent complex (2) gives rise to an exact sequence in cohomology

$$(5) \quad H^1(\Omega_{\mathbb{P}^3}^1 \otimes \mathcal{O}_S) \cong \mathbb{C} \xrightarrow{\gamma} H^1(\Omega_S^1) \xrightarrow{\delta} H^2(\mathcal{O}_S(-s)) \simeq H^0(\mathcal{O}_S(2s-4))^*$$

and one knows that $\gamma(1) = \eta_H$, so that the kernel of δ is the \mathbb{C} -line spanned by η_H . From $mC + nD + pH \sim 0$ we then deduce $m\alpha(C) = -n\alpha(D)$. Since m and n are nonzero, the linear forms $\alpha(C)$ and $\alpha(D)$ have the same kernel, hence $I_{\alpha(C)} = I_{\alpha(D)}$.

In the other direction, suppose $I_{\alpha(C)} = I_{\alpha(D)}$, that is, $\alpha(C)$ and $\alpha(D)$ have the same kernel. Then $\alpha(C) = c\alpha(D)$ for a nonzero complex number c . Using (5) and the intersection pairing we deduce that are integers m, n, p , with m and n nonzero, such that $mC + nD + pH$ is linearly equivalent to zero. Finally, m and n can be taken relatively prime because $\text{Pic}(S)/\mathbb{Z}H$ has no torsion (see for example [B78, Theorem B]). In particular, when $D = 0$, one can take $m = 1$. \square

As noted in [EP93] and [MS19, Lemma 2.3], the ideal $I_{\alpha(C)}$ contains the ideal of C in S . This follows from the remark of [EP93] that $\alpha(C) \in H^0(\mathcal{O}_S(2s-4))^*$ is the pull-back of a linear form $\beta(C) \in H^0(\mathcal{O}_C(2s-4))^*$.

For the benefit of the reader and for later use, we give a proof of this fact. The linear form $\beta(C)$ arises from the normal bundles exact sequence:

$$(6) \quad 0 \rightarrow \mathcal{N}_{C/S} \cong \omega_C(4-s) \rightarrow \mathcal{N}_{C/\mathbb{P}^3} \rightarrow \mathcal{N}_{S/\mathbb{P}^3} \otimes \mathcal{O}_C \cong \mathcal{O}_C(s) \rightarrow 0.$$

Tensoring (6) with $\mathcal{O}_C(-s)$ and taking cohomology we obtain a map $H^0(\mathcal{O}_C) \rightarrow H^1(\omega_C(4-2s))$ and we let

$$\beta(C) \in H^0(\mathcal{O}_C(2s-4))^* \cong H^1(\omega_C(4-2s))$$

denote the image of $1 \in H^0(\mathcal{O}_C)$.

Proposition 2.4. [EP93, Construction 1.8] *The linear form $\alpha(C)$ is the pull-back of $\beta(C)$ to S , that is, $\alpha(C) = \rho^*(\beta(C))$ where ρ^* is the transpose of the natural map $\rho : H^0(\mathcal{O}_S(2s-4)) \rightarrow H^0(\mathcal{O}_C(2s-4))$.*

Proof. Observe that Ω_S^1 is a rank two vector bundle with determinant ω_S , hence the tangent bundle $\mathcal{T}_S = (\Omega_S^1)^\vee$ is isomorphic to $\Omega_S^1 \otimes \omega_S^{-1} = \Omega_S^1(4-s)$. The tangent complex of $S \subseteq \mathbb{P}^3$ and the normal bundle sequence (6) give rise to a commutative diagram

$$\begin{array}{ccccccc} 0 & \longrightarrow & \Omega_S^1 \cong \mathcal{T}_S(s-4) & \longrightarrow & \mathcal{T}_{\mathbb{P}^3} \otimes \mathcal{O}_S(s-4) & \longrightarrow & \mathcal{O}_S(2s-4) \longrightarrow 0 \\ & & \downarrow & & \downarrow & & \downarrow \\ 0 & \longrightarrow & \omega_C & \longrightarrow & \mathcal{N}_{C/\mathbb{P}^3}(s-4) & \longrightarrow & \mathcal{O}_C(2s-4) \longrightarrow 0 \end{array}$$

Taking cohomology and dualizing one sees that $\alpha(C)$ is the pull back of $\beta(C)$ to S . \square

The following Lemma in [EP93] gives an effective method to compute I_α in many cases.

Lemma 2.5. [EP93, Lemma 1.10] *Let $N(C)$ denote the image of the map $H_*^0 \mathcal{N}_{C/\mathbb{P}^3}(-s) \rightarrow H_*^0(\mathcal{O}_C)$ arising from the normal bundle sequence (6). Let $\pi : R = H_*^0(\mathcal{O}_{\mathbb{P}^3}) \rightarrow H_*^0(\mathcal{O}_C)$ be the natural map. Then, for every integer n ,*

$$\pi^{-1}(N(C)_n) \subseteq I_{\alpha(C),n}$$

with equality if π_{2s-4-n} is surjective.

Proof. The exact sequence

$$H_*^0 \mathcal{N}_{C/\mathbb{P}^3}(-s) \longrightarrow H_*^0(\mathcal{O}_C) \xrightarrow{1 \mapsto \beta} (H_*^0(\mathcal{O}_C(2s-4)))^*$$

shows $N(C) = \text{Ann}_{H_*^0(\mathcal{O}_C)}(\beta)$.

The map $\pi : R \rightarrow H_*^0(\mathcal{O}_C)$ factors through $\rho : H_*^0(\mathcal{O}_S) \rightarrow H_*^0(\mathcal{O}_C)$. To simplify notation, write $T = H_*^0(\mathcal{O}_S)$ and $e = 2s-4$. As α is an element of the T -module T^* , the ideal I_α , which by definition is the annihilator of α in R , is the inverse image of $\text{Ann}_T(\alpha)$ under the surjective map $R \rightarrow T$. Hence what we have to prove is that $\rho^{-1}(N(C)_n) \subseteq \text{Ann}_T(\alpha)_n$ for every integer n , with equality holding when ρ_{e-n} is surjective. Now

$$\text{Ann}_T(\alpha = \rho^*(\beta))_n = \{g \in T_n : g\rho^*(\beta)(v) = \beta(\rho(g)\rho(v)) = 0 \quad \forall v \in T_{e-n}\}$$

while the inverse image $\rho^{-1}(N(C)_n)$ of the n^{th} graded piece of the annihilator of $\beta(C)$ in $H_*^0(\mathcal{O}_C)$ is equal to

$$\{g \in T_n : (\rho(g)\beta)(w) = \beta(\rho(g)w) = 0 \quad \forall w \in H^0(\mathcal{O}_C(e - n))\}.$$

The thesis is now evident. \square

Corollary 2.6. *The annihilator $I_{\alpha(C)}$ of $\alpha(C)$ contains both the homogeneous ideal of C and the Jacobian ideal of the surface S .*

To exemplify the scope of this construction, we remark that it immediately yields the following well known corollary (originally due to Griffiths and Harris, see [EGPSm85] for more details).

Corollary 2.7. *Suppose S is a smooth surface in \mathbb{P}^3 and C is an effective divisor on S . Then C is a complete intersection of S and another surface if and only if the sequence 6 of normal bundles splits.*

Proof. If C is a complete intersection of S and another surface, it is clear that the sequence splits. Conversely, if the sequence splits, then $\beta(C) = 0$. Therefore $\alpha(C) = 0$, and the thesis follows from Proposition 2.3. \square

3. RECONSTRUCTION OF THE IDEAL

Motivated by [MS19], we want to compare I_C and $I_{\alpha(C)}$. The following proposition gives rather sharp sufficient conditions for the curve C to be reconstructed at level p by $I_{\alpha(C)}$.

Proposition 3.1. *Let S be a smooth surface of degree s in \mathbb{P}^3 , and let C be an effective Cartier divisor on S . Assume that the homogeneous ideal I_C is generated by its forms of degree $\leq p$ and that the following vanishing conditions are satisfied*

- (1) $h^1(\mathcal{I}_C(2s - 4 - p)) = 0$
- (2) $h^0(\mathcal{N}_{C/\mathbb{P}^3}(p - s)) = 0$

then $I_{\alpha(C),p} = I_{C,p}$, therefore C is reconstructed at level p by $I_{\alpha(C)}$.

Proof. Since $h^0(\mathcal{N}_{C/\mathbb{P}^3}(p - s)) = 0$, the annihilator of $\beta(C)$ in degree p vanishes. Since $\pi_{2s-4-p} : R_{2s-4-p} \rightarrow H^0(\mathcal{O}_C(2s - 4 - p))$ is surjective, by Lemma 2.5

$$I_{\alpha(C),p} = \pi_p^{-1}(\text{Ann}(\beta_C)_p) = I_{C,p}.$$

\square

We can now answer a question raised in [MS19, Section 2.3.1] about twisted cubics contained in quartic surfaces: if C is a twisted cubic contained in a smooth quartic surface $S \subset \mathbb{P}^3$, then C is cut out by quadrics in $I_{\alpha(C)}$. More generally:

Corollary 3.2. *Suppose $C \subset \mathbb{P}^3$ is a general rational curve of degree $d \geq 3$ and let n_0 be the round up of $\sqrt{6d - 2} - 3$, that is, the smallest positive integer n such that $\binom{n+3}{3} - nd - 1 \geq 0$. If C is contained in a smooth surface S of degree $s \geq n_0 + 3$, then C is reconstructed at level $n_0 + 1$ by $I_{\alpha(C,S)}$.*

Proof. By [HH85] a general rational curve is a curve of maximal rank, that is, $h^0(\mathcal{I}_C(n)) = 0$ for $n \leq n_0 - 1$ and $h^1(\mathcal{I}_C(n)) = 0$ for $n \geq n_0$. Hence C is $n_0 + 1$ regular in the sense of Castelnuovo-Mumford, and I_C is generated by its forms of degree $\leq n_0 + 1$. Furthermore, by [EVdV81] the normal bundle of the immersion $\mathbb{P}^1 \rightarrow C \subset \mathbb{P}^3$ is isomorphic to $\mathcal{O}_{\mathbb{P}^1}(2d-1) \oplus \mathcal{O}_{\mathbb{P}^1}(2d-1)$. Hence $h^0(\mathcal{N}_{C/\mathbb{P}^3}(-m)) = 0$ for every $m \leq -2$. Thus we can apply Proposition 3.1 with $p = n_0 + 1$. \square

Remark 3.3. If C is a smooth irreducible curve of degree d , then $h^1(\mathcal{I}_C(n)) = 0$ for every $n \geq d - 3 - e$ (see [GP87] and [Han00]), where $e := e(C) = \max\{n \mid h^1(\mathcal{O}(n)) > 0\}$ is the *index of speciality* of C .

Corollary 3.4. *Let S be a smooth surface of degree s in \mathbb{P}^3 , and let C be an effective Cartier divisor on S . Suppose \mathcal{I}_C is r -regular in the sense of Castelnuovo-Mumford. If $s \geq 2r + 1$, then C is reconstructed at level r by $I_{\alpha(C)}$.*

Proof. Since \mathcal{I}_C is r -regular, the ideal I_C is generated by its forms of degree $\leq r$ and $H^1(\mathcal{I}_C(n)) = 0$ for every $n \geq r - 1$. As $s \geq 2r + 1$ and $r \geq 1$, the first condition $h^1 \mathcal{I}_C(2s - 4 - r) = 0$ in Proposition 3.1 is satisfied for $p = r$.

We are left to check that $h^0 \mathcal{N}_{C/\mathbb{P}^3}(r - s) = 0$.

By [PS74, Prop 4.1], there are two surfaces S_1 and S_2 of degree r meeting properly in a complete intersection

$$X = S_1 \cap S_2 = C \cup D$$

so that C and D have no common component. Consider the exact sequence

$$0 \rightarrow \mathcal{I}_X \rightarrow \mathcal{I}_C \rightarrow \mathcal{I}_{C,X} \rightarrow 0.$$

Applying $\text{Hom}(-, \mathcal{O}_C)$ we get

$$0 \rightarrow \text{Hom}(\mathcal{I}_X, \mathcal{O}_C) \rightarrow \mathcal{N}_C \rightarrow \mathcal{N}_{X|C}$$

and $\text{Hom}(\mathcal{I}_X, \mathcal{O}_C) = 0$ since C and D have no common component. Therefore, there is an inclusion

$$\mathcal{N}_C \hookrightarrow (\mathcal{N}_X)_{|C} = \mathcal{O}_C(r) \oplus \mathcal{O}_C(r)$$

hence $h^0 \mathcal{N}_C(m) = 0$ for $m \leq -r - 1$. In particular $h^0 \mathcal{N}_C(r - s) = 0$ because $s \geq 2r + 1$. \square

4. ARITHMETICALLY COHEN-MACAULAY CURVES

In this section we explain how Example 1.15.3 in [EP93] extends the result about the perfection of complete intersections to the much larger class of arithmetically Cohen-Macaulay curves (from now on, ACM curves). Recall that a curve $C \subset \mathbb{P}^3$ is called ACM if its homogeneous ring $R_C = R/I_C$ is Cohen-Macaulay, or, equivalently, if C is locally Cohen-Macaulay of pure dimension 1 and $H_*^1(\mathcal{I}_C) = 0$. A smooth ACM curve is what classically was referred to as a *projectively normal curve*. We refer the reader to [HS11] for a detailed study of ACM curves on a surface in \mathbb{P}^3 .

If $C \subset \mathbb{P}^3$ is an ACM curve, then I_C has a free graded resolution of the form

$$(7) \quad 0 \rightarrow E = \bigoplus_{j=0}^r R(-b_j) \xrightarrow{\phi} F = \bigoplus_{i=0}^{r+1} R(-a_i) \rightarrow I_C \rightarrow 0$$

and I_C coincides with the ideal generated by the $r \times r$ minors of ϕ by the Hilbert-Burch theorem - cf. [MDP90, Proposition II.1.1 p. 37].

Applying the functor $\text{Hom}_R(\bullet, R/I_C)$ to (7) as in [El175, p. 428] one obtains a long exact sequence

$$(8) \quad 0 \rightarrow H_*^0(\mathcal{N}_C) \rightarrow \bigoplus_{i=0}^{r+1} R_C(a_i) \rightarrow \bigoplus_{j=0}^r R_C(b_j) \rightarrow H_*^0(\omega_C(4)) \rightarrow 0$$

The importance of this sequence for our purposes is that it allows to compute the Hilbert function $n \mapsto h^0(\mathcal{N}_{C, \mathbb{P}^3}(n))$ of $\mathcal{N}_{C, \mathbb{P}^3}$ as a function of the Hilbert function $n \mapsto h^0(\mathcal{O}_C(n))$ of C ; we can then compute the dimension of $\text{Ann}(\beta(C))_n$ and of $I_{\alpha(C), n}$ in terms solely of the Hilbert function of C and of the degree s of S . To justify our assertion, one needs to observe that to compute $h^0(\mathcal{N}_{C, \mathbb{P}^3}(n))$ out of (8) one does not need to know the numbers a_i 's and b_j 's, but only for each n the difference

$$\#\{i : a_i = n\} - \#\{j : b_j = n\}$$

which depends only on the Hilbert function of C .

As an application of this argument, we can give for ACM curves a sharp bound for the smallest integer n such that $I_{\alpha(C), n} = I_{C, n}$. For this we will not need the full Hilbert function of C , but just its index of speciality $e := e(C) = \max\{n \mid h^1(\mathcal{O}_C(n)) = h^2(\mathcal{I}_C(n)) > 0\}$ and the minimum degree $s(C)$ of a surface containing C : $s(C) = \min\{n \mid h^0(\mathcal{I}_C(n)) > 0\}$. For an ACM curve C , the ideal \mathcal{I}_C is $e+3$ -regular because $H_*^1(\mathcal{I}_C) = 0$. In particular, the ideal I_C is generated in degrees $\leq e+3$, and $s(C) \leq e+3$.

Theorem 4.1. *Let S be a smooth surface of degree s in \mathbb{P}^3 . Let $C \subset S$ be an ACM curve, let $s(C)$ be the minimum degree of a surface containing C and let $e(C)$ be the index of speciality of C .*

If $s \geq 2e(C) + 8 - s(C)$ then $I_{\alpha(C), (e+3)} = I_{C, (e+3)}$. Therefore C is reconstructed at level $e+3$ by $I_{\alpha(C)}$.

Proof. The statement follows from Proposition 3.1 with $p = e+3$ provided we can show that $h^0(\mathcal{N}_{C/\mathbb{P}^3}(e+3-s)) = 0$. For this we use the exact sequence (8), which shows that the maximum n for which $h^0 \mathcal{N}_{C, \mathbb{P}^3}(n) = 0$ is $n = s(C) - e(C) - 5$. \square

Remark 4.2. A twisted cubic curve C is ACM with invariants $s(C) = 2$ and $e(C) = -1$. Hence from Theorem 4.1 it follows once more that, if C is contained in a smooth quartic surface S , then C is cut out by quadrics in $I_{\alpha(C), S}$.

Remark 4.3. Theorem 4.1 improves for ACM curves the bound of Corollary 3.4 because, since $r = e+3$, then $2e+8-s(C) = 2r+2-s(C)$.

In [MS19, Sec 2.3], motivated by the case of complete intersections, formulate the notion of a *perfect class*:

Definition 4.4. Let S be a smooth surface of degree s in \mathbb{P}^3 . A class $\alpha \in H^1(\Omega_S)/\mathbb{C}\eta_H \subseteq H^0(\mathcal{O}_S(2s-4))^*$ is *perfect at level m* if there exist effective divisors D_1, \dots, D_q in S such that $I_{\alpha(D_i)} = I_\alpha$ for every $i = 1, \dots, q$ and

$$I_{\alpha,j} = \sum_{i=1}^q I_{D_i,j} + J_{S,j} \quad \text{for every } j \leq m.$$

We say the class is *perfect* if $I_\alpha = \sum_{i=1}^q I_{D_i} + J_S$. We make the convention that the zero class is perfect - geometrically, this amounts to consider the empty set as a (empty) curve, and is consistent with regarding the zero divisor as an effective divisor.

Example 4.5. If $C \subset S$ is the complete intersection of two surfaces meeting properly, then $\alpha(C)$ is perfect (see [MS19, Ex 2.11], [EP93, Ex 1.15.2], [Dan17, Prop. 2.14]). If one does not agree that the zero class is perfect, then one needs to add the condition that C is cut out by two surfaces of degrees $< s = \deg(S)$.

We now wish to generalize the previous example to the class of ACM curves showing that, if C is ACM, then the class $\alpha(C)$ is perfect. For this we need to recall more facts from [EP93]. Suppose the ACM curve C is contained in a smooth surface S of degree s and equation $f = 0$. Then the polynomial f can be written in the form

$$f = \sum_{i=1}^{r+1} g_i h_i$$

where the h_i 's are the images of the generators of the free module F in the resolution (7) of I_C . Since the h_i 's are the signed $r \times r$ minors of ϕ , then polynomial f is the determinant of the morphism $\psi : E \oplus R(-s) \rightarrow F$ obtained adding the column $[g_1, \dots, g_{r+1}]^T$ to the matrix of ϕ : in other words, ψ coincides with ϕ on E , and sends $1 \in R(-s)$ to $\sum_{i=1}^{p+1} g_i e_i$, where the e_i 's are the generators of F . We thus obtain a resolution of I_C/I_S :

$$(9) \quad 0 \rightarrow E \oplus R(-s) \xrightarrow{\psi} F \rightarrow I_C/I_S \rightarrow 0$$

Since S is smooth, the curve C is Cartier on S so that I_C/I_S can locally be generated by one element. It follows that the ideal $I_r(\psi)$ generated by the $r \times r$ minors of ψ is irrelevant, that is, its radical is the irrelevant maximal ideal (x, y, z, w) of the polynomial ring R .

Proposition 4.6. [EP93, Prop. 1.16] *Let $C \subset \mathbb{P}^3$ be an ACM curve contained in the smooth surface S . Suppose I_C has the resolution (7). Then*

- (1) *if ψ is as in exact sequence (9) the presentation of $I_{C,S}$, then*

$$I_{\alpha(C)} = I_r(\psi)$$

is the ideal generated by the $r \times r$ minors of ψ ;

- (2) the n^{th} -graded piece $\text{Ann}(\alpha(C))_n$ of the annihilator of $\alpha(C)$ in $H_*^0(\mathcal{O}_S)$ is the image of the natural map

$$\bigoplus_{m \in \mathbb{Z}} H^0 \mathcal{O}_S(C + (n + m)H) \otimes H^0 \mathcal{O}_S(-C - mH) \longrightarrow H^0(\mathcal{O}_S(n))$$

Remark 4.7. Note that $\text{Ann}(\alpha(C)) = I_{\alpha(C)}/I_S$. The equality $I_{\alpha(C)} = I_r(\psi)$ is a non trivial fact that is not given a full proof in [EP93]; a complete proof can be found in [KMR09, Proposition 4.3 and p. 382].

We can now prove that the class $\alpha(C)$ of an ACM curve in a smooth surface S is perfect:

Theorem 4.8. *Let $C \subset S$ be an ACM curve and let S be a smooth surface. Then the class $\alpha(C)$ of C in S is perfect.*

Proof. Fix an integer n . By Proposition 4.6 $\text{Ann}(\alpha(C))_n$ is the image of the natural map

$$\bigoplus_{m \in \mathbb{Z}} H^0 \mathcal{O}_S(C + (n + m)H) \otimes H^0 \mathcal{O}_S(-C - mH) \longrightarrow H^0(\mathcal{O}_S(n)).$$

Note the sum on the left hand side is finite, and consists of those m for which the linear systems $|C + (m + n)H|$ and $|-C - mH|$ are both non-empty. For such an m we pick a basis g_1, \dots, g_{r_m} of $H^0 \mathcal{O}_S(-C - mH)$ and corresponding effective divisors $D_k = (g_k)_0 \in |-C - mH|$. The image of

$$H^0 \mathcal{O}_S(C + (n + m_k)H) \otimes g_k$$

in $H^0(\mathcal{O}_S(n))$ is $H^0 \mathcal{I}_{D_k/S}(n)$. (if $C \sim tH$ is a complete intersection of S and another surface, taking $m_k = -t$ and $n = 0$ we get D_k the empty curve, and in this case $\alpha(C) = 0$ is perfect by our definition). Note that $\mathbb{Q}\alpha(D_k) = \mathbb{Q}\alpha(C)$ by Proposition 2.3. Now letting k and m vary we see that $\alpha(C)$ is perfect at level n , for every n . Since $\text{Ann}(\alpha(C))$ is finitely generated, we can let n vary up to the maximum degree of a generator of $\text{Ann}(\alpha(C))$, and recover the whole $\text{Ann}(\alpha(C))$ as the sum of finitely many $\mathcal{I}_{D_k/S}$ with $D_k \sim C + (n + m)H$ for some m and n . Therefore α_C in S is perfect. \square

5. EXAMPLE OF A NON PERFECT CLASS

Theorem 5.1. *Let $C \subset \mathbb{P}^3$ be a smooth rational curve of degree 4 contained in a smooth surface S of degree $s = 4$. The class $\alpha(C)$ in S is not perfect at level 3.*

Proof. A smooth rational quartic curve $C \subset \mathbb{P}^3$ is contained in a unique quadric surface Q , and Q is necessarily smooth (all curves on the quadric cone are arithmetically Cohen-Macaulay by [Har77, Chapter V Ex. 2.9]). We may assume C is a divisor of type $(3, 1)$ on Q . The ideal sheaf of C is 3-regular, hence I_C generated by quadrics and cubics.

Suppose C is contained in a smooth quartic surface S . Then $Q \cap S$ is the union of C and an effective divisor D_0 of type $(1, 3)$ on Q . Note that D_0 is a curve of degree 4 and arithmetic genus 0; as the divisor class of D_0 is different from that of C and C is irreducible, we conclude that C and D_0 have no common component.

The curves C and D_0 don't move in their linear system on the quartic surface S : for C this follows from $C^2 = -2$, and in any case for both D_0 and C_0 one might argue that

$$h^0(\mathcal{O}_S(D_0)) = h^0(\mathcal{O}_S(2H - C)) = h^0(\mathcal{I}_C(2)) = 1.$$

Having established the geometric set-up, we proceed to show that $I_{\alpha(C)}$ contains too many cubics for $\alpha(C)$ to be perfect at level 3. To compute the dimension of $I_{\alpha(C),3}$, we use the fact that $R/I_{\alpha(C)}$ is a Gorenstein ring with socle in degree $2s - 4 = 4$, hence

$$\dim I_{\alpha(C),3} = \dim I_{\alpha(C),1} + \dim R_3 - \dim R_1 = \dim I_{\alpha(C),1} + 16 \geq 16.$$

This estimate is good enough for us to prove the theorem, but let us show anyway that $\dim I_{\alpha(C),3} = 16$: as C is a divisor of type $(3, 1)$ on Q , $h^1(\mathcal{I}_C(3)) = 0$ hence by Lemma 2.5 $I_{\alpha(C),1}$ is the pull back to R_1 of $N(C)_1$, the image of $H^0(\mathcal{N}_{C/\mathbb{P}^3}(-3))$ in $H^0(\mathcal{O}_C(1))$; as the normal bundle of C pulls-back on \mathbb{P}^1 to $\mathcal{O}_{\mathbb{P}^1}(7) \oplus \mathcal{O}_{\mathbb{P}^1}(7)$ by [EVdV81, Proposition 6]), we conclude that $I_{\alpha(C),1} = 0$, hence $\dim I_{\alpha(C),3} = 16$. The same argument shows that $I_{\alpha(C),2} = I_{C,2}$ as well.

To check whether $I_{\alpha,3}$ is perfect, we need to determine curves D in S with $I_{\alpha(D)} = I_{\alpha(C)}$ and $h^0(\mathcal{I}_D(3)) \geq 1$ so that D can contribute to $I_{\alpha,3}$. Thus suppose D is such a curve. By Proposition 2.3, there exist $m, n, p \in \mathbb{Z}$, $m, n \neq 0$ and relatively prime, such that $pH + mC + nD$ is linearly equivalent to zero. By assumption $3H - D$ is effective; as C is not linearly equivalent to tH for any t , neither is D , hence $1 \leq \deg(D) = D \cdot H \leq 11$. Replacing D with $D' = 3H - D$ we can even assume $D \cdot H \leq 6$.

Now consider the matrix

$$M = \begin{bmatrix} H^2 & C \cdot H & H \cdot D \\ C \cdot H & C^2 & C \cdot D \\ H \cdot D & C \cdot D & D^2 \end{bmatrix} = \begin{bmatrix} 4 & 4 & x \\ 4 & -2 & y \\ x & y & z \end{bmatrix}$$

As $pH + mC + nD$ is linearly equivalent to zero, the vector $v = [p, m, n]^T$ is in the kernel of M . Set $x = H \cdot D$, $y = C \cdot D$ and $z = D^2$. Note that $z = D^2 = 2(p_a(D) - 1) = 2q$ is even.

The determinant of M must vanish, so

$$x^2 + 4xy - 2y^2 - 24q = 0$$

From this we deduce first that x and y must be even, and then that 4 divides x . As $1 \leq x \leq 6$, we must have $x = 4$. Thus D is a curve of degree 4, and either $D = C$ or C is not a component of D , hence $y = C \cdot D \geq 0$. Assume that $D \neq C$. Writing $y = 2t$ with $t \geq 0$, we obtain the equation

$$t^2 - 4t + 3q - 2 = 0$$

Looking at the discriminant of this quadratic equation in t we deduce $6 - 3q$ is a perfect square, so that $q = 2 - 3a^2$ for an integer $a \geq 0$. Then solving for t and imposing $t \geq 0$ we obtain $t = 2 + 3a$. So $H \cdot D = x = 4$, $C \cdot D = y = 4 + 6a$ and $D^2 = 4 - 6a^2$. Then solving the linear system $Mv = 0$ for $v = [p, m, n]^T$ we find $m = an$ and $p = -(a + 1)n$. Since m and n are relatively prime and non zero and $a \geq 0$, the only possibility is that $a = 1$. Then we can take $m = n = 1$ and conclude $C + D \sim 2H$, so that $C + D$ is the complete

intersection of the unique quadric Q containing C with S , and $D = D_0$ is the residual to C in the complete intersection $Q \cap S$.

We conclude that the only curves D in S that are contained in a cubic surface and satisfy $I_{\alpha(D)} = I_{\alpha(C)}$ are C , the residual D_0 to C in the complete intersection $Q \cap S$, and the effective divisors linearly equivalent to either $3H - C$ or $3H - D_0$. But observe that, if $D' \sim 3H - D_0 \sim C + H$ is effective, then

$$h^0 \mathcal{I}_{D'}(3) = h^0 \mathcal{I}_C(2) = 1.$$

Therefore there is a unique cubic containing D' , whose equation is contained in the ideal of D_0 . Similarly, if $D'' \sim 3H - C$ is effective, there is a unique cubic containing D'' , whose equation is contained in the ideal of C . Hence any cubic form that belongs to the ideal of a curve D on S satisfying $I_{\alpha(D)} = I_{\alpha(C)}$ is in the vector space spanned by $I_{C,3}$ and $I_{D_0,3}$.

To show $\alpha(C)$ is not perfect at level 3 it is now enough to show that cubics containing either C or D_0 plus the cubics in the Jacobian ideal J_S do not span $I_{\alpha(C),3}$.

To this end, note that cubic surfaces that contain both C and D_0 are in the ideal of the complete intersection of S and Q , and so form a vector space of dimension 4. By Grassmann's formula

$$\dim I_{C,3} + \dim I_{D_0,3} = 7 + 7 - 4 = 10$$

There are four independent cubics in the Jacobian ideal, so

$$\dim I_{C,3} + \dim I_{D_0,3} + \dim J_{S,3} \leq 14 < 16 = \dim I_{\alpha(C),3}$$

and this shows that $\alpha(C)$ in S is not perfect at level 3. \square

REFERENCES

- [Bř8] L. Bădescu. A remark on the Grothendieck-Lefschetz theorem about the Picard group. *Nagoya Math. J.*, 71:169–179, 1978.
- [Dan17] A. Dan. Noether-Lefschetz locus and a special case of the variational Hodge conjecture: using elementary techniques. In *Analytic and algebraic geometry*, pages 107–115. Hindustan Book Agency, New Delhi, 2017.
- [EGPSm85] G. Ellingsrud, L. Gruson, C. Peskine, and S. A. Strømme. On the normal bundle of curves on smooth projective surfaces. *Invent. Math.*, 80(1):181–184, 1985.
- [Ell75] G. Ellingsrud. Sur le schéma de Hilbert des variétés de codimension 2 dans \mathbf{P}^e à cône de Cohen-Macaulay. *Ann. Sci. École Norm. Sup. (4)*, 8(4):423–431, 1975.
- [EP93] G. Ellingsrud and C. Peskine. Anneau de Gorenstein associé à un fibré inversible sur une surface de l'espace et lieu de Noether-Lefschetz. In *Proceedings of the Indo-French Conference on Geometry (Bombay, 1989)*, pages 29–42. Hindustan Book Agency, Delhi, 1993.
- [EVdV81] D. Eisenbud and A. Van de Ven. On the normal bundles of smooth rational space curves. *Math. Ann.*, 256(4):453–463, 1981.
- [GH83] P. Griffiths and J. Harris. Infinitesimal variations of Hodge structure. II. An infinitesimal invariant of Hodge classes. *Compositio Math.*, 50(2-3):207–265, 1983.
- [GP87] L. Gruson and C. Peskine. Space curves: complete series and speciality. In *Space curves (Rocca di Papa, 1985)*, volume 1266 of *Lecture Notes in Math.*, pages 108–123. Springer, Berlin, 1987.
- [Han00] F. Han. Geometrically reducible line complexes. *Math. Ann.*, 316(4):819–824, 2000.

- [Har77] R. Hartshorne. *Algebraic geometry*. Springer-Verlag, New York-Heidelberg, 1977. Graduate Texts in Mathematics, No. 52.
- [HH85] R. Hartshorne and A. Hirschowitz. Courbes rationnelles et droites en position générale. *Ann. Inst. Fourier (Grenoble)*, 35(4):39–58, 1985.
- [HS11] R. Hartshorne and E. Schlesinger. Gonality of a general ACM curve in \mathbb{P}^3 . *Pacific J. Math.*, 251(2):269–313, 2011.
- [KMR09] J. O. Kleppe and R. M. Miró-Roig. Ideals generated by submaximal minors. *Algebra Number Theory*, 3(4):367–392, 2009.
- [MDP90] M. Martin-Deschamps and D. Perrin. *Sur la classification des courbes gauches*. Astérisque 184-185. Société Mathématique de France, 1990.
- [MS19] H. Movasati and E. C. Sertöz. On reconstructing subvarieties from their periods, 2019.
- [PS74] C. Peskine and L. Szpiro. Liaison des variétés algébriques. I. *Invent. Math.*, 26:271–302, 1974.
- [SGA73] *Groupes de monodromie en géométrie algébrique. II*. Lecture Notes in Mathematics, Vol. 340. Springer-Verlag, Berlin-New York, 1973. Séminaire de Géométrie Algébrique du Bois-Marie 1967–1969 (SGA 7 II), Dirigé par P. Deligne et N. Katz.
- [Voi02] C. Voisin. *Hodge theory and complex algebraic geometry. I*, volume 76 of *Cambridge Studies in Advanced Mathematics*. Cambridge University Press, Cambridge, 2002. Translated from the French original by Leila Schneps.

(M.G.C.) DIPARTIMENTO DI MATEMATICA E FISICA, UNIVERSITÀ DI ROMA 3, LARGO SAN LEONARDO MURIALDO, 1 – 00146 ROMA
Email address: `mariagioia.cifani@uniroma3.it`

(G.P.P.) DEPARTMENT OF MATHEMATICS 'F. CASORATI', UNIVERSITY OF PAVIA, VIA FERRATA 5, 27100 PAVIA, ITALY
Email address: `gianpietro.pirola@unipv.it`

(E.S.) DIPARTIMENTO DI MATEMATICA 'F. BRIOSCHI', POLITECNICO DI MILANO, PIAZZA LEONARDO DA VINCI 32, 20133 MILANO, ITALY
Email address: `enrico.schlesinger@polimi.it`