

THE BAR INVOLUTION FOR QUANTUM SYMMETRIC PAIRS – HIDDEN IN PLAIN SIGHT

STEFAN KOLB

To Jasper Stokman on his 50th birthday in gratitude and friendship

ABSTRACT. We show that all quantum symmetric pair coideal subalgebras $\mathcal{B}_{\mathbf{c}}$ of Kac-Moody type have a bar involution for a suitable choice of parameters \mathbf{c} . The proof relies on a generalized notion of quasi K -matrix. The proof does not involve an explicit presentation of $\mathcal{B}_{\mathbf{c}}$ in terms of generators and relations.

1. INTRODUCTION

Let \mathfrak{g} be a symmetrizable Kac-Moody algebra and $\theta : \mathfrak{g} \rightarrow \mathfrak{g}$ an involutive Lie algebra automorphism of the second kind. The theory of quantum symmetric pairs, as developed by G. Letzter in the finite setting [Let99], studies certain coideal subalgebras $\mathcal{B}_{\mathbf{c}}$ of the quantized enveloping algebra $U = U_q(\mathfrak{g})$. The coideal subalgebras $\mathcal{B}_{\mathbf{c}}$ depend on a family of parameters \mathbf{c} . They are quantum group analogs of $U(\mathfrak{g}^{\theta})$ where $\mathfrak{g}^{\theta} = \{x \in \mathfrak{g} \mid \theta(x) = x\}$.

In March 2012, C. Stroppel asked the author if there exists a bar involution for $\mathcal{B}_{\mathbf{c}}$. She conjectured that $\mathcal{B}_{\mathbf{c}}$ should allow an algebra automorphism $\overline{}^B : \mathcal{B}_{\mathbf{c}} \rightarrow \mathcal{B}_{\mathbf{c}}$ which is the identity on the Letzter generators $B_i \in \mathcal{B}_{\mathbf{c}}$ and maps q to q^{-1} . Unfortunately, back in 2012, this author did not sufficiently appreciate the question.

This changed drastically with the appearance of the preprint versions of [BW18] and [ES18] in October 2013. Both these papers contained a bar involution for an explicit example of $\mathcal{B}_{\mathbf{c}}$ of type AIII. For this example, H. Bao and W. Wang used the bar involution to construct an element \mathfrak{X} (denoted by Υ in [BW18]), now called the quasi K -matrix for $\mathcal{B}_{\mathbf{c}}$, which intertwines the Lusztig bar involution $\overline{}$ for U and the new bar involution $\overline{}^B$ for $\mathcal{B}_{\mathbf{c}}$ as follows

$$(1.1) \quad \overline{b}^B \mathfrak{X} = \mathfrak{X} \overline{b} \quad \text{for all } b \in \mathcal{B}_{\mathbf{c}}.$$

Moreover, Bao and Wang used the quasi K -matrix to construct a natural family of $\mathcal{B}_{\mathbf{c}}$ -module automorphisms \mathcal{T} on the category of finite-dimensional U -modules. At this point it became clear that the family \mathcal{T} was a hot candidate for a universal K -matrix which the author had been chasing previously in joint work with J. Stokman [KS09]. Subsequently, this was worked out jointly with M. Balagović [BK19].

The line of argument, first proposed in [BW18] and then performed in [BK15], [BK19] in a general Kac-Moody setting, was as follows. First the existence of a bar involution on $\mathcal{B}_{\mathbf{c}}$ was established via an explicit presentation of $\mathcal{B}_{\mathbf{c}}$ in terms of

2010 *Mathematics Subject Classification.* Primary: 17B37, Secondary: 53C35, 16T05, 17B67.
Key words and phrases. Quantum symmetric pairs, bar involution.

generators and relations. Then it was proved that the defining relation (1.1) of the quasi K -matrix has an essentially unique solution.

Writing down $\mathcal{B}_{\mathbf{c}}$ in terms of generators and relations is a difficult problem in the general Kac-Moody setting. The quest for the bar involution for $\mathcal{B}_{\mathbf{c}}$ was often cited as an important motivation to find such a presentation of $\mathcal{B}_{\mathbf{c}}$, see for example the introductions of the papers [CLW20], [dC19]. However, G. Lusztig asked the author in January 2019 if there is a construction of the bar involution for $\mathcal{B}_{\mathbf{c}}$ which does not rely on a presentation of $\mathcal{B}_{\mathbf{c}}$ in terms of generators and relations. Lusztig pointed out that the construction of the bar involution for the positive part U^+ in [Lus94, 1.2.12] does not involve explicit knowledge of the quantum Serre relations for U .

The quasi K -matrix is the quantum symmetric pair analog of the quasi R -matrix for U . The quasi R -matrix for U has a description which does not involve Lusztig's bar involution, see [Lus94, Theorem 4.1.2 (b)]. Hence it is natural to ask for a bar involution free description of the quasi K -matrix. Such a construction was given in the quasi-split case (and in a more general setting of Nichols algebras of diagonal type) in [KY20]. For suitable parameters \mathbf{c} there exists an element \mathfrak{X} satisfying a relation similar to (1.1), and the map

$$b \mapsto \mathfrak{X} \bar{b} \mathfrak{X}^{-1} \quad \text{for all } b \in \mathcal{B}_{\mathbf{c}}$$

is a bar involution on $\mathcal{B}_{\mathbf{c}}$. The element \mathfrak{X} in [KY20] is given by $\mathfrak{X} = (\varepsilon \otimes \text{id})(\Theta^\theta)$ where ε is the counit and the element Θ^θ is defined by [KY20, (6.1)], see also [KY20, Remark 6.3].

Along a slightly different line, A. Appel and B. Vlaar observed in [AV20] that the construction of the quasi K -matrix in [BK19] can be reformulated without the bar involution $\bar{}$. Recall that $\mathcal{B}_{\mathbf{c}}$ is defined for a (generalized) Satake diagram (X, τ) , where X is a subset of the nodes I of the Dynkin diagram of \mathfrak{g} , and $\tau : I \rightarrow I$ is an involutive diagram automorphism. The pair (X, τ) needs to satisfy certain compatibility conditions, see Section 2.2. The Letzter generators of $\mathcal{B}_{\mathbf{c}}$ are given by $B_i = F_i$ for $i \in X$ and $B_i = F_i - c_i T_{w_X}(E_{\tau(i)}) K_i^{-1}$ for $i \in I \setminus X$, see Section 3.1. With this notation, condition (3.9) is equivalent to the condition

$$B_i \mathfrak{X} = \mathfrak{X} \overline{B'_i} \quad \text{for all } i \in I$$

where B_i are the Letzter generators corresponding to the parameter family \mathbf{c} , and B'_i are Letzter generators corresponding to a different parameter family \mathbf{c}' . One then obtains an algebra isomorphism $\Psi : \mathcal{B}_{\mathbf{c}'} \rightarrow \mathcal{B}_{\mathbf{c}}$ defined by $\Psi(b) = \mathfrak{X} \bar{b} \mathfrak{X}^{-1}$. This isomorphism satisfies $\Psi(B'_i) = B_i$ and $\Psi(q) = q^{-1}$.

In [AV20] Appel and Vlaar are mostly concerned with the bar involution free formulation of the quasi K -matrix \mathfrak{X} . The main point of the present note, which we still believe has novelty to it, is the observation that the above argument provides a proof of the existence of the bar involution for all symmetrizable Kac-Moody algebras and all (generalized) Satake diagrams if the parameters are chosen such that $\mathbf{c}' = \mathbf{c}$, see Corollary 4.2. Although Appel and Vlaar get very close to this observation, they stop just short of the formulation of the existence of the bar involution in [AV20, Remark 7.2]. In particular, with the new perspective on the quasi K -matrix, the study of the quantum Serre relations for $\mathcal{B}_{\mathbf{c}}$ mentioned in [AV20, Remark 7.2] is not necessary anymore.

This note is organised as follows. In Section 2 we fix notation for quantized enveloping algebras and recall the notion of a generalized Satake diagram as formulated in [RV20] which generalizes the notion of an admissible pair in [Kol14, Definition 2.3]. We promote the view that future papers on quantum symmetric pairs should be written in the setting of generalized Satake diagrams. In Section 3 we recall the definition of $\mathcal{B}_{\mathbf{c}}$ and reprove the existence of the quasi K -matrix \mathfrak{X} . While this result is contained in [AV20, Theorem 7.4], we feel that it is worthwhile to reproduce it in the precise notations and setting of [BK19]. Section 4 contains the existence of the isomorphism Ψ . As a consequence we obtain the main message of this note, namely the general, relation-free existence proof for the bar involution on $\mathcal{B}_{\mathbf{c}}$, see Corollary 4.2

2. PRELIMINARIES

2.1. Quantum groups. Let \mathfrak{g} be a symmetrizable Kac-Moody algebra with generalized Cartan matrix $(a_{ij})_{i,j \in I}$ where I is a finite set and Cartan subalgebra $\mathfrak{h} \subset \mathfrak{g}$. Let $\{d_i \mid i \in I\}$ be a set of relatively prime positive integers such that the matrix $(d_i a_{ij})$ is symmetric. Let $\Pi = \{\alpha_i \mid i \in I\}$ be the set of simple roots for \mathfrak{g} , let Φ be the root system, and let $Q = \mathbb{Z}\Pi$ be the root lattice. For $\beta = \sum_{i \in I} n_i \alpha_i \in Q$ we write $\text{ht}(\beta) = \sum_{i \in I} n_i$. Consider the symmetric bilinear form $(\cdot, \cdot) : Q \times Q \rightarrow \mathbb{Z}$ defined by $(\alpha_i, \alpha_j) = d_i a_{ij}$ for all $i, j \in I$. We denote $Q^+ = \mathbb{N}_0 \Pi$ where $\mathbb{N}_0 = \{0, 1, 2, \dots\}$. Let W be the Weyl group of \mathfrak{g} which is generated by the simple reflections s_i for $i \in I$.

Throughout this paper let k be a field of characteristic zero and let $\mathbb{K} = k(q)$ be the field of rational functions in a variable q . We consider the quantized enveloping algebra $U = U_q(\mathfrak{g})$ as the \mathbb{K} -algebra with generators $E_i, F_i, K_i^{\pm 1}$ for $i \in I$ subject to the relations given in [Lus94, 3.1.1]. For $\beta = \sum_{i \in I} n_i \alpha_i \in Q$ we write $K_\beta = \prod_{i \in I} K_i^{n_i}$. Let U^+ and U^- denote the subalgebras of U generated by $\{E_i \mid i \in I\}$ and $\{F_i \mid i \in I\}$, respectively. For any $\mu \in Q^+$ we write $U_\mu^+ = \text{span}_{\mathbb{K}}\{E_{i_1} \dots E_{i_\ell} \mid \sum_{j=1}^{\ell} \alpha_{i_j} = \mu\}$ where $\text{span}_{\mathbb{K}}$ denotes the \mathbb{K} -linear span. Moreover, define $U_{-\mu}^- = \omega(U_\mu^+)$ where $\omega : U \rightarrow U$ denotes the algebra automorphism given in [Lus94, 3.1.3]. For any $i \in I$ let $T_i : U \rightarrow U$ denote the algebra automorphism denoted by $T''_{i,1}$ in [Lus94, 37.1]. By [Lus94, 39.4.3] the automorphisms T_i satisfy braid relations. Hence, for $w \in W$ with reduced expression $w = s_{i_1} \dots s_{i_\ell}$ there is a well-defined automorphism $T_w = T_{i_1} \dots T_{i_\ell}$ of U . For any $i \in I$ let $r_i, {}_i r : U^+ \rightarrow U^+$ denote the Lusztig-Kashiwara skew-derivations which are defined uniquely by the relation

$$(2.1) \quad xF_i - F_i x = \frac{1}{q_i - q_i^{-1}} (r_i(x)K_i - K_i^{-1}{}_i r(x)) \quad \text{for all } x \in U^+$$

where $q_i = q^{d_i}$, see [Lus94, 3.1.6]. The maps r_i and ${}_i r$ satisfy the relations $r_i(E_j) = {}_i r(E_j) = \delta_{i,j}$ and the skew-derivation properties

$$(2.2) \quad r_i(xy) = q^{(\alpha_i, \nu)} r_i(x)y + x r_i(y), \quad {}_i r(xy) = {}_i r(x)y + q^{(\alpha_i, \mu)} x {}_i r(y)$$

for all $x \in U_\mu^+, y \in U_\nu^+$.

Recall that the bar involution for U is the k -algebra automorphism $\bar{} : U \rightarrow U$, $x \mapsto \bar{x}$ defined by

$$\overline{E_i} = E_i, \quad \overline{F_i} = F_i, \quad \overline{K_i} = K_i^{-1}, \quad \overline{q} = q^{-1}$$

for all $i \in I$.

To formulate the properties of the quasi K -matrix we need to work in an over-algebra \mathcal{U} of U . Let \mathcal{O}_{int} denote the category of integrable U -modules in category \mathcal{O} and let $\mathcal{F}or : \mathcal{O}_{\text{int}} \rightarrow \mathcal{Vect}$ be the forgetful functor into the category of \mathbb{K} -vector spaces. We define $\mathcal{U} = \text{End}(\mathcal{F}or)$ to be the set of natural transformations from $\mathcal{F}or$ to itself. By construction, \mathcal{U} is an algebra which contains U as a subalgebra. See [BK19, Section 3] for precise definitions and further details. Consider the elements of the product $\widehat{U^+} = \prod_{\mu \in Q^+} U_{\mu}^+$ as additive infinite sums of elements in U^+ . Then $\widehat{U^+}$ is a subalgebra of \mathcal{U} , see [BK19, Example 3.2].

2.2. Generalized Satake diagrams. In [Kol14] quantum symmetric pairs were defined for involutive automorphisms of \mathfrak{g} of the second kind. Such involutions are classified in terms of Satake diagrams (X, τ) where $X \subset I$ is a subset of finite type and $\tau : I \rightarrow I$ is an involutive diagram automorphism satisfying the compatibility conditions given in [Kol14, Definition 2.3]. Associated to the finite type subset X is a finite type root system $\Phi_X \subset \Phi$ with parabolic Weyl group $W_X \subset W$. Let $w_X \in W_X$ be the longest element and let $\rho_X^{\vee} \in \mathfrak{h}$ be the half sum of positive coroots of Φ_X . To be a Satake diagram, the pair (X, τ) has to satisfy the two conditions

- (1) $\tau(\alpha_j) = -w_X(\alpha_j)$ for all $j \in X$;
- (2) If $i \in I \setminus X$ and $\tau(i) = i$ then $\alpha_i(\rho_X^{\vee}) \in \mathbb{Z}$.

Condition (1) implies that the map $\Theta = -w_X \circ \tau : Q \rightarrow Q$ is an involutive automorphism of the root lattice. It was observed by V. Regelskis and B. Vlaar that condition (2) can be replaced by the weaker condition

- (2') If $\tau(i) = i$ and $a_{ji} = -1$ then $\Theta(\alpha_i) \neq -\alpha_i - \alpha_j$.

Following [RV20], we call pairs (X, τ) as above, satisfying conditions (1), (2') generalized Satake diagrams. As explained in [RV20, Section 4] the construction of quantum symmetric pairs and the structure theory developed in [Kol14] remains valid for generalized Satake diagrams.

The diagram automorphism τ lifts to an Hopf-algebra automorphism of U . As in [Lus94, 3.1.3] let $\sigma : U \rightarrow U$ denote the \mathbb{K} -algebra antiautomorphism defined by

$$\sigma(E_i) = E_i, \quad \sigma(F_i) = F_i, \quad \sigma(K_i) = K_i^{-1} \quad \text{for all } i \in I.$$

The following proposition was conjectured in [BK15, Conjecture 2.7] and was proved in [BK15, Proposition 2.5] up to a sign. The sign was confirmed by H. Bao and W. Wang in [BW21] by a subtle argument involving canonical bases for U . The arguments in [BK15], [BW21] do not involve condition (2) above, and hence also hold in the setting of generalized Satake diagrams, see also [AV20, Lemma 7.14].

Proposition 2.1. *Let (X, τ) be a generalized Satake diagram for the symmetrizable Kac-Moody algebra \mathfrak{g} . Then the relation*

$$\sigma \circ \tau(r_i(T_{w_X}(E_i))) = r_i(T_{w_X}(E_i))$$

holds in U for all $i \in I \setminus X$.

3. THE QUASI K -MATRIX, REVISITED

3.1. Quantum symmetric pairs. Let (X, τ) be an generalized Satake diagram. Set $\mathcal{M}_X = \mathbb{K}\langle E_j, F_j, K_j^{\pm 1} \mid j \in X \rangle$ and set $U_{\Theta}^0 = \mathbb{K}\langle K_{\beta} \mid \beta \in Q, \Theta(\beta) = \beta \rangle$. Define

a set of parameters

$$\mathcal{C} = \{\mathbf{c} = (c_i)_{i \in I \setminus X} \in (\mathbb{K}^\times)^{I \setminus X} \mid c_i = c_{\tau(i)} \text{ if } (\alpha_i, \Theta(\alpha_i)) = 0\}.$$

For any $\mathbf{c} = (c_i)_{i \in I \setminus X} \in \mathcal{C}$ we consider the subalgebra $\mathcal{B}_{\mathbf{c}}$ of $U_q(\mathfrak{g})$ generated by \mathcal{M}_X , $U_{\mathfrak{g}}^0$ and the elements

$$(3.1) \quad B_i = F_i - c_i T_{w_X}(E_{\tau(i)}) K_i^{-1} \quad \text{for all } i \in I \setminus X.$$

To unify notation, we set $c_i = 0$ for $i \in X$ and extend (3.1) by writing $B_i = F_i$ if $i \in X$. Following [Let99], [Kol14] we call $\mathcal{B}_{\mathbf{c}}$ the quantum symmetric pair coideal subalgebra of $U_q(\mathfrak{g})$ corresponding to the generalized Satake diagram (X, τ)

Remark 3.1. The parameters c_i were denoted by $c_i s(\tau(i))$ in [BK19]. The additional parameters $s(\tau(i))$ appeared in the construction of involutive automorphisms of \mathfrak{g} of the second kind corresponding to the Satake diagram (X, τ) . For the construction of quantum symmetric pairs, however, it is advantageous to suppress the parameters $s(\tau(i))$ in the notation.

Remark 3.2. In [Kol14], quantum symmetric pairs depend on a second family of parameters \mathbf{s} in a certain subset $\mathcal{S} \subset \mathbb{K}^{I \setminus X}$. The corresponding coideal subalgebras are then denoted by $\mathcal{B}_{\mathbf{c}, \mathbf{s}}$. By [Kol14, Theorem 7.1] the algebra $\mathcal{B}_{\mathbf{c}}$ is isomorphic to $\mathcal{B}_{\mathbf{c}, \mathbf{s}}$ as an algebra for all \mathbf{s} . As we only aim to establish the existence of a bar involution for $\mathcal{B}_{\mathbf{c}, \mathbf{s}}$, it suffices to consider the case where $\mathbf{s} = (0, 0, \dots, 0)$. Moreover, it was noted in [DK19, 3.5], that for the construction of the quasi K -matrix there is no loss of generality in the restriction to the case $\mathbf{s} = (0, 0, \dots, 0)$.

3.2. The quasi K -matrix \mathfrak{X} . In [AV20, Section 7] A. Appel and V. Vlaar gave a reformulation of the results of [BK19, Section 6] which does not rely on the existence of a bar involution for $\mathcal{B}_{\mathbf{c}}$. As notations and conventions in [AV20] somewhat differ from those of the present note, we feel it is beneficial to recall this reformulation in the present setting. We begin with a reformulation of [BK19, Proposition 6.1]. Let $2\rho_X$ be the sum of positive roots in Φ_X .

Proposition 3.3. *Let $\mathfrak{X} = \sum_{\mu \in Q^+} \mathfrak{X}_{\mu} \in \widehat{U}^+$ with $\mathfrak{X}_{\mu} \in U_{\mu}^+$. The following are equivalent:*

(1) *For all $i \in I$ the relation*

$$B_i \mathfrak{X} = \mathfrak{X} (F_i - (-1)^{2\alpha_i(\rho_X^\vee)} q^{-(\alpha_i, \Theta(\alpha_i) - 2\rho_X)} c_{\tau(i)} \overline{T_{w_X}(E_{\tau(i)})} K_i)$$

holds in \mathcal{U} .

(2) *The element \mathfrak{X} satisfies the relations*

$$(3.2) \quad r_i(\mathfrak{X}_{\mu}) = (q_i - q_i^{-1}) (-1)^{2\alpha_i(\rho_X^\vee)} q^{-(\alpha_i, \Theta(\alpha_i) - 2\rho_X)} c_{\tau(i)} \mathfrak{X}_{\mu + \Theta(\alpha_i) - \alpha_i} \overline{T_{w_X}(E_{\tau(i)})},$$

$$(3.3) \quad {}_i r(\mathfrak{X}_{\mu}) = (q_i - q_i^{-1}) q^{-(\Theta(\alpha_i), \alpha_i)} c_i T_{w_X}(E_{\tau(i)}) \mathfrak{X}_{\mu + \Theta(\alpha_i) - \alpha_i}$$

for all $i \in I$.

If the equivalent conditions (1) and (2) hold, then the following also hold:

(3) *For all $x \in U_{\mathfrak{g}}^0 \mathcal{M}_X$ one has $x\mathfrak{X} = \mathfrak{X}x$.*

(4) *For all $\mu \in Q^+$ such that $\mathfrak{X}_{\mu} \neq 0$ one has $\Theta(\mu) = -\mu$.*

Proof. The equivalence of (1) and (2) follows from the property (2.1) of the skew derivations r_i and ${}_i r$. Property (4) follows by induction over the height of μ and in turn implies property (3) just as in the proof of [BK19, Proposition 6.1]. \square

Next we need [BK19, Proposition 6.3]. This proposition is a general statement about solving recursive relations of the type (3.2), (3.3) simultaneously, and hence holds in our setting. Let $\langle \cdot, \cdot \rangle : U^- \otimes U^+ \rightarrow \mathbb{K}$ be the nondegenerate pairing considered in [BK19, Section 2.3].

Proposition 3.4. ([BK19, Proposition 6.3]) *Let $\mu \in Q^+$ with $\text{ht}(\mu) \geq 2$ and fix elements $A_i, {}_iA \in U_{\mu-\alpha_i}^+$ for all $i \in I$. Then the following are equivalent:*

(1) *There exists an element $\underline{X} \in U_\mu^+$ such that*

$$(3.4) \quad r_i(\underline{X}) = A_i \quad \text{and} \quad {}_i r(\underline{X}) = {}_i A \quad \text{for all } i \in I.$$

(2) *The elements $A_i, {}_iA$ have the following two properties:*

(a) *For all $i, j \in I$ one has*

$$r_i({}_j A) = {}_j r(A_i).$$

(b) *For all $i \neq j \in I$ one has*

$$(3.5) \quad \frac{-1}{q_i - q_i^{-1}} \sum_{s=1}^{1-a_{ij}} \begin{bmatrix} 1 - a_{ij} \\ s \end{bmatrix}_{q_i} (-1)^s \langle F_i^{1-a_{ij}-s} F_j F_i^{s-1}, A_j \rangle \\ - \frac{1}{q_j - q_j^{-1}} \langle F_i^{1-a_{ij}}, A_j \rangle = 0.$$

Moreover, if the system of equation (3.4) has a solution \underline{X} then this solution is uniquely determined.

We now translate [BK19, Section 6.4] into our setting. This is the crucial step where we also need Proposition 2.1. We hence give all the details. Fix $\mu \in Q^+$ and assume that a collection $(\mathfrak{X}_{\mu'})_{\mu' < \mu}$ with $\mathfrak{X}_{\mu'} \in U_{\mu'}^+$ and $\mathfrak{X}_0 = 1$ has already been constructed, and that this collection satisfies the relations (3.2),(3.3) for all $\mu' < \mu$ and all $i \in I$. Define

$$(3.6) \quad A_i = (q_i - q_i^{-1})(-1)^{2\alpha_i(\rho_X^\vee)} q^{-(\alpha_i, \Theta(\alpha_i) - 2\rho_X)} c_{\tau(i)} \mathfrak{X}_{\mu + \Theta(\alpha_i) - \alpha_i} \overline{T_{w_X}(E_{\tau(i)})}$$

$$(3.7) \quad {}_i A = (q_i - q_i^{-1}) q^{-(\Theta(\alpha_i), \alpha_i)} c_i T_{w_X}(E_{\tau(i)}) \mathfrak{X}_{\mu + \Theta(\alpha_i) - \alpha_i}.$$

We have the following analog of [BK19, Lemma 6.7].

Lemma 3.5. *The relation $r_i({}_j A) = {}_j r(A_i)$ holds for all $i, j \in I$.*

Proof. Using the skew-derivation properties (2.2) and the assumptions on ${}_i A, A_i$ we calculate

$$\begin{aligned} r_i({}_j A) &= (q_j - q_j^{-1}) q^{-(\Theta(\alpha_j), \alpha_j)} c_j T_{w_X}(E_{\tau(j)}) r_i(\mathfrak{X}_{\mu + \Theta(\alpha_j) - \alpha_j}) \\ &\quad + (q_j - q_j^{-1}) q^{-(\Theta(\alpha_j), \alpha_j)} c_j q^{(\alpha_i, \mu + \Theta(\alpha_j) - \alpha_j)} r_i(T_{w_X}(E_{\tau(j)})) \mathfrak{X}_{\mu + \Theta(\alpha_j) - \alpha_j} \\ &= (q_j - q_j^{-1})(q_i - q_i^{-1})(-1)^{2\alpha_i(\rho_X^\vee)} q^{-(\Theta(\alpha_j), \alpha_j)} q^{-(\alpha_i, \Theta(\alpha_i) - 2\rho_X)} c_j c_{\tau(i)} \\ &\quad \cdot T_{w_X}(E_{\tau(j)}) \mathfrak{X}_{\mu + \Theta(\alpha_j) - \alpha_j + \Theta(\alpha_i) - \alpha_i} \overline{T_{w_X}(E_{\tau(i)})} \\ &\quad + (q_j - q_j^{-1}) q^{-(\Theta(\alpha_j), \alpha_j)} c_j q^{(\alpha_i, \mu + \Theta(\alpha_j) - \alpha_j)} r_i(T_{w_X}(E_{\tau(j)})) \mathfrak{X}_{\mu + \Theta(\alpha_j) - \alpha_j} \end{aligned}$$

and similarly

$$\begin{aligned}
{}_j r(A_i) &= (q_i - q_i^{-1})(-1)^{2\alpha_i(\rho_X^\vee)} q^{-(\alpha_i, \Theta(\alpha_i) - 2\rho_X)} c_{\tau(i)} {}_j r(\mathfrak{X}_{\mu + \Theta(\alpha_i) - \alpha_i} \overline{T_{w_X}(E_{\tau(i)})}) \\
&\quad + (q_i - q_i^{-1})(-1)^{2\alpha_i(\rho_X^\vee)} q^{(\alpha_j, \mu + \Theta(\alpha_i) - \alpha_i) - (\alpha_i, \Theta(\alpha_i) - 2\rho_X)} \times \\
&\quad \quad \quad \times c_{\tau(i)} \mathfrak{X}_{\mu + \Theta(\alpha_i) - \alpha_i} {}_j r(\overline{T_{w_X}(E_{\tau(i)})}) \\
&= (q_i - q_i^{-1})(q_j - q_j^{-1})(-1)^{2\alpha_i(\rho_X^\vee)} q^{-(\alpha_i, \Theta(\alpha_i) - 2\rho_X)} q^{-(\Theta(\alpha_j), \alpha_j)} c_{\tau(i)} c_j \times \\
&\quad \quad \quad \times T_{w_X}(E_{\tau(j)}) \mathfrak{X}_{\mu + \Theta(\alpha_i) - \alpha_i + \Theta(\alpha_j) - \alpha_j} \overline{T_{w_X}(E_{\tau(i)})} \\
&\quad + (q_i - q_i^{-1})(-1)^{2\alpha_i(\rho_X^\vee)} q^{(\alpha_j, \mu + \Theta(\alpha_i) - \alpha_i) - (\alpha_i, \Theta(\alpha_i) - 2\rho_X)} \times \\
&\quad \quad \quad \times c_{\tau(i)} \mathfrak{X}_{\mu + \Theta(\alpha_i) - \alpha_i} {}_j r(\overline{T_{w_X}(E_{\tau(i)})}).
\end{aligned}$$

Comparing the above two expressions we see that the relation $r_i({}_j A) = {}_j r(A_i)$ is equivalent to the relation

$$\begin{aligned}
(3.8) \quad (q_j - q_j^{-1}) q^{-(\Theta(\alpha_j), \alpha_j)} c_j q^{(\alpha_i, \mu + \Theta(\alpha_j) - \alpha_j)} r_i(T_{w_X}(E_{\tau(j)})) \mathfrak{X}_{\mu + \Theta(\alpha_j) - \alpha_j} \\
= (q_i - q_i^{-1})(-1)^{2\alpha_i(\rho_X^\vee)} q^{(\alpha_j, \mu + \Theta(\alpha_i) - \alpha_i) - (\alpha_i, \Theta(\alpha_i) - 2\rho_X)} \times \\
\quad \quad \quad \times c_{\tau(i)} \mathfrak{X}_{\mu + \Theta(\alpha_i) - \alpha_i} {}_j r(\overline{T_{w_X}(E_{\tau(i)})}).
\end{aligned}$$

By [Lus94, 1.2.14] we have ${}_j r(\overline{T_{w_X}(E_{\tau(i)})}) = q^{(\alpha_j, w_X(\alpha_{\tau(i)}) - \alpha_j)} r_j(\overline{T_{w_X}(E_{\tau(i)})})$. Hence both sides of (3.8) vanish unless $\tau(i) = j \in I \setminus X$, and in this case (3.8) can be rewritten as

$$\begin{aligned}
q^{(\alpha_j, \Theta(\alpha_i) - \Theta(\alpha_j) - \alpha_i) + (\alpha_i, \mu)} r_i(T_{w_X}(E_i)) \mathfrak{X}_{\mu + \Theta(\alpha_j) - \alpha_j} \\
= (-1)^{2\alpha_i(\rho_X^\vee)} q^{(\alpha_j, \mu - \alpha_i - \alpha_j) - (\alpha_i, \Theta(\alpha_i) - 2\rho_X)} \mathfrak{X}_{\mu + \Theta(\alpha_j) - \alpha_j} \overline{r_j(T_{w_X}(E_j))}
\end{aligned}$$

From now on we assume $\tau(i) = j$ and hence $\Theta(\alpha_i) - \alpha_i = \Theta(\alpha_j) - \alpha_j$. Hence we may assume that $\mathfrak{X}_{\mu + \Theta(\alpha_j) - \alpha_j} \neq 0$ and in this case the above equation is equivalent to the equation

$$q^{(\alpha_i, \mu)} r_i(T_{w_X}(E_i)) = (-1)^{2\alpha_i(\rho_X^\vee)} q^{(\alpha_{\tau(i)}, \mu - \alpha_i) - (\alpha_i, \Theta(\alpha_i) - 2\rho_X)} \overline{r_{\tau(i)}(T_{w_X}(E_{\tau(i)}))}.$$

By Proposition 3.3.(4) we have $(\alpha_i - \alpha_{\tau(i)}, \mu) = 0$ and hence the above equation can be rewritten as

$$\overline{r_i(T_{w_X}(E_i))} = (-1)^{2\alpha_i(\rho_X^\vee)} q^{(\alpha_i, \alpha_i - w_X(\alpha_i) - 2\rho_X)} r_{\tau(i)}(T_{w_X}(E_{\tau(i)})).$$

By [BK15, Lemma 2.9], which also holds for generalized Satake diagrams, and by Proposition 2.1 the above equation does indeed hold. \square

Next we obtain an analog of [BK19, Lemma 6.8]. We include the proof which is simplified along the lines of [BK19, Remark 6.9].

Lemma 3.6. *For all $i \neq j \in I$ the elements A_i, A_j given by (3.6) satisfy the relation (3.5).*

Proof. As $A_i \in U_{\mu - \alpha_i}^+$ we may assume that $\mu = (1 - a_{ij})\alpha_i + \alpha_j$ and $\Theta(\mu) = -\mu$ as otherwise both terms of (3.5) vanish. By [BK19, Lemma 6.4] it suffices to consider the following two cases:

Case 1: $\Theta(\alpha_i) = -\alpha_j$ and $a_{ij} = 0$. In this case $\mu = \alpha_i + \alpha_j$ and $(\alpha_i, \alpha_k) = (\alpha_j, \alpha_k) = 0$ for all $k \in X$, and hence

$$A_i = (q_i - q_i^{-1}) c_j E_j, \quad A_j = (q_i - q_i^{-1}) c_i E_i.$$

Hence the left hand side of (3.5) vanishes.

Case 2: $\Theta(\alpha_i) = -\alpha_i$ and $\Theta(\alpha_j) = -\alpha_j$. In this case induction on μ' for $\mu' < \mu$ shows that $\mathfrak{X}_{\mu'} \neq 0$ implies that $\mu' \in \text{span}_{\mathbb{N}_0} \{2\alpha_i, 2\alpha_j\}$, see [BK19, Lemma 6.5]. Hence for $\mu = (1 - a_{ij})\alpha_i + \alpha_j$ we have $\mathfrak{X}_{\mu+\Theta(\alpha_i)-\alpha_i} = \mathfrak{X}_{\mu-2\alpha_i} = 0$ and similarly $\mathfrak{X}_{\mu+\Theta(\alpha_j)-\alpha_j} = 0$. Hence $A_i = A_j = 0$ which implies (3.5) also in this case. \square

With the above preparations the following theorem is proved just as in [BK19, Theorem 6.10] using Proposition 3.4.

Theorem 3.7. *There exists a uniquely determined element $\mathfrak{X} = \sum_{\mu \in Q^+} \mathfrak{X}_\mu \in \widehat{U^+}$ with $\mathfrak{X}_0 = 1$ and $\mathfrak{X}_\mu \in U_\mu^+$ such that the equality*

$$(3.9) \quad B_i \mathfrak{X} = \mathfrak{X} (F_i - (-1)^{2\alpha_i(\rho_X^\vee)} q^{-(\alpha_i, \Theta(\alpha_i) - 2\rho_X)} c_{\tau(i)} \overline{T_{w_X}(E_{\tau(i)})} K_i)$$

holds in \mathcal{U} for all $i \in I$. Moreover, the element \mathfrak{X} commutes with all elements of $U_{\Theta}^0 \mathcal{M}_X$.

Remark 3.8. We call the element $\mathfrak{X} \in \mathcal{U}$ from the above theorem the quasi K -matrix corresponding to the quantum symmetric pair coideal subalgebra $\mathcal{B}_{\mathbf{c}}$. The quasi K -matrix is invertible in $\widehat{U^+} = \prod_{\mu \in Q^+} U_\mu^+$ because $\mathfrak{X}_0 = 1 \neq 0$.

4. THE BAR INVOLUTION FOR QUANTUM SYMMETRIC PAIRS, REVISITED

Recall that $\mathbb{K} = k(q)$. Following [AV20, Section 7.3] for any $\mathbf{c} = (c_i)_{i \in I \setminus X} \in \mathbb{K}^{I \setminus X}$ we define $\mathbf{c}' = (c'_i)_{i \in I \setminus X} \in \mathbb{K}^{I \setminus X}$ by

$$c'_i = (-1)^{2\alpha_i(\rho_X^\vee)} q^{(\alpha_i, \Theta(\alpha_i) - 2\rho_X)} \overline{c_{\tau(i)}}.$$

Observe that $\mathbf{c} \in \mathcal{C}$ if and only if $\mathbf{c}' \in \mathcal{C}$. For $\mathbf{c} \in \mathcal{C}$ let $\mathcal{B}_{\mathbf{c}'}$ denote the quantum symmetric pair coideal subalgebra corresponding to the parameters \mathbf{c}' . We denote the generators (3.1) for $\mathcal{B}_{\mathbf{c}'}$ by B'_i to distinguish them from the corresponding generators B_i of $\mathcal{B}_{\mathbf{c}}$. We immediately obtain the following consequence of Theorem 3.7.

Corollary 4.1. *For any $\mathbf{c} \in \mathcal{C}$ there exists a k -algebra isomorphism $\Psi : \mathcal{B}_{\mathbf{c}'} \rightarrow \mathcal{B}_{\mathbf{c}}$ such that*

$$\Psi|_{\mathcal{M}_X U_{\mathfrak{g}}^0} = \overline{\quad}|_{\mathcal{M}_X U_{\mathfrak{g}}^0} \quad \text{and} \quad \Psi(B'_i) = B_i \quad \text{for all } i \in I \setminus X.$$

In particular, we have $\Psi(q) = q^{-1}$ and $\Psi(K_\beta) = K_{-\beta}$ for all $\beta \in Q^\Theta$.

Proof. With the above notation, Equation (3.9) can be rewritten as

$$(4.1) \quad B_i \mathfrak{X} = \mathfrak{X} \overline{B'_i} \quad \text{for all } i \in I \setminus X$$

Hence there is a well-defined k -algebra homomorphism $\Psi : \mathcal{B}_{\mathbf{c}'} \rightarrow \mathcal{B}_{\mathbf{c}}$ such that $\Psi(b) = \mathfrak{X} b \mathfrak{X}^{-1}$ for all $b \in \mathcal{B}_{\mathbf{c}'}$. Equation (4.1) implies that $\Psi(B'_i) = B_i$ for all $i \in I \setminus X$. Property (3) in Proposition 3.3 implies that $\Psi|_{\mathcal{M}_X U_{\mathfrak{g}}^0} = \overline{\quad}|_{\mathcal{M}_X U_{\mathfrak{g}}^0}$. \square

In the special case that $\mathbf{c} = \mathbf{c}'$ the above corollary provides the desired bar involution for $\mathcal{B}_{\mathbf{c}}$.

Corollary 4.2. *Assume that the parameters $\mathbf{c} = (c_i)_{i \in I \setminus X} \in \mathbb{K}^{I \setminus X}$ satisfy the relation*

$$(4.2) \quad \overline{c_i} = (-1)^{2\alpha_i(\rho_X^\vee)} q^{-(\alpha_i, \Theta(\alpha_i) - 2\rho_X)} c_{\tau(i)}$$

for all $i \in I \setminus X$. Then there exists a k -algebra automorphism

$$\overline{}^B : \mathcal{B}_{\mathbf{c}} \rightarrow \mathcal{B}_{\mathbf{c}}, \quad b \mapsto \overline{b}^B$$

such that $\overline{}^B|_{U_{\mathfrak{g}}^0 \mathcal{M}_X} = \overline{}|_{U_{\mathfrak{g}}^0 \mathcal{M}_X}$ and $\overline{B_i}^B = B_i$ for all $i \in I \setminus X$. In particular, the automorphism $\overline{}^B$ of $\mathcal{B}_{\mathbf{c}}$ satisfies the relation $\overline{q}^B = q^{-1}$.

REFERENCES

- [AV20] A. Appel and B. Vlaar, *Universal k -matrix for quantum Kac-Moody algebras*, Preprint, [arXiv:2007.09218v2](https://arxiv.org/abs/2007.09218v2) (2020), 67 pp.
- [BK15] M. Balagović and S. Kolb, *The bar involution for quantum symmetric pairs*, Represent. Theory **19** (2015), 186–210.
- [BK19] ———, *Universal K -matrix for quantum symmetric pairs*, J. reine angew. Math. **89** (2019), 299–353.
- [BW18] H. Bao and W. Wang, *A new approach to Kazhdan-Lusztig theory of type B via quantum symmetric pairs*, Asterisque **402** (2018), vii+134pp.
- [BW21] ———, *Canonical bases arising from quantum symmetric pairs of Kac-Moody type*, Compositio Math. in press, [arXiv:1811.09848v2](https://arxiv.org/abs/1811.09848v2) (2021), 31pp.
- [CLW20] X. Chen, M. Lu, and W. Wang, *A Serre presentation for the iq quantum groups*, Transform. Groups (online first), <https://doi.org/10.1007/s00031-020-09581-5> (2020), 31pp.
- [dC19] H. de Clercq, *Generators and relations for quantum symmetric pair coideals of Kac-Moody type*, preprint, [arXiv:1912.05368](https://arxiv.org/abs/1912.05368) (2019), 51pp.
- [DK19] L. Dobson and S. Kolb, *Factorisation of quasi K -matrices for quantum symmetric pairs*, Selecta Math. (N.S.) **25** (2019), no. 4, Paper No. 63, 55pp.
- [ES18] M. Ehrig and C. Stroppel, *Nazarov-Wenzl algebras, coideal subalgebras and categorified skew Howe duality*, Adv. Math. **331** (2018), 58–142.
- [Kol14] S. Kolb, *Quantum symmetric Kac-Moody pairs*, Adv. Math. **267** (2014), 395–469.
- [KS09] S. Kolb and J. Stokman, *Reflection equation algebras, coideal subalgebras, and their centres*, Selecta Math. (N.S.) **15** (2009), 621–664.
- [KY20] S. Kolb and M. Yakimov, *Symmetric pairs for Nichols algebras of diagonal type via star products*, Adv. Math. **365** (2020), art. 107042, 69pp.
- [Let99] G. Letzter, *Symmetric pairs for quantized enveloping algebras*, J. Algebra **220** (1999), 729–767.
- [Lus94] G. Lusztig, *Introduction to quantum groups*, Birkhäuser, Boston, 1994.
- [RV20] V. Regelskis and B. Vlaar, *Quasitriangular coideal subalgebras of $U_q(\mathfrak{g})$ in terms of generalized Satake diagrams*, Bull. Lond. Math. Soc. **52** (2020), no. 4, 693–715.

SCHOOL OF MATHEMATICS, STATISTICS AND PHYSICS, NEWCASTLE UNIVERSITY, NEWCASTLE UPON TYNE NE1 7RU, UNITED KINGDOM

Email address: stefan.kolb@newcastle.ac.uk