

A comprehensive study of vector leptoquark with $U(1)_{B_3-L_2}$ on the B -meson and Muon $g-2$ anomalies

Kayoung Ban,^{1,*} Yongsoo Jho,^{2,†} Youngjoon Kwon,^{1,‡} Seong Chan Park,^{1,§} Seokhee Park,^{3,¶} and Po-Yan Tseng^{4,**}

¹*Department of Physics and IPAP, Yonsei University, Seoul 03722, Republic of Korea*

²*Weizmann Institute, Department of Particle Physics and Astrophysics, Rehovot, Israel 7610001*

³*High Energy Accelerator Research Organization (KEK), Tsukuba 305-0801, Japan*

⁴*Department of Physics, National Tsing Hua University, Hsinchu 300, Taiwan*

Recently reported anomalies in various B meson decays and also in the anomalous magnetic moment of muon $(g-2)_\mu$ motivate us to consider a particular extension of the standard model incorporating new interactions in lepton and quark sectors simultaneously. Our minimal choice would be leptoquark. In particular, we take vector leptoquark (U_1) and comprehensively study all related observables including $(g-2)_\mu$, $R_{K^{(*)}}$, $R_{D^{(*)}}$, $B \rightarrow (K)\ell\ell'$ where $\ell\ell'$ are various combinations of μ and τ , and also lepton flavor violation in the τ decays. We find that a hybrid scenario with additional $U(1)_{B_3-L_2}$ gauge boson provides a common explanation of all these anomalies.

I. INTRODUCTION

Physics is a data-driven science, and we are keen to modify the standard model (SM) when experimental results deviate from the theoretical predictions. Over the past several years, the B -physics experiments BaBar, Belle, and LHCb have reported several anomalous results in the $b \rightarrow s\ell\ell$ and $b \rightarrow c\ell\nu$ processes, which are not properly explained within the SM thus call for new physics. In particular, the lepton flavor universality (LFU), which is one of the approximate symmetries in the SM, seems to be broken beyond the expected range according to the observables of R_D , R_{D^*} , R_K , and R_{K^*} , which measured the ratios of different lepton flavors

$$R_{D^{(*)}} \equiv \frac{\mathcal{B}(B \rightarrow D^{(*)}\tau\nu)}{\mathcal{B}(B \rightarrow D^{(*)}\ell\nu)}, R_{K^{(*)}} \equiv \frac{\mathcal{B}(B \rightarrow K^{(*)}\mu^+\mu^-)}{\mathcal{B}(B \rightarrow K^{(*)}e^+e^-)}. \quad (1)$$

The precise measurement of those quantities would test the basic structure of the SM since LFU is only violated by the lepton masses in the SM.

The world averaged experimental values [1] based on measurements from BaBar [2], Belle [3–5], and LHCb [6, 7] are

$$R_D = 0.340 \pm 0.027 \pm 0.013 \quad R_{D^*} = 0.295 \pm 0.011 \pm 0.008, \quad (2)$$

and the combined discrepancy to SM prediction is at the 3.1σ level [1, 8].

The most precise measurement to date of the R_K has been performed by LHCb [9]

$$R_K = 0.846_{-0.041}^{+0.044}, \quad q^2 \subseteq [1.1, 6.0] \text{ GeV}^2, \quad (3)$$

which has 3.1σ deviation from the SM expectation. For the R_{K^*} , LHCb Run-1 provides [10]

$$R_{K^*} = \begin{cases} 0.66_{-0.07}^{+0.11} \pm 0.03, & q^2 \subseteq [0.045, 1.1] \text{ GeV}^2, \\ 0.69_{-0.07}^{+0.11} \pm 0.05, & q^2 \subseteq [1.1, 6.0] \text{ GeV}^2. \end{cases} \quad (4)$$

* ban94gy@yonsei.ac.kr

† yongsoo.jho@weizmann.ac.il

‡ yjkwon63@yonsei.ac.kr

§ sc.park@yonsei.ac.kr

¶ seokhee.park@kek.jp

** pytseng@phys.nthu.edu.tw

Combining both q^2 bins, it has 2.5σ tension with the SM. On the other side, the R_{K^*} and R_K measurements from Belle [11, 12]

$$R_{K^*} = \begin{cases} 0.90_{-0.21}^{+0.27} \pm 0.10, & q^2 \subseteq [0.1, 8.0] \text{ GeV}^2, \\ 1.18_{-0.32}^{+0.52} \pm 0.10, & q^2 \subseteq [15, 19] \text{ GeV}^2, \end{cases}$$

$$R_K = \begin{cases} 0.98_{-0.23}^{+0.27} \pm 0.06, & q^2 \subseteq [1.0, 6.0] \text{ GeV}^2, \\ 1.11_{-0.26}^{+0.29} \pm 0.07, & 14.18 \text{ GeV} < q^2, \end{cases} \quad (5)$$

are still compatible with SM predictions within their large uncertainties. In the near future, Belle II is expected to significantly improve the uncertainties [13].

The other long-standing problem is the anomalous magnetic moment of muon. Recently, the Muon ($g-2$) experiment at Fermilab reported the value, $a_\mu^{\text{FNAL}} = (116592040 \pm 54) \times 10^{-11}$ [14], or, the discrepancy from the SM

$$\Delta a_\mu^{\text{FNAL}} = a_\mu^{\text{FNAL}} - a_\mu^{\text{SM}} = (230 \pm 69) \times 10^{-11}, \quad (6)$$

which is a 3.3σ deviation. Since the value is compatible with the earlier value from BNL [15, 16], the significance is now strengthened to 4.2σ level:

$$\Delta a_\mu^{\text{BNL+FNAL}} = a_\mu^{\text{exp}} - a_\mu^{\text{SM}} = (251 \pm 59) \times 10^{-11}. \quad (7)$$

Even though it is not completely settled down from the lattice calculations [17], certainly it is worth considering new physics as its solution.

Finding a common origin of the B -meson and $(g-2)_\mu$ anomalies is non-trivial, but it is appealing from the theoretical point of view.¹ In early attempts [16, 30–40], the $U_1 = (3, 1)_{2/3}$ singlet vector Leptoquark, in general couples to both left-handed (LH) and right-handed (RH) SM fermions, as single-mediator accounts for all the low-energy data. Its simultaneous explanations of the $R_{K^{(*)}}$ and $R_{D^{(*)}}$ anomalies only require the LH couplings between second and third-generation quarks and leptons. However, the LH couplings cannot produce large enough muon magnetic moment for $(g-2)_\mu$ anomaly. In this work, we further extend non-zero RH couplings of U_1 , such that it substantially enhances the contribution to $(g-2)_\mu$. In addition, we will also consider $U(1)_{B_3-L_2}$ gauge boson (X) [41–43] to improve our fit to the experimental data. We explored the plausible parameter space and search the common solution for both B -meson and $(g-2)_\mu$ anomalies.²

II. MODEL

In this section, we set our theoretical model to explain B -anomalies and $(g-2)_\mu$.

A. Vector Leptoquark $U_1 = (3, 1)_{2/3}$

Leptoquark is a natural candidate of new physics linking quark sector and lepton sector [16]. In particular, we focus on the $U_1 = (3, 1)_{2/3}$ weak singlet vector leptoquark because it could provide simultaneous explanations for $R_{K^{(*)}}$ and $R_{D^{(*)}}$ anomalies with its coupling with LH fermions [30–33]. However, the general Lagrangian includes the U_1 couplings to both LH and RH fermion under the SM gauge symmetry. Including the most relevant interactions with the LH couplings to the 2nd and 3rd generations of leptons and quarks and also the RH couplings, we consider the model Lagrangian:

$$\mathcal{L} \supset U_{1\mu} \sum_{i,j=1,2,3} \left[x_L^{ij} (\bar{d}_L^i \gamma^\mu e_L^j) + (V_{\text{CKM}} x_L U_{\text{PMNS}}^*)_{ij} (\bar{u}_L^i \gamma_\mu \nu_L^j) + x_R^{ij} (\bar{d}_R^i \gamma^\mu e_R^j) \right] + h.c. \quad (8)$$

¹ See [18–29] for some of the earlier attempts to account $(g-2)$ and also various anomalies and possible experimental probes.

² When we are finishing our paper, similar idea has been considered in [44]. Unfortunately, however, they have missed some relevant constraints from low-energy experiments. In particular, the experimental $B_s \rightarrow \mu\mu$ data conflicts with their preferred parameter region.

We adopt the real parts of CKM and PMNS matrices which is conveniently written as [16]

$$V_{\text{CKM}} x_L U_{\text{PMNS}}^* = \begin{pmatrix} 0.974 & 0.225 & 0.001 \\ -0.224 & 0.974 & 0.042 \\ 0.009 & -0.041 & 0.999 \end{pmatrix} \begin{pmatrix} x_L^{11} & x_L^{12} & x_L^{13} \\ x_L^{21} & x_L^{22} & x_L^{23} \\ x_L^{31} & x_L^{32} & x_L^{33} \end{pmatrix} \begin{pmatrix} 0.821 & 0.551 & -0.150 \\ -0.307 & 0.600 & 0.739 \\ 0.481 & -0.580 & 0.657 \end{pmatrix},$$

above expression omits the imaginary parts in V_{CKM} , but we adopt the full CKM parameterization from Ref [45]. in our numerical computation. The couplings x_L^{ij} to the first generation leptons and quarks are strongly constrained from $\mu - e$ conversion on nuclei, and atomic parity violation on $\mathcal{B}(K \rightarrow \pi \nu \bar{\nu})$, therefore we simply set them zero.

The most relevant Wilson coefficients of the effective Lagrangian in $R_{K^{(*)}}$, $R_{D^{(*)}}$, and $\mathcal{B}(B_s \rightarrow \mu\mu)$ are $C_9^{\mu\mu}$ ($= -C_{10}^{\mu\mu}$) [30], and they are induced from the LH couplings

$$C_9^{\mu\mu} = -C_{10}^{\mu\mu} = -\frac{\pi v^2}{V_{tb} V_{ts}^* \alpha_{\text{EM}}} \frac{x_L^{22} (x_L^{32})^*}{m_{U_1}^2}, \quad (9)$$

where $v = 246$ GeV is the vacuum expectation value of the Higgs, and α_{EM} is the fine structure constant. From the fit to R_K , R_{K^*} and $\mathcal{B}(B_s \rightarrow \mu\mu)$ data, we find the parameter window for couplings and the mass of U_1

$$C_9^{\mu\mu} = -C_{10}^{\mu\mu} \subseteq [-0.85, -0.50] \Rightarrow -\frac{x_L^{22} (x_L^{32})^*}{m_{U_1}^2} \subseteq [0.83, 1.41] \times 10^{-3} \text{ TeV}^{-2}. \quad (10)$$

The interactions from Eq.(8) also give rise to the effective coefficient [30]

$$g_{V_L} = \frac{v^2}{2m_{U_1}^2} (x_L^{b\ell})^* \left[x_L^{b\ell'} + \frac{V_{cs}}{V_{cb}} x_L^{s\ell'} + \frac{V_{cd}}{V_{cb}} x_L^{d\ell'} \right], \quad (11)$$

and contribute to $b \rightarrow c\ell\bar{\nu}_{\ell'}$. It becomes one solution for the R_D and R_{D^*} anomalies, and the 1σ region for $b \rightarrow c\tau\bar{\nu}_\tau$ requires

$$g_{V_L} \subseteq [0.09, 0.13] \Rightarrow \frac{(V_{cs}^{\text{CKM}} x_L^{23} + V_{cb}^{\text{CKM}} x_L^{33})(x_L^{33})^*}{m_{U_1}^2} \subseteq [0.12, 0.18] \text{ TeV}^{-2}. \quad (12)$$

The RH couplings x_R , combining with x_L^{22} and x_L^{32} , contribute to the Wilson coefficients $(C_S)' = (C_P)'$, thus their magnitudes are bounded by the $B \rightarrow K\tau\mu$ and $B_s \rightarrow \mu\mu$ data, which hampers from generating large enough muon magnetic dipole for $(g-2)_\mu$ anomaly. Therefore, we are motivated to further extend our model.

B. Vector Leptoquark with $U(1)_{B_3-L_2}$ X boson

In the following comprehensive analysis, with the preferred parameter region for $R_{K^{(*)}}$ and $R_{D^{(*)}}$ anomalies, the U_1 leptoquark itself may still not be able to provide large enough contribution for $(g-2)_\mu$ anomaly. Therefore, we invoke the additional particle X , $U(1)_{B_3-L_2}$ gauge boson to enhance the contribution, which is alternatively plausible candidate for R_K anomaly and significantly change the preferred parameter space. We take the most relevant interactions with X boson in the effective Lagrangian

$$\begin{aligned} \mathcal{L}_{\text{eff}}^X \supseteq & -g_X \bar{\mu} \gamma^\alpha \mu X_\alpha - g_X \bar{\nu}_\mu \gamma^\alpha P_L \nu_\mu X_\alpha \\ & + \frac{g_X}{3} \bar{\mathbf{u}}_L \gamma^\alpha \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 1 \end{pmatrix} \mathbf{u}_L X_\alpha + \frac{g_X}{3} \bar{\mathbf{d}}_L \gamma^\alpha \underbrace{\begin{pmatrix} |V_{td}|^2 & V_{ts} V_{td}^* & V_{tb} V_{td}^* \\ V_{td} V_{ts}^* & |V_{ts}|^2 & V_{tb} V_{ts}^* \\ V_{td} V_{tb}^* & V_{ts} V_{tb}^* & |V_{tb}|^2 \end{pmatrix}}_{=V_{\text{CKM}}^\dagger V_{B_3} V_{\text{CKM}}} \mathbf{d}_L X_\alpha \\ & + \frac{1}{2} m_X^2 X^\mu X_\mu \end{aligned} \quad (13)$$

where $\mathbf{u}_L^T \equiv (u_L, c_L, t_L)$ and $\mathbf{d}_L^T \equiv (d_L, s_L, b_L)$. Here, we chose the configuration that the left-handed down-quark mixing contributes entire CKM matrix to avoid the constraint from $D - \bar{D}$ mixing. We assume that the m_X is induced by a new Higgs mechanism but the additional contribution from the new Higgs can be neglected. The flavor-changing neutral current (FCNC) in down quarks from the X boson contributes to $B \rightarrow K\mu\mu$ and $B_s \rightarrow \mu\mu$. At the same time, the muon coupling modifies the $(g-2)_\mu$. In the following, we assume the $U(1)_{B_3-L_2}$ are broken under energy scale $\mathcal{O}(100)$ GeV, which justifies the non-zero low-energy effective couplings at $\mathcal{O}(m_b)$ scale in Eq. (8).

III. LOW-ENERGY OBSERVABLES

In this section, we summarize the low-energy observables with the U_1 vector leptoquark contributions.

A. $(g-2)_\mu$

The previous result of Muon $(g-2)$ experiment with the BNL E821 was 3.7σ from the SM. After the FNAL result, the difference between experiment and SM has become [14, 46]

$$\Delta a_\mu = a_\mu^{\text{exp}} - a_\mu^{\text{SM}} = (251 \pm 59) \times 10^{-11}, \quad (14)$$

which is a deviation of 4.2σ significance from the SM prediction and this enhances the motivation for SM extensions for new couplings with leptons. In this paper, we present the single leptoquark which is described in Sec. II A to explain not only $(g-2)_\mu$ anomaly but also B -meson anomalies.

For the large mass of U_1 leptoquark, it contributes to the $(g-2)_\mu$ anomaly as³

$$\begin{aligned} \Delta a_\mu = \frac{N_c}{16\pi^2} \sum_i \left[2Q_U \tilde{\kappa}_Y \text{Im}(x_L^{i2} (x_R^{i2})^*) \frac{m_{d_i} m_\mu}{m_{U_1}^2} \left(\ln \left(\frac{\Lambda_{UV}^2}{M_{U_1}^2} \right) + \frac{5}{2} \right) \right. \\ \left. + 2\text{Re}(x_L^{i2} (x_R^{i2})^*) \frac{m_{d_i} m_\mu}{m_{U_1}^2} \left(2Q_d + Q_{U_1} \left((1 - \kappa_Y) \ln \left(\frac{\Lambda_{UV}^2}{M_{U_1}^2} \right) + \frac{1 - 5\kappa_Y}{2} \right) \right) \right. \\ \left. - (|x_L^{i2}|^2 + |x_R^{i2}|^2) \frac{m_\mu^2}{m_{U_1}^2} \left(\frac{4}{3} Q_d + Q_{U_1} \left((1 - \kappa_Y) \ln \left(\frac{\Lambda_{UV}^2}{M_{U_1}^2} \right) - \frac{1 + 9\kappa_Y}{6} \right) \right) \right]. \quad (15) \end{aligned}$$

If $\kappa_Y \neq 1$ and $\tilde{\kappa}_Y \neq 0$, the dipole moment exhibits logarithmic dependence on the cut-off scale Λ_{UV} not far above the leptoquark mass. So, the leptoquark contribution to $(g-2)_\mu$ anomaly becomes

$$\begin{aligned} \Delta a_\mu = \frac{N_c}{16\pi^2} \sum_i \left[2\text{Re}(x_L^{i2} (x_R^{i2})^*) \frac{m_{d_i} m_\mu}{m_{U_1}^2} \left(2Q_d + Q_{U_1} \left(\frac{1 - 5\kappa_Y}{2} \right) \right) \right. \\ \left. - (|x_L^{i2}|^2 + |x_R^{i2}|^2) \frac{m_\mu^2}{m_{U_1}^2} \left(\frac{4}{3} Q_d + Q_{U_1} \left(-\frac{1 + 9\kappa_Y}{6} \right) \right) \right], \quad (16) \end{aligned}$$

where we use $\kappa_Y = 1$, $\tilde{\kappa}_Y = 0$, $N_C = 3$, $Q_b = -1/3$, and $Q_{U_1} = 2/3$. We handle the renormalization group running from leptoquark scale down to muon mass by evaluating the quark masses at $\mathcal{O}(\text{TeV})$ scale in Eq.(16), e.g. $m_b(\text{TeV}) \simeq 2.4$ GeV [8].

The X boson also contributes to $(g-2)_\mu$ as [18]

$$\Delta a_\mu^X = \frac{g_X^2}{8\pi^2} \int_0^1 dz \frac{2z(1-z)^2}{(1-z)^2 + z(m_X/m_\mu)^2} \quad (17)$$

$$\simeq (3g_X^2/4\pi)(m_\mu^2/m_X^2) \quad \text{when } m_X \gg m_\mu \quad (18)$$

$$\simeq 2.7 \times 10^{-11} \times \left(\frac{g_X}{0.01} \right)^2 \left(\frac{100 \text{ GeV}}{m_X} \right)^2, \quad (19)$$

³ We use the same κ and $\tilde{\kappa}$ parameters defined in Ref. [8], which are the tri-gauge boson couplings between U_1 leptoquarks and B_μ gauge boson from $U(1)_Y$. In Eq. (16), we set $\kappa = 1$ and $\tilde{\kappa} = 0$, in such a way, the dipole moment becomes independent on the logarithmic term and UV theory.

thus it is negligible compared to the experimental value for $m_X \simeq 100$ GeV and $g_X \simeq 0.01$. However, with X boson contributions, the preferred parameter space for B -meson anomalies would move and our fit to the data can be significantly improved as we will describe below.

B. $R_{K^{(*)}}, R_{D^{(*)}}$

To explain the experimental result of $R_{K^{(*)}}$, it requires the Wilson coefficients as

$$\Delta C_9^{\mu\mu}|_{\text{exp}} = -0.40 \pm 0.12, \quad \Delta C_9^U|_{\text{exp}} = -0.50 \pm 0.38, \quad (20)$$

with correlation -0.5 [32, 33, 47] between them. And the ratio of the SM predictions and experimental observation is

$$\frac{R_D^{\text{exp}}}{R_D^{\text{SM}}} = 1.14 \pm 0.10, \quad \frac{R_{D^*}^{\text{exp}}}{R_{D^*}^{\text{SM}}} = 1.14 \pm 0.05, \quad (21)$$

with correlation -0.37 [33]. For the U_1 leptoquark, the contribution to Wilson coefficients is [32, 33]

$$\Delta C_9^{\mu\mu} = -\Delta C_{10}^{\mu\mu} = -\frac{4\pi^2}{e^2} \frac{v^2}{m_{U_1}^2} \frac{x_L^{32}(x_L^{22})^*}{V_{ts}^* V_{tb}}, \quad (22)$$

$$\Delta C_9^U \simeq -\frac{1}{V_{tb} V_{ts}^*} \frac{2}{3} \frac{v^2}{m_{U_1}^2} x_L^{23}(x_L^{33})^* \ln \left(\frac{m_b^2}{m_{U_1}^2} \right), \quad (23)$$

where ΔC_9^U is the lepton-universal contribution to $b \rightarrow s\ell\ell$ and originates from the x_L^{23} contributing through a log-enhanced photon penguin diagram [33, 40]. Similarly, the U_1 contribution to R_D and R_{D^*} are [33]

$$\begin{aligned} \frac{R_D}{R_D^{\text{SM}}} &\simeq \left[1 + \frac{v^2}{m_{U_1}^2} \text{Re} \left\{ (x_L^{33} - 1.5\eta_S(x_R^{33})^*) \frac{(V_{cb}x_L^{33} + V_{cs}x_L^{23} + V_{cd}x_L^{13})}{V_{cb}} \right\} \right], \\ \frac{R_{D^*}}{R_{D^*}^{\text{SM}}} &\simeq \left[1 + \frac{v^2}{m_{U_1}^2} \text{Re} \left\{ (x_L^{33} - 0.14\eta_S(x_R^{33})^*) \frac{(V_{cb}x_L^{33} + V_{cs}x_L^{23} + V_{cd}x_L^{13})}{V_{cb}} \right\} \right], \end{aligned} \quad (24)$$

where $\eta_S \simeq 1.8$ accounts for the running of the scalar operator from $m_{U_1} = 4$ TeV to m_b .

The additional correction from X boson to the $\Delta C_9^{\mu\mu}$ is given as [43]

$$\begin{aligned} \Delta C_9^X &= \left(\frac{g_X^2}{3} V_{tb} V_{ts}^* \right) \left(\frac{36 \text{ TeV}}{m_X} \right)^2 \\ &\approx -0.18 \times \left(\frac{g_X}{0.01} \right)^2 \left(\frac{100 \text{ GeV}}{m_X} \right)^2 \end{aligned} \quad (25)$$

It is important to notice that the negative value from V_{ts} makes it trend toward the experimental value for B -meson anomalies but does not contribute to $(g-2)_\mu$.

C. $B_c^- \rightarrow \tau^- \bar{\nu}_\tau, B^+ \rightarrow \tau^+ \nu_\tau$

The bound from observation [48, 49] and SM prediction for $B_c \rightarrow \tau\nu$ are [8, 50]

$$\begin{aligned} \mathcal{B}(B_c^- \rightarrow \tau^- \bar{\nu}_\tau)_{\text{exp}} &\leq 0.60 \\ \mathcal{B}(B_c^- \rightarrow \tau^- \bar{\nu}_\tau)_{\text{SM}} &= (2.21 \pm 0.09) \times 10^{-2} \end{aligned} \quad (26)$$

The proportion of experimental value and SM prediction for $B^+ \rightarrow \tau^+ \nu_\tau$ is [8]

$$\frac{\mathcal{B}(B^+ \rightarrow \tau^+ \nu_\tau)_{\text{exp}}}{\mathcal{B}(B^+ \rightarrow \tau^+ \nu_\tau)_{\text{SM}}} = 1.30 \pm 0.29. \quad (27)$$

And the U_1 leptoquark contributions to each of observables are [8]

$$\frac{\mathcal{B}(B_c^- \rightarrow \tau^- \bar{\nu}_\tau)}{\mathcal{B}(B_c^- \rightarrow \tau^- \bar{\nu}_\tau)_{\text{SM}}} = \left| 1 - \frac{(V_{cd}x_L^{13} + V_{cs}x_L^{23} + V_{cb}x_L^{33})}{V_{cb}} \frac{v^2}{m_{U_1}^2} \left(\frac{(x_L^{33})^*}{2} + \frac{(x_R^{33})^* m_{B_c}^2}{m_\tau(m_b + m_c)} \right) \right|^2, \quad (28)$$

$$\frac{\mathcal{B}(B^+ \rightarrow \tau^+ \nu_\tau)}{\mathcal{B}(B^+ \rightarrow \tau^+ \nu_\tau)_{\text{SM}}} = \left| 1 - \frac{(V_{ud}x_L^{13} + V_{us}x_L^{23} + V_{ub}x_L^{33})}{V_{ub}} \frac{v^2}{m_{U_1}^2} \left(\frac{(x_L^{33})^*}{2} + \frac{(x_R^{33})^* m_{B^\pm}^2}{m_\tau m_b} \right) \right|^2. \quad (29)$$

D. $B_s^0 \rightarrow \tau^+ \tau^-$, $B_s^0 \rightarrow \mu^+ \mu^-$, $B_s^0 \rightarrow \tau^\pm \mu^\mp$ and $B^+ \rightarrow K^+ \tau^+ \tau^-$

The experimental value from LHCb is [51] and SM prediction is [52]

$$\mathcal{B}(B_s^0 \rightarrow \tau^+ \tau^-)_{\text{exp}} < 6.8 \times 10^{-3} \quad \text{at 95\% C.L.}, \quad (30)$$

$$\mathcal{B}(B_s^0 \rightarrow \tau^+ \tau^-)_{\text{SM}} = (7.73 \pm 0.49) \times 10^{-7}. \quad (31)$$

And the related contribution for U_1 leptoquark is [8]

$$\begin{aligned} \frac{\mathcal{B}(B_s^0 \rightarrow \tau^+ \tau^-)}{\mathcal{B}(B_s^0 \rightarrow \tau^+ \tau^-)_{\text{SM}}} &= \frac{16\pi^4}{e^4 (C_{10}^{\text{SM}})^2} \frac{v^4}{m_{U_1}^4} \frac{m_{B_s}^4}{m_\tau^2 m_b^2} \left| \frac{(x_L^{33})^* x_R^{23} - (x_R^{33})^* x_L^{23}}{V_{ts}^* V_{tb}} \right|^2 \left(1 - \frac{4m_\tau^2}{m_{B_s}^2} \right) \\ &+ \left| 1 + \frac{4\pi^2}{e^2 C_{10}^{\text{SM}}} \frac{v^2}{m_{U_1}^2} \left(\frac{(x_L^{33})^* x_L^{23} + (x_R^{33})^* x_R^{23}}{V_{ts}^* V_{tb}} - \frac{m_{B_s}^2}{m_\tau m_b} \frac{(x_L^{33})^* x_R^{23} + (x_R^{33})^* x_L^{23}}{V_{ts}^* V_{tb}} \right) \right|^2, \end{aligned} \quad (32)$$

where $C_{10}^{\text{SM}} \simeq -4.1$ which we use for a normalization such that the SM value for the Wilson coefficient [53].

The ratio between the SM prediction and experimental value is [8, 32]

$$\frac{\mathcal{B}(B_s^0 \rightarrow \mu^+ \mu^-)_{\text{exp}}}{\mathcal{B}(B_s^0 \rightarrow \mu^+ \mu^-)_{\text{SM}}} = 0.73_{-0.10}^{+0.13}. \quad (33)$$

And the following U_1 vector leptoquark and X -boson total contribution is written by [8, 54]

$$\begin{aligned} \frac{\mathcal{B}(B_s^0 \rightarrow \mu^+ \mu^-)}{\mathcal{B}(B_s^0 \rightarrow \mu^+ \mu^-)_{\text{SM}}} &= \frac{16\pi^4}{e^4 (C_{10}^{\text{SM}})^2} \frac{v^4}{m_{U_1}^4} \frac{m_{B_s}^4}{m_\mu^2 m_b^2} \left| \frac{(x_L^{32})^* x_R^{22} - (x_R^{32})^* x_L^{22}}{V_{ts}^* V_{tb}} \right|^2 \\ &+ \left| 1 + \frac{4\pi^2}{e^2 C_{10}^{\text{SM}}} \frac{v^2}{m_{U_1}^2} \left(\frac{(x_L^{32})^* x_L^{22} + (x_R^{32})^* x_R^{22}}{V_{ts}^* V_{tb}} - \frac{m_{B_s}^2}{m_\mu m_b} \frac{(x_L^{32})^* x_R^{22} + (x_R^{32})^* x_L^{22}}{V_{ts}^* V_{tb}} \right) \right. \\ &\quad \left. - \left[\frac{\alpha}{2\pi \sin^2 \theta_W} Y \left(\frac{m_t^2}{m_W^2} \right) \right]^{-1} \left(\frac{2g_X^2 m_Z^2}{3g^2 m_X^2} \right) \right|^2, \end{aligned} \quad (34)$$

where the last term in the second absolute bracket comes from the X -boson. Here we take $Y(m_t^2/m_W^2) = 1.05$, and $g \simeq 0.652$ the $SU(2)_L$ coupling constant.

LHCb search on $B_s^0 \rightarrow \tau^\pm \mu^\mp$ provides an upper limit

$$\mathcal{B}(B_s^0 \rightarrow \tau^\pm \mu^\mp)_{\text{exp}} < 2.1 \times 10^{-5}, \quad (35)$$

at 95 % confidence level [55]. SM prediction of this branching fraction is extremely small as $\mathcal{O}(10^{-54})$ [56]. The expression of U_1 contribution to $B_s^0 \rightarrow \tau^\pm \mu^\mp$ is [33]

$$\begin{aligned} \mathcal{B}(B_s^0 \rightarrow \tau^\pm \mu^\mp) &= \frac{1}{\Gamma_{B_s}} \frac{m_{B_s} f_{B_s}^2 G_F^2}{8\pi} m_\tau^2 \left(1 - \frac{m_\tau^2}{m_{B_s}^2} \right)^2 \\ &\quad \times \frac{v^4}{4m_{U_1}^4} \left| x_L^{22} (x_L^{33})^* - \frac{2\eta_S m_{B_s}^2}{m_\tau (m_s + m_b)} x_L^{22} (x_R^{33})^* \right|^2. \end{aligned} \quad (36)$$

where $G_F = 1.166 \times 10^{-5} \text{ GeV}^{-2}$ is the Fermi constant, $f_{B_s} = 0.225 \text{ GeV}$ [57] is the leptonic decay constant of B_s^0 , and $\Gamma_{B_s} = 4.34 \times 10^{-13} \text{ GeV}$ is the total width of B_s^0 .

BaBar experiment measured the branching fraction [58]

$$\mathcal{B}(B^+ \rightarrow K^+ \tau^+ \tau^-)_{\text{exp}} = (1.31 \pm 0.71) \times 10^{-3}. \quad (37)$$

with an upper limit of $\text{Br}(B^+ \rightarrow K^+ \tau^+ \tau^-) < 2.25 \times 10^{-3}$ at the 90% confidence level. The expression of U_1 leptoquark contribution to this process is given by [33]

$$\begin{aligned} \mathcal{B}(B^+ \rightarrow K^+ \tau^+ \tau^-) \simeq & 1.5 \times 10^{-7} + 10^{-3} \frac{v^2}{2m_{U_1}^2} [1.4 \text{Re}(x_L^{23}(x_L^{33})^*) - 3.3 \text{Re}(x_L^{23}(x_R^{33})^*)] \\ & + \frac{v^4}{4m_{U_1}^4} |x_L^{23}|^2 [3.5 |x_L^{33}|^2 - 16.4 \text{Re}(x_R^{33}(x_L^{33})^*) + 95.0 |x_R^{33}|^2]. \end{aligned} \quad (38)$$

where $v = 246 \text{ GeV}$ is the electroweak vacuum expectation value.

E. $B^+ \rightarrow K^+ \tau^+ \mu^-$, $B^+ \rightarrow K^+ \tau^- \mu^+$

From BaBar experiment, we obtain upper limits [59]

$$\mathcal{B}(B^+ \rightarrow K^+ \tau^+ \mu^-) < 2.8 \times 10^{-5}, \quad (39)$$

$$\mathcal{B}(B^+ \rightarrow K^+ \tau^- \mu^+) < 4.5 \times 10^{-5} \quad (40)$$

at 90% confidence level. The leptoquark contribution is given by [33, 60]

$$\begin{aligned} \mathcal{B}(B^+ \rightarrow K^+ \tau^+ \mu^-) \simeq & \frac{v^4}{4m_{U_1}^4} |x_L^{22}|^2 [8.3 |x_L^{33}|^2 + 155.2 |x_R^{33}|^2 - 42.3 \text{Re}((x_L^{33})^* x_R^{33})], \\ \mathcal{B}(B^+ \rightarrow K^+ \tau^- \mu^+) \simeq & 8.3 \frac{v^4}{4m_{U_1}^4} |x_L^{32}(x_L^{23})^*|^2. \end{aligned} \quad (41)$$

F. $\tau \rightarrow \mu\gamma$, $\mu\phi$ and LFU in τ decays

Due to its sizeable couplings to muon and tau leptons, U_1 leptoquark can significantly affect the Lepton-flavor-violation in τ decays. The experimental upper limits are [61, 62]

$$\mathcal{B}(\tau \rightarrow \mu\gamma) < 3.0 \times 10^{-8}. \quad (42)$$

$$\mathcal{B}(\tau \rightarrow \mu\phi) < 5.1 \times 10^{-8} \quad (43)$$

at 90% confidence level. In addition, the LFU in the decay of charged leptons can give stringent bounds on the leptoquark couplings. The experimentally measured values are [61, 63]

$$(g_\tau/g_\mu)_{\text{exp}} = 1.0000 \pm 0.0014, \quad (44)$$

$$(g_\tau/g_\mu)_\ell = 1.0010 \pm 0.0015, \quad (45)$$

$$(g_\tau/g_\mu)_\pi = 0.9961 \pm 0.0027, \quad (46)$$

$$(g_\tau/g_\mu)_K = 0.9860 \pm 0.0070. \quad (47)$$

For the U_1 leptoquark contribution to $\mathcal{B}(\tau \rightarrow \mu\gamma)$, we adopt complete formula of the decay width [64]

$$\Gamma(\tau \rightarrow \mu\gamma) = \frac{\alpha_{\text{EM}}(m_\tau^2 - m_\mu^2)^3}{4m_\tau^3} (|\sigma_L^{32}|^2 + |\sigma_R^{32}|^2), \quad (48)$$

where

$$\begin{aligned}
\sigma_L^{32} &= -\frac{iN_c}{16\pi^2 m_{U_1}^2} \sum_{k=1,2,3} \left\{ \frac{2}{3} \left[\left(x_R^{k2*} x_R^{k3} m_\tau + x_L^{k2*} x_L^{k3} m_\mu \right) g(t_k) + x_R^{k2*} x_L^{k3} m_{d_k} j(t_k) \right] \right. \\
&\quad \left. - \frac{1}{3} \left[\left(x_R^{k2*} x_R^{k3} m_\tau + x_L^{k2*} x_L^{k3} m_\mu \right) f(t_k) + x_R^{k2*} x_L^{k3} m_{d_k} h(t_k) \right] \right\} \\
\sigma_R^{32} &= -\frac{iN_c}{16\pi^2 m_{U_1}^2} \sum_{k=1,2,3} \left\{ \frac{2}{3} \left[\left(x_L^{k2*} x_L^{k3} m_\tau + x_R^{k2*} x_R^{k3} m_\mu \right) g(t_k) + x_L^{k2*} x_R^{k3} m_{d_k} j(t_k) \right] \right. \\
&\quad \left. - \frac{1}{3} \left[\left(x_L^{k2*} x_L^{k3} m_\tau + x_R^{k2*} x_R^{k3} m_\mu \right) f(t_k) + x_L^{k2*} x_R^{k3} m_{d_k} h(t_k) \right] \right\}
\end{aligned} \tag{49}$$

with $t_k \equiv m_{d_k}^2/m_{U_1}^2$ and f, g, h, j are loop functions [64].

For $\mathcal{B}(\tau \rightarrow \mu\phi)$, it is given by [33, 65]

$$\mathcal{B}(\tau \rightarrow \mu\phi) = \frac{1}{\Gamma_\tau} \frac{f_\phi^2 G_F^2}{16\pi} m_\tau^3 \left(1 - \frac{m_\phi^2}{m_\tau^2} \right)^2 \left(1 + \frac{2m_\phi^2}{m_\tau^2} \right) \frac{v^4}{4m_{U_1}^4} |x_L^{23} (x_L^{22})^*|^2, \tag{50}$$

where ϕ is the $s\bar{s}$ vector meson with $f_\phi = 0.225$ GeV, $m_\phi = 1.019$ GeV [65]. For LFU in lepton decays, we use the expression [33, 63]

$$\left(\frac{g_\tau}{g_\mu} \right)_{\ell, \pi, K} \simeq 1 - 0.08 \times \frac{(x_L^{33})^2 v^2}{4m_{U_1}^2}. \tag{51}$$

More specifically, they can be written in terms of the effective Lagrangian for leptonic decay [63]

$$\mathcal{L}_{\ell \rightarrow \ell' \nu \bar{\nu}} = -\frac{4G_F}{\sqrt{2}} \left([C_{\nu e}^{V,LL}]_{\rho\sigma\alpha\beta} (\bar{\nu}_L^\rho \gamma^\mu \nu_L^\sigma) (\bar{\ell}_L^\alpha \gamma^\mu \ell_L^\beta) + [C_{\nu e}^{V,LR}]_{\rho\sigma\alpha\beta} (\bar{\nu}_L^\rho \gamma^\mu \nu_L^\sigma) (\bar{\ell}_R^\alpha \gamma^\mu \ell_R^\beta) \right), \tag{52}$$

and for hadronic decay [63]

$$\mathcal{L}_{\tau \rightarrow h\nu} = -\frac{4G_F}{\sqrt{2}} \sum_\rho \left\{ \left(\delta_{\rho 3} (V_{CKM})_{ji}^* [C_{\nu edu}^{V,LL}]_{\rho 3ij} \right) (\bar{\nu}_L^\rho \gamma^\mu \tau_L) (\bar{d}_L^i \gamma^\mu u_L^j) + [C_{\nu edu}^{S,RL}]_{\rho 3ij} (\bar{\nu}_L^\rho \tau_L) (\bar{d}_R^i u_R^j) \right\} \tag{53}$$

result in the following expressions for couplings with τ and μ

$$\begin{aligned}
\left(\frac{g_\tau}{g_\mu} \right)_\ell &= \left[\frac{\sum_{\rho\sigma} \left(\left| \delta_{\rho 3} \delta_{\sigma 1} + [C_{\nu e}^{V,LL}]_{\rho\sigma 13} \right|^2 + \left| [C_{\nu e}^{V,LR}]_{\rho\sigma 13} \right|^2 \right)}{\sum_{\rho\sigma} \left(\left| \delta_{\rho 2} \delta_{\sigma 1} + [C_{\nu e}^{V,LL}]_{\rho\sigma 12} \right|^2 + \left| [C_{\nu e}^{V,LR}]_{\rho\sigma 12} \right|^2 \right)} \right]^{1/2}, \\
\left(\frac{g_\tau}{g_\mu} \right)_\pi &= \left(\frac{\sum_\rho \left| \delta_{\rho 3} V_{ud}^* + [C_{\nu edu}^{V,LL}]_{\rho 311} + \frac{m_\pi^2}{m_\tau(m_d+m_u)} [C_{\nu edu}^{S,RL}]_{\rho 311} \right|^2}{\sum_\rho \left| \delta_{\rho 2} V_{ud} + [C_{\nu edu}^{V,LL}]_{\rho 211}^* + \frac{m_\pi^2}{m_\mu(m_d+m_u)} [C_{\nu edu}^{S,RL}]_{\rho 211}^* \right|^2} \right)^{1/2}, \\
\left(\frac{g_\tau}{g_\mu} \right)_K &= \left(\frac{\sum_\rho \left| \delta_{\rho 3} V_{us}^* + [C_{\nu edu}^{V,LL}]_{\rho 321} + \frac{m_K^2}{m_\tau(m_d+m_u)} [C_{\nu edu}^{S,RL}]_{\rho 321} \right|^2}{\sum_\rho \left| \delta_{\rho 2} V_{us} + [C_{\nu edu}^{V,LL}]_{\rho 221}^* + \frac{m_K^2}{m_\mu(m_d+m_u)} [C_{\nu edu}^{S,RL}]_{\rho 221}^* \right|^2} \right)^{1/2}.
\end{aligned} \tag{54}$$

For U_1 leptoquark, by using the Fierz transformation in Eq.(8),

$$[\bar{u}_{1L} \gamma^\mu u_{2L}] [\bar{u}_{3L} \gamma_\mu u_{4L}] = -[\bar{u}_{1L} \gamma^\mu u_{4L}] [\bar{u}_{3L} \gamma_\mu u_{2L}],$$

TABLE I. The list of observables for the χ^2 scanning with the measured values and predictions from SM. The equations of the $U_1(+X)$ model are also referenced.

| Observable | Experiment | SM predict | $U_1(+X)$ predict |
|---|---|--|-------------------|
| $R_{D^{(*)}}$ | $\frac{R_D^{\text{exp}}}{R_D^{\text{SM}}} = 1.14 \pm 0.10,$ | $\frac{R_{D^*}^{\text{exp}}}{R_{D^*}^{\text{SM}}} = 1.14 \pm 0.05$ | (24) |
| $\Delta C_9^{\mu\mu} = -\Delta C_{10}^{\mu\mu} (R_{K^{(*)}})$ | -0.40 ± 0.12 | 0 | (22) |
| ΔC_9^U | -0.50 ± 0.38 | 0 | (23), (25) |
| $B_c^- \rightarrow \tau^- \bar{\nu}_\tau$ | < 0.60 | $(2.21 \pm 0.09) \times 10^{-2}$ | (28) |
| $B^+ \rightarrow \tau^+ \nu_\tau$ | $(1.09 \pm 0.24) \times 10^{-4}$ | $(8.8 \pm 0.6) \times 10^{-5}$ | (29) |
| $B_s^0 \rightarrow \tau^+ \tau^-$ | $< 6.8 \times 10^{-3}$ | $(7.73 \pm 0.49) \times 10^{-7}$ | (32) |
| $B_s^0 \rightarrow \mu^+ \mu^-$ | $\frac{BR(B_s^0 \rightarrow \mu^+ \mu^-)^{\text{exp}}}{BR(B_s^0 \rightarrow \mu^+ \mu^-)^{\text{SM}}} = 0.73_{-0.10}^{+0.13}$ | | (34) |
| $B_s^0 \rightarrow \tau^\pm \mu^\mp$ | $< 2.1 \times 10^{-5}$ | | (36) |
| $B^+ \rightarrow K^+ \tau^+ \tau^-$ | $(1.31 \pm 0.71) \times 10^{-3}$ | $(1.20 \pm 0.12) \times 10^{-7}$ | (41) |
| $B^+ \rightarrow K^+ \tau^+ \mu^-$ | $\leq 2.8 \times 10^{-5}$ | | (41) |
| $B^+ \rightarrow K^+ \tau^- \mu^+$ | $\leq 4.5 \times 10^{-5}$ | | (41) |
| $\tau \rightarrow \mu \gamma$ | $< 3.0 \times 10^{-8}$ | | (48) |
| $\tau \rightarrow \mu \phi$ | $< 5.1 \times 10^{-8}$ | | (50) |
| LFU in τ decay | $(g_\tau/g_\mu)_\ell = 1.0010 \pm 0.0015$ | | (54) |
| | $(g_\tau/g_\mu)_\pi = 0.9961 \pm 0.0027$ | | |
| | $(g_\tau/g_\mu)_K = 0.9860 \pm 0.0070$ | | |

we replace the Wilson coefficients as

$$\begin{aligned}
 [C_{\nu edu}^{\text{V,LL}}]_{\rho\sigma ij} &\Leftrightarrow -\frac{2v^2}{4m_{U_1}^2} (V_{\text{CKM}} x_L U_{\text{PMNS}}^*)_{j\rho} (x_L)_{i\sigma}, \\
 [C_{\nu edu}^{\text{S,RL}}] &= [C_{\nu e}^{\text{V,LL}}] = [C_{\nu e}^{\text{V,LR}}] = 0.
 \end{aligned} \tag{55}$$

IV. PARAMETER SCANNING

We perform the χ^2 parameter scanning, with the values of all 17 observables constituting the χ^2 listed in Table I, which includes anomalies of $R_{K^{(*)}}$, $R_{D^{(*)}}$, constraints from other B -meson decay channels, and constraints from τ decays. Under the null hypothesis (SM only), we have $\chi_{\text{SM}}^2 = 26.0$ along with $p_{\text{SM}} = 0.074$ where p_{SM} is the P -value of the null hypothesis. We compare this result with the following three scenarios:

- **scan-1:** $P_{\text{scan-1}} = (x_L^{22}, x_L^{23}, x_L^{32}, x_L^{33})$, with $m_{U_1} = 2, 5$ TeV.
Results are in Fig. 1.
- **scan-2:** $P_{\text{scan-2}} = P_{\text{scan-1}} \oplus x_R^{32}$, with $m_{U_1} = 2$ TeV.
Results are in Fig. 2.
- **scan-3:** $P_{\text{scan-3}} = P_{\text{scan-2}} \oplus g_X$, with $m_{U_1} = 2$ TeV, and $m_X = 100$ GeV.
Results are in Fig. 3.




FIG. 1. **Scan-1:** The region satisfies $\Delta\chi^2 \equiv \chi^2 - \chi_{\min,1}^2 \leq 2.3$, and $\chi_{\min,1}^2 = 9.23$.

For **scan-1**, we choose a set of relevant couplings, x_L^{22} , x_L^{23} , x_L^{32} , x_L^{33} , that is sufficient to explain the B -meson anomalies and satisfies all the low-energy observables. It gives rise to the best fit $\chi_{\min,1}^2 = 9.23$ and $p_1 = 0.755$ with $(x_L^{22}, x_L^{23}, x_L^{32}, x_L^{33}) = (7.90 \times 10^{-2}, 0.328, -3.83 \times 10^{-2}, 0.862)$ and $m_{U_1} = 2$ TeV. Figure 1 shows the 1σ region for the **scan-1**. The favored region of the (x_L^{23}, x_L^{33}) plane, shown in the upper-left panel of Fig. 1, is mainly determined by the $R_{D^{(*)}}$, meanwhile the $R_{K^{(*)}}$ dictates the favored region of the (x_L^{22}, x_L^{32}) plane. There is contribution to Δa_μ from the LH couplings, as shown in the upper-right panels of Fig. 1. However, it is not large enough to explain the recent measurement from Fermilab, thus providing the motivation to turn on the RH coupling in the **scan-2**.

In the **scan-2** ($\chi_{\min,2}^2 = 9.06$ and $p_2 = 0.698$), according to Eq.(16), we found the most efficient way to enhance Δa_μ is to use x_R^{32} , due to the fact that the multiplicity of $(x_L^{32} x_R^{32})$ has milder mass suppression factor ($m_b m_\mu / m_{U_1}$) than that of pure LH couplings or RH couplings. In the $(x_L^{32}, \Delta a_\mu)$ and $(x_R^{32}, \Delta a_\mu)$ planes of Fig. 2, they show that the value of Δa_μ can be increased by more than an order of magnitude in contrast to the **scan-1**. However, it still cannot explain the BNL and FNAL results without additional contribution. For example, incorporating a muon-philic vector boson X with coupling $g_X = 0.2$ and mass $m_X = 100$ GeV, which is consistent with current experimental bounds, endows $\Delta a_\mu|_X = 38 \times 10^{-11}$ shown by the green hatched regions overlapping with U_1 2σ -allowed region in Fig. 2.




FIG. 2. **Scan-2:** The 1σ (2σ) region satisfies $\Delta\chi^2 \equiv \chi^2 - \chi_{\min,2}^2 \leq 2.30$ (5.99), and $\chi_{\min,2}^2 = 9.06$. Here we fix $m_{U_1} = 2$ TeV. The green hatched regions show the effect of a muon-philic X vector boson with coupling $g_X = 0.2$ and mass $m_X = 100$ GeV, which ends $\Delta a_\mu|_X = 38 \times 10^{-11}$.

In the **scan-3**, we include a specific $U(1)_{B_3-L_2}$ X boson in addition to the U_1 leptoquark framework, and demonstrate that under this framework it can alleviate the $(g-2)_\mu$ and B -physics tensions within 2σ . Not only the $(g-2)_\mu$, but this X boson also contributes to both ΔC_9 (see Eq.(25)) and $B_s \rightarrow \mu\mu$ (see Eq.(34)). In particular, we vary the coupling $0.01 \leq g_X \leq 0.05$, and fix $m_X = 100$ GeV, which is partially allowed under current experimental constraints from neutrino-trident [66], $B_s^0 - \bar{B}_s^0$ mixing [67], $K_L \rightarrow \mu^+\mu^-$ [68], ATLAS [69], and CMS [70, 71].

It is worth to mention the reliable range of X boson parameters consistent with the measurement of i) $B_s^0 - \bar{B}_s^0$ mixing (ΔM_s), ii) $Br(K_L \rightarrow \mu^+\mu^-)$ and iii) $K^0 - \bar{K}^0$ mixing parameters. The mass difference ΔM_s has been precisely measured by CDF2, LHCb and CMS collaborations [72, 73], and its theoretical predictions has been improved by developed sum rules and lattice calculations. We use the weighted average of the latest results given in Ref. [67] as our ΔM_s^{SM} . In the presence of X boson, new physics contribution to ΔM_s is approximated by $\frac{\Delta M_s^{SM+NP}}{\Delta M_s^{NP}} \approx \left| 1 + \frac{1}{360} \left(\frac{\delta C_9^\mu}{-0.53} \right)^2 \left(\frac{m_X/g_X}{1 \text{ TeV}} \right)^2 \right|$ [67]. Putting $\delta C_9^\mu (= -\delta C_{10}^\mu) = -0.40 \pm 0.12$ [32, 33, 47] into the expression and requiring




FIG. 3. **Scan-3:** Same as Fig. 2, but includes the contribution of the additional $U(1)_{B_3-L_2}$ X boson. Here we vary $0.01 \leq g_X \leq 0.05$, meanwhile fix $m_{U_1} = 2$ TeV and $m_X = 100$ GeV.

$\Delta M_s = (1.04_{-0.07}^{+0.04})\Delta M_s^{\text{exp}}$, the lower limit on the coupling is $g_X > 1.01 \times 10^{-2}$ within 1σ for $m_X = 100$ GeV.⁴ For $\text{Br}(K_L \rightarrow \mu^+ \mu^-)$, requiring that new physics contribution to dimension-six $\Delta F = 1$ operator $(\bar{s}_L \gamma^\mu d_L)(\bar{\mu} \gamma_\mu \mu)$ is smaller than SM contribution [68, 75], we impose the upper limit on the coupling $g_X < 4.61 \times 10^{-2}$ for $m_X = 100$ GeV. Focusing on the upper limit on the short-distance contribution $\text{Br}(K_L \rightarrow \mu^+ \mu^-)_{\text{SD}}$, a dedicate analysis gives $g_X|_{m_X=100 \text{ GeV}} < 3.55 \times 10^{-2}$ [76]. For the neutral Kaon mixing, the most stringent bound comes from the measurement of ϵ_K . Following the approach in Ref. [76], we get the allowed range on the coupling as $-4.13 \times 10^{-2} < g_X < 4.24 \times 10^{-2}$ for $m_X = 100$ GeV. We assume the left-handed coupling to X boson is dominant for flavor-violating vertices in the hadronic sector.

Such X boson has negligible effect on $(g-2)_\mu$, but gives modest contribution to ΔC_9 . Due to the negative value of $V_{ts} \simeq -0.40$, the X boson trends to the observed value ($C_9^{\mu\mu} = -0.40 \pm 0.12$). Consequently, the U_1 leptoquark in

⁴ Using the global fit $\delta C_9^\mu (= -\delta C_{10}^\mu) = -0.39 \pm 0.07$ based on Moriond 2021 result [74], the lower limit on the coupling is $g_X > 1.15 \times 10^{-2}$ within 1σ for $m_X = 100$ GeV.

conjunction with X boson explain the $R_{K^{(*)}}$ anomaly, but the former solely contributes to the $(g-2)_\mu$. Comparing to **Scan-2**, the $\chi^2_{\min} = 9.06$ at $(x_L^{22}, x_L^{23}, x_L^{32}, x_L^{33}, x_R^{32}, g_X) = (4.04 \times 10^{-2}, 0.324, -9.30 \times 10^{-3}, 0.810, 3.73 \times 10^{-4}, 1.47 \times 10^{-2})$ is no further reduced by including the X boson. However, the 2σ chi-square regions (yellow regions in Fig. 3) extend overlapping with $(g-2)_\mu$ 2σ region. Two representative points in the overlapped regions are

$$(x_L^{22}, x_L^{23}, x_L^{32}, x_L^{33}, x_R^{32}, g_X) = \begin{cases} (+3.28 \times 10^{-4}, 0.167, -0.846, 0.644, +0.520, 1.32 \times 10^{-2}), \\ (-3.08 \times 10^{-4}, 0.227, +0.831, 0.668, -0.554, 1.44 \times 10^{-2}), \end{cases}$$

respectively give $(\chi^2, \Delta a_\mu) = (13.9, 193 \times 10^{-11})$ and $(\chi^2, \Delta a_\mu) = (13.7, 201 \times 10^{-11})$. As a result, this hybrid scenario, U_1 leptoquark in conjunction with $U(1)_{B_3-L_2}$ gauge boson, explains the B -physics and $(g-2)_\mu$ anomalous within 2σ .

V. SUMMARY

The recent observational anomalies lead us to consider a vector leptoquark whose couplings with both left and right chiral fermions are essential. It affects various channels of B -meson decays and generate lepton flavor universality breaking. At the same time, the leptoquark can contribute to $(g-2)_\mu$ with significant enhancement. We perform the global analysis to several low-energy observables and encounter both left- and right-handed U_1 leptoquark couplings. Unfortunately, we have not found common parameter region for B and $(g-2)_\mu$ anomalies without additional muonphilic vector boson. Motivated by this, we found the U_1 leptoquark in conjunction with additional $U(1)_{B_3-L_2}$ gauge boson is able to reconcile the B -physics and $(g-2)_\mu$ anomalies within 2σ . We expect the experimental measurements will be much more improved in the future and the leptoquark and the new gauge boson will be better tested accordingly.

ACKNOWLEDGMENTS

The work is supported by the National Research Foundation of Korea NRF-2021R1A4A2001897, NRF-2019R1A2C1089334 (SCP), and NRF-2020R1I1A1A01066413 (PYT). We thank Andreas Crivellin for discussion on the bottom quark mass in the computation of Muon $(g-2)$. SCP is thankful to Michael Peskin for inviting to the symposium on the muon $g-2$ anomaly and discussion. SCP and YJK also acknowledges the support from Yonsei U. and KIAS during his sabbatical period. We acknowledge the hospitality at APCTP where part of this work was done.

-
- [1] HFLAV collaboration, Y. S. Amhis et al., *Averages of b -hadron, c -hadron, and τ -lepton properties as of 2018*, **1909.12524**.
 - [2] BABAR collaboration, J. P. Lees et al., *Evidence for an excess of $\bar{B} \rightarrow D^{(*)}\tau^-\bar{\nu}_\tau$ decays*, *Phys. Rev. Lett.* **109** (2012) 101802, [1205.5442].
 - [3] BELLE collaboration, M. Huschle et al., *Measurement of the branching ratio of $\bar{B} \rightarrow D^{(*)}\tau^-\bar{\nu}_\tau$ relative to $\bar{B} \rightarrow D^{(*)}\ell^-\bar{\nu}_\ell$ decays with hadronic tagging at Belle*, *Phys. Rev.* **D92** (2015) 072014, [1507.03233].
 - [4] BELLE collaboration, S. Hirose et al., *Measurement of the τ lepton polarization and $R(D^*)$ in the decay $\bar{B} \rightarrow D^*\tau^-\bar{\nu}_\tau$* , *Phys. Rev. Lett.* **118** (2017) 211801, [1612.00529].
 - [5] BELLE collaboration, G. Caria et al., *Measurement of $\mathcal{R}(D)$ and $\mathcal{R}(D^*)$ with a semileptonic tagging method*, *Phys. Rev. Lett.* **124** (2020) 161803, [1910.05864].
 - [6] LHCb collaboration, R. Aaij et al., *Measurement of the ratio of branching fractions $\mathcal{B}(\bar{B}^0 \rightarrow D^{*+}\tau^-\bar{\nu}_\tau)/\mathcal{B}(\bar{B}^0 \rightarrow D^{*+}\mu^-\bar{\nu}_\mu)$* , *Phys. Rev. Lett.* **115** (2015) 111803, [1506.08614].
 - [7] LHCb collaboration, R. Aaij et al., *Measurement of the ratio of the $B^0 \rightarrow D^{*-}\tau^+\nu_\tau$ and $B^0 \rightarrow D^{*-}\mu^+\nu_\mu$ branching fractions using three-prong τ -lepton decays*, *Phys. Rev. Lett.* **120** (2018) 171802, [1708.08856].

- [8] W. Altmannshofer, S. Gori, H. H. Patel, S. Profumo and D. Tuckler, *Electric dipole moments in a leptoquark scenario for the B-physics anomalies*, *JHEP* **05** (2020) 069, [2002.01400].
- [9] LHCb collaboration, R. Aaij et al., *Test of lepton universality in beauty-quark decays*, **2103.11769**.
- [10] LHCb collaboration, R. Aaij et al., *Test of lepton universality with $B^0 \rightarrow K^{*0} \ell^+ \ell^-$ decays*, *JHEP* **08** (2017) 055, [1705.05802].
- [11] BELLE collaboration, A. Abdesselam et al., *Test of lepton flavor universality in $B \rightarrow K^* \ell^+ \ell^-$ decays at Belle*, **1904.02440**.
- [12] BELLE collaboration, S. Choudhury et al., *Test of lepton flavor universality and search for lepton flavor violation in $B \rightarrow K \ell \ell$ decays*, *JHEP* **03** (2021) 105, [1908.01848].
- [13] E. Kou, P. Urquijo, W. Altmannshofer, F. Beaujean, G. Bell, M. Beneke et al., *The Belle II Physics Book, Progress of Theoretical and Experimental Physics* **2019** (12, 2019) , [<https://academic.oup.com/ptep/article-pdf/2019/12/123C01/32693980/ptz106.pdf>].
- [14] MUON $g - 2$ collaboration, B. Abi, T. Albahri, S. Al-Kilani, D. Allspach, L. P. Alonzi, A. Anastasi et al., *Measurement of the positive muon anomalous magnetic moment to 0.46 ppm*, *Phys. Rev. Lett.* **126** (Apr, 2021) 141801.
- [15] MUON G-2 collaboration, G. W. Bennett et al., *Final Report of the Muon E821 Anomalous Magnetic Moment Measurement at BNL*, *Phys. Rev.* **D73** (2006) 072003, [hep-ex/0602035].
- [16] PARTICLE DATA GROUP collaboration, P. A. Zyla et al., *Review of Particle Physics*, *PTEP* **2020** (2020) 083C01.
- [17] S. Borsanyi et al., *Leading hadronic contribution to the muon magnetic moment from lattice QCD*, *Nature* **593** (2021) 51–55, [2002.12347].
- [18] Y. Jho, Y. Kwon, S. C. Park and P.-Y. Tseng, *Search for muon-philic new light gauge boson at Belle II*, *JHEP* **10** (2019) 168, [1904.13053].
- [19] Y. Jho, S. M. Lee, S. C. Park, Y. Park and P.-Y. Tseng, *Light gauge boson interpretation for $(g - 2)$ and the $K_L \rightarrow \pi^0 +$ (invisible) anomaly at the J-PARC KOTO experiment*, *JHEP* **04** (2020) 086, [2001.06572].
- [20] K. Ban, Y. Jho, Y. Kwon, S. C. Park, S. Park and P.-Y. Tseng, *Search for new light vector boson using J/Ψ at BESIII and Belle II*, **2012.04190**.
- [21] Y. Jho, J.-C. Park, S. C. Park and P.-Y. Tseng, *Leptonic New Force and Cosmic-ray Boosted Dark Matter for the XENON1T Excess*, *Phys. Lett.* **B811** (2020) 135863, [2006.13910].
- [22] S. C. Park and H. S. Song, *Brane fluctuation and anomalous muon magnetic moment*, *Phys. Lett.* **B523** (2001) 161–164, [hep-ph/0109121].
- [23] S. C. Park and H. S. Song, *Muon anomalous magnetic moment and the stabilized Randall-Sundrum scenario*, *Phys. Lett.* **B506** (2001) 99–102, [hep-ph/0103072].
- [24] S. Kanemura, K. Nishiwaki, H. Okada, Y. Orikasa, S. C. Park and R. Watanabe, *LHC 750 GeV diphoton excess in a radiative seesaw model*, *PTEP* **2016** (2016) 123B04, [1512.09048].
- [25] B. Zhu and X. Liu, *Probing light dark matter with scalar mediator: muon $(g - 2)$ deviation, the proton radius puzzle*, **2104.03238**.
- [26] P. Athron, C. Balázs, D. H. Jacob, W. Kotlarski, D. Stöckinger and H. Stöckinger-Kim, *New physics explanations of a_μ in light of the FNAL muon $g - 2$ measurement*, **2104.03691**.
- [27] A. Crivellin, D. Müller and F. Saturnino, *Flavor Phenomenology of the Leptoquark Singlet-Triplet Model*, *JHEP* **06** (2020) 020, [1912.04224].
- [28] M. Lindner, M. Platscher and F. S. Queiroz, *A Call for New Physics : The Muon Anomalous Magnetic Moment and Lepton Flavor Violation*, *Phys. Rept.* **731** (2018) 1–82, [1610.06587].
- [29] H. Terazawa, *All the hadronic contributions to the anomalous magnetic moment of the muon and the lamb shift in the hydrogen atom*, *Prog. Theor. Phys.* **39** (1968) 1326–1332.
- [30] A. Angelescu, D. Bečirević, D. A. Faroughy and O. Sumensari, *Closing the window on single leptoquark solutions to the B-physics anomalies*, *JHEP* **10** (2018) 183, [1808.08179].
- [31] D. Buttazzo, A. Greljo, G. Isidori and D. Marzocca, *B-physics anomalies: a guide to combined explanations*, *JHEP* **11** (2017) 044, [1706.07808].
- [32] J. Aebischer, W. Altmannshofer, D. Guadagnoli, M. Reboud, P. Stangl and D. M. Straub, *B-decay discrepancies after Moriond 2019*, *Eur. Phys. J.* **C80** (2020) 252, [1903.10434].
- [33] C. Cornella, J. Fuentes-Martin and G. Isidori, *Revisiting the vector leptoquark explanation of the B-physics anomalies*, *JHEP* **07** (2019) 168, [1903.11517].
- [34] N. Assad, B. Fornal and B. Grinstein, *Baryon Number and Lepton Universality Violation in Leptoquark and Diquark Models*, *Phys. Lett.* **B777** (2018) 324–331, [1708.06350].

- [35] R. Alonso, B. Grinstein and J. Martin Camalich, *Lepton universality violation and lepton flavor conservation in B -meson decays*, *JHEP* **10** (2015) 184, [1505.05164].
- [36] L. Calibbi, A. Crivellin and T. Ota, *Effective Field Theory Approach to $b \rightarrow s\ell\ell^{(\prime)}$, $B \rightarrow K^{(*)}\nu\bar{\nu}$ and $B \rightarrow D^{(*)}\tau\nu$ with Third Generation Couplings*, *Phys. Rev. Lett.* **115** (2015) 181801, [1506.02661].
- [37] A. Bhaskar, D. Das, T. Mandal, S. Mitra and C. Neeraj, *Precise limits on the charge-2/3 U_1 vector leptoquark*, 2101.12069.
- [38] P. S. Bhupal Dev, R. Mohanta, S. Patra and S. Sahoo, *Unified explanation of flavor anomalies, radiative neutrino masses, and ANITA anomalous events in a vector leptoquark model*, *Phys. Rev.* **D102** (2020) 095012, [2004.09464].
- [39] M. Blanke and A. Crivellin, *B Meson Anomalies in a Pati-Salam Model within the Randall-Sundrum Background*, *Phys. Rev. Lett.* **121** (2018) 011801, [1801.07256].
- [40] A. Crivellin, C. Greub, D. Müller and F. Saturnino, *Importance of Loop Effects in Explaining the Accumulated Evidence for New Physics in B Decays with a Vector Leptoquark*, *Phys. Rev. Lett.* **122** (2019) 011805, [1807.02068].
- [41] C. Bonilla, T. Modak, R. Srivastava and J. W. F. Valle, *$U(1)_{B_3-3L_\mu}$ gauge symmetry as a simple description of $b \rightarrow s$ anomalies*, *Phys. Rev. D* **98** (2018) 095002, [1705.00915].
- [42] R. Alonso, P. Cox, C. Han and T. T. Yanagida, *Flavoured $B-L$ local symmetry and anomalous rare B decays*, *Phys. Lett.* **B774** (2017) 643–648, [1705.03858].
- [43] B. C. Allanach, *$U(1)_{B_3-L_2}$ explanation of the neutral current B -anomalies*, *Eur. Phys. J. C* **81** (2021) 56, [2009.02197].
- [44] M. Du, J. Liang, Z. Liu and V. Q. Tran, *A vector leptoquark interpretation of the muon $g-2$ and B anomalies*, 2104.05685.
- [45] PARTICLE DATA GROUP collaboration, R. L. Workman and Others, *Review of Particle Physics*, *PTEP* **2022** (2022) 083C01.
- [46] W. J. Marciano, A. Masiero, P. Paradisi and M. Passera, *Contributions of axionlike particles to lepton dipole moments*, *Phys. Rev.* **D94** (2016) 115033, [1607.01022].
- [47] M. Algueró, B. Capdevila, A. Crivellin, S. Descotes-Genon, P. Masjuan, J. Matias et al., *Emerging patterns of New Physics with and without Lepton Flavour Universal contributions*, *Eur. Phys. J. C* **79** (2019) 714, [1903.09578].
- [48] A. G. Akeroyd and C.-H. Chen, *Constraint on the branching ratio of $B_c \rightarrow \tau\bar{\nu}$ from LEP1 and consequences for $R(D^{(*)})$ anomaly*, *Phys. Rev.* **D96** (2017) 075011, [1708.04072].
- [49] R. Alonso, B. Grinstein and J. Martin Camalich, *Lifetime of B_c^- Constrains Explanations for Anomalies in $B \rightarrow D^{(*)}\tau\nu$* , *Phys. Rev. Lett.* **118** (2017) 081802, [1611.06676].
- [50] M. Blanke, A. Crivellin, T. Kitahara, M. Moscati, U. Nierste and I. Nišandžić, *Addendum to “Impact of polarization observables and $B_c \rightarrow \tau\nu$ on new physics explanations of the $b \rightarrow c\tau\nu$ anomaly”*, 1905.08253.
- [51] LHCb collaboration, R. Aaij et al., *Search for the decays $B_s^0 \rightarrow \tau^+\tau^-$ and $B^0 \rightarrow \tau^+\tau^-$* , *Phys. Rev. Lett.* **118** (2017) 251802, [1703.02508].
- [52] C. Bobeth, M. Gorbahn, T. Hermann, M. Misiak, E. Stamou and M. Steinhauser, *$B_{s,d} \rightarrow l^+l^-$ in the Standard Model with Reduced Theoretical Uncertainty*, *Phys. Rev. Lett.* **112** (2014) 101801, [1311.0903].
- [53] W. Altmannshofer, P. Ball, A. Bharucha, A. J. Buras, D. M. Straub and M. Wick, *Symmetries and Asymmetries of $B \rightarrow K^*\mu^+\mu^-$ Decays in the Standard Model and Beyond*, *JHEP* **01** (2009) 019, [0811.1214].
- [54] K. Cheung, C.-W. Chiang, N. G. Deshpande and J. Jiang, *Constraints on flavor-changing Z' models by $B(s)$ mixing, Z' production, and $B(s) \rightarrow \mu\nu + \mu\nu$* , *Phys. Lett. B* **652** (2007) 285–291, [hep-ph/0604223].
- [55] LHCb collaboration, R. Aaij et al., *Search for the lepton-flavour-violating decays $B_s^0 \rightarrow \tau^\pm\mu^\mp$ and $B^0 \rightarrow \tau^\pm\mu^\mp$* , *Phys. Rev. Lett.* **123** (2019) 211801, [1905.06614].
- [56] L. Calibbi and G. Signorelli, *Charged Lepton Flavour Violation: An Experimental and Theoretical Introduction*, *Riv. Nuovo Cim.* **41** (2018) 71–174, [1709.00294].
- [57] C. McNeile, C. T. H. Davies, E. Follana, K. Hornbostel and G. P. Lepage, *High-Precision f_{B_s} and HQET from Relativistic Lattice QCD*, *Phys. Rev.* **D85** (2012) 031503, [1110.4510].
- [58] BABAR collaboration, J. P. Lees et al., *Search for $B^+ \rightarrow K^+\tau^+\tau^-$ at the BaBar experiment*, *Phys. Rev. Lett.* **118** (2017) 031802, [1605.09637].
- [59] BABAR collaboration, J. P. Lees et al., *A search for the decay modes $B^{+-} \rightarrow h^{+-}\tau^{+-}l$* , *Phys. Rev.* **D86** (2012) 012004, [1204.2852].
- [60] D. Bečirević, O. Sumensari and R. Zukanovich Funchal, *Lepton flavor violation in exclusive $b \rightarrow s$ decays*, *Eur. Phys. J. C* **76** (2016) 134, [1602.00881].
- [61] HFLAV collaboration, Y. Amhis et al., *Averages of b -hadron, c -hadron, and τ -lepton properties as of summer 2016*, *Eur. Phys. J. C* **77** (2017) 895, [1612.07233].

- [62] BELLE collaboration, Y. Miyazaki et al., *Search for Lepton-Flavor-Violating tau Decays into a Lepton and a Vector Meson*, *Phys. Lett.* **B699** (2011) 251–257, [1101.0755].
- [63] M. Bordone, C. Cornella, J. Fuentes-Martín and G. Isidori, *Low-energy signatures of the PS³ model: from B-physics anomalies to LFV*, *JHEP* **10** (2018) 148, [1805.09328].
- [64] C. Hati, J. Kriewald, J. Orloff and A. M. Teixeira, *A nonunitary interpretation for a single vector leptoquark combined explanation to the B-decay anomalies*, *JHEP* **12** (2019) 006, [1907.05511].
- [65] B. Bhattacharya, A. Datta, J.-P. Guévin, D. London and R. Watanabe, *Simultaneous Explanation of the R_K and $R_{D^{(*)}}$ Puzzles: a Model Analysis*, *JHEP* **01** (2017) 015, [1609.09078].
- [66] W. Altmannshofer, S. Gori, M. Pospelov and I. Yavin, *Neutrino Trident Production: A Powerful Probe of New Physics with Neutrino Beams*, *Phys. Rev. Lett.* **113** (2014) 091801, [1406.2332].
- [67] L. Di Luzio, M. Kirk, A. Lenz and T. Rauh, *ΔM_s theory precision confronts flavour anomalies*, *JHEP* **12** (2019) 009, [1909.11087].
- [68] A. J. Buras and R. Fleischer, *Quark mixing, CP violation and rare decays after the top quark discovery*, *Adv. Ser. Direct. High Energy Phys.* **15** (1998) 65–238, [hep-ph/9704376].
- [69] ATLAS, CMS collaboration, F. Pandolfi, *Searches for New Heavy Resonances in Final States with Leptons and Photons in ATLAS and CMS*, 9, 2017, 1709.06119.
- [70] CMS collaboration, A. M. Sirunyan et al., *Search for electroweak production of charginos and neutralinos in multilepton final states in proton-proton collisions at $\sqrt{s} = 13$ TeV*, *JHEP* **03** (2018) 166, [1709.05406].
- [71] CMS collaboration, A. M. Sirunyan et al., *Search for resonant and nonresonant new phenomena in high-mass dilepton final states at $\sqrt{s} = 13$ TeV*, *JHEP* **07** (2021) 208, [2103.02708].
- [72] HFLAV collaboration, Y. Amhis et al., *Averages of b-hadron, c-hadron, and τ -lepton properties as of summer 2016*, *Eur. Phys. J. C* **77** (2017) 895, [1612.07233].
- [73] HFLAV collaboration, Y. Amhis et al., *Averages of b-hadron, c-hadron, and τ -lepton properties as of 2021*, 2206.07501.
- [74] W. Altmannshofer and P. Stangl, *New physics in rare B decays after Moriond 2021*, *Eur. Phys. J. C* **81** (2021) 952, [2103.13370].
- [75] G. D’Ambrosio, A. M. Iyer, F. Mahmoudi and S. Neshatpour, *Anatomy of kaon decays and prospects for lepton flavour universality violation*, *JHEP* **09** (2022) 148, [2206.14748].
- [76] M. Endo, T. Kitahara, S. Mishima and K. Yamamoto, *Revisiting Kaon Physics in General Z Scenario*, *Phys. Lett. B* **771** (2017) 37–44, [1612.08839].