

# RIGID MANIFOLDS OF GENERAL TYPE WITH NON-CONTRACTIBLE UNIVERSAL COVER

DAVIDE FRAPPORTI, CHRISTIAN GLEISSNER

ABSTRACT. For each  $n \geq 3$  we give examples of infinitesimally rigid projective manifolds of general type of dimension  $n$  with non-contractible universal cover. We provide examples with projective and examples with non-projective universal cover.

## INTRODUCTION

In [BC18] several notions of *rigidity* have been discussed, the relations among them have been studied and many questions and conjectures have been proposed. In particular the authors showed that a rigid compact complex surface has Kodaira dimension  $-\infty$  or 2, and observed that all known examples of rigid surfaces of general type are  $K(\pi, 1)$  spaces. Recall that a CW complex with fundamental group  $\pi$  is called  $K(\pi, 1)$  *space* if its universal cover is contractible, and that these spaces have the property that their homotopy type is uniquely determined by their fundamental group (cf. [Hat02, §1.B]). This implies that the topological invariants, such as homology and cohomology, are determined by  $\pi$ . In [BC18] the following natural question has been posed.

**Question.** *Do there exist infinitesimally rigid surfaces of general type with non-contractible universal cover?*

The aim of this paper is to give a positive answer for the analogous question in higher dimensions. More precisely, we construct for each  $n \geq 3$  an infinitesimally rigid manifold of general type of dimension  $n$  with non-contractible universal cover. For surfaces the question remains open. We recall now the notions of rigidity that are relevant for our purposes.

### Definition 1.

Let  $X$  be a compact complex manifold of dimension  $n$ .

- (1) A *deformation of  $X$*  is a proper smooth holomorphic map of pairs

$$f: (\mathfrak{X}, X) \rightarrow (\mathcal{B}, b_0),$$

where  $(\mathcal{B}, b_0)$  is a connected (possibly not reduced) germ of a complex space.

- (2)  $X$  is said to be *rigid* if for each deformation of  $X$ ,  $f: (\mathfrak{X}, X) \rightarrow (\mathcal{B}, b_0)$  there is an open neighbourhood  $U \subset \mathcal{B}$  of  $b_0$  such that  $X_t := f^{-1}(t) \simeq X$  for all  $t \in U$ .

---

*Date:* March 8, 2023.

*2010 Mathematics Subject Classification.* 32G05, 14J10, 14L30, 14J40, 32Q30, 14B05.

*Key words and phrases.* Rigid complex manifolds, deformation theory, fundamental group, classifying space.

The authors thank I. Bauer, F. Catanese, S. Coughlan and F. Fallucca for their interest and fruitful conversations.

The first author is member of G.N.S.A.G.A. of I.N.d.A.M. and acknowledges support of the ERC Advanced grant n. 340258-TADMICAMT.

- (3)  $X$  is said to be *infinitesimally rigid* if  $H^1(X, \Theta_X) = 0$ , where  $\Theta_X$  is the sheaf of holomorphic vector fields on  $X$ .
- (4)  $X$  is said to be (infinitesimally) étale rigid if all finite étale covers  $f: Y \rightarrow X$  are (infinitesimally) rigid.

*Remark 2.* i) By Kodaira-Spencer-Kuranishi theory every infinitesimally rigid manifold is rigid. The converse does not hold in general as it was shown in [BP18] and [BGP20] (cf. also [MK71]).

ii) Beauville surfaces are examples of rigid, but not étale rigid manifolds (see [Cat00]).

Both the examples constructed in [BP18] and Beauville surfaces are product quotient varieties, i.e. (resolutions of singularities of) finite quotients of product of curves with respect to a holomorphic group action. In recent years, product quotients turned out to be a very fruitful source of examples of rigid complex manifolds with additional properties. Besides the examples above, we mention [BG20], where the authors construct the first examples of rigid complex manifolds with Kodaira dimension 1 in arbitrary dimension  $n \geq 3$ , and [BG21] where they constructed new rigid three- and four-folds with Kodaira dimension 0. We refer to [CF18, FGP20, FG20, GPR18, LP16, LP20] for other interesting examples of product quotient varieties.

The manifolds we construct are also product quotients. More precisely, inspired by the construction in [BP18] in Section 1 we consider for each  $n \geq 3$  and  $d \geq 4$ , even and not divisible by 3 the  $n$ -fold product  $C^n$  of the Fermat curve  $C$  of degree  $d$  together with a suitable action of  $\mathbb{Z}_d^2$ . The quotient  $X_{n,d} := C^n / \mathbb{Z}_d^2$  is a normal projective variety with isolated cyclic quotient singularities of type  $\frac{1}{2}(1, \dots, 1)$ , Kodaira dimension  $n$  and

$$H^1(X_{n,d}, \Theta_{X_{n,d}}) = H^1(C^n, \Theta_{C^n})^{\mathbb{Z}_d^2} = 0.$$

Blowing up the singular points, we obtain a resolution  $\widehat{X}_{n,d} \rightarrow X_{n,d}$  such that  $H^1(X_{n,d}, \Theta_{X_{n,d}}) = H^1(\widehat{X}_{n,d}, \Theta_{\widehat{X}_{n,d}})$ . Therefore,  $\widehat{X}_{n,d}$  is an infinitesimally rigid projective manifold of general type.

In Section 2 we show that the universal cover  $U_{n,d}$  of  $\widehat{X}_{n,d}$  is non-contractible since it contains several  $\mathbb{P}^{n-1}$  (see Proposition 10). We then discuss the finiteness of the fundamental group  $\pi_1(X_{n,d}) = \pi_1(\widehat{X}_{n,d})$ . The crucial ingredient here is Armstrong's description of the fundamental group of a quotient space [Arm68] adapted to product quotients by [BCGP12]. The finiteness of  $\pi_1(\widehat{X}_n)$  is equivalent to the finiteness of certain groups (Proposition 16: Finiteness criterion). This allows us to prove the following.

**Theorem 3.** *For each  $n \geq 3$ ,  $d \geq 4$ , even and not divisible by 3 there exists an infinitesimally rigid projective  $n$ -dimensional manifold of general type  $\widehat{X}_{n,d}$ , whose universal cover  $U_{n,d}$  is non-contractible. Moreover, the universal cover  $U_{n,d}$  is projective if and only if  $d = 4$ .*

The construction actually works also for  $n = 2$ : the surface  $\widehat{X}_{2,4}$  is not rigid, whereas the surface  $\widehat{X}_{2,d}$  for  $d \geq 8$  is rigid but not infinitesimally rigid (see [BP18]), and its universal cover is non-contractible.

**Notation.** We work over the field of complex numbers, and we denote by  $\mathbb{Z}_n$  the cyclic group of order  $n$  and by  $\zeta_n$  a primitive  $n$ -th root of unity. The rest of the notation is standard in complex algebraic geometry.

## 1. THE FAMILIES

Let  $C_d := \{x_0^d + x_1^d + x_2^d = 0\} \subset \mathbb{P}^2$  be the Fermat curve of degree  $d$ . Consider the group action

$$\phi_1: \mathbb{Z}_d^2 \rightarrow \text{Aut}(C), \quad (a, b) \mapsto [(x_0 : x_1 : x_2) \mapsto (\zeta_d^a x_0 : \zeta_d^b x_1 : x_2)].$$

There are  $3d$  points on  $C_d$  with non-trivial stabilizer. They form three orbits of length  $d$ . A representative of each orbit and a generator of the corresponding stabilizer is given in the table below:

point	$(0 : 1 : \zeta_{2d})$	$(1 : 0 : \zeta_{2d})$	$(1 : \zeta_{2d} : 0)$
generator	$(1, 0)$	$(0, 1)$	$(1, 1)$

Hence the quotient map

$$f: C_d \rightarrow \mathbb{P}^1, \quad (x_0 : x_1 : x_2) \mapsto (x_0^d : x_1^d)$$

is branched in  $(0 : 1)$ ,  $(1 : 0)$  and  $(1 : -1)$ , each with branch index  $d$ .

**1.1. The singular quotients  $X_{n,d}$ .** From now on we fix  $d \geq 4$ , even and not divisible by 3, and denote  $C_d$  simply by  $C$ . Let  $A$  be the automorphism of  $\mathbb{Z}_d^2$  given by the matrix

$$\begin{pmatrix} 1 & -2 \\ 2 & -1 \end{pmatrix} \in \text{GL}(2, \mathbb{Z}_d),$$

and let  $\phi_2 := \phi_1 \circ A^{-1}$ . For each  $n \geq 2$  consider the  $\mathbb{Z}_d^2$  diagonal action on  $C^n$  defined by

$$g(z_1, \dots, z_n) := (\phi_1(g) \cdot z_1, \phi_2(g) \cdot z_2, \phi_2(g) \cdot z_3, \dots, \phi_2(g) \cdot z_n)$$

and let  $X_{n,d}$  be the quotient variety  $X_{n,d} := C^n / \mathbb{Z}_d^2$ .

*Remark 4.* The diagonal action is not free, indeed

$$\text{Fix}(\phi_1(g)) \neq \emptyset \text{ and } \text{Fix}(\phi_2(g)) \neq \emptyset \iff g \in H := \left\langle \left( \frac{d}{2}, 0 \right), \left( 0, \frac{d}{2} \right) \right\rangle.$$

Noting that  $\phi_1|_H = \phi_2|_H$ , we see that a point  $(z_1, \dots, z_n) \in C^n$  has a non-trivial stabilizer if and only if all its coordinates  $z_i$  belong to one and only one of the three  $\mathbb{Z}_d^2$ -orbits displayed in the table above.

**Proposition 5.** *For  $n \geq 3$  the projective variety  $X_{n,d}$  is infinitesimally rigid and of general type. The singular locus of  $X_{n,d}$  consists of  $6 \cdot d^{n-2}$  cyclic quotient singularities of type  $\frac{1}{2}(1, \dots, 1)$ .*

*Proof.* By Remark 4 there are  $3 \cdot d^n$  points on  $C^n$  with non-trivial stabilizer, each generated by one of the order 2 elements in  $\mathbb{Z}_d^2$ . Thus,  $X_{n,d}$  has  $(3 \cdot d^n) / (d^2/2) = 6 \cdot d^{n-2}$  singularities of type  $\frac{1}{2}(1, \dots, 1)$ .

These singularities are terminal if  $n \geq 3$ , see [Rei87, p. 376 Theorem]. Since the quotient map  $C^n \rightarrow X_{n,d}$  is quasi-étale,  $g(C) = (d-1)(d-2)/2 \geq 3$  and  $X_{n,d}$  is terminal, its Kodaira dimension is  $\kappa(X_{n,d}) = \kappa(C^n) = n$  (cf. [Cat07, p. 51]).

According to Schlessinger [Sch71], isolated quotient singularities in dimension at least three are rigid, i.e.  $\mathcal{E}xt^1(\Omega_{X_{n,d}}^1, \mathcal{O}_{X_{n,d}}) = 0$ . Thus the local-to-global Ext spectral sequence yields

$$H^1(X_{n,d}, \Theta_{X_{n,d}}) \simeq \text{Ext}^1(\Omega_{X_{n,d}}^1, \mathcal{O}_{X_{n,d}}).$$

Hence it suffices to verify that  $X_{n,d}$  has no equisingular deformations. Since  $g(C) \geq 3$  we have  $H^0(C, \Theta_C) = 0$ , hence by Künneth formula we get

$$H^1(C^n, \Theta_{C^n}) = \bigoplus_{i=1}^n H^1(C, \Theta_C).$$

Using the fact that the quotient map  $C^n \rightarrow X_{n,d}$  is quasi-étale and the action is diagonal, we obtain

$$H^1(X_{n,d}, \Theta_{X_{n,d}}) = H^1(C^n, \Theta_{C^n})^{\mathbb{Z}_d^2} = \bigoplus_{i=1}^n H^1(C, \Theta_C)^{\mathbb{Z}_d^2}.$$

The branch locus  $B$  of  $f: C \rightarrow C/\mathbb{Z}_d^2 \simeq \mathbb{P}^1$  consists of 3 points  $p_i$  with branch indices  $m_{p_i} = d$ , thus by [Bea78, Ex. VI.12] we have

$$\begin{aligned} \dim H^1(C, \Theta_C)^{\mathbb{Z}_d^2} &= \dim H^0(C, 2K_C)^{\mathbb{Z}_d^2} = h^0(\mathbb{P}^1, 2K_{\mathbb{P}^1} + \sum_{p_i \in B} p_i \cdot [2(1 - \frac{1}{m_{p_i}})]) \\ &= h^0(\mathbb{P}^1, \mathcal{O}(-1)) = 0. \end{aligned}$$

□

## 1.2. Resolution of singularities of type $\frac{1}{2}(1, \dots, 1)$ .

**Proposition 6.** *A singularity  $U := \mathbb{C}^n/\mathbb{Z}_2$  of type  $\frac{1}{2}(1, \dots, 1)$  admits a resolution  $\rho: \widehat{U} \rightarrow U$  by a single blow-up, with exceptional prime divisor  $\mathbb{P}^{n-1}$ . If  $n \geq 3$ ,*

$$\rho_* \Theta_{\widehat{U}} = \Theta_U \quad \text{and} \quad R^1 \rho_* \Theta_{\widehat{U}} = 0.$$

For a proof we refer to [Sch71, proof of Theorem 4], see also [BG20, Corollary 5.9, Proposition 5.10].

*Remark 7* (see [BG20, Remark 5.4]). Both properties are not obvious and in general even false. For any resolution  $\rho: Z' \rightarrow Z$  of a normal variety  $Z$ , the direct image  $\rho_* \Theta_{Z'}$  is a subsheaf of the reflexive sheaf  $\Theta_Z$ , and this inclusion is in general strict: e.g. take the blow-up of the origin of  $\mathbb{C}^2$ .

The vanishing of  $R^1 \rho_* \Theta_{Z'}$  is also not automatic: take the resolution of an  $A_1$  surface singularity (i.e.  $\frac{1}{2}(1, 1)$ ) by a  $-2$  curve, then  $R^1 \rho_* \Theta_{Z'}$  is a skyscraper sheaf at the singular point with value  $H^1(\mathbb{P}^1, \mathcal{O}(-2)) \cong \mathbb{C}$ . More generally, for canonical ADE surface singularities  $R^1 \rho_* \Theta_{Z'}$  is never zero, cf. [BW74, Pin81, Sch71].

**Corollary 8.** *Let  $Z_n$  be a projective variety of dimension  $n \geq 3$  with only singularities of type  $\frac{1}{2}(1, \dots, 1)$ . Then there exists a resolution  $\rho: \widehat{Z}_n \rightarrow Z_n$ , such that*

$$H^1(Z_n, \Theta_{Z_n}) \simeq H^1(\widehat{Z}_n, \Theta_{\widehat{Z}_n}).$$

*In particular, if  $Z_n$  is infinitesimally rigid, so is  $\widehat{Z}_n$ .*

*Proof.* Since the singularities of  $Z_n$  are isolated, we resolve them simultaneously using Proposition 6 and we get a resolution  $\rho: \widehat{Z}_n \rightarrow Z_n$  having the same properties:

$$\rho_* \Theta_{\widehat{Z}_n} = \Theta_{Z_n} \quad \text{and} \quad R^1 \rho_* \Theta_{\widehat{Z}_n} = 0.$$

Leray's spectral sequence implies  $H^1(\widehat{Z}_n, \Theta_{\widehat{Z}_n}) \simeq H^1(Z_n, \Theta_{Z_n})$ . □

By the corollary, for  $n \geq 3$  there exists a resolution  $\widehat{X}_{n,d} \rightarrow X_{n,d}$  of the singularities of  $X_{n,d}$ , which is infinitesimally rigid. By Remark 7, for  $n = 2$  the minimal resolution  $\widehat{X}_{2,d}$  of  $X_{2,d}$  is not infinitesimally

rigid, nevertheless the main theorem of [BP18] shows that  $\widehat{X}_{2,d}$  is rigid for  $d \geq 8$ , whereas  $\widehat{X}_{2,4}$  is a numerical Campedelli surface, whose Kuranishi family has dimension 6.

**1.3. Non-étale infinitesimally rigidity.** We conclude this section constructing an étale cover of  $\widehat{X}_{n,d}$  which is not infinitesimally rigid, thus  $\widehat{X}_{n,d}$  is not étale infinitesimally rigid.

Let  $H := \langle (\frac{d}{2}, 0), (0, \frac{d}{2}) \rangle$  be as in Remark 4.

**Lemma 9.** *Let  $Y_{n,d} := C^n/H$  be the quotient with respect to the restricted diagonal action, then:*

- (1) *The natural morphism  $\psi: Y_{n,d} \rightarrow X_{n,d}$  is an unramified Galois cover with group  $\mathbb{Z}_d^2/H$ .*
- (2)  $h^1(Y_{n,d}, \Theta_{Y_{n,d}}) = 3n \cdot \left(\frac{d^2-2d}{8}\right)$ .

*Proof.* (1) Since  $H$  is a normal subgroup of  $\mathbb{Z}_d^2$  the map  $\psi$  is a Galois cover with group  $\mathbb{Z}_d^2/H \cong \mathbb{Z}_{d/2}^2$ . By Remark 4 the stabilizer of a point  $z \in C^n$  with respect to the  $\mathbb{Z}_d^2$ -action is contained in  $H$ , whence the map  $\psi$  is unramified.

(2) Since  $C \rightarrow C/H$  is branched in  $\frac{3d}{2}$  points and  $g(C/H) = \frac{(d-2)(d-4)}{8}$ , we have

$$\dim(H^1(C^n, \Theta_{C^n})^H) = n \cdot \dim(H^1(C, \Theta_C)^H) = 3n \left(\frac{d^2-2d}{8}\right)$$

arguing as in Proposition 5. □

## 2. THE UNIVERSAL COVER OF $\widehat{X}_{n,d}$

In this section we prove that the universal cover  $U_{n,d}$  of  $\widehat{X}_{n,d}$  is non-contractible, and then we discuss whether it is projective or not.

**Proposition 10.** *Let  $X$  be a compact Kähler manifold, containing a  $\mathbb{P}^m$ . Then the universal cover  $U$  of  $X$  is non-contractible.*

*Proof.* Since  $\mathbb{P}^m$  is simply connected, the inclusion map  $i: \mathbb{P}^m \hookrightarrow X$  lifts to a map  $f: \mathbb{P}^m \rightarrow U$ . Looking for a contradiction, assume that  $U$  is contractible, then  $f$  is homotopic to a constant map, therefore the inclusion  $i$  is also homotopic to a constant map. In particular we see that the induced linear map  $i^*: H^2(X, \mathbb{C}) \rightarrow H^2(\mathbb{P}^m, \mathbb{C})$  is the zero map. Now let  $[\omega]$  be a Kähler class of  $X$ . Its restriction  $i^*([\omega])$  is a Kähler class of  $\mathbb{P}^m$ , whence non zero, contradiction. □

**Corollary 11.** *The universal cover  $U_{n,d}$  of  $\widehat{X}_{n,d}$  is non-contractible.*

*Proof.* By Proposition 6  $\widehat{X}_{n,d}$  contains several  $\mathbb{P}^{n-1}$ . □

*Remark 12.* By Lemma 9 the universal cover  $U_{n,d}$  of  $\widehat{X}_{n,d}$  is not infinitesimally rigid.

### 2.1. The Fundamental Group.

In this section we discuss the finiteness of the fundamental group  $\pi_1(\widehat{X}_{n,d})$ . In order to do this we use the main theorem of [Arm68] in the case of product quotient varieties following [BCGP12, DP12]. We briefly recall their strategy and we refer to them for further details.

Let  $G$  be a finite group acting diagonally on a product  $Z := C_1 \times \dots \times C_n$  of curves of genus at least 2, and consider the group  $\mathbb{G}$  of all possible lifts of automorphisms induced by the action of  $G$  on  $Z$  to the universal cover  $u: \mathbb{H}^n \rightarrow Z$ . The group  $\mathbb{G}$  acts properly discontinuously on  $\mathbb{H}^n$  and  $u$  is

equivariant with respect to the natural map  $\mathbb{G} \rightarrow G$ , hence we have an isomorphism  $\mathbb{H}^n/\mathbb{G} \cong Z/G$ . Since  $\mathbb{H}^n$  is simply connected we can apply Armstrong's results (see [Arm68]) and get the following.

**Proposition 13.** *Let  $\text{Fix}(\mathbb{G})$  be the normal subgroup of  $\mathbb{G}$  generated by the elements having non-empty fixed locus. Then*

$$\pi_1(Z/G) = \mathbb{G}/\text{Fix}(\mathbb{G}).$$

Assume that the  $G$ -action on  $Z$  restricts to a faithful action  $\phi_i$  on each factor  $C_i$ . Let  $\mathbb{T}_i$  be the group of all possible lifts of automorphisms induced by the action of  $G$  on  $C_i$  to the universal cover  $\mathbb{H}$  of  $C_i$ , and let  $\varphi_i : \mathbb{T}_i \rightarrow G$  be the natural map. In this setting, the above group  $\mathbb{G}$  is the preimage of the diagonal subgroup  $\Delta_G \subset G^n$  under  $\varphi_1 \times \dots \times \varphi_n$ :

$$\mathbb{G} = \{(x_1, \dots, x_n) \in \mathbb{T}_1 \times \dots \times \mathbb{T}_n \mid \varphi_1(x_1) = \dots = \varphi_n(x_n)\}.$$

There is also a similar description of  $\mathbb{G}$  in the non-faithful case, see [DP12, Proposition 3.3].

*Remark 14.* i) The group  $\mathbb{T}_i$  has a simple presentation (see also [Cat15, Example 29]): let  $g'$  be the genus of  $C_i/G$  and  $m_1, \dots, m_r$  be the ramification indices of the branch points of the covering map  $C_i \rightarrow C_i/G$ , then

$$\mathbb{T}_i = \mathbb{T}(g'; m_1, \dots, m_r) := \langle a_1, b_1, \dots, a_{g'}, b_{g'}, c_1, \dots, c_r \mid c_1^{m_1}, \dots, c_r^{m_r}, \prod_{i=1}^{g'} [a_i, b_i] \cdot c_1 \cdots c_r \rangle.$$

ii) The group  $\mathbb{T}(g'; m_1, \dots, m_r)$  is called the *orbifold surface group* of type  $[g'; m_1, \dots, m_r]$ .

The non-trivial stabilizers of the  $\mathbb{T}_i$ -action on  $\mathbb{H}$  are cyclic and generated by the conjugates of the elements  $c_k$ . The restriction of  $\varphi_i$  to each one of these subgroups is an isomorphism onto its image, which is the stabilizer of a point in  $C_i$ . Conversely, all non-trivial stabilizers of the  $G$ -action on  $C_i$  are of this form (see [BCGP12]).

**Definition 15.** Let  $L_i \subset \mathbb{T}_i$  be set of the elements  $c_j^{l_j} \in \mathbb{T}_i$  such that  $\varphi_i(c_j^{l_j}) \in G$  has non-empty fixed locus on  $Z = C_1 \times \dots \times C_n$ , where  $j \in \{1, \dots, r\}$  and  $l_j \in \{1, \dots, m_j - 1\}$ .

We denote by  $\langle\langle L_i \rangle\rangle_{\mathbb{T}_i}$  the normal subgroup of  $\mathbb{T}_i$  generated by  $L_i$ .

**Proposition 16** (Finiteness criterion). *The group  $\pi_1(Z/G) = \mathbb{G}/\text{Fix}(\mathbb{G})$  is finite if and only if the groups  $\mathbb{T}_i/\langle\langle L_i \rangle\rangle_{\mathbb{T}_i}$  are finite.*

*Proof.* According to [BCGP12, pag.1018-1019] the group  $\mathbb{G}/\text{Fix}(\mathbb{G})$  fits in an exact sequence

$$1 \rightarrow E \rightarrow \mathbb{G}/\text{Fix}(\mathbb{G}) \rightarrow \mathbf{H} \rightarrow 1,$$

where  $E$  is a finite group and  $\mathbf{H}$  is a subgroup of finite index of the product

$$\mathbb{T}_1/\langle\langle L_1 \rangle\rangle_{\mathbb{T}_1} \times \dots \times \mathbb{T}_n/\langle\langle L_n \rangle\rangle_{\mathbb{T}_n}. \quad \square$$

*Remark 17.* Let  $X$  be a normal variety with only quotient singularities, and let  $\rho: \widehat{X} \rightarrow X$  be a resolution of singularities. Then  $\rho_*: \pi_1(\widehat{X}) \rightarrow \pi_1(X)$  is an isomorphism, by [Kol93, Theorem 7.8].

In particular,  $\pi_1(\widehat{X}_{n,d}) \simeq \pi_1(X_{n,d})$ .

According to the description of  $X_{n,d}$  given in the previous section its associated orbifold surface groups  $\mathbb{T}_i$  are all of type  $[0; d, d, d]$ , and applying this discussion to our situation we get the following.

**Theorem 18.** *The universal cover  $U_{n,d}$  of  $\widehat{X}_{n,d}$  is projective if and only if  $d = 4$ .*

*Proof.* The universal cover  $U_{n,d}$  of  $\widehat{X}_{n,d}$  is projective if and only if the fundamental group  $\pi_1(\widehat{X}_{n,d})$  is finite. Therefore, by Proposition 16  $U_{n,d}$  is projective if and only if the groups  $\mathbb{T}_i/\langle\langle L_i \rangle\rangle_{\mathbb{T}_i}$  are finite. Let  $k := \frac{d}{2}$ . Since the elements in  $\mathbb{Z}_d^2$  fixing points on  $C^m$  are exactly the elements in  $H = \langle(k, 0), (0, k)\rangle$ , by Remark 14 ii) we see that  $L_i = \{c_1^k, c_2^k, c_3^k\}$ , whence

$$\mathbb{T}_i/\langle\langle L_i \rangle\rangle_{\mathbb{T}_i} \cong \mathbb{T}(0; d, d, d)/\langle\langle c_1^k, c_2^k, c_3^k \rangle\rangle = \langle c_1, c_2, c_3 | c_1^k, c_2^k, c_3^k, c_1 c_2 c_3 \rangle \cong \mathbb{T}(0; k, k, k).$$

The statement follows since the group  $\mathbb{T}(0; 2, 2, 2) \cong \mathbb{Z}_2^2$  is finite, whereas  $\mathbb{T}(0; k, k, k)$  is infinite for  $k > 2$ .  $\square$

*Remark 19.* i) The first Betti number  $b_1$  of  $Y_{n,4}$  is zero, because the quotient  $C/H$  is isomorphic to the projective line. Indeed by Künneth formula and [Mac62, §1.2] we have

$$H^1(Y_{n,4}, \mathbb{C}) = H^1(C^m, \mathbb{C})^H = \bigoplus H^1(C, \mathbb{C})^H = \bigoplus H^1(\mathbb{P}^1, \mathbb{C}) = 0.$$

Assuming  $d = 4$ , we can actually prove that  $g^2 = 1$  for all  $g \in \pi_1(Y_{n,4}) = \mathbb{G}/\text{Fix}(\mathbb{G})$ . This tells us that  $\pi_1(Y_{n,4}) = \pi_1(\widehat{Y}_{n,4}) \cong \mathbb{Z}_2^s$  for some  $s \in \mathbb{N}$ .

The element  $g$  is represented by an  $n$ -tuple

$$(w_1, \dots, w_n) \in \mathbb{G} = \mathbb{T}_1 \times_H \dots \times_H \mathbb{T}_n$$

where  $\mathbb{T}_k = \mathbb{T}(0; 2, 2, 2, 2, 2, 2)$  and all the maps  $\varphi_k : \mathbb{T}_k \rightarrow H$  are equal, as we consider the same action on each factor (see Remark 4). Since  $\varphi_k(w_k^2) = (0, 0) \in H = \mathbb{Z}_2^2$ , the tuple

$$(1, \dots, 1, w_k^2, 1, \dots, 1)$$

belongs to  $\mathbb{G}$ , and to prove the claim it suffices to show that this tuple is contained in  $\text{Fix}(\mathbb{G})$ .

Note that the number of occurrences  $n_i$  of the letter  $c_i$  in the word  $w_k^2$  is even. Observe now, that in any group a product  $a \cdot b$  can be written as  $b \cdot (b^{-1} \cdot a \cdot b)$ , hence we can write  $w_k^2$  as

$$(2.1) \quad w_k^2 = \left( \prod_{i=1}^{n_1} g_i^{-1} c_1 g_i \right) \cdot \dots \cdot \left( \prod_{j=1}^{n_6} h_j^{-1} c_6 h_j \right),$$

for certain  $g_i, \dots, h_j \in \mathbb{T}_k$ .

By Remark 14 ii) and since  $H$  is abelian, we get  $(c_1, \dots, c_1, g_i^{-1} c_1 g_i, c_1, \dots, c_1) \in \text{Fix}(\mathbb{G})$ . We conclude that

$$(1, \dots, 1, \prod_{i=1}^{n_1} g_i^{-1} c_1 g_i, 1, \dots, 1) = \prod_{i=1}^{n_1} (c_1, \dots, c_1, g_i^{-1} c_1 g_i, c_1, \dots, c_1) \in \text{Fix}(\mathbb{G}).$$

The same applies to each factor in the RHS of (2.1) and so  $(1, \dots, 1, w_k^2, 1, \dots, 1) \in \text{Fix}(\mathbb{G})$ . This shows  $g^2 = 1$ , whence  $\pi_1(Y_{n,4})$  is abelian, and it is finite since  $\pi_1(Y_{n,4}) = \pi_1(Y_{n,4})^{ab} = H_1(Y_{n,4}, \mathbb{Z})$  has rank 0.

ii) We implemented Proposition 13 using the computer algebra system MAGMA [BCP97], and we found  $\pi_1(Y_{n,4}) = \mathbb{Z}_2^{n-1}$  and  $\pi_1(X_{n,4}) = \mathbb{Z}_2^{n+1}$  for  $n = 2, 3, 4, 5$ . In particular, the universal cover of the varieties  $X_{n,4}$  and  $Y_{n,4}$  has  $3 \cdot 2^{3n-2}$  singularities of type  $\frac{1}{2}(1, \dots, 1)$ . We expect the above to generalize to any dimension.

*Remark 20.* The surfaces  $\widehat{X}_{2,d}$  with  $d \geq 8$  are rigid but not infinitesimally rigid (see [BP18]), and their universal cover is non-contractible. This answer partially the question posed in the Introduction in the case of surfaces.

## REFERENCES

- [Arm68] M.A. Armstrong, *The fundamental group of the orbit space of a discontinuous group*, Proc. Cambridge Phil. Soc. **64** (1968), 299–301.
- [BC18] I. Bauer and F. Catanese, *On rigid compact complex surfaces and manifolds*, Adv. Math. **333** (2018), 620–669.
- [BCGP12] I. Bauer, F. Catanese, F. Grunewald, and R. Pignatelli, *Quotients of products of curves, new surfaces with  $p_g = 0$  and their fundamental groups*, American Journal of Mathematics **134** (2012), no. 4, 993–1049.
- [BCP97] W. Bosma, J. Cannon, and C. Playoust, *The Magma algebra system. I. The user language*, J. Symbolic Comput. **24** (1997), no. 3-4, 235–265. Computational algebra and number theory (London, 1993).
- [Bea78] A. Beauville, *Surfaces algébriques complexes*, Asterisque **54**, Soc.Math. France, 1978.
- [BG20] I. Bauer and C. Gleissner, *Fermat’s cubic, Klein’s quartic and rigid complex manifolds of Kodaira dimension one*, Doc. Math. **25** (2020), 1241–1262.
- [BG21] ———, *Towards a Classification of Rigid Product Quotient Varieties of Kodaira Dimension 0* (2021), available at [ArXiv:2101.06925](https://arxiv.org/abs/2101.06925).
- [BGP20] C. Böhning, H.-C. Graf von Bothmer, and R. Pignatelli, *A rigid, not infinitesimally rigid surface with  $K$  ample* (2020), available at [ArXiv:2010.14371](https://arxiv.org/abs/2010.14371).
- [BP18] I. Bauer and R. Pignatelli, *Rigid but not infinitesimally rigid compact complex manifolds* (2018), available at [ArXiv:1805.02559](https://arxiv.org/abs/1805.02559). To appear in Duke Mathematical Journal.
- [BW74] D. M. Burns Jr. and J. M. Wahl, *Local contributions to global deformations of surfaces*, Invent. Math. **26** (1974), 67–88.
- [Cat00] F. Catanese, *Fibred surfaces, varieties isogenous to a product and related moduli spaces.*, American Journal of Mathematics **122** (2000), no. 1, 1–44.
- [Cat07] ———, *Q.E.D. for algebraic varieties.*, J. Differential Geom. **77** (2007), no. 1, 43–75.
- [Cat15] ———, *Topological methods in moduli theory*, Bull. Math. Sci. **5** (2015), no. 3, 287–449.
- [CF18] N. Cancian and D. Frapporti, *On semi-isogenous mixed surfaces*, Math. Nachr. **291** (2018), no. 2-3, 264–283.
- [DP12] T. Dedieu and F. Perroni, *The fundamental group of a quotient of a product of curves*, J. Group Theory **15** (2012), no. 3, 439–453.
- [FG20] D. Frapporti and C. Gleißner, *A family of threefolds of general type with canonical map of high degree*, Taiwanese J. Math. **24** (2020), no. 5, 1107–1115.
- [FGP20] F. Favale, C. Gleissner, and R. Pignatelli, *The pluricanonical systems of a product-quotient variety*, Galois covers, grothendieck-teichmüller theory and dessins d’enfants, 2020, pp. 89–119.
- [GPR18] C. Gleissner, R. Pignatelli, and C. Rito, *New surfaces with canonical map of high degree* (2018), available at [ArXiv:1807.11854](https://arxiv.org/abs/1807.11854). To appear in Commun. Anal. Geom.
- [Hat02] A. Hatcher, *Algebraic topology*, Cambridge university press, 2002.
- [Kol93] J. Kollár, *Shafarevich maps and plurigenera of algebraic varieties*, Invent. Math. **113** (1993), no. 1, 177–215.
- [LP16] M. Lönne and M. Penegini, *On asymptotic bounds for the number of irreducible components of the moduli space of surfaces of general type II*, Doc. Math. **21** (2016), 197–204.
- [LP20] M. Lönne and M. Penegini, *On Zariski Multiplets of Branch Curves from Surfaces Isogenous to a Product*, Michigan Math. J. **69** (2020), no. 4, 779–792.
- [Mac62] I. G. Macdonald, *Symmetric products of an algebraic curve*, Topology **1** (1962), 319–343.
- [MK71] J. Morrow and K. Kodaira, *Complex manifolds*, Holt, Rinehart and Winston, Inc., New York-Montreal, Que.-London, 1971.
- [Pin81] H. Pinkham, *Some local obstructions to deforming global surfaces*, Nova Acta Leopoldina (N.F.) **52** (1981), no. 240, 173–178. Leopoldina Symposium: Singularities (Thüringen, 1978).
- [Rei87] M. Reid, *Young person’s guide to canonical singularities*, Algebraic geometry, Bowdoin, 1985 (Brunswick, Maine, 1985), 1987, pp. 345–414.

[Sch71] M. Schlessinger, *Rigidity of quotient singularities*, Invent. Math. **14** (1971), 17–26.

DAVIDE FRAPPORTI, UNIVERSITÀ DEGLI STUDI DI GENOVA, DIPARTIMENTO DI MATEMATICA - DIMA;  
VIA DODECANESO 35, I-16146 GENOVA, ITALY.

*Email address:* `daide.frapporti@edu.unige.it`

CHRISTIAN GLEISSNER , UNIVERSITY OF BAYREUTH, LEHRSTUHL MATHEMATIK VIII;  
UNIVERSITÄTSSTRASSE 30, D-95447 BAYREUTH, GERMANY.

*Email address:* `Christian.Gleissner@uni-bayreuth.de`