

Causal Inference with Invalid Instruments: Post-selection Problems and A Solution Using Searching and Sampling

Zijian Guo *

Abstract

Instrumental variable methods are among the most commonly used causal inference approaches to account for unmeasured confounders in observational studies. The presence of invalid instruments is a significant concern for practical applications, and a fast-growing area of research is inference for the causal effect with possibly invalid instruments. The existing inference methods rely on correctly separating valid and invalid instruments in a data-dependent way. This paper illustrates that the existing confidence intervals may substantially undercover due to the post-selection problem. To address this, we construct uniformly valid confidence intervals for the causal effect, robust to the mistakes in separating valid and invalid instruments. Our proposal is to search for a range of effect values that lead to sufficiently many valid instruments. We further devise a novel sampling method, which, together with searching, leads to a more precise confidence interval. Our proposed searching and sampling confidence intervals are uniformly valid and achieve the parametric length under the finite-sample majority and plurality rules. We compare our proposed methods with existing inference methods over a large set of simulation studies and apply them to study the effect of education on earnings.

Keywords: Unmeasured confounders; Uniform inference; Majority rule; Plurality rule; Mendelian Randomization.

1 Introduction

Unmeasured confounders are a major concern for causal inference with observational studies. The instrumental variable (IV) method is one of the most commonly used causal inference approaches to deal with unmeasured confounders. The IVs are required to satisfy three identification conditions: conditioning on the baseline covariates,

- (A1) the IVs are associated with the treatment;
- (A2) the IVs are independent with the unmeasured confounders;
- (A3) the IVs have no direct effect on the outcome.

*Z. Guo is an assistant professor at Department of Statistics, Rutgers University. The research of Z. Guo was supported in part by the NSF DMS 1811857, 2015373 and NIH R01GM140463-01.

The main challenge of IV-based methods is identifying instruments satisfying (A1), (A2), and (A3) simultaneously. The exclusion restriction assumptions (A2) and (A3) are crucial for identifying the causal effect as they assume that the IVs can only affect the outcome through the treatment. However, assumptions (A2) and (A3) may be violated in applications and cannot even be tested in a data-dependent way. We define an IV as “invalid” if it violates assumptions (A2) or (A3). If an invalid IV is mistakenly taken as valid, it generally leads to a biased estimator of the causal effect. A fast-growing literature is to conduct causal inference with possibly invalid IVs [9, 11, 21, 25, 31–33, 47, 52, 53, e.g.]. Many of these works are motivated by Mendelian Randomization studies using genetic variants as IVs [12]. The adopted genetic variants are possibly invalid since they may affect both treatment and outcome due to the pleiotropy effect [16, 17].

In this paper, we focus on the setting with multiple IVs, which are possibly invalid. Under this framework, the effect identification requires extra conditions such as the majority rule [11, 32] and plurality rule [25, 28]; see the exact definitions in Conditions 1 and 2 in Section 2. Both conditions require that there exist enough valid IVs even though the validity of any IV is not known a priori. The existing works [25, 32, 52, 53, e.g.] leveraged the majority and plurality rules to select valid IVs in a data-dependent way. The selected valid IVs were used for the following-up causal inference, where the invalid IVs are included as the baseline covariates. However, there are chances that we make mistakes in separating valid and invalid IVs, especially when there is only a finite amount of data. The IV selection error leads to unreliable confidence intervals; see Section 3 for an illustration.

It is of practical importance to develop uniform inference methods robust to the IV selection error. Certain invalid IVs can be hard to detect in applications with the given data set. We refer to such invalid IVs as “locally invalid IVs” and provide a formal definition in the following Definition 1 in Section 3. The current paper aims to propose uniformly valid confidence intervals robust to IV selection error.

1.1 Results and contributions

The current paper is focused on the linear outcome model with possibly invalid IVs and heteroscedastic errors. In Section 3, we demonstrate that even if the majority rule (Condition 1) or plurality rule (Condition 2) holds, the existing inference methods TSHT [25] and CIIV [53] produce unreliable confidence intervals with a finite amount of data. To better accommodate finite-sample inferential properties, we introduce the finite-sample majority and plurality rule in the following Conditions 3 and 4, respectively.

We start by explaining our methods under the finite-sample majority rule. Our first proposed confidence interval (CI) is based on the *searching* idea. For every treatment effect, we implement a hard thresholding step to decide which candidate IVs are valid. We propose to search for a range of the treatment effects such that the majority of candidate IVs can be taken as valid. We further propose a novel *sampling* method to improve the precision of the searching CI.

For settings only assuming the finite-sample plurality rule (Condition 4), we construct an initial estimator $\hat{\mathcal{V}}$ of the set of valid IVs and then apply the searching and sampling method

over $\widehat{\mathcal{V}}$. Our proposed searching CI works even if $\widehat{\mathcal{V}}$ does not correctly recover the set of valid IVs. This contrasts sharply with the existing works [25, 53], where the validity of these inference methods relies on $\widehat{\mathcal{V}}$ to recover the set of valid IVs correctly.

Our proposed searching and sampling CIs are shown to achieve the desired coverage under the finite-sample majority or plurality rule. The CIs are uniformly valid in the sense that the coverage is guaranteed even if some invalid IVs only violate (A2) and (A3) locally; see the following Definition 1 in Section 3. We also establish that the searching and sampling CIs achieve the $1/\sqrt{n}$ length.

The proposed CIs are computationally efficient as the searching method searches over one-dimension space, and we only resample the reduced-form estimators instead of the full data. In Section 7, we conduct a large set of simulation studies to compare our proposed methods with existing CI construction methods: TSHT [25], CIIV [53], and Union method [31]. In Section 8, we apply our proposed method to study the effect of education on earnings.

To sum up, the contribution of the current paper is two-folded.

1. We introduce the finite-sample majority and plurality rules to better accommodate the effect identification with finite samples.
2. We propose novel searching and sampling methods to construct uniform CIs for the treatment effect. Our proposed CIs are robust to the mistakes in separating valid and invalid IVs.

1.2 Literature comparison

The most relevant papers to the current work are [25, 53]. The validity of the CIs in [25, 53] requires the invalid IVs to be correctly identified. However, having a perfect separation between valid and invalid IVs is challenging, especially in the presence of locally invalid instruments; see its definition in Definition 1 in Section 3. More detailed comparison with [53] is presented in Section 5.2. In Figure 3 in Section 3, we observe that the CIs by [25, 53] may undercover in finite samples when locally invalid IVs are mistakenly selected as valid IVs. In contrast, our proposed searching and sampling CIs achieve the desired coverage level.

Another closely related work [31] proposed constructing CIs by taking a union of intervals constructed by a given number of candidate IVs and not rejected by the Sargan test [27, 42]. An upper bound for the number of invalid IVs is required for the construction in [31]. Our proposed searching and sampling CIs do not rely on such information and are computationally more efficient than [31] since our proposed methods avoid searching over many sub-models. As illustrated in Figure 3 in Section 3, our proposed sampling and searching CIs are typically much shorter than the CIs in [31].

Different identifiability conditions have been proposed to identify the causal effect when the IV assumptions (A2) and (A3) fail to hold. [10] and [33] assumed that the IVs' direct effect on the outcome and the IVs' association with the treatment are nearly orthogonal. The majority rule was applied to identify the treatment effect in both the linear outcome model [11, 32, 52] and

the nonlinear outcome model [38]. [44] proposed to identify the treatment effect by requiring the interaction of the IVs and an environmental factor to satisfy the IV assumptions (A1)-(A3). [36,37,40,47] leveraged the heteroscedastic covariance restriction to identify the treatment effect. In Mendelian Randomization studies, much progress has been made in inference with summary statistics, which is not the main focus of the current paper; see [10,11,57] for examples. In the GMM setting, [13,14,39] leveraged a set of pre-specified valid moment conditions for the model identification and further tested the validity of another set of moment conditions. The current paper is entirely different in the sense that there is no prior knowledge of the validity of any given IV.

The construction of uniformly valid CIs after model selection is a major focus in statistics under the name of post-selection inference. Many methods [5,15,30,34,35,49,50,56] have been proposed and the focus is on (but not limited to) inference for regression coefficients after some variables or sub-models are selected. This paper considers a different problem, post-selection inference for causal effect with possibly invalid instruments. To our best knowledge, this problem has not been carefully investigated in the post-selection inference literature. Furthermore, our proposed sampling method is different from other existing post-selection inference methods. The sampling method can be of independent interest and find applications in other post-selection inference problems.

Notations. For a set S and a vector $x \in \mathbb{R}^p$, $|S|$ denotes the cardinality of S and x_S is the sub-vector of x with indices in S . The ℓ_q norm of a vector x is defined as $\|x\|_q = (\sum_{l=1}^p |x_l|^q)^{\frac{1}{q}}$ for $q \geq 0$ with $\|x\|_0 = |\{1 \leq l \leq p : x_l \neq 0\}|$ and $\|x\|_\infty = \max_{1 \leq l \leq p} |x_l|$. We use $\mathbf{0}_q$ and $\mathbf{1}_q$ to denote the q -dimension vector with all entries equal to 0 and 1, respectively. For a matrix X , $X_{i\cdot}$, $X_{\cdot j}$ and X_{ij} are used to denote its i -th row, j -th column and (i, j) entry, respectively. For a sequence of random variables X_n indexed by n , we use $X_n \xrightarrow{d} X$ to denote that X_n converges to X in distribution. We use c and C to denote generic positive constants that may vary from place to place. For two positive sequences a_n and b_n , $a_n \lesssim b_n$ means that $\exists C > 0$ such that $a_n \leq Cb_n$ for all n ; $a_n \asymp b_n$ if $a_n \lesssim b_n$ and $b_n \lesssim a_n$, and $a_n \ll b_n$ if $\limsup_{n \rightarrow \infty} a_n/b_n = 0$. For a matrix A , we use $\|A\|_2$ and $\|A\|_\infty$ to denote its spectral norm and element-wise maximum norm, respectively. For a symmetric matrix A , we use $\lambda_{\max}(A)$ and $\lambda_{\min}(A)$ to denote its maximum and minimum eigenvalues, respectively.

2 Models and Reduced-form Estimators

We consider the i.i.d. data $\{Y_i, D_i, X_i, Z_i\}_{1 \leq i \leq n}$, where $Y_i \in \mathbb{R}$, $D_i \in \mathbb{R}$ and $X_i \in \mathbb{R}^{p_x}$ and $Z_i \in \mathbb{R}^{p_z}$ denote the outcome, the treatment, the baseline covariates, and candidate IVs, respectively. We consider the following outcome model with possibly invalid IVs [25,32,43,53],

$$Y_i = D_i\beta^* + Z_i^\top \pi^* + X_i^\top \phi^* + e_i \quad \text{with} \quad \mathbf{E}(e_i Z_i) = \mathbf{0} \quad \text{and} \quad \mathbf{E}(e_i X_i) = \mathbf{0}, \quad (1)$$

where $\beta^* \in \mathbb{R}$ denotes the treatment effect, $\pi^* \in \mathbb{R}^{p_z}$, and $\phi^* \in \mathbb{R}^{p_x}$. We set the first element of X_i as the constant 1. If the IVs satisfy (A2) and (A3), this leads to $\pi^* = \mathbf{0}$ in (1). A non-zero

vector π^* indicates that the IVs violate the classical IV assumptions (A2) and (A3); see the following Figure 1 for an illustration.

We consider the association model for the treatment D_i ,

$$D_i = Z_i^\top \gamma^* + X_i^\top \psi^* + \delta_i \quad \text{with} \quad \mathbf{E}(\delta_i | Z_i) = \mathbf{0} \quad \text{and} \quad \mathbf{E}(\delta_i | X_i) = \mathbf{0}. \quad (2)$$

The model (2) can be viewed as the best linear approximation of D_i by Z_i and X_i instead of a casual model. In (2), $\gamma_j^* \neq 0$ indicates that the j -th IV satisfies assumption (A1). Due to the existence of unmeasured confounders, e_i and δ_i can be correlated and the treatment D_i is endogenous with $\mathbf{E}(D_i e_i) \neq 0$.

Following [32], we discuss the causal interpretation of the model (1) and explain why $\pi^* \neq 0$ represents the violation of assumptions (A2) and (A3). For two treatment values $d, d' \in \mathbb{R}$ and two realizations of IVs $\mathbf{z}, \mathbf{z}' \in \mathbb{R}^{p_z}$, define the following potential outcome model:

$$Y_i^{(d', \mathbf{z}')} - Y_i^{(d, \mathbf{z})} = (d' - d)\beta^* + (\mathbf{z}' - \mathbf{z})^\top \kappa^* \quad \text{and} \quad \mathbf{E}(Y_i^{(0, \mathbf{0})} | Z_i, X_i) = Z_i^\top \eta^* + X_i^\top \phi^*$$

where $\beta^* \in \mathbb{R}$ is the treatment effect, $\kappa^*, \eta^* \in \mathbb{R}^{p_z}$ and $\phi^* \in \mathbb{R}^{p_x}$. As illustrated in Figure 1, $\kappa_j^* \neq 0$ indicates that the j -th candidate IV has a direct effect on outcome, which violates the assumption (A3); $\eta_j \neq 0$ indicates that the j -th candidate IV is associated with the unmeasured confounder, which violates the assumption (A2). Under the consistency condition $Y_i = Y_i^{(D_i, Z_i)}$, the above potential outcome model implies (1) with the invalidity vector $\pi^* = \kappa^* + \eta^*$ and $e_i = Y_i^{(0, \mathbf{0})} - \mathbf{E}(Y_i^{(0, \mathbf{0})} | Z_i, X_i)$.




Figure 1: Illustration of violations of (A2) and (A3) in the model (1) with $\pi^* = \kappa^* + \eta^*$.

We define the set \mathcal{S} of relevant instruments and the set \mathcal{V} of valid instruments as,

$$\mathcal{S} = \{1 \leq j \leq p_z : \gamma_j^* \neq 0\} \quad \text{and} \quad \mathcal{V} = \{j \in \mathcal{S} : \pi_j^* = 0\}. \quad (3)$$

The IVs belonging to \mathcal{S} satisfy the IV assumption (A1). The set \mathcal{V} is a subset of \mathcal{S} and the IVs belonging to \mathcal{V} satisfy the classical IV assumptions (A1)-(A3) simultaneously.

We now review the identification strategy under models (1) and (2). We plug in the treatment model (2) into the outcome model (1) and obtain the reduced-form model,

$$\begin{aligned} Y_i &= Z_i^\top \Gamma^* + X_i^\top \Psi^* + \epsilon_i \quad \text{with} \quad \mathbf{E}(Z_i \epsilon_i) = \mathbf{0}, \quad \mathbf{E}(X_i \epsilon_i) = \mathbf{0}, \\ D_i &= Z_i^\top \gamma^* + X_i^\top \psi^* + \delta_i \quad \text{with} \quad \mathbf{E}(Z_i \delta_i) = \mathbf{0}, \quad \mathbf{E}(X_i \delta_i) = \mathbf{0}, \end{aligned} \quad (4)$$

where $\Gamma^* = \beta^*\gamma^* + \pi^* \in \mathbb{R}^{p_z}$, $\Psi^* = \beta^*\psi^* + \phi^* \in \mathbb{R}^{p_x}$, and $\epsilon_i = \beta^*\delta_i + e_i$. We shall devise our methods under the model (4), which is induced by the models (1) and (2).

Since Z_i and X_i are uncorrelated with ϵ_i and δ_i , we can identify the reduced-form parameters Γ^* and γ^* in the model (4). However, since $\pi^* \neq \mathbf{0}$, the identification of β^* through solving the equation $\Gamma^* = \beta^*\gamma^* + \pi^* \in \mathbb{R}^{p_z}$ requires extra assumptions. The following majority rule has been proposed for identifying the causal effect with invalid IVs [11, 25, 32, 52].

Condition 1 (Population Majority Rule) *More than half of the relevant IVs are valid, that is, $|\mathcal{V}| > |\mathcal{S}|/2$, where \mathcal{V} and \mathcal{S} are defined in (3).*

Condition 1 requires that more than half of the relevant IVs are valid but does not directly require the knowledge of \mathcal{V} . For the j -th IV, we may identify the effect as $\beta^{[j]} = \Gamma_j^*/\gamma_j^*$. If the j -th IV is valid, $\beta^{[j]} = \beta^*$; otherwise, $\beta^{[j]} \neq \beta^*$. Under Condition 1, β^* can be identified using the majority of $\{\beta^{[j]}\}_{j \in \mathcal{S}}$. [25, 28] proposed the following plurality rule as a weaker identification condition.

Condition 2 (Population Plurality Rule) *The number of valid IVs is larger than the number of invalid IVs with any given invalidity level $\nu \neq 0$, that is,*

$$|\mathcal{V}| > \max_{\nu \neq 0} |\mathcal{I}_\nu| \quad \text{with} \quad \mathcal{I}_\nu = \{j \in \mathcal{S} : \pi_j^*/\gamma_j^* = \nu\},$$

where the set \mathcal{V} of valid IVs is defined in (3).

The term π_j^*/γ_j^* represents the invalidity level: $\pi_j^*/\gamma_j^* \neq 0$ indicates that the j -th IV violates assumptions (A2) and (A3). \mathcal{I}_ν denotes the set of all IVs with the same invalidity level ν . Note that $\mathcal{V} = \mathcal{I}_0$. Under the plurality rule, β^* can be identified using the largest cluster of $\{\beta^{[j]}\}_{j \in \mathcal{S}}$. We refer to Conditions 1 and 2 as *population* identification conditions since they are used to identify β^* with an infinite amount of data. To account for the finite-sample uncertainty, we shall modify Conditions 1 and 2 and propose the finite-sample rules in the following Conditions 3 and 4.

We now present the data-dependent estimators of γ^* and Γ^* in the model (4). Define $p = p_x + p_z$, $W = (Z, X) \in \mathbb{R}^{n \times p}$, and $\widehat{\Sigma} = \frac{1}{n} \sum_{i=1}^n W_i W_i^\top$. We estimate γ^* and Γ^* by the Ordinary Least Squares (OLS) estimators, defined as

$$(\widehat{\Gamma}^\top, \widehat{\Psi}^\top)^\top = (W^\top W)^{-1} W^\top Y \quad \text{and} \quad (\widehat{\gamma}^\top, \widehat{\psi}^\top)^\top = (W^\top W)^{-1} W^\top D. \quad (5)$$

Under regularity conditions, the OLS estimators satisfy

$$\sqrt{n} \begin{pmatrix} \widehat{\Gamma} - \Gamma^* \\ \widehat{\gamma} - \gamma^* \end{pmatrix} \xrightarrow{d} N(\mathbf{0}, \text{Cov}) \quad \text{with} \quad \text{Cov} = \begin{pmatrix} \mathbf{V}^\Gamma & \mathbf{C} \\ \mathbf{C}^\top & \mathbf{V}^\gamma \end{pmatrix}, \quad (6)$$

where $\mathbf{V}^\Gamma \in \mathbb{R}^{p_z \times p_z}$, $\mathbf{V}^\gamma \in \mathbb{R}^{p_x \times p_x}$, and $\mathbf{C} \in \mathbb{R}^{p_z \times p_x}$ are explicitly defined in Lemma 1 in the supplement. Define the residues $\widehat{\epsilon}_i = Y_i - Z_i^\top \widehat{\Gamma} - X_i^\top \widehat{\Psi}$ and $\widehat{\delta}_i = D_i - Z_i^\top \widehat{\gamma} - X_i^\top \widehat{\psi}$ for $1 \leq i \leq n$.

We estimate \mathbf{V}^Γ , \mathbf{V}^γ , and \mathbf{C} in (6) by

$$\begin{aligned} \widehat{\mathbf{V}}^\Gamma &= \left[\widehat{\Sigma}^{-1} \left(\frac{1}{n} \sum_{i=1}^n \widehat{\epsilon}_i^2 W_i W_i^\top \right) \widehat{\Sigma}^{-1} \right]_{1:p_z, 1:p_z}, \quad \widehat{\mathbf{V}}^\gamma = \left[\widehat{\Sigma}^{-1} \left(\frac{1}{n} \sum_{i=1}^n \widehat{\delta}_i^2 W_i W_i^\top \right) \widehat{\Sigma}^{-1} \right]_{1:p_z, 1:p_z}, \\ \widehat{\mathbf{C}} &= \left[\widehat{\Sigma}^{-1} \left(\frac{1}{n} \sum_{i=1}^n \widehat{\epsilon}_i \widehat{\delta}_i W_i W_i^\top \right) \widehat{\Sigma}^{-1} \right]_{1:p_z, 1:p_z}. \end{aligned} \quad (7)$$

The variance estimators in (7) are robust to the heteroscedastic errors and referred to as the sandwich estimators [20, 29, 51]; see Chapter 4.2.3 of [54] for details. We use the OLS estimators in (5) and the sandwich estimators in (7) as a prototype to discuss our proposed methods. Our proposed methods are effective for any reduced-form estimators satisfying (6). In Section 7.3, we consider the high-dimensional setting with $p > n$ and demonstrate the effectiveness of our proposal by constructing the debiased Lasso estimators $\widehat{\Gamma}$ and $\widehat{\gamma}$ satisfying (6).

3 IV Selection Errors and Nonuniform Inference

In this section, we demonstrate that even if the majority rule (Condition 1) or plurality rule (Condition 2) holds, the CIs by TSHT [25] and CIIV [53] may be unreliable. The main idea of [25] and [53] is to estimate \mathcal{V} by a set estimator $\widehat{\mathcal{V}}$ and then identify β^* through the following expression or its weighted version,

$$\beta(\widehat{\mathcal{V}}) = \sum_{j \in \widehat{\mathcal{V}}} \Gamma_j^* \gamma_j^* / \sum_{j \in \widehat{\mathcal{V}}} (\gamma_j^*)^2. \quad (8)$$

When there are no selection errors (i.e. $\widehat{\mathcal{V}} = \mathcal{V}$), we have $\beta(\widehat{\mathcal{V}}) = \beta^*$. The validity of the CIs by TSHT and CIIV requires $\widehat{\mathcal{V}}$ to recover \mathcal{V} correctly. However, we likely make mistakes in estimating \mathcal{V} in finite samples, and the CIs by TSHT and CIIV are unreliable when invalid IVs are included in $\widehat{\mathcal{V}}$. The IV selection error leads to a bias in identifying β^* with $\beta(\widehat{\mathcal{V}})$. We shall point out that the targeted effect β^* is fixed, but the set $\widehat{\mathcal{V}}$ used for identifying β in (8) is selected in a data-dependent way.

In the following, we define ‘‘locally invalid IVs’’ as invalid IVs that are hard to be separated from valid IVs with a given amount of data. To facilitate the discussion, we introduce the following uncertainty measure of estimating $\pi_j^*/\gamma_j^* - \pi_k^*/\gamma_k^*$ for $j, k \in \mathcal{S}$,

$$\mathbf{T}_{j,k} := \min\{\mathbf{T}_{j,k}^0, \mathbf{T}_{k,j}^0\} \quad \text{with} \quad \mathbf{T}_{j,k}^0 = \sqrt{\frac{1}{n} \left(\mathbf{R}_{k,k}^{[j]} / [\gamma_k^*]^2 + \mathbf{R}_{j,j}^{[j]} / [\gamma_j^*]^2 - 2\mathbf{R}_{j,k}^{[j]} / [\gamma_k^* \gamma_j^*] \right)}, \quad (9)$$

where $\mathbf{R}^{[j]} = \mathbf{V}^\Gamma + (\beta^{[j]})^2 \mathbf{V}^\gamma - 2\beta^{[j]} \mathbf{C}$ and $\beta^{[j]} = \Gamma_j^* / \gamma_j^*$. We can consistently estimate $\mathbf{T}_{j,k}$ by replacing $\Gamma^*, \gamma^*, \mathbf{V}^\Gamma, \mathbf{V}^\gamma, \mathbf{C}$ in (9) with estimators in (5) and (7). We now formally define locally invalid IVs and illustrate this definition in the following Example 1.

Definition 1 (Locally invalid IV) For $j \in \mathcal{S}$, the j -th IV is locally invalid if

$$0 < |\pi_j^* / \gamma_j^*| < s_j(n) \quad \text{with} \quad s_j(n) := 2\sqrt{\log n} \cdot \max_{k \in \mathcal{V}} |\mathbf{T}_{j,k}|.$$

The definition of locally invalid IVs depends on the invalidity level π_j^*/γ_j^* and the separation level $s_j(n)$, which stands for the uncertainty level in separating the j -th IV and valid IVs. We show in Proposition 2 in Section 6.2 that the j -th IV, if invalid, can be separated from valid IVs if $|\pi_j^*/\gamma_j^*| \geq s_j(n)$. The separation level $s_j(n)$ is of the order $\sqrt{\log n/n}$ if all of $\{\gamma_j^*\}_{j \in \mathcal{S}}$ are constants. For a sufficiently large n , the set of locally invalid IVs becomes empty since $s_j(n) \rightarrow 0$. The large sample size enhances the power of detecting invalid IVs. Definition 1 is related to the local violation of IV exogeneity assumptions and valid moment conditions studied in [6, 13, 22, 26].

The theoretical results of TSHT [25] or CIIV [53] essentially assume that there are no locally invalid IVs; see Assumption 8 in [25]. However, the absence of locally invalid IVs may not properly accommodate real data analysis with a finite sample. For a given sample size n , there are chances that an invalid IV with $|\pi_j^*/\gamma_j^*| < s_j(n)$ is taken as valid by mistake. We now consider a numerical example and demonstrate that the coverage levels of CIs by TSHT and CIIV may be below the nominal level when locally invalid IVs exist.

Example 1 (Setting S2 in Section 7) For the models (1) and (2), set $\gamma^* = 0.5 \cdot \mathbf{1}_{10}$ and $\pi^* = (\mathbf{0}_4^\top, \tau/2, \tau/2, -1/3, -2/3, -1, -4/3)^\top$. The plurality rule is satisfied with $|\mathcal{V}| = 4 > \max_{v \neq 0} |\mathcal{I}_v| = 2$. We vary τ across $\{0.025, 0.05, 0.075, 0.1, 0.2, 0.3, 0.4, 0.5\}$. $s_j(n)$ for $j = 5, 6$ in Definition 1 takes the value 0.96 for $n = 500$ and 0.53 for $n = 2000$.

The fifth and sixth IVs (with a small τ) are locally invalid IVs, which are hard to detect for $n = 500, 2000$. In Figure 2, we plot the histogram of the TSHT and CIIV estimators for $\tau = 0.1$. The top and bottom panels correspond to $n = 500, 2000$, respectively. Since locally invalid IVs (with $\tau = 0.1$) are likely to be selected as valid IVs, the sample averages of the 500 estimates (the dashed line) deviate from the true value $\beta^* = 1$ (the solid line).




Figure 2: Histogram of 500 TSHT and CIIV estimates for **Example 1** with $\tau = 0.1$ and $n = 500$ (top panel) and $n = 2000$ (bottom panel).

The empirical coverage is reported in Figure 3. For a small n and τ , the estimated sets $\hat{\mathcal{V}}$ by TSHT and CIIV often contain locally invalid IVs, and the coverage levels of TSHT and CIIV are below 95%. For $n = 2000$ and $\tau = 0.5$, there are nearly no locally invalid IVs, and the CIs by TSHT and CIIV achieve the nominal level 95%. For $\tau = 0.1$, with n increasing from 500 to 2000, the fifth and sixth IVs are still locally invalid. Consequently, the bias due to the locally invalid

IVs remains unchanged, and the empirical coverage levels of TSHT and CIIV decrease since the standard errors get smaller with a larger n ; see Figure 2. In contrast, our proposed sampling CI in the following Algorithm 3 and the **Union** interval in [31] achieve the desired coverage at the expense of wider intervals. Our proposed sampling CIs are significantly shorter than the **Union** intervals.




Figure 3: Empirical coverage and average lengths for **Example 1** with $\tau \in \{0.025, 0.05, 0.075, 0.1, 0.2, 0.3, 0.4, 0.5\}$. TSHT, CIIV, and **Union** stand for the CIs by [25], [53], and [31], respectively. **Oracle BA** stands for the oracle bias-aware CI in (10), which is the benchmark. **Sampling** corresponds to our proposed sampling CI in Algorithm 3 in Section 5.1.

We introduce an oracle bias-aware CI as the benchmark when there exists IV selection error. We argue that the oracle bias-aware CI serves as a better benchmark than the oracle CI assuming the knowledge of \mathcal{V} , since it accounts for the IV selection error. For the TSHT estimator $\hat{\beta}$ and its standard error estimator $\widehat{\text{SE}}(\hat{\beta})$, we assume $(\hat{\beta} - \beta^*)/\widehat{\text{SE}}(\hat{\beta}) \xrightarrow{d} N(b, 1)$ with $|b| \leq |\mathbf{E}\hat{\beta} - \beta^*|/\widehat{\text{SE}}(\hat{\beta})$. Following (7) in [2], we leverage the oracle knowledge of $|\mathbf{E}\hat{\beta} - \beta^*|$ and form the oracle bias-aware CI as

$$\left(\hat{\beta} - \chi, \hat{\beta} + \chi\right) \quad \text{with} \quad \chi = \widehat{\text{SE}}(\hat{\beta}) \cdot \sqrt{cv_\alpha(|\mathbf{E}\hat{\beta} - \beta^*|^2/\widehat{\text{SE}}^2(\hat{\beta}))}, \quad (10)$$

where $cv_\alpha(B^2)$ is the $1 - \alpha$ quantile of the χ^2 distribution with 1 degree of freedom and non-centrality parameter B^2 . In Figure 3, the oracle bias-aware CI achieves the desired coverage and the sampling CI has a comparable length to the oracle bias-aware CI.

4 Searching and Sampling: Robust Inference Methods under Majority Rule

In this section, we focus on the majority rule's setting and generalize the proposed methods to the setting under the plurality rule in Section 5. As an analogy to TSLS and TSHT, our proposed procedure consists of two stages. In the first stage, we fit the treatment model and follow [25]

to select a set of strong IVs; see Section 4.1. In the second stage, we propose novel searching and sampling CIs in Sections 4.2 to 4.4, which are robust to the mistakes in separating valid and invalid IVs.

4.1 First-stage selection and the finite-sample majority rule

Following [25], we estimate \mathcal{S} by applying the first-stage hard thresholding to the reduced-form estimator $\hat{\gamma}$ defined in (5),

$$\hat{\mathcal{S}} = \left\{ 1 \leq j \leq p_z : |\hat{\gamma}_j| \geq \sqrt{\log n} \cdot \sqrt{\hat{\mathbf{V}}_{jj}^\gamma/n} \right\}, \quad (11)$$

where $\hat{\mathbf{V}}^\gamma$ is defined in (7). The main purpose of (11) is to estimate the set of relevant IVs and screen out IVs which are weakly associated with the treatment. The term $\sqrt{\log n}$ is introduced to adjust for the multiplicity of testing p_z hypothesis; see more discussions in the following Remark 1. The screening step in (11) guarantees the robustness of our proposal when some IVs are weakly associated with the treatment; see Section 4.5 for more discussions. With a high probability, our estimated set $\hat{\mathcal{S}}$ in (11) belongs to \mathcal{S} and contains the set \mathcal{S}_{str} of strongly relevant IVs, defined as

$$\mathcal{S}_{\text{str}} = \left\{ 1 \leq j \leq p_z : |\gamma_j^*| \geq 2\sqrt{\log n} \cdot \sqrt{\mathbf{V}_{jj}^\gamma/n} \right\} \quad \text{with } \mathbf{V}^\gamma \text{ defined in (6)}. \quad (12)$$

The set \mathcal{S}_{str} contains the j -th IV if its individual strength $|\gamma_j^*|$ is well above its estimation accuracy $\sqrt{\mathbf{V}_{jj}^\gamma/n}$. The factor 2 in (12) ensures that the individual IV strength is sufficiently large such that the j -th IV can be included into the set $\hat{\mathcal{S}}$ defined in (11).

We now modify Condition 1 to accommodate the finite-sample uncertainty.

Condition 3 (Finite-sample Majority Rule) *More than half of the relevant IVs are strongly relevant and valid, that is, $|\mathcal{V} \cap \mathcal{S}_{\text{str}}| > |\mathcal{S}|/2$, where \mathcal{S} and \mathcal{V} are defined in (3) and \mathcal{S}_{str} is defined in (12).*

Condition 3 is slightly stronger than Condition 1 for a relatively small sample size n . When $n \rightarrow \infty$ and the IV strengths $\{\gamma_j^*\}_{1 \leq j \leq p_z}$ do not change with n , the set \mathcal{S}_{str} converges to \mathcal{S} and Condition 3 is the same as Condition 1.

Remark 1 The hard thresholding [18, 19, 56] is a prevalent tool in support recovery, which is recovering the set of relevant IVs in our problem. There are other choices of the threshold level in (11). We can replace $\sqrt{\log n}$ in (11) and (12) by any $f(n)$ satisfying $f(n) \rightarrow \infty$ with $n \rightarrow \infty$ and $f(n) > \sqrt{2 \log p_z}$. [25] used $f(n) = \sqrt{2.01 \log \max\{n, p_z\}}$, which is applicable in both low and high dimensions. In Section A.1 in the supplement, we consider the settings with weak IVs and demonstrate that our proposed methods have nearly the same performance with both the thresholds in (11) and that in [25].

4.2 The searching confidence interval

In the following, we restrict our attention to the estimated set $\widehat{\mathcal{S}}$ of relevant IVs and leverage Condition 3 to devise the searching CI for β^* . The main idea is to search for β values that lead to the majority of the instruments being detected as valid. The searching idea is highly relevant to the Anderson-Rubin test [1] for the weak IV problem, which searches for a range of β values leading to a sufficiently small χ^2 test statistic; see the following Remark 2 for more discussions.

For any given $\beta \in \mathbb{R}$, we apply the relation $\Gamma_j^* = \beta^* \gamma_j^* + \pi_j^*$ and construct the initial estimator of π_j^* as $\widehat{\Gamma}_j - \beta \widehat{\gamma}_j$. The estimation error of $\widehat{\Gamma}_j - \beta \widehat{\gamma}_j$ is decomposed as,

$$(\widehat{\Gamma}_j - \beta \widehat{\gamma}_j) - \pi_j^* = \widehat{\Gamma}_j - \Gamma_j^* - \beta(\widehat{\gamma}_j - \gamma_j^*) + (\beta^* - \beta)\gamma_j^*. \quad (13)$$

When $\beta = \beta^*$, the above estimation error is further simplified as $\widehat{\Gamma}_j - \Gamma_j^* - \beta(\widehat{\gamma}_j - \gamma_j^*)$. We quantify the uncertainty of $\{\widehat{\Gamma}_j - \Gamma_j^* - \beta(\widehat{\gamma}_j - \gamma_j^*)\}_{j \in \widehat{\mathcal{S}}}$ by the following union bound,

$$\liminf_{n \rightarrow \infty} \mathbf{P} \left[\max_{j \in \widehat{\mathcal{S}}} \frac{|\widehat{\Gamma}_j - \Gamma_j^* - \beta(\widehat{\gamma}_j - \gamma_j^*)|}{\sqrt{(\widehat{\mathbf{V}}_{jj}^\Gamma + \beta^2 \widehat{\mathbf{V}}_{jj}^\gamma - 2\beta \widehat{\mathbf{C}}_{jj})/n}} \leq \Phi^{-1} \left(1 - \frac{\alpha}{2|\widehat{\mathcal{S}}|} \right) \right] \geq 1 - \alpha, \quad (14)$$

where $\alpha \in (0, 1)$ and Φ^{-1} is the inverse CDF of the standard normal distribution. We rescale the upper bound in (14) and construct the following threshold for $\widehat{\Gamma}_j - \Gamma_j^* - \beta(\widehat{\gamma}_j - \gamma_j^*)$ with $j \in \widehat{\mathcal{S}}$,

$$\widehat{\rho}_j(\beta) := \Phi^{-1} \left(1 - \frac{\alpha}{2|\widehat{\mathcal{S}}|} \right) \cdot \sqrt{(\widehat{\mathbf{V}}_{jj}^\Gamma + \beta^2 \widehat{\mathbf{V}}_{jj}^\gamma - 2\beta \widehat{\mathbf{C}}_{jj})/n}. \quad (15)$$

We further apply the hard thresholding to $\{\widehat{\Gamma}_j - \beta \widehat{\gamma}_j\}_{j \in \widehat{\mathcal{S}}}$ and estimate π_j^* by

$$\widehat{\pi}_j(\beta) = \left(\widehat{\Gamma}_j - \beta \widehat{\gamma}_j \right) \cdot \mathbf{1} \left(\left| \widehat{\Gamma}_j - \beta \widehat{\gamma}_j \right| \geq \widehat{\rho}_j(\beta) \right) \quad \text{for } j \in \widehat{\mathcal{S}}. \quad (16)$$

If $\beta = \beta^*$, the hard thresholding in (16) guarantees $\widehat{\pi}_j(\beta^*) = 0$ for $j \in \mathcal{V}$ with probability larger than $1 - \alpha$. For any β , we can construct the vector $\widehat{\pi}_{\widehat{\mathcal{S}}}(\beta) = (\widehat{\pi}_j(\beta))_{j \in \widehat{\mathcal{S}}}$ and calculate the number of non-zero entries in $\widehat{\pi}_{\widehat{\mathcal{S}}}(\beta)$, denoted as $\|\widehat{\pi}_{\widehat{\mathcal{S}}}(\beta)\|_0$. Due to the hard thresholding, $\widehat{\pi}_{\widehat{\mathcal{S}}}(\beta)$ can be a sparse vector and $\|\widehat{\pi}_{\widehat{\mathcal{S}}}(\beta)\|_0$ stands for the number of invalid IVs for the given β .

We search for a range of β such that the number of invalid IVs is below $|\widehat{\mathcal{S}}|/2$,

$$\text{CI}^{\text{sear}} := \left\{ \beta \in \mathbb{R} : \|\widehat{\pi}_{\widehat{\mathcal{S}}}(\beta)\|_0 < |\widehat{\mathcal{S}}|/2 \right\}. \quad (17)$$

The above searching CI is valid since under Condition 3, $\|\widehat{\pi}_{\widehat{\mathcal{S}}}(\beta^*)\|_0 < |\widehat{\mathcal{S}}|/2$ with probability $1 - \alpha$; see equation (54) in the supplement. We consider the following example and illustrate the construction of CI^{sear} in Figure 4.

Example 2 Generate the models (1) and (2) with no baseline covariates, set $\beta^* = 1$, $n = 2000$, $\gamma_j^* = 0.5 \cdot \mathbf{1}_{10}$ and $\pi^* = (\mathbf{0}_6^\top, 0.05, 0.05, -0.5, -1)^\top$. In Figure 4, we plot $|\widehat{\mathcal{S}}| - \|\widehat{\pi}_{\widehat{\mathcal{S}}}(\beta)\|_0$ over different β and the red interval (0.914, 1.112) covers $\beta^* = 1$.




Figure 4: The x-axis plots a range of β values and the y-axis plots the number of valid IVs (i.e. $|\widehat{\mathcal{S}}| - \|\widehat{\pi}_{\widehat{\mathcal{S}}}(\beta)\|_0$) for every given β . The red interval (0.914, 1.112) denotes CI^{sear} in (17).

Remark 2 The proposed searching idea is related to but different from the Anderson-Rubin test [1, 41] for the weak IV problem. In (17), we use the sparsity as the test statistic and invert the sparsity level to determine the confidence region for β^* . Our inverted test statistics $\|\widehat{\pi}_{\widehat{\mathcal{S}}}(\beta)\|_0$ is a discrete function of β and its limiting distribution is not standard, which is fundamentally different from the χ^2 statistics used in AR test; see Figure 4. Due to the discreteness of the test statistics, the searching interval in (17) can be a union of disjoint intervals.

4.3 Efficient implementation of the searching CI

We now propose a computationally efficient implementation of CI^{sear} in (17). To implement (17), we can enumerate all values of β and calculate $\|\widehat{\pi}_{\widehat{\mathcal{S}}}(\beta)\|_0$. However, the enumeration method for constructing CI^{sear} can be time consuming if there is a huge amount of β values. To improve the computational efficiency, we construct an initial set $[L, U]$ such that $\mathbf{P}(\beta^* \in [L, U]) \rightarrow 1$ and construct a grid set with the grid size n^{-a} for $a > 1/2$. Then we discretize $[L, U]$ into the following grid set,

$$\mathcal{B} = \{\beta_1, \beta_2, \dots, \beta_K\} \quad \text{with } \beta_1 = L, \beta_K = U, |\beta_{j+1} - \beta_j| = n^{-a} \text{ for } 1 \leq j \leq K - 2, \quad (18)$$

and $|\beta_K - \beta_{K-1}| \leq n^{-a}$. The reason of requiring $a > 1/2$ is to ensure that the approximation error due to interval discretization is smaller than the parametric rate $n^{-1/2}$. As an important remark, the constructed confidence interval is almost invariant to the choices of $[L, U]$ and the grid size n^{-a} as long as $[L, U]$ contains β^* and $a > 1/2$. We demonstrate the invariance in Table 2 in Section 7.

Throughout the paper, we choose the default grid size $n^{-0.6}$ and present the following default construction of the initial range $[L, U]$. For $j \in \widehat{\mathcal{S}}$, we estimate β^* by the ratio $\widehat{\Gamma}_j/\widehat{\gamma}_j$ and estimate its variance by $\widehat{\text{Var}}\left(\widehat{\Gamma}_j/\widehat{\gamma}_j\right) = \frac{1}{n} \left(\widehat{\mathbf{V}}_{jj}^{\Gamma}/\widehat{\gamma}_j^2 + \widehat{\mathbf{V}}_{jj}^{\gamma}\widehat{\Gamma}_j^2/\widehat{\gamma}_j^4 - 2\widehat{\mathbf{C}}_{jj}\widehat{\Gamma}_j/\widehat{\gamma}_j^3 \right)$. We then construct L and U as

$$L = \min_{j \in \widehat{\mathcal{S}}} \left\{ \widehat{\Gamma}_j/\widehat{\gamma}_j - \sqrt{\log n \cdot \widehat{\text{Var}}\left(\widehat{\Gamma}_j/\widehat{\gamma}_j\right)} \right\}, \quad U = \max_{j \in \widehat{\mathcal{S}}} \left\{ \widehat{\Gamma}_j/\widehat{\gamma}_j + \sqrt{\log n \cdot \widehat{\text{Var}}\left(\widehat{\Gamma}_j/\widehat{\gamma}_j\right)} \right\}, \quad (19)$$

where $\sqrt{\log n}$ is used to adjust for multiplicity. Such an initial range has been discussed in Section 3 of [53].

With the grid set defined in (18), we modify (17) and propose the following computationally efficient implementation,

$$\widehat{\text{CI}}^{\text{sear}} = \left[\min_{\{\beta \in \mathcal{B}: \|\widehat{\pi}_{\widehat{\mathcal{S}}}(\beta)\|_0 < |\widehat{\mathcal{S}}|/2\}} \beta, \max_{\{\beta \in \mathcal{B}: \|\widehat{\pi}_{\widehat{\mathcal{S}}}(\beta)\|_0 < |\widehat{\mathcal{S}}|/2\}} \beta \right]. \quad (20)$$

Instead of searching over the real line as in (17), we restrict our attention to the grid set \mathcal{B} defined in (18) and (19) and search for the smallest and largest grid value such that more than half of the relevant IVs are valid. In contrast to CI^{sear} in (17), $\widehat{\text{CI}}^{\text{sear}}$ is guaranteed to be an interval. When the majority rule is violated, there is no β such that $\|\widehat{\pi}_{\widehat{\mathcal{S}}}(\beta)\|_0 < |\widehat{\mathcal{S}}|/2$ and hence $\widehat{\text{CI}}^{\text{sear}}$ in (20) is empty. This indicates that the majority rule is violated, which can be used as a partial check of the majority rule.

We summarize the construction of searching CI in Algorithm 1. As a remark, Algorithm 1 can be implemented with the summary statistics $\widehat{\Gamma}, \widehat{\gamma}$ and $\widehat{\mathbf{V}}^\Gamma, \widehat{\mathbf{V}}^\gamma, \widehat{\mathbf{C}}$, which are the inputs for steps 2 to 6 of Algorithm 1.

Algorithm 1 Searching CI (Uniform Inference under Majority Rule)

Input: Outcome $Y \in \mathbb{R}^n$; Treatment $D \in \mathbb{R}^n$; IVs $Z \in \mathbb{R}^{n \times p_z}$; Covariates $X \in \mathbb{R}^{n \times p_x}$; Significance level $\alpha \in (0, 1)$.

Output: Confidence interval $\widehat{\text{CI}}^{\text{sear}}$; Majority rule check $R \in \{0, 1\}$.

- 1: Construct $\widehat{\Gamma} \in \mathbb{R}^{p_z}, \widehat{\gamma} \in \mathbb{R}^{p_z}$ as in (5) and $\widehat{\mathbf{V}}^\Gamma, \widehat{\mathbf{V}}^\gamma$ and $\widehat{\mathbf{C}}$ as in (7);
 - 2: Construct $\widehat{\mathcal{S}}$ as in (11); ▷ First-stage selection
 - 3: Construct L and U as in (19);
 - 4: Construct the grid set \mathcal{B} as in (18) with $a = 0.6$;
 - 5: Construct $\{\widehat{\pi}_j(\beta)\}_{j \in \widehat{\mathcal{S}}, \beta \in \mathcal{B}}$ as in (16);
 - 6: Construct $\widehat{\text{CI}}^{\text{sear}}$ as in (20) and set $R = \mathbf{1}(\widehat{\text{CI}}^{\text{sear}} \neq \emptyset)$. ▷ Searching CI
-

4.4 The Sampling CI

In this subsection, we devise a novel sampling method to improve the precision of the searching CI. Conditioning on the observed data, we resample $\{\widehat{\Gamma}^{[m]}, \widehat{\gamma}^{[m]}\}_{1 \leq m \leq M}$ as

$$\begin{pmatrix} \widehat{\Gamma}^{[m]} \\ \widehat{\gamma}^{[m]} \end{pmatrix} \stackrel{\text{i.i.d.}}{\sim} N \left[\begin{pmatrix} \widehat{\Gamma} \\ \widehat{\gamma} \end{pmatrix}, \begin{pmatrix} \widehat{\mathbf{V}}^\Gamma/n & \widehat{\mathbf{C}}/n \\ \widehat{\mathbf{C}}^\top/n & \widehat{\mathbf{V}}^\gamma/n \end{pmatrix} \right] \quad \text{for } 1 \leq m \leq M, \quad (21)$$

where M denotes the resampling size (with the default value 1000), $\widehat{\gamma}$ and $\widehat{\Gamma}$ are defined in (5), and $\widehat{\mathbf{V}}^\Gamma, \widehat{\mathbf{C}}$ and $\widehat{\mathbf{V}}^\gamma$ are defined in (7).

For $1 \leq m \leq M$, we use $\{\widehat{\Gamma}^{[m]}, \widehat{\gamma}^{[m]}\}$ to replace $\{\widehat{\Gamma}, \widehat{\gamma}\}$ and implement Algorithm 1 to construct a searching CI. We refer to this CI as the m -th sampled searching CI. In implementing Algorithm 1, we decrease the thresholding level used in the hard thresholding step of estimating π^* . Consequently, each sampled searching CI can be much shorter than the original searching

CI; see the following Figure 5. Our proposal of decreasing the threshold relies on the following observation: there exists $1 \leq m^* \leq M$ such that

$$\max_{j \in \widehat{\mathcal{S}}} \frac{|\widehat{\Gamma}_j^{[m^*]} - \Gamma_j^* - \beta(\widehat{\gamma}_j^{[m^*]} - \gamma_j^*)|}{\sqrt{(\widehat{\mathbf{V}}_{jj}^\Gamma + \beta^2 \widehat{\mathbf{V}}_{jj}^\gamma - 2\beta \widehat{\mathbf{C}}_{jj})/n}} \leq \lambda \cdot \Phi^{-1} \left(1 - \frac{\alpha}{2|\widehat{\mathcal{S}}|} \right) \quad \text{with} \quad \lambda \asymp \left(\frac{\log n}{M} \right)^{\frac{1}{2|\widehat{\mathcal{S}}|}}. \quad (22)$$

The rigorous statement is provided in Proposition 1 in Section 6.1. Compared with (14), the upper bound in (22) is shrunk by a factor λ , which is close to zero with a large M (e.g., $M = 1000$). If we had access to $\{\widehat{\Gamma}^{[m^*]}, \widehat{\gamma}^{[m^*]}\}$, we could use a much smaller threshold level in constructing the searching CI.

With the above observation, we now provide the complete details about constructing the sampling CI. We shall consider that the tuning parameter λ in (22) is given to facilitate the discussion. We will present a data-dependent way of choosing λ in the following Remark 3. Motivated by (22), we multiply the threshold level by λ and implement Algorithm 1 for each of $\{\widehat{\Gamma}^{[m]}, \widehat{\gamma}^{[m]}\}_{1 \leq m \leq M}$. Particularly, for $1 \leq m \leq M$, we modify (16) and estimate π^* by

$$\widehat{\pi}_j^{[m]}(\beta, \lambda) = \left(\widehat{\Gamma}_j^{[m]} - \beta \widehat{\gamma}_j^{[m]} \right) \cdot \mathbf{1} \left(\left| \widehat{\Gamma}_j^{[m]} - \beta \widehat{\gamma}_j^{[m]} \right| \geq \lambda \cdot \widehat{\rho}_j(\beta) \right) \quad \text{for} \quad 1 \leq j \leq |\widehat{\mathcal{S}}|, \quad (23)$$

with $\widehat{\rho}_j(\beta)$ defined in (15). We apply (20) and construct the m -th sampled searching CI as $[\beta_{\min}^{[m]}(\lambda), \beta_{\max}^{[m]}(\lambda)]$ with

$$\beta_{\min}^{[m]}(\lambda) = \min_{\{\beta \in \mathcal{B}: \|\widehat{\pi}_{\widehat{\mathcal{S}}}^{[m]}(\beta, \lambda)\|_0 < |\widehat{\mathcal{S}}|/2\}} \beta \quad \text{and} \quad \beta_{\max}^{[m]}(\lambda) = \max_{\{\beta \in \mathcal{B}: \|\widehat{\pi}_{\widehat{\mathcal{S}}}^{[m]}(\beta, \lambda)\|_0 < |\widehat{\mathcal{S}}|/2\}} \beta. \quad (24)$$

Similarly to (20), we set $[\beta_{\min}^{[m]}(\lambda), \beta_{\max}^{[m]}(\lambda)] = \emptyset$ if there is no β such that $\|\widehat{\pi}^{[m]}(\beta, \lambda)\|_0 < |\widehat{\mathcal{S}}|/2$. We aggregate the M searching CIs and propose the following sampling CI:

$$\text{CI}^{\text{samp}} = \left[\min_{m \in \mathcal{M}} \beta_{\min}^{[m]}(\lambda), \max_{m \in \mathcal{M}} \beta_{\max}^{[m]}(\lambda) \right], \quad (25)$$

where the index set $\mathcal{M} = \{1 \leq m \leq M : [\beta_{\min}^{[m]}(\lambda), \beta_{\max}^{[m]}(\lambda)] \neq \emptyset\}$ denotes all indexes corresponding to non-empty $[\beta_{\min}^{[m]}(\lambda), \beta_{\max}^{[m]}(\lambda)]$.

In Figure 5, we demonstrate the sampling CI in (25) using Example 2. Many of the M sampled searching CIs are empty, and the non-empty ones can be much shorter than the searching CI. As a consequence, the sampling CI (in red) is much shorter than the searching CI (in blue); see Table 1 and Figure 6 in Section 7 for details. An important reason for the sampling method improving the precision is that the test statistics used in constructing the searching CI is a discrete function of β and the limiting distribution of the test statistics is not standard.




Figure 5: The axis corresponds to sampling indexes $\{1, 2, \dots, 102\}$ (after re-ordering) and the y-axis reports the sampled CIs. The red interval is $\text{CI}^{\text{samp}} = (0.955, 1.060)$, the blue interval is $\widehat{\text{CI}}^{\text{sear}} = (0.914, 1.112)$, and the blue dashed interval is the oracle CI $(0.983, 1.052)$ with prior information on valid IVs.

We summarize the sampling CI in Algorithm 2 and will provide two remarks about the implementation of Algorithm 2.

Algorithm 2 Sampling CI (Uniform Inference under Majority Rule)

Input: Outcome $Y \in \mathbb{R}^n$; Treatment $D \in \mathbb{R}^n$; IVs $Z \in \mathbb{R}^{n \times p_z}$; Covariates $X \in \mathbb{R}^{n \times p_x}$;
Sampling number $M=1000$; $\lambda = c_* (\log n/M)^{1/(2|\widehat{\mathcal{S}}|)}$; Significance level $\alpha \in (0, 1)$

Output: Confidence interval CI^{samp}

- 1: Implement steps 1 to 4 as in Algorithm 1;
 - 2: **for** $m \leftarrow 1$ to M **do**
 - 3: Sample $\widehat{\Gamma}^{[m]}$ and $\widehat{\gamma}^{[m]}$ as in (21);
 - 4: Compute $\{\widehat{\pi}_j^{[m]}(\beta, \lambda)\}_{j \in \widehat{\mathcal{S}}, \beta \in \mathcal{B}}$ as in (23);
 - 5: Compute $\beta_{\min}^{[m]}(\lambda)$ and $\beta_{\max}^{[m]}(\lambda)$ in (24);
 - 6: **end for**
 - 7: Construct CI^{samp} as in (25) ▷ Sampling CI
-

Remark 3 (Tuning parameter selection for sampling) We demonstrate in Section A.2 in the supplement that the sampling CIs in Algorithm 2 do not materially change with different choices of L, U, a and the resampling number M . The following Theorem 2 suggests the form of the tuning parameter λ as $c_* (\log n/M)^{1/(2|\widehat{\mathcal{S}}|)}$. We present a data-dependent way to specify the constant c_* . If the λ value is too small, very few of the resampled reduced-form estimators will pass the majority rule and most of the $M = 1000$ sampled intervals will be empty. Hence, the proportion of the non-empty intervals is used as an indicator for whether λ is large enough. We start with a small value $\lambda = 1/6 \cdot (\log n/M)^{1/(2|\widehat{\mathcal{S}}|)}$ and increase the value of λ by a factor of 1.25 until more than **prop** = 10% of the $M = 1000$ intervals are non-empty. We choose the smallest λ value achieving this and use this λ value to implement Algorithm 2. In Figure 7 in Section 7, we demonstrate the robustness of our procedure to different λ values.

Remark 4 (Alternative aggregation) We may combine the sampled CIs as $\text{CI}_0^{\text{samp}} = \cup_{m=1}^M [\beta_{\min}^{[m]}(\lambda), \beta_{\max}^{[m]}(\lambda)]$. Since $\text{CI}_0^{\text{samp}}$ may not be an interval due to its definition, we focus on CI^{samp} defined in (25). In addition, we may filter out $\{\widehat{\gamma}^{[m]}, \widehat{\Gamma}^{[m]}\}_{1 \leq m \leq M}$ near the boundary of the sampling distribution in (21). Particularly, define

$$\mathcal{M}_0 = \left\{ 1 \leq m \leq M : \max_{j \in \widehat{\mathcal{S}}} \max \left\{ \left| \widehat{\gamma}_j^{[m]} - \widehat{\gamma}_j \right| / \sqrt{\widehat{\mathbf{V}}_{jj}^\gamma / n}, \left| \widehat{\Gamma}_j^{[m]} - \widehat{\Gamma}_j \right| / \sqrt{\widehat{\mathbf{V}}_{jj}^\Gamma / n} \right\} \leq 1.1 \Phi^{-1} \left(1 - \frac{\alpha_0}{4|\widehat{\mathcal{S}}|} \right) \right\}$$

with $\alpha_0 = 0.05$. We modify (25) and construct the sampling CI as

$$\text{CI}^{\text{samp}}(\mathcal{M}_0) = \left[\min_{m \in \mathcal{M}'} \beta_{\min}^{[m]}(\lambda), \max_{m \in \mathcal{M}'} \beta_{\max}^{[m]}(\lambda) \right], \quad (26)$$

with the index set $\mathcal{M}' = \{m \in \mathcal{M}_0 : [\beta_{\min}^{[m]}(\lambda), \beta_{\max}^{[m]}(\lambda)] \neq \emptyset\}$. In Table A.6 in the supplement, we show that $\text{CI}^{\text{samp}}(\mathcal{M}_0)$ is nearly the same as CI^{samp} in (25).

4.5 Robustness to the existence of weak IVs

Our proposed searching and sampling CIs in Sections 4.2 to 4.4 are robust to both weak and locally invalid IVs. The validity of our proposed CIs relies on Condition 3, which requires more than half of the IVs to be strongly relevant and valid but allows the remaining IVs to be arbitrarily weak and/or invalid. In Section A.1 in the supplement, we demonstrate that our proposed CIs achieve the desired coverage even if some IVs are weakly associated with the treatment; see Table A.2 in the supplement for details.

In the following, we make two important remarks about weak IVs. Firstly, our focused setting is completely different from the classical weak IV framework [45, 46, e.g.], where all IVs as a total are weak. The F test for the treatment model and the concentration parameters can provide evidence on whether the IVs are weak [46, e.g.]. Similarly, the estimated set $\widehat{\mathcal{S}}$ in (11) can also be used to check whether the given data set falls into the classical weak IV framework. If $\widehat{\mathcal{S}}$ is non-empty, this indicates the existence of strong IVs. This paper focuses on the regime where $\widehat{\mathcal{S}}$ is non-empty. In studying the education returns in Section 8, we observe that the selected set of IVs have significant concentration parameters, indicating that we are not facing the classical weak IV problem.

Secondly, the challenge due to weak IVs is fundamentally different from locally invalid IVs. In the first-stage selection, the set of relatively weak IVs $\mathcal{S} \setminus \mathcal{S}_{\text{str}}$ are not guaranteed to be selected by $\widehat{\mathcal{S}}$ in (11). The uncertainty of selecting IVs inside $\mathcal{S} \setminus \mathcal{S}_{\text{str}}$ is of a different nature from the uncertainty of detecting locally invalid IVs. In Section A.1 in the supplement, we demonstrate that both TSHT and CIIV perform well even if there exists uncertainty of selecting IVs belonging to $\mathcal{S} \setminus \mathcal{S}_{\text{str}}$.

5 Uniform Inference Methods under Plurality Rule

This section considers the more challenging setting under the plurality rule. We introduce the finite-sample plurality rule and then generalize the method proposed in Section 4. For a given

invalidity level $v \in \mathbb{R}$, we define the set of IVs,

$$\mathcal{I}(v, \tau) = \{j \in \mathcal{S} : |\pi_j^*/\gamma_j^* - v| \leq \tau\} \quad \text{with } \tau \in \mathbb{R}. \quad (27)$$

When τ is small, $\mathcal{I}(v, \tau)$ denotes the set of IVs with the invalidity level around v . With $\mathbf{T}_{j,k}$ defined in (9), we define the separation level as

$$\text{sep}(n) := 2\sqrt{\log n} \max_{j,k \in \widehat{\mathcal{S}}} \mathbf{T}_{j,k}. \quad (28)$$

For a given n , $\text{sep}(n)$ denotes the separation level such that the IVs can be separated into groups based on the identified effects $\{\beta^{[j]}\}_{j \in \widehat{\mathcal{S}}}$ with $\beta^{[j]} = \Gamma_j^*/\gamma_j^*$. Since $\beta^{[j]} = \beta^* + \pi_j^*/\gamma_j^*$, we have $\beta^{[j]} - \beta^{[k]} = \pi_j^*/\gamma_j^* - \pi_k^*/\gamma_k^*$. In (28), $\max_{j,k \in \widehat{\mathcal{S}}} \mathbf{T}_{j,k}$ denotes the largest error of estimating the pairwise difference $\{\beta^{[j]} - \beta^{[k]}\}_{j,k \in \widehat{\mathcal{S}}}$ or equivalently $\{\pi_j^*/\gamma_j^* - \pi_k^*/\gamma_k^*\}_{j,k \in \widehat{\mathcal{S}}}$. The term $\sqrt{\log n}$ is used to adjust the multiplicity of hypothesis testing and the factor 2 in front of $\sqrt{\log n}$ is to ensure that the difference $\pi_j^*/\gamma_j^* - \pi_k^*/\gamma_k^*$ is sufficiently large such that it can be detected from the data; see more discussion after Proposition 2 in Section 6.2. When $\{\gamma_j^*\}_{j \in \widehat{\mathcal{S}}}$ are non-zero constants, $\text{sep}(n)$ is of order $\sqrt{\log n/n}$.

We now introduce the finite-sample plurality rule.

Condition 4 (Finite-sample Plurality Rule) For $\tau_n = 3\text{sep}(n)$,

$$|\mathcal{V} \cap \mathcal{S}_{\text{str}}| > \max_{v \in \mathbb{R}} |\mathcal{I}(v, \tau_n)/\mathcal{V}| \quad (29)$$

where $\text{sep}(n)$, \mathcal{V} , \mathcal{S}_{str} and $\mathcal{I}(v, \tau_n)$ are defined in (28), (3), (12) and (27), respectively.

In (29), the set $\mathcal{V} \cap \mathcal{S}_{\text{str}}$ consists of the strongly relevant and valid instruments, which we can rely on to make inference for β^* . Since $\tau_n \rightarrow 0$, the set $\mathcal{I}(v, \tau_n)/\mathcal{V}$ contains all invalid instruments with invalidity levels $\pi_j^*/\gamma_j^* \approx v$. When $v = 0$, the set $\mathcal{I}(0, \tau_n)/\mathcal{V}$ is the set of invalid IVs whose invalidity levels are below τ_n . Condition 4 requires that the cardinality of $\mathcal{V} \cap \mathcal{S}_{\text{str}}$ is larger than the number of invalid IVs with $\pi_j^*/\gamma_j^* \approx v$.

In comparison to the population plurality rule (Condition 2), Condition 4 accommodates for the finite-sample approximation error by grouping together invalid IVs with similar invalidity levels. For a large sample size, Condition 4 is the same as Condition 2. Specifically, if $n \rightarrow \infty$ and $\{\pi_j^*\}_{1 \leq j \leq p_z}$ and $\{\gamma_j^*\}_{1 \leq j \leq p_z}$ do not change with n , then

$$\lim_{n \rightarrow \infty} |\mathcal{V} \cap \mathcal{S}_{\text{str}}| = |\mathcal{V}|, \quad \lim_{n \rightarrow \infty} |\mathcal{I}(v, \tau_n)/\mathcal{V}| = |\mathcal{I}_\nu/\mathcal{V}| = \begin{cases} 0 & \text{if } \nu = 0 \\ |\mathcal{I}_\nu| & \text{if } \nu \neq 0 \end{cases}$$

with \mathcal{I}_ν defined in Condition 2. In the simulation, we also test the robustness of our proposed methods when the Condition 4 does not hold; see settings **S3** and **S5** in Section 7 for details. We shall emphasize that $\text{sep}(n)$ defined in (28) is not required in the following confidence interval construction.

5.1 Searching and sampling CIs under the plurality rule

In this subsection, we propose a two-step inference procedure for β^* under Condition 4. In the first step, we construct an initial set $\widehat{\mathcal{V}}$ satisfying

$$\mathcal{V} \cap \mathcal{S}_{\text{str}} \subset \widehat{\mathcal{V}} \subset \mathcal{I}(0, 3\text{sep}(n)). \quad (30)$$

Since Condition 4 implies that $\mathcal{V} \cap \mathcal{S}_{\text{str}}$ is the majority of the set $\mathcal{I}(0, 3\text{sep}(n))$, we apply (30) and establish that the set $\mathcal{V} \cap \mathcal{S}_{\text{str}}$ becomes the majority of the initial set $\widehat{\mathcal{V}}$. In the second step, we restrict our attention to $\widehat{\mathcal{V}}$ and generalize the methods in Section 4. Importantly, the initial set $\widehat{\mathcal{V}}$ is allowed to include locally invalid IVs.

We now construct an initial set $\widehat{\mathcal{V}}$ satisfying (30) by modifying TSHT [25]. Without loss of generality, we set $\widehat{\mathcal{S}} = \{1, 2, \dots, |\widehat{\mathcal{S}}|\}$. For any $j \in \widehat{\mathcal{S}}$, we construct an estimator of β^* and π^* as

$$\widehat{\beta}^{[j]} = \widehat{\Gamma}_j / \widehat{\gamma}_j \quad \text{and} \quad \widehat{\pi}_k^{[j]} = \widehat{\Gamma}_k - \widehat{\beta}^{[j]} \widehat{\gamma}_k \quad \text{for } k \in \widehat{\mathcal{S}}, \quad (31)$$

where the super index j stands for the model identification by assuming the j -th IV to be valid. We define $\widehat{\mathbf{R}}^{[j]} = \widehat{\mathbf{V}}^\Gamma + (\widehat{\beta}^{[j]})^2 \widehat{\mathbf{V}}^\gamma - 2\widehat{\beta}^{[j]} \widehat{\mathbf{C}}$, and further estimate the standard error of $\widehat{\pi}_k^{[j]}$ with $k \in \widehat{\mathcal{S}}$ by

$$\widehat{\text{SE}}(\widehat{\pi}_k^{[j]}) = \sqrt{\left(\widehat{\mathbf{R}}_{k,k}^{[j]} + (\widehat{\gamma}_k / \widehat{\gamma}_j)^2 \widehat{\mathbf{R}}_{j,j}^{[j]} - 2\widehat{\gamma}_k / \widehat{\gamma}_j \widehat{\mathbf{R}}_{k,j}^{[j]} \right) / n}. \quad (32)$$

For $1 \leq k, j \leq |\widehat{\mathcal{S}}|$, we apply the following hard thresholding and construct the (k, j) entry of the voting matrix $\widehat{\Pi} \in \mathbb{R}^{|\widehat{\mathcal{S}}| \times |\widehat{\mathcal{S}}|}$ as

$$\widehat{\Pi}_{k,j} = \mathbf{1} \left(|\widehat{\pi}_k^{[j]}| \leq \widehat{\text{SE}}(\widehat{\pi}_k^{[j]}) \cdot \sqrt{\log n} \quad \text{and} \quad |\widehat{\pi}_j^{[k]}| \leq \widehat{\text{SE}}(\widehat{\pi}_j^{[k]}) \cdot \sqrt{\log n} \right), \quad (33)$$

with $\widehat{\pi}_k^{[j]}$ and $\widehat{\pi}_j^{[k]}$ defined in (31), $\widehat{\text{SE}}(\widehat{\pi}_k^{[j]})$ and $\widehat{\text{SE}}(\widehat{\pi}_j^{[k]})$ defined in (32), and $\sqrt{\log n}$ used to adjust for multiplicity. In (33), $\widehat{\Pi}_{k,j} = 1$ represents that the k -th and j -th IVs support each other to be valid while $\widehat{\Pi}_{k,j} = 0$ represents that they do not. The voting matrix in (33) is symmetric version of the voting matrix proposed in [25].

We now construct the initial set by leveraging the voting matrix in (33). Define $\widehat{\mathcal{W}} = \arg \max_{1 \leq j \leq |\widehat{\mathcal{S}}|} \|\widehat{\Pi}_j\|_0$ as the set of IVs receiving the largest number of votes. We construct the following initial set,

$$\widehat{\mathcal{V}}^{\text{TSHT}} := \{1 \leq l \leq |\widehat{\mathcal{S}}| : \text{there exists } 1 \leq k \leq |\widehat{\mathcal{S}}| \text{ such that } \widehat{\Pi}_{j,k} \widehat{\Pi}_{k,l} = 1 \text{ for } j \in \widehat{\mathcal{W}}\}. \quad (34)$$

In words, if the l -th IV from $\widehat{\mathcal{S}}$ and the j -th IV from $\widehat{\mathcal{W}}$ are claimed to be valid by any IV from $\widehat{\mathcal{S}}$, then the l -th IV is also included in $\widehat{\mathcal{V}}^{\text{TSHT}}$. We show in Proposition 2 in Section 6.2 that $\widehat{\mathcal{V}}^{\text{TSHT}}$ is guaranteed to satisfy (30).

Since $\mathcal{V} \cap \mathcal{S}_{\text{str}}$ is the majority of the initial set $\widehat{\mathcal{V}}^{\text{TSHT}}$ in (34), we restrict to $\widehat{\mathcal{V}}^{\text{TSHT}}$ and apply Algorithms 1 and 2 with $\widehat{\mathcal{S}}$ replaced by $\widehat{\mathcal{V}}^{\text{TSHT}}$. We summarize our proposed searching and sampling CIs in Algorithm 3.

The tuning parameters of Algorithm 3 are selected in the same way as that in Remark 3. In Section 7 and Section A in the supplement, we demonstrate that the constructed CIs do not

materially change for a wide range of tuning parameters. In Algorithm 3 can be implemented without requiring the raw data, but with $\widehat{\Gamma}, \widehat{\gamma}$ and $\widehat{\mathbf{V}}^\Gamma, \widehat{\mathbf{V}}^\gamma, \widehat{\mathbf{C}}$ as the inputs. We have demonstrated our method by constructing $\widehat{\mathcal{V}} = \widehat{\mathcal{V}}^{\text{TSHT}}$ as in (34). However, as long as the constructed $\widehat{\mathcal{V}}$ satisfies (30), our proposed CIs in Algorithm 3 are effective under Condition 4. In simulation studies, we investigate the finite-sample performance using $\widehat{\mathcal{V}}^{\text{CIIV}}$, the set of valid IVs selected by the CIIV method proposed in [53].

Algorithm 3 Uniform Inference with Searching and Sampling (Plurality Rule)

Input: Outcome $Y \in \mathbb{R}^n$; Treatment $D \in \mathbb{R}^n$; IVs $Z \in \mathbb{R}^{n \times p_z}$; Covariates $X \in \mathbb{R}^{n \times p_x}$;

Sampling number $M=1000$; $\lambda = c_* (\log n/M)^{1/(2|\widehat{\mathcal{S}}|)}$; Significance level $\alpha \in (0, 1)$.

Output: Confidence intervals $\widehat{\text{CI}}^{\text{sear}}$ and CI^{samp} ; Plurality rule check R .

- 1: Construct $\widehat{\Gamma} \in \mathbb{R}^{p_z}, \widehat{\gamma} \in \mathbb{R}^{p_z}$ as in (5) and $\widehat{\mathbf{V}}^\Gamma, \widehat{\mathbf{V}}^\gamma$ and $\widehat{\mathbf{C}}$ as in (7);
 - 2: Construct $\widehat{\mathcal{S}}$ as in (11);
 - 3: Construct the voting matrix $\widehat{\Pi} \in \mathbb{R}^{|\widehat{\mathcal{S}}| \times |\widehat{\mathcal{S}}|}$ as in (33);
 - 4: Construct $\widehat{\mathcal{V}} = \widehat{\mathcal{V}}^{\text{TSHT}}$ as in (34); ▷ Construction of $\widehat{\mathcal{V}}$
 - 5: Construct L and U as in (19) with $\widehat{\mathcal{S}} = \widehat{\mathcal{V}}$;
 - 6: Construct the grid set $\mathcal{B} \subset [L, U]$ as in (18) with the grid size $n^{-0.6}$;
 - 7: Compute $\{\widehat{\rho}_j(\beta)\}_{j \in \widehat{\mathcal{V}}, \beta \in \mathcal{B}}$ as in (15) with $\widehat{\mathcal{S}} = \widehat{\mathcal{V}}$;
 - 8: Compute $\{\widehat{\pi}_j(\beta)\}_{j \in \widehat{\mathcal{V}}, \beta \in \mathcal{B}}$ as in (16) with $\widehat{\mathcal{S}} = \widehat{\mathcal{V}}$;
 - 9: Construct $\widehat{\text{CI}}^{\text{sear}}$ as in (20) with $\widehat{\mathcal{S}} = \widehat{\mathcal{V}}$ and set $R = \mathbf{1}(\widehat{\text{CI}}^{\text{sear}} \neq \emptyset)$; ▷ Searching CI
 - 10: **for** $m \leftarrow 1$ to M **do**
 - 11: Sample $\widehat{\Gamma}^{[m]}$ and $\widehat{\gamma}^{[m]}$ as in (21);
 - 12: Compute $\{\widehat{\pi}_j^{[m]}(\beta, \lambda)\}_{j \in \widehat{\mathcal{V}}, \beta \in \mathcal{B}}$ as in (23) with $\widehat{\mathcal{S}} = \widehat{\mathcal{V}}$;
 - 13: Construct $\beta_{\min}^{[m]}(\lambda)$ and $\beta_{\max}^{[m]}(\lambda)$ as in (24) with $\widehat{\mathcal{S}} = \widehat{\mathcal{V}}$;
 - 14: **end for**
 - 15: Construct CI^{samp} as in (25). ▷ Sampling CI
-

5.2 Comparison with the CIIV method [53]

The idea of searching has been developed in [53] to select valid IVs using confidence interval methods. We now follow [53] and sketch the intuitive idea of the CIIV method. For any grid value $\delta_g \in [L, U]$, define the set

$$\widehat{\mathcal{V}}(\delta_g) = \{j \in \mathcal{S} : \Gamma_j^*/\gamma_j^* = \delta_g \text{ is not rejected}\}.$$

Here, $\widehat{\mathcal{V}}(\delta_g)$ denotes a subset of IVs such that the corresponding hypothesis $\Gamma_j^*/\gamma_j^* = \delta_g$ is not rejected. As explained in [53], the CIIV method examines all values of δ_g and selects the largest set $\widehat{\mathcal{V}}(\delta_g)$ as the set of valid IVs, that is,

$$\widehat{\mathcal{V}}^{\text{CIIV}} = \widehat{\mathcal{V}}(\widehat{\delta}_g) \quad \text{with} \quad \widehat{\delta}_g = \arg \max_{\delta_g \in [L, U]} |\widehat{\mathcal{V}}(\delta_g)|. \quad (35)$$

See Section 3 of [53] for more discussion.

Our proposed searching CI is different from [53] since we directly construct CIs by searching for a range of suitable β values, while the CIIV method applies the searching idea to select the set of valid IVs. As illustrated in Figure 3, the selected IVs may include locally invalid IVs, which leads to unreliable CIs.

We provide some intuitions on why our proposal is more robust to the IV selection error. We first screen out the strongly invalid IVs and construct an initial set estimator $\widehat{\mathcal{V}} = \widehat{\mathcal{V}}^{\text{TSHI}}$; then, we restrict to the set $\widehat{\mathcal{V}}$ and apply searching and sampling CIs developed under the majority rule. The majority rule in the second step explains the robustness: we compare the number of votes to $|\widehat{\mathcal{V}}|/2$, which is fixed after computing $\widehat{\mathcal{V}}$. However, the optimization in (35) chooses δ_g giving the largest number of votes, which can be more vulnerable to selection/testing errors. The validity of the CIIV method requires that $\widehat{\mathcal{V}}^{\text{CIIV}}$ does not contain any invalid IVs; in contrast, our method is still effective even if the initial set $\widehat{\mathcal{V}}$ contains the invalid IVs but satisfies (30).

6 Theoretical Justification

We focus on the low-dimensional setting with heteroscedastic errors and introduce the following regularity conditions.

- (C1) For $1 \leq i \leq n$, $W_i = (X_i^\top, Z_i^\top)^\top \in \mathbb{R}^p$ are i.i.d. Sub-gaussian random vectors with $\Sigma = \mathbf{E}(W_i W_i^\top)$ satisfying $c_0 \leq \lambda_{\min}(\Sigma) \leq \lambda_{\max}(\Sigma) \leq C_0$ for some positive constants $C_0 \geq c_0 > 0$.
- (C2) For $1 \leq i \leq n$, the errors $(\epsilon_i, \delta_i)^\top$ in (4) are i.i.d Sub-gaussian random vectors; the conditional covariance matrix satisfying $c_1 \leq \lambda_{\min}(\mathbf{E}[(\epsilon_i, \delta_i)^\top (\epsilon_i, \delta_i) | W_i]) \leq \lambda_{\max}(\mathbf{E}[(\epsilon_i, \delta_i)^\top (\epsilon_i, \delta_i) | W_i]) \leq C_1$ for some positive constants $C_1 \geq c_1 > 0$.

Conditions (C1) and (C2) are imposed on the reduced-form model (4), which includes the outcome model (1) and the treatment model (2) as a special case. We assume that the covariance matrix of W_i is well conditioned and also the covariance matrix of the errors is well conditioned. Condition (A2) in general holds if e_i in (1) and δ_i in (2) are not perfectly correlated. Our setting allows for the heteroscedastic errors. If we further assume $(\epsilon_i, \delta_i)^\top$ to be independent of W_i , Condition (C2) assumes the covariance matrix of $(\epsilon_i, \delta_i)^\top$ to be well conditioned. As a remark, the Sub-gaussian conditions on both W_i and the errors might be relaxed to the moment conditions in low dimensions.

6.1 Majority rule

In this subsection, we present the theory under the majority rule and the more general results under the plurality rule are presented in Section 6.2. The following theorem justifies the searching CI under the majority rule.

Theorem 1 *Consider the model (4). Suppose that Condition 3, Conditions (C1) and (C2) hold, and $\alpha \in (0, 1/4)$ is the significance level. Then CI^{sear} defined in (17) and $\widehat{\text{CI}}^{\text{sear}}$ defined*

in (20) satisfy

$$\liminf_{n \rightarrow \infty} \mathbf{P}(\beta^* \in \text{CI}^{\text{sear}}) \geq 1 - \alpha \quad \text{and} \quad \liminf_{n \rightarrow \infty} \mathbf{P}(\beta^* \in \widehat{\text{CI}}^{\text{sear}}) \geq 1 - \alpha.$$

There exists a positive constant $C > 0$ such that

$$\liminf_{n \rightarrow \infty} \mathbf{P} \left(\max\{\mathbf{L}(\text{CI}^{\text{sear}}), \mathbf{L}(\widehat{\text{CI}}^{\text{sear}})\} \leq \frac{C}{\min_{j \in \widehat{\mathcal{S}} \cap \mathcal{V}} |\gamma_j^*| \cdot \sqrt{n}} \right) \geq 1 - \alpha,$$

where $\mathbf{L}(\cdot)$ denotes the interval length.

If $\min_{j \in \widehat{\mathcal{S}} \cap \mathcal{V}} |\gamma_j^*|$ is of a constant order, then Theorem 1 implies that the searching CI is of the parametric rate $1/\sqrt{n}$. In contrast to TSHT [25] and CIIV [53], the inference result in Theorem 1 is uniformly valid. Particularly, the theoretical justification of TSHT and CIIV requires the invalidity levels of invalid IVs to be well separated from zero, which leads to all invalid IVs being correctly identified; see Assumption 8 in [25]. However, the searching CI guarantees the coverage property without requiring such well separation conditions.

We now turn to the sampling CI. Before presenting the theory for the sampling CI, we justify in the following Proposition 1 why we are able to decrease the thresholding level in constructing the sampling CI. For $\alpha_0 \in (0, 1/4)$, define the positive constant

$$c^*(\alpha_0) = \frac{1}{[3\pi \cdot \lambda_{\min}(\text{Cov})]^{|\widehat{\mathcal{S}}|}} \exp \left(-\frac{|\widehat{\mathcal{S}}| \cdot 3\lambda_{\max}(\text{Cov})}{\lambda_{\min}(\text{Cov})} \cdot \left[\Phi^{-1} \left(1 - \frac{\alpha_0}{4|\widehat{\mathcal{S}}|} \right) \right]^2 \right), \quad (36)$$

with Cov defined in (6) and Φ^{-1} denoting the inverse CDF of the standard normal distribution. Note that $c_1/C_0 \leq \lambda_{\min}(\text{Cov}) \leq \lambda_{\max}(\text{Cov}) \leq C_1/c_0$; see its proof in equations (42) and (41) in the supplement. Hence, for a fixed p_z and $\alpha_0 \in (0, 1)$, $c^*(\alpha_0)$ is a positive constant independent of n .

Proposition 1 *Suppose Conditions (C1) and (C2) hold and $\alpha_0 \in (0, 1/4)$. If $\text{err}_n(M, \alpha_0) = \left[\frac{2 \log n}{c^*(\alpha_0)M} \right]^{\frac{1}{2|\widehat{\mathcal{S}}|}} \rightarrow 0$ with $c^*(\alpha_0)$ defined in (36), then there exists a positive constant $C > 0$ such that*

$$\liminf_{n \rightarrow \infty} \mathbf{P} \left(\min_{1 \leq m \leq M} \left[\max_{\beta \in \mathcal{U}(a)} \max_{j \in \widehat{\mathcal{S}}} \frac{|\widehat{\Gamma}_j^{[m]} - \Gamma_j^* - \beta(\widehat{\gamma}_j^{[m]} - \gamma_j^*)|}{\sqrt{(\widehat{\mathbf{V}}_{jj}^\Gamma + \beta^2 \widehat{\mathbf{V}}_{jj}^\gamma - 2\beta \widehat{\mathbf{C}}_{jj})/n}} \right] \leq C \text{err}_n(M, \alpha_0) \right) \geq 1 - \alpha_0, \quad (37)$$

with $\mathcal{U}(a) := \{\beta \in \mathbb{R} : |\beta - \beta^*| \leq n^{-a}\}$ for any $a > 1/2$.

Since $c^*(\alpha_0)$ is of a constant order, $\text{err}_n(M, \alpha_0) \rightarrow 0$ if the resampling size M is much larger than $\log n$. The above proposition states that, there exists $1 \leq m^* \leq M$ such that

$$\max_{\beta \in \mathcal{U}(a)} \max_{j \in \widehat{\mathcal{S}}} \frac{|\widehat{\Gamma}_j^{[m^*]} - \Gamma_j^* - \beta(\widehat{\gamma}_j^{[m^*]} - \gamma_j^*)|}{\sqrt{(\widehat{\mathbf{V}}_{jj}^\Gamma + \beta^2 \widehat{\mathbf{V}}_{jj}^\gamma - 2\beta \widehat{\mathbf{C}}_{jj})/n}} \leq C \text{err}_n(M, \alpha_0) \rightarrow 0.$$

In comparison to (14), the threshold decreases from $\Phi^{-1}\left(1 - \frac{\alpha}{2|\widehat{\mathcal{S}}|}\right)$ to $C\text{err}_n(M, \alpha_0)$. A related sampling property was established in [23] to address a different nonstandard inference problem.

We now apply Proposition 1 to justify the sampling CI under the majority rule.

Theorem 2 *Suppose that the conditions of Proposition 1 hold and λ used in (24) satisfies $\lambda \geq 2C\text{err}_n(M, \alpha_0)/\Phi^{-1}[1 - \alpha/(2|\widehat{\mathcal{S}}|)]$ and $\lambda \gg n^{1/2-a}$ with the constant C used in (37) and $a > 1/2$. Then CI^{samp} defined in (25) and $\text{CI}^{\text{samp}}(\mathcal{M}_0)$ defined in (26) satisfy*

$$\liminf_{n \rightarrow \infty} \mathbf{P}(\beta^* \in \text{CI}^{\text{samp}}) \geq \liminf_{n \rightarrow \infty} \mathbf{P}(\beta^* \in \text{CI}^{\text{samp}}(\mathcal{M}_0)) \geq 1 - \alpha_0.$$

There exists a positive constant $C > 0$ such that

$$\liminf_{n \rightarrow \infty} \mathbf{P}\left(\max\left\{\frac{\mathbf{L}(\text{CI}^{\text{samp}})}{\sqrt{\log|\mathcal{M}|}}, \mathbf{L}(\text{CI}^{\text{samp}}(\mathcal{M}_0))\right\} \leq \frac{C}{\min_{j \in \widehat{\mathcal{S}} \cap \mathcal{V}} |\gamma_j^*| \cdot \sqrt{n}}\right) \geq 1 - \alpha_0.$$

Note that $2C\text{err}_n(M, \alpha_0)/\Phi^{-1}[1 - \alpha/(2|\widehat{\mathcal{S}}|)] = c(\log n/M)^{1/(2|\widehat{\mathcal{S}}|)}$ for some positive constant $c > 0$. Motivated by the condition $\lambda \geq 2C\text{err}_n(M, \alpha_0)/\Phi^{-1}[1 - \alpha/(2|\widehat{\mathcal{S}}|)]$, we choose the tuning parameter λ in the form $\lambda = c_*(\log n/M)^{1/(2|\widehat{\mathcal{S}}|)}$ in Remark 3. Similar to Theorem 1, Theorem 2 shows that our proposed searching CI does not require the well-separation condition on the invalidity levels. If the IV strengths $\{\gamma_j^*\}_{j \in \mathcal{S}}$ are assumed to be of a constant order, then the length of $\text{CI}^{\text{samp}}(\mathcal{M}_0)$ defined in (26) is $1/\sqrt{n}$. We can only establish the upper bound $\sqrt{\log|\mathcal{M}|/n}$ for the length of CI^{samp} defined in (25). However, we believe that this is mainly a technical artifact since $\text{CI}^{\text{samp}}(\mathcal{M}_0)$ and CI^{samp} are nearly the same in the numerical studies; see Section A.2 in the supplement.

6.2 Plurality Rule

We now consider the more challenging setting only assuming the finite-sample plurality rule (Condition 4). The main extra step is to show that our constructed initial set $\widehat{\mathcal{V}} = \widehat{\mathcal{V}}^{\text{TSHT}}$ satisfies (30). To establish this, we provide a careful finite-sample analysis of the voting scheme described in (33) and $\widehat{\mathcal{V}}^{\text{TSHT}}$ defined in (34).

Proposition 2 *Suppose that Conditions (C1) and (C2) hold. Consider the indexes $j \in \widehat{\mathcal{S}}$ and $k \in \widehat{\mathcal{S}}$.*

1. *If $\pi_k^*/\gamma_k^* = \pi_j^*/\gamma_j^*$, then $\liminf_{n \rightarrow \infty} \mathbf{P}\left(\widehat{\Pi}_{k,j} = \widehat{\Pi}_{j,k} = 1\right) = 1$.*
2. *If $\left|\pi_k^*/\gamma_k^* - \pi_j^*/\gamma_j^*\right| \geq 2\sqrt{\log n} \cdot \mathbf{T}_{j,k}$ with $\mathbf{T}_{j,k}$ defined in (9), then*

$$\liminf_{n \rightarrow \infty} \mathbf{P}\left(\widehat{\Pi}_{k,j} = \widehat{\Pi}_{j,k} = 0\right) = 1.$$

Under the additional Condition 4, the constructed $\widehat{\mathcal{V}} = \widehat{\mathcal{V}}^{\text{TSHT}}$ in (34) satisfies (30)

$$\liminf_{n \rightarrow \infty} \mathbf{P} \left(\mathcal{V} \cap \mathcal{S}_{\text{str}} \subset \widehat{\mathcal{V}} \subset \mathcal{I}(0, 3\text{sep}(n)) \right) = 1,$$

with $\text{sep}(n)$ defined in (28).

The above proposition shows that if two IVs are of the same invalidity level, then they vote for each other with a high probability. If two candidate IVs are well-separated (i.e. $|\pi_k^*/\gamma_k^* - \pi_j^*/\gamma_j^*| \geq 2\sqrt{\log n} \cdot \mathbf{T}_{j,k}$), then they vote against each other with a high probability. If $0 < |\pi_k^*/\gamma_k^* - \pi_j^*/\gamma_j^*| < 2\sqrt{\log n} \cdot \mathbf{T}_{j,k}$, there is no theoretical guarantee on how the two candidate IVs will vote. The above proposition explains why we are likely to make a mistake in detecting some locally invalid IVs defined in Definition 1.

With Proposition 2, we connect the finite-sample plurality rule to the finite-sample majority rule. We apply the voting algorithm to remove all strongly invalid IVs and the set $\widehat{\mathcal{V}}^{\text{TSHT}}$ only consists of valid IVs and the locally invalid IVs. The finite-sample plurality condition (Condition 4) assumes that the number of valid IVs is more than that of the locally invalid IVs, that is, the finite-sample majority rule is satisfied if we restrict to $\widehat{\mathcal{V}}^{\text{TSHT}}$. Then it is sufficient to apply the theoretical analysis of the majority rule by replacing $\widehat{\mathcal{S}}$ with $\widehat{\mathcal{V}}^{\text{TSHT}}$ or any $\widehat{\mathcal{V}}$ satisfying (30). The following theorem justifies sampling and searching CIs for the plurality rule setting.

Theorem 3 *Consider the model (4). Suppose that Condition 4, Conditions (C1) and (C2) hold. Then with $\widehat{\mathcal{S}}$ replaced by $\widehat{\mathcal{V}}^{\text{TSHT}}$, the coverage and precision properties in Theorem 1 hold for the searching interval in Algorithm 3 and the coverage and precision properties in Theorem 2 hold for the sampling interval in Algorithm 3.*

7 Simulation Studies

Throughout the numerical studies, we implement our proposed $\widehat{\text{CI}}^{\text{sear}}$ and CI^{samp} detailed in Algorithm 3. We use two different initial estimators of \mathcal{V} : $\widehat{\mathcal{V}}^{\text{TSHT}}$ defined in (34) and the set of valid IVs $\widehat{\mathcal{V}}^{\text{CIIV}}$ outputted by CIIV [53]. The code for replicating all numerical results in the current paper is available at <https://github.com/zijguo/Searching-Sampling-Replication>.

We compare with three existing CIs allowing for invalid IVs: TSHT [25], CIIV [53], and the Union method [31]. TSHT and CIIV are implemented with the codes on the Github websites¹ while the Union method is implemented with the code shared by the authors of [31]. The Union method takes a union of intervals that are constructed by a given number of candidate IVs and are not rejected by the Sargan test. An upper bound \bar{s} for the number of invalid IVs is required for the implementation. We consider two specific upper bounds: $\bar{s} = p_z - 1$ corresponds to the existence of two valid IVs, and $\bar{s} = \lceil p_z/2 \rceil$ corresponds to the majority rule being satisfied. We conduct 500 replications of simulations and compare different CIs in terms of empirical coverage and average lengths.

¹The code for TSHT is obtained from <https://github.com/hyunseungkang/invalidIV> and for CIIV is obtained from <https://github.com/xlbristol/CIIV>.

We implement two oracle methods as the benchmark. Firstly, we implement the oracle TSLS assuming the prior knowledge of \mathcal{V} . This method serves as a benchmark when the set \mathcal{V} of valid IVs are correctly recovered. Secondly, we implement the oracle bias-aware confidence interval in (10) assuming the oracle knowledge of the bias of TSHT estimator. We argue that the oracle bias-aware confidence interval serves as a better benchmark, especially when \mathcal{V} might not be correctly recovered in finite samples.

We generate the i.i.d. data $\{Y_i, D_i, Z_i, X_i\}_{1 \leq i \leq n}$ using the outcome model (1) and the treatment model (2). We generate $\gamma^* \in \mathbb{R}^{p_z}$ and $\pi^* \in \mathbb{R}^{p_z}$ as follows,

S1 (Majority rule): set $\gamma^* = \gamma_0 \cdot \mathbf{1}_{10}$ and $\pi^* = (\mathbf{0}_6, \tau \cdot \gamma_0, \tau \cdot \gamma_0, -0.5, -1)^\top$;

S2 (Plurality rule): set $\gamma^* = \gamma_0 \cdot \mathbf{1}_{10}$ and $\pi^* = (\mathbf{0}_4, \tau \cdot \gamma_0, \tau \cdot \gamma_0, -\frac{1}{3}, -\frac{2}{3}, -1, -\frac{4}{3})^\top$;

S3 (Plurality rule): set $\gamma^* = \gamma_0 \cdot \mathbf{1}_{10}$ and $\pi^* = (\mathbf{0}_4, \tau \cdot \gamma_0, \tau \cdot \gamma_0, -\frac{1}{6}, -\frac{1}{3}, -\frac{1}{2}, -\frac{2}{3})^\top$;

S4 (Plurality rule): set $\gamma^* = \gamma_0 \cdot \mathbf{1}_6$ and $\pi^* = (\mathbf{0}_2, -0.8, -0.4, \tau \cdot \gamma_0, 0.6)^\top$;

S5 (Plurality rule): set $\gamma^* = \gamma_0 \cdot \mathbf{1}_6$ and $\pi^* = (\mathbf{0}_2, -0.8, -0.4, \tau \cdot \gamma_0, \tau \cdot \gamma_0 + 0.1)^\top$.

The parameter γ_0 denotes the IV strength and is set as 0.5. The parameter τ denotes the invalidity level and is varied across $\{0.2, 0.4\}$. The setting **S1** satisfies the population majority rule while the settings **S2** to **S5** only satisfy the population plurality rule. Settings **S4** and **S5** represent the challenging settings where there are only two valid IVs. We introduce settings **S3** and **S5** to test the robustness of our proposed method when the finite-sample plurality rule might be violated. For example, for the setting **S5** with small n (e.g. $n = 500$), the invalid IVs with π_j^* values $\tau \cdot \gamma_0, \tau \cdot \gamma_0 + 0.1$ have similar invalidity levels and may violate Condition 4.

We now specify the remaining details for the generating models (1) and (2). Set $p_x = 10$, $\psi^* = (0.6, 0.7, \dots, 1.5)^\top \in \mathbb{R}^{10}$ in (2) and $\phi^* = (1.1, 1.2, \dots, 2)^\top \in \mathbb{R}^{10}$ in (1). We vary n across $\{500, 1000, 2000, 5000\}$. For $1 \leq i \leq n$, generate the covariates $W_i = (Z_i^\top, X_i^\top)^\top \in \mathbb{R}^p$ following a multivariate normal distribution with zero mean and covariance $\Sigma \in \mathbb{R}^{p \times p}$ where $\Sigma_{jl} = 0.5^{|j-l|}$ for $1 \leq j, l \leq p$; generate the errors $(e_i, \delta_i)^\top$ following bivariate normal with zero mean, unit variance and $\text{Cov}(e_i, \delta_i) = 0.8$.

In Table 1, we report the empirical coverage and interval length for $\tau = 0.2$. The CIs by TSHT and CIIV undercover for $n = 500, 1000$ and 2000 and only achieve the 95% coverage level for a large sample size $n = 5000$. Our proposed searching and sampling CIs achieve the desired coverage levels in most settings. For settings **S1** to **S4**, both initial estimates of set of valid IVs $\hat{\mathcal{V}}^{\text{TSHT}}$ and $\hat{\mathcal{V}}^{\text{CIIV}}$ lead to CIs achieving the 95% coverage level. For the more challenging settings **S5**, the empirical coverage level of our proposed searching and sampling CIs achieve the desired coverage with sample sizes above 2000. For $n = 500$ and $n = 1000$, our proposed searching and sampling methods improve the coverage of TSHT and CIIV. The undercoverage happens mainly due to the fact that the finite-sample plurality rule might fail for setting **S5** with a relatively small sample size. The CIs by the Union method [31] with $\bar{s} = p_z - 1$ (assuming there are two valid IVs) achieve the desired coverage level while those with $\bar{s} = \lceil p_z/2 \rceil$ (assuming the majority rule) do not achieve the desired coverage level for settings **S2** to **S5**.

Empirical Coverage of Confidence Intervals for $\tau = 0.2$

		Oracle				Searching		Sampling		Union		
Set	n	TSLS	BA	TSHT	CIIV	$\hat{\mathcal{V}}^{\text{TSHT}}$	$\hat{\mathcal{V}}^{\text{CIIV}}$	$\hat{\mathcal{V}}^{\text{TSHT}}$	$\hat{\mathcal{V}}^{\text{CIIV}}$	Check	$p_z - 1$	$\lceil p_z/2 \rceil$
S1	500	0.95	0.97	0.53	0.61	1.00	1.00	1.00	1.00	1.00	1.00	0.99
	1000	0.94	0.97	0.45	0.69	1.00	1.00	1.00	1.00	1.00	1.00	0.98
	2000	0.96	0.95	0.63	0.79	1.00	1.00	1.00	1.00	1.00	1.00	0.96
	5000	0.94	0.95	0.89	0.92	1.00	1.00	1.00	1.00	1.00	1.00	0.96
S2	500	0.95	0.96	0.56	0.51	1.00	1.00	1.00	0.99	1.00	1.00	0.27
	1000	0.94	0.95	0.45	0.58	0.99	0.99	1.00	0.97	1.00	1.00	0.03
	2000	0.94	0.93	0.51	0.74	0.98	0.98	0.98	0.97	1.00	1.00	0.00
	5000	0.96	0.93	0.85	0.95	0.99	1.00	1.00	1.00	1.00	1.00	0.00
S3	500	0.95	0.95	0.63	0.63	0.99	0.99	0.99	0.99	1.00	1.00	0.62
	1000	0.94	0.97	0.62	0.60	0.99	0.99	0.99	0.97	1.00	1.00	0.18
	2000	0.94	0.93	0.62	0.73	0.97	0.98	0.97	0.96	1.00	1.00	0.01
	5000	0.96	0.93	0.85	0.95	0.99	1.00	0.99	1.00	1.00	1.00	0.00
S4	500	0.95	0.96	0.72	0.66	0.94	0.95	0.94	0.93	0.98	0.98	0.00
	1000	0.94	0.95	0.65	0.58	1.00	0.97	0.99	0.96	0.95	0.98	0.00
	2000	0.93	0.99	0.68	0.58	0.98	0.97	0.97	0.93	0.99	0.95	0.00
	5000	0.95	0.95	0.91	0.88	0.98	0.95	0.98	0.95	1.00	0.94	0.00
S5	500	0.95	0.97	0.49	0.51	0.81	0.89	0.88	0.86	0.98	0.97	0.14
	1000	0.94	0.96	0.31	0.50	0.68	0.89	0.76	0.86	0.91	0.98	0.00
	2000	0.93	0.98	0.46	0.57	0.86	0.96	0.86	0.92	0.84	0.95	0.00
	5000	0.95	0.95	0.90	0.88	0.98	0.95	0.97	0.94	0.98	0.94	0.00

Average Lengths of Confidence Intervals for $\tau = 0.2$

		Oracle				Searching		Sampling		Union		
Set	n	TSLS	BA	TSHT	CIIV	$\hat{\mathcal{V}}^{\text{TSHT}}$	$\hat{\mathcal{V}}^{\text{CIIV}}$	$\hat{\mathcal{V}}^{\text{TSHT}}$	$\hat{\mathcal{V}}^{\text{CIIV}}$	Check	$p_z - 1$	$\lceil p_z/2 \rceil$
S1	500	0.10	0.16	0.08	0.08	0.59	0.61	0.34	0.34	-	1.16	0.25
	1000	0.07	0.14	0.06	0.06	0.39	0.43	0.24	0.24	-	0.63	0.16
	2000	0.05	0.10	0.05	0.05	0.27	0.30	0.17	0.17	-	0.42	0.09
	5000	0.03	0.04	0.03	0.03	0.17	0.18	0.10	0.10	-	0.27	0.04
S2	500	0.13	0.24	0.13	0.10	0.58	0.59	0.37	0.36	-	2.46	0.07
	1000	0.09	0.24	0.13	0.08	0.37	0.41	0.26	0.26	-	1.45	0.02
	2000	0.06	0.26	0.14	0.06	0.25	0.29	0.19	0.18	-	0.76	0.00
	5000	0.04	0.13	0.08	0.04	0.16	0.17	0.10	0.10	-	0.28	0.00
S3	500	0.13	0.22	0.10	0.10	0.62	0.62	0.45	0.38	-	1.77	0.13
	1000	0.09	0.21	0.10	0.08	0.38	0.41	0.29	0.27	-	1.36	0.03
	2000	0.06	0.25	0.13	0.06	0.26	0.29	0.19	0.18	-	0.86	0.00
	5000	0.04	0.13	0.08	0.04	0.16	0.17	0.10	0.10	-	0.35	0.00
S4	500	0.23	0.62	0.24	0.18	0.56	0.56	0.48	0.44	-	0.87	0.00
	1000	0.16	0.56	0.17	0.13	0.44	0.36	0.38	0.29	-	0.42	0.00
	2000	0.11	0.32	0.14	0.10	0.27	0.24	0.22	0.18	-	0.20	0.00
	5000	0.07	0.13	0.08	0.07	0.14	0.13	0.11	0.10	-	0.09	0.00
S5	500	0.23	0.63	0.27	0.17	0.42	0.51	0.41	0.42	-	1.01	0.05
	1000	0.16	0.61	0.18	0.13	0.32	0.36	0.30	0.29	-	0.50	0.00
	2000	0.11	0.38	0.12	0.10	0.28	0.24	0.25	0.18	-	0.22	0.00
	5000	0.07	0.15	0.08	0.07	0.15	0.13	0.12	0.10	-	0.09	0.00

Table 1: Empirical coverage and average lengths for **S1** to **S5** with $\tau = 0.2$. The columns indexed with TSLS, BA TSHT and CIIV represent the oracle TSLS CI with the knowledge of \mathcal{V} , the oracle bias-aware CI in (10), the CI by [25], and the CI by [53], respectively. Under the columns indexed with “Searching” (or “Sampling”), the columns indexed with $\hat{\mathcal{V}}^{\text{TSHT}}$ and $\hat{\mathcal{V}}^{\text{CIIV}}$ represent our proposed searching (or sampling) CI in Algorithm 3 with $\hat{\mathcal{V}}^{\text{TSHT}}$ and $\hat{\mathcal{V}}^{\text{CIIV}}$, respectively. The column indexed with “Check” reports the proportion of simulations passing the plurality rule check in Algorithm 3. The columns indexed with **Union** represent the method by [31]. The columns indexed with $p_z - 1$ and $\lceil p_z/2 \rceil$ correspond to the **Union** methods assuming two valid IVs and the majority rule, respectively.

We now compare the CI lengths. When the CIs by TSHT and CIIV are valid, their lengths are similar to the length of the CI by oracle TSLS, which match with the theory in [25, 53]. For the CIs achieving valid coverage, our proposed sampling CI is in general shorter than the searching CI and the **Union** CI with $p_z - 1$. We shall point out that our proposed sampling CI can be even shorter than the oracle bias-aware CI (the benchmark), which requires the oracle knowledge of the bias size. As an important remark, the oracle bias-aware CI, the sampling CI, the searching CI, and the **Union** CI are in general longer than the CI by the `oracle` TSLS, which is a price to pay for constructing uniformly valid CIs.

In Section E.1 in the supplement, we report the results for $\tau = 0.4$ and compare our proposed methods with existing methods for settings with heteroscedastic errors. In Section E.2 in the supplement, we further explore the settings considered in [53]. The results for these settings are similar to those in Table 1.

7.1 Varying violation strength

We compare our proposed methods with existing methods over different IV invalidity levels. Particularly, we focus on **S2** and vary $\tau \in \{0.025, 0.05, 0.075, 0.1, 0.2, 0.3, 0.4, 0.5\}$. We consider the heteroscedastic errors: for $1 \leq i \leq n$, generate $\delta_i \sim N(0, 1)$ and $e_i = 0.3\delta_i + \sqrt{[1 - 0.3^2]/[0.86^4 + 1.38^2]}(1.38 \cdot \tau_{1,i} + 0.86^2 \cdot \tau_{2,i})$, where conditioning on Z_i , $\tau_{1,i} \sim N(0, [0.5 \cdot Z_{i,1}^2 + 0.25]^2)$, $\tau_{2,i} \sim N(0, 1)$, and $\tau_{1,i}$ and $\tau_{2,i}$ are independent of δ_i . The above error generating method follows from [4].

As reported in Figure 6, our proposed searching and sampling CIs achieve the desired coverage (95%) while TSHT and CIIV only achieve the desired coverage for $\tau = 0.5$ with $n = 2000$ and $\tau \geq 0.3$ with $n = 5000$. In terms of length, we observe that the sampling CI is shorter than the searching CI. The oracle bias-aware CI is used as a benchmark, which requires the oracle bias level of the TSHT method. The sampling CI can be even shorter than this oracle benchmark. We do not plot the length of the **Union** CI, which is three to six times longer than our proposed sampling CI; see Figure 3 for details.

An interesting observation is that the empirical coverage of TSHT and CIIV is around 90% for $\tau = 0.025$. This happens since the invalidity levels of the IVs are small, and even the inclusion of such invalid IVs does not significantly worsen the empirical coverage. We present the results for homoscedastic errors in Figure E.1 in the supplement.

7.2 Tuning parameter selection

We now explore the effect of the tuning parameters used in Algorithm 3. For the searching CI, we observe in Table 2 that the empirical coverage and the average lengths of the searching CIs are almost invariant to different choices of L , U and a . For two intervals (a_1, b_1) and (a_2, b_2) , we define its difference as $|a_1 - a_2| + |b_1 - b_2|$. In Table 2, we report the difference between the searching CI constructed with the default choice of L, U, a in Algorithm 3 and searching CIs with other choices of L, U, a . As reported in Table 2, the average interval difference is smaller than twice the default grid size $n^{-0.6}$.




Figure 6: Empirical coverage and average lengths for the setting **S2** with $\tau \in \{0.025, 0.05, 0.075, 0.1, 0.2, 0.3, 0.4, 0.5\}$ and heteroscedastic errors. **Oracle-BA**, **TSHT**, and **CIIV** represent the oracle bias-aware CI in (10), the CI by [25], and the CI by [53], respectively. The searching and sampling CIs are implemented as in Algorithm 3.

In Section A.2 in the supplement, we demonstrate that the sampling CIs have nearly the same empirical coverage and length for different choices of L, U, a and the resampling size M . The choice of the shrinkage parameter $\lambda > 0$ has a more obvious effect on the sampling CI. As explained in Remark 3, we choose the smallest λ such that more than a pre-specified proportion (denoted as **prop**) of the $M = 1000$ intervals are non-empty. In Figure 7, we compare the coverage and length properties of the sampling CIs by varying **prop** across $\{1\%, 5\%, 10\%, 20\%, 30\%\}$. If **prop** $\geq 5\%$, the empirical coverage reaches the nominal level and vary slightly with **prop**. In terms of length, the intervals get slightly longer with a larger value of **prop**. The empirical coverage and average lengths of the sampling CIs are robust to a wide range of λ values, as long as the corresponding λ guarantees a sufficient proportion of non-empty sampled searching CIs.

		S1 ($n = 2000$)			S2 ($n = 2000$)			S3 ($n = 2000$)			S4 ($n = 2000$)		
[L,U]	a	Cov	Len	Diff	Cov	Len	Diff	Cov	Len	Diff	Cov	Len	Diff
As in (19)	0.6	1	0.287	0.000	1	0.268	0.000	0.994	0.275	0.000	0.980	0.274	0.000
	0.8	1	0.295	0.008	1	0.276	0.008	0.996	0.283	0.008	0.982	0.282	0.008
	1.0	1	0.297	0.010	1	0.278	0.010	0.996	0.284	0.010	0.984	0.284	0.010
[-10,10]	0.6	1	0.287	0.007	1	0.268	0.008	0.994	0.275	0.008	0.978	0.274	0.007
	0.8	1	0.295	0.008	1	0.277	0.009	0.996	0.283	0.009	0.984	0.282	0.009
	1.0	1	0.297	0.010	1	0.278	0.010	0.996	0.285	0.010	0.984	0.284	0.010
[-20,20]	0.6	1	0.287	0.007	1	0.268	0.007	0.992	0.275	0.007	0.982	0.274	0.007
	0.8	1	0.295	0.008	1	0.277	0.009	0.996	0.283	0.009	0.982	0.282	0.009
	1.0	1	0.297	0.010	1	0.278	0.010	0.996	0.285	0.010	0.984	0.284	0.010

Table 2: Comparison of searching CIs with different choices of L, U, a for settings **S1** to **S4** with $\tau = 0.4$ and $n = 2000$. The columns indexed with “Cov” and “Len” denote the empirical coverage and the average lengths, respectively. The column indexed with “Diff” represents the average difference between the searching CI in Algorithm 3 and the searching CIs with other choices of L, U , and n^{-a} .




Figure 7: Comparison of the sampling CIs with different λ for $\tau = 0.4$. The resampling size M is set as 1000. We choose the smallest value of λ such that $1000 \cdot \text{prop}$ sampled intervals are non-empty. For example, for $\text{prop} = 10\%$, we choose the smallest λ leading to 100 non-empty sampled intervals.

7.3 High dimensional IVs and covariates

In this subsection, we extend our proposed searching and sampling method to the settings with high-dimensional IVs and covariates. We modify Algorithm 3 to accommodate for the high-dimensional setting. Particularly, for step 1 of Algorithm 3, we construct $\hat{\Gamma} \in \mathbb{R}^{p_z}, \hat{\gamma} \in \mathbb{R}^{p_z}$ by applying the debiased lasso estimators [30, 49, 56]. The detailed discussions about the reduced-form estimators and its covariance estimators can be found in Section 4.1 of [24].

For step 2 of Algorithm 3, we modify the definition of $\widehat{\mathcal{S}}$ in (11) by replacing $\sqrt{\log n}$ with $\sqrt{2.01 \cdot \log \max\{p_z, n\}}$; for step 3 of Algorithm 3, we modify the construction of $\widehat{\Pi}$ in (33) by replacing $\sqrt{\log n}$ with $2.01 \cdot \sqrt{\log \max\{p_z, n\}}$; see Remark 1 and [25] for more discussions.

We follow the same high-dimensional setting as in Section 5.3 of [25]. Particularly, we set $p_x = 150, p_z = 200$ and generate the sparse vector $\gamma^* = 0.5 \cdot (\mathbf{1}_7^\top, \mathbf{0}_{p_z-7}^\top)^\top$, $\pi^* = (\mathbf{0}_5^\top, \tau/2, \tau/2, \mathbf{0}_{p_z-7}^\top)^\top$. That is, $\mathcal{S} = \{1, 2, \dots, 7\}$, $\mathcal{V} = \{1, 2, \dots, 5\}$, and the 6-th and 7-th IVs are invalid. As reported in Table E.6, the CIs by TSHT can only achieve the desired coverage for a sufficiently large sample size n and invalidity level τ . In contrast, our proposed searching and sampling CIs achieve the desired coverage across all settings. The sampling CIs are helpful in reducing the length of the searching CIs. The sampling CIs are in general longer than the oracle bias-aware CIs (the benchmark), but the sampling CIs have a comparable length to the oracle bias-aware CIs. More simulation results are presented in Section E.3 in the supplement.




Figure 8: Empirical coverage and average lengths for high-dimensional setting with $p_z = 200$ and $p_x = 150$. We vary the violation strength τ across $\{0.5, 1\}$. **Oracle-BA** and **TSHT** represent the oracle bias-aware CI in (10) and the CI by [25], respectively. The searching and sampling CIs are implemented as in Algorithm 3.

8 Real Data Analysis

We study the effect of education on earnings by analyzing the 2018 survey of China Family Panel Studies [55] available at <https://www.issf.pku.edu.cn/cfps/en/data/public/index.htm>. The outcome is the logarithm transformation of salary in the past 12 months, and the treatment

is the years of education. We include three binary baseline covariates: gender, urban (whether the subject lives in the urban area), and hukou (whether the subject’s hukou is agricultural or non-agricultural). Following the literature [3, 7, 8, 48], we have induced the following nine IVs:

1. Family background variables [3, 8, 48, e.g.]: the father’s education level, the mother’s education level, the spouse’s education level, the family size, and the log transformation of the education expenditure in the past 12 months.
2. The group-level years of education [7], where the groups are formulated by age, gender, and region,
3. Personal traits [3]: the statement on the fair competition, the statement on the talent pay-off, and whether the subject reads some books or not in the past 12 months.

The statement on fair competition measures an individual’s willingness to compete through ability. The statement on the talent pay-off is about an individual’s viewpoint on whether their educational endeavors will pay off. After removing the missing values, the data consists of 3758 observations. In the supplement, we report the summary statistics of all IVs and baseline covariates in Table E.8.

In Table 3, we compare our proposed searching and sampling CIs with existing methods. By applying TSHT, we identified six relevant instruments: the father’s education level, the mother’s education level, the spouse’s education level, the group-level years of education, the family size, whether to read some books or not in the past 12 months. Out of these IVs, the family size is detected as the invalid IV. CIIV outputs the same set of valid IVs. When we include all 9 IVs, the concentration parameter is 2906.36. If we only include the 5 IVs selected by TSHT and CIIV, the concentration parameter is 2850.57. This indicates that the whole set of IVs are strongly associated with the treatment, but some IVs (e.g., the family size) are possibly invalid.

Method	CI	Method	CI
OLS	(0.0305, 0.0503)	Searching CI	(0.0409, 0.1698)
TOLS	(0.0959, 0.1190)	Sampling CI	(0.0552, 0.1268)
TSHT	(0.0946, 0.1178)	Union ($\bar{s} = p_z - 1$)	(-0.4915, 1.6043)
CIIV	(0.0948, 0.1175)	Union($\bar{s} = \lceil p_z/2 \rceil$)	(0.0409, 0.1342)

Table 3: Confidence intervals for the effect of education on earnings.

As reported in Table 3, CIs by TSHT and CIIV are relatively short but they may undercover due to the IV selection error. We compare the lengths of **Union** method and our proposed searching and sampling CIs, which are all robust to the IV selection error. The sampling CI is the shortest among these CIs. The validity of the **Union** CI with $\bar{s} = \lceil p_z/2 \rceil$ requires half of the candidate IVs to be valid; if we can only assume that two of candidate IVs are valid, then the CI by **Union** with $\bar{s} = \lceil p_z/2 \rceil$ may not be valid but the CI by **Union** with $\bar{s} = p_z - 1$ is valid.

We plot the searching CI (in blue) and the sampling CI (in red) in Figure 9. Out of the 1000 sampled intervals, 103 of them are non-empty and the union of these 103 intervals is shorter than the searching CI.




Figure 9: The axis corresponds to sampling indexes $\{1, 2, \dots, 103\}$ (after re-ordering) and the y-axis reports the sampled CIs. Along the y-axis, the red interval is $CI^{\text{samp}} = (0.0552, 0.1268)$ and the blue interval is $\widehat{CI}^{\text{sear}} = (0.0409, 0.1698)$.

9 Conclusion and Discussion

Causal inference from observational studies is a challenging task. Typically, stringent identification conditions are required to facilitate various causal inference approaches. The valid IV assumption is one of such assumptions to handle unmeasured confounders. In the current paper, we devise uniformly valid confidence intervals for the causal effect when the candidate IVs are possibly invalid. Our proposed searching and sampling confidence intervals add to the fast-growing literature on robust inference with possibly invalid IVs. The proposed method has the advantage of being more robust to the mistakes in separating the valid and invalid IVs at the expense of a wider confidence interval. The proposed intervals are computationally efficient and less conservative than existing uniformly valid confidence intervals.

Acknowledgement

The research of Z. Guo was partly supported by the NSF grants DMS 1811857 and 2015373 and NIH grants R01GM140463 and R01LM013614. Z. Guo is grateful to the participants at Penn Causal Reading Group and CUHK econometrics seminar for their helpful discussion, to Dr. Frank Windmeijer for pointing out the connection in Section 5.2, and to Dr. Hyunseung Kang for sharing the code for replicating the `Union` method. Z. Guo thanks the editors, two anonymous referees, and Drs. Hyunseung Kang, Zhonghua Liu, and Molei Liu for constructive comments on a previous draft. Z. Guo thanks Mr. Zhenyu Wang for the help with the numerical implementation and Dr. Junhui Yang and Mr. Zhenyu Wang for cleaning the CFHS data set.

References

- [1] Theodore W Anderson and Herman Rubin. Estimation of the parameters of a single equation in a complete system of stochastic equations. *Annals of Mathematical statistics*,

20(1):46–63, 1949.

- [2] Timothy B Armstrong, Michal Kolesár, and Soonwoo Kwon. Bias-aware inference in regularized regression models. *arXiv preprint arXiv:2012.14823*, 2020.
- [3] Jere R Behrman, Olivia S Mitchell, Cindy K Soo, and David Bravo. How financial literacy affects household wealth accumulation. *American Economic Review*, 102(3):300–304, 2012.
- [4] Paul A Bekker and Federico Crudu. Jackknife instrumental variable estimation with heteroskedasticity. *Journal of econometrics*, 185(2):332–342, 2015.
- [5] Richard Berk, Lawrence Brown, Andreas Buja, Kai Zhang, and Linda Zhao. Valid post-selection inference. *The Annals of Statistics*, 41(2):802–837, 2013.
- [6] Daniel Berkowitz, Mehmet Caner, and Ying Fang. The validity of instruments revisited. *Journal of Econometrics*, 166(2):255–266, 2012.
- [7] Sajjad Haider Bhatti, Jean Bourdon, and Muhammad Aslam. Economic returns to education in france: Ols and instrumental variable estimations. *The Lahore Journal of Economics*, 18(2):51, 2013.
- [8] McKinley L Blackburn and David Neumark. Omitted-ability bias and the increase in the return to schooling. *Journal of labor economics*, 11(3):521–544, 1993.
- [9] Jack Bowden, George Davey Smith, and Stephen Burgess. Mendelian randomization with invalid instruments: effect estimation and bias detection through egger regression. *International journal of epidemiology*, 44(2):512–525, 2015.
- [10] Jack Bowden, George Davey Smith, and Stephen Burgess. Mendelian randomization with invalid instruments: effect estimation and bias detection through egger regression. *International journal of epidemiology*, 44(2):512–525, 2015.
- [11] Jack Bowden, George Davey Smith, Philip C Haycock, and Stephen Burgess. Consistent estimation in mendelian randomization with some invalid instruments using a weighted median estimator. *Genetic epidemiology*, 40(4):304–314, 2016.
- [12] Stephen Burgess, Dylan S Small, and Simon G Thompson. A review of instrumental variable estimators for mendelian randomization. *Statistical methods in medical research*, 26(5):2333–2355, 2017.
- [13] Mehmet Caner, Xu Han, and Yoonseok Lee. Adaptive elastic net gmm estimation with many invalid moment conditions: Simultaneous model and moment selection. *Journal of Business & Economic Statistics*, 36(1):24–46, 2018.
- [14] Xu Cheng and Zhipeng Liao. Select the valid and relevant moments: An information-based lasso for gmm with many moments. *Journal of Econometrics*, 186(2):443–464, 2015.

- [15] Victor Chernozhukov, Christian Hansen, and Martin Spindler. Post-selection and post-regularization inference in linear models with many controls and instruments. 2015.
- [16] George Davey Smith and Shah Ebrahim. Mendelian randomization: can genetic epidemiology contribute to understanding environmental determinants of disease? *International journal of epidemiology*, 32(1):1–22, 2003.
- [17] George Davey Smith and Gibran Hemani. Mendelian randomization: genetic anchors for causal inference in epidemiological studies. *Human molecular genetics*, 23(R1):R89–R98, 2014.
- [18] David L Donoho and Iain M Johnstone. Minimax risk over p -balls for p -error. *Probability Theory and Related Fields*, 99(2):277–303, 1994.
- [19] David L Donoho and Jain M Johnstone. Ideal spatial adaptation by wavelet shrinkage. *biometrika*, 81(3):425–455, 1994.
- [20] F Eicker. Limit theorems for regression with unequal and dependent errors. In *Proc. Fifth Berkeley Symp. Statist. Probab.*, pages 59–82, 1967.
- [21] Qingliang Fan and Yaqian Wu. Endogenous treatment effect estimation with some invalid and irrelevant instruments. *arXiv preprint arXiv:2006.14998*, 2020.
- [22] Patrik Guggenberger. On the asymptotic size distortion of tests when instruments locally violate the exogeneity assumption. *Econometric Theory*, 28(2):387–421, 2012.
- [23] Zijian Guo. Inference for high-dimensional maximin effects in heterogeneous regression models using a sampling approach. *arXiv preprint arXiv:2011.07568*, 2020.
- [24] Zijian Guo, Hyunseung Kang, T Tony Cai, and Dylan S Small. Testing endogeneity with high dimensional covariates. *Journal of Econometrics*, 207(1):175–187, 2018.
- [25] Zijian Guo, Hyunseung Kang, T Tony Cai, and Dylan S Small. Confidence intervals for causal effects with invalid instruments by using two-stage hard thresholding with voting. *Journal of the Royal Statistical Society: Series B (Statistical Methodology)*, 80(4):793–815, 2018.
- [26] Jinyong Hahn and Jerry Hausman. Estimation with valid and invalid instruments. *Annales d’Economie et de Statistique*, pages 25–57, 2005.
- [27] Lars Peter Hansen. Large sample properties of generalized method of moments estimators. *Econometrica: Journal of the Econometric Society*, pages 1029–1054, 1982.
- [28] Fernando Pires Hartwig, George Davey Smith, and Jack Bowden. Robust inference in summary data mendelian randomization via the zero modal pleiotropy assumption. *International journal of epidemiology*, 46(6):1985–1998, 2017.

- [29] Peter J Huber. Under nonstandard conditions. In *Proceedings of the Fifth Berkeley Symposium on Mathematical Statistics and Probability: Weather Modification; University of California Press: Berkeley, CA, USA*, page 221, 1967.
- [30] Adel Javanmard and Andrea Montanari. Confidence intervals and hypothesis testing for high-dimensional regression. *The Journal of Machine Learning Research*, 15(1):2869–2909, 2014.
- [31] Hyunseung Kang, Youjin Lee, T Tony Cai, and Dylan S Small. Two robust tools for inference about causal effects with invalid instruments. *Biometrics*, 2020.
- [32] Hyunseung Kang, Anru Zhang, T Tony Cai, and Dylan S Small. Instrumental variables estimation with some invalid instruments and its application to mendelian randomization. *Journal of the American Statistical Association*, 111(513):132–144, 2016.
- [33] Michal Kolesár, Raj Chetty, John Friedman, Edward Glaeser, and Guido W Imbens. Identification and inference with many invalid instruments. *Journal of Business & Economic Statistics*, 33(4):474–484, 2015.
- [34] Jason D Lee, Dennis L Sun, Yuekai Sun, and Jonathan E Taylor. Exact post-selection inference, with application to the lasso. *Annals of Statistics*, 44(3):907–927, 2016.
- [35] Hannes Leeb and Benedikt M Pötscher. Model selection and inference: Facts and fiction. *Econometric Theory*, pages 21–59, 2005.
- [36] Arthur Lewbel. Using heteroscedasticity to identify and estimate mismeasured and endogenous regressor models. *Journal of Business & Economic Statistics*, 30(1):67–80, 2012.
- [37] Arthur Lewbel. Identification and estimation using heteroscedasticity without instruments: The binary endogenous regressor case. *Economics Letters*, 165:10–12, 2018.
- [38] Sai Li and Zijian Guo. Causal inference for nonlinear outcome models with possibly invalid instrumental variables. *arXiv preprint arXiv:2010.09922*, 2020.
- [39] Zhipeng Liao. Adaptive gmm shrinkage estimation with consistent moment selection. *Econometric Theory*, 29(5):857–904, 2013.
- [40] Zhonghua Liu, Ting Ye, Baolu Sun, Mary Schooling, and Eric Tchetgen Tchetgen. On mendelian randomization mixed-scale treatment effect robust identification (mr misteri) and estimation for causal inference. *arXiv preprint arXiv:2009.14484*, 2020.
- [41] Anna Mikusheva. Survey on statistical inferences in weakly-identified instrumental variable models. *Applied Econometrics*, 29(1):117–131, 2013.
- [42] John D Sargan. The estimation of economic relationships using instrumental variables. *Econometrica: Journal of the Econometric Society*, pages 393–415, 1958.

- [43] Dylan S Small. Sensitivity analysis for instrumental variables regression with overidentifying restrictions. *Journal of the American Statistical Association*, 102(479):1049–1058, 2007.
- [44] Wes Spiller, David Slichter, Jack Bowden, and George Davey Smith. Detecting and correcting for bias in mendelian randomization analyses using gene-by-environment interactions. *International journal of epidemiology*, 48(3):702–712, 2019.
- [45] Douglas Staiger and James H Stock. Instrumental variables regression with weak instruments. *Econometrica: Journal of the Econometric Society*, pages 557–586, 1997.
- [46] James H Stock, Jonathan H Wright, and Motohiro Yogo. A survey of weak instruments and weak identification in generalized method of moments. *Journal of Business & Economic Statistics*, 20(4):518–529, 2002.
- [47] Eric J Tchetgen Tchetgen, BaoLuo Sun, and Stefan Walter. The genius approach to robust mendelian randomization inference. *arXiv preprint arXiv:1709.07779*, 2017.
- [48] Philip Trostel, Ian Walker, and Paul Woolley. Estimates of the economic return to schooling for 28 countries. *Labour economics*, 9(1):1–16, 2002.
- [49] Sara van de Geer, Peter Bühlmann, Yaacov Ritov, and Ruben Dezeure. On asymptotically optimal confidence regions and tests for high-dimensional models. *The Annals of Statistics*, 42(3):1166–1202, 2014.
- [50] Peng Wang and Minge Xie. Repro sampling method for statistical inference of high dimensional linear models. *Research Manuscript*, 2020.
- [51] Halbert White. A heteroskedasticity-consistent covariance matrix estimator and a direct test for heteroskedasticity. *Econometrica: journal of the Econometric Society*, pages 817–838, 1980.
- [52] Frank Windmeijer, Helmut Farbmacher, Neil Davies, and George Davey Smith. On the use of the lasso for instrumental variables estimation with some invalid instruments. *Journal of the American Statistical Association*, 114(527):1339–1350, 2019.
- [53] Frank Windmeijer, Xiaoran Liang, Fernando P Hartwig, and Jack Bowden. The confidence interval method for selecting valid instrumental variables. *Journal of the Royal Statistical Society: Series B (Statistical Methodology)*, 83(4):752–776, 2021.
- [54] Jeffrey M Wooldridge. *Econometric analysis of cross section and panel data*. MIT press, 2010.
- [55] Yu Xie. China family panel studies (2010) user’s manual. *Beijing: Institute of Social Science Survey, Peking University*, 2012.

- [56] Cun-Hui Zhang and Stephanie S Zhang. Confidence intervals for low dimensional parameters in high dimensional linear models. *Journal of the Royal Statistical Society: Series B (Statistical Methodology)*, 76(1):217–242, 2014.
- [57] Qingyuan Zhao, Jingshu Wang, Gibran Hemani, Jack Bowden, and Dylan S Small. Statistical inference in two-sample summary-data mendelian randomization using robust adjusted profile score. *The Annals of Statistics*, 48(3):1742–1769, 2020.

A Extra Discussions on Tuning Parameter Selection

We demonstrate in the following that our proposed searching and sampling methods are invariant to different choices of tuning parameters, in terms of both empirical coverage and average lengths.

A.1 First-stage selection

In selecting the valid instruments, the first stage is to construct an initial set $\widehat{\mathcal{S}}$ of relevant IVs. Such a first-stage selection is required for TSHT [25], CIIV [53], and our current proposal. We shall first show that the existing TSHT and CIIV can still achieve the desired coverage level even in the presence of individually weak IVs. Particularly, we consider the following simulation setting,

$$\mathbf{S0} : \text{set } \gamma^* = (\gamma_1 \cdot \mathbf{1}_2, 0.5 \cdot \mathbf{1}_4, \gamma_1, 0.5, 0.5, 0.5, 0.5) \text{ and } \pi^* = (\mathbf{0}_2, \mathbf{0}_4, \tau/2, \tau/2, -0.5, -1)^\top.$$

We first set $\tau = 0.5$ such that the invalid IVs have a large invalidity level. The first, second, and seventh IVs are generated as individually weak IVs, where the individual IV strength is controlled by γ_1 . We vary γ_1 across $\{0.05, 0.075, 0.10\}$.

As reported in Table A.1, the three individually weak IVs are only included into $\widehat{\mathcal{S}}$ for a proportion of the 500 simulations. However, even if there are uncertainties of including these relatively weak IVs, the existing CIs by TSHT and CIIV in general achieve the desired coverage.

By comparing the results in Table A.1 and Figure 3 in the main paper, we reach the conclusion that the challenge of locally invalid IVs is fundamentally different from that of the weak IVs. Even if there are individually weak IVs, the existing CIs by TSHT and CIIV can still achieve the desired coverage level as long as there are enough strongly associated IVs. In contrast, as demonstrated in Figure 3 in the main paper, the existence of locally invalid IVs poses great challenges for inference with TSHT and CIIV.

We shall construct $\widehat{\mathcal{S}}$ with different threshold levels and compare the performance of our proposed searching and sampling CIs with the possibly different initial sets $\widehat{\mathcal{S}}$. We demonstrate the performance using the setting **S0** since the change of the thresholds in the construction of $\widehat{\mathcal{S}}$ does not affect the performance over the settings **S1** to **S5**. We vary the invalidity level τ across $\{0.2, 0.4, 0.5\}$ and the IV strength γ_1 across $\{0.05, 0.075, 0.1\}$. Particularly, we consider the threshold $\sqrt{\log n}$ in construction of $\widehat{\mathcal{S}}$ as in (11). In addition, we follow [25] and replace $\sqrt{\log n}$ with $\sqrt{2.01 \log \max\{n, p_z\}}$ in constructing the $\widehat{\mathcal{S}}$. As reported in Table A.2, the empirical coverage and the average lengths vary slightly with different first-stage threshold levels.

A.2 Searching and sampling: dependence on L, U, a, M and λ

We further explore the effect of tuning parameters on the construction of the sampling CI. In Section 7.2 in the main paper, we have shown that the searching CI is almost invariant to the choices of the initial range $[L, U]$ and the grid size n^{-a} . In the following Table A.3, we show that

		TSHT		CIIV		Prop of Belonging to $\widehat{\mathcal{S}}$		
γ_1	n	Cov	Len	Cov	Len	First IV	Second IV	Seventh IV
0.050	500	0.902	0.156	0.912	0.136	0.144	0.088	0.116
	1000	0.928	0.105	0.932	0.096	0.188	0.180	0.174
	2000	0.932	0.079	0.934	0.067	0.404	0.310	0.352
0.075	500	0.908	0.160	0.900	0.134	0.226	0.190	0.190
	1000	0.934	0.107	0.930	0.094	0.454	0.398	0.364
	2000	0.936	0.077	0.936	0.065	0.794	0.672	0.698
0.100	500	0.902	0.156	0.898	0.132	0.396	0.334	0.340
	1000	0.930	0.113	0.928	0.091	0.710	0.624	0.594
	2000	0.936	0.081	0.940	0.062	0.944	0.902	0.916

Table A.1: Empirical coverage (Cov) and average lengths (Len) of CIs for the setting **S0**. The columns indexed with TSHT and CIIV represent the confidence intervals by [25] and [53], respectively. The columns indexed with “first IV”, “second IV”, and “seventh IV” report the proportion of the first IV, the second IV, and the seventh IV being included into the set $\widehat{\mathcal{S}}$.

the sampling CI only changes slightly with different choices of $[L, U]$, n^{-a} and the resampling size M , in terms of the empirical coverage and the average lengths.

			$\sqrt{\log n}$				$\sqrt{2.01 \log \max\{n, p_z\}}$			
			Searching		Sampling		Searching		Sampling	
τ	γ_1	n	Cov	Len	Cov	Len	Cov	Len	Cov	Len
0.2	0.050	500	0.998	0.640	0.996	0.418	0.998	0.637	0.996	0.417
		1000	1.000	0.442	0.996	0.292	1.000	0.441	0.996	0.291
		2000	1.000	0.309	0.996	0.203	1.000	0.306	0.998	0.199
	0.075	500	1.000	0.648	0.998	0.424	0.998	0.636	0.996	0.418
		1000	1.000	0.446	1.000	0.295	1.000	0.441	0.996	0.291
		2000	1.000	0.320	0.998	0.216	1.000	0.307	0.998	0.201
	0.100	500	1.000	0.660	0.996	0.426	0.998	0.636	0.994	0.420
		1000	1.000	0.458	1.000	0.303	1.000	0.441	1.000	0.292
		2000	1.000	0.337	1.000	0.229	1.000	0.316	1.000	0.210
0.4	0.050	500	0.998	0.610	0.992	0.397	0.998	0.603	0.992	0.390
		1000	1.000	0.409	0.998	0.255	1.000	0.407	1.000	0.252
		2000	1.000	0.284	1.000	0.174	1.000	0.283	1.000	0.170
	0.075	500	1.000	0.617	0.998	0.400	0.998	0.604	0.994	0.392
		1000	1.000	0.417	0.998	0.259	1.000	0.408	1.000	0.251
		2000	1.000	0.295	1.000	0.185	1.000	0.283	1.000	0.174
	0.100	500	1.000	0.628	0.996	0.404	0.998	0.606	0.994	0.391
		1000	0.998	0.434	0.998	0.273	1.000	0.410	1.000	0.254
		2000	1.000	0.311	1.000	0.201	1.000	0.294	1.000	0.187
0.5	0.050	500	0.996	0.596	0.996	0.378	0.996	0.590	0.996	0.372
		1000	1.000	0.401	0.998	0.245	1.000	0.400	1.000	0.242
		2000	1.000	0.284	1.000	0.175	1.000	0.283	1.000	0.172
	0.075	500	0.998	0.603	0.996	0.382	0.996	0.591	0.994	0.374
		1000	1.000	0.407	0.996	0.252	1.000	0.402	1.000	0.246
		2000	1.000	0.295	1.000	0.188	1.000	0.284	1.000	0.175
	0.100	500	0.998	0.616	0.996	0.388	0.996	0.593	0.992	0.374
		1000	0.998	0.424	0.998	0.267	1.000	0.403	1.000	0.249
		2000	1.000	0.318	1.000	0.206	1.000	0.302	1.000	0.192

Table A.2: Empirical coverage (Cov) and average lengths (Len) of searching and sampling CIs for the setting **S0**. The columns indexed with “Searching” and “Sampling” represent our proposed searching and sampling CIs in Algorithm 3. The columns indexed with $\sqrt{\log n}$ and $\sqrt{2.01 \log \max\{n, p_z\}}$ stands for the construction of \hat{S} in (11) and the construction of \hat{S} in (11) with replacing $\sqrt{\log n}$ by $\sqrt{2.01 \log \max\{n, p_z\}}$, respectively.

			$M = 500$		$M = 1000$		$M = 1500$		$M = 2000$	
set	n	LUa	Cov	Len	Cov	Len	Cov	Len	Cov	Len
S1	500	1	1.000	0.301	1.000	0.333	1.000	0.349	1.000	0.356
		2	1.000	0.309	1.000	0.338	1.000	0.357	1.000	0.363
		3	0.998	0.303	1.000	0.331	1.000	0.349	1.000	0.357
	1000	1	0.998	0.204	0.998	0.221	0.998	0.233	0.998	0.240
		2	0.998	0.209	0.998	0.227	0.998	0.237	0.998	0.244
		3	0.996	0.204	0.998	0.222	0.998	0.233	0.998	0.241
	2000	1	1.000	0.147	1.000	0.159	1.000	0.166	1.000	0.171
		2	1.000	0.151	1.000	0.162	1.000	0.169	1.000	0.174
		3	1.000	0.147	1.000	0.159	1.000	0.166	1.000	0.171
S2	500	1	0.958	0.377	0.958	0.413	0.960	0.429	0.968	0.441
		2	0.964	0.389	0.956	0.418	0.960	0.436	0.968	0.447
		3	0.958	0.378	0.956	0.413	0.960	0.430	0.968	0.442
	1000	1	0.976	0.217	0.980	0.238	0.980	0.246	0.982	0.257
		2	0.976	0.224	0.982	0.244	0.982	0.251	0.982	0.259
		3	0.976	0.216	0.982	0.237	0.982	0.246	0.982	0.255
	2000	1	0.996	0.149	1.000	0.162	1.000	0.171	1.000	0.176
		2	0.996	0.154	1.000	0.166	0.998	0.174	1.000	0.178
		3	0.996	0.149	1.000	0.162	1.000	0.170	1.000	0.176

Table A.3: Comparison of the sampling CIs with different choices of M and L, U, a for settings **S1** to **S5** with $\tau = 0.4$. For column indexed with “LUa”, 1 stands for construction of $[L, U]$ as in (19) with $a = 0.6$; 2 stands for construction of $[L, U]$ as in (19) with $a = 0.8$; 3 stands for setting $[L, U] = [-5, 5]$ and $a = 0.6$. The columns indexed with “Cov” and “Len” denote the empirical coverage and the average lengths, respectively. We shall increase the value of λ such that there are $10\%M$ sampled intervals are non-empty. For example, for $M = 1000$, we increase λ until 100 sampled intervals are non-empty.

In the following Tables A.4 and A.5, we consider $\tau = 0.2$ and $\tau = 0.4$ and report the extra results for the dependence of the sampling on the choice of λ .

In the following Table A.6, we compare CI^{samp} defined in (25) and $\text{CI}^{\text{samp}}(\mathcal{M}_0)$ defined in (26) for settings **S1** to **S5** with $\tau = 0.2$ and $\tau = 0.4$. We observe that these two sampling CIs are nearly the same and the extra filtering step used for constructing $\text{CI}^{\text{samp}}(\mathcal{M}_0)$ in (26) only decreases the average lengths slightly.

		prop = 1%		prop = 5%		prop = 10%		prop = 20%		prop = 30%	
Set	n	Cov	Len	Cov	Len	Cov	Len	Cov	Len	Cov	Len
S1	500	0.982	0.231	1.000	0.298	1.000	0.336	1.000	0.383	1.000	0.412
	1000	0.966	0.155	0.994	0.210	0.996	0.240	0.998	0.272	1.000	0.294
	2000	0.954	0.099	0.998	0.142	0.998	0.167	0.998	0.194	1.000	0.210
S2	500	0.952	0.257	0.984	0.321	0.996	0.370	0.998	0.426	1.000	0.460
	1000	0.904	0.172	0.978	0.228	0.988	0.260	0.992	0.294	0.992	0.311
	2000	0.870	0.109	0.974	0.162	0.988	0.188	0.988	0.213	0.984	0.226
S3	500	0.928	0.265	0.980	0.389	0.984	0.446	0.986	0.521	0.986	0.554
	1000	0.884	0.184	0.980	0.253	0.990	0.286	0.990	0.322	0.990	0.339
	2000	0.864	0.114	0.966	0.168	0.976	0.195	0.976	0.223	0.976	0.240
S4	500	0.946	0.471	0.954	0.484	0.950	0.484	0.944	0.493	0.942	0.510
	1000	0.988	0.375	0.986	0.375	0.978	0.380	0.988	0.398	0.994	0.416
	2000	0.954	0.217	0.952	0.222	0.970	0.226	0.976	0.245	0.974	0.259
S5	500	0.698	0.306	0.852	0.379	0.876	0.407	0.844	0.424	0.836	0.430
	1000	0.688	0.280	0.802	0.308	0.760	0.310	0.696	0.304	0.688	0.305
	2000	0.870	0.254	0.866	0.254	0.858	0.256	0.856	0.266	0.856	0.277

Table A.4: Choices of the tuning parameter λ of the sampling method for $\tau = 0.2$. The resampling size M is set as 1000. We shall increase the value of λ such that there are $1000 \cdot \text{prop}$ sampled intervals are non-empty. For example, for $\text{prop} = 10\%$, we increase λ until 100 sampled intervals are non-empty.

		prop = 1%		prop = 5%		prop = 10%		prop = 20%		prop = 30%	
Set	n	Cov	Len	Cov	Len	Cov	Len	Cov	Len	Cov	Len
S1	500	0.978	0.214	1.000	0.290	1.000	0.330	1.000	0.389	1.000	0.423
	1000	0.980	0.168	0.998	0.198	1.000	0.222	1.000	0.254	1.000	0.277
	2000	0.992	0.133	1.000	0.143	1.000	0.159	1.000	0.179	1.000	0.194
S2	500	0.884	0.268	0.954	0.354	0.958	0.410	0.964	0.467	0.964	0.495
	1000	0.968	0.190	0.980	0.214	0.980	0.234	0.982	0.269	0.982	0.288
	2000	0.986	0.141	0.996	0.148	0.996	0.163	1.000	0.185	1.000	0.200
S3	500	0.864	0.345	0.916	0.478	0.906	0.518	0.902	0.560	0.904	0.583
	1000	0.878	0.229	0.938	0.287	0.936	0.316	0.936	0.347	0.936	0.364
	2000	0.986	0.148	0.996	0.158	0.994	0.172	0.994	0.197	0.994	0.212
S4	500	0.890	0.791	0.882	0.787	0.888	0.799	0.896	0.827	0.898	0.849
	1000	0.944	0.563	0.962	0.572	0.970	0.589	0.982	0.612	0.988	0.630
	2000	0.942	0.199	0.948	0.206	0.974	0.223	0.978	0.244	0.982	0.257
S5	500	0.620	0.697	0.660	0.711	0.614	0.701	0.590	0.702	0.584	0.705
	1000	0.940	0.784	0.948	0.786	0.962	0.795	0.962	0.808	0.966	0.819
	2000	0.942	0.283	0.958	0.291	0.968	0.303	0.980	0.321	0.984	0.333

Table A.5: Comparison of the sampling CIs with different λ for $\tau = 0.4$. The resampling size M is set as 1000. The columns indexed with "Cov" and "Len" denote the empirical coverage and the average lengths, respectively. We shall increase the value of λ such that there are $1000 \cdot \text{prop}$ sampled intervals are non-empty. For example, for $\text{prop} = 10\%$, we increase λ until 100 sampled intervals are non-empty.

			$\text{CI}^{\text{samp}}(\mathcal{M}_0)$ in (26)		CI^{samp} in (25)	
set	τ	n	Cov	Len	Cov	Len
S1	0.2	500	1.000	0.339	1.000	0.344
		1000	0.998	0.240	0.998	0.243
		2000	0.998	0.167	1.000	0.168
	0.4	500	1.000	0.332	1.000	0.339
		1000	0.998	0.225	0.998	0.228
		2000	1.000	0.160	1.000	0.161
S2	0.2	500	0.998	0.374	0.998	0.377
		1000	0.996	0.263	0.996	0.266
		2000	0.984	0.188	0.988	0.190
	0.4	500	0.960	0.413	0.960	0.417
		1000	0.980	0.234	0.980	0.238
		2000	1.000	0.163	1.000	0.164
S3	0.2	500	0.986	0.453	0.986	0.460
		1000	0.990	0.286	0.988	0.291
		2000	0.972	0.196	0.976	0.197
	0.4	500	0.908	0.527	0.906	0.533
		1000	0.936	0.316	0.936	0.320
		2000	0.994	0.173	0.994	0.176
S4	0.2	500	0.940	0.487	0.946	0.497
		1000	0.990	0.381	0.990	0.386
		2000	0.968	0.227	0.962	0.233
	0.4	500	0.892	0.799	0.888	0.807
		1000	0.968	0.587	0.976	0.593
		2000	0.974	0.224	0.966	0.226
S5	0.2	500	0.880	0.408	0.882	0.418
		1000	0.756	0.307	0.766	0.316
		2000	0.860	0.254	0.854	0.261
	0.4	500	0.608	0.697	0.620	0.709
		1000	0.958	0.794	0.962	0.798
		2000	0.974	0.303	0.978	0.307

Table A.6: Comparison of CI^{samp} defined in (25) and $\text{CI}^{\text{samp}}(\mathcal{M}_0)$ defined in (26) for settings **S1** to **S5** with $\tau = 0.2$ and $\tau = 0.4$. The columns indexed with “Cov” and “Len” denote the empirical coverage and the average lengths, respectively.

B Additional Method and Theory

B.1 Homoscedastic setting

We assume the homoscedastic regression errors: $\mathbf{E}(e_i^2 | Z_i, X_i) = \sigma_\epsilon^2$ and $\mathbf{E}(\delta_i^2 | Z_i, X_i) = \sigma_\delta^2$. We estimate the matrices \mathbf{V}^Γ , \mathbf{V}^γ and \mathbf{C} in (6) by

$$\widehat{\mathbf{V}}^\Gamma = \widehat{\sigma}_\epsilon^2 \widehat{\Omega}, \quad \widehat{\mathbf{V}}^\gamma = \widehat{\sigma}_\delta^2 \widehat{\Omega}, \quad \widehat{\mathbf{C}} = \widehat{\sigma}_{\epsilon,\delta} \widehat{\Omega} \quad \text{with} \quad \widehat{\Omega} = [(W^\top W/n)^{-1}]_{1:p_z, 1:p_z},$$

where $\widehat{\sigma}_\epsilon^2 = \|Y - Z\widehat{\Gamma} - X\widehat{\Psi}\|_2^2/(n-1)$, $\widehat{\sigma}_\delta^2 = \|D - Z\widehat{\gamma} - X\widehat{\psi}\|_2^2/(n-1)$, and

$$\widehat{\sigma}_{\epsilon,\delta} = (Y - Z\widehat{\Gamma} - X\widehat{\Psi})^\top (D - Z\widehat{\gamma} - X\widehat{\psi}) / (n-1).$$

In the homoscedastic setting, we can simplify $\widehat{\text{SE}}(\widehat{\pi}_k^{[j]})$ defined in (32) as

$$\widehat{\text{SE}}(\widehat{\pi}_k^{[j]}) = \sqrt{(\widehat{\sigma}_\epsilon^2 + (\widehat{\beta}^{[j]})^2 \widehat{\sigma}_\delta^2 - 2\widehat{\beta}^{[j]} \widehat{\sigma}_{\epsilon,\delta})/n \cdot \sqrt{\widehat{\Omega}_{kk} - 2\widehat{\gamma}_k/\widehat{\gamma}_j \cdot \widehat{\Omega}_{jk} + (\widehat{\gamma}_k/\widehat{\gamma}_j)^2 \widehat{\Omega}_{jj}}}.$$

B.2 Equivalent Definition of $\widehat{\mathcal{V}}^{\text{TSHT}}$ in (34)

Recall $\widehat{\mathcal{W}} = \arg \max_{1 \leq j \leq |\widehat{\mathcal{S}}|} \|\widehat{\Pi}_j\|_0$. With $\widehat{\mathcal{W}}$, we further define the index set $\widetilde{\mathcal{V}}$ as

$$\widetilde{\mathcal{V}} = \cup_{j \in \widehat{\mathcal{W}}} \left\{ 1 \leq k \leq |\widehat{\mathcal{S}}| : \widehat{\Pi}_{j,k} = 1 \right\}. \quad (38)$$

The set $\widetilde{\mathcal{V}}$ denotes the set of IVs who support (and are also supported by) at least one element in $\widehat{\mathcal{W}}$. We finally construct the index set $\widehat{\mathcal{V}} \subset \{1, 2, \dots, |\widehat{\mathcal{S}}|\}$ as

$$\widehat{\mathcal{V}}^{\text{TSHT}} = \cup_{k \in \widetilde{\mathcal{V}}} \left\{ 1 \leq l \leq |\widehat{\mathcal{S}}| : \widehat{\Pi}_{k,l} = 1 \right\}. \quad (39)$$

This set $\widehat{\mathcal{V}}^{\text{TSHT}}$ contains all candidate IVs that are claimed to be valid by at least one element from $\widetilde{\mathcal{V}}$. The set defined in (39) is equivalent to that in (34). We illustrate the definitions of the voting matrix, $\widehat{\Pi}$, $\widehat{\mathcal{W}}$ and $\widehat{\mathcal{V}}^{\text{TSHT}}$ using the following example.

Example 3 We consider an example with $p_z = 8$ IVs, where $\{Z_1, Z_2, Z_3, Z_4\}$ are valid IVs, $\{Z_5, Z_6, Z_7\}$ are invalid IVs sharing the same invalidity level and Z_8 is invalid with a different invalidity level. The left panel of Table B.1 corresponds to a favorable scenario where the valid IVs $\{Z_1, Z_2, Z_3, Z_4\}$ only vote for each other. On the right panel of Table B.1, the candidate IV Z_5 receives the votes (by mistake) from three valid IVs $\{Z_2, Z_3, Z_4\}$. This might happen when the IV Z_5 is a locally invalid IV.

On the left panel of Table B.1, we have $\widehat{\mathcal{V}}^{\text{TSHT}} = \widehat{\mathcal{W}} = \{1, 2, 3, 4\} = \mathcal{V}$ and the property (30) is satisfied. On the right panel of Table B.1, we have $\widehat{\mathcal{W}} = \{5\}$ and $\widehat{\mathcal{V}}^{\text{TSHT}} = \{1, 2, 3, 4, 5, 6, 7\}$. Only $\widehat{\mathcal{V}}^{\text{TSHT}}$ satisfies (30) but not $\widehat{\mathcal{W}}$.

	Z ₁	Z ₂	Z ₃	Z ₄	Z ₅	Z ₆	Z ₇	Z ₈
Z ₁	✓	✓	✓	✓	×	×	×	×
Z ₂	✓	✓	✓	✓	×	×	×	×
Z ₃	✓	✓	✓	✓	×	×	×	×
Z ₄	✓	✓	✓	✓	×	×	×	×
Z ₅	×	×	×	×	✓	✓	✓	×
Z ₆	×	×	×	×	✓	✓	✓	×
Z ₇	×	×	×	×	✓	✓	✓	×
Z ₈	×	×	×	×	×	×	×	✓
Votes	4	4	4	4	3	3	3	1

	Z ₁	Z ₂	Z ₃	Z ₄	Z ₅	Z ₆	Z ₇	Z ₈
Z ₁	✓	✓	✓	✓	×	×	×	×
Z ₂	✓	✓	✓	✓	✓	×	×	×
Z ₃	✓	✓	✓	✓	✓	×	×	×
Z ₄	✓	✓	✓	✓	✓	×	×	×
Z ₅	×	✓	✓	✓	✓	✓	✓	×
Z ₆	×	×	×	×	✓	✓	✓	×
Z ₇	×	×	×	×	✓	✓	✓	×
Z ₈	×	×	×	×	×	×	×	✓
Votes	4	5	5	5	6	3	3	1

Table B.1: The left voting matrix $\widehat{\Pi}$ denotes that all valid IVs $\{Z_1, Z_2, Z_3, Z_4\}$ support each other but not any other invalid IV. The right voting matrix $\widehat{\Pi}$ denotes that the (weakly) invalid IV Z_5 receives support from valid IVs $\{Z_2, Z_3, Z_4\}$ and invalid IVs $\{Z_6, Z_7\}$.

C Proofs

C.1 Proof Preparation

Throughout the proof, we focus on the low-dimensional setting with heteroscedastic errors. Define $\Omega = \Sigma^{-1}$ and $\widehat{\Omega} = \widehat{\Sigma}^{-1}$. The OLS estimators $\widehat{\gamma}$ and $\widehat{\Gamma}$ defined in (5) satisfy the following expression,

$$\widehat{\gamma}_j - \gamma_j^* = \widehat{\Omega}_j^\top \frac{1}{n} W^\top \delta \quad \text{and} \quad \widehat{\Gamma}_j - \Gamma_j^* = \widehat{\Omega}_j^\top \frac{1}{n} W^\top \epsilon \quad \text{for } 1 \leq j \leq p. \quad (40)$$

The following lemma states the asymptotic properties of the reduced-form estimators. This lemma was proved in the proof of Theorem 4.2 in [54]; see also Section 4.2.3 in [54].

Lemma 1 *Consider the model (4). Suppose that Conditions (C1) and (C2) hold, then we have*

$$\sqrt{n} \begin{pmatrix} \widehat{\Gamma} - \Gamma^* \\ \widehat{\gamma} - \gamma^* \end{pmatrix} \xrightarrow{d} N(\mathbf{0}, \text{Cov}) \quad \text{with} \quad \text{Cov} = \begin{pmatrix} \mathbf{V}^\Gamma & \mathbf{C} \\ \mathbf{C}^\top & \mathbf{V}^\gamma \end{pmatrix},$$

where $\mathbf{V}^\Gamma = [\Sigma^{-1} (\mathbf{E} \epsilon_i^2 W_i W_i^\top) \Sigma^{-1}]_{1:p_z, 1:p_z}$, $\mathbf{V}^\gamma = [\Sigma^{-1} (\mathbf{E} \delta_i^2 W_i W_i^\top) \Sigma^{-1}]_{1:p_z, 1:p_z}$, and $\mathbf{C} = [\Sigma^{-1} (\mathbf{E} \epsilon_i \delta_i W_i W_i^\top) \Sigma^{-1}]_{1:p_z, 1:p_z}$.

Define the augmented covariance matrix

$$\text{Cov}^A = \begin{pmatrix} \Sigma^{-1} (\mathbf{E} \epsilon_i^2 W_i W_i^\top) \Sigma^{-1} & \Sigma^{-1} (\mathbf{E} \epsilon_i \delta_i W_i W_i^\top) \Sigma^{-1} \\ \Sigma^{-1} (\mathbf{E} \epsilon_i \delta_i W_i W_i^\top) \Sigma^{-1} & \Sigma^{-1} (\mathbf{E} \delta_i^2 W_i W_i^\top) \Sigma^{-1} \end{pmatrix} \in \mathbb{R}^{2p \times 2p}.$$

For any $u, v \in \mathbb{R}^p$, we have

$$(u^\top, v^\top) \text{Cov}^A (u^\top, v^\top)^\top = \mathbf{E}(U_i, V_i) (\epsilon_i, \delta_i)^\top (\epsilon_i, \delta_i) (U_i, V_i)^\top,$$

where $U_i = W_i^\top \Sigma^{-1} u$ and $V_i = W_i^\top \Sigma^{-1} v$. Since $\lambda_{\min}(\mathbf{E}[(\epsilon_i, \delta_i)^\top (\epsilon_i, \delta_i) | W_i]) \geq c_1 > 0$, $\mathbf{E} U_i^2 = u^\top \Sigma^{-1} u$, and $\mathbf{E} V_i^2 = v^\top \Sigma^{-1} v$, we have

$$(u^\top, v^\top) \text{Cov}^A (u^\top, v^\top)^\top \geq c_1 (u^\top \Sigma^{-1} u + v^\top \Sigma^{-1} v) \geq c_1 \lambda_{\min}(\Sigma^{-1}) (\|u\|_2^2 + \|v\|_2^2).$$

This further implies

$$\lambda_{\min}(\text{Cov}) \geq \lambda_{\min}(\text{Cov}^A) \geq c_1 \cdot \lambda_{\min}(\Sigma^{-1}) \geq c_1/C_0. \quad (41)$$

With a similar argument, we establish

$$\lambda_{\max}(\text{Cov}) \leq \lambda_{\max}(\text{Cov}^A) \leq C_1 \cdot \lambda_{\max}(\Sigma^{-1}) \leq C_1/c_0. \quad (42)$$

Define the matrix

$$\mathbf{R}(\beta) = \mathbf{V}^\Gamma + \beta^2 \mathbf{V}^\gamma - 2\beta \mathbf{C} \quad \text{and} \quad \widehat{\mathbf{R}}(\beta) = \widehat{\mathbf{V}}^\Gamma + \beta^2 \widehat{\mathbf{V}}^\gamma - 2\beta \widehat{\mathbf{C}}. \quad (43)$$

We apply (41) and establish

$$\lambda_{\min}(\mathbf{R}(\beta)) \geq \lambda_{\min}(\text{Cov})(1 + \beta^2) \geq c_1 \lambda_{\min}(\Sigma^{-1}). \quad (44)$$

Define $\mathcal{U}(a) = \{\beta \in \mathbb{R} : |\beta - \beta^*| \leq n^{-a}\}$ for some $a > 0.5$ and the events

$$\begin{aligned} \mathcal{E}_0(\alpha) &= \left\{ \max_{\beta \in \mathcal{U}(a)} \max_{j \in \widehat{\mathcal{S}}} \frac{|\widehat{\Gamma}_j - \Gamma_j^* - \beta(\widehat{\gamma}_j - \gamma_j^*)|}{\sqrt{(\widehat{\mathbf{V}}_{jj}^\Gamma + \beta^2 \widehat{\mathbf{V}}_{jj}^\gamma - 2\beta \widehat{\mathbf{C}}_{jj})/n}} \leq \Phi^{-1} \left(1 - \frac{\alpha}{2|\widehat{\mathcal{S}}|} \right) \right\}, \\ \widetilde{\mathcal{E}}_0(\alpha) &= \left\{ \max_{\beta \in \mathcal{U}(a)} \max_{j \in \widehat{\mathcal{S}}} \frac{|\widehat{\Gamma}_j - \beta \widehat{\gamma}_j - \pi_j^*|}{\sqrt{(\widehat{\mathbf{V}}_{jj}^\Gamma + \beta^2 \widehat{\mathbf{V}}_{jj}^\gamma - 2\beta \widehat{\mathbf{C}}_{jj})/n}} \leq \Phi^{-1} \left(1 - \frac{\alpha}{2|\widehat{\mathcal{S}}|} \right) \right\}. \end{aligned} \quad (45)$$

We present the following lemma to control the probability of the above events, whose proof is postponed to Section D.2.

Lemma 2 *Suppose that the conditions (C1) and (C2) hold, then the events $\mathcal{E}_0(\alpha)$ and $\widetilde{\mathcal{E}}_0(\alpha)$ defined in (45) satisfy*

$$\liminf_{n \rightarrow \infty} \mathbf{P}(\mathcal{E}_0(\alpha)) \geq 1 - \alpha \quad \text{and} \quad \liminf_{n \rightarrow \infty} \mathbf{P}(\widetilde{\mathcal{E}}_0(\alpha)) \geq 1 - \alpha.$$

Define the following subset of relevant IVs,

$$\mathcal{S}^0 = \left\{ 1 \leq j \leq p_z : |\gamma_j^*| \geq (\sqrt{\log n} - C(\log n)^{1/4}) \cdot \sqrt{\widehat{\mathbf{V}}_{jj}^\gamma/n} \right\}. \quad (46)$$

Define the following events

$$\begin{aligned}
\mathcal{G}_1 &= \left\{ \max \left\{ \left\| \frac{1}{n} W^\top \epsilon \right\|_\infty, \left\| \frac{1}{n} W^\top \delta \right\|_\infty \right\} \leq C \frac{(\log n)^{1/4}}{\sqrt{n}} \right\} \\
\mathcal{G}_2 &= \left\{ \max_{1 \leq j \leq p} \max \left\{ |\hat{\gamma}_j - \gamma_j^*| / \sqrt{\mathbf{V}_{jj}^\gamma / n}, |\hat{\Gamma}_j - \Gamma_j^*| / \sqrt{\mathbf{V}_{jj}^\Gamma / n} \right\} \leq C (\log n)^{1/4} \right\} \\
\mathcal{G}_3 &= \left\{ \|\hat{\Omega} - \Sigma^{-1}\|_2 \leq C \sqrt{\log n / n} \right\} \\
\mathcal{G}_4 &= \left\{ \max \left\{ \|\hat{\mathbf{V}}^\Gamma - \mathbf{V}^\Gamma\|_2, \|\hat{\mathbf{V}}^\gamma - \mathbf{V}^\gamma\|_2, \|\hat{\mathbf{C}} - \mathbf{C}\|_2 \right\} \leq C (\log n)^{3/2} / \sqrt{n} \right\} \\
\mathcal{G}_5 &= \left\{ \mathcal{S}_{\text{str}} \subset \hat{\mathcal{S}} \subset \mathcal{S}^0 \subset \mathcal{S} \right\} \\
\mathcal{G}_6 &= \left\{ \max_{j, k \in \hat{\mathcal{S}}} \left| \frac{\hat{\gamma}_k / \hat{\gamma}_j}{\gamma_k^* / \gamma_j^*} - 1 \right| \leq C \frac{1}{(\log n)^{1/4}} \right\} \\
\mathcal{G}_7 &= \left\{ \max_{j \in \hat{\mathcal{S}}} \left| \frac{\hat{\Gamma}_j}{\hat{\gamma}_j} - \frac{\Gamma_j^*}{\gamma_j^*} \right| \leq C \left(1 + \left| \frac{\Gamma_j^*}{\gamma_j^*} \right| \right) \frac{1}{(\log n)^{1/4}} \right\}.
\end{aligned} \tag{47}$$

Define $\mathcal{G} = \bigcap_{j=1}^7 \mathcal{G}_j$. The following lemmas control the probability of \mathcal{G} , whose proof is presented in Section D.1.

Lemma 3 *Suppose that the conditions (C1) and (C2) hold, then for a sufficiently large n ,*

$$\mathbf{P}(\mathcal{G}) \geq 1 - \exp(-c\sqrt{\log n}),$$

for some positive constant $c > 0$.

Note that

$$\max_{\beta \in \mathcal{U}(a)} \|\mathbf{R}(\beta^*) - \mathbf{R}(\beta)\|_2 \leq C n^{-a}. \tag{48}$$

On the event \mathcal{G}_4 , we have

$$\max_{1 \leq j \leq p_z} \left| \hat{\mathbf{R}}_{jj}(\beta) - \mathbf{R}_{jj}(\beta) \right| \leq C \frac{(1 + \beta^2)(\log n)^{3/2}}{\sqrt{n}}, \tag{49}$$

for some positive constant $C > 0$. Together with (44), we have

$$\min_{1 \leq j \leq p_z} \mathbf{R}_{jj}(\beta) \geq \lambda_{\min}(\text{Cov}) / 2(1 + \beta^2). \tag{50}$$

On the event \mathcal{G} ,

$$\max_{j \in \hat{\mathcal{S}}} \hat{\mathbf{R}}_{jj}(\beta) = \max_{j \in \hat{\mathcal{S}}} \left[\hat{\mathbf{V}}_{jj}^\Gamma + \beta^2 \hat{\mathbf{V}}_{jj}^\gamma - 2\beta \hat{\mathbf{C}}_{jj} \right] \leq C(1 + \beta^2).$$

and hence

$$\max_{j \in \hat{\mathcal{S}}} \hat{\rho}_j(\beta) \lesssim \sqrt{\frac{1 + \beta^2}{n}}. \tag{51}$$

C.2 Proof of Theorem 1

C.2.1 The coverage property of CI^{sear} defined in (17)

By the decomposition (13), if β is taken as β^* , then

$$(\widehat{\Gamma}_j - \beta^* \widehat{\gamma}_j) - \pi_j^* = \widehat{\Gamma}_j - \Gamma_j^* - \beta^* (\widehat{\gamma}_j - \gamma_j^*).$$

Hence, on the event $\mathcal{E}_0(\alpha)$ defined in (45), for all $j \in \mathcal{V} \cap \widehat{\mathcal{S}}$,

$$\left| \widehat{\Gamma}_j - \beta^* \widehat{\gamma}_j \right| \leq \widehat{\rho}_j(\beta^*)$$

where $\widehat{\rho}_j(\beta^*)$ is defined in (15). This leads to

$$\left| \left\{ j \in \widehat{\mathcal{S}} : \left| \widehat{\Gamma}_j - \beta^* \widehat{\gamma}_j \right| \leq \widehat{\rho}_j(\beta^*) \right\} \right| \geq \left| \mathcal{V} \cap \widehat{\mathcal{S}} \right|. \quad (52)$$

On the event \mathcal{G}_5 defined in (47), we have

$$\left| \mathcal{V} \cap \widehat{\mathcal{S}} \right| \geq \left| \mathcal{V} \cap \mathcal{S}_{\text{str}} \right| > \frac{|\mathcal{S}|}{2} \geq \frac{|\widehat{\mathcal{S}}|}{2}, \quad (53)$$

where the second inequality follows from the finite-sample majority rule and the last inequality follows from the definition of \mathcal{G}_5 in (47).

By combining (52) and (53), we show that, on the event $\mathcal{E}_0(\alpha) \cap \mathcal{G}_5$,

$$\|\widehat{\pi}_{\widehat{\mathcal{S}}}(\beta^*)\|_0 \leq |\widehat{\mathcal{S}}| - \left| \mathcal{V} \cap \widehat{\mathcal{S}} \right| < \frac{|\widehat{\mathcal{S}}|}{2}. \quad (54)$$

Hence, on the event $\mathcal{E}_0(\alpha) \cap \mathcal{G}_5$, $\beta^* \in \text{CI}^{\text{sear}}$, that is,

$$\mathbf{P}(\beta^* \in \text{CI}^{\text{sear}}) \geq \mathbf{P}(\mathcal{E}_0(\alpha) \cap \mathcal{G}_5).$$

We establish the coverage property by applying Lemmas 2 and 3.

C.2.2 The coverage property of $\widehat{\text{CI}}^{\text{sear}}$ defined in (20)

If $j \in \mathcal{V} \cap \widehat{\mathcal{S}}$, we have

$$\mathbf{P} \left(\widehat{\Gamma}_j / \widehat{\gamma}_j - \sqrt{\log n \cdot \widehat{\text{Var}} \left(\widehat{\Gamma}_j / \widehat{\gamma}_j \right)} \leq \beta^* \leq \widehat{\Gamma}_j / \widehat{\gamma}_j + \sqrt{\log n \cdot \widehat{\text{Var}} \left(\widehat{\Gamma}_j / \widehat{\gamma}_j \right)} \right) \rightarrow 1,$$

where $\widehat{\text{Var}} \left(\widehat{\Gamma}_j / \widehat{\gamma}_j \right) = \frac{1}{n} \left(\widehat{\mathbf{V}}_{jj}^{\Gamma} / \widehat{\gamma}_j^2 + \widehat{\mathbf{V}}_{jj}^{\gamma} \widehat{\Gamma}_j^2 / \widehat{\gamma}_j^4 - 2 \widehat{\mathbf{C}}_{jj} \widehat{\Gamma}_j / \widehat{\gamma}_j^3 \right)$. Then this implies

$$\mathbf{P}(\beta^* \in [L, U]) \rightarrow 1.$$

We consider two cases

1. $\beta^* \in \mathcal{B}$;

2. $\beta^* \notin \mathcal{B}$. By the construction of \mathcal{B} , there exists $\beta^L, \beta^U \in \mathcal{B}$ such that $\beta^L \leq \beta^* \leq \beta^U$ and $\beta^U - \beta^L \leq n^{-a}$ for $a > 0.5$.

For the case (a), we can apply the same proof as that in Section C.2.1. In the following, we shall modify the proof in Section C.2.1 and establish the coverage property of $\widehat{\text{CI}}^{\text{sear}}$ for the case (b). It follows from (13) that

$$(\widehat{\Gamma}_j - \beta^L \widehat{\gamma}_j) - \pi_j^* = \widehat{\Gamma}_j - \Gamma_j^* - \beta^L (\widehat{\gamma}_j - \gamma_j^*) + (\beta^* - \beta^L) \gamma_j^*.$$

On the event $\widetilde{\mathcal{E}}_0(\alpha)$ defined in (45), we have

$$\left| \widehat{\Gamma}_j - \beta^L \widehat{\gamma}_j \right| \leq \widehat{\rho}_j(\beta^L) \quad \text{for all } j \in \mathcal{V} \cap \widehat{\mathcal{S}},$$

which implies

$$\left| \left\{ j \in \widehat{\mathcal{S}} : \left| \widehat{\Gamma}_j - \beta^L \widehat{\gamma}_j \right| \leq \widehat{\rho}_j(\beta^L) \right\} \right| \geq \left| \mathcal{V} \cap \widehat{\mathcal{S}} \right|. \quad (55)$$

By the same argument as in (53) and (54), we establish that, on the event $\widetilde{\mathcal{E}}_0(\alpha) \cap \mathcal{G}$,

$$\|\widehat{\pi}_{\widehat{\mathcal{S}}}(\beta^L)\|_0 \leq |\widehat{\mathcal{S}}| - |\mathcal{V} \cap \widehat{\mathcal{S}}| < \frac{|\widehat{\mathcal{S}}|}{2}.$$

That is, $\beta^L \in (\beta_{\min}, \beta_{\max})$. With a similar argument, on the event $\widetilde{\mathcal{E}}_0(\alpha) \cap \mathcal{G}$, we have $\beta^U \in (\beta_{\min}, \beta_{\max})$. Then we establish

$$\mathbf{P}(\beta^* \in (\beta^L, \beta^U) \subset (\beta_{\min}, \beta_{\max})) \geq \mathbf{P}(\widetilde{\mathcal{E}}_0(\alpha) \cap \mathcal{G}).$$

We establish the coverage property by applying Lemmas 2 and 3.

C.2.3 Length of CI^{sear} and $\widehat{\text{CI}}^{\text{sear}}$

For the j -th IV with $\pi_j^* = 0$, we simplify the decomposition in (13) as

$$\widehat{\Gamma}_j - \beta \widehat{\gamma}_j = \widehat{\Gamma}_j - \Gamma_j^* - \beta (\widehat{\gamma}_j - \gamma_j^*) + (\beta^* - \beta) \gamma_j^*.$$

In the remaining of the proof, we assume that the event $\mathcal{E}_0(\alpha) \cap \mathcal{G}$ happens. For β satisfying $|\gamma_j^*| \cdot |\beta - \beta^*| \geq 2\widehat{\rho}_j(\beta)$, we have $\widehat{\pi}_j(\beta) \neq 0$. Consequently, if β satisfies

$$|\beta - \beta^*| \geq \max_{j \in \widehat{\mathcal{S}} \cap \mathcal{V}} \frac{2\widehat{\rho}_j(\beta)}{|\gamma_j^*|}, \quad (56)$$

then

$$\|\widehat{\pi}_{\widehat{\mathcal{S}}}(\beta)\|_0 \geq |\widehat{\mathcal{S}} \cap \mathcal{V}| > \frac{|\widehat{\mathcal{S}}|}{2},$$

where the second inequality follows from (53). That is, for β satisfying (56), $\beta \notin \text{CI}^{\text{sear}}$ and $\beta \notin \widehat{\text{CI}}^{\text{sear}}$. If $\beta \in \text{CI}^{\text{sear}}$ or $\beta \in \widehat{\text{CI}}^{\text{sear}}$, then we apply (56) and obtain

$$|\beta - \beta^*| \leq \max_{j \in \widehat{\mathcal{S}} \cap \mathcal{V}} \frac{2\widehat{\rho}_j(\beta)}{|\gamma_j^*|} \leq C \frac{\sqrt{1 + \beta^2}}{\sqrt{n} \min_{j \in \widehat{\mathcal{S}} \cap \mathcal{V}} |\gamma_j^*|}, \quad (57)$$

where the last inequality follows from (51). By the definition of \mathcal{S}^0 and the event \mathcal{G}_5 , we further simplify the above inequality as

$$|\beta - \beta^*| \leq C \frac{\sqrt{1 + \beta^2}}{\sqrt{\log n}}.$$

If $|\beta| \geq 2|\beta^*|$, then the above inequality implies $|\beta| \leq 2C/\sqrt{\log n - 4C^2}$. This implies that, if $\beta \in \text{CI}^{\text{sear}}$ or $\beta \in \widehat{\text{CI}}^{\text{sear}}$, then there exists some positive constant $C > 0$ such that

$$|\beta| \leq \max\{2|\beta^*|, \frac{2C}{\log n}\} \leq C. \quad (58)$$

We apply (57) again and establish that, if $\beta \in \text{CI}^{\text{sear}}$ or $\beta \in \widehat{\text{CI}}^{\text{sear}}$, then

$$|\beta - \beta^*| \leq \max_{|\beta| \leq C} \max_{j \in \widehat{\mathcal{S}} \cap \mathcal{V}} \frac{4\widehat{\rho}_j(\beta)}{|\gamma_j^*|}.$$

That is,

$$\max\left\{\mathbf{L}(\text{CI}^{\text{sear}}), \mathbf{L}(\widehat{\text{CI}}^{\text{sear}})\right\} \leq \max_{|\beta| \leq C} \max_{j \in \widehat{\mathcal{S}} \cap \mathcal{V}} \frac{4\widehat{\rho}_j(\beta)}{|\gamma_j^*|},$$

where $C > 0$ is some positive constant. We apply (51) and establish

$$\max_{|\beta| \leq C} \max_{j \in \widehat{\mathcal{S}} \cap \mathcal{V}} \widehat{\rho}_j(\beta) \lesssim 1/\sqrt{n}.$$

Then we establish the upper bound for the length of CI^{sear} and $\widehat{\text{CI}}^{\text{sear}}$.

C.3 Proof of Proposition 1

We shall establish the following result, which will imply Proposition 1.

$$\liminf_{n \rightarrow \infty} \mathbf{P} \left(\min_{m \in \mathcal{M}_0} \left[\max_{\beta \in \mathcal{U}(a)} \max_{j \in \widehat{\mathcal{S}}} \frac{|\widehat{\Gamma}_j^{[m]} - \Gamma_j^* - \beta(\widehat{\gamma}_j^{[m]} - \gamma_j^*)|}{\sqrt{(\widehat{\mathbf{V}}_{jj}^\Gamma + \beta^2 \widehat{\mathbf{V}}_{jj}^\gamma - 2\beta \widehat{\mathbf{C}}_{jj})/n}} \right] \leq C \text{err}_n(M, \alpha_0) \right) \geq 1 - \alpha_0, \quad (59)$$

where \mathcal{M}_0 is defined as

$$\mathcal{M}_0 = \left\{ 1 \leq m \leq M : \max_{j \in \widehat{\mathcal{S}}} \left\{ |\widehat{\gamma}_j^{[m]} - \widehat{\gamma}_j| / \sqrt{\widehat{\mathbf{V}}_{jj}^\gamma/n}, |\widehat{\Gamma}_j^{[m]} - \widehat{\Gamma}_j| / \sqrt{\widehat{\mathbf{V}}_{jj}^\Gamma/n} \right\} \leq 1.1 \Phi^{-1} \left(1 - \frac{\alpha_0}{4|\widehat{\mathcal{S}}|} \right) \right\}. \quad (60)$$

Denote the observed data by \mathcal{O} , that is, $\mathcal{O} = \{Y_i, D_i, Z_i, X_i\}_{1 \leq i \leq n}$. Define

$$\widehat{U} = \sqrt{n} \left[\begin{pmatrix} \widehat{\Gamma}_{\widehat{\mathcal{S}}} \\ \widehat{\gamma}_{\widehat{\mathcal{S}}} \end{pmatrix} - \begin{pmatrix} \Gamma_{\widehat{\mathcal{S}}}^* \\ \gamma_{\widehat{\mathcal{S}}}^* \end{pmatrix} \right] \quad \text{and} \quad U^{[m]} = \sqrt{n} \left[\begin{pmatrix} \widehat{\Gamma}_{\widehat{\mathcal{S}}} \\ \widehat{\gamma}_{\widehat{\mathcal{S}}} \end{pmatrix} - \begin{pmatrix} \widehat{\Gamma}_{\widehat{\mathcal{S}}}^{[m]} \\ \widehat{\gamma}_{\widehat{\mathcal{S}}}^{[m]} \end{pmatrix} \right] \quad \text{for } 1 \leq m \leq M.$$

By slightly abusing the notation, we use the following notations throughout the rest of the proof of Proposition 1,

$$\text{Cov} = \begin{pmatrix} \mathbf{V}_{\widehat{\mathcal{S}}, \widehat{\mathcal{S}}}^\Gamma & \mathbf{C}_{\widehat{\mathcal{S}}, \widehat{\mathcal{S}}} \\ \mathbf{C}_{\widehat{\mathcal{S}}, \widehat{\mathcal{S}}}^\top & \mathbf{V}_{\widehat{\mathcal{S}}, \widehat{\mathcal{S}}}^\gamma \end{pmatrix} \quad \text{and} \quad \widehat{\text{Cov}} = \begin{pmatrix} \widehat{\mathbf{V}}_{\widehat{\mathcal{S}}, \widehat{\mathcal{S}}}^\Gamma & \widehat{\mathbf{C}}_{\widehat{\mathcal{S}}, \widehat{\mathcal{S}}} \\ \widehat{\mathbf{C}}_{\widehat{\mathcal{S}}, \widehat{\mathcal{S}}}^\top & \widehat{\mathbf{V}}_{\widehat{\mathcal{S}}, \widehat{\mathcal{S}}}^\gamma \end{pmatrix}.$$

Recall that \widehat{U} and $\widehat{\mathcal{S}}$ are functions of the observed data \mathcal{O} , and

$$U^{[m]} | \mathcal{O} \stackrel{i.i.d.}{\sim} N(\mathbf{0}, \widehat{\text{Cov}}) \quad \text{for } 1 \leq m \leq M.$$

Let $f(\cdot | \mathcal{O})$ denote the conditional density function of $U^{[m]}$ given the data \mathcal{O} , that is,

$$f(U^m = U | \mathcal{O}) = \frac{1}{\sqrt{(2\pi)^{2|\widehat{\mathcal{S}}|} \det(\widehat{\text{Cov}})}} \exp\left(-\frac{1}{2} U^\top \widehat{\text{Cov}}^{-1} U\right).$$

We define the following event for the data \mathcal{O} ,

$$\mathcal{E}_1 = \left\{ \|\widehat{\text{Cov}} - \text{Cov}\|_2 < c_2 \right\} \quad (61)$$

where $\|\widehat{\text{Cov}} - \text{Cov}\|_2$ denotes the spectral norm of the matrix $\widehat{\text{Cov}} - \text{Cov}$ and $0 < c_2 < \lambda_{\min}(\text{Cov})/2$ is a small positive constant.

We define the following function to facilitate the proof,

$$g(U) = \frac{1}{\sqrt{(2\pi)^{2|\widehat{\mathcal{S}}|} \det(\text{Cov} + c_2 \mathbf{I})}} \exp\left(-\frac{1}{2} U^\top (\text{Cov} + c_2 \mathbf{I})^{-1} U\right). \quad (62)$$

On the event $\mathcal{O} \in \mathcal{E}_1$, we have $\text{Cov} + c_2 \mathbf{I} \succ \widehat{\text{Cov}} \succ \text{Cov} - c_2 \mathbf{I} \succ \frac{1}{2} \lambda_{\min}(\text{Cov}) \cdot \mathbf{I}$, where two matrices A and B satisfy $A \succ B$ if $A - B$ is a positive definite matrix. Hence

$$f(U^{[m]} = U | \mathcal{O}) \cdot \mathbf{1}_{\mathcal{O} \in \mathcal{E}_1} \geq g(U) \cdot \mathbf{1}_{\mathcal{O} \in \mathcal{E}_1}. \quad (63)$$

We define the following event for the data \mathcal{O} ,

$$\mathcal{E}_2 = \left\{ \max_{j \in \widehat{\mathcal{S}}} \max \left\{ |\widehat{\gamma}_j - \gamma_j^*| / \sqrt{\widehat{\mathbf{V}}_{jj}^\gamma / n}, |\widehat{\Gamma}_j - \Gamma_j^*| / \sqrt{\widehat{\mathbf{V}}_{jj}^\Gamma / n} \right\} \leq \Phi^{-1} \left(1 - \frac{\alpha_0}{4|\widehat{\mathcal{S}}|} \right) \right\} \quad (64)$$

By the definition of \mathcal{G} and Lemma 1, we apply the union bound and establish

$$\liminf_{n \rightarrow \infty} \mathbf{P}(\mathcal{E}_1 \cap \mathcal{E}_2) \geq 1 - \alpha_0. \quad (65)$$

By the definition of the event $\mathcal{E}_1 \cap \mathcal{E}_2$, we now establish a lower bound for $g(\widehat{U})$. On the event $\mathcal{E}_1 \cap \mathcal{E}_2$, we have

$$\begin{aligned} \widehat{U}^\top (\text{Cov} - c_2 \mathbf{I})^{-1} \widehat{U} &\leq \frac{\sum_{j \in \widehat{\mathcal{S}}} \widehat{\mathbf{V}}_{jj}^\gamma + \sum_{j \in \widehat{\mathcal{S}}} \widehat{\mathbf{V}}_{jj}^\Gamma}{\lambda_{\min}(\text{Cov})/2} \cdot \left[\Phi^{-1} \left(1 - \frac{\alpha_0}{4|\widehat{\mathcal{S}}|} \right) \right]^2 \\ &\leq \frac{|\widehat{\mathcal{S}}| \cdot 3\lambda_{\max}(\text{Cov})}{\lambda_{\min}(\text{Cov})} \cdot \left[\Phi^{-1} \left(1 - \frac{\alpha_0}{4|\widehat{\mathcal{S}}|} \right) \right]^2. \end{aligned}$$

Hence, we apply the definition of the g function in (62) and establish

$$g(\widehat{U}) \cdot \mathbf{1}_{\mathcal{E}_1 \cap \mathcal{E}_2} \geq c^*(\alpha_0) \quad \text{with } c^*(\alpha_0) \text{ defined in (36)}. \quad (66)$$

If $m \notin \mathcal{M}_0$, there exists $j \in \widehat{\mathcal{S}}$ such that

$$\frac{|\widehat{\gamma}_j^{[m]} - \widehat{\gamma}_j|}{\sqrt{\widehat{\mathbf{V}}_{jj}^\gamma/n}} \geq 1.1\Phi^{-1}\left(1 - \frac{\alpha_0}{4|\widehat{\mathcal{S}}|}\right) \quad \text{or} \quad \frac{|\widehat{\Gamma}_j^{[m]} - \widehat{\Gamma}_j|}{\sqrt{\widehat{\mathbf{V}}_{jj}^\Gamma/n}} \geq 1.1\Phi^{-1}\left(1 - \frac{\alpha_0}{4|\widehat{\mathcal{S}}|}\right).$$

On the event \mathcal{E}_2 , we further establish that, there exists $j \in \widehat{\mathcal{S}}$ such that

$$\frac{|\widehat{\gamma}_j^{[m]} - \gamma_j^*|}{\sqrt{\widehat{\mathbf{V}}_{jj}^\gamma/n}} \geq 0.1\Phi^{-1}\left(1 - \frac{\alpha_0}{4|\widehat{\mathcal{S}}|}\right) \quad \text{or} \quad \frac{|\widehat{\Gamma}_j^{[m]} - \Gamma_j^*|}{\sqrt{\widehat{\mathbf{V}}_{jj}^\Gamma/n}} \geq 0.1\Phi^{-1}\left(1 - \frac{\alpha_0}{4|\widehat{\mathcal{S}}|}\right).$$

This further implies that, on the event $\mathcal{E}_1 \cap \mathcal{E}_2$,

$$\min_{m \notin \mathcal{M}_0} \|U^{[m]} - \widehat{U}\|_\infty \geq 0.1 \min_{j \in \widehat{\mathcal{S}}} \min\{\widehat{\mathbf{V}}_{jj}^\Gamma, \widehat{\mathbf{V}}_{jj}^\gamma\} \Phi^{-1}\left(1 - \frac{\alpha_0}{4|\widehat{\mathcal{S}}|}\right) > \text{err}_n(M, \alpha_0),$$

where the last inequality holds since $\text{err}_n(M, \alpha_0) \rightarrow 0$. That is, on the event $\mathcal{E}_1 \cap \mathcal{E}_2$,

$$\left\{ \min_{m \notin \mathcal{M}_0} \|U^{[m]} - \widehat{U}\|_\infty \right\} = \left\{ \min_{1 \leq m \leq M} \|U^{[m]} - \widehat{U}\|_\infty \right\}. \quad (67)$$

We use $\mathbf{P}(\cdot | \mathcal{O})$ to denote the conditional probability with respect to the observed data \mathcal{O} . Note that

$$\begin{aligned} & \mathbf{P}\left(\min_{1 \leq m \leq M} \|U^{[m]} - \widehat{U}\|_\infty \leq \text{err}_n(M, \alpha_0) \mid \mathcal{O}\right) \\ &= 1 - \mathbf{P}\left(\min_{1 \leq m \leq M} \|U^{[m]} - \widehat{U}\|_\infty \geq \text{err}_n(M, \alpha_0) \mid \mathcal{O}\right) \\ &= 1 - \prod_{m=1}^M \left[1 - \mathbf{P}\left(\|U^{[m]} - \widehat{U}\|_\infty \leq \text{err}_n(M, \alpha_0) \mid \mathcal{O}\right)\right] \\ &\geq 1 - \exp\left[-\sum_{m=1}^M \mathbf{P}\left(\|U^{[m]} - \widehat{U}\|_\infty \leq \text{err}_n(M, \alpha_0) \mid \mathcal{O}\right)\right], \end{aligned}$$

where the second equality follows from the conditional independence of $\{U^{[m]}\}_{1 \leq m \leq M}$ given the data \mathcal{O} and the last inequality follows from $1 - x \leq e^{-x}$. By applying the above inequality and (67), we establish

$$\begin{aligned} & \mathbf{P}\left(\min_{m \in \mathcal{M}_0} \|U^{[m]} - \widehat{U}\|_\infty \leq \text{err}_n(M, \alpha_0) \mid \mathcal{O}\right) \cdot \mathbf{1}_{\mathcal{O} \in \mathcal{E}_1 \cap \mathcal{E}_2} \\ &= \mathbf{P}\left(\min_{1 \leq m \leq M} \|U^{[m]} - \widehat{U}\|_\infty \leq \text{err}_n(M, \alpha_0) \mid \mathcal{O}\right) \cdot \mathbf{1}_{\mathcal{O} \in \mathcal{E}_1 \cap \mathcal{E}_2} \\ &\geq \left(1 - \exp\left[-\sum_{m=1}^M \mathbf{P}\left(\|U^{[m]} - \widehat{U}\|_\infty \leq \text{err}_n(M, \alpha_0) \mid \mathcal{O}\right)\right]\right) \cdot \mathbf{1}_{\mathcal{O} \in \mathcal{E}_1 \cap \mathcal{E}_2} \\ &= 1 - \exp\left[-\sum_{m=1}^M \mathbf{P}\left(\|U^{[m]} - \widehat{U}\|_\infty \leq \text{err}_n(M, \alpha_0) \mid \mathcal{O}\right) \cdot \mathbf{1}_{\mathcal{O} \in \mathcal{E}_1 \cap \mathcal{E}_2}\right]. \end{aligned} \quad (68)$$

For the remaining of the proof, we establish a lower bound for

$$\mathbf{P} \left(\|U^{[m]} - \widehat{U}\|_\infty \leq \text{err}_n(M, \alpha_0) \mid \mathcal{O} \right) \cdot \mathbf{1}_{\mathcal{O} \in \mathcal{E}_1 \cap \mathcal{E}_2}, \quad (69)$$

and then apply (68) to establish a lower bound for

$$\mathbf{P} \left(\min_{m \in \mathcal{M}_0} \|U^{[m]} - \widehat{U}\|_\infty \leq \text{err}_n(M, \alpha_0) \mid \mathcal{O} \right).$$

We apply (63) and further lower bound the targeted probability in (69) as

$$\begin{aligned} & \mathbf{P} \left(\|U^{[m]} - \widehat{U}\|_\infty \leq \text{err}_n(M, \alpha_0) \mid \mathcal{O} \right) \cdot \mathbf{1}_{\mathcal{O} \in \mathcal{E}_1 \cap \mathcal{E}_2} \\ &= \int f(U^{[m]} = U \mid \mathcal{O}) \cdot \mathbf{1}_{\{\|U - \widehat{U}\|_\infty \leq \text{err}_n(M, \alpha_0)\}} dU \cdot \mathbf{1}_{\mathcal{O} \in \mathcal{E}_1 \cap \mathcal{E}_2} \\ &\geq \int g(U) \cdot \mathbf{1}_{\{\|U - \widehat{U}\|_\infty \leq \text{err}_n(M, \alpha_0)\}} dU \cdot \mathbf{1}_{\mathcal{O} \in \mathcal{E}_1 \cap \mathcal{E}_2} \\ &= \int g(\widehat{U}) \cdot \mathbf{1}_{\{\|U - \widehat{U}\|_\infty \leq \text{err}_n(M, \alpha_0)\}} dU \cdot \mathbf{1}_{\mathcal{O} \in \mathcal{E}_1 \cap \mathcal{E}_2} \\ &\quad + \int [g(U) - g(\widehat{U})] \cdot \mathbf{1}_{\{\|U - \widehat{U}\|_\infty \leq \text{err}_n(M, \alpha_0)\}} dU \cdot \mathbf{1}_{\mathcal{O} \in \mathcal{E}_1 \cap \mathcal{E}_2}. \end{aligned} \quad (70)$$

By (66), we establish

$$\begin{aligned} & \int g(\widehat{U}) \cdot \mathbf{1}_{\{\|U - \widehat{U}\|_\infty \leq \text{err}_n(M, \alpha_0)\}} dU \cdot \mathbf{1}_{\mathcal{O} \in \mathcal{E}_1 \cap \mathcal{E}_2} \\ &\geq c^*(\alpha_0) \cdot \int \mathbf{1}_{\{\|U - \widehat{U}\|_\infty \leq \text{err}_n(M, \alpha_0)\}} dU \cdot \mathbf{1}_{\mathcal{O} \in \mathcal{E}_1 \cap \mathcal{E}_2} \\ &\geq c^*(\alpha_0) \cdot [2\text{err}_n(M, \alpha_0)]^{2|\widehat{\mathcal{S}}|} \cdot \mathbf{1}_{\mathcal{O} \in \mathcal{E}_1 \cap \mathcal{E}_2}. \end{aligned} \quad (71)$$

There exists $t \in (0, 1)$ such that

$$g(U) - g(\widehat{U}) = [\nabla g(\widehat{U} + t(U - \widehat{U}))]^\top (U - \widehat{U}),$$

with

$$\nabla g(u) = \frac{1}{\sqrt{(2\pi)^{2|\widehat{\mathcal{S}}|} \det(\text{Cov} + c_2 \mathbf{I})}} \exp \left(-\frac{1}{2} u^\top (\text{Cov} - c_2 \mathbf{I})^{-1} u \right)^{-1} (\text{Cov} - c_2 \mathbf{I})^{-1} u.$$

Since $\lambda_{\min}(\text{Cov} - c_2 \mathbf{I}) \geq \lambda_{\min}(\text{Cov})/2$, then ∇g is bounded and there exists a positive constant $C > 0$ such that

$$\left| g(U) - g(\widehat{U}) \right| \leq C \sqrt{2|\widehat{\mathcal{S}}|} \|U - \widehat{U}\|_\infty.$$

Then we establish

$$\begin{aligned} & \left| \int [g(U) - g(\widehat{U})] \cdot \mathbf{1}_{\{\|U - \widehat{U}\|_\infty \leq \text{err}_n(M, \alpha_0)\}} dU \cdot \mathbf{1}_{\mathcal{O} \in \mathcal{E}_1 \cap \mathcal{E}_2} \right| \\ &\leq C \sqrt{2|\widehat{\mathcal{S}}|} \cdot \text{err}_n(M, \alpha_0) \cdot \int \mathbf{1}_{\{\|U - \widehat{U}\|_\infty \leq \text{err}_n(M, \alpha_0)\}} dU \cdot \mathbf{1}_{\mathcal{O} \in \mathcal{E}_1 \cap \mathcal{E}_2} \\ &= C \sqrt{2|\widehat{\mathcal{S}}|} \cdot \text{err}_n(M, \alpha_0) \cdot [2\text{err}_n(M, \alpha_0)]^{2|\widehat{\mathcal{S}}|} \cdot \mathbf{1}_{\mathcal{O} \in \mathcal{E}_1 \cap \mathcal{E}_2}. \end{aligned} \quad (72)$$

Since $\text{err}_n(M, \alpha_0) \rightarrow 0$ and $c^*(\alpha_0)$ is a positive constant, then we have

$$C\sqrt{2|\widehat{\mathcal{S}}|} \cdot \text{err}_n(M, \alpha_0) \leq \frac{1}{2}c^*(\alpha_0).$$

We combine the above inequality, (70), (71) and (72) and obtain

$$\mathbf{P} \left(\|U - \widehat{U}\|_\infty \leq \text{err}_n(M, \alpha_0) \mid \mathcal{O} \right) \cdot \mathbf{1}_{\mathcal{O} \in \mathcal{E}_1 \cap \mathcal{E}_2} \geq \frac{1}{2}c^*(\alpha_0) \cdot [2\text{err}_n(M, \alpha_0)]^{2|\widehat{\mathcal{S}}|} \cdot \mathbf{1}_{\mathcal{O} \in \mathcal{E}_1 \cap \mathcal{E}_2}.$$

Together with (68), we establish

$$\begin{aligned} & \mathbf{P} \left(\min_{m \in \mathcal{M}_0} \|U^{[m]} - \widehat{U}\|_\infty \leq \text{err}_n(M, \alpha_0) \mid \mathcal{O} \right) \cdot \mathbf{1}_{\mathcal{O} \in \mathcal{E}_1 \cap \mathcal{E}_2} \\ & \geq 1 - \exp \left[-M \cdot \frac{1}{2}c^*(\alpha_0) \cdot [2\text{err}_n(M, \alpha_0)]^{2|\widehat{\mathcal{S}}|} \cdot \mathbf{1}_{\mathcal{O} \in \mathcal{E}_1 \cap \mathcal{E}_2} \right] \\ & = \left(1 - \exp \left[-M \cdot \frac{1}{2}c^*(\alpha_0) \cdot [2\text{err}_n(M, \alpha_0)]^{2|\widehat{\mathcal{S}}|} \right] \right) \cdot \mathbf{1}_{\mathcal{O} \in \mathcal{E}_1 \cap \mathcal{E}_2}. \end{aligned} \quad (73)$$

With $\mathbf{E}_{\mathcal{O}}$ denoting the expectation taken with respect to the observed data \mathcal{O} , we further integrate with respect to \mathcal{O} and establish

$$\begin{aligned} & \mathbf{P} \left(\min_{m \in \mathcal{M}_0} \|U^{[m]} - \widehat{U}\|_\infty \leq \text{err}_n(M, \alpha_0) \right) \\ & = \mathbf{E}_{\mathcal{O}} \left[\mathbf{P} \left(\min_{m \in \mathcal{M}_0} \|U^{[m]} - \widehat{U}\|_\infty \leq \text{err}_n(M, \alpha_0) \mid \mathcal{O} \right) \right] \\ & \geq \mathbf{E}_{\mathcal{O}} \left[\mathbf{P} \left(\min_{m \in \mathcal{M}_0} \|U^{[m]} - \widehat{U}\|_\infty \leq \text{err}_n(M, \alpha_0) \mid \mathcal{O} \right) \cdot \mathbf{1}_{\mathcal{O} \in \mathcal{E}_1 \cap \mathcal{E}_2} \right] \\ & \geq \mathbf{E}_{\mathcal{O}} \left[\left(1 - \exp \left[-M \cdot \frac{1}{2}c^*(\alpha_0) \cdot [2\text{err}_n(M, \alpha_0)]^{2|\widehat{\mathcal{S}}|} \right] \right) \cdot \mathbf{1}_{\mathcal{O} \in \mathcal{E}_1 \cap \mathcal{E}_2} \right]. \end{aligned}$$

We choose

$$\text{err}_n(M, \alpha_0) = \frac{1}{2} \left[\frac{2 \log n}{c^*(\alpha_0)M} \right]^{\frac{1}{2|\widehat{\mathcal{S}}|}},$$

and establish

$$\mathbf{P} \left(\min_{m \in \mathcal{M}_0} \|U^{[m]} - \widehat{U}\|_\infty \leq \text{err}_n(M, \alpha_0) \right) \geq (1 - n^{-1}) \cdot \mathbf{P}(\mathcal{E}_1 \cap \mathcal{E}_2).$$

We further apply (65) and establish

$$\liminf_{n \rightarrow \infty} \mathbf{P} \left(\min_{m \in \mathcal{M}_0} \|U^{[m]} - \widehat{U}\|_\infty \leq \text{err}_n(M, \alpha_0) \right) \geq \mathbf{P}(\mathcal{E}_1 \cap \mathcal{E}_2) \geq 1 - \alpha_0.$$

We use m^* to denote the index such that

$$\|U^{[m^*]} - \widehat{U}\|_\infty = \min_{m \in \mathcal{M}_0} \|U^{[m]} - \widehat{U}\|_\infty.$$

Then we have

$$\max_{j \in \widehat{\mathcal{S}}} |\widehat{\gamma}_j^{[m^*]} - \gamma_j^*| \leq \frac{\text{err}_n(M, \alpha_0)}{\sqrt{n}}, \quad \max_{j \in \widehat{\mathcal{S}}} \left| \widehat{\Gamma}_j^{[m^*]} - \Gamma_j^* - \beta(\widehat{\gamma}_j^{[m^*]} - \gamma_j^*) \right| \leq (1 + |\beta|) \frac{\text{err}_n(M, \alpha_0)}{\sqrt{n}}. \quad (74)$$

By combining the above inequality and (50), we establish that there exists some positive constant $C > 0$ such that

$$\max_{j \in \widehat{\mathcal{S}}} \frac{|\widehat{\Gamma}_j^{[m^*]} - \Gamma_j^* - \beta(\widehat{\gamma}_j^{[m^*]} - \gamma_j^*)|}{\sqrt{(\widehat{\mathbf{V}}_{jj}^\Gamma + \beta^2 \widehat{\mathbf{V}}_{jj}^\gamma - 2\beta \widehat{\mathbf{C}}_{jj})/n}} \leq 0.9 C \text{err}_n(M, \alpha_0). \quad (75)$$

Note that

$$\begin{aligned} & \left| \max_{j \in \widehat{\mathcal{S}}} \frac{|\widehat{\Gamma}_j^{[m^*]} - \Gamma_j^* - \beta(\widehat{\gamma}_j^{[m^*]} - \gamma_j^*)|}{\sqrt{(\widehat{\mathbf{V}}_{jj}^\Gamma + \beta^2 \widehat{\mathbf{V}}_{jj}^\gamma - 2\beta \widehat{\mathbf{C}}_{jj})/n}} - \max_{j \in \widehat{\mathcal{S}}} \frac{|\widehat{\Gamma}_j^{[m^*]} - \Gamma_j^* - \beta^*(\widehat{\gamma}_j^{[m^*]} - \gamma_j^*)|}{\sqrt{(\widehat{\mathbf{V}}_{jj}^\Gamma + [\beta^*]^2 \widehat{\mathbf{V}}_{jj}^\gamma - 2\beta^* \widehat{\mathbf{C}}_{jj})/n}} \right| \\ & \leq \max_{j \in \widehat{\mathcal{S}}} \left| \frac{|\widehat{\Gamma}_j^{[m^*]} - \Gamma_j^* - \beta(\widehat{\gamma}_j^{[m^*]} - \gamma_j^*)|}{\sqrt{(\widehat{\mathbf{V}}_{jj}^\Gamma + \beta^2 \widehat{\mathbf{V}}_{jj}^\gamma - 2\beta \widehat{\mathbf{C}}_{jj})/n}} - \frac{|\widehat{\Gamma}_j^{[m^*]} - \Gamma_j^* - \beta^*(\widehat{\gamma}_j^{[m^*]} - \gamma_j^*)|}{\sqrt{(\widehat{\mathbf{V}}_{jj}^\Gamma + [\beta^*]^2 \widehat{\mathbf{V}}_{jj}^\gamma - 2\beta^* \widehat{\mathbf{C}}_{jj})/n}} \right| \\ & \leq \max_{j \in \widehat{\mathcal{S}}} \left| \frac{(\beta - \beta^*)(\widehat{\gamma}_j^{[m^*]} - \gamma_j^*)}{\sqrt{\widehat{\mathbf{R}}_{jj}(\beta)/n}} \right| + \max_{j \in \widehat{\mathcal{S}}} \left| \frac{|\widehat{\Gamma}_j^{[m^*]} - \Gamma_j^* - \beta^*(\widehat{\gamma}_j^{[m^*]} - \gamma_j^*)|}{\sqrt{\widehat{\mathbf{R}}_{jj}(\beta)/n}} \left(\sqrt{\frac{\widehat{\mathbf{R}}_{jj}(\beta)}{\widehat{\mathbf{R}}_{jj}(\beta^*)}} - 1 \right) \right| \\ & \leq C \text{err}_n(M, \alpha_0) \left(|\beta - \beta^*| + \left| \frac{\widehat{\mathbf{R}}_{jj}(\beta)}{\widehat{\mathbf{R}}_{jj}(\beta^*)} - 1 \right| \right) \leq 0.1 \cdot C \text{err}_n(M, \alpha_0), \end{aligned}$$

where the third inequality follows from (74) and the last inequality follows from the fact $|\beta - \beta^*| \leq n^{-a}$ and (48), (44), and (49). Together with (75), we have

$$\max_{\beta \in \mathcal{U}^{(a)}} \max_{j \in \widehat{\mathcal{S}}} \frac{|\widehat{\Gamma}_j^{[m^*]} - \Gamma_j^* - \beta(\widehat{\gamma}_j^{[m^*]} - \gamma_j^*)|}{\sqrt{(\widehat{\mathbf{V}}_{jj}^\Gamma + \beta^2 \widehat{\mathbf{V}}_{jj}^\gamma - 2\beta \widehat{\mathbf{C}}_{jj})/n}} \leq C \text{err}_n(M, \alpha_0).$$

Hence, we have established (59).

C.4 Proof of Theorem 2

We apply the definitions of CI^{samp} in (25) and $\text{CI}^{\text{samp}}(\mathcal{M}_0)$ in (26) and establish

$$\liminf_{n \rightarrow \infty} \mathbf{P}(\beta^* \in \text{CI}^{\text{samp}}) \geq \liminf_{n \rightarrow \infty} \mathbf{P}(\beta^* \in \text{CI}^{\text{samp}}(\mathcal{M}_0)).$$

Hence, it is sufficient to control the probability $\mathbf{P}(\beta^* \in \text{CI}^{\text{samp}}(\mathcal{M}_0))$.

C.4.1 The coverage property of $\text{CI}^{\text{samp}}(\mathcal{M}_0)$ in (26)

Recall the definition of $\text{CI}^{\text{samp}}(\mathcal{M}_0)$ in (26),

$$\text{CI}^{\text{samp}}(\mathcal{M}_0) = \left(\min_{m \in \mathcal{M}'} \beta_{\min}^{[m]}(\lambda), \max_{m \in \mathcal{M}'} \beta_{\max}^{[m]}(\lambda) \right),$$

where $\mathcal{M}' = \{m \in \mathcal{M}_0 : [\beta_{\min}^{[m]}(\lambda), \beta_{\max}^{[m]}(\lambda)] \neq \emptyset\}$, with \mathcal{M}_0 defined in (60).

We define the event

$$\mathcal{E}_3 = \left\{ \text{there exists } m^* \in \mathcal{M}_0 \text{ such that } \max_{\beta \in \mathcal{U}(a)} \max_{j \in \widehat{\mathcal{S}}} \frac{|\widehat{\Gamma}_j^{[m^*]} - \Gamma_j^* - \beta(\widehat{\gamma}_j^{[m^*]} - \gamma_j^*)|}{\sqrt{(\widehat{\mathbf{V}}_{jj}^\Gamma + \beta^2 \widehat{\mathbf{V}}_{jj}^\gamma - 2\beta \widehat{\mathbf{C}}_{jj})/n}} \leq \frac{\lambda}{2} \cdot \Phi^{-1} \left(1 - \frac{\alpha}{2|\widehat{\mathcal{S}}|} \right) \right\}.$$

By assuming

$$\frac{1}{2} \lambda \cdot \Phi^{-1} \left(1 - \frac{\alpha}{2|\widehat{\mathcal{S}}|} \right) \geq C_{\text{err}_n}(M, \alpha_0),$$

we apply Proposition 1 and establish that

$$\liminf_{n \rightarrow \infty} \mathbf{P}(\mathcal{E}_3) \geq 1 - \alpha_0. \quad (76)$$

Using a similar decomposition as (13), we have

$$\left(\widehat{\Gamma}_j^{[m]} - \beta \widehat{\gamma}_j^{[m]} \right) - \pi_j^* = \widehat{\Gamma}_j^{[m]} - \Gamma_j^* - \beta(\widehat{\gamma}_j^{[m]} - \gamma_j^*) + (\beta^* - \beta)\gamma_j^* \quad \text{for } 1 \leq m \leq M. \quad (77)$$

We consider two cases

- (a) $\beta^* \in \mathcal{B}$;
- (b) $\beta^* \notin \mathcal{B}$. By the construction of \mathcal{B} , there exists $\beta^L, \beta^U \in \mathcal{B}$ such that $\beta^L \leq \beta^* \leq \beta^U$ and $\beta^U - \beta^L \leq n^{-a}$ for $a > 0.5$.

Case (a). If β is taken as β^* , then

$$\left(\widehat{\Gamma}_j^{[m]} - \beta^* \widehat{\gamma}_j^{[m]} \right) - \pi_j^* = \widehat{\Gamma}_j^{[m]} - \Gamma_j^* - \beta^*(\widehat{\gamma}_j^{[m]} - \gamma_j^*). \quad (78)$$

On the event \mathcal{E}_3 , we show that there exists $1 \leq m^* \leq M$ such that

$$\left| \widehat{\Gamma}_j^{[m^*]} - \beta^* \widehat{\gamma}_j^{[m^*]} \right| \leq \lambda \widehat{\rho}_j(\beta^*) \quad \text{for any } j \in \mathcal{V} \cap \widehat{\mathcal{S}}. \quad (79)$$

The above inequality implies

$$\left| \left\{ j \in \widehat{\mathcal{S}} : \left| \widehat{\Gamma}_j^{[m^*]} - \beta^* \widehat{\gamma}_j^{[m^*]} \right| \leq \lambda \widehat{\rho}_j(\beta^*) \right\} \right| \geq \left| \mathcal{V} \cap \widehat{\mathcal{S}} \right|. \quad (80)$$

Together with the definition in (23) and (53), we show that, on the event $\mathcal{G}_5 \cap \mathcal{E}_3$,

$$\|\widehat{\pi}_{\widehat{\mathcal{S}}}^{[m^*]}(\beta^*, \lambda)\|_0 \leq |\widehat{\mathcal{S}}| - |\mathcal{V} \cap \widehat{\mathcal{S}}| < \frac{|\widehat{\mathcal{S}}|}{2}.$$

Then on the event $\mathcal{G}_5 \cap \mathcal{E}_3$,

$$\beta^* \in \left(\beta_{\min}^{[m^*]}, \beta_{\max}^{[m^*]} \right) \subset \text{CI}^{\text{samp}}(\mathcal{M}_0). \quad (81)$$

Hence the definition of $\text{CI}^{\text{samp}}(\mathcal{M}_0)$ in (26) implies

$$\mathbf{P}(\beta^* \in \text{CI}^{\text{samp}}(\mathcal{M}_0)) \geq \mathbf{P}(\mathcal{G}_5 \cap \mathcal{E}_3).$$

Together with Lemma 3 and (76), we establish $\mathbf{P}(\beta^* \in \text{CI}^{\text{samp}}(\mathcal{M}_0)) \geq 1 - \alpha_0$.

Case (b). It follows from (77) that

$$\widehat{\Gamma}_j^{[m]} - \beta^L \widehat{\gamma}_j^{[m]} - \pi_j^* = \widehat{\Gamma}_j^{[m]} - \Gamma_j^* - \beta^L (\widehat{\gamma}_j^{[m]} - \gamma_j^*) + (\beta^* - \beta^L) \gamma_j^* \quad \text{for } 1 \leq m \leq M. \quad (82)$$

On the event \mathcal{E}_3 , there exists $1 \leq m^* \leq M$ such that, for any $j \in \mathcal{V} \cap \widehat{\mathcal{S}}$,

$$\left| \widehat{\Gamma}_j^{[m^*]} - \Gamma_j^* - \beta^L (\widehat{\gamma}_j^{[m^*]} - \gamma_j^*) \right| \leq \frac{1}{2} \lambda \widehat{\rho}_j(\beta^L). \quad (83)$$

Note that $\lambda \gg n^{1/2-a}$ implies

$$|(\beta^* - \beta^L) \gamma_j^*| \leq n^{-a} |\gamma_j^*| \leq \frac{1}{2} \lambda \widehat{\rho}_j(\beta^L).$$

Combined with (82) and (83), we establish

$$\left| \widehat{\Gamma}_j^{[m^*]} - \beta^L \widehat{\gamma}_j^{[m^*]} \right| \leq \lambda \widehat{\rho}_j(\beta^L) \quad \text{for any } j \in \mathcal{V} \cap \widehat{\mathcal{S}}.$$

This is similar to the result in (79) with replacing β^* by β^L . Then we apply a similar argument as that of (81) and establish that, on the event $\mathcal{G} \cap \mathcal{E}_3$,

$$\beta^L \in \left(\beta_{\min}^{[m^*]}, \beta_{\max}^{[m^*]} \right) \subset \text{CI}^{\text{samp}}(\mathcal{M}_0).$$

Similarly, we can show that, on the event $\mathcal{G} \cap \mathcal{E}_3$, $\beta^U \in \text{CI}^{\text{samp}}(\mathcal{M}_0)$. It follows from the definition of $\text{CI}^{\text{samp}}(\mathcal{M}_0)$ in (26) that on the event $\mathcal{G} \cap \mathcal{E}_3$, $\beta^* \in \text{CI}^{\text{samp}}(\mathcal{M}_0)$. Hence

$$\mathbf{P}(\beta^* \in \text{CI}^{\text{samp}}(\mathcal{M}_0)) \geq \mathbf{P}(\mathcal{G} \cap \mathcal{E}_3).$$

Together with Lemma 3 and (76), we establish $\mathbf{P}(\beta^* \in \text{CI}^{\text{samp}}(\mathcal{M}_0)) \geq 1 - \alpha_0$.

C.4.2 Length of CI^{samp} in (25) and $\text{CI}^{\text{samp}}(\mathcal{M}_0)$ in (26)

Recall the definition of the index sets

$$\mathcal{M}' = \{m \in \mathcal{M}_0 : [\beta_{\min}^{[m]}(\lambda), \beta_{\max}^{[m]}(\lambda)] \neq \emptyset\}, \quad \mathcal{M} = \{1 \leq m \leq M : [\beta_{\min}^{[m]}(\lambda), \beta_{\max}^{[m]}(\lambda)] \neq \emptyset\},$$

with \mathcal{M}_0 defined in (60). We define the following events

$$\begin{aligned} \mathcal{E}_4 &= \left\{ \max_{m \in \mathcal{M}'} \max_{j \in \widehat{\mathcal{S}}} \left(\left| \widehat{\Gamma}_j^{[m]} - \Gamma_j^* \right| + \left| \widehat{\gamma}_j^{[m]} - \gamma_j^* \right| \right) \leq C \frac{1}{\sqrt{n}} \right\}, \\ \mathcal{E}_5 &= \left\{ \max_{m \in \mathcal{M}} \max_{j \in \widehat{\mathcal{S}}} \left(\left| \widehat{\Gamma}_j^{[m]} - \Gamma_j^* \right| + \left| \widehat{\gamma}_j^{[m]} - \gamma_j^* \right| \right) \leq C \frac{\sqrt{\log |\mathcal{M}|}}{\sqrt{n}} \right\}, \end{aligned}$$

where for some positive constant $C > 0$ independent of n . For any $1 \leq m \leq M$, we have

$$\max_{j \in \widehat{\mathcal{S}}} \left| \widehat{\Gamma}_j^{[m]} - \Gamma_j^* \right| \leq \max_{j \in \widehat{\mathcal{S}}} \left| \widehat{\Gamma}_j^{[m]} - \widehat{\Gamma}_j \right| + \max_{j \in \widehat{\mathcal{S}}} \left| \widehat{\Gamma}_j - \Gamma_j^* \right|. \quad (84)$$

On the event \mathcal{E}_2 and the definition of \mathcal{M}_0 , we apply (84) and establish

$$\max_{m \in \mathcal{M}'} \max_{j \in \widehat{\mathcal{S}}} \left| \widehat{\Gamma}_j^{[m]} - \Gamma_j^* \right| \lesssim \max_{j \in \widehat{\mathcal{S}}} \sqrt{\widehat{\mathbf{V}}_{jj}^\Gamma / n} \cdot \Phi^{-1} \left(1 - \frac{\alpha_0}{4|\widehat{\mathcal{S}}|} \right).$$

By a similar argument, we can establish that, on the event $\mathcal{E}_2 \cap \mathcal{G}$,

$$\max_{m \in \mathcal{M}'} \max_{j \in \widehat{\mathcal{S}}} \left| \widehat{\gamma}_j^{[m]} - \gamma_j^* \right| \lesssim \max_{j \in \widehat{\mathcal{S}}} \sqrt{\widehat{\mathbf{V}}_{jj}^\gamma / n} \cdot \Phi^{-1} \left(1 - \frac{\alpha_0}{4|\widehat{\mathcal{S}}|} \right).$$

The above two inequalities imply

$$\liminf_{n \rightarrow \infty} \mathbf{P}(\mathcal{E}_4) \geq \liminf_{n \rightarrow \infty} \mathbf{P}(\mathcal{E}_2 \cap \mathcal{G}) \geq 1 - \alpha_0. \quad (85)$$

We now consider the event \mathcal{E}_5 and the only difference is that

$$\mathbf{P} \left(\max_{m \in \mathcal{M}} \max_{j \in \widehat{\mathcal{S}}} \left| \widehat{\Gamma}_j^{[m]} - \widehat{\Gamma}_j \right| \geq C \max_{j \in \widehat{\mathcal{S}}} \sqrt{\widehat{\mathbf{V}}_{jj}^\Gamma / n} \sqrt{\log |\mathcal{M}|} \right) \leq |\mathcal{M}|^{-c}$$

for some positive constants $c > 0$ and $C > 0$. Hence we establish

$$\liminf_{n \rightarrow \infty} \mathbf{P}(\mathcal{E}_5) \geq 1 - \alpha_0. \quad (86)$$

We now control the length of the sampling interval. For $1 \leq m \leq M$ and $j \in \mathcal{V} \cap \widehat{\mathcal{S}}$, we have

$$\widehat{\Gamma}_j^{[m]} - \beta \widehat{\gamma}_j^{[m]} = \widehat{\Gamma}_j^{[m]} - \Gamma_j^* - \beta(\widehat{\gamma}_j^{[m]} - \gamma_j^*) + (\beta^* - \beta)\gamma_j^*.$$

For β satisfying

$$|\gamma_j^*| \cdot |\beta - \beta^*| \geq \left| \widehat{\Gamma}_j^{[m]} - \Gamma_j^* - \beta(\widehat{\gamma}_j^{[m]} - \gamma_j^*) \right| + \lambda \widehat{\rho}_j(\beta),$$

we have $\widehat{\pi}_j^{[m]}(\beta, \lambda) \neq 0$ for $j \in \mathcal{V} \cap \widehat{\mathcal{S}}$. Consequently, if β satisfies

$$|\beta - \beta^*| \geq \max_{j \in \widehat{\mathcal{S}} \cap \mathcal{V}} \frac{\left| \widehat{\Gamma}_j^{[m]} - \Gamma_j^* - \beta(\widehat{\gamma}_j^{[m]} - \gamma_j^*) \right| + \lambda \widehat{\rho}_j(\beta)}{|\gamma_j^*|}, \quad (87)$$

then we have

$$\|\widehat{\pi}_{\widehat{\mathcal{S}}}^{[m]}(\beta, \lambda)\|_0 \geq |\widehat{\mathcal{S}} \cap \mathcal{V}| > |\widehat{\mathcal{S}}|/2,$$

where the second inequality follows from (53). The above inequality implies that if β satisfies (87), then $\beta \notin [\beta_{\min}^{[m]}, \beta_{\max}^{[m]}]$. That is,

$$\left| \beta_{\min}^{[m]} - \beta^* \right| \leq \max_{j \in \widehat{\mathcal{S}} \cap \mathcal{V}} \frac{\left| \widehat{\Gamma}_j^{[m]} - \Gamma_j^* - \beta_{\min}^{[m]}(\widehat{\gamma}_j^{[m]} - \gamma_j^*) \right| + \lambda \widehat{\rho}_j(\beta_{\min}^{[m]})}{|\gamma_j^*|},$$

and

$$\left| \beta_{\max}^{[m]} - \beta^* \right| \leq \max_{j \in \widehat{\mathcal{S}} \cap \mathcal{V}} \frac{\left| \widehat{\Gamma}_j^{[m]} - \Gamma_j^* - \beta_{\max}^{[m]}(\widehat{\gamma}_j^{[m]} - \gamma_j^*) \right| + \lambda \widehat{\rho}_j(\beta_{\max}^{[m]})}{|\gamma_j^*|}.$$

On the event $\mathcal{G} \cap \mathcal{E}_4$, we apply (51) and establish

$$\left| \beta_{\min}^{[m]} - \beta^* \right| \leq C \max_{j \in \widehat{\mathcal{S}} \cap \mathcal{V}} \frac{(1 + \lambda) \sqrt{1 + (\beta_{\min}^{[m]})^2}}{\sqrt{n} |\gamma_j^*|}, \quad \left| \beta_{\max}^{[m]} - \beta^* \right| \leq C \max_{j \in \widehat{\mathcal{S}} \cap \mathcal{V}} \frac{(1 + \lambda) \sqrt{1 + (\beta_{\max}^{[m]})^2}}{\sqrt{n} |\gamma_j^*|}. \quad (88)$$

The above inequalities are similar to (57) for the searching CI. By a similar argument as that of (58), we establish that $|\beta_{\max}^{[m]}| \leq C$ and $|\beta_{\min}^{[m]}| \leq C$. On the event $\mathcal{E}_4 \cap \mathcal{G}$, we apply (88) and establish

$$\max_{m \in \mathcal{M}'} \max \left\{ \left| \beta_{\max}^{[m]} - \beta^* \right|, \left| \beta_{\min}^{[m]} - \beta^* \right| \right\} \leq \max_{j \in \widehat{\mathcal{S}} \cap \mathcal{V}} \frac{C}{\sqrt{n} |\gamma_j^*|}.$$

Similarly, on the event $\mathcal{E}_5 \cap \mathcal{G}$, we have

$$\max_{m \in \mathcal{M}} \max \left\{ \left| \beta_{\max}^{[m]} - \beta^* \right|, \left| \beta_{\min}^{[m]} - \beta^* \right| \right\} \leq \max_{j \in \widehat{\mathcal{S}} \cap \mathcal{V}} \frac{C \sqrt{\log |\mathcal{M}|}}{\sqrt{n} |\gamma_j^*|}.$$

C.5 Proof of Proposition 2

C.5.1 Finite-sample analysis of the voting method

For $j, k \in \widehat{\mathcal{S}}$, we have the following error decomposition of $\widehat{\beta}^{[j]} = \widehat{\Gamma}_j / \widehat{\gamma}_j$ and $\widehat{\pi}_k^{[j]}$ defined in (31),

$$\widehat{\beta}^{[j]} - \frac{\Gamma_j^*}{\gamma_j^*} = \frac{1}{\gamma_j^*} \widehat{\Omega}_j \cdot \frac{1}{n} W^\top \left(\epsilon - \frac{\Gamma_j^*}{\gamma_j^*} \cdot \delta \right) + \frac{1}{\gamma_j^*} \left(\frac{\widehat{\Gamma}_j}{\widehat{\gamma}_j} - \frac{\Gamma_j^*}{\gamma_j^*} \right) (\gamma_j^* - \widehat{\gamma}_j), \quad (89)$$

and

$$\begin{aligned} \widehat{\pi}_k^{[j]} - \left(\Gamma_k^* - \frac{\Gamma_j^*}{\gamma_j^*} \gamma_k^* \right) &= \left(\widehat{\Gamma}_k - \widehat{\beta}^{[j]} \widehat{\gamma}_k \right) - \left(\Gamma_k^* - \frac{\Gamma_j^*}{\gamma_j^*} \gamma_k^* \right) \\ &= \left(\widehat{\Gamma}_k - \Gamma_k^* \right) - \frac{\Gamma_j^*}{\gamma_j^*} (\widehat{\gamma}_k - \gamma_k^*) - \gamma_k^* \left(\widehat{\beta}^{[j]} - \frac{\Gamma_j^*}{\gamma_j^*} \right) - \left(\widehat{\beta}^{[j]} - \frac{\Gamma_j^*}{\gamma_j^*} \right) (\widehat{\gamma}_k - \gamma_k^*). \end{aligned} \quad (90)$$

Note that

$$\Gamma_k^* - \frac{\Gamma_j^*}{\gamma_j^*} \gamma_k^* = \pi_k^* - \frac{\pi_j^*}{\gamma_j^*} \gamma_k^*. \quad (91)$$

By plugging (91) and (89) into (90), we have the following decomposition of $\widehat{\pi}_k^{[j]} - \pi_k^*$

$$\widehat{\pi}_k^{[j]} - \left(\pi_k^* - \frac{\pi_j^*}{\gamma_j^*} \gamma_k^* \right) = \mathcal{M}_k^{[j]} + \mathcal{A}_k^{[j]}, \quad (92)$$

where

$$\mathcal{M}_k^{[j]} = \left(\widehat{\Omega}_k - \frac{\gamma_k^*}{\gamma_j^*} \widehat{\Omega}_j \right)^\top \frac{1}{n} W^\top \left(\epsilon - \frac{\Gamma_j^*}{\gamma_j^*} \delta \right),$$

and

$$\mathcal{A}_k^{[j]} = -\frac{\gamma_k^*}{\gamma_j^*} \left(\widehat{\beta}^{[j]} - \frac{\Gamma_j^*}{\gamma_j^*} \right) (\gamma_j^* - \widehat{\gamma}_j) - \left(\widehat{\beta}^{[j]} - \frac{\Gamma_j^*}{\gamma_j^*} \right) (\widehat{\gamma}_k - \gamma_k^*). \quad (93)$$

The remaining proof requires the following lemma, whose proof can be found in Section D.3.

Lemma 4 Suppose that Conditions (C1) and (C2) hold, then for any $j, k \in \mathcal{S}^0$, we have

$$\frac{1}{\text{SE}(\widehat{\pi}_k^{[j]})} \left(\widehat{\Omega}_k - \frac{\gamma_k^*}{\gamma_j^*} \widehat{\Omega}_j \right)^\top \frac{1}{n} W^\top \left(\epsilon - \frac{\Gamma_j^*}{\gamma_j^*} \delta \right) \xrightarrow{d} N(0, 1) \quad (94)$$

where \mathcal{S}^0 is defined in (46) and

$$\text{SE}(\widehat{\pi}_k^{[j]}) = |\gamma_k^*| \cdot \mathbf{T}_{j,k}^0 = \sqrt{\frac{1}{n} \left(\mathbf{R}_{k,k}^{[j]} + \frac{[\gamma_k^*]^2}{[\gamma_j^*]^2} \mathbf{R}_{j,j}^{[j]} - \frac{2\gamma_k^*}{\gamma_j^*} \mathbf{R}_{j,k}^{[j]} \right)}$$

with $\mathbf{T}_{j,k}^0$ defined in (9). In addition, we have

$$\text{SE}(\widehat{\pi}_k^{[j]}) \geq \sqrt{\frac{c_1 \cdot \lambda_{\min}(\Sigma^{-1})}{n}} \sqrt{1 + \left(\frac{\Gamma_j^*}{\gamma_j^*} \right)^2} \cdot \sqrt{1 + \left(\frac{\gamma_k^*}{\gamma_j^*} \right)^2}. \quad (95)$$

On the event \mathcal{G} , $\widehat{\text{SE}}(\widehat{\pi}_k^{[j]})$ defined in (32) satisfies

$$\left| \left(\frac{\widehat{\text{SE}}(\widehat{\pi}_k^{[j]})}{\text{SE}(\widehat{\pi}_k^{[j]})} \right)^2 - 1 \right| \lesssim \frac{1}{(\log n)^{1/4}}. \quad (96)$$

Define the event

$$\mathcal{F} = \cap_{j,k \in \widehat{\mathcal{S}}} \mathcal{F}_k^{[j]} \quad \text{with} \quad \mathcal{F}_k^{[j]} = \left\{ \left| \mathcal{M}_k^{[j]} \right| \leq 0.9 \sqrt{\log n} \cdot \widehat{\text{SE}}(\widehat{\pi}_k^{[j]}) \right\}. \quad (97)$$

By (94) and (96), we apply the union bound and establish

$$\liminf_{n \rightarrow \infty} \mathbf{P}(\mathcal{F}) = 1.$$

On the event \mathcal{G} , we apply the expression (93) and establish

$$\left| \mathcal{A}_k^{[j]} \right| \lesssim \left(1 + \left| \frac{\gamma_k^*}{\gamma_j^*} \right| \right) \left(1 + \left| \frac{\Gamma_j^*}{\gamma_j^*} \right| \right) \frac{1}{(\log n)^{1/4}} \cdot \frac{(\log n)^{1/4}}{\sqrt{n}}.$$

On the event \mathcal{G} , we further apply (95) and (96) and establish that

$$\max_{j,k \in \widehat{\mathcal{S}}} \left| \mathcal{A}_k^{[j]} \right| \leq 0.05 \sqrt{\log n} \widehat{\text{SE}}(\widehat{\pi}_k^{[j]}), \quad (98)$$

for a sufficiently large n .

We shall consider two cases in the following.

Case (a). When

$$\left| \frac{\pi_k^*}{\gamma_k^*} - \frac{\pi_j^*}{\gamma_j^*} \right| = 0,$$

we simplify (92) as

$$\widehat{\pi}_k^{[j]} = \mathcal{M}_k^{[j]} + \mathcal{A}_k^{[j]}. \quad (99)$$

By (98), on the event $\mathcal{G} \cap \mathcal{F}$,

$$\max_{j,k \in \widehat{\mathcal{S}}} \left| \mathcal{M}_k^{[j]} + \mathcal{A}_k^{[j]} \right| \leq 0.95 \sqrt{\log n} \cdot \widehat{\text{SE}}(\widehat{\pi}_k^{[j]}). \quad (100)$$

Similarly, we establish on the event $\mathcal{G} \cap \mathcal{F}$,

$$\max_{j,k \in \widehat{\mathcal{S}}} \left| \mathcal{M}_j^{[k]} + \mathcal{A}_j^{[k]} \right| \leq 0.95 \sqrt{\log n} \cdot \widehat{\text{SE}}(\widehat{\pi}_j^{[k]}). \quad (101)$$

By the definition in (33), we have

$$\liminf_{n \rightarrow \infty} \mathbf{P} \left(\widehat{\Pi}_{k,j} = \widehat{\Pi}_{j,k} = 1 \right) \geq \liminf_{n \rightarrow \infty} \mathbf{P} (\mathcal{F} \cap \mathcal{G}) = 1.$$

Case (b). By (92), the event $\{\widehat{\Pi}_{k,j} = \widehat{\Pi}_{j,k} = 0\}$ is equivalent to that at least one of the following two events happens

$$\left| \pi_k^* - \frac{\pi_j^*}{\gamma_j^*} \gamma_k^* + \mathcal{M}_k^{[j]} + \mathcal{A}_k^{[j]} \right| > \sqrt{\log n} \cdot \widehat{\text{SE}}(\widehat{\pi}_k^{[j]}); \quad (102)$$

$$\left| \pi_j^* - \frac{\pi_k^*}{\gamma_k^*} \gamma_j^* + \mathcal{M}_j^{[k]} + \mathcal{A}_j^{[k]} \right| > \sqrt{\log n} \cdot \widehat{\text{SE}}(\widehat{\pi}_j^{[k]}). \quad (103)$$

By the upper bound in (100), on the event $\mathcal{G} \cap \mathcal{F}$, the event in (102) happens if

$$\left| \frac{\pi_k^*}{\gamma_k^*} - \frac{\pi_j^*}{\gamma_j^*} \right| \geq 1.95 \sqrt{\log n} \cdot \frac{\widehat{\text{SE}}(\widehat{\pi}_k^{[j]})}{\gamma_k^*};$$

the event (103) happens if

$$\left| \frac{\pi_j^*}{\gamma_j^*} - \frac{\pi_k^*}{\gamma_k^*} \right| \geq 1.95 \sqrt{\log n} \cdot \frac{\widehat{\text{SE}}(\widehat{\pi}_j^{[k]})}{\gamma_j^*}.$$

That is, the event $\{\widehat{\Pi}_{k,j} = \widehat{\Pi}_{j,k} = 0\}$ happens if

$$\left| \frac{\pi_k^*}{\gamma_k^*} - \frac{\pi_j^*}{\gamma_j^*} \right| \geq 1.95 \sqrt{\log n} \cdot \min \left\{ \frac{\widehat{\text{SE}}(\widehat{\pi}_j^{[k]})}{\gamma_j^*}, \frac{\widehat{\text{SE}}(\widehat{\pi}_k^{[j]})}{\gamma_k^*} \right\}.$$

Together with (96), we obtain that, on the event $\mathcal{G} \cap \mathcal{F}$, the event $\{\widehat{\Pi}_{k,j} = \widehat{\Pi}_{j,k} = 0\}$ happens if

$$\left| \frac{\pi_k^*}{\gamma_k^*} - \frac{\pi_j^*}{\gamma_j^*} \right| \geq 2 \sqrt{\log n} \cdot \min \left\{ \frac{\text{SE}(\widehat{\pi}_j^{[k]})}{\gamma_j^*}, \frac{\text{SE}(\widehat{\pi}_k^{[j]})}{\gamma_k^*} \right\}.$$

Hence, we establish

$$\liminf_{n \rightarrow \infty} \mathbf{P} \left(\widehat{\Pi}_{k,j} = \widehat{\Pi}_{j,k} = 0 \right) \geq \liminf_{n \rightarrow \infty} \mathbf{P} (\mathcal{G} \cap \mathcal{F}) = 1.$$

C.5.2 Analysis of $\widehat{\mathcal{V}}^{\text{TSHT}}$ in (34)

We apply the definition of $\widehat{\mathcal{V}}^{\text{TSHT}}$ in (34) and write $\widehat{\mathcal{V}}^{\text{TSHT}}$ as $\widehat{\mathcal{V}}$ in the following proof. By the construction, we have $\widehat{\mathcal{W}} \subset \widehat{\mathcal{V}}$. Throughout the proof, we condition on the event $\mathcal{G} \cap \mathcal{F}$, which has asymptotic probability 1. Then we can directly apply case (a) and case (b) of Proposition 2. In particular, we use the following results: for any $l \in \widehat{\mathcal{V}}$, there exists $j \in \widehat{\mathcal{W}}$ such that

$$\left| \frac{\pi_l^*}{\gamma_l^*} - \frac{\pi_j^*}{\gamma_j^*} \right| \leq 2\text{sep}(n). \quad (104)$$

Similarly, for any $j \in \widehat{\mathcal{W}}$, there exists $l \in \widehat{\mathcal{V}}$ such that (104) holds. In the following, we apply (104) and consider two cases.

Case 1. We first consider the case where $\widehat{\mathcal{W}} \subset \mathcal{V}$. It follows from (104) that

$$\widehat{\mathcal{V}} \subset \mathcal{I}(0, 2\text{sep}(n)). \quad (105)$$

For $k \in \mathcal{V} \cap \widehat{\mathcal{S}}$ and $j \in \widehat{\mathcal{W}} \subset \mathcal{V}$, we have $\frac{\pi_k^*}{\gamma_k^*} = \frac{\pi_j^*}{\gamma_j^*} = 0$. Case (a) of Proposition 2 implies

$$\widehat{\Pi}_{k,j} = \widehat{\Pi}_{j,k} = 1.$$

Then we apply the definition of $\widehat{\mathcal{V}} = \widehat{\mathcal{V}}^{\text{TSHT}}$ in (34) and establish that $k \in \widehat{\mathcal{V}}$, which implies

$$\mathcal{V} \cap \widehat{\mathcal{S}} \subset \widehat{\mathcal{V}}. \quad (106)$$

Since $\mathcal{V} \cap \mathcal{S}_{\text{str}} \subset \mathcal{V} \cap \widehat{\mathcal{S}}$, we combine (106) and (105) and establish

$$\mathcal{V} \cap \mathcal{S}_{\text{str}} \subset \widehat{\mathcal{V}} \subset \mathcal{I}(0, 2\text{sep}(n)). \quad (107)$$

Case 2. We then consider the case where $\widehat{\mathcal{W}} \not\subset \mathcal{V}$. Then there exists j such that $j \in \widehat{\mathcal{W}}$ but $j \notin \mathcal{V}$. Let $\text{supp}(\widehat{\Pi}_j)$ denote the support of $\widehat{\Pi}_j$.

We define $l \in \mathcal{V} \cap \widehat{\mathcal{S}}$. Case (a) of Proposition 2 implies

$$\mathcal{V} \cap \widehat{\mathcal{S}} \subset \text{supp}(\widehat{\Pi}_l). \quad (108)$$

Case (b) of Proposition 2 implies

$$\text{supp}(\widehat{\Pi}_j) \subset \mathcal{I}\left(\frac{\pi_j^*}{\gamma_j^*}, 2\text{sep}(n)\right) \cap \widehat{\mathcal{S}}. \quad (109)$$

In the following, we show by contradiction that $\text{supp}(\widehat{\Pi}_j) \cap \mathcal{V} \neq \emptyset$. Assume that

$$\text{supp}(\widehat{\Pi}_j) \cap \mathcal{V} = \emptyset. \quad (110)$$

The definition of $\widehat{\mathcal{W}}$ implies $|\text{supp}(\widehat{\Pi}_l)| \leq |\text{supp}(\widehat{\Pi}_j)|$. We apply (108) and establish

$$|\mathcal{V} \cap \widehat{\mathcal{S}}| \leq |\text{supp}(\widehat{\Pi}_j)| = |\text{supp}(\widehat{\Pi}_j) \setminus \mathcal{V}| \leq \left| \mathcal{I}\left(\frac{\pi_j^*}{\gamma_j^*}, 2\text{sep}(n)\right) \setminus \mathcal{V} \right|,$$

where the first equality follows from the assumption (110) and the last inequality follows from (109). The above inequality implies that

$$|\mathcal{V} \cap \mathcal{S}_{\text{str}}| \leq |\mathcal{V} \cap \widehat{\mathcal{S}}| \leq \left| \mathcal{I} \left(\frac{\pi_j^*}{\gamma_j^*}, 2\text{sep}(n) \right) \setminus \mathcal{V} \right|,$$

which contradicts the finite-sample plurality rule and hence the conjecture (110) does not hold. That is, for any $j \in \widehat{\mathcal{W}}$,

$$\text{there exists } k \in \text{supp}(\widehat{\Pi}_j) \cap \mathcal{V}. \quad (111)$$

Together with case (b) of Proposition 2, we establish

$$|\pi_j^*/\gamma_j^*| \leq \text{sep}(n) \quad \text{for } j \in \widehat{\mathcal{W}}.$$

Combined with (104), we establish that

$$\widehat{\mathcal{V}} \subset \mathcal{I}(0, 3\text{sep}(n)). \quad (112)$$

For any $k \in \mathcal{V} \cap \widehat{\mathcal{S}}$, we have $\mathcal{V} \cap \widehat{\mathcal{S}} \subset \text{supp}(\widehat{\Pi}_k)$. By combining this with (111) and the definition of $\widehat{\mathcal{V}} = \widehat{\mathcal{V}}^{\text{TSHT}}$ in (34), we establish

$$\mathcal{V} \cap \widehat{\mathcal{S}} \subset \widehat{\mathcal{V}}. \quad (113)$$

Since $\mathcal{V} \cap \mathcal{S}_{\text{str}} \subset \mathcal{V} \cap \widehat{\mathcal{S}}$, we combine (113) and (112) and establish

$$\mathcal{V} \cap \mathcal{S}_{\text{str}} \subset \widehat{\mathcal{V}} \subset \mathcal{I}(0, 3\text{sep}(n)). \quad (114)$$

We combine (107) and (114) and establish that $\widehat{\mathcal{V}} = \widehat{\mathcal{V}}^{\text{TSHT}}$ defined in (34) satisfies (30).

C.6 Proofs of Theorem 3

The proof of the searching interval follows from that of Theorem 1, and Proposition 2. Note that, on the event \mathcal{G} , we have $\mathcal{V} \cap \mathcal{S}_{\text{str}} \subset \widehat{\mathcal{V}} \subset \mathcal{I}(0, 3\text{sep}(n))$. By the finite sample plurality rule (Condition 4), $\mathcal{V} \cap \mathcal{S}_{\text{str}}$ is the majority of the set $\mathcal{I}(0, 3\text{sep}(n))$ and also the majority of $\widehat{\mathcal{V}}$. We then apply the same argument for Theorem 1 by replacing $\widehat{\mathcal{S}}$ with $\widehat{\mathcal{V}}$ and \mathcal{S} with $\mathcal{I}(0, 3\text{sep}(n))$.

The proof of the sampling interval follows from that of Theorem 2 and Proposition 2. Note that, on the event \mathcal{G} , we have $\mathcal{V} \cap \mathcal{S}_{\text{str}} \subset \widehat{\mathcal{V}} \subset \mathcal{I}(0, 3\text{sep}(n))$. By the finite sample plurality rule (Condition 4), $\mathcal{V} \cap \mathcal{S}_{\text{str}}$ is the majority of the set $\mathcal{I}(0, 3\text{sep}(n))$ and also the majority of $\widehat{\mathcal{V}}$. We then apply the same argument for Theorem 2 by replacing $\widehat{\mathcal{S}}$ with $\widehat{\mathcal{V}}$ and \mathcal{S} with $\mathcal{I}(0, 3\text{sep}(n))$.

D Proofs of Lemmas

D.1 Proof of Lemma 3

Control of \mathcal{G}_1 . We present the proof of controlling $\left\| \frac{1}{n} W^\top \epsilon \right\|_\infty$ in the following. The proof of $\left\| \frac{1}{n} W^\top \delta \right\|_\infty$ is similar to that of $\left\| \frac{1}{n} W^\top \epsilon \right\|_\infty$. Note that $\mathbf{E} W_{ij} \delta_i = 0$ for any $1 \leq i \leq n$ and

$1 \leq j \leq p$ and $W_{ij}\delta_i$ is Sub-exponential random variable. Since $(W^\top \delta)_j = \sum_{i=1}^n W_{ij}\delta_i$, we apply Proposition 5.16 of [?] with the corresponding $t = C\sqrt{n\sqrt{\log n}}$ and establish

$$\mathbf{P}\left(\left|\sum_{i=1}^n W_{ij}\delta_i\right| \leq C\sqrt{n\sqrt{\log n}}\right) \geq 1 - \exp(-c'\sqrt{\log n}),$$

where C and c' are positive constants independent of n . For the fixed p setting, we apply the union bound and establish that

$$\begin{aligned} \mathbf{P}\left(\|W^\top \delta\|_\infty = \max_{1 \leq j \leq p_z} \left|\sum_{i=1}^n W_{ij}\delta_i\right| \leq C\sqrt{n\sqrt{\log n}}\right) &\geq 1 - p_z \exp(-c'\sqrt{\log n}) \\ &\geq 1 - \exp(-c\sqrt{\log n}), \end{aligned} \quad (115)$$

for some positive constant $c > 0$.

Control of \mathcal{G}_3 . Since $\{W_i\}_{1 \leq i \leq n}$ are i.i.d Sub-gaussian random vectors and the dimension p is fixed, then we apply equation (5.25) in [?] and establish the following concentration results for $\widehat{\Sigma} - \Sigma$: with probability larger than $1 - n^{-c}$,

$$\|\widehat{\Sigma} - \Sigma\|_2 \leq C\sqrt{\log n/n}, \quad (116)$$

where c and C are positive constants independent of n . As a consequence, we have

$$|\lambda_{\min}(\widehat{\Sigma}) - \lambda_{\min}(\Sigma)| \leq \|\widehat{\Sigma} - \Sigma\|_2 \leq C\sqrt{\log n/n}. \quad (117)$$

Since $\widehat{\Sigma}^{-1} - \Sigma^{-1} = \widehat{\Sigma}^{-1}(\Sigma - \widehat{\Sigma})\Sigma^{-1}$, we have

$$\|\widehat{\Sigma}^{-1} - \Sigma^{-1}\|_2 \leq \frac{1}{\lambda_{\min}(\widehat{\Sigma}) \cdot \lambda_{\min}(\Sigma)} \|\Sigma - \widehat{\Sigma}\|_2 \leq \frac{C\sqrt{\log n/n}}{(\lambda_{\min}(\Sigma) - C\sqrt{\log n/n}) \cdot \lambda_{\min}(\Sigma)}.$$

Then we have

$$\mathbf{P}(\mathcal{G}_3) \geq 1 - n^{-c}. \quad (118)$$

Control of \mathcal{G}_2 . We shall focus on the analysis of $\widehat{\gamma}_j - \gamma_j^*$ and the analysis of $\widehat{\Gamma}_j - \Gamma_j^*$ is similar. We apply the expression (40) and establish

$$\frac{\widehat{\gamma}_j - \gamma_j^*}{\sqrt{\mathbf{V}_{jj}^\gamma/n}} = \frac{\widehat{\Omega}_{j \cdot}^\top \frac{1}{n} W^\top \delta}{\sqrt{\mathbf{V}_{jj}^\gamma/n}} = \frac{\Omega_{j \cdot}^\top \frac{1}{n} W^\top \delta}{\sqrt{\mathbf{V}_{jj}^\gamma/n}} + \frac{(\widehat{\Omega}_{j \cdot} - \Omega_{j \cdot})^\top \frac{1}{n} W^\top \delta}{\sqrt{\mathbf{V}_{jj}^\gamma/n}}. \quad (119)$$

By Lemma 1, we have

$$\Omega_{j \cdot}^\top \frac{1}{n} W^\top \delta / \sqrt{\mathbf{V}_{jj}^\gamma/n} \xrightarrow{d} N(0, 1).$$

On the event $\mathcal{G}_1 \cap \mathcal{G}_3$, we apply the decomposition (117) together with (41), (115) and (119) and establish

$$\mathbf{P}\left(\max_{1 \leq j \leq p_z} |\widehat{\gamma}_j - \gamma_j^*| / \sqrt{\mathbf{V}_{jj}^\gamma/n} \leq C(\log n)^{1/4}\right) \geq 1 - \exp(-c\sqrt{\log n}).$$

We can apply a similar argument to control $\widehat{\Gamma}_j - \Gamma_j^*$ and then establish

$$\mathbf{P}(\mathcal{G}_2) \geq 1 - \exp(-c\sqrt{\log n}). \quad (120)$$

By a similar argument, for the fixed p setting, we can establish

$$\mathbf{P}\left(\|\Psi^* - \widehat{\Psi}\|_1 + \|\psi^* - \widehat{\psi}\|_1 \leq C \frac{(\log n)^{1/4}}{\sqrt{n}}\right) \geq 1 - \exp(-c\sqrt{\log n}). \quad (121)$$

Control of \mathcal{G}_4 . We shall control the difference $\|\widehat{\mathbf{C}} - \mathbf{C}\|_2$ and the upper bounds for $\|\widehat{\mathbf{V}}^\Gamma - \mathbf{V}^\Gamma\|_2$ and $\|\widehat{\mathbf{V}}^\gamma - \mathbf{V}^\gamma\|_2$ can be established in a similar way. Define $A = \mathbf{E}\epsilon_i \delta_i W_i W_i^\top$ and $\widehat{A} = \frac{1}{n} \sum_{i=1}^n \widehat{\epsilon}_i \widehat{\delta}_i W_i W_i^\top$. We first control $\|\widehat{A} - A\|_2$. Note that $\widehat{\epsilon}_i = \epsilon_i + Z_i^\top(\Gamma^* - \widehat{\Gamma}) + X_i^\top(\Psi^* - \widehat{\Psi})$ and $\widehat{\delta}_i = \delta_i + Z_i^\top(\gamma^* - \widehat{\gamma}) + X_i^\top(\psi^* - \widehat{\psi})$.

For $u \in \mathbb{R}^p$, we define $U_i = u^\top W_i$ and decompose the error $u^\top(\widehat{A} - A)u$ as

$$\begin{aligned} & \frac{1}{n} \sum_{i=1}^n \epsilon_i \delta_i U_i^2 - \mathbf{E}\epsilon_i \delta_i U_i^2 + \frac{1}{n} \sum_{i=1}^n \epsilon_i \left(Z_i^\top(\gamma^* - \widehat{\gamma}) + X_i^\top(\psi^* - \widehat{\psi}) \right) U_i^2 \\ & + \frac{1}{n} \sum_{i=1}^n \delta_i \left(Z_i^\top(\Gamma^* - \widehat{\Gamma}) + X_i^\top(\Psi^* - \widehat{\Psi}) \right) U_i^2 \\ & + \frac{1}{n} \sum_{i=1}^n \left(Z_i^\top(\gamma^* - \widehat{\gamma}) + X_i^\top(\psi^* - \widehat{\psi}) \right) \left(Z_i^\top(\Gamma^* - \widehat{\Gamma}) + X_i^\top(\Psi^* - \widehat{\Psi}) \right) U_i^2. \end{aligned}$$

Since ϵ_i, δ_i and Z_i and X_i are Sub-gaussian random variables, then with probability larger than $1 - n^{-c}$, $\max_{1 \leq i \leq n} \max\{|\epsilon_i|, |\delta_i|, \|Z_i\|_\infty, \|X_i\|_\infty\} \leq C\sqrt{\log n}$ where $C > 0$ and $c > 0$ are positive constants independent of n . Together with (121), we establish that, with probability larger than $1 - \exp(-c\sqrt{\log n})$, we have

$$\left| u^\top(\widehat{A} - A)u - \left(\frac{1}{n} \sum_{i=1}^n \epsilon_i \delta_i U_i^2 - \mathbf{E}\epsilon_i \delta_i U_i^2 \right) \right| \leq C \frac{(\log n)^{3/2}}{\sqrt{n}} \cdot \frac{1}{n} \sum_{i=1}^n U_i^2 \leq C \frac{(\log n)^{3/2}}{\sqrt{n}}, \quad (122)$$

where the last inequality follows from (116).

We define $\mathcal{H}_i = \{\max\{|\epsilon_i|, |\delta_i|\} \leq 2\sqrt{\log n}\}$ and we have $\mathbf{P}(\cap_{i=1}^n \mathcal{H}_i) \geq 1 - Cn^{-1}$ for some positive constant $C > 0$. We decompose $\frac{1}{n} \sum_{i=1}^n \epsilon_i \delta_i U_i^2 - \mathbf{E}\epsilon_i \delta_i U_i^2$ as

$$\frac{1}{n} \sum_{i=1}^n \epsilon_i \delta_i U_i^2 \cdot \mathcal{H}_i - \mathbf{E}\epsilon_i \delta_i U_i^2 \cdot \mathcal{H}_i + \frac{1}{n} \sum_{i=1}^n \epsilon_i \delta_i U_i^2 \cdot \mathcal{H}_i^c - \mathbf{E}\epsilon_i \delta_i U_i^2 \cdot \mathcal{H}_i^c.$$

Since ϵ_i and δ_i are sub-gaussian, $\mathbf{P}(\frac{1}{n} \sum_{i=1}^n \epsilon_i \delta_i U_i^2 \cdot \mathcal{H}_i^c \neq 0) \leq Cn^{-1}$, for some positive constant $C > 0$. Note that $\mathbf{E}\epsilon_i \delta_i U_i^2 \cdot \mathcal{H}_i^c \leq \sqrt{\mathbf{E}\epsilon_i^2 \delta_i^2 U_i^4} \sqrt{\mathbf{P}(\mathcal{H}_i^c)} \leq Cn^{-1}$ for some positive constant $C > 0$. Since $\frac{1}{4\log n} |\epsilon_i \delta_i| \cdot \mathcal{H}_i \leq 1$, we generalize (5.25) in [?] and establish

$$\mathbf{P}\left(\max_{u \in \mathbb{R}^p, \|u\|_2=1} \frac{1}{4\log n} \left(\frac{1}{n} \sum_{i=1}^n \epsilon_i \delta_i U_i^2 \cdot \mathcal{H}_i - \mathbf{E}\epsilon_i \delta_i U_i^2 \cdot \mathcal{H}_i \right) \leq C \sqrt{\frac{p + \log n}{n}}\right) \geq 1 - n^{-c},$$

for some positive constant $c > 0$. Combined with (122), we have

$$\mathbf{P} \left(\|\widehat{\mathbf{C}} - \mathbf{C}\|_2 \leq C \frac{(\log n)^{3/2}}{\sqrt{n}} \right) \geq 1 - \exp(-c\sqrt{\log n}), \quad (123)$$

for some positive constants $C > 0$ and $c > 0$. Note that

$$\begin{aligned} \|\widehat{\mathbf{C}} - \mathbf{C}\|_2 &\leq \left\| \widehat{\Sigma}^{-1} \widehat{A} \widehat{\Sigma}^{-1} - \Sigma^{-1} A \Sigma^{-1} \right\|_2 \\ &\leq \|\widehat{\Sigma}^{-1} - \Sigma^{-1}\|_2 \|\widehat{A}\|_2 \|\widehat{\Sigma}^{-1}\|_2 + \|\Sigma^{-1}\|_2 \|\widehat{A} - A\|_2 \|\widehat{\Sigma}^{-1}\|_2 + \|\Sigma^{-1}\|_2 \|A\|_2 \|\widehat{\Sigma}^{-1} - \Sigma^{-1}\|_2. \end{aligned}$$

On the event \mathcal{G}_3 , we apply (123) and (42) and establish

$$\mathbf{P}(\mathcal{G}_3) \geq 1 - \exp(-c\sqrt{\log n}). \quad (124)$$

Control of \mathcal{G}_5 . Recall that $\widehat{\mathcal{S}}$ is defined in (11) and \mathcal{S}_{str} is defined in (12). Then for $j \in \mathcal{S}_{\text{str}}$, on the event $\mathcal{G}_2 \cap \mathcal{G}_4$, if $\sqrt{\log n} > C(\log n)^{1/4}$, we have

$$|\widehat{\gamma}_j| \geq \sqrt{\log n} \cdot \sqrt{\widehat{\mathbf{V}}_{jj}^\gamma/n},$$

that is $\mathcal{S}_{\text{str}} \subset \widehat{\mathcal{S}}$. Furthermore, for $j \in \widehat{\mathcal{S}}$, on the event $\mathcal{G}_2 \cap \mathcal{G}_4$, we have

$$|\gamma_j^*| \geq (\sqrt{\log n} - C(\log n)^{1/4}) \cdot \sqrt{\widehat{\mathbf{V}}_{jj}^\gamma/n}$$

that is $\widehat{\mathcal{S}} \subset \mathcal{S}^0$. Hence, we have

$$\mathbf{P}(\mathcal{G}_4 \cap \mathcal{G}_5) \geq \mathbf{P}(\mathcal{G}_2 \cap \mathcal{G}_4 \cap \mathcal{G}_3). \quad (125)$$

Control of events \mathcal{G}_6 and \mathcal{G}_7 . On the event $\mathcal{G}_2 \cap \mathcal{G}_4 \cap \mathcal{G}_5$, for $j \in \mathcal{S}^0$

$$\left| \frac{\widehat{\gamma}_j}{\gamma_j^*} - 1 \right| \leq \frac{C(\log n)^{1/4} \sqrt{\mathbf{V}_{jj}^\gamma/n}}{(\sqrt{\log n} - C(\log n)^{1/4}) \sqrt{\widehat{\mathbf{V}}_{jj}^\gamma/n}} \lesssim \frac{1}{(\log n)^{1/4}},$$

and hence

$$\max_{j \in \mathcal{S}^0} \left| \frac{\gamma_j^*}{\widehat{\gamma}_j} - 1 \right| \lesssim \frac{1}{(\log n)^{1/4}}. \quad (126)$$

By the decomposition

$$\begin{aligned} (\widehat{\gamma}_k/\widehat{\gamma}_j - \gamma_k^*/\gamma_j^*)\gamma_j^* &= (\widehat{\gamma}_k - \gamma_k^*) \frac{\gamma_j^*}{\widehat{\gamma}_j} + \gamma_k^* \left(\frac{\gamma_j^*}{\widehat{\gamma}_j} - 1 \right) \\ &= \left(\frac{\widehat{\gamma}_k}{\gamma_k^*} - 1 \right) \cdot \gamma_k^* \cdot \frac{\gamma_j^*}{\widehat{\gamma}_j} + \gamma_k^* \left(\frac{\gamma_j^*}{\widehat{\gamma}_j} - 1 \right), \end{aligned}$$

we apply (126) and establish

$$\left| (\widehat{\gamma}_k/\widehat{\gamma}_j - \gamma_k^*/\gamma_j^*)\gamma_j^* \right| \lesssim \frac{|\gamma_k^*|}{(\log n)^{1/4}} \quad \text{and} \quad \left| \frac{\widehat{\gamma}_k}{\widehat{\gamma}_j} - \frac{\gamma_k^*}{\gamma_j^*} \right| \lesssim \left| \frac{\gamma_k^*}{\gamma_j^*} \right| \cdot \frac{1}{(\log n)^{1/4}}.$$

That is, the event \mathcal{G}_6 holds and

$$\mathbf{P}(\mathcal{G}_6) \geq \mathbf{P}(\mathcal{G}_2 \cap \mathcal{G}_4 \cap \mathcal{G}_5). \quad (127)$$

By the decomposition

$$(\widehat{\Gamma}_j/\widehat{\gamma}_j - \Gamma_j^*/\gamma_j^*)\gamma_j^* = (\widehat{\Gamma}_j - \Gamma_j^*)\frac{\gamma_j^*}{\widehat{\gamma}_j} + \Gamma_j^* \left(\frac{\gamma_j^*}{\widehat{\gamma}_j} - 1 \right),$$

we apply (126) and establish on the event \mathcal{G}_2 that

$$\left| (\widehat{\Gamma}_j/\widehat{\gamma}_j - \Gamma_j^*/\gamma_j^*)\gamma_j^* \right| \leq C\sqrt{\mathbf{V}_{jj}^\Gamma/n}(\log n)^{1/4} + |\Gamma_j^*| \frac{1}{(\log n)^{1/4}}.$$

Hence, for $j \in \mathcal{S}^0$, we have

$$\left| \widehat{\Gamma}_j/\widehat{\gamma}_j - \Gamma_j^*/\gamma_j^* \right| \leq C(1 + |\Gamma_j^*/\gamma_j^*|) \frac{1}{(\log n)^{1/4}}.$$

That is, the event \mathcal{G}_7 holds and

$$\mathbf{P}(\mathcal{G}_7) \geq \mathbf{P}(\mathcal{G}_2 \cap \mathcal{G}_4 \cap \mathcal{G}_5). \quad (128)$$

We establish the lemma by combining (118), (120), (124), (125) and (128).

D.2 Proof of Lemma 2

Recall the definitions of $\widehat{\mathbf{R}}(\beta)$ and $\mathbf{R}(\beta)$ in (43). Note that

$$\frac{\widehat{\Gamma}_j - \Gamma_j^* - \beta^*(\widehat{\gamma}_j - \gamma_j^*)}{\sqrt{\mathbf{R}_{jj}(\beta^*)/n}} \xrightarrow{d} N(0, 1).$$

Together with (44) and (49), we establish

$$\frac{\widehat{\Gamma}_j - \Gamma_j^* - \beta^*(\widehat{\gamma}_j - \gamma_j^*)}{\sqrt{\widehat{\mathbf{R}}_{jj}(\beta^*)/n}} \xrightarrow{d} N(0, 1). \quad (129)$$

For any $\beta \in \mathcal{U}(a)$, we have

$$\begin{aligned} & \frac{\widehat{\Gamma}_j - \Gamma_j^* - \beta(\widehat{\gamma}_j - \gamma_j^*)}{\sqrt{\widehat{\mathbf{R}}_{jj}(\beta)/n}} - \frac{\widehat{\Gamma}_j - \Gamma_j^* - \beta^*(\widehat{\gamma}_j - \gamma_j^*)}{\sqrt{\mathbf{R}_{jj}(\beta^*)/n}} \\ &= \frac{\widehat{\Gamma}_j - \Gamma_j^* - \beta^*(\widehat{\gamma}_j - \gamma_j^*)}{\sqrt{\widehat{\mathbf{R}}_{jj}(\beta^*)/n}} \cdot \left(1 - \frac{\sqrt{\widehat{\mathbf{R}}_{jj}(\beta)/n}}{\sqrt{\mathbf{R}_{jj}(\beta^*)/n}} \right) - \frac{(\beta - \beta^*)(\widehat{\gamma}_j - \gamma_j^*)}{\sqrt{\widehat{\mathbf{R}}_{jj}(\beta)/n}}. \end{aligned} \quad (130)$$

By the triangle inequality, we further establish

$$\begin{aligned} & \left| \max_{\beta \in \mathcal{U}_a} \frac{|\widehat{\Gamma}_j - \Gamma_j^* - \beta(\widehat{\gamma}_j - \gamma_j^*)|}{\sqrt{\widehat{\mathbf{R}}_{jj}(\beta)/n}} - \frac{|\widehat{\Gamma}_j - \Gamma_j^* - \beta^*(\widehat{\gamma}_j - \gamma_j^*)|}{\sqrt{\mathbf{R}_{jj}(\beta^*)/n}} \right| \\ & \leq \max_{\beta \in \mathcal{U}_a} \left| \frac{|\widehat{\Gamma}_j - \Gamma_j^* - \beta^*(\widehat{\gamma}_j - \gamma_j^*)|}{\sqrt{\widehat{\mathbf{R}}_{jj}(\beta^*)/n}} \cdot \left(1 - \frac{\sqrt{\widehat{\mathbf{R}}_{jj}(\beta)/n}}{\sqrt{\mathbf{R}_{jj}(\beta^*)/n}} \right) - \frac{(\beta - \beta^*)(\widehat{\gamma}_j - \gamma_j^*)}{\sqrt{\widehat{\mathbf{R}}_{jj}(\beta)/n}} \right|. \end{aligned}$$

By applying (48), (44), (49), and (129), we obtain

$$\left| \max_{\beta \in \mathcal{U}_a} \frac{|\widehat{\Gamma}_j - \Gamma_j^* - \beta(\widehat{\gamma}_j - \gamma_j^*)|}{\sqrt{\widehat{\mathbf{R}}_{jj}(\beta)/n}} - \frac{|\widehat{\Gamma}_j - \Gamma_j^* - \beta^*(\widehat{\gamma}_j - \gamma_j^*)|}{\sqrt{\mathbf{R}_{jj}(\beta^*)/n}} \right| \xrightarrow{p} 0 \quad (131)$$

Together with (129), we establish

$$\mathbf{P} \left[\max_{\beta \in \mathcal{U}(a)} \frac{|\widehat{\Gamma}_j - \Gamma_j^* - \beta(\widehat{\gamma}_j - \gamma_j^*)|}{\sqrt{(\widehat{\mathbf{V}}_{jj}^\Gamma + \beta^2 \widehat{\mathbf{V}}_{jj}^\gamma - 2\beta \widehat{\mathbf{C}}_{jj})/n}} \geq \Phi^{-1} \left(1 - \frac{\alpha}{2|\widehat{\mathcal{S}}|} \right) \right] \leq \alpha/|\widehat{\mathcal{S}}|.$$

We further apply the union bound and establish

$$\begin{aligned} & \mathbf{P} \left[\max_{\beta \in \mathcal{U}(a)} \max_{j \in \widehat{\mathcal{S}}} \frac{|\widehat{\Gamma}_j - \Gamma_j^* - \beta(\widehat{\gamma}_j - \gamma_j^*)|}{\sqrt{(\widehat{\mathbf{V}}_{jj}^\Gamma + \beta^2 \widehat{\mathbf{V}}_{jj}^\gamma - 2\beta \widehat{\mathbf{C}}_{jj})/n}} \geq \Phi^{-1} \left(1 - \frac{\alpha}{2|\widehat{\mathcal{S}}|} \right) \right] \\ & \leq \sum_{j \in \widehat{\mathcal{S}}} \mathbf{P} \left[\max_{\beta \in \mathcal{U}(a)} \frac{|\widehat{\Gamma}_j - \Gamma_j^* - \beta(\widehat{\gamma}_j - \gamma_j^*)|}{\sqrt{(\widehat{\mathbf{V}}_{jj}^\Gamma + \beta^2 \widehat{\mathbf{V}}_{jj}^\gamma - 2\beta \widehat{\mathbf{C}}_{jj})/n}} \geq \Phi^{-1} \left(1 - \frac{\alpha}{2|\widehat{\mathcal{S}}|} \right) \right] \leq \alpha. \end{aligned} \quad (132)$$

This implies $\liminf_{n \rightarrow \infty} \mathbf{P}(\mathcal{E}_0(\alpha)) \geq 1 - \alpha$.

Note that

$$\widehat{\Gamma}_j - \beta \widehat{\gamma}_j - \pi_j^* = \widehat{\Gamma}_j - \Gamma_j^* - \beta(\widehat{\gamma}_j - \gamma_j^*) - (\beta^* - \beta)\gamma_j^*.$$

Similarly to (130), we have

$$\begin{aligned} & \frac{\widehat{\Gamma}_j - \beta \widehat{\gamma}_j - \pi_j^*}{\sqrt{\widehat{\mathbf{R}}_{jj}(\beta)/n}} - \frac{\widehat{\Gamma}_j - \Gamma_j^* - \beta^*(\widehat{\gamma}_j - \gamma_j^*)}{\sqrt{\mathbf{R}_{jj}(\beta^*)/n}} \\ & = \frac{\widehat{\Gamma}_j - \Gamma_j^* - \beta^*(\widehat{\gamma}_j - \gamma_j^*)}{\sqrt{\widehat{\mathbf{R}}_{jj}(\beta^*)/n}} \cdot \left(1 - \frac{\sqrt{\widehat{\mathbf{R}}_{jj}(\beta)/n}}{\sqrt{\mathbf{R}_{jj}(\beta^*)/n}} \right) - \frac{(\beta - \beta^*)(\widehat{\gamma}_j - \gamma_j^*)}{\sqrt{\widehat{\mathbf{R}}_{jj}(\beta)/n}} + \frac{(\beta - \beta^*)\gamma_j^*}{\sqrt{\widehat{\mathbf{R}}_{jj}(\beta)/n}}. \end{aligned}$$

Applying the similar argument as (131) and (132), we establish $\liminf_{n \rightarrow \infty} \mathbf{P}(\widetilde{\mathcal{E}}_0(\alpha)) \geq 1 - \alpha$ since $n^{1/2-a} \ll \Phi^{-1} \left(1 - \frac{\alpha}{2|\widehat{\mathcal{S}}|} \right)$.

D.3 Proof of Lemma 4

We shall establish the lower bound for $\text{SE}(\widehat{\pi}_k^{[j]})$ in (95).

$$\mathbf{R}_{k,k}^{[j]} + \frac{[\gamma_k^*]^2}{[\gamma_j^*]^2} \mathbf{R}_{j,j}^{[j]} - \frac{2\gamma_k^*}{\gamma_j^*} \mathbf{R}_{j,k}^{[j]} \geq \lambda_{\min}(\mathbf{R}_{\mathcal{J},\mathcal{J}}^{[j]}) \left(1 + \frac{[\gamma_k^*]^2}{[\gamma_j^*]^2}\right), \quad (133)$$

where $\mathcal{J} = \{j, k\}$. For any $\|u\|_2 = 1$ and $u \in \mathbb{R}^2$, we have

$$u^\top \mathbf{R}_{\mathcal{J},\mathcal{J}}^{[j]} u = u^\top \mathbf{V}_{\mathcal{J},\mathcal{J}}^\Gamma u + \left(\frac{\Gamma_j^*}{\gamma_j^*}\right)^2 u^\top \mathbf{V}_{\mathcal{J},\mathcal{J}}^\gamma u - 2\frac{\Gamma_j^*}{\gamma_j^*} u^\top \mathbf{C}_{\mathcal{J},\mathcal{J}} u \geq \left[1 + \left(\frac{\Gamma_j^*}{\gamma_j^*}\right)^2\right] \lambda_{\min}(\text{Cov}).$$

Combined with (133) and (41), we establish (95).

We now establish the asymptotic normality of $\mathcal{M}_k^{[j]}$. Note that

$$\begin{aligned} & \left(\widehat{\Omega}_k - \frac{\gamma_k^*}{\gamma_j^*} \widehat{\Omega}_j\right)^\top \frac{1}{n} W^\top \left(\epsilon - \frac{\Gamma_j^*}{\gamma_j^*} \delta\right) = \left(\Omega_{k,\cdot} - \frac{\gamma_k^*}{\gamma_j^*} \Omega_{j,\cdot}\right)^\top \frac{1}{n} W^\top \left(\epsilon - \frac{\Gamma_j^*}{\gamma_j^*} \delta\right) \\ & + \left(\widehat{\Omega}_k - \Omega_{k,\cdot}\right)^\top \frac{1}{n} W^\top \left(\epsilon - \frac{\Gamma_j^*}{\gamma_j^*} \delta\right) - \left(\widehat{\Omega}_j - \Omega_{j,\cdot}\right)^\top \frac{1}{n} W^\top \left(\epsilon - \frac{\Gamma_j^*}{\gamma_j^*} \delta\right) \cdot \frac{\gamma_k^*}{\gamma_j^*}. \end{aligned} \quad (134)$$

On the event \mathcal{G} , we have

$$\begin{aligned} & \left| \left(\widehat{\Omega}_k - \Omega_{k,\cdot}\right)^\top \frac{1}{n} W^\top \left(\epsilon - \frac{\Gamma_j^*}{\gamma_j^*} \delta\right) + \frac{\gamma_k^*}{\gamma_j^*} \left(\widehat{\Omega}_j - \Omega_{j,\cdot}\right)^\top \frac{1}{n} W^\top \left(\epsilon - \frac{\Gamma_j^*}{\gamma_j^*} \delta\right) \right| \\ & \lesssim \frac{\log n}{n} \left(1 + \left|\frac{\gamma_k^*}{\gamma_j^*}\right|\right) \left(1 + \left|\frac{\Gamma_j^*}{\gamma_j^*}\right|\right). \end{aligned}$$

Combined with (95), we have

$$\frac{1}{\text{SE}(\widehat{\pi}_k^{[j]})} \left| \left(\widehat{\Omega}_k - \Omega_{k,\cdot}\right)^\top \frac{1}{n} W^\top \left(\epsilon - \frac{\Gamma_j^*}{\gamma_j^*} \delta\right) + \frac{\gamma_k^*}{\gamma_j^*} \left(\widehat{\Omega}_j - \Omega_{j,\cdot}\right)^\top \frac{1}{n} W^\top \left(\epsilon - \frac{\Gamma_j^*}{\gamma_j^*} \delta\right) \right| \lesssim \frac{\log n}{\sqrt{n}}.$$

Together with

$$\frac{1}{\text{SE}(\widehat{\pi}_k^{[j]})} \left(\Omega_{k,\cdot} - \frac{\gamma_k^*}{\gamma_j^*} \Omega_{j,\cdot}\right)^\top \frac{1}{n} W^\top \left(\epsilon - \frac{\Gamma_j^*}{\gamma_j^*} \delta\right) \xrightarrow{d} N(0, 1),$$

we apply the decomposition (134) and establish (94).

We now establish the consistency of $\widehat{\text{SE}}(\widehat{\pi}_k^{[j]})$. We control the difference between $\widehat{\mathbf{R}}^{[j]}$ and $\mathbf{R}^{[j]} = \mathbf{V}^\Gamma + (\beta^{[j]})^2 \mathbf{V}^\gamma - 2\beta^{[j]} \mathbf{C}$. On the event \mathcal{G}_4 , we have

$$\max_{j,k \in \widehat{\mathcal{S}}} \left| \widehat{\mathbf{R}}_{j,k}^{[j]} - \mathbf{R}_{j,k}^{[j]} \right| \lesssim \frac{(\log n)^{3/2}}{\sqrt{n}} \left((\beta^{[j]})^2 + 1 \right) + \left| \widehat{\beta}^{[j]} - \beta^{[j]} \right| \cdot \left(1 + |\beta^{[j]}| \right) + (\widehat{\beta}^{[j]} - \beta^{[j]})^2. \quad (135)$$

Then we have the following upper bound for $n \cdot \left| \widehat{\text{SE}}^2(\widehat{\pi}_k^{[j]}) - \text{SE}^2(\widehat{\pi}_k^{[j]}) \right|$

$$C \max_{j,k \in \widehat{\mathcal{S}}} \left| \widehat{\mathbf{R}}_{j,k}^{[j]} - \mathbf{R}_{j,k}^{[j]} \right| \cdot \left(1 + \left(\frac{\widehat{\gamma}_k}{\widehat{\gamma}_j} \right)^2 \right) + C \max_{j,k \in \widehat{\mathcal{S}}} \left| \mathbf{R}_{j,k}^{[j]} \right| \cdot \left(\left| \left(\frac{\widehat{\gamma}_k}{\widehat{\gamma}_j} \right)^2 - \left(\frac{\gamma_k^*}{\gamma_j^*} \right)^2 \right| + \left| \frac{\widehat{\gamma}_k}{\widehat{\gamma}_j} - \frac{\gamma_k^*}{\gamma_j^*} \right| \right). \quad (136)$$

On the event $\mathcal{G}_6 \cap \mathcal{G}_7$, we apply (135) and establish

$$\max_{j,k \in \widehat{\mathcal{S}}} \left| \widehat{\mathbf{R}}_{j,k}^{[j]} - \mathbf{R}_{j,k}^{[j]} \right| \cdot \left(1 + \left(\frac{\widehat{\gamma}_k}{\widehat{\gamma}_j} \right)^2 \right) \lesssim \left(1 + \left(\frac{\Gamma_j^*}{\gamma_j^*} \right)^2 \right) \cdot \left(1 + \left(\frac{\gamma_k^*}{\gamma_j^*} \right)^2 \right) \frac{1}{(\log n)^{1/4}}. \quad (137)$$

Note that

$$\begin{aligned} \mathbf{R}_{j,k}^{[j]} &= u^\top \mathbf{V}_{j,k}^\Gamma + \left(\frac{\Gamma_j^*}{\gamma_j^*} \right)^2 \mathbf{V}_{j,k}^\gamma - 2 \frac{\Gamma_j^*}{\gamma_j^*} \mathbf{C}_{j,k} \leq \left[1 + \left(\frac{\Gamma_j^*}{\gamma_j^*} \right)^2 \right] \lambda_{\max}(\text{Cov}) \\ &\leq \left[1 + \left(\frac{\Gamma_j^*}{\gamma_j^*} \right)^2 \right] C_1 \cdot \lambda_{\max}(\Sigma^{-1}), \end{aligned}$$

where the least inequality follows from (42). On the event \mathcal{G}_6 , we have

$$\max_{j,k \in \widehat{\mathcal{S}}} \left| \mathbf{R}_{j,k}^{[j]} \right| \cdot \left(\left| \left(\frac{\widehat{\gamma}_k}{\widehat{\gamma}_j} \right)^2 - \left(\frac{\gamma_k^*}{\gamma_j^*} \right)^2 \right| + \left| \frac{\widehat{\gamma}_k}{\widehat{\gamma}_j} - \frac{\gamma_k^*}{\gamma_j^*} \right| \right) \lesssim \left[1 + \left(\frac{\Gamma_j^*}{\gamma_j^*} \right)^2 \right] \left(1 + \left(\frac{\gamma_k^*}{\gamma_j^*} \right)^2 \right) \frac{1}{(\log n)^{1/4}}.$$

Together with the error decomposition (136) and the inequalities (137) and (95), we establish (96).

E Additional Numerical Results

E.1 Additional Simulation Results for Settings S1 to S5

We present the extra simulation results for settings **S1** to **S5**. In Table E.1, we report the coverage and length properties of different CIs for Settings **S1** to **S5** with $\tau = 0.4$.

We also consider the heteroscedastic errors demonstrate the robustness of our method. For $1 \leq i \leq n$, generate $\delta_i \sim N(0, 1)$ and

$$e_i = 0.3\delta_i + \sqrt{[1 - 0.3^2]/[0.86^4 + 1.38072^2]}(1.38072 \cdot \tau_{1,i} + 0.86^2 \cdot \tau_{2,i}), \quad (138)$$

where conditioning on Z_i , $\tau_{1,i} \sim N(0, [0.5 \cdot Z_{i,1}^2 + 0.25]^2)$, $\tau_{2,i} \sim N(0, 1)$, and $\tau_{1,i}$ and $\tau_{2,i}$ are independent of δ_i . In Table E.2, we report the coverage and length properties of different CIs for Settings **S1** to **S5** with heteroscedastic errors and $\tau = 0.2$. In Table E.3, we report the coverage and length properties of different CIs for Settings **S1** to **S5** with heteroscedastic errors and $\tau = 0.4$.

The main observation is similar to those in Section 7 in the main paper, which is summarized in the following.

1. The CIs by TSHT [25] and CIIV [53] achieve the 95% coverage level for a large sample size and a relatively large violation level, such as $n = 5000$ and $\tau = 0.4$. For many settings with $\tau = 0.2$, the CIs by TSHT and CIIV do not even have coverage when $n = 5000$. The CI by CIIV is more robust in the sense that its validity may require a smaller sample size than TSHT.

Empirical Coverage of Confidence Intervals for $\tau = 0.4$

		Oracle				Searching		Sampling		Union		
Set	n	TSLS	BA	TSHT	CIIV	$\hat{\mathcal{V}}^{\text{TSHT}}$	$\hat{\mathcal{V}}^{\text{CIIV}}$	$\hat{\mathcal{V}}^{\text{TSHT}}$	$\hat{\mathcal{V}}^{\text{CIIV}}$	Check	$p_z - 1$	$\lceil p_z/2 \rceil$
S1	500	0.95	0.95	0.79	0.82	1.00	1.00	1.00	1.00	1.00	1.00	0.97
	1000	0.94	0.95	0.89	0.89	1.00	1.00	1.00	1.00	1.00	1.00	0.95
	2000	0.96	0.96	0.96	0.95	1.00	1.00	1.00	1.00	1.00	1.00	0.95
	5000	0.94	0.95	0.94	0.94	1.00	1.00	1.00	1.00	1.00	1.00	0.96
S2	500	0.95	0.95	0.72	0.72	0.97	0.98	0.96	0.96	0.98	1.00	0.00
	1000	0.94	0.93	0.79	0.87	0.98	1.00	0.98	0.99	0.99	1.00	0.00
	2000	0.94	0.96	0.92	0.93	1.00	1.00	1.00	1.00	1.00	1.00	0.00
	5000	0.96	0.95	0.96	0.95	1.00	1.00	1.00	1.00	1.00	1.00	0.00
S3	500	0.95	0.93	0.67	0.71	0.90	0.96	0.91	0.94	0.97	1.00	0.26
	1000	0.94	0.93	0.71	0.83	0.94	0.99	0.94	0.98	0.94	1.00	0.01
	2000	0.94	0.93	0.88	0.91	0.99	1.00	0.99	1.00	1.00	1.00	0.00
	5000	0.96	0.98	0.96	0.95	1.00	1.00	1.00	1.00	1.00	1.00	0.00
S4	500	0.95	0.96	0.53	0.53	0.90	0.83	0.89	0.81	0.75	0.95	0.00
	1000	0.94	0.91	0.64	0.84	0.98	0.93	0.97	0.92	0.75	0.94	0.00
	2000	0.93	0.95	0.88	0.92	0.98	0.98	0.97	0.96	0.94	0.94	0.00
	5000	0.95	0.99	0.94	0.94	0.98	0.98	0.98	0.98	0.99	0.94	0.00
S5	500	0.95	0.96	0.19	0.45	0.59	0.74	0.61	0.73	0.61	0.95	0.00
	1000	0.94	0.89	0.30	0.77	0.96	0.85	0.96	0.84	0.44	0.94	0.00
	2000	0.93	0.94	0.74	0.91	0.99	0.98	0.97	0.97	0.78	0.94	0.00
	5000	0.95	0.99	0.94	0.94	0.98	0.98	0.98	0.98	0.99	0.94	0.00

Average Lengths of Confidence Intervals for $\tau = 0.4$

		Oracle				Searching		Sampling		Union		
Set	n	TSLS	BA	TSHT	CIIV	$\hat{\mathcal{V}}^{\text{TSHT}}$	$\hat{\mathcal{V}}^{\text{CIIV}}$	$\hat{\mathcal{V}}^{\text{TSHT}}$	$\hat{\mathcal{V}}^{\text{CIIV}}$	Check	$p_z - 1$	$\lceil p_z/2 \rceil$
S1	500	0.10	0.16	0.09	0.09	0.57	0.62	0.33	0.35	-	1.28	0.17
	1000	0.07	0.10	0.07	0.07	0.40	0.41	0.22	0.23	-	0.70	0.08
	2000	0.05	0.05	0.05	0.05	0.29	0.29	0.16	0.16	-	0.41	0.05
	5000	0.03	0.03	0.03	0.03	0.18	0.18	0.10	0.10	-	0.25	0.03
S2	500	0.13	0.48	0.27	0.12	0.56	0.59	0.41	0.37	-	2.58	0.00
	1000	0.09	0.33	0.16	0.09	0.37	0.39	0.23	0.24	-	1.49	0.00
	2000	0.06	0.09	0.07	0.07	0.27	0.27	0.16	0.16	-	0.73	0.00
	5000	0.04	0.04	0.04	0.04	0.17	0.17	0.10	0.10	-	0.26	0.00
S3	500	0.13	0.53	0.21	0.13	0.61	0.64	0.53	0.41	-	1.90	0.04
	1000	0.09	0.44	0.21	0.09	0.42	0.41	0.31	0.26	-	1.40	0.00
	2000	0.06	0.24	0.14	0.07	0.27	0.27	0.17	0.16	-	0.83	0.00
	5000	0.04	0.05	0.04	0.04	0.17	0.17	0.10	0.10	-	0.33	0.00
S4	500	0.23	1.35	0.31	0.24	0.86	0.54	0.79	0.40	-	0.80	0.00
	1000	0.16	1.14	0.18	0.16	0.64	0.32	0.58	0.23	-	0.34	0.00
	2000	0.11	0.46	0.11	0.11	0.27	0.21	0.22	0.16	-	0.14	0.00
	5000	0.07	0.12	0.07	0.07	0.14	0.14	0.11	0.11	-	0.08	0.00
S5	500	0.23	1.05	0.24	0.23	0.70	0.52	0.69	0.39	-	0.98	0.00
	1000	0.16	1.01	0.14	0.16	0.83	0.31	0.79	0.23	-	0.40	0.00
	2000	0.11	0.52	0.11	0.11	0.34	0.21	0.30	0.16	-	0.15	0.00
	5000	0.07	0.14	0.07	0.07	0.14	0.14	0.11	0.10	-	0.08	0.00

Table E.1: Settings **S1** to **S5** with $\tau = 0.4$ and homoscedastic errors. The columns indexed with TSLS, BA TSHT and CIIV represent the oracle TSLS CI with the knowledge of \mathcal{V} , the oracle bias-aware confidence interval, the CI by TSHT, and the CI by CIIV, respectively. Under the columns indexed with “Searching” (or “Sampling”), the columns indexed with $\hat{\mathcal{V}}^{\text{TSHT}}$ and $\hat{\mathcal{V}}^{\text{CIIV}}$ represent our proposed searching (or sampling) CI in Algorithm 3 with $\hat{\mathcal{V}}^{\text{TSHT}}$ and $\hat{\mathcal{V}}^{\text{CIIV}}$, respectively. The column indexed with “Check” reports the proportion of simulations passing the Plurality rule check in Algorithm 3. The columns indexed with Union represent the union of TSLS estimators, which pass the Sargan test. The columns indexed with $p_z - 1$ and $\lceil p_z/2 \rceil$ correspond to the Union methods assuming two valid IVs and the majority rule, respectively.

2. The CIs by the **Union** method [31] with $\bar{s} = p_z - 1$ (assuming there are two valid IVs) achieve the desired coverage levels for all settings. The CIs by the **Union** method with $\bar{s} = \lceil p_z/2 \rceil$ (assuming the majority rule) do not achieve the desired coverage level, except for the setting **S1** where the majority rule holds.
3. Our proposed searching and sampling CIs achieve the desired coverage levels in most settings. Settings **S1**, **S2** and **S4** are relatively easier as the corresponding finite-sample majority and plurality rules hold more plausibly.
4. When the CIs by **TSHT** [25] and **CIIV** [53] are valid, their lengths are similar to the length of the CI by **oracle** TSLS, which has been justified in [25, 53]. The sampling CI, searching CI and CI by the **Union** are in general longer than the CI by the **oracle** TSLS, which is a price to pay for constructing uniformly valid CIs.
5. Among all CIs achieving the desired coverage level, the sampling CIs are typically the shortest CIs achieving the desired coverage levels. Both searching and sampling CIs are in general shorter than the CIs by the **Union** method.
6. The sampling CI can have a comparable length to the oracle bias-aware confidence interval. The oracle bias-aware confidence interval serves as a benchmark method in the settings with the IV selection error.

In Figure E.1, we investigate the performance of our proposed methods by varying the violation strength. We present the results for homoscedastic regression errors with different τ , which is the counterpart of the results reported in Figure 6 in the main paper. We generate the errors $(e_i, \delta_i)^\top$ following bivariate normal with zero mean, unit variance and $\text{Cov}(e_i, \delta_i) = 0.8$.

E.2 CIIV setting

Similar to the setting in CIIV paper [53], we further consider the following settings.

CIIV-1 (Plurality rule): set $\gamma^* = 0.4 \cdot \mathbf{1}_{21}$ and $\pi^* = (\mathbf{0}_9, \tau \cdot \mathbf{1}_6, \frac{\tau}{2} \cdot \mathbf{1}_6)^\top$.

CIIV-2 (Plurality rule): set $\gamma^* = 0.4 \cdot \mathbf{1}_{21}$ and $\pi^* = (\mathbf{0}_9, \tau \cdot \mathbf{1}_3, -\tau \cdot \mathbf{1}_3, \frac{\tau}{2} \cdot \mathbf{1}_3, -\frac{\tau}{2} \cdot \mathbf{1}_3)^\top$.

We vary τ across $\{0.2, 0.4\}$ where τ represents the invalidity level. The setting **CIIV-1** with $\tau = 0.4$ corresponds to the exact setting considered in [53]. For a small τ and sample size n , the setting **CIIV-1** does not necessarily satisfy the finite-sample plurality rule (Condition 4) since τ and $\tau/2$ are close to each other for a small $\tau > 0$. For the setting **CIIV-2**, the invalidity levels are more spread out and the finite-sample plurality rule may hold more plausibly.

In Table E.4, we consider the setting **CIIV-1** and compare different CIs in terms of empirical coverage and average lengths. In terms of coverage, our proposed searching and sampling CIs attain the desired coverage level (95%); CIs by the **Union** method achieve the desired coverage level; **CIIV** achieves the desired 95% coverage level for $\tau = 0.2$ with $n = 5000$ and $\tau = 0.4$ with $n = 2000, 5000$; and **TSHT** achieves the desired 95% coverage level only for $\tau = 0.4$ with

Empirical Coverage of Confidence Intervals for $\tau = 0.2$

		Oracle				Searching		Sampling		
Set	n	TSLS	BA	TSHT	CIIV	$\hat{\mathcal{V}}^{\text{TSHT}}$	$\hat{\mathcal{V}}^{\text{CIIV}}$	$\hat{\mathcal{V}}^{\text{TSHT}}$	$\hat{\mathcal{V}}^{\text{CIIV}}$	Check
S1	500	0.92	0.96	0.49	0.51	1.00	1.00	1.00	1.00	1.00
	1000	0.94	0.98	0.38	0.61	1.00	1.00	1.00	0.99	1.00
	2000	0.95	0.95	0.52	0.75	1.00	1.00	1.00	1.00	1.00
	5000	0.94	0.94	0.83	0.90	1.00	1.00	1.00	1.00	1.00
S2	500	0.95	0.95	0.58	0.41	1.00	1.00	0.99	1.00	1.00
	1000	0.92	0.95	0.47	0.45	1.00	1.00	0.98	0.98	1.00
	2000	0.94	0.94	0.52	0.61	0.99	0.97	0.99	0.94	1.00
	5000	0.94	0.94	0.76	0.86	0.98	0.99	0.99	0.99	1.00
S3	500	0.95	0.96	0.65	0.52	0.99	0.99	0.98	0.98	1.00
	1000	0.92	0.96	0.57	0.46	0.98	1.00	0.97	0.97	1.00
	2000	0.94	0.95	0.60	0.61	0.98	0.97	0.97	0.94	1.00
	5000	0.94	0.95	0.76	0.86	0.98	0.99	0.99	0.99	1.00
S4	500	0.94	0.96	0.68	0.63	0.97	0.97	0.97	0.95	0.99
	1000	0.96	0.99	0.54	0.45	0.98	0.94	0.98	0.94	0.99
	2000	0.94	0.98	0.48	0.42	0.99	0.93	0.97	0.92	1.00
	5000	0.94	0.97	0.72	0.75	0.97	0.93	0.96	0.92	1.00
S5	500	0.94	0.97	0.47	0.37	0.83	0.86	0.84	0.81	0.98
	1000	0.96	0.97	0.33	0.33	0.55	0.75	0.67	0.73	0.98
	2000	0.94	0.97	0.19	0.36	0.51	0.85	0.61	0.84	0.89
	5000	0.94	0.95	0.57	0.75	0.97	0.92	0.96	0.92	0.84

Average Lengths of Confidence Intervals for $\tau = 0.2$

		Oracle				Searching		Sampling		
Set	n	TSLS	BA	TSHT	CIIV	$\hat{\mathcal{V}}^{\text{TSHT}}$	$\hat{\mathcal{V}}^{\text{CIIV}}$	$\hat{\mathcal{V}}^{\text{TSHT}}$	$\hat{\mathcal{V}}^{\text{CIIV}}$	Check
S1	500	0.10	0.18	0.09	0.09	0.63	0.65	0.37	0.37	-
	1000	0.08	0.16	0.06	0.07	0.44	0.47	0.27	0.28	-
	2000	0.05	0.13	0.05	0.05	0.31	0.33	0.20	0.20	-
	5000	0.03	0.05	0.03	0.03	0.19	0.20	0.11	0.11	-
S2	500	0.15	0.28	0.17	0.11	0.64	0.65	0.41	0.41	-
	1000	0.11	0.27	0.15	0.09	0.43	0.47	0.30	0.31	-
	2000	0.08	0.27	0.16	0.07	0.30	0.33	0.24	0.23	-
	5000	0.05	0.21	0.13	0.05	0.18	0.20	0.13	0.13	-
S3	500	0.15	0.27	0.13	0.11	0.63	0.68	0.45	0.42	-
	1000	0.11	0.25	0.13	0.09	0.43	0.46	0.32	0.31	-
	2000	0.08	0.26	0.15	0.07	0.31	0.33	0.24	0.23	-
	5000	0.05	0.21	0.13	0.05	0.18	0.20	0.13	0.13	-
S4	500	0.32	0.60	0.31	0.21	0.69	0.70	0.61	0.59	-
	1000	0.23	0.40	0.20	0.16	0.51	0.49	0.45	0.42	-
	2000	0.16	0.30	0.17	0.13	0.36	0.32	0.32	0.28	-
	5000	0.10	0.22	0.13	0.10	0.21	0.18	0.19	0.15	-
S5	500	0.32	0.58	0.32	0.20	0.54	0.61	0.50	0.51	-
	1000	0.23	0.57	0.27	0.16	0.34	0.46	0.36	0.40	-
	2000	0.16	0.49	0.18	0.13	0.27	0.32	0.28	0.28	-
	5000	0.10	0.31	0.12	0.10	0.25	0.18	0.24	0.15	-

Table E.2: Settings **S1** to **S5** with $\tau = 0.2$ and heteroscedastic errors. The columns indexed with TSLS, BA TSHT and CIIV represent the oracle TSLS CI with the knowledge of \mathcal{V} , the oracle bias-aware confidence interval, the CI by TSHT, and the CI by CIIV, respectively. Under the columns indexed with “Searching” (or “Sampling”), the columns indexed with $\hat{\mathcal{V}}^{\text{TSHT}}$ and $\hat{\mathcal{V}}^{\text{CIIV}}$ represent our proposed searching (or sampling) CI in Algorithm 3 with $\hat{\mathcal{V}}^{\text{TSHT}}$ and $\hat{\mathcal{V}}^{\text{CIIV}}$, respectively. The column indexed with “Check” reports the proportion of simulations passing the Plurality rule check in Algorithm 3. The columns indexed with **Union** represent the union of TSLS estimators, which pass the Sargan test. The columns indexed with $p_z - 1$ and $\lceil p_z/2 \rceil$ correspond to the **Union** methods assuming two valid IVs and the majority rule, respectively.

Empirical Coverage of Confidence Intervals for $\tau = 0.4$

		Oracle				Searching		Sampling		
Set	n	TSLs	BA	TSHT	CIIV	$\hat{\mathcal{V}}^{\text{TSHT}}$	$\hat{\mathcal{V}}^{\text{CIIV}}$	$\hat{\mathcal{V}}^{\text{TSHT}}$	$\hat{\mathcal{V}}^{\text{CIIV}}$	Check
S1	500	0.92	0.94	0.63	0.72	1.00	1.00	1.00	1.00	1.00
	1000	0.94	0.94	0.80	0.86	1.00	1.00	1.00	1.00	1.00
	2000	0.95	0.95	0.94	0.95	1.00	1.00	1.00	1.00	1.00
	5000	0.94	0.94	0.94	0.93	1.00	1.00	1.00	1.00	1.00
S2	500	0.95	0.96	0.68	0.60	0.97	0.98	0.97	0.96	1.00
	1000	0.92	0.95	0.77	0.78	0.98	0.98	0.98	0.97	1.00
	2000	0.94	0.92	0.88	0.94	0.99	1.00	0.99	1.00	1.00
	5000	0.94	0.97	0.93	0.94	1.00	1.00	1.00	1.00	1.00
S3	500	0.95	0.94	0.72	0.60	0.92	0.97	0.92	0.95	0.97
	1000	0.92	0.94	0.82	0.72	0.89	0.98	0.89	0.97	0.99
	2000	0.94	0.93	0.86	0.93	0.98	1.00	0.98	1.00	0.99
	5000	0.94	0.97	0.92	0.94	1.00	1.00	1.00	1.00	1.00
S4	500	0.94	0.93	0.52	0.38	0.91	0.86	0.91	0.84	0.92
	1000	0.96	0.92	0.69	0.70	0.98	0.93	0.97	0.91	0.92
	2000	0.94	0.98	0.89	0.91	0.98	0.96	0.97	0.96	0.98
	5000	0.94	0.97	0.93	0.93	0.97	0.97	0.98	0.97	1.00
S5	500	0.94	0.99	0.28	0.21	0.38	0.60	0.49	0.58	0.91
	1000	0.96	1.00	0.17	0.52	0.55	0.73	0.57	0.72	0.64
	2000	0.94	0.98	0.33	0.84	0.97	0.89	0.96	0.89	0.40
	5000	0.94	0.83	0.77	0.93	0.98	0.97	0.98	0.97	0.82

Average Lengths of Confidence Intervals for $\tau = 0.4$

		Oracle				Searching		Sampling		
Set	n	TSLs	BA	TSHT	CIIV	$\hat{\mathcal{V}}^{\text{TSHT}}$	$\hat{\mathcal{V}}^{\text{CIIV}}$	$\hat{\mathcal{V}}^{\text{TSHT}}$	$\hat{\mathcal{V}}^{\text{CIIV}}$	Check
S1	500	0.10	0.23	0.10	0.10	0.62	0.68	0.39	0.39	-
	1000	0.08	0.13	0.08	0.07	0.42	0.45	0.25	0.26	-
	2000	0.05	0.06	0.06	0.05	0.30	0.31	0.17	0.18	-
	5000	0.03	0.04	0.04	0.03	0.20	0.20	0.11	0.11	-
S2	500	0.15	0.51	0.34	0.14	0.62	0.68	0.49	0.45	-
	1000	0.11	0.44	0.29	0.11	0.41	0.45	0.31	0.29	-
	2000	0.08	0.26	0.14	0.08	0.29	0.31	0.19	0.20	-
	5000	0.05	0.06	0.05	0.05	0.19	0.19	0.12	0.12	-
S3	500	0.15	0.47	0.26	0.13	0.61	0.68	0.54	0.46	-
	1000	0.11	0.43	0.30	0.11	0.41	0.46	0.34	0.31	-
	2000	0.08	0.33	0.20	0.08	0.30	0.31	0.20	0.20	-
	5000	0.05	0.09	0.06	0.05	0.19	0.19	0.12	0.12	-
S4	500	0.32	0.99	0.37	0.26	0.79	0.68	0.72	0.57	-
	1000	0.23	0.92	0.29	0.21	0.59	0.41	0.53	0.35	-
	2000	0.16	0.40	0.18	0.16	0.30	0.27	0.26	0.22	-
	5000	0.10	0.13	0.10	0.10	0.17	0.17	0.14	0.14	-
S5	500	0.32	1.04	0.37	0.25	0.52	0.64	0.58	0.56	-
	1000	0.23	0.97	0.23	0.21	0.55	0.42	0.57	0.35	-
	2000	0.16	0.78	0.14	0.16	0.57	0.27	0.55	0.22	-
	5000	0.10	0.41	0.10	0.10	0.25	0.17	0.23	0.14	-

Table E.3: Settings **S1** to **S5** with $\tau = 0.4$ and heteroscedastic errors. The columns indexed with TSLs, BA TSHT and CIIV represent the oracle TSLs CI with the knowledge of \mathcal{V} , the oracle bias-aware confidence interval, the CI by TSHT, and the CI by CIIV, respectively. Under the columns indexed with ‘‘Searching’’ (or ‘‘Sampling’’), the columns indexed with $\hat{\mathcal{V}}^{\text{TSHT}}$ and $\hat{\mathcal{V}}^{\text{CIIV}}$ represent our proposed searching (or sampling) CI in Algorithm 3 with $\hat{\mathcal{V}}^{\text{TSHT}}$ and $\hat{\mathcal{V}}^{\text{CIIV}}$, respectively. The column indexed with ‘‘Check’’ reports the proportion of simulations passing the Plurality rule check in Algorithm 3. The columns indexed with **Union** represent the union of TSLs estimators, which pass the Sargan test. The columns indexed with $p_z - 1$ and $\lceil p_z/2 \rceil$ correspond to the **Union** methods assuming two valid IVs and the majority rule, respectively.




Figure E.1: Empirical coverage and average lengths for the setting **S2** with $\tau \in \{0.025, 0.05, 0.075, 0.1, 0.2, 0.3, 0.4, 0.5\}$ and homoscedastic errors. **TSHT**, **CIIV**, and **Union** stand for the CIs by [25], [53], and [31], respectively. **Oracle BA** stands for the oracle bias-aware CI in (10), which is the benchmark. **Searching** and **Sampling** correspond to our proposed searching and sampling CIs in Algorithm 3 in the main paper.

Empirical Coverage of Confidence Intervals For **CIIV-1**

		Oracle				Searching		Sampling			Union
τ	n	TSLS	BA	TSHT	CIIV	$\hat{\mathcal{V}}^{\text{TSHT}}$	$\hat{\mathcal{V}}^{\text{CIIV}}$	$\hat{\mathcal{V}}^{\text{TSHT}}$	$\hat{\mathcal{V}}^{\text{CIIV}}$	Check	$p_z - 1$
0.2	500	0.94	0.96	0.00	0.16	1.00	0.99	0.92	0.62	1.00	1.00
	1000	0.93	0.98	0.00	0.42	1.00	0.91	0.94	0.69	1.00	1.00
	2000	0.95	0.93	0.00	0.72	0.58	0.94	0.58	0.93	1.00	1.00
	5000	0.95	0.94	0.01	0.91	0.66	1.00	0.66	1.00	0.80	1.00
0.4	500	0.94	0.95	0.00	0.69	0.60	0.90	0.60	0.87	1.00	1.00
	1000	0.93	0.95	0.00	0.89	0.36	0.98	0.36	0.98	0.98	1.00
	2000	0.95	0.90	0.40	0.94	0.98	1.00	0.98	1.00	0.90	1.00
	5000	0.95	0.95	0.95	0.95	1.00	1.00	1.00	1.00	1.00	1.00

Average Lengths of Confidence Intervals for **CIIV-1**

		Oracle				Searching		Sampling			Union
τ	n	TSLS	BA	TSHT	CIIV	$\hat{\mathcal{V}}^{\text{TSHT}}$	$\hat{\mathcal{V}}^{\text{CIIV}}$	$\hat{\mathcal{V}}^{\text{TSHT}}$	$\hat{\mathcal{V}}^{\text{CIIV}}$	Check	$p_z - 1$
0.2	500	0.09	0.50	0.06	0.09	0.90	0.88	0.52	0.40	-	1.44
	1000	0.07	0.51	0.04	0.07	0.60	0.60	0.47	0.29	-	1.10
	2000	0.05	0.49	0.03	0.05	0.39	0.42	0.39	0.21	-	0.90
	5000	0.03	0.51	0.05	0.03	0.18	0.27	0.12	0.13	-	0.73
0.4	500	0.09	0.96	0.06	0.10	0.88	0.90	0.88	0.42	-	2.08
	1000	0.07	1.02	0.06	0.07	0.39	0.61	0.34	0.28	-	1.68
	2000	0.05	1.08	0.20	0.05	0.40	0.43	0.20	0.21	-	1.25
	5000	0.03	0.03	0.03	0.03	0.27	0.27	0.13	0.13	-	0.80

Table E.4: The setting **CIIV-1**. The columns indexed with TSLS, BA TSHT and CIIV represent the oracle TSLS CI with the knowledge of \mathcal{V} , the oracle bias-aware confidence interval, the CI by TSHT, and the CI by CIIV, respectively. Under the columns indexed with ‘‘Searching’’ (or ‘‘Sampling’’), the columns indexed with $\hat{\mathcal{V}}^{\text{TSHT}}$ and $\hat{\mathcal{V}}^{\text{CIIV}}$ represent our proposed searching (or sampling) CI in Algorithm 3 with $\hat{\mathcal{V}}^{\text{TSHT}}$ and $\hat{\mathcal{V}}^{\text{CIIV}}$, respectively. The column indexed with ‘‘Check’’ reports the proportion of simulations passing the Plurality rule check in Algorithm 3. The column indexed with $p_z - 1$ corresponds to the Union methods assuming two valid IVs.

$n = 5000$. We shall point out that the searching and sampling CIs with inaccurate initial estimators $\hat{\mathcal{V}}^{\text{TSHT}}$ tend to perform badly in terms of coverage, see the settings with $\tau = 0.2$ and $n = 5000$ or $\tau = 0.4$ and $n = 2000$. The corresponding searching and sampling CIs with the initial estimators $\hat{\mathcal{V}}^{\text{CIIV}}$ are more reliable in these settings.

In terms of interval lengths, the sampling and searching CIs are much shorter than the CIs by the **Union** method. The sampling CI can have a comparable length with the oracle bias-aware confidence interval.

The results for the setting **CIIV-2** is reported in Table E.5.

τ	n	Oracle				Searching		Sampling		Check	Union
		TSLS	BA	TSHT	CIIV	$\hat{\mathcal{V}}^{\text{TSHT}}$	$\hat{\mathcal{V}}^{\text{CIIV}}$	$\hat{\mathcal{V}}^{\text{TSHT}}$	$\hat{\mathcal{V}}^{\text{CIIV}}$		
0.2	500	0.94	0.94	0.60	0.55	1.00	1.00	1.00	0.97	1.00	1.00
	1000	0.93	0.94	0.55	0.80	1.00	1.00	1.00	0.98	1.00	1.00
	2000	0.95	0.95	0.60	0.84	0.95	1.00	0.95	1.00	1.00	1.00
	5000	0.95	0.93	0.88	0.91	1.00	1.00	1.00	1.00	0.98	1.00
0.4	500	0.94	0.95	0.59	0.83	0.96	1.00	0.96	1.00	1.00	1.00
	1000	0.93	0.94	0.70	0.90	0.96	1.00	0.96	1.00	0.98	1.00
	2000	0.95	0.98	0.71	0.94	1.00	1.00	1.00	1.00	1.00	1.00
	5000	0.95	0.96	0.95	0.95	1.00	1.00	1.00	1.00	1.00	1.00

τ	n	Oracle				Searching		Sampling		Check	Union
		TSLS	BA	TSHT	CIIV	$\hat{\mathcal{V}}^{\text{TSHT}}$	$\hat{\mathcal{V}}^{\text{CIIV}}$	$\hat{\mathcal{V}}^{\text{TSHT}}$	$\hat{\mathcal{V}}^{\text{CIIV}}$		
0.2	500	0.09	0.14	0.06	0.09	0.94	0.92	0.55	0.43	-	1.78
	1000	0.07	0.14	0.05	0.07	0.60	0.61	0.50	0.30	-	1.46
	2000	0.05	0.10	0.04	0.05	0.30	0.42	0.30	0.21	-	1.30
	5000	0.03	0.09	0.06	0.03	0.23	0.27	0.13	0.13	-	1.04
0.4	500	0.09	0.18	0.08	0.09	0.72	0.91	0.70	0.42	-	2.91
	1000	0.07	0.15	0.07	0.07	0.52	0.61	0.34	0.29	-	2.48
	2000	0.05	0.41	0.21	0.05	0.40	0.43	0.19	0.21	-	1.81
	5000	0.03	0.03	0.03	0.03	0.27	0.27	0.13	0.13	-	1.11

Table E.5: The setting **CIIV-2**. The columns indexed with TSLS, BA TSHT and CIIV represent the oracle TSLS CI with the knowledge of \mathcal{V} , the oracle bias-aware confidence interval, the CI by TSHT, and the CI by CIIV, respectively. Under the columns indexed with ‘‘Searching’’ (or ‘‘Sampling’’), the columns indexed with $\hat{\mathcal{V}}^{\text{TSHT}}$ and $\hat{\mathcal{V}}^{\text{CIIV}}$ represent our proposed searching (or sampling) CI in Algorithm 3 with $\hat{\mathcal{V}}^{\text{TSHT}}$ and $\hat{\mathcal{V}}^{\text{CIIV}}$, respectively. The column indexed with ‘‘Check’’ reports the proportion of simulations passing the Plurality rule check in Algorithm 3. The column indexed with $p_z - 1$ corresponds to the **Union** methods assuming two valid IVs.

E.3 Additional high-dimensional results

As reported in Table E.6, the CIs by TSHT can only achieve the desired coverage for a sufficiently large sample size n and invalidity level τ . In contrast, our proposed searching and sampling CIs achieve the desired coverage across all settings. The sampling CI is helpful in reducing the

length of the searching CI. The sampling CI is in general longer than the oracle bias-aware CI (the benchmark), but the sampling CI can have a comparable length to the oracle bias-aware CI in certain scenarios. In Table E.7, we report the high-dimensional simulation results with $p_z = 100$ and the results are similar to those in Table E.6.

		Oracle TSLS		Oracle BA		TSHT			Searching		Sampling	
τ	n	Cov	Len	Cov	Len	Bias	Cov	Len	Cov	Len	Cov	Len
0.5	200	0.96	0.26	0.97	0.53	0.17	0.10	0.18	0.99	1.52	0.97	0.80
	300	0.94	0.21	0.95	0.47	0.16	0.04	0.15	1.00	1.28	0.99	0.66
	1000	0.95	0.11	0.96	0.39	0.15	0.01	0.09	1.00	0.71	1.00	0.35
	2500	0.95	0.07	0.92	0.17	0.02	0.72	0.07	1.00	0.51	1.00	0.26
1.0	200	0.96	0.26	0.98	0.84	0.25	0.10	0.18	0.99	1.48	0.96	0.83
	300	0.94	0.21	0.98	0.74	0.20	0.15	0.16	1.00	1.23	1.00	0.66
	1000	0.95	0.11	0.89	0.27	0.03	0.81	0.12	1.00	0.69	1.00	0.38
	2500	0.95	0.07	0.94	0.07	0.00	0.95	0.07	1.00	0.51	1.00	0.27
2.0	200	0.96	0.26	0.96	0.37	0.05	0.80	0.23	1.00	1.44	0.99	0.81
	300	0.94	0.21	0.95	0.24	0.03	0.88	0.19	1.00	1.21	1.00	0.67
	1000	0.95	0.11	0.96	0.13	0.01	0.93	0.12	1.00	0.70	1.00	0.38
	2500	0.95	0.07	0.94	0.07	0.00	0.95	0.07	1.00	0.51	1.00	0.27

Table E.6: High-dimensional setting with $p_z = 200$ and $p_x = 150$. The columns indexed with “Cov”, “Len”, and “Bias” denote the empirical coverage, the average lengths, and the absolute bias, respectively. The columns indexed with **Oracle TSLS**, **Oracle BA**, and **TSHT** represent the oracle TSLS CI with the knowledge of \mathcal{V} and relevant IV and covariates, the oracle bias-aware CI in (10), the CI by [25], respectively.

E.4 Extra real data results

We present the summary statistics of the candidate IVs and baseline covariates in Table E.8. We shall provide some explanations on “Statement about fairness” and “Statement about Talent”. The CFPS survey has a few questions about subjects’ view on the fair competition and talent pay-off, which are measures of subjects’ personality traits. The statement on fair competition measures an individual’s willingness to compete through ability or education. A strong sense of competing for opportunities through fair competition signals a willingness to improve personal skills and knowledge through education. The statement on the talent indicates if an individual thinks their educational endeavors will pay off. A high score on the pay-off manifests that one would like to strive for higher education because they believe that their education will reward with their income.

		Oracle TSLS		Oracle BA		TSHT			Searching		Sampling	
τ	n	Cov	Len	Cov	Len	Bias	Cov	Len	Cov	Len	Cov	Len
0.5	200	0.94	0.26	0.96	0.52	0.16	0.11	0.19	1.00	1.46	0.97	0.78
	300	0.95	0.21	0.97	0.48	0.15	0.05	0.15	1.00	1.30	0.99	0.69
	1000	0.94	0.11	0.99	0.40	0.13	0.08	0.09	1.00	0.68	1.00	0.36
	2500	0.94	0.07	0.92	0.12	0.01	0.85	0.07	1.00	0.48	1.00	0.26
1.0	200	0.94	0.26	0.98	0.82	0.23	0.14	0.19	1.00	1.42	0.97	0.83
	300	0.95	0.21	0.99	0.75	0.19	0.18	0.17	1.00	1.27	1.00	0.69
	1000	0.94	0.11	0.97	0.15	0.01	0.91	0.11	1.00	0.67	1.00	0.37
	2500	0.94	0.07	0.99	0.09	0.00	0.94	0.07	1.00	0.47	1.00	0.26
2.0	200	0.94	0.26	0.96	0.38	0.04	0.81	0.24	1.00	1.40	0.99	0.78
	300	0.95	0.21	0.97	0.27	0.02	0.87	0.20	1.00	1.27	1.00	0.70
	1000	0.94	0.11	0.96	0.12	0.01	0.93	0.11	1.00	0.68	1.00	0.37
	2500	0.94	0.07	0.96	0.07	0.00	0.94	0.07	1.00	0.48	1.00	0.26

Table E.7: High-dimensional setting with $p_z = 100$ and $p_x = 150$. The columns indexed with **Oracle TSLS**, **Oracle BA**, and **TSHT** represent the oracle TSLS CI with the knowledge of \mathcal{V} and relevant IV and covariates, the oracle bias-aware CI in (10), the CI by [25], respectively.

Variables	Statistics
Father's Education Level	
Illiteracy	1051 (27.97%)
Primary School	1098 (29.22%)
Middle School	996 (26.50%)
High School	487 (12.96%)
College and Above	126 (3.53%)
Mother's Education Level	
Illiteracy	1863 (49.57%)
Primary School	948 (25.23%)
Middle School	653 (17.38%)
High School	258 (6.87%)
College and Above	36 (0.96%)
Spouse's Education Level	
Illiteracy	347 (9.23%)
Primary School	573 (15.25%)
Middle School	1306 (34.75%)
Higher School	687 (18.28%)
College and Above	845 (22.49%)
Group-level Education Years	Mean (9.45) SD (2.47)
The Family Size	Mean (4.30) SD (1.96)
Statement about Fairness	
Strongly Disagree	49 (1.30%)
Disagree	475 (12.64%)
Neutral	2658 (70.73%)
Agree	489 (13.01%)
Strongly Agree	87 (2.32%)
Statement about Talent	
Strongly Disagree	23 (0.61%)
Disagree	494 (13.15%)
Neutral	2683 (71.39%)
Agree	432 (11.50%)
Strongly Agree	126 (3.35%)
Reading books in the past 12 months	
No	2396 (63.76%)
Yes	1362 (36.24%)
Log Education Expenditure in the past 12 months	Mean (6.20) SD (4.03)
Urban	
Rural	1400 (37.25%)
Urban	2358 (62.75%)
Hukou	
Agricultural	2445 (65.06%)
Non-agricultural	1313 (34.94%)
Gender	
Female	1425 (37.92%)
Male	2333 (62.08%)

Table E.8: The summary statistics of instrumental variables and baseline covariates.