

Semiparametric Sensitivity Analysis: Unmeasured Confounding in Observational Studies

Daniel Scharfstein

daniel.scharfstein@hsc.utah.edu
Department of Population Health Sciences
University of Utah School of Medicine
Salt Lake City, UT, USA

Razieh Nabi

rnabi@jhu.edu
Department of Computer Science
Johns Hopkins University
Baltimore, MD, USA.

Edward H. Kennedy

edward@stat.cmu.edu
Department of Statistics & Data Science
Carnegie Mellon University
Pittsburgh, PA, USA

Ming-Yueh Huang

myh0728@stat.sinica.edu.tw
Institute of Statistical Science
Academia Sinica
Nankang, Taipei, TAIWAN

Matteo Bonvini

mbonvini@stat.cmu.edu
Department of Statistics & Data Science
Carnegie Mellon University
Pittsburgh, PA, USA

Marcela Smid

marcela.smid@hsc.utah.edu
Department of Obstetrics and Gynecology
University of Utah School of Medicine
Salt Lake City, UT, USA

December 3, 2021

Abstract

Establishing cause-effect relationships from observational data often relies on untestable assumptions. It is crucial to know whether, and to what extent, the conclusions drawn from non-experimental studies are robust to potential unmeasured confounding. In this paper, we focus on the average causal effect (ACE) as our target of inference. We build on the work of Franks et al. [2019] and Robins et al. [2000] by specifying non-identified sensitivity parameters that govern a contrast between the conditional (on measured covariates) distributions of the outcome under treatment (control) between treated and untreated individuals. We use semiparametric theory to derive the non-parametric efficient influence function of the ACE, for fixed sensitivity parameters. We use this influence function to construct a one-step bias-corrected estimator of the ACE. Our estimator depends on semiparametric models for the distribution of the observed data; importantly, these models do not impose any restrictions on the values of sensitivity analysis parameters. We establish sufficient conditions ensuring that our estimator has \sqrt{n} asymptotics. We use our methodology to evaluate the causal effect of smoking during pregnancy on birth weight. We also evaluate the performance of estimation procedure in a simulation study.

1 Introduction

In causal inference, we often seek to make inferences about the population effect of a binary treatment on an outcome variable by contrasting means of potential outcomes $Y(1)$ and $Y(0)$ (i.e., counterfactuals), where $Y(t)$ represents the outcome of a random individual under treatment t , [Neyman, 1923, Rubin, 1974]. Identification of this contrast, called the *average causal effect (ACE)*, from observational data (i.e., non-experimental studies) requires untestable assumptions. Standard assumptions include:

1. *Consistency*: The observed outcome Y is equal to the potential outcome $Y(t)$ when the treatment received is t , i.e., $T = t$;
2. *Conditional ignorability*: There exists a set of measured pre-treatment covariates X such that treatment is conditionally independent of the potential outcomes given X , i.e.,

$$\{Y(1), Y(0)\} \perp T \mid X; \tag{1}$$

3. *Positivity*: For each level of the covariates X , the probability of receiving either treatment is greater than zero, i.e., $P(T = t \mid X = x) > 0$, for all x in the state space of X .

Under these assumptions, the ACE is identified from the observed data distribution via the *adjustment formula*:

$$ACE = \int_x \left\{ \mathbb{E}[Y \mid T = 1, X = x] - \mathbb{E}[Y \mid T = 0, X = x] \right\} dF(x), \tag{2}$$

where $F(\cdot)$ denotes the cumulative distribution function of X . Using n independent and identically distributed copies of $O = (X, T, Y)$, many methods have been developed to draw inference about the ACE functional, e.g., propensity score matching [Rosenbaum and Rubin, 1983b], g-computation [Robins, 1986], (stabilized) inverse probability weighting [Hernán and Robins, 2006], augmented inverse probability weighting [Robins et al., 1994], and targeted maximum likelihood [Van Der Laan and Rubin, 2006].

The conditional independence expressed in (1) implies that there are no unmeasured confounders between treatment and outcome. Assessing the robustness of inferences to potential unmeasured confounding is considered crucial. The goal of this manuscript is to provide a methodology for evaluating the sensitivity of inferences about the ACE to deviations from (1). In the next section, we provide a brief overview of the literature on sensitivity analysis in causal inference. In Section 3, we present a specific class of assumptions indexed by sensitivity analysis parameters that quantify departures from (1) and provide an identification formula for the ACE. This class of assumptions was proposed by Robins et al. [2000] and studied more recently by Franks et al. [2019]. In Section 4, we present our unique contribution, which is efficient inference for the ACE under this class of assumptions. In Section 5, we use our methodology to evaluate the causal effect of smoking during pregnancy on birth weight. Section 6 provides the results of a realistic simulation study. Section 7 is devoted to a discussion.

2 Prior Work on Sensitivity Analysis

Sensitivity analysis to the “no unmeasured confounders” assumption is designed to probe the impact of residual unmeasured confounding on causal effect estimates. One of the earliest works on sensitivity analysis is attributed to Cornfield et al. [1959]. In this work, the authors show that, for a binary unmeasured factor to explain away the observed association between smoking and lung cancer, the relative prevalence of the unmeasured factor (e.g., gene) among smokers and non-smokers must be greater than the observed relative risk of developing lung cancer for smokers vs non-smokers; since the observed relative risk was 9, they concluded that the “magnitude of the excess lung-cancer risk among cigarette smokers is so great that the results cannot be interpreted as arising from an indirect association of cigarette smoking with some other agent or characteristic, since this hypothetical agent would have to be at least as strongly associated with lung cancer as cigarette use; no such agent has been found or suggested.” The work of Cornfield et al. [1959] is not immediately relevant for our setup as it focuses on relative risks and does not incorporate measured covariates.

Recently, Ding and VanderWeele [2016] extended the approach of Cornfield et al. [1959] by allowing for low-dimensional measured covariates and introducing two sensitivity analysis parameters that govern the impact of unmeasured confounding on the outcome and treatment, respectively. They derive a bound on the relative risk in terms of the observed relative risk and the two sensitivity analysis parameters. Other sensitivity analysis methods have been developed for relative risks (see E-Appendix 4 of Ding and VanderWeele [2016]). In addition to focusing on relative risks, these methods do not accommodate complex measured confounders.

We provide a more extensive review on sensitivity analysis for the ACE of a binary treatment in the next section. Recent reviews on this topic include Liu et al. [2013] and Richardson et al. [2014]. Our review divides approaches into two types: those that seek set identification and those that seek point identification of the ACE (at each sensitivity parameter value).

2.1 Set Identification

2.1.1 Bounds without Sensitivity Parameters

If no assumptions are made on the unmeasured confounders and the outcome is bounded, the ACE can be restricted to an interval informed by the observed distribution [Robins, 1989, Manski, 1990]. The lower and upper bounds of this interval are computed under extreme instances of residual confounding. For this reason, the interval tends to be wide and necessarily includes zero. Robins [1989] and Manski [1990] derive tighter bounds by imposing additional non-identifiable assumptions.

2.1.2 Bounds with Sensitivity Parameters

To achieve better control over departures from the no-unmeasured-confounding assumption, sensitivity analysis procedures have been proposed that bound the impact of an unmeasured confounder on the treatment and/or the outcome. One way is to assume that, for units sharing the same value of measured covariates but a different value of the unmeasured confounder, the odds ratio of the

probabilities of receiving treatment differs by at most Γ . For matched studies, Section 4 of Rosenbaum [2002] describes methods to find the minimum Γ such that inference about the ACE is not “statistically significant”. Gastwirth et al. [1998] extended this idea by incorporating a bound on the impact of the unmeasured confounder on the outcome. Yadlowsky et al. [2018] extended the idea of Rosenbaum [2002] to general study designs. Shen et al. [2011] and Zhao et al. [2019] derived bounds on ACE by bounding the ratio of the probability of receiving given measured and unmeasured confounders to the probability of receiving given measured confounders.

In other work, Díaz and van der Laan [2013] and Díaz et al. [2018] derived bounds on the ACE by bounding the difference of the mean potential outcome had patients received treatment or control, given covariates, among those who actually received treatment versus control. Bonvini and Kennedy [2021] took a contamination model approach, giving bounds on the ACE by constraining the proportion of units affected by unmeasured confounding.

2.2 Point Identification

An alternative approach is to posit sensitivity analysis parameters that admit identification of the ACE. A number of authors have proposed using sensitivity analysis parameters to govern the relationship among unmeasured confounder(s), outcome and treatment. Rosenbaum and Rubin [1983a] developed a methodology that handles low-dimensional measured covariates, binary treatment, binary outcome, and a binary unmeasured confounder. This approach has been extended to accommodate normally distributed outcomes [Imbens, 2003], continuous treatments and a normally distributed unmeasured confounder [Carnegie et al., 2016], and a semiparametric Bayesian approach when the treatment and unmeasured confounder are binary [Dorie et al., 2016].

In order to avoid positing a marginal distribution for the unmeasured confounder(s), Zhang and Tchetgen Tchetgen [2019] devised a semiparametric approach to sensitivity analysis that requires models for the conditional probability of receiving treatment and the conditional mean of the outcomes; a sensitivity analysis parameter in each of these models governs the influence of the unmeasured confounder. Veitch and Zaveri [2020] circumvented the need to model the marginal distribution of the unmeasured confounders by specifying a propensity score model depending on measured and unmeasured covariates; the model is anchored at a propensity score model that is only conditional on measured confounders and is indexed by a sensitivity analysis parameter governing the influence of the unmeasured confounder. They also introduced a sensitivity analysis parameter that governs the influence of the propensity score on the conditional mean of outcome given treatment and measured covariates.

VanderWeele and Arah [2011] introduced a general “bias” formula for the difference between the possibly incorrect expression for the ACE under no unmeasured confounding and the correct expression for the ACE when accounting for both measured and unmeasured confounding in terms of many sensitivity parameters. They introduce simplifying assumptions in order to require fewer sensitivity parameters in their formalization and make it easier to use in practice. Cinelli and Hazlett [2020] focused on the linear model setting and introduced an “omitted-variable” bias formula that depends on partial R^2 values that govern the association between the unmeasured confounder, the outcome, and the treatment; these R^2 values are specified as sensitivity analysis parameters.

Brumback et al. [2004] and Robins [1999] discussed a sensitivity analysis methodology for un-

measured confounding in the setting of the time-varying treatment regimens. In the context of a point exposure, their approach is tantamount to specifying a sensitivity analysis function that governs the difference in the conditional (on measured covariates) means of the outcome under treatment (control) between treated and untreated individuals. Robins et al. [2000] and Franks et al. [2019] specified sensitivity parameters that govern a contrast between the conditional (on measured covariates) distributions of the outcome under treatment (control) between treated and untreated individuals; Robins et al. [2000] and Franks et al. [2019] adopted frequentist and Bayesian approaches to inference, respectively. An attractive feature of the approaches of Robins [1999], Brumback et al. [2004], Robins et al. [2000] and Franks et al. [2019] is that the sensitivity analysis specification does not impose any restrictions on the distribution of the observed data while yielding identification of the ACE. In this paper, we focus on the sensitivity analysis approach of Robins et al. [2000] and Franks et al. [2019], and propose an estimator for the ACE using semiparametric efficiency theory.

3 Sensitivity Analysis Model and Identification

Robins et al. [2000] and Franks et al. [2019] propose the following class of assumptions:

$$dF(Y(t)|T = 1 - t, X) = dF(Y(t)|T = t, X) \frac{\exp\{\gamma_t s_t(Y(t))\}}{C_t(\gamma_t; X)}, \quad (3)$$

where $s_t(\cdot)$ is a specified “tilting” function of its argument, γ_t is a sensitivity parameter, and $C_t(\gamma_t; X) = \mathbb{E}[\exp\{\gamma_t s_t(Y(t))\} | T = t, X]$. The choice of γ_t and $s_t(\cdot)$ allows the user to flexibly model the unobserved density $dF(Y(t) | T = 1 - t, X)$ through a modification of the observed density $dF(Y(t) | T = t, X)$, as opposed to just assuming that all confounders have been measured so that the two densities are equal. The case $\gamma_t = 0$ corresponds to the no unmeasured confounding assumption. For fixed γ_t , it can be shown that $E[Y(t)]$ is identified via the following formula:

$$\begin{aligned} \mathbb{E}[Y(t)] &= \int_x \left\{ \mathbb{E}[Y | T = t, X = x] \times P(T = t | X = x) + \right. \\ &\quad \left. \frac{\mathbb{E}[Y \exp\{\gamma_t s_t(Y)\} | T = t, X = x]}{\mathbb{E}[\exp\{\gamma_t s_t(Y)\} | T = t, X = x]} \times P(T = 1 - t | X = x) \right\} dF(x). \end{aligned} \quad (4)$$

Then, the ACE is identified by the following formula:

$$\begin{aligned} ACE &= \int_x \left\{ \mathbb{E}[Y | T = 1, X = x] \times P(T = 1 | X = x) + \right. \\ &\quad \left. \frac{\mathbb{E}[Y \exp\{\gamma_1 s_1(Y)\} | T = 1, X = x]}{\mathbb{E}[\exp\{\gamma_1 s_1(Y)\} | T = 1, X = x]} \times P(T = 0 | X = x) \right\} dF(x) - \\ &\quad \int_x \left\{ \mathbb{E}[Y | T = 0, X = x] \times P(T = 0 | X = x) + \right. \\ &\quad \left. \frac{\mathbb{E}[Y \exp\{\gamma_0 s_0(Y)\} | T = 0, X = x]}{\mathbb{E}[\exp\{\gamma_0 s_0(Y)\} | T = 0, X = x]} \times P(T = 1 | X = x) \right\} dF(x). \end{aligned} \quad (5)$$

Note that when $\gamma_1 = \gamma_0 = 0$, (5) reduces to (2).

Assumption (3) can be re-written, using Bayes rule, as

$$\text{logit}\{P(T = 1 - t | X, Y(t))\} = h(X; \gamma_t) + \gamma_t s_t(Y(t)), \quad (6)$$

where $h(X; \gamma_t) = \text{logit}\{P(T = 1 - t | X)\} - \log\{C_t(\gamma_t; X)\}$. This representation of Assumption (3) shows that $\exp(-\gamma_t)$ can be interpreted as the conditional (on X) odds ratio of receiving treatment $1 - t$ for patients who differ by one unit in $s_t(Y(t))$. For example, if $Y(t)$ is binary and $s_t(Y(t) = Y(t))$, then $\exp(-\gamma_t)$ is the conditional (on X) odds ratio of receiving treatment $1 - t$ for patients with $Y(t) = 1$ versus those with $Y(t) = 0$.

4 Inference

We derive efficient estimators for $\mathbb{E}[Y(t)]$ for $t = 0, 1$ and then combine the resulting estimators to draw inference about the ACE. Our estimation strategy is based on semiparametric efficiency theory. In what follows, we let P denote the true distribution of the observed data O and define our target of inference as $\psi_t(P) := \mathbb{E}[Y(t)]$.

Theorem 1. Non-Parametric Efficient Influence Function

Given the target of inference $\mathbb{E}[Y(t)]$ in (4), the corresponding non-parametric efficient influence function, denoted by $\phi_t(P)(O)$, is of the form:

$$\begin{aligned} \phi_t(P)(O) &= \mathbb{I}(T = t) \left\{ Y + Y \times \frac{P(T = 1 - t | X)}{P(T = t | X)} \times \frac{\exp\{\gamma_t s_t(Y)\}}{\mathbb{E}[\exp\{\gamma_t s_t(Y)\} | T = t, X]} - \right. \\ &\quad \left. \frac{P(T = 1 - t | X)}{P(T = t | X)} \times \frac{\exp\{\gamma_t s_t(Y)\} \times \mathbb{E}[Y \exp\{\gamma_t s_t(Y)\} | T = t, X]}{\mathbb{E}^2[\exp\{\gamma_t s_t(Y)\} | T = t, X]} \right\} \\ &\quad + \mathbb{I}(T = 1 - t) \times \frac{\mathbb{E}[Y \exp\{\gamma_t s_t(Y)\} | T = t, X]}{\mathbb{E}[\exp\{\gamma_t s_t(Y)\} | T = t, X]} - \psi_t(P) \end{aligned} \quad (7)$$

Proof. See Appendix A. □

4.1 Inferential Strategy

In this section, we outline the approach we take to estimation and inference, which comes from semiparametric theory and has recently been termed double machine learning [Bickel et al., 1993, Robins and van der Laan, 2003, Tsiatis, 2006]. Consider a statistical model \mathcal{M} composed of distributions \tilde{P} , with P denoting the true distribution. Define the functional $\psi_t : \tilde{P} \rightarrow \mathbb{R}$. Under sufficient smoothness conditions, we can write down the following von Mises expansion for $\psi_t(\tilde{P})$,

$$\psi_t(\tilde{P}) = \psi_t(P) - \int \phi_t(\tilde{P})(o) dP(o) + \text{Rem}_t(\tilde{P}, P), \quad (8)$$

where $\phi_t(\tilde{P})$ is influence function corresponding to $\psi_t(\tilde{P})$ and satisfies $\int \phi_t(P)(o) dP(o) = 0$. The last term in (8) is a *second-order* remainder term such that $\text{Rem}_t(\tilde{P}, P) = \mathcal{O}(\|\tilde{P} - P\|^2)$, see [Bickel et al., 1993] for more details. This distributional Taylor expansion is, under regularity

conditions, equivalent to pathwise differentiability [Robins and van der Laan, 2003, Kennedy et al., 2021]. If we estimate P by \hat{P} , the *first-order* bias of the corresponding estimator $\psi_t(\hat{P})$ is equal to

$$\psi_t(\hat{P}) - \psi_t(P) = \underbrace{- \int \phi_t(\hat{P})(o) dP(o)}_{\text{Bias}_t} + \underbrace{\text{Rem}_t(\hat{P}, P)}_{\mathcal{O}(\|\hat{P}-P\|^2)}. \quad (9)$$

The *bias* can be estimated using the empirical distribution, P_n , in place of P ,

$$\widehat{\text{Bias}}_t = - \int \phi_t(\hat{P})(o) dP_n(o).$$

This suggests a corrected plug-in estimator (one-step estimator) of the form:

$$\hat{\psi}_t = \psi_t(\hat{P}) - \widehat{\text{Bias}}_t = \psi_t(\hat{P}) + \int \phi_t(\hat{P})(o) dP_n(o). \quad (10)$$

In order to analyze the asymptotic properties of $\hat{\psi}_t$, we need to understand the behavior of $\sqrt{n}(\hat{\psi}_t - \psi_t)$ as $n \rightarrow \infty$, where $\psi_t := \psi_t(P)$. Note that

$$\begin{aligned} \hat{\psi}_t - \psi_t &= \left\{ \psi_t(\hat{P}) + \int \phi_t(\hat{P})(o) dP_n(o) \right\} - \psi_t \\ &= \left\{ \psi_t(\hat{P}) - \psi_t \right\} + \int \phi_t(\hat{P})(o) dP_n(o) \\ &= \left\{ - \int \phi_t(\hat{P})(o) dP(o) + \text{Rem}_t(\hat{P}, P) \right\} + \int \phi_t(\hat{P})(o) dP_n(o) \\ &= \int \phi_t(\hat{P})(o) d\{P_n(o) - P(o)\} + \text{Rem}_t(\hat{P}, P) \\ &= \int \phi_t(P)(o) dP_n(o) + \int \{\phi_t(\hat{P})(o) - \phi_t(P)(o)\} d\{P_n(o) - P(o)\} + \text{Rem}_t(\hat{P}, P), \end{aligned} \quad (11)$$

where the third equality follows by (9). Therefore $\sqrt{n}(\hat{\psi}_t - \psi_t)$ can be expressed as:

$$\begin{aligned} \sqrt{n}(\hat{\psi}_t - \psi_t) &= \\ &= \underbrace{\sqrt{n} \int \phi_t(P)(o) dP_n(o)}_{CLT} + \underbrace{\sqrt{n} \int \{\phi_t(\hat{P})(o) - \phi_t(P)(o)\} d\{P_n(o) - P(o)\}}_{\text{Empirical Process}} + \underbrace{\sqrt{n} \text{Rem}_t(\hat{P}, P)}_{\text{Second-order remainder}}. \end{aligned} \quad (12)$$

By the central limit theorem, the first term on the right hand side of (12) is asymptotically normal. Therefore, if the second and third terms are $o_p(1)$, then $\sqrt{n}(\hat{\psi}_t - \psi_t) \xrightarrow{D(P)} \mathcal{N}(0, \mathbb{E}[\phi_t(P)^2])$. The second term is a centered empirical process, and under certain Donsker conditions will be $o_p(1)$. In Subsection 4.1.2, we use sample-splitting procedure to assure that the second term is $o_p(1)$, even if Donsker conditions are not met. The third term is a second-order remainder term and usually needs to be studied on a case-by-case basis.

4.1.1 Second Order Remainder Term

In this subsection, we characterize the second-order remainder error described in the previous section (essentially the estimation bias of the one-step estimator) and give sufficient conditions

under which it is asymptotically negligible. For concreteness, we focus on nuisance estimators built from generalized additive and single index models, but the results hold more generally for any $n^{-1/4}$ consistent estimators.

Lemma 1. *Given the target of inference $\mathbb{E}[Y(t)]$ in (4) and its corresponding influence function $\phi_t(P)$ in Theorem 1, $\text{Rem}_t(\tilde{P}, P)$, given in (8), is equal to*

$$\begin{aligned} & \text{Rem}_t(\tilde{P}, P) \\ &= \mathbb{E} \left[\frac{\mu_t(Y \exp(\gamma_t s_t(Y)); X) \tilde{\mu}_t(\exp(\gamma_t s_t(Y)); X) - \tilde{\mu}_t(Y \exp(\gamma_t s_t(Y)); X) \mu_t(\exp(\gamma_t s_t(Y)); X)}{\tilde{\pi}_t(X) \tilde{\mu}_t(\exp(\gamma_t s_t(Y)); X)} \right. \\ & \quad \left. \times \left\{ \frac{\tilde{\pi}_{1-t}(X) \pi_t(X)}{\tilde{\mu}_t(\exp(\gamma_t s_t(Y)); X)} - \frac{\pi_{1-t}(X) \tilde{\pi}_t(X)}{\mu_t(\exp(\gamma_t s_t(Y)); X)} \right\} \right], \end{aligned}$$

where $\tilde{\pi}_t(X) = \tilde{P}(T = t \mid X)$, $\tilde{\mu}_t(g(y); X) = \mathbb{E}_{\tilde{P}}[g(Y) \mid T = t, X]$, $\pi_t(X) = P(T = t \mid X)$, and $\mu_t(g(y); X) = \mathbb{E}_P[g(Y) \mid T = t, X]$.

Proof. See Appendix B. □

Let $\hat{\pi}_t(X)$, $\hat{\mu}_t(\exp(\gamma_t s_t(Y)); X)$ and $\hat{\mu}_t(Y \exp(\gamma_t s_t(Y)); X)$ be estimators of $\pi_t(X)$, $\mu_t(\exp(\gamma_t s_t(Y)); X)$ and $\mu_t(Y \exp(\gamma_t s_t(Y)); X)$, respectively. If $\pi_t(X)$ and $\mu_t(\exp(\gamma_t s_t(Y)); X)$ are bounded away from zero with probability one, Lemma 1 can be used to show that,

$$\begin{aligned} & |\text{Rem}_t(\hat{P}, P)| \\ & \leq \hat{C}_1 \times \left\| \hat{\mu}_t(\exp(\gamma_t s_t(Y)); X) - \mu_t(\exp(\gamma_t s_t(Y)); X) \right\|_{L_2} \times \\ & \quad \left\| \hat{\mu}_t(Y \exp(\gamma_t s_t(Y)); X) - \mu_t(Y \exp(\gamma_t s_t(Y)); X) \right\|_{L_2} + \\ & \hat{C}_2 \times \left\| \hat{\mu}_t(\exp(\gamma_t s_t(Y)); X) - \mu_t(\exp(\gamma_t s_t(Y)); X) \right\|_{L_2} \times \left\| \hat{\pi}_t(X) - \pi_t(X) \right\|_{L_2} + \\ & \hat{C}_3 \times \left\| \hat{\mu}_t(Y \exp(\gamma_t s_t(Y)); X) - \mu_t(Y \exp(\gamma_t s_t(Y)); X) \right\|_{L_2} \times \left\| \hat{\pi}_t(X) - \pi_t(X) \right\|_{L_2} + \\ & \hat{C}_4 \times \left\{ \left\| \hat{\mu}_t(\exp(\gamma_t s_t(Y)); X) - \mu_t(\exp(\gamma_t s_t(Y)); X) \right\|_{L_2} \right\}^2, \end{aligned} \tag{13}$$

where $\hat{C}_1, \dots, \hat{C}_4$ are $\mathcal{O}_p(1)$, and $\|f\|_{L_2} := \sqrt{\mathbb{E}[f^2]}$. Whether $\sqrt{n} |\text{Rem}_t(\hat{P}, P)|$ is $o_p(1)$ relies on rates of convergence for the models involved in (13).

We estimate $\pi_t(X)$ using a generalized additive model (GAM). Let $\hat{\pi}_t(X)$ be the GAM estimator of $\pi_t(X)$. Horowitz et al. [2004] showed that

$$\left\| \hat{\pi}_t(X) - \pi_t(X) \right\|_{L_2} = O_P(n^{-2/5}).$$

Note that $\mu_t(Y \exp(\gamma_t s_t(Y)); X)$ and $\mu_t(\exp(\gamma_t s_t(Y)); X)$ are both defined in terms of the conditional distribution of Y given $T = t$ and X . We will use the single-index model [Chiang and

Huang, 2012] for the conditional cumulative function of Y given $T = t$ and $X = x$. This model posits that

$$P(Y \leq y \mid T = t, X) = F_t(y, X' \beta_t; \beta_t),$$

where $F_t(y, u; \beta_t)$ is a cumulative distribution function in y for each u , $\beta_t = (\beta_{1t}, \dots, \beta_{pt})$ is a vector of unknown parameters and, for purposes of identifiability, β_{1t} is set to 1. We estimate $F_t(y, x' \beta_t; \beta_t)$ by

$$\widehat{F}_t(y, X' \widehat{\beta}_t; \widehat{\beta}_t) = \frac{\sum_{T_i=t} \mathbb{I}(Y_i \leq y) K_{\widehat{h}_t}(X'_i \widehat{\beta}_t - X' \widehat{\beta}_t)}{\sum_{T_i=t} K_{\widehat{h}_t}(X'_i \widehat{\beta}_t - X' \widehat{\beta}_t)},$$

where $K_h(v) = K(v/h)/h$, K is a q th-order kernel, $\widehat{\beta}_t$ is an estimator of β_t and \widehat{h}_t is an estimator of the bandwidth h (Note: $\widehat{\beta}_t$ are computed using a cross-validation procedure described in Appendix C, and the choice of q and \widehat{h}_t will be discussed later). We estimate $\mu_t(Y \exp(\gamma_t s_t(Y)); X)$ and $\mu_t(\exp(\gamma_t s_t(Y)); X)$ by

$$\widehat{\mu}_t(Y \exp(\gamma_t s_t(Y)); X) = \int y \exp(\gamma_t s_t(y)) d\widehat{F}_t(y, X' \widehat{\beta}_t; \widehat{\beta}_t), \text{ and} \quad (14)$$

$$\widehat{\mu}_t(\exp(\gamma_t s_t(Y)); X) = \int \exp(\gamma_t s_t(y)) d\widehat{F}_t(y, X' \widehat{\beta}_t; \widehat{\beta}_t). \quad (15)$$

The single-index model fitting used for estimating $\mu_t(Y \exp(\gamma_t s_t(Y)); X)$ and $\mu_t(\exp(\gamma_t s_t(Y)); X)$ involves two-stage kernel smoothing. In the first stage, we estimate β_t through the criterion of Chiang and Huang [2012] with a fourth-order kernel function, and obtain a $n^{1/2}$ -consistent estimator $\widehat{\beta}_t$ and an optimal bandwidth \widetilde{h}_t of rate $O_P(n^{-1/9})$. Second, with $\widehat{\beta}_t$, a second-order kernel function, and another adjusted bandwidth $\widehat{h} = \widetilde{h} n^{-4/45}$, we estimate $\mu_t(Y \exp(\gamma_t s_t(Y)); X)$ and $\mu_t(\exp(\gamma_t s_t(Y)); X)$ through (14)–(15).

Given that $\left\| \widehat{\pi}_t(X) - \pi_t(X) \right\|_{L_2} = O_P(n^{-2/5})$, we know that $\sqrt{n} |\text{Rem}_t(\widehat{P}, P)|$ will be $o_P(1)$ if $\left\| \widehat{\mu}_t(\exp(\gamma_t s_t(Y)); X) - \mu_t(\exp(\gamma_t s_t(Y)); X) \right\|_{L_2} = o_P(n^{-1/4})$ and $\left\| \widehat{\mu}_t(Y \exp(\gamma_t s_t(Y)); X) - \mu_t(Y \exp(\gamma_t s_t(Y)); X) \right\|_{L_2} = O_P(n^{-1/4})$. In Appendix C, we prove that this will be case if $\widehat{\beta}_t$ is $n^{1/2}$ -consistent, the order of the kernel is greater than $1/2$, and the bandwidth is of $O_P(n^{-1/(2q+1)})$. The choice of the order of the kernel q implicitly encodes a smoothness condition on $F_t(y, u; \beta_t)$ as q is such that the $(q+1)^{\text{th}}$ derivative of $F_t(y, u; \beta_t)$ with respect to u is Lipschitz continuous, with Lipschitz constant independent of (y, β_t) . In our data analysis and simulation study, we use $q = 2$ and $\widehat{h}_t = \widetilde{h}_t n^{-4/45}$. With this choice, $\left\| \widehat{\mu}_t(\exp(\gamma_t s_t(Y)); X) - \mu_t(\exp(\gamma_t s_t(Y)); X) \right\|_{L_2}$ and $\left\| \widehat{\mu}_t(Y \exp(\gamma_t s_t(Y)); X) - \mu_t(Y \exp(\gamma_t s_t(Y)); X) \right\|_{L_2}$ are $O_P(n^{-2/5})$, which satisfies the needed rates of convergence.

4.1.2 Empirical Process Term

Here, we discuss conditions under which the empirical process term in the earlier decomposition is asymptotically negligible, and give our final result regarding the *sqrtn* consistency and asymptotic normality of the proposed estimator.

As discussed above, the second term in (12), i.e.,

$$\sqrt{n} \int \{\phi_t(\hat{P})(o) - \phi_t(P)(o)\} d\{P_n(o) - P(o)\},$$

is $o_p(1)$, if $\phi_t(P)$ belongs to a Donsker class. A more convenient way to ensure that this term disappears in large sample sizes, is using *sample splitting* in order to separate \hat{P} and P_n and avoid overfitting. The idea is to randomly split the observations into K disjoint sets. Let S_i denote the split membership of the i th observation (i.e, $S_i \in \{1, \dots, K\}$). Let $\hat{P}^{(-k)}$ denote the estimator of P using observations from all the splits except that of k th split, and $P_{n_k}^{(k)}$ be the empirical distribution based on the $n_k (\approx n/K)$ observations in the k th split. The sample splitting estimator of ψ_t is

$$\tilde{\psi}_t = \frac{1}{K} \sum_{k=1}^K \hat{\psi}_t^{(k)}, \quad (16)$$

where $\hat{\psi}_t^{(k)} = \int \nu_t(\hat{P}^{(-k)})(o) dP_{n_k}^{(k)}(o) = \frac{1}{n_k} \sum_{i: S_i=k} \nu_t(\hat{P}^{(-k)})(O_i)$ and $\nu_t(P)(O) = \phi_t(P)(O) + \psi_t(P)$.

In order to understand the asymptotic properties of the new estimator $\tilde{\psi}_t$, we follow similar steps as above to write

$$\sqrt{n}(\tilde{\psi}_t - \psi_t) = \sqrt{n} \int \phi_t(P)(o) dP_n(o) + \frac{1}{\sqrt{K}} \sum_{k=1}^K \sqrt{n_k} \left(R_{t,k}^{(1)} + R_{t,k}^{(2)} \right), \quad (17)$$

where $R_{t,k}^{(1)} = \int \left\{ \phi_t(\hat{P}^{(-k)})(o) - \phi_t(P)(o) \right\} d\{P_{n_k}^{(k)}(o) - P(o)\}$ and $R_{t,k}^{(2)} = \text{Rem}_t(\hat{P}^{(-k)}, P)$.

Replacing \hat{P} with $\hat{P}^{(-k)}$ in the arguments made in the previous subsection, it can be easily shown that $\sqrt{n_k} |R_{t,k}^{(2)}|$ will be $o_p(1)$. It is useful to note that $R_{t,k}^{(1)} = \int \left\{ \nu_t(\hat{P}^{(-k)})(o) - \nu_t(P)(o) \right\} d\{P_{n_k}^{(k)}(o) - P(o)\}$, where $\nu_t(P)(o) = \phi_t(P)(o) + \psi_t(P)$. Kennedy et al. [2020] showed that $\sqrt{n_k} |R_{t,k}^{(1)}| = O_P\left(\|\nu_t(\hat{P}^{(-k)}) - \nu_t(P)\|_{L_2}\right)$. To show $\sqrt{n_k} |R_{t,k}^{(1)}|$ is $o_p(1)$, it is sufficient to prove, as we do in the following lemma, that $\|\nu_t(\hat{P}^{(-k)}) - \nu_t(P)\|_{L_2}$ is $o_p(1)$.

Lemma 2. *Given the target of inference $\mathbb{E}[Y(t)]$ in (4) and its corresponding sample-splitting estimator in (16), we have that $\sqrt{n_k} |R_{1k}| = o_p(1)$.*

Proof. See Appendix D. □

Now, we are in the position to establish the asymptotic normality of the sample splitting estimator.

Theorem 2. *Assume the following conditions hold:*

1. $|Y|$ and $|\exp(\gamma_t s_t(Y))|$ are bounded in probability
2. $P(\epsilon < \hat{\pi}_t(X) < 1 - \epsilon) = 1$ for some $\epsilon > 0$;
3. $\|\phi_t(\hat{P}) - \phi_t(P)\|_{L_2} = o_p(1)$;

4. $|Rem_t(\hat{P}, P)| = o_P(n^{-1/2})$ for $|Rem_t(\hat{P}, P)|$ defined in (13).

Given the target of inference $\mathbb{E}[Y(t)]$ in (4) and its corresponding sample-splitting estimator in (16), we have that $\sqrt{n}(\tilde{\psi}_t - \psi_t) \xrightarrow{D(P)} \mathcal{N}(0, \mathbb{E}[\phi_t(P)^2])$.

Proof. The second term on the right hand side of (17) is $o_P(1)$ because it is a constant times the finite sum of $o_P(1)$ terms. By the central limit theorem, the first term on the right hand side of (17) converges to a normal distribution with mean zero and variance $\mathbb{E}[\phi_t(P)^2]$. \square

Condition 1 is a mild boundedness assumption that we rely on in the proof of Lemma 2. Condition 2 requires that the estimated propensity score is bounded away from 0 and 1. Condition 3 is a mild consistency condition requiring that $\phi_t(\hat{P})$ converges to the truth in L_2 at any rate. Finally, condition 4 is the more stringent condition, but it still allows estimation in flexible nonparametric models. In particular, we have shown that this condition is satisfied if $\pi_t(X)$ follows a GAM model and $P(Y \leq y | T = t, X)$ follows a single-index model.

The asymptotic variance of the sample-splitting estimator can be consistently estimated by $\frac{1}{n} \sum_{k=1}^K \sum_{i:S_i=k} \phi_t(\hat{P}^{(-k)})(O_i)^2$. ACE can be estimated by $\tilde{\psi}_1 - \tilde{\psi}_0$. The asymptotic variance can be consistently estimated by $\frac{1}{n} \sum_{k=1}^K \sum_{i:S_i=k} \{\phi_1(\hat{P}^{(-k)})(O_i) - \phi_0(\hat{P}^{(-k)})(O_i)\}^2$.

5 Implementation Details

In executing our methodological approach, there are two main issues that require attention. First, as with all augmented inverse weighted estimation procedures, there can be some numerical instability due to small predicted probabilities of treatment received. To address this issue, we use the tuning-free Huberization procedure developed by Wang et al. [2012]. In this approach, the k th split estimator to be taken to be

$$\hat{\psi}_t^{(k)} = \frac{1}{n_k} \sum_{i:S_i=k} \min \left\{ |\nu_t(\hat{P}^{(-k)})(O_i)|, \tau_t^{(k)} \right\} \text{sign} \left\{ \nu_t(\hat{P}^{(-k)})(O_i) \right\},$$

where $\tau_t^{(k)}$ is the non-negative solution to

$$\sum_{i:S_i=k} \frac{\min \left\{ \nu_t(\hat{P}^{(-k)})(O_i)^2, \left(\tau_t^{(k)} \right)^2 \right\}}{\left(\tau_t^{(k)} \right)^2} = \log(n_k).$$

This procedure effectively truncates $\nu_t(\hat{P}^{(-k)})(O_i)$ to be in $[-\tau_t^{(k)}, \tau_t^{(k)}]$.

Second, we have found that, depending on the value of the sensitivity analysis parameters, coverage of 95% normality-based and percentile bootstrap confidence intervals can have coverage below nominal levels. To address this problem, we suggest using symmetric-t percentile bootstrap confidence intervals that are calibrated using a nested level of re-sampling; this is called iterated or double bootstrap [Hall, 1986, Efron and Tibshirani, 1994, Tu and Shao, 1995, Beran, 1987, Hall and Martin, 1988]. In our data analysis and simulated study, we used 250 re-samples are the first and second levels of the double bootstrap.

6 Data Analysis

Almond et al. [2005] and Cattaneo [2010] conducted an analysis to evaluate the causal effect of smoking during pregnancy on birth weight. Their analyses were based on data on singleton births of approximately 500,000 births in Pennsylvania between 1989 and 1991 and assumed no unmeasured confounding. After adjustment for measured “pretreatment variables” in a regression model, Almond et al. [2005] reported a reduction of 203.2 grams in birthweight for smokers versus non-smokers¹. Cattaneo [2010] analyzed smoking status as a six-level treatment variable and adjusted for the same covariates as Almond et al. [2005]. They concluded that “there is a large reduction of about 150 grams when the mother starts to smoke (1-5 cigarettes), an additional reduction of approximately 70 grams when changing from 1-5 to 6-10 cigarettes-per-day, and no additional effects once the mother smokes at least 11 cigarettes”. Key factors that have not been controlled for in their analyses are maternal nutrition, social determinants of health, use of substances other than alcohol, genetics and epigenetics.

We analyzed a sample of 4,996 of these births that was available online.² In our analysis, we accounted for the following maternal covariates: (1) age, (2) education (less than high school, high school, greater than high school), (3) white (yes/no), (4) hispanic (yes/no), (5) foreign (yes/no), (6) alcohol use, (7) married (yes/no), (8) liver birth order (one, two, greater than 2), (9) number of prenatal visits.³ Columns 2 and 3 of Table 1 shows summary statistics for these covariates, stratified by maternal smoking status. There are striking differences between smokers and non-smokers, especially with respect to education, marital status, alcohol use and live birth order.

Columns 4-6 of Table 1 provides an assessment of goodness-of-fit of our models for the observed data. To evaluate goodness of fit, we simulated two large datasets, each comprised of observed data on 100,000 individuals. Both datasets were generated using the empirical distribution of covariates. The first dataset (“semiparametric”) is generated using the estimated fits of the GAM and single-index models. The second dataset (“parametric”) is generated using the estimated fits of a logistic regression model for the probability of $T = 1$ given X and normal regression models for conditional distribution of Y given $T = t$ and X . We computed empirical distributions for various subgroups (e.g., smokers aged 13-22) based on the original (“nonparametric”) dataset, the semiparametric dataset and the parametric dataset. For each subgroup, we then computed Kolmogorov-Smirnov statistics of differences between the distributions estimated from the nonparametric dataset and the semiparametric/parametric datasets. Using this metric, our analysis demonstrates that our semiparametric model provides a better fit to the observed data than the parametric model for most subgroups considered. With that said, there are some sizable subgroups where the Kolmogorov-Smirnov statistics for the semiparametric model are sub-optimal (≈ 0.10).

¹The following variables were included: mother’s age, education, race, ethnicity, marital status, foreign-born status; father’s age, education, race, and ethnicity; dummies for trimester of first prenatal care visit, number of prenatal visits, and adequacy of care; controls for alcohol use and number of drinks per week; pregnancy history variables (parity indicator, live birth-order, number of previous births where newborn died, interval since last live birth, indicators for previous birth over 4000 grams and previous birth preterm or small-for-gestational-age); maternal medical risk factors that are not believed to be affected by smoking during pregnancy (anemia, cardiac disease, lung disease, diabetes, genital herpes, hyrdamnios/oligohydramnios, hemoglobinopathy, chronic hypertension, eclampsia, incompetent cervix, renal disease, Rh sensitization, uterine bleeding); month of birth and county of residence indicators.

²https://github.com/mdcattaneo/replication-C_2010_JOE. The dataset has information on 5,000 individuals. We removed four individuals whose live birth order was zero.

³This is a subset of the variables used by Almond et al. [2005] and Cattaneo [2010]. Alcohol use and number of prenatal visits are considered a proxy for maternal health seeking behavior.

	Smoker $n = 942$	Non-smoker $n = 4054$	K-S Statistic		K-S Statistic	
			Cond. Prob. of Smoker (GAM/LR)	Cond. Dist. of Birthweight Smokers (SI/Normal)	Cond. Prob. of Smoker (GAM/LR)	Cond. Dist. of Birthweight Non-Smokers (SI/Normal)
Age (Mean/IQR)	25.4/8.0	26.9/8.0				
13 – 22 (%)	32.5	23.6	0.001/0.02	0.04/0.06	0.08/0.07	
23 – 27 (%)	32.7	28.8	0.000/0.03	0.05/0.08	0.03/0.09	
28 – 31 (%)	20.9	26.3	0.001/0.004	0.05/0.06	0.02/0.11	
31 – 45 (%)	13.9	21.1	0.002/0.02	0.09/0.12	0.03/0.12	
Education (%)						
Less than HS (%)	32.5	14.6	0.004/0.006	0.07/0.06	0.04/0.08	
HS (%)	50.8	43.3	0.0001/0.002	0.02/0.07	0.04/0.08	
Greater than HS (%)	16.7	42.1	0.001/0.001	0.05/0.06	0.02/0.11	
Race (%)						
White (%)	79.6	84.0	0.001/0.003	0.02/0.06	0.02/0.12	
Non-White (%)	20.4	16.0	0.001/0.002	0.10/0.06	0.10/0.15	
Ethnicity (%)						
Hispanic (%)	3.0	4.0	0.007/0.006	0.11/0.15	0.07/0.10	
Non-Hispanic (%)	97.0	96.0	0.001/0.001	0.01/0.05	0.03/0.08	
Foreign (%)						
Yes (%)	2.7	6.2	0.002/0.0005	0.19/0.17	0.06/0.08	
No (%)	97.3	93.8	0.001/0.001	0.01/0.05	0.03/0.08	
Alcohol (%)						
Yes (%)	9.6	1.9	0.006/0.002	0.10/0.08	0.05/0.10	
No (%)	90.4	98.1	0.001/0.002	0.01/0.06	0.03/0.08	
Marital Status (%)						
Married (%)	47.3	74.6	0.002/0.001	0.05/0.06	0.09/0.08	
Not Married (%)	52.7	36.4	0.001/0.002	0.04/0.07	0.02/0.12	
Live Birth Order (%)						
One	34.3	43.0	0.001/0.0002	0.05/0.08	0.06/0.04	
Two	32.8	31.9	0.003/0.003	0.04/0.06	0.03/0.12	
Greater than Two	32.9	25.1	0.003/0.003	0.04/0.07	0.03/0.12	
Prenatal Visits (Mean/IQR)	9.7 (5.0)	10.8 (4.0)				
0 – 9 (%)	41.0	28.4	0.003/0.01	0.05/0.06	0.09/0.09	
10 – 11 (%)	22.4	23.9	0.0003/0.002	0.07/0.10	0.05/0.08	
12 – 13 (%)	22.5	29.6	0.003/0.003	0.07/0.10	0.05/0.15	
14 – 49 (%)	14.1	18.2	0.003/0.003	0.07/0.13	0.04/0.17	

Table 1: Columns 2 and 3: Distribution of covariates by smoking status; Columns 4: Kolmogorov-Smirnov statistics for non-parametric vs model-based estimates (GAM/Logistic Regression) of conditional probability of smoker. Columns 5 and 6: Kolmogorov-Smirnov statistics for non-parametric vs. model-based estimates (Single Index/Normal) of conditional distribution of birthweight for smokers and non-smokers.




Figure 1: Distribution of observed birthweight, by smoking status. Dash lines represent observed means.


Figure 1 shows the distribution functions of birth weight for smokers and non-smokers. The average birth weight is 3133 grams and 3412 grams for smokers and non-smokers, respectively. The naive estimated difference (smokers minus non-smokers) is -278 grams (95% CI: -319 to -238). After adjustment for measured covariates listed in Table 1, the estimated effect of smoking on birth weight is -223 grams (95% CI: -274 to -172).

Figure 2 displays our choices for $s_1(Y(1))$ and $s_0(Y(0))$. Our choice for $s_1(Y(1))$ is motivated by the clinical belief that, within levels of measured covariates, the distribution of birth weight under smoking for non-smokers will be concentrated at “better” values than the distribution of birth weight under smoking for smokers. Here “better” means values between 2,500 and 4,000 grams; these are the thresholds for low and high birth weight, respectively. This is accomplished with the tilting function $s_1(y) = y * \Phi((4000 - y)/200) + (4000 - y) * (1 - \Phi((4000 - y)/200))^4$ coupled with a positive value of γ_1 . Our choice for $s_0(Y(0))$ is motivated by the clinical belief that, within levels of measured covariates, the distribution of birth weight under not smoking for smokers will be concentrated at lower values than the distribution of birth weight under not smoking for non-smokers. This is accomplished with the tilting function $s_0(Y(0)) = y * \Phi((y - 2000)/2000)$ coupled with negative values for γ_0 .


Figure 3 displays the estimated means (solid lines) of $Y(1)$ and $Y(0)$ as a function of γ_1 and γ_0 , respectively. The figures include pointwise 95% confidence intervals (dashed lines). The range of γ_1 and γ_0 were selected to be wide enough to include extreme values, as we will discuss below. Figure 5 display a contour plot of the difference between the average causal effect, $E[Y(1)] - E[Y(0)]$ as a function of the sensitivity parameters γ_1 and γ_0 . The region on the top left denoted as “Smoking Worse” has 95% confidence intervals where the upper limit is negative and the region on the bottom right denoted as “Smoking Better” has 95% confidence intervals where the lower limit is positive.

To understand the choice of sensitivity parameters, consider Figure 4. In this figure, we present

⁴ $\Phi()$ is the cumulative distribution function of a standard normal random variable.




(a) Smoking




(b) Non-Smoking

Figure 2: Tilting functions (a) $s_1(Y(1))$ and (b) $s_0(Y(0))$




(a) Smoking



(b) Non-Smoking

Figure 3: Estimated means (solid lines) of (a) $Y(1)$ and (b) $Y(0)$ as a function of γ_1 and γ_0 , respectively. Dashed lines represent 95% pointwise confidence intervals.



(a) Mean of $Y(1)$ among non-smokers

(b) Mean of $Y(0)$ among smokers

Figure 4: Induced estimated means of $Y(t)$ given $T = 1 - t$ as function of γ_t .

the induced estimated mean of $Y(t)$ given $T = 1 - t$ as a function of γ_t . For fixed γ_t , the induced estimated mean of $Y(t)$ given $T = 1 - t$ is computed as $(\tilde{\psi}_t - E_n[Y|T = t]P_n[T = t])/P_n[T = 1 - t]$, where $E_n[Y|T = t]$ is the observed mean birth weight among individuals with $T = t$ and $P_n[T = t']$ is the observed proportion of individuals with smoking status t' . Figure 4(a) focuses on the induced mean of $Y(1)$ among non-smokers. It makes clinical sense that $E[Y(1)|T = 0] \leq E[Y(0)|T = 0]$. This would restrict the value of $\gamma_1 \leq 0.003$. Figure 4(b) focuses on the induced mean of $Y(0)$ among smokers. Here, it makes clinical sense that $E[Y(0)|T = 1] \geq E[Y(1)|T = 1]$. This would restrict the value of $\gamma_0 \geq -0.006$. Constraining γ_1 and γ_0 to this region, we see, in Figure 5, there are values of sensitivity parameters that lead to inferences that are consistent with no effect of smoking on birth weight. However, there is a strong biological rationale for a detrimental effect of smoking on birth weight. Nicotine and the other substances involved in tobacco combustion (e.g., carbon monoxide) have been shown to impair fetal oxygen delivery and blood flow which leads to fetal growth restriction and low birth weight [Burton et al., 1989, Larsen et al., 2002, Bush et al., 2000, Lehtovirta and Forss, 1978, Caravati et al., 1988, Gozubuyuk et al., 2017, Albuquerque et al., 2004]. While there is strong reason to believe that there is negative impact of maternal smoking on birthweight, the average causal effect is likely lower than 200 grams.

7 Simulation Study

We used the empirical distribution of X , the estimated distributions of $P[Y \leq y|T = t, X]$ and $P[T = 1|X]$ from the data analysis above as the true observed data generating mechanisms. We used the functional forms of $s_t(Y(t))$ specified in the data analysis and considered values of γ_1 ranging from 0.0 to 0.01 and γ_0 ranging from -0.0025 to 0.0. Using the observed data distribution, $s_t(Y(t))$ and γ_t , we can use (4) to compute the true value of $E[Y(t)]$. In our simulation, we considered sample sizes of 1000, 1500 and 2000. We evaluated estimation bias (on percentage scale) and confidence interval coverage (normality-based, percentile bootstrap, double bootstrap).

The simulation results for smoking and non-smoking are presented in Tables 2 and 3, respec-




Figure 5: Contour plot for estimated average casual effect as a function of γ_1 and γ_0 .

tively. Bias is very low. In terms of confidence interval coverage, double bootstrap performs better than normality-based and percentile bootstrap.

8 Discussion

In this paper, we developed a semiparametric method for assessing sensitivity to the “no unmeasured confounding” assumption, which when true, implies that the average causal effect is identified via covariate adjustment. However, the adjustment functional is not the only way the ACE can be identified. For instance, in the “front door” model where we have measured a mediator with no unmeasured causes that captures all the effect of the treatment on outcome, the ACE is identifiable via a more complicated functional [Pearl, 2009]. There exist more examples of identification despite the presence of unmeasured confounders; see [Shpitser and Pearl, 2006, Bhattacharya et al., 2020]. The semiparametric sensitivity analysis procedure described here can be adapted to assess the underlying assumptions in those models in a robust way. This opens up several interesting directions for future work.

In the interest of simplicity and to fix ideas, we chose to use GAMs and single index models to estimate the nuisance functions. We used cross-validation to select tuning parameters that minimize estimates of the nuisance functions’ mean-square-errors and assessed the quality of our model fit

γ_1	n	Percent Bias	Coverage Normal	Coverage Percentile	Coverage Double
0.000	1000	0.110	0.956	0.949	0.966
	1500	0.011	0.959	0.955	0.967
	2000	0.070	0.950	0.937	0.955
0.001	1000	-0.011	0.970	0.959	0.972
	1500	-0.037	0.966	0.959	0.966
	2000	0.012	0.956	0.944	0.956
0.002	1000	-0.039	0.967	0.954	0.967
	1500	-0.030	0.966	0.958	0.966
	2000	0.012	0.958	0.952	0.958
0.003	1000	-0.031	0.966	0.953	0.966
	1500	-0.020	0.963	0.951	0.963
	2000	0.016	0.960	0.952	0.960
0.004	1000	-0.021	0.958	0.943	0.959
	1500	-0.015	0.965	0.952	0.965
	2000	0.021	0.958	0.946	0.958
0.005	1000	-0.011	0.953	0.937	0.954
	1500	-0.012	0.962	0.957	0.962
	2000	0.025	0.959	0.948	0.959
0.006	1000	-0.004	0.947	0.940	0.948
	1500	-0.010	0.960	0.954	0.960
	2000	0.027	0.955	0.949	0.956
0.007	1000	0.002	0.944	0.938	0.946
	1500	-0.008	0.959	0.946	0.959
	2000	0.029	0.950	0.940	0.951
0.008	1000	0.006	0.945	0.937	0.946
	1500	-0.007	0.954	0.939	0.954
	2000	0.031	0.951	0.943	0.951
0.009	1000	0.010	0.942	0.931	0.942
	1500	-0.005	0.949	0.938	0.949
	2000	0.031	0.949	0.937	0.949
0.01	1000	0.013	0.947	0.931	0.948
	1500	-0.004	0.948	0.934	0.948
	2000	0.033	0.948	0.944	0.949

Table 2: Results of simulation study: Smokers

γ_0	n	Percent Bias	Coverage Normal	Coverage Percentile	Coverage Double
-0.00250	1000	0.107	0.905	0.917	0.944
	1500	-0.060	0.909	0.910	0.946
	2000	-0.124	0.915	0.913	0.945
-0.00225	1000	0.060	0.919	0.919	0.949
	1500	-0.088	0.925	0.913	0.955
	2000	-0.146	0.921	0.923	0.951
-0.00200	1000	-0.002	0.932	0.927	0.950
	1500	-0.113	0.931	0.926	0.957
	2000	-0.167	0.931	0.927	0.951
-0.00175	1000	-0.065	0.935	0.922	0.953
	1500	-0.134	0.942	0.930	0.957
	2000	-0.174	0.933	0.912	0.949
-0.00150	1000	-0.094	0.946	0.927	0.961
	1500	-0.131	0.944	0.933	0.963
	2000	-0.161	0.936	0.922	0.953
-0.00125	1000	-0.091	0.953	0.941	0.960
	1500	-0.109	0.947	0.931	0.964
	2000	-0.133	0.944	0.932	0.961
-0.00100	1000	-0.072	0.950	0.941	0.964
	1500	-0.078	0.945	0.939	0.957
	2000	-0.103	0.956	0.940	0.968
-0.00075	1000	-0.048	0.954	0.937	0.957
	1500	-0.049	0.946	0.938	0.956
	2000	-0.075	0.960	0.950	0.969
-0.00050	1000	-0.027	0.950	0.946	0.959
	1500	-0.025	0.947	0.934	0.950
	2000	-0.049	0.962	0.953	0.968
-0.00025	1000	-0.011	0.951	0.943	0.958
	1500	-0.006	0.948	0.937	0.954
	2000	-0.029	0.957	0.948	0.964
0.00000	1000	0.001	0.955	0.944	0.961
	1500	0.007	0.948	0.937	0.954
	2000	-0.014	0.957	0.948	0.965

Table 3: Results of simulation study: Non-smokers

in Table 1. Due to our use of sample splitting and influence function-based estimators, all the asymptotic results will continue to hold for any nuisance estimators that converge to the truth faster than $n^{-1/4}$ rates. For example, we could have used sufficient dimension reduction techniques (see, e.g., [Ma and Zhu, 2012, 2013, Huang and Chiang, 2017]). Alternatively, we could have used a black-box ensemble method such as Super Learner [Van der Laan et al., 2007]; inference would then still be asymptotically valid, as long as at least one learner has root mean squared error that scales faster than $n^{-1/4}$. Achieving root mean square error of smaller order than $n^{-1/4}$ requires that the nuisance functions belong in a function class that is not too complex. Under certain conditions, this requirement could be weakened by incorporating higher order influence function terms that would estimate the remainder $\text{Rem}_t(\tilde{P}, P)$ [Robins et al., 2008, 2017]. This results in estimators that are considerably more complex than the (first-order) influence-function based estimator that we have constructed, and is thus beyond the scope of this paper.

While we imposed the consistency assumption in our data analysis, we realize that this may be implausible. That is, the amount of smoking during pregnancy may yield different outcomes, as suggested by Cattaneo [2010]. In future work, it would be useful to extend the method developed here to a continuous treatment. Our analysis, like those of Almond et al. [2005] and Cattaneo [2010], conditions on live birth. This is problematic because it is known that smoking increases the risk of miscarriage and stillbirth [Pineles et al., 2014]. This can lead to ill-defined counterfactuals, i.e., birthweight under smoking of a non-smoking mother may not be defined if the mother was to miscarry or have a stillborn child had she smoked. To address this issue, one could target the survivor average causal effect [Frangakis and Rubin, 2002, Egleston et al., 2007]. For identification, it would be reasonable to impose a monotonicity assumption in which it is assumed that if a mother had a live birth under smoking she would also have a live birth under not smoking. Nonetheless, sensitivity analysis to address unmeasured confounding would still be necessary. Thus, it would be useful to extend the ideas presented here to target this estimand.

Our approach relies on the scientific judgement of clinical experts through the selection of the key measured confounders, identification of the key unmeasured confounders, specification of the tilting functions, constraining the sensitivity parameters and interpretation of the results of the sensitivity analysis. The key advantage of our approach relative to other approaches is its reliance on flexible semiparametric models for the observed data that are disentangled from the sensitivity parameters. We hope that the methods (i.e., model-specification, derivation of the efficient influence function and asymptotic theory derived using sample-splitting) and clinician-guided data analysis discussed here can serve as a template for future methods development and the presentation of results in the presence of potential unmeasured confounding.

References

- Cheryl A Albuquerque, Karen R Smith, Cherie Johnson, Rex Chao, and Richard Harding. Influence of maternal tobacco smoking during pregnancy on uterine, umbilical and fetal cerebral artery blood flows. *Early human development*, 80(1):31–42, 2004.
- Douglas Almond, Kenneth Y Chay, and David S Lee. The costs of low birth weight. *The Quarterly Journal of Economics*, 120(3):1031–1083, 2005.
- Rudolf Beran. Prepivoting to reduce level error of confidence sets. *Biometrika*, 74(3):457–468, 1987.
- Rohit Bhattacharya, Raziieh Nabi, and Ilya Shpitser. Semiparametric inference for causal effects in graphical models with hidden variables. *arXiv preprint arXiv: 2003.12659*, 2020.
- P.J. Bickel, C.A.J. Klaassen, Y. Ritov, and J.A. Wellner. *Efficient and adaptive estimation for semiparametric models*. Johns Hopkins University Press Baltimore, 1993.
- Matteo Bonvini and Edward H Kennedy. Sensitivity analysis via the proportion of unmeasured confounding. *Journal of the American Statistical Association*, pages 1–11, 2021.
- Babette A Brumback, Miguel A Hernán, Sebastien JPA Haneuse, and James M Robins. Sensitivity analyses for unmeasured confounding assuming a marginal structural model for repeated measures. *Statistics in medicine*, 23(5):749–767, 2004.
- Graham J Burton, Marion E Palmer, and Kevin J Dalton. Morphometric differences between the placental vasculature of non-smokers, smokers and ex-smokers. *BJOG: An International Journal of Obstetrics & Gynaecology*, 96(8):907–915, 1989.
- PG Bush, TM Mayhew, DR Abramovich, PJ Aggett, MD Burke, and KR Page. A quantitative study on the effects of maternal smoking on placental morphology and cadmium concentration. *Placenta*, 21(2-3):247–256, 2000.
- E Martin Caravati, Carol J Adams, Steven M Joyce, and Nathan C Schafer. Fetal toxicity associated with maternal carbon monoxide poisoning. *Annals of emergency medicine*, 17(7):714–717, 1988.
- Nicole Bohme Carnegie, Masataka Harada, and Jennifer L Hill. Assessing sensitivity to unmeasured confounding using a simulated potential confounder. *Journal of Research on Educational*, 9(3): 395–420, 2016.
- Matias D Cattaneo. Efficient semiparametric estimation of multi-valued treatment effects under ignorability. *Journal of Econometrics*, 155(2):138–154, 2010.
- Chin-Tsang Chiang and Ming-Yueh Huang. New estimation and inference procedures for a single-index conditional distribution model. *Journal of Multivariate Analysis*, 111:271–285, 2012.
- Carlos Cinelli and Chad Hazlett. Making sense of sensitivity: Extending omitted variable bias. *Journal of the Royal Statistical Society: Series B (Statistical Methodology)*, 82(1):39–67, 2020.
- Jerome Cornfield, William Haenszel, E Cuyler Hammond, Abraham M Lilienfeld, Michael B Shimkin, and Ernst L Wynder. Smoking and lung cancer: recent evidence and a discussion of some questions. *Journal of the National Cancer institute*, 22(1):173–203, 1959.

- Iván Díaz and Mark J van der Laan. Sensitivity analysis for causal inference under unmeasured confounding and measurement error problems. *The international journal of biostatistics*, 9(2): 149–160, 2013.
- Iván Díaz, Alexander R Luedtke, and Mark J van der Laan. Sensitivity analysis. In *Targeted Learning in Data Science*, pages 511–522. Springer, 2018.
- Peng Ding and Tyler J VanderWeele. Sensitivity analysis without assumptions. *Epidemiology (Cambridge, Mass.)*, 27(3):368, 2016.
- Vincent Dorie, Masataka Harada, Nicole Bohme Carnegie, and Jennifer Hill. A flexible, interpretable framework for assessing sensitivity to unmeasured confounding. *Statistics in medicine*, 35(20):3453–3470, 2016.
- Bradley Efron and Robert J Tibshirani. *An introduction to the bootstrap*. CRC press, 1994.
- Brian L Egleston, Daniel O Scharfstein, Ellen E Freeman, and Sheila K West. Causal inference for non-mortality outcomes in the presence of death. *Biostatistics*, 8(3):526–545, 2007.
- Constantine E Frangakis and Donald B Rubin. Principal stratification in causal inference. *Biometrics*, 58(1):21–29, 2002.
- Alexander Franks, Alexander D’Amour, and Avi Feller. Flexible sensitivity analysis for observational studies without observable implications. *Journal of the American Statistical Association*, 2019.
- Joseph L Gastwirth, Abba M Krieger, and Paul R Rosenbaum. Dual and simultaneous sensitivity analysis for matched pairs. *Biometrika*, 85(4):907–920, 1998.
- Atilla Alp Gozubuyuk, Huseyin Dag, Alper Kaçar, Yakup Karakurt, and Vefik Arica. Epidemiology, pathophysiology, clinical evaluation, and treatment of carbon monoxide poisoning in child, infant, and fetus. *Northern clinics of Istanbul*, 4(1):100, 2017.
- Peter Hall. On the bootstrap and confidence intervals. *The Annals of Statistics*, pages 1431–1452, 1986.
- Peter Hall and Michael A Martin. On bootstrap resampling and iteration. *Biometrika*, 75(4): 661–671, 1988.
- Miguel A Hernán and James M Robins. Estimating causal effects from epidemiological data. *Journal of Epidemiology & Community Health*, 60(7):578–586, 2006.
- Joel L Horowitz, Enno Mammen, et al. Nonparametric estimation of an additive model with a link function. *The Annals of Statistics*, 32(6):2412–2443, 2004.
- Ming-Yueh Huang and Chin-Tsang Chiang. An effective semiparametric estimation approach for the sufficient dimension reduction model. *Journal of the American Statistical Association*, 112(519):1296–1310, 2017.
- Guido W Imbens. Sensitivity to exogeneity assumptions in program evaluation. *American Economic Review*, 93(2):126–132, 2003.
- Edward H Kennedy, Sivaraman Balakrishnan, Max G’Sell, et al. Sharp instruments for classifying compliers and generalizing causal effects. *Annals of Statistics*, 48(4):2008–2030, 2020.

- Edward H Kennedy, Sivaraman Balakrishnan, and Larry Wasserman. Semiparametric counterfactual density estimation. *arXiv preprint arXiv:2102.12034*, 2021.
- Lise G Larsen, Helle V Clausen, and Lisbeth Jønsson. Stereologic examination of placentas from mothers who smoke during pregnancy. *American journal of obstetrics and gynecology*, 186(3): 531–537, 2002.
- P Lehtovirta and M Forss. The acute effect of smoking on intervillous blood flow of the placenta. *BJOG: An International Journal of Obstetrics & Gynaecology*, 85(10):729–731, 1978.
- Weiwei Liu, S Janet Kuramoto, and Elizabeth A Stuart. An introduction to sensitivity analysis for unobserved confounding in nonexperimental prevention research. *Prevention science*, 14(6): 570–580, 2013.
- Yanyuan Ma and Liping Zhu. A semiparametric approach to dimension reduction. *Journal of the American Statistical Association*, 107(497):168–179, 2012.
- Yanyuan Ma and Liping Zhu. Efficient estimation in sufficient dimension reduction. *Annals of statistics*, 41(1):250, 2013.
- Charles F Manski. Nonparametric bounds on treatment effects. *The American Economic Review*, 80(2):319–323, 1990.
- Jerzy Neyman. Sur les applications de la thar des probabilités aux expériences agricoles: Essai des principe. excerpts reprinted (1990) in English. *Statistical Science*, 5:463–472, 1923.
- Judea Pearl. *Causality: Models, Reasoning, and Inference*. Cambridge University Press, 2 edition, 2009. ISBN 978-0521895606.
- Beth L Pineles, Edward Park, and Jonathan M Samet. Systematic review and meta-analysis of miscarriage and maternal exposure to tobacco smoke during pregnancy. *American journal of epidemiology*, 179(7):807–823, 2014.
- Amy Richardson, Michael G Hudgens, Peter B Gilbert, and Jason P Fine. Nonparametric bounds and sensitivity analysis of treatment effects. *Statistical science: a review journal of the Institute of Mathematical Statistics*, 29(4):596, 2014.
- J. M. Robins. A new approach to causal inference in mortality studies with sustained exposure periods – application to control of the healthy worker survivor effect. *Mathematical Modeling*, 7: 1393–1512, 1986.
- James Robins, Lingling Li, Eric Tchetgen, Aad van der Vaart, et al. Higher order influence functions and minimax estimation of nonlinear functionals. In *Probability and statistics: essays in honor of David A. Freedman*, pages 335–421. Institute of Mathematical Statistics, 2008.
- James M Robins. The analysis of randomized and non-randomized aids treatment trials using a new approach to causal inference in longitudinal studies. *Health service research methodology: a focus on AIDS*, pages 113–159, 1989.
- James M Robins. Association, causation, and marginal structural models. *Synthese*, pages 151–179, 1999.
- James M. Robins and Mark van der Laan. *Unified Methods for Censored Longitudinal Data and Causality*. Springer-Verlag New York, Inc., 2003.

- James M. Robins, Andrea Rotnitzky, and Lue P. Zhao. Estimation of regression coefficients when some regressors are not always observed. *Journal of the American Statistical Association*, 89: 846–866, 1994.
- James M Robins, Andrea Rotnitzky, and Daniel O Scharfstein. Sensitivity analysis for selection bias and unmeasured confounding in missing data and causal inference models. In *Statistical models in epidemiology, the environment, and clinical trials*, pages 1–94. Springer, 2000.
- James M Robins, Lingling Li, Rajarshi Mukherjee, Eric Tchetgen Tchetgen, Aad van der Vaart, et al. Minimax estimation of a functional on a structured high-dimensional model. *Annals of Statistics*, 45(5):1951–1987, 2017.
- Paul R. Rosenbaum. *Observational Studies*. Springer-Verlag New York, 2002.
- Paul R Rosenbaum and Donald B Rubin. Assessing sensitivity to an unobserved binary covariate in an observational study with binary outcome. *Journal of the Royal Statistical Society: Series B (Methodological)*, 45(2):212–218, 1983a.
- Paul R. Rosenbaum and Donald B. Rubin. The central role of the propensity score in observational studies for causal effects. *Biometrika*, 70:41–55, 1983b.
- Donald B. Rubin. Estimating causal effects of treatments in randomized and non-randomized studies. *Journal of Educational Psychology*, 66:688–701, 1974.
- Changyu Shen, Xiaochun Li, Lingling Li, and Martin C Were. Sensitivity analysis for causal inference using inverse probability weighting. *Biometrical journal*, 53(5):822–837, 2011.
- Ilya Shpitser and Judea Pearl. Identification of joint interventional distributions in recursive semi-Markovian causal models. In *Proceedings of the Twenty-First National Conference on Artificial Intelligence (AAAI-06)*. AAAI Press, Palo Alto, 2006.
- Anastasios Tsiatis. *Semiparametric Theory and Missing Data*. Springer-Verlag New York, 1st edition edition, 2006.
- Dongsheng Tu and Jun Shao. *The Jackknife and bootstrap*. Springer Series in Statistics, New York, 1995.
- Mark J Van Der Laan and Daniel Rubin. Targeted maximum likelihood learning. *The International Journal of Biostatistics*, 2(1), 2006.
- Mark J Van der Laan, Eric C Polley, and Alan E Hubbard. Super learner. *Statistical applications in genetics and molecular biology*, 6(1), 2007.
- Tyler J VanderWeele and Onyebuchi A Arah. Bias formulas for sensitivity analysis of unmeasured confounding for general outcomes, treatments, and confounders. *Epidemiology*, pages 42–52, 2011.
- Victor Veitch and Anisha Zaveri. Sense and sensitivity analysis: Simple post-hoc analysis of bias due to unobserved confounding. *arXiv preprint arXiv:2003.01747*, 2020.
- Lili Wang, Chao Zheng, wen Zhou, and Wen-Xin Zhou. A new principle for tuning-free huber regression. *Journal of Multivariate Analysis*, 111:271–285, 2012.

Steve Yadlowsky, Hongseok Namkoong, Sanjay Basu, John Duchi, and Lu Tian. Bounds on the conditional and average treatment effect with unobserved confounding factors. *arXiv preprint arXiv:1808.09521*, 2018.

Bo Zhang and Eric J Tchetgen Tchetgen. A semiparametric approach to model-based sensitivity analysis in observational studies. *arXiv preprint arXiv:1910.14130*, 2019.

Qingyuan Zhao, Dylan S Small, and Bhaswar B Bhattacharya. Sensitivity analysis for inverse probability weighting estimators via the percentile bootstrap. *Journal of the Royal Statistical Society: Series B (Statistical Methodology)*, 81(4):735–761, 2019.

Appendix

(A) Theorem 1. Non-Parametric Efficient Influence Function

A distribution $\tilde{P} \in \mathcal{M}$ is characterized by $\tilde{F}_t(y | x) = \tilde{P}(Y \leq y | T = t, X = x)$, $\tilde{\pi}_t(x) = \tilde{P}(T = t | X = x)$, and $\tilde{F}(x) = \tilde{P}(X \leq x)$. Let $\{\tilde{P}_\theta : \tilde{P}_\theta \in \mathcal{M}\}$. Let $s(O)$ be the score for θ evaluated at $\theta = 0$. We consider parametric submodels of the following form:

$$\begin{aligned} d\tilde{F}_\theta(x) &= dF(x)\{1 + \epsilon h(x)\} \\ d\tilde{F}_{t,\theta}(y|x) &= dF_t(y | x)\{1 + \eta_t k_t(y, x)\} \\ \tilde{\pi}_{t,\theta}(x) &= \frac{\{\pi_1(x) \exp\{\delta l(x)\}\}^t \pi_0(x)^{1-t}}{\pi_1(x) \exp\{\delta l(x)\} + \pi_0(x)} \end{aligned}$$

where $\theta = (\epsilon, \eta_0, \eta_1, \delta)$, $\mathbb{E}[h(X)] = 0$, $\mathbb{E}[k_t(Y, X) | T = t, X] = 0$ and $l(X)$ is any function of X . The associated score functions are $h(X)$, $Tk_1(Y, X) + (1 - T)k_0(Y, X)$, and $\{T - \pi_1(X)\}l(X)$.

The target parameter as a function of \tilde{P}_θ , $\psi_t(\tilde{P}_\theta)$, is

$$\psi_t(\tilde{P}_\theta) = \int_x \int_y y d\tilde{F}_{t,\theta}(y | x) \tilde{\pi}_{t,\theta}(x) d\tilde{F}_\theta(x) + \int_x \frac{\int_y y \exp(\gamma_t s_t(y)) d\tilde{F}_{t,\theta}(y | x)}{\int_y \exp(\gamma_t s_t(y)) d\tilde{F}_{t,\theta}(y | x)} \tilde{\pi}_{1-t,\theta}(x) d\tilde{F}_\theta(x).$$

The derivative of $\psi_t(\tilde{P}_\theta)$ with respect to ϵ evaluated at $\theta = 0$ is

$$\int_x \left\{ \mu_t(Y; x) \pi_t(x) + \frac{\mu_t(Y \exp(\gamma_t s_t(Y)); x)}{\mu_t(\exp(\gamma_t s_t(Y)); x)} \pi_{1-t}(x) \right\} h(x) dF(x).$$

The derivative of $\psi_t(\tilde{P}_\theta)$ with respect to η_t evaluated at $\theta = 0$ is

$$\begin{aligned} \int_x \left\{ \left[\frac{\mu_t(Y \exp(\gamma_t s_t(Y))) k_t(Y, x); x}{\mu_t(\exp(\gamma_t s_t(Y)); x)} - \frac{\mu_t(\exp(\gamma_t s_t(Y))) k_t(Y, x); x}{\mu_t(\exp(\gamma_t s_t(Y)); x)^2} \frac{\mu_t(Y \exp(\gamma_t s_t(Y)); x)}{\mu_t(\exp(\gamma_t s_t(Y)); x)} \right] \pi_{1-t}(x) \right. \\ \left. + \mu_t(Y k_t(Y, x); x) \pi_t(x) \right\} dF(x). \end{aligned}$$

The derivative of $\psi_t(\tilde{P}_\theta)$ with respect to η_{1-t} evaluated at $\theta = 0$ is 0.

The derivative of $\psi_t(\tilde{P}_\theta)$ with respect to δ evaluated at $\theta = 0$ is

$$\int_x (-1)^{t+1} \left\{ \mu_t(Y; x) - \frac{\mu_t(Y \exp(\gamma_t s_t(Y)); x)}{\mu_t(\exp(\gamma_t s_t(Y)); x)} \right\} \pi_1(x) \pi_0(x) l(x) dF(x).$$

Any mean zero observed data random variable can be expressed as

$$d(O) = a(X) + \sum_{t=0}^1 I(T = t) b_t(Y, X) + (T - \pi_1(X)) c(X),$$

where $E[a(X)] = 0$, $E[b_t(Y, X) | T = t, X] = 0$ and $c(X)$ is an unspecified function of X . The set of all $d(O)$ is the non-parametric tangent space. To find the non-parametric efficient influence function,

we need to find choices of $a(X)$, $b_t(Y, X)$ and $c(X)$ such that $E[a(X)h(X)] = \partial\psi_t(\tilde{P}_\theta)/\partial\epsilon|_{\theta=0}$, $E[I(T=t)b_t(Y, X)k_t(Y, X)] = \partial\psi_t(P_\theta)/\partial\eta_t|_{\theta=0}$ and $E[(T - \pi_1(X))^2c(X)l(X)] = \psi_t(\tilde{P}_\theta)/\partial\delta|_{\theta=0}$. It can be shown that

$$\begin{aligned} a(X) &= \mu_t(Y; X)\pi_t(X) + \frac{\mu_t(Y \exp(\gamma_t s_t(Y)); X)}{\mu_t(\exp(\gamma_t s_t(Y)); X)}\pi_{1-t}(X) - \psi_t(P) \\ b_t(Y, X) &= Y + \frac{Y \exp(\gamma_t s_t(Y))}{\mu_t(\exp(\gamma_t s_t(Y)); X)} \frac{\pi_{1-t}(X)}{\pi_t(X)} - \exp(\gamma_t s_t(Y)) \frac{\mu_t(Y \exp(\gamma_t s_t(Y)); X)}{\mu_t(\exp(\gamma_t s_t(Y)); X)^2} \frac{\pi_{1-t}(X)}{\pi_t(X)} - \mu_t(Y; X) \\ b_{1-t}(Y, X) &= 0 \\ c(X) &= (-1)^{t+1} \left\{ \mu_t(Y; X) - \frac{\mu_t(Y \exp(\gamma_t s_t(Y)); X)}{\mu_t(\exp(\gamma_t s_t(Y)); X)} \right\}. \end{aligned}$$

Hence, the non-parametric efficient influence function that corresponds to ψ_t is as follows:

$$\begin{aligned} \phi_t(P) &= \mathbb{I}(T=t) \left\{ Y + \frac{Y \exp(\gamma_t s_t(Y))}{\mu_t(\exp(\gamma_t s_t(Y)); X)} \frac{\pi_{1-t}(X)}{\pi_t(X)} - \exp(\gamma_t s_t(Y)) \frac{\mu_t(Y \exp(\gamma_t s_t(Y)); X)}{\mu_t(\exp(\gamma_t s_t(Y)); X)^2} \frac{\pi_{1-t}(X)}{\pi_t(X)} \right\} \\ &\quad + \mathbb{I}(T=1-t) \frac{\mu_t(Y \exp(\gamma_t s_t(Y)); X)}{\mu_t(\exp(\gamma_t s_t(Y)); X)} - \psi_t(P). \end{aligned}$$

(B) Lemma 1. Second-Order Remainder Term

$\text{Rem}_t(\tilde{P}, P)$ is defined as $\psi_t(\tilde{P}) - \psi_t(P) + \mathbb{E}[\phi_t(\tilde{P})(O)]$

$$\psi_t(P) = \mathbb{E} \left[\mu_t(Y; X)\pi_t(X) + \frac{\mu_t(Y \exp(\gamma_t s_t(Y)); X)}{\mu_t(\exp(\gamma_t s_t(Y)); X)}\pi_{1-t}(X) \right],$$

and

$$\begin{aligned} \mathbb{E}[\phi_t(\tilde{P})(O)] &= \mathbb{E} \left[\pi_t(X)\mu_t(Y; X) + \pi_t(X) \left\{ \frac{\mu_t(Y \exp(\gamma_t s_t(Y)), X)}{\tilde{\mu}_t(\exp(\gamma_t s_t(Y)); X)} \times \frac{\tilde{\pi}_{1-t}(X)}{\tilde{\pi}_t(X)} \right. \right. \\ &\quad \left. \left. - \frac{\mu_t(\exp(\gamma_t s_t(Y)); X)}{\tilde{\mu}_t(\exp(\gamma_t s_t(Y)); X)^2} \frac{\tilde{\mu}_t(Y \exp(\gamma_t s_t(Y)); X)}{\tilde{\pi}_t(X)} \right\} \right. \\ &\quad \left. + \pi_{1-t}(X) \times \frac{\tilde{\mu}_t(Y \exp(\gamma_t s_t(Y)); X)}{\tilde{\mu}_t(\exp(\gamma_t s_t(Y)); X)} \right] - \psi_t(\tilde{P}). \end{aligned}$$

Therefore,

$$\begin{aligned} \text{Rem}_t(\tilde{P}, P) &= \mathbb{E} \left[\pi_t(X) \left\{ \frac{\mu_t(Y \exp(s_t(Y)); X)}{\tilde{\mu}_t(\exp(s_t(Y)); X)} \times \frac{\tilde{\pi}_{1-t}(X)}{\tilde{\pi}_t(X)} - \frac{\mu_t(\exp(\gamma_t s_t(Y)); X)\tilde{\mu}_t(Y \exp(\gamma_t s_t(Y)); X)}{\tilde{\mu}_t(\exp(\gamma_t s_t(Y)); X)^2} \times \frac{\tilde{\pi}_{1-t}(X)}{\tilde{\pi}_t(X)} \right\} + \right. \\ &\quad \left. \pi_{1-t}(X) \times \frac{\tilde{\mu}_t(Y \exp(\gamma_t s_t(Y)); X)}{\tilde{\mu}_t(\exp(\gamma_t s_t(Y)); X)} - \pi_{1-t}(X) \times \frac{\mu_t(Y \exp(\gamma_t s_t(Y)); X)}{\mu_t(\exp(\gamma_t s_t(Y)); X)} \right] \\ &= \mathbb{E} \left[\pi_t(X) \frac{\tilde{\pi}_{1-t}(X)}{\tilde{\pi}_t(X)} \left\{ \frac{\mu_t(Y \exp(s_t(Y)); X)\tilde{\mu}_t(\exp(s_t(Y)); X) - \mu_t(\exp(\gamma_t s_t(Y)); X)\tilde{\mu}_t(Y \exp(\gamma_t s_t(Y)); X)}{\tilde{\mu}_t(\exp(\gamma_t s_t(Y)); X)^2} \right\} - \right. \end{aligned}$$

$$\begin{aligned}
& \pi_{1-t}(X) \left\{ \frac{-\tilde{\mu}_t(Y \exp(\gamma_t s_t(Y)); X) \mu_t(\exp(\gamma_t s_t(Y)); X) + \mu_t(Y \exp(\gamma_t s_t(Y)); X) \tilde{\mu}_t(\exp(\gamma_t s_t(Y)); X)}{\tilde{\mu}_t(\exp(\gamma_t s_t(Y)); X) \mu_t(\exp(\gamma_t s_t(Y)); X)} \right\} \\
= & \mathbb{E} \left[\frac{\pi_t(X) \tilde{\pi}_{1-t}(X)}{\tilde{\mu}_t(\exp(\gamma_t s_t(Y)); X)} \left\{ \frac{\mu_t(Y \exp(s_t(Y)); X) \tilde{\mu}_t(\exp(s_t(Y)); X) - \mu_t(\exp(\gamma_t s_t(Y)); X) \tilde{\mu}_t(Y \exp(\gamma_t s_t(Y)); X)}{\tilde{\pi}_t(X) \tilde{\mu}_t(\exp(\gamma_t s_t(Y)); X)} \right\} - \right. \\
& \left. \frac{\tilde{\pi}_t(X) \pi_{1-t}(X)}{\mu_t(\exp(\gamma_t s_t(Y)); X)} \left\{ \frac{-\tilde{\mu}_t(Y \exp(\gamma_t s_t(Y)); X) \mu_t(\exp(\gamma_t s_t(Y)); X) + \mu_t(Y \exp(\gamma_t s_t(Y)); X) \tilde{\mu}_t(\exp(\gamma_t s_t(Y)); X)}{\tilde{\pi}_t(X) \tilde{\mu}_t(\exp(\gamma_t s_t(Y)); X)} \right\} \right] \\
= & \mathbb{E} \left[\left\{ \frac{\mu_t(Y \exp(s_t(Y)); X) \tilde{\mu}_t(\exp(s_t(Y)); X) - \mu_t(\exp(\gamma_t s_t(Y)); X) \tilde{\mu}_t(Y \exp(\gamma_t s_t(Y)); X)}{\tilde{\pi}_t(X) \tilde{\mu}_t(\exp(\gamma_t s_t(Y)); X)} \right\} \times \right. \\
& \left. \left\{ \frac{\pi_t(X) \tilde{\pi}_{1-t}(X)}{\tilde{\mu}_t(\exp(\gamma_t s_t(Y)); X)} - \frac{\tilde{\pi}_t(X) \pi_{1-t}(X)}{\mu_t(\exp(\gamma_t s_t(Y)); X)} \right\} \right]
\end{aligned}$$

(C) Single Index Model

Without loss of generality, we assume that we are within the $T = t$ stratum and drop the dependence of the notation on t . Let p be the dimension of X . The single index model for the conditional distribution of Y given $X = x$ posits that

$$P[Y \leq y \mid X = x] = F(y, x' \beta; \beta),$$

where $F(y, u; \beta)$ is a cumulative distribution function in y for each u and $\beta = (\beta_1, \dots, \beta_p)$ is a p -dimensional vector of unknown parameters. For purposes of identifiability, β_1 is set to 1.

Let

$$\hat{F}(y, u; \beta, h) = \frac{\sum_{i=1}^n I(Y_i \leq y) K_h(X_i' \beta - u)}{\sum_{i=1}^n K_h(X_i' \beta - u)}$$

where $K_h(v) = K(v/h)/h$, K is a q^* -th order kernel and h is a positive bandwidth. Let

$$\hat{F}^{(-i)}(y, X_i' \beta; \beta, h) = \frac{\sum_{j \neq i} I(Y_j \leq y) K_h(X_j' \beta - X_i' \beta)}{\sum_{j \neq i} K_h(X_j' \beta - X_i' \beta)}$$

and

$$CV(\beta, h) = \frac{1}{n} \sum_{i=1}^n \int \{I(Y_i \leq y) - \hat{F}^{(-i)}(y, X_i' \beta; \beta, h)\}^2 d\hat{F}(y),$$

where $\hat{F}(\cdot)$ is the empirical distribution of Y .

When $F(y, u; \beta)$ and the density function $f_{X' \beta}(u)$ of $X' \beta$ have Lipschitz $(q+1)$ th-order derivatives, standard nonparametric smoothing results can be used to show that

$$\sup_{y, u, \beta} |\hat{F}(y, u; \beta, h_n) - F(y, u; \beta)| = O_p(h_n^{q^*} + \{\log n / (nh_n)\}^{1/2})$$

Assume $h_n \in [h_l n^{-\delta}, h_u n^{-\delta}]$ for some positive constants h_l and h_u and $\delta \in (1/(4q^*), 1/5)$. Let $I_n = \cup_{\delta \in (1/(4q^*), 1/5)} [h_l n^{-\delta}, h_u n^{-\delta}]$, $\tilde{h}_\beta = \operatorname{argmin}_{h \in I_n} CV(\beta, h)$, and $\hat{\beta}_h = \operatorname{argmin}_\beta CV(\beta, h)$. Chiang and Huang [2012] showed the following:

1. $\tilde{h}_\beta = O_P(n^{-1/(2q^*+1)})$ uniformly in β .
2. $\sqrt{n}(\hat{\beta}_{h_n} - \beta) \xrightarrow{D} N(0, V^{-1}\Sigma V^{-1})$.

Now let $(\hat{\beta}, \tilde{h})$ be $\operatorname{argmin}_{\beta \in \mathbb{R}^{p-1}, h \in I_n} CV(\beta, h)$. To ensure $\tilde{h} \in I_n$ with probability one, it is required that $1/(4q^*) < 1/(2q^* + 1) < 1/5$. Thus, we choose $q^* = 4$. This implies that $\tilde{h} = O_P(n^{-1/(2q^*+1)})$ and satisfies the assumption on h_n . Since $\hat{\beta} = \hat{\beta}_{\tilde{h}}$, we can conclude that $\sqrt{n}(\hat{\beta} - \beta) \xrightarrow{D} N(0, V^{-1}\Sigma V^{-1})$. That is, the estimator $\hat{\beta}$ is $n^{1/2}$ -consistent under this semiparametric modeling framework.

To estimate $\mu_t(\exp(\gamma_t s_t(Y)); X) \mu_t(Y \exp(\gamma_t s_t(Y)); X)$ under this single-index model setting, we consider a general transformation on Y : $G_\tau(x) = E[\tau(Y)|X = x]$, which we estimate by

$$\hat{G}_\tau(x; h_n) = \int \tau(y) d\hat{F}(y, x' \hat{\beta}, \hat{\beta}, h_n).$$

Let

$$\hat{G}_\tau(u, \beta, h) = \int \tau(y) d\hat{F}(y, u, \beta, h) = \frac{\sum_{i=1}^n \tau(Y_i) K_h(X_i' \beta - u)}{\sum_{i=1}^n K_h(X_i' \beta - u)},$$

where now K is a q -th order kernel function. Standard nonparametric smoothing theory guarantees

$$\sup_{u, \beta} |\hat{G}_\tau(u, \beta, h_n) - G_\tau(u; \beta)| = O_P(h_n^q + \{\log n/(nh_n)\}^{1/2}),$$

where $G_\tau(u; \beta) = E[\tau(Y)|X' \beta = u]$. Since $\hat{\beta}$ is $n^{1/2}$ -consistent to β , we can show that

$$\begin{aligned} \sup_x |\hat{G}_\tau(x; h_n) - G_\tau(x)| &\leq \sup_x |\hat{G}_\tau(x' \hat{\beta}; \hat{\beta}, h_n) - \hat{G}_\tau(x' \beta; \beta, h_n)| + \sup_x |\hat{G}_\tau(x' \beta; \beta, h_n) - G_\tau(x' \beta; \beta)| \\ &= \sup_x \left| \frac{\partial}{\partial \beta} \hat{G}_\tau(x' \beta; \beta, h_n) \Big|_{\beta = \hat{\beta}^*} (\hat{\beta} - \beta) \right| + \sup_x |\hat{G}_\tau(x' \beta; \beta, h_n) - G_\tau(x' \beta; \beta)| \\ &\leq \underbrace{\sup_x \left| \frac{\partial}{\partial \beta} \hat{G}_\tau(x' \beta; \beta, h_n) \Big|_{\beta = \hat{\beta}^*} \right|}_{O_P(1)} \underbrace{|\hat{\beta} - \beta|}_{O_P(n^{-1/2})} + \underbrace{\sup_x |\hat{G}_\tau(x' \beta; \beta, h_n) - G_\tau(x' \beta; \beta)|}_{O_P(h_n^q + \{\log n/(nh_n)\}^{1/2})} \end{aligned} \quad (18)$$

where $\hat{\beta}^*$ lies on the line segment between $\hat{\beta}$ and β . In the first term of (18),

$$\begin{aligned} &\sup_x \left| \frac{\partial}{\partial \beta} \hat{G}_\tau(x' \beta; \beta, h_n) \Big|_{\beta = \hat{\beta}^*} \right| \\ &\leq \underbrace{\sup_{x, \beta} \left| \frac{\partial}{\partial \beta} \hat{G}_\tau(x' \beta; \beta, h_n) - G_\tau^{[1]}(x; \beta) \right|}_{O_P(h_n^q + \{\log n/(nh_n^2)\}^{1/2})} + \underbrace{\sup_{x, \beta} |G_\tau^{[1]}(x; \beta)|}_M = O_P(1) \end{aligned} \quad (19)$$

for some deterministic function $G_\tau^{[1]}(x; \beta)$ that we assume to be continuously differentiable with respect its arguments and to have compact support. Thus, $\sup_{x, \beta} |G_\tau^{[1]}(x; \beta)|$ will be less than or equal to a constant, say M , that is finite. In addition, Chiang and Huang [2012] showed that

$$\sup_{x, \beta} \left| \frac{\partial}{\partial \beta} \hat{G}_\tau(x' \beta; \beta, h_n) - G_\tau^{[1]}(x; \beta) \right| = O_P(h_n^q + \{\log n/(nh_n^2)\}^{1/2}).$$

If $h_n \rightarrow 0$ and $nh_n^2 \rightarrow \infty$, then $O_P(h_n^q + \{\log n/(nh_n^2)\}^{1/2}) = o_P(1)$. It can also be shown that $n^{-1/2}/(h_n^q + \{\log n/(nh_n^2)\}^{1/2}) \rightarrow 0$. Thus, we may conclude that

$$\sup_x |\widehat{G}_\tau(x, h_n) - G_\tau(x)| = O_P(h_n^q + \{\log n/(nh_n^2)\}^{1/2}).$$

Further, if h_n is of the optimal rate $O(n^{-1/(2q+1)})$ and $q > 1/2$, we have

$$\sup_x |\widehat{G}_\tau(x, h_n) - G_\tau(x)| = O_P(n^{-1/4}).$$

In practice, we use $q = 2$ and $\widehat{h} = \widetilde{h}n^{-4/45}$. Note that $\widetilde{h} = O_P(n^{-1/9})$ from the above procedure of estimating $\widehat{\beta}$, and hence \widehat{h} is of the optimal rate $O(n^{-1/5})$

(D) Lemma 2. Sample Splitting as an Alternative to Donsker Conditions

It is sufficient to show that $\|\nu_t(\widehat{P}^{(-k)}) - \nu_t(P)\|_{L_2}$ is $o_P(1)$. Note that

$$\begin{aligned} & \nu_t(\widehat{P}^{(-k)})(O) - \nu_t(P)(O) \\ &= I(T = t) \left\{ \frac{\widehat{\pi}_{1-t}^{(-k)}(X)}{\widehat{\pi}_t^{(-k)}(X)\widehat{\mu}_t^{(-k)}(\exp(\gamma_t s_t(Y)), X)} - \frac{\pi_{1-t}(X)}{\pi_t(X)\mu_t(\exp(\gamma_t s_t(Y)), X)} \right\} Y \exp(\gamma_t s_t(Y)) - \\ & \quad I(T = t) \left\{ \frac{\widehat{\pi}_{1-t}^{(-k)}(X)\widehat{\mu}_t^{(-k)}(Y \exp(\gamma_t s_t(Y)), X)}{\widehat{\pi}_t^{(-k)}(X)\widehat{\mu}_t^{(-k)}(\exp(\gamma_t s_t(Y)), X)^2} - \frac{\pi_{1-t}(X)\mu_t(Y \exp(\gamma_t s_t(Y)), X)}{\pi_t(X)\mu_t(\exp(\gamma_t s_t(Y)), X)^2} \right\} \exp(\gamma_t s_t(Y)) + \\ & \quad I(T = 1 - t) \left\{ \frac{\widehat{\mu}_t^{(-k)}(Y \exp(\gamma_t s_t(Y)), X)}{\widehat{\mu}_t^{(-k)}(\exp(\gamma_t s_t(Y)), X)} - \frac{\mu_t(Y \exp(\gamma_t s_t(Y)), X)}{\mu_t(\exp(\gamma_t s_t(Y)), X)} \right\} \end{aligned}$$

We assume that $|Y|$ and $|\exp(\gamma_t s_t(Y))|$ are bounded in probability. Using the triangle and Cauchy-Schwarz inequalities, it can be shown that

$$\begin{aligned} & \|\nu_t(\widehat{P}^{(-k)}) - \nu_t(P)\|_{L_2} \\ & \leq \widehat{D}_1^{(-k)} \underbrace{\|\widehat{\pi}_t^{(-k)}(X) - \pi_t(X)\|_{L_2}}_{o_P(1)} + \widehat{D}_2^{(-k)} \underbrace{\|\widehat{\mu}_t^{(-k)}(\exp(\gamma_t s_t(Y)), X) - \mu_t(\exp(\gamma_t s_t(Y)), X)\|_{L_2}}_{o_P(1)} + \\ & \quad \widehat{D}_3^{(-k)} \underbrace{\|\widehat{\mu}_t^{(-k)}(Y \exp(\gamma_t s_t(Y)), X) - \mu_t(Y \exp(\gamma_t s_t(Y)), X)\|_{L_2}}_{o_P(1)}, \end{aligned}$$

where $\widehat{D}_1^{(-k)}$, $\widehat{D}_2^{(-k)}$, and $\widehat{D}_3^{(-k)}$ are $O_P(1)$. As a result, $\|\nu_t(\widehat{P}^{(-k)}) - \nu_t(P)\|_{L_2}$ is $o_P(1)$.