

PERIODICITY OF QUANTUM WALKS DEFINED BY MIXED PATHS AND MIXED CYCLES

SHO KUBOTA, HIROTO SEKIDO, AND HARUNOBU YATA

ABSTRACT. In this paper, we determine periodicity of quantum walks defined by mixed paths and mixed cycles. By the spectral mapping theorem of quantum walks, consideration of periodicity is reduced to eigenvalue analysis of η -Hermitian adjacency matrices. First, we investigate coefficients of the characteristic polynomials of η -Hermitian adjacency matrices. We show that the characteristic polynomials of mixed trees and their underlying graphs are same. We also define $n+1$ types of mixed cycles and show that every mixed cycle is switching equivalent to one of them. We use these results to discuss periodicity. We show that the mixed paths are periodic for any η . In addition, we provide a necessary and sufficient condition for a mixed cycle to be periodic and determine their periods.

1. INTRODUCTION

Quantum walks are quantum analogues of classical random walks [1, 3, 10]. In the last two decades, a great deal of research on quantum walks has been carried out, and they have strong connections with various fields. In quantum information, quantum walk models can be seen as a generalization of Grover's search algorithm [9, 31]. An important fact in mathematics is the spectral mapping theorem of quantum walks [14, 23]. The spectral mapping theorems reduce eigenvalue analysis of time evolution operators to eigenvalue analysis of other self-adjoint operators. They bring quantum walks into close connection with functional analysis [34] and spectral graph theory [20]. In the study of periodicity of discrete-time quantum walk, field theory and algebraic number theory have also been leveraged [18, 35]. Studies of perfect state transfer in continuous-time quantum walks have also been done by algebraic graph theory and algebraic combinatorics. We refer to Godsil's survey [8]. Recent studies on state transfer are in [6, 26, 27, 29, 36, 39].

1.1. Related works to periodicity. The topic discussed in this paper is periodicity of discrete-time quantum walks. The works of [15, 19] triggered off studies of periodicity on various graphs. For example, the studies done on Grover walks can be summarized in Table 1. In other models, periodicity of Fourier walks has been considered by Saito

2010 *Mathematics Subject Classification.* 05C50; 81Q99; 05C20; 05C81.

Key words and phrases. Quantum walk, periodicity, mixed graph, Hermitian adjacency matrix, spectral graph theory .

Graphs	Ref.
Complete graphs, complete bipartite graphs, SRGs	[15]
Generalized Bethe trees	[21]
Distance regular graphs	[38]
Cycle (3-state)	[18]
Complete graphs with self loops	[16]

TABLE 1. Prior works on periodicity of Grover walks on undirected graphs

[32], and periodicity of staggered walks has been studied in [22]. Recently, periodicity of quantum walks with generalized Grover coins has been considered by Sarkar et al [35].

1.2. Main Results. In this paper, we study periodicity of mixed graphs. There are three main theorems. See later sections for more detailed terms and definitions. First, we generalize the result in [4] related to classification of mixed cycles to η -Hermitian adjacency matrices. In [4], Akbari et al provided several typical switching functions and classified mixed cycles into three types. We give similar considerations in η -Hermitian adjacency matrices. Among the four types of switching defined in [4], one switching cannot be used in η -Hermitian adjacency matrices. Due to this, we show that there are at most $n + 1$ switching equivalence classes of mixed cycles in η -Hermitian adjacency matrices. The claim is as follows:

Theorem 1.1. *Let $G = (V, \mathcal{A})$ be a mixed cycle of length n . Then, there exists $j \in \{0, 1, \dots, n\}$ such that G and C_n^j is H_η -cospectral. Moreover, we have*

$$\det H_\eta(C_n^j) = (-1)^{n+1} 2 \cos(\eta j) + (-1)^{\lfloor \frac{n}{2} \rfloor} (1 + (-1)^n).$$

The second main theorem is to determine periodicity of mixed paths. Using the model defined in [20], we study periodicity of quantum walks defined by mixed graphs. This model is defined by both a mixed graph and a real number $\eta \in [0, 2\pi)$. The claim is as follows:

Theorem 1.2. *Let $G = (V, \mathcal{A})$ be a mixed path on n vertices equipped with an η -function θ . Then G is periodic for any $\eta \in [0, 2\pi)$, and the period is $2(n - 1)$.*

The third main theorem is to determine periodicity of mixed cycles. Since periodicity is determined by eigenvalues, it is sufficient to consider the only $n + 1$ types of mixed cycles by the first main theorem. The claim is as follows:

Theorem 1.3. *Let $G = (V, \mathcal{A})$ be a mixed cycle on n vertices equipped with an η -function θ . Then, G is periodic if and only if $\eta \in \mathbb{Q}\pi$. In addition, we suppose that $\eta \in \mathbb{Q}\pi$ and the mixed cycle G is H_η -cospectral with C_n^j . Let $\eta = \frac{p}{q}\pi$, where p and q are*

coprime. Then, the period τ of G is the following:

$$\tau = \begin{cases} \frac{2qn}{(j,2q)} & \text{if } p \text{ is odd,} \\ \frac{qn}{(j,q)} & \text{if } p \text{ is even.} \end{cases} \quad (1.1)$$

This paper is organized as follows. In Section 2, we prepare terminologies on spectral graph theory. The definitions of mixed graphs and their η -Hermitian adjacency matrices are provided. In Section 3, coefficients of the characteristic polynomials of η -Hermitian adjacency matrices are discussed. We focus on permutations that contribute to values of determinants. Relationship between the characteristic polynomials of a mixed graph and its underlying graph is clarified. In Section 4, we carry out classification of mixed cycles by η -Hermitian adjacency matrices. We introduce $n + 1$ types of mixed cycles and show that every mixed cycle is switching equivalent to one of them. On the other hand, we show that the $n + 1$ types of mixed cycles have different eigenvalues except for a finite number of η . In Section 5, we prepare quantum walks defined by mixed graphs. We define periodicity of mixed graphs and provide some characterizations of it. In Section 6, we discuss periodicity of mixed paths. We observe action of time evolution matrices on the unit vectors and provide a visual proof. In Section 7, we discuss periodicity of mixed cycles. It is easy to provide a necessary and sufficient condition for a mixed cycle to be periodic, but determination of the period is a bit complicated.

2. PRELIMINARIES ON SPECTRAL GRAPH THEORY

Let $\Gamma = (V, E)$ be a finite simple and connected graph with the vertex set V and the edge set E . For $x \in V$, the set of neighbors of x is denoted by $N(x)$. Define $\mathcal{A} = \mathcal{A}(\Gamma) = \{(x, y), (y, x) \mid xy \in E(\Gamma)\}$, which is the set of the symmetric arcs of Γ . The origin and terminus of $a = (x, y) \in \mathcal{A}$ are denoted by $o(a), t(a)$, respectively. We express a^{-1} as the inverse arc of a .

A *mixed graph* G consists of a finite set $V(G)$ of vertices together with a subset $\mathcal{A}(G) \subset V(G) \times V(G) \setminus \{(x, x) \mid x \in V\}$ of ordered pairs called *arcs*. Let G be a mixed graph. Define $\mathcal{A}^{-1}(G) = \{a^{-1} \mid a \in \mathcal{A}(G)\}$ and $\mathcal{A}^{\pm}(G) = \mathcal{A}(G) \cup \mathcal{A}^{-1}(G)$. If there is no danger of confusion, we write $\mathcal{A}(G)$ as $\mathcal{A}$ simply. If $(x, y) \in \mathcal{A} \cap \mathcal{A}^{-1}$, we say that the unordered pair $\{x, y\}$ is a *digon* of G . For a vertex $x \in V(G)$, define $\deg_G x = \deg_{G^{\pm}} x$. A mixed graph G is k -regular if $\deg_G x = k$ for any vertex $x \in V(G)$. The graph $G^{\pm} = (V(G), \mathcal{A}^{\pm})$ is so-called the *underlying graph* of a mixed graph G , and this is regarded as an undirected graph depending on context. On the other hand, we equate an undirected graph with a mixed graph by considering undirected edges xy as bidirectional arcs $(x, y), (y, x)$. Throughout this paper, we assume that mixed graphs are weakly connected, i.e., we assume that $G^{\pm}$ is connected.

Let $G = (V, \mathcal{A})$ be a mixed graph. For $\eta \in [0, 2\pi)$, the η -Hermitian adjacency matrix $H_\eta = H_\eta(G) \in \mathbb{C}^{V \times V}$ is defined by

$$(H_\eta)_{x,y} = \begin{cases} 1 & \text{if } (x, y) \in \mathcal{A} \cap \mathcal{A}^{-1}, \\ e^{\eta i} & \text{if } (x, y) \in \mathcal{A} \setminus \mathcal{A}^{-1}, \\ e^{-\eta i} & \text{if } (x, y) \in \mathcal{A}^{-1} \setminus \mathcal{A}, \\ 0 & \text{otherwise.} \end{cases}$$

When $\eta = \frac{\pi}{2}$, the matrix $H_{\frac{\pi}{2}}$ is nothing but the Hermitian adjacency matrix. This is introduced by Guo–Mohar [12] and Li–Liu [24], independently. When $\eta = \frac{\pi}{3}$, the matrix $H_{\frac{\pi}{3}}$ is called the Hermitian adjacency matrix of the second kind. This is introduced by Mohar [28]. We refer to [2, 13, 25] as recent studies on Hermitian adjacency matrices. Note that $H_\eta(G^\pm)$ coincides with the ordinary adjacency matrix of $G^\pm$. Define the degree matrix $D = D(G) \in \mathbb{C}^{V \times V}$ by $D_{x,y} = (\deg_G x) \delta_{x,y}$ for vertices $x, y \in V(G)$, where $\delta_{x,y}$ is the Kronecker delta. For $\eta \in [0, 2\pi)$, the normalized η -Hermitian adjacency matrix $\tilde{H}_\eta$ is defined by

$$\tilde{H}_\eta = D^{-\frac{1}{2}} H_\eta D^{-\frac{1}{2}}.$$

Note that if a mixed graph G is k -regular, we have $\tilde{H}_\eta = \frac{1}{k} H_\eta$.

Let G be a mixed graph. The list of the eigenvalues of $H_\eta(G)$ together with their multiplicities, denoted by $\text{Spec}(H_\eta(G))$, is called H_η -spectrum of G . We say that mixed graphs G and G' are H_η -cospectral if they have the same H_η -spectrum. The same is on $\tilde{H}_\eta$.

3. PERMUTATIONS AND CHARACTERISTIC POLYNOMIALS

Let Γ be an undirected graph. A mixed graph G is said to be a *mixed* Γ if $G^\pm$ is isomorphic to Γ . Similarly, we say that G is a *mixed tree* if $G^\pm$ is a tree. Let $G = (V, \mathcal{A})$ be a mixed graph, and let $x_1, x_2, \dots, x_l \in V$. We say that a sequence $C = (x_1, x_2, \dots, x_l)$ is an l -cycle in G if C is an l -cycle in $G^\pm$. The *girth* of G is defined by the girth of $G^\pm$. Note that if the girth of G is $s + 1$, then it has no l -cycle for $l \in \{1, 2, \dots, s\}$.

Let $G = (V, \mathcal{A})$ be a mixed graph. Put $X = \lambda I - H_\eta(G)$. Then

$$\det X = \sum_{\sigma \in \text{Sym}(V)} \text{sgn}(\sigma) \prod_{x \in V} X_{x, \sigma(x)},$$

where $\text{Sym}(V)$ is the set of all permutations of V . Let

$$\mathcal{P}(X) := \left\{ \sigma \in \text{Sym}(V) \mid \prod_{x \in V} X_{x, \sigma(x)} \neq 0 \right\}.$$

In addition, we define

$$\mathcal{P}_m(X) := \left\{ \sigma \in \mathcal{P}(X) \mid |\{x \in V \mid \sigma(x) \neq x\}| = m \right\}$$

for $m \in \mathbb{N}$. Any permutation $\sigma \in \text{Sym}(V)$ can be expressed as the product of disjoint cyclic permutations, say $\sigma = \sigma_1 \sigma_2 \cdots \sigma_l$ for some $l \in \mathbb{N}$. We call each σ_i a *factor* of σ .

Lemma 3.1. *Let $G = (V, \mathcal{A})$ be a mixed graph with girth $s+1$, and let $X = \lambda I - H_\eta(G)$. For $l \in \{1, 2, \dots, s\}$, all factors of $\sigma \in \mathcal{P}_l(X)$ are transpositions, where $n = |V|$.*

Proof. Let $H_\eta = H_\eta(G)$. Suppose $\sigma \in \mathcal{P}_l(X)$ has a factor σ_i of length $t > 2$. Display as $\sigma_i = (x_1 x_2 \cdots x_t)$. Then $(H_\eta)_{x_1, x_2} (H_\eta)_{x_2, x_3} \cdots (H_\eta)_{x_t, x_1} \neq 0$. However, G has no t -cycles since $t \leq l \leq s$. Thus, at least one of $(H_\eta)_{x_1, x_2}, (H_\eta)_{x_2, x_3}, \dots, (H_\eta)_{x_t, x_1}$ is 0. This is a contradiction. $\square$

On the other hand, for distinct vertices x, y in a mixed graph G , we have

$$(H_\eta)_{x,y} (H_\eta)_{y,x} = |(H_\eta)_{x,y}|^2 = \begin{cases} 1 & \text{if } x \text{ is adjacent to } y \text{ in } G^\pm, \\ 0 & \text{otherwise,} \end{cases} \quad (3.1)$$

since H_η is Hermitian. Remarking this, we have the following.

Proposition 3.2. *Let $G = (V, \mathcal{A})$ be a mixed graph with girth $s+1$. Let*

$$\begin{aligned} \det(\lambda I - H_\eta(G)) &= \lambda^n + a_1 \lambda^{n-1} + \cdots + a_{n-1} \lambda + a_n, \\ \det(\lambda I - H_\eta(G^\pm)) &= \lambda^n + b_1 \lambda^{n-1} + \cdots + b_{n-1} \lambda + b_n, \end{aligned}$$

where $n = |V|$. Then we have $a_l = b_l$ for any $l \in \{1, 2, \dots, s\}$.

Proof. Put $X = \lambda I - H_\eta(G)$ and $Y = \lambda I - H_\eta(G^\pm)$. For $x, y \in V$, $X_{x,y} \neq 0$ if and only if $Y_{x,y} \neq 0$, so $\mathcal{P}(X) = \mathcal{P}(Y)$. In particular, $\mathcal{P}_l(X) = \mathcal{P}_l(Y)$. If $\mathcal{P}_l(X) = \emptyset$, we have $a_l = b_l = 0$. We consider $\mathcal{P}_l(X) \neq \emptyset$. Let $\sigma \in \mathcal{P}_l(X)$. By Lemma 3.1, all factors of σ are transpositions. Display as $\sigma = (x_1 y_1)(x_2 y_2) \cdots (x_t y_t)$, where $t = l/2$. By (3.1),

$$\begin{aligned} \text{sgn}(\sigma) \prod_{x \in V} X_{x, \sigma(x)} &= \lambda^{n-2t} (H_\eta(G))_{x_1, y_1} (H_\eta(G))_{y_1, x_1} \cdots (H_\eta(G))_{x_t, y_t} (H_\eta(G))_{y_t, x_t} \\ &= \lambda^{n-2t} \cdot 1 \\ &= \lambda^{n-2t} (H_\eta(G^\pm))_{x_1, y_1} (H_\eta(G^\pm))_{y_1, x_1} \cdots (H_\eta(G^\pm))_{x_t, y_t} (H_\eta(G^\pm))_{y_t, x_t} \\ &= \text{sgn}(\sigma) \prod_{x \in V} Y_{x, \sigma(x)}. \end{aligned}$$

Therefore,

$$a_l = \sum_{\sigma \in \mathcal{P}_l(X)} \text{sgn}(\sigma) \prod_{x \in V} X_{x, \sigma(x)} = \sum_{\sigma \in \mathcal{P}_l(Y)} \text{sgn}(\sigma) \prod_{x \in V} Y_{x, \sigma(x)} = b_l.$$

$\square$

Corollary 3.3. *Let G be a mixed tree. Then*

$$\det(\lambda I - H_\eta(G)) = \det(\lambda I - H_\eta(G^\pm)),$$

i.e., G and $G^\pm$ are H_η -cospectral.

Proof. Since G is a mixed tree, the girth is ∞ . By Proposition 3.2, all coefficients of both characteristic polynomials are equal. $\square$

For distinct vertices x, y in a mixed graph G , we also have

$$(\tilde{H}_\eta)_{x,y}(\tilde{H}_\eta)_{y,x} = |(\tilde{H}_\eta)_{x,y}|^2 = \begin{cases} \frac{1}{\deg x \deg y} & \text{if } x \text{ is adjacent to } y \text{ in } G^\pm, \\ 0 & \text{otherwise.} \end{cases}$$

Therefore, the same result as Proposition 3.2 holds for $\tilde{H}_\eta$.

Proposition 3.4. *Let $G = (V, \mathcal{A})$ be a mixed graph with girth $s + 1$. Let*

$$\begin{aligned} \det(\lambda I - \tilde{H}_\eta(G)) &= \lambda^n + a_1 \lambda^{n-1} + \cdots + a_{n-1} \lambda + a_n, \\ \det(\lambda I - \tilde{H}_\eta(G^\pm)) &= \lambda^n + b_1 \lambda^{n-1} + \cdots + b_{n-1} \lambda + b_n, \end{aligned}$$

where $n = |V|$. Then we have $a_l = b_l$ for any $l \in \{1, 2, \dots, s\}$. In particular, if G is a mixed tree, then

$$\det(\lambda I - \tilde{H}_\eta(G)) = \det(\lambda I - \tilde{H}_\eta(G^\pm)),$$

i.e., G and $G^\pm$ are $\tilde{H}_\eta$ -cospectral.

4. CLASSIFICATION OF MIXED CYCLES BY H_η -SPECTRA

The relation that two mixed graphs are H_η -cospectral is an equivalence relation. We call its equivalence class H_η -cospectral class. In this section, we determine the equivalence classes in the mixed cycles.


Let $G = (V, \mathcal{A})$ be a mixed graph. A function $\alpha : V \rightarrow \{1, e^{\pm i\eta}\}$ is called a *switching function*. For a switching function α , we define the matrix $D(\alpha) \in \mathbb{C}^{V \times V}$ by $D(\alpha)_{x,y} = \alpha(x)\delta_{x,y}$. Taking a switching function α well, the matrix $D(\alpha)H_\eta(G)D(\alpha)^*$ is the η -Hermitian adjacency matrix of another mixed graph G' . Then, we say that G' is obtained by switching with respect to α from G . Clearly, G and G' are H_η -cospectral. Note that if a mixed graph G' is obtained by switching with respect to α from a mixed graph G , we also say that G and G' are *switching equivalent*. Recent studies related to switching equivalence of mixed graphs are in [17, 37].

4.1. Switching functions. In [4], Akbari et al defined the four typical switching functions and determined the $H_{\frac{\pi}{2}}$ -cospectral classes in the mixed cycles. We generalize their result to general $\eta \in [0, 2\pi)$. Let $G = (V, \mathcal{A})$ be a mixed cycle. First, we define the three typical switching functions as follows:

Sw.2'. For a vertex $x \in V$, define the switching function

$$\alpha(v) = \begin{cases} e^{i\eta} & \text{if } v = x, \\ 1 & \text{otherwise.} \end{cases}$$


Let $N_{G^\pm}(x) = \{v_1, v_2\}$. If $(v_1, x), (v_2, x) \in \mathcal{A} \setminus \mathcal{A}^{-1}$, then we have the mixed graph $(V, \mathcal{A} \cup \{(v_1, x)^{-1}, (v_2, x)^{-1}\})$ by switching.

FIGURE 1. Sw.2' on the vertex x

Sw.3'. For a vertex $x \in V$, define the switching function

$$\alpha(v) = \begin{cases} e^{-i\eta} & \text{if } v = x, \\ 1 & \text{otherwise.} \end{cases}$$


Let $N_{G^\pm}(x) = \{v_1, v_2\}$. If $(x, v_1), (x, v_2) \in \mathcal{A} \setminus \mathcal{A}^{-1}$, then we have the mixed graph $(V, \mathcal{A} \cup \{(x, v_1)^{-1}, (x, v_2)^{-1}\})$ by switching.

FIGURE 2. Sw.3' on the vertex x

Sw.4'. For a vertex $x \in V$, define the switching function

$$\alpha(v) = \begin{cases} e^{i\eta} & \text{if } v = x, \\ 1 & \text{otherwise.} \end{cases}$$


Let $N_{G^\pm}(x) = \{v_1, v_2\}$. If $(v_1, x) \in \mathcal{A} \setminus \mathcal{A}^{-1}$ and $(x, v_2) \in \mathcal{A} \cap \mathcal{A}^{-1}$, then we have the mixed graph $(V, (\mathcal{A} \setminus \{(v_2, x)\}) \cup \{(v_1, x)^{-1}\})$ by switching.

FIGURE 3. Sw.4' on the vertex x

See also Figures 1, 2 and 3. The above switching functions are named after [4]. The lack of “Sw.1'” is due to the generalization of η .

4.2. The $n + 1$ types of mixed cycles. Let $n \in \mathbb{N}$ and let $j \in \{0, 1, \dots, n\}$. We define the *mixed cycle* C_n^j of type j by

$$\begin{aligned} V(C_n^j) &= \{x_1, x_2, \dots, x_n\}, \\ \mathcal{A}(C_n^j) &= \{(x_1, x_2), \dots, (x_j, x_{j+1})\} \cup \{(x_{j+1}, x_{j+2}), \dots, (x_n, x_{n+1})\}^\pm, \end{aligned}$$

FIGURE 4. The mixed cycles C_8^3 and C_8^8

where we set $x_{n+1} = x_1$. We provide the mixed cycles C_8^3 and C_8^8 in Figure 4 as examples.

Note that C_n^0 is the undirected cycle of length n . We will show that any mixed cycle is H_η -cospectral with some type of the mixed cycle, and different types of mixed cycles have different spectra except for a finite number of η .

Lemma 4.1. *Let $G = (V, \mathcal{A})$ be a mixed cycle of length n , and let $x, v_1, v_2, \dots, v_l, y \in V$. If $(x, v_1) \in \mathcal{A} \setminus \mathcal{A}^{-1}$, $(v_1, v_2), (v_2, v_3), \dots, (v_{l-1}, v_l) \in \mathcal{A} \cap \mathcal{A}^{-1}$, and $(v_l, y) \in \mathcal{A}^{-1} \setminus \mathcal{A}$. Then, the mixed cycle $(V, \mathcal{A} \cup \{(x, v_1)^{-1}, (v_l, y)\})$ is obtained by switching from G .*

Proof. We prove by induction on l . Consider $l = 1$. Applying Sw.2' with respect to v_1 , we have the statement. We suppose that the statement follows in the case of $l - 1$. We apply Sw.4' with respect to v_1 . The switched mixed cycle is in the situation of the case of $l - 1$. By the assumption of the induction, we have the statement. $\square$

If $C_n^j = (V, \mathcal{A})$, then we let $(C_n^j)^{-1} = (V, \mathcal{A}^{-1})$.

Proposition 4.2. *Let $G = (V, \mathcal{A})$ be a mixed cycle of length n . Then, there exists $j \in \{0, 1, \dots, n\}$ such that G and C_n^j is H_η -cospectral.*

Proof. By Lemma 4.1, we have a switched mixed cycle G' such that the directions of all arcs are aligned clockwise or anticlockwise. Applying Sw.4' many times, arcs in the graph are replaced consecutively. This graph is C_n^j or $(C_n^k)^{-1}$ for some $j, k \in \{0, 1, \dots, n\}$. Note that $(C_n^k)^{-1}$ is isomorphic to C_n^k . $\square$

Figure 5 shows that switching yields the mixed graph of type 3 from a mixed cycle. Note that Sw.3' is actually unnecessary. However, it can be used to obtain C_n^j with less switching in some cases.

4.3. Determinants of $H_\eta(C_n^j)$. Next, we show that different types of mixed cycles have different spectra. We focus on the constant term of the characteristic polynomial of $H_\eta(C_n^j)$. Let P_n be the undirected path graph on n vertices.

Lemma 4.3. *We have*

$$\det H_\eta(P_n) = (-1)^{\lfloor \frac{n}{2} \rfloor} \frac{1 + (-1)^n}{2}.$$




FIGURE 5. Switching a mixed cycle

Proof. We will show that

$$\det H_\eta(P_n) = \begin{cases} 1 & \text{if } n \equiv 0 \pmod{4}, \\ -1 & \text{if } n \equiv 2 \pmod{4}, \\ 0 & \text{if } n \equiv 1, 3 \pmod{4}. \end{cases}$$

The determinant is

$$\det(H_\eta(P_n)) = \begin{vmatrix} 0 & 1 & 0 & \cdots & \cdots & \cdots & 0 \\ 1 & 0 & 1 & 0 & \cdots & \cdots & 0 \\ 0 & 1 & 0 & 1 & 0 & \cdots & 0 \\ \vdots & \ddots & \ddots & \ddots & \ddots & & \vdots \\ \vdots & & \ddots & \ddots & \ddots & \ddots & 0 \\ 0 & \cdots & \cdots & 0 & 1 & 0 & 1 \\ 0 & \cdots & \cdots & \cdots & 0 & 1 & 0 \end{vmatrix}.$$

We first apply the cofactor expansion along the first row, and we then apply it again along the first column. We have $\det(H_\eta(P_n)) = -\det(H_\eta(P_{n-2}))$. Since $\det H_\eta(P_1) = 0$ and $\det H_\eta(P_2) = -1$, we have the statement. $\square$

Proposition 4.4. *We have*

$$\det H_\eta(C_n^j) = (-1)^{n+1} 2 \cos(\eta j) + (-1)^{\lfloor \frac{n}{2} \rfloor} (1 + (-1)^n).$$

We apply the cofactor expansion along the first column to D_1 . By Corollary 3.3, we have

$$\begin{aligned} D_1 &= (-1)^{1+1} e^{-\eta i} \det H_\eta(P_{n-2}) + (-1)^{(n-1)+1} e^{(j-1)\eta i} \\ &= e^{-\eta i} \det H_\eta(P_{n-2}) + (-1)^n e^{(j-1)\eta i}. \end{aligned}$$

Applying the cofactor expansion along the first column to D_2 , we have

$$\begin{aligned} D_2 &= (-1)^{1+1} e^{-\eta i} \cdot e^{-(j-1)\eta i} + (-1)^{(n-1)+1} \det H_\eta(P_{n-2}) \\ &= e^{-j\eta i} + (-1)^n \det H_\eta(P_{n-2}). \end{aligned}$$

Therefore, by Lemma 4.3,

$$\begin{aligned} \det H_\eta(C_n^j) &= -e^{\eta i} D_1 + (-1)^{n+1} D_2 \\ &= (-1)^{n+1} 2 \cos(\eta j) + (-2) \det H_\eta(P_{n-2}) \\ &= (-1)^{n+1} 2 \cos(\eta j) + (-2) \cdot (-1)^{\lfloor \frac{n-2}{2} \rfloor} \frac{1 + (-1)^{n-2}}{2} \\ &= (-1)^{n+1} 2 \cos(\eta j) + (-1)^{\lfloor \frac{n}{2} \rfloor} (1 + (-1)^n). \end{aligned}$$

□

Proposition 4.2 and Proposition 4.4 derive Theorem 1.1, which is our first main theorem. In addition, Proposition 4.4 yields that, except for a finite number of η ,

$$\det(\lambda I - H_\eta(C_n^j)) \neq \det(\lambda I - H_\eta(C_n^k))$$

for $j \neq k$. We supplement the phrase “except for a finite number of η ” here. For example, we consider $\eta = \frac{\pi}{2}$ and the mixed cycles of length 4. We have

$$\begin{aligned} \det H_{\frac{\pi}{2}}(C_4^j) &= 2 - 2 \cos \frac{j\pi}{2} \\ &= \begin{cases} 0 & \text{if } j \in \{0, 4\}, \\ 2 & \text{if } j \in \{1, 3\}, \\ 4 & \text{if } j = 2. \end{cases} \end{aligned}$$

As this example points out, when $\eta \in \mathbb{Q}\pi$ and the denominator of η/π is smaller than a given n , the determinants could be equal for different types of mixed cycles. There are only a finite number of such η for given n . This is the reason why the only three types of mixed cycles appeared in the study of [4].

4.4. Characteristic polynomials of $H_\eta(C_n^j)$. On the other hand, if the determinants are same, the characteristic polynomials of mixed cycles are actually equal. We find the characteristic polynomial of C_n^j . An *elementary subgraph* of an undirected graph Γ is a subgraph of Γ such that every component is either K_2 or an undirected cycle. Let $\mathcal{H}(\Gamma)$ denote the set of all the elementary subgraphs of Γ , and let $\mathcal{H}_l(\Gamma)$ denote the set of all the elementary subgraphs of Γ on l vertices. The *rank* and *corank* of an

undirected graph $\Gamma = (V, E)$ are, respectively, $r(\Gamma) = |V| - c$ and $s(\Gamma) = |E| - |V| + c$, where c is the number of components of Γ . Let C_n be the undirected cycle graph on n vertices.

Proposition 4.5. *We have*

$$\det(\lambda I - H_\eta(C_n^j)) = \sum_{k=0}^{\lfloor \frac{n-1}{2} \rfloor} (-1)^k \frac{n}{n-k} \binom{n-k}{k} \lambda^{n-2k} - 2 \cos(\eta j) + (-1)^{\lfloor \frac{n}{2} \rfloor} (1 + (-1)^n).$$

Proof. Let

$$\begin{aligned} \det(\lambda I - H_\eta(C_n^j)) &= \lambda^n + a_1 \lambda^{n-1} + \cdots + a_{n-1} \lambda + a_n, \\ \det(\lambda I - H_\eta(C_n)) &= \lambda^n + b_1 \lambda^{n-1} + \cdots + b_{n-1} \lambda + b_n. \end{aligned}$$

Fix $l \in \{1, \dots, n-1\}$. Since the girth of C_n^j is n , we have $a_l = b_l$ by Proposition 3.2. Also,

$$\mathcal{H}_l(C_n) = \begin{cases} \{\Gamma' \in \mathcal{H}(C_n) \mid \Gamma' \simeq \frac{l}{2} K_2\} & \text{if } l \text{ is even,} \\ \emptyset & \text{if } l \text{ is odd,} \end{cases} \quad (4.1)$$

where $\Gamma' \simeq \frac{l}{2} K_2$ denotes that the graph Γ' is isomorphic to the disjoint union of the $\frac{l}{2}$ complete graphs K_2 . Thus, $b_l = 0$ if l is odd. In addition, if l is even, by Proposition 7.1 in [5] and (4.1),

$$\begin{aligned} b_l &= \sum_{\Gamma' \in \mathcal{H}_l(C_n)} (-1)^{r(\Gamma')} 2^{s(\Gamma')} \\ &= \sum_{\Gamma' \in \mathcal{H}_l(C_n)} (-1)^{\frac{l}{2}} \\ &= (-1)^{\frac{l}{2}} |\{\frac{l}{2}\text{-matching in } C_n\}| \\ &= (-1)^{\frac{l}{2}} \frac{n}{n - \frac{l}{2}} \binom{n - \frac{l}{2}}{\frac{l}{2}}. \end{aligned}$$

The last equality is given by Exercises in p.14 of [7]. By Proposition 4.4, the characteristic polynomial $\det(\lambda I - H_\eta(C_n^j))$ is

$$\begin{aligned} & \sum_{k=0}^{\lfloor \frac{n-1}{2} \rfloor} (-1)^k \frac{n}{n-k} \binom{n-k}{k} \lambda^{n-2k} + (-1)^n \{(-1)^{n+1} 2 \cos(\eta j) + (-1)^{\lfloor \frac{n}{2} \rfloor} (1 + (-1)^n)\} \\ &= \sum_{k=0}^{\lfloor \frac{n-1}{2} \rfloor} (-1)^k \frac{n}{n-k} \binom{n-k}{k} \lambda^{n-2k} - 2 \cos(\eta j) + (-1)^{\lfloor \frac{n}{2} \rfloor + n} (1 + (-1)^n) \\ &= \sum_{k=0}^{\lfloor \frac{n-1}{2} \rfloor} (-1)^k \frac{n}{n-k} \binom{n-k}{k} \lambda^{n-2k} - 2 \cos(\eta j) + (-1)^{\lfloor \frac{n}{2} \rfloor} (1 + (-1)^n). \end{aligned}$$

We note that $(-1)^{\lfloor \frac{n}{2} \rfloor + n} \neq (-1)^{\lfloor \frac{n}{2} \rfloor}$, but $(-1)^{\lfloor \frac{n}{2} \rfloor + n}(1 + (-1)^n) = (-1)^{\lfloor \frac{n}{2} \rfloor}(1 + (-1)^n)$ in the last calculation. We have the statement. $\square$

4.5. Mixed graphs as $\mathbb{T}$ -gain graphs. In this subsection, we briefly touch gain graphs. The η -Hermitian adjacency matrix is also seen as an adjacency matrix of a $\mathbb{T}$ -gain graph. We refer to [30, 33] for readers. We use notations and terminologies in [30, 33]. The η -Hermitian adjacency matrix of a mixed graph $G = (V, \mathcal{A})$ is the adjacency matrix of the $\mathbb{T}$ -gain graph $\Phi = (G^\pm, \mathbb{T}, \varphi)$ defined by the $\mathbb{T}$ -gain $\varphi : \mathcal{A}^\pm \rightarrow \mathbb{T}$ such that

$$\varphi(a) = \begin{cases} 1 & \text{if } a \in \mathcal{A} \cap \mathcal{A}^{-1}, \\ e^{\eta i} & \text{if } a \in \mathcal{A} \setminus \mathcal{A}^{-1}, \\ e^{-\eta i} & \text{if } a \in \mathcal{A}^{-1} \setminus \mathcal{A}, \\ 0 & \text{otherwise.} \end{cases}$$

In [11, 30], the authors mentioned the determinants of the adjacency matrices of $\mathbb{T}$ -gain graphs. In addition, the coefficients of the characteristic polynomials are also found by using principal minors. The discussions in this section can also be carried out in terms of $\mathbb{T}$ -gain graphs.

5. PRELIMINARIES ON QUANTUM WALKS DEFINED BY MIXED GRAPHS

Let $\eta \in [0, 2\pi)$, and let $G = (V, \mathcal{A})$ be a mixed graph. The η -function $\theta : \mathcal{A}^\pm \rightarrow \mathbb{R}$ of a mixed graph G is defined by

$$\theta(a) = \begin{cases} \eta & \text{if } a \in \mathcal{A} \setminus \mathcal{A}^{-1}, \\ -\eta & \text{if } a \in \mathcal{A}^{-1} \setminus \mathcal{A}, \\ 0 & \text{if } a \in \mathcal{A} \cap \mathcal{A}^{-1}. \end{cases}$$

Note that $\theta(a^{-1}) = -\theta(a)$ for any $a \in \mathcal{A}^\pm$.

5.1. Several matrices on quantum walks defined by mixed graphs. In [20], the authors provided a quantum walk defined by a mixed graph. Let $G = (V, \mathcal{A})$ be a mixed graph equipped with an η -function θ . We define several matrices (operators) on quantum walks. The *boundary operator* $K = K(G) \in \mathbb{C}^{V \times \mathcal{A}^\pm}$ is defined by

$$K_{x,a} = \frac{1}{\sqrt{\deg x}} \delta_{x,t(a)}.$$

The *coin operator* $C = C(G) \in \mathbb{C}^{\mathcal{A}^\pm \times \mathcal{A}^\pm}$ is defined by $C = 2K^*K - I$. The *shift operator* $S_\theta = S_\theta(G) \in \mathbb{C}^{\mathcal{A}^\pm \times \mathcal{A}^\pm}$ is defined by $(S_\theta)_{a,b} = e^{\theta(b)i} \delta_{a,b^{-1}}$. Define the *time evolution matrix* $U_\theta = U_\theta(G) \in \mathbb{C}^{\mathcal{A}^\pm \times \mathcal{A}^\pm}$ by $U_\theta = S_\theta C$.

Lemma 5.1. *Let $G = (V, \mathcal{A})$ be a mixed graph equipped with an η -function θ . We have*

$$(U_\theta)_{a,b} = e^{-\theta(a)i} \left(\frac{2}{\deg_G t(b)} \delta_{o(a),t(b)} - \delta_{a,b^{-1}} \right)$$

for any $a, b \in \mathcal{A}^\pm$.

Proof. Indeed,

$$\begin{aligned}
(U_\theta)_{a,b} &= (2S_\theta K^* K - S_\theta)_{a,b} \\
&= 2(S_\theta K^* K)_{a,b} - (S_\theta)_{a,b} \\
&= 2 \sum_{z \in \mathcal{A}} \sum_{x \in V} (S_\theta)_{a,z} (K^*)_{z,x} K_{x,b} - e^{\theta(b)i} \delta_{a,b^{-1}} \\
&= 2 \sum_{z \in \mathcal{A}} \sum_{x \in V} e^{\theta(z)i} \frac{1}{\sqrt{\deg x}} \frac{1}{\sqrt{\deg x}} \delta_{a,z^{-1}} \delta_{x,t(z)} \delta_{x,t(b)} - e^{\theta(a^{-1})i} \delta_{a,b^{-1}} \\
&= 2 \sum_{x \in V} e^{\theta(a^{-1})i} \frac{1}{\deg x} \delta_{x,t(a^{-1})} \delta_{x,t(b)} - e^{\theta(a^{-1})i} \delta_{a,b^{-1}} \\
&= \frac{2e^{\theta(a^{-1})i}}{\deg_G t(b)} \delta_{o(a),t(b)} - e^{\theta(a^{-1})i} \delta_{a,b^{-1}} \\
&= e^{-\theta(a)i} \left(\frac{2}{\deg_G t(b)} \delta_{o(a),t(b)} - \delta_{a,b^{-1}} \right).
\end{aligned}$$

□

The following is an important theorem that links quantum walks and spectral graph theory. We cite [20]. In [14], Higuchi et al proved a similar claim in more general models.

Theorem 5.2 ([20]). *Let $G = (V, \mathcal{A})$ be a mixed graph equipped with an η -function θ , and let U_θ be the time evolution matrix. Then we have*

$$\text{Spec}(U_\theta) = \{e^{\pm i \cos^{-1}(\lambda)} \mid \lambda \in \text{Spec}(\tilde{H}_\eta(G))\} \cup \{1\}^{M_1} \cup \{-1\}^{M_{-1}},$$

where

$$\begin{aligned}
M_1 &= \frac{1}{2} |\mathcal{A}^\pm| - |V| + \dim \ker(\tilde{H}_\eta(G) - I), \\
M_{-1} &= \frac{1}{2} |\mathcal{A}^\pm| - |V| + \dim \ker(\tilde{H}_\eta(G) + I).
\end{aligned}$$

The operators (matrices) used in our quantum walks are summarized in Table 2, where $G = (V, \mathcal{A})$ is a mixed graph equipped with an η -function θ .

Notation	Name	Indices of rows and columns	Definition
K	Boundary	$V \times \mathcal{A}^\pm$	$K_{x,a} = \frac{1}{\sqrt{\deg x}} \delta_{x,t(a)}$
C	Coin	$\mathcal{A}^\pm \times \mathcal{A}^\pm$	$C = 2K^* K - I$
S_θ	Shift	$\mathcal{A}^\pm \times \mathcal{A}^\pm$	$(S_\theta)_{ab} = e^{\theta(b)i} \delta_{a,b^{-1}}$
U_θ	Time evolution	$\mathcal{A}^\pm \times \mathcal{A}^\pm$	$U_\theta = S_\theta C$

TABLE 2. The operators (matrices) used in our quantum walk

5.2. Necessary and sufficient conditions on periodicity. Let U_θ be a time evolution matrix of a mixed graph G equipped with an η -function θ . We say that G is *periodic* if there exists $\tau \in \mathbb{N}$ such that $U_\theta^\tau = I$. When the mixed graph G is periodic, the *period* is defined by $\min\{\tau \in \mathbb{N} \mid U_\theta^\tau = I\}$.

Lemma 5.3. *Let U_θ be a time evolution matrix of a mixed graph G equipped with an η -function θ . Then, we have*

$$\{\tau \in \mathbb{N} \mid U_\theta^\tau = I\} = \{\tau \in \mathbb{N} \mid \lambda^\tau = 1 \text{ for any } \lambda \in \text{Spec}(U_\theta)\}. \quad (5.1)$$

In particular, G is periodic if and only if there exists $\tau \in \mathbb{N}$ such that $\lambda^\tau = 1$ holds for any eigenvalue λ of U_θ .

Proof. Since U_θ is unitary, there exists a unitary matrix Q such that

$$Q^* U_\theta Q = \text{diag}(\lambda_1, \dots, \lambda_{2m}),$$

where m is the number of edges of $G^\pm$. Thus for $\tau \in \mathbb{N}$,

$$Q^* U_\theta^\tau Q = \text{diag}(\lambda_1^\tau, \dots, \lambda_{2m}^\tau).$$

This implies (5.1). □

Define $\mathbf{e}^{(a)} \in \mathbb{C}^{\mathcal{A}^\pm}$ by $(\mathbf{e}^{(a)})_z = \delta_{a,z}$. Let $\mathcal{E}_{\mathcal{A}^\pm} = \{\mathbf{e}^{(a)} \mid a \in \mathcal{A}^\pm\}$. This is the canonical basis of $\mathbb{C}^{\mathcal{A}^\pm}$.

Lemma 5.4. *Let U_θ be a time evolution matrix of a mixed graph G equipped with an η -function θ . Then, we have*

$$\{\tau \in \mathbb{N} \mid U_\theta^\tau = I\} = \{\tau \in \mathbb{N} \mid U_\theta^\tau \mathbf{e}^{(a)} = \mathbf{e}^{(a)} \text{ for any } \mathbf{e}^{(a)} \in \mathcal{E}_{\mathcal{A}^\pm}\}.$$

In particular, G is periodic if and only if there exists $\tau \in \mathbb{N}$ such that $U_\theta^\tau \mathbf{e}^{(a)} = \mathbf{e}^{(a)}$ holds for any vector $\mathbf{e}^{(a)} \in \mathcal{E}_{\mathcal{A}^\pm}$.

Proof. It is clear that the left-hand side is included in the right-hand side. We show the reverse inclusion. Suppose $U_\theta^\tau \mathbf{e}^{(a)} = \mathbf{e}^{(a)}$ for any vector $\mathbf{e}^{(a)} \in \mathcal{E}_{\mathcal{A}^\pm}$ and for some $\tau \in \mathbb{N}$. Number the arc set $\mathcal{A}^\pm$ as $a_1, a_2, \dots, a_{2m}$, where m is the number of edges of $G^\pm$. Let $P = [\mathbf{e}^{(a_1)} \mathbf{e}^{(a_2)} \dots \mathbf{e}^{(a_{2m})}]$. Then, we have $U_\theta^\tau P = P$. Since P is invertible, $U_\theta^\tau = I$ holds. □

6. PERIODICITY OF MIXED PATHS

In this section, we discuss periodicity of mixed paths. As a preparation, we introduce notations for expressing dynamics of quantum walk.

Let $G = (V, \mathcal{A})$ be a mixed graph equipped with an η -function θ . We write the components of a vector $\Psi \in \mathbb{C}^{\mathcal{A}^\pm}$ on the arcs of the graph as in Figure 6. If a component of Ψ is 0, we omit the arc itself corresponding to the component. If a component of Ψ is 1, we may omit the value on the arc corresponding to the component.







FIGURE 6. The components of a vector written on the arcs of the graph


FIGURE 7. Mixed graph G

We provide an example. Let $G = (V, \mathcal{A})$ be the mixed graph in Figure 7. We consider the η -function θ for $\eta \in [0, 2\pi)$. We focus on the vector $e^{((v_3, v_2))} \in \mathcal{E}_{\mathcal{A}^\pm}$. The actions of the coin operator C and the shift operator S_θ are shown in Figure 8 and 9. Since $U_\theta = S_\theta C$, the action of the time evolution matrix U_θ is as in Figure 10.

FIGURE 8. Action of C FIGURE 9. Action of S_θ

Lemma 6.1. *Let $G = (V, \mathcal{A})$ be a mixed graph equipped with an η -function θ , and let $a \in \mathcal{A}^\pm$. We have the following:*

- (1) *If $\deg t(a) = 1$, then $U_\theta e^{(a)} = e^{\theta(a)i} e^{(a^{-1})}$; and*

FIGURE 10. Action of U_θ

(2) If $\deg t(a) = 2$, then $U_\theta \mathbf{e}^{(a)} = e^{-\theta(b)i} \mathbf{e}^{(b)}$, where b is the arc in $\{z \in \mathcal{A}^\pm \mid o(z) = t(a)\} \setminus \{a^{-1}\}$.

Proof. Let $U_\theta = U_\theta(G)$. First, by Lemma 5.1, we have

$$\begin{aligned} (U_\theta \mathbf{e}^{(a)})_z &= \sum_{w \in \mathcal{A}^\pm} (U_\theta)_{z,w} (\mathbf{e}^{(a)})_w \\ &= (U_\theta)_{z,a} \\ &= e^{-\theta(z)i} \left(\frac{2}{\deg_G t(a)} \delta_{o(z), t(a)} - \delta_{z, a^{-1}} \right). \end{aligned}$$

Consider the case of $\deg t(a) = 1$. Then $o(z) = t(a)$ if and only if $z = a^{-1}$. Thus,

$$(U_\theta \mathbf{e}^{(a)})_z = e^{-\theta(a^{-1})i} (2\delta_{z, a^{-1}} - \delta_{z, a^{-1}}) = e^{\theta(a)i} \delta_{z, a^{-1}}.$$

We have $U_\theta \mathbf{e}^{(a)} = e^{\theta(a)i} \mathbf{e}^{(a^{-1})}$. We next consider the case of $\deg t(a) = 2$. Then,

$$\begin{aligned} (U_\theta \mathbf{e}^{(a)})_z &= e^{-\theta(z)i} (\delta_{o(z), t(a)} - \delta_{z, a^{-1}}) \\ &= \begin{cases} 0 & \text{if } z = a^{-1}, \\ e^{-\theta(b)i} & \text{if } z = b, \\ 0 & \text{otherwise.} \end{cases} \end{aligned}$$

Therefore, we have $U_\theta \mathbf{e}^{(a)} = e^{-\theta(b)i} \mathbf{e}^{(b)}$. □

The above lemma is illustrated as in Figure 11 and 12. Now, we discuss periodicity of mixed paths. The following is our second main theorem.




FIGURE 11. Illustration of Lemma 6.1 (1)

Theorem 6.2. Let $G = (V, \mathcal{A})$ be a mixed path on n vertices equipped with an η -function θ . Then G is periodic for any $\eta \in [0, 2\pi)$, and the period is $2(n-1)$.





FIGURE 12. Illustration of Lemma 6.1 (2)

Proof. By Proposition 3.4, G and $G^\pm$ are $\tilde{H}_\eta$ -cospectral since G is a mixed tree. By Theorem 5.2, $U_\theta(G)$ and $U_\theta(G^\pm)$ have the same eigenvalues. From Lemma 5.3, periodicity of G and its period are determined by the eigenvalues of the time evolution matrix. Thus, it is sufficient to discuss only periodicity of $G^\pm$, which is the undirected path graph P_n on n vertices. By Lemma 6.1, we have $U_\theta(P_n)^{2(n-1)} \mathbf{e}^{(a)} = \mathbf{e}^{(a)}$ for any vector $\mathbf{e}^{(a)} \in \mathcal{E}_{\mathcal{A}^\pm}$. The dynamics is as follows:



By Lemma 5.4, we see that G is periodic whose period is $2(n-1)$. $\square$

Note that the periodicity of the undirected paths has actually studied in [21] by eigenvalue analysis. In this paper, we have proven the same fact in a different way.

7. PERIODICITY OF MIXED CYCLES

Finally, we discuss periodicity of mixed cycles. Strategy is similar to the mixed paths, but the discussion is more complicated. Recall that a mixed cycle G on n vertices is H_η -cospectral with C_n^j for some $j \in \{0, 1, \dots, n\}$ by Proposition 4.2. For $m_1, m_2 \in \mathbb{Z}$, the greatest common divisor of m_1 and m_2 is denoted by (m_1, m_2) . Note that $(0, m_1) = m_1$. The following is our third main theorem.

Theorem 7.1. *Let $G = (V, \mathcal{A})$ be a mixed cycle on n vertices equipped with an η -function θ . Then, G is periodic if and only if $\eta \in \mathbb{Q}\pi$. In addition, we suppose that*

$\eta \in \mathbb{Q}\pi$ and the mixed cycle G is H_η -cospectral with C_n^j . Let $\eta = \frac{p}{q}\pi$, where p and q are coprime. Then, the period τ of G is the following:

$$\tau = \begin{cases} \frac{2qn}{(j, 2q)} & \text{if } p \text{ is odd,} \\ \frac{qn}{(j, q)} & \text{if } p \text{ is even.} \end{cases} \quad (7.1)$$

Proof. By Proposition 4.2, G is H_η -cospectral with C_n^j for some $j \in \{0, 1, \dots, n\}$. Since G and C_n^j are 2-regular, they are also $\tilde{H}_\eta$ -cospectral. By Theorem 5.2, $U_\theta(G)$ and $U_\theta(C_n^j)$ have the same eigenvalues. From Lemma 5.3, periodicity is determined by the eigenvalues of the time evolution matrices, so it is sufficient to discuss only periodicity of C_n^j for $j \in \{0, 1, \dots, n\}$. Let $U_\theta = U_\theta(C_n^j)$, and let a be an arbitrary arc of C_n^j . By Lemma 6.1, we have

$$U_\theta^n \mathbf{e}^{(a)} = e^{\pm jn} \mathbf{e}^{(a)}. \quad (7.2)$$

If $\eta \notin \mathbb{Q}\pi$, then $jl\eta \notin 2\pi\mathbb{Z}$ for any $l \in \mathbb{N}$, so the mixed graph is not periodic. On the other hand, we suppose $\eta \in \mathbb{Q}\pi$, say $\eta = \frac{p}{q}\pi$. Then,

$$U_\theta^{2qn} \mathbf{e}^{(a)} = e^{\pm 2qj \cdot \frac{p}{q}\pi i} \mathbf{e}^{(a)} = e^{\pm 2pj\pi i} \mathbf{e}^{(a)} = \mathbf{e}^{(a)}.$$

Thus, the mixed graph is periodic.

Next, we determine the period τ . Let $\eta \in \mathbb{Q}\pi$ and let $\eta = \frac{p}{q}\pi$, where p and q are coprime. By Lemma 6.1, the vector $U_\theta^k \mathbf{e}^{(a)}$ coincides with a complex multiple of $\mathbf{e}^{(a)}$ if and only if k is a multiple of n . Thus, the period τ is a multiple of n , namely, $\tau = ln$ for some $l \in \mathbb{N}$. Then,

$$\mathbf{e}^{(a)} = U_\theta^{ln} \mathbf{e}^{(a)} = e^{\pm lj \cdot \frac{p}{q}\pi i} \mathbf{e}^{(a)},$$

so $\frac{pjl}{q}\pi \in 2\pi\mathbb{Z}$, i.e., $pjl \in 2q\mathbb{Z}$. We would like to find $\min\{l \in \mathbb{N} \mid pjl \in 2q\mathbb{Z}\}$. If $j = 0$, we have $\min\{l \in \mathbb{N} \mid pjl \in 2q\mathbb{Z}\} = 1$, so the period is n . This satisfies (7.1). We assume that $j > 0$ in the discussion below.

First, we consider the case where p is odd. We will show that

$$\min\{l \in \mathbb{N} \mid pjl \in 2q\mathbb{Z}\} = \frac{2q}{(j, 2q)}.$$

Since p is odd and $(p, q) = 1$, we have

$$(p, 2q) = 1. \quad (7.3)$$

Let $d = (j, 2q)$. There exists $j' \in \mathbb{N}$ such that

$$j = j'd \quad (7.4)$$

and

$$(j', 2q) = 1. \quad (7.5)$$

Therefore, by (7.3), (7.4) and (7.5), we have

$$\begin{aligned}
\min\{l \in \mathbb{N} \mid pjl \in 2q\mathbb{Z}\} &= \min\{l \in \mathbb{N} \mid jl \in 2q\mathbb{Z}\} \\
&= \min\{l \in \mathbb{N} \mid j'dl \in 2q\mathbb{Z}\} \\
&= \min\{l \in \mathbb{N} \mid dl \in 2q\mathbb{Z}\} \\
&= \frac{2q}{d}.
\end{aligned}$$

The last equality is obtained from the fact that d is a divisor of $2q$.

Next, we consider the case where p is even. We will show that

$$\min\{l \in \mathbb{N} \mid pjl \in 2q\mathbb{Z}\} = \frac{q}{(j, q)}. \quad (7.6)$$

If $p = 0$, then $q = 1$ since p and q are coprime. We have $\min\{l \in \mathbb{N} \mid pjl \in 2q\mathbb{Z}\} = 1$ and $\frac{q}{(j, q)} = 1$. Equality (7.6) is satisfied in this case. We assume that $p > 0$. Since p is even, we have

$$p = 2p' \quad (7.7)$$

for some $p' \in \mathbb{N}$. Since p and q are coprime,

$$(p', q) = 1. \quad (7.8)$$

Let $d = (j, q)$. There exists $j' \in \mathbb{N}$ such that

$$j = j'd \quad (7.9)$$

and

$$(j', q) = 1. \quad (7.10)$$

Therefore, by (7.7), (7.8), (7.9) and (7.10) we have

$$\begin{aligned}
\min\{l \in \mathbb{N} \mid pjl \in 2q\mathbb{Z}\} &= \min\{l \in \mathbb{N} \mid p'jl \in q\mathbb{Z}\} \\
&= \min\{l \in \mathbb{N} \mid jl \in q\mathbb{Z}\} \\
&= \min\{l \in \mathbb{N} \mid j'dl \in q\mathbb{Z}\} \\
&= \min\{l \in \mathbb{N} \mid dl \in q\mathbb{Z}\} \\
&= \frac{q}{d}.
\end{aligned}$$

We have the statement. □

ACKNOWLEDGEMENTS

This paper is based on the graduation theses of the second and third authors. We would like to thank their advisor, Professor Norio Konno, for his fruitful comments and helpful advice. The first author is supported by JSPS KAKENHI (Grant No. 20J01175).

REFERENCES

- [1] D. Aharonov, A. Ambainis, J. Kempe, U. Vazirani, *Quantum walks on graphs*, Proc. 33rd STOC, ACM (2001), pp. 50–59.
- [2] M. Abudayah, O. Alomari, T. Sander, *Hermitian Adjacency Matrices of Mixed Graphs*, arXiv preprint arXiv:2103.16969 (2021).
- [3] Y. Aharonov, L. Davidovich, N. Zagury, *Quantum random walks*, Physical Review A, 48, (1993), 1687–1690.
- [4] S. Akbari, A. Ghafaria, M. Nahvib, M.A. Nematollahia, *Mixed paths and cycles determined by their spectrum*, Linear Algebra Appl., 586 (2020) 325–346.
- [5] N. Biggs, *Algebraic Graph Theory*, Cambridge University Press, Cambridge (1974).
- [6] L. C. Brown, W. J. Martin, D. Wright, *Continuous Time Quantum Walks on Graphs: Group State Transfer*, arXiv preprint 2103.08837 (2021).
- [7] C. Godsil, *Algebraic Combinatorics*, Chapman & Hall, 1993.
- [8] C. Godsil, *State transfer on graphs*, Discrete Math., 312(1):129–147, 2012.
- [9] L. Grover, *A fast quantum search mechanical algorithm for database search*, Proceedings of the 28th annual ACM symposium on theory of computing, (1996), 212–219.
- [10] S. P. Gudder, *Quantum probability*, Academic Press Inc., (1988).
- [11] Shahul Hameed K, K. A. Germina, *Balance in gain graphs - a spectral analysis*, Linear Algebra and its Applications 436 (2012), no. 5, 1114–1121.
- [12] K. Guo, B. Mohar, *Hermitian adjacency matrix of digraphs and mixed graphs*, J. Graph Theory (2016).
- [13] A. L. Gavriluk, S. Suda, *On the multiplicities of digraph eigenvalues*, arXiv preprint arXiv:1911.11055 (2020).
- [14] Y. Higuchi, N. Konno, I. Sato, E. Segawa, *Spectral and asymptotic properties of Grover walks on crystal lattices*, J. Funct. Anal. 267, 4197–4235 (2014).
- [15] Y. Higuchi, N. Konno, I. Sato, E. Segawa, *Periodicity of the discrete-time quantum walk on a finite graph*, Interdiscip. Inf. Sci., 23, 75–86 (2017).
- [16] N. Ito, T. Matsuyama, T. Tsurii, *Periodicity of Grover walks on complete graphs with self-loops*, Linear Algebra and its Applications, Volume 599, (2020), Pages 121–132.
- [17] M. Kadyan, B. Bhattacharjya, *Switching equivalence of Hermitian adjacency matrix of mixed graphs*, arXiv preprint 2103.13632 (2021).
- [18] T. Kajiwarra, N. Konno, S. Koyama, K. Saito, *Periodicity for the 3-state quantum walk on cycles*, Quantum Inf. Comp. 19, 1081–1088 (2019).
- [19] N. Konno, Y. Shimizu and M. Takei, *Periodicity for the Hadamard walk on cycles*, Interdiscip. Inf. Sci., 23 (1) (2017), pp. 1–8.
- [20] S. Kubota, E. Segawa, T. Taniguchi, *Quantum walks defined by digraphs and generalized Hermitian adjacency matrices*, Quantum Information Processing volume 20, Article number: 95 (2021).
- [21] S. Kubota, E. Segawa, T. Taniguchi, Y. Yoshie, *Periodicity of Grover walks on generalized Bethe trees*, Linear Algebra Its Appl., 554, 371–391 (2018).
- [22] S. Kubota, E. Segawa, T. Taniguchi, Y. Yoshie, *A quantum walk induced by Hoffman graphs and its periodicity*, Linear Algebra and its Application, 579, (2019), 217–236.
- [23] S. Kubota, K. Saito, Y. Yoshie, *A new type of spectral mapping theorem for quantum walks with a moving shift on graphs*, arXiv preprint arXiv:2103.05235 (2021).
- [24] J. Liu, X. Li, *Hermitian-adjacency matrices and Hermitian energies of mixed graphs*, Linear Algebra Appl 466 (2015), 182–207.
- [25] S. Li, Y. Yu, *Hermitian adjacency matrix of the second kind for mixed graphs*, arXiv preprint arXiv:2102.03760 (2021).

- [26] X. Liu, Q. Wang, *Laplacian state transfer in total graphs*, Discrete Mathematics Volume 344, Issue 1, 112139, (2021).
- [27] Y. Li, X. Liu, S. Zhang, S. Zhou, *Perfect state transfer in NEPS of complete graphs*, Discrete Applied Mathematics, Volume 289, 31, Pages 98–114, (2021).
- [28] B. Mohar, *A new kind of Hermitian matrices for digraphs*, Linear Algebra and its Applications 584 (2020): 343–352.
- [29] G. Mograby, M. Derevyagin, G. V Dunne, A. Teplyaev, *Spectra of perfect state transfer Hamiltonians on fractal-like graphs*, Journal of Physics A: Mathematical and Theoretical 54.12 (2021): 125301.
- [30] R. Mehatari, M.R. Kannan, A. Samanta, *On the adjacency matrix of a complex unit gain graph*, Linear Multilinear Algebra (2020).
- [31] Portugal, R., *Quantum Walks and Search Algorithm*, Springer (2013).
- [32] K. Saito, *Periodicity for the Fourier quantum walk on regular graphs*, Quantum Inf. Comput., 19 (2019), pp. 23–34.
- [33] A. Samanta, M. R. Kannan, *On the spectrum of complex unit gain graph*, arXiv preprint arXiv:1908.10668 (2019).
- [34] E. Segawa, A. Suzuki, *Spectral mapping theorem of an abstract quantum walk*, Quantum Information Processing. 18 (11), (2019), 333.
- [35] R. S. Sarkar, A. Mandal, B. Adhikari, *Periodicity of lively quantum walks on cycles with generalized Grover coin*, Linear Algebra and its Applications 604, 399–424 (2020).
- [36] J. Wang, X. Liu, *Laplacian state transfer in edge complemented coronas*, Discrete Applied Mathematics 293 (2021): 1–14.
- [37] Y. Wang, B. Yuan, *On graphs whose orientations are determined by their Hermitian spectra*, The Elect. J. Commin 26.3 (2019).
- [38] Y. Yoshie, *Periodicity of Grover walks on distance-regular graphs*, Graphs Comb., 35 (2019), pp.1305–1321.
- [39] H. Zhan, *An infinite family of circulant graphs with perfect state transfer in discrete quantum walks*, Quantum Information Processing 18.12 (2019): 1–26.

DEPARTMENT OF APPLIED MATHEMATICS, FACULTY OF ENGINEERING, YOKOHAMA NATIONAL UNIVERSITY, HODOGAYA, YOKOHAMA 240-8501, JAPAN

Email address: kubota-sho-bp@ynu.ac.jp

DEPARTMENT OF APPLIED MATHEMATICS, GRADUATE SCHOOL OF ENGINEERING SCIENCE, YOKOHAMA NATIONAL UNIVERSITY, HODOGAYA, YOKOHAMA, 240-8501, JAPAN

Email address: sekido-hiroto-zk@ynu.jp

DEPARTMENT OF APPLIED MATHEMATICS, GRADUATE SCHOOL OF ENGINEERING SCIENCE, YOKOHAMA NATIONAL UNIVERSITY, HODOGAYA, YOKOHAMA, 240-8501, JAPAN

Email address: yata-harunobu-dx@ynu.jp