

# 3-Coloring on Regular, Planar, and Ordered Hamiltonian Graphs

Dario Cavallaro

Till Fluschnik\*

Technische Universität Berlin, Faculty IV,  
Algorithmics and Computational Complexity, Germany.  
cavallaro@campus.tu-berlin.de, till.fluschnik@tu-berlin.de

**Abstract.** We prove that 3-COLORING remains NP-hard on 4- and 5-regular planar Hamiltonian graphs, strengthening the results of Dailey [Disc. Math.'80] and Fleischner and Sabidussi [J. Graph. Theor.'02]. Moreover, we prove that 3-COLORING remains NP-hard on  $p$ -regular Hamiltonian graphs for every  $p \geq 6$  and  $p$ -ordered regular Hamiltonian graphs for every  $p \geq 3$ .

*Keywords.* Hamiltonian cycle, NP-hardness, 2-factor, Hamiltonian-ordered

## 1 Introduction

We study the computational complexity of the following classic NP-complete problem [11] on Hamiltonian graphs (graphs admitting a cycle that visits every vertex exactly once) that are additionally restricted to be planar, regular, and ordered.

**Problem 1.** 3-COLORING (3-COL)

**Input:** An undirected graph  $G = (V, E)$ .

**Question:** Is  $G$  3-colorable?

Figure 1 gives an overview of our results. Dailey [5] proved that 3-COLORING is NP-complete on 4- and 5-regular planar graphs, and Fleischner and Sabidussi [8] proved that 3-COLORING is NP-complete and on 4-regular Hamiltonian graphs. We strengthen these two results by proving that 3-COLORING is NP-hard on 4- and 5-regular planar Hamiltonian graphs (Section 3). Note that 3-COLORING is polynomial-time solvable on graphs of maximum degree three [3]. Moreover, we prove that 3-COLORING is NP-hard on  $p$ -regular Hamiltonian graphs for every  $p \geq 6$  (Section 4). Finally, we prove that 3-COLORING is NP-hard on  $p$ -ordered regular Hamiltonian graphs for every  $p \geq 3$ , which implies NP-hardness on regular  $(p - 1)$ -connected Hamiltonian graphs (Section 5). All our NP-hardness results still hold true if a Hamiltonian cycle is additionally provided as part of the input (recall that computing a Hamiltonian cycle is NP-complete in general [11]).

---

\*Supported by DFG, project TORE (NI/369-18).




Figure 1: Overview of our results. An arrow from a box  $A$  to a box  $B$  describes that  $A$ 's graph class is included in  $B$ 's graph class. All shown NP-hardness results hold true even if a Hamiltonian cycle is provided as part of the input. \* (holds still true if the graph is additionally regular)

**Related Work.** In our previous work [4], we proved that FEEDBACK VERTEX SET remains NP-complete on 4- and 5-regular planar Hamiltonian graphs,  $p$ -regular Hamiltonian graphs for every  $p \in \mathbb{N}_{\geq 6}$ , and  $p$ -Hamiltonian-ordered graphs for every  $p \in \mathbb{N}_{\geq 3}$ . Several classic graph problems are studied on planar regular graphs. For instance, INDEPENDENT SET and DOMINATING SET are proven to be NP-complete on planar 3-regular graphs [12, 14]. Moreover, INDEPENDENT SET is proven to be NP-hard on 3- and 4-regular Hamiltonian graphs [9].  $p$ -Hamiltonian-ordered graphs were introduced by Ng and Schultz [16].

## 2 Preliminaries

We denote by  $\mathbb{N}$  and  $\mathbb{N}_0$  the natural numbers excluding and including zero, respectively. We denote by  $\mathbb{N}_{\geq p}$  with  $p \in \mathbb{N}$  the set  $\mathbb{N}_0 \setminus \{0, \dots, p-1\}$ .

**Graph Theory.** We use basic notations from graph theory [6]. We call graph  $G = (V, E)$  even if  $|V|$  is even, and odd otherwise. The neighborhood  $N_G(v)$  of a vertex  $v \in V$  in  $G$  is the vertex set  $\{w \in V \mid \{v, w\} \in E\}$ . The degree of a vertex  $v$  is  $|N_G(v)|$ . A graph is  $(k)$ -regular if every vertex has the same degree  $(k)$ . A cycle is a 2-regular connected graph. We represent a cycle also by a tuple, that is, the cycle  $C = (v_0, \dots, v_{n-1}, v_0)$  is the cycle with vertex set  $\{v_0, \dots, v_{n-1}\}$  and edge set  $\{\{v_i, v_{i+1 \bmod n}\} \mid i \in \{0, \dots, n-1\}\}$ . Let  $v, w$  be two distinct vertices in  $G = (V, E)$ . The graph obtained by *identifying*  $v$  with  $w$  has vertex set  $(V \setminus \{v, w\}) \cup \{vw\}$ , where  $vw$  is a new vertex, and edge set  $(E \setminus \{e \in E \mid \{v, w\} \cap e \neq \emptyset\}) \cup \{\{vw, x\} \mid x \in (N_G(v) \cup N_G(w)) \setminus \{v, w\}\}$ . A graph is planar if it can be drawn on the two-dimensional plane with no two edges crossing except at their endpoints. A graph is  $p$ -(Hamiltonian)-ordered,  $p \in \mathbb{N}_{\geq 3}$  if for each  $p$ -tuple  $(v_1, \dots, v_p)$  of vertices there exists a (Hamiltonian) cycle that visits the vertices  $x_1, \dots, x_p$  in this order.

A  $k$ -coloring of  $G = (V, E)$  is a function  $f: V \rightarrow \{1, \dots, k\}$ . A  $k$ -coloring  $f$  of  $G$  is valid if for every  $\{v, w\} \in E$  it holds that  $f(v) \neq f(w)$ . Graph  $G$  is  $k$ -colorable if there is a valid  $k$ -




Figure 2: The graphs (a)  $L$ , (b)  $H$ , and (c)  $S$ . For each graph, an extendable partial (without  $y, y'$ ) 3-coloring is given. Moreover, for graph  $H$ , an  $x$ - $x'$  path and a  $y$ - $y'$  path (blue) is depicted covering all vertices from the graph. For graph  $S$ , an  $x$ - $y'$  path and an  $x'-y$  path (blue) is depicted covering all vertices from the graph.

coloring of  $G$ . For  $\emptyset \neq V' \subset V$ , a  $k$ -coloring  $f': V' \rightarrow \{1, \dots, k\}$  is called partial. A partial  $k$ -coloring  $f': V' \rightarrow \{1, \dots, k\}$  is called extendable if there is a partial  $k$ -coloring  $g: V \setminus V' \rightarrow \{1, \dots, k\}$  such that the  $k$ -coloring  $f: V \rightarrow \{1, \dots, k\}$  with  $f(v) = f'(v)$  if  $v \in V'$ , and  $f(v) = g(v)$  if  $v \in V \setminus V'$ , is valid.

### 3 Planar Regular Hamiltonian Graphs

In this section, we prove that 3-COLORING is NP-hard on 4- and 5-regular planar Hamiltonian graphs with a Hamiltonian cycle provided. Note that no planar  $p$ -regular graph with  $p \geq 6$  exists.

#### 3.1 4-regular planar Hamiltonian

We prove the following.

**Theorem 1.** 3-COLORING on 4-regular planar Hamiltonian graphs is NP-hard, even if a Hamiltonian cycle is provided.

Following the idea of Fleischner and Sabidussi [8], we first compute a 2-factor in polynomial time. A 2-factor of a graph  $G$  is a set  $\{Q_1, \dots, Q_t\}$ ,  $t \in \mathbb{N}$ , of cycles such that  $V(G) = \bigcup_{i \in \{1, \dots, t\}} V(Q_i)$ . Then, we iteratively “merge” two cycles from the 2-factor via one of the graphs  $H$  and  $S$  (see Figure 2). By this, in each repetition, we reduce the number of cycles in the 2-factor while preserving a 2-factor. Finally, we obtain a Hamiltonian cycle.

Each of the graphs  $H$  and  $S$ , contains two disjoint subgraphs isomorphic to the graph  $L$  (see Figure 2(a)). We have the following important property of  $L$ .

**Observation 1.** Graph  $L$  is 3-colorable and for every valid 3-coloring  $f: V(L) \rightarrow \{1, 2, 3\}$ , it holds that  $f(x) \neq f(y)$ ,  $f(x) = f(y_1)$ , and  $f(x) \cup f(x_1) \cup f(y_2) = \{1, 2, 3\}$ .

*Proof.* An extendable partial 3-coloring of  $L$  is depicted in Figure 2(a). Let  $f: V(L) \rightarrow \{1, 2, 3\}$  be an arbitrary valid 3-coloring of  $L$ . We have that  $f(x) \notin \{f(x_1), f(x_2)\}$ , and hence  $f(z) = f(x)$ .




Figure 3: Illustration to **Construction 1** of (a) Case 1 and (b) Case 2. Indicated are two different 2-factor components in each upper part (magenta and green) and the “merged” cycle, being part of the new 2-factor with one less component, in the lower part (blue).

Since  $f(z) \notin \{f(y_2), f(y_3)\}$ , we have that  $f(y_1) = f(x)$  and hence  $f(y) \neq f(x)$ . Since  $f(x_2) \neq f(y_3)$ , we have  $f(x_1) \neq f(y_2)$  and thus  $f(x) \cup f(x_1) \cup f(y_2) = \{1, 2, 3\}$ .  $\square$

We have the following important properties on  $H$  and  $S$ .

**Lemma 1.** *Graph  $X \in \{H, S\}$  is 3-colorable and for every valid 3-coloring  $f: V(X) \rightarrow \{1, 2, 3\}$ , it holds that  $f(x) \neq f(y)$ ,  $f(x') \neq f(y')$ , and  $f(x) \neq f(x')$ .*

*Proof.* We distinguish whether  $X = H$  or  $X = S$ .

*Case 1:  $X = H$ .* An extendable partial 3-coloring of  $X$  is depicted **Figure 2(b)**. Let  $f: V(X) \rightarrow \{1, 2, 3\}$  be a valid 3-coloring of  $X$ . Due to **Observation 1**, we know that  $f(x) \neq f(y)$  and  $f(x') \neq f(y')$ . Moreover, we know that  $f(x) = f(y_1)$  and  $f(x') = f(y'_1)$ , and since  $\{y_1, y'_1\} \in E(X)$ , it follows that  $f(x) \neq f(x')$ .

*Case 2:  $X = S$ .* An extendable partial 3-coloring of  $X$  is depicted **Figure 2(c)**. Let  $f: V(X) \rightarrow \{1, 2, 3\}$  be a valid 3-coloring of  $X$ . Due to **Observation 1**, we know that  $f(x) \neq f(y)$  and  $f(x') \neq f(y')$ . Moreover, we know that  $f(x') \cup f(x'_1) \cup f(y'_2) = \{1, 2, 3\}$ . Since  $\{y_1, x'_1, y'_2\} \subseteq N_X(z)$ , it follows that  $f(y_1) \neq f(x')$ . Since  $f(x) = f(y_1)$ , we thus have that  $f(x) \neq f(x')$ .  $\square$

**Definition 1** (Insertion). Let  $G$  be a graph and  $(v', v, u, u') \in V(G)^4$  be a quadruple with  $\{v', v\}$ ,  $\{v, u\}$ ,  $\{u, u'\} \in E(G)$ . An  $X$ -insertion at  $(v', v, u, u')$  with  $X \in \{H, S\}$  results in the graph obtained from  $G$  by deleting the edges  $\{v', v\}$ ,  $\{v, u\}$ , and  $\{u, u'\}$ , adding a copy of  $X$  to  $G$ , and identifying  $y'$  with  $v'$ ,  $x'$  with  $v$ ,  $x$  with  $u$ , and  $y$  with  $u'$ .

**Construction 1.** Let  $G$  be a connected planar 4-regular graph, and let  $\mathcal{Q}$  be a 2-factor of  $G$  with  $|\mathcal{Q}| > 1$ . Then there are two distinct cycles  $Q_i, Q_j \in \mathcal{Q}$  with two adjacent vertices  $u$  and  $v$  with  $u \in V(Q_i)$  and  $v \in V(Q_j)$ . We distinguish two cases (see **Figure 3** for an illustration).

**Case 1:** There are two edges  $\{u', u\} \in E(Q_i)$  and  $\{v, v'\} \in E(Q_j)$  sharing the same face (see **Figure 3(a)**). Make an  $H$ -insertion at  $(v', v, u, u')$ .

**Case 2:** There are no two edges  $e \in E(Q_i)$  and  $e' \in E(Q_j)$  with  $u \in e$  and  $v \in e'$  sharing the same face (see [Figure 3\(b\)](#)). Since  $u$  and  $v$  are of degree four, there are two edges  $\{u, u'\} \in E(Q_i)$  and  $\{v, v'\} \in E(Q_j)$  such that each share a face with  $e$ . Make an  $S$ -insertion at  $(v', v, u, u')$ .  $\diamond$

[Lemma 1](#) implies the following.

**Observation 2.** *Let  $G'$  be the graph obtained from a graph  $G$  using [Construction 1](#). Then,  $G$  is 3-colorable if and only if  $G'$  is 3-colorable.*

Garey et al. [10, Theorems 2.1–2.3] proved that 3-COLORING is NP-hard on *connected* planar graphs of maximum vertex degree four. Dailey [5] proved that 3-COLORING remains NP-hard on connected 4-regular planar graphs. We are set to prove the main result of this section.

*Proof of [Theorem 1](#).* Let  $(G)$  with  $G = (V, E)$  be an instance of 3-COL on connected 4-regular planar graphs. Compute a 2-factor  $\mathcal{Q}$  of  $G$  in polynomial time [8, 15]. If  $\mathcal{Q} = \{Q_1\}$ , then return  $(G, Q_1)$ . Otherwise, apply [Construction 1](#) iteratively. In each iteration  $i \geq 1$ , we obtain an equivalent instance ([Observation 2](#)) and a 2-factor, say  $\mathcal{Q}'$ , with  $|\mathcal{Q}'| = |\mathcal{Q}| - i$ . Thus, after at most  $n/3$  steps [8], we obtain a graph  $G'$  and a Hamiltonian cycle  $\mathcal{C}$  of  $G'$ . Finally, return the instance  $(G', \mathcal{C})$ .  $\square$

In the next section, we prove that 3-COLORING is NP-hard on 5-regular planar Hamiltonian graphs. For this, we need an even input graph as we will connect disjoint pairs of vertices with some gadget. Building on [Theorem 1](#), we prove that we can further turn a 4-regular planar Hamiltonian graph into an even 4-regular planar Hamiltonian graph.

**Proposition 1.** *3-COLORING is NP-hard on even 4-regular planar Hamiltonian graphs, even if a Hamiltonian cycle is given.*

To prove [Proposition 1](#), we will add a disjoint copy of the input graph to itself and connect the two graphs using two copies of the graph  $W$  (see [Figure 4\(a\)](#)). The graph  $W$  is inspired by a graph of Dailey [5, Fig. 3]. It is not difficult to see that following holds true for graph  $W$ .

**Observation 3.** *Graph  $W$  is 3-colorable, for every valid 3-coloring  $f: V(W) \rightarrow \{1, 2, 3\}$ , it holds true that  $f(x) = f(x_0) = f(y_0) = f(y)$ , and for every  $c \in \{1, 2, 3\}$ , there is a valid 3-coloring with  $f(x) = c$ .*

**Construction 2.** Let  $(G, \mathcal{C})$  be an instance of 3-COL on 4-regular planar Hamiltonian graphs with graph  $G = (V, E)$  and Hamiltonian cycle  $\mathcal{C}$ . Construct a graph  $G^*$  and a Hamiltonian cycle  $H^*$  of  $G^*$  as follows (see [Figure 4](#) for an illustration). Take  $G$  and a disjoint copy  $G'$  of  $G$ . Let  $e = \{u, v\}$  be an edge in  $\mathcal{C}$  of  $G$ , and let  $e' = \{u', v'\}$  its copy in  $G'$ . Let  $G$  be embedded such that  $e$  is incident with the outer face [1]. Let  $G'$  have the same embedding as  $G$  but mirrored along the  $y$ -axis (see [Figure 4\(b\)](#); we assume such an embedding from now on). Connect the graphs  $G$  and  $G'$  through the vertices  $u, u', v, v'$  as follows (see [Figure 4\(c\)](#)).

Add a copy of the graph  $W$ , and identify  $x$  with  $u$  (call the vertex again  $u$ ) and  $y$  with  $u'$  (call the vertex again  $u'$ ) and rename  $x_0$  by  $u_0$  and  $y_0$  by  $u'_0$ . Let  $v, u_1, u_2, u_3$  be the neighbors of  $u$  in clockwise order. Remove the edges  $\{u, u_2\}$  and  $\{u, u_3\}$ , and add the edges  $\{u_0, u_2\}$  and  $\{u_0, u_3\}$ . Let  $v', u'_1, u'_2, u'_3$  be the neighbors of  $u'$  in counter-clockwise order. Remove the edges  $\{u', u'_2\}$  and  $\{u', u'_3\}$ , and add the edges  $\{u'_0, u'_2\}$  and  $\{u'_0, u'_3\}$ .

Add another copy of the graph  $W$ , and identify  $y$  with  $v$  (call the vertex again  $v$ ) and  $x$  with  $v'$  (call the vertex again  $v'$ ) and rename  $y_0$  by  $v_0$  and  $x_0$  by  $v'_0$ . Let  $u, v_1, v_2, v_3$  be the neighbors of  $v$  in counter-clockwise order. Remove the edges  $\{v, v_2\}$  and  $\{v, v_3\}$ , and add the




Figure 4: Illustration to the proof of **Proposition 1**. (a) the graph  $W$  with a valid 3-coloring, (b) excerpts of the graphs  $G$  and  $G'$  with Hamiltonian cycle  $C$  (magenta) and  $C'$  (green), and (c) excerpt of the graph  $G^*$  with Hamiltonian cycle  $C^*$  (blue).

edges  $\{v_0, v_2\}$  and  $\{v_0, v_3\}$ . Let  $u', v'_1, v'_2, v'_3$  be the neighbors of  $v'$  in clockwise order. Remove the edges  $\{v', v'_2\}$  and  $\{v', v'_3\}$ , and add the edges  $\{v'_0, v'_2\}$  and  $\{v'_0, v'_3\}$ .

We can merge  $C$  and  $C'$  through the two added  $W$ 's as depicted in **Figure 4(c)** to a Hamiltonian cycle  $C^*$  of  $G^*$ .  $\diamond$

*Proof of Proposition 1.* Let  $(G, C)$  be an instance of the NP-hard (**Theorem 1**) 3-COL on 4-regular planar Hamiltonian graphs with graph  $G = (V, E)$  and Hamiltonian cycle  $C$ . If  $G$  is even, then we return  $(G, C)$ . Assume that  $G$  is odd. Construct the graph  $G^*$  with Hamiltonian cycle  $H^*$  from  $G$  using **Construction 2**. Clearly,  $G^*$  is Hamiltonian, 4-regular, planar, and even (note that  $|V(G^*)| = 2|V(G)| + 26$ ). Finally, due to **Observation 3**, we have that  $G$  is 3-colorable if and only if  $G^*$  is 3-colorable.  $\square$

### 3.2 5-regular planar Hamiltonian

In this section, we prove that 3-COLORING is also NP-hard on 5-regular planar Hamiltonian graphs with provided Hamiltonian cycle.

**Theorem 2.** 3-COLORING on 5-regular planar Hamiltonian graphs is NP-hard, even if a Hamiltonian cycle is given.

We will use multiple copies of the graph  $D$  (see **Figure 5**) to turn a planar 4-regular Hamiltonian graph into a planar 5-regular Hamiltonian graph. The graph  $D$  is inspired by a graph of Dailey [5, Fig. 4]. We have the following.

**Observation 4.** Graph  $D$  is 3-colorable and for every distinct  $c, c' \in \{1, 2, 3\}$ , there is a valid 3-coloring  $f: V(D) \rightarrow \{1, 2, 3\}$  with  $f(x) = c$  and  $f(y) = c'$ .




Figure 5: The graph  $D$ . A valid 3-coloring and an  $x$ - $y$  Hamiltonian path (blue) are depicted.




Figure 6: Illustration to the proof of [Theorem 2](#). The magenta path depicts the Hamiltonian cycle before the  $D$ -insertions, and the blue path depicts the Hamiltonian cycle after the  $D$ -insertions.

We will perform a series of  $D$ -insertions defined as follows.

**Definition 2** ( $D$ -insertion). Let  $G$  be a graph and  $\{v, u\} \in E(G)$ . A  $D$ -insertion at  $\{v, u\}$  results in the graph obtained from  $G$  by adding a copy of  $D$  to  $G$ , and identifying  $x$  with  $v$  and  $y$  with  $u$ .

*Proof of [Theorem 2](#).* Let  $(G, \mathcal{C})$  be an instance of the NP-hard ([Proposition 1](#)) 3-COL on even 4-regular planar Hamiltonian graphs with graph  $G = (V, E)$  and Hamiltonian cycle  $\mathcal{C} = (v_1, \dots, v_n)$  of  $G$ . Construct a graph  $G'$  with Hamiltonian cycle  $\mathcal{C}'$  as follows (see [Figure 6](#) for an illustration). For each  $i \in \{1, \dots, n/2\}$ , make a  $D$ -insertion at  $\{v_{2i-1}, v_{2i}\}$ . For  $\mathcal{C}'$ , replace each edge  $\{v_{2i-1}, v_{2i}\}$  by the  $x$ - $y$  Hamiltonian path through the  $D$  inserted at  $\{v_{2i-1}, v_{2i}\}$ . Note that  $G'$  is 5-regular and planar. Finally, due to [Observation 4](#),  $G'$  is 3-colorable if and only if  $G$  is 3-colorable.  $\square$




Figure 7: Illustration to the proof of **Theorem 3**. The input graph  $G$  (left-hand side) and the output graph  $G^*$  (right-hand side). For  $G$  and  $G^*$  are Hamiltonian cycles depicted in magenta and blue, respectively.

## 4 Regular Hamiltonian Graphs

In this section, we prove that 3-COLORING remains NP-hard on  $p$ -regular Hamiltonian graphs for every  $p \in \mathbb{N}_{\geq 4}$ .

**Theorem 3.** *For every  $p \in \mathbb{N}_{\geq 4}$ , 3-COLORING on  $p$ -regular Hamiltonian graphs is NP-hard, even if a Hamiltonian cycle is provided.*

*Proof.* We prove the statement via induction. Due to **Theorems 1** and **2**, we know that 3-COL is NP-hard on  $p$ -regular Hamiltonian graphs for  $p \in \{4, 5\}$ . Assume that the statement is true for  $p' \in \{3, 4, \dots, p\}$ ,  $p \in \mathbb{N}_{\geq 4}$ . Let  $(G, \mathcal{C})$  be an instance of the NP-hard (by induction) 3-COL on  $p$ -regular Hamiltonian graphs with graph  $G = (V, E)$  and Hamiltonian cycle  $\mathcal{C} = (v_0, \dots, v_{n-1})$  of  $G$ , where  $n = |V|$ . We construct a graph  $G^*$  from  $G$  as follows (see **Figure 7**). Add a disjoint copy  $G' = (V', E')$  of  $G$  to  $G$ . Denote the vertices of  $G'$  by  $v'_0, \dots, v'_{n-1}$ , and the copy of  $\mathcal{C}$  in  $G'$  by  $\mathcal{C}'$ . Finally, add the edge set  $\{\{v_i, v'_{i+1 \bmod n}\} \mid i \in \{0, \dots, n-1\}\}$ . Observe that  $G^*$  admits the following Hamiltonian cycle  $\mathcal{C}^* = (v_{n-1}, v'_0, v'_{n-1}, \dots, v'_1, v_0, v_1, \dots, v_{n-2}, v_{n-1})$ . Clearly,  $G^*$  is  $(p+1)$ -regular. Moreover, since  $G \subseteq G^*$ , every valid 3-coloring of  $G^*$  induces a valid 3-coloring of  $G$ . We claim that if  $G$  is 3-colorable, then  $G^*$  is 3-colorable.

Let  $f: V(G) \rightarrow \{1, 2, 3\}$  be a valid 3-coloring of  $G$ . We claim that  $f^*: V(G^*) \rightarrow \{1, 2, 3\}$  with  $f^*(v_i) = f(v_i)$  and  $f^*(v'_i) = f(v_i)$  for all  $i \in \{0, \dots, n-1\}$  is a valid 3-coloring of  $G^*$ . Clearly, for every  $Z \in \{V, V'\}$  and for every  $\{v, w\} \in E(G^*) \cap \binom{Z}{2}$ , we have  $f^*(v) \neq f^*(w)$ . Moreover, for each  $i \in \{0, \dots, n-1\}$ , we have that  $f^*(v_i) \neq f^*(v_{(i+1) \bmod n}) = f^*(v'_{(i+1) \bmod n})$ , and thus each edge  $\{v_i, v'_{(i+1) \bmod n}\}$  has differently colored endpoints. It follows that  $f^*$  is a valid 3-coloring of  $G^*$ .  $\square$

## 5 Ordered and Connected Hamiltonian Graphs

In this section, we prove 3-COLORING to remain NP-hard on  $p$ -ordered Hamiltonian graphs for every  $p \geq 3$ .

**Theorem 4.** *3-COLORING is NP-hard on  $p$ -ordered regular Hamiltonian graphs for every  $p \in \mathbb{N}_{\geq 3}$ , even if a Hamiltonian cycle is given.*

Every Hamiltonian graph is 3-Hamiltonian-ordered and hence 3-ordered Hamiltonian. Starting from here, we will inductively prove **Theorem 4**. In each inductive step, we will construct a graph of connectivity high enough and employ the following.

**Fact 1** ([2, 7]). *If  $G$  is  $22p$ -connected graph for some  $p \in \mathbb{N}_{\geq 3}$ , then  $G$  is  $p$ -ordered.*






Figure 8: Illustration of  $C_q(G)$  with  $q = 3$ .  $G^1, G^2, G^3$  denote the three copies of  $G$ , illustrated through a Hamiltonian cycle of  $G$ . Orange edges connect vertices from  $G^1$  and  $G^2$ , green edges connect vertices from  $G^1$  and  $G^3$ , and gray edges connect vertices from  $G^2$  and  $G^3$ . Only adjacencies along the Hamiltonian cycle are depicted (not all edges are shown). Moreover, a Hamiltonian cycle in  $C_3(G)$  is depicted (blue).

**Construction 3** ( $q$ -complete graph). Let  $G = (V, E)$  and let  $q \in \mathbb{N}$ . Construct the  $q$ -complete graph  $C_q(G) = (V_q, E_q)$  as follows (see Figure 8 for an illustration). Let  $V_q := V^1 \uplus \dots \uplus V^q$  where  $V^i := \{v^i \mid v \in V\}$  for all  $i \in \{1, \dots, q\}$ . Let  $E_q := \bigcup_{i,j \in \{1, \dots, q\}} \{\{u^i, v^j\} \mid \{u, v\} \in E\}$ .  $\diamond$

By construction, we have the following.

**Observation 5.** Let  $q \in \mathbb{N}$ . Then,  $G$  is 3-colorable if and only if  $C_q(G)$  is 3-colorable.

Moreover, the  $q$ -complete graph preserves Hamiltonicity.

**Lemma 2.** Let  $q \in \mathbb{N}$ . If  $G$  admits a Hamiltonian cycle, then  $C_q(G)$  admits a Hamiltonian cycle computable in polynomial time.

*Proof.* Let  $\mathcal{C} = (v_0, \dots, v_{n-1}, v_0)$  be a Hamiltonian cycle of  $G$ . Then

$$\mathcal{C}_q := (v_0^1, \dots, v_{n-1}^1, v_0^2, \dots, v_{n-1}^2, v_0^3, \dots, v_{n-1}^q, v_0^1)$$

is a Hamiltonian cycle in  $C_q(G)$ .  $\square$

Now we argue about the connectivity.

**Lemma 3.** Let  $q \in \mathbb{N}$ . If  $G$  is  $p$ -ordered for some  $p \in \mathbb{N}_{\geq 3}$ , then  $C_q(G)$  is  $q(p-1)$ -connected.

*Proof.* Let  $G$  be  $p$ -ordered for some  $p \in \mathbb{N}_{\geq 3}$ . Let  $v, w \in V$  be two arbitrary vertices of  $G$  and let  $x, y \in \{1, \dots, q\}$  be arbitrary. Since  $G$  is  $p$ -ordered,  $G$  is  $(p-1)$ -connected [16], and thus there are  $p-1$  mutually internally vertex-disjoint  $v$ - $w$  paths  $P_1, \dots, P_{p-1}$ . For all  $i \in \{1, \dots, p-1\}$  and  $j \in \{1, \dots, q\}$ , let  $P_i^j$  denote the copy of path  $P_i$  in  $G^j = (V^j, E^j)$ . In each path  $P_i^j$ , replace  $v^j$  with  $v^x$  and  $w^j$  with  $w^y$ . By this, we obtain  $q(p-1)$  mutually internally vertex-disjoint  $v^x$ - $w^y$ -paths. It follows that  $C_p(G)$  is  $q(p-1)$ -connected [13].  $\square$

We are set to prove the main result of this section.

*Proof of Theorem 4.* We prove the statement via induction on  $p \in \mathbb{N}_{\geq 3}$ . We know that 3-COL is NP-hard on 3-ordered regular Hamiltonian graphs with a Hamiltonian cycle provided. Let the statement hold true for all  $p' \in \{1, \dots, p\}$ ,  $p \in \mathbb{N}_{\geq 3}$ . Let  $(G, \mathcal{C})$  be an instance of the NP-hard (by induction) 3-COL on  $p$ -ordered regular Hamiltonian graphs with Hamiltonian cycle  $\mathcal{C}$ . Let  $q := \lceil 22(p+1)/(p-1) \rceil$ . Compute graph  $G' := C_q(G)$  using Construction 3 in polynomial time. Clearly,  $G'$  is regular. Due to Observation 5, we know that  $G$  is 3-colorable if and only if  $G'$  is 3-colorable. Due to Lemma 2,  $G'$  is Hamiltonian and we can compute a Hamiltonian cycle  $\mathcal{C}'$  of  $G'$  in polynomial time. Finally, due to Lemma 3,  $G'$  is  $q(p-1) \geq 22(p+1)$ -connected, and hence due to Fact 1,  $G'$  is  $(p+1)$ -ordered.  $\square$

## 6 Conclusion

3-COLORING remains NP-hard when requiring a Hamiltonian input graph and a witnessing Hamiltonian cycle in the input, even if the input graph is already restricted to 4- or 5-regular planar, to  $p$ -regular for any  $p \geq 6$ , or to  $p$ -ordered regular for any  $p \geq 3$  graphs. We close with the following.

**Question 1.** Is 3-COLORING NP-hard on  $p$ -Hamiltonian-ordered graphs for every  $p \geq 4$ ?

## References

- [1] Rangaswami Balakrishnan and Kanna Ranganathan. *A textbook of graph theory*. Springer Science & Business Media, 2012. 5
- [2] Béla Bollobás and Andrew Thomason. Highly linked graphs. *Comb.*, 16(3):313–320, 1996. [URL](#) [8](#)
- [3] Rowland Leonard Brooks. On colouring the nodes of a network. In *Mathematical Proceedings of the Cambridge Philosophical Society*, volume 37, pages 194–197. Cambridge University Press, 1941. [1](#), [2](#)
- [4] Dario Cavallaro and Till Fluschnik. Feedback vertex set on Hamiltonian graphs. *CoRR*, abs/2104.05322, 2021. [URL](#) [2](#)
- [5] David P Dailey. Uniqueness of colorability and colorability of planar 4-regular graphs are NP-complete. *Discrete Mathematics*, 30(3):289–293, 1980. [1](#), [5](#), [6](#)
- [6] Reinhard Diestel. *Graph Theory*, volume 173 of *Graduate Texts in Mathematics*. Springer, 4th edition, 2010. 2
- [7] Jill R. Faudree and Ralph J. Faudree. Forbidden subgraphs that imply  $k$ -ordered and  $k$ -ordered hamiltonian. *Discret. Math.*, 243(1-3):91–108, 2002. [URL](#) [8](#)
- [8] Herbert Fleischner and Gert Sabidussi. 3-colorability of 4-regular Hamiltonian graphs. *Journal of Graph Theory*, 42(2):125–140, 2003. [1](#), [3](#), [5](#)
- [9] Herbert Fleischner, Gert Sabidussi, and Vladimir I. Sarvanov. Maximum independent sets in 3- and 4-regular Hamiltonian graphs. *Discrete Mathematics*, 310(20):2742–2749, 2010. [URL](#) [2](#)

- [10] M. R. Garey, David S. Johnson, and Larry J. Stockmeyer. Some simplified NP-complete graph problems. *Theoretical Computer Science*, 1(3):237–267, 1976. [↗](#). URL [↗](#). 5
- [11] Richard M. Karp. Reducibility among combinatorial problems. In *Proceedings of a Symposium on the Complexity of Computer Computations*, The IBM Research Symposia Series, pages 85–103. Plenum Press, New York, 1972. [↗](#). 1
- [12] Tohru Kikuno, Noriyoshi Yoshida, and Yoshiaki Kakuda. The NP-completeness of the dominating set problem in cubic planer graphs. *IEICE TRANSACTIONS (1976-1990)*, 63(6):443–444, 1980. 2
- [13] Karl Menger. Über reguläre Baumkurven. *Mathematische Annalen*, 96(1):572–582, 1927. 9
- [14] Bojan Mohar. Face covers and the genus problem for apex graphs. *Journal of Combinatorial Theory, Series B*, 82(1):102–117, 2001. [↗](#). URL [↗](#). 2
- [15] Henry Martyn Mulder. Julius Petersen’s theory of regular graphs. *Discrete Mathematics*, 100(1-3):157–175, 1992. [↗](#). URL [↗](#). 5
- [16] Lenhard Ng and Michelle Schultz.  $k$ -ordered Hamiltonian graphs. *Journal of Graph Theory*, 24(1):45–57, 1997. [↗](#). URL [↗](#). 2, 9