

Extremal digraphs avoiding distinct walks of length 3 with the same endpoints

Zejun Huang*, Zhenhua Lyu†

Abstract

In this paper, we determine the maximum size of digraphs on n vertices in which there are no two distinct walks of length 3 with the same initial vertex and the same terminal vertex. The digraphs attaining this maximum size are also characterized. Combining this with previous results, we obtain a full solution to a problem proposed by X. Zhan in 2007 .

Key words: digraph, Turán problem, walk

2010 Mathematics Subject Classifications: 05C20, 05C35

1 Introduction and main results

The Turán-type extremal problem is an important topics in extremal graph theory, which concerns the maximum number of edges in graphs containing no given subgraphs and the extremal graphs achieving this maximum. Most of the previous results on Turán problems concern undirected graphs and only a few Turán problems on digraphs have been investigated; see [2, 4, 5, 8, 9, 14, 15] and the references therein.

A *simple digraph* is a digraph that does not contain multiple arcs but allows loops. A *strict digraph* is a loopless simple digraph. Digraphs in this paper are simple unless otherwise stated. We follow the terminology and notations in [1]. Directed walks, directed paths and directed cycles are abbreviate as walks, paths and cycles, respectively. The number of vertices in a digraph is called its *order* and the number of arcs its *size*. Given a digraph D and a set of digraphs F , D is said to be F -free if D contains no digraph in F as its subgraph.

*College of Mathematics and Statistics, Shenzhen University, Shenzhen, 518060, P.R. China. (mathzejun@gmail.com)

†School of Science, Shenyang Aerospace University, Shenyang, 110136, P.R. China. (lyuzhh@outlook.com)

Denote by \vec{K}_n the complete digraph on n vertices. A natural Turán problem on digraphs is determining the maximum size of \vec{K}_r -free digraphs of a given order, which has been solved in [12]. Brown and Harary [4] determined the precise extremal size of digraphs as well as the extremal digraphs that avoids a tournament. They also studied digraphs avoiding a direct sum of two tournaments, or a digraph on at most 4 vertices where any two vertices are joined by at least one arc. In [9], we determined the extremal size of digraphs avoiding an orientation of the 4-cycle as well as the extremal digraphs. By adopting dense matrices, Brown, Erdős, and Simonovits [2, 3] presented asymptotic results on extremal digraphs avoiding a family of digraphs. Howalla, Dabboucy and Tout [6, 7] determined the extremal sizes of the \vec{K}_2 -free digraphs avoiding k paths with the same initial vertex and terminal vertex for $k = 2, 3$. Maurer, Rabinovitch and Trotter [14] studied the extremal \vec{K}_2 -free transitive digraphs which contain at most one path from x to y for any two distinct vertices x, y .

Denote by F_k the family of digraphs consisting of two different walks of length k with the same initial vertex and the same terminal vertex. Let $ex(n, F_k)$ be the maximum size of F_k -free digraphs of order n , and let $EX(n, F_k)$ be the set of F_k -free digraphs of order n with size $ex(n, F_k)$. We are interested in the following Turán-type problem.

Problem 1. *Given positive integers n and k , determine $ex(n, F_k)$ and $EX(n, F_k)$.*

Let $D = (V, A)$ be a digraph with vertex set $V = \{v_1, v_2, \dots, v_n\}$ and arc set A . Its *adjacency matrix* $A_D = (a_{ij})$ is defined by

$$a_{ij} = \begin{cases} 1, & \text{if } (v_i, v_j) \in A; \\ 0, & \text{otherwise.} \end{cases} \quad (1.1)$$

Conversely, given an 0-1 matrix $A = (a_{ij})_{n \times n}$, we can define its digraph $D(A) = (V, A)$ on vertices v_1, v_2, \dots, v_n by (1.1), whose adjacency matrix is A . Considering the entries of $A^k(D)$, Problem 1 is equivalent to the following matrix problem, which was proposed by X. Zhan in 2007; see [17, p.234].

Problem 1'. *Determine the maximum number of ones in a 0-1 matrix A of order n such that A^k is also a 0-1 matrix. Characterize the extremal 0-1 matrices attaining this maximum.*

In 2010, Wu [16] solved Problem 1 for the case $k = 2$. Huang and Zhan [11] solved the case $k \geq n - 1 \geq 4$ and determined $ex(k + 2, F_k)$, $ex(k + 3, F_k)$ when $k \geq 4$. Huang, Lyu and Qiao [10] determined $ex(n, F_k)$ for $4 \leq k \leq n - 4$, and characterized $EX(n, F_k)$ for $5 \leq k \leq n - 4$. They also characterize $EX(k + 2, F_k)$ and $EX(k + 3, F_k)$ when $k \geq 4$. Lyu [13] characterized $EX(n, F_4)$ when $n \geq 8$. In this paper, we solve the last remained case $k = 3$ when $n \geq 16$.

To state our results, we need the following notations and definitions. For a digraph $D = (V, A)$, we denote by $a(D)$ the size of D . For $i, j \in V$, if D contains an arc from i to j ,

we say that j is a *successor* of i , and i is a *predecessor* of j , denoted ij or $i \rightarrow j$. A directed walk $xv_1v_2 \cdots v_k y$ is called a *xy-walk*, where x is its *initial vertex* and y is its *terminal vertex*. For $S, T \subset V$, we denote by $D[S]$ the subgraph of D induced by S , $A(S, T)$ the set of arcs from S to T , $a(S, T)$ the cardinality of $A(S, T)$. Let

$$N^+(u) = \{x \in V | ux \in A\} \quad \text{and} \quad N^-(u) = \{x \in V | xu \in A\}$$

be the out-neighbour set and in-neighbour set a vertex u , respectively. The *out-degree* and *in-degree* of u are $d^+(u) \equiv |N^+(u)|$ and $d^-(u) \equiv |N^-(u)|$, respectively. Let $\Delta^+(D)$ denote the maximum out-degree of D , which are abbreviate as Δ^+ if there is no confusion. Given $X \subseteq V$, $d_X^+(u)$ and $d_X^-(u)$ are the numbers of the successors and predecessors of u from X , respectively.

Two digraphs $D_1 = (V_1, A_1)$ and $D_2 = (V_2, A_2)$ are *isomorphic*, written $D_1 \cong D_2$, if there is a bijection $\sigma : V_1 \rightarrow V_2$ such that $(u, v) \in A_1$ if and only if $(\sigma(u), \sigma(v)) \in A_2$.

For a digraph $D = (V, A)$ with $V = \{v_1, v_2, \dots, v_n\}$, a *blow-up* of D is obtained by replacing every vertex v_i with a finite collection of copies of v_i , denoted V_i , so that xy is an arc for $x \in V_i$ and $y \in V_j$ if and only if $v_i v_j \in A(D)$. A blow-up of a digraph is said to be *balanced* if $|V_i|$ and $|V_j|$ differ by at most one for any pair $i \neq j$. Denote by $B(V_1, V_2)$ the blow-up of an arc with vertex set $V_1 \cup V_2$ such that xy is an arc in $B(V_1, V_2)$ if and only if $x \in V_1, y \in V_2$, and $T(V_1, V_2, V_3)$ the blow-up of the transitive tournament of order 3 such that xy is an arc in $T(V_1, V_2, V_3)$ if and only if $x \in V_i, y \in V_j$ with $i < j$.

Given a digraph D , denote by $D + e$ the digraph obtained from D by adding an arc e , where the arc e is allowed to be a loop. We define $H(V_1, V_2)$ to be the digraph obtained from $B(V_1, V_2)$ by adding an arc or a loop with both ends from V_1 . Let

$$T_{3,n} = \left\{ T(V_1, V_2, V_3) + e : |V_1| + |V_2| + |V_3| = n, |V_i| = \left\lfloor \frac{n}{3} \right\rfloor \text{ or } \left\lceil \frac{n}{3} \right\rceil \text{ for } i = 1, 2, 3, \right. \\ \left. e \text{ is an arc with both ends from } V_2 \right\},$$

which is the set of digraphs obtained from a balanced blow-up of the transitive tournament of order 3 by embedding an arc in the middle partite set. Then each digraph in $T_{3,n}$ has one of the diagrams in Figure 1.




Figure 1: The diagrams of the digraphs in $T_{3,n}$.

Now we state our main result as follows.

Theorem 2. *Let $n \geq 16$ be an integer. Then*

$$ex(n, F_3) = \left\lfloor \frac{n^2}{3} \right\rfloor + 1 \quad \text{and} \quad EX(n, F_3) = T_{3,n}.$$

Notice that $T_{3,n}$ contains some loopless digraphs. By Theorem 2 we have the following result for strict digraphs.

Corollary 3. *Let D be an F_3 -free strict digraphs on $n \geq 16$ vertices. Then*

$$a(D) \leq \left\lfloor \frac{n^2}{3} \right\rfloor + 1$$

with equality if and only if $D \in T_{3,n}$.

Remark. Combining Theorem 2 with previous results, we now obtain a full solution to Problem 1 (Problem 1').

2 Proofs

We will need the following lemmas to prove Theorem 2.

Lemma 4. *Let D be a digraph on $n \geq 8$ vertices such that there are no two walks of length 2 with the same terminal vertex. Then*

$$a(D) \leq \left\lfloor \frac{n^2}{4} \right\rfloor + 1 \tag{2.1}$$

with equality if and only if D is isomorphic to $H(V_1, V_2)$ with $\{|V_1|, |V_2|\} = \{\lfloor n/2 \rfloor, \lceil n/2 \rceil\}$.

Proof. Suppose $D = (V, A)$. Let $V_1 = \{u \in V \mid d^+(u) > 0\}$ and let $V_2 = V \setminus V_1$. Then we have

$$a(V_2, V) = 0 \tag{2.2}$$

and

$$d^-(u) \leq 1 \quad \text{for all } u \in V_1.$$

Let $V'_1 = \{u \in V_1 \mid d^-(u) = 0\}$ and $V''_1 = \{u \in V_1 \mid d^-(u) = 1\}$. Then $V_1 = V'_1 \cup V''_1$ and $V'_1 \cap V''_1 = \emptyset$. Moreover,

$$a(V_1, V_1) = \sum_{u \in V_1} d^-(u) = |V''_1|. \tag{2.3}$$

If V''_1 is empty, we have

$$\begin{aligned} a(D) &= a(V_1, V_2) + a(V_1, V_1) + a(V_2, V) \\ &= a(V_1, V_2) + |V''_1| + 0 \\ &\leq |V_1||V_2| \leq \left\lfloor \frac{n^2}{4} \right\rfloor, \end{aligned} \tag{2.4}$$

which implies (2.1).

Now suppose $|V_1''| \geq 1$. Then we have

$$a(V_1'', V_2) \leq |V_2|.$$

Otherwise $a(V_1'', V_2) > |V_2|$ implies that there exist $u_1, u_2 \in V_1''$ sharing a common successor in V_2 , which contradicts the given condition. Notice that

$$|V_1''||V_2| + 1 \geq |V_2| + |V_1''|.$$

We have

$$\begin{aligned} a(D) &= a(V_1, V_2) + a(V_1, V_1) + a(V_2, V) \\ &= a(V_1', V_2) + a(V_1'', V_2) + |V_1''| \\ &\leq |V_1'||V_2| + |V_2| + |V_1''| \\ &= |V_1||V_2| - |V_1''||V_2| + |V_2| + |V_1''| \\ &\leq |V_1||V_2| + 1 \\ &\leq \left\lfloor \frac{n^2}{4} \right\rfloor + 1. \end{aligned} \tag{2.5}$$

Hence, (2.1) holds.

If equality in (2.1) holds, then $|V_1''| \geq 1$, and all inequalities in (2.5) are equalities. We can deduce that $||V_1| - |V_2|| \leq 1$, $a(V_1', V_2) = |V_1'||V_2|$, $a(V_1'', V_2) = |V_2|$ and $|V_1''| = 1$. Therefore, D is isomorphic to $H(V_1, V_2)$ with $\{|V_1|, |V_2|\} = \{\lfloor n/2 \rfloor, \lceil n/2 \rceil\}$. \square

Corollary 5. *Let D be a digraph on $n \geq 8$ vertices without two walks of length 2 sharing the same terminal vertex such that*

$$a(D) = \left\lfloor \frac{n^2}{4} \right\rfloor. \tag{2.6}$$

Then D is isomorphic to a digraph obtained from $H(V_1, V_2)$ with $\{|V_1|, |V_2|\} = \{\lfloor n/2 \rfloor, \lceil n/2 \rceil\}$ by deleting an arbitrary arc, or $D = H(V_1, V_2)$ with $\{|V_1|, |V_2|\} = \{n/2 - 1, n/2 + 1\}$ provided n is even.

Proof. Define V_1, V_2, V_1', V_1'' as in Lemma 4. If V_1'' is empty, then we have (2.4). By (2.2) and (2.3), equality in (2.4) holds if and only if D is isomorphic to $B(V_1, V_2)$ with $\{|V_1|, |V_2|\} = \{\lfloor n/2 \rfloor, \lceil n/2 \rceil\}$.

If $|V_1''| \geq 1$, then we have (2.5). The condition (2.6) leads to $|V_1''| = 1$ and

$$||V_1| - |V_2|| = \begin{cases} 1, & \text{if } n \text{ is odd;} \\ 0 \text{ or } 2, & \text{if } n \text{ is even.} \end{cases}$$

Moreover, (2.2) and (2.3) implies that there is exactly one arc e in $D[V_1]$ and all the other arcs in D have tails in V_1 and heads in V_2 . If n is odd, then $||V_1| - |V_2|| = 1$ implies that D is isomorphic to a digraph obtained from $B(V_1, V_2) + e \in H(V_1, V_2)$ by deleting an arc in $B(V_1, V_2)$. If n is even, then $|V_1| = |V_2| = n/2$ implies that D is isomorphic to the digraphs obtained from $B(V_1, V_2) + e$ by deleting an arc in $B(V_1, V_2)$, and $||V_1| - |V_2|| = 2$ implies that D is isomorphic to $H(V_1, V_2)$ with $\{|V_1|, |V_2|\} = \{n/2 - 1, n/2 + 1\}$.

□

Now we are ready to present the proof of Theorem 2.

Proof of Theorem 2. Let $D \in T_{3,n}$. Then $D = T(V_1, V_2, V_3) + e$ with $|V_i| = \lfloor n/3 \rfloor$ or $\lceil n/3 \rceil$ for $i = 1, 2, 3$, and e is an arc with both ends from V_2 . It is clear that all walks in D have length ≤ 3 . Moreover, for any walk $v_1 v_2 v_3 v_4$ of length 3, we have $v_2 v_3 = e$. Therefore, there are no two distinct walks of length 3 with the same initial vertex and the same terminal vertex, i.e., D is F_3 -free. Hence,

$$ex(n, F_3) \geq a(D) = \left\lfloor \frac{n^2}{3} \right\rfloor + 1. \quad (2.7)$$

Now let $D = (V, A) \in EX(n, F_3)$. Suppose v_0 is a vertex with $d^+(v_0) = \Delta^+$. It follows from (2.7) that

$$\Delta^+ \geq \left\lfloor \frac{n}{3} \right\rfloor + 1. \quad (2.8)$$

Let $V_1 = N^+(v_0)$ and $V_2 = V \setminus V_1$. We count $a(D)$ in the following way:

$$a(D) = a(V_1, V_1) + a(V_1, V_2) + a(V_2, V) = a(V_1, V_1) + \sum_{u \in V_2} [d^+(u) + d_{V_1}^-(u)]. \quad (2.9)$$

For any $u \in V_2$, if $d_{V_1}^-(u) \geq 2$, then $d^+(u) = 0$ as D is F_3 -free. Therefore,

$$d^+(u) + d_{V_1}^-(u) \leq \Delta^+ + 1 \quad (2.10)$$

for all $u \in V_2$. Moreover, equality in (2.10) holds only if $d^+(u) = \Delta^+$ and $d_{V_1}^-(u) = 1$. Now we prove the following claim.

Claim 1. $d^+(u) + d_{V_1}^-(u) \leq \Delta^+$ for all $u \in V_2$.

Proof of Claim 1. We first assert that there are at most two vertices in V_2 satisfying the equality in (2.10). Suppose otherwise there are three vertices $u_1, u_2, u_3 \in V_2$ satisfying the equality in (2.10). Then we have

$$d_{V_1}^-(u_1) = d_{V_1}^-(u_2) = d_{V_1}^-(u_3) = 1$$

and

$$d^+(u_1) = d^+(u_2) = d^+(u_3) = \Delta^+.$$

Hence, there are $u'_1, u'_2, u'_3 \in V_1$ such that $\{u'_1 u_1, u'_2 u_2, u'_3 u_3\} \subseteq A$. By (2.8), there are two vertices in $\{u_1, u_2, u_3\}$ sharing a common successor. Without loss of generality, we assume $u_1 \rightarrow u$ and $u_2 \rightarrow u$. Then D contains the walks $v_0 \rightarrow u'_1 \rightarrow u_1 \rightarrow u$ and $v_0 \rightarrow u'_2 \rightarrow u_2 \rightarrow u$, a contradiction.

Since D is F_3 -free and V_1 is the out-neighbour set of v_0 , $D[V_1]$ does not contain two distinct walks of length 2 with a common terminal vertex. Applying Lemma 4 on $D[V_1]$, we have

$$a(D[V_1]) = a(V_1, V_1) \leq \left\lfloor \frac{(\Delta^+)^2}{4} \right\rfloor + 1.$$

Combining this with (2.7) and (2.9), we obtain

$$\left\lfloor \frac{(\Delta^+)^2}{4} \right\rfloor + \Delta^+(n - \Delta^+) + 3 \geq a(D) \geq \left\lfloor \frac{n^2}{3} \right\rfloor + 1,$$

which leads to

$$\frac{(\Delta^+)^2}{4} + \Delta^+(n - \Delta^+) + 3 \geq \frac{n^2}{3}.$$

It follows that

$$\left\lceil \frac{2n}{3} \right\rceil - 2 \leq \Delta^+ \leq \left\lfloor \frac{2n}{3} \right\rfloor + 2. \quad (2.11)$$

Suppose that there are $u_1, u_2 \in V_2$ such that

$$d^+(u_1) + d_{V_1}^-(u_1) = d^+(u_2) + d_{V_1}^-(u_2) = \Delta^+ + 1.$$

Then $d^+(u_1) = d^+(u_2) = \Delta^+$ and $d_{V_1}^-(u_1) = d_{V_1}^-(u_2) = 1$, which implies that there are $u'_1, u'_2 \in V_1$ such that $\{u'_1 u_1, u'_2 u_2\} \subseteq A$. By (2.11), u_1 and u_2 share a common successor u . Therefore, D contains the walks $v_0 \rightarrow u'_1 \rightarrow u_1 \rightarrow u$ and $v_0 \rightarrow u'_2 \rightarrow u_2 \rightarrow u$, a contradiction.

Note that $\Delta^+(n - \Delta^+) + \lfloor (\Delta^+)^2/4 \rfloor$ is the size of a complete 3-partite (undirected) graph on n vertices and $\lfloor n^2/3 \rfloor$ is the size of a balanced 3-partite complete (undirected) graph on n vertices. We always have

$$\lfloor n^2/3 \rfloor \geq \Delta^+(n - \Delta^+) + \lfloor (\Delta^+)^2/4 \rfloor. \quad (2.12)$$

Suppose there is only one vertex $u \in V_2$ satisfying the equality in (2.10). Then we get

$$d^+(u) = \Delta^+. \quad (2.13)$$

Moreover, u has exactly one predecessor w in V_1 . By (2.7), (2.9), (2.10) and (2.12), we get

$$\begin{aligned} a(V_1, V_1) &= a(D) - \sum_{u \in V_2} [d^+(u) + d_{V_1}^-(u)] \\ &\geq \left\lfloor \frac{n^2}{3} \right\rfloor - \Delta^+(n - \Delta^+) \\ &\geq \left\lfloor \frac{(\Delta^+)^2}{4} \right\rfloor. \end{aligned}$$

Therefore, we have either

$$a(V_1, V_1) = \left\lfloor \frac{(\Delta^+)^2}{4} \right\rfloor + 1 \quad (2.14)$$

or

$$a(V_1, V_1) = \left\lfloor \frac{(\Delta^+)^2}{4} \right\rfloor. \quad (2.15)$$

If (2.14) holds, then $D[V_1]$ isomorphic to $H(U_1, U_2)$ with $\{|U_1|, |U_2|\} = \{\lfloor \Delta^+/2 \rfloor, \lceil \Delta^+/2 \rceil\}$. If (2.15) holds, applying Corollary 5 on $D[V_1]$, $D[V_1]$ is isomorphic to a digraph obtained from $H(U_1, U_2)$ with $\{|U_1|, |U_2|\} = \{\lfloor \Delta^+/2 \rfloor, \lceil \Delta^+/2 \rceil\}$ by deleting an arbitrary arc, or $D = H(U_1, U_2)$ with $\{|U_1|, |U_2|\} = \{\Delta^+/2 - 1, \Delta^+/2 + 1\}$ provided Δ^+ is even. In all the above cases, there is a subset of V_1 with cardinality $\lceil \Delta^+/2 \rceil - 1$, say V_3 , in which any pair of vertices has a common successor in V_1 .

Since D is F_3 -free, u has at most one successor in V_3 . Now $n \geq 16$, $d^+(u) = \Delta^+$ and (2.11) enforce that u has at least three successors u_1, u_2, u_3 in V_2 . On the other hand, we assert that there is at most one vertex $v \in V_2$ such that

$$d^+(v) + d_{V_1}^-(v) \leq \Delta^+ - 1.$$

Otherwise we have

$$\begin{aligned} a(D) &= a(V_1, V_1) + \sum_{v \in V_2} [d^+(v) + d_{V_1}^-(v)] \\ &\leq \left(\left\lfloor \frac{(\Delta^+)^2}{4} \right\rfloor + 1 \right) + (n - \Delta^+) \Delta^+ - 1 \\ &< \left\lfloor \frac{n^2}{3} \right\rfloor + 1, \end{aligned}$$

a contradiction. Hence, two of u_1, u_2, u_3 , say u_1 and u_2 , satisfy $d^+(u_i) + d_{V_1}^-(u_i) \geq \Delta^+ - 1$. It follows that $d^+(u_i) \geq \Delta^+ - 2$ for $i = 1, 2$. Since u_1 and u_2 have at most one successor in V_3 , they have a common successor x . Therefore, D contains the walks $w \rightarrow u \rightarrow u_1 \rightarrow x$ and $w \rightarrow u \rightarrow u_2 \rightarrow x$, a contradiction.

Therefore, there is no vertex u in V_2 such that the equality in (2.12) holds. This completes the proof of Claim 1. \square

By Claim 1, (2.9) and (2.12), we get

$$\begin{aligned} a(D) &= a(V_1, V_1) + \sum_{u \in V_2} [d^+(u) + d_{V_1}^-(u)] \\ &\leq \left\lfloor \frac{(\Delta^+)^2}{4} \right\rfloor + 1 + (n - \Delta^+) \Delta^+ \end{aligned} \quad (2.16)$$

$$\leq \left\lfloor \frac{n^2}{3} \right\rfloor + 1. \quad (2.17)$$

Combining this with (2.7), we have

$$ex(n, F_3) = a(D) = \left\lfloor \frac{n^2}{3} \right\rfloor + 1. \quad (2.18)$$

Now we characterize the structure of D . The equality (2.18) implies that (2.16) and (2.17) are both equalities. The equality in (2.17) leads to

$$\Delta^+ = \begin{cases} 2n/3, & \text{if } n \equiv 0 \pmod{3}; \\ \lfloor 2n/3 \rfloor \text{ or } \lfloor 2n/3 \rfloor + 1, & \text{otherwise.} \end{cases} \quad (2.19)$$

The equality in (2.16) implies

$$a(V_1, V_1) = \left\lfloor \frac{(\Delta^+)^2}{4} \right\rfloor + 1$$

and

$$d^+(u) + d_{V_1}^-(u) = \Delta^+ \quad \text{for all } u \in V_2.$$

Applying Lemma 4 on $D[V_1]$, $D[V_1]$ is isomorphic to $H(V_3, V_4)$ with $\{|V_3|, |V_4|\} = \{\lfloor \Delta^+/2 \rfloor, \lceil \Delta^+/2 \rceil\}$.

Next we prove

$$a(V_1, V_2) = 0. \quad (2.20)$$

If there is a vertex $u \in V_2$ such that $d_{V_1}^-(u) \geq 2$, then u has no successor, which implies $d_{V_1}^-(u) = \Delta^+$. Therefore, all vertices in V_4 are predecessors of u . Choose any $u_1, u_2 \in V_3$ and $u_3 \in V_4$. We find two walks $v_0 \rightarrow u_1 \rightarrow u_3 \rightarrow u$ and $v_0 \rightarrow u_2 \rightarrow u_3 \rightarrow u$ in D , a contradiction. Therefore, we have

$$d_{V_1}^-(u) \leq 1 \quad \text{and} \quad d^+(u) \geq \Delta^+ - 1 \quad \text{for all } u \in V_2. \quad (2.21)$$

Suppose $d_{V_1}^-(u) = 1$ with w being the predecessor of u in V_1 . Then $d^+(u) = \Delta^+ - 1$. We assert that u has at most one successor in V_3 . Otherwise if u has two successors u_1, u_2 in V_3 , then D contains two walks $w \rightarrow u \rightarrow u_1 \rightarrow u_3$ and $w \rightarrow u \rightarrow u_2 \rightarrow u_3$ for any $u_3 \in V_4$, a contradiction. By (2.21), u has at least two successors in V_2 , say u_1 and u_2 , which have a common successor u_3 . Then D contains two walks $w \rightarrow u \rightarrow u_1 \rightarrow u_3$ and $w \rightarrow u \rightarrow u_2 \rightarrow u_3$, a contradiction. Therefore, we have

$$d_{V_1}^-(u) = 0 \quad \text{and} \quad d^+(u) = \Delta^+ \quad \text{for all } u \in V_2, \quad (2.22)$$

and (2.20) holds.

Finally, we assert

$$a(V_2, V_2) = 0. \quad (2.23)$$

Otherwise suppose $u_1 u_2 \in A(D[V_2])$. If u_2 has two successors in V_2 , say u_3 and u_4 , by (2.19) and (2.22), u_3 and u_4 share a common successor u_5 . Then D contains two walks

$u_1 \rightarrow u_2 \rightarrow u_3 \rightarrow u_5$ and $u_1 \rightarrow u_2 \rightarrow u_4 \rightarrow u_5$, a contradiction. If u_2 has two successors $u_3, u_4 \in V_3$, then those successors share a common successor u_5 in V_4 , and D also contains the above walks, a contradiction again. It follows that

$$d^+(u_2) \leq |V_2| + 2 < \Delta^+,$$

which contradicts (2.22). Hence, we have (2.23).

By (2.20), (2.22) and (2.23), we see that $D[V_2]$ is an empty digraph and all vertices in V_1 are successors of u for all $u \in V_2$. Therefore, $D = T(V_2, V_3, V_4) + e \in T_{3,n}$, where both ends of the arc e are from V_3 . This completes the proof. \square

Acknowledgement

This work was supported by Science and Technology Foundation of Shenzhen City (No. JCYJ20190808174211224)

References

- [1] J.A. Bondy, U.S.R. Murty, Graph Theory with Applications, The Macmillan Press, London, 1976.
- [2] W.G. Brown, P. Erdős, M. Simonovits, Extremal problems for directed graphs, J. Combin. Theory Ser. B 15 (1973) 77-93.
- [3] W.G. Brown, P. Erdős, M. Simonovits, Algorithmic solution of extremal digraph problems, Trans. Amer. Math. Soc. 292 (1985) 421-449.
- [4] W.G. Brown, F. Harary, Extremal digraphs. Combinatorial Theory and Its Applications, I, Proc. Colloq., Balatonfred, 1969, North-Holland, Amsterdam, 1970, 135-198.
- [5] W.G. Brown, M. Simonovits, Extremal multigraph and digraph problems, Paul Erdős and His Mathematics, II (Budapest, 1999), 157-203, Bolyai Soc. Math. Stud., 11, János Bolyai Math. Soc., Budapest, 2002.
- [6] K. Howalla, A. N Dabboucy, R. Tout, On the maximum number of arcs in some classes of graphs, Časopis Pěst. Mat. 107 (1982) 388-392.
- [7] K. Howalla, A. N Dabboucy, R. Tout, An extremal problem for some classes of oriented graphs, Časopis Pěst. Mat. 108 (1983) 53-69.

- [8] Z. Huang, Z. Lyu, 0-1 matrices whose k -th powers have bounded entries, *Linear Multilinear Algebra* 68 (2020) 1972-1982.
- [9] Z. Huang, Z. Lyu, Extremal digraphs avoiding an orientation of C_4 , *Discrete Math.* 343 (2020) 111827.
- [10] Z. Huang, Z. Lyu, P. Qiao, A Turán problem on digraphs avoiding distinct walks of a given length with the same endpoints, *Discrete Math.* 342 (2019) 1703-1717.
- [11] Z. Huang, X. Zhan, Digraphs that have at most one walk of a given length with the same endpoints, *Discrete Math.* 311 (2011) 70-79.
- [12] H. Jacob, H. Meyniel, Extension of Turán's and Brooks' theorems and new notions of stability and coloring in digraphs, *Combinatorial mathematics (Marseille-Luminy, 1981)*, 365-370, *North-Holland Math. Stud.*, 75, *Ann. Discrete Math.*, 17, North-Holland, Amsterdam, 1983.
- [13] Z. Lyu, Extremal digraphs avoiding distinct walks of length 4 with the same endpoints, *Discuss. Math. Graph Theory*, doi:10.7151/dmgt.2321.
- [14] S.B. Maurer, I. Rabinovitch, W.T. Trotter, Jr., A generalization of Turán's theorem to directed graphs, *Discrete Math.* 32 (1980) 167-189.
- [15] A.D. Scott, Subdivisions of transitive tournaments, *European J. Combin.* 21 (2000) 1067-1071.
- [16] H. Wu, On the 0-1 matrices whose squares are 0-1 matrices, *Linear Algebra Appl.* 432 (2010) 2909-2924.
- [17] X. Zhan, *Matrix Theory*, Graduate Studies in Mathematics, vol. 147, American Mathematical Society, Providence, RI, 2013.