

ROUGH SETS IN GRAPHS USING SIMILARITY RELATIONS

IMRAN JAVAID*, SHAHROZ ALI, SHAHID UR REHMAN, AQSA SHAH

ABSTRACT. In this paper, we use theory of rough set to study graphs using the concept of orbits. We investigate the indiscernibility partitions and approximations of graphs induced by orbits of graphs. We also study rough membership functions, essential sets, discernibility matrix and their relationships for graphs.

INTRODUCTION

Rough set theory (RST), introduced by Pawlak [14], provides elegant and powerful techniques to extract information from data associated with various structures. It also provides terminology to minimize the number of significant attributes in data tables, also called information systems. It is always interesting and insightful to know about the objects having similar characteristics to simplify the study of objects under consideration. The objects which are indistinguishable from each other are called information granules. Zadeh introduced and studied information granularity in [27]. Information granules are objects placed together due to similarity of features of interest and are dealt together yielding partition of objects based upon features. Many applications of granular computing appear in connection with fuzzy set theory [9, 18], information science [12, 26], data mining [8, 11, 25], formal concept analysis [10, 25], database theory [7, 19] and rough set theory [1, 15, 16, 17, 22] etc. in fields, such as medicine, economy, finance, business, environment, electrical and computer engineering. Granular computing and rough sets have been studied in context of graphs [4, 5], digraph [13] and hyper graphs [2]. The reader is referred to the following [3, 20, 21, 23] for further reading in this area of study. Chiaselloti et al. [4, 5, 6] have studied well known families of graphs using the notion of neighbourhood of vertices. They have remarked two vertices x, y as indiscernible with respect to a vertex subset \mathcal{A} when these vertices have same neighbours with respect to \mathcal{A} . Symbolically, $x \equiv_{\mathcal{A}} y \Leftrightarrow N(x) \cap \mathcal{A} = N(y) \cap \mathcal{A}$. In [6], they have remarked \mathcal{A} as symmetry axis and two vertices are indiscernible with respect to \mathcal{A} if they are in a ‘symmetrical’ position with respect to all vertices of \mathcal{A} . Equivalence classes induced by indiscernibility relation of \mathcal{A} are the granules and yield partition of the vertex set of the graph, denoted by $\gamma_{\mathcal{A}}(\mathcal{G})$. Rough set theory can be thought of as a particular type of granular computing. Chiaselloti et al. [6] referred the triplet $(\mathcal{G}, \mathcal{A}, \gamma_{\mathcal{A}}(\mathcal{G}))$ as \mathcal{A} -granular reference system. Intersection of granular computing and rough set theory provides tools to analyse graphs, digraphs and hypergraphs. The information about granules will simplify studies associated to graphs and can give important insights about graphs, enriching their applications in various fields and introduce new perspectives.

Key words and phrases. Rough Sets, Automorphisms, Orbit, Reduct, Essential, Discernibility Matrix.

* Corresponding author: imran.javaid@bzu.edu.pk.

By associating rough set and graphs, properties of rough sets can be understood with the help of graphs and those of graphs can be explained by using RST. Graphs have been studied using rough set [3, 5, 13] and rough sets may yield graphs [21]. A strategy developed in one domain can give insights in another domain. It is worth noting that the partition in the granular reference system may be induced using different paradigms and it will give useful insights in various directions. Making comparisons, identifying similarities and studying differences are canonical approaches of study. In this paper, we study graphs using the idea of orbit of vertices and we remark two vertices u, v as indiscernible when orbits intersect with \mathcal{A} in the same set or equivalently two vertices u, v as indiscernible if they belong to same orbits of elements of \mathcal{A} . Chiaselotti et al. [4] remarked two vertices as indiscernible if their open neighbourhoods intersect with a given set \mathcal{A} in the same set. We consider a different approach from that of Chiaselotti et al. [4] and consider similarity relation yielded by graph automorphisms to fill the entries of information table. We will study indiscernibility relations introduced with the help of orbits and prove several results including parallel to those of Chiaselotti et al. [4, 5, 6] using the concept of orbits of vertices. We present introductory terminology needed for the paper first.

A graph \mathcal{G} is an ordered pair consists on two finite sets $\mathcal{V}(\mathcal{G})$ and $\mathcal{E}(\mathcal{G})$ called vertex set and edge set respectively. In this paper, all considered graphs are simple and non-trivial, represented by $\mathcal{G} = (\mathcal{V}(\mathcal{G}), \mathcal{E}(\mathcal{G}))$, when there is no doubt, we simply write $\mathcal{G} = (\mathcal{V}, \mathcal{E})$. Two vertices are called adjacent or neighbours of each other if there is an edge between them. The cardinality of vertex set \mathcal{V} and edge set \mathcal{E} of \mathcal{G} is termed as *order* and *size* of \mathcal{G} respectively. For $x, y \in \mathcal{V}$, $x \sim y$ means x, y are adjacent to each other by an edge and $x \not\sim y$ means x, y are non-adjacent. *Open neighbour* of $x \in \mathcal{V}$ is defined as $\mathcal{N}_{\mathcal{G}}(x) = \{y \in \mathcal{V} : x \sim y\}$. Similarly, *closed neighbour* of x is defined as $\mathcal{N}_{\mathcal{G}}[x] = \mathcal{N}_{\mathcal{G}}(x) \cup \{x\}$.

A *permutation* of a set is a bijection from the set to itself. A graph *automorphism* is a permutation of vertex set that preserves adjacency and non-adjacency of the vertices. Alternatively, $\pi : \mathcal{V}(\mathcal{G}) \rightarrow \mathcal{V}(\mathcal{G})$ is an automorphism of a graph \mathcal{G} if for all $x, y \in \mathcal{V}(\mathcal{G})$, $\pi(x) \sim \pi(y)$ if and only if $x \sim y$. The collection of all automorphisms of graph \mathcal{G} forms a group, named as the *automorphism group* of the graph \mathcal{G} . We use $\Gamma(\mathcal{G})$ or Γ if \mathcal{G} is clear from context, to represent the group of automorphisms of graph \mathcal{G} . For $y \in \mathcal{V}$, the *orbit* of y , denoted by $\mathcal{O}(y)$ is defined as $\mathcal{O}(y) = \{\pi(y) : \pi \in \Gamma\}$ and for any set \mathcal{A} , the orbit of \mathcal{A} is represented by $\mathcal{O}(\mathcal{A})$ is define as $\mathcal{O}(\mathcal{A}) = \cup \mathcal{O}(x_i) \forall x_i \in \mathcal{A}$. Two vertices in same orbit are called *similar* vertices.

A partition γ on a finite set \mathcal{V} is a family of non-empty subsets B_1, B_2, \dots, B_n of \mathcal{V} such that $B_i \cap B_j = \emptyset$ for all $i \neq j$ and $\cup_{i=1}^n B_i = \mathcal{V}$. The subsets B_1, B_2, \dots, B_n are termed as blocks of γ , we write $\gamma := B_1 | B_2 | \dots | B_n$ to represent that γ is a partition of sets with blocks B_1, B_2, \dots, B_n . If $x \in \mathcal{V}$, we denote $\gamma(x)$ as the block of γ containing the element x . An *information table* is denoted by $\mathcal{I} = \langle \mathcal{U}, \mathbb{A}, \text{Val}, \mathcal{F} \rangle$ is a quadruple, where \mathcal{U} represents the universal set of objects, \mathbb{A} represents the attribute set, Val is a set of outcomes and $\mathcal{F} : \mathcal{U} \times \mathbb{A} \rightarrow \text{Val}$ represents an information map. The information table is called *Boolean* if $\text{Val} = \{0, 1\}$. For the purpose of our paper, both universal set \mathcal{U} and attribute set \mathbb{A} are the vertex set \mathcal{V} of the graph \mathcal{G} and the *indiscernibility* relation $\equiv_{\mathcal{A}}$ between the two vertices of the graphs is an equivalence relation which depends upon a set of attributes $\mathcal{A} \subseteq \mathcal{V}$ defined as: $x \equiv_{\mathcal{A}} y$ if and only if for all $z \in \mathcal{A}$, if $z \in \mathcal{O}(x)$ then $z \in \mathcal{O}(y)$. If there exists $z \in \mathcal{A}$

such that z does not belong to $\mathcal{O}(x)$ and $\mathcal{O}(y)$ simultaneously, then we say x and y do not belong the same equivalence class and write it as $\neg(x \equiv_{\mathcal{A}} y)$. Equivalence classes also called \mathcal{A} -granules of object x under consideration of $\equiv_{\mathcal{A}}$ are represented by $\mathcal{O}_{\mathcal{A}}(x)$ and the indiscernibility partition of \mathcal{I} with respect to \mathcal{A} is the family of all equivalence classes, *i.e.*, $\gamma_{\mathcal{A}}(\mathcal{G}) := \{\mathcal{O}_{\mathcal{A}}(x) : x \in \mathcal{V}\}$. Given an attribute subset \mathcal{A} and object subset \mathcal{Q} , lower and upper approximations of \mathcal{Q} by considering the information presented by \mathcal{A} are:

$$\begin{aligned}\mathbb{L}_{\mathcal{A}}(\mathcal{Q}) &= \{x \in \mathcal{V} : \mathcal{O}_{\mathcal{A}}(x) \subseteq \mathcal{Q}\} = \cup\{\mathcal{D} \in \gamma_{\mathcal{A}}(\mathcal{G}) : \mathcal{D} \subseteq \mathcal{Q}\} \\ \mathbb{U}_{\mathcal{A}}(\mathcal{Q}) &= \{x \in \mathcal{V} : \mathcal{O}_{\mathcal{A}}(x) \cap \mathcal{Q} \neq \emptyset\} = \cup\{\mathcal{D} \in \gamma_{\mathcal{A}}(\mathcal{G}) : \mathcal{D} \cap \mathcal{Q} \neq \emptyset\}\end{aligned}$$

In general, $\mathbb{L}_{\mathcal{A}}(\mathcal{Q})$ is subset of \mathcal{Q} while $\mathbb{U}_{\mathcal{A}}(\mathcal{Q})$ is a superset of \mathcal{Q} . A subset \mathcal{Q} is termed as \mathcal{A} -exact if and only if $\mathbb{L}_{\mathcal{A}}(\mathcal{Q}) = \mathbb{U}_{\mathcal{A}}(\mathcal{Q})$, \mathcal{A} -rough otherwise. Let $\mathcal{A}, \mathcal{Q} \subseteq \mathcal{V}$ then rough membership function is defined by:

$$\mu_{\mathcal{Q}}^{\mathcal{A}}(x) = \frac{|\mathcal{O}_{\mathcal{A}}(x) \cap \mathcal{Q}|}{|\mathcal{O}_{\mathcal{A}}(x)|} = \frac{|[x]_{\mathcal{A}} \cap \mathcal{Q}|}{|[x]_{\mathcal{A}}|}.$$

For $\mathcal{A}, \mathcal{D} \subseteq \mathcal{V}$ the positive region is defined as $POS_{\mathcal{A}}(\mathcal{D}) = \{x \in \mathcal{V} : \mathcal{O}_{\mathcal{A}}(x) \subseteq \mathcal{O}_{\mathcal{D}}(x)\}$, and the number $deg_{\mathcal{A}}(\mathcal{D}) = \frac{|POS_{\mathcal{A}}(\mathcal{D})|}{|\mathcal{V}|}$ is called \mathcal{A} -degree dependency of \mathcal{D} .

Important terms associated with rough set theory include the indiscernibility relations, lower and upper approximations, reducts, essential sets and the discernibility matrix. The reduct [14] of an information table consists of the set of attributes which provide same information and characterization as does the set of all attributes. Essential sets and the core [14] of an information table usually consists of the set of attributes which are considered most important in characterization of an information table. The discernibility matrix [20] is a square matrix with i_j^{th} entry consists of attributes for which the attribute value is different for objects x_i and x_j .

This paper is organized as follows: In section 1, we introduce and study the indiscernibility partitions and we also discuss the action of automorphisms in connection with indiscernibility relations. In section 2, we study lower and upper approximations of graphs in terms of orbits. We also examine the rough membership function and dependency measures for graphs. In section 3, we discuss essential sets, discernibility matrix and their relationships. Conclusions are given as last section of the paper.

1. INDISCERNIBILITY PARTITIONS OF GRAPHS

An undirected simple graph \mathcal{G} is described by an information table represented by $\mathcal{I}(\mathcal{G})$. Suppose that the vertex set $\mathcal{V}(\mathcal{G}) = \{x_1, x_2, \dots, x_n\}$ and both universal and attribute sets of information table $\mathcal{I}(\mathcal{G})$ are equal to \mathcal{V} and characterize the information map as follows: for $x_i \in \mathcal{A}$, $\mathcal{F}(x_i, x_j) = 1$ if $x_j \in \mathcal{O}(x_i)$ and $\mathcal{F}(x_i, x_j) = 0$ if $x_j \notin \mathcal{O}(x_i)$.

In theorem 1.0.1, we show how indiscernibility relation $\equiv_{\mathcal{A}}$ and the concept of orbit are related.

Theorem 1.0.1. *For an attribute subset $\mathcal{A} \subseteq \mathcal{V}$, any two vertices $x, y \in \mathcal{V}$ are indiscernible if and only if $\mathcal{O}(x) \cap \mathcal{A} = \mathcal{O}(y) \cap \mathcal{A}$.*

Proof. If $z \in \mathcal{O}(x) \cap \mathcal{A}$ then $\mathcal{F}(x, z) = 1 = \mathcal{F}(y, z)$ by definition it implies that $z \in \mathcal{O}(y) \cap \mathcal{A}$. Hence $\mathcal{O}(x) \cap \mathcal{A} \subseteq \mathcal{O}(y) \cap \mathcal{A}$. Using similar argument, $\mathcal{O}(y) \cap \mathcal{A} \subseteq \mathcal{O}(x) \cap \mathcal{A}$. Hence $\mathcal{O}(x) \cap \mathcal{A} = \mathcal{O}(y) \cap \mathcal{A}$.

Now suppose $\mathcal{O}(x) \cap \mathcal{A} = \mathcal{O}(y) \cap \mathcal{A}$, then $z \in \mathcal{O}(x) \cap \mathcal{A}$ implies that $\mathcal{F}(x, z) = 1$ which implies that $\mathcal{F}(y, z) = 1$. Hence $\mathcal{F}(x, z) = \mathcal{F}(y, z)$, which implies that $x \equiv_{\mathcal{A}} y$. \square

Before, we provide a sketch of indiscernibility partitions of a graph \mathcal{G} in the form of orbits, please note the relationship of similarity of vertices of graph with indiscernibility of vertices. If $\mathcal{O}(x) = \mathcal{O}(y)$ in \mathcal{G} then $x \equiv_{\mathcal{A}} y$ for any attribute subset \mathcal{A} . Moreover, converse is not true in general. Consider a path graph P_5 with the vertex set $\{x_1, x_2, x_3, x_4, x_5\}$ and x_i is adjacent to x_{i+1} for $i = 1, 2, 3, 4$. The orbits of P_5 are $\{x_1, x_5\}$, $\{x_2, x_4\}$ and $\{x_3\}$. For $\mathcal{A} = \{x_1\}$ be an attribute set the \mathcal{A} -equivalence classes are $\{x_1, x_5\}$, $\{x_2, x_3, x_4\}$. Check that $x_2 \equiv_{\mathcal{A}} x_3$ but x_2 is not similar to x_3 in P_5 , hence converse is not true in general.

For a subset \mathcal{A} of \mathcal{V} , set $\mathcal{O}(\mathcal{A}) = \cup_{x \in \mathcal{A}} \mathcal{O}(x)$. In the following proposition, we give a complete sketch for the indiscernibility partitions of graph \mathcal{G} in the form of orbits.

Proposition 1.0.2. *Let \mathcal{G} be a graph and $\mathcal{A} \subseteq \mathcal{V}$ be a given attribute subset. If $\mathcal{B}_{\mathcal{A}}(\mathcal{G}) = (\mathcal{O}(\mathcal{A}))^c$ and $\mathcal{A} = \{x_1, x_2, \dots, x_k\}$ such that $\mathcal{O}(x_i) \cap \mathcal{O}(x_j) = \emptyset$ for each $i \neq j$ and $\mathcal{O}(\mathcal{A}) = \cup_{i=1}^k \mathcal{O}(x_i)$, then $\gamma_{\mathcal{A}}(\mathcal{G}) = \mathcal{B}_{\mathcal{A}}(\mathcal{G})|\mathcal{O}(x_1)|\mathcal{O}(x_2)|\dots|\mathcal{O}(x_k)$.*

Proof. Let $x, y \in \mathcal{V}$ such that $x \equiv_{\mathcal{A}} y$ which gives that $\mathcal{O}(x) \cap \mathcal{A} = \mathcal{O}(y) \cap \mathcal{A}$ and we have the following two cases.

Case 1. Suppose $\mathcal{O}(x) \cap \mathcal{A} = \mathcal{O}(y) \cap \mathcal{A} = \emptyset$, clearly $x, y \in \mathcal{B}_{\mathcal{A}}(\mathcal{G})$ which gives that $\mathcal{B}_{\mathcal{A}}(\mathcal{G})$ is an \mathcal{A} -equivalence class in \mathcal{V} .

Case 2. Suppose $\mathcal{O}(x) \cap \mathcal{A} = \mathcal{O}(y) \cap \mathcal{A} \neq \emptyset$ implies that $x, y \in \mathcal{O}(\mathcal{A}) = \cup_{i=1}^k \mathcal{O}(x_i)$ and $\mathcal{O}(x_i) \cap \mathcal{O}(x_j) = \emptyset$ gives that $x, y \in \mathcal{O}(x_i)$ for some $1 \leq i \leq k$ because if $x \in \mathcal{O}(x_i)$ and $y \in \mathcal{O}(x_j)$ for $i \neq j$ then $\mathcal{O}(x) \cap \mathcal{A} \neq \mathcal{O}(y) \cap \mathcal{A}$. Note that $\mathcal{O}(x) = \mathcal{O}(y)$ implies $x \equiv_{\mathcal{A}} y$, hence $\mathcal{O}_{\mathcal{A}}(x) = \mathcal{O}_{\mathcal{A}}(y) = \mathcal{O}_{\mathcal{A}}(x_i) = \mathcal{O}(x_i)$ and $\mathcal{O}_{\mathcal{A}}(x_i) = \mathcal{O}(x_i)$ for each i .

Concluding above two cases and using the fact that $\mathcal{B}_{\mathcal{A}}(\mathcal{G}), \mathcal{O}(x_1), \mathcal{O}(x_2), \dots, \mathcal{O}(x_k)$ gives a partition of \mathcal{V} , we have $\gamma_{\mathcal{A}}(\mathcal{G}) = \mathcal{B}_{\mathcal{A}}(\mathcal{G})|\mathcal{O}(x_1)|\mathcal{O}(x_2)|\dots|\mathcal{O}(x_k)$. \square

Example 1.0.3. *Consider a graph \mathcal{G} in FIGURE 1.*




FIGURE 1. A Graph \mathcal{G}

In FIGURE 1, subscripts of the vertices are used as labels. We have $\mathcal{O}(0) = \{0, 5, 1\} = \mathcal{O}(5) = \mathcal{O}(1)$, $\mathcal{O}(4) = \{2, 4\} = \mathcal{O}(2)$, $\mathcal{O}(3) = \{3\}$. Fix a vertex subset

$\mathcal{A} = \{1, 5\}$, then we have $0 \equiv_{\mathcal{A}} 1 \equiv_{\mathcal{A}} 5$, and similarly $2 \equiv_{\mathcal{A}} 3 \equiv_{\mathcal{A}} 4$ so $\gamma_{\mathcal{A}}(\mathcal{G}) = 015|234$.

It is an interesting question to identify conditions under which for given two attribute sets \mathcal{A}_1 and \mathcal{A}_2 , $\gamma_{\mathcal{A}_1}(\mathcal{G}) = \gamma_{\mathcal{A}_2}(\mathcal{G})$. We now provide a necessary and sufficient condition under which the indiscernible partitions associated to two different attribute sets are same.

Proposition 1.0.4. *For an attribute subset $\mathcal{A} \subseteq \mathbb{A}$, define $\mathcal{B}_{\mathcal{A}}(\mathcal{G}) = (\mathcal{O}(\mathcal{A}))^c$. Let $\mathcal{A}_1, \mathcal{A}_2$ be two attribute subsets, then $\gamma_{\mathcal{A}_1}(\mathcal{G}) = \gamma_{\mathcal{A}_2}(\mathcal{G})$ if and only if one of the following hold;*

- (1) $\mathcal{B}_{\mathcal{A}_1}(\mathcal{G}) = \mathcal{B}_{\mathcal{A}_2}(\mathcal{G})$
- (2) $\mathcal{B}_{\mathcal{A}_1}(\mathcal{G}), \mathcal{B}_{\mathcal{A}_2}(\mathcal{G})$ are either empty or consists of only one orbit.

Proof. (1) Suppose that $\mathcal{B}_{\mathcal{A}_1}(\mathcal{G}) = \mathcal{B}_{\mathcal{A}_2}(\mathcal{G})$ then $\mathcal{O}(\mathcal{A}_1) = \mathcal{O}(\mathcal{A}_2)$. Hence by Proposition 1.0.2, $\gamma_{\mathcal{A}_1}(\mathcal{G}) = \gamma_{\mathcal{A}_2}(\mathcal{G})$.

(2) Suppose that $\mathcal{B}_{\mathcal{A}_1}(\mathcal{G}) = \emptyset$ and $\mathcal{B}_{\mathcal{A}_2}(\mathcal{G})$ consists of only one orbit. Then the orbit of $\mathcal{B}_{\mathcal{A}_2}(\mathcal{G})$ must coincide with at least one orbit of $\gamma_{\mathcal{A}_1}(\mathcal{G})$. Hence by Proposition 1.0.2, $\gamma_{\mathcal{A}_1}(\mathcal{G}) = \gamma_{\mathcal{A}_2}(\mathcal{G})$. Similarly, if $\mathcal{B}_{\mathcal{A}_2}(\mathcal{G}) = \emptyset$ and $\mathcal{B}_{\mathcal{A}_1}(\mathcal{G})$ consists of only one orbit, then $\gamma_{\mathcal{A}_1}(\mathcal{G}) = \gamma_{\mathcal{A}_2}(\mathcal{G})$.

Now, suppose contrary that $\mathcal{B}_{\mathcal{A}_1}(\mathcal{G}) \neq \mathcal{B}_{\mathcal{A}_2}(\mathcal{G})$ and neither $\mathcal{B}_{\mathcal{A}_1}(\mathcal{G})$ nor $\mathcal{B}_{\mathcal{A}_2}(\mathcal{G})$ is empty or consists of only one orbit. By proposition 1.0.2, this implies that $\gamma_{\mathcal{A}_1}(\mathcal{G}) \neq \gamma_{\mathcal{A}_2}(\mathcal{G})$ but $\gamma_{\mathcal{A}_1}(\mathcal{G}) = \gamma_{\mathcal{A}_2}(\mathcal{G})$, a contradiction. \square

Suppose that a graph \mathcal{G} have n orbits then following propositions are obvious.

Proposition 1.0.5. *If a graph \mathcal{G} has n orbits and attribute subsets \mathcal{A}_1 and \mathcal{A}_2 contain elements of n or $n - 1$ orbits then $\gamma_{\mathcal{A}_1}(\mathcal{G}) = \gamma_{\mathcal{A}_2}(\mathcal{G})$.*

Proof. Since graph \mathcal{G} has n orbits. If $\mathcal{O}(\mathcal{A}_1) = \mathcal{O}(\mathcal{A}_2)$, then $\mathcal{B}_{\mathcal{A}_1}(\mathcal{G}) = \mathcal{B}_{\mathcal{A}_2}(\mathcal{G})$ hence by proposition 1.0.4, $\gamma_{\mathcal{A}_1}(\mathcal{G}) = \gamma_{\mathcal{A}_2}(\mathcal{G})$. Now suppose that \mathcal{A}_1 contains elements of n orbits and \mathcal{A}_2 contains elements of $n - 1$ orbits, then by second part of proposition 1.0.4, $\gamma_{\mathcal{A}_1}(\mathcal{G}) = \gamma_{\mathcal{A}_2}(\mathcal{G})$. \square

Proposition 1.0.6. *If \mathcal{A}_1 and \mathcal{A}_2 are two subsets of attributes with elements from at most $n - 2$ orbits then $\gamma_{\mathcal{A}_1}(\mathcal{G}) = \gamma_{\mathcal{A}_2}(\mathcal{G})$ if and only if $\mathcal{O}(\mathcal{A}_1) = \mathcal{O}(\mathcal{A}_2)$.*

Proof. As $\mathcal{O}(\mathcal{A}_1) = \mathcal{O}(\mathcal{A}_2)$, by proposition 1.0.4, $\gamma_{\mathcal{A}_1}(\mathcal{G}) = \gamma_{\mathcal{A}_2}(\mathcal{G})$. Suppose contrary that $\mathcal{O}(\mathcal{A}_1) \neq \mathcal{O}(\mathcal{A}_2)$ but $\gamma_{\mathcal{A}_1}(\mathcal{G}) = \gamma_{\mathcal{A}_2}(\mathcal{G})$. Since \mathcal{A}_1 and \mathcal{A}_2 contain elements of at most $n - 2$ orbits. Hence their complement will consist of the elements of at least two different orbits which implies that $\mathcal{B}_{\mathcal{A}_1}(\mathcal{G}) \neq \mathcal{B}_{\mathcal{A}_2}(\mathcal{G})$, yielding $\gamma_{\mathcal{A}_1}(\mathcal{G}) \neq \gamma_{\mathcal{A}_2}(\mathcal{G})$, a contradiction. \square

In the next proposition, we show that the automorphisms of \mathcal{G} preserve the structure of blocks in indiscernibility partition.

Proposition 1.0.7. *For a graph \mathcal{G} , let $\mathcal{A} \subseteq \mathcal{V}(\mathcal{G})$ be any attribute subset. If $\eta \in \Gamma(\mathcal{G})$ then $\gamma_{\mathcal{A}}(\mathcal{G}) = \gamma_{\eta(\mathcal{A})}(\mathcal{G})$.*

Proof. Suppose $\gamma_{\mathcal{A}}(\mathcal{G}) = \mathcal{B}_1|\mathcal{B}_2|\dots|\mathcal{B}_m$ and $\gamma_{\eta(\mathcal{A})}(\mathcal{G}) = \mathcal{K}_1|\mathcal{K}_2|\dots|\mathcal{K}_n$. Let $\mathcal{B}_i = \mathcal{O}_{\mathcal{A}}(x_i) = \{x_j : \mathcal{O}(x_j) \cap \mathcal{A} = \mathcal{O}(x_i) \cap \mathcal{A}\}$ and $\mathcal{K}_i = \mathcal{O}_{\eta(\mathcal{A})}(x_i) = \{x_j : \mathcal{O}(x_j) \cap \eta(\mathcal{A}) = \mathcal{O}(x_i) \cap \eta(\mathcal{A})\}$.

It is easy to see that for $\eta \in \Gamma(\mathcal{G})$, $\mathcal{O}(x_i) \cap \mathcal{A} \neq \emptyset$ if and only if $\mathcal{O}(x_i) \cap \eta(\mathcal{A}) \neq \emptyset$ for $1 \leq i \leq k$ because $x_k \in \mathcal{O}(x_i) \cap \mathcal{A}$ if and only if $\eta(x_k) \in \mathcal{O}(x_i) \cap \eta(\mathcal{A})$. By 1.0.2, $\mathcal{B}_i = \mathcal{K}_i = \mathcal{O}(x_i)$ or $\mathcal{B}_i = \mathcal{K}_i = (\mathcal{O}(\mathcal{A}))^c$ which implies that $\gamma_{\mathcal{A}}(\mathcal{G}) = \gamma_{\eta(\mathcal{A})}(\mathcal{G})$. \square

For a set of attributes $\mathcal{A} \subseteq \mathcal{V}$, $IND_{\mathcal{A}}(\mathcal{V}) = \{(x, y) \in \mathcal{V}^2 \mid \mathcal{O}(x) \cap \mathcal{A} = \mathcal{O}(y) \cap \mathcal{A}\}$. An attribute $x \in \mathcal{A}$ is called dispensable in \mathcal{A} if $IND_{\mathcal{A}}(\mathcal{V}) = IND_{\mathcal{A} \setminus \{x\}}(\mathcal{V})$, otherwise x is called indispensable with respect to \mathcal{A} . A minimal subset \mathcal{R} of an attribute set \mathcal{A} that yields the same partition as provided by the set of all attributes is called *reduct*.

Proposition 1.0.8. *For a graph \mathcal{G} with a reduct \mathcal{R} and orbits $\mathcal{O}_1, \mathcal{O}_2, \dots, \mathcal{O}_k$, $|\mathcal{R} \cap \mathcal{O}_i| = 1$ for all $i = 1, 2, \dots, k$ except one i .*

Proof. Suppose there exists two orbits for which $\mathcal{R} \cap \mathcal{O}_i$ and $\mathcal{R} \cap \mathcal{O}_j$ is empty then $\mathcal{O}_i \cup \mathcal{O}_j$ forms one class. Hence, \mathcal{R} must have empty intersection with at most one orbit. Now, suppose that there exists an orbit \mathcal{O}_k such that $|\mathcal{R} \cap \mathcal{O}_k| > 1$. Let $x_1, x_2 \in \mathcal{R} \cap \mathcal{O}_k$ then $\gamma_{\mathcal{R}}(\mathcal{I}) = \gamma_{\mathcal{R} \setminus \{x_2\}}(\mathcal{I})$. This implies \mathcal{R} is not minimal (a contradiction). Hence, $|\mathcal{R} \cap \mathcal{O}_i| = 1$ for all $i = 1, 2, \dots, k$ except one i . \square

It is obvious that any attribute subset \mathcal{A} may have more than one reducts, the set of all reducts of an attribute subset is denoted by $\mathcal{RED}_{\mathcal{A}}(\mathcal{G})$. Let $\mathcal{K} \in \mathcal{RED}_{\mathcal{A}}(\mathcal{G})$ then $\gamma_{\mathcal{A}}(\mathcal{G}) = \gamma_{\mathcal{K}}(\mathcal{G}) = \mathcal{B}_1 | \mathcal{B}_2 | \dots | \mathcal{B}_n$. By proposition 1.0.7, $\gamma_{\eta(\mathcal{A})}(\mathcal{G}) = \gamma_{\eta(\mathcal{K})}(\mathcal{G}) = \mathcal{B}_1 | \mathcal{B}_2 | \dots | \mathcal{B}_n$. We note that the automorphism of \mathcal{G} preserves the structure of reducts. Hence we have the following proposition:

Proposition 1.0.9. *For a graph \mathcal{G} , let $\mathcal{A} \subseteq \mathcal{V}$. If $\eta \in \Gamma(\mathcal{G})$, then $\mathcal{K} \in \mathcal{RED}_{\mathcal{A}}(\mathcal{G})$ if and only if $\eta(\mathcal{K}) \in \mathcal{RED}_{\eta(\mathcal{A})}(\mathcal{G})$.*

According to Proposition 1.0.9, for a graph \mathcal{G} , attribute set \mathcal{A} and $\eta \in \Gamma(\mathcal{G})$, $\mathcal{K} \in \mathcal{RED}_{\mathcal{A}}(\mathcal{G})$ implies that $\eta(\mathcal{K}) \in \mathcal{RED}_{\eta(\mathcal{A})}(\mathcal{G})$. If \mathcal{K} is a reduct for attribute set \mathcal{A} then $\eta(\mathcal{K})$ does not necessarily belong to $\mathcal{RED}_{\mathcal{A}}(\mathcal{G})$.




FIGURE 2. A Graph \mathcal{G}

Consider a graph \mathcal{G} in FIGURE 2. Note that $\mathcal{O}(1) = \mathcal{O}(4) = \{1, 4\}$, $\mathcal{O}(2) = \mathcal{O}(3) = \{2, 3\}$ and $\mathcal{O}(5) = \mathcal{O}(6) = \mathcal{O}(7) = \mathcal{O}(8) = \{5, 6, 7, 8\}$. Let $\mathcal{A} = \{1, 2, 5, 6\}$ and take $\eta = (14)(23)(57)(68)$, then $\mathcal{K} = \{1, 2\}$ is a reduct of \mathcal{A} and $\eta(\mathcal{K}) = \{3, 4\}$ is a reduct of $\eta(\mathcal{A})$ but not that of \mathcal{A} .

Let $\mathcal{P}(\mathcal{V})$ denotes the set of all partitions of \mathcal{V} . For two partitions γ_1 and γ_2 of \mathcal{V} , γ_1 is called finer than γ_2 if all blocks of γ_1 are subset of the blocks of γ_2 , denoted by $\gamma_1 \preceq \gamma_2$. It is straightforward to see that for two attribute sets, \mathcal{A}_1 and \mathcal{A}_2 of graph \mathcal{G} with $\mathcal{A}_1 \subseteq \mathcal{A}_2$, then $\gamma_{\mathcal{A}_2}(\mathcal{G}) \preceq \gamma_{\mathcal{A}_1}(\mathcal{G})$. For a graph \mathcal{G} with orbits $\mathcal{O}_1, \mathcal{O}_2, \dots, \mathcal{O}_k$ and attribute sets $\mathcal{A}_1 \subseteq \mathcal{A}_2$, $\gamma_{\mathcal{A}_2}(\mathcal{G}) = \gamma_{\mathcal{A}_1}(\mathcal{G})$, if \mathcal{A}_1 and \mathcal{A}_2 have non-empty intersection with same orbits \mathcal{O}_i for $1 \leq i \leq k$. If $\mathcal{A}_1 \subseteq \mathcal{A}_2$ and there exists some \mathcal{O}_i ($1 \leq i \leq k$) such that $\mathcal{A}_1 \cap \mathcal{O}_i = \emptyset$ and $\mathcal{A}_2 \cap \mathcal{O}_i \neq \emptyset$ then $\gamma_{\mathcal{A}_2}(\mathcal{G}) \prec \gamma_{\mathcal{A}_1}(\mathcal{G})$. From this discussion, we conclude that a partition consisting of orbits of the graphs is the finest partition of the vertex set in $\mathcal{P}(\mathcal{V})$.

Now, we introduce three sets, namely \mathcal{A} -interior set, \mathcal{A} -exterior set and \mathcal{A} -delimiting set as follows:

Definition 1.0.10. For an attribute subset $\mathcal{A} \subseteq \mathbb{A}$, let $x \in \mathcal{V}$. Then

- x is called \mathcal{A} -interior, if $\mathcal{O}(x) \subseteq \mathcal{A}$.
- x is called \mathcal{A} -exterior, if $\mathcal{O}(x) \subseteq \mathcal{V} \setminus \mathcal{A}$.
- x is called \mathcal{A} -delimiting, if $\mathcal{O}_{\mathcal{A}}(x) \neq \emptyset$ and $\mathcal{O}_{\mathcal{V} \setminus \mathcal{A}}(x) \neq \emptyset$.

Represent by $Int(\mathcal{A})$, $Ext(\mathcal{A})$ and $Del(\mathcal{A})$ respectively, the subset of all \mathcal{A} -interior, \mathcal{A} -exterior and \mathcal{A} -delimiting vertices of graph \mathcal{G} .

Note that the family of all subsets of definition 1.0.10 is a set partition of $\mathcal{V}(\mathcal{G})$ and for an attribute subset $\mathcal{A} \subseteq \mathcal{V}(\mathcal{G})$, and $\mathcal{A}^c = \mathcal{V} \setminus \mathcal{A}$ then $Int(\mathcal{A}) = Ext(\mathcal{A}^c)$ and $Del(\mathcal{A}) = Del(\mathcal{A}^c)$. For an attribute subset $\mathcal{A} \subseteq \mathcal{V}(K_n)$, following is straightforward.

- (i): If $|\mathcal{A}| < n$ then $Int(\mathcal{A}) = Ext(\mathcal{A}) = \emptyset$ and $Del(\mathcal{A}) = \mathcal{V}(K_n)$.
- (ii): If $|\mathcal{A}| = n$ then $Del(\mathcal{A}) = Ext(\mathcal{A}) = \emptyset$ and $Int(\mathcal{A}) = \mathcal{V}(K_n)$.
- (iii): If $\mathcal{A} = \emptyset$ then $Int(\mathcal{A}) = Del(\mathcal{A}) = \emptyset$ and $Ext(\mathcal{A}) = \mathcal{V}(K_n)$.

Proposition 1.0.11. For an attribute subset $\mathcal{A} \subseteq \mathbb{A}$, a non-empty $Ext(\mathcal{A})$ is a block of \mathcal{A} -indiscernible partition $\gamma_{\mathcal{A}}(\mathcal{G})$.

Proof. Let $x, x' \in Ext(\mathcal{A})$ then for all $y \in \mathcal{A}$, we have $\mathcal{F}(x, y) = \mathcal{F}(x', y) = 0$, so $x \equiv_{\mathcal{A}} x'$. If $z \in \mathcal{V}(\mathcal{G})$ such that for some $z' \in Ext(\mathcal{A})$, $z \equiv_{\mathcal{A}} z'$ then $\mathcal{O}_{\mathcal{A}}(z) = \emptyset$ implying that $z \in Ext(\mathcal{A})$. Because if there exists a vertex $y \in \mathcal{O}_{\mathcal{A}}(z)$ then $1 = \mathcal{F}(y, z) \neq \mathcal{F}(y, z') = 0$, which is contradiction to the assumption that $z \equiv_{\mathcal{A}} z'$. Therefore, $\mathcal{O}_{\mathcal{A}}(z) = \emptyset$ and $z \in Ext(\mathcal{A})$. \square

Remark 1.0.12. It is straightforward to observe that the $Int(\mathcal{A})$ and $Del(\mathcal{A})$ need not be a block of $\gamma_{\mathcal{A}}(\mathcal{G})$. Further, $Int(\mathcal{A})$ and $Del(\mathcal{A})$ are blocks of $\gamma_{\mathcal{A}}(\mathcal{G})$ if and only if it consists of only one orbit.

Proposition 1.0.13. For an attribute subset \mathcal{A} , if $x, y \in Int(\mathcal{A}) \cup Del(\mathcal{A})$, then $x \equiv_{\mathcal{A}} y$ if and only if $\mathcal{O}(x) = \mathcal{O}(y)$.

Proof. Suppose that $x, y \in Int(\mathcal{A}) \cup Del(\mathcal{A})$ and $\mathcal{O}(x) = \mathcal{O}(y)$, then by definition of indiscernibility, $x \equiv_{\mathcal{A}} y$. Conversely, suppose that $x \equiv_{\mathcal{A}} y$ and $\mathcal{O}(x) \neq \mathcal{O}(y)$. Then there exists $z \in \mathcal{A}$ which will belong to at most one of $\mathcal{O}(x)$ or $\mathcal{O}(y)$ because if no such $z \in \mathcal{A}$ belongs to $\mathcal{O}(x)$ or $\mathcal{O}(y)$ then $x, y \in Ext(\mathcal{A})$. Therefore, $\neg(x \equiv_{\mathcal{A}} y)$, contradiction. Hence $\mathcal{O}(x) = \mathcal{O}(y)$. \square

2. UPPER AND LOWER APPROXIMATIONS IN GRAPHS

In this section, lower and upper approximations of various subsets of vertices are studied using similarity relations.

Theorem 1.0.1, provides information about the behavior of objects in the indiscernibility relation $\equiv_{\mathcal{A}}$. Indiscernibility relations can be considered as a type of symmetry relations as for the attribute subset \mathcal{A} . Using the fact that for any two distinct vertices $x, y \in \mathcal{V}(\mathcal{G})$ either $\mathcal{O}(x) = \mathcal{O}(y)$ or $\mathcal{O}(x) \cap \mathcal{O}(y) = \emptyset$, it is straightforward to note that every proper subset X of $\mathcal{V}(\mathcal{G})$ is rough if $\mathcal{O}(x) = \mathcal{V}(\mathcal{G})$. Also note that for $x, y \in \mathcal{V}(\mathcal{G})$ with $\mathcal{O}(x) \neq \mathcal{O}(y)$ and $\mathcal{O}(x) \cup \mathcal{O}(y) = \mathcal{V}(\mathcal{G})$, a subset X of $\mathcal{V}(\mathcal{G})$ is rough if and only if $X \neq \mathcal{O}(x)$, $X \neq \mathcal{O}(y)$ and $X \neq \mathcal{V}(\mathcal{G})$.

Note that the indiscernibility partition of complete graph on n vertices consists of only one orbit yielding one block. If $\mathcal{V}(K_n) = \{x_1, \dots, x_n\}$, then $\gamma_{\mathcal{A}}(K_n) =$

x_1, \dots, x_n for any attribute subset $\mathcal{A} \subseteq K_n$ and every proper subset \mathcal{Q} of $\mathcal{V}(K_n)$ is \mathcal{A} -rough because $\mathbb{L}_{\mathcal{A}}(\mathcal{Q}) = \emptyset$ and $\mathbb{U}_{\mathcal{A}}(\mathcal{Q}) = \mathcal{V}(K_n)$. A subset \mathcal{Q} of $\mathcal{V}(K_n)$ is \mathcal{A} -exact if and only if $\mathcal{Q} = \mathcal{V}(K_n)$.

A given graph \mathcal{G} in which $\mathcal{V} = \mathcal{V}_1 \cup \mathcal{V}_2$, $\mathcal{V}_1 \cap \mathcal{V}_2 = \emptyset$, $x \sim y$ for each $x \in \mathcal{V}_1, y \in \mathcal{V}_2$ and $x \approx y$ if and only if $x, y \in \mathcal{V}_1$ or $x, y \in \mathcal{V}_2$, is called a *complete bipartite graph*. For a complete bipartite graph $\mathcal{K}_{m,n} = (\mathcal{V}_1 | \mathcal{V}_2)$ where $\mathcal{V}_1 = \{x_1, x_2, \dots, x_m\}$ and $\mathcal{V}_2 = \{y_1, y_2, \dots, y_n\}$, $\gamma_{\mathcal{A}}(\mathcal{K}_{m,n}) = x_1, x_2, \dots, x_m | y_1, y_2, \dots, y_n$, for any non-empty attribute subset \mathcal{A} .

In the following proposition, we study rough sets in complete bipartite graphs.

Proposition 2.0.1. *If \mathcal{G} is a complete bipartite graph then a subset \mathcal{Q} of \mathcal{V} is rough with respect to attributes \mathcal{A} if and only if $\mathcal{Q} \neq \mathcal{V}_1$, $\mathcal{Q} \neq \mathcal{V}_2$ and $\mathcal{Q} \neq \mathcal{V}$, where $\mathcal{V}_1, \mathcal{V}_2$ are partites of \mathcal{G} .*

Proof. Let \mathcal{Q} be a rough set in \mathcal{G} . If $\mathcal{Q} = \mathcal{V}$ then clearly \mathcal{Q} is exact, hence $\mathcal{Q} \neq \mathcal{V}$. Now if $\mathcal{Q} = \mathcal{V}_1$ then for each $x \in \mathcal{V}_2$, $\mathcal{O}(x) \cap \mathcal{Q} = \emptyset$ which gives that \mathcal{Q} is exact. Similar arguments holds for $\mathcal{Q} = \mathcal{V}_2$. Hence, $\mathcal{Q} \neq \mathcal{V}_1$, $\mathcal{Q} \neq \mathcal{V}_2$ and $\mathcal{Q} \neq \mathcal{V}$.

Conversely, suppose $\mathcal{Q} \neq \mathcal{V}_1$, $\mathcal{Q} \neq \mathcal{V}_2$ and $\mathcal{Q} \neq \mathcal{V}$, we need to prove that $\mathbb{L}_{\mathcal{A}}(\mathcal{Q}) \neq \mathbb{U}_{\mathcal{A}}(\mathcal{Q})$. Since $\mathcal{Q} \neq \mathcal{V}_1$ so we have three cases:

(1) Suppose $\mathcal{V}_1 \cap \mathcal{Q} = \emptyset$ and $\mathcal{V}_2 \cap \mathcal{Q} \neq \emptyset$ then $\mathcal{Q} \neq \mathcal{V}_2$ which implies that $\mathcal{Q} \subset \mathcal{V}_2$. Now for any $x \in \mathcal{V}$, if $x \in \mathcal{V}_1$ then $\mathcal{O}(x) \cap \mathcal{Q} = \emptyset$ because $\mathcal{O}(x) = \mathcal{V}_1$. Moreover, for $x \in \mathcal{V}_2$, $\mathcal{O}(x) = \mathcal{V}_2$ which implies that $\mathbb{L}_{\mathcal{A}}(\mathcal{Q}) = \emptyset$ and $\mathbb{U}_{\mathcal{A}}(\mathcal{Q}) = \mathcal{V}_2$. Hence, \mathcal{Q} is a rough set.

(2) For the case $\mathcal{V}_1 \cap \mathcal{Q} \neq \emptyset$ and $\mathcal{V}_2 \cap \mathcal{Q} = \emptyset$ then $\mathcal{Q} \neq \mathcal{V}_1$ which implies that $\mathcal{Q} \subset \mathcal{V}_1$. Now for any $x \in \mathcal{V}$, if $x \in \mathcal{V}_1$ then $\mathcal{O}(x) \cap \mathcal{Q} \neq \emptyset$ because $\mathcal{O}(x) = \mathcal{V}_1$. Moreover, for $x \in \mathcal{V}_2$, $\mathcal{O}(x) = \mathcal{V}_2$ which implies that $\mathbb{L}_{\mathcal{A}}(\mathcal{Q}) = \emptyset$ and $\mathbb{U}_{\mathcal{A}}(\mathcal{Q}) = \mathcal{V}_1$. Hence, \mathcal{Q} is a rough set.

(3) Now suppose $\mathcal{V}_1 \cap \mathcal{Q} \neq \emptyset$ and $\mathcal{V}_2 \cap \mathcal{Q} \neq \emptyset$ then we have three sub-cases:

i) Suppose $\mathcal{V}_1 \cap \mathcal{Q} = \mathcal{V}_1$ then $\mathbb{L}_{\mathcal{A}}(\mathcal{Q}) = \mathcal{V}_1$ and $\mathbb{U}_{\mathcal{A}}(\mathcal{Q}) = \mathcal{V}$ which gives that \mathcal{Q} is rough.

ii) Suppose $\mathcal{V}_2 \cap \mathcal{Q} = \mathcal{V}_2$ then for every $x \in \mathcal{V}_2$ we have $x \in \mathbb{L}_{\mathcal{A}}(\mathcal{Q}) = \mathcal{V}_2$ and $\mathbb{U}_{\mathcal{A}}(\mathcal{Q}) = \mathcal{V}$ which gives that \mathcal{Q} is rough.

iii) Suppose $\mathcal{V}_1 \cap \mathcal{Q} \neq \mathcal{V}_1$ and $\mathcal{V}_2 \cap \mathcal{Q} \neq \mathcal{V}_2$. Let $x \in \mathcal{V}$ such that $x \in \mathcal{V}_1$ then $x \in \mathbb{U}_{\mathcal{A}}(\mathcal{Q})$ because $\mathcal{O}(x) = \mathcal{V}_1$ and $\mathcal{O}(x) \cap \mathcal{Q} \neq \emptyset$ but $\mathcal{O}(x) \not\subseteq \mathcal{Q}$. Similarly, if $x \in \mathcal{V}_2$ then $x \in \mathbb{U}_{\mathcal{A}}(\mathcal{Q})$. Also, for every $x \in \mathcal{V}$ we have $\mathcal{O}(x) \not\subseteq \mathcal{Q}$ which gives that $\mathbb{L}_{\mathcal{A}}(\mathcal{Q}) = \emptyset$ so \mathcal{Q} is a rough set in \mathcal{G} . \square

In the following proposition, we discuss lower and upper approximation of a subset \mathcal{Q} of complete bipartite graph $\mathcal{K}_{m,n}$.

Proposition 2.0.2. *Let $\mathcal{K}_{m,n} = (\mathcal{V}_1 | \mathcal{V}_2)$ is a complete bipartite graph where $\mathcal{V}_1 = \{x_1, x_2, \dots, x_m\}$ and $\mathcal{V}_2 = \{y_1, y_2, \dots, y_n\}$. Let \mathcal{Q} and \mathcal{A} are two subsets of $\mathcal{V}(\mathcal{K}_{m,n})$ such that $\mathcal{Q} \neq \mathcal{V}$. Then*

$$\begin{aligned} (1) \quad \mathbb{L}_{\mathcal{A}}(\mathcal{Q}) &= \begin{cases} \mathcal{V}_1 & \text{if } \mathcal{V}_1 \subseteq \mathcal{Q} \\ \mathcal{V}_2 & \text{if } \mathcal{V}_2 \subseteq \mathcal{Q} \\ \emptyset & \text{otherwise.} \end{cases} \\ (2) \quad \mathbb{U}_{\mathcal{A}}(\mathcal{Q}) &= \begin{cases} \mathcal{V}_1 & \text{if } \mathcal{Q} \subseteq \mathcal{V}_1 \\ \mathcal{V}_2 & \text{if } \mathcal{Q} \subseteq \mathcal{V}_2 \\ \mathcal{V} & \text{otherwise.} \end{cases} \\ (3) \quad \mathcal{Q} \text{ is } \mathcal{A}\text{-exact} &\text{ if and only if } \mathcal{Q} = \mathcal{V}_1 \text{ or } \mathcal{V}_2. \end{aligned}$$

Proof.

- (1) Let $\mathcal{V}_1 \subseteq \mathcal{Q}$. If $x \in \mathcal{V}_1$, then it follow by Theorem 1.0.1 that $\mathcal{O}_{\mathcal{A}}(x) \subseteq \mathcal{Q}$, thus by definition of the lower approximation, we get $\mathcal{V}_1 \subseteq \mathbb{L}_{\mathcal{A}}(\mathcal{Q})$. Moreover, if $x \in \mathcal{V}_2 \cap \mathbb{L}_{\mathcal{A}}(\mathcal{Q})$, for some $x \in \mathcal{V}$, then, again by Theorem 1.0.1 and definition of the lower approximation, we get $\mathcal{V}_2 = \mathcal{O}_{\mathcal{A}}(x) \subseteq \mathcal{Q}$. Because $\mathcal{V}_1 | \mathcal{V}_2$ are set partition for \mathcal{V} , so last inclusion implies $\mathcal{Q} = \mathcal{V}$, a contradiction to our assumption. Thus, $\mathcal{V}_1 \subseteq \mathbb{L}_{\mathcal{A}}(\mathcal{Q})$ and $\mathcal{V}_2 \cap \mathbb{L}_{\mathcal{A}}(\mathcal{Q}) = \emptyset$. Similarly, if $\mathcal{V}_2 \subseteq \mathcal{Q}$ then $\mathbb{L}_{\mathcal{A}}(\mathcal{Q}) = \mathcal{V}_2$. Hence let $\mathcal{V}_1 \not\subseteq \mathcal{Q}$ and $\mathcal{V}_2 \not\subseteq \mathcal{Q}$, but each vertex $x \in \mathcal{V}$ is either in \mathcal{V}_1 or \mathcal{V}_2 . Therefore, by Theorem 1.0.1, we have $\mathcal{O}_{\mathcal{A}}(x) = \mathcal{V}_1 \not\subseteq \mathcal{Q}$ and $\mathcal{O}_{\mathcal{A}}(x) = \mathcal{V}_2 \not\subseteq \mathcal{Q}$, i.e., $x \notin \mathbb{L}_{\mathcal{A}}(\mathcal{Q})$. Hence $\mathbb{L}_{\mathcal{A}}(\mathcal{Q}) = \emptyset$.
- (2) Let $\mathcal{Q} \subseteq \mathcal{V}_1$. If $x \in \mathcal{V}_1$, then it follow by Theorem 1.0.1 that $\mathcal{O}_{\mathcal{A}}(x) = \mathcal{V}_1 \cap \mathcal{Q} \neq \emptyset$, because \mathcal{Q} is non-empty subset of \mathcal{Q}_1 . Hence $x \in \mathbb{U}_{\mathcal{A}}(\mathcal{Q})$. Moreover, if $x \in \mathbb{U}_{\mathcal{A}}(\mathcal{Q})$ then by definition of upper approximation we get $\mathcal{O}_{\mathcal{A}}(x) \cap \mathcal{Q} \neq \emptyset$. Let $y \in \mathcal{O}_{\mathcal{A}}(x) \cap \mathcal{Q}$. Since $y \in \mathcal{Q} \subseteq \mathcal{V}_1$, thus by Theorem 1.0.1 we have $\mathcal{V}_1 = \mathcal{O}_{\mathcal{A}}(x) = \mathcal{O}_{\mathcal{A}}(y)$, therefore again by Theorem 1.0.1 we deduce that $x \in \mathcal{V}_1$. Hence $\mathbb{U}_{\mathcal{A}}(\mathcal{Q}) = \mathcal{V}_1$. The case of $\mathcal{Q} \subseteq \mathcal{V}_2$ is similar. Finally, $\mathcal{Q} \not\subseteq \mathcal{V}_1$ and $\mathcal{Q} \not\subseteq \mathcal{V}_2$. Since $\mathcal{V}_1 | \mathcal{V}_2$ is a partition of \mathcal{V} , which implies that $\mathcal{V}_1 \cap \mathcal{Q} \neq \emptyset$ and $\mathcal{V}_2 \cap \mathcal{Q} \neq \emptyset$. Now select arbitrary a vertex $x \in \mathcal{V}$, then either $x \in \mathcal{V}_1$ or \mathcal{V}_2 . If $x \in \mathcal{V}_1$, then by Theorem 1.0.1 it follows that $\mathcal{O}_{\mathcal{A}}(x) \cap \mathcal{Q} = \mathcal{V}_1 \cap \mathcal{Q} \neq \emptyset$. Thus, $x \in \mathbb{U}_{\mathcal{A}}(\mathcal{Q})$. Analogously, if $x \in \mathcal{V}_2$. Then it shows that $\mathcal{V} \subseteq \mathbb{U}_{\mathcal{A}}(\mathcal{Q})$. i.e., $\mathcal{V} = \mathbb{U}_{\mathcal{A}}(\mathcal{Q})$.
- (3) This follow from the definition of exactness and Theorem 1.0.1.

□

In proposition 2.0.3, we provide complete description for the lower approximations of graph \mathcal{G} .

Proposition 2.0.3. *For a graph \mathcal{G} , let $\mathcal{A}, \mathcal{Q} \subseteq \mathcal{V}$. Then*

$$\mathbb{L}_{\mathcal{A}}(\mathcal{Q}) = \begin{cases} \mathcal{O}(\mathcal{A}) & \text{where } x \in \mathcal{A} \wedge (\mathcal{O}(x) \setminus \mathcal{Q}) = \emptyset \\ \emptyset & \text{otherwise} \end{cases}$$

Proof. Let a vertex $x \in \mathcal{A}$ such that $\mathcal{O}_{\mathcal{A}}(x) \setminus \mathcal{Q} = \emptyset$, which give $\mathcal{O}_{\mathcal{A}}(x) \subseteq \mathcal{Q}$, implies that $\mathcal{O}_{\mathcal{A}}(x) \in \mathbb{L}_{\mathcal{A}}(\mathcal{Q})$. Since x is chosen arbitrary, hence $\mathbb{L}_{\mathcal{A}}(\mathcal{Q}) = \mathcal{O}(\mathcal{A})$. For $\mathcal{O}_{\mathcal{A}}(x) \setminus \mathcal{Q} \neq \emptyset$, $\mathcal{O}_{\mathcal{A}}(x) \not\subseteq \mathcal{Q}$ giving that $\mathcal{O}_{\mathcal{A}}(x) \notin \mathbb{L}_{\mathcal{A}}(\mathcal{Q})$ implies that $\mathbb{L}_{\mathcal{A}}(\mathcal{Q}) = \emptyset$. □

In proposition 2.0.4, we provide a necessary and sufficient condition for a subset $\mathcal{Q} \subseteq \mathcal{V}$ to be exact.

Proposition 2.0.4. *For a graph \mathcal{G} , let $\mathcal{A}, \mathcal{Q} \subseteq \mathcal{V}$. Then \mathcal{Q} is exact with respect to \mathcal{A} , if and only if $\mathcal{Q} = \cup_{x \in \mathcal{Q}} \mathcal{O}_{\mathcal{A}}(x)$.*

Proof. Suppose \mathcal{Q} is exact then for $x \in \mathcal{Q}$, $\mathcal{O}_{\mathcal{A}}(x) \subseteq \mathcal{Q}$. As x is arbitrary, $\cup_{x \in \mathcal{Q}} \mathcal{O}_{\mathcal{A}}(x) \subseteq \mathcal{Q}$. For each $x \in \mathcal{Q}$, $x \in \mathcal{O}_{\mathcal{A}}(x)$ which yields that $\mathcal{Q} \subseteq \cup_{x \in \mathcal{Q}} \mathcal{O}_{\mathcal{A}}(x)$. Hence $\mathcal{Q} = \cup_{x \in \mathcal{Q}} \mathcal{O}_{\mathcal{A}}(x)$. Conversely, suppose $\mathcal{Q} = \cup_{x \in \mathcal{Q}} \mathcal{O}_{\mathcal{A}}(x)$ which clearly implies $\mathbb{L}_{\mathcal{A}}(\mathcal{Q}) = \mathbb{U}_{\mathcal{A}}(\mathcal{Q})$ giving that \mathcal{Q} is exact. □

A graph \mathcal{G} in which no two vertices are similar is called *rigid* graph. In such graphs, $\gamma_{\mathcal{A}}(\mathcal{G})$ consists of blocks having singleton elements only, which implies $\mathbb{L}_{\mathcal{A}}(\mathcal{Q}) = \mathbb{U}_{\mathcal{A}}(\mathcal{Q})$. Hence every subset of the vertex set of a rigid graph is exact.

2.1. Rough Membership Function and Dependency. Now, we introduce and present results on rough membership function, positive region and degree of dependency of graphs using orbits of the graphs.

Let $\mathcal{G} = (\mathcal{V}, \mathcal{E})$ and $\mathcal{A}, \mathcal{Q} \subseteq \mathcal{V}$. The rough membership function on the vertex set \mathcal{V} is defined by:

$$\mu_{\mathcal{Q}}^{\mathcal{A}}(x) = \frac{|\{y \in \mathcal{Q} : \mathcal{O}(x) \cap \mathcal{A} = \mathcal{O}(y) \cap \mathcal{A}\}|}{|\{y \in \mathcal{V} : \mathcal{O}(x) \cap \mathcal{A} = \mathcal{O}(y) \cap \mathcal{A}\}|}.$$

For $\mathcal{A}, \mathcal{Q} \subseteq \mathcal{V}$, $POS_{\mathcal{A}}(\mathcal{Q}) = \{x \in \mathcal{V} : (y \in \mathcal{V} \wedge \mathcal{O}(x) \cap \mathcal{A} = \mathcal{O}(y) \cap \mathcal{A}) \Rightarrow (\mathcal{O}(x) \cap \mathcal{Q} = \mathcal{O}(y) \cap \mathcal{Q})\}$. The number $deg_{\mathcal{A}}(\mathcal{Q}) = \frac{|POS_{\mathcal{A}}(\mathcal{Q})|}{|\mathcal{V}|}$ is referred to as \mathcal{A} -degree dependency of \mathcal{Q} . It is easy to note that for two different attribute sets \mathcal{A} and \mathcal{Q} which yield the same partition, $deg_{\mathcal{A}}(\mathcal{Q}) = 1$. Further, if \mathcal{G} has k orbits and \mathcal{A} has non-empty intersection with k or $k - 1$ orbits of \mathcal{G} , $deg_{\mathcal{A}}(\mathcal{V}) = 1$.

In proposition 2.1.1, we examine the rough membership function of complete bipartite graph $\mathcal{K}_{m,n}$ and compute \mathcal{A} -positive region of \mathcal{Q} and the degree of dependency.

Proposition 2.1.1. *For a complete bipartite graph $\mathcal{K}_{m,n}$ with bipartition $(\mathcal{V}_1 | \mathcal{V}_2)$, let \mathcal{A} and \mathcal{Q} are two subset of $\mathcal{V}(\mathcal{K}_{m,n})$. Then*

$$\begin{aligned} (i). \quad \mu_{\mathcal{Q}}^{\mathcal{A}}(x) &= \begin{cases} |\mathcal{V}_1 \cap \mathcal{Q}|/|\mathcal{V}_1| & \text{if } x \in \mathcal{V}_1 \\ |\mathcal{V}_2 \cap \mathcal{Q}|/|\mathcal{V}_2| & \text{if } x \in \mathcal{V}_2 \end{cases} \\ (ii). \quad POS_{\mathcal{A}}(\mathcal{Q}) &= \begin{cases} \emptyset & \text{if } \mathcal{A} = \emptyset \wedge \mathcal{Q} \neq \emptyset \\ \mathcal{V} & \text{otherwise} \end{cases} \\ (iii). \quad deg_{\mathcal{A}}(\mathcal{Q}) &= \begin{cases} 0 & \text{if } \mathcal{A} = \emptyset \wedge \mathcal{Q} \neq \emptyset \\ 1 & \text{otherwise} \end{cases} \end{aligned}$$

Proof. (i). According to proposition 2.0.2, we know that $\mathcal{O}_{\mathcal{A}}(x) = \mathcal{V}_i$, if and only if $x \in \mathcal{V}_i$, for $i=1,2$, therefore,

$$\mathcal{O}_{\mathcal{A}}(x) \cap \mathcal{Q} = \begin{cases} |\mathcal{V}_1 \cap \mathcal{Q}| & \text{if } x \in \mathcal{V}_1 \\ |\mathcal{V}_2 \cap \mathcal{Q}| & \text{if } x \in \mathcal{V}_2 \end{cases}$$

Hence the proof follow directly by definition of rough membership.

(ii). If $\mathcal{A} = \emptyset$ and $\mathcal{Q} \neq \emptyset$ then $\gamma_{\mathcal{A}}(\mathcal{K}_{m,n}) = \mathcal{V}$. As $\gamma_{\mathcal{Q}}(\mathcal{K}_{m,n}) = \mathcal{V}_1 | \mathcal{V}_2$. Therefore, $\mathcal{O}_{\mathcal{A}}(x) = \mathcal{V}$ and $\mathcal{O}_{\mathcal{Q}}(x) = \mathcal{V}_i$ for some $i = 1, 2$ which implies that $\mathcal{O}_{\mathcal{A}}(x) \not\subseteq \mathcal{O}_{\mathcal{Q}}(x) \forall x \in \mathcal{V}$, hence $POS_{\mathcal{A}}(\mathcal{Q}) = \emptyset$ if $\mathcal{A} = \emptyset$ and $\mathcal{Q} \neq \emptyset$ then $\gamma_{\mathcal{A}}(\mathcal{K}_{m,n}) = \gamma_{\mathcal{Q}}(\mathcal{K}_{m,n}) = \mathcal{V}$, therefore $\mathcal{O}_{\mathcal{A}}(x) = \mathcal{V} \subseteq \mathcal{O}_{\mathcal{Q}}(x) = \mathcal{V} \forall x \in \mathcal{V}$. Hence $POS_{\mathcal{A}}(\mathcal{Q}) = \mathcal{V}$. If $\mathcal{A} \neq \emptyset$ and $\mathcal{Q} = \emptyset$ then $\gamma_{\mathcal{A}}(\mathcal{K}_{m,n}) = \mathcal{V}_1 | \mathcal{V}_2$ by Proposition 2.0.2, and $\gamma_{\mathcal{Q}}(\mathcal{K}_{m,n}) = \mathcal{V}$. Which implies that $\mathcal{O}_{\mathcal{A}}(x) = \mathcal{V}_i$ for some $i = 1, 2$ and $\mathcal{O}_{\mathcal{Q}}(x) = \mathcal{V}$ for all $x \in \mathcal{V}$. Hence $POS_{\mathcal{A}}(\mathcal{Q}) = \mathcal{V}$. Finally, if $\mathcal{A} \neq \emptyset$ and also $\mathcal{Q} \neq \emptyset$ then by Proposition 2.0.2, we have $\gamma_{\mathcal{A}}(\mathcal{K}_{m,n}) = \gamma_{\mathcal{Q}}(\mathcal{K}_{m,n}) = \mathcal{V}_1 | \mathcal{V}_2$. This implies that $\mathcal{O}_{\mathcal{A}}(x) = \mathcal{O}_{\mathcal{Q}}(x) = \mathcal{V}_i$ for all $x \in \mathcal{V}$ and for some $i=1,2$. Thus in this case we have $POS_{\mathcal{A}}(\mathcal{Q}) = \mathcal{V}$.

(iii). It is following directly from definition and (ii). \square

Note that, for complete graphs K_n and cycle graphs C_n , $|POS_{\mathcal{A}}(\mathcal{Q})| = |\mathcal{V}|$ yields that $deg_{\mathcal{A}}(\mathcal{Q}) = 1$. It is observed that for K_n, C_n , we have $\gamma_{\mathcal{V}} = \mathcal{O}_1$ and for complete bipartite graphs $\mathcal{K}_{m,n}$, we have $\gamma_{\mathcal{V}} = \mathcal{O}_m | \mathcal{O}_n$. Therefore, $deg_{\mathcal{A}}(\mathcal{Q}) = 1$ when $m = n$ and $deg_{\mathcal{A}}(\mathcal{Q}) < 1$ when $m \neq n$. For a path graph P_n , with k orbits and $\mathcal{A}, \mathcal{Q} \subseteq \mathcal{V}$ such that $\mathcal{A} \subseteq \mathcal{Q}$ and $\mathcal{A} \cap \mathcal{O}_{k-i}$ and $\mathcal{Q} \cap \mathcal{O}_{k-1}$ is non-empty $\forall i > 1$ then $deg_{\mathcal{A}}(\mathcal{Q}) < 1$ and $deg_{\mathcal{Q}}(\mathcal{A}) = 1$.

It is easy to see that for a graph \mathcal{G} with $\mathcal{A}, \mathcal{Q} \subseteq \mathcal{V}$, $\mu_{\mathcal{Q}}^{\mathcal{A}}(x) = 0$ if $\gamma(x) \cap \mathcal{Q} = \emptyset$, $\mu_{\mathcal{Q}}^{\mathcal{A}}(x) < 1$ if $\gamma(x) \cap \mathcal{Q} \neq \emptyset$ and $\mu_{\mathcal{Q}}^{\mathcal{A}}(x) = 1$ if $\gamma(x) \subset \mathcal{Q}$ where $\gamma(x)$ is the block of partition γ containing x .

3. ESSENTIAL SETS AND DISCERNIBILITY MATRIX OF GRAPHS

In this section, we will introduce essential sets as well as discernibility matrices of graphs. We will also study the relationship between these two important concepts.

3.1. Essential Sets of Graphs. Chiaselotti et al. [6] presented classical model of Pawlak's core in a more general way. The core of an information system \mathcal{I} associated to \mathcal{G} is the intersection of all reducts, represented by $\mathcal{CORE}(\mathcal{G})$. Hence removal of any attribute belonging to core of a graph leads to change in the indiscernibility partition. In case, core is empty, this extension becomes important because it provides with a set on minimum number of vertices whose removal yields a partition different from the one yielded by all attributes. Concepts of definition 3.1.1 were introduced by Chiaselotti et al. in [6].

Definition 3.1.1. A subset $\mathcal{S} \subseteq \mathcal{A}$, is called \mathcal{I} -essential of \mathcal{G} , if $\gamma_{\mathcal{A} \setminus \mathcal{S}}(\mathcal{G}) \neq \gamma_{\mathcal{A}}(\mathcal{G})$ and $\forall \mathcal{Q} \subsetneq \mathcal{S}$ we have $\gamma_{\mathcal{A} \setminus \mathcal{Q}}(\mathcal{G}) = \gamma_{\mathcal{A}}(\mathcal{G})$.

$\text{ESS}(\mathcal{I})$ represents the collection of all \mathcal{I} -essential subsets of \mathcal{G} . For $l \in \{1, 2, \dots, n\}$, set

$$\text{ESS}_l(\mathcal{I}) = \{\mathcal{S} \in \text{ESS}(\mathcal{I}) : |\mathcal{S}| = l\}$$

and essential numerical sequence of \mathcal{I} is defined by

$$\text{ens}(\mathcal{I}) = (|\text{ESS}_1(\mathcal{I})|, |\text{ESS}_2(\mathcal{I})|, \dots, |\text{ESS}_n(\mathcal{I})|).$$

Finally, essential dimension of \mathcal{I} is defined as the positive integer

$$\text{Edim}(\mathcal{I}) = \min\{l : |\text{ESS}_l(\mathcal{I})| \neq 0\}.$$

Note that, for a graph with at most two orbits, $\text{ESS}(\mathcal{I}) = \emptyset$. For complete graphs K_n , cycles C_n and complete bipartite graphs $\mathcal{K}_{m,n}$, $\text{ESS}(\mathcal{I}) = \emptyset$. As there exists only one orbit for K_n and C_n , but for $\mathcal{K}_{m,n}$, there exists one orbit when $m = n$ and two orbits when $m \neq n$. Also, it is observed that if $\mathcal{A} \subseteq \mathcal{V}$, then there does not exist any set \mathcal{S} for complete graphs, cycles and complete bipartite graphs such that $\gamma_{\mathcal{A}}(\mathcal{I}) \neq \gamma_{\mathcal{A} \setminus \mathcal{S}}(\mathcal{I})$. Therefore, $\text{ens}(K_n) = (0, 0, 0, \dots, 0) \forall n$, $\text{ens}(C_n) = (0, 0, 0, \dots, 0) \forall n$ and $\text{ens}(\mathcal{K}_{m,n}) = (0, 0, 0, \dots, 0) \forall m, n$.

For a path P_n on $n \geq 5$ vertices with vertex set $\mathcal{V} = \{x_1, x_2, \dots, x_n\}$ and edge set $\mathcal{E} = \{x_i x_{i+1} | 1 \leq i \leq n-1\}$. It is observed that $\{\{x_i, x_{n-i+1}\} | 1 \leq i \leq \lfloor \frac{n}{2} \rfloor\}$ is the set of orbits of P_n . Note that each orbit has at most two elements. It can be easily seen that $\text{ESS}_l(\mathcal{I}) = \emptyset$ for $l < 3$. For odd n , $\{x_{\frac{n+1}{2}}\}$ forms an orbit on one vertex and $\text{ESS}_3(\mathcal{I})$ is the union of orbit $\{x_{\frac{n+1}{2}}\}$ with any other orbit of P_n . Also, $\text{ESS}_4(\mathcal{I})$ is the union of any two orbits of P_n . Further, observe that if $n = 2k$ for $k > 2$, then each orbit of path graph P_n will consist of exactly two vertices. Therefore, $\text{Edim}(P_n) = 4$. Similarly, if $n = 2k+1$ for $k \geq 2$, then P_n will contain exactly one orbit of order one and all other orbits of order two. Then $\text{Edim}(P_n) = 3$. Also note that, for a rigid graph, $\text{Edim}(\mathcal{G}) = 2$ as the cardinality of each orbit of a rigid graph is one.

Hence, we have the following straightforward proposition for path graphs P_n .

Proposition 3.1.2. *If \mathcal{G} is a path graph then we have the following:*

- (i) $ESS(P_n) = \emptyset$ for $n \leq 4$.
- (ii) $|ESS_3(P_{2k+1})| = k; k \geq 2$ and $|ESS_3(P_{2k})| = 0; k \geq 2$.
- (iii) $|ESS_4(P_{2k})| = \binom{k}{2}; n \geq 4$
- (iv) $ens(P_{2k+1}) = (0, 0, k, \binom{k}{2}, 0, \dots)$ if $k \geq 2$ and $ens(P_{2k}) = (0, 0, 0, \binom{k}{2}, 0, \dots)$ if $k \geq 3$.

In the next proposition, we prove that essential set \mathcal{S} for any graph \mathcal{G} is exactly union of two orbits.

Proposition 3.1.3. *\mathcal{I} – essential set \mathcal{S} is always union of two orbits of graph \mathcal{G} .*

Proof. Assume contrary that set $\mathcal{S} \subseteq \mathcal{A}$ is a union of more than two orbits of \mathcal{G} then there exists $\mathcal{Q} \subset \mathcal{S}$ such that $\gamma_{\mathcal{A} \setminus \mathcal{Q}}(\mathcal{G}) \neq \gamma_{\mathcal{A}}(\mathcal{G})$, i.e., there exists $x, x' \in \mathcal{A} \setminus \mathcal{Q}$ such that $x \equiv_{\mathcal{A} \setminus \mathcal{Q}} x'$ and $\neg(x \equiv_{\mathcal{A}} x')$ which does not satisfy the minimality condition for \mathcal{S} to be essential. Hence, \mathcal{S} is a union of two orbits of a graph \mathcal{G} . \square

3.2. Discernibility Matrix of Graph. In rough set theory, discernibility matrix is a tool to study information system. Here, we concentrate on the structural concept of discernibility matrix in the context of graph theory.

Let $\mathcal{I} = \langle \mathcal{V}, \mathcal{V}, \mathcal{Val}, \mathcal{F} \rangle$ is an information system with $\mathcal{V} = \{x_1, x_2, \dots, x_n\}$. The discernibility matrix $\Delta[\mathcal{I}]$ of \mathcal{I} is an $n \times n$ matrix with ij th entry, $\Delta_{\mathcal{G}}(x_i, x_j)$, of the matrix is the attribute subset corresponding to the pair (x_i, x_j) given as:

$$\Delta_{\mathcal{G}}(x_i, x_j) = \{a \in \mathcal{A} : \mathcal{F}(x_i, a) \neq \mathcal{F}(x_j, a)\}.$$

Note that in context of granular referencing system, \mathcal{A} serves as reference set and helps in identification whether given two vertices are identical. Two elements in an orbit are similar for any choice of \mathcal{A} . Therefore, two vertices in different orbits are discernible by any of the vertices in those orbits. We define the entries of the discernibility matrix in the following way:

For a graph \mathcal{G} , if $x_i, x_j \in \mathcal{V}$, then

$$\Delta_{\mathcal{G}}(x_i, x_j) = \begin{cases} \mathcal{O}(x_i) \cup \mathcal{O}(x_j) & \text{if } x_i \notin \mathcal{O}(x_j) \\ \emptyset & \text{if } x_i \in \mathcal{O}(x_j) \end{cases}$$

In the next theorem, we show that the discernibility matrices of graphs always characterize graphs uniquely.

Theorem 3.2.1. *For two graphs \mathcal{G}_1 and \mathcal{G}_2 such that $\mathcal{V}(\mathcal{G}) = \mathcal{V}(\mathcal{G}_1) = \mathcal{V}(\mathcal{G}_2)$. $\Delta[\mathcal{G}_1] = \Delta[\mathcal{G}_2]$ if and only if $\mathcal{G}_1 \cong \mathcal{G}_2$.*

Proof. For any two graphs \mathcal{G}_1 and \mathcal{G}_2 , if $\mathcal{G}_1 \cong \mathcal{G}_2$ then without loss of generality, $\Delta[\mathcal{G}_1] = \Delta[\mathcal{G}_2]$ (which is a trivial case). Conversely, suppose that $\Delta_{\mathcal{G}_1}(x_i, x_j) = \Delta_{\mathcal{G}_2}(x_i, x_j)$ for $x_i, x_j \in \mathcal{V}(\mathcal{G})$, then by definition, it follows that $\mathcal{F}_{\mathcal{G}_1}(x_i, x_j) = 0$ if $x_i \notin \mathcal{O}(x_j)$ and $\mathcal{F}_{\mathcal{G}_2}(x_i, x_j) = 1$ if $x_i \in \mathcal{O}(x_j)$. We claim that if $\mathcal{F}_{\mathcal{G}_1}(x_i, x_j) = 1 = \mathcal{F}_{\mathcal{G}_2}(x_i, x_j)$, then $x_i \in \mathcal{O}(x_j)$ and $x_j \in \mathcal{O}(x_i)$ which yields that $\mathcal{O}(x_i) = \mathcal{O}(x_j)$. Thus, preserving the adjacency and degrees of vertices resulting into $\mathcal{E}(\mathcal{G}_1) = \mathcal{E}(\mathcal{G}_2)$. Hence $\mathcal{G}_1 \cong \mathcal{G}_2$. \square

Please note that rows and columns of discernibility matrix corresponding to similar vertices are equal. Hence instead of considering individual elements, their orbits can be considered. Therefore, we first evaluate the classes of orbits of graphs and then consider the rows and columns of discernibility matrix in terms of the

orbits of graphs. We name the discernibility matrix obtained by considering orbits instead of vertices as Quotient Discernibility Matrix (QDM). QDM has order equal to the number of orbits of the graphs. Rows and columns of QDM are labeled by orbits of the graph and entries of QDM are defined as follows: For a graph \mathcal{G} , with two arbitrary orbits \mathcal{O}_i and \mathcal{O}_j

$$\Delta_{\mathcal{G}}(\mathcal{O}_i, \mathcal{O}_j) = \begin{cases} \mathcal{O}_i \cup \mathcal{O}_j & \text{if } i \neq j \\ \emptyset & \text{if } i = j. \end{cases}$$

Next, we represent by $EQDM(\mathcal{I}(\mathcal{G}))$, the collection of all distinct entries of $\Delta[\mathcal{I}]$.

In Proposition 3.2.2, we describe a relationship between $EQDM(\mathcal{I}(\mathcal{G}))$ and $ESS(\mathcal{I}(\mathcal{G}))$, the family of \mathcal{I} -essential subsets.

Proposition 3.2.2. *Let \mathcal{G} be a graph then $EQDM(\mathcal{I}(\mathcal{G})) = ESS(\mathcal{I}(\mathcal{G}))$.*

Proof. Let $S \in ESS(\mathcal{I}(\mathcal{G}))$ then by Proposition 3.1.3, $S = \mathcal{O}_i \cup \mathcal{O}_j$. Note that $\Delta_{\mathcal{G}}(\mathcal{O}_i, \mathcal{O}_j) = \mathcal{O}_i \cup \mathcal{O}_j$ hence $S \subseteq EQDM(\mathcal{I}(\mathcal{G}))$ and $ESS(\mathcal{I}(\mathcal{G})) \subseteq EQDM(\mathcal{I}(\mathcal{G}))$. Conversely, suppose that $S \in EQDM(\mathcal{I}(\mathcal{G}))$ then there exist \mathcal{O}_i and \mathcal{O}_j such that $\Delta_{\mathcal{G}}(\mathcal{O}_i, \mathcal{O}_j) = \mathcal{O}_i \cup \mathcal{O}_j$. Note that $\gamma_{\mathcal{V} \setminus \{\mathcal{O}_i \cup \mathcal{O}_j\}} \neq \gamma_{\mathcal{V}}$ which satisfies first condition for S to be an essential set. Let $T \subset \mathcal{O}_i \cup \mathcal{O}_j$ then we have three possibilities. (i). $T \subseteq \mathcal{O}_i$, (ii). $T \subseteq \mathcal{O}_j$, (iii). $T \cap \mathcal{O}_i \neq \emptyset$ and $T \cap \mathcal{O}_j \neq \emptyset$. It is easy to see in all these three cases that $\gamma_{\mathcal{V} \setminus T}(\mathcal{G}) = \gamma_{\mathcal{V}}(\mathcal{G})$ which satisfies the second condition for S to be an essential set. Hence $S \in ESS(\mathcal{I}(\mathcal{G}))$ and $EQDM(\mathcal{I}(\mathcal{G})) \subseteq ESS(\mathcal{I}(\mathcal{G}))$ which gives required result. \square

Example 3.2.3. *Consider the graph \mathcal{G} shown in FIGURE 1 and its discernibility matrix is given in the following TABLE 1:*

	$\mathcal{O}(0)$	$\mathcal{O}(2)$	$\mathcal{O}(3)$
$\mathcal{O}(0)$	\emptyset	*	*
$\mathcal{O}(2)$	$\mathcal{O}(0) \cup \mathcal{O}(2)$	\emptyset	*
$\mathcal{O}(3)$	$\mathcal{O}(0) \cup \mathcal{O}(3)$	$\mathcal{O}(2) \cup \mathcal{O}(3)$	\emptyset

TABLE 1. Discernibility Matrix $\Delta[\mathcal{I}]$

$$ESS(\mathcal{I}(\mathcal{G})) = EQDM(\mathcal{I}(\mathcal{G})) = \{\mathcal{O}(1)\mathcal{O}(2), \mathcal{O}(1)\mathcal{O}(3), \mathcal{O}(2)\mathcal{O}(3)\}$$

CONCLUSIONS

We have used orbits of graphs to construct information systems to study graphs using rough set theory. We have studied indiscernibility partition, lower and upper approximations of subsets of vertices of graphs, the rough membership function and rough positive region for some well-known families of graphs like cycle, complete and complete bipartite graphs. We have also studied the essential sets of graphs and the discernibility matrix of graphs in terms of orbit partitions. Identifying vertices having similar characteristics simplifies the structures of graphs. The terminology emerging from the merger of rough set theory and graph theory will be useful for exploring new problems associated to symmetries of graphs.

REFERENCES

- [1] Cattaneo, G. (2011). An investigation about rough set theory: Some foundational and mathematical aspects. *Fundamenta Informaticae*, 108(3-4), 197-221.
- [2] Cattaneo, G., Chiaselotti, G., Ciucci, D., and Gentile, T. (2016). On the connection of hypergraph theory with formal concept analysis and rough set theory. *Information Sciences*, 330, 342-357.
- [3] Chen, J., and Li, J. (2012). An application of rough sets to graph theory. *Information Sciences*, 201, 114-127.
- [4] Chiaselotti, G., Ciucci, D., and Gentile, T. (2015). Simple undirected graphs as formal contexts. *Formal Concept Analysis*, 9113, 287-302.
- [5] Chiaselotti, G., Ciucci, D., Gentile, T., and Infusino, F. (2015). Rough set theory applied to simple undirected graphs. *Rough Sets and Knowledge Technology*, 423-434.
- [6] Chiaselotti, G., Ciucci, D., and Gentile, T. (2016). Simple graphs in granular computing. *Information Sciences*, 340, 279-304.
- [7] Hońko, P. (2012). Relational pattern updating. *Information Sciences*, 189, 208-218.
- [8] Hońko, P. (2013). Association discovery from relational data via granular computing. *Information Sciences*, 234, 136-149.
- [9] Huang, B., Zhuang, Y., Li, H., and Wei, D. (2013). A dominance intuitionistic fuzzy-rough set approach and its applications. *Applied Mathematical Modelling*, 37(12-13), 7128-7141.
- [10] Kang, X., Li, D., Wang, S., and Qu, K. (2012). Formal concept analysis based on fuzzy granularity base for different granulations. *Fuzzy Sets and Systems*, 203, 33-48.
- [11] Lin, T. Y. (1999). Data Mining: Granular computing approach. *Methodologies for Knowledge Discovery and Data Mining*, 1574, 24-33.
- [12] Lin, T. Y. (2000). Data mining and machine oriented modeling: a granular approach. *Applied Intelligence*, 13(2), 113-124.
- [13] Midelfart, H., and Komorowski, J. (2002). A rough set framework for learning in a directed acyclic graph. *Rough Sets and Current Trends in Computing*, 2475, 144-155.
- [14] Pawlak, Z. (1991). *Rough sets: Theoretical aspects of reasoning about data*. Kluwer Academic Publishers.
- [15] Pawlak, Z., and Skowron, A. (2007a). Rough sets and boolean reasoning. *Information Sciences*, 177(1), 41-73.
- [16] Pawlak, Z., and Skowron, A. (2007b). Rough sets: Some extensions. *Information Sciences*, 177(1), 28-40.
- [17] Pawlak, Z., and Skowron, A. (2007c). Rudiments of rough sets. *Information Sciences*, 177(1), 3-27.
- [18] Pedrycz, A., Hirota, K., Pedrycz, W., and Dong, F. (2012). Granular representation and granular computing with fuzzy sets. *Fuzzy Sets and Systems*, 203, 17-32.
- [19] Qiu, T., Chen, X., Liu, Q., and Huang, H. (2010). Granular computing approach to finding association rules in relational database. *International Journal of Intelligent Systems*, 25, 165-179.
- [20] Skowron, A., and Rauszer, C. (1992). The discernibility matrices and functions in information systems. *Intelligent Decision Support*, 11, 331-362.
- [21] Tang, J., She, K., and Zhu, W. (2012). Matroidal structure of rough sets from the viewpoint of graph theory. *Journal of Applied Mathematics*, 2012, 1-27.
- [22] Wang, C., and Chen, D. (2010). A short note on some properties of rough groups. *Computers and Mathematics with Applications*, 59(1), 431-436.
- [23] Wang, S., Zhu, Q., Zhu, W., and Min, F. (2013). Equivalent characterizations of some graph problems by covering-based rough sets. *Journal of Applied Mathematics*, 2013, 1-7.
- [24] Wei-Zhi Wu, Yee Leung, and Ju-Sheng Mi. (2009). Granular computing and knowledge reduction in formal contexts. *IEEE Transactions on Knowledge and Data Engineering*, 21(10), 1461-1474.
- [25] Yao, Y. Y. (2001). On modeling data mining with granular computing. *25th Annual International Computer Software and Applications Conference. COMPSAC 2001*, 2001, 638-643.
- [26] Yao, Y. Y. (2002). A granular computing approach to machine learning. *Proceedings of the 1st International Conference on Fuzzy Systems and Knowledge Discovery*, 732-736.
- [27] Zadeh, L. A. (1996). Fuzzy set and information granularity. *Advances in Fuzzy Systems — Applications and Theory*, 433-448.

CENTRE FOR ADVANCED STUDIES IN PURE AND APPLIED MATHEMATICS, BAHAUDDIN ZAKARIYA
UNIVERSITY, MULTAN, PAKISTAN.

Email address: `imran.javaid@bzu.edu.pk`, `shahroz.ali0304@gmail.com`, `shahidurrehman1982@gmail.com`,
`aqsa6977@gmail.com`