

Cosmology and Warped Space Times in Dynamical String Tension Theories

E.I. Guendelman

Department of Physics, Ben-Gurion University of the Negev, Beer-Sheva, Israel

Frankfurt Institute for Advanced Studies, Giersch Science Center, Campus Riedberg,
Frankfurt am Main, Germany

Bahamas Advanced Studies Institute and Conferences, 4A Ocean Heights, Hill View
Circle, Stella Maris, Long Island, The Bahamas

E-mail: guendel@bgu.ac.il,

Abstract

The string and brane tensions do not have to be put in by hand, they can be dynamically generated, as in the case when we formulate string and brane theories in the modified measure formalism. Then string and brane tensions appears, but as an additional dynamical degree of freedom. It can be seen however that these string or brane tensions are not universal, but rather each string and each brane generates its own tension, which can have a different value for each string or brane. The consequences of this for the spectrum of these string and brane theories is profound both in the ultraviolet behavior as in the low energy physics. There should be also a considerable effect for the effective gravity theories derived from these theories. We consider new background fields that can couple to these new types of extended objects, one of them, the "tension scalar" is capable of changing locally along the world sheet the value of the tension of the extended object. When many strings probing the same region of space are considered this tension scalar is constrained by the requirement of quantum conformal invariance. For the case of two strings probing the same region of space with different dynamically generated tensions, there are two different metrics, associated to the different strings, that have to satisfy vacuum Einsteins equations and the consistency of these two Einstein's equation determines the tension scalar. The universal metric, common to both strings generically does not satisfy Einstein's equation. The problem is analyzed in the case of a Schwarzschild background and for the cosmological case of a Kasner type solution. In the case of the flat space for the string associated metrics, in the Milne representation, for the case of two types of string tensions, there are solutions with negative string tension at the early universe that whose tension approaches zero in the late universe and a positive string tension type of strings appears for the late universe with its tension approaching a constant value at the late universe. The universal metric is not flat, instead it represents a non singular bounce cosmology. The case in a warped space time where positive and negative string tensions are separated by a spontaneously generated wall is also studied.

1 Introduction

String and Brane Theories have been studied as candidates for the theory of all matter and interactions including gravity, in particular string theories [1]. But string theory has a dimensionful parameter, the tension of the string, in its standard formulation, the same is true for brane theories in their better known formulations.

The appearance of a dimensionful string tension and brane theories from the start appears somewhat unnatural. Previously however, in the framework of a Modified Measure Theory, a formalism originally used for gravity theories, see for example [2, 3, 5, 4, 6, 7, 8, 9], the tension was derived as an additional degree of freedom [10, 11, 12, 13, 14, 15, 16]. See also the treatment by Townsend and collaborators [17, 18].

A floating cosmological is a generic feature of the modified measure theories of gravity [2, 3, 5, 4, 6, 7, 8, 9], including the covariant formulation of the unimodular theory [19], which is in fact a particular case of a modified measure theory, as reviewed in [20].

The tension of the string plays a very similar role to the cosmological constant in four dimensional gravity, but the analogous situation and the role of the cosmological constant is quite different to that of the string tension, because while several world sheets of strings can exist in the same universe and in this way many strings can probe at the same time the same region of space time, but the same is not the case for the cosmological constant, where every cosmological constant defines necessarily a different universe.

This paper is organized as follows. In Section 2 we review the modified-measure theory in the string context. In Section 3 we review the modified-measure theory in the brane context. The modified-measure theories of strings and branes rely on two basic elements, the modified measure and the existence of internal gauge fields in the strings and branes, the equations of motion of these gauge fields lead to an equation of motion whose integration constant is the Tension of the extended object. In Section 4 we discuss the fact that this tension generation (the integration constant) could take place independently for each world sheet separately, which would mean that the string or brane tension is not a fundamental coupling in nature and it could be different for different strings or branes. In section 5 we discuss possible consequences for physics that are derived from these string theories with dynamical string tension generation from general considerations. In section 6 we discuss possible background fields that can be introduced into the theory for the bosonic case, we find that for strings and branes a new field can be introduced that changes the value of the tension of the extended object along its world sheet, we call this the tension scalar. Section 6, the study of this tension scalar and its effects is the bulk of the contents of this paper. We present first the coupling of gauge fields in the extended objects to currents in the world sheet of the extended object that couple to the gauge fields, as a consequence this coupling induces variations of the tension along the world sheet of the extended object. Then we consider a bulk scalar and how this scalar naturally can induce this world sheet current that couples to the internal gauge fields, the equation of motion of the internal gauge field lead to the remarkably simple equation that the local value of the tension along the string is given by $T = g\phi + T_i$, where g is a coupling constant that defines the coupling of the bulk scalar to the world sheet gauge fields and T_i is an integration constant which can be different for each string in the universe. Then each

string is considered as an independent system that can be quantized. We take into account the string generation by introducing the tension as a function of the scalar field as a factor inside a Polyakov type action with such string tension, then the metric and the factor $g\phi + T_i$ enter together in this effective action, so if there was just one string the factor could be incorporated into the metric and the condition of conformal invariance will not say very much about the scalar ϕ , but if many strings are probing the same regions of space time, then considering a background metric $g_{\mu\nu}$, for each string the "string dependent metric" $(\phi + T_i)g_{\mu\nu}$ appears and in the absence of other background fields, like dilaton and antisymmetric tensor fields, Einstein's equations apply for each of the metrics $(\phi + T_i)g_{\mu\nu}$, considering two types of strings with $T_1 \neq T_2$. We call $g_{\mu\nu}$ the universal metric, which in fact does not necessarily satisfy Einstein's equations. The problem is analyzed in the case of a Schwarzschild background and for the cosmological case of a Kasner type solution. In the case of the flat space for the string associated metrics, in the Milne representation, for the case of two types of string tensions, there are solutions with negative string tension at the early universe that whose tension approaches zero in the late universe and a positive string tension type of strings appears for the late universe with its tension approaching a constant value at the late universe. The universal metric is not flat, instead it represents a non singular bounce cosmology. The case where positive and negative string tensions are separated by a spontaneously generated wall is also studied, in warp space times of the type considered by Wesson and collaborators. Other possible effects are discussed at a more qualitative level.

2 The Modified Measure Theory String Theory

The standard world sheet string sigma-model action using a world sheet metric is [21], [22], [23]

$$S_{sigma-model} = -T \int d^2\sigma \frac{1}{2} \sqrt{-\gamma} \gamma^{ab} \partial_a X^\mu \partial_b X^\nu g_{\mu\nu}. \quad (1)$$

Here γ^{ab} is the intrinsic Riemannian metric on the 2-dimensional string worldsheet and $\gamma = \det(\gamma_{ab})$; $g_{\mu\nu}$ denotes the Riemannian metric on the embedding spacetime. T is a string tension, a dimension full scale introduced into the theory by hand.

From the variations of the action with respect to γ^{ab} and X^μ we get the following equations of motion:

$$T_{ab} = (\partial_a X^\mu \partial_b X^\nu - \frac{1}{2} \gamma_{ab} \gamma^{cd} \partial_c X^\mu \partial_d X^\nu) g_{\mu\nu} = 0, \quad (2)$$

$$\frac{1}{\sqrt{-\gamma}} \partial_a (\sqrt{-\gamma} \gamma^{ab} \partial_b X^\mu) + \gamma^{ab} \partial_a X^\nu \partial_b X^\lambda \Gamma_{\nu\lambda}^\mu = 0, \quad (3)$$

where $\Gamma_{\nu\lambda}^\mu$ is the affine connection for the external metric.

There are no limitations on employing any other measure of integration different than $\sqrt{-\gamma}$. The only restriction is that it must be a density under arbitrary diffeomorphisms (reparametrizations) on the underlying spacetime manifold. The modified-

measure theory is an example of such a theory.

In the framework of this theory two additional worldsheet scalar fields $\varphi^i (i = 1, 2)$ are introduced. A new measure density is

$$\Phi(\varphi) = \frac{1}{2} \epsilon_{ij} \epsilon^{ab} \partial_a \varphi^i \partial_b \varphi^j. \quad (4)$$

Then the modified bosonic string action is (as formulated first in [10] and latter discussed and generalized also in [11])

$$S = - \int d^2 \sigma \Phi(\varphi) \left(\frac{1}{2} \gamma^{ab} \partial_a X^\mu \partial_b X^\nu g_{\mu\nu} - \frac{\epsilon^{ab}}{2\sqrt{-\gamma}} F_{ab}(A) \right), \quad (5)$$

where F_{ab} is the field-strength of an auxiliary Abelian gauge field A_a : $F_{ab} = \partial_a A_b - \partial_b A_a$.

It is important to notice that the action (5) is invariant under conformal transformations of the intrinsic measure combined with a diffeomorphism of the measure fields,

$$\gamma_{ab} \rightarrow J \gamma_{ab}, \quad (6)$$

$$\varphi^i \rightarrow \varphi'^i = \varphi'^i(\varphi^i) \quad (7)$$

such that

$$\Phi \rightarrow \Phi' = J \Phi \quad (8)$$

Here J is the jacobian of the diffeomorphism in the internal measure fields which can be an arbitrary function of the world sheet space time coordinates, so this can be called indeed a local conformal symmetry.

To check that the new action is consistent with the sigma-model one, let us derive the equations of motion of the action (5).

The variation with respect to φ^i leads to the following equations of motion:

$$\epsilon^{ab} \partial_b \varphi^i \partial_a (\gamma^{cd} \partial_c X^\mu \partial_d X^\nu g_{\mu\nu} - \frac{\epsilon^{cd}}{\sqrt{-\gamma}} F_{cd}) = 0. \quad (9)$$

It implies

$$\gamma^{cd} \partial_c X^\mu \partial_d X^\nu g_{\mu\nu} - \frac{\epsilon^{cd}}{\sqrt{-\gamma}} F_{cd} = M = \text{const}. \quad (10)$$

The equations of motion with respect to γ^{ab} are

$$T_{ab} = \partial_a X^\mu \partial_b X^\nu g_{\mu\nu} - \frac{1}{2} \gamma_{ab} \frac{\epsilon^{cd}}{\sqrt{-\gamma}} F_{cd} = 0. \quad (11)$$

We see that these equations are the same as in the sigma-model formulation (2), (3). Namely, taking the trace of (11) we get that $M = 0$. By solving $\frac{\epsilon^{cd}}{\sqrt{-\gamma}} F_{cd}$ from (24) (with $M = 0$) we obtain (2).

A most significant result is obtained by varying the action with respect to A_a :

$$\epsilon^{ab}\partial_b\left(\frac{\Phi(\varphi)}{\sqrt{-\gamma}}\right)=0. \quad (12)$$

Then by integrating and comparing it with the standard action it is seen that

$$\frac{\Phi(\varphi)}{\sqrt{-\gamma}}=T. \quad (13)$$

That is how the string tension T is derived as a world sheet constant of integration opposite to the standard equation (1) where the tension is put ad hoc. The variation with respect to X^μ leads to the second sigma-model-type equation (3). The idea of modifying the measure of integration proved itself effective and profitable. This can be generalized to incorporate super symmetry, see for example [11], [13], [12], [14]. For other mechanisms for dynamical string tension generation from added string world sheet fields, see for example [17] and [18]. However the fact that this string tension generation is a world sheet effect and not a universal uniform string tension generation effect for all strings has not been sufficiently emphasized before. Now we go and review the Modified Measure Brane Theory

3 The Modified Measure Brane Theory

The standard world sheet string sigma-model action using a world sheet metric is [24]:

$$S_{sigma-model} = -T \int d^d\sigma \frac{1}{2} \sqrt{-\gamma} (\gamma^{ab} \partial_a X^\mu \partial_b X^\nu g_{\mu\nu} + 2\Lambda). \quad (14)$$

Notice that now a cosmological term has been added as well, that was not needed in the usual formulation of the string theory or in the modified measure formulation of the string theory. In the modified measure formulation of the brane theory such term will not be required. As it is well known, and as we will review, in the standard formulation, this cosmological term needs to be fine tuned. Here again γ^{ab} is the intrinsic Riemannian metric on the d-dimensional brane worldsheet and $\gamma = \det(\gamma_{ab})$; $g_{\mu\nu}$ denotes the Riemannian metric on the embedding spacetime. T is a brane tension, a dimension full scale introduced into the theory by hand.

From the variations of the action with respect to γ^{ab} and X^μ we get the following equations of motion:

$$T_{ab} = (\partial_a X^\mu \partial_b X^\nu - \frac{1}{2} \gamma_{ab} \gamma^{cd} \partial_c X^\mu \partial_d X^\nu) g_{\mu\nu} - \gamma_{ab} \Lambda = 0, \quad (15)$$

$$\frac{1}{\sqrt{-\gamma}} \partial_a (\sqrt{-\gamma} \gamma^{ab} \partial_b X^\mu) + \gamma^{ab} \partial_a X^\nu \partial_b X^\lambda \Gamma_{\nu\lambda}^\mu = 0, \quad (16)$$

where $\Gamma_{\nu\lambda}^\mu$ is the affine connection for the external metric.

From (15) we see that without the cosmological constant term, or if this term is not fine tuned, the equations of motion are inconsistent upon considering the trace for example. This feature is absent in the modified measure brane theory, where

an integration constant, that takes the role of the cosmological term is determined dynamically by the consistency of the equations of motion.

To show this, again, as in the string case, we consider other measure of integration different than $\sqrt{-\gamma}$. The only restriction is that it must be a density under arbitrary diffeomorphisms (reparametrizations) on the underlying spacetime manifold, as we have seen before while reviewing the modified measure string theory, a metric independent measure build along the lines of what we did in the previous section is an example for that

In the framework of a d dimensional brane d additional worldsheet scalar fields $\varphi^i (i = 1, 2, \dots, d)$ are introduced. The new measure density is defined now as

$$\Phi(\varphi) = \epsilon_{ijk\dots m} \epsilon^{abc\dots d} \partial_a \varphi^i \partial_b \varphi^j \dots \partial_d \varphi^m \quad (17)$$

Then the modified bosonic brane action is

$$S = - \int d^d \sigma \Phi(\varphi) \left(\frac{1}{2} \gamma^{ab} \partial_a X^\mu \partial_b X^\nu g_{\mu\nu} - \frac{\epsilon^{abcd\dots}}{2\sqrt{-\gamma}} F_{abcd\dots}(A) \right), \quad (18)$$

where $F_{abcd\dots}$ is the rank d totally antisymmetric field-strength derived from an auxiliary abelian gauge field $A_{acd\dots}$ of rank $d-1$: $F_{ab\dots cd} = \partial_a A_{bcd\dots} + \dots$, the totally antisymmetrized curl of a $d-1$ potential $A_{bcd\dots}$.

In the brane modified theory there is a scale invariance, although it is only a global one, as opposed to the string case, where it is local,

$$\gamma_{ab} \rightarrow J \gamma_{ab}, \quad (19)$$

$$\varphi^i \rightarrow \varphi'^i = \lambda^{ij}(\varphi^j) \quad (20)$$

such that

$$\Phi \rightarrow \Phi' = J \Phi \quad (21)$$

That is if $\det(\lambda^{ij}) = J$, where J, λ^{ij} are just a constant number and a constant matrix now, and finally the antisymmetric tensor field $A_{bcd\dots}$ must transform as

$$A_{bcd\dots} \rightarrow J^{\frac{d-2}{2}} A_{bcd\dots} \quad (22)$$

To check that the new action is consistent with the sigma-model one, let us derive the equations of motion of the action (14). To start, the variation with respect to φ^i leads to the following equations of motion:

$$K_b^a \partial_a (\gamma^{cd} \partial_c X^\mu \partial_d X^\nu g_{\mu\nu} - \frac{\epsilon^{cdef\dots}}{\sqrt{-\gamma}} F_{cdef\dots}) = 0. \quad (23)$$

Where the matrix K_b^a is just like the measure, except that one factor of a derivative of a measure field is missing, so that there are two free indices therefore the determinant of K_b^a is a power of the measure, so if the measure is non vanishing, this implies

$$\gamma^{cd} \partial_c X^\mu \partial_d X^\nu g_{\mu\nu} - \frac{\epsilon^{cdef\dots}}{\sqrt{-\gamma}} F_{cdef\dots} = M = \text{const.} \quad (24)$$

The equations of motion with respect to γ^{ab} are

$$T_{ab} = \partial_a X^\mu \partial_b X^\nu g_{\mu\nu} - \frac{1}{2} \gamma_{ab} \frac{\epsilon^{cdef\dots}}{\sqrt{-\gamma}} F_{cdef\dots} = 0. \quad (25)$$

We will see now that the final equations are the same as in the sigma-model formulation (2), (3). Namely, taking the trace of (25), using also (24), and then reinserting into (25), we get that $M \neq 0$ now. By solving $\frac{\epsilon^{cdef\dots}}{\sqrt{-\gamma}} F_{cdef\dots}$ from (24) we obtain a relation between the intrinsic metric and the induced metric if $p \neq 1$, (where p is the number of spacelike world sheet coordinates ($p = d - 1$)) that if we do not deal with the string case ($p = 1$),

$$\gamma_{ab} = \frac{1-p}{M} \partial_a X^\mu \partial_b X^\nu g_{\mu\nu} \quad (26)$$

Using the global scale symmetry of the modified measure brane theory we can set $\frac{1-p}{M} = 1$ if M is negative if we wish, for $p \neq 1$. That is, the intrinsic metric in the world sheet becomes equal to the induced metric from the embedding metric, while in the string case there is only a conformal equivalence between the intrinsic metric in the world sheet and the induced metric. Varying the action with respect to $A_{abc\dots}$:

$$\epsilon^{abc\dots d} \partial_d \left(\frac{\Phi(\varphi)}{\sqrt{-\gamma}} \right) = 0. \quad (27)$$

Then by integrating and comparing it with the standard action it is seen that

$$\frac{\Phi(\varphi)}{\sqrt{-\gamma}} = T. \quad (28)$$

That is how the brane tension T is derived as a world sheet constant of integration, opposite to the standard equation (1) where the tension is put in from the start. One can see indeed that the string energy excitations will be proportional to T in exactly the same way as it is the case in the standard case. The variation with respect to X^μ leads to the second sigma-model-type equation. The idea of modifying the measure of integration proved itself effective and profitable.

4 Each String and Each Brane in its own world sheet determines its own tension. Therefore the tension is not universal for all strings or branes

If we look at a single string, the dynamical string tension theories and the standard string theories appear indeed indistinguishable, there are however more than one string and/or one brane in the universe then, let us now observe indeed that it does not appear that the string tension or the brane tension derived in the sections above correspond to "the" string or brane tensions of the theory. The derivation of the string or brane tensions in the previous sections holds for a given string or brane, there is no obstacle that for another string or brane these could acquire a different string or brane tension. In other words, the string or brane tension is a world sheet constant, but it does not appear to be a universal constant same for all strings and for all branes. Similar situation takes place in the dynamical string generation proposed by Townsend for example [17], in that paper worldsheet fields include an electromagnetic gauge potential. Its

equations of motion are those of the Green-Schwarz superstring but with the string tension given by the circulation of the worldsheet electric field around the string. So again, in [17] also a string will determine a given tension, but another string may determine another tension. If the tension is a universal constant valid for all strings, that would require an explanation in the context of these dynamical tension string theories, for example some kind of interactions that tend to equalize string tensions, or that all strings in the universe originated from the splittings of one primordial string or some other mechanism.

In any case, if one believes for example in strings, on the light of the dynamical string tension mechanism being a process that takes place at each string independently, we must ask whether all strings have the same string tension.

5 Equations for the Background fields and a new background field

As discussed by Polchinski for example in [27], gravity can be introduced in two different ways in string theory. One way is by recognizing the graviton as one of the fundamental excitations of the string, the other is by considering the effective action of the embedding metric, by integrating out the string degrees of freedom and then the embedding metric and other originally external fields acquire dynamics which is enforced by the requirement of a zero beta function. These equations fortunately appear to be string tension independent for the critical dimension $D=26$ in the bosonic string for example, so they will not be changed by introducing different strings with different string tensions, if these tensions are constant along the world sheet.

However, in addition to the traditional background fields usually considered in conventional string theory, one may consider as well an additional scalar field that induces currents in the string world sheet and since the current couples to the world sheet gauge fields, this produces a dynamical tension controlled by the external scalar field as shown at the classical level in [25]. In the next two subsections we will study how this comes about in two steps, first we introduce world sheet currents that couple to the internal gauge fields in Strings and Branes and second we define a coupling to an external scalar field by defining a world sheet currents that couple to the internal gauge fields in Strings and Branes that is induced by such external scalar field. This is very much in accordance to the philosophy of Schwinger [28] that proposed long time ago that a field theory must be understood by probing it with external sources.

As we will see however, there will be a fundamental difference between this background field and the more conventional ones (the metric, the dilaton field and the two index anti symmetric tensor field) which are identified with some string excitations as well. Instead, here we will see that a single string does not provide dynamics for this field, but rather when the condition for world sheet conformal invariance is implemented for two strings which sample the same region of space time, so it represents a collective effect instead.

5.1 Introducing world sheet currents that couple to the internal gauge fields in Strings and Branes

If to the action of the brane (18) we add a coupling to a world-sheet current $j^{a_2 \dots a_{p+1}}$, $p = d - 1$ (the case $p = 1$ represents a string) i.e. a term

$$S_{\text{current}} = \int d^{p+1} \sigma A_{a_2 \dots a_{p+1}} j^{a_2 \dots a_{p+1}}, \quad (29)$$

then the variation of the total action with respect to $A_{a_2 \dots a_{p+1}}$ gives

$$\epsilon^{a_1 \dots a_{p+1}} \partial_{a_1} \left(\frac{\Phi}{\sqrt{-\gamma}} \right) = j^{a_2 \dots a_{p+1}}. \quad (30)$$

We thus see indeed that, in this case, the dynamical character of the brane is crucial here.

5.2 Coupling to a bulk scalar field, the tension field

Suppose that we have an external scalar field $\phi(x^\mu)$ defined in the bulk. From this field we can define the induced conserved world-sheet current

$$j^{a_1 \dots a_{p+1}} = e \partial_\mu \phi \frac{\partial X^\mu}{\partial \sigma^a} \epsilon^{a a_2 \dots a_{p+1}} \equiv e \partial_a \phi \epsilon^{a a_2 \dots a_{p+1}}, \quad (31)$$

where e is some coupling constant.

Then (30) can be integrated to obtain

$$T = \frac{\Phi}{\sqrt{-\gamma}} = e\phi + T_i, \quad (32)$$

or equivalently

$$\Phi = \sqrt{-\gamma}(e\phi + T_i), \quad (33)$$

The constant of integration T_i may vary from one extended object to the other. Notice that for the string case the interaction is metric independent since the internal gauge field does not transform under the conformal transformations. This interaction does not therefore spoil the world sheet conformal transformation invariance in the case the field ϕ does not transform under this transformation. One may interpret (33) as the result of integration out classically (through integration of equations of motion) or quantum mechanically (by functional integration of the internal gauge field, respecting the boundary condition that characterizes the constant of integration T_i for a given string or brane). Then replacing $\Phi = \sqrt{-\gamma}(e\phi + T_i)$ back into the remaining terms in the action gives a correct effective action for each string. Each string is going to be quantized with each a different T_i . The consequences of an independent quantization of many strings with different T_i covering the same region of space time will be studied in the next section.

5.3 Constraints from quantum conformal invariance for the Tension field in the string case, when several strings share the same region of space for some gravitational backgrounds. The emergence of a Multi Metric Gravity Theory, or a Multi Measures Gravity Theory

If we have a scalar field coupled to a string or a brane in the way described in the sub section above, i.e. through the current induced by the scalar field in the extended object, according to eq. (33), so we have two sources for the variability of the tension when going from one string to the other: one is the integration constant T_i which varies from string to string and the other the local value of the scalar field, which produces also variations of the tension even within the string or brane world sheet.

As we discussed in the previous section, we can incorporate the result of the tension as a function of scalar field ϕ , given as $e\phi + T_i$, for a string with the constant of integration T_i by defining the action that produces the correct equations of motion for such string, adding also other background fields, the anti symmetric two index field $A_{\mu\nu}$ that couples to $\epsilon^{ab}\partial_a X^\mu \partial_b X^\nu$ and the dilaton field φ that couples to the topological density $\sqrt{-\gamma}R$

$$S_i = - \int d^2\sigma (e\phi + T_i) \frac{1}{2} \sqrt{-\gamma} \gamma^{ab} \partial_a X^\mu \partial_b X^\nu g_{\mu\nu} + \int d^2\sigma A_{\mu\nu} \epsilon^{ab} \partial_a X^\mu \partial_b X^\nu + \int d^2\sigma \sqrt{-\gamma} \varphi R. \quad (34)$$

Notice that if we had just one string, or if all strings will have the same constant of integration $T_i = T_0$.

In any case, it is not our purpose here to do a full generic analysis of all possible background metrics, antisymmetric two index tensor field and dilaton fields, instead, we will take cases where the dilaton field is a constant or zero, and the antisymmetric two index tensor field is pure gauge or zero, then the demand of conformal invariance for $D = 26$ becomes the demand that all the metrics

$$g_{\mu\nu}^i = (e\phi + T_i) g_{\mu\nu} \quad (35)$$

will satisfy simultaneously the vacuum Einstein's equations,

5.3.1 The case where all all string tensions are the same, i.e., $T_i = T_0$, and the appearance of a target space conformal invariance

If all $T_i = T_0$, we just redefine our background field so that $e\phi + T_0 \rightarrow e\phi$ and then in the effective action for all the strings the same combination $e\phi g_{\mu\nu}$, and only this combination will be determined by the requirement that the conformal invariance in the world sheet of all strings be preserved quantum mechanically, that is, that the beta function be zero. So in this case we will not be able to determine $e\phi$ and $g_{\mu\nu}$ separately, just the product $e\phi g_{\mu\nu}$, so the equation obtained from equating the beta function to zero will have the target space conformal invariance $e\phi \rightarrow F(x)e\phi$, $g_{\mu\nu} \rightarrow F(x)^{-1} g_{\mu\nu}$.

That is, there is no independent dynamics for the Tension Field in this case. So in conclusion, if we just look at one string, or if we look at a set of strings, all of them equal string tensions, then the tension field is not observable, can be gauged away by the target space conformal invariance explained above. Another way to see this is

to realize that there is no string excitation of a single string that corresponds to the tension field, unlike the metric, the dilaton or the two index antisymmetric potential.

On the other hand, if there are at least two types of string tensions, that symmetry will not exist and there is the possibility of determining separately $e\phi$ and $g_{\mu\nu}$ (setting the dilaton and the two index antisymmetric potential to be trivial).

So the tension field dynamics corresponds to a non trivial collective dynamics of strings that cannot be manifest for the equal string tensions case or the single string case. Of course collective effects are very familiar in physics, for example the statistics of fermions and bosons can be understood only when we consider many particles. Here we will see in the next subsection how the Tension field gets dynamics in the case of two strings with different tensions.

5.3.2 The case where not all string tensions are the same, with special emphasis of two types of strings with $T_1 \neq T_2$,

The interesting case to consider is therefore many strings with different T_i , let us consider the simplest case of two strings, labeled 1 and 2 with $T_1 \neq T_2$, then we will have two Einstein's equations, for $g_{\mu\nu}^1 = (e\phi + T_1)g_{\mu\nu}$ and for $g_{\mu\nu}^2 = (e\phi + T_2)g_{\mu\nu}$,

$$R_{\mu\nu}(g_{\alpha\beta}^1) = 0 \quad (36)$$

and, at the same time,

$$R_{\mu\nu}(g_{\alpha\beta}^2) = 0 \quad (37)$$

These two simultaneous conditions above impose a constraint on the tension field ϕ , because the metrics $g_{\alpha\beta}^1$ and $g_{\alpha\beta}^2$ are conformally related, but Einstein's equations are not conformally invariant, so the condition that Einstein's equations hold for both $g_{\alpha\beta}^1$ and $g_{\alpha\beta}^2$ is highly non trivial.

Let us consider the case that one of the metrics, say $g_{\alpha\beta}^2$ is a Schwarzschild solution, either a 4 D Schwarzschild solution or a product flat of Torus compactified extra dimensions

$$ds_2^2 = -(1 - 2GM/r)dt^2 + \frac{dr^2}{1 - 2GM/r} + r^2 d\Omega^2 + dy_4^2 + dy_5^2 + \dots + dy_{25}^2 \quad (38)$$

or just a 26 D Schwarzschild solution, in this case, it does not appear possible to have a conformally transformed $g_{\alpha\beta}^2$ for anything else than in the case that the conformal factor that transforms the two metrics is a constant, let us call it c , in that case $g_{\alpha\beta}^1$ is a Schwarzschild solution of the same type, just with a different mass parameter and different sizes of extra dimensions if the compactified solution is considered. Similar consideration holds for the case the 2 metric is a Kasner solution,

$$ds_2^2 = -dt^2 + t^{2p_1} dx_1^2 + t^{2p_2} dx_2^2 + \dots + t^{2p_{25}} dx_{25}^2 \quad (39)$$

with the well known constraints on the powers p_i that the sum of the p_i s is 1 and also that the sum of the squares of the p_i s is also 1. Then in this case also, it does not appear possible to have a conformally transformed $g_{\alpha\beta}^2$ for anything else than in the case that the conformal factor that transforms the two metrics is a constant, we will find other cases where the conformal factor will not be a constant, let us call the conformal factor c in general. One can also study metrics used to describe gravitational

radiation, then again, multiplying by a constant both the background flat space and the perturbation gives us also a solution of vacuum Einstein's equations.

Then for these situations, we have,

$$e\phi + T_1 = c(e\phi + T_2) \quad (40)$$

which leads to a solution for $e\phi$

$$e\phi = \frac{cT_2 - T_1}{1 - c} \quad (41)$$

which leads to the tensions of the different strings to be

$$e\phi + T_1 = \frac{c(T_2 - T_1)}{1 - c} \quad (42)$$

and

$$e\phi + T_2 = \frac{(T_2 - T_1)}{1 - c} \quad (43)$$

It is important that we were forced to consider a multi metric situation. One must also realize that the constant c is physical, because both metrics live in the same spacetime, so even if c is a constant, we are not allowed to perform a coordinate transformation, consisting for example of a rescaling of coordinates for one of the metrics and not do the same transformation for the other metric.

Other way to see that c is physical consist of considering the scalar consisting of the ratio of the two measures $\sqrt{-g^1}$ and $\sqrt{-g^2}$ where $g^1 = \det(g_{\alpha\beta}^1)$ and $g^2 = \det(g_{\alpha\beta}^2)$, and we find that the scalar $\frac{\sqrt{-g^1}}{\sqrt{-g^2}} = c^{D/2}$, showing that c is a coordinate invariant.

Let us study now a case where c is not a constant, we will also focus on a cosmological case. To find this it is useful to consider flat space in the Milne representation, $D = 4$ this reads,

$$ds^2 = -dt^2 + t^2(d\chi^2 + \sinh^2\chi d\Omega_2^2) \quad (44)$$

where $d\Omega_2^2$ represent the contribution of the 2 angles to the metric when using spherical coordinates, that is, it represents the metric of a two dimensional sphere of unit radius. In D dimensions we will have a similar expression but now we must introduce the metric of a $D - 2$ unit sphere $d\Omega_{D-2}^2$ so we end up with the following metric that we will take as the metric 2

$$ds_2^2 = -dt^2 + t^2(d\chi^2 + \sinh^2\chi d\Omega_{D-2}^2) \quad (45)$$

For the metric 1 we will take the metric that we would obtain from the coordinate $t \rightarrow 1/t$ and we furthermore multiply by a constant σ , so

$$ds_1^2 = \frac{\sigma}{t^4}(-dt^2 + t^2(d\chi^2 + \sinh^2\chi d\Omega_{D-2}^2)) \quad (46)$$

Then the equations (40), (41), (42), (43), with $c = \frac{\sigma}{t^4}$. If we want that the only possible singularities take place at $t = 0$ only, we must take σ negative, and therefore the two strings have string tensions of opposite signs.

Whether strings 1 or 2 are the ones with negative tensions depends on the sign of $T_2 - T_1$. If we want the strings with negative tension to exist only in the early universe, we must take $T_2 - T_1$ to be negative. At the same time there will not be positive tension

strings in the early universe, but in the late universe approaches a constant value. The positive string tension are the strings 1, with zero tension in the early universe and the tension $T_1 - T_2$ in the late universe. The negative string tension are the strings 2, with $T_1 - T_2$ tension in the early universe and the tension zero tension in the late universe.

5.4 The Non Singular Bouncing Solution for the Universal Metric

As we have seen when the space time is probed by two types of strings, there are two metrics that have to satisfy the vacuum Einstein's equations, this is enough to solve the problem, The interesting thing however is that the universal metric $g_{\mu\nu}$ does not have to satisfy Einstein's equation. We can see this by solving $g_{\mu\nu}$ in terms of one of the metrics, for example from $g_{\mu\nu}^2 = (e\phi + T_2)g_{\mu\nu}$, we have that

$$ds^2 = g_{\mu\nu}dx^\mu dx^\nu = \left(\frac{1-c}{T_2-T_1}\right)(-dt^2 + t^2(d\chi^2 + \sinh^2\chi d\Omega_{D-2}^2)) \quad (47)$$

and considering that $c = \frac{\sigma}{t^4} = \frac{-K}{t^4}$, where K is positive. So the coefficient of the hyperbolic D -1 dimensional metric $d\chi^2 + \sinh^2\chi d\Omega_{D-2}^2$ is $\frac{t^2 + \frac{-K}{t^4}}{T_2-T_1}$, showing a contraction, a bounce and a subsequent expansion. The initial and final spacetimes are flat and satisfy vacuum Einstein's equations, but not the full space, with most appreciable deviations from Einstein's equations at the bouncing time, $t = t_* = K^{1/4}$.

One can bring this metric in terms of cosmic time, where the 00 of metric component of $g_{\mu\nu}$ is normalized to 1. One has to notice however that we cannot do the transformation of coordinates only on one of the three metrics we have discussed $g_{\mu\nu}^2, g_{\mu\nu}^1$ and $g_{\mu\nu}$, and if we bring the the 00 metric component of the metric $g_{\mu\nu}$ is normalized to 1, it will not happen simultaneously for $g_{\mu\nu}^2, g_{\mu\nu}^1$. Having this in mind, the cosmic time coordinate T where where the 00 of metric component of $g_{\mu\nu}$ is normalized to 1 is defined by

$$dT = \sqrt{\frac{1 + \frac{K}{t^4}}{T_1 - T_2}} dt \quad (48)$$

So, we see that as $t \rightarrow \infty$, $T \rightarrow c_1 t$, while for $t \rightarrow 0$, $T \rightarrow -c_2/t$, here c_1, c_2 are positive constants. So at large negative cosmic time we have a contracting Milne space, a bounce and the evolution towards an expanding Milne space at large positive cosmic time.

5.5 The case where positive and negative string tensions are separated by a spontaneously generated wall in Wesson warped spaces

One may wonder if there are similar solutions to the vacuum Einstein's equations similar to the Milne space but where instead of time some spacial coordinate would play a similar way. The answer to this question is yes, and these are the solutions in higher dimensional vacuum General Relativity discovered by Wesson and collaborators, see [29] and references there. In five dimensions for example the following warped solution

is found,

$$ds^2 = l^2 dt^2 - l^2 \cosh^2 t \left(\frac{dr^2}{1-r^2} + r^2 d\Omega_2^2 \right) - dl^2 \quad (49)$$

where l is the fourth dimension, so we see that as in the fourth dimension l such a solution is homogeneous of degree two, just as the Milne space time was homogeneous of degree two with respect to the time. Notice that maximally symmetric de Sitter space times sub spaces $l = \text{constant}$ appear for instead of euclidean spheres that appear in the Milne Universe for $t = \text{constant}$.

The list of space times of this type is quite large, for example, one can find solutions of empty GR with Schwarzschild de Sitter subspaces for $l = \text{constant}$, as in

$$ds^2 = \frac{\Lambda l^2}{3} \left(dt^2 \left(1 - \frac{2M}{r} - \frac{\Lambda r^2}{3} \right) - \frac{dr^2}{1 - \frac{2M}{r} - \frac{\Lambda r^2}{3}} - r^2 d\Omega_2^2 \right) - dl^2 \quad (50)$$

This of course can be extended to D dimensions, where we choose one dimension l to have a factor l^2 warp factor for the other dimensions, generically for D dimensions as in

$$ds_2^2 = l^2 \bar{g}_{\mu\nu}(x) dx^\mu dx^\nu - dl^2 \quad (51)$$

where $\bar{g}_{\mu\nu}(x)$ is a $D-1$ Schwarzschild de Sitter metric for example [29]. This we will take as our 2 metric,

In any case, working with this generic metric of the form (51), but now in D dimensions, we can perform the inversion transformation $l \rightarrow \frac{1}{l}$, and multiplying also by a factor σ and obtain the conformally transformed metric 1 that also satisfies the vacuum Einstein's equations

$$ds_1^2 = \sigma l^{-2} \bar{g}_{\mu\nu}(x) dx^\mu dx^\nu - \sigma \frac{dl^2}{l^4} = \sigma l^{-4} ds_2^2 \quad (52)$$

From this point on, the equations the solutions for the tensions of the 1 and 2 strings are the same as in the cosmological case, just that $t \rightarrow l$, so now $c = \sigma l^{-4}$, so that we now insert this expression for c in (42) and in (43), obtaining that on one direction in l negative string tensions dominate, while in the other direction positive string tensions dominate, and we still take $\sigma = -K$, where K is positive.

The universal metric, following the steps done for the cosmological case is now,

$$ds^2 = \left(\frac{1-c}{T_2 - T_1} \right) (l^2 \bar{g}_{\mu\nu}(x) dx^\mu dx^\nu - dl^2) \quad (53)$$

looking at the coefficient of $\bar{g}_{\mu\nu}(x) dx^\mu dx^\nu$, the function is $(\frac{l^2 + K/l^2}{T_2 - T_1})$, so the space time is expanded or contracted as we move in the dimension l by this factor. This factor is minimized at $l_* = K^{1/4}$. We can define a proper length coordinate L where where the ll of metric component of the metric is normalized to -1 is defined by

$$dL = \sqrt{\frac{1 + \frac{K}{l^4}}{T_1 - T_2}} dl \quad (54)$$

So, we see that as $l \rightarrow \infty$, $L \rightarrow c_1 l$, while for $l \rightarrow 0$, $L \rightarrow -c_2/l$, here c_1, c_2 are positive constants.

Finally we can compare the resulting gravity theories resulting from these multi string effects with known gravity theories discussed in the literature. We have found

in the examples discussed so far, the need to define of two conformally related metrics. Since these two metrics are conformally related, it is not exactly correct to say that we have generated two independent metrics, more accurate is to say that there is a metric say, take the metric $g_{\mu\nu}^2$, and that there is a measure independent of that metric, $\sqrt{-\det(g_{\mu\nu}^1)}$, so what most resembles this structure are the modified measure theories of gravity [2, 3, 5, 4, 6, 7, 8, 9], where there is a metric and a measure that is independent a priori of that metric.

6 General Discussion of the Perspectives on Dynamically Generated String and Brane Tension Theories for Cosmology, Warped Space times and other possible applications

The consideration of dynamical tension generation lead us to theories where each extended object, either strings or branes generates dynamically its own string tension. This has profound consequences. We in particular discuss a new background scalar field, the "tension" field that can change dynamically the value of the tension along the world sheet of the extended objects. This new field introduced in the theory of extended objects appears to have very important consequences in various cosmological and other scenarios.

We have studied in particular how from the integration of the world sheet gauge fields, that control the dynamics of the dynamical tension, that the tension in a given string labeled by i is given by $e\phi + T_i$, where T_i is a constant of integration that may differ for different extended objects.

Then we consider the constraint from the quantum conformal invariance in a situation where at least two strings with constants of integration T_1 and T_2 , with $T_1 \neq T_2$ are considered and both strings probe the same region of spacetime. For simplicity we consider cases where only the background metric is non trivial, while the dilaton is taken as a constant and the anti symmetric tensor field is zero or pure gauge.

We construct actions that reproduce the equations with tensions $e\phi + T_1$ and $e\phi + T_2$, then the requirement of quantum conformal invariance, independently for the string 1 or the string 2 are that the auxiliary metrics $g_{\mu\nu}^1 = (e\phi + T_1)g_{\mu\nu}$ and $g_{\mu\nu}^2 = (e\phi + T_2)g_{\mu\nu}$ both satisfy the vacuum Einstein's equations. The metric $g_{\mu\nu}$ we call the Universal metric. These three metrics, $g_{\mu\nu}^1, g_{\mu\nu}$ and $g_{\mu\nu}^2$ are conformally related, the conformal relation between $g_{\mu\nu}^1 = (e\phi + T_1)g_{\mu\nu}$ and $g_{\mu\nu}^2 = (e\phi + T_2)g_{\mu\nu}$, implies $g_{\mu\nu}^1 = c(e\phi + T_2)g_{\mu\nu}$ where $c \neq 1$ if $T_1 \neq T_2$. c could be a constant. There are many ways to satisfy this, for example scaling a Schwarzschild metric gives us again a Schwarzschild metric, scaling a Kasner solution gives us again a Kasner solution. Notice that c is physical, since any coordinate transformation has to be done on the two metrics, but then the ratio of the densities $\frac{\sqrt{-g^1}}{\sqrt{-g^2}} = c^{D/2}$, is a scalar, showing that c is a coordinate invariant. In this case when expressing the equations of motion of the two strings with the same coordinates, if c is a constant, they both satisfy the same geodesic equation (the factor c comes out of the action and does not affect the geodesic equation).

We have obtained the case c not a constant in two types of situations, one consid-

ering the Milne Universe in D dimensions for the 2 metric and for the 1 metric, the one that is obtained by the transformation $t \rightarrow 1/t$ and then multiplying by a constant σ , so $c = \sigma/t^4$, This allows us to solve for the tensions of the two types of strings. At the early Universe we have negative string tensions, which then go to zero tension and at the same time the positive string tensions appear and those positive string tensions become constant in the late universe. The universal metric is non singular and experiences a bounce. Negative and positive string tensions appear separated by time.

Considering warped Universes of the Wesson type in D dimensions for the 2 metric and for the 1 metric, the one that is obtained by the transformation $l \rightarrow 1/l$ and then multiplying by a constant σ , so $c = \sigma/l^4$, This allows us to solve for the tensions of the two types of strings. On one direction of the warp coordinate we have negative string tensions, which then go to zero tension and at the same time the positive string tensions appear in the other direction of the warp coordinate and those positive string tensions become constant tension as get deep far in that direction. The universal metric is non singular and experiences a wormhole behavior. Negative and positive string tensions appear separated by spacial direction.

While the 1 and 2 metrics are the metrics with which the strings 1 and 2 follow geodesics (which are the same equations if the conformal factor between these two metrics is a constant) and for $g_{\mu\nu}$, neither strings 1 or 2 follow geodesics with respect to this metric, unless the conformal factor between metrics 1 and 2 is a constant, but a bound state of strings 1 and 2 may follow geodesics with respect to this metric. Whether such bound states exist and whether such conjecture is true is an open question at this point. The universal metric $g_{\mu\nu}$ does not satisfy Einstein's equations.

Notice also that the warped Wesson type space times are cosmological spacetimes, since the warp de Sitter or Schwarzschild de Sitter spaces. So they may provide a different perspective on the cosmological constant problem, as Wesson suggested.

One very interesting problem could be to consider the effect of the warp of the spacetime, as it has been done in more conventional scenarios in [30]

Although our main emphasis here has been on strings, many formulas have been given for the dynamical tension brane theory as well, but then quantum considerations for the case of branes are not so reliable, nevertheless some progress in this direction should be explored also.

Concerning carrying out the program carried out here, but using alternative formulations of the dynamical tension strings, like the formulation by Townsend and collaborators [17], [18], this may be possible, but preliminary calculations indicate that the analysis seems considerable more complicated.

We have in this paper started the study of the effective gravity theory generated by these dynamical string theories with many strings probing the same region of space. The analysis here has been concentrated on simple special cases, clearly a more generic approach is necessary, in particular one should study solutions with non trivial dilaton and antisymmetric tensor fields, as well as cosmological and warped solutions with all kind of matter.

Comparing the resulting gravity theories resulting from these multi string effects with known gravity theories discussed in the literature. We have found, in the examples discussed so far, the need to define of two conformally related metrics. Since these two are metrics are conformally related, it is not exactly correct to say that we have generated two independent metrics, more accurate is to say that there is a metric say, take

the metric $g_{\mu\nu}^2$, and that there is a measure independent of that metric, $\sqrt{-\det(g_{\mu\nu}^1)}$, so what most resembles this structure are the modified measure theories of gravity [2, 3, 5, 4, 6, 7, 8, 9], where there is a metric and a measure that is independent a priori of that metric. The tension of the strings enters not only in the effective gravity the string theory generates, but also on the spectrum of the strings themselves, these aspects have to be further investigated.

We can see also that in the case that both the metric 1 and the metric 2 are two flat metrics, like in our studies of cosmological solutions, or for the warped spaces in the case the warped space 51 is defined with $\bar{g}_{\mu\nu}(x)$ being de Sitter and not the more generic Schwarzschild de Sitter case, the solutions are most likely exact and not just to first order in the slope, since not only the Ricci tensor vanishes but also the Riemann curvature and so will all higher curvature corrections to the beta function. In other paper we will address other scenarios for cosmology and warped spaces where we will emphasize the effect of dynamical string tensions on the possibility of escaping the Hagedorn temperature, this could be possible if the string tensions become very large at the early Universe or at some position in the warped coordinate [31]. In this case in certain regions of space both type of string tensions are positive and go to infinity in such a way that their ratio is one and for equal tension strings the more conventional string interactions (besides the one governed by the tension field), have a chance to operate. Notice that a multi string tension scenario formulated phenomenologically in order to avoid the Hagedorn temperature was formulated by Oleg Andreev [32], which seems to be the only other place in the literature, besides our research, where a multi string tension scenario has been discussed.

Another subject that this paper can be related to is the conformal cyclic cosmology of Penrose and collaborators [33], where the role of conformal invariance is studied in the cyclic scenarios, which contain bounces that avoid the singularities. We do not have target space conformal invariance in the ordinary sense, but a kind of dynamical target space conformal invariance in the sense that there is the 1 metric that satisfies Einstein's equations and then we demand the existence of a second metric 2, conformally related to 1 that also satisfies Einstein's equations. This type of dynamical target space conformal invariance also appears to produce bounces, so that may be there is a point of contact with [33].

Finally, we notice that the tension field has been determined by the effects of more than one string probing a region of space time, which suggest a formal presentation of the dynamical tension theories in the context of string field theory could be very useful.

Acknowledgments I thank Oleg Andreev, David Andriot, Stefano Ansoldi, David Benisty, Emil Nissimov, Hitoshi Nishino, Svetlana Pacheva, Subhash Rajpoot, Euro Spallucci and Tatiana Vulfs for useful discussions. I also want to thank the Foundational Questions Institute (FQXi) and the COST actions Quantum Gravity Phenomenology in the multi messenger approach, CA18108 and Gravitational waves, Black Holes and Fundamental Physics, CA16104 for support.

References

- [1] "Superstrings", John H. Schwarz, Vols- 1 and 2, World Scientific, 1985. ; M. B. Green, J. H. Schwarz and E. Witten, Superstring Theory, Cambridge University

Press, 1987.

- [2] E.I. Guendelman, A.B. Kaganovich, Phys.Rev.D55:5970-5980 (1997)
- [3] E.I. Guendelman, Mod.Phys.Lett.A14, 1043-1052 (1999)
- [4] E.I. Guendelman, O. Katz, Class.Quant.Grav. 20 (2003) 1715-1728 • e-Print: gr-qc/0211095 [gr-qc]
- [5] Frank Gronwald, Uwe Muench, Alfredo Macias, Friedrich W. Hehl, Phys.Rev.D 58 (1998) 084021 • e-Print: gr-qc/9712063 [gr-qc]
- [6] Eduardo Guendelman, Ramón Herrera, Pedro Labrana, Emil Nissimov, Svetlana Pacheva, Gen.Rel.Grav. 47 (2015) 2, 10 • e-Print: 1408.5344 [gr-qc]
- [7] Eduardo Guendelman, Douglas Singleton, Nattapong Yongram, JCAP 11 (2012) 044 • e-Print: 1205.1056 [gr-qc]
- [8] R. Cordero, O.G. Miranda, M. Serrano-Crivelli, JCAP 07 (2019) 027 • e-Print: 1905.07352 [gr-qc]
- [9] Eduardo Guendelman, Emil Nissimov, Svetlana Pacheva, Eur.Phys.J.C 75 (2015) 10, 472 • e-Print: 1508.02008 [gr-qc]
- [10] E.I. Guendelman, Class.Quant.Grav. 17, 3673-3680 (2000)
- [11] E.I. Guendelman, A.B. Kaganovich, E.Nissimov, S. Pacheva, Phys.Rev.D66:046003 (2002)
- [12] E.I. Guendelman, Phys.Rev.D 63 (2001) 046006 • e-Print: hep-th/0006079 [hep-th]
- [13] Hitoshi Nishino, Subhash Rajpoot, Phys.Lett.B 736 (2014) 350-355 e-Print: 1411.3805 [hep-th].
- [14] T.O. Vulf, E.I. Guendelman, Annals Phys. 398 (2018) 138-145 • e-Print: 1709.01326 [hep-th]
- [15] T.O. Vulf, E.I. Guendelman, Int.J.Mod.Phys.A 34 (2019) 31, 1950204 • e-Print: 1802.06431 [hep-th]
- [16] T.O. Vulf, Ben Gurion University Ph.D Thesis, (2021), arXiv:2103.08979.
- [17] P.K. Townsend, Phys.Lett.B 277 (1992) 285-288.
- [18] E. Bergshoeff, L.A.J. London, P.K. Townsend, Class.Quant.Grav. 9 (1992) 2545-2556, Class. Quantum Grav. 9 (1992) 2545-2556 • e-Print: hep-th/9206026 [hep-th]
- [19] M. Henneaux, C. Teitelboim, Phys.Lett.B 222 (1989) 195-199
- [20] David Bensity, Eduardo I. Guendelman, Alexander Kaganovich, Emil Nissimov, Svetlana Pacheva, Eur.Phys.J.Plus 136 (2021) 1, 46 • e-Print: 2006.04063 [gr-qc]
- [21] Deser, S. and Zumino, Phys. Lett. B65 , 369, (1976)
- [22] Brink , L., Di Vecchia, P and Howe, S. ,Phys. Lett. B65, 471 , (1976)
- [23] Polyakov, A. M., ,Phys. Lett. B103 ,207, (1980).
- [24] For a review see, Ne'eman , Y and Eizenberg, E. Membranes and other Extendons, World Scientific, 1995.
- [25] S. Ansoldi, E. I. Guendelman, E. Spallucci, Mod.Phys.Lett.A 21 (2006) 2055-2065 • e-Print: hep-th/0510200 [hep-th]

- [26] A. Vilenkin, E.P. S. Shellard, Cosmic Strings and Other Topological Defects, Cambridge Monographs on Mathematical Physics, Cambridge University Press, 2000, 580 pages, ISBN: 9780521654760.
- [27] Joseph Polchinski, String Theory, vol. 1 , Cambridge University Press (1998). Some papers on strings with background fields are C. G. Callan, D. Friedan, E. J. Martinec and M. J. Perry, Nucl. Phys. B 262 (1985) 593; T. Banks, D. Nemeschansky and A. Sen, Nucl. Phys. B 277 (1986) 67
- [28] Particles and Sources, Julian Schwinger, Phys.Rev. 152 (1966) 1219-1226 DOI: 10.1103/PhysRev.152.1219
- [29] Paul Wesson, "Space -Time-Matter. World Scientific, (1999)"; Paul S. Wesson, Hongya Liu, Int.J.Mod.Phys. D10 (2001) 905-912, • e-Print: gr-qc/0104045.
- [30] Warp factor and the gravitational wave spectrum, David Andriot, Paul Marconnet, Dimitrios Tsimpis, arXiv:2103.09240 [hep-th]
- [31] E.I. Guendelman, Escaping the Hagedorn Temperature in Cosmology and Warped Spaces with Dynamical String Tension, e-Print: 2105.02279 [hep-th]
- [32] Oleg Andreev, JHEP 0903:098,2009, e.print: 0807.1017
- [33] See for example, V.G. Gurzadyan, R. Penrose, Eur.Phys.J.Plus 128 (2013) 22 • e-Print: 1302.5162 [astro-ph.CO]