

PATHLIKE CO/BIALGEBRAS AND THEIR ANTIPODES WITH APPLICATIONS TO BI- AND HOPF ALGEBRAS APPEARING IN TOPOLOGY, NUMBER THEORY AND PHYSICS.

RALPH M. KAUFMANN AND YANG MO

Dedicated to Professor Dirk Kreimer on the occasion of his 60th birthday

ABSTRACT. We develop an algebraic theory of flavored and pathlike co-, bi- and Hopf algebras. This is the right framework in which to discuss antipodes for bialgebras naturally appearing in topology, number theory and physics. In particular, we can precisely give conditions the invertibility of characters needed for renormalization in the formulation of Connes and Kreimer and in the canonical examples these conditions are met. To construct the antipodes directly, we discuss formal localization constructions and quantum deformations. These allow to construct and naturally explain the appearance Brown style coactions. Using previous results, we can tie all the relevant coalgebras to categorical constructions and the bialgebra structures to Feynman categories.

INTRODUCTION

Although the use of Hopf algebras has a long history, the seminal paper [CK98] led to a turbocharged development for their use which has penetrated deeply into mathematical physics, number theory and also topology, their original realm—see [AFS08] for the early beginnings. The important realization in [CK98, CK00, CK01] was that the renormalization procedure for quantum field theory can be based on a character theory for Hopf algebras via a so-called Birkhoff factorization and convolution inverses. The relevant Hopf algebras are those of trees with a coproduct given by a sum over so-called admissible cuts—with the factors of the coproduct being the left over stump and the collection of cut off branches—and Hopf algebras of graphs in which the factors of the summands of the coproduct are given by a subgraph and the quotient graph. A planar version of the tree formalism was pioneered in [Foi02]. The appearance in number theory of this type of coproduct goes back to [Gon05] and was developed further and applied with great success in [Bro15]. In all these cases, the product structure is free. A precursor to these considerations can be identified in [Bau81].

2010 *Mathematics Subject Classification.* 18A30, 18A40.

Key words and phrases. Feynman Category, bialgebra, Hopf algebra.

In [GCKT20a, GCKT20b], we gave the details to prove the realization announced in [KW17] that all these structures stem from a categorical framework where the coproduct corresponds to deconcatenation of morphisms. Such coproduct structures can be traced back to [Ler75] and appear in combinatorics [JR79]. In [KW17], we developed a theory of so-called Feynman categories, which are essentially a special type of monoidal category and could connect the product structure to the monoidal structure on the level of morphisms. Special cases are related to operads, the simplicial category and categories whose morphisms are classified by graphs. In [GCKT20b], we could show that the monoidal product in these categories is compatible with the deconcatenation co-product thus yielding bialgebras. This generalizes the Connes–Kreimer tree examples and those of Baues and Goncharov, which have operadic origin [GCKT20a] and comprises the Connes–Kreimer graph examples. The co- and bialgebras which have a categorical interpretation also include path coalgebras and incidence coalgebras, see §2. There are two versions of the story, one is symmetric and yields cocommutative bialgebras and the second is non-symmetric and in general yields to non-cocommutative bialgebras, such as those from planar structures.

In this paper, we address the question of constructing antipodes in order to make the bialgebras into Hopf algebras. In [GCKT20a, GCKT20b] this was done by taking a quotient which made the bialgebras connected. This quotient is taken implicitly in the examples discussed above, see [GCKT20a, §1]. The questions that we can now answer is: What structures precisely exist already on the bialgebra level, and what is their role?

To this end, we give an *algebraic characterization* of the bi-algebras in terms of what we call *flavored* and *pathlike* coalgebras, which capture and generalize the essential features of categorical coalgebras, especially path algebras for quivers. This allows us effective criteria for the existence of antipodes and convolution invertibility of characters. We also work over an arbitrary ground ring. The origins for the theory of flavored coalgebras can be traced back to [TW74] and with hindsight, several of the quotient construction are foreshadowed in [Sch87] for the special case of incidence bialgebras.¹ The existence of antipodes is usually proven via some sort of connectivity of a filtration [Tak71, Qui67]. The problem is that the bialgebras are usually not connected for the standard filtrations, which is why quotients are taken.

Concretely, after reviewing the basic structures the generality we need in §1, and providing an extended list of examples, to illustrate and motivate the constructions, in §2, we introduce the concept of a QT-filtration in §3, which generalizes the commonly used filtrations. This is what is needed to make Takeuchi’s argument work and results in:

Theorem 3.28 *Consider a coalgebra C with counit ϵ and an algebra A with unit η . Given a QT-sequence $\{F_i\}_{i=0}^\infty$ on C , such that $\text{Ext}_R^1(C/F_0C, A) =$*

¹We thank D. Kreimer for pointing this out

0, for any element f of $\text{Hom}(C, A)$, f is \star -invertible if and only if the restriction $f|_{F_0C}$ is \star -invertible in $\text{Hom}(F_0C, A)$.

A special type of QT-filtration, which we call the bivariate Quillen filtration (see §4), arises by considering skew primitive elements. These are the essential actors for so-called flavored coalgebras and are key to understanding the structures, before taking quotients. In the applications a slightly stronger condition is satisfied for the level 0 part of the filtration, which leads to the notion of pathlike co/bialgebras.

Theorem 4.16 *Let C be pathlike coalgebra and A algebra and $f \in \text{Hom}(C, A)$. If $\text{Ext}(C/\bar{F}_0C^Q, A) = 0$, then f has an \star -inverse in the convolution algebra $\text{Hom}(C, A)$ if and only if for every grouplike element g , $f(g)$ has an inverse as ring element in A . In particular every grouplike central character is \star -invertible.*

These results are then directly applicable to renormalization via Rota-Baxter algebras and Birkhoff decompositions, which are briefly reviewed in §5. A closer inspection of this framework yields two paths of actions. The first is to realize that the bialgebras need not necessarily be Hopf algebras, but only the characters need to be \star -invertible. Thus restricting the characters by mandating certain properties on them or the target algebra will ensure their invertibility in the pathlike case. The second is to formally invert the grouplike elements. Combining the two approaches yields several other universal constructions, through which the characters with special properties factor. These are natural construction spelled out in §5 which also naturally lead to *quantum deformations* of the algebras. This generalizes and explains similar constructions in [GCKT20a]. it also gives an astonishing connection to *Brown's coactions* [Bro15] and allows to generalize them to the full framework, see §4.4.1 and §5.2.2. With hindsight, the special role and treatment of $\zeta^m(2)$ is precisely the difference between the two approaches. Setting it to 0 implements the quotient and keeping is as an invertible central elements defines the q -deformation.

We then apply the general framework to the case of pathlike bialgebras arising from Feynman categories in §6, giving the details for the simplicial setup of Goncharov and Baues—including Joyal duality—the Connes–Kreimer tree bialgebras (with leaves, planar and non-planar), their generalization to operads and finally graphical Feynman categories that are related to Kreimer's core Hopf algebra. To be more self-contained, these are given in a new set-theoretical presentation in an appendix. We close with technical conditions on Feynman categories, which are met in all the examples and guarantee that the bialgebras associated to the Feynman categories are pathlike.

Theorem 6.15 *The bialgebra $B^{\text{iso}}(\mathcal{F})$, is pathlike if \mathcal{F} is generated by isomorphisms and a set of generators which are essentially skew primitive or essentially indecomposable. In case that there are homogenous relations, the coalgebra will be graded.*

For a non-symmetric Feynman category $B(\mathcal{F})$ is pathlike, and \mathcal{F} is generated by identities and skew primitives, that is indecomposables, which is equivalent to a set of generators which are skew primitive. In case that there are homogenous relations, the coalgebra will be graded. It is necessary that \mathcal{V} is discrete.

Here \mathcal{F} and \mathcal{V} are part of the data for a Feynman category, see §2.6.

Acknowledgments. This paper is the outgrowth of a long conversation with Dirk Kreimer whom we would like to congratulate and profoundly thank for his interest and support. Much of the impetus for this work has come through conversations with him and his group. We wish to thank the Humboldt University, the KMPB, and the Humboldt Foundation for making these visits possible.

We furthermore would like to thank F. Brown, K. Yeats and C. Berger for helpful discussions and the organizers of the conference *Algebraic Structures in Perturbative Quantum Field Theory* at the IHES, in November 2020. The wonderful interactions that took place there led to part of the results of this paper.

1. PRELIMINARIES ON HOPF ALGEBRA

We collect the relevant results on Hopf algebras to fix a common notation and nomenclature — see e.g, [Car07, MP12] . Throughout this paper, we work over a commutative unital ground ring R (and k be a field). Tensor are understood to be over the ground ring, thus \otimes means \otimes_R . For a R -module M there is a canonical isomorphism $M \otimes R \simeq M$ given by the R -module structure on M and we will simply implement this isomorphism as an identification—effectively making R a strict monoidal unit.

We will either work in the category of R -modules or graded R -modules. In the second case, we assume R is concentrated in degree 0 and the grading by the semi-group \mathbb{N} unless otherwise stated.

1.1. Basic definitions and structures.

Definition 1.1. An associative and unital algebra (graded algebra) A is an R -module (R -graded module) equipped with two R -linear maps (k -homogeneous maps) multiplication denoted by m and unit denoted by η such that the following diagrams commute on the nose, where Id_A means the identity map on A :

$$\begin{array}{ccc} A \otimes A \otimes A & \xrightarrow{m \otimes \text{Id}_A} & A \otimes A \\ \downarrow \text{Id}_A \otimes m & & \downarrow m \\ A \otimes A & \xrightarrow{m} & A \end{array} \quad \text{and} \quad \begin{array}{ccccc} A \otimes R & \xrightarrow{\text{Id}_A \otimes \eta} & A \otimes A & \xleftarrow{\eta \otimes \text{Id}_A} & R \otimes A \\ & \searrow & \downarrow m & \swarrow & \\ & & A & & \end{array}$$

Let τ be an endomorphism on $A \otimes A$ that permutes entries $\tau(x \otimes y) = (y \otimes x)$. A is commutative if the following diagram commutes.

$$\begin{array}{ccc} A \otimes A & \xrightarrow{\tau} & A \otimes A \\ & \searrow m & \downarrow m \\ & & A \end{array}$$

Remark 1.2. If A is a R -algebra, then $A \otimes A$ is also naturally a R -algebra with the identity $1_{A \otimes A} = 1_A \otimes 1_A \cong 1_A$ and the multiplication $m_{A \otimes A} = (m_A \otimes m_A) \circ \tau$ (This simply means $a \otimes b \cdot c \otimes d = ac \otimes bd$).

Definition 1.3. A coassociative and counital *coalgebra* (*graded coalgebra*) C is an R -module (graded R -module) with two R -linear maps (homogeneous k -linear maps): Δ , the comultiplication, and ϵ the counit, such that the following diagrams commute on the nose, where Id_C means the identity map on C :

$$\begin{array}{ccc} C & \xrightarrow{\Delta \otimes \text{Id}_A} & C \otimes C \\ \downarrow \text{Id}_A \otimes \Delta & & \downarrow \Delta \\ C \otimes C & \xrightarrow{\Delta} & C \otimes C \otimes C \end{array} \quad \text{and} \quad \begin{array}{ccccc} C \otimes R & \xleftarrow{\text{Id}_A \otimes \epsilon} & C \otimes C & \xrightarrow{\epsilon \otimes \text{Id}_C} & R \otimes C \\ & \searrow & \uparrow \Delta & \swarrow & \\ & & C & & \end{array}$$

Moreover, let τ be an endomorphism on $C \otimes C$ such that it permutes entries $\tau(x \otimes y) = (y \otimes x)$. Then we say A is cocommutative if the following diagram commute.

$$\begin{array}{ccc} C \otimes C & \xrightarrow{\tau} & C \otimes C \\ \uparrow \Delta & \nearrow \Delta & \\ C & & \end{array}$$

Note that due to the coassociativity, there is a unique morphism $\Delta^{[n]} : C \rightarrow C^{\otimes n}$ which is given by any iteration of diagonals.

Definition 1.4.

- (1) A *coaugmented* coalgebra is a coalgebra equipped with the unit map $\eta : R \rightarrow C$ as coalgebra morphism. We will write $e = \eta(1_R)$.
- (2) A *augmented* coalgebra is a coalgebra equipped with the counit map $\eta : C \rightarrow R$ as algebra morphism.
- (3) A graded coalgebra is said to be *connected* if the degree zero part is isomorphic to R .

Remark 1.5. In a coaugmented coalgebra, we have that $\epsilon \circ \eta = \text{id}_R$ as η preserves the counit. Setting $e = \eta(1)$, then $\Delta(e) = e \otimes e$ and there is a splitting $C = C_0 \oplus \overline{C}$ where $C_0 = Re$ and $\overline{C} = \ker(\epsilon)$.

In this case, one also regards the reduced diagonal $\overline{\Delta}$ defined by $\overline{\Delta}(x) = \Delta(x) - e \otimes x - x \otimes e$.

Definition 1.6. A *bialgebra* (*graded bialgebra*) B is both an algebra (graded algebra) (B, m, η) and coalgebra (graded coalgebra) (B, Δ, ϵ) over R s. t. counit and comultiplication map are algebra/graded algebra morphisms.

Definition 1.7. A Hopf algebra H is a bialgebra (graded bialgebra) with an R -linear (R -homogeneous) map S called antipode (also sometimes called coinverse) s. t.

$$\eta \circ \epsilon = m \circ (S \otimes \text{Id}) \circ \Delta = m \circ (\text{Id} \otimes S) \circ \Delta$$

or equivalently the following diagram commutes

$$\begin{array}{ccccc}
 & H \otimes H & \xrightarrow{S \otimes \text{Id}} & H \otimes H & \\
 & \uparrow \Delta & & \downarrow m & \\
 H & \xrightarrow{\epsilon} & R & \xrightarrow{\eta} & H \\
 & \downarrow \Delta & & \uparrow m & \\
 & H \otimes H & \xrightarrow{\text{Id} \otimes S} & H \otimes H &
 \end{array}$$

Alternatively, S is characterized by being the convolution inverse of the identity map, see below. As such an antipode is unique, if it exists. An antipode is an antihomomorphism $S(ab) = S(b)S(a)$.

NB: An antipode is not always bijective, see [Tak71].

Example 1.8. By requiring R to be unital, R turns out to be a coalgebra over R itself, by setting $\Delta(1) = 1 \otimes 1$ and $\epsilon(1) = 1$. In fact, it is a Hopf algebra over itself in a unique way with $S(1) = 1$.

Definition 1.9. Given a coalgebra C , a element g in C is said to be *semigrouplike* if $\Delta(g) = g \otimes g$ and of additionally $\epsilon(g) = 1$, then g is said to be *grouplike*. The set of semigrouplike elements in C is denoted by $sgpl(C)$ and the set of grouplike elements in C by $gpl(C)$.

Remark 1.10.

- (1) If the coalgebra has a counit ϵ , then for $g \in sgpl(G)$ $(1 - \epsilon(g))g = 0$ that is $1 - \epsilon(g) \in \text{Ann}(g)$.
- (2) In a bialgebra the semigrouplike elements form a monoid as $\Delta(gh) = gh \otimes gh$ for two semigrouplike elements g, h . The grouplike elements form a submonoid: $\epsilon(gh) = \epsilon(g)\epsilon(h) = 1$. These structures are unital with unit $e = \eta(1)$.
- (3) In a Hopf algebra for any semigrouplike element g : $S(g)g = gS(g) = \epsilon(g)e$. In particular if the element is grouplike, $S(g) = g^{-1}$.
- (4) In a graded coalgebra all grouplike elements are necessarily in degree 0 as $2i = i$ implies $i = 0$.
- (5) In a connected graded coalgebra, $e = \eta(1)$, which is the generator of C_0 , is the unique grouplike element.

Definition 1.11. Let g and h be semigrouplike elements in C , an element x in C is called (g, h) -skew primitive if $\Delta(x) = x \otimes g + h \otimes x$. And when $g = h$,

it is called *g-primitive element*. If g, h are only assumed to be semigrouplike, then we obtain the notion of semiskew primitive.

1.2. Modules/comodules relative tensor products.

Definition 1.12. A left/right module for an R -algebra A is an R -module M and a R -module map $\rho : A \otimes M \rightarrow M, a \otimes m \rightarrow am$ respectively $\lambda : M \otimes A \rightarrow M : m \otimes a \rightarrow ma$. These have to satisfy the condition $(ab)m = a(bm)$ respectively $m(ab) = (ma)b$.

Dually a left/right comodule for a coalgebra C is an R -module M with maps $\rho : M \rightarrow M \otimes C$ respectively $\lambda : M \rightarrow C \otimes M$. Satisfying $\rho(\rho(m)) = (id \otimes \Delta)(m)$ respectively $\lambda(\lambda(m)) = (\Delta \otimes id)\lambda(m)$.

For a right A module M and a left A module N there is the relative tensor product $M \otimes_A N$ which is the quotient of $M \otimes_R N$ by the relation $m \otimes am = ma \otimes n$.

The dual notion is the where the in general cotensor product of a right comodule M and a left comodule N . This is the subset of $M \otimes_R N$ for which $a \otimes b \in C \otimes C$ such that $\rho(a) = \lambda(b)$.

More formally it is the coequalizer:

$$M_{\lambda \square \rho} N \longrightarrow M \otimes_R N \xrightarrow[id_M \otimes_R \rho_L]{\rho_R \otimes_R id_N} M \otimes_R C \otimes_R N \quad (1.1)$$

1.3. Duals. The dual of a coalgebra C is the convolution algebra $Hom(C, R)$ which is unital if C has a counit and R has a unit. The dual of a *finite* dimensional algebra $A^* = Hom(A, R)$ is a coalgebra with the multiplication.

$$\Delta(f)(a \otimes b) := f(ab) \quad (1.2)$$

In general, this is a morphism $A^* \rightarrow (A \otimes A)^*$ and there is an injection $A^* \otimes A^* \rightarrow (A \otimes A)^*$, which is an isomorphism in the case of a finite dimensional algebra. One can look at the graded version and the graded dual in the case of a graded algebra with finite dimensional pieces. One can also restrict to the so-called finite dual A° which is the maximal subspace of A^* such that $\Delta(A^\circ) \subset A^\circ \otimes A^\circ$. For a field, this is:

$$A^\circ = \{f \in A^* | Ker(f) \text{ contains an ideal of finite codimension}\} \quad (1.3)$$

The Hopf-dual of a Hopf algebra over a field is defined to be H° with the induced structures.

2. EXAMPLES

2.1. Set-based examples. Given any set S , we can associate a coalgebra $R[S]$, where every element is linear formal sum of S and every element in S is grouplike. This is what is called the set-like coalgebra given by the set S .

If S is a (unital) monoid then $R[S]$ is a (unital) algebra by R -linear extension of the monoid structure.

It is easy to check for a monoid M that $R[M]$ is a bialgebra, this is Hopf, with the antipode $S(g) = g^{-1}$, precisely when M is a group.

2.2. Path/Quiver coalgebra. A quiver Q is a graph with an orientation of each edge. A path on a graph is a sequence of oriented edges. An oriented edge \vec{e} has a source and a target vertex $s(\vec{e})$ and $t(\vec{e})$. A sequence of directed edges $p = (\vec{e}_1, \dots, \vec{e}_n)$ is a path if $t(\vec{e}_i) = s(\vec{e}_{i+1})$. The start of p is $v = s(e_1)$ and the end of p is $w = t(e_n)$ and we say p is a path from v to w . The length of p is n . By definition for each vertex v there is an empty path, i.e. a path of length 0 at v , which we denote by v .

Let $P(Q)$ be the set of paths of Q , then $R[P(Q)]$ has the coalgebra structure

$$\Delta(\vec{e}_1, \dots, \vec{e}_n) = s(\vec{e}_1) \otimes (\vec{e}_1, \dots, \vec{e}_n) + \sum_{i=1}^{n-1} (\vec{e}_1, \dots, \vec{e}_i) \otimes (\vec{e}_{i+1}, \dots, \vec{e}_n) + (\vec{e}_1, \dots, \vec{e}_n) \otimes t(e_n) \quad (2.1)$$

and $\Delta(v) = v \otimes v$. It has counit $\epsilon(v) = \epsilon(\vec{e}_1, \dots, \vec{e}_n) = 0$.

The grouplike elements are hence exactly the length 0 paths v and the paths of length 1 are skew primitive with (\vec{e}) being $(s(\vec{e}), t(\vec{e}))$ -skew primitive.

2.3. Colored monoid/algebra and dual coalgebras. Paths actually form a partial algebra structure by concatenation. The partial structure dictates that the end of the first path is the start of the second path. This is a particular example of a colored algebra.

Definition 2.1. A colored monoid A is a set together with the following data. A set X of colors, two morphisms $s, t : A \rightarrow X$, and a partial product

$$\circ : A_s \times_t A \rightarrow A \quad (2.2)$$

which is associative in the obvious sense. If $(ab)c$ exists, then so does $a(bc)$ and they coincide. It is decomposition or locally finite, if all the fibers $\circ^{-1}(a)$ for $a \in A$ are finite. It is unital if there is morphism $id : X \rightarrow A$, which is a section of both s and t such that $a \circ id(t(a)) = id(s(a)) \circ a = a$.

Note if $X = \{*\}$ this is simply a (unital) monoid structure.

The free R -module on A , i.e. $R[A] = \bigoplus_{a \in A} Ra$ is an R -algebra with the product $a \cdot b = \delta_{t(a), s(b)} ab$.

Consider $A^* = Hom(A, R)$ which is naturally an R -module by $(rf)(a) = r(f(a))$. Consider the characteristic functions $\delta_a \in Hom(A, R)$, $a \in A$, defined by $\delta_a(b) = \delta_{a,b}$. One can identify $R[A]$, with the sub-module of A^* of finite sums by sending $a \rightarrow \delta_a$. Notice that by definition $Hom(A, R) = Hom_R(R[A], R)$.

Proposition 2.2. *The free R -module of a colored locally finite monoid is a coalgebra, with coproduct (1.2) for $f \in Hom_R(A, R)$. If the colored monoid is unital then the dual coalgebra is counital with counit for which $\epsilon(\delta_{id(X)}) = 1$ and 0 else.*

Proof. Then $\Delta(\delta_a) = \sum_{(a_1, a_2): a_1 a_2 = a} \delta_{a_1} \otimes \delta_{a_2}$ which as a finite sum, the index set $\circ^{-1}(a)$ is finite, is in $A^* \otimes A^*$. The counit property follows from a direct computation. \square

Example 2.3 (Path coalgebra of a Quiver). The path coalgebra is a special case. It is the free colored algebra whose set of colors is the set of vertices and whose morphism are freely generated by the directed edges with the source and target maps mapping an directed edge to its source vertex and its target vertex.

Remark 2.4. In the coalgebra, the $\delta_{id(a)}$ are the grouplike elements and if a is indecomposable, δ_a is $(\delta_{id(s(a))}, \delta_{id(t(a))})$ skew primitive. This gives a shadow of the colored original colored structure in the coalgebra. A more general way to “see” the coloring is via cotensor products, see §4. The generalization that only takes the intrinsic notion of skew primitives, viz. the shadow, into account is the notion of a flavored coalgebra of §4.

2.3.1. Colored coalgebras. To recover the type of algebra obtained from a category, we introduce the notion of a colored coalgebra.

Definition 2.5. Let X be a set and consider the setlike coalgebra $C(X) = R[X]$. A colored colgebra C with colors X is a coalgebra C together with a bi-comodule structure of C over $C[X]$, that is $\lambda : C \rightarrow C[X] \otimes C, \rho : C \rightarrow C[X] \otimes C$ such that the coalgebra map factors through the cotensor product.

$$\begin{array}{ccc} C & \xrightarrow{\quad} & C_{\lambda \square \rho} C \\ & \searrow \Delta & \downarrow \\ & & C \otimes C \end{array} \quad (2.3)$$

The colored coalgebra is called simply colored if $C = \bigoplus_{(x,y) \in X \times X} C_{x,y}$ where for all $c \in C_{x,y} : \lambda(c) \subset R[X] \otimes C \subset C[X] \otimes C$ and $\rho(c) = c \otimes R[y]$.

In a simply colored coalgebra: $\Delta(C_{x,y}) \subset \bigoplus_z C_{x,z} \otimes C_{z,y}$.

Example 2.6. The coalgebra of colored algebra is a simply colored coalgebra with the same colors. The coaction has another description, which is as follows: The maps s, t induce a pull-back maps $s^*, t^* : R[A] \rightarrow R[X]$ which are split by the map induced by id^* the coactions are then given by $\lambda = (s^* \otimes id) \circ \Delta$ and $\rho = (id \otimes t^*) \circ \Delta$. In particular $\lambda(\delta_a) = \delta_{s(a)} \otimes \delta_a, \rho(\delta_a) = \delta_a \otimes \delta_{t(a)}$. Together with $id^* : R[X] \rightarrow R[A]$ they also provides the splitting.

2.4. Categories as colored algebras and categorical path (co)algebras.

Categories are an equivalent way of viewing colored coalgebras. The bijection between these two notions is given as follows: This is almost tautological if one uses the definition of a (small) category given by two sets $Mor(C)$

and $Obj(\mathcal{C})$ together with maps, see e.g. [GM03].

$$\begin{array}{ccc} N_1(\mathcal{C}) = Mor(\mathcal{C}) & & \\ \downarrow s \uparrow id \downarrow t & \circ : Mor_s \times_t Mor \rightarrow Mor & (2.4) \\ N_0(\mathcal{C}) = Obj(\mathcal{C}) & & \end{array}$$

where \circ is associative, id and provides the identity maps.

In this correspondence, the colors are the objects and the algebra elements are the morphisms. A path is now a composable sequence of morphisms.

Given a locally finite category, that is it is locally finite as a colored algebra, we obtain a coproduct on the free algebra of morphisms.

$$\Delta(\phi) = \sum_{(\phi_0, \phi_1): \phi_1 \circ \phi_0 = \phi} \phi_0 \otimes \phi_1 \quad (2.5)$$

which was considered in [KW17, GCKT20a, GCKT20b], and goes back to [Ler75] cf. also [JR79].

Remark 2.7. Note that there are two ways to write down the composition maps.

$$\begin{array}{ccc} Hom(X, Y) \times Hom(Y, Z) & \rightarrow & Hom(X, Z) \\ (\phi_0, \phi_1) & \mapsto & \phi_1 \circ \phi_0 \\ Hom(Z, Y) \times Hom(X, Y) & \rightarrow & Hom(X, Z) \\ (\phi_1, \phi_0) & \mapsto & \phi_1 \circ \phi_0 \end{array} \quad (2.6)$$

which correspond to the two ways to write the compositions

$$Mor_s \times_t Mor \rightarrow Mor \text{ or } Mor_t \times_s Mor \rightarrow Mor \quad (2.7)$$

The two coproducts are opposites Δ and $\Delta^{op} = \tau \circ \Delta$ where τ is the flip. In the example below, the first version of this for the coproduct, is what is used for posets and quivers, fits with the Connes–Kreimer coproduct [CK98, GCKT20a, GCKT20b], where subgraph is on the left and the cograph is on the right, while the second one is more natural from a category point of view and corresponds to the tree Hopf algebra where the stump is on the left and the branches on the right.

Note that this ambiguity is non-essential for a coalgebra or a bialgebra. For a Hopf algebra, it *a priori* might make a difference, see Remark 4.18 but if the bialgebra is pathlike it does not, see Corollary 4.19.

2.4.1. Path (co)algebra for a quiver or graph. In the case of the path (co)algebra, the category is the category generated by the quiver. That is the (co)algebra for the category $\mathcal{C}(Q)$ freely generated by the arrows. The objects of the category are the vertices. There is a generating morphism for each directed edge. A path is the composition of the composable morphisms.

The quiver for a set-like coalgebra is given by a rose or tadpole graph whose set of loops is S . One can fix any orientation for a loop to make it a quiver.

More generally, for a graph without direction of edges one chooses one generator for each of the two orientations for the edge and sets $\vec{e} \circ \bar{e} = id_v$ where v is the source of \vec{e} and hence the target of the edge with opposite orientation obtaining a groupoid, see below.

2.4.2. Incidence coalgebras. Another example of a categorical coalgebra is a so-called incidence coalgebra [JR79, Sch87]. The main observation is that a poset $(S, <)$ defines a category. whose objects are S and whose morphisms are as follows. There is one exactly one morphism $\phi_{x \rightarrow y}$ between x and y iff $x \leq y$. The category is locally finite when the poset is. The coproduct is then given as follows.

$$\Delta(\phi_{x \rightarrow y}) = \sum_{z \in [x, y]} \phi_{x \rightarrow y} \otimes \phi_{y \rightarrow z} \quad (2.8)$$

or in the usual poset notation where one identifies $\phi_{x \rightarrow y}$ with the interval $[x, y]$ and $id_x = [x, x]$, see e.g. [JR79]:

$$\Delta([x, y]) = \sum_{z \in [x, y]} [x, z] \otimes [z, y] \quad (2.9)$$

This is a special case of a quotient of a quiver coalgebra. The quiver is given by the x, y primitives. These are the elements $x < y$ where there is no $z : x < z < y$ as then $\Delta(\phi_{x \rightarrow y}) = id_x \otimes \phi_{x \rightarrow y} + \phi_{x \rightarrow y} \otimes id_y$. The quotient is by the coideal that $p = q$ whenever $s(p) = s(q)$ and $t(p) = t(q)$. This corresponds to modding out the category by the relation obtained from any diagram of two chains of morphisms with the same start and end.

2.4.3. Monoid/Groupoid/Setlike coalgebra. In the case of a unital monoid M , the corresponding category is often denoted by \underline{M} . It has one object $\{*\}$ and morphisms M . The coproduct is then simply

$$\Delta : R[M] \rightarrow R[M] \otimes R[M] : \Delta(m) = \sum_{(m_1, m_2) : m_1 m_2 = m} m_1 \otimes m_2 \quad (2.10)$$

and in case that M is finite, this is indeed the coproduct on characters, viz. on $M^* = Hom(M, R) = Hom_R(R[M], R)$. The identity e is grouplike, if and only if there are pairs m_1, m_2 , such that $m_1 m_2 = e$ that is no elements with a right inverse or equivalently no elements with a left inverse. An element $m \neq e$ is primitive if is indecomposable and is semigrouplike, with $\epsilon(m) = 0$ if it is idempotent.

The setlike coalgebras are just the coalgebras for the free unital monoid generated by the set S . This is $S^\times = \coprod_{n \in \mathbb{N}_0} S^{\times n}$ with merging stings as the product. $(s_1, \dots, s_n, s'_1, \dots, s'_m) = (s_1, \dots, s_n) \cdot (s'_1, \dots, s'_m)$ with the identity being the empty collection $() = \emptyset \in S^{\times 0} = \emptyset$.

2.4.4. Group(oid) coalgebras and Isomorphism. If all elements are invertible, that is $M = G$ is a group then \underline{G} is often called the associated groupoid and the coproduct is given by

$$\Delta(g) = \sum_{h \in g} h \otimes h^{-1}g = \sum_{h \in G} gh \otimes h^{-1} \quad (2.11)$$

and the counit is $\epsilon(e) = 1, \epsilon(g) = 0$ for $g \neq e$.

A category is a groupoid if all its morphisms are invertible. Given a category \mathcal{C} , $Iso(\mathcal{C})$, that is \mathcal{C} with only the isomorphisms is a groupoid.

The groupoid acts from the left and right on the $Hom(X, Y)$ by $\phi \rightarrow (\sigma \Downarrow \sigma')(\phi) := \sigma' \circ \phi \circ \sigma^{-1}$. This action is reflected in the coproduct: for $\phi \in Hom(X, Y)$

$$\Delta(\phi) = \sum_{\sigma_x \in Iso(X, \cdot), \sigma_y \in Iso(Y, \cdot)} [\sigma_x \otimes \phi \circ \sigma_x^{-1} + \sigma_y \circ \phi \otimes \sigma_y^{-1}] + \dots \quad (2.12)$$

Thus we will not have any primitive or skew primitive elements in the presence of isomorphisms.

Definition 2.8. We will call a morphism ϕ *essentially skew primitive* if $\Delta(\phi)$ is only given by the first two terms in (2.12). This is synonymous with essentially indecomposable, which means that the only decompositions into two factors have at one factor being an isomorphism.

Remark 2.9.

- (1) If \mathcal{C} is skeletal, then the sum will only be over automorphisms.
- (2) Using Δ^{iso} as in §2.6.2 turns the essentially skew primitive morphisms into skew primitives.
- (3) There is also a left and right comodule structure of $\mathcal{C}(C)$ over $C(Iso(\mathcal{C}))$ which captures this structure: This is given by

$$\lambda(\phi) = \sum_{\sigma_x \in Iso(s(\phi), \cdot)} \sigma_x \otimes \phi \circ \sigma_x^{-1}, \rho(\phi) = \sum_{\sigma_y \in Iso(t(\phi), \cdot)} \sigma_y \circ \phi \otimes \sigma_y^{-1} \quad (2.13)$$

2.4.5. A complete graph groupoid, iterated integrals and multi-zetas and polylogs. In [GCKT20a, 4], we considered the simplicial object coming from the path-algebra of a complete graphs. The elements of the groupoid algebra can be translated into sequence of vertices or numbers, by noticing that there is exactly one morphism between any two vertices. Thus, the path is fixed once the vertex sequence is fixed. These translate in bi-algebras, also see [GCKT20b, §3.3.1]. In particular the complete graph on two vertices $\{0, 1\}$, which yields the groupoid $\hookrightarrow 0 \rightrightarrows 1 \hookleftarrow$ and whose path algebra is responsible for Goncharov's and Brown's Hopf algebras. This is also the fundamental path groupoid for $\mathbb{C} \setminus \{0, 1\}$ with tangential base points and is directly linked, [GCKT20a, §1], to iterated integrals [Che73]. In particular, the bi-algebra structure is founded in the simplicial structure, see §6.1.2.

2.5. Algebras from strict monoidal categories. If a category has a strict monoidal structure, then the morphisms become a unital monoid and hence $R[Mor]$. Recall that a monoidal structure $\otimes : \mathcal{C} \times \mathcal{C} \rightarrow \mathcal{C}$ on a category defines a monoid on objects $(X, Y) \rightarrow X \otimes Y$ and on morphisms $(f, g) \rightarrow f \otimes g$, which extends to a monoid structure on $R[Mod]$. in particular $Hom(X, Y) \times Hom(X', Y') \rightarrow Hom(X \otimes X', Y \otimes Y')$. There is also a unit object $\mathbb{1}$ for which $X \otimes \mathbb{1} = \mathbb{1} \otimes X = X$ in the strict case. It follows that $f \otimes id_{\mathbb{1}} = id_{\mathbb{1}} \otimes f = f$, is a unit for the algebra structure.

Example 2.10. For posets, this is simply the product of the posets, with the order $(x, x') < (y, y')$ iff $x < y \wedge x' < y'$.

2.6. Categorical Bialgebras from Feynman categories. Notice that for a locally finite strict monoidal category, although there is a unital algebra and a counital coalgebra structure, the bialgebra equation does not need to hold. It does hold if the monoidal category has extra structure, namely that of a Feynman category, see [KW17]. It roughly says that the objects are a free \otimes -monoid on basic objects and that the morphisms are a free monoid on the so-called basic morphisms, which have \otimes indecomposables as targets. By definition a Feynman category is a triple $(\mathcal{V}, \mathcal{F}, \iota)$ of a groupoid \mathcal{V} , of basic objects aka. vertices, a (symmetric) monoidal category \mathcal{F} and a functor $\iota : \mathcal{V} \rightarrow \mathcal{F}$, such that

- (i) The free (symmetric) monoidal category \mathcal{V}^{\otimes} is equivalent to $Iso(\mathcal{F})$ via ι , where $Iso(\mathcal{F})$ is \mathcal{F} with only its isomorphisms as morphisms.
- (ii) The groupoid of morphisms $Iso(ar(\mathcal{F}))$ is equivalent to the free (symmetric) monoidal category on the morphisms from objects from \mathcal{V}^{\otimes} to \mathcal{V} . In formulas $Iso(ar(\mathcal{F})) \simeq Iso(\mathcal{V}^{\otimes} \downarrow \mathcal{V})$.
- (iii) The category of objects over a given object of \mathcal{F} is essentially small.

The category $Iso(ar(\mathcal{F}))$ has the morphisms of \mathcal{F} as objects and the morphisms are given by pairs of isomorphisms (σ, σ') which map $f \mapsto \sigma^{-1} f \sigma'$ when the compositions are defined. The first condition says that every object of \mathcal{F} decomposes $X \simeq \iota(*_1) \otimes \cdots \otimes \iota(*_n)$ essentially uniquely, where this means up to isomorphisms on the indecomposables and in the symmetric case up to permutations. Dropping the ι the second condition means that any morphisms $\phi : Y \rightarrow X$ of \mathcal{F} decomposes into indecomposables as well $\phi = \phi_1 \otimes \cdots \otimes \phi_n$ where each the $\phi_i : Y_i \rightarrow *_i$ with $X \simeq *_1 \otimes \cdots \otimes *_n$ and $Y \simeq Y_1 \otimes \cdots \otimes Y_n$, that is into multi-to-one morphisms. Again this is essentially unique, that is up to isomorphisms on the ϕ_i and permutations in the symmetric case. The third condition is technical and used for the existence of certain colimits.

Remark 2.11. The following considerations also works in an enriched setting, that is if the morphisms themselves are objects of a symmetric monoidal category, e.g. R -modules. This is handled in [KW17], but for brevity, we will not consider this situation here.

For instance for operads \otimes for the monoidal structure in which the algebra or operad takes values. If it is set valued, as in all the examples, $\otimes = \amalg$ is simply the disjoint union. In full generality, the notion of algebra or operad with values in \mathcal{C} etc. is replaced by $\mathcal{F}\text{-Ops}_{\mathcal{C}} = \text{Fun}_{\otimes}[\mathcal{F}, \mathcal{C}]$, the category of strong monoidal functors from \mathcal{F} to \mathcal{C} . For readers not familiar with operads, this and the examples below will give a definition.

Example 2.12 (Finite sets/Surjection/Injection). A good simple, but not too simple Feynman category is given by $\mathcal{F} = \text{FinSet}$, with disjoint union \amalg . There is only one vertex type, that is $\mathcal{V} = \ast$ the category with one object and its identity, which is $\underline{1}$ for the trivial group $G = 1$ and any choice of ι . Fix $\iota(\ast) = \{\ast\}$, this is a choice of an atom in set theoretical terms. The first condition is fulfilled as for any set $T \simeq \amalg_{t \in T} \{\ast\}$. Now any morphisms $S \rightarrow T$ decomposes into fibers $\amalg_{t \in T} f|_{f^{-1}(t)}$ with $f|_{f^{-1}(t)} : f^{-1}(t) \rightarrow \{t\} \simeq \{\ast\}$.

Note FinSet is not locally finite. However the subcategories FI and FS of finite sets and injections and finite sets and surjections are both locally finite Feynman categories, see [KW17, GCKT20b].

Example 2.13 (Ordered finite sets/Simplicial). Adding an order to the finite sets and considering only order preserving morphisms, one obtains a non-symmetric Feynman category $\text{FinSet}_{>}$ by the same construction.

Restricting to a skeleton, one obtains the category whose objects are only the natural numbers. This category is also known as Δ_+ or the simplicial category with unit. To make the translation, one can use the notation $\bar{n} = \{1, \dots, n\}$, with the natural order so that $S \simeq |\bar{S}|$ —the bijection is fixed by the order. In the usual simplicial notation $[n] = \{0, \dots, n\}$ so that $[n] \simeq \overline{n+1}$ and $\emptyset = [-1] = \bar{0}$. The disjoint union becomes addition $\bar{n} \amalg \bar{m} = \overline{n+m}$ in the first notation and join in the second as $[n] \ast [m] = [n+m+1]$ as $n+1+m+1 = n+m+2$.

The permutations $\sigma : \bar{n} \rightarrow \bar{n}$ are not order preserving. There are two ways to add them. The first retrieves finite sets as above and the second non-commutative sets as the crossed simplicial group ΔS , see [Kau21b] for details.

Again $\text{FinSet}_{>}$ or even its skeleton Δ is not locally finite, but the subcategories OI and OS of finite ordered sets and order preserving injections both have locally finite skeleta which are non-symmetric Feynman categories. This means that in Δ we only face maps or degeneracies, see [Kau21b] for details.

Example 2.14 (Hereditary collection of posets). A hereditary collection of posets \mathcal{P} , following [Sch87], is a collection of posets, that is closed under Cartesian products. The underlying \mathcal{V} is discrete, that is there are no non-trivial morphisms. It is given as the disjoint union of all the posets that are not a product of two other posets. For instance the free hereditary poset based on a poset $P = (S, <)$ given by the free monoid $S^{\times} = \amalg_{n \in \mathbb{N}_0} S^{\times n}$ with the product order on each of the $S^{\times n}$.

Example 2.15 (Free Feynman category on a category, Quivers, Posets). Given a category \mathcal{C} , then setting $\mathcal{V} = \text{Iso}(\mathcal{C})$ and $\mathcal{F} = \mathcal{C}^\otimes$ yields a Feynman category for the inclusion of $\mathcal{V} \rightarrow \mathcal{F}$. The basic morphisms are those of \mathcal{C} .

- (1) If $\text{Iso}(\mathcal{C})$ is discrete, viz. the identities are the only isomorphisms, then \mathcal{V} are just the objects under the identification $X \rightarrow id_X$.
- (2) Quivers and Posets are examples of discrete categories whose free symmetric monoidal categories yield Feynman categories.
- (3) The simplest example with a given \mathcal{V} is $\mathcal{F} = \mathcal{V}^\otimes$, which is a trivial Feynman category on \mathcal{V} and an initial object in the category of Feynman categories with fixed \mathcal{V} . The case of $\mathcal{V} = \underline{G}$ is of interest as it covers the representation theory of G , see [Kau21b].

Example 2.16 (Operads/Trees/Connes-Kreimer). Given a (non-symmetric) operad \mathcal{O} , for simplicity, let's assume $\mathcal{O}(1) = \mathbb{1}$, that is the $\mathcal{O}(1)$ is reduced for simplicity, see [Kau21b] and [KW17] for details about the non-reduced case. Here $\mathbb{1}$ is the monoidal unit, which is R for R -modules and the one point set $\{*\}$ for sets or topological spaces with Cartesian product and \emptyset for sets with disjoint union. Then letting the objects be \mathbb{N}_0 , setting $\text{Hom}(n, 1) = \mathcal{O}(n)$ and $\text{Hom}(n, m) = \bigotimes_{f:n \rightarrow m} \mathcal{O}(|f^{-1}(i)|)$ defines a (non-symmetric) Feynman category. Setting $\mathcal{O}_{CK}^{(pl)}(n)$ the set of (planar) rooted (planar) trees with n leaves, which are labeled in the non-planar case. We obtain the (non-symmetric) Feynman categories for (planar) rooted forests. More details are in §6.2.

Remark 2.17. There is a slight generalization in the case of operads, where one can consider operads with multiplication, [GCKT20a, §3]. This adds an associative multiplication $\mathcal{O}(n_1) \otimes \cdots \otimes \mathcal{O}(n_k) \rightarrow \mathcal{O}(n)$. This can be handled more generally in a non-connected setting [KW17, §3.2] which has relations to the B_+ operator.

2.6.1. Non-symmetric, non-commutative case.

Theorem 2.18. [GCKT20b] *If a locally finite monoidal category \mathcal{F} is part of a Feynman non-symmetric category \mathfrak{F} , then the algebra and coalgebra structures give a bialgebra structure on $R[\text{Mor}(\mathcal{F})]$. This bialgebra is not cocommutative in general.*

Example 2.19 (Leaf labelled rooted trees aka. planar Connes-Kreimer with legs). The bialgebras of leaf labelled rooted forests is an example. The bialgebra is the free algebra on $\bigoplus_{n \in \mathbb{N}_0} R[\mathcal{O}_{CK}^{(pl)}(n)]$ that is simply symmetric planar forests, where n is now the total number of leaves. The coproduct is given by admissible cuts, analogously to [CK98], but with the difference that upon a cut, the edge is not removed, but is split into a new root tail and a leaf, see [GCKT20a] and Figure 2. A full treatment is in §6.2.

Example 2.20 (Non-Sigma operads and Posets). In fact all non-Sigma operads including binary trees, the cyclic, associative operads, planar, non-rooted trees, Stasheff polyhedra, etc. The bialgebra will be on the free

algebra generated by $\bigoplus_{n \in \mathbb{N}_0} \mathcal{O}(n)$, that is $R[\mathcal{O}^{nc}] = \bigoplus_n R[\mathcal{O}^{nc}(n)]$ with $R[\mathcal{O}^{nc}(n)] = \bigoplus_{(n_1, \dots, n_k) | \sum n_i = n} R[\mathcal{O}(n_1)] \otimes \cdots \otimes R[\mathcal{O}(n_k)]$ —here nc stands for *non-connected*.

This also includes the free monoid on a poset as examples as considered in [Sch87].

2.6.2. Symmetric/commutative case. If the category \mathcal{F} is symmetric monoidal, being Feynman entails using a free *symmetric* monoidal category on a groupoid. This means that there are extra symmetries permuting objects and morphisms. The co-invariants are then yield a commutative monoids. To achieve this one considers isomorphism classes. This is the R -module $R[Mor(\mathcal{F})]$ with $f \sim g$ if there are isomorphisms σ, σ' , s.t. $f = \sigma' f \sigma^{-1}$. Note the product descends since if $\phi \sim \phi'$ and $\psi \sim \psi'$ then $\phi \otimes \psi \sim \phi' \otimes \psi'$. We denote the quotient by $R[Mor(\mathcal{F})]^{iso} = R[Mor(\mathcal{F})]/\sim$.

In order to give a formula, we need a few definitions. These are needed to avoid an over-counting of decompositions due to the isomorphisms given by the permutations. Otherwise, the coproduct will not be coassociative, see [KW17, Example 2.11].

Definition 2.21. [GCKT20b] A weak decomposition of a morphism ϕ is a pair of morphisms (ϕ_0, ϕ_1) for which there exist isomorphisms $\sigma, \sigma', \sigma''$ such that $\phi = \sigma \circ \phi_1 \circ \sigma' \circ \phi_0 \circ \sigma''$. In particular, a decomposition (ϕ_0, ϕ_1) is weakly composable. We introduce an equivalence relation on weakly composable morphisms, which says that $(\phi_0, \phi_1) \sim (\psi_0, \psi_1)$ if they are weak decompositions of the same morphism. An equivalence class of weak decompositions will be called a decomposition channel which is denoted by $[(\phi_0, \phi_1)]$.

The following is contained in Theorem 2.1.5 of [KW17]:

Theorem 2.22. *If a monoidal category \mathcal{F} is part of a Feynman category \mathfrak{F} , then the algebra and coalgebra structures give a bialgebra structure on $R[Mor(\mathcal{F})]^{iso}$, which is given by*

$$\Delta^{iso}([\phi]) = \sum_{[(\phi_0, \phi_1)]} [\phi_0] \otimes [\phi_1] \quad (2.14)$$

where the sum is over a complete system of representative for the decomposition channels for a fixed representative ϕ . The counit is given by $\epsilon([id_X]) = 1$ and $\epsilon([\phi]) = 0$ if $[\phi] \neq [id_{s(\phi)}]$.

Corollary 2.23. *A class $[\phi]$ is skew primitive if and only if ϕ is essentially skew primitive. Its coproduct is given by*

$$\Delta([\phi]) = [id_{s(\phi)}] \otimes [\phi] + [\phi] \otimes [id_{t(\phi)}] \quad (2.15)$$

The semigrouplike elements are the classes of the identities $[id_X]$ and they are grouplike.

Proof. Applying the definition, of the coproduct, we see that the two first terms in the coproduct of any ϕ for which $[\phi] \neq [id_{s(\phi)}] = [id_{t(\phi)}]$ will be

those of (2.15). These are the only two terms if ϕ is essentially skew primitive. If $[\phi] = [id_X]$, then there is only one term since then the decomposition channels coincide and $\Delta([id_X])$. Since $\epsilon([\phi]) = 1$, these are grouplike. \square

Example 2.24 (Leaf labelled rooted trees (Connes-Kreimer)). The bialgebras will be given by $R[\mathcal{O}_{CK}^{pl}(n)]$ and the \mathbb{S}_n coinvariants $R[\mathcal{O}_{CK}(n)]^{\mathbb{S}_n}$. The \mathbb{S}_n permute the labels, so the coinvariants can be identified with unlabelled trees. The coproduct is given by admissible cuts, analogously to [CK98], but with the difference that upon a cut, the edge is not removed but is split into a new root tail and a leaf, see [GCKT20a] and Figure 2. A full treatment is given in §6.2.

Example 2.25 (Bialgebra for operads). More generally, this construction works for operads and recovers the bialgebra for an operad with free multiplication of [GCKT20a]. The bialgebra structure is on the coinvariants. In particular on $\bigoplus_n R[\mathcal{O}^{nc}(n)]^{\mathbb{S}_n}$ which is the analogue of a symmetric forest. That is $\mathcal{O}^{nc}(n)^{\mathbb{S}_n} = \bigoplus_{(n_1, \dots, n_k) | \sum n_i = n} \text{Sym}(\mathcal{O}(n_1)^{\mathbb{S}_{n_1}} \otimes \dots \otimes \mathcal{O}(n_k)^{\mathbb{S}_{n_k}})$ where Sym is the symmetric product sometimes also denoted by \odot .

Example 2.26 (Connes-Kreimers graph bialgebra with legs). [GCKT20b, §3.4] There are Feynman categories whose isomorphism classes are graphs with external legs. That is given a morphism ϕ there is a graph $\mathbb{T}(\phi)$ which characterizes the isomorphism class. The coproduct is the usual CK-Coproduct

$$\Delta^{op}[\Gamma] = \sum_{\gamma \subset \Gamma} \gamma \otimes \Gamma/\gamma \quad (2.16)$$

where γ is a not necessarily connected subgraph and Γ/γ is the co-graph. Note that in this version the subgraph contains the half-edges of its vertices. See Figure 3 for a concrete example and §6.4 for more details.

Example 2.27. As being isomorphic is an order preserving equivalence relation for a category of a poset $\mathcal{C}(P)$, This construction recovers that of [Sch87] in this particular case.

2.7. The Drinfel'd double. There is a general construction, see [Kas95, IX.4] of a Drinfel'd double for a finite dimensional Hopf algebra with invertible antipode, we will concentrate on the case $k[G]$. This is an interesting Hopf algebra whose algebra part comes from a category.

Definition 2.28. For a finite group G a the *Drinfel'd double* $D(k[G])$ is the quasi-triangular quasi-Hopf algebra whose underlying *vector space* has the basis $g_{\frac{\iota}{x}}$ with $x, g \in G$

$$D(k[G]) = \bigoplus k g_{\frac{\iota}{x}} \quad (2.17)$$

With algebra structure is given by

$$g_{\frac{\iota}{x}} h_{\frac{\iota}{y}} = \delta_{g, xhx^{-1}} g_{\frac{\iota}{xy}} \quad (2.18)$$

and coalgebra structure is given by

$$\Delta(g_{\frac{\iota}{x}}) = \sum_{g_1 g_2 = g} g_{1\frac{\iota}{x}} \otimes g_{2\frac{\iota}{x}} \quad (2.19)$$

The antipode S is given by

$$S(g_{\frac{\iota}{x}}) = x^{-1} g^{-1} x_{\frac{\iota}{x^{-1}}} \quad (2.20)$$

The unit is $\sum_{g \in G} g_{\frac{\iota}{e}}$ and the counit is $\epsilon(e_{\frac{\iota}{x}}) = 1, \epsilon(g_{\frac{\iota}{x}}) = 0$ else.

Remark 2.29.

- (1) The algebra structure is the algebra of morphisms of the so-called loop groupoid, $\Lambda \underline{G}$. This is the category which has object set G and morphisms set $g_{\frac{\iota}{x}} \in \text{Hom}(g, x^{-1}gx)$, see e.g. [Wil08, KP09]. It is also the category $\text{Iso}(\underline{G})$.
- (2) The coalgebra structure is the categorical coalgebra for the groupoid $\coprod_{x \in G} \underline{G}$ which has $X = G$ as objects and has morphisms $g_{\frac{\iota}{x}} : x \rightarrow x$. Accordingly, the coalgebra is simply colored for the comodule structures for the set-like coalgebra $k[G]$: $\lambda(g_{\frac{\iota}{x}}) = x \otimes g_{\frac{\iota}{x}}, \rho(g_{\frac{\iota}{x}}) = g_{\frac{\iota}{x}} \otimes x$.
- (3) The notation $g_{\frac{\iota}{x}}$ goes back to [DPR90] where these are partition functions on a torus with for a field theory with finite gauge group, also see [Kau03, KP09].

Proposition 2.30. *The dual of $D(k[G])$ is a Hopf algebra, whose coproduct is the coalgebra structure coming from the morphisms of a category. In particular let $\delta_{g_{\frac{\iota}{x}}}$ be the dual basis then the product is given by*

$$\delta_{g_{\frac{\iota}{x}}} \delta_{h_{\frac{\iota}{y}}} = \delta_{x,y} g h_{\frac{\iota}{x}} \quad (2.21)$$

and the coproduct is given by

$$\Delta(\delta_{g_{\frac{\iota}{x}}}) = \sum_y \delta_{g_{\frac{\iota}{y}}} \otimes \delta_{y^{-1}gy} \quad y^{-1}x \quad (2.22)$$

and the antipode is

$$S(\delta_{g_{\frac{\iota}{x}}}) = \delta_{xg^{-1}x^{-1}} \quad x^{-1} \quad (2.23)$$

Proof. The first statement is clear from the Remark. For the second statement: Let $\delta_{g_{\frac{\cdot}{x}}}$ be the dual basis, then

$$\begin{aligned} \delta_{g_{\frac{\cdot}{x}}} \delta_{h_{\frac{\cdot}{y}}} (k_{\frac{\cdot}{z}}) &= \sum_{(k_1, k_2): k_1 k_2 = k} \delta_{g_{\frac{\cdot}{x}}} (k_{\frac{\cdot}{z}}) \delta_{h_{\frac{\cdot}{y}}} (k_{\frac{\cdot}{z}}) \\ &= \sum_{(k_1, k_2): k_1 k_2 = k} \delta_{g, k_1} \delta_{h, k_2} \delta_{x, z} \delta_{y, z} = \delta_{gh, k} \delta_{x, z} \delta_{y, z} \end{aligned} \quad (2.24)$$

For the coproduct,

$$\begin{aligned} \Delta(\delta_{g_{\frac{\cdot}{x}}})(h_{\frac{\cdot}{y}} \otimes k_{\frac{\cdot}{z}}) &= \delta_{g_{\frac{\cdot}{x}}}(h_{\frac{\cdot}{y}} k_{\frac{\cdot}{z}}) = \delta_{h, yky^{-1}} \delta_{g_{\frac{\cdot}{x}}}(h_{\frac{\cdot}{yz}}) \\ &= \delta_{g, h} \delta_{x, yz} \delta_{h, yky^{-1}} \end{aligned} \quad (2.25)$$

Lastly,

$$S(\delta_{g_{\frac{\cdot}{x}}})(h_{\frac{\cdot}{y}}) = \delta_{g_{\frac{\cdot}{x}}}(S(h_{\frac{\cdot}{y}})) = \delta_{g_{\frac{\cdot}{x}}}(y^{-1} h^{-1} y_{\frac{\cdot}{y^{-1}}}) = \delta_{g, y^{-1} h^{-1} y} \delta_{x, y^{-1}} \quad (2.26)$$

□

Remark 2.31. The product structure uses the underlying monoid structure of the objects/colors. This is not quite a monoidal structure on the category $\Lambda \underline{G}$. The definition of the antipode uses that the objects (as colors) are invertible. But note, that $g_{\frac{\cdot}{x}} \otimes h_{\frac{\cdot}{y}}$ would need to be a morphism from gh to $x^{-1}gxy^{-1}hy$ and if $x = y$ indeed $gh_{\frac{\cdot}{x}}$ is such a morphism. In general, if G is not Abelian no such morphism need exist.

In fact, the correction notion is that the Drinfel'd double is a subcategory of a natural double category structure on $Iso(\mathcal{C})$, see [Kau21b, Appendix C], where in this particular case $\mathcal{C} = \underline{G}$. This generalizes the dictum: “A monoidal category is a two category with one object”. The composability of horizontal 2-cells is then exactly the condition that $x = y$ in the above calculation. In particular the symbols $g_{\frac{\cdot}{x}}$ will two-morphisms with the following horizontal and vertical composition.

$$\begin{array}{ccccc} \begin{array}{c} * \xrightarrow{g} * \\ x \downarrow \Downarrow g_{\frac{\cdot}{x}} \downarrow \\ * \xrightarrow{xgx^{-1}} * \end{array} & \begin{array}{c} * \xrightarrow{g} * \\ x \downarrow \Downarrow g_{\frac{\cdot}{x}} \downarrow \\ * \xrightarrow{xgx^{-1}} * \end{array} & \begin{array}{c} * \xrightarrow{g} * \\ xy \downarrow \Downarrow g_{\frac{\cdot}{xy}} \downarrow \\ * \xrightarrow{xyg(xy)^{-1}} * \end{array} & \begin{array}{c} * \xrightarrow{g} * \xrightarrow{h} * \\ x \downarrow \Downarrow g_{\frac{\cdot}{x}} \downarrow \quad x \downarrow \Downarrow k_{\frac{\cdot}{x}} \downarrow \\ * \xrightarrow{xgx^{-1}} * \xrightarrow{hxx^{-1}} * \end{array} & \begin{array}{c} * \xrightarrow{gh} * \\ x \downarrow \Downarrow g_{\frac{\cdot}{x}} \downarrow \\ * \xrightarrow{xghx^{-1}} * \end{array} \\ & \begin{array}{c} * \xrightarrow{g} * \\ y \downarrow \Downarrow g_{\frac{\cdot}{y}} \downarrow \\ * \xrightarrow{xygy^{-1}x^{-1}} * \end{array} & & & \end{array} \quad (2.27)$$

The algebra structure is given the category algebra for the vertical composition, that is by \circ_h as the colored multiplication with $s_v(g_{\frac{\cdot}{x}}) = t_v(g_{\frac{\cdot}{x}}) = x$.

$$g_{\frac{\cdot}{x}} \cdot h_{\frac{\cdot}{y}} = \delta_{s(g_{\frac{\cdot}{x}})=t(h_{\frac{\cdot}{y}})} g_{\frac{\cdot}{x}} \circ_v h_{\frac{\cdot}{y}} \quad (2.28)$$

and Δ will be given as the dual of the horizontal composition with $s_h(g_{\frac{\cdot}{x}}) = g, t_h(g_{\frac{\cdot}{x}}) = xgx^{-1}$

$$\Delta(g_{\frac{\cdot}{x}}) = \sum_{(h_{\frac{\cdot}{y}}, k_{\frac{\cdot}{z}}): h_{\frac{\cdot}{y}} \circ_h k_{\frac{\cdot}{z}} = g_{\frac{\cdot}{x}}} h_{\frac{\cdot}{y}} \otimes k_{\frac{\cdot}{z}} \quad (2.29)$$

where we omitted the dual notation δ .

We will consider the generalization of the bialgebra and Hopf structures along these lines in [KY21].

3. ANTIPODES, CONVOLUTION INVERSES AND FILTRATIONS

3.1. Convolution algebra, inverses and antipodes. When C is a coalgebra and A is an algebra, the set of R -linear map $\text{Hom}(C, A)$ is not just a R -module but also a commutative unital k -algebra.

Definition 3.1. Given C a coalgebra and A an algebra, the k -module $\text{Hom}(C, A)$ is a R -algebra with unit $\eta \circ \epsilon$ and multiplication \star given by $(f \star g)(x) = m \circ (f \otimes g) \circ \Delta(x) = \sum_x f(x_{(1)})g(x_{(2)})$. We say $\{\text{Hom}(C, A), \star, \eta \circ \epsilon\}$ is the *convolution algebra* (associated to coalgebra C and algebra A).

If for $f, g \in \text{Hom}(C, A)$ there is $f \star g = g \star f = \eta \circ \epsilon$, we say f is the \star -inverse (or convolution inverse) to g and *vice versa* g to f .

Remark 3.2.

- (1) Because C is coassociative and A is associative, $\text{Hom}(C, A)$ is associative. In particular: $(f \star g) \star h = f \star (g \star h) = \sum_x f(x_{(1)})g(x_{(2)})h(x_{(3)})$.
- (2) A convolution inverse (or \star -inverse) of $f \in \text{Hom}(C, A)$ is the unique inverse $f^{\star^{-1}}$ under the convolution product \star since $\text{Hom}(C, A)$ is a monoid. This then implies antipode has to be unique as it is the convolution inverse of Id in $\text{Hom}(H, H)$.

3.1.1. (Co)unit preserving submonoid. If C is coaugmented with unit map η_C and A is augmented with counit ϵ_A . Let $G(C, A) \subset \text{Hom}(C, A)$: $G(C, A)$, where \cdot is a subset consisting of $f \in \text{Hom}(C, A)$ s.t. $\epsilon_A \circ f = \epsilon_C$ and $f \circ \eta_C = \eta_A$ where we added subscripts to clarify which unit/counit we are referring to.

Lemma 3.3. *If it exists, the convolution inverse of an element f in $G(C, A)$ is also in $G(C, A)$.*

Proof. Assume $f \in G(C, A)$ has a \star -inverse and denote it by g . For the compatibility with the unit: $f(1_C) = 1_A$ is given and we need to show that: $g \circ \eta_C = \eta_C$. Since $1_C = \eta_C(1_R)$ this means we need to show $g(1_C) = 1_A$. On one hand

$$\begin{aligned} (f \star g)(1_C) &= m_A \circ (f \otimes g) \circ \Delta_C(1_C) = (m_A \circ (f \otimes g))(1_C \otimes 1_C) \\ &= f(1_C)g(1_C) = g(1_C) \end{aligned}$$

On the other hand,

$$\begin{aligned} (f \star g)(1_C) &= (\eta_A \circ \epsilon_C)(1_C) = (\eta_A \circ \epsilon_C \circ \eta_C)(1_R) \\ &= \eta_A \circ (\epsilon_C \circ \eta_C)(1_R) = \eta_A(1_R) = 1_A \end{aligned}$$

For the compatibility with the counit we need to show that $\epsilon_A \circ g = \epsilon_C$.

$$\begin{aligned} \epsilon_C(a) &= \epsilon_C(a)1_R = \epsilon_C(a)\epsilon_A(1_A) = \epsilon_A(\epsilon_C(a)1_A) = \epsilon_A(\epsilon_C(a)\eta_A(1_R)) \\ &= \epsilon_A(\eta_A(\epsilon_C(a))) = \epsilon_A\left(\sum_a f(a_{(1)})g(a_{(2)})\right) \\ &= \sum_a \epsilon_A(f(a_{(1)}))\epsilon_A(g(a_{(2)})) = \sum_a \epsilon_C(a_{(1)})\epsilon_A(g(a_{(2)})) \\ &= \epsilon_A\left(\sum_a \epsilon_C(a_{(1)})g(a_{(2)})\right) = \epsilon_A\left(g\left(\sum_a \epsilon_C(a_{(1)})a_{(2)}\right)\right) = \epsilon_A(g(a)) \end{aligned}$$

This completes the proof. \square

Corollary 3.4. *For a bialgebra B to be a Hopf algebra, it is sufficient to prove $G(B, B)$ is a group. Because when $C = A = B$ are bialgebras, by definition the antipode S is two-sided \star -inverse to id_B which is an element of $G(B, B) \subset \text{Hom}(B, B)$.*

3.1.2. Characters.

Definition 3.5. If B is a bialgebra over R and A is an algebra over R , then the set of *characters* of B with values in A are the algebra homomorphisms $\text{Hom}_{R\text{-alg}}(B, A)$.

An algebra homomorphism preserves the unit, hence $\phi(1_C) = 1_A$, the characters are a subset of $G(B, A)$.

Proposition 3.6.

- (1) *If A is commutative, the characters form an algebra under convolution.*
- (2) *If B is a Hopf algebra, then $\phi \circ S = \phi^{-1}$ is the convolution inverse.*
- (3) *If B is Hopf and A is commutative, then the characters form a group.*

Proof. For the first statement: $(f \star g)(ab) = (f \otimes g)(\sum a_{(1)}b_{(1)} \otimes a_{(2)}b_{(2)}) = \sum f(a_{(1)})f(b_{(1)})g(a_{(2)})g(b_{(2)}) = \sum f(a_{(1)})g(a_{(2)})f(b_{(1)})g(b_{(2)}) = (f \star g)(a)(f \star g)(b)$.

For the second statement: $((\phi \circ S) \star \phi)(a) = \sum_a \phi(S(a_{(1)}))\phi(a_{(2)}) = \sum_a \phi(a_{(1)}S(a_{(2)})) = \phi(\eta \circ \epsilon(a)) = (\eta \circ \epsilon)(a)$.

For the last statement, we need to show that the inverse of a character is a character, but $\phi(S(ab)) = \phi(S(b)S(a)) = \phi(S(b)\phi(S(a)) = \phi(S(a))\phi(S(b))$, where the last equation holds since A is commutative. \square

Remark 3.7.

- (1) The commutativity can be generalized to a braided commutativity and a compatibility of the bradding operations on the bialgebra and algebra sides.
- (2) The second statement says that an antipode guarantees the convolution inverse. But, an antipode is not strictly necessary for a convolution inverse to exist. If a \star -inverse for a character exists, it constrains a putative antipode. Lastly a partially defined antipode may suffice to \star -invert a particular set of characters.

3.2. Product of submodules. There are several complication when working over a commutative ring R . First, there is no canonical way to identify a tensor product of submodules with a submodule. This may lead to different subcoalgebra structures. Second, a submodule with the restriction of the coalgebra structure may not yield a coalgebra. It may not even be possible to give a coalgebra structure.

Example 3.8. [NS82] Take $C = \mathbb{Z} \oplus \frac{\mathbb{Z}}{4\mathbb{Z}}$ as a module over \mathbb{Z} . We give C a coalgebra structure by making $g = (1, 0)$ grouplike and $x = (0, 1)$ g -primitive. Notice $\epsilon(x) = 0$ because it is a torsion-element. Consider a submodule V spanned by $2x$ and g . Then, by our notation $\Delta(V) \subset V \boxtimes V$. For instance, $\Delta(5x + 2g) = 5x \otimes x + x \otimes 2g + 2g \otimes x$ with $2g, x, 5x \in V$. Nicolas and Sweedler point out that one can give two non-isomorphic coalgebra structures on V , which make inclusion of V to C a coalgebra morphism.

Example 3.9. [NS82] Start with the \mathbb{Z} -module $C = \frac{\mathbb{Z}}{8\mathbb{Z}} \oplus \frac{\mathbb{Z}}{2\mathbb{Z}}$. Let $x = (1, 0)$ and $z = (0, 1)$ and endow C with a coalgebra structure by setting $\Delta(x) = 0$ and $\Delta(z) = 4x \otimes x$. Since $4x \otimes x$ has order of 2 in $C \otimes C$ the coproduct is well-defined. Also, $\epsilon(x) = \epsilon(z) = 0$, because they are torsion elements. Let $y = 2x$ and consider $V = \mathbb{Z}y + \mathbb{Z}z \subset C$. As $\Delta(z) = y \otimes y$, $\Delta(V) \subset V \boxtimes V$. For instance, $\Delta(z + y) = y \otimes y \in V \boxtimes V$. But unlike to the previous example, there is no coalgebra structure on V s.t. the natural inclusion is a coalgebra morphism. One can check the map $\Delta : V \rightarrow C \otimes C$ has no lifting to $V \otimes V$ because every preimage of $\Delta(z)$ has order 4.

The complication arises due to torsion elements, which is one of the difficulties of handling \mathbb{Z} -bialgebras. To avoid these issues, we will use an internal product denoted by \boxtimes for the underlying R -submodules.

Remark 3.10. If the inclusion $C' \subset C$ split, then C' has a unique coalgebra structure inherited from C . In this sense, we can consider C' as a subcoalgebra.

Definition 3.11. Given a coalgebra (C, Δ, η) over R and two submodules M, N of C , their product $M \boxtimes N$ is defined to be the following submodule

of $C \otimes C$

$$M \boxtimes N := \{\text{finite sum of } x_{(1)} \otimes x_{(2)} \in C \otimes C \mid x_{(1)} \in M, x_{(2)} \in N\}$$

For two families of submodules $\{N_j\}, \{M_i\}$, we will also use the notation $\Delta(x) \in \sum_j M_j \boxtimes N_i$. This means $\Delta(x)$ can be decomposed to finite sum of $x_{(1)} \otimes x_{(2)}$ such that every summand belongs to $M_j \boxtimes N_i$ for some i, j .

We will simply regard the structures on a submodule as the restriction $\Delta|_V : V \rightarrow C \otimes C$ for $V \subset C$. Then one can say that V is closed under comultiplication, if $\Delta(V) \subset V \boxtimes V$.

The following property is straightforward:

Lemma 3.12. *If $M \subset M'$ and $N \subset N'$ as submodules, then $M \boxtimes N \subset M' \boxtimes N'$. \square*

Remark 3.13. In the case that $R = k$ is a field, we can work with the usual monoidal product \otimes .

Definition 3.14. We call a submodule S semigrouplike, if $\Delta(S) \subset S \boxtimes S$.

Lemma 3.15. *Given a semigrouplike S , $\text{Hom}(S, A)$ is a unital subalgebra of the convolution algebra with unit $(\eta \otimes \epsilon)|_S = \eta \circ \epsilon|_S$.*

Proof. Since $\Delta(S) \subset S \boxtimes S$, the formula $(f \star g)(s) = m(f \otimes g) \circ \Delta(s) = \sum f(s')g(s_{(2)})$ makes sense and hence $\text{Hom}(S, C)$ is a subalgebra. The unit restricts. \square

Proposition 3.16. *Given a coaugmented coalgebra, for every $x \in \overline{C}$, $\Delta(x)$ can be written as $e \otimes x + x \otimes e + w$ with $w \in \overline{C} \boxtimes \overline{C}$, that is $w = \overline{\Delta}(X)$.*

Moreover, if $C = \bigoplus_{i \geq 0} C_i$ (thus $\overline{C} = \bigoplus_{i > 0} C_i$) is graded and coaugmented and $x \in C_n$ for $n > 0$, then $w \in \bigoplus_{i=1}^{n-1} C_i \boxtimes C_{n-i}$.

Proof. Since $C = Re \oplus \overline{C}$, using the distributive law

$$\begin{aligned} C \otimes C &= (Re \oplus \overline{C}) \otimes (Re \oplus \overline{C}) \\ &= (Re \boxtimes Re) \oplus (e \boxtimes \overline{C}) \oplus (\overline{C} \boxtimes Re) \oplus (\overline{C} \boxtimes \overline{C}) \end{aligned} \quad (3.1)$$

Now, if $x \in \overline{C}$ then decomposing $\Delta(x) = \mu' \otimes \mu'' + \lambda' \otimes y'' + y' \otimes \lambda'' + z \otimes z'$ according to (3.1). Applying the left counit constraint $(\epsilon \otimes id) \circ \Delta = id$ to x , we find $x = \mu\mu + \lambda y''$. Since $\epsilon(x) = 0$, $\mu'\mu'' = 0$, so the first term vanishes and hence $\lambda'y' = x$ and the second term must equal $e \otimes x$. Similarly from the right counit constraint, the third term is $x \otimes e$ proving the claim. In the graded case $C \otimes C = \bigoplus_{i \geq 0} C_i \otimes \bigoplus_{j \geq 0} C_j = \bigoplus_{i,j \geq 0} C_i \boxtimes C_j$, which also shows that the sum $\sum_{i=0}^n C_i \otimes C_{n-i}$ is direct. We can consider $\overline{C} \boxtimes \overline{C} = \bigoplus_{i,j > 0} C_i \boxtimes C_j$ as R -submodule of $C \otimes C$. The comultiplication is graded, so $\Delta x \in \bigoplus_{i=0}^n C_i \boxtimes C_{n-i}$ and in this particular case it equals $\bigoplus_{i \geq 0} C_i \boxtimes C_{n-i}$. Therefore, $w \in \bigoplus_{i \geq 0} C_i \boxtimes C_{n-i} \subset \bigoplus_{i,j \geq 0} C_i \boxtimes C_j$. But $w \in \bigoplus_{i,j > 0} C_i \boxtimes C_j$, and in particular, the projections of w onto $C_0 \boxtimes C_n$ and $C_n \boxtimes C_0$ are zero. Therefore, we conclude $\overline{\Delta}(x) \in \bigoplus_{i=1}^{n-1} C_i \boxtimes C_{n-i}$. \square

3.3. Filtrations. In order to define antipodes it is useful to regard filtrations, as in good cases one can recursively build up the antipode from its value on the degree 0 part.

We recall classical filtrations used to study coalgebras and Hopf algebras before we give the definition of a QT-filtration (Quillen Takeuchi) which generalizes all of them. This is what is needed to prove a general version of Takeuchi's lemma for the existence of convolution inverses.

3.3.1. Coalgebra filtration.

Definition 3.17. A *coalgebra filtration* $\{C_n\}_{n \geq 0}$ of C over a field R is a family of R -submodules $\{C_n\}$ of C which satisfies $C_0 \subset C_1 \subset C_2 \dots = \bigcup_{i \geq 0} C_i = C$ and $\Delta(C_n) \subset \sum_{l=0}^n C_{n-l} \boxtimes C_l$ for all $n \geq 0$.

Every *graded* coalgebra has a natural coalgebra filtration given by $C_r := \{x \mid \deg x \leq r\}$.

In general, this is too strong a condition. Weaker conditions are the following ones.

3.3.2. Coradical filtration and pointed coalgebras. We briefly recall the concept of pointed coalgebra and coalgebra filtration [Rad12] [Mon93].

Definition 3.18. Let C be a coalgebra over a field k . The *coradical* of C is sum of the simple coalgebras of C and denoted by C_0 , where *simple coalgebra* is a coalgebra whose subcoalgebras are only 0 and itself.

Every coalgebra has a natural coalgebra filtration called *coradical filtration*, see e.g. [Rad12], which is defined recursively by the following

- (1) C_0 = sum of all simple subcoalgebras of C .
- (2) $C_n = \Delta^{-1}(C \otimes C_0 + C_{n-1} \otimes C)$

Definition 3.19. A coalgebra over a field k is called *pointed* if all simple subcoalgebras of C are 1-dimensional.

Example 3.20. A path coalgebra over k is pointed, with the 1-dim subcoalgebras being the $k[v], v \in V$. In general in a pointed coalgebra the 1-dimensional subalgebras are precisely those generated by a grouplike element. $C_0 = \bigoplus_{g \in GLE} k[g]$. A conilpotent coalgebra or a connected coalgebra, see below, are also pointed.

There is a nice structure theorem of pointed coalgebras, see [TW74] and its improved version [Rad12, Proposition 4.3.], which essentially motivates our work on flavored coalgebras. This can be seen as a generalization of Proposition 3.16.

Theorem 3.21. Let C be a pointed coalgebra over a field k with coradical filtration $\{C_i\}_{i=0}^{\infty}$ and denote the set of grouplike elements in C as $gpl(C)$. For any $c \in C_n$,

$$c = \sum_{g,h \in gpl(C)} c_{g,h}, \text{ where } \Delta(c_{g,h}) = c_{g,h} \otimes g + h \otimes c_{g,h} + w$$

For some $w \in C_{n-1} \otimes C_{n-1}$.

3.3.3. Conilpotent coradical filtration. For a coaugmented coalgebra, with $e = \eta(1)$ and $\bar{C} = \ker(\epsilon)$ as above, there is a filtration called *conilpotent coradical filtration*, see e.g. [Lod98], which is recursively defined as follows: Note that $\bar{\Delta} : \bar{C} \rightarrow \bar{C} \boxtimes \bar{C}$. The iteration of $\bar{\Delta}$ on \bar{C} is defined by $\bar{\Delta}^0 = 0$, $\bar{\Delta}^1 = id_{\bar{C}}$ and $\bar{\Delta}^{n+1} = (\bar{\Delta}_{\bar{C}} \otimes id_{\bar{C}} \otimes \cdots \otimes id_{\bar{C}}) \circ \bar{\Delta}^{n-1}$. This yields a filtration of \bar{C} and in turn the following filtration on C :

- (1) $F_0 C = Re$
- (2) $F_r C = Re \oplus \{x \in C \mid \bar{\Delta}^i(x) = 0, i \geq r\}$

A coaugmented coalgebra is called conilpotent if the above filtration is exhaustive. This means that for every element $x \in \bar{C}$ there is some n such that $\bar{\Delta}^n(x) = 0$ or equivalently letting $p = (id_C - \eta \circ \epsilon)$ be the projection to \bar{C} for any x there is an n such that $x \in \ker(p^{\otimes n+1} \circ \Delta^{[n]})$.

Lemma 3.22. *A connected graded coaugmented coalgebra C is conilpotent.*

Proof. Given $x \in \ker(\epsilon) \cap C_r$, $n \geq 0$, by Proposition 3.16, $\bar{\Delta}(x) \in \bigoplus_{i=1}^r C_i \times C_{r-i}$ and hence $\bar{\Delta}^r(x) = 0$ and moreover if $x \in C_r$, we have $\bar{\Delta}^r(x) = 0$ for all $i \geq r$ and thus the conilpotent coradical filtration is exhaustive. \square

3.3.4. Quillen filtration. For a coaugmented coalgebra Quillen [Qui67] introduced a filtration which can be written as follows. Setting $e = \eta(1)$, see Remark 1.5, the *Quillen filtration of a coaugmented coalgebra C* over a ring R is defined as:

- (1) $F_0^Q C = Re$
- (2) $F_r^Q C = \{x \in C \mid \bar{\Delta} \in F_{r-1}^Q C \boxtimes F_{r-1}^Q C\}$

Furthermore, if this filtration is exhaustive $C = \bigcup_{i \geq 0} F_i C$, then the coalgebra will be called Quillen connected.

If k is a field, one can replace \boxtimes by \otimes which agrees with original definition when R is a DG coalgebra over k , see [Qui67].

Lemma 3.23. *A Quillen connected coalgebra is conilpotent coalgebra and if kernel and tensor commute a conilpotent coalgebra is Quillen connected.*

Proof. The zeroth part of the two filtrations agree. For the higher components, observe that $F_r^Q C \subset F_r C$ and there is an equality if tensor and kernel commute. \square

Remark 3.24. A Quillen connected coalgebra over a field k is the same as an irreducibly pointed coalgebra. More specifically, a coalgebra C over a field is said to be irreducibly pointed if its coradical is one dimensional and spanned by the unique grouplike element e . By Proposition 3.16, for every $x \in C_n$, $\bar{\Delta}(x) \in C_{n-1} \otimes C_{n-1}$. Due to the fact $F_0^Q C = C_0 = Re$, it is easy to see by induction $C_n = F_n C$ and thus the Quillen filtration and coradical filtration agree.

3.3.5. Connecteness. The different notions of connectedness are compatible.

Lemma 3.25. *A connected graded coaugmented coalgebra C is Quillen connected.*

Proof. Let $e = \eta(1)$ be the unique grouplike element. One can identify $F_0C = Re = C_0$ and $F_n^Q C = \bigoplus_{i=1}^n C_i$ is the Quillen filtration by induction. The cases of $n = 0, 1$ are trivial. Suppose $F_{n-1}C = \bigoplus_{i=1}^{n-1} C_i$ is true (also $F_{n-1} \boxtimes F_{n-1} \subset C \otimes C$ due to the grading). By definition, $F_nC = \{x \in C \mid \Delta(x) - x \otimes e - e \otimes x \in F_{n-1}C \otimes F_{n-1}C = \bigoplus_{i=1}^{n-1} C_i \otimes \bigoplus_{i=1}^{n-1} C_i\}$. Then $C_i \subset F_nC$ for $0 \geq i \geq n$ and $C_i \cap F_nC = \emptyset$ for $i > n$ by previous proposition. Then we conclude $F_nC = \bigoplus_{i=1}^n C_i$ by induction, from which the claim follows. \square

3.3.6. QT-filtrations. Motivated by Quillen and Takeuchi [Tak71], we consider the following general setup, which contains all the previous examples.

Definition 3.26. A *QT-filtration* of a coalgebra C is a filtration of C by R -submodules $F_p(C)$, $p \geq 0$ such that

- (1) $F_0C = S$ is semigrouplike, that is $\Delta(S) \subset S \boxtimes S$.
- (2) $\Delta(F_pC) \subset F_{p-1}C \boxtimes C + C \boxtimes F_{p-1}C$.

We call a QT-filtration a *QT-sequence* if the filtration is exhaustive $C = \bigcup_{i=0}^{\infty} F_iC$ and call the QT-sequence *split*, if F_0C is a direct summand, that is $0 \rightarrow F_0C \rightarrow C \rightarrow F_0C/C \rightarrow 0$ is split. A coaugmented coalgebra is *QT-connected* if it has a QT-sequence whose degree 0 part is Re .

Example 3.27. One can construct QT-filtrations in both graded coalgebra and coalgebra over a field.

- (1) The filtration associated with the grading of a \mathbb{N} -graded coalgebra ($F_r := \{x \mid \deg x \leq r\}$) is a QT-filtration based on F_0 , because $F_i \subset F_r$ if $i \leq r$. $F_i \boxtimes F_j \subset F_r \boxtimes F_q$ if $i \leq r$ and $j \leq q$.
- (2) More generally, in the category of k -coalgebras, every coalgebra filtration gives rise to a QT-filtration with S being the degree 0 part.
- (3) A special case is given by specifying a base, that is a semigrouplike submodule S as setting:
 - (a) $F_0C = S$
 - (b) $F_rC = \{x \mid \Delta(x) \in C \boxtimes F_0C + F_{p-1}C \boxtimes F_{p-1}C + F_0C \boxtimes C\}$.
Quillens original filtration is of this type.
- (4) Given a \mathbb{Z} graded filtration, with C_0 split semigrouplike and with $\Delta(F_p) \subset F_{p-1} \boxtimes C + C \boxtimes F_{p-1}$, $p > 0$ and $\Delta(F_p) \subset F_{p+1} \boxtimes C + C \boxtimes F_{p+1}$, $p < 0$. Also see Remark 3.24. Setting $\hat{F}_p = F_p + F_{-p}$, $p \geq 0$ produces a QT-filtration

We are going to prove a version of Takeuchi's lemma for QT-sequences. It was proven originally for the coradical filtration in [Tak71] and later generalized to any graded coalgebra. We now generalize to the context of QT-sequences.

Theorem 3.28. *Consider a coalgebra C with counit ϵ and an algebra A with unit η . Given a QT-sequence $\{F_i\}_{i=0}^\infty$ on C , such that $\text{Ext}_R^1(C/F_0C, A) = 0$, for any element f of $\text{Hom}(C, A)$, f is \star -invertible if and only if the restriction $f|_{F_0C}$ is \star -invertible in $\text{Hom}(F_0C, A)$.*

Proof. First assume that f is convolution invertible in $\text{Hom}(C, A)$, and let g be \star -inverse of f . By restriction to F_0C , see Lemma 3.15, it follows that $(\eta \circ \epsilon)|_{F_0C} = (f \star g)|_{F_0C} = f|_{F_0C} \star g|_{F_0C}$ and $(\eta \circ \epsilon)|_{F_0C}$ is the unit of the restricted convolution algebra $\text{Hom}(F_0C, A)$. Thus $g|_{F_0C}$ is a right \star -inverse of $f|_{F_0C}$. By symmetry, it is also a left \star -inverse and thus the \star -inverse in the restricted algebra.

Conversely, assume that $f|_{F_0C}$ is convolution invertible in $\text{Hom}(F_0C, A)$. Thus, there is an element g of $\text{Hom}(F_0C, A)$ such that $f \star g = \eta \circ \epsilon = g \star f$ on F_0C . We can extend the R -linear map g to C as the restriction morphisms $\text{Hom}(C, A) \rightarrow \text{Hom}(F_0C, A)$ is surjective due to the assumption that $\text{Ext}(C/F_0C, A) = 0$.

Because the convolution algebra $(\text{Hom}(C, A), \star, \eta \circ \epsilon)$ is a monoid, if we show that both $f \star g$ and $g \star f$ have a \star -inverse this will imply f has a \star -inverse. Thus, replacing f with $f \star g$, we may assume $f|_{F_0C} = \eta \circ \epsilon|_{F_0C}$ that is $\eta \circ \epsilon|_{F_0C} - f|_{F_0C} = 0$.

Since the filtration is exhaustive, for every $x \in C$, there exist a natural number N such that $x \in F_N$. Then we claim that $(f - \eta \circ \epsilon)^{\star(N+1)}(x) = 0$. This is because, after using the coassociativity for the diagonal to decrease the filtration degree, every summand from $\Delta^{[N+1]}(x) = \sum_x x_{(1)}x_{(2)}x_{(3)}\dots, x_{(N+1)}$ has to contain a factor in F_0C according to the definition of the QT-filtration, and thus $\eta \circ \epsilon - f$ annihilates each term. So, the infinite sum $h = \sum_{i=0}^\infty (\eta \circ \epsilon - f)^{\star i}$ is locally finite and well defined —where we set $(\eta \circ \epsilon - f)^{\star 0} = \eta \circ \epsilon$. It is easy to verify that h is R -linear, so $h \in \text{Hom}(C, A)$. That h is the inverse of f follows by computation: $f \star h = (f - \eta \circ \epsilon) \star h + \eta \circ \epsilon \star h = (f - \eta \circ \epsilon) \star h + h = -\sum_{i=1}^\infty (\eta \circ \epsilon - f)^{\star i} + \sum_{i=0}^\infty (\eta \circ \epsilon - f)^{\star i} = (\eta \circ \epsilon - f)^{\star 0} = \eta \circ \epsilon$. The fact that $h \star f = \eta \circ \epsilon$ can be checked in the same way. Therefore, we conclude h is \star -inverse of f . \square

This applies in all the situations above. Note that if $F_0^Q C$ is a direct summand, i.e. the sequence $0 \rightarrow F_0^Q C \rightarrow C \rightarrow C/F_0^Q C \rightarrow 0$ splits and the Ext group vanishes. In particular, we recover Takeuchi's and Quillen's results.

We say that a bialgebra has a QT-sequence, if the coalgebra has a QT-sequence.

Theorem 3.29. *A bialgebra with QT-sequence and $\text{Ext}_R^1(C/F_0^Q C, C) = 0$ is Hopf, i.e. has an antipode if and only if $\text{id}|_{F_0^Q C}$ has a \star inverse.*

Proof. By Corollary 3.4 the existence of an antipode is equivalent to the existence of a \star -inverse of id which by the Theorem 4.16 exists as soon as the Ext group vanishes and the restriction of id to F_0^Q has an inverse, which hold by assumption. \square

The following corollary recovers many of the existence theorems for antipodes.

Corollary 3.30. *A bialgebra with a split QT-sequence is Hopf if $\text{id}|_{F_0^Q C}$ has an inverse. In particular, if C is a QT-connected bialgebra, then it is Hopf* \square .

Proof. The first part directly follows from Theorem 3.29. If C is QT-connected then $S(e) = e$ is the required antipode on $F_0^Q C$. \square

4. FLAVORED COALGEBRAS

In this section, we will build a coalgebra which in principle should capture the essence of “pointedness” we mention at the introduction. It is based on a decomposition according to (g, h) -primitives and is motivated by the structure theorem for pointed coalgebra, we introduced at the end of section 1. It generalizes connected graded coalgebras and May’s component coalgebras [MP12]. The following construction has origins going back to [TW74].

4.1. Reduced comultiplication.

Definition 4.1. Let C be a R -coalgebra and g, h semigrouplike. We define the (g, h) reduced comultiplication to be $\bar{\Delta}_{g,h}(x) = \Delta(x) - x \otimes g - h \otimes x$. We will write $\bar{\Delta}_g := \bar{\Delta}_{g,g}$ and call it the reduced comultiplication associated to g . If there is a fixed grouplike element, like in Quillen’s work or if there is a unique such element, we will further abbreviate to $\bar{\Delta}$ and refer to it as the reduced comultiplication.

Proposition 4.2. *The new binary operation $\bar{\Delta}_{g,h}$ is coassociative if $g = h$, i.e. $(\bar{\Delta}_{g,g} \otimes \text{Id}) \circ \bar{\Delta}_{g,g} = (\text{Id} \otimes \bar{\Delta}_{g,g}) \circ \bar{\Delta}_{g,g}$. Furthermore, if g is grouplike and $x \in \ker \epsilon$, then $(\epsilon \otimes \text{Id})\bar{\Delta}_{g,g} = (\text{Id} \otimes \epsilon)\bar{\Delta}_{g,g} = 0$*

Proof. This is a straightforward computation. For the first part, the left hand side yields

$$\begin{aligned} (\bar{\Delta}_{g,h} \otimes \text{Id})\bar{\Delta}_{g,h}(x) &= \sum_x x_{(11)} \otimes x_{(12)} \otimes x_{(2)} - x_{(1)} \otimes g \otimes x_{(2)} - h \otimes x_{(1)} \otimes x_{(2)} \\ &\quad - x_{(1)} \otimes x_{(2)} \otimes g + x \otimes g \otimes g + h \otimes x \otimes g + h \otimes g \otimes x \end{aligned}$$

The right hand side

$$\begin{aligned} (\text{Id} \otimes \overline{\Delta}_{g,h})\overline{\Delta}_{g,h}(x) &= \sum_x x_{(11)} \otimes x_{(21)} \otimes x_{(22)} - x_{(1)} \otimes x_{(2)} \otimes g - x_{(1)} \otimes h \otimes x_{(2)} \\ &\quad + x \otimes h \otimes g - h \otimes x_{(1)} \otimes x_{(2)} + h \otimes x \otimes g + h \otimes h \otimes x \end{aligned}$$

The difference of both sides is

$$x_{(1)} \otimes (h - g) \otimes x_{(2)} - x \otimes (h - g) \otimes g - h \otimes (h - g) \otimes x$$

which vanishes when $h = g$.

For the second part,

$$(\epsilon \otimes \text{Id})\overline{\Delta}_{g,g}(x) = x - \epsilon(x)g - \epsilon(g)x = 0$$

Similar for other direction. This completes the proof. \square

As a result, the iterated reduced comultiplication $\overline{\Delta}_g^{[n]}(x)$ is well-defined.

4.2. Main construction. Given a coalgebra C , we will inductively define sequences of R -modules each depending on two elements $g, h \in \text{sgpl}(C)$. These will be simultaneously be defined for all pairs of such elements and will yield steps of filtrations.

(1) For any two $g, h \in \text{sgpl}(C)$:

$$F_0^Q C_{g,h} := Rg \cap Rh \text{ is the base } R\text{-module.}$$

(2) Inductively over all pairs:

$$F_n^Q C_{g,h} := \{x | \overline{\Delta}_{g,h}(x) = \sum_{t \in \text{sgpl}(C)} F_{n-1}^Q C_{g,t} \boxtimes F_{n-1}^Q C_{t,h}\}.$$

That is $F_n^Q C_{g,h}$ is the subset in which every element x in the set has the property $\overline{\Delta}_{g,h}(x) = \sum x_{(1)} \otimes x_{(2)}$ such that for each summand $x_{(1)} \otimes x_{(2)}$ there exists a semigrouplike element t with $x_{(1)} \in F_{n-1}^Q C_{g,t}$ and $x_{(2)} \in F_{n-1}^Q C_{t,h}$. This is also a R -module as every $F_n^Q C_{g_1,g_2}$ with $g_1, g_2 \in \text{sgpl}(C)$ is a R -module inductively. This can be written in our previous introduced notation

Remark 4.3. $F_0^Q C_{g,h}$ may be non-empty, if g, h are two distinct semigrouplike elements, as it may happen that there exist $s, t \in R$ such that $sg = th$ so that. $kg \cap kh \neq 0$.

As a concrete example, consider the set-like coalgebra $\mathbb{Z}\{x, y\}$, see example §2.1. Let I be the module generated by $5x - 5y$. We see $\epsilon(I) = 0$ and $\Delta(I) \subset C \otimes I + I \otimes C$ because $\Delta(5x - 5y) = (5x - 5y) \otimes x + y \otimes (5x - 5y)$. Therefore, I is a coideal and we have a quotient coalgebra $C = \mathbb{Z}\{x, y\}/I$. The image \bar{x}, \bar{y} of x, y are still grouplike elements in C . While $\bar{x} \neq \bar{y}$, $5\bar{x} = 5\bar{y}$. Therefore, $\mathbb{Z}\bar{x} \cap \mathbb{Z}\bar{y} \neq 0$.

This complication it will not happen in the case of coalgebra over a ground field k , because it implies $kg = kh$. Similarly, if the underlying modules are free over \mathbb{Z} : $sg = th$ also means $\mathbb{Z}g = \mathbb{Z}h$ as they are linearly independent. In fact, for both cases $M \otimes N$ is indeed submodule of $C \otimes C$.

Proposition 4.4. *The $F_i C_{g,k}^Q$ yield filtrations: $F_i C_{g,h}^Q \subset F_{i+1} C_{g,h}^Q$ for $i \geq 0$.*

Proof. We use induction. Base case: to show that $Rg \cap Rh = F_0 C_{g,h}^Q \subset F_1 C_{g,h}^Q$ pick $x \in F_0 C_{g,h}^Q$, say $x = rg$, then $\bar{\Delta}_{g,h}(x) = -x \otimes h \in F_0 C_{g,h}^Q \boxtimes F_0 C_{h,h}^Q$.

By induction assume that $F_{n-1} C_{g,h}^Q \subset F_n C_{g,h}^Q$ is true. Let $x \in F_n C_{g,h}^Q$. Then $\bar{\Delta}_{g,h}(x) \in \sum_{t \in \text{sgpl}(C)} F_{n-1} C_{g,t}^Q \boxtimes F_{n-1} C_{t,h}^Q \subset \sum_{t \in \text{sgpl}(C)} F_n C_{g,t}^Q \boxtimes F_n C_{t,h}^Q$ by induction hypothesis. Therefore, $x \in F_{n+1} C_{g,h}^Q$ and we conclude the proof by induction. \square

We set $C_{g,h}^Q = \sum_{i \geq 0} F_i C_{g,h}^Q = \bigcup_{i \geq 0} F_i C_{g,h}^Q$ and call it the (g, h) -Quillen component of C . The filtrations of $C_{g,h}^Q$ yield a common filtration $\bar{F}_r C := \sum_{g,h} F_r C_{g,h}^Q$ and call it the bivariate Quillen filtration. Note all semigrouplike elements of C lie in $\bar{F}_0 C$ and all skew primitive elements C lie in $\bar{F}_1 C$.

Remark 4.5. $C_{g,h}^Q$ is never empty, this is clear for $g = h$ and for $g \neq h$ $\Delta(g-h) = g \otimes (g-h) + (g-h) \otimes h$ and hence $\bar{\Delta}_{g,h}(g-h) = 0 \in F_0 C_{g,t}^Q \boxtimes F_0 C_{t,h}^Q$ for any t . Thus the sum in $C = \sum_{g,h} C_{g,h}^Q$ can never be direct, if there is more than one semigrouplike elements, as $g-h$ is in $C_{g,g}^Q + C_{h,h}^Q \cap C_{g,h}^Q$ and $g-h \in C_{g,h}^Q$.

In the case of a field $R = k$, $g-h$ generates the intersection and in fact over a field one can make the sum direct by reducing $C_{g,h}$ by $k(g-h)$ following the line of arguments presented in [Mon93].

Definition 4.6. We say a coalgebra C is *flavored* if the bivariate Quillen filtration is exhaustive, that is $C = \sum_{(g,h), g,h \in \text{sgpl}(C)} C_{g,h}^Q$. A bi- or Hopf-algebra is flavored if the underlying coalgebra is flavored.

Corollary 4.7. The $\bar{F}_r C := \sum_{g,h} F_r C_{g,h}^Q$ for $r \geq 0$, form a QT-filtration, which is a QT-sequence if C is flavored.

Proof. By definition the elements in $\bar{F}_0 C$ are multiples of semigrouplike elements and hence F_0 is semigrouplike. Condition (2) for a QT-filtration is satisfied by construction. In the flavored case, the filtration is exhaustive by definition. \square

Example 4.8. A connected graded coalgebra C is a flavored coalgebra. In fact, a Quillen connected coalgebra is a flavored coalgebra. There is only one component $C_{e,e}^Q$ with e the unique grouplike element and it actually equal the whole coalgebra C .

Example 4.9. A coalgebra C generated by semigrouplike elements is flavored. In fact, a bialgebra B generated as an algebra by semigrouplike elements is flavored. This is easy to see because multiplication of semigrouplike elements is semigrouplike. Thus, we can write $B = \sum_{g \in \text{sgpl}(B)} C_{g,g}^Q$ as every element of B is a finite sum of semigrouplike elements.

Example 4.10. Due to the symmetry of the definition of the filtration, the opposite of flavored coalgebra also still flavored. Moreover, $sgpl(H^{op}) = sgpl(H)$.

Corollary 4.11. *Given a flavored coalgebra then if F_0C^Q is a coideal, and $C^{red} := C/F_0^Q$ is QT-connected. For a flavored bialgebra it follows that $B/(F_0B^Q)$ is QT-connected and hence Hopf.*

Proof. The first statement follows directly from the definition as $F_0C_{g,h}^Q$ is already semigrouplike and the second from Corollary 3.30. \square

4.3. Pathlike coalgebras. For most of application, we need a stronger notion than being flavored, which restricts \bar{F}_0C , but weaker than being QT-connected.

Definition 4.12. A flavored coalgebra is said to be *pathlike* if

- (1) $sgpl(C) = gpl(C)$
- (2) $\bar{F}_0C = \bigoplus_{g \in gpl(C)} Rg$

We say a pathlike coalgebra is split, if F_0C is a direct summand, i.e. the sequence of R modules $0 \rightarrow F_0C \rightarrow C \rightarrow C/F_0C \rightarrow 0$ is split.

A bi- or Hopf-algebra is pathlike or split pathlike if the underlying coalgebra is.

Pathlike coalgebras are the right type for a generalization of path coalgebras to a categorical setting, see §6.

Remark 4.13.

- (1) At the moment, we will disregard semigrouplike elements that are not grouplike. Handling then latter elements is more tricky, they only appear in torsion, but they might be of interest in the situation with modified counits, [GCKT20b, 2.2] and for applications like [KMM20].
- (2) The freeness assumption of \bar{F}_0C mimics the fact that in the case of a field for a pointed coalgebra, the coradical filtration satisfies $C_0 = \bigoplus_{g \in gpl} k[g]$.

Example 4.14. The path algebra $C = R(P(Q))$ of a quiver is pathlike. The $sgpl = gpl = V(Q)$, the set of vertices. $Rv \cap Rw = \delta_{v,w}Rv$. A path of length n from v to w is in $F_nC_{v,w}^Q$ and hence the filtration is exhaustive. Finally, $C_0F^qC = \bigoplus_{v \in V} Rv$. Note that $C_1C_{g,h}^Q = R\vec{e} + R(g-h)$, if there is a directed edge \vec{e} from g to h .

More generally:

Proposition 4.15. *A simply colored algebra is flavored. Hence, the dual of a colored algebra is flavored. It is also pathlike if it is locally decomposition finite.*

Proof. Straightforward from the definitions. \square

Theorem 4.16. *Let C be pathlike coalgebra and A algebra and $f \in \text{Hom}(C, A)$. If $\text{Ext}(C/\bar{F}_0 C^Q, A) = 0$, then f has a \star -inverse in the convolution algebra $\text{Hom}(C, A)$ if and only if for every grouplike element g , $f(g)$ has an inverse as ring element in A . In particular every grouplike central character is \star -invertible.*

Proof. Using the filtration $\bar{F}_r C$ for a flavored coalgebra, by Theorem 3.28 f having a \star -inverse is equivalent to the restriction of f on $\sum_{g \in \text{gpl}(C)} Rg = \bigoplus_{g \in \text{gpl}(C)} Rg, A$ considered as element in $\text{Hom}(\bigoplus_{g \in \text{gpl}(C)} Rg, A)$ having a star inverse.

If for every $g \in \text{gpl}(C)$ every $f(g)$ has an inverse, we can define $h(g) = f(g)^{-1}$ yielding a convolution inverse as $(f \star h)(g) = f(g)h(g) = 1_A = \eta_A(\epsilon_C(g))$ and symmetrically. On the other hand, if f has a convolution inverse h , then for $g \in \text{gpl}(C)$ $(f \star h)(g) = f(g)h(g)\eta_A(\epsilon_C(g)) = 1_A$ and likewise $h(g)f(g) = 1_A$, so that $h(g) = f(g)^{-1}$. \square

4.4. Pathlike bi- and Hopf algebras. Pathlike bi- and Hopf algebras enjoy several properties, similar to those of pointed coalgebras.

Corollary 4.17. *A pathlike bialgebra B is a Hopf algebra if and only if the set of grouplike elements a group under multiplication, or equivalently they form a set-like bialgebra.*

Proof. Using Corollary 3.30, and the Theorem above, having an antipode on B is equivalent to having an antipode on $F_0^Q C = \bigoplus_{g \in \text{gpl}} Rg$. The only possible antipode on grouplike elements is $S(g) = g^{-1}$ if the inverses exist. Hence the existence is equivalent to the existence of inverses. \square

Remark 4.18. Notice that for a bialgebra B switching both multiplication and comultiplication, that is $B^{op, cop}$ in the usual notation, yields a bialgebra and in the Hopf case a Hopf algebra. The opposite or coopposite of Hopf algebra alone may not be a Hopf algebra, however.

Corollary 4.19. *The antipode of a split pathlike Hopf algebra H is bijective.*

Proof. Using Example 4.10 and Theorem 4.16, it follows that H^{op} is also Hopf algebra. Let T be the antipode for H^{op} and S the antipode for H . Then for $h \in H$, we have $h_{(2)}T(h_{(1)}) = \epsilon(h)1_H = T(h_{(2)})h_{(1)}$ (notice the difference from axiom of normal antipode). Applying S and using the fact S is an algebra antihomomorphism, it follows that

$$(S \circ T)(h_{(1)})S(h_{(2)}) = \epsilon(h)1_H = S(h_{(1)})(S \circ T)(h_{(2)})$$

We see $(S \circ T)$ is both a left and a right convolution inverse to S . Thus, $S \circ T = \text{Id}_H$ since convolution inverse is unique. Replacing h by $S(h)$ in $h_{(2)}T(h_{(1)}) = \epsilon(h)1_H = T(h_{(2)})h_{(1)}$, it follows that

$$S(h_{(1)})(T \circ S)(h_{(2)}) = \epsilon(h)1_H = (T \circ S)(h_{(1)})S(h_{(2)})$$

Thus, $T \circ S = \text{Id}_H$, and we conclude that S is a bijection. \square

Remark 4.20. This means that there is a Drinfel'd double for Hopf pathlike Hopf algebras, see e.g. [Kas95, IX,4].

4.4.1. Connected quotient and Brown coaction. The following generalizes the coaction of [Bro15] to pathlike bialgebras. Let B be a pathlike bialgebra, then the ideal spanned by $F_0^Q(B) = \bigoplus_{g \in \text{gpl}(B)} Rg$ is a coideal and $B^{\text{red}} = B/F_0^Q B$ is a Hopf algebra.

Corollary 4.21. B^{red} is a Hopf algebra.

Proof. This follows from Corollary 4.11. \square

Proposition 4.22. Let $\pi : B \rightarrow B^{\text{red}}$ be the projection. The Brown map

$$\Delta_B : B \rightarrow B \otimes_R B^{\text{red}}, \quad \Delta_B(x) = \sum x_{(1)} \otimes \pi(x_{(2)}) \quad (4.1)$$

is a coaction.

Proof. The result of both iterations is $\sum x_{(1)} \otimes \pi(x_{(2)}) \otimes \pi(x_{(3)})$. \square

Example 4.23. The example of Brown is essentially the one based on (the skeleton of) $OI_{*,*}$, see 6.1.2. In Brown's notation, up to extra decoration, we have $\mathcal{H} = B = B^{\text{comm}}(OI_{*,*})$ and $\mathcal{A} = B^{\text{red}} = \mathcal{H}/(\zeta(2)^{\mathfrak{m}}$ and he uses the map Δ^{op} and hence switches the factors of \mathcal{H} and \mathcal{A} and $\zeta(2)^{\mathfrak{m}}$ is the grouplike element corresponding to $\text{id}_{[1]}$.

4.5. Further examples of flavored coalgebras. We now give further examples of classes of flavored algebras that appear naturally.

4.5.1. Flavored coalgebras generated by grouplike and skew primitives. Similar to pointed coalgebra over k , a bialgebra generated by grouplike and skew primitive elements is flavored. In the categorical setting this lets us generalize from quivers to presentation in terms of irreducibility, see §6 for the application to Feynman categories.

To prove the result, we need the following lemma.

Lemma 4.24. Let B be a bialgebra and g, h, p, q be grouplikes and x be (p, q) -skew primitive, then $pC_{g,h}^Q \subset C_{pg,ph}^Q$ and $xC_{g,h}^Q \subset C_{pg,qh}^Q$.

Proof. In fact, $pF_n B_{g,h}^Q \subset F_n B_{pg,ph}^Q$ and $xF_n B_{g,h}^Q \subset F_{n+1} C_{pg,qh}^Q$. We prove these relations by induction. For the base case, we use the fact that the multiplication of two grouplike elements is grouplike and the multiplication of a grouplike element and a skew primitive element is skew primitive. In particular, we have gx is a (gp, gq) -skew primitive element. Now we assume $pF_{n-1} B_{g,h}^Q \subset F_{n-1} B_{pg,ph}^Q$. Observe for $z \in F_n B_{g,h}^Q$ and $\bar{\Delta}_{g,h}(z) = \sum z_{(1)} \otimes z_{(2)} \in \sum_{t \in \text{sgpl}(C)} F_{n-1} B_{g,t}^Q \boxtimes F_{n-1} B_{t,h}^Q$. By algebraic manipulations, it follows

$$\bar{\Delta}_{pg,ph}(pz) = \sum pz_{(1)} \otimes pz_{(2)}$$

where $\bar{\Delta}_{pg,ph}(pz) \in \sum_{t \in \text{sgpl}(C)} pF_{n-1} B_{g,t}^Q \boxtimes pF_{n-1} B_{t,h}^Q$. By induction hypothesis: $\bar{\Delta}_{pg,ph}(pz) \in \sum_{t \in \text{sgpl}(C)} F_{n-1} B_{pg,pt}^Q \boxtimes F_{n-1} B_{pt,ph}^Q$. Therefore, $pz \in$

$F_n B_{pg,ph}^Q$ and $pF_n B_{g,h}^Q \subset F_n B_{pg,ph}^Q$. Now, we assume $xF_{n-1} B_{g,h}^Q \subset F_n C_{pg,qh}^Q$. Observe

$$\bar{\Delta}_{pg,qh}(xz) = xg \otimes qz + \sum pz_{(1)} \otimes xz_{(2)} + \sum xz_{(1)} \otimes qz_{(2)} + pz \otimes xh$$

so

$$\begin{aligned} \bar{\Delta}_{pg,qh}(xz) \in F_1 B_{pg,qg}^Q \boxtimes qF_n B_{g,h}^Q + \sum_{t \in \text{sgpl}(C)} pF_{n-1} B_{g,t}^Q \boxtimes xF_{n-1} B_{t,h}^Q + \\ \sum_{t \in \text{sgpl}(C)} xF_{n-1} B_{g,t}^Q \boxtimes qF_{n-1} B_{t,h}^Q + pF_n B_{g,h}^Q \boxtimes F_1 B_{ph,qh}^Q \end{aligned}$$

Using the previous result for grouplike elements and Lemma 3.12:

$$\begin{aligned} \bar{\Delta}_{pg,qh}(xz) \in F_n B_{pg,qg}^Q \boxtimes F_n B_{qg,qh}^Q + \sum_{t \in \text{sgpl}(C)} F_n B_{pg,pt}^Q \boxtimes F_n B_{xt,xh}^Q + \\ \sum_{t \in \text{sgpl}(C)} F_n B_{xg,xt}^Q \boxtimes F_n B_{qt,qh}^Q + F_n B_{pg,ph}^Q \boxtimes F_n B_{ph,qh}^Q. \end{aligned}$$

Hence, $xz \in F_{n+1} B_{pg,qh}^Q$. \square

Theorem 4.25. *If a bialgebra B is generated by grouplike and skew primitive elements, then it is flavored.*

Proof. (NB: Sweedler notation is not used in this argument) Observe that if g, h are grouplike elements and x a skew primitive element, then gh is grouplike and gx is a skew primitive. Therefore, we focus on the products of skew primitives $x_n \cdots x_1$. Observe that if x is a (g, h) -skew primitive and x' is a (g', h') -skew primitive. It follows

$$\Delta(xx') - (gg' \otimes xx' + xx' \otimes hh') = xg' \otimes hx' + gx' \otimes xh'$$

where xg' is a (gg', hg') -skew primitive, hx' a (hg', hh') -skew primitive, gx' is a (gg', gh') -skew primitive and xh' is a (gh', hh') -skew primitive. Therefore, $\Delta(xx') - (gg' \otimes xx' + xx' \otimes hh') \in F_1 C_{gg',hg'} \boxtimes F_1 C_{hg',hh'} + F_1 C_{gg',gh'} \boxtimes F_1 C_{gh',hh'}$ and $xx' \in F_2 C_{gg',hh'}$. We claim every words of skew primitive elements $x_n \cdots x_1$ belongs to a component. We proceed by induction. The above observation forms a base case. Suppose $x_1 \cdots x_{n-1}$ belongs to some component for every possible product of skew primitive elements. Using previous lemma and considering $x_n(x_{n-1} \cdots x_1)$, we are done. \square

Example 4.26. Consider a free R -algebra generated by x and t . We give a coalgebra structure by giving two maps Δ and ϵ defined by (and extended linearly by) $\Delta(x) = 1 \otimes x + x \otimes t$, $\epsilon(x) = 0$ and $\Delta(t) = t \otimes t$, $\epsilon(t) = 1$. We will call this bialgebra B . Let I be a two-sided ideal generated by $t^2 - 1, x^2, xt + tx$. This is a coideal so it becomes an bi-ideal. Let $H = B/I$ be the quotient bialgebra, then H as a module is isomorphic to $R1 \oplus Rx \oplus Rt \oplus Rxt$. We see $H = H_{1,1}^Q + H_{1,t}^Q + H_{t,1}^Q + H_{t,t^2}^Q$.

Example 4.27. The path algebra of a quiver is generated by grouplike elements and skew primitives. Indeed the generators are the id_v which are grouplike and the \vec{e} which are skew primitive.

Example 4.28. Consider the monoid \mathbb{Z}^\times . The coproduct is given by $\Delta(n) = \sum_{n_1, n_2 = n} n_1 \otimes n_2$. Then this is generated by skew primitives, namely the prime numbers.

Corollary 4.29. *In particular:*

- (1) *The coalgebra of a locally finite category generated by irreducibles is pathlike. This includes the examples of quiver and incidence coalgebras.*
- (2) *The bialgebra $B(\mathcal{F})$ stemming from a non-symmetric Feynman category is pathlike if \mathcal{F} is generated by irreducibles as these are skew primitives, cf. §2.4.4. A necessary condition is that \mathcal{V} is discrete, cf. §2.4.4. This includes the simplicial examples as well as the Connes-Kreimer bialgebra of planted planar forests, see §6.2.1.*
- (3) *The bialgebra B^{iso} coming from a Feynman category with is pathlike, if it is generated by essentially skew primitives, cf. §2.4.4. This includes those coming from operads such as the Connes-Kreimer rooted labelled forest bialgebra, see §6.2.2 as well as those coming from graphs, like the core bialgebra for graphs with legs, see §6.4.*

Other concrete examples and details are given in 6.

4.5.2. Component and looplike coalgebras. One motivational example for the results on antipodes is given by component graded coalgebras, cf. [MP12], whose definitions we recall here.

Definition 4.30.

- (1) A graded augmented algebra A is said to be *grouplike* if the set $\epsilon^{-1}(1)$ of degree 0 elements is a group under the product of A .
- (2) Let C be a graded coalgebra such that C_0 is R -free and define $\pi C = \{g | \Delta(g) = g \otimes g \text{ and } g \neq 0\} \subset C_0$. For $g \in \pi C$, $g = \epsilon(g)g$ by the counit property and thus $\epsilon(g) = 1$ since C_0 is assumed to be k -free. Define the component C_g of g by letting $C_g = Rg \oplus \bar{C}_g$, where the R -module \bar{C}_g of positive degree elements of \bar{C}_g is

$$\{x | \Delta(x) = x \otimes g + \sum x_{(1)} \otimes x_{(2)} + g \otimes x, \deg x_{(1)} > 0 \text{ and } \deg x_{(2)} > 0\}$$

- (3) Say that C is a *component graded coalgebra* if C_0 is R -free, each C_g is a subcoalgebra of C , and C is the direct sum of the C_g .

Not that there is a grading and a grading by components.

Example 4.31. This definition is motivated by $H_*(X, R)$ and $H_*(\Omega X, R)$, where X is a based space and R is a ring, see [MP12]. If both modules are R -flat, the first homology ring is a unital component coalgebras and the second ring is a grouplike component Hopf algebra. Moreover, the condition

for second ring to be connected in the algebra sense is equivalent to X being connected in the topology sense.

Proposition 4.32. *A grouplike component bialgebra is a Hopf algebra.*

Proof. There are two proofs, the first uses the coalgebra grading and a version of Takeuchi's lemma for this filtration. We can alternatively construct a QT-sequence. In fact, the component coalgebra can be treated a direct sum of coaugmented conilpotent graded coalgebras except that there are different choices for the coaugmentations. One can see directly there is a filtration $F_n C = \bigoplus_{g \in \text{gpl}(C)} \bigoplus_{i=0}^n C_g^{(i)}$. It is clear that $F_n C \subset F_{n+1} C$ and this filtration is exhaustive. In addition, the inclusion $F_n C \subset C$ split so $F_n C \otimes F_n C \subset C \otimes C$. If $x \in F_n C$, $x = \sum_g x_g$ with $x_g \in C_g^{(n)}$. By definition $\Delta(x_g) = g \otimes x_g + x'_g \otimes x''_g + x_g \otimes g$ s.t. $x'_g, x''_g \in \bigoplus_{i=0}^n C_g^{(i)}$. Hence, $\Delta(x) \in C_0 \otimes C + F_{n-1} C \otimes F_{n-1} C + C \otimes C_0$. F_n is then a QT-sequence, and the issue of antipode is reduced to investigate the structure of $F_0 C = \bigoplus_{g \in G} Rg$. \square

One can generalize the definition of a component coalgebra by dropping the assumption of grading. We start with the observation $F_n C_{g,g}^Q = \bigoplus_{i=0}^n C_g^{(i)}$ where $C_g^{(n)}$ means component of grade n of C_g in the perspective of bivariate Quillen component. In fact C_g is a Quillen connected coalgebra. This is captured by the following definition of looplike coalgebra.

Definition 4.33. A coalgebra C over R is said to be *looplike* if it has a decomposition into subcoalgebras,

$$C = \bigoplus_{g \in \text{gpl}(C)} C_g \quad (4.2)$$

where each C_g is Quillen connected subcoalgebra for a choice of coaugmentation. A *looplike* coalgebra is such a coalgebra with a fixed choice of coaugmentations.

Proposition 4.34. *A looplike coalgebra is a pathlike coalgebra. And hence, a looplike is bialgebra is Hopf the grouplike elements form a group.*

Proof. Observe that $C_{g,g}^Q \supset C_g$. \square

Remark 4.35. Counterintuitively, even for a looplike coalgebra, the bivariate Quillen components contains a non-zero $C_{g,h}^Q$ as $g - h$ is a (g, h) -skew primitive, as mentioned above, see Remark 4.5.

Remark 4.36. The name looplike stems from the fact that when k is a field such a coalgebra is exactly a path coalgebra for a category that is totally disconnected, that is for any two objects $X \neq Y$, $\text{Hom}(X, Y) = \emptyset$ and hence all paths of composable morphisms are loops, viz. they have the same source and target.

In the particular case of a quiver, the vertex set V is indexed by grouplike element g . For the arrow set Q , we set that no arrow between any distinct

vertices i.e. $Q(x, y) = \emptyset$ if $x \neq y \in V$ and $Q(x, x)$ is underlying set of a free group generated by one element.

5. RENORMALIZATION, QUOTIENTS AND LOCALIZATION

The obstruction to having a Hopf algebra structure on a pathlike bialgebra, is the invertibility of the grouplike elements. There are basically two approaches to remedy this perceived deficiency. One is adding formal inverses, which is possible by universal constructions But, as at the end of the day for renormalization in à la Connes–Kreimer one actually only needs convolution inverses for characters, there is another option. This is to restrict the target and place restrictions on the characters. For instance that the target algebra is commutative, or that the characters restricted to grouplike elements have special properties. In this case there are universal quotients that the characters factor through. We first briefly explain the setup to give the motivation for these constructions.

5.1. Renormalization. A renormalization scheme in the Connes–Kreimer formalization of the BPHZ renormalization [CK00], is defined on the convolution algebra of a bialgebra with a Rota-Baxter algebra. see also [MN16, DEEFG16, EFK05, Guo12].

5.1.1. Rota-Baxter operators and algebras.

Definition 5.1. A Rota–Baxter (RB) operator of weight λ on a associative R -algebra A is a linear operator $T : A \rightarrow A$ that satisfies the equation

$$T(X)T(Y) = T(T(X)(Y)) + T(XT(Y)) + \lambda T(XY) \quad (5.1)$$

A Rota-Baxter (RB) algebra is a unital associative algebra with a choice of Rota–Baxter operator.

Note that if T is an RB-operator, so is $(1 - T)$, furthermore (5.1) says that $Im(T)$, and similarly $Im(1 - T)$, are subalgebras of A , which will be called A_+ and A_- as $(1 - T) + T = 1$, we see that $A = A_+ + A_-$.

NB: This sum does not need to be direct.

The pair (A_+, A_-) is called the *Birkhoff decomposition* of A defined by T .

Remark 5.2 (Scaling). Scaling the RB operator by μ , we see that μT is a RB algebra of weight $\lambda\mu$, so that if $\lambda \in R^*$ is invertible, we can always scale to have $\lambda = 0$ or $\lambda = -1$.

Example 5.3. The standard example are Laurent series $k[x, x^{-1}]$ with T the projection onto the pole part is an RB-algebra with $\lambda = -1$.

A generalization of this is uses projectors, which are operators with $T^2 =$. In this case, we also have $1 - T$ is a projector and these are orthogonal $T(1 - T) = 0$. For a projector T on an algebra A , set $(1 - T)A = A_+$ and $T(A) = A_-$ then $A = A_+ \oplus A_-$.

Proposition 5.4 (Projectors and subalgebras). *Given a projector T , if A_+ and A_- are subalgebras then T is an RB-operator with weight $\lambda = -1$. Furthermore, $T(1)$ is an idempotent, $T(1)^2 = T(1)$, and multiplication by $T(1)$ is a projector onto a subspace of A_- .*

In particular, if $T(1) = 0$, then A_+ is a unital subalgebra, and, if $T(1)$ is regular, then $T(1) = 1$ and A_- is a unital subalgebra.

Proof. The conditions means that $(1 - T)$ and T are the projectors onto $A = A_+ \oplus A_-$. Since A_- is a subalgebra, we have that

$$0 = (1 - T)[T(X)T(Y)] = T(X)T(Y) - T(T(X)T(Y)) \quad (5.2)$$

Since A_+ we have that

$$\begin{aligned} 0 &= T[(1 - T)(X)(1 - T)(Y)] = \\ &\quad T(XY) - T(T(X)Y) - T(XT(Y)) + T(T(X)T(Y)) \end{aligned} \quad (5.3)$$

Plugging (5.2) into (5.3), we obtain (5.1) for $\lambda = -1$. Plugging in $X = Y = 1$ and using that $T^2(1) = T(1)$ we obtain that $T(1)^2 = T(1)$ and hence multiplication by $T(1)$ is a projector. Plugging in $Y = 1$ we obtain $T(1)(1 - T)(X) = 0$. \square

Remark 5.5.

- (1) If A is regular, e.g. a domain, then there are only two possibilities $T(1) = 0$ or $T(1) = 1$. Thus after possibly switching T and $1 - T$, we have that $A = A_+ \oplus A_-$ with A_+ a unital subalgebra and A_- a non-unital subalgebra.

Vice-versa an RB-operator for $\lambda = -1$, comes from a subdirect sum.

Definition 5.6. [Atk63] Two subalgebras (A_-, A_+) form subdirect sum of A if there exists a subalgebra $C \in A \times B$ which satisfies that for any $a \in A$ there is a unique representation $a = a_- + a_+$, with $(a_-, a_+) \in C$.

Given such a decomposition defines $T(a) = a_-$ and checks that it satisfies (5.1) for $\lambda = -1$. Vice-versa given an RB T with $\lambda = -1$, setting $A_- = T(A)$, $A_+ = (1 - T)(A)$, $C = \{(T(a), (1 - T)(a)) : a \in A\} \subset A_- \times A_+$ gives the subdirect sum.

Theorem 5.7. [Atk63] *The set of RB-operators T with weight $\lambda = -1$ is in bijection with the subdirect sum decomposition of A under the above correspondence.*

5.1.2. Birkhoff decomposition and renormalization for a Feynman rule. A Feynman rule is an a character in the convolution algebra $\phi \in \text{Hom}_{R\text{-alg}}(B, A)$ the renormalization of ϕ based on an Rota–Baxter (RB)–operator is a pair of characters $\phi_{\pm} \in \text{Hom}_{R\text{-alg}}(B, A_{\pm})$ such that $\phi = \phi_-^{-1} \star \phi_+$. The character ϕ_+ is then the renormalized Feynman rule. It was shown in [CK00, CK01] that there is a unique such decomposition for the Connes–Kreimer Hopf algebra of graphs and a commutative RB algebra, if ϕ is the exponential of an infinitesimal character. The solution is

then given by a recursive formula. This was generalized in [EFGK05] for B any connected bialgebra and a character that is the exponential of an infinitesimal character. The following analysis provides the background for a generalization of these results to the case of a bialgebra with a QT-sequence.

Remark 5.8.

- (1) If B is a Hopf algebra, then the inverse is given by $\phi_-^{-1} = \phi_- \circ S$, see Proposition 3.6.
- (2) If ϕ is grouplike central then it passes through the simpler construction $B_{\mathbf{q}}$, see §5.2.1.
- (3) if A is commutative, then ϕ passes through the quotient B^{com} , see §5.2.1 and Proposition 3.6 applies.

5.2. Quotients. We have already seen that taking the somewhat drastic quotient by $F_0^Q(C)$ makes a flavored coalgebra connected. There are, however intermediary quotients, which make characters with special properties invertible.

Definition 5.9. We say a character is *grouplike invertible*, if for every grouplike element $g: \phi(g) \in A^\times$, *grouplike central* if for all $g \in gpl: \phi(g) \in Z(A)$, *grouplike scalar* if $\phi(g) \in R^\times$ and *grouplike normalized* if $\phi(g) = 1$ for all grouplike g .

Recall that in a graded setting, the grouplike elements are in degree 0, and if A is graded and ϕ preserves the grading, then the grouplike elements have to land in A_0 . It is quite common, that even if A is not commutative, A_0 is. In several application the characters are scalar and take values $(\frac{1}{2\pi i})^k$. This is also a form of grading.

5.2.1. Quotients for “grouplike” characters. The previous paragraph and the natural examples, see e.g. [CK98, CK00, CK01, EFK05, MN16], motivates the study of characters that have special properties.

Proposition 5.10. *A grouplike normalized character factors through the quotient B/I_N where I_N is the ideal spanned by $1 - g$ for $g \in gpl(B)$.*

Proof. We need to check that I_N is a coideal. Indeed $\Delta(1 - g) = 1 \otimes 1 - g \otimes g = (1 - g) \otimes (1 + g)$ and $\epsilon(1 - g) = 0$, since g is grouplike. Furthermore if ϕ is normalized the $\phi(1 - g) = 0$ so that $I \subset \ker(\phi)$. \square

NB: $R(g - h) \subset I$ since $g - h = (1 - h) - (1 - g) = g - h$. This is the line which keeps the sum in a pathlike bialgebra from being direct, see Remark 4.5.

Proposition 5.11. *Let $J = ([B, B])$ the ideal spanned by commutators. Then J is a codideal, and If A is commutative, then any character $\phi \in \text{Hom}_{R\text{-alg}}(B, A)$, factors through $B^{com} = B/([B, B])$.*

Proof. J is a coideal: $\Delta(ab - ba) = \sum a_{(1)}b_{(1)} \otimes a_{(2)}b_{(2)} - b_{(1)}a_{(1)} \otimes b_{(2)}a_{(2)} = (a_{(1)}b_{(1)} - b_{(1)}a_{(1)}) \otimes a_{(2)}b_{(2)} + b_{(1)}a_{(1)}(a_{(2)}b_{(2)} - b_{(2)}a_{(2)})$ and $\epsilon(ab - ba) = 0$. Since $\phi: I \subset \ker(\phi)$, the statement follows. \square

There is a further quotient that is of interest which was studied in [GCKT20a]. Consider the ideal I_C generated by $(ag - ga)$ for $a \in B, g \in gpl(B)$.

Proposition 5.12. *I_C is a coideal and the bialgebra $B/I_C \simeq B_{\mathbf{q}}/I$, where $B_{\mathbf{q}}$ is defined below. Furthermore any grouplike central character factors through B/I_C .*

Proof. The fact that I_C is a coideal follows as in Proposition 5.11. For and grouplike central character, $\phi(ag - ga) = 0$, so $\ker(\phi) \supset I_C$. \square

5.2.2. Q(uantum) deformation and coaction. For a grouplike invertible central characters, there is a universal construction which can be viewed as a quantum deformation and gives rise to a Brown type coaction. Consider the algebra $B_{\mathbf{q}} = B[q_g, q_g^{-1}] : g \in gpl$, of Laurent polynomials with coefficients in the possibly noncommutative B . Note the q_g lie in the center of $B_{\mathbf{q}}$. Endow $B_{\mathbf{q}}$ with the bialgebra structure where the q_g, q_g^{-1} are grouplike. Consider the ideal I generated by $q_g - g$.

Proposition 5.13. *I is a coideal and any grouplike central character ϕ can be lifted to a character $\hat{\phi} : B_{\mathbf{q}} \rightarrow A$ and $\hat{\phi}$ factors through the quotient $B_{\mathbf{q}}/I$. Embedding $B \rightarrow B_{\mathbf{q}} \rightarrow B_{\mathbf{q}}/I$ the $\bar{\phi}(b) = \phi(b)$.*

Proof. By definition $\epsilon(q_g - g) = 1 - 1 = 0$ and $\Delta(q_g - g) = q_g \otimes q_g - g \otimes g = (q_g - g) \otimes (q_g + g)$, so I is a coideal. The lift is given by $\hat{\phi}(q_g) = \phi(g)$ and consequentially $\hat{\phi}(q_g^{-1}) = \phi(g)^{-1}$. The other statements follow from $\phi(g) = \hat{\phi}(q_g)$. \square

Note that the image of g in $B_{\mathbf{q}}/I$ lies in the center, since q_g does.

Remark 5.14. If the character is only grouplike invertible it will likewise factor through the noncommutative Laurent polynomials.

- (1) The bialgebra $B_{\mathbf{q}}$ can be viewed as a multi-parameter quantum deformation. We have that as $q_g \rightarrow 1$ $B_{\mathbf{q}} \rightarrow B$.
- (2) If e is not the only grouplike There is a single parameter deformation given by $B_q = B[q, q^{-1}]$ which is the quotient $B_{\mathbf{q}}/J$, where J is generated by the $(q_g - q_h)$, $g, h \in gpl, g, h \neq e$.
- (3) The polynomial rings $B[gpl]$ and $B[q]$ are sub-bialgebras and I is a coideal for this sub-bialgebra as well. Furthermore $B_{\mathbf{q}}/I \simeq B/I_C$ which is the universal quotient for grouplike central characters. This was the constuction considered in [GCKT20a, §2.6] for a bialgebra coming from a pointed cooperad with mutliplication.
- (4) The factors of q give a natural grading, which is important for the infinitesimal versions, see [GCKT20a, §2.7]

This quantum deformation also lets us make the coaction of Brown, see §4.4.1 commutative and give a nice interpretation in terms of polynomials and Laurent series,

Definition 5.15. The commutative Hopf coaction is given by

$$\Delta_B : B_{\mathbf{q}}/I \rightarrow B_{\mathbf{q}}/I \otimes B^{\text{red}}, \Delta_B(x) = \sum x_{(1)} \otimes \pi(x_{(2)}) \quad (5.4)$$

where now both factors are Hopf algebras and π simply sets the factors $q_g = 1$ on the left to 1.

Proposition 5.16. *Let B be pathlike such that $\bigoplus_{g \in \text{gpl}}$ is a free subalgebra generated by elements $g_j, j \in J$, then $\mathcal{B}_{\mathbf{q}}/I \simeq B^{\text{red}} \otimes_R R[q_g, q_g^{-1}] \simeq B^{\text{red}}[q_{g_j}, q_{g_j}^{-1}]$ and similarly for the polynomial subrings.*

Proof. The correspondence is given by collecting the factors of q_g respectively q_g^{-1} on the right, as they are central, and expressing them in terms of the $q_{g_j}, q_{g_j}^{-1}$. \square

Remark 5.17. For the polynomial version in the operad case, the isomorphism in the proposition was realized as the quotient by the ideal $(| - q)$ in [GCKT20a, §2.6], see also §6.2. The proposition can also be used to describe the coaction restricted to the quotient $B[\text{gpl}]/I$ in the case of a free $F_0^Q B$ subalgebra of a pathlike algebra, generalizing Example 4.23,

5.3. Adding formal inverses. In general one can formally invert the grouplike elements, this is a universal construction called the group completion if one is speaking of a monoid or the localization if one is inverting a multiplicative subset. In the current R -linear context this becomes a subalgebra.

Given a subalgebra S of an R algebra A , the localization at S , S^{-1} is an object, together with a morphism $j : A \rightarrow S^{-1}A$ with the universal property that if $f : A \rightarrow A'$ is an algebra homomorphism such that for all $s \in S$, $f(s)$ is invertible then f factors as $g \circ j$, with $g : S^{-1}A \rightarrow A'$. This is an abstract construction the general case, but it is not effective if there are no other conditions. This can be realized as a quotient of the free algebra of R -linear alternating words in A and S . Assuming that A is unital, set

$$T_R(A \otimes S^{\text{op}}) = R \oplus A \otimes S^{\text{op}} \oplus A \otimes S^{\text{op}} \otimes A \otimes S^{\text{op}} \oplus \dots \quad (5.5)$$

which is an algebra by concatenation. The unit is 1_R , where we identify $R \otimes_R M = M$ for any R -module M . Then $AS^{-1} = (A \otimes S^{\text{op}})/I$, where I is the ideal spanned by the following relations

- (1) $1 \otimes \bar{1} = 1_R, \bar{1} \otimes 1 = 1_R$ which makes $1 \otimes \bar{1}$ a unit.
- (2) $s \otimes \bar{s} = 1 \otimes \bar{1}$, which inverts the elements of s ,
- (3) $a \otimes \bar{1} \otimes a' \otimes \bar{s} = aa' \otimes \bar{s}$
- (4) $a \otimes \bar{s} \otimes 1 \otimes \bar{s}' = a \otimes \bar{s}\bar{s}'$.

where we write \bar{s} to indicate the factors of S^{op} . In this notation $\overline{s_1 s_2} = \bar{s}_2 \bar{s}_1$ by the definition of the opposite multiplication. We let $\epsilon(a \otimes \bar{s}) := \epsilon(a)\epsilon(s)^{-1}$.

If A is also a bialgebra and S is also semigrouplike and $\epsilon(S) \subset R^*$ then we define the comultiplication:

$$\Delta(a \otimes \bar{s}) = \sum (a' \otimes \bar{s}) \otimes (a_{(2)} \otimes \bar{s}) \quad (5.6)$$

This fixes the comultiplication on the free algebra via the bialgebra equation. Defining the counit as $\epsilon(a \otimes \bar{s}) = \epsilon(a)\epsilon(s)^{-1}$, defines the counit on the free algebra.

Proposition 5.18. *I is a coideal and hence AS^{-1} is a bialgebra. Similarly one can define a left localization $S^{-1}A$ by using $T_R(S^{op} \otimes A)$.*

Proof. This is a straightforward calculation:

- (1) $\Delta(1 \otimes \bar{1} - 1_R) = 1 \otimes \bar{1} \otimes 1 \otimes \bar{1} - 1_R \otimes 1_R = (1 \otimes \bar{1} - 1_R) \otimes (1 \otimes \bar{1} + 1_R) \in I \otimes A.$
- (2) $\Delta(s \otimes \bar{s} - 1_R) = (s \otimes \bar{s}) \otimes (s \otimes \bar{s}) - 1_R = (s \otimes \bar{s} + 1_R) \otimes (s \otimes \bar{s} - 1_R) \in A \otimes I$
- (3) $\Delta(a \otimes \bar{1} \otimes a' \otimes \bar{s} - aa' \otimes \bar{s}) = \sum a_{(1)} \otimes \bar{1} \otimes \sum a'_{(1)} \otimes \bar{s} \otimes a_{(2)} \otimes \bar{1} \otimes a'_2 \otimes \bar{s} - \sum a_{(1)} a'_{(1)} \otimes \bar{s} \otimes a_{(2)} a'_2 \otimes \bar{s} \equiv 0 \text{ mod } (I \otimes A + A \otimes I)$
- (4) $\Delta(a \otimes \bar{s} \otimes 1 \otimes \bar{s}' - a \otimes \bar{s} \bar{s}') = \sum a_{(1)} \otimes \bar{s} \otimes 1 \otimes \bar{s}' \otimes a_{(2)} \otimes \bar{s} \otimes 1 \otimes \bar{s}' - a_{(1)} \otimes \bar{s} \bar{s}' \otimes a_2 \otimes \bar{s} \bar{s}' \equiv 0 \text{ mod } (I \otimes A + A \otimes I)$

For the counit, it is a straightforward check that $\epsilon(I) \equiv 0$. \square

5.4. Calculus of fractions. There is a more concise version of this construction if the Ore condition and cancellability are met. This gives a calculus of fraction, similar to the calculus of roofs [GM03]. The right Ore condition states that

$$\forall a \in A, s \in S, \exists a' \in A, s' \in S : as = s'a' \quad (5.7)$$

One also needs cancellability, or S regularity. That is if $as = 0$ or $sa = 0$ then $a = 0$. This means that $sa = sb$ implies $a = b$ as does $as = bs$.

Proposition 5.19. *If these conditions are met any element of AS^{-1} can be written as as^{-1} and an injective algebra homomorphisms $A \rightarrow AS^{-1}$, such that every element of s is invertible. Also the left and right fractions become isomorphic.*

Proof. See [Art99]. \square

Remark 5.20.

- (1) For a free non-commutative algebra $R\{x, y\}$ and $S = \langle y \rangle$ the condition is not met.
- (2) For the Weyl algebra $R\{x, y\}/(xy - yx = 1)$ the condition is met [Art99].
- (3) In similar spirit, if the product is the tensor product the Ore condition is usually not met, as $X \otimes Y$ is rarely $Y \otimes X$. In particular in the case of interest, where one needs to invert the identities id_X one would need a Y and a morphism g such that for a given morphism $f \otimes id_X = id_Y \otimes g$.
- (4) In the commutative case this always holds. In particular this holds for the symmetric monoidal setting when using coinvariants \mathcal{B}^{iso} .
- (5) For a QT-sequence, one does not need full commutativity, but only a central character. In this case, one can alternatively also invert the ideal making the grouplike elements central.

- (6) A further route of exploration is to use higher homotopy commutativity, i.e. an E_1 for the Ore condition.

Lemma 5.21. *In the special case of $B[\text{gpl}(B)]$ the multiplicative set S spanned by the q_g is Ore, since these elements are central, and the localization $B[\text{gpl}(B)]S^{-1}$ is the Laurent-series ring $B_{\mathbf{q}}$.*

Proof. The isomorphism is given by $\bar{q} \rightarrow q_g^{-1}$ which exists by the universal property. \square

Remark 5.22 (Abelian case, Grothendieck construction). In the fully commutative case, this concretely boils down to the Grothendieck construction and localization. Given a monoid (M, \cdot) , the Grothendieck construction gives the group completion. $K(M) = M \times M / \sim$ where $(m, n) \sim (m', n')$ if $m \cdot n' = m' \cdot n$. There is an injection $M \rightarrow K(M)$ given by $m \mapsto (m, 1)$, with inverses given by $(0, m)$. More generally if R is a commutative ring and S a multiplicatively closed subset the localization $S^{-1}R$ is given by $S^{-1}R = R \times S / \sim$ where $(r, s) \sim (r', s')$ if there is a $t \in S$ such that $t(rs' - r's) = 0$.

Similar constructions can be found in [Sch87].

Remark 5.23 (Braided/crossed case). The Ore condition is clearly met when S is central as exploited above. It is more lax though. It for instance allows for a crossed product types and essentially formalizes bicrossed products. These appear naturally for isomorphisms [FL91], [KW17, §6.2].

For the example of the algebra of morphisms of a category, the Ore condition is guaranteed if there is a commutative diagram for every pair (a, s)

$$\begin{array}{ccc} X & \xrightarrow{s} & Y \\ a' \downarrow & & \downarrow a \\ X' & \xrightarrow{s'} & Y' \end{array} \quad (5.8)$$

5.5. Application to pathlike bialgebras. The following results follow in a straightforward fashion from the previous ones.

Theorem 5.24. *Let B be a pathlike bialgebra:*

- (1) *Every grouplike invertible character has a \star -inverse.*
- (2) *The quotient bialgebra B/I_N of Proposition 5.10 is connected and hence Hopf. In particular, every grouplike normalized character $\phi \in \text{Hom}(B, A)$ when factored through to $\bar{\phi} \in \text{Hom}(B/I_N, A)$ has an inverse computed by $\bar{\phi}^{-1} = \bar{\phi} \circ S$.*
- (3) *The bialgebra $B_{\mathbf{q}}/I$ is Hopf and the \star inverse of a grouplike central character $\phi \in \text{Hom}(B, A)$ when lifted and factored through $\hat{\phi} \in \text{Hom}(B_{\mathbf{q}}/I, A)$ has inverse computed by $\hat{\phi}^{-1} = \hat{\phi} \circ S$.*

\square

6. EXAMPLES FROM FEYNMAN CATEGORIES

We will consider the bialgebras $\mathcal{B}(\mathcal{F})$, for non-symmetric, and $B^{iso}(\mathcal{F})$ for symmetric Feynman categories in particular cases special interest. The examples will all be generated by grouplike elements and skew primitive elements and hence will be pathlike by Theorem 4.25, see Corollary 4.29. As they are categorical coalgebras they are also split. The grouplike elements will be the objects id_X represented by their identities id_X in the absence of isomorphisms and by their classes for B^{iso} . Note these are usually not \otimes -invertible. For this they would have to live in $Pic(\mathcal{F})$, which is not usually the case. It is, however, a natural assumption, that their images under characters are invertible as this happens in concrete applications.

In general there is a grading given by $|X| = \text{word length}$. That is if $X \simeq *_1 \otimes \cdots \otimes *_n$ then $|X| = n$. This is well defined due to axiom (i). This give a grading on the morphisms as $|\phi| = |X| - |Y|$. There may be more intrinsic gradings, for instance by assigning a degree to the generators. We will concentrate of Feynman categories with generators and relations, see [KW17, §5] for more details. All the algebras/coalgebra structures are graded by $|\cdot|$.

The characters need not be graded, but what is common is that map the additive degree to a multiplicative one as in $n \mapsto q^n$, where q can be of degree 0, which can also be achieved by shifting degrees or by inverting elements in the target. One typical such q would be $\frac{1}{2\pi i}$, see [KY], or factors of $\zeta^m(2)$ in the framework of Brown, see [Bro15].

6.1. Surjections and double basepoint preserving injections.

6.1.1. Non-symmetric case. In the non-symmetric case, $B = B(sk(OS))$ —where $sk(OS) \subset \Delta$ is the subcategory of surjection—there is one basic object $[0]$, with $[n] = [0] * \cdots * [0] = [0]^{*n}$, the monoidal unit is $\mathbb{1} = \emptyset = [0]^{*0}$. We use the simplicial notation of Example 2.13. The generating morphisms are $id_{\mathbb{1}}, id_{[0]}$ and $\mu : [1] \rightarrow [0]$ the unique map from $\{0, 1\}$ to $\{0\}$. The relations for morphisms are that $id_{\mathbb{1}}$ is the unit $id_{\mathbb{1}} * f = f$ and $(\mu * id_{[0]} \circ \mu) = (id_{[0]} * \mu) \circ \mu \in Hom([2], [0])$, which we call associativity of μ . Indeed any surjection can be factored through these maps. If $f \in Hom([n], [m])$ then $|f| = n + 1 - m + 1 = n - m$, hence $|id_{[n]}| = 0$ and $|\mu| = 1$. The target degree $|t(f)|$ is sometimes corresponds to the *depth* in applications to polyzeta functions and the degree $|f|$ is the *weight*, [GCKT20a].

$$\Delta(id_{\mathbb{1}}) = id_{\mathbb{1}} \otimes id_{\mathbb{1}}, \Delta(id_{[0]}) = id_{[0]} \otimes id_{[0]}, \Delta(\mu) = id_{[2]} \otimes \mu + \mu \otimes id_{[0]} \quad (6.1)$$

The inclusion of $id_{\mathbb{1}}$ is the coaugmentation. $id_{[0]}$ is grouplike, and hence also all $id_{[n]} = id_{[0]}^{*n}$. It is an easy check that these are the only grouplike elements. and μ is $(id_{[2]}, id_{[0]})$ skew primitive. Hence $B(OS)$ is flavored. Moreover, $F_0^Q B = \bigoplus_n R id_{[n]}$ and this is also split, so it is split pathlike.

B is also a graded coalgebra with the grading $|f|$. The unique maps $\pi_{n+1} : [n] \rightarrow [0]$ are algebra generators. Notice that due to the associativity n -th

iterates, aka. bracketings, coincide and are in fact are the π_{n+1} . $|\pi_{n+1}| = n$ and

$$\Delta(\pi_n) = \sum_{k, (n_1, \dots, n_k) : \sum_{i=1}^k n_i = n} \pi_k \otimes \pi_{n_1} * \cdots * \pi_{n_k} \quad (6.2)$$

Corresponding to all factorizations $[n-1] \rightarrow [k-1] \rightarrow [0]$ Notice that the degree of the l.h.s. is the weight and that $|f|$ is also filtration degree for the bivariate Quillen filtration.

The $id_{[n]}$ are not $*$ invertible, so the bialgebra is not Hopf, but any character $\chi : B \rightarrow A$ for which $\chi(id_{[0]})$ is invertible has a convolution inverse. Quotienting by $id_{\mathbb{1}} - id_{[0]}$, the bialgebra becomes connected. The q deformation is $B[q]/(id_{[0]} - q)$. B^{red} is spanned by the $\pi_i, i \geq 2$ and $\pi_1 = q \in B_q/I$ and the Brown coaction reads

$$\Delta_B(\pi_n) = \pi_n q + \sum \pi_k \otimes \pi_{n_1} * \cdots * \pi_{n_l} \quad (6.3)$$

where the sum runs over the $k, 2 \leq l \leq k, (n_1, \dots, n_l), n_i \geq 2, \sum n_i \leq n$.

Any character passes through the Laurent series.

There is a pictorial representation in terms of planted planar corollas and gluing at leaves, see [GCKT20b, 3.6.3]. A morphism is a forest of planar planted corollas, with $|s(f)|$ leaves and $|t(f)|$ roots or components.

6.1.2. Joyal dual case/Polyzeta/Baues. There is a duality of Δ_+ and the double base point preserving subcategory $\Delta_{*,*} \subset \Delta$. This means that for $f \in Hom([n], [m])$ $f(0) = 0, f(n) = (m)$. Joyal duality is an equivalence of monoidal categories $(\Delta_+, *)$ and $(\Delta_{*,*}^{op}, * \otimes *)$, where $[n]_* \otimes_* [m] = [n+m]$ obtained by identifying $0 \in [m]$ with $n \in [n]$. [GCKT20b, §3.6] for details. The monoidal unit is $[0]$. This maps $[n]$ to $[n+1]$, $id_{\mathbb{1}}$ to $id_{[0]}$, $id_{[0]}$ to $id_{[1]}$ and μ to $*i_* \in Hom([1], [2])$ which is the unique double base point preserving injection. Let $sk(OI_{*,*})$ be the subcategory double base point preserving ordered injections. Then these morphisms generate and have the same properties as above. It is in this form that the diagonal of the coproduct of Baues and Goncharov are best understood, see [GCKT20a, GCKT20b], and in which the cubical structure becomes apparent, see in particular [GCKT20a, §4.3]. The pictorial realization is in terms of Goncharov's half circle diagrams, see [GCKT20b, 3.6.3].

More generally, for simplicial objects this leads to the following terms for the coproduct, see equation (4.29) of [GCKT20a].

$$x_{(0, i_1, \dots, i_{k-1}, n)} \times x_{(0, 1, \dots, i_1)} x_{(i_1, i_1+1, \dots, i_2)} \cdots x_{(i_{k-1}, i_{k-1}+1, \dots, n)}. \quad (6.4)$$

which are Joyal duals to (6.2).

Remark 6.1.

- (1) Goncharov's symbols $\hat{I}(a_0 : a_1, \dots, a_n : a_{n+1})$, see e.g. [GCKT20a, §1.1.5] can be understood as formal characters. The equation that $\hat{I}(a : \emptyset : b) = 0$ then means that these characters factor through

the connected quotient given by coideal $id_1 - id_{[1]}$ in $\Delta_{*,*}$. The bialgebra in question is actually stems for a decoration of the Feynman category by sequences, see [GCKT20b, 3.6.5].

- (2) Not insisting on this relation, one can actually assign any invertible value to $\hat{I}(a : \emptyset : b)$, which is then captured by the Laurent series quotient and Brown's coaction, see §4.4.1, and §5.17.
- (3) In fact, the symbols of Goncharov are characters to a commutative ring, so that we are actually working with $B^{comm}(OI_{*,*})$.
- (4) Contrary to this the image of the characters corresponding to the coproduct of Baues, which is also a decoration, is not commutative.
- (5) The condition that the character passes through the connected quotient is related condition that the underlying space is simply connected, see the discussion in [GCKT20a].

6.1.3. Symmetric case. Now consider FS . This category has isomorphisms $Aut(S) \simeq \mathbb{S}_{|S|}$. Up to isomorphisms it is generated by the same morphisms as above. The bialgebra $B^{iso}(FS) = B^{iso}(sk(FS))$ is generated by $[id_\emptyset]$ which is the class of the unit as above $[id_\emptyset] \amalg [f] = [f]$, $[id_{[0]}]$ and $[\mu]$ where μ is as above, with the relation $([\mu \amalg id_{[0]}] \circ [\mu]) = id_{[0]} \amalg \mu \circ \mu \in Hom([2], [0])$. Indeed up to isomorphism any surjection can be factored through these maps. Here we used the disjoint union for monoidal structure, which extends the join to all finite sets.

For the generators:

$$\Delta(id_1) = id_1 \otimes id_1, \Delta(id_{[0]}) = id_{[0]} \otimes id_{[0]}, \Delta(\mu) = id_{[1]} \otimes \mu + \mu \otimes id_{[0]} \quad (6.5)$$

The rest of the story is as analogous to the above. The inclusion of $[id_1]$ is the coaugmentation. $[id_{[0]}]$ is grouplike, and hence also all $[id_{[n]}] = [id_{[0]}^{*n}]$. It is an easy check that these are the only semigrouplike elements. The generating morphism μ is $([id_{[2]}], [id_{[0]}])$ skew primitive. Note that B^{iso} is commutative, however, since the commutator $f \amalg g \rightarrow g \amalg f$ is an isomorphism.

Hence $B^{iso}(FS)$ is flavored. Moreover, $F_0^Q B = \bigoplus_n R[id_{[n]}]$ and this is also split, so it is split pathlike. The $id_{[n]}$ are not \amalg -invertible, so the bialgebra is not Hopf, but any character $\chi : B \rightarrow A$ for which $\chi([id_{[0]}])$ is invertible has a convolution inverse. Quotienting by $[id_1] - [id_{[0]}]$, the bialgebra becomes connected and looplike. The q deformation is $B[q]/([id_{[0]}] - q)$. And any character passes through the Laurent series.

The pictorial representation is given by unlabelled rooted corollas, see [GCKT20b, §3.6.3].

6.1.4. Summary.

Theorem 6.2.

- (1) *The bialgebra $B(sk(OS))$ is commutative, non-cocommutative and split pathlike. It is generated by the unit, the grouplike $id_{[0]}$ and the $(id_{[1]}, id_{[0]})$ -skew primitive μ .*

- (2) *Joyal duality takes on the following form: $B(sk(OS))^{op} = B(sk(OS)^{op}) = B(Inj_{*,*})$ where $Inj_{*,*} \subset \Delta_{*,*}$ is the subcategory of double base point preserving injections.*
- (3) *The bialgebra $B^{iso}(FS)$, is commutative, cocommutative and split pathlike. It is generated by the unit, the grouplike $[id_{[0]}]$ and the $([id_{[1]}], [id_{[0]}])$ -skew primitive $[\mu]$.*
- (4) *In both cases, the bivariate Quillen filtration is indexed by $|f|$ and $|f|$ makes the coalgebra into a non-connected graded coalgebra.*

Remark 6.3. In the localized version $S(\mu) = -id_{[1]}^{-1} * \mu * id_{[0]}^{-1}$, with $id_{[1]}^{-1} = id_{[0]}^{-2}$ and in Laurent series $S(\mu) = -q^{-3}\mu$.

6.2. CK-bialgebras for (planar) trees with leaves, and operads. For these bialgebras, the underlying Feynman categories are enrichments of the examples above. Concretely, the objects are again finite sets, or natural numbers in the skeletal version. We will use the non-simplicial notation. It suffices to give the $Hom(\bar{n}, \bar{k}) = \mathcal{O}^{nc}(k)$ and their compositions where

$$\mathcal{O}^{nc}(n, k) = \bigoplus_{\substack{(n_1, \dots, n_k) \\ \sum_{i=1}^k n_i = n}} \mathcal{O}(n_1) \otimes \dots \otimes \mathcal{O}(n_k) \quad (6.6)$$

Here and in the following \bigoplus stands for colimit and we will use \otimes for the monoidal product. If the $\mathcal{O}(n)$ are sets, then these are simply \amalg and \times . In case they are, e.g. vector spaces, where the coproduct is \oplus , then one has to work with enriched categories. The $\mathcal{O}(n)$ are then elements of an enrichment category \mathcal{E} and $\otimes = \otimes_{\mathcal{E}}$ is the monoidal product on \mathcal{E} , see [Kau21b, §4] for a detailed account. In our current context, we are working with $\mathcal{E} = R\text{-Mod}$.

$$\gamma_{k, n_1, \dots, n_k} : \mathcal{O}(k) \otimes \mathcal{O}(n_1) \otimes \dots \otimes \mathcal{O}(n_k) \rightarrow \mathcal{O}(\sum n_i) \quad (6.7)$$

which correspond to the components of the composition

$$Hom(\bar{k}, \bar{1}) \times Hom(\bar{n}, \bar{k}) \rightarrow Hom(\bar{n}, \bar{1}) \quad (6.8)$$

This is equivalent to the statement that the $\mathcal{O}(n)$ form an operad, see in particular [Kau21b, Proposition 3.20]. In the non-symmetric case, this is a non-symmetric operad and in the symmetric case, the automorphisms \mathbb{S}_n act on the objects yielding a compatible action on the $\mathcal{O}^{nc}(n, k)$ which is compatible with composition. Concretely $\mathcal{O}(n)$ will be rooted trees or planar planted trees with n leaves and the composition is given by gluing on leaves.

In the particular case of trees, the $\mathcal{O}^{nc}(n, k)$ are the forests with a total of n leaves made up of forests with k trees and with n_1, \dots, n_k leaves.

Each such \mathcal{O} gives a Feynman category $\mathfrak{F}_{\mathcal{O}}$. This holds more generally in the context of plus constructions, see [Kau21b]. We will denote the bialgebras by $B(\mathcal{O})$ or $\mathcal{B}^{iso}(\mathcal{O})$ respectively.

Remark 6.4. In the framework of cooperads with multiplication, [GCKT20a] the underlying cooperad is the dual of $\mathcal{O}^{nc}(n) = \bigoplus_k \mathcal{O}^{nc}(n, k)$ if the cooperad comes from an operad, which has free multiplication \otimes . In the current context that theory can be understood as considering relations imposed on the algebra part and which structures of the free model survive and which are recoverable.

6.2.1. Non-symmetric case. In the non-symmetric case, taking $\mathcal{O}(n)$ to be planted planar trees with n leaves, then $\mathcal{O}^{nc}(n)$ are the planted planar forests, where this includes special elements, namely the identities $id_n = id_1^{\otimes n} \in \mathcal{O}^{nc}(n)$ where id_1 can be depicted by a vertexless leaf, denoted by $|$ in [GCKT20a]. The algebra structure is given by juxtaposition and the coalgebra structure is dual to gluing the roots to the vertices.

In general, the formula for the coproduct is given by

$$\Delta(\tau) = \sum_{\tau_0 \subseteq \tau} \tau_0 \otimes \tau \setminus \tau_0 \quad (6.9)$$

where τ_0 is a rooted subtree —stump— including all its tails and $\tau \setminus \tau_0$ removes all the vertices and half edges of τ_0 , viz, the set of cut off branches, see [GCKT20a] for details. A summand is depicted in Figure 2.

The element id_1 and hence all the id_n are grouplike and the τ_n are (id_1, id_n) skew primitive.

$$\Delta(\tau_n) = id_1 \otimes \tau_n + \tau_n \otimes id_n \quad (6.10)$$

The operad \mathcal{O}^{nc} is freely generated by $1 = id_1$, which is the empty forest id_1 the $\tau_n \in \mathcal{O}(n)$ —which are the corollas with n leaves and id_n — under monoidal product and composition. This reflects that any tree can be decomposed into the rooted corollas $\tau_{ar(v)}$ by cutting all the edges. Here v runs through the vertices of the tree and $ar(v)$ is the arity of v that is the number of incoming edges, where edges are oriented toward the root. The degree of a forest ϕ is $|s(\phi)|$ that is the total number of leaves, which is n for $\phi \in \mathcal{O}^{nc}(n)$. The depth $|t(\phi)|$ is the number of roots, and $|\tau|$ is the weight which is the number of leaves minus the number of roots.

Proposition 6.5. *The bialgebra $B(\mathcal{O})$ is commutative and non-cocommutative split pathlike. It is graded by $|V(\tau)|$ and $|\tau|$. With the filtration level for the bivariate Quillen filtration given by $|V\tau|$ The subalgebra generated by the grouplike elements is freely generated by id_1 and id_1 and hence $\mathcal{B}_{\mathbf{q}}/I \simeq \mathcal{B}^{red}[q, q^{-1}]$.*

NB: It is neither connected with respect to the grading $|\tau|$ nor that by $|V(\tau)|$.

Proof. If $\tau \neq id_1^{\otimes n} = |^n$ then applying the coproduct $|E(\tau)|$ times, dissects the tree into its corolla components, which are skew primitive and hence any tree lives in $F_{|E(\tau)|+1}^Q$ which shows that the filtration is exhaustive. This also shows that the $|E(\tau)| + 1 = |V|$ is the filtration degree. This extends to

the special case of $|$ which has 0 vertices. Moreover, the filtration actually gives a grading with $\deg(\tau) = V(\tau)$. The $id_n = id_1^{\otimes n}$ are easily seen to be the only grouplike elements, since such elements may not have any internal edges and they have no relations. \square

This recovers the grading discussion of [GCKT20a, 2.12] and explains the lift of the coradical grading of the original leaf-less Connes–Kreimer Hopf algebra.

6.2.2. Symmetric case. In the symmetric case, the algebra generators $\mathcal{O}(n)$ are rooted trees with labelled leaves. These are permuted by an \mathbb{S}_n action. Their isomorphism classes are the elements of $[\tau] \in \mathcal{O}(n)^{\mathbb{S}_n}$ which are unlabelled rooted trees and as an algebra $B^{iso} = R[[\tau]]$ is a polynomial, aka. symmetric algebra in the unlabelled rooted trees. The generators are unlabelled unordered rooted forests. The comultiplication which is dual to the composition by gluing roots to leaves is given by

$$\Delta([\tau]) = \sum_{\tau_0 \subset \tau} [\tau_0] \otimes [\tau \setminus \tau_0] \quad (6.11)$$

on the isomorphism classes. See Figure 2 for a summand.

All the previous comments and proof apply *mutatis mutandi*.

Proposition 6.6. *For $\mathcal{O}(n)$ given by rooted labelled trees, the bialgebra $B^{iso}(\mathcal{O})$ is commutative, cocommutative and split pathlike. It is graded by $|V(\tau)|$ and $|\tau|$. The subalgebra generated by the grouplike elements is freely generated by $[id_{\mathbb{1}}]$ and $[id_1]$ and hence $\mathcal{B}_{\mathbf{q}}/I \simeq \mathcal{B}^{red}[q, q^{-1}]$. \square*

6.2.3. Observations for the tree algebras.

Remark 6.7. In this language a particular property of the CK-Hopf algebra becomes apparent. It is a pathlike coalgebra whose grouplike elements are generated by $e = id_{\mathbb{1}}$ and $| = id_{\overline{1}}$ and it has a unique $id_{\overline{1}}$ primitive \blacklozenge which generates a split line $R\blacklozenge$. This yields the special properties explored in [Moe01] and its generalizations, [GCKT20a, Example 2.50], where, via coloring, many of these are introduced as additional direct components, so that the bialgebras are still pathlike.

Remark 6.8. In the localized Hopf algebra, $S(\blacklozenge) = -id_{\overline{1}}^{-1} \otimes \blacklozenge \otimes id_{\overline{1}}^{-1}$ and hence in the Laurent series $S(\blacklozenge) = -\blacklozenge q^{-2}$. More generally, for the skew primitive τ_n , $S(\tau_n) = -id_{\overline{1}}^{-1} \otimes \tau_n \otimes id_n^{-1}$ respectively $S(\tau_n) = -q^{-(n+1)}\tau_n$.

Remark 6.9. The Brown coaction which given by setting the q terms on the right in Δ to 1. Combined with the previous remark, this explains the fact that the infinitesimal structure can be given by a q -shifted comultiplication and a residue, see [GCKT20a, Corollary 2.2.5]. This generalizes Brown's infinitesimal structure to the level of operads. More generally, there might be more than one deformation parameter.

6.3. General operads. For general operads the proposition needs to include a condition on $\mathcal{O}(1)$. Note that $\mathcal{O}(1)$ is a monoid. The corresponding $\mathcal{O}(1)$ is a subcategory of $\mathcal{F}_{\mathcal{O}}$. The results of [KW17] in the current language yield a refinement:

Theorem 6.10. *For a non-symmetric operad \mathcal{O} with $\mathcal{O}(0)$ spanned by $\mathbb{1}$. $B(\mathcal{O})$ is flavored if and only if the coalgebra of $\mathcal{O}(1)$ is flavored. Similarly, for an operad \mathcal{O} , B^{iso} is flavored if and only if $[\mathcal{O}(1)]$ is flavored.*

They are pathlike if and only if the corresponding restriction to $\mathcal{O}(1)$ are pathlike. In this case, the degree 0 part of the bivariate Quillen filtration for the whole bialgebra is the free (symmetric) algebra on the restriction to $\mathcal{O}(1)$.

Proof. This is a translation of the proofs. The main point is that iterating the coproduct, lowers n in $\mathcal{O}^{nc}(n)$ and produces factors in $\mathcal{O}^{nc}(1) = \mathcal{O}(1)$. In the end only $\mathcal{O}(1)$ factors survive. The condition ensures that the QT-filtration is exhaustive and vice-versa if it is, it follows that the restriction is.

For the last part, in one direction this follows by restriction. Assume now that the coalgebra for $\mathcal{O}(1)$ is pathlike. The semigrouplike elements have to lie, or have representatives, in $\mathcal{O}^{nc}(n, n) = \mathcal{O}(1) \otimes \cdots \otimes \mathcal{O}(1)$. Since the product is free, this means that they are products of semigrouplike elements stemming from $\mathcal{O}(1)$. If the latter are all grouplike, so are the former. Likewise, since the multiplication is free, the degree 0 part of the bivariate Quillen filtration is then a direct sum if and only if the former is. \square

Remark 6.11. Note that these bialgebras are always graded by $|\phi|$, but they are usually not connected in this grading.

6.4. Feynman categories of graphs. There is a basic Feynman category of graphs called \mathfrak{G} , see [KW17, §2.1]. We review salient features here. In the appendix there is a purely set-theoretical version, which is of independent interest and makes things even more combinatorial.

6.4.1. Objects, morphisms and underlying graphs. The category \mathcal{V} has S -labelled corollas $*_S$ as objects with $Aut(*_S) \simeq Aut(S)$. This means that a general object is a forest or aggregate of such corollas. The connected version \mathfrak{G}^{ctd} has morphisms generated by simple edge contractions: $_{s \circ t} : *_S \amalg *_T \rightarrow *_S \setminus_{\sigma \amalg T} \{t\}$ and simple loop contractions $\circ_{ss'} : *_S \rightarrow *_S \setminus_{\{s, s'\}}$. These satisfy quadratic relations and are crossed with respect to the isomorphisms. For any ϕ and any isomorphism σ there are unique ϕ' and σ' such that $\phi \circ \sigma = \sigma' \circ \phi'$. Each morphism has an underlying graph $\mathbb{F}(\phi)$ which has one edge for each contraction. It is defined in a way such that $\mathbb{F}(\phi \circ \sigma) = \mathbb{F}(\phi')$ and its automorphisms are those of the source. The isomorphism class of this graph gives the isomorphism class of ϕ : $[\mathbb{F}(\phi)] = [\phi]$. A morphism ϕ in \mathfrak{G}^{ctd} is basic if and only if $\mathbb{F}(\phi)$ is connected. See Figure 1.

For the non-connected version, there is one more generator ${}_S\Box_T : {}_S\amalg {}_T \rightarrow {}_{S\amalg T}$. In this case, there is a non-homogeneous relation ${}_s\circ_t = \circ_{s,t} {}_S\Box_T$ which allow to replace the edge contraction and only retain the loop contractions and mergers as generators. The basic morphisms now need not to have connected underlying graph anymore.

6.4.2. Isomorphism classes. Each corolla ${}_S$ is isomorphic to ${}_{|\overline{S}|}$, and hence $[id_{{}_S}] = [id_{|\overline{S}|}]$. Furthermore fixing isomorphisms $\sigma : S \rightarrow \overline{S}$, with $\sigma(s) = 1, \sigma(s') = 2$ and $\sigma' : T \rightarrow \overline{T}$ with $\sigma'(t) = 1, [{}_s\circ_t] = [{}_0\circ_1], \circ_{s,s'} = [{}_0\circ_1, 2]$ and $[{}_S\Box_T] = [|\overline{S}|] \Box [|\overline{T}|]$. That is there is only one isomorphism class per generating type.

6.4.3. Grading. Beside the grading $|\cdot|$ which is 1 for an edge contraction or a merger and 0 for an isomorphism or a loop contraction, there is an additional degree given by $deg({}_S) = |S|$ which is additively continued $deg({}_S \amalg {}_T) = |S| + |T|$, to count the total number of legs in the forest. This yields $deg(\phi) := \frac{1}{2}(\deg(s(\phi)) - \deg(t(\phi)))$, which is integer, since both loop and edge contractions reduce the set of legs by an even number, while isomorphisms and mergers keep this number fixed. The degree of an isomorphism or a merger is hence 0 and that of a simple loop or edge contraction is 1. In this way $deg(\phi) = |E(\mathbb{T}(\phi))|$. The sum $wt(\phi) = deg(\phi) + |\phi|$ is 1 for the mergers and loop contractions and 2 for the edge contractions.

Remark 6.12 (Loop number and Euler characteristic). In the graphical interpretation for a basic morphism $wt(\phi) = |E(\mathbb{T}(\phi))| + b_0(\phi)$ where b_0 is the number of components of $\mathbb{T}(\phi)$, aka. zeroth Betti number. This is related to the Euler characteristic as follows $\chi(\mathbb{T}(\phi)) = wt(\phi) - |\phi|$, where $\chi(\mathbb{T}(\phi)) = b_0(\mathbb{T}(\phi)) - b_1(\mathbb{T}(\phi)) = |V(\mathbb{T}(\phi))| - |E(\mathbb{T}(\phi))|$ and b_1 is the loop number, aka. first Betti number.

6.4.4. Bi and Hopf algebra structures. The coproduct in B^{iso} is given by [GCKT20b, §3.6].

$$\Delta[\phi] = \Delta[\mathbb{T}(\phi)] = \sum_{\gamma \subset \mathbb{T}(\phi)} [\gamma] \otimes [\mathbb{T}(\phi)/\gamma] \quad (6.12)$$

This is a version of the core Hopf algebra [Kre10], but with legs and at a bialgebra level. See Figure 3 for an example of a summand.

Theorem 6.13. *Both $B^{iso}(\mathfrak{G}^{ctd})$ and $B^{iso}(\mathfrak{G})$ are split pathlike. They are both graded by $deg, |\cdot|$ and wt . For $B^{iso}(\mathfrak{G}^{ctd})$ the bivariate Quillen level is given by deg and for $B^{iso}(\mathfrak{G})$ the bivariate Quillen level is given by wt .*

More precisely, the grouplike elements are $[id_1]$ and $[id_X]$ which are generated by the $[id_{{}_S}] = [id_{|\overline{S}|}]$. In the connected case, the skew primitive elements are generated by the $[{}_s\circ_t]$ and the $[\circ_{s,s'}]$ while in the connected case these are generated by the $[\circ_{s,s'}]$ and the $[{}_S\Box_T]$ via products with grouplikes.

Proof. The fact about the skew primitives and the grouplikes is straightforward to check. All semigrouplikes are grouplike and they split by definition of the algebra structure. Up to isomorphism any morphism can be decomposed into chains of morphisms in the indecomposable generators of length $wt(\phi)$ respectively $deg(\phi)$. Applying the reduced coproduct, the wt respectively deg drops at least on one side. This proves by induction that the bivariate Quillen filtration is exhaustive. Since the coalgebras are categorical the structures are split. \square

Remark 6.14. The antipodes for the skew primitives in the localization and the single parameter Laurent polynomials are given as follows: $S(\overline{n\Box_m}) = -[id_n]^{-1}[id_m]^{-1}[id_{n+m}]^{-1}n\Box_m$, $S(\overline{n\Box_m}) = -q^{-2(n+m)}n\Box_m$ where we dropped the overline. $S([{}_s\circ_t]) = [id_{|S|}]^{-1}[id_{|T|}]^{-1}[id_{|S|+|T|-2}]^{-1}[{}_s\circ_t]$, respectively $[{}_s\circ_t] = -q^{-2(|S|+|T|+1)}{}_s\circ_t$. Similarly $S([\circ_{s,s'}]) = [id_{|S|}]^{-1}[id_{|T|}]^{-1}[id_{|S|+|T|-2}]^{-1}[\circ_{s,s'}]$, respectively $[\circ_{s,s'}] = -q^{-2(|S|+|T|+1)}[\circ_{s,s'}]$.

The connected quotient identifies all the objects with $\mathbb{1}$ and hence all grouplikes id_X with $1 = id_{\mathbb{1}}$. The Brown coaction again sets the q variables on the right to 1, which halves the q degrees in the above formulae.

6.5. General Feynman categories. The preceding arguments outline a general approach. The input is a Feynman category with a given presentation, that is a subset of generators Φ of basic morphisms under composition and monoidal product. Recall from Corollary 2.23 that the only semigrouplike elements for B^{iso} are the identities which are grouplike, and $[\phi]$ being skew primitive means that ϕ is essentially primitive.

Theorem 6.15. *The bialgebra $B^{iso}(\mathcal{F})$, is pathlike if \mathcal{F} is generated by isomorphisms and a set of generators which are essentially skew primitive or essentially indecomposable. In case that there are homogenous relations, the coalgebra will be graded.*

For a non-symmetric Feynman category $B(\mathcal{F})$ is pathlike, and \mathcal{F} is generated by identities and skew primitives, that is indecomposables, which is equivalent to a set of generators which are skew primitive. In case that there are homogenous relations, the coalgebra will be graded. It is necessary that \mathcal{V} is discrete.

Proof. Give a morphism ϕ , it will have be expressible in indecomposables. If n is the maximal number of such an expression, then there is a chain of reduced coproducts which will have the indecomposables as factors, and hence ϕ will live in $F_n^Q(B^{iso}(\mathcal{F}))$. If the relations are homogenous, then there is a function l defined to be 1 on the generators, so that $l(\phi)$ as above is n . The semigrouplike elements are the classes of the identities, see Remark 2.9, these are grouplike, and they split off.

In the non-sigma case, the condition is that the generators are indecomposable, then the rest follows as above. Notice that if there would be isomorphisms besides the identities, then these isomorphisms would not be in the bivariate Quillen filtration. \square

Remark 6.16. In these cases the localization and the Laurent series will give Hopf algebras. The q deformations will have parameters given by the isomorphism classes of objects. The $[id_X]$ are also the classes that will be formally inverted. This can be thought of as extending \mathcal{F} to $S^{-1}\mathcal{F}$ such that the X to live in the $Pic(S^{-1}\mathcal{F})$. And the same observations as in Remark 6.14 apply.

Remark 6.17. In the above cases, we had generators with quadratic relations. The function l was deg in the connected case and wt in the non-connected case. Further examples are those from Feynman categories with a generating set that is at most quadratic [KW17, Definition 5.6].

Example 6.18. The Feynman category for operads and that for operads with multiplication are examples as are all the examples coming from restriction and decoration listed in [KW17, 2.6.4]. For the former the generators are simply edge contractions. It is interesting to note that for operads with multiplication there are two sets of skew primitives τ_b^m and τ_w^n which play an important role in the theory of E_2 operads, [KZ17].

CONCLUSION AND OUTLOOK

We have shown that pathlike coalgebras in addition to quiver and incidence coalgebras capture the two types of bialgebras stemming from Feynman categories, such as those of Baues, Goncharov and those of Connes–Kreimer. This realization provides several outcomes.

The structure of path-like coalgebras exactly answers the question of what exactly the harbinger of the antipode is in the various constructions which we could house in a common categorical context. It moreover precisely identifies the grouplike elements as the obstruction to the existence of the antipode. For characters, their values determine the \star -invertibility. This also explains the appearance of quotients. These correspond to restrictions on the characters. The most common, which says that all grouplike elements have character value 1, leads to the connected quotient. Other restrictions yield different quotients and explain the universality of the quantum deformation, which in turn explain the naturality of Brown’s coaction.

Furthermore, we identified the skew primitive elements as indecomposables as being of central importance. This parallels the role of primitives in the usual connected case. In particular, there should be B_+ operators associated to them. These are indeed explicitly present and key elements for E_2 -operads and hence for Gerstenhaber brackets and Deligne’s conjecture [KZ17]. Moreover, there is a connection between these elements and the terms of the master equations [KMZ96] and [KW17, §7.5]. The general structure of B_+ operators will be taken up in [Kau21a], where we will also make the connection to Hochschild complexes.

The example of the Drinfel’d double which is resolved by going to the double-categorical level in §2.7 and is related to the localization construction, leads the way to a new research direction which will be pursued further,

especially in view of [Kau21b]. In particular, this example gives a new method to deal with isomorphisms. On the double categorical side, the antipode, in a sense, switches the horizontal and vertical composition.

Another direction, is the connection to the cubical structure, whose analysis in this context appeared in [GCKT20a, §4.5] and appeared again in the cubical chain complex of [BK21]. These are intimately related to Cutkosky rules and the work of [KY]. In particular, as discussed in [GCKT20a] the simplicial coproduct (6.4) has a cubical structure which can be understood comes via the Serre diagonal:

$$\delta : P = [0, 1]^{n-1} \xrightarrow{\cong} \bigcup_{K \cup L = \{1, \dots, n-1\}} \partial_K^- [0, 1]^{n-1} \times \partial_L^+ [0, 1]^{n-1} \xrightarrow{\subset} P \times P \quad (6.13)$$

The term given in (6.4) is identified with the subset $L = \{i_1, \dots, i_{k-1}\}$. In view of [KY] one has an additional interpretation. The right hand side (1.2) respectively 6.13 can be seen as cuts (or stops) in the path from 0 to n on the quiver $[n] = 0 \rightarrow 1 \rightarrow \dots \rightarrow n$ and these can be viewed as cuts of a linear graph. This is in line with point of view of the simplicial structure of iterated integrals. Furthermore, there is a symmetry in the Serre diagonal of the cube given by interchanging 0 and 1 in the standard presentation. This switches the coproduct to its opposite. These two observations should generalize to the categorical setting, and when applied to graphs may explain some of the results of [KY]. A last aspect is functoriality of the construction, see [GCKT20b, §1.7], which is likely related the structure of cointeracting coalgebras [Foi, KY] in the present context.

FIGURES




FIGURE 1. The basic morphisms: Edge contraction loop contraction (excluded for trees/forests) and merger (excluded for connected version)




FIGURE 2. A summand of the coproduct for (planar) trees with leaves and roots. In the planar case, the r.h.s. is either in the order depicted and in the non-planar it is a symmetric product.




FIGURE 3. A summand of the coproduct. Note for the morphisms all flags and vertices are labelled and tracked here only marked the vertices to avoid clutter. For the coproduct only the isomorphism classes of the morphism are involved.

APPENDIX A. SET VERSION OF GRAPHS

We give a non-graphical version here, a reference for the graphical version is [KW17, §2.1] and [KW17, Appendix A]. The basic objects will be $\mathcal{V} = \text{Iso}(\text{FinSet})$. In the graphical version this is given by identifying a finite set S by an S -labelled corolla $*_S$. This fixes the $\text{Iso}(\mathcal{V}^{\otimes})$ to finite tuples of sets and the inner and outer permutations. The outer permutations act as follows: Let $p \in \mathbb{S}_n$ the acting by $p(S_1, \dots, S_n) = (S_{p(1)}, \dots, S_{p(n)})$. The inner permutations $\sigma \in \text{Aut}(S_i)$ act as $(id, \dots, id, \sigma, id, \dots, id)$.

In graphical form these are aggregates, that is a disjoint union, aka. forests, of corollas, and permutation of the corollas and the flags of each corolla. These will be written as finite tuples of finite sets (S_1, \dots, S_n) . There are two versions. The connected and the non-connected version.

The for the connected connected version the morphisms are generated monoidally and by concatenation by the isomorphisms and the following morphisms

$$s \circ_t : (S, T) \rightarrow (S \setminus \sigma \amalg T \setminus \{t\}), \quad \circ_{ss'} : (S) \rightarrow (S \setminus \{s, s'\}), \quad (\text{A.1})$$

called simple edge contractions, simple loop contractions.

To write the relations in a concise fashion, we abuse notation and write ϕ for $(id, \dots, id, \phi, id, \dots, id)$.

- (1) The basic morphisms are equivariant with respect to isomorphisms.

- (a) Internal action for $\sigma \in Aut(S), \sigma' \in Aut(T)$:

$$\begin{aligned} \sigma(s) \circ_{\sigma'(t)}(\sigma, \sigma') &= \sigma \amalg \sigma' |_{S \setminus \{s\} \amalg T \setminus \{t\}} s \circ_t \\ \circ_{\sigma(s), \sigma(s')} \sigma &= \sigma |_{S \setminus \{s, s'\}} \circ_{s, s'} \end{aligned} \quad (\text{A.2})$$

- (b) External action commutes with loop contractions.

$$p \circ_{s, s'} = \circ_{s, s'} p \quad (\text{A.3})$$

- (c) External action is equivariant with edge contraction. If $s \in S_i$ and $t \in S_{i+1}$ then

$$s \circ_t = t \circ_s \tau_{i, i+1} \text{ and } p_{s \circ_t} = s \circ_t p' \quad (\text{A.4})$$

where p' is the image of the permutation of in the embedding $\mathbb{S}_{n-1} \rightarrow \mathbb{S}_n$ given by the block permutation.

- (2) Loop contractions commute:

$$\circ_{s_1, s'_1} \circ_{s_2, s'_2} = \circ_{s_2, s'_2} \circ_{s_1, s'_1} \quad (\text{A.5})$$

- (3) Loop and edge contractions commute

$$s_1 \circ_{t_1} \circ_{s_2, s'_2} = \circ_{s_2, s'_2} s_1 \circ_{t_1} \quad (\text{A.6})$$

- (4) Edge contraction commute if the “edges” do not form a cycle. For $s_1 \in S_1, t_1 \in T_1, s_2 \in S_2, t_2 \in T_2$ and $|\{S_1, T_1, S_2, T_2\}| \geq 3$

$$s_1 \circ_{t_1} s_2 \circ_{t_2} = s_1 \circ_{t_1} s_2 \circ_{t_2} \quad (\text{A.7})$$

- (5) Edge contraction commute even if they form a cycle, only that after one edge contraction, the second edge contraction is a loop contraction. For $s_1, s_2 \in S, t_1, t_2 \in T$,

$$\circ_{s_2, t_2} s_1 \circ_{t_1} = \circ_{s_1, t_1} s_2 \circ_{t_2} \quad (\text{A.8})$$

In the non-connected case, there is an additional generating morphism called simple merger:

$$s \boxminus_T : (S, T) \rightarrow (S \amalg T) \quad (\text{A.9})$$

These satisfies the relations

- (1) Mergers are equivariant with respect to the internal action

$$s \boxminus_T(\sigma, id) = (\sigma \amalg id) s \boxminus_T \quad (\text{A.10})$$

- (2) Mergers are equivariant with respect to the outer action $\text{On} (S_1, \dots, S_n)$:

$$s_{i+1} \boxminus s_i = s_i \boxminus_{S_{i+1}} \tau_{i,i+1} \text{ and } p s_i \boxminus_{S_{i+1}} = s_{p'(i)} \boxminus_{S_{p'(i+1)}} p' \quad (\text{A.11})$$

where p' is the image of the permutation p under the embedding $\mathbb{S}_{n-1} \rightarrow \mathbb{S}_n$ given by the block permutation with block $(i, i+1)$.

- (3) Simple mergers commute

$$s_1 \boxminus_{T_1} s_2 \boxminus_{T_2} = s_2 \boxminus_{T_2} s_1 \boxminus_{T_1} \quad (\text{A.12})$$

- (4) Mergers and edge/loop contractions on different sets commute.

- (5) If $s \in S$ and $t \in T$, then

$$s \circ t = \circ_{st} s \boxminus_T \quad (\text{A.13})$$

REFERENCES

- [AFS08] Nicolás Andruskiewitsch and Walter Ferrer Santos. The beginnings of the theory of hopf algebras. *Acta Applicandae Mathematicae*, 108(1):3, 2008.
- [Art99] M. Artin. Noncommutative rings. Course notes. Available at <http://www-math.mit.edu/~etingof/artinnotes.pdf>, 1999.
- [Atk63] F. V. Atkinson. Some aspects of Baxter’s functional equation. *J. Math. Anal. Appl.*, 7:1–30, 1963.
- [Bau81] H. J. Baues. The double bar and cobar constructions. *Compositio Math.*, 43(3):331–341, 1981.
- [BK21] Clemens Berger and Ralph M. Kaufmann. Derived Decorated Feynman Categories. *In preparation*, 2021.
- [Bro15] Francis Brown. Motivic periods and the projective line minus three points. *Preprint*, [arxiv:1407.5165](https://arxiv.org/abs/1407.5165), 2015.
- [Car07] Pierre Cartier. A primer of Hopf algebras. In *Frontiers in number theory, physics, and geometry. II*, pages 537–615. Springer, Berlin, 2007.
- [Che73] Kuo-tsai Chen. Iterated integrals of differential forms and loop space homology. *Ann. of Math. (2)*, 97:217–246, 1973.
- [CK98] Alain Connes and Dirk Kreimer. Hopf algebras, renormalization and non-commutative geometry. *Comm. Math. Phys.*, 199(1):203–242, 1998.
- [CK00] Alain Connes and Dirk Kreimer. Renormalization in quantum field theory and the Riemann-Hilbert problem. I. The Hopf algebra structure of graphs and the main theorem. *Comm. Math. Phys.*, 210(1):249–273, 2000.
- [CK01] Alain Connes and Dirk Kreimer. Renormalization in quantum field theory and the Riemann-Hilbert problem. II. The β -function, diffeomorphisms and the renormalization group. *Comm. Math. Phys.*, 216(1):215–241, 2001.
- [DEEFG16] Luis A. Duffaut Espinosa, Kurusch Ebrahimi-Fard, and W. Steven Gray. A combinatorial Hopf algebra for nonlinear output feedback control systems. *J. Algebra*, 453:609–643, 2016.
- [DPR90] R. Dijkgraaf, V. Pasquier, and P. Roche. Quasi Hopf algebras, group cohomology and orbifold models. volume 18B, pages 60–72 (1991). 1990. Recent advances in field theory (Annecy-le-Vieux, 1990).
- [EFGK05] Kurusch Ebrahimi-Fard, Li Guo, and Dirk Kreimer. Integrable renormalization. II. The general case. *Ann. Henri Poincaré*, 6(2):369–395, 2005.
- [EFK05] Kurusch Ebrahimi-Fard and Dirk Kreimer. The Hopf algebra approach to Feynman diagram calculations. *J. Phys. A*, 38(50):R385–R407, 2005.
- [FL91] Zbigniew Fiedorowicz and Jean-Louis Loday. Crossed simplicial groups and their associated homology. *Trans. Amer. Math. Soc.*, 326(1):57–87, 1991.

- [Foi] Loïc Foissy. Cointeracting bialgebras. talk given at Higher Structures Emerging from Renormalisation, ESI, Vienna, October 2020 (<https://www.esi.ac.at/events/t304/>) and at Algebraic Structures in Perturbative Quantum Field Theory, IHES, November 2020.
- [Foi02] L. Foissy. Les algèbres de Hopf des arbres enracinés décorés. II. *Bull. Sci. Math.*, 126(4):249–288, 2002.
- [GCKT20a] Imma Gálvez-Carrillo, Ralph M. Kaufmann, and Andrew Tonks. Three Hopf Algebras from Number Theory, Physics & Topology, and their Common Background I: operadic & simplicial aspects. *Comm. in Numb. Th. and Physics*, 14(1):1–90, 2020.
- [GCKT20b] Imma Gálvez-Carrillo, Ralph M. Kaufmann, and Andrew Tonks. Three hopf algebras from number theory, physics & topology, and their common background ii: general categorical formulation. *Comm. in Numb. Th. and Physics*, 14(1):91–169, 2020.
- [GM03] Sergei I. Gelfand and Yuri I. Manin. *Methods of homological algebra*. Springer Monographs in Mathematics. Springer-Verlag, Berlin, second edition, 2003.
- [Gon05] A. B. Goncharov. Galois symmetries of fundamental groupoids and noncommutative geometry. *Duke Math. J.*, 128(2):209–284, 2005.
- [Guo12] Li Guo. *An introduction to Rota-Baxter algebra*, volume 4 of *Surveys of Modern Mathematics*. International Press, Somerville, MA; Higher Education Press, Beijing, 2012.
- [JR79] S. A. Joni and G.-C. Rota. Coalgebras and bialgebras in combinatorics. *Stud. Appl. Math.*, 61(2):93–139, 1979.
- [Kas95] Christian Kassel. *Quantum groups*, volume 155 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, 1995.
- [Kau03] Ralph M. Kaufmann. Orbifolding Frobenius algebras. *Internat. J. Math.*, 14(6):573–617, 2003.
- [Kau21a] Ralph M. Kaufmann. B-plus operators in Operators in Field Theory, Feynman Categories and Hopf Algebras. *In preparation*, 2021.
- [Kau21b] Ralph M. Kaufmann. Feynman categories and representation theory. *Commun. Contemp. Math. to appear preprint arXiv:1911.10169*, 2021.
- [KMM20] Ralph M. Kaufmann and Anibal M. Medina-Mardones. Chain level Steenrod operations. *arXiv:2010.02571*, 2020.
- [KMZ96] R. Kaufmann, Yu. Manin, and D. Zagier. Higher Weil-Petersson volumes of moduli spaces of stable n -pointed curves. *Comm. Math. Phys.*, 181(3):763–787, 1996.
- [KP09] Ralph M. Kaufmann and David Pham. The Drinfel’d double and twisting in stringy orbifold theory. *Internat. J. Math.*, 20(5):623–657, 2009.
- [Kre10] Dirk Kreimer. The core Hopf algebra. *Clay Math. Proc.*, 11:313–322, 2010, 0902.1223.
- [KW17] Ralph M. Kaufmann and Benjamin C. Ward. Feynman Categories. *Astérisque*, (387):vii+161, 2017. arXiv:1312.1269.
- [KY] Dirk Kreimer and Karen Yeats. Algebraic Interplay between Renormalization and Monodromy. preprint.
- [KY21] Ralph M. Kaufmann and Mo Yang. Higher categorical hopf algebras. *In preparation*, 2021.
- [KZ17] Ralph M. Kaufmann and Yongheng Zhang. Permutohedral structures on E_2 -operads. *Forum Math.*, 29(6):1371–1411, 2017.
- [Ler75] Pierre Leroux. Les catégories de Möbius. *Cahiers Topologie Géom. Différentielle*, 16(3):280–282, 1975.

- [Lod98] Jean-Louis Loday. *Cyclic homology*, volume 301 of *Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences]*. Springer-Verlag, Berlin, second edition, 1998. Appendix E by María O. Ronco, Chapter 13 by the author in collaboration with Teimuraz Pirashvili.
- [MN16] Matilde Marcolli and Xiang Ni. Rota-Baxter algebras, singular hypersurfaces, and renormalization on Kausz compactifications. *J. Singul.*, 15:80–117, 2016.
- [Moe01] I. Moerdijk. On the Connes-Kreimer construction of Hopf algebras. In *Homotopy methods in algebraic topology (Boulder, CO, 1999)*, volume 271 of *Contemp. Math.*, pages 311–321. Amer. Math. Soc., Providence, RI, 2001.
- [Mon93] Susan Montgomery. *Hopf algebras and their actions on rings*, volume 82 of *CBMS Regional Conference Series in Mathematics*. Published for the Conference Board of the Mathematical Sciences, Washington, DC; by the American Mathematical Society, Providence, RI, 1993.
- [MP12] J. P. May and K. Ponto. *More concise algebraic topology*. Chicago Lectures in Mathematics. University of Chicago Press, Chicago, IL, 2012. Localization, completion, and model categories.
- [NS82] Warren Nichols and Moss Sweedler. Hopf algebras and combinatorics. In *Umbral calculus and Hopf algebras (Norman, Okla., 1978)*, volume 6 of *Contemp. Math.*, pages 49–84. Amer. Math. Soc., Providence, R.I., 1982.
- [Qui67] Daniel G. Quillen. *Homotopical algebra*. Lecture Notes in Mathematics, No. 43. Springer-Verlag, Berlin, 1967.
- [Rad12] David E. Radford. *Hopf algebras*, volume 49 of *Series on Knots and Everything*. World Scientific Publishing Co. Pte. Ltd., Hackensack, NJ, 2012.
- [Sch87] William R. Schmitt. Antipodes and incidence coalgebras. *J. Combin. Theory Ser. A*, 46(2):264–290, 1987.
- [Tak71] Mitsuhiro Takeuchi. Free Hopf algebras generated by coalgebras. *J. Math. Soc. Japan*, 23:561–582, 1971.
- [TW74] Earl J. Taft and Robert Lee Wilson. On antipodes in pointed Hopf algebras. *J. Algebra*, 29:27–32, 1974.
- [Wil08] Simon Willerton. The twisted Drinfeld double of a finite group via gerbes and finite groupoids. *Algebr. Geom. Topol.*, 8(3):1419–1457, 2008.

DEPARTMENT OF MATHEMATICS, DEPARTMENT OF PHYSICS AND ASTRONOMY, PURDUE UNIVERSITY, WEST LAFAYETTE, IN, USA

Email address: `rkaufman@purdue.edu`

DEPARTMENT OF MATHEMATICS, PURDUE UNIVERSITY, WEST LAFAYETTE, IN, USA

Email address: `mo12@purdue.edu`