

Some Identities Related to the Second-Order Eulerian Numbers

Amy. M. Fu

School of Mathematics, Shanghai University of Finance and Economics

Shanghai, 200433, China

Email: fu.mei@sufe.edu.cn

Abstract: We express the Nörlund polynomials in terms of the second-order Eulerian numbers. Based on this expression, we derive several identities related to the Bernoulli numbers. In particular, we present a short proof of the problem raised by Rządowski and Urlińska.

Keywords: second-order Eulerian numbers; Nörlund polynomials; Bernoulli numbers.

AMS subject classifications: 05A19; 05A15

1 Introduction

The Stirling permutations were introduced by Gessel and Stanley [6]. For some related results on this subject, we refer to [1, 4, 7, 9]. Let Q_n be the multiset $\{1, 1, 2, 2, \dots, n, n\}$. A Stirling permutation of order n is a permutation of Q_n such that for each $1 \leq m \leq n$, the elements lying between two occurrences of m are greater than m .

The second-order Eulerian numbers $C_{n,k}$ count the Stirling permutations of order n with k decents, which satisfy the recurrence relation:

$$C_{n,k} = kC_{n-1,k} + (2n - k)C_{n-1,k-1} \quad (1.1)$$

with $C_{1,1} = 1$ and $C_{1,0} = 0$.

By exhibiting a bijection between the set of partitions of $[n + m]$ with m blocks and the bar permutations on the elements of Q_n with m bars, Gessel and Stanley [6] proved that

$$\sum_{m=0}^{\infty} S(n + m, m)x^m = \sum_{k=1}^n C_{n,k}x^k / (1 - x)^{2n+1}, \quad (1.2)$$

where $S(n, m)$ are the Stirling numbers of the second kind.

The Nörlund polynomials $B_n^{(z)}$ can be defined by the exponential generating function:

$$\sum_{n=0}^{\infty} B_n^{(z)} \frac{x^n}{n!} = \left(\frac{x}{e^x - 1} \right)^z. \quad (1.3)$$

Note that for fixed n , $B_n^{(z)}$ are polynomials in z with degree n . If $z = 0$, then we have $B_0^{(0)} = 1$ and $B_n^{(0)} = 0$ for $n \geq 1$. If $z = 1$, then $B_n^{(1)}$ are the classical Bernoulli numbers B_n , i.e.,

$$\sum_{n=0}^{\infty} B_n \frac{x^n}{n!} = \frac{x}{e^x - 1}. \quad (1.4)$$

If $z = n$, we have $B_n^{(n)} = (-1)^n c_n^{(2)}$, which are the Cauchy numbers of the second kind [3]:

$$\sum_{n=0}^{\infty} c_n^{(2)} \frac{x^n}{n!} = \frac{-x}{(1 - x) \ln(1 - x)}. \quad (1.5)$$

In [2], Carlitz showed that the Nörlund polynomials and the Stirling numbers of the second kind satisfy the relation:

$$S(m+n, m) = \binom{m+n}{n} B_n^{(-m)}. \quad (1.6)$$

Let $\langle z \rangle_n$ be the rising factorial defined by $\langle z \rangle_n = z(z+1) \cdots (z+n-1)$. The following result expresses the Nörlund polynomials in terms of the second-order Eulerian numbers.

Theorem 1.1 *We have*

$$B_n^{(z)} = \frac{n!}{(2n)!} \sum_{k=1}^n (-1)^k C_{n,k} \langle z \rangle_k \langle -z+n+1 \rangle_{n-k}. \quad (1.7)$$

Based on Theorem 1.1, we obtain several identities involving the convolutions of Bernoulli numbers. In particular, we give a short proof of the problem raised by Rządowski and Urlińska:

$$\int_0^1 \sum_{k=0}^{n-1} C_{n,k+1} u^{k+1} (u-1)^{2n-k} du = \frac{B_{n+1}}{n+1}. \quad (1.8)$$

By computing the integral

$$\int_0^1 u^{k+1} (u-1)^{2n-k} du = \frac{(-1)^k}{2(n+1)} \binom{2n+1}{k}^{-1},$$

we may restate (1.8) as follows.

Theorem 1.2 *We have*

$$\sum_{k=1}^n (-1)^{k-1} \binom{2n+1}{k}^{-1} C_{n,k} = 2B_{n+1}. \quad (1.9)$$

Notice that, by considering the partial derivative equation $\partial_t^n(\psi) = v_n(\psi)$, where ψ is defined by the Lambert W-function $\psi(t, x) = W(xe^{x+t})$ and $v_n(x)$ is given by

$$v_n(x) = -x(1+x)^{-2n+1} \sum_{k \geq 1} (-1)^k C_{n,k} x^k,$$

an alternative proof of (1.9) was discussed on Mathoverflow [11] recently.

Let H_n be the harmonic number defined by $H_n = \sum_{i=1}^n 1/i$. By using the p -adic method, Miki [8] proved that

$$\sum_{k=2}^{n-2} \frac{B_k}{k} \frac{B_{n-k}}{n-k} = 2H_n \frac{B_n}{n} + \sum_{k=2}^{n-2} \binom{n}{k} \frac{B_k}{k} \frac{B_{n-k}}{n-k}. \quad (1.10)$$

By computing the second derivative of the both sides of (1.7), then employing Miki's identity (1.10), we derive the following result involving the harmonic numbers.

Theorem 1.3 *We have*

$$\sum_{k=1}^n (-1)^k \binom{2n-1}{k-1}^{-1} (H_{2n-k} - H_{k-1}) C_{n,k} = \frac{n^2}{n-1} B_{n-1} + n \sum_{k=2}^{n-2} \frac{B_k}{k} \frac{B_{n-k}}{n-k}. \quad (1.11)$$

For $n \geq N$, Dilcher [5] proved that

$$\sum_{k_1+k_2+\dots+k_N=n} \binom{n}{k_1, k_2, \dots, k_N} B_{k_1} B_{k_2} \cdots B_{k_N} = N \binom{n}{N} \sum_{k=0}^{N-1} (-1)^{N-1-k} s(N, N-k) \frac{B_{n-k}}{n-k}, \quad (1.12)$$

where $s(n, k)$ is the Stirling number of the first kind. By setting $z = N$ in (1.7), then employing (1.12), we obtain the following identity.

Theorem 1.4 *Let N, n be nonnegative integers with $n \geq N$. We have*

$$\sum_{k=1}^n (-1)^k \binom{2n-1}{N+k-1}^{-1} C_{n,k} = 2n \sum_{k=0}^{N-1} (-1)^{N-1-k} s(N, N-k) \frac{B_{n-k}}{n-k}. \quad (1.13)$$

In particular, if we let $n = N$ in (1.13), the Cauchy numbers can be related by the second-order Eulerian numbers:

$$2c_n^{(2)} = \sum_{k=1}^n (-1)^{n-k} \binom{2n-1}{n+k-1}^{-1} C_{n,k}. \quad (1.14)$$

2 Proofs

If two polynomials in a single variable z agree for every nonnegative integer z , then they agree as polynomials. Therefore, to prove Theorem 1.1, it suffices to prove the following lemma.

Lemma 2.5 *Given a nonnegative integer m , we have*

$$B_n^{(-m)} = \frac{n!}{(2n)!} \sum_{k=1}^n (-1)^k C_{n,k} \langle -m \rangle_k \langle m+n+1 \rangle_{n-k}. \quad (2.1)$$

Proof: Equating the coefficients of x^m on the both sides of (1.2) leads to

$$S(m+n, m) = \sum_{k=1}^n \binom{2n+m-k}{2n} C_{n,k}. \quad (2.2)$$

Comparing (1.6) and (2.2) for any integer $m \geq 0$, we have

$$B_n^{(-m)} = \frac{m!n!}{(m+n)!} \sum_{k=1}^n \binom{2n+m-k}{2n} C_{n,k} = \frac{n!}{(2n)!} \sum_{k=1}^n (-1)^k C_{n,k} \langle -m \rangle_k \langle m+n+1 \rangle_{n-k}, \quad (2.3)$$

as desired. We complete the proof of Lemma 2.5, and the proof of Theorem 1.1 as well. \blacksquare

Observe that

$$\frac{d}{dx} \ln \left(\frac{e^x - 1}{x} \right) = \frac{1}{x} \left(\frac{-x}{e^{-x} - 1} - 1 \right) = \sum_{n=1}^{\infty} (-1)^n B_n \frac{x^{n-1}}{n!}.$$

Thus,

$$\ln \left(\frac{e^x - 1}{x} \right) = \sum_{n=1}^{\infty} (-1)^n \frac{B_n}{n} \frac{x^n}{n!} = \frac{x}{2} + \sum_{n=2}^{\infty} \frac{B_n}{n} \frac{x^n}{n!}.$$

Note that $B_1 = -\frac{1}{2}$ and $B_n = 0$ when n is odd and greater than 1.

Since

$$\frac{d^\ell}{dz^\ell} \left(\frac{x}{e^x - 1} \right)^z = \frac{d^\ell}{dz^\ell} e^{z \ln \left(\frac{x}{e^x - 1} \right)} = (-1)^\ell e^{z \ln \left(\frac{x}{e^x - 1} \right)} \left[\ln \left(\frac{e^x - 1}{x} \right) \right]^\ell,$$

we have

$$\frac{d}{dz} B_n^{(z)}|_{z=0} = -\frac{B_n}{n} \quad (2.4)$$

and

$$\frac{d^2}{dz^2} B_n^{(z)}|_{z=0} = \frac{n}{n-1} B_{n-1} + \sum_{k=2}^{n-2} \binom{n}{k} \frac{B_k}{k} \frac{B_{n-k}}{n-k}. \quad (2.5)$$

Using Miki's identity (1.10), we can rewrite (2.5) as

$$\frac{d^2}{dz^2} B_n^{(z)}|_{z=0} = \frac{n}{n-1} B_{n-1} + \sum_{k=2}^{n-2} \frac{B_k}{k} \frac{B_{n-k}}{n-k} - 2H_n \frac{B_n}{n}. \quad (2.6)$$

Proof of Theorem 1.2: Applying the derivation $d/dz|_{z=0}$ to both sides of (1.7), then using (2.4), we have

$$-\frac{B_n}{n} = \frac{1}{2n} \sum_{k=1}^n (-1)^k \binom{2n-1}{k-1}^{-1} C_{n,k}. \quad (2.7)$$

Mutiplied both sides by $-2n$, then replacing n by $n+1$, we have

$$\sum_{k=1}^{n+1} (-1)^{k-1} \binom{2n+1}{k-1}^{-1} C_{n+1,k} = 2B_{n+1}. \quad (2.8)$$

By the recurrence relation (1.1), the left-hand side of (2.8) can be rewritten as

$$\begin{aligned} & \sum_{k=1}^{n+1} (-1)^{k-1} \binom{2n+1}{k-1}^{-1} (kC_{n,k} + (2n+2-k)C_{n,k-1}) \\ &= \sum_{k=1}^n (-1)^{k-1} \frac{k!(2n+2-k)!}{(2n+1)!} C_{n,k} - \sum_{k=1}^n (-1)^{k-1} \frac{k!(2n+1-k)!(2n+1-k)}{(2n+1)!} C_{n,k} \\ &= \sum_{k=1}^n (-1)^{k-1} \frac{k!(2n+1-k)!}{(2n+1)!} C_{n,k} = \sum_{k=1}^n (-1)^{k-1} \binom{2n+1}{k}^{-1} C_{n,k}, \end{aligned}$$

as desired. We complete the proof of Theorem 1.2. ■

Proof of Theorem 1.3: Apply the derivation $d^2/dz^2|_{z=0}$ to both sides of (1.7). By (2.7), we have

$$\begin{aligned}\frac{d^2}{dz^2}B_n^{(z)}|_{z=0} &= \frac{1}{n} \sum_{k=1}^n (-1)^k \binom{2n-1}{k-1}^{-1} C_{n,k}(H_{k-1} - (H_{2n-k} - H_n)) \\ &= \frac{1}{n} \sum_{k=1}^n (-1)^k \binom{2n-1}{k-1}^{-1} C_{n,k}(H_{k-1} - H_{2n-k}) - 2H_n \frac{B_n}{n}.\end{aligned}$$

Combining with (2.6), after some arrangements, we complete the proof of Theorem 1.3. \blacksquare

We conclude this paper by the *proof of Theorem 1.4*. Setting $z = N$ in (1.3) leads to

$$\begin{aligned}\sum_{k=0}^{\infty} B_k^{(N)} \frac{x^k}{k!} &= \left(\frac{x}{e^x - 1} \right)^N = \left(\sum_{k=0}^{\infty} B_k \frac{x^k}{k!} \right)^N \\ &= \sum_{n=0}^{\infty} \frac{x^n}{n!} \sum_{k_1+k_2+\dots+k_N=n} \binom{n}{k_1, k_2, \dots, k_N} B_{k_1} B_{k_2} \cdots B_{k_N},\end{aligned}$$

where k_1, k_2, \dots, k_N are nonnegative integers.

Hence,

$$B_n^{(N)} = \sum_{k_1+k_2+\dots+k_N=n} \binom{n}{k_1, k_2, \dots, k_N} B_{k_1} B_{k_2} \cdots B_{k_N}.$$

By Theorem 1.1, for $1 \leq N \leq n$, we have

$$\begin{aligned}B_n^{(N)} &= \frac{n!}{(2n)!} \sum_{k=1}^n (-1)^k C_{n,k} \langle N \rangle_k \langle -N + n + 1 \rangle_{n-k} \\ &= \frac{N}{2n} \binom{n}{N} \sum_{k=1}^n (-1)^k C_{n,k} \binom{2n-1}{N+k-1}^{-1}.\end{aligned}$$

Combining with Dilcher's identity (1.12), we have

$$\frac{N}{2n} \binom{n}{N} \sum_{k=1}^n (-1)^k C_{n,k} \binom{2n-1}{N+k-1}^{-1} = N \binom{n}{N} \sum_{k=0}^{N-1} (-1)^{N-1-k} s(N, N-k) \frac{B_{n-k}}{n-k}.$$

Divided both sides by $\frac{N}{2n} \binom{n}{N}$, we complete the proof of Theorem 1.4. \blacksquare

References

- [1] M. Bóna, Real zeros and normal distribution for statistics on Stirling permutations defined by Gessel and Stanley, SIAM J. Discrete Math. **23**(2008), 401–406.
- [2] L. Carlitz, Note on Nörlund polynomials $B_n^{(z)}$, Proc. Amer. Math. Soc. **11**(1960), 452–455.
- [3] L. Comtet, Advanced Combinatorics, Reidel, Dordrecht, (1974).

- [4] W.Y.C. Chen and A.M. Fu, Context-free grammars for permutations and increasing trees, *Adv. in Appl. Math.* **82**(2017), 58–82.
- [5] K. Dilcher, Sums of products of Bernoulli numbers, *J. Number Theory* **60**(1996), 23–41.
- [6] I. Gessel and R.P. Stanley, Stirling polynomials, *J. Combin. Theory Ser. A* **24**(1978), 24–33.
- [7] S.-M. Ma and Y.-N. Yeh, Stirling permutations, cycle structure of permutations and perfect matchings, *Electron. J. Comb.* **22**(2015), #P4.43.
- [8] H. Miki, A relation between Bernoulli numbers, *J. Number Theory* **10**(1978), 297–302.
- [9] S. Janson, M. Kuba and A. Panholzer. Generalized Stirling permutations, families of increasing trees and urn models, *J. Combin. Theory Ser. A* **118**(2011), 94–114.
- [10] G. Rządowski and M. Urlińska, Some applications of the generalized Eulerian numbers, *J. Combin. Theory Ser. A* **63**(2019), 85–97.
- [11] Mathoverflow, <https://mathoverflow.net/questions/45756>.