

Understanding $P_{cs}(4459)$ as a hadronic molecule in the $\Xi_b^- \rightarrow J/\psi \Lambda K^-$ decay

Jun-Xu Lu,^{1,2} Ming-Zhu Liu,^{1,2} Rui-Xiang Shi,² and Li-Sheng Geng^{2,3,4,*}

¹*School of Space and Environment, Beihang University, Beijing 102206, China*

²*School of Physics, Beihang University, Beijing 102206, China*

³*Beijing Key Laboratory of Advanced Nuclear Materials and Physics,
Beihang University, Beijing 102206, China*

⁴*School of Physics and Microelectronics,
Zhengzhou University, Zhengzhou, Henan 450001, China*

Abstract

Recently, the LHCb Collaboration reported on the evidence for a hidden charm pentaquark state with strangeness, i.e., $P_{cs}(4459)$, in the $J/\psi \Lambda$ invariant mass distribution of the $\Xi_b^- \rightarrow J/\psi \Lambda K^-$ decay. In this work, assuming that $P_{cs}(4459)$ is a $\bar{D}^* \Xi_c$ molecular state, we study this decay via triangle diagrams $\Xi_b \rightarrow \bar{D}_s^{(*)} \Xi_c \rightarrow (\bar{D}^{(*)} \bar{K}) \Xi_c \rightarrow P_{cs} \bar{K} \rightarrow (J/\psi \Lambda) \bar{K}$. Our study shows that the production yield of a spin 3/2 $\bar{D}^* \Xi_c$ state is approximately one order of magnitude larger than that of a spin 1/2 state due to the interference of $\bar{D}_s \Xi_c$ and $\bar{D}_s^* \Xi_c$ intermediate states. We obtain a model independent constraint on the product of couplings $g_{P_{cs} \bar{D}^* \Xi_c}$ and $g_{P_{cs} J/\psi \Lambda}$. With the predictions of two particular molecular models as inputs, we calculate the branching ratio of $\Xi_b^- \rightarrow (P_{cs} \rightarrow) J/\psi \Lambda K^-$ and compare it with the experimental measurement. We further predict the lineshape of this decay which could be useful to future experimental studies.

* Corresponding author: lisheng.geng@buaa.edu.cn

I. INTRODUCTION

In 2015, two pentaquark states $P_c(4380)$ and $P_c(4450)$ were observed in the $J/\psi p$ invariant mass distribution of the $\Lambda_b \rightarrow J/\psi p K^-$ decay by the LHCb Collaboration [1], which have long been anticipated theoretically [2–11]. Since then, a large amount of theoretical works have been performed to understand their nature. The most popular interpretations include $\bar{D}^{(*)}\Sigma_c^{(*)}$ molecular states [12–20], compact pentaquark states [21–23], and kinematical effects [24, 25]. The experimental results were updated in 2019 with a data sample of almost ten times larger [26]. A new narrow state, $P_c(4312)$ was discovered. More interestingly, the original $P_c(4450)$ state splits into two states, $P_c(4440)$ and $P_c(4457)$. The masses and widths of these states are tabulated in Table I. After the 2019 update, the pentaquark states look more like $\bar{D}^{(*)}\Sigma_c^{(*)}$ molecules [27–40], but again non-molecular interpretations are possible, such as compact pentaquark states [41–43] and even double triangle singularities [44].

Most recently, the LHCb Collaboration reported on the first evidence for a structure in the $J/\psi \Lambda$ invariant mass distribution of the $\Xi_b^- \rightarrow J/\psi \Lambda K^-$ decay [45], hinting at the existence of a pentaquark state with strangeness, i.e., $P_{cs}(4459)$. It should be noted that the existence of pentaquark states with strangeness was predicted together with their non-strange counterparts in the molecular picture [2, 3]. The $P_{cs}(4459)$ state is located close to the $\bar{D}^*\Xi_c$ threshold, leading naturally to a molecular interpretation [46–51]. One interesting point to be noted is that in addition to the four P_c 's discovered experimentally, there may be three more candidates which strongly couple to $\bar{D}^*\Sigma_c^*$ with $J^P = \frac{1}{2}^-, \frac{3}{2}^-, \frac{5}{2}^-$, as dictated by the heavy quark spin symmetry (HQSS) [30, 35, 36, 39]. As for pentaquark states with strangeness, one expects 10 of them [47, 50, 52, 53].

In addition to the masses and widths of the pentaquark states, the LHCb Collaboration also reported the production yields of $P_c(4312)$, $P_c(4380)$, $P_c(4440)$, $P_c(4457)$, and $P_{cs}(4459)$, which are collected in Table I. One notes that the production yield for $P_c(4380)$ is one order of magnitude larger than that of $P_c(4312)$, which provides an explanation why $P_c(4312)$ was not observed in 2015. However, we note that the sum of the production yields of $P_c(4440)$ and $P_c(4457)$ is only half of that of $P_c(4450)$, which may indicate that something is missing, maybe a new resonance as suggested in several works [19, 54–56]. Clearly, understanding the production yields, particularly, the pattern shown in Table I, will greatly improve our understanding of the pentaquark states.

In the present work, we study the branching ratio of $\Xi_b^- \rightarrow P_{cs} K^-$.¹ The present work differs

¹ In Ref. [57], a similar mechanism has been applied to study the $D_s^+ \rightarrow \pi^+ \pi^0 \eta$ decay and it is shown that both

TABLE I. Resonance parameters of the newly discovered pentaquark states and their production ratios, defined as $R = \frac{\mathcal{B}(\Lambda_b(\Xi_b) \rightarrow P_c(P_{cs})\bar{K})\mathcal{B}(P_c(P_{cs}) \rightarrow J/\psi p(\Lambda))}{\mathcal{B}(\Lambda_b(\Xi_b) \rightarrow K J/\psi p(\Lambda))}$.

State	Mass (MeV)	Width (MeV)	R(%)
$P_c(4312)$	$4311.9 \pm 0.7^{+6.8}_{-0.6}$	$9.8 \pm 2.7^{+3.7}_{-4.5}$	$0.30 \pm 0.07^{+0.34}_{-0.09}$ [26]
$P_c(4380)$	$4380 \pm 8 \pm 29$	$205 \pm 18 \pm 86$	$8.4 \pm 0.7 \pm 4.2$ [1]
$P_c(4440)$	$4440.3 \pm 1.3^{+4.1}_{-4.7}$	$20.6 \pm 4.9^{+8.7}_{-10.1}$	$1.11 \pm 0.33^{+0.22}_{-0.10}$ [26] $4.1 \pm 0.5 \pm 1.1$ [1]
$P_c(4457)$	$4457.3 \pm 0.6^{+4.1}_{-1.7}$	$6.4 \pm 2.0^{+5.7}_{-1.9}$	$0.53 \pm 0.16^{+0.15}_{-0.13}$ [26]
$P_{cs}(4459)$	$4458.8 \pm 2.9^{+4.7}_{-1.1}$	$17.3 \pm 6.5^{+8.0}_{-5.7}$	$2.7^{+1.9+0.7}_{-0.6-1.3}$ [45]

from those of Refs. [58, 59] in two ways. First, the weak production formalism is different from that of Ref. [58] (see also Ref. [60]), which allows for a prediction of the absolute branching ratio of the $\Xi_b^- \rightarrow P_{cs}K^- \rightarrow J/\psi \Lambda K^-$ decay, within a molecular model. Compared to Ref. [59], we use different parameterizations of form factors and predict the lineshape of the $\Xi_b^- \rightarrow P_{cs}K^- \rightarrow J/\psi \Lambda K^-$ decay. Taking two molecular models for the $P_{cs}(4459)$ state [47], we compare the so-obtained branching ratios with the experimental data.

The present work is organized as follows. In Sec. II, we explain in detail the mechanism for the $P_{cs}(4459)$ production in the Ξ_b^- decay, which involves a weak interaction part and a strong interaction part. In Sec. III, we present the numerical results and compare with the experimental data, followed by a short summary in Sec. IV.

II. THEORETICAL FRAMEWORK

Assuming that $P_{cs}(4459)$ is a molecule mainly composed of $\bar{D}^*\Xi_c$, the $\Xi_b^- \rightarrow P_{cs}K^-$ decay can proceed as shown in diagram (a) of Fig. 1. The Ξ_b state first decays into Ξ_c by emitting a W^- boson which is then converted into a pair of $\bar{c}s$, which after hadronization turns into a $D_s^{(*)}$. Next the $D_s^{(*)}$ meson emits a kaon and a $\bar{D}^*$. The final state interaction of $\bar{D}^*\Xi_c$ dynamically generates the $P_{cs}(4459)$ state which then decays into $J/\psi \Lambda$, as shown in Fig. 2.

In addition to the W -emission diagram discussed above, the Ξ_b^- decay can also proceed via the branching ratio and the $\pi^{+(0)}\eta$ lineshape are well described. In particular, the large branching ratio of $D_s^+ \rightarrow a_0^+\pi^0(a_0^0\pi^+)$ is naturally explained, while for a pure W -annihilation process one would expect a much smaller value.




FIG. 1. External W -emission (a) and internal W -conversion (b) mechanism for the Ξ_b decay.

the internal W -exchange mechanism shown in diagram (b) of Fig. 1. The ssd cluster can either directly hadronize into a Ξ^- or, by picking up a pair of $q\bar{q}$ from the vacuum, hadronizes into ΛK^- . The former is indeed a main decay channel of Ξ_b^- [61], while the latter has been studied in Ref. [58].



FIG. 2. Triangle diagrams for the $\Xi_b^- \rightarrow P_{cs} K^-$ decay.

A. Branching ratio of $\Xi_b \rightarrow P_{cs} K^-$

In the following, we describe how to calculate the diagrams of Fig. 2. The effective Lagrangian responsible for the $\Xi_b \rightarrow \Xi_c \bar{D}_s^{(*)}$ decay reads

$$\begin{aligned}\mathcal{L}_{\Xi_b \Xi_c D_s} &= i \bar{\Xi}_c (A + B \gamma_5) \Xi_b D_s, \\ \mathcal{L}_{\Xi_b \Xi_c D_s^*} &= \bar{\Xi}_c (A_1 \gamma_\mu \gamma_5 + A_2 \frac{p_{2\mu}}{m} \gamma_5 + B_1 \gamma_\mu + B_2 \frac{p_{2\mu}}{m}) \Xi_b D_s^{*\mu}.\end{aligned}\quad (1)$$

The A_1 , A_2 , B_1 , B_2 , A , and B can be expressed with the six form factors describing the $\Xi_b \rightarrow \Xi_c$ transition [62] as ²

$$\begin{aligned}A &= \lambda f_{D_s} [(m - m_2) f_1^V + \frac{m_1^2}{m} f_3^V], & B &= \lambda f_{D_s} [(m + m_2) f_1^A - \frac{m_1^2}{m} f_3^A], \\ A_1 &= -\lambda f_{D_s^*} m_1 [f_1^A - f_2^A \frac{m - m_2}{m}], & B_1 &= \lambda f_{D_s^*} m_1 [f_1^V + f_2^V \frac{m + m_2}{m}], \\ A_2 &= 2\lambda f_{D_s^*} m_1 f_2^A, & B_2 &= -2\lambda f_{D_s^*} m_1 f_2^V,\end{aligned}\quad (2)$$

² Here we adopt the convention for the form factors of Ref. [63] in which there exists an extra minus in front of f_2^A and f_2^V .

TABLE II. Parameters $F(0)$, Λ_1 , Λ_2 in the form factors of the $\Xi_b \rightarrow \Xi_c$ transition form factors.

	F_1^V	F_2^V	F_3^V	F_1^A	F_2^A	F_3^A
$F(0)$	0.467	0.145	0.086	0.447	-0.035	-0.278
$\Lambda_1(\text{GeV})$	5.10	4.89	6.14	4.69	4.97	4.58
$\Lambda_2(\text{GeV})$	9.03	5.46	6.28	12.20	5.05	7.08

where $\lambda = \frac{G_F}{\sqrt{2}} V_{cb} V_{cs} a_1$ with $a_1 = 1.07$ [64]. The decay constants $f_{D_s^{(*)}}$ for $\bar{D}_s$ and $\bar{D}_s^*$ are set to be 0.247 GeV and m, m_1, m_2 refer to the masses of $\Xi_b, \bar{D}_s^{(*)}$, and Ξ_c respectively.

Following the double-pole parametrization proposed in Ref. [65], one can rewrite the form factors as

$$f_i^{V/A}(q^2) = F_i^{V/A}(0) \frac{\Lambda_1^2}{q^2 - \Lambda_1^2} \frac{\Lambda_2^2}{q^2 - \Lambda_2^2}. \quad (3)$$

Fitting to the results of the relativistic quark-diquark model [63], we can obtain the values of $F(0)$, Λ_1 , and Λ_2 , which are tabulated in Table II.

The effective Lagrangians for the $P_{cs} \rightarrow \bar{D}^* \Xi_c$ and $\bar{D}_s^{(*)} \rightarrow \bar{D}^* \bar{K}$ read

$$\begin{aligned} \mathcal{L}_{P_{cs1} \Xi_c \bar{D}^*} &= g_{P_{cs1} \Xi_c \bar{D}^*} \bar{\Xi}_c \gamma_5 (g_{\mu\nu} - \frac{p_\mu p_\nu}{m_{P_{cs}}^2}) \gamma^\nu P_{cs} D^{*\mu}, \\ \mathcal{L}_{P_{cs2} \Xi_c \bar{D}^*} &= g_{P_{cs2} \Xi_c \bar{D}^*} \bar{\Xi}_c P_{cs2\mu} D^{*\mu}, \\ \mathcal{L}_{KD_s D^*} &= i g_{KD_s D^*} D^{*\mu} [\bar{D}_s \partial_\mu K - (\partial_\mu \bar{D}_s) K] + H.c., \\ \mathcal{L}_{KD_s^* D^*} &= -g_{KD_s^* D^*} \epsilon^{\mu\nu\alpha\beta} [\partial_\mu \bar{D}_\nu^* \partial_\alpha D_{s\beta}^* \bar{K} + \partial_\mu D_\nu^* \partial_\alpha \bar{D}_{s\beta}^* K] + H.c., \end{aligned} \quad (4)$$

where P_{cs1} and P_{cs2} denote the $P_{cs}(4459)$ state with $J^P = \frac{1}{2}^-$ and $\frac{3}{2}^-$, respectively. The $g_{KD_s D^*}$ and $g_{KD_s^* D^*}$ are the kaon meson couplings to $D_s D^*$ and $D_s^* D^*$, respectively. We take $g_{KD_s D^*} = 5.0$ and $g_{KD_s^* D^*} = 7.0 \text{ GeV}^{-1}$ in the present work, which are extracted from Ref. [66]. The $g_{P_{cs1} \Xi_c \bar{D}^*}$ and $g_{P_{cs2} \Xi_c \bar{D}^*}$ are the couplings between P_{cs} and its components, whose values are not known a priori, but can be computed with the compiteness conditions [67–69] or in molecular models, e.g., Refs. [47], or in lattice QCD.

With the effective Lagrangians above, the decay amplitudes for $\Xi_b(p) \rightarrow \bar{D}_s^{(*)}(p_1) \Xi_c(p_2) [\bar{D}^*(q)] \rightarrow$

$\bar{K}(p_3)P_{cs1}(p_4)$ read

$$\begin{aligned}
\mathcal{M}_{P_{cs1}} &= \mathcal{M}_{\bar{D}_s}^{P_{cs1}} + \mathcal{M}_{\bar{D}_s^*}^{P_{cs1}}, \\
\mathcal{M}_{\bar{D}_s}^{P_{cs1}} &= i^3 \int \frac{d^4 q}{(2\pi)^4} [g_{p_{cs1}\Xi_c \bar{D}^*} \bar{u}(p_4) \gamma^\nu \gamma_5 (g_{\mu\nu} - \frac{p_{4\mu} p_{4\nu}}{m_4^2})] (\not{p}_2 + m_2) \\
&\quad \times [i(A + B\gamma_5)u(p)] [-g_{KD^*D_s}(p_1 + p_3)_\alpha] (-g^{\mu\alpha} + \frac{q^\mu q^\alpha}{m_E^2}) \\
&\quad \times \frac{1}{p_1^2 - m_1^2} \frac{1}{p_2^2 - m_2^2} \frac{1}{q^2 - m_E^2} \mathcal{F}(q^2, m_E^2), \\
\mathcal{M}_{\bar{D}_s^*}^{P_{cs1}} &= i^3 \int \frac{d^4 q}{(2\pi)^4} [g_{p_{cs1}\Xi_c \bar{D}^*} \bar{u}(p_4) \gamma^\nu \gamma_5 (g_{\mu\nu} - \frac{p_{4\mu} p_{4\nu}}{m_4^2})] (\not{p}_2 + m_2) \\
&\quad \times [(A_1 \gamma_\alpha \gamma_5 + A_2 \frac{p_{2\alpha}}{m} \gamma_5 + B_1 \gamma_\alpha + B_2 \frac{p_{2\alpha}}{m}) u(p)] \\
&\quad \times [-g_{KD^*D_s^*} \varepsilon_{\rho\lambda\eta\tau} q^\rho p_1^\eta] (-g^{\mu\lambda} + \frac{q^\mu q^\lambda}{m_E^2}) (-g^{\alpha\tau} + \frac{p_1^\alpha p_1^\tau}{m_1^2}) \\
&\quad \times \frac{1}{p_1^2 - m_1^2} \frac{1}{p_2^2 - m_2^2} \frac{1}{q^2 - m_E^2} \mathcal{F}(q^2, m_E^2). \tag{5}
\end{aligned}$$

The decay amplitudes of $\Xi_b(p) \rightarrow \bar{D}_s^{(*)}(p_1) \Xi_c(p_2) [\bar{D}^*(q)] \rightarrow \bar{K}(p_3) P_{cs2}(p_4)$ read

$$\begin{aligned}
\mathcal{M}_{P_{cs2}} &= \mathcal{M}_{\bar{D}_s}^{P_{cs2}} + \mathcal{M}_{\bar{D}_s^*}^{P_{cs2}}, \\
\mathcal{M}_{\bar{D}_s}^{P_{cs2}} &= i^3 \int \frac{d^4 q}{(2\pi)^4} [-ig_{p_{cs2}\Xi_c \bar{D}^*} \bar{u}_\mu(p_4)] (\not{p}_2 + m_2) [i(A + B\gamma_5) \\
&\quad u(p)] [-g_{KD^*D_s}(p_1 + p_3)_\nu] (-g^{\mu\nu} + \frac{q^\mu q^\nu}{m_E^2}) \\
&\quad \times \frac{1}{p_1^2 - m_1^2} \frac{1}{p_2^2 - m_2^2} \frac{1}{q^2 - m_E^2} \mathcal{F}(q^2, m_E^2), \\
\mathcal{M}_{\bar{D}_s^*}^{P_{cs2}} &= i^3 \int \frac{d^4 q}{(2\pi)^4} [-ig_{p_{cs2}\Xi_c \bar{D}^*} \bar{u}_\sigma(p_4)] (\not{p}_2 + m_2) \\
&\quad \times [(A_1 \gamma_\rho \gamma_5 + A_2 \frac{p_{2\rho}}{m} \gamma_5 + B_1 \gamma_\rho + B_2 \frac{p_{2\rho}}{m}) u(p)] \\
&\quad \times [-g_{KD^*D_s^*} \varepsilon_{\mu\nu\alpha\beta} q^\mu p_1^\alpha] (-g^{\sigma\nu} + \frac{q^\sigma q^\nu}{m_E^2}) (-g^{\rho\beta} + \frac{p_1^\rho p_1^\beta}{m_1^2}) \\
&\quad \times \frac{1}{p_1^2 - m_1^2} \frac{1}{p_2^2 - m_2^2} \frac{1}{q^2 - m_E^2} \mathcal{F}(q^2, m_E^2). \tag{6}
\end{aligned}$$

We follow Ref. [37] and introduce a monopole form factor to depict the off-shell effect of the exchanged $\bar{D}^*$ mesons,

$$\mathcal{F}(q^2, m^2) = \frac{m^2 - \Lambda^2}{q^2 - \Lambda^2}, \tag{7}$$

where $\Lambda = m + \alpha \Lambda_{QCD}$ with $\Lambda_{QCD} = 220$ MeV, and α is a model parameter. In this way, the triangle diagrams are free of any ultraviolet divergence. Collecting all the pieces together, the

decay width for $\Xi_b \rightarrow P_{cs} \bar{K}$ could be expressed as

$$\Gamma = \frac{1}{2J+1} \frac{1}{8\pi} \frac{1}{m_{\Xi_b}^2} |\vec{p}| \sum |\mathcal{M}_{P_{cs1}/P_{cs2}}|^2, \quad (8)$$

where $|\vec{p}|$ denotes the momentum of $\bar{K}$ or P_{cs} in the rest frame of Ξ_b .




FIG. 3. Feynman diagrams for the $\Xi_b^- \rightarrow (P_{cs} \rightarrow) J/\psi \Lambda K^-$ decay.

B. $J/\psi \Lambda$ invariant mass distribution of the $\Xi_b^- \rightarrow K^- P_{cs} \rightarrow K^- J/\psi \Lambda$

With the weak decay vertices described in Eq. (1) and Eq. (2), we can further work out the invariant mass distribution of the $\Xi_b^- \rightarrow J/\psi \Lambda K^-$ decay. Parameterizing the intermediate $P_{cs}(4459)$ state with a Breit-Wigner resonance, the amplitudes of Fig. 3 read

$$\begin{aligned} \mathcal{A} &= \mathcal{A}_{\bar{D}_s} + \mathcal{A}_{\bar{D}_s^*}, \\ \mathcal{A}_{\bar{D}_s} &= i \int \frac{d^4 q}{(2\pi)^4} \bar{u}(p_5) \epsilon^\mu(p_4) \mathcal{A}_{\mu\alpha P_{cs1}/P_{cs2}} \cdot (\not{p}_2 + m_2) \cdot \mathcal{A}_{\Xi_b \Xi_c \bar{D}_s} u(P) \\ &\quad \times \frac{\mathcal{A}_{K D_s D^*}^\alpha \cdot (-g_{\mu\alpha} + \frac{q^\mu q^\alpha}{m_E^2})}{(p_2^2 - m_2^2)(p_1^2 - m_1^2)(q^2 - m_E^2)} \mathcal{F}(q^2, m_E^2), \\ \mathcal{A}_{\bar{D}_s^*} &= i \int \frac{d^4 q}{(2\pi)^4} \bar{u}(p_5) \epsilon^\mu(p_4) \mathcal{A}_{\mu\alpha P_{cs1}/P_{cs2}} \cdot (\not{p}_2 + m_2) \cdot \mathcal{A}_{\Xi_b \Xi_c \bar{D}_s^*}^\beta u(P) \\ &\quad \times \frac{\mathcal{A}_{K D_s^* D^*}^{\nu\beta} \cdot (-g_{\mu\nu} + \frac{q^\mu q^\nu}{m_E^2})(-g_{\alpha\beta} + \frac{q^\alpha q^\beta}{m_1^2})}{(p_2^2 - m_2^2)(p_1^2 - m_1^2)(q^2 - m_E^2)} \mathcal{F}(q^2, m_E^2), \end{aligned} \quad (9)$$

where $(p_4 + p_5)^2 = p_{45}^2 = M_{45}^2$ denotes the invariant mass of the $J/\psi\Lambda$ final state and

$$\begin{aligned}
\mathcal{A}_{\Xi_b \Xi_c \bar{D}_s} &= i(A + B\gamma_5), \\
\mathcal{A}_{\Xi_b \Xi_c \bar{D}_s^*}^\rho &= A_1 \gamma_\rho \gamma_5 + A_2 \frac{p_{2\rho}}{m} \gamma_5 + B_1 \gamma_\rho + B_2 \frac{p_{2\rho}}{m}, \\
\mathcal{A}_{K D_s D^*}^\alpha &= -g_{K D_s D^*} (p_1^\alpha + p_3^\alpha), \\
\mathcal{A}_{K D_s^* D^*}^{\nu\beta} &= -g_{K D_s^* D^*} \epsilon_{\mu\nu\alpha\beta} (q^\mu p_1^\alpha), \\
\mathcal{A}_{P_{cs1}}^{ua} &= \frac{g_{P_{cs1} \Xi_c \bar{D}^*} g_{P_{cs1} J/\psi \Lambda}}{p_{45}^2 - m_{P_{cs}}^2 + i\Gamma_{P_{cs}} m_{P_{cs}}} \gamma_5 (g^{\mu\nu} - \frac{p_{45}^\mu p_{45}^\nu}{m_{P_{cs}}^2}) \gamma_\nu \cdot (\not{p}_{45} + m_{P_{cs}}) \cdot \gamma_5 (g_{ab} - \frac{p_{45}^a p_{45}^b}{m_{P_{cs}}^2}) \gamma_b, \\
\mathcal{A}_{P_{cs2}}^{\mu\nu} &= \frac{g_{P_{cs2} \Xi_c \bar{D}^*} g_{P_{cs2} J/\psi \Lambda}}{p_{45}^2 - m_{P_{cs}}^2 + i\Gamma_{P_{cs}} m_{P_{cs}}} (\not{p}_{45} + m_{P_{cs}}) \cdot (-g^{\mu\nu} + \frac{\gamma^\mu \gamma^\nu}{d-1} + \frac{\gamma^\mu p_{45}^\nu - \gamma^\nu p_{45}^\mu}{(d-1)m_{P_{cs}}} + \frac{d-2}{(d-1)m_{P_{cs}}^2} p_{45}^\mu p_{45}^\nu).
\end{aligned}$$

The partial decay rate for $\Xi_b \rightarrow J/\psi \Lambda \bar{K}$ as a function of the invariant mass $M_{J/\psi \Lambda}$ then reads

$$\frac{d\Gamma}{dM_{J/\psi \Lambda}} = \frac{1}{2J+1} \frac{1}{64\pi^3} \frac{1}{m_{\Xi_b}^2} |p_3^*| |p_4| \sum |\mathcal{A}|^2, \quad (10)$$

with

$$\begin{aligned}
p_3^* &= \frac{\sqrt{(m_{\Xi_b}^2 - (m_K - M_{J/\psi \Lambda})^2)(m_{\Xi_b}^2 - (m_K + M_{J/\psi \Lambda})^2)}}{2m_{\Xi_b}}, \\
p_4 &= \frac{\sqrt{(M_{J/\psi \Lambda}^2 - (m_{J/\psi} - m_\Lambda)^2)(M_{J/\psi \Lambda}^2 - (m_{J/\psi} + m_\Lambda)^2)}}{2M_{J/\psi \Lambda}}.
\end{aligned} \quad (11)$$

III. RESULTS AND DISCUSSIONS

In this section, we explore the decay mechanism proposed in this work. We divide our discussions into two categories, those only depend on the decay mechanism explored and those depend on a particular molecular model.

In our framework, the parameter α is not known, though its value is often assumed to be about 1 [70–73]. Therefore, we first study how the calculated branching ratios depend on the value of α . Varying α from 0.8 to 1.2, we plot the values of $\text{Br}[\Xi_b \rightarrow P_{cs1}(P_{cs2})\bar{K}]/g_{P_{cs1/2} \Xi_c \bar{D}^*}^2$ in Fig. 4. One can see that the branching ratios for P_{cs1} and P_{cs2} are moderately sensitive to the value of α in the range studied. As a consequence, in the following, we will take $\alpha = 1.0 \pm 0.1$ to take into account the uncertainties from α .




FIG. 4. Dependence of the branching ratios $\text{Br}[\Xi_b \rightarrow P_{cs} \bar{K}]$ on α .

A. Model independent predictions

To compute the absolute branching ratio $\text{Br}[\Xi_b \rightarrow P_{cs} \bar{K}]$, we need to know the coupling constants $g_{P_{cs1}\Xi_c \bar{D}^*}$ and $g_{P_{cs2}\Xi_c \bar{D}^*}$. They can be determined model independently with the compositeness conditions [67–69], as was done in, e.g., Ref. [32] for the pentaquark states. With the experimental mass of P_{cs} , the couplings read $g_{P_{cs1}\Xi_c \bar{D}^*} = 1.59$ and $g_{P_{cs2}\Xi_c \bar{D}^*} = 2.76$, corresponding to a cutoff $\Lambda = 1.0$ GeV (more details can be found in the Appendix). With these couplings, we find, surprisingly, that the branching ratio for the P_{cs} state with $J^P = 3/2^-$ is approximately one order of magnitude larger than that for the P_{cs} state with $J^P = 1/2^-$, which are

$$\begin{aligned} \text{Br}[\Xi_b \rightarrow P_{cs1} \bar{K}] &= (9.84 \pm 1.04) \times 10^{-5}, \\ \text{Br}[\Xi_b \rightarrow P_{cs2} \bar{K}] &= (9.48 \pm 1.08) \times 10^{-4}. \end{aligned} \tag{12}$$

In addition, using the experimental branching ratio $\text{Br}[\Xi_b \rightarrow J/\psi \Lambda \bar{K}] = (2.31 \pm 1.37) \times 10^{-4}$ (see the Appendix how to derive this), the mass, width, and branching ratio R of the P_{cs} state given in Table I as inputs, we can provide a model independent constraint on the product of the two couplings in Eq. (4), $g_{P_{cs} \bar{D}^* \Xi_c}$ and $g_{P_{cs} J/\psi \Lambda}$, within the decay mechanism studied in the present work. The experimental branching ratio given in Table I is $R = 2.7^{+2.0}_{-1.4}\%$. Using the formalism

detailed in Sec.IIB, we obtain

$$g_{P_{cs}\Xi_c\bar{D}^*}g_{P_{cs}J/\psi\Lambda} = \begin{cases} 0.18_{-0.08}^{+0.10} & \text{for } J^P = \frac{1}{2}^- \\ 0.17_{-0.08}^{+0.10} & \text{for } J^P = \frac{3}{2}^- \end{cases}. \quad (13)$$

The above product can be used to constrain molecular models.

B. Comparison with models

In order to produce the branching ratio R defined in the introduction, in addition to the information derived above, we need to know the partial decay width of P_{cs} into $J/\psi\Lambda$. For this, we turn to specific molecular models. In the following, we study the unitary approach of Ref. [47] and the one-boson-exchange (OBE) model of Ref. [51], calculate the branching ratio R , and compare with the LHCb measurement.

First, we focus on Ref. [47]. Note that the difference between the definition of their couplings and ours (see the Appendix for details) and with the branching ratios $\text{Br}[P_{cs} \rightarrow J/\psi\Lambda] = 3.31\%$ for P_{cs1} and 14.68% for P_{cs2} from Ref. [47], we obtain the couplings as $g_{P_{cs1}J/\psi\Lambda} = 0.07$ and $g_{P_{cs2}J/\psi\Lambda} = 0.27$. The branching ratios R for the spin-parity assignment $1/2^-$ and $3/2^-$ are found to be

$$\begin{aligned} R_{P_{cs1}} &= \frac{\text{Br}[\Xi_b \rightarrow P_{cs1}\bar{K}]\text{Br}[P_{cs1} \rightarrow J/\psi\Lambda]}{\text{Br}[\Xi_b \rightarrow J/\psi\Lambda\bar{K}]} = 1.4 \pm 0.8\%, \\ R_{P_{cs2}} &= \frac{\text{Br}[\Xi_b \rightarrow P_{cs2}\bar{K}]\text{Br}[P_{cs2} \rightarrow J/\psi\Lambda]}{\text{Br}[\Xi_b \rightarrow J/\psi\Lambda\bar{K}]} = 60.3 \pm 36.4\%. \end{aligned} \quad (14)$$

In the one-boson-exchange (OBE) model of Ref. [51], the $P_{cs}(4459)$ state is interpreted as a $J^P = 3/2^-$ molecular state and the partial decay width of $P_{cs} \rightarrow J/\psi\Lambda$ is estimated to be $0.06 \sim 0.2$ MeV. The main decay mode is found to be $P_{cs} \rightarrow K^*\Xi(\omega\Lambda)$, which accounts for 80% of the total decay width. These numbers lead to an even smaller branching ratio $\text{Br}[P_{cs} \rightarrow J/\psi\Lambda] = 0.6\% - 0.8\%$ corresponding to the total decay width ranging from 10 to 25 MeV. For the coupling between the P_{cs} state with $J^P = 3/2^-$ and its components, we adopt the value $g_{P_{cs2}\Xi_c\bar{D}^*} = 2.76$ obtained from the compositeness condition. Using 0.7% as the central value for $\text{Br}[P_{cs} \rightarrow J/\psi\Lambda]$ and 0.1% as its error, we obtain

$$R_{P_{cs2}} = \frac{\text{Br}[\Xi_b \rightarrow P_{cs2}\bar{K}]\text{Br}[P_{cs2} \rightarrow J/\psi\Lambda]}{\text{Br}[\Xi_b \rightarrow J/\psi\Lambda\bar{K}]} = 2.87 \pm 1.75\%. \quad (15)$$

All these numbers are compared with the LHCb measurement in Fig. 5. It is clear that the result of the OBE model seems to agree with the experimental measurement, as well as the $J^P = 1/2^-$




FIG. 5. Branching ratios R for $P_{cs1}(J^P = 1/2^-)$ and $P_{cs2}(J^P = 3/2^-)$. The red square and blue circle denote our results given in Eq. (14), while the black diamond denotes the LHCb measurement [45]. The results with the partial decay width obtained from the OBE model of Ref. [51] (green triangle) is also shown for comparison.

case of the unitary approach. The predicted branching ratio for $J^P = 3/2^-$ in the unitary approach is however much larger than the experimental number.




FIG. 6. Invariant mass distribution of $\Xi_b^- \rightarrow P_{cs} K^- \rightarrow J/\psi \Lambda K^-$ for P_{cs} with $J^P = 1/2^-$ (left) and $J^P = 3/2^-$ (right).

Finally, in Fig. 6, we show the $J/\psi \Lambda$ invariant mass distribution of the $\Xi_b^- \rightarrow J/\psi \Lambda K^-$ decay with all the relevant couplings provided by the unitary approach of Ref. [47] (see the Appendix

for more details). They might be useful for future experimental searches.

IV. SUMMARY

In this work, we studied the decay of $\Xi_b^- \rightarrow P_{cs}K^- \rightarrow J/\psi\Lambda K^-$ via a triangle mechanism. The decay consists of three steps. First, Ξ_b^- decays weakly into $\bar{D}_s^{(*)}$ and Ξ_c via the external W -emission diagram. Using the relevant form factors determined in the relativistic quark-diquark model, this weak interaction part can be computed without any free parameters. Followed by the creation of $\bar{D}_s^{(*)}$ and Ξ_c in the first step, the $\bar{D}_s^{(*)}$ state then emits a kaon and a $\bar{D}^*$. The $\bar{D}^*$ and Ξ_c interact with each other to dynamically generate the $P_{cs}(4459)$ state, which then decays into $J/\psi\Lambda$. From such a decay mechanism, we derived a constraint on the product of couplings of the $P_{cs}(4459)$ state to the $\bar{D}^*\Xi_c$ and $J/\psi\Lambda$ channels. Determining the coupling between P_{cs} and the $\bar{D}^*\Xi_c$ channel using the compositeness condition, we predicted the branching ratio $\text{Br}[\Xi_b^- \rightarrow P_{cs}K^-]$. These can be useful to understanding the nature of $P_{cs}(4459)$ as a molecular state.

Using the predicted couplings by the unitary approach [47] and the one-boson exchange model of Ref. [51], we calculated the branching ratios $\text{Br}[\Xi_b^- \rightarrow (P_{cs} \rightarrow J/\psi\Lambda)K^-]$. We found that in the unitary approach, the $J^P = 1/2$ assignment is preferred, while the $J^P = 3/2$ assignment gives a branching ratio much larger than the experimental measurement. On the other hand, the $3/2$ assignment in the one-boson-exchange model of Ref. [51] yields a branching ratio in agreement with the LHCb data. This can be traced back to the drastically different partial decay width of $P_{cs} \rightarrow J/\psi\Lambda$.

In principle, the present formalism can also be utilized to study the $\Lambda_b \rightarrow J/\psi p K^-$ decay, where the four pentaquark states, $P_c(4312)$, $P_c(4380)$, $P_c(4440)$, and $P_c(4457)$, were discovered. This has been explored in Ref. [37], which, however, suffers from the fact that the weak decay $\Lambda_b \rightarrow \bar{D}_s^{(*)}\Sigma_c$ is suppressed because the ud quark pair in Λ_b has spin 0, but that in Σ_c has spin 1. As a result, the relevant transition form factors are not known and therefore one could not arrive at a quantitative determination of the branching ratios. In addition, compared to the present case, the suppression of the $\Lambda_b \rightarrow \bar{D}_s^{(*)}\Sigma_c$ transition indicates that other mechanisms may play a role than the external W -emission studied in the present work, which complicates the study a lot.

V. ACKNOWLEDGEMENTS

We thank Qi Wu, Dian-Yong Chen, and Li-Ming Zhang for useful communications. This work is partly supported by the National Natural Science Foundation of China under Grants No.11735003, No.11975041, and No.11961141004, and the fundamental Research Funds for the Central Universities.

VI. APPENDIX

A. Couplings from the compositeness conditions

With the assumption that the P_{cs} state observed by the LHCb Collaboration can be interpreted as a molecular state of $\bar{D}^*\Xi_c$ with $J^P = 1/2^-$ or $J^P = 3/2^-$, we can calculate the couplings between the P_{cs} state and its components with the compositeness condition, which is quite similar to what was done in Refs. [32, 74].




FIG. 7. Mass operators of the P_c

According to the compositeness rule [67–69], the coupling constant $g_{P_{cs1/2}\Xi_c\bar{D}^*}$ can be determined from the fact that the renormalization constant of the wave function of a composite particle should be zero. That is,

$$Z_{P_{cs}} = 1 - \frac{d\Sigma_{P_{cs}}(k_0)}{dk_0} \Big|_{k_0=m_{P_{cs}}} = 0, \quad (16)$$

where $\Sigma_{P_{cs}}$ denotes the self-energy of P_{cs1} and P_{cs2} . Applying the effective Lagrangians listed in Eq. (4), the self-energy $\Sigma_{P_{cs1/2}}$ reads

$$\begin{aligned} \Sigma_{P_{cs1}}(k_0) &= g_{P_{cs1}\Xi_c\bar{D}^*}^2 \int \frac{d^4k_1}{i(2\pi)^4} \{ \Phi^2[-(k_1 - k_0\omega_{\Xi_c})^2] \mathcal{A}_{P_{cs1}}^\mu \frac{1}{k_1 - m_{\Xi_c}} \mathcal{A}_{P_{cs1}}^\nu \frac{-g^{\mu\nu} + \frac{k_2^\mu k_2^\nu}{m_{D^*}^2}}{k_2^2 - m_{D^*}^2}, \\ \Sigma_{P_{cs2}}^{\mu\nu}(k_0) &= g_{P_{cs2}\Xi_c\bar{D}^*}^2 \int \frac{d^4k_1}{i(2\pi)^4} \{ \Phi^2[-(k_1 - k_0\omega_{\Xi_c})^2] \frac{1}{k_1 - m_{\Xi_c}} \frac{-g^{\mu\nu} + \frac{k_2^\mu k_2^\nu}{m_{D^*}^2}}{k_2^2 - m_{D^*}^2}, \end{aligned} \quad (17)$$

with

$$\begin{aligned}\omega_{\Xi_c} &= \frac{m_{\Xi_c}}{m_{\Xi_c} + m_{D^*}}, \\ \mathcal{A}_{P_{cs1}}^\mu &= \gamma_5(g^{\mu\nu} - \frac{k_0^\mu k_0^\nu}{m_{P_{cs}}^2})\gamma_\nu.\end{aligned}\tag{18}$$

The $\Phi[-p^2] = \exp(p^2/\Lambda^2)$ is the Fourier transformation of the correlation in the Gaussian form with Λ being the size parameter which characterizes the distribution of components inside the molecule. With all the formula above and taking $\Lambda = 1.0$ GeV, we obtain the couplings between the P_{cs} states and $\bar{D}^*\Xi_c$, which are $g_{P_{cs1}\Xi_c\bar{D}^*} = 1.59$ for $J^P = 1/2^-$ and $g_{P_{cs2}\Xi_c\bar{D}^*} = 2.76$ for $J^P = 3/2^-$.

B. Determination of the branching ratio $\text{Br}[\Xi_b \rightarrow J/\psi\Lambda\bar{K}]$

Experimentally, the branching ratio of $\Xi_b \rightarrow J/\psi\Lambda\bar{K}$ has been measured to be [75]

$$\frac{f_{\Xi_b}}{f_{\Lambda_b}} \times \frac{\text{Br}[\Xi_b \rightarrow J/\psi\Lambda\bar{K}]}{\text{Br}[\Lambda_b \rightarrow J/\psi\Lambda]} = (4.19 \pm 0.29 \pm 0.15) \times 10^{-2},\tag{19}$$

where f_{Ξ_b} and f_{Λ_b} refer to the b quark fragmentation fractions into Ξ_b^- and Λ_b^0 , the ratio of which is [76]

$$\frac{f_{\Xi_b}}{f_{\Lambda_b}} = (6.7 \pm 0.5 \pm 0.5 \pm 2.0) \times 10^{-2},\tag{20}$$

while the branching ratio of $\Lambda_b \rightarrow J/\psi\Lambda$ has been measured by the CDF Collaboration [77]

$$\text{Br}[\Lambda_b \rightarrow J/\psi\Lambda] = (3.7 \pm 1.7 \pm 0.7) \times 10^{-4}.\tag{21}$$

With all the ratios given above, one can compute the branching ratio of $\Xi_b \rightarrow J/\psi\Lambda\bar{K}$

$$\text{Br}[\Xi_b \rightarrow J/\psi\Lambda\bar{K}] = (2.31 \pm 1.37) \times 10^{-4}.\tag{22}$$

The large uncertainty can be traced back to the experimental uncertainty in the braching ratio $\text{Br}[\Lambda_b \rightarrow J/\psi\Lambda]$, which accounts for about 50%, and the large uncertainty in the ratio of fragmentation fractions coming from the estimation of SU(3) breaking effects [76].

C. Couplings from the unitary approach

In our convention, the $J/\psi\Lambda$ partial decay widths of the P_{cs} state with $J^P = 1/2^-$ and $3/2^-$ are expressed as

$$\begin{aligned}\Gamma_{P_{cs1} \rightarrow J/\psi\Lambda} &= \frac{1}{2} \frac{g_{P_{cs1}J/\psi\Lambda}^2}{8\pi} \frac{1}{m_{P_{cs}}^2} |q| \sum |\mathcal{A}_{P_{cs1}}|^2, \\ \Gamma_{P_{cs2} \rightarrow J/\psi\Lambda} &= \frac{1}{4} \frac{g_{P_{cs2}J/\psi\Lambda}^2}{8\pi} \frac{1}{m_{P_{cs}}^2} |q| \sum |\mathcal{A}_{P_{cs2}}|^2,\end{aligned}\tag{23}$$

where the modules of amplitude squared are

$$\begin{aligned}\sum |\mathcal{A}_{P_{cs1}}|^2 &= \frac{\left((m_\Lambda + m_{P_{cs}})^2 - m_{J/\psi}^2\right) \left(\left(m_{J/\psi}^2 - m_\Lambda^2\right)^2 + 2m_{P_{cs}}^2 \left(5m_{J/\psi}^2 - m_\Lambda^2\right) + m_{P_{cs}}^4\right)}{2m_{J/\psi}^2 m_{P_{cs}}^2}, \\ \sum |\mathcal{A}_{P_{cs2}}|^2 &= \frac{\left((m_\Lambda + m_{P_{cs}})^2 - m_{J/\psi}^2\right) \left(\left(m_{J/\psi}^2 - m_\Lambda^2\right)^2 + 2m_{P_{cs}}^2 \left(5m_{J/\psi}^2 - m_\Lambda^2\right) + m_{P_{cs}}^4\right)}{3m_{J/\psi}^2 m_{P_{cs}}^2},\end{aligned}\tag{24}$$

in which q denotes the momentum of J/ψ in the rest frame of the P_{cs} state. Using the partial decay widths from Ref. [47], we obtain $g_{P_{cs1}J/\psi\Lambda} = 0.07$ and $g_{P_{cs2}J/\psi\Lambda} = 0.27$. Similarly, we obtain $g_{P_{cs1}\Xi_c\bar{D}^*} = 1.25$ and $g_{P_{cs2}\Xi_c\bar{D}^*} = 2.17$.

-
- [1] R. Aaij et al. (LHCb), Phys. Rev. Lett. **115**, 072001 (2015), arXiv:1507.03414 [hep-ex].
 - [2] J.-J. Wu, R. Molina, E. Oset, and B. S. Zou, Phys. Rev. Lett. **105**, 232001 (2010), arXiv:1007.0573 [nucl-th].
 - [3] J.-J. Wu, R. Molina, E. Oset, and B. S. Zou, Phys. Rev. **C84**, 015202 (2011), arXiv:1011.2399 [nucl-th].
 - [4] W. L. Wang, F. Huang, Z. Y. Zhang, and B. S. Zou, Phys. Rev. **C84**, 015203 (2011), arXiv:1101.0453 [nucl-th].
 - [5] Z.-C. Yang, Z.-F. Sun, J. He, X. Liu, and S.-L. Zhu, Chin. Phys. **C36**, 6 (2012), arXiv:1105.2901 [hep-ph].
 - [6] S. G. Yuan, K. W. Wei, J. He, H. S. Xu, and B. S. Zou, Eur. Phys. J. **A48**, 61 (2012), arXiv:1201.0807 [nucl-th].
 - [7] J.-J. Wu, T. S. H. Lee, and B. S. Zou, Phys. Rev. C **85**, 044002 (2012), arXiv:1202.1036 [nucl-th].

- [8] C. Garcia-Recio, J. Nieves, O. Romanets, L. L. Salcedo, and L. Tolos, *Phys. Rev. D* **87**, 074034 (2013), arXiv:1302.6938 [hep-ph].
- [9] C. W. Xiao, J. Nieves, and E. Oset, *Phys. Rev.* **D88**, 056012 (2013), arXiv:1304.5368 [hep-ph].
- [10] T. Uchino, W.-H. Liang, and E. Oset, *Eur. Phys. J. A* **52**, 43 (2016), arXiv:1504.05726 [hep-ph].
- [11] M. Karliner and J. L. Rosner, *Phys. Rev. Lett.* **115**, 122001 (2015), arXiv:1506.06386 [hep-ph].
- [12] L. Roca, J. Nieves, and E. Oset, *Phys. Rev.* **D92**, 094003 (2015), arXiv:1507.04249 [hep-ph].
- [13] J. He, *Phys. Lett.* **B753**, 547 (2016), arXiv:1507.05200 [hep-ph].
- [14] C. W. Xiao and U. G. Meißner, *Phys. Rev.* **D92**, 114002 (2015), arXiv:1508.00924 [hep-ph].
- [15] R. Chen, X. Liu, X.-Q. Li, and S.-L. Zhu, *Phys. Rev. Lett.* **115**, 132002 (2015), arXiv:1507.03704 [hep-ph].
- [16] H.-X. Chen, W. Chen, X. Liu, T. G. Steele, and S.-L. Zhu, *Phys. Rev. Lett.* **115**, 172001 (2015), arXiv:1507.03717 [hep-ph].
- [17] T. J. Burns, *Eur. Phys. J. A* **51**, 152 (2015), arXiv:1509.02460 [hep-ph].
- [18] C.-W. Shen, F.-K. Guo, J.-J. Xie, and B.-S. Zou, *Nucl. Phys. A* **954**, 393 (2016), arXiv:1603.04672 [hep-ph].
- [19] L. Geng, J. Lu, and M. P. Valderrama, *Phys. Rev.* **D97**, 094036 (2018), arXiv:1704.06123 [hep-ph].
- [20] M.-Z. Liu, F.-Z. Peng, M. Sánchez Sánchez, and M. P. Valderrama, *Phys. Rev. D* **98**, 114030 (2018), arXiv:1811.03992 [hep-ph].
- [21] L. Maiani, A. D. Polosa, and V. Riquer, *Phys. Lett.* **B749**, 289 (2015), arXiv:1507.04980 [hep-ph].
- [22] R. F. Lebed, *Phys. Lett.* **B749**, 454 (2015), arXiv:1507.05867 [hep-ph].
- [23] Z.-G. Wang, *Eur. Phys. J. C* **76**, 70 (2016), arXiv:1508.01468 [hep-ph].
- [24] F.-K. Guo, U.-G. Meißner, W. Wang, and Z. Yang, *Phys. Rev. D* **92**, 071502 (2015), arXiv:1507.04950 [hep-ph].
- [25] X.-H. Liu, Q. Wang, and Q. Zhao, *Phys. Lett. B* **757**, 231 (2016), arXiv:1507.05359 [hep-ph].
- [26] R. Aaij *et al.* (LHCb), *Phys. Rev. Lett.* **122**, 222001 (2019), arXiv:1904.03947 [hep-ex].
- [27] H.-X. Chen, W. Chen, and S.-L. Zhu, *Phys. Rev.* **D100**, 051501 (2019), arXiv:1903.11001 [hep-ph].
- [28] R. Chen, Z.-F. Sun, X. Liu, and S.-L. Zhu, (2019), arXiv:1903.11013 [hep-ph].
- [29] J. He, *Eur. Phys. J. C* **79**, 393 (2019), arXiv:1903.11872 [hep-ph].
- [30] M.-Z. Liu, Y.-W. Pan, F.-Z. Peng, M. Sánchez Sánchez, L.-S. Geng, A. Hosaka, and M. Pavon Valderrama, *Phys. Rev. Lett.* **122**, 242001 (2019), arXiv:1903.11560 [hep-ph].
- [31] Y. Shimizu, Y. Yamaguchi, and M. Harada, (2019), arXiv:1904.00587 [hep-ph].

- [32] C.-J. Xiao, Y. Huang, Y.-B. Dong, L.-S. Geng, and D.-Y. Chen, Phys. Rev. D **100**, 014022 (2019), arXiv:1904.00872 [hep-ph].
- [33] Z.-H. Guo and J. A. Oller, Phys. Lett. **B793**, 144 (2019), arXiv:1904.00851 [hep-ph].
- [34] C. Fernández-Ramírez, A. Pilloni, M. Albaladejo, A. Jackura, V. Mathieu, M. Mikhasenko, J. A. Silva-Castro, and A. P. Szczepaniak (JPAC), Phys. Rev. Lett. **123**, 092001 (2019), arXiv:1904.10021 [hep-ph].
- [35] C. W. Xiao, J. Nieves, and E. Oset, Phys. Rev. **D100**, 014021 (2019), arXiv:1904.01296 [hep-ph].
- [36] Y. Yamaguchi, H. Garcia-Tecocoatz, A. Giachino, A. Hosaka, E. Santopinto, S. Takeuchi, and M. Takizawa, (2019), arXiv:1907.04684 [hep-ph].
- [37] Q. Wu and D.-Y. Chen, Phys. Rev. D **100**, 114002 (2019), arXiv:1906.02480 [hep-ph].
- [38] M. Pavon Valderrama, (2019), arXiv:1907.05294 [hep-ph].
- [39] M.-Z. Liu, T.-W. Wu, M. Sánchez Sánchez, M. P. Valderrama, L.-S. Geng, and J.-J. Xie, (2019), arXiv:1907.06093 [hep-ph].
- [40] Y.-H. Lin and B.-S. Zou, Phys. Rev. D **100**, 056005 (2019), arXiv:1908.05309 [hep-ph].
- [41] A. Ali, I. Ahmed, M. J. Aslam, A. Y. Parkhomenko, and A. Rehman, JHEP **10**, 256 (2019), arXiv:1907.06507 [hep-ph].
- [42] Z.-G. Wang, Int. J. Mod. Phys. A **35**, 2050003 (2020), arXiv:1905.02892 [hep-ph].
- [43] A. Ali and A. Y. Parkhomenko, Phys. Lett. B **793**, 365 (2019), arXiv:1904.00446 [hep-ph].
- [44] S. X. Nakamura, (2021), arXiv:2103.06817 [hep-ph].
- [45] R. Aaij et al. (LHCb), (2020), arXiv:2012.10380 [hep-ex].
- [46] F.-Z. Peng, M.-J. Yan, M. Sánchez Sánchez, and M. P. Valderrama, (2020), arXiv:2011.01915 [hep-ph].
- [47] C. W. Xiao, J. J. Wu, and B. S. Zou, Phys. Rev. D **103**, 054016 (2021), arXiv:2102.02607 [hep-ph].
- [48] J.-T. Zhu, L.-Q. Song, and J. He, (2021), arXiv:2101.12441 [hep-ph].
- [49] X.-K. Dong, F.-K. Guo, and B.-S. Zou, Progr. Phys. **41**, 65 (2021), arXiv:2101.01021 [hep-ph].
- [50] M.-Z. Liu, Y.-W. Pan, and L.-S. Geng, Phys. Rev. D **103**, 034003 (2021), arXiv:2011.07935 [hep-ph].
- [51] R. Chen, Eur. Phys. J. C **81**, 122 (2021), arXiv:2101.10614 [hep-ph].
- [52] C. W. Xiao, J. Nieves, and E. Oset, Phys. Lett. B **799**, 135051 (2019), arXiv:1906.09010 [hep-ph].
- [53] B. Wang, L. Meng, and S.-L. Zhu, Phys. Rev. D **101**, 034018 (2020), arXiv:1912.12592 [hep-ph].
- [54] T. J. Burns and E. S. Swanson, Phys. Rev. D **100**, 114033 (2019), arXiv:1908.03528 [hep-ph].

- [55] F.-Z. Peng, J.-X. Lu, M. Sánchez Sánchez, M.-J. Yan, and M. Pavon Valderrama, Phys. Rev. D **103**, 014023 (2021), arXiv:2007.01198 [hep-ph].
- [56] H. Xu, Q. Li, C.-H. Chang, and G.-L. Wang, Phys. Rev. D **101**, 054037 (2020), arXiv:2001.02980 [hep-ph].
- [57] X.-Z. Ling, M.-Z. Liu, J.-X. Lu, L.-S. Geng, and J.-J. Xie, (2021), arXiv:2102.05349 [hep-ph].
- [58] H.-X. Chen, L.-S. Geng, W.-H. Liang, E. Oset, E. Wang, and J.-J. Xie, Phys. Rev. C **93**, 065203 (2016), arXiv:1510.01803 [hep-ph].
- [59] Q. Wu, D.-Y. Chen, and R. Ji, (2021), arXiv:2103.05257 [hep-ph].
- [60] J.-X. Lu, E. Wang, J.-J. Xie, L.-S. Geng, and E. Oset, Phys. Rev. D **93**, 094009 (2016), arXiv:1601.00075 [hep-ph].
- [61] P. A. Zyla et al. (Particle Data Group), PTEP **2020**, 083C01 (2020).
- [62] H.-Y. Cheng, Phys. Rev. D **56**, 2799 (1997), [Erratum: Phys.Rev.D **99**, 079901 (2019)], arXiv:hep-ph/9612223.
- [63] R. N. Faustov and V. O. Galkin, Phys. Rev. D **98**, 093006 (2018), arXiv:1810.03388 [hep-ph].
- [64] H.-n. Li, C.-D. Lu, and F.-S. Yu, Phys. Rev. D **86**, 036012 (2012), arXiv:1203.3120 [hep-ph].
- [65] T. Gutsche, M. A. Ivanov, J. G. Körner, V. E. Lyubovitskij, P. Santorelli, and N. Habył, Phys. Rev. D **91**, 074001 (2015), [Erratum: Phys.Rev.D **91**, 119907 (2015)], arXiv:1502.04864 [hep-ph].
- [66] R. S. Azevedo and M. Nielsen, Phys. Rev. C **69**, 035201 (2004), arXiv:nucl-th/0310061.
- [67] S. Weinberg, Phys. Rev. **130**, 776 (1963).
- [68] A. Salam, Nuovo Cim. **25**, 224 (1962).
- [69] K. Hayashi, M. Hirayama, T. Muta, N. Seto, and T. Shirafuji, Fortsch. Phys. **15**, 625 (1967).
- [70] N. A. Tornqvist, Nuovo Cim. A **107**, 2471 (1994), arXiv:hep-ph/9310225.
- [71] N. A. Tornqvist, Z. Phys. **C61**, 525 (1994), arXiv:hep-ph/9310247 [hep-ph].
- [72] M. P. Locher, Y. Lu, and B. S. Zou, Z. Phys. A **347**, 281 (1994), arXiv:nucl-th/9311021.
- [73] X.-Q. Li, D. V. Bugg, and B.-S. Zou, Phys. Rev. D **55**, 1421 (1997).
- [74] c.-J. Xiao, D.-Y. Chen, and Y.-L. Ma, Phys. Rev. D **93**, 094011 (2016), arXiv:1601.06399 [hep-ph].
- [75] R. Aaij et al. (LHCb), Phys. Lett. B **772**, 265 (2017), arXiv:1701.05274 [hep-ex].
- [76] R. Aaij et al. (LHCb), Phys. Rev. D **99**, 052006 (2019), arXiv:1901.07075 [hep-ex].
- [77] F. Abe et al. (CDF), Phys. Rev. D **55**, 1142 (1997).

