

REPRESENTATIONS AND TENSOR PRODUCT GROWTH

MICHAEL LARSEN, ANER SHALEV, AND PHAM HUU TIEP

ABSTRACT. The deep theory of approximate subgroups establishes 3-step product growth for subsets of finite simple groups G of Lie type of bounded rank. In this paper we obtain 2-step growth results for representations of such groups G (including those of unbounded rank), where products of subsets are replaced by tensor products of representations.

Let G be a finite simple group of Lie type and χ a character of G . Let $|\chi|$ denote the sum of the squares of the degrees of all (distinct) irreducible characters of G which are constituents of χ . We show that for all $\delta > 0$ there exists $\epsilon > 0$, independent of G , such that if χ is an irreducible character of G satisfying $|\chi| \leq |G|^{1-\delta}$, then $|\chi^2| \geq |\chi|^{1+\epsilon}$. We also obtain results for reducible characters, and establish faster growth in the case where $|\chi| \leq |G|^\delta$.

In another direction, we explore covering phenomena, namely situations where every irreducible character of G occurs as a constituent of certain products of characters. For example, we prove that if $|\chi_1| \cdots |\chi_m|$ is a high enough power of $|G|$, then every irreducible character of G appears in $\chi_1 \cdots \chi_m$. Finally, we obtain growth results for compact semisimple Lie groups.

1. INTRODUCTION

In the past two decades there has been intense interest in growth phenomena in groups and in finite simple groups in particular; see [He1, Hr, BGT, He2, PS, Br]. The celebrated Product Theorem, proved independently in [BGT] and in [PS], shows that, if G is a finite simple group of Lie type of rank r , and $A \subset G$ is a generating set, then either $A^3 = G$ or $|A^3| \geq |A|^{1+\epsilon}$ where $\epsilon > 0$ depends only on r . Here A^k denotes the set of all products $a_1 a_2 \cdots a_k$ where $a_1, \dots, a_k \in A$.

The main goal of this paper is to study analogous power growth phenomena in representation theory, with emphasis on (complex) representations of finite simple groups G of Lie type. Here products of subsets of G are replaced by tensor products of representations. Our results on tensor product growth are somewhat stronger than the Product Theorem in two senses: we establish 2-step (instead of 3-step) growth, as well as uniform growth when the rank of G tends to infinity.

Let G be any finite group. If $X = \{\chi_1, \dots, \chi_k\}$ is a set of (pairwise distinct) irreducible characters of G , we define

$$|X| = \sum_{i=1}^k \chi_i(1)^2.$$

This is the Plancherel measure, normalized so that $|\text{Irr}(G)| = |G|$. If χ is any character of G , we define $|\chi| = |X_\chi|$, where X_χ denotes the set of distinct irreducible constituents of χ . Recall that G is *quasisimple* if $G = [G, G]$ and $G/\mathbf{Z}(G)$ is simple.

Our first growth results are as follows:

ML was partially supported by the NSF grant DMS-2001349. AS was partially supported by ISF grant 686/17 and the Vinik Chair of mathematics. PT was partially supported by the NSF grant DMS-1840702, the Joshua Barlaz Chair in Mathematics, and the Charles Simonyi Endowment at the Institute for Advanced Study (Princeton). All three authors were partially supported by BSF grant 2016072.

Theorem 1. *For all $\delta > 0$, there exists $\epsilon > 0$ such that if G is a finite quasisimple group of Lie type and χ is an irreducible character of G with $|\chi| \leq |G|^{1-\delta}$, then $|\chi^2| \geq |\chi|^{1+\epsilon}$ and $|\chi\bar{\chi}| \geq |\chi|^{1+\epsilon}$.*

We also have a version of this result for general characters in groups of high rank:

Theorem 2. *For all $\delta > 0$, there exist $\epsilon > 0$ and $R > 0$ such that if G is a finite quasisimple group of Lie type and rank $\geq R$, and χ is any (not necessarily irreducible) character of G with $|\chi| \leq |G|^{1-\delta}$, then $|\chi^2| \geq |\chi|^{1+\epsilon}$ and $|\chi\bar{\chi}| \geq |\chi|^{1+\epsilon}$.*

The proofs of these results present ϵ as an explicit function of δ , e.g. $\epsilon = \frac{c\delta}{4+2c(1-\delta)}$ in Theorem 1, where $c > 0$ is the absolute constant in [LaT, Theorem A]. Moreover, if G is sufficiently large but of bounded rank r , and χ is irreducible, then $\epsilon = \frac{\delta}{2-2\delta}$ will do; for example, any irreducible character χ of G with $|\chi| \leq |G|^{1/2}$ satisfies $|\chi^2| \geq |\chi|^{3/2}$.

Power growth of conjugacy classes A of arbitrary finite simple groups G was studied before the Product Theorem was proved. It is shown in [Sh, 2.7] there that for any $\delta > 0$ there exists $\epsilon > 0$, depending only on δ , such that $|A| \leq |G|^{1-\delta}$ implies $|A^3| \geq |A|^{1+\epsilon}$. Furthermore, if G is of Lie type then $|A^2| \geq |A|^{1+\epsilon}$ where $\epsilon > 0$ depends only on the rank of G [Sh, 10.4].

Subsequently, growth of general normal subsets (namely, union of conjugacy classes) was also studied. It is shown in [GPSS, 1.5] that there are absolute constants $N \in \mathbb{N}$ and $\epsilon > 0$ such that for any normal subset A of a finite simple group G , either $A^N = G$ or $|A^2| \geq |A|^{1+\epsilon}$.

In [LSS] faster growth of the form $|A^2| \geq |A|^{2-\epsilon}$ for small normal subsets A of arbitrary finite simple groups is established. Our next result gives a character-theoretic analogue of [LSS, Theorem 1.3]:

Theorem 3. *For any $\epsilon > 0$, there exists an explicit $\delta > 0$ such that the following statement holds. If G is a finite quasisimple group of Lie type and χ_1, χ_2 are any (not necessarily irreducible) characters of G with $|\chi_1|, |\chi_2| \leq |G|^\delta$, then*

$$|\chi_1\chi_2| \geq (|\chi_1| \cdot |\chi_2|)^{1-\epsilon}.$$

In particular, if χ is a character of G satisfying $|\chi| \leq |G|^\delta$ then $|\chi^2| \geq |\chi|^{2-2\epsilon}$.

We note that $|\chi_1\chi_2| \leq |\chi_1| \cdot |\chi_2|$ (see Lemma 2.1 below), hence the growth established in Theorem 3 is almost best possible. As a consequence of Theorem 3, we obtain:

Corollary 4. *For any $\epsilon > 0$ and any integer $k \geq 2$, there exists an explicit $\gamma = \gamma(\epsilon, k) > 0$ such that the following statement holds. If G is a finite quasisimple group of Lie type and $\chi_1, \chi_2, \dots, \chi_k$ are any (not necessarily irreducible) characters of G with $|\chi_1|, |\chi_2|, \dots, |\chi_k| \leq |G|^\gamma$, then*

$$|\chi_1\chi_2 \cdots \chi_k| \geq (|\chi_1| \cdot |\chi_2| \cdots |\chi_k|)^{1-\epsilon}.$$

In particular, if χ is a character of G satisfying $|\chi| \leq |G|^\gamma$ then $|\chi^k| \geq |\chi|^{k-k\epsilon}$.

The above result shows that, for any $\epsilon > 0$ and any integer $k \geq 2$ there exists an explicit $\delta = \delta(\epsilon, k) > 0$ such that, for G as above and any (not necessarily irreducible) character χ of G satisfying $|\chi| \leq |G|^\delta$ we have $|\chi^k| \geq |\chi|^{k-k\epsilon}$; indeed, define $\delta(\epsilon, k) = \gamma(\epsilon/k, k)$.

Applying Theorem 3 we deduce the following result, which is a character-theoretic analogue of [LSS, Theorem 1.1]:

Theorem 5. *For all $\epsilon > 0$, there exists an explicit positive integer N such that if G is a finite simple group of Lie type and χ is any (not necessarily irreducible) character of G , then either χ^N contains every irreducible character of G , or $|\chi^2| \geq |\chi|^{2-\epsilon}$.*

The analogy with Gowers' theorem raises the question of whether $N = 3$ suffices in Theorem 5 when $|\chi|$ is sufficiently large. A recent theorem of Sellke [Se, Theorem 1.2] shows that the answer to this question is affirmative for large G if χ is so large that $\frac{|\chi|}{|G|}$ is bounded away from 0. We therefore ask the following:

Question. If G is a finite simple group of Lie type and χ is an arbitrary character of G such that $\frac{\log |\chi|}{\log |G|}$ is sufficiently close to 1, is it true that $|\chi^3| = |G|$?

We remark that the example of $\text{PSU}_{2n+1}(q)$ [HSTZ, Theorem 1.2] shows, in general, it would be too much to ask for $|\chi^2| = |G|$. On the other hand, for certain simple groups of Lie type, we can bring N down to 6 or 7.

Theorem 6. *If $G = \text{PSL}_n(q)$ and q is sufficiently large in terms of n , then $|\chi| \geq |G|^{11/12}$ implies $|\chi^6| = |G|$. If $G = \text{PSU}_n(q)$ and q is sufficiently large in terms of n , then $|\chi| \geq |G|^{11/12}$ implies $|\chi^7| = |G|$.*

We also offer a character-theoretic analogue of the Rodgers-Saxl theorem on products of conjugacy classes in [RS]. In the case that the characters are irreducible, this analogue was conjectured by Gill in [Gi] and proved in [LaT, Theorem 8.5].

Theorem 7. *There exists an explicit constant $c > 0$ such that the following statement holds. If G is a finite simple group of Lie type, $m \geq 1$ any integer, and $\chi_1, \chi_2, \dots, \chi_m$ are any (not necessarily irreducible) characters of G with $\prod_{i=1}^m |\chi_i| \geq |G|^c$, then $|\chi_1 \chi_2 \dots \chi_m| = |G|$ and thus $\chi_1 \chi_2 \dots \chi_m$ contains every irreducible character of G .*

Finally, we prove an analogue of Theorem 1 for compact semisimple Lie groups.

Theorem 8. *Let G be a compact semisimple Lie group. Then there exists $\epsilon > 0$ such that for each irreducible character χ of G , we have $|\chi^2| \geq |\chi|^{1+\epsilon}$.*

It is possible that some of the presented results also hold for alternating groups (and likewise, Theorem 2 may hold for Lie-type groups of bounded rank as well). However, the techniques developed in the paper do not seem to apply to these open cases.

Some words on the structure of this paper. In Section 2 we prove some preliminary results for finite groups. Section 3 is devoted to the proofs of Theorem 1 and Theorem 2. In Section 4 we prove Theorems 3, 5, 7, and Corollary 4, while the proof of Theorem 6 is carried out in Section 5. In Section 6 we study tensor product growth of representations of semisimple compact Lie groups and prove Theorem 8.

2. PRELIMINARIES

We begin with general inequalities for finite groups G .

Lemma 2.1. *Let α, β be any (not necessarily irreducible) characters of G . Then*

- (a) $|\alpha + \beta| \leq |\alpha| + |\beta|$.
- (b) $|\alpha\beta| \leq |\alpha| \cdot |\beta|$.

Proof. Without any loss of generality we may assume that $\alpha = \sum_i \alpha_i$ and $\beta = \sum_j \beta_j$ are multiplicity-free, with $\alpha_i, \beta_j \in \text{Irr}(G)$. Then

$$|\alpha + \beta| = \left| \sum_i \alpha_i + \sum_j \beta_j \right| \leq \sum_i \alpha_i(1)^2 + \sum_j \beta_j(1)^2 = |\alpha| + |\beta|,$$

proving (a). Next, $|\alpha_i \beta_j| \leq (\alpha_i \beta_j(1))^2 = \alpha_i(1)^2 \beta_j(1)^2$. It follows from (a) that

$$|\alpha \beta| = \left| \sum_{i,j} \alpha_i \beta_j \right| \leq \sum_{i,j} |\alpha_i \beta_j| \leq \sum_{i,j} \alpha_i(1)^2 \beta_j(1)^2 = \left(\sum_i \alpha_i(1)^2 \right) \cdot \left(\sum_j \beta_j(1)^2 \right) = |\alpha| \cdot |\beta|,$$

proving (b). □

Proposition 2.2. *Let $\chi_1, \dots, \chi_n$ be irreducible characters of G , $n \geq 2$.*

(i) *In general, we have*

$$|\chi_1 \dots \chi_n| \geq \max_{1 \leq i \leq n} |\chi_i|.$$

(ii) *If G is quasisimple and $\chi_i \neq 1_G$ for $1 \leq i \leq n$, then*

$$|\chi_1 \dots \chi_n| > \max_{1 \leq i \leq n} |\chi_i|.$$

(iii) *If G is perfect, $\chi_1 \neq 1_G$, and $\chi_i \in \{\chi_1, \overline{\chi}_1\}$ for all i , then $|\chi_1 \dots \chi_n| > |\chi_1| = \dots = |\chi_n|$.*

Proof. (i) Without any loss of generality, assume $\chi_1(1) = \max_i \chi_i(1)$. Note that

$$\begin{aligned} \langle \chi_1 \dots \chi_n, \chi_1 \dots \chi_n \rangle &= \frac{1}{|G|} \sum_{g \in G} |\chi_1(g) \dots \chi_n(g)|^2 \\ &\leq \frac{1}{|G|} \max_{h \in G} |\chi_2(h) \dots \chi_n(h)|^2 \sum_{g \in G} |\chi_1(g)|^2 \\ &= \max_{h \in G} |\chi_2(h) \dots \chi_n(h)|^2 \langle \chi_1, \chi_1 \rangle \\ &= \chi_2(1)^2 \dots \chi_n(1)^2, \end{aligned}$$

with equality only if, for every $g \in G$,

$$(2.1) \quad \text{either } \chi_1(g) = 0 \text{ or } |\chi_2(g) \dots \chi_n(g)| = \chi_2(1) \dots \chi_n(1).$$

If $d_1, d_2, \dots, d_k$ are the degrees of the distinct irreducible constituents of $\chi_1 \dots \chi_n$ and $m_1, \dots, m_k$ their multiplicities in $\chi_1 \dots \chi_n$, then

$$\sum_i m_i^2 = \langle \chi_1 \dots \chi_n, \chi_1 \dots \chi_n \rangle \leq \chi_2(1)^2 \dots \chi_n(1)^2$$

and

$$\sum_i d_i m_i = \chi_1(1) \dots \chi_n(1).$$

So by the Cauchy-Schwarz inequality,

$$|\chi_1 \dots \chi_n| = \sum_{i=1}^k d_i^2 \geq \frac{\chi_1(1)^2 \dots \chi_n(1)^2}{\sum_{i=1}^k m_i^2} \geq \chi_1(1)^2 = |\chi_1|,$$

as stated.

(ii) Suppose G is quasisimple, $\chi_i \neq 1_G$ for all i , but $|\chi_1 \dots \chi_n| = |\chi_1| = \max_i |\chi_i|$. Note that $K_i := \{g \in G \mid |\chi_i(g)| = \chi_i(1)\}$ is a normal subgroup of G , which is contained in $\mathbf{Z}(G)$. Setting $K := \bigcap_{i=2}^n K_i$ and using (2.1), we see that $\chi_1(g) = 0$ for all $g \notin K$. As $K \leq \mathbf{Z}(G)$, we have $|\chi_1(g)| = \chi_1(1)$ for $g \in K$. It follows that

$$|G| = \sum_{g \in G} |\chi_1(g)|^2 = |K| \chi_1(1)^2,$$

and thus $\chi_1(1) = |G/K|^{1/2}$, i.e. χ_1 is of central type character for $G/\text{Ker}(\chi)$. By the Howlett-Isaacs theorem [HI], $G/\text{Ker}(\chi_1)$ is solvable, which is impossible, since it maps onto the simple group $G/\mathbf{Z}(G)$.

(iii) Suppose G is perfect and $\chi_1 \neq 1$, but $|\chi_1^{n-j}\overline{\chi_1}^j| = |\chi_1|$ for some $0 \leq j \leq n$. We again have that $K := \{g \in G \mid |\chi_1(g)| = \chi_1(1)\}$ is a normal subgroup of G , and $\chi_1(g) = 0$ for all $g \notin K$ by (2.1). It follows that

$$|G| = \sum_{g \in G} |\chi_1(g)|^2 = |K| \chi_1(1)^2,$$

and thus $\chi_1(1) = |G/K|^{1/2}$, i.e. χ_1 is of central type character for $G/\text{Ker}(\chi_1)$. By the Howlett-Isaacs theorem, $G/\text{Ker}(\chi_1)$ is solvable, so $\text{Ker}(\chi_1) \geq [G, G] = G$ and thus $\chi_1 = 1_G$, a contradiction. $\square$

The following example shows the absence of growth in G in a class of groups G which are far from being simple.

Example 2.3. Let p be any prime, $n \geq 1$ any integer, and G be an extraspecial p -group of order p^{1+2n} . Then G has p^{2n} linear characters and $p-1$ irreducible characters of degree p^n ; let χ be one of the latter. Now, if $k \geq 1$ be any integer, then χ^k is the sum of p^{2n} linear characters if $p|k$, and a multiple of a single irreducible character of degree p^n if $p \nmid k$. Thus we always have $|\chi^k| = p^{2n} = |\chi| = |G|^{1-1/(2n+1)}$.

Question. When χ_i are not assumed irreducible, can it ever happen that

$$|\chi_1 \cdots \chi_n| < \max(|\chi_1|, \dots, |\chi_n|)?$$

3. PROOFS OF THEOREMS 1 AND 2

We begin this section by proving Theorem 2 in the case that χ is irreducible, which is also the high rank case of Theorem 1.

Proposition 3.1. *For all $\delta > 0$, there exist $\epsilon > 0$ and $R > 0$ such that if G is a finite quasisimple group of Lie type and rank $\geq R$, and $\chi \in \text{Irr}(G)$ satisfies $|\chi| \leq |G|^{1-\delta}$, then $|\chi^2| \geq |\chi|^{1+\epsilon}$ and $|\chi\overline{\chi}| \geq |\chi|^{1+\epsilon}$.*

Proof. Let $r \geq R$ be the rank of G and q the cardinality of the field of definition (for Suzuki and Ree groups, q is defined, as usual, so that q^2 is an odd power of 2 or 3.) By the main theorem of [LaT], there exists $c > 0$ such that

$$|\chi(g)| \leq \chi(1)^{1-c \frac{\log |g^G|}{\log |G|}}.$$

Given $\alpha > 0$, let $S = S_\alpha \subset G$ denote the set of elements such that $|g^G| \leq |G|^{\alpha/2}$. Then, by a well known result of Fulman and Guralnick [FG], we have

$$|S| \leq k(G) |G|^{\alpha/2} \leq 27.2 q^r |G|^{\alpha/2} \leq |G|^\alpha,$$

if R is taken sufficiently large in terms of α . Thus,

$$\begin{aligned}
\langle \chi^2, \chi^2 \rangle &= \langle \chi \bar{\chi}, \chi \bar{\chi} \rangle = \frac{1}{|G|} \sum_{g \in G} |\chi(g)|^4 \\
&= \frac{1}{|G|} \sum_{g \in S} |\chi(g)|^4 + \frac{1}{|G|} \sum_{g \in G \setminus S} |\chi(g)|^4 \\
&\leq \frac{|S|}{|G|} \chi(1)^4 + \frac{1}{|G|} \max_{g \in G \setminus S} |\chi(g)|^2 \sum_{g \in G \setminus S} |\chi(g)|^2 \\
&\leq |G|^{\alpha-1} \chi(1)^4 + \frac{1}{|G|} \chi(1)^{2(1-c\alpha/2)} \sum_{g \in G \setminus S} |\chi(g)|^2 \\
&\leq |G|^{\alpha-1} \chi(1)^4 + \chi(1)^{2-c\alpha}.
\end{aligned}$$

By hypothesis, $|\chi| \leq |G|^{1-\delta}$, so $\chi(1) \leq |G|^{1/2-\delta/2}$, which implies

$$\langle \chi^2, \chi^2 \rangle \leq \chi(1)^{4-\frac{2-2\alpha}{1-\delta}} + \chi(1)^{2-c\alpha} = (\chi(1)^{-\frac{2\delta-2\alpha}{1-\delta}} + \chi(1)^{-c\alpha}) \chi(1)^2.$$

Choosing

$$\alpha := \frac{2\delta}{2+c(1-\delta)}, \quad \epsilon := \frac{c\alpha}{4},$$

we get

$$\langle \chi^2, \chi^2 \rangle \leq 2\chi(1)^{-c\alpha} \chi(1)^2 \leq \chi(1)^{2-2\epsilon}$$

if R is taken sufficiently large. Applying the Cauchy-Schwarz inequality, we get

$$|\chi^2| \geq \frac{\chi(1)^4}{\langle \chi^2, \chi^2 \rangle} \geq \chi(1)^{2+2\epsilon} = |\chi|^{1+\epsilon}.$$

Likewise,

$$|\chi \bar{\chi}| \geq \frac{\chi(1)^4}{\langle \chi \bar{\chi}, \chi \bar{\chi} \rangle} \geq \chi(1)^{2+2\epsilon} = |\chi|^{1+\epsilon}.$$

□

To complete the proof of Theorem 1, we need only treat the bounded rank case. Using Proposition 2.2(iii), we may take $|G|$ to be as large as we wish.

Proposition 3.2. *Let R be fixed. For all $\delta > 0$, there exists $\epsilon > 0$ such that if G is a quasisimple group of Lie type of rank $r < R$ and χ is an irreducible character of G with $|\chi| \leq |G|^{1-\delta}$, then $|\chi^2|$ and $|\chi \bar{\chi}|$ are both at least $|\chi|^{1+\epsilon}$.*

Proof. We have already remarked that G can be assumed arbitrarily large. Since the rank r is bounded, this means we may take q arbitrarily large. By an estimate of Gluck [Gl], for $g \neq 1$ we have $|\chi(g)| \leq C/\sqrt{q}$, where C is an absolute constant. On the other hand, our upper bound on the rank gives an upper bound on $|G|$ of the form q^N for some N . Taking q sufficiently large, we therefore have

$$\frac{|\chi(g)|}{\chi(1)} \leq Cq^{-1/2} \leq q^{-1/3} \leq |G|^{-\frac{1}{3N}} \leq \chi(1)^{-\frac{1}{3N}}$$

for all $g \neq 1$.

Setting $S = \{1\}$, we proceed as in the proof of Proposition 3.1, obtaining

$$\begin{aligned} \langle \chi^2, \chi^2 \rangle &= \langle \chi \bar{\chi}, \chi \bar{\chi} \rangle = \frac{1}{|G|} \sum_{g \in G} |\chi(g)|^4 \\ &= \frac{\chi(1)^4}{|G|} + \frac{1}{|G|} \sum_{g \neq 1} |\chi(g)|^4 \\ &\leq \frac{\chi(1)^4}{|G|} + \frac{1}{|G|} \max_{g \neq 1} |\chi(g)|^2 \sum_{g \neq 1} |\chi(g)|^2 \\ &\leq \chi(1)^{4 - \frac{2}{1-\delta}} + \chi(1)^{2(1 - \frac{1}{3N})}. \end{aligned}$$

Assuming $\delta \leq \frac{1}{3N+1}$, we have

$$\langle \chi^2, \chi^2 \rangle \leq 2\chi(1)^{2 - \frac{2\delta}{1-\delta}} \leq \chi(1)^{2 - \frac{\delta}{1-\delta}}$$

if $|G|$, and therefore $\chi(1)$, is sufficiently large. Taking $\epsilon := \frac{\delta}{2-2\delta}$ and using Cauchy-Schwarz as before, the proposition follows. $\square$

Now we prove Theorem 2.

Proof. By [LiSh2, Theorem 1.2], for all N , there exists R such that if G has rank $r \geq R$, then

$$\sum_{\chi \in \text{Irr}(G)} \chi(1)^{-1/N} \leq \frac{3}{2}.$$

This implies that the number of irreducible characters of G of degree $\leq D$ is at most $D^{1/N}$. Therefore, if χ is a character of G with distinct irreducible factors $\chi_1, \dots, \chi_k$ of degrees $d_1 \leq d_2 \leq \dots \leq d_k$, then

$$|\chi| = \sum_{i=1}^k d_i^2 \leq d_k^{1/N} d_k^2,$$

so

$$|\chi_k| = d_k^2 \geq |\chi|^{\frac{N}{2N+1}}.$$

On the other hand, $|\chi^2| \geq |\chi_k^2|$ and $|\chi \bar{\chi}| \geq |\chi_k \bar{\chi}_k|$. Theorem 1 now follows by applying Proposition 3.1 to χ_k . $\square$

4. PROOF OF THEOREMS 3, 5 AND 7

The notion of the *level* $\mathfrak{l}(\chi)$ of an irreducible character $\chi \in \text{Irr}(G)$ of a finite classical group G was introduced in [GLT1] for groups of type A , and [GLT2, Definition 3.2] for other classical types. For brevity, we use SL^ϵ to denote SL when $\epsilon = +$ and SU when $\epsilon = -$. Using the results of [GLT1, GLT2], we can prove the following bound (which is useful only when $L < \sqrt{n}$).

Proposition 4.1. *Let $G = \text{SL}_n^\epsilon(q)$ with $n \geq 7$, or $\text{Sp}_{2n}(q)$, $\Omega_{2n+1}(q)$, $\Omega_{2n}^\pm(q)$ with $n \geq 6$, and let $L \geq 1$ be any real number.*

- (a) *Suppose $G = \text{SL}_n^\epsilon(q)$ and $L \leq n/6$. Then the number $N(L)$ of $\chi \in \text{Irr}(G)$ with $\chi(1) \leq q^{nL}$ is at most q^{12L^2} .*
- (b) *Suppose $G = \text{Sp}_{2n}(q)$ with $2 \nmid q$. Then the number $N(L)$ of $\chi \in \text{Irr}(G)$ with $\chi(1) \leq q^{nL}$ is at most q^{18L^2} .*
- (c) *Suppose $G = \text{Sp}_{2n}(q)$ with $2|q$, or $\Omega_{2n+1}(q)$ with $2 \nmid q$, or $\Omega_{2n}^+(q)$ for any q . Then the number $N(L)$ of $\chi \in \text{Irr}(G)$ with $\chi(1) \leq q^{nL}$ is at most q^{50L^2} .*

(d) Suppose $G = \Omega_{2n}^-(q)$. Then the number $N(L)$ of $\chi \in \text{Irr}(G)$ with $\chi(1) \leq q^{(n-1)L}$ is at most q^{41L^2} .

Proof. (a) First we bound $j := \ell(\chi)$ in terms of L . If $j > n/2$, then by [GLT1, Theorem 1.3(ii)] we have

$$\chi(1) \geq (2/3)q^{n^2/4-3} > q^{n^2/6}$$

since $n \geq 7$, contradicting the condition $L \leq n/6$. Hence $j \leq n/2$. Now using [GLT1, Theorem 1.3(i)] we get

$$q^{nL} \geq \chi(1) \geq \frac{1}{2(q+1)}q^{j(n-j)} > q^{j(n-j)-2.6} \geq q^{jn/2-2.6},$$

which implies

$$(4.1) \quad j \leq \frac{nL + 2.6}{n/2} = 2L + \frac{5.2}{n} < \frac{11}{4}L.$$

Let $\delta = +$ and consider the reducible *Weil character* $\tau = \tau_{n,q}$ of $\text{SL}_n(q)$, see [GLT1, (1.1)]. Note that τ is just the permutation character of $G = \text{SL}_n(q)$ acting on the point set of $V := \mathbb{F}_q^n$; in particular τ contains 1_G , and so τ^j contains all irreducible constituents of $\sum_{i=0}^j \tau^i$. By definition, $j = \ell(\chi)$ is the smallest non-negative integer such that χ is a constituent of τ^j . Hence, (4.1) implies that $N(L) \leq N^*(l)$ with $l := \lfloor 11L/4 \rfloor$, where $N^*(l)$ is the number of distinct irreducible constituents of τ^l . Next,

$$N^*(l) \leq \langle \tau^l, \tau^l \rangle = \langle \tau^{2l}, 1_G \rangle,$$

the number of G -orbits on V^{2l} . By [GLT1, Lemma 2.4], the number of $\text{GL}_n(q)$ -orbits on V^{2l} is at most $8q^{l^2}$. Since G has index $q-1$ in $\text{GL}_n(q)$, we have

$$N(L) \leq 8(q-1)q^{l^2} \leq q^{l^2+3} \leq q^{(11L/4)^2+3} < q^{11L^2}.$$

Now suppose $\delta = -$ and consider the reducible *Weil character* $\zeta = \zeta_{n,q}$ of $\text{SU}_n(q)$, see [GLT1, (1.1)]. Note that $\zeta^2 = (\tau_{n,q^2})|_G$ is just the permutation character of $G = \text{SU}_n(q)$ acting on the point set of $U := \mathbb{F}_{q^2}^n$; in particular ζ^2 contains 1_G , and so $\zeta^j + \zeta^{j-1}$ contains all irreducible constituents of $\sum_{i=0}^j \zeta^i$. Hence, (4.1) implies that $N(L) \leq N^*(l) + N^*(l-1)$ with $l := \lfloor 11L/4 \rfloor$, where $N^*(l)$ is the number of distinct irreducible constituents of ζ^l . Next,

$$N^*(l) \leq \langle \zeta^l, \zeta^l \rangle = \langle \zeta^{2l}, 1_G \rangle = \langle (\tau_{n,q^2})|_G, 1_G \rangle,$$

the number of G -orbits on U^{2l} . By [GLT1, Lemma 2.4], the number of $\text{GU}_n(q)$ -orbits on U^{2l} is at most $2q^{l^2}$. Since G has index $q+1$ in $\text{GU}_n(q)$, we have

$$N(L) \leq 4(q+1)q^{l^2} < q^{l^2+4} \leq q^{(11L/4)^2+4} < q^{12L^2}.$$

(b) In the remaining cases, we set $k := \lfloor (j+2)/3 \rfloor$ for $j := \ell(\chi)$, so that

$$j/3 \leq k \leq (j+2)/3.$$

Consider the case of $G = \text{Sp}_{2n}(q)$ with $2 \nmid q$. By [GLT2, Lemma 3.4], $j \leq 2n+1$, so

$$(k+1)/2 \leq (2n+6)/6 \leq n/2$$

when $n \geq 6$. Now, applying [GLT2, Theorem 1.5] we have

$$q^{nL} \geq \chi(1) \geq q^{k(n-(k+1)/2)} \geq q^{kn/2},$$

whence $k \leq 2L$ and so

$$(4.2) \quad j \leq 3k \leq 6L.$$

By definition, $\mathbf{l}(\chi) = j$ means that χ is an irreducible constituent of $(\omega + \omega^*)^j$, where ω and ω^* are the two reducible *Weil characters* of $\mathrm{Sp}_{2n}(q)$, see [GLT2, §3]. Next, $(\omega + \omega^*)^2$ always contains $(\tau_{2n,q})|_G$, and hence 1_G as well, by [GLT2, Proposition 3.1]. It follows that $(\omega + \omega^*)^j + (\omega + \omega^*)^{j-1}$ contains all irreducible constituents of $\sum_{i=0}^j (\omega + \omega^*)^i$. Hence, (4.2) implies that $N(L) \leq N^*(l) + N^*(l-1)$ with $l := \lfloor 6L \rfloor$, where $N^*(l)$ is the number of distinct irreducible constituents of $(\omega + \omega^*)^l$. By [GLT2, Proposition 3.2], $\omega^2 = (\omega^*)^2$, so in fact any irreducible constituent of $(\omega + \omega^*)^l$ is an irreducible constituent of ω^l or of $\omega^{l-1}\omega^*$. If $q \equiv 3 \pmod{4}$, then by [GLT2, Proposition 3.2] for $0 \leq i \leq l$ we have

$$\langle \omega^{l-i}(\omega^*)^i, \omega^{l-i}(\omega^*)^i \rangle = \langle \omega^{l-i}\bar{\omega}^i, \omega^{l-i}\bar{\omega}^i \rangle = \langle (\omega\bar{\omega})^{2l}, 1_G \rangle = \langle (\tau_{2n,q}^l)|_G, 1_G \rangle,$$

the number of G -orbits on V^l with $V := \mathbb{F}_q^{2n}$, which is at most $6q^{l(l-1)/2}$ by [GLT2, Lemma 2.6]. It follows that $N^*(l) \leq 12q^{l(l-1)/2}$.

If $q \equiv 1 \pmod{4}$, then by [GLT2, Proposition 3.2] for $0 \leq i \leq l$ we have

$$\langle \omega^{l-i}(\omega^*)^i, \omega^{l-i}(\omega^*)^i \rangle = \langle (\omega)^{2l-2i}(\omega^*)^{2i}, 1_G \rangle = \langle (\tau_{2n,q}^l)|_G, 1_G \rangle,$$

which is again the number of G -orbits on V^l with $V := \mathbb{F}_q^{2n}$ and so at most $6q^{l(l-1)/2}$ by [GLT2, Lemma 2.6]. Thus we also have $N^*(l) \leq 12q^{l(l-1)/2}$ in this case.

Thus $N(L) \leq 12q^{l(l-1)/2} + 12q^{(l-1)(l-2)/2} < q^{6L(6L-1)/2+3} < q^{18L^2}$.

(c) Now consider the case of $G = \mathrm{Sp}_{2n}(q)$ with $2|q$, or $\Omega_{2n+1}(q)$ with $2 \nmid q$, or $\Omega_{2n}^+(q)$. By [GLT2, Lemma 3.4], $j \leq n+1$, so

$$k+1 \leq (n+6)/3 \leq 2n/3$$

when $n \geq 6$. Applying [GLT2, Theorem 1.5] we have

$$q^{nL} \geq \chi(1) \geq q^{2k(n-(k+1))} \geq q^{2kn/3},$$

whence $k \leq 3L/2$ and so

$$(4.3) \quad j \leq 3k \leq 9L/2.$$

By definition, $\mathbf{l}(\chi) = j$ means that χ is an irreducible constituent of $((\tau + \zeta)|_G)^j$, where τ and ζ are the reducible Weil characters of $\mathrm{SL}_D(q)$ and $\mathrm{SU}_D(q)$, with $D = 2n$ or $2n+1$, the dimension of the natural module $V := \mathbb{F}_q^D$ of G . In particular, τ_G contains 1_G . Hence, (4.3) implies that $N(L) \leq N^*(l)$ with $l := \lfloor 9L/2 \rfloor$, where $N^*(l)$ is the number of distinct irreducible constituents of $(\tau|_G + \zeta|_G)^l$. Since $(\tau|_G)^2 = (\zeta|_G)^2$, any irreducible constituent of $(\tau|_G + \zeta|_G)^l$ is an irreducible constituent of $(\tau|_G)^l$ or of $(\tau|_G)^{l-1}(\zeta|_G)$. Now, for $0 \leq i \leq l$ we have

$$\langle (\tau|_G)^{l-i}(\zeta|_G)^i, (\tau|_G)^{l-i}(\zeta|_G)^i \rangle = \langle (\tau|_G)^{2l-2i}(\zeta|_G)^{2i}, 1_G \rangle = \langle (\tau^{2l})|_G, 1_G \rangle,$$

the number of G -orbits on V^{2l} . By [GLT2, Lemma 2.6], the number of $\tilde{G}$ -orbits on V^{2l} is at most $6q^{l(2l+1)}$, for $\tilde{G} = \mathrm{Sp}_D(q)$ or $\mathrm{GO}_D(q)$, respectively. Note that $[\tilde{G} : G]$ is 1 in the Sp -case, 2 in the Ω -case with $2 \nmid D$ or with $2|q$, and 4 otherwise. It follows that

$$N^*(l) \leq 12q^{l(2l+1)/2} \cdot [\tilde{G} : G] < q^{l(2l+1)+5} < q^{50L^2}.$$

(d) Finally, assume that $G = \Omega_{2n}^-(q)$. In this case, $j \leq n+1$, so

$$k+1 \leq (n+3)/3 \leq 3(n-1)/5$$

when $n \geq 6$. Applying [GLT2, Theorem 1.5] we have

$$q^{(n-1)L} \geq \chi(1) \geq q^{2k(n-1-k)} \geq q^{4k(n-1)/5},$$

whence $k \leq 5L/4$ and so $j \leq 3k \leq 15L/4$. Now we can repeat the arguments in (c) to get $N(l) \leq N^*(l) \leq q^{l(2l+1)+5} < q^{41L^2}$. $\square$

We can now prove Theorem 3 and Corollary 4.

Proof of Theorem 3. Let $\epsilon > 0$ be given. We claim there exists $\delta > 0$ such that if G is a finite quasisimple group of Lie type, and χ_1 and χ_2 are (possibly reduced) characters of G with $|\chi_1|, |\chi_2| \leq |G|^\delta$, then

$$|\chi_1 \chi_2| \geq (|\chi_1| \cdot |\chi_2|)^{1-\epsilon}.$$

It suffices to prove the statement in the case $|\chi_1|, |\chi_2| > 1$; in particular, the χ_i contain nontrivial irreducible constituents α_1 and α_2 , respectively. We may also assume that each χ_i is multiplicity-free. If G is an exceptional group of Lie type, or a finite classical group of rank ≤ 7 , then [LaSe] implies that $\alpha_i(1) > |G|^{1/20}$. Taking $\delta \leq 1/10$, we may assume G is a classical group of rank ≥ 8 . By [GLT2, Proposition 6.3], if G is of orthogonal type, the bounds on $|\chi_i|$ imply that they arise from characters of $\Omega_{2n+1}^+(q)$, $\Omega_{2n}^+(q)$, or $\Omega_{2n}^-(q)$ by composition with a central quotient map.

We will view χ_1, χ_2 as characters of $\hat{G} = \mathrm{SL}_n^\pm(q)$, $\mathrm{Sp}_{2n}(q)$, $\Omega_{2n+1}^+(q)$, $\Omega_{2n}^+(q)$, or $\Omega_{2n}^-(q)$, respectively, where $n \geq 8$. In all cases, $\hat{G}$ contains a subgroup $H \cong \mathrm{SL}_m^\pm(q)$ with $m := n$ in the first case, and $H \cong \mathrm{SL}_m(q)$ (inside a split Levi subgroup) with $m := n, n, n$, or $n-1$, respectively, in the other four cases. One checks that $q^{m^2/2} < |G| < q^{4m^2}$ in all cases; furthermore, $\alpha_i(1) > q^{m/2}$. Writing

$$(4.4) \quad q^{mL_i/2} \leq \chi_i(1) = q^{mD_i} \leq q^{mL_i}$$

for some (real) numbers $L_i \geq 1$ and $L_i/2 \leq D_i \leq L_i$, we will choose δ so that

$$q^{mL_i} \leq |\chi_i| \leq |G|^\delta \leq q^{4m^2\delta};$$

in particular,

$$(4.5) \quad L_i \leq 4m\delta.$$

Writing

$$\chi_1 \chi_2 = \sum_{\gamma \in \mathrm{Irr}(G)} m_\gamma \gamma, \quad \chi_1^2 = \sum_{\gamma \in \mathrm{Irr}(G)} a_\gamma \gamma, \quad \chi_2^2 = \sum_{\gamma \in \mathrm{Irr}(G)} b_\gamma \gamma,$$

we have by the Cauchy-Schwarz inequality

$$\begin{aligned} \left(\sum_{\gamma} m_\gamma^2 \right)^2 &= \langle \chi_1 \chi_2, \chi_1 \chi_2 \rangle^2 \\ &= \langle \chi_1 \bar{\chi}_1, \chi_2 \bar{\chi}_2 \rangle^2 \\ &\leq \langle \chi_1 \bar{\chi}_1, \chi_1 \bar{\chi}_1 \rangle \cdot \langle \chi_2 \bar{\chi}_2, \chi_2 \bar{\chi}_2 \rangle \\ &= \langle \chi_1^2, \chi_1^2 \rangle \cdot \langle \chi_2^2, \chi_2^2 \rangle \\ &= \sum_{\gamma} a_\gamma^2 \cdot \sum_{\gamma} b_\gamma^2 \\ &\leq \left(\sum_{\gamma} a_\gamma \right)^2 \cdot \left(\sum_{\gamma} b_\gamma \right)^2, \end{aligned}$$

and thus

$$(4.6) \quad \sum_{\gamma} m_\gamma^2 \leq \sum_{\gamma} a_\gamma \cdot \sum_{\gamma} b_\gamma = \sigma(\chi_1^2, G) \cdot \sigma(\chi_2^2, G),$$

where, $\sigma(\cdot, \cdot)$, as defined in [GLT2, (2.1)], denotes the sum of the multiplicities of all the irreducible constituents of the first argument, regarded as a character on the group specified in the second argument.

First we consider the case $\hat{G} = \mathrm{SL}_n^\pm(q)$. Then each irreducible constituent θ of χ_i lies under some $\tilde{\theta} \in \mathrm{Irr}(\mathrm{GL}_n^\pm(q))$ of degree less than $(q \mp 1)\theta(1)$. Hence χ_i lies under some character $\tilde{\chi}_i$ of $\mathrm{GL}_n^\pm(q)$ of degree at most $q^{mL_i+2} \leq q^{5mL_i/4}$ since $m \geq 8$. Applying [GLT2, Corollary 5.2(i)] to $\tilde{\chi}_i$ and restricting further down to $\mathrm{SL}_n^\pm(q)$, we have

$$\sigma(\chi_i^2, G) \leq q^{(125/4)L_i^2+2} \sigma(\chi_i, G)^2 < q^{46L_i^2}.$$

Here, we use the fact that χ_i is multiplicity-free, and so $\sigma(\chi_i, G)$ is at most the number of irreducible characters of G of degree $\leq q^{nL_i}$, which is at most $q^{12L_i^2}$ by Proposition 4.1(a).

Next suppose we are in the symplectic-orthogonal case. Taking $\delta > 0$ small enough, we may assume that $L_i \leq n/9$. Applying Theorem 5.8 and Corollary 5.9 of [GLT2] to χ and restricting further down from a normal subgroup of $\mathrm{GL}_m(q)$ containing $H = \mathrm{SL}_m(q)$ to H if necessary, we have

$$\sigma(\chi_i^2, G) \leq q^{A\sqrt{mL_i^3}+60L_i^2} \sigma(\chi_i, G)^2 \leq q^{A\sqrt{mL_i^3}+110L_i^2}$$

for some explicit A (which can be taken to be 70). Here, we again use the fact that χ_i is multiplicity-free, and so $\sigma(\chi_i, G)$ is at most the number of irreducible characters of G of degree $\leq q^{mL_i}$, which is at most $q^{50L_i^2}$ by Proposition 4.1(b)–(d).

Using (4.6), in either case we have

$$\sum_{\gamma} m_{\gamma}^2 \leq q^{A\sqrt{mL_1^3}+A\sqrt{mL_2^3}+BL_1^2+BL_2^2}$$

with $B = 110$. It follows that

$$|\chi_1 \chi_2| = \sum_{m_{\gamma} > 0} \gamma(1)^2 \geq \frac{(\sum_{\gamma} m_{\gamma} \gamma(1))^2}{\sum_{\gamma} m_{\gamma}^2} = \frac{\chi_1(1)^2 \chi_2(1)^2}{\sum_{\gamma} m_{\gamma}^2} \geq q^{2m(D_1+D_2)-A\sqrt{mL_1^3}-A\sqrt{mL_2^3}-BL_1^2-BL_2^2}.$$

Recalling (4.5), we observe

$$\frac{\sqrt{mL_1^3} + \sqrt{mL_2^3}}{m(D_1 + D_2)} \leq \frac{2mL_1\sqrt{\delta} + 2mL_2\sqrt{\delta}}{m(L_1 + L_2)/2} = 4\sqrt{\delta},$$

and

$$\frac{L_1^2 + L_2^2}{m(D_1 + D_2)} \leq \frac{2mL_1\delta + 2mL_2\delta}{m(L_1 + L_2)/2} = 4\delta.$$

Hence

$$|\chi_1 \chi_2| \geq q^{2m(D_1+D_2)(1-2A\sqrt{\delta}-2B\delta)}.$$

Note that $|\chi_1| \cdot |\chi_2| \leq \chi_1(1)^2 \chi_2(1)^2 = q^{2m(D_1+D_2)}$. Hence, taking $\delta > 0$ so that

$$2A\sqrt{\delta} + 2B\delta \leq \epsilon,$$

we have $|\chi_1 \chi_2| \geq (|\chi_1| \cdot |\chi_2|)^{1-\epsilon}$, as desired. $\square$

Proof of Corollary 4. We prove by induction on $k \geq 2$ the equivalent statement:

For any $\epsilon > 0$ and any $k \geq 2$, there exists an explicit $\gamma > 0$ (depending on both ϵ and k) such that the following statement holds. If G is a finite quasisimple group of Lie type and $\chi_1, \chi_2, \dots, \chi_k$ are any characters of G with $|\chi_1|, |\chi_2|, \dots, |\chi_k| \leq |G|^{\gamma}$, then

$$|\chi_1 \chi_2 \cdots \chi_k| \geq (|\chi_1| \cdot |\chi_2| \cdots |\chi_k|)^{1-k\epsilon}.$$

We will show that this statement holds with $\gamma := \delta/(k-1)$, where δ the constant in Theorem 3. The case $k = 2$ already established in Theorem 3. For the inductive step, note by Lemma 2.1 and the induction hypothesis that

$$(|\chi_2| \cdots |\chi_k|)^{1-(k-1)\epsilon} \leq |\chi_2 \cdots \chi_k| \leq \prod_{i=2}^k |\chi_i| \leq |G|^{\gamma(k-1)} \leq |G|^\delta.$$

Since $|\chi_1| \leq |G|^\delta$, by Theorem 3 we have

$$|\chi_1 \chi_2 \cdots \chi_k| \geq (|\chi_1| \cdot |\chi_2 \cdots \chi_k|)^{1-\epsilon} \geq (|\chi_1| \cdot (|\chi_2| \cdots |\chi_k|)^{1-(k-1)\epsilon})^{1-\epsilon} \geq (|\chi_1| \cdot |\chi_2| \cdots |\chi_k|)^{1-k\epsilon}.$$

□

The next result is required in our proof of Theorem 5.

Theorem 4.2. *For any $\delta > 0$, there exists an explicit integer N such that the following statement holds. If G is a finite simple group of Lie type, and $\chi_1, \dots, \chi_N$ are any (not necessarily irreducible) characters of G with $|\chi_i| \geq |G|^\delta$ for all i , then $|\chi_1 \chi_2 \cdots \chi_N| = |G|$ and thus $\chi_1 \chi_2 \cdots \chi_N$ contains every irreducible character of G .*

Proof. For any character χ of G , let χ^* denote (some) irreducible constituent of largest degree of χ .

First we consider the case $k(G) \geq |G|^{\delta/3}$. If G is of rank r over $\mathbb{F}_q$, then by [FG] we have $k(G) \leq (27.2)q^r$, whereas $|G| \geq q^{r^2}$. It follows that $r \leq r_0$ is bounded (in terms of δ). On the other hand, $|\chi_i| > 1$ implies that $\chi_i^* \neq 1_G$, and so $\chi_i^*(1) > q^r/3$ by [LaSe]. As $r \leq r_0$, $\chi_i^*(1) \geq |G|^{\delta_0}$ for some δ_0 depending on δ . Applying [LaT, Theorem 8.5] and taking $N \geq \epsilon/\delta_0$ (with ϵ the constant in [LaT, Theorem 8.5]), we see that $\chi_1^* \cdots \chi_N^*$ contains $\text{Irr}(G)$.

Now suppose that $k(G) \leq |G|^{\delta/3}$. Then $\chi_i^*(1) \geq (|\chi_i|/k(G))^{1/2} \geq |G|^{\delta/3}$. Applying [LaT, Theorem 8.5] and taking $N \geq 3\epsilon/\delta$, we again see that $\chi_1^* \cdots \chi_N^*$ contains $\text{Irr}(G)$. □

Proof of Theorem 5. Fix $\epsilon > 0$. It follows from Theorem 3, applied for $\chi_1 = \chi_2 = \chi$ with ϵ replaced by $\epsilon/2$, that there exists some $\delta > 0$ depending only on ϵ such that if $|\chi| \leq |G|^\delta$, then $|\chi^2| \geq |\chi|^{2-\epsilon}$. By Theorem 4.2, there is some integer $N > 0$ depending only on ϵ , such that if $|\chi| \geq |G|^\delta$ then χ^N contains $\text{Irr}(G)$. The result follows. □

Proof of Theorem 7. For any character χ of G , again let χ^* denote (some) irreducible constituent of largest degree of χ . It suffices to prove the statement in the case $|\chi_i| > 1$, and thus $\chi_i^* \neq 1_G$. By the results of [FG] and [LaSe], the degree of any non-trivial irreducible character of G is at least $k(G)^{1/6}$; in particular,

$$\chi_i^*(1) \geq k(G)^{1/10} \geq m_i^{1/6},$$

if m_i denotes the number of distinct irreducible constituents of χ_i . It follows that

$$|\chi_i| \leq m_i \chi_i^*(1)^2 \leq \chi_i^*(1)^8,$$

and so $\chi_i^*(1) \geq |\chi_i|^{1/8}$. Now applying [LaT, Theorem 8.5] and taking $c = 8\epsilon$ (with ϵ the constant in [LaT, Theorem 8.5]), we have that $\prod_{i=1}^m \chi_i^*(1) \geq |G|^\epsilon$, and so $\chi_1^* \cdots \chi_N^*$ contains $\text{Irr}(G)$. □

5. PROOF OF THEOREM 6

First we prove the theorem when $G = \mathrm{PSL}_2(q)$. Let χ_1 and χ_2 denote any irreducible characters of G . Then $|\chi_i(g)| \leq q^{1/2}$ for all $g \in G$, $|\chi_i(g)| \leq 2$ for all non-trivial semisimple $g \in G$, and the total number of non-semisimple elements of G is less than q . Therefore

$$\sum_{g \neq 1} |\chi_1^6(g) \chi_2(g)| \leq \chi_2(1)q^4 + 128q^3.$$

If χ_1 is non-trivial, it has degree at least $\frac{q-1}{2}$, so for q sufficiently large,

$$\sum_{g \in G} \chi_1^6(g) \chi_2(g) \neq 0,$$

and applying this to any non-trivial constituent χ_1 or χ_2 , we get the desired result.

As before, we use the notation PSL^ϵ to denote PSL when $\epsilon = +$ and PSU when $\epsilon = -$. Next we consider the case $G = \mathrm{PSL}_3^\epsilon(q)$. Again, the generic character table has been computed explicitly [SF]. There are $O(q)$ characters of degree $\leq q^2 + q + 1$, and all other characters have degree at least $q^3/4$ (when q is not too small). Therefore, the sum of the squares of the degrees of irreducible characters of G of degree $< q^3/4$ is smaller than $|G|^{11/12}$ for large values of q . We therefore assume χ has an irreducible constituent χ_1 of degree $\geq q^3/4$. We have $|\chi_i(g)| = O(q)$ for all irreducible characters and all non-trivial elements, so

$$\sum_{g \neq 1} |\chi_1^6(g) \bar{\theta}(g)| = O(q^6 |G| \theta(1)) = O(q^{14} \theta(1)) = o(\chi_1(1)^6 \theta(1)).$$

Therefore, if $\chi_1(q) \geq q^3/4$ and q is sufficiently large, any $\theta \in \mathrm{Irr}(G)$ is a constituent of χ_1^6 .

We may therefore assume $G = \mathrm{PSL}_n^\epsilon(q)$ with $n \geq 4$. We claim that for q sufficiently large, $|\chi| \geq |G|^{11/12}$ implies χ has an irreducible constituent χ_1 with

$$(5.1) \quad \chi_1(1) \geq q^{\frac{44n^2}{131}}.$$

Indeed, the number of characters of G which do not satisfy this inequality is less than $k(G) = O(q^{n-1})$ and $|G| = O(q^{n^2-1})$, so it suffices to note that

$$\frac{88n^2}{131} + (n-1) < \frac{11(n^2-1)}{12}$$

for $n \geq 4$.

We claim that if q sufficiently large in terms of n , (5.1) implies that the Steinberg character St is a constituent of χ_1^3 . If $\epsilon = +$ or if $\epsilon = -$ but $2 \nmid n$, then every irreducible character of G appears in St^2 [HSTZ, Theorem 1.2], and so it follows that $|\chi_1^6| = |G|$. Consider the case $\epsilon = -$ and $2 \nmid n$. Then by [HSTZ, Theorem 1.2], St^2 contains all but one irreducible character α of $\mathrm{PSU}_n(q)$, which has smallest degree $(q^n - q)/(q + 1)$ among all nontrivial irreducible characters of G . Let $\theta \in \mathrm{Irr}(G)$. By Proposition 2.2(i) and (5.1),

$$|\theta \bar{\chi}_1| \geq |\chi_1(1)| > |\alpha|,$$

and so $\theta \bar{\chi}_1 \neq m\alpha$ for any $m \in \mathbb{N}$ by (5.1). Hence $\theta \bar{\chi}_1$ contains some irreducible character $\beta \neq \alpha$, whence $\langle \mathrm{St}^2, \beta \rangle > 0$ and so

$$\langle \chi_1^7, \theta \rangle = \langle \chi_1^6, \theta \bar{\chi}_1 \rangle \geq \langle \mathrm{St}^2, \beta \rangle > 0,$$

and thus χ_1^7 contains $\mathrm{Irr}(G)$, as desired.

It suffices to show that for large enough q ,

$$(5.2) \quad \sum_{g \neq 1} \frac{|\mathrm{St}(g)\chi_1(g)^3|}{|\mathrm{St}(1)\chi_1(1)^3|} < 1.$$

As St vanishes at all non-semisimple elements, we need only consider semisimple elements g in (5.2). Let X_s denote the set of conjugacy classes of semisimple elements in G of support s , i.e., for which the largest dimension of any $\overline{\mathbb{F}}_q$ -eigenspace is $n - s$. We can rewrite (5.2) as

$$(5.3) \quad \sum_{s=1}^{n-1} \sum_{C \in X_s} |C| \frac{|\mathrm{St}(C)|}{|\mathrm{St}(1)|} \left(\frac{|\chi_1(C)|}{|\chi_1(1)|} \right)^3 < 1.$$

We consider the factors of the summand in the left hand side of (5.3), one by one. If $C \in X_s$ has an element g represented by a matrix M with eigenvalue multiplicities $m_1, \dots, m_k$, then the centralizer of M in $\mathrm{GL}_n^\epsilon(q)$ is the group of $\mathbb{F}_q$ points (possibly twisted) of a connected algebraic group of dimension $\sum_i m_i^2$, so its order is $(1 + o(1))q^{\sum_i m_i^2}$. Therefore, C , which is contained in the $\mathrm{PSL}_n^\epsilon(q)$ -conjugacy class of g , has cardinality

$$O(\mathrm{gcd}(n, q - \epsilon)q^{n^2 - \sum_i m_i^2}) = O(q^{n^2 - \sum_i m_i^2}).$$

The absolute value of $\mathrm{St}(C) = \mathrm{St}(g)$ is the order of a p -Sylow subgroup of the centralizer of M , which is $q^{\sum_i \binom{m_i}{2}}$, and $\mathrm{St}(1) = q^{\binom{n}{2}}$. Therefore,

$$\frac{|\mathrm{St}(C)|}{|\mathrm{St}(1)|} = q^{-\frac{n^2 - \sum_i m_i^2}{2}},$$

and

$$|C| \frac{|\mathrm{St}(C)|}{|\mathrm{St}(1)|} = O\left(q^{\frac{n^2 - \sum_i m_i^2}{2}}\right).$$

As some m_i equals $n - s$, we have

$$\sum_i m_i^2 \geq (n - s)^2 + s \cdot 1^2,$$

so

$$|C| \frac{|\mathrm{St}(C)|}{|\mathrm{St}(1)|} = O\left(q^{\frac{2sn - s^2 - s}{2}}\right).$$

By [TT, Theorem 1.13] and [LiST, Theorem 3.1], for q large enough we have

$$\frac{|\chi_1(g)|}{|\chi_1(1)|} = O(\chi_1(1)^{-s/n}) = O(q^{-\frac{44sn}{131}}),$$

where the implicit constant depends only on n . Thus, the summand as a whole is $O(q^{-\frac{sn}{131} - \frac{s^2 + s}{2}})$.

Finally, we claim that $|X_s| = O(q^s)$. Since each conjugacy class of $\mathrm{PGL}_n^\epsilon(q)$ decomposes into at most $\mathrm{gcd}(n, q - \epsilon)$ conjugacy classes of G , it suffices to prove the same thing for $\mathrm{PGL}_n^\epsilon(q)$ conjugacy classes containing semisimple elements of $\mathrm{PSL}_n^\epsilon(q)$ or, indeed, $\mathrm{GL}_n^\epsilon(\mathbb{F}_q)$ conjugacy classes containing semisimple elements of $\mathrm{SL}_n^\epsilon(\mathbb{F}_q)$. Since any such element has connected centralizer in $\mathrm{GL}_n(\overline{\mathbb{F}}_q)$, any such class is uniquely determined by the spectrum of its representatives, equivalently, by their common characteristic polynomial.

Let $Y_{s,i}$ denote the set of conjugacy classes in $\mathrm{GL}_n^\epsilon(\mathbb{F}_q)$ of semisimple elements M in $\mathrm{SL}_n^\epsilon(\mathbb{F}_q)$ which have exactly i eigenvalues of multiplicity $n - s$ (and all other eigenvalues of lower multiplicity), where $1 \leq i \leq t := \lfloor n/(n - s) \rfloor$. Let $P(x)$ be the monic polynomial of degree i

whose roots are the i eigenvalues of M of multiplicity $n-s$, and $Q(x)$ be the monic polynomial of degree $n-i(n-s)$, such that $P(x)^i Q(x)$ is the characteristic polynomial of M . Then

$$\det P(0)^i Q(0) = (-1)^n \det(M) = (-1)^n.$$

So the constant term of $P(x)$ is determined up to at most $i \leq n$ possibilities by the constant term of $Q(x)$, and so there are at most nq^{i-1} possibilities for $P(x)$ for any fixed (constant term of) $Q(x)$. Since there are at most $q^{n-i(n-s)}$ possibilities for $Q(x)$, we see that the total number of possibilities for $P(x)^i Q(x)$, which, as explained above, gives an upper bound for $|Y_{s,i}|$, is at most

$$nq^{i-1}q^{n-i(n-s)} = nq^{s+(1-i)(n-s-1)} = O(q^s).$$

Since q is large compared to n , it follows that $|X_s| = \sum_{i=1}^t |Y_{s,i}| = O(q^s)$, as claimed.

We conclude that the left hand side of (5.3) is $O(q^{-sn/131})$. Taking q large enough, the statement now follows.

6. SEMISIMPLE COMPACT LIE GROUPS

In this section, we briefly consider the situation when G is a compact (connected) semisimple Lie group instead of a finite simple group of Lie type. For χ a character of G , the definition of $|\chi|$ works as before. We have the following analogue of Proposition 2.2, which no longer requires irreducibility.

Proposition 6.1. *Let G be a semisimple compact Lie group, $n \geq 2$, and $\chi_1, \chi_2, \dots, \chi_n$ non-trivial characters. Then $|\chi_1 \chi_2 \cdots \chi_n| > \max(|\chi_1|, |\chi_2|, \dots, |\chi_n|)$.*

Proof. It suffices to treat the case $n = 2$ and to prove in this case that if χ_1 is non-trivial, then $|\chi_1 \chi_2| > |\chi_2|$. It is enough to treat the case that χ_1 is irreducible. Let λ_1 denote the highest weight of χ_1 . Let $\{\varphi_1, \dots, \varphi_k\}$ denote the irreducible constituents of χ_2 , and let μ_i denote the highest weight of φ_i .

Then $\lambda_1 + \mu_i$ is the highest weight in $\chi_1 \varphi_i$, so there is an irreducible constituent ψ_i of $\chi_1 \chi_2$ with highest weight $\lambda_1 + \mu_i$. By the Weyl dimension formula,

$$(6.1) \quad |\chi_1 \chi_2| \geq \sum_{i=1}^k |\psi_i|^2 = \sum_{i=1}^k \prod_{\alpha \succ 0} \frac{(\delta + \lambda_1 + \mu_i, \alpha)^2}{(\delta, \alpha)^2} = \sum_{i=1}^k \prod_{\alpha \succ 0} \frac{((\delta + \mu_i, \alpha) + (\lambda_1, \alpha))^2}{(\delta, \alpha)^2},$$

where δ denotes half the sum of the positive roots. As λ_1 is a non-zero dominant weight, we have (λ_1, α) non-negative for all positive roots α and strictly positive for at least one of them. Therefore, the right hand side of (6.1) is strictly greater than

$$\sum_{i=1}^k \prod_{\alpha \succ 0} \frac{(\delta + \mu_i, \alpha)^2}{(\delta, \alpha)^2} = \sum_{i=1}^k |\varphi_i|^2 = |\chi_2|.$$

□

When G is of positive dimension, it no longer makes sense to compare $|\chi|$ to $|G|$, so we do not have an analogue of Theorem 3 or any of the subsequent results proved above for groups of Lie type. We can still ask about power growth of $|\chi|$. The case of $\text{SU}(2)$ illustrates the situation: we get uniform power growth when χ is irreducible, but there is no such growth for general characters.

Proposition 6.2. *Let $G = \text{SU}(2)$.*

(i) If χ ranges over irreducible characters of G , then

$$\lim_{\chi} \frac{\log |\chi^2|}{\log |\chi|} = \frac{3}{2}.$$

(ii) If χ ranges over all characters of G , then

$$\liminf_{\chi} \frac{\log |\chi^2|}{\log |\chi|} = 1.$$

Proof. If χ_n denotes the unique irreducible character of G of degree $n+1$, then by the Clebsch-Gordan formula ([Hu, §22, Ex. 7]),

$$\chi_n^2 = \sum_{i=0}^n \chi_{2i}.$$

Thus,

$$|\chi_n^2| = \sum_{i=0}^n (2i+1)^2 = \binom{2n+3}{3},$$

while $|\chi_n| = (n+1)^2$. This gives (i).

Again by Clebsch-Gordan, χ_n^4 is a linear combination of $\chi_0, \chi_2, \chi_4, \dots, \chi_{4n}$ with all coefficients positive, so

$$|\chi_n^4| = \binom{4n+3}{3}.$$

Therefore, as χ ranges over $\{\chi_n, | n \in \mathbb{N}\}$, the limit of $\frac{|\chi^2|}{|\chi|}$ is 8, and the limit of $\frac{\log |\chi^2|}{\log |\chi|}$ is 1. $\square$

Next, we prove Theorem 8 establishing uniform power growth for irreducible characters of any fixed compact semisimple Lie group.

Proof of Theorem 8. Let $\varpi_1, \dots, \varpi_r$ denote the fundamental weights of G . Let χ_λ be the irreducible character with highest weight $\lambda = a_1\varpi_1 + \dots + a_r\varpi_r$. We define

$$\langle \lambda, \alpha \rangle := \frac{2(\lambda, \alpha)}{(\alpha, \alpha)}.$$

By [Hu, §22, Ex. 1], $\lambda - k\alpha_i$ is a weight of χ_λ for $0 \leq k \leq \langle \lambda, \alpha_i \rangle = a_i$.

We claim that $\delta + 2\lambda - k\alpha_i$ is dominant for $0 \leq k \leq a_i$. It suffices to check the non-negativity of $\langle \delta + 2\lambda - k\alpha_i, \alpha_j \rangle$ for $1 \leq j \leq r$. For $j \neq i$, we have

$$\langle \delta + 2\lambda - k\alpha_i, \alpha_j \rangle = 1 + 2a_j - \langle \alpha_i, \alpha_j \rangle \geq 1,$$

and for $j = i$, we have

$$\langle \delta + 2\lambda - k\alpha_i, \alpha_i \rangle = 1 + 2a_i - 2k \geq 1.$$

Therefore, by a theorem of Brauer [Hu, §24, Ex. 9], for each k in this range, χ_λ^2 contains the irreducible character with highest weight $2\lambda - k\alpha_i$.

By the Weyl dimension formula [Hu, §24.3],

$$\chi_\lambda(1) = \prod_{\alpha > 0} \frac{(\lambda + \delta, \alpha)}{(\delta, \alpha)},$$

where the product is taken over the set Φ^+ of positive roots α . Now, for $1 \leq i \leq r$, $(\varpi_i, \alpha) \geq 0$, so regarded as a function in λ , $\chi_\lambda(1)$ is a product of affine linear functions in λ which take positive values in the cone of dominant weights. Thus, if $a_i \geq 1$ and $0 \leq k \leq a_i$, then

$$(6.2) \quad \chi_{2\lambda - k\alpha_i}(1) \geq (1 - k/a_i)^{|\Phi^+|} \chi_{2\lambda}(1) \geq (1 - k/a_i)^{|\Phi^+|} \chi_\lambda(1).$$

We fix i such that $a_i = \max(a_1, \dots, a_r)$. Thus,

$$|\chi_\lambda| = \chi_\lambda(1)^2 = O(a_i^{2|\Phi^+|}) = O(a_i^{|\Phi|}).$$

On the other hand, by (6.2),

$$|\chi_\lambda^2| \geq \sum_{k=0}^{a_i} |\chi_{2\lambda - k\alpha_i}| \geq |\chi_\lambda| \sum_{k=0}^{a_i} (1 - k/a_i)^{|\Phi|}.$$

As $a_i \rightarrow \infty$, the sum $\sum_k (1 - k/a_i)^{|\Phi|}$ can be bounded below by a positive constant multiple of a_i and therefore by a positive constant multiple of $|\chi_\lambda|^{1/|\Phi|}$. If $\epsilon < 1/|\Phi|$, therefore, $|\chi_\lambda^2| \geq |\chi_\lambda|^{1+\epsilon}$ with finitely many exceptions λ . The theorem is trivial for $\lambda = 0$, and for each $\lambda \neq 0$, it holds when $\epsilon > 0$ is small enough by Proposition 6.1. $\square$

Note that, unlike Theorem 1, Theorem 8 does not guarantee that power growth is uniform in G . This is unavoidable, as the following proposition shows.

Proposition 6.3. *There exists a sequence $\chi_2, \chi_3, \dots$ of non-trivial irreducible characters of the Lie groups $SU(2), SU(3), \dots$ respectively, such that*

$$\liminf_n \frac{\log |\chi_n^2|}{\log |\chi_n|} = 1.$$

Proof. We choose for each $n \geq 2$ a positive integer k_n and define χ_n to be the character of the irreducible representation of $SU(n)$ with highest weight $\lambda_n := k_n \delta_n$, where δ_n is half the sum of the set Φ_n^+ of positive roots of $SU(n)$. By the Weyl dimension formula,

$$\chi_n(1) = \prod_{\alpha \in \Phi_n^+} \frac{(\lambda_n + \delta_n, \alpha)}{(\delta_n, \alpha)} = (k_n + 1)^{|\Phi_n^+|} = (k_n + 1)^{n(n-1)/2}.$$

On the other hand, for each fixed n , the weights of χ_n are contained in the convex hull of $\{w(\lambda_n) \mid w \in S_n\}$. This can be expressed as $k_n X_n$, where X_n denotes the convex hull of $\{w(\delta_n) \mid w \in S_n\}$. Thus, the weights of χ_n^2 are contained in $2k_n X_n$. Since the highest weight of every irreducible constituent of χ_n^2 is a lattice point in the fixed polytope $2X_n \subset \mathbb{R}^{n-1}$, scaled by k_n , the number of such highest weights is $O(k_n^{n-1})$, where the implicit constant depends only on n .

On the other hand, by the Weyl dimension formula, the degree of each irreducible factor of χ_n^2 is $O(k_n^{n(n-1)/2})$. Therefore, assuming that each k_n is sufficiently large, we can guarantee

$$\frac{\log |\chi_n^2|}{\log |\chi_n|} < \frac{n+1}{n},$$

which implies the proposition. $\square$

REFERENCES

- [Br] Breuillard, Emmanuel: Lectures on approximate groups and Hilbert's 5th problem. *Recent trends in combinatorics*. 369–404, IMA Vol. Math. Appl., **159**, Springer, 2016.
- [BGT] Breuillard, Emmanuel; Green, Ben; Tao, Terence: Approximate subgroups of linear groups, *Geom. Funct. Anal.* **21** (2011), no. 4, 774–819.
- [FG] Fulman, Jason; Guralnick, Robert: Bounds on the number and sizes of conjugacy classes in finite Chevalley groups with applications to derangements. *Trans. Amer. Math. Soc.* **364** (2012), no. 6, 3023–3070.
- [Gi] Gill, Nick: blog post, <https://nickpgill.github.io/a-rodgers-saxl-conjecture-for-characters>.

- [GPSS] Gill, Nick; Pyber, László; Short, Ian; Endre Szabó: On the product decomposition conjecture for finite simple groups, *Groups Geom. Dyn.* **7** (2013), no. 4, 867–882.
- [Gl] Gluck, David, Character value estimates for nonsemisimple elements. *J. Algebra* **155** (1993), no. 1, 221–237.
- [GLT1] Guralnick, Robert M.; Larsen, Michael; Tiep, Pham Huu: Character levels and character bounds, *Forum of Math. Pi* **8** (2020), e2, 81 pages.
- [GLT2] Guralnick, Robert M.; Larsen, Michael; Tiep, Pham Huu: Character levels and character bounds for finite classical groups, arXiv:1904.08070.
- [HSTZ] Heide, Gerhard; Saxl, Jan; Tiep, Pham Huu; Zalesski, Alexandre E.: Conjugacy action, induced representations and the Steinberg square for simple groups of Lie type. *Proc. Lond. Math. Soc.* (3) **106** (2013), no. 4, 908–930.
- [He1] Helfgott, Harald: Growth and generation in $SL_2(\mathbb{Z}/p\mathbb{Z})$. *Annals of Math.* **167** (2008), no. 2, 601–623.
- [He2] Helfgott, Harald: Growth in $SL_3(\mathbb{Z}/p\mathbb{Z})$, *J. Eur. Math. Soc. (JEMS)* **13** (2011), no. 3, 761–851.
- [HI] Howlett, Robert, B; Isaacs, I. Martin: On groups of central type, *Math. Z.* **179** (1982), 555–569.
- [Hr] Hrushovski, Ehud: Stable group theory and approximate subgroups, *J. Amer. Math. Soc.* **25** (2012), no. 1, 189–243.
- [Hu] Humphreys, James E.: Introduction to Lie algebras and representation theory. Second printing, revised. Graduate Texts in Mathematics, 9. Springer-Verlag, New York-Berlin, 1978.
- [LaSe] Landazuri, Vicente; Seitz, Gary M.: On the minimal degrees of projective representations of the finite Chevalley groups. *J. Algebra* **32** (1974), 418–443.
- [LaT] Larsen, Michael; Tiep, Pham Huu: Uniform character bounds for finite classical groups, preprint, 2021.
- [LiSh] Liebeck, Martin; Shalev, Aner: Diameters of finite simple groups: sharp bounds and applications, *Annals of Math.* **154** (2001), 383–406.
- [LiSh2] Liebeck, Martin; Shalev, Aner: Character degrees and random walks in finite groups of Lie type. *Proc. London Math. Soc.* **90** (2005), 61–86.
- [LSS] Liebeck, Martin; Schul, Gili; Shalev, Aner: Rapid growth in finite simple groups. *Trans. Amer. Math. Soc.* **369** (2017), no. 12, 8765–8779.
- [LiST] Liebeck, Martin; Shalev, Aner; Tiep, Pham Huu: On the diameter of McKay graphs for finite simple groups. *Isr. J. Math.* **241** (2021), no. 1, 449–464.
- [PS] Pyber, László; Szabó, Endre: Growth in finite simple groups of Lie type, *J. Amer. Math. Soc.* **29** (2016), no. 1, 95–146.
- [RS] Rodgers, D. M.; Saxl, Jan: Products of conjugacy classes in the special linear groups, *Comm. Algebra* **31** (2003), 4623–4638.
- [Se] Sellke, Mark: Tensor quasi-random groups, arXiv:2103.11048.
- [SF] Simpson, William A.; Frame, J. Sutherland: The character tables for $SL(3, q)$, $SU(3, q^2)$, $PSL(3, q)$, $PSU(3, q^2)$. *Canadian J. Math.* **25** (1973), 486–494.
- [Sh] Shalev, Aner: Word maps, conjugacy classes, and a noncommutative Waring-type theorem, *Annals of Math.* **170** (2009), no. 3, 1383–1416.
- [TT] Taylor, Jay; Tiep, Pham Huu: Lusztig induction, unipotent supports, and character bounds. *Trans. Amer. Math. Soc.* **373** (2020), no. 12, 8637–8676.

Email address: mjlarsen@indiana.edu

DEPARTMENT OF MATHEMATICS, INDIANA UNIVERSITY, BLOOMINGTON, IN 47405, U.S.A.

Email address: shalev@math.huji.ac.il

EINSTEIN INSTITUTE OF MATHEMATICS, HEBREW UNIVERSITY, GIVAT RAM, JERUSALEM 91904, ISRAEL

Email address: tiep@math.rutgers.edu

DEPARTMENT OF MATHEMATICS, RUTGERS UNIVERSITY, PISCATAWAY, NJ 08854-8019, U.S.A.