

On the two and one-half dimensional Vlasov–Poisson system with an external magnetic field: Global well-posedness and stability of confined steady states

Patrik Knopf ^{*} Jörg Weber [†]

December 1, 2021

This is a preprint version of the paper. Please cite as:

P. Knopf and J. Weber, Nonlinear Anal. Real World Appl. 65:103460 (2022)

<https://doi.org/10.1016/j.nonrwa.2021.103460>

Abstract

The time evolution of a two-component collisionless plasma is modeled by the Vlasov–Poisson system. In this work, the setting is two and one-half dimensional, that is, the distribution functions of the particles species are independent of the third space dimension. We consider the case that an external magnetic field is present in order to confine the plasma in a given infinitely long cylinder. After discussing global well-posedness of the corresponding Cauchy problem, we construct stationary solutions whose support stays away from the wall of the confinement device. Then, in the main part of this work we investigate the stability of such steady states, both with respect to perturbations of the initial data, where we employ the energy-Casimir method, and also with respect to perturbations of the external magnetic field.

Keywords: magnetic confinement, nonlinear partial differential equations, stationary solutions, Vlasov–Poisson equation, energy-Casimir method.

Mathematics Subject Classification: 35B35; 35Q83; 82D10.

1 Introduction

The investigation of a high-temperature plasma that is influenced by an exterior magnetic field is one of the most important aspects in the research on thermonuclear fusion. In this context, the *magnetic confinement* of plasmas is of particular interest. The idea of magnetic confinement is to apply magnetic fields to influence the plasma in such a way that it keeps a certain distance to the reactor wall. This is necessary because a plasma colliding with the wall will cool down (which interrupts the fusion process) and might even damage the

^{*}Department of Mathematics, University of Regensburg, 93053 Regensburg, Germany
(patrik.knopf@ur.de)

[†]Centre of Mathematical Sciences, Lund University, 221 00 Lund, Sweden
(jorg.weber@math.lu.se)

reactor. The most common reactor types for magnetic confinement are toroidal devices, for instance, Tokamaks, Stellarators and reversed field pinches. However, there are also linear (usually cylindrical) confinement devices, such as z -pinch or θ -pinch configurations. Their advantage is that they can be described more easily by mathematical models. For more details on thermonuclear fusion from a physics point of view we refer to [38].

In this paper, we intend to study the behavior of a plasma in a long cylindrical reactor. As a simplification for the mathematical analysis, we assume that the length of this cylinder is infinite. Moreover, we suppose that the volume charge density of the plasma is constant along lines parallel to the cylinder axis.

There are different possibilities of describing the time evolution of plasmas by means of mathematical equations. A very popular ansatz is to regard the plasma as an electrically conducting fluid. Its behavior is then described by the *magneto-hydrodynamic (MHD) equations*, which are a combination of the Navier–Stokes equations of fluid dynamics and Maxwell’s equations of electro-magnetism.

Another way to describe the motion of a plasma are kinetic equations. These models are particularly suitable if collisions among the ions and electrons can be neglected. It has been observed that such collisions only play a minor role in very hot and sufficiently diluted plasmas. The motion of a plasma can then be described by a *Vlasov equation* which will be the model of choice in our work.

To describe the time evolution of a two-component plasma (consisting of both electrons and positively charged ions) in a cylinder of infinite length, we consider the following Vlasov–Poisson system that is equipped with an external magnetic field $B = \text{curl}_x A$:

$$\partial_t f^\pm + p \cdot \partial_x f^\pm \pm (-\partial_x U + p \times B) \cdot \partial_p f^\pm = 0 \quad \text{on } \mathbb{R}_0^+ \times \mathbb{R}^2 \times \mathbb{R}^3, \quad (1.1a)$$

$$-\Delta_x U = 4\pi\rho, \quad \text{on } \mathbb{R}_0^+ \times \mathbb{R}^2, \quad (1.1b)$$

$$\lim_{|x| \rightarrow \infty} (U(t, x) + 2M \ln |x|) = 0 \quad \text{on } \mathbb{R}_0^+, \quad (1.1c)$$

$$\rho^\pm = \int_{\mathbb{R}^3} f^\pm(\cdot, \cdot, p) dp, \quad \rho = \rho^+ - \rho^- \quad \text{on } \mathbb{R}_0^+ \times \mathbb{R}^2, \quad (1.1d)$$

$$B = \text{curl}_x A \quad \text{on } \mathbb{R}_0^+ \times \mathbb{R}^2, \quad (1.1e)$$

$$f^\pm(0) = \mathring{f}^\pm \quad \text{on } \mathbb{R}^2 \times \mathbb{R}^3. \quad (1.1f)$$

Here, the plasma is represented by the electron density $f^- = f^-(t, x, p) \geq 0$ and the ion density $f^+ = f^+(t, x, p) \geq 0$, which both depend on time $t \in \mathbb{R}_0^+ := [0, \infty[$, position $x \in \mathbb{R}^2$ and velocity $p \in \mathbb{R}^3$. In the standard three-dimensional Vlasov–Poisson system, the position coordinate x is also a vector in \mathbb{R}^3 . However, as we consider the plasma in a cylinder of infinite length, we assume that the density function is constant along the direction of the cylinder axis, which in turn is supposed to point in the x_3 direction. This means that the distribution function f does not depend on the x_3 variable and it thus suffices to consider $x = (x_1, x_2) \in \mathbb{R}^2$. Since f may still depend on velocity vectors in \mathbb{R}^3 , the system (1.1) is referred to as the *two and one-half dimensional Vlasov–Poisson system*. The initial condition for f is given in (1.1f). In the following, we write $z = x_3$ to denote the third spatial component and thus, the x_3 -axis will also be referred to as the z -axis.

The scalar function U denotes the electrostatic potential which is induced by the charge of the particles. We point out that in the Vlasov–Poisson system, electromagnetic effects such as the induction of an internal magnetic field are neglected. Hence, the field equation for U is the Poisson equation (1.1b) with the charge density $\rho = \rho^+ - \rho^-$, given by (1.1d), as its right-hand side. The internal self-consistent electric field, which drives the movement of the charged particles, is then given as $E = -\partial_x U$. In order to determine U uniquely for given ρ , we impose the boundary condition (1.1c) at spatial infinity. Here, $M = \int_{\mathbb{R}^2} \rho \, dx$ is the total charge in any cross section of the cylinder. Note that M is a conserved quantity, i.e., it does not depend on time. We point out that in our scenario, the electric potential U does not vanish at spatial infinity as it would be the case in a fully three-dimensional setting. This is because the fundamental solution $-\frac{1}{2\pi} \ln |\cdot|$ of Poisson’s equation in $2D$ does not vanish at infinity. For a more detailed discussion on this issue, we refer to [22]. Throughout this paper we use modified Gaussian units such that all physical constants (mass and modulus of the charge of a particle) are normalized to unity.

In [22], confined steady states for the two-dimensional analogue of (1.1) (meaning that $x, p \in \mathbb{R}^2$) were constructed. The same has been done in [39] for the two and one-half dimensional relativistic Vlasov–Maxwell system. In both papers an additional symmetry, namely rotational symmetry about the z -axis, is assumed, and the ansatz

$$f_0^\pm = \eta^\pm(\mathcal{E}^\pm, \mathcal{F}^\pm, \mathcal{G}^\pm)$$

for a steady state is made, where \mathcal{E}^\pm , \mathcal{F}^\pm , and \mathcal{G}^\pm are invariants of the characteristic flow. For the steady states in our setting, the particle energy $\mathcal{E}^\pm = \frac{1}{2}|p|^2 \pm U$, which is (for the variables $(x, p \pm A)$) the Hamiltonian governing the motion of the particles (ions or electrons, respectively), is the invariant corresponding to time symmetry. Moreover, from the Lagrangian

$$\mathcal{L}^\pm = -\sqrt{1 - |(\dot{x}, \dot{z})|^2} \mp U \pm (\dot{x}, \dot{z}) \cdot A \quad \text{on } \mathbb{R}^3 \times \mathbb{R}^3,$$

we derive the invariant quantities corresponding to the spatial symmetries: the canonical angular momentum $\mathcal{F}^\pm = \partial_{\dot{\varphi}} \mathcal{L}^\pm = r(p_\varphi \pm A_\varphi)$ corresponds to rotational invariance, and $\mathcal{G}^\pm = \partial_{\dot{z}} \mathcal{L}^\pm = p_3 \pm A_3$, that is the third component of the canonical momentum, corresponds to translational invariance. Here and throughout this paper, we frequently use cylindrical coordinates (r, φ, z) on \mathbb{R}^3 . In this context, v_r , v_φ , and v_3 denote the coordinates of a vector v in the corresponding local coordinate system.

In this paper we study the stability properties of steady states of the separated form

$$f_0^\pm = \vartheta^\pm(\mathcal{E}^\pm) \psi^\pm(\mathcal{F}^\pm, \mathcal{G}^\pm)$$

with respect to both perturbations of the initial data and perturbations of the external magnetic field. As for stability with respect to perturbations of the initial data, it is well-known in the study of Vlasov type systems (both in the gravitational and plasma physics case) that a certain monotonicity assumption on ϑ^\pm is important; see (S4).

Comparison to related results in the literature. The mathematical investigation of stationary solutions in kinetic theory and their stability has an extensive history. A detailed study of steady states of the three-dimensional Vlasov–Poisson system, both in the gravitational case (without any external field) and in the plasma physics case (where no

confinement properties are taken into account), can be found in [35] and the large amount of references therein. In particular, we refer to [18, 25, 31–34]. Within this family of works, we especially want to point out [6], which is closest to our strategy. For the mathematical investigation of steady states in similar Vlasov type models, we refer to [2, 14–17] to mention but a few.

In order to investigate steady states of plasmas described by Vlasov type models, the system has to be equipped with a suitable external field. Besides the aforementioned papers [22] and [39], confined steady states in various settings under the influence of an external magnetic field were constructed in [3, 4, 37]. In the non-stationary case, an external magnetic field becoming infinite at the boundary of the confinement device was used in [7–10] to confine a Vlasov–Poisson plasma. More information on the confinement problem in various configurations with different Vlasov type plasma models (in some cases addressing linear stability), can be found in [12, 19, 28, 29, 40, 41]. We further want to mention the paper [5] dealing with instability of z -pinches in an MHD model.

Structure of this paper. After stating some important auxiliary results in Section 2, we discuss the global-in-time classical well-posedness of system (1.1) in Section 3. Here, we employ some of the strategies that have already been used in the literature (see [1, 23, 27, 30, 36] for results on local and global well-posedness) to tackle the three-dimensional Vlasov–Poisson system. In Section 4, we briefly outline the construction of confined steady states in our setting proceeding similarly to [22, 39]. The main novelties of the present paper are the stability results for such confined steady states, which are presented in Section 5. Restricting ourselves to a θ -pinch configuration (i.e., the external magnetic field points in the direction of the cylinder axis), we first prove stability of a previously constructed confined steady state with respect to perturbations of the initial data (see Theorem 5.1). This is done by suitably adapting the strategy of [6]. Furthermore, for perturbations of the external magnetic field, we also obtain a stability estimate (see Theorem 5.3). However, this estimate has the drawback of not being uniform in time. Eventually both results can be combined to obtain a stability estimate with respect to perturbations of both the initial data and the external magnetic field (see Theorem 5.4).

2 Preliminaries

First, we recall the following classical result on the Newtonian potential in two dimensions (see, e.g., [26, Theorem 10.2 and Theorem 10.3]).

Lemma 2.1. *Let $\varrho \in C_c^1(\mathbb{R}^2)$ be arbitrary. We set $M_\varrho := \int_{\mathbb{R}^2} \varrho \, dx$.*

Then the function

$$U_\varrho : \mathbb{R}^2 \rightarrow \mathbb{R}, \quad U_\varrho(x) := -2 \int_{\mathbb{R}^2} \varrho(y) \ln |x - y| \, dy \quad (2.1)$$

belongs to $C^2(\mathbb{R}^2)$ and is the unique classical solution of Poisson’s equation

$$\begin{aligned} -\Delta_x U_\varrho &= 4\pi\varrho \quad \text{on } \mathbb{R}^2, \\ \lim_{|x| \rightarrow \infty} (U_\varrho(t, x) + 2M_\varrho \ln |x|) &= 0. \end{aligned}$$

The gradient of U_ϱ is given as

$$\nabla U_\varrho(x) = -2 \int_{\mathbb{R}^2} \varrho(y) \frac{x-y}{|x-y|^2} dy, \quad x \in \mathbb{R}^2.$$

In particular, this entails that $U_\varrho(x) = \mathcal{O}(|x|^{-1})$ as $|x| \rightarrow \infty$. Moreover, we obtain the estimates

$$\|\nabla U_\varrho\|_\infty \leq c_1 \|\varrho\|_q^{q/2} \|\varrho\|_\infty^{1-q/2} \quad \text{for all } q \in [1, 2[, \quad (2.2)$$

$$\|D^2 U_\varrho\|_\infty \leq c_2 \left[(1 + \|\varrho\|_\infty) \left(1 + \ln_+ (\|\nabla \varrho\|_\infty) \right) + \|\varrho\|_1 \right] \quad (2.3)$$

with positive constants $c_1 = c_1(q)$ and c_2 that do not depend on ϱ .

Next, we state two important properties of the potential energy: an upper bound is provided by the logarithmic Hardy–Littlewood–Sobolev inequality (see [11, Theorem 1]), and positive definiteness under the assumption that the total charge is zero is ensured by [24, Theorem 1.16].

Lemma 2.2. *Suppose that $\rho \in L^1(\mathbb{R}^2)$.*

(a) *If ρ is non-negative almost everywhere on \mathbb{R}^2 , and the integral*

$$\int_{\mathbb{R}^2} \rho(x) \ln(1 + |x|^2) dx$$

is finite, then there exists a constant $C > 0$ independent of ρ such that

$$- \int_{\mathbb{R}^2} \int_{\mathbb{R}^2} \ln|x-y| \rho(y) \rho(x) dy dx \leq \frac{1}{2} \|\rho\|_1 \int_{\mathbb{R}^2} \rho \ln \frac{\rho}{\|\rho\|_1} dx + C \|\rho\|_1^2.$$

(b) *If $\int_{\mathbb{R}^2} \rho dx = 0$, it holds that*

$$\int_{\mathbb{R}^2} \int_{\mathbb{R}^2} (-\ln|x-y|) \rho(y) \rho(x) dy dx \geq 0,$$

provided that this integral exists.

3 Global classical well-posedness

We first want to establish the existence of a unique global classical solution $f = (f^+, f^-)$ with $f^\pm \in C^1([0, \infty[; \mathbb{R}^5)$ to the system (1.1).

To this end, we first recall that for any functions $F, G \in C_b([0, \infty[; C_b^1(\mathbb{R}^2))$, f is a classical solution to the equations

$$\partial_t f^\pm + p \cdot \partial_x f^\pm \pm (F + p \times G) \cdot \partial_p f^\pm = 0 \quad \text{on } \mathbb{R}_0^+ \times \mathbb{R}^5 \quad (3.1)$$

if and only if the functions f^\pm are constant along solutions of the characteristic systems

$$\dot{x} = p, \quad \dot{p} = \pm (F(\cdot, x) + p \times G(\cdot, x)). \quad (3.2)$$

This equivalence is a direct consequence of [21, Lemma 4] and [21, Lemma 5] (which are stated in the three-dimensional setting but can easily be transferred to the two-and-one-half dimensional case). We directly infer that any classical solution f to (3.1) can be expressed as

$$f^\pm(t, z) = \mathring{f}^\pm(Z^\pm(0, t, z)) \quad \text{for all } t \geq 0, z = (x, p) \in \mathbb{R}^5,$$

where $Z^\pm(\cdot, t, z) = (X^\pm, P^\pm)(\cdot, t, z)$ denotes the unique solution of (3.2) satisfying the condition $Z^\pm(t, t, z) = z$ for any given $z = (x, p) \in \mathbb{R}^5$. One can easily see that the characteristic flow is measure preserving, meaning that

$$\det \left(\frac{\partial Z}{\partial z} \right) \equiv 1.$$

For more details, we refer to [21, Lemma 4]. The above results directly imply that any classical solution $f = (f^+, f^-)$ to (3.1) satisfies

$$\text{supp } f^\pm(t) = Z^\pm(t, 0, \text{supp } \mathring{f}^\pm) \quad \text{and} \quad \|f^\pm(t)\|_{L^q(\mathbb{R}^5)} = \|\mathring{f}^\pm\|_{L^q(\mathbb{R}^5)}$$

for all $q \in [1, \infty]$ and all $t \geq 0$ for which the solution exists (cf. [21, Lemma 5]).

For any classical solution f to the system (1.1), we further define the quantities

$$\begin{aligned} \mathbb{P}^\pm(t) &:= \sup \{ |P^\pm(s, 0, z)| \mid z \in \text{supp } \mathring{f}^\pm, 0 \leq s \leq t \} \\ &= \sup \{ |p| \mid \exists s \in [0, t] : (x, p) \in \text{supp } f^\pm(s) \}, \\ \mathbb{X}^\pm(t) &:= \sup \{ |X^\pm(s, 0, z)| \mid z \in \text{supp } \mathring{f}^\pm, 0 \leq s \leq t \} \\ &= \sup \{ |x| \mid \exists s \in [0, t] : (x, p) \in \text{supp } f^\pm(s) \}, \\ \mathbb{P}(t) &:= \max_{\pm} \mathbb{P}^\pm(t), \quad \mathbb{X}(t) := \max_{\pm} \mathbb{X}^\pm(t), \end{aligned}$$

for all $t \geq 0$ at which the solution exists. Here, (X^\pm, P^\pm) denotes the solutions of the characteristic systems (3.2) for the choices $G = B$ and $F = -\partial_x U$, where U is the self-consistent electric potential satisfying (1.1b)–(1.1d) and B is given by (1.1e). We point out that the functions \mathbb{P}^\pm , \mathbb{X}^\pm , \mathbb{P} and \mathbb{X} are non-decreasing but we cannot a priori exclude the possibility that they become infinite already in finite time. However, to show that the classical solutions exist globally in time, it will be crucial to show that $\mathbb{P}(t)$ actually remains bounded on finite time intervals.

Furthermore, we consider the total energy

$$\mathcal{H}(f(t)) = E_{\text{kin}}(f(t)) + E_{\text{pot}}(f(t)),$$

where the kinetic energy E_{kin} and the potential energy E_{pot} are defined as

$$\begin{aligned} E_{\text{kin}}(f(t)) &:= \frac{1}{2} \sum_{\pm} \int_{\mathbb{R}^2} \int_{\mathbb{R}^3} |p|^2 f^\pm(t, x, p) \, dp \, dx, \\ E_{\text{pot}}(f(t)) &:= \frac{1}{2} \int_{\mathbb{R}^2} U(t, x) \rho(t, x) \, dx. \end{aligned}$$

It is straightforward to check that the total energy $\mathcal{H}(f(t))$ is constant in time, meaning that $\mathcal{H}(f(t)) = \mathcal{H}(\mathring{f})$, as long as the solution exists.

The global well-posedness result reads as follows:

Theorem 3.1. Let $\mathring{f} = (\mathring{f}^+, \mathring{f}^-)$ with $\mathring{f}^\pm \in C_c^1(\mathbb{R}^5; \mathbb{R}_0^+)$ be arbitrary initial data, and let $A \in C_b([0, \infty[; C_b^2(\mathbb{R}^2; \mathbb{R}^3))$ be any given external magnetic vector potential.

Then there exists a unique classical solution $f = (f^+, f^-)$ with $f^\pm \in C^1([0, \infty[\times \mathbb{R}^5)$ such that the supports $\text{supp } f^\pm \subset \mathbb{R}^5$ remain compact for all $t \geq 0$.

Moreover, for any $\varepsilon > 0$, there exists a constant $c > 0$ depending only on ε , $\mathcal{H}(\mathring{f})$, $\|\mathring{f}^\pm\|_1$ and $\|\mathring{f}^\pm\|_\infty$ such that

$$\mathbb{P}(t) \leq c(1+t)^{2+\varepsilon} \quad \text{for all } t \geq 0. \quad (3.3)$$

Proof. *Step 1: Local well-posedness.* For the standard three-dimensional Vlasov–Poisson system (i.e., without any external magnetic field), the existence and uniqueness of a local classical solution was first established by R. Kurth [23]. As the external magnetic field $B = \text{curl}_x A \in C_b([0, \infty[; C_b^1(\mathbb{R}^2; \mathbb{R}^3))$ is sufficiently regular, and the electric potential can be bounded by means of the estimates (2.2) and (2.3), we can proceed completely analogously to Kurth’s proof to establish the existence of a local classical solution. A sketch of the proof for the corresponding local existence result in three dimensions (with external magnetic field) can be found in [21, Theorem 1]. In this way, we obtain a unique local classical solution $f = (f^+, f^-)$ to the system (1.1) existing on a right-maximal time interval $[0, T_*[$.

Step 2: Continuation criterion. For the local classical solution of the standard three-dimensional Vlasov–Poisson system (i.e., $B \equiv 0$), a continuation criterion was established by J. Batt [1]. It states that the local solution can be extended in time as long as the velocity support $\mathbb{P}(t)$ remains under control. Proceeding analogously, we conclude that for the maximal local classical solution $f = (f^+, f^-)$ constructed in Step 1, it either holds that

$$\lim_{\substack{t \rightarrow T_* \\ 0 \leq t < T_*}} \mathbb{P}(t) = \infty \quad (3.4)$$

or $T_* = \infty$. Hence, in order to show that our unique maximal solution actually exists globally in time, we must ensure that $\mathbb{P}(t)$ remains bounded on finite time intervals as long as the solution exists.

Step 3: Extension on $[0, \infty[$. Based on Batt’s continuation criterion, two different proofs for global existence of classical solutions for the standard three-dimensional Vlasov–Poisson system (i.e., $B \equiv 0$) were virtually obtained simultaneously: one of them by P.L. Lions and B. Perthame [27], and the other one by K. Pfaffelmoser [30]. Pfaffelmoser’s proof was later greatly simplified by J. Schaeffer [36]. We point out that the Pfaffelmoser–Schaeffer proof was further generalized in [21, Theorem 1] to handle the three-dimensional Vlasov–Poisson system endowed with an external magnetic field.

Although our very basic strategy is the same, we will see that for our two-and-one-half dimensional system, the proof of global existence differs greatly from the three-dimensional case. Fortunately, we have much better estimates for the electric field $-\partial_x U$ (see Lemma 2.1) than it would be the case in three dimensions. This means that the main difficulties of the proofs by Lions and Perthame or Pfaffelmoser and Schaeffer do not occur in our situation. Therefore, our proof will be much more straightforward. However, we will face different problems as the two-dimensional electric potential is given by a logarithmic convolution kernel and thus, it is non-trivial to establish that the potential energy is bounded from below.

We argue by contradiction and assume that $T_* < \infty$. Hence, the goal is to show that $\mathbb{P}(t)$ remains bounded for all $t \in [0, T_*[$. This would be a contradiction to (3.4) and we thus conclude that actually $T_* = \infty$. To this end, let $t \in [0, T_*[$ be arbitrary. In the following, the letter C will denote generic positive constants that depend only $\|\mathring{f}^\pm\|_1$ and $\|\mathring{f}^\pm\|_\infty$, and may change their value from line to line.

It is crucial that an external magnetic field does not change the modulus of a particle's velocity, but only its direction. For any $s \in [0, t]$, we thus obtain the estimate

$$\frac{1}{2}|P^\pm(s)|^2 \leq \frac{1}{2}|P^\pm(0)|^2 + \int_0^s |\partial_x U(\tau, X(\tau))| |P^\pm(\tau)| \, d\tau$$

for all $s \in [0, t]$. Invoking a quadratic version of Gronwall's lemma (see [13, Theorem 5]), and taking the maximum of the components \pm , we conclude that

$$\mathbb{P}(s) \leq \mathbb{P}(0) + \int_0^s \|\partial_x U(\tau)\|_\infty \, d\tau \quad (3.5)$$

for all $s \in [0, t]$. Hence, the next goal is to establish a suitable bound for the $L^\infty(\mathbb{R}^2)$ -norm of $\partial_x U(\tau)$. Therefore, we recall that the total energy $\mathcal{H}(f(\tau))$ is conserved, meaning that $\mathcal{H}(f(\tau)) = \mathcal{H}(\mathring{f})$ for all $\tau \in [0, T_*[$. Using the convolution representation of U (cf. (2.1)), we obtain

$$E_{\text{kin}}(f(\tau)) = \mathcal{H}(\mathring{f}) + \int_{\mathbb{R}^2} \int_{\mathbb{R}^2} \ln|x-y| \rho(\tau, y) \rho(\tau, x) \, dy \, dx \quad (3.6)$$

for all $\tau \in [0, T_*[$. Unfortunately, as the total charge is generally not zero, we cannot directly use Lemma 2.2 to bound second summand on the right-hand side, as it has the “wrong” sign. We thus expand this term as follows:

$$\begin{aligned} & \int_{\mathbb{R}^2} \int_{\mathbb{R}^2} \ln|x-y| \rho(\tau, y) \rho(\tau, x) \, dy \, dx \\ &= \int_{\mathbb{R}^2} \int_{\mathbb{R}^2} \ln|x-y| \rho^+(\tau, y) \rho^+(\tau, x) \, dy \, dx \\ & \quad + \int_{\mathbb{R}^2} \int_{\mathbb{R}^2} \ln|x-y| \rho^-(\tau, y) \rho^-(\tau, x) \, dy \, dx \\ & \quad - 2 \int_{\mathbb{R}^2} \int_{\mathbb{R}^2} \ln|x-y| \rho^+(\tau, y) \rho^-(\tau, x) \, dy \, dx \end{aligned}$$

for all $\tau \in [0, T_*[$. Let now $\ln_+ \geq 0$ and $\ln_- \geq 0$ denote the positive and the negative part of the logarithm \ln , meaning that $\ln = \ln_+ - \ln_-$. Moreover, we define the function $\bar{\rho} := \max\{\rho^+, \rho^-\}$. We thus obtain the estimate

$$\begin{aligned} & \int_{\mathbb{R}^2} \int_{\mathbb{R}^2} \ln|x-y| \rho(\tau, y) \rho(\tau, x) \, dy \, dx \\ & \leq \int_{\mathbb{R}^2} \int_{\mathbb{R}^2} \ln_+ |x-y| \rho^+(\tau, y) \rho^+(\tau, x) \, dy \, dx \\ & \quad + \int_{\mathbb{R}^2} \int_{\mathbb{R}^2} \ln_+ |x-y| \rho^-(\tau, y) \rho^-(\tau, x) \, dy \, dx \end{aligned}$$

$$+ 2 \int_{\mathbb{R}^2} \int_{\mathbb{R}^2} \ln_- |x - y| \bar{\rho}(\tau, y) \bar{\rho}(\tau, x) \, dy \, dx \quad (3.7)$$

for all $\tau \in [0, T_*[$. We next recall that $\rho^\pm(\tau, x) = 0$ for all $x \in \mathbb{R}^2$ with $|x| > \mathbb{X}(\tau)$. Hence, the first two summands on the right-hand side can be bounded in the following way:

$$\begin{aligned} & \int_{\mathbb{R}^2} \int_{\mathbb{R}^2} \ln_+ |x - y| \rho^\pm(\tau, y) \rho^\pm(\tau, x) \, dy \, dx \\ & \leq \ln(1 + 2\mathbb{X}(\tau)) \|\rho^\pm(\tau)\|_1^2 = \ln(1 + 2\mathbb{X}(\tau)) \|\mathring{f}^\pm\|_1^2 \end{aligned} \quad (3.8)$$

for all $\tau \in [0, T_*[$. To bound the third summand on the right-hand side of (3.7), we recall that $\ln_- |x - y| = -\ln |x - y|$ if $|x - y| < 1$, and $\ln_- |x - y| = 0$ if $|x - y| \geq 1$.

$$\begin{aligned} & \int_{\mathbb{R}^2} \int_{\mathbb{R}^2} \ln_- |x - y| \bar{\rho}(\tau, y) \bar{\rho}(\tau, x) \, dy \, dx \\ & = - \iint_{\{|x-y|<1\}} \ln |x - y| \bar{\rho}(\tau, y) \bar{\rho}(\tau, x) \, dy \, dx \\ & = - \int_{\mathbb{R}^2} \int_{\mathbb{R}^2} \ln |x - y| \bar{\rho}(\tau, y) \bar{\rho}(\tau, x) \, dy \, dx \\ & \quad + \iint_{\{|x-y|\geq 1\}} \ln |x - y| \bar{\rho}(\tau, y) \bar{\rho}(\tau, x) \, dy \, dx \\ & \leq - \int_{\mathbb{R}^2} \int_{\mathbb{R}^2} \ln |x - y| \bar{\rho}(\tau, y) \bar{\rho}(\tau, x) \, dy \, dx \\ & \quad + \int_{\mathbb{R}^2} \int_{\mathbb{R}^2} \ln_+ |x - y| \bar{\rho}(\tau, y) \bar{\rho}(\tau, x) \, dy \, dx \end{aligned} \quad (3.9)$$

for all $\tau \in [0, T_*[$. Obviously, the second integral in the last line can be bounded by proceeding as in (3.8). The first integral in the last line of (3.9) now has the “correct” sign such that Lemma 2.2(a) can be applied. Without loss of generality, we assume that $\mathring{f}^+ \geq 0$ is nontrivial, and we thus have

$$0 < \|\mathring{f}^+\|_1 = \|\rho^+(\tau)\|_1 \leq \|\bar{\rho}(\tau)\|_1 \leq \|\rho^+(\tau)\|_1 + \|\rho^-(\tau)\|_1 \leq \|\mathring{f}^+\|_1 + \|\mathring{f}^-\|_1 \leq C.$$

By means of Lemma 2.2, we now obtain

$$\begin{aligned} & - \int_{\mathbb{R}^2} \int_{\mathbb{R}^2} \ln |x - y| \bar{\rho}(\tau, y) \bar{\rho}(\tau, x) \, dy \, dx \\ & \leq \frac{1}{2} \|\bar{\rho}(\tau)\|_1 \int_{\mathbb{R}^2} \bar{\rho}(\tau, x) \ln \left(\frac{\bar{\rho}(\tau, x)}{\|\bar{\rho}(\tau)\|_1} \right) \, dx + C \|\bar{\rho}(\tau)\|_1^2 \\ & \leq C + C \ln(1 + \|\rho^+(\tau)\|_\infty + \|\rho^-(\tau)\|_\infty) \end{aligned} \quad (3.10)$$

for all $\tau \in [0, T_*[$. Combining the estimates (3.7)–(3.10), we can estimate the right-hand side of (3.6). This gives

$$E_{\text{kin}}(f(\tau)) \leq C + C \ln(1 + 2\mathbb{X}(\tau)) + C \ln(1 + \|\rho^+(\tau)\|_\infty + \|\rho^-(\tau)\|_\infty) \quad (3.11)$$

for all $\tau \in [0, T_*[$. Since any solutions (X^\pm, P^\pm) of the characteristic systems (3.2) (written for $F = -\partial_x U$ and $G = B$) satisfy $\dot{X}^\pm = P^\pm$, it is straightforward to check that

$$\mathbb{X}(\tau) \leq \mathbb{X}(0) + \int_0^\tau \mathbb{P}(s) \, ds \leq C + \tau \mathbb{P}(\tau) \quad \text{for all } \tau \in [0, T_*[.$$

Furthermore, we have

$$\|\rho^\pm(\tau)\|_{L^\infty(\mathbb{R}^2)} \leq \frac{4\pi}{3} \|\mathring{f}^\pm\|_\infty \mathbb{P}(\tau)^3 \leq C \mathbb{P}(\tau)^3 \quad \text{for all } \tau \in [0, T_*[. \quad (3.12)$$

Let now $0 < \delta < \frac{1}{2}$ be arbitrary. In the following, let C_δ denote generic positive constants that depend only on δ , $\|\mathring{f}^\pm\|_1$ and $\|\mathring{f}^\pm\|_\infty$, and may change their value from line to line. In combination with (3.11), we conclude that

$$\begin{aligned} E_{\text{kin}}(f(\tau)) &\leq C + C \ln(C(1+\tau)(1+\mathbb{P}(\tau))) + C \ln(C(1+\mathbb{P}(\tau))^3) \\ &\leq C_\delta(1+\tau)^{2\delta}(1+\mathbb{P}(\tau))^{2\delta} \end{aligned} \quad (3.13)$$

for all $\tau \in [0, T_*[$. By means of an interpolation argument, we further obtain the estimate

$$\rho^\pm(\tau, x) \leq C \left(\int_{\mathbb{R}^3} |p|^2 f^\pm(\tau, x, p) \, dp \right)^{\frac{3}{5}}$$

for all $\tau \in [0, T_*[$ and $x \in \mathbb{R}^2$. Along with (3.13), this leads to the estimate

$$\begin{aligned} \|\rho(\tau)\|_{L^{5/3}(\mathbb{R}^2)}^{5/3} &= \int_{\mathbb{R}^2} |\rho(\tau, x)|^{5/3} \, dx \leq C E_{\text{kin}}(f(\tau)) \\ &\leq C_\delta(1+\tau)^{2\delta}(1+\mathbb{P}(\tau))^{2\delta} \end{aligned} \quad (3.14)$$

for all $\tau \in [0, T_*[$. Invoking the estimate (2.2) from Lemma 2.1, as well as the estimates (3.12) and (3.14), we thus get

$$\begin{aligned} \|\partial_x U(\tau)\|_{L^\infty(\mathbb{R}^2)} &\leq C \|\rho(\tau)\|_{L^{5/3}(\mathbb{R}^2)}^{5/6} \|\rho(\tau)\|_{L^\infty(\mathbb{R}^2)}^{1/6} \\ &\leq C_\delta(1+\tau)^\delta (1+\mathbb{P}(\tau))^{\delta+1/2} \end{aligned}$$

for all $\tau \in [0, T_*[$. Using this estimate to bound the right-hand side of (3.5), we obtain

$$(1 + \mathbb{P}(s)) \leq (1 + \mathbb{P}(0)) + \int_0^s C_\delta(1+\tau)^\delta (1 + \mathbb{P}(\tau))^{\delta+1/2} \, d\tau$$

for all $s \in [0, t]$. Applying a nonlinear generalization of Gronwall's lemma (see [13, Theorem 21] with $\alpha = \frac{1}{2} + \delta$), and recalling that $\delta < \frac{1}{2}$, we eventually conclude that

$$1 + \mathbb{P}(t) \leq \left[(1 + \mathbb{P}(0))^{\frac{1}{2}-\delta} + C_\delta \left(\frac{1}{2} - \delta \right) \int_0^t (1+s)^\delta \, ds \right]^{\frac{2}{1-2\delta}}.$$

Since $t \in [0, T_*[$ was arbitrary, this yields

$$\mathbb{P}(t) \leq C_\delta(1+t)^{\frac{2\delta+2}{1-2\delta}} = C_\delta(1+t)^{2+\frac{6\delta}{1-2\delta}} \quad \text{for all } t \in [0, T_*[. \quad (3.15)$$

This contradicts (3.4) and thus, it actually holds that $T_* = \infty$. For any arbitrary $\varepsilon > 0$, we choose

$$\delta := \frac{\varepsilon}{6 + 2\varepsilon}, \quad \text{i.e.,} \quad \varepsilon = \frac{6\delta}{1 - 2\delta},$$

and thus, the estimate (3.3) follows directly from (3.15). Hence, all assertions are established and the proof is complete. \square

In addition, we can show that the classical solution depends continuously on the external magnetic field, respectively. This is established by the following theorem.

Theorem 3.2. *Let $\mathring{f} = (\mathring{f}^+, \mathring{f}^-)$ with $\mathring{f}^\pm \in C_c^1(\mathbb{R}^5; \mathbb{R}_0^+)$ be any initial datum, let furthermore $A^1, A^2 \in C_b([0, \infty[; C_b^2(\mathbb{R}^2; \mathbb{R}^3))$ be any given external magnetic vector potentials, and let B^1 and B^2 denote the corresponding magnetic fields. Moreover, for $i \in \{1, 2\}$, let $f_i = (f_i^+, f_i^-)$ denote the global-in-time classical solution of the system (1.1) to the initial data f_0 corresponding to the magnetic vector potential A^i .*

Then, for any $\gamma > 4$, there exists a real number $q \in (2, \infty)$ depending only on γ as well as a constant $c > 0$ depending only on $\mathcal{H}(\mathring{f})$, $\|\mathring{f}^\pm\|_1$, $\|\mathring{f}^\pm\|_\infty$ and γ such that for all $t \geq 0$,

$$\sum_{\pm} \|f_1^\pm(t) - f_2^\pm(t)\|_2 \leq a_{f_2}(t) \exp(b_{f_2}(t) c(1+t)^\gamma) \|B^1 - B^2\|_{L^1(0,t; L^2(\mathbb{R}^2))} \quad (3.16)$$

where

$$\begin{aligned} a_{f_2}(t) &:= 2 \max_{\pm} \max_{s \in [0, t]} \|p \times \partial_p f_2^\pm(s)\|_{L^\infty(\mathbb{R}^2; L^2(\mathbb{R}^3))}, \\ b_{f_2}(t) &:= 2 \max_{\pm} \max_{s \in [0, t]} \|\partial_p f_2^\pm(s)\|_{L^q(\mathbb{R}^2; L^2(\mathbb{R}^3))}. \end{aligned}$$

Proof. For brevity, we write

$$\begin{aligned} \bar{f}^\pm &:= f_1^\pm - f_2^\pm, \quad \bar{U} := U_1 - U_2, \quad \bar{B} := B^1 - B^2, \\ \bar{\rho}^\pm &:= \rho_1^\pm - \rho_2^\pm, \quad \bar{\rho} := \rho_1 - \rho_2 = \bar{\rho}^+ - \bar{\rho}^-. \end{aligned}$$

Subtracting the two equations

$$\begin{aligned} \partial_t f_1^\pm + p \cdot \partial_x f_1^\pm \pm (-\partial_x U_1 + p \times B^1) \cdot \partial_p f_1^\pm &= 0, \\ \partial_t f_2^\pm + p \cdot \partial_x f_2^\pm \pm (-\partial_x U_2 + p \times B^2) \cdot \partial_p f_2^\pm &= 0, \end{aligned}$$

we obtain

$$\partial_t \bar{f}^\pm + p \cdot \partial_x \bar{f}^\pm \pm (-\partial_x U_1 + p \times B^1) \cdot \partial_p \bar{f}^\pm = \mp (-\partial_x \bar{U} + p \times \bar{B}) \cdot \partial_p f_2^\pm. \quad (3.17)$$

To estimate the L^2 -norm of \bar{f}^\pm , we multiply (3.17) by \bar{f}^\pm and integrate with respect to (x, p) . This yields

$$\frac{1}{2} \frac{d}{dt} \|\bar{f}^\pm(t)\|_2^2 = -\frac{1}{2} \int_{\mathbb{R}^2} \int_{\mathbb{R}^3} \left(\operatorname{div}_x (p(\bar{f}^\pm)^2) \pm \operatorname{div}_p ((-\partial_x U_1 + p \times B^1)(\bar{f}^\pm)^2) \right) dp dx$$

$$\begin{aligned}
& \mp \int_{\mathbb{R}^2} \int_{\mathbb{R}^3} \bar{f}^\pm(-\partial_x \bar{U} + p \times \bar{B}) \cdot \partial_p f_2^\pm \, dp \, dx \\
&= \mp \int_{\mathbb{R}^2} \int_{\mathbb{R}^3} \bar{f}^\pm(-\partial_x \bar{U} + p \times \bar{B}) \cdot \partial_p f_2^\pm \, dp \, dx
\end{aligned}$$

for all $t \geq 0$. In the following, let $\gamma > 4$ be arbitrary, and let $c > 0$ denote a generic constant that may depend on $\mathcal{H}(\bar{f})$, $\|\bar{f}^\pm\|_1$, $\|\bar{f}^\pm\|_\infty$ and γ and may change its value from line to line. We further choose any real numbers $r \in (2, \infty)$, $q \in (1, 2)$ and $\varepsilon > 0$ (all depending on γ) such that

$$\frac{1}{r} + \frac{1}{q} = \frac{1}{2} \quad \text{and} \quad \frac{6+2\varepsilon}{r} + \frac{3\varepsilon}{2} + 4 < \gamma. \quad (3.18)$$

Since $\bar{f}^\pm(0) = 0$, we have

$$\begin{aligned}
\frac{1}{2} \|\bar{f}^\pm(t)\|_2^2 &= \mp \int_0^t \int_{\mathbb{R}^2} \int_{\mathbb{R}^3} \bar{f}^\pm(-\partial_x \bar{U} + p \times \bar{B}) \cdot \partial_p f_2^\pm \, dp \, dx \, ds \\
&\leq \int_0^t \frac{1}{2} (b_{f_2}(t) \|\partial_x \bar{U}(s)\|_r + a_{f_2}(t) \|\bar{B}(s)\|_2) \|\bar{f}^\pm(s)\|_2 \, ds
\end{aligned}$$

for all $t \geq 0$. By the quadratic Gronwall lemma (see [13, Theorem 5]), it follows that

$$\|\bar{f}^\pm(t)\|_2 \leq \frac{1}{2} b_{f_2}(t) \int_0^t \|\partial_x \bar{U}(s)\|_r \, ds + \frac{1}{2} a_{f_2}(t) \int_0^t \|\bar{B}(s)\|_2 \, ds. \quad (3.19)$$

We now have a closer look at the term $\|\partial_x \bar{U}(t)\|_r$. Using the Hardy–Littlewood–Sobolev inequality, we obtain the estimate

$$\|\partial_x \bar{U}(t)\|_r \leq c \|\bar{\rho}(t)\|_{r_*} \quad \text{for all } t \geq 0, \quad (3.20)$$

where $r_* \in (1, 2)$ is chosen such that $\frac{1}{r_*} = \frac{1}{2} + \frac{1}{r}$. For any $t \geq 0$, we now write

$$\begin{aligned}
\bar{\mathbb{P}}^\pm(t) &:= \sup \{ |p| \mid \exists s \in [0, t] : (x, p) \in \text{supp } \bar{f}^\pm(s) \}, \\
\bar{\mathbb{X}}^\pm(t) &:= \sup \{ |x| \mid \exists s \in [0, t] : (x, p) \in \text{supp } \bar{f}^\pm(s) \}, \\
\bar{\mathbb{P}}(t) &:= \max_{\pm} \bar{\mathbb{P}}^\pm(t), \quad \bar{\mathbb{X}}(t) := \max_{\pm} \bar{\mathbb{X}}^\pm(t).
\end{aligned}$$

As a consequence of Theorem 3.1, it holds that

$$\bar{\mathbb{P}}(t) \leq c(1+t)^{2+\varepsilon} \quad \text{and} \quad \bar{\mathbb{X}}(t) \leq c(1+t)^{3+\varepsilon}$$

for all $t \geq 0$. Since

$$|\bar{\rho}(t, x)| = \left| \int_{\mathbb{R}^3} (\bar{f}^+ - \bar{f}^-)(t, x, p) \, dp \right| \leq \sqrt{4\pi/3} \bar{\mathbb{P}}(t)^{3/2} \left(\int_{\mathbb{R}^3} |(\bar{f}^+ - \bar{f}^-)(t, x, p)|^2 \, dp \right)^{1/2},$$

we thus infer that

$$\|\bar{\rho}(t)\|_{r_*} \leq \pi^{1/r} \bar{\mathbb{X}}(t)^{2/r} \|\bar{\rho}(t)\|_2 \leq \sqrt{8\pi/3} \pi^{1/r} \bar{\mathbb{X}}(t)^{2/r} \bar{\mathbb{P}}(t)^{3/2} \sum_{\pm} \|\bar{f}^\pm(t)\|_2$$

$$\leq c(1+t)^{\frac{6+2\varepsilon}{r}+\frac{3\varepsilon}{2}+3} \sum_{\pm} \|\bar{f}^{\pm}(t)\|_2$$

for all $t \geq 0$. In combination with (3.20) and (3.19), we get

$$\sum_{\pm} \|\bar{f}^{\pm}(t)\|_2 \leq a_{f_2}(t) \|\bar{B}\|_{L^1(0,t;L^2(\mathbb{R}^2))} + b_{f_2}(t) c \int_0^t (1+s)^{\frac{6+2\varepsilon}{r}+\frac{3\varepsilon}{2}+3} \sum_{\pm} \|\bar{f}^{\pm}(s)\|_2 \, ds,$$

and thus, Gronwall's lemma implies that

$$\sum_{\pm} \|\bar{f}^{\pm}(t)\|_2 \leq a_{f_2}(t) \|\bar{B}\|_{L^1(0,t;L^2(\mathbb{R}^2))} \exp \left(b_{f_2}(t) c (1+t)^{\frac{6+2\varepsilon}{r}+\frac{3\varepsilon}{2}+4} \right)$$

for all $t \geq 0$. Due to (3.18), this proves (3.16) and thus, the proof is complete. \square

4 Existence of confined steady states

We now show the existence of a confined steady state of the system (1.1) under the influence of an external magnetic field $B = \text{curl}_x A$. Such steady states are defined as follows:

Definition 4.1 (Steady states). *A pair of functions $f_0 = (f_0^+, f_0^-)$ is called a steady state of the Vlasov–Poisson system (1.1) if f_0^{\pm} , the induced electric potential U_0 , the external vector potential A^0 and its induced magnetic field B^0 have the regularity*

$$f_0^{\pm} \in C^1(\mathbb{R}^5; \mathbb{R}_0^+), \quad U_0 \in C^2(\mathbb{R}^2), \quad A^0 \in C^2(\mathbb{R}^2; \mathbb{R}^3), \quad B^0 \in C^1(\mathbb{R}^2; \mathbb{R}^3),$$

and satisfy the following equations:

$$p \cdot \partial_x f_0^{\pm} \pm (-\partial_x U_0 + p \times B^0) \cdot \partial_p f_0^{\pm} = 0 \quad \text{on } \mathbb{R}^2 \times \mathbb{R}^3, \quad (4.1a)$$

$$-\Delta_x U_0 = 4\pi \rho_0 \quad \text{on } \mathbb{R}^2, \quad (4.1b)$$

$$\lim_{|x| \rightarrow \infty} (U_0(x) + 2M \ln |x|) = 0, \quad (4.1c)$$

$$\rho_0^{\pm} = \int_{\mathbb{R}^3} f_0^{\pm}(\cdot, \cdot, p) \, dp, \quad \rho_0 = \rho_0^+ - \rho_0^- \quad \text{on } \mathbb{R}^2, \quad (4.1d)$$

$$\text{curl}_x A^0 = B^0 \quad \text{on } \mathbb{R}^2, \quad (4.1e)$$

where $M = \int_{\mathbb{R}^2} \rho_0 \, dx$. A steady state f_0^{\pm} is called confined in a cylinder with radius $R > 0$ if $f_0^{\pm}(x, p) = 0$ for all $|x| > R$, and it is called nontrivial if both f_0^{\pm} are not identically zero.

We are looking for steady states which are axially symmetric about the z -axis, meaning that

$$f_0^{\pm}(x, p) = f_0^{\pm}(Rx, R\bar{p}, p_3), \quad U_0(x) = U_0(Rx), \quad A^0(x) = RA^0(Rx), \quad B^0(x) = RB^0(Rx)$$

for all $x \in \mathbb{R}^2$, $p = (\bar{p}, p_3) \in \mathbb{R}^3$, and any rotation matrix $R \in \text{SO}(2)$. In the following, we will sometimes write

$$f_0^{\pm}(r, p) = f_0^{\pm}(x, p), \quad U_0(r) = U_0(x), \quad A^0(r) = A^0(x), \quad B^0(r) = B^0(x)$$

with some abuse of notation.

To construct a nontrivial, confined steady state, we make the general ansatz

$$f_0^\pm = \eta^\pm(\mathcal{E}^\pm, \mathcal{F}^\pm, \mathcal{G}^\pm)$$

where $\eta^\pm \in C^1(\mathbb{R}^3)$, and

$$\mathcal{E}^\pm(x, p) := \frac{1}{2}|p|^2 \pm U_0, \quad \mathcal{F}^\pm(x, p) := r(p_\varphi \pm A_\varphi^0), \quad \mathcal{G}^\pm(x, p) := p_3 \pm A_3^0$$

are invariants of the characteristic flow.

Furthermore, also due to the axial symmetry, we need some further assumptions on the external magnetic vector potential.

General assumption on A^0 .

(A0) The external magnetic vector potential $A^0 \in C^2(\mathbb{R}^2; \mathbb{R}^3)$ satisfies $A_r^0 = 0$ everywhere on \mathbb{R}^2 and $A_\varphi^0 = A_3^0 = 0$ on the z -axis.

To make this assumption plausible, we point out three facts: First, A_r^0 does not affect B and should satisfy $\frac{1}{r}\partial_r(rA_r^0) = 0$ in view of the gauge condition $\operatorname{div}_x A^0 = 0$, so that the choice $A_r^0 = 0$ is no loss of generality. Second, A_φ^0 necessarily has to vanish on the z -axis in order for $B^0 = \operatorname{curl}_x A^0$ to be well-defined on the z -axis in the classical sense. Third, since adding constants to A^0 does not affect B^0 , the choice $A_3^0 = 0$ on the z -axis is also no loss of generality.

For any given radius $R_c > 0$ of the reaction chamber and any radius $0 < R < R_c$, the following theorem ensures the existence of a nontrivial steady state that is confined in the infinite cylinder about the z -axis with radius R . Note that both a θ -pinch and a z -pinch configuration are considered.

Theorem 4.2 (Existence of confined steady states). *Suppose that (A0) holds and let $R, R_c \in \mathbb{R}$ with $0 < R < R_c$ be arbitrary. We further assume that the ansatz functions $\eta^\pm \in C^1(\mathbb{R}^3)$ and the axially symmetric external magnetic vector potential $A^0 \in C^2(\mathbb{R}^2; \mathbb{R}^3)$ satisfy the following assumptions:*

(A1) *There exist functions $\eta_*^\pm \in L^1(\mathbb{R}^2)$ such that*

$$\eta^\pm(\mathcal{E}, \mathcal{F}, \mathcal{G}) \leq \eta_*^\pm(\mathcal{E}, \mathcal{G}) \quad \text{for all } (\mathcal{E}, \mathcal{F}, \mathcal{G}) \in \mathbb{R}^3.$$

(A2) *There exist functions $\eta_\#^\pm \in L^1(\mathbb{R})$ such that*

$$|\partial_{\mathcal{E}} \eta^\pm(\mathcal{E}, \mathcal{F}, \mathcal{G})| \leq \eta_\#^\pm(\mathcal{E}, \mathcal{G}) \quad \text{for all } (\mathcal{E}, \mathcal{F}, \mathcal{G}) \in \mathbb{R}^3,$$

(A3) *There exist cut-off energies $\mathcal{E}_{\max}^\pm > 0$ such that*

$$\eta^\pm(\mathcal{E}, \mathcal{F}, \mathcal{G}) = 0 \quad \text{if } \mathcal{E} \geq \mathcal{E}_{\max}^\pm,$$

(A4) *There exist real numbers $a > 0$, $b > 0$ and $c > 0$ such that each of the functions η^\pm is strictly positive on $]0, a[\times]-b, 0[\times]-c, c[$ or on $]0, a[\times]0, b[\times]-c, c[$.*

(A5) If a θ -pinch configuration is considered, it holds that

$$\eta^\pm(\mathcal{E}, \mathcal{F}, \mathcal{G}) = 0 \quad \text{if } \pm \mathcal{F} \geq 0 \quad (4.2)$$

and A_φ^0 satisfies

$$A_\varphi^0(r) \geq \max_{\pm} \sqrt{2\mathcal{E}_{\max}^\pm + 4\pi^2 \|\eta_*^\pm\|_1 r^2} \quad \text{for all } r \geq R. \quad (4.3)$$

If a z -pinch configuration is considered, there exist $\mathcal{G}_0^\pm \in \mathbb{R}$ with $\pm \mathcal{G}_0 > 0$ such that

$$\eta^\pm(\mathcal{E}, \mathcal{F}, \mathcal{G}) = 0 \quad \text{if } \pm \mathcal{G} \geq \pm \mathcal{G}_0, \quad (4.4)$$

and that A_3^0 satisfies

$$A_3^0(r) \geq \max_{\pm} \left(|\mathcal{G}_0^\pm| + \sqrt{2\mathcal{E}_{\max}^\pm + 4\pi^2 \|\eta_*^\pm\|_1 r^2} \right) \quad \text{for all } r \geq R. \quad (4.5)$$

Then the ansatz

$$f_0^\pm = \eta^\pm(\mathcal{E}^\pm, \mathcal{F}^\pm, \mathcal{G}^\pm), \quad (4.6)$$

defines a nontrivial steady state of the Vlasov–Poisson system (1.1) in the sense of Definition 4.1. In particular, $f_0 = (f_0^+, f_0^-)$ is compactly supported in $\overline{\mathcal{B}_R} \times \mathbb{R}^3 \subset \mathbb{R}^5$ and thus confined in the cylinder with radius $R < R_c$.

Here and in the following, we denote by \mathcal{B}_R the ball in \mathbb{R}^2 about the origin with radius R . For the construction of such a steady state, we follow the reasoning in [22] and [39]. In [22] the existence of confined steady states for the two dimensional Vlasov–Poisson system was established, whereas in [39] confined steady states for the two and one-half-dimensional Vlasov–Maxwell system were constructed. As we are investigating steady states of the two and one-half dimensional Vlasov–Poisson system, the arguments of [22] and [39] can be combined. Indeed, since our line of argument is very similar, we only outline the strategy and sketch the most important steps.

Sketch of the proof. As in [22], we first observe that the densities ρ_0^\pm can be expressed by means of the ansatz functions η^\pm via

$$\begin{aligned} \rho_0^\pm(r) &= \int_{\mathbb{R}^3} f_0^\pm(r, p) \, dp = \int_{\mathbb{R}^3} \eta^\pm \left(\frac{1}{2}|p|^2 \pm U_0(r), r(p_\varphi \pm A_\varphi^0(r)), p_3 \pm A_3^0(r) \right) \, dp \\ &= \int_{\mathbb{R}} \int_0^\infty \int_0^{2\pi} u \eta^\pm \left(\frac{1}{2}u^2 + \frac{1}{2}p_3^2 \pm U_0(r), r(u \sin \theta \pm A_\varphi^0(r)), p_3 \pm A_3^0(r) \right) \, d\theta \, du \, dp_3 \\ &= \int_{\mathbb{R}} \int_{\frac{1}{2}(\mathcal{G} \mp A_3^0(r))^2 \pm U_0(r)}^\infty \int_0^{2\pi} \eta^\pm \left(\mathcal{E}, r \left(\sqrt{2(\mathcal{E} \mp U_0(r)) - (\mathcal{G} \mp A_3^0(r))^2} \sin \theta + A_\varphi^0(r) \right), \mathcal{G} \right) \\ &\quad d\theta \, d\mathcal{E} \, d\mathcal{G}. \end{aligned} \quad (4.7)$$

Note that the integral is well defined due to the assumption (A1). Under our symmetry assumptions, the Laplacian simplifies to $\Delta_x = \frac{1}{r}\partial_r(r\partial_r)$. Thus, (4.1b) can be integrated with respect to r , which yields

$$U_0(r) = -4\pi \int_0^r \frac{1}{s} \int_0^s \sigma \rho_0(\sigma) \, d\sigma \, ds. \quad (4.8)$$

We point out that in this formula, the boundary condition $U_0(0) = 0$ is incorporated. Thus, in general, the boundary condition (4.1c) cannot be satisfied. However, under the assumption that ρ_0 is compactly supported, it is easy to see that the limit in (4.1c) exists and is finite, but not necessarily vanishes. Since adding constants to the electric potential does not affect the electric field, (4.1c) can be replaced by the boundary condition $U_0(0) = 0$ which corresponds to (4.8). For a more detailed discussion see [22, Sect. 3].

After merging (4.7) and (4.8) the problem of finding a steady state can then be regarded as a fixed point problem for U_0 . Under the assumption (A2), on each bounded interval $[0, \delta]$, $\delta > 0$, this fixed point problem (in the space of continuous functions) can be tackled by Schaefer's fixed point theorem (cf. [39]) or a direct fixed point iteration (cf. [22]) to obtain a unique solution, which gains regularity a posteriori by means of (4.8). Now, that U_0 is constructed, the quantity \mathcal{E}^\pm is known explicitly. Since \mathcal{E}^\pm , \mathcal{F}^\pm and \mathcal{G}^\pm are invariants of the characteristic flow, we conclude that the ansatz (4.6) defines a steady state $f_0 = (f_0^+, f_0^-)$ of the Vlasov–Poisson system (1.1) (with (4.1c) being replaced by the equivalent boundary condition $U_0(0) = 0$) in the sense of Definition 4.1.

Having the steady state f_0 at hand, it remains to establish its claimed properties. Firstly, the assumption (A3) suffices to ensure that f_0 is compactly supported with respect to the velocity variable p (cf. [39, Theorem 4.8(i)]). Secondly, the assumption (A4) implies that f_0 is nontrivial (cf. [39, Theorem 4.8(ii)]). Thirdly, and most importantly, we need to show that the steady state is confined within a cylinder with radius $R < R_c$. Proceeding as in the proof of [39, Theorem 5.1] this assertion can be established by means of the assumption (A5) which ensures that the magnetic field B^0 is sufficiently strong to confine the plasma. We point out that, in order to satisfy (4.3) (or (4.5)), the function A_φ^0 (or A_3^0) cannot be chosen completely arbitrarily on $[0, R]$ due to the condition $A_\varphi^0(0) = 0$ (or $A_3^0(0) = 0$) resulting from the normalization at $r = 0$. Now all claims are established. \square

Let us make some concluding remarks.

- Remark 4.3.** (a) The assumptions (A1)–(A5) entail that $a \leq \min\{\mathcal{E}_{\max}^+, \mathcal{E}_{\max}^-\}$, and that $c \leq \min\{\mathcal{G}_0^+, -\mathcal{G}_0^-\}$ in the z -pinch case.
- (b) In the same fashion we could also consider the corresponding the ‘sign-reversed’ cases where the conditions $\pm\mathcal{F} \geq 0$ in (4.2) or $\pm\mathcal{G} \geq \pm\mathcal{G}_0 > 0$ in (4.5) are replaced by $\pm\mathcal{F} \leq 0$ or $\pm\mathcal{G} \leq \pm\mathcal{G}_0 < 0$, respectively. In these alternative cases, the results of Theorem 4.2 could be established analogously.
- (c) It would certainly also be possible to construct a nontrivial, confined steady state in a *screw-pinch configuration* that is a combination of a θ -pinch and a z -pinch. In this case the external magnetic vector potential would be given as a superposition of two vector-potentials A_φ^0 and A_3^0 , i.e., $A^0 = A_\varphi^0 + A_3^0$, such that (A5) is satisfied in the

sense that (4.2) and (4.4) hold, and (4.3) or (4.5) hold for all $r \geq R$. However, for simplicity, we restrict ourselves to the investigation of the pure θ -pinch and z -pinch configurations.

- (d) It is important to point out that we slightly differ from [39] as follows: In [39] the ansatz (4.6) is only made in the region $r \leq R_c$ and outside f_0^\pm is defined to be zero (that is to say, possible shells in the region $r > R_c$ are left out). This causes no problem thanks to $f_0^\pm = 0$ on $[R, R_c]$. As a consequence, (4.3) or (4.5) only need to be imposed on $[R, R_c]$, but it is unclear whether the relation (4.6) is also satisfied for $r > R_c$. However, in this paper it will be important—in particular, for (5.9)—that (4.6) holds for all $r \geq 0$. Thus, (4.3) or (4.5) have to be imposed on $[R, \infty[$.

5 Stability of confined steady states

We split the investigation of stability properties into two parts: First, the external magnetic potential is fixed and the initial data are perturbed. Second, the initial data are fixed and the external magnetic potential is perturbed. In the end we show that both results can also be combined.

Throughout this section, we will use the notation

$$\mathcal{R}(h) := \min\{r \geq 0 \mid h(x, v) = 0 \text{ if } |x| \geq r\}$$

for any axially symmetric $h \in C_c(\mathbb{R}^5)^j$ with $j \in \{1, 2\}$.

In Subsection 5.1, we will mainly follow the ideas of [6]. We fix an external magnetic potential A^0 satisfying (A0) and make the following ansatz for the steady state f_0^\pm :

$$f_0^\pm = \eta^\pm(\mathcal{E}^\pm, \mathcal{F}^\pm, \mathcal{G}^\pm) = \vartheta^\pm(\mathcal{E}^\pm) \psi^\pm(\mathcal{F}^\pm, \mathcal{G}^\pm). \quad (5.1)$$

In the following, we consider a θ -pinch configuration. In order to satisfy the assumptions made in Section 4, we assume

- (S1) $\vartheta^\pm \in C^1(\mathbb{R}) \cap W^{1,1}(\mathbb{R})$, and there exists a $\mathcal{E}_{\max}^\pm > 0$ such that $\vartheta^\pm(\tau) = 0$ for all $\tau \geq \mathcal{E}_{\max}^\pm$ and $\vartheta^\pm(\tau) > 0$ for all $\tau < \mathcal{E}_{\max}^\pm$;
- (S2) $\psi^\pm \in C^1(\mathbb{R}^2)$, $\psi^\pm(\sigma, \mu) > 0$ for $\pm\sigma < 0$, and $\psi^\pm(\sigma, \mu) = 0$ for $\pm\sigma \geq 0$, and there exist $\psi_*^\pm \in L^1(\mathbb{R})$ such that $\psi^\pm(\sigma, \mu) \leq \psi_*^\pm(\mu)$ for all $(\sigma, \mu) \in \mathbb{R}^2$;
- (S3) there exists $0 < \tilde{R} < R_c$ such that $A_\varphi^0(r) \geq \max_\pm \sqrt{2\mathcal{E}_{\max}^\pm + 4\pi^2 \|\vartheta^\pm\|_1 \|\psi_*^\pm\|_1} r^2$ for all $r \geq \tilde{R}$.

Here, $R_c > 0$ stands again for the fixed radius of the cylindrical reaction chamber.

Since the assumptions (S1)–(S3) imply (A1)–(A5) (with \tilde{R} instead of R in (A5)), Theorem 4.2 ensures that the steady state f_0 in (5.1) actually exists, is nontrivial and confined in a cylinder with radius $R = \mathcal{R}(f_0) \leq \tilde{R} < R_c$. In particular, R is thus strictly smaller than the radius R_c of the reaction chamber.

We let $R^\pm := \mathcal{R}(f_0^\pm)$, $\mathcal{E}_0^\pm := \frac{1}{2}|p|^2 \pm U_0$, and

$$\mathcal{E}_{\min}^\pm := \inf_{\mathcal{B}_{R^\pm} \times \mathbb{R}^3} \mathcal{E}_0^\pm = \inf_{\mathcal{B}_{R^\pm}} \pm U_0.$$

Notice that $\mathcal{E}_{\min}^\pm < \mathcal{E}_{\max}^\pm$ since the steady state $f_0 = (f_0^+, f_0^-)$ is nontrivial.

Moreover, we assume

$$(S4) \quad (\vartheta^\pm)'(\tau) < 0 \text{ for } \tau \in [\mathcal{E}_{\min}^\pm, \mathcal{E}_{\max}^\pm[,$$

which is crucial in order to obtain stability.

For $f = (f^+, f^-) \in C_c(\mathbb{R}^5)^2$, we introduce the kinetic energy

$$E_{\text{kin}}(f) = \sum_{\pm} \frac{1}{2} \int_{\mathbb{R}^5} |p|^2 f^\pm \, dz$$

and the potential energy

$$E_{\text{pot}}(f) = \frac{1}{2} \int_{\mathbb{R}^2} U_f \rho_f \, dx = - \int_{\mathbb{R}^2} \int_{\mathbb{R}^2} \ln |x - y| \rho_f(y) \rho_f(x) \, dy \, dx,$$

where

$$\rho_f = \int_{\mathbb{R}^3} f^+ \, dp - \int_{\mathbb{R}^3} f^- \, dp, \quad U_f = -2 \ln |\cdot| * \rho_f.$$

It is very important to notice that we cannot use integration by parts to equivalently describe the potential energy as an integral over $|\partial_x U_f|^2$ as it would be the case in a three dimensional setting. This is because, in contrast to a 3D situation with kernel $|\cdot|^{-1}$, the kernel $\ln |\cdot|$ does not vanish at infinity. We further recall that the total energy

$$\mathcal{H}(f) = E_{\text{kin}}(f) + E_{\text{pot}}(f)$$

is conserved along classical solutions of (1.1).

5.1 Stability with respect to perturbations of the initial data

We consider perturbations from the following set of functions:

$$X := \left\{ g = (g^+, g^-) \in C_c^1(\mathbb{R}^5)^2 \left| \begin{array}{l} g^\pm \geq 0, g^\pm \text{ are axially symmetric,} \\ g^\pm / (\psi^\pm \circ (\mathcal{F}^\pm, \mathcal{G}^\pm)) \in L^1(\{\pm \mathcal{F}^\pm < 0\}), \\ \int_{\mathbb{R}^2} \rho_g \, dx = \int_{\mathbb{R}^2} \rho_0 \, dx \end{array} \right. \right\}.$$

Notice that, on the one hand, the assumption that g has the same total charge as f_0 is reasonable since physically meaningful perturbations preserve the total charge, and, on the other hand, important for the positive definiteness of the potential energy induced by a difference $g - f_0$; cf. Lemma 2.2(b).

Before we proceed, we first recall that the potential U_0 satisfies (4.8). We now define $M(r) := 2\pi \int_0^r s \rho_0(s) \, ds$, i.e., $M(r)$ is the charge in a sliced circle with radius r , and the

total charge in each slice can be expressed as $M = M(R)$. Then the electric potential U_0 can be written as

$$U_0(r) = \begin{cases} -2 \int_0^r \frac{M(s)}{s} ds, & 0 \leq r \leq R, \\ -2 \int_0^R \frac{M(s)}{s} ds - 2M(\ln r - \ln R), & r > R. \end{cases}$$

Therefore, the function

$$\xi(r) := \sup_{B_r} |U_0|, \quad r > 0 \quad (5.2)$$

grows logarithmically on $[R, \infty[$ if $M \neq 0$ and is constant on $[R, \infty[$ if $M = 0$.

5.1.1 Construction of the energy-Casimir functional

To prove stability with respect to perturbations of the initial data, we employ the *energy-Casimir method*. The idea is to construct a *Casimir functional*

$$\mathcal{C}(f) = \sum_{\pm} \int_{\mathbb{R}^2} \int_{\mathbb{R}^3} \Phi^{\pm}(f^{\pm}, \mathcal{F}^{\pm}, \mathcal{G}^{\pm}) dp dx \quad (5.3)$$

in such a way that its derivative at f_0 matches exactly or at least dominates the negative of the linear part of the expansion

$$\begin{aligned} \mathcal{H}(f) &= \mathcal{H}(f_0) + \sum_{\pm} \int_{\mathbb{R}^2} \int_{\mathbb{R}^3} \mathcal{E}_0^{\pm}(f^{\pm} - f_0^{\pm}) dp dx \\ &\quad - \int_{\mathbb{R}^2} \int_{\mathbb{R}^2} \ln|x-y|(\rho_f - \rho_0)(y)(\rho_f - \rho_0)(x) dy dx. \end{aligned} \quad (5.4)$$

Note that for any choice of Φ any $f \in X$, the expression $\mathcal{C}(f)$ is constant in time. For more background on the energy-Casimir method we refer to [6, 18, 20, 32, 33].

Let $\vartheta_{\max}^{\pm} := \vartheta^{\pm}(\mathcal{E}_{\min}^{\pm})$. By assumption, the map $\vartheta^{\pm}: [\mathcal{E}_{\min}^{\pm}, \mathcal{E}_{\max}^{\pm}] \rightarrow [0, \vartheta_{\max}^{\pm}]$ is strictly decreasing and onto. We now construct the functions $\Phi^{\pm}: [0, \infty[\times \mathbb{R} \times \mathbb{R} \rightarrow \mathbb{R}$ which defines the Casimir functional $\mathcal{C}(f)$. Note that $\Phi^{\pm} = \Phi^{\pm}(\tau, \sigma, \mu)$ can be chosen arbitrarily (e.g., zero) if $\sigma = 0$ since the set $\{\mathcal{F}^{\pm} = 0\}$ has Lebesgue measure zero. First, we consider $\pm\sigma < 0$. For $\mu \in \mathbb{R}$ and $\tau \in [0, \vartheta_{\max}^{\pm} \psi^{\pm}(\sigma, \mu)]$ we define

$$\Phi^{\pm}(\tau, \sigma, \mu) := -\psi^{\pm}(\sigma, \mu) \int_0^{\tau/\psi^{\pm}(\sigma, \mu)} (\vartheta^{\pm})^{-1}(s) ds.$$

Clearly, $\Phi^{\pm}(\cdot, \sigma, \mu) \in C^1([0, \vartheta_{\max}^{\pm} \psi^{\pm}(\sigma, \mu)]) \cap C^2([0, \vartheta_{\max}^{\pm} \psi^{\pm}(\sigma, \mu)])$ with

$$\partial_{\tau}^{\pm} \Phi(\tau, \sigma, \mu) = -(\vartheta^{\pm})^{-1} \left(\frac{\tau}{\psi^{\pm}(\sigma, \mu)} \right)$$

for $\tau \in [0, \vartheta_{\max}^{\pm} \psi^{\pm}(\sigma, \mu)]$ and

$$\partial_{\tau}^2 \Phi^{\pm}(\tau, \sigma, \mu) = -\frac{1}{(\vartheta^{\pm})'((\vartheta^{\pm})^{-1}(\tau/\psi^{\pm}(\sigma, \mu))) \psi^{\pm}(\sigma, \mu)}$$

$$\geq -\frac{1}{\left(\inf_{[\mathcal{E}_{\min}^{\pm}, \mathcal{E}_{\max}^{\pm}]}(\vartheta^{\pm})'\right)\psi^{\pm}(\sigma, \mu)} =: \frac{c_{\vartheta^{\pm}}}{\psi^{\pm}(\sigma, \mu)}$$

for $\tau \in]0, \vartheta_{\max}^{\pm}\psi^{\pm}(\sigma, \mu)]$; notice that $c_{\vartheta^{\pm}} > 0$. Next, for $\tau > \vartheta_{\max}^{\pm}\psi^{\pm}(\sigma, \mu)$ we choose

$$\begin{aligned} \Phi^{\pm}(\tau, \sigma, \mu) &:= -\frac{(\tau - \vartheta_{\max}^{\pm}\psi^{\pm}(\sigma, \mu))^2}{2(\vartheta^{\pm})'(\mathcal{E}_{\min}^{\pm})\psi^{\pm}(\sigma, \mu)} - \mathcal{E}_{\min}^{\pm}(\tau - \vartheta_{\max}^{\pm}\psi^{\pm}(\sigma, \mu)) \\ &\quad - \psi^{\pm}(\sigma, \mu) \int_0^{\vartheta_{\max}^{\pm}} (\vartheta^{\pm})^{-1}(s) \, ds \end{aligned}$$

to extend Φ^{\pm} to a function $\Phi^{\pm} \in C^1([0, \infty[\times \mathbb{R}_{\mp} \times \mathbb{R})$ with $\Phi^{\pm}(\cdot, \sigma, \mu) \in C^2([0, \infty[)$, and

$$\begin{aligned} |\Phi^{\pm}(\tau, \sigma, \mu)| &\leq c_{\pm} \left(\tau + \frac{\tau^2}{\psi^{\pm}(\sigma, \mu)} \right), \quad \tau \geq 0, \\ |\partial_{\tau} \Phi^{\pm}(\tau, \sigma, \mu)| &\leq c_{\pm} \left(1 + \frac{\tau}{\psi^{\pm}(\sigma, \mu)} \right), \quad \tau \geq 0, \end{aligned} \tag{5.5}$$

$$\partial_{\tau}^2 \Phi^{\pm}(\tau, \sigma, \mu) \geq \frac{c_{\vartheta^{\pm}}}{\psi^{\pm}(\sigma, \mu)}, \quad \tau > 0. \tag{5.6}$$

for positive constants c_{\pm} and $c_{\vartheta^{\pm}}$ that may depend on ψ^{\pm} and ϑ^{\pm} but not on (τ, σ, μ) .

Let us now consider $\pm\sigma > 0$. We fix $r_0 > 0$ such that

$$\min\{A_{\varphi}^0(r), r\} \geq \sqrt{2|U_0(r)|} \quad \text{for all } r \geq r_0. \tag{5.7}$$

Notice that there exists such an r_0 due to (S3) and the fact that $|U_0|$ grows at most logarithmically for large r . For all $\tau \geq 0$ and $\mu \in \mathbb{R}$, we define

$$\Phi^{\pm}(\tau, \sigma, \mu) := (\pm\sigma + \xi(r_0))\tau,$$

where ξ is the function introduced in (5.2). Thus, $\Phi^{\pm} \in C^{\infty}([0, \infty[\times \mathbb{R}_{\pm} \times \mathbb{R})$.

Now, for any $f \in X$, we define $\mathcal{C}(f)$ by (5.3). Notice that $\mathcal{C}(f)$ is well-defined because of

$$\begin{aligned} &\int_{\mathbb{R}^2} \int_{\mathbb{R}^3} |\Phi^{\pm}(f^{\pm}, \mathcal{F}^{\pm}, \mathcal{G}^{\pm})| \, dp \, dx \\ &= \iint_{\{\pm\mathcal{F}^{\pm} > 0\}} |\Phi^{\pm}(f^{\pm}, \mathcal{F}^{\pm}, \mathcal{G}^{\pm})| \, dp \, dx + \iint_{\{\pm\mathcal{F}^{\pm} < 0\}} |\Phi(f^{\pm}, \mathcal{F}^{\pm}, \mathcal{G}^{\pm})| \, dp \, dx \\ &\leq \|\mathcal{F}^{\pm} f^{\pm}\|_1 + (\xi(r_0) + c_{\pm})\|f^{\pm}\|_1 + c_{\pm}\|f^{\pm}\|_{\infty} \iint_{\{\pm\mathcal{F}^{\pm} < 0\}} \frac{f^{\pm}}{\psi^{\pm}(\mathcal{F}^{\pm}, \mathcal{G}^{\pm})} \, dp \, dx < \infty. \end{aligned}$$

Since \mathcal{F}^{\pm} and \mathcal{G}^{\pm} are invariants of the characteristic flow associated with $B^0 = \text{curl}_x A^0$, \mathcal{C} is conserved along classical solutions of (1.1) to the external vector potential A^0 . Hence, the same holds for the *energy-Casimir functional* $\mathcal{H}_{\mathcal{C}} := \mathcal{H} + \mathcal{C}$. In view of (5.4), we have the expansion

$$\begin{aligned} &\mathcal{H}_{\mathcal{C}}(f) - \mathcal{H}_{\mathcal{C}}(f_0) \\ &= \sum_{\pm} \int_{\mathbb{R}^2} \int_{\mathbb{R}^3} (\Phi^{\pm}(f^{\pm}, \mathcal{F}^{\pm}, \mathcal{G}^{\pm}) - \Phi^{\pm}(f_0^{\pm}, \mathcal{F}^{\pm}, \mathcal{G}^{\pm}) + \mathcal{E}_0^{\pm}(f^{\pm} - f_0^{\pm})) \, dp \, dx \end{aligned}$$

$$- \int_{\mathbb{R}^2} \int_{\mathbb{R}^2} \ln |x - y| (\rho_f - \rho_0)(y) (\rho_f - \rho_0)(x) \, dy \, dx. \quad (5.8)$$

In the next two subsections, we establish lower and upper estimates of the right-hand side of this expansion.

5.1.2 Lower estimates on the expansion

First, we infer from Lemma 2.2(b) that

$$\begin{aligned} & \mathcal{H}_{\mathcal{C}}(f) - \mathcal{H}_{\mathcal{C}}(f_0) \\ & \geq \sum_{\pm} \int_{\mathbb{R}^2} \int_{\mathbb{R}^3} (\Phi^{\pm}(f^{\pm}, \mathcal{F}^{\pm}, \mathcal{G}^{\pm}) - \Phi^{\pm}(f_0^{\pm}, \mathcal{F}^{\pm}, \mathcal{G}^{\pm}) + \mathcal{E}_0^{\pm}(f^{\pm} - f_0^{\pm})) \, dp \, dx \\ & = \sum_{\pm} \left(\iint_{\{\pm \mathcal{F}^{\pm} < 0, f_0^{\pm} > 0\}} + \iint_{\{\pm \mathcal{F}^{\pm} < 0, f_0^{\pm} = 0\}} + \iint_{\{\pm \mathcal{F}^{\pm} > 0, f_0^{\pm} = 0\}} \right) Q^{\pm} \, dp \, dx, \end{aligned}$$

where Q^{\pm} is defined as

$$Q^{\pm} := \Phi^{\pm}(f^{\pm}, \mathcal{F}^{\pm}, \mathcal{G}^{\pm}) - \Phi(f_0^{\pm}, \mathcal{F}^{\pm}, \mathcal{G}^{\pm}) + \mathcal{E}_0^{\pm}(f^{\pm} - f_0^{\pm});$$

notice that $\{\mathcal{F}^{\pm} = 0\} \subset \mathbb{R}^5$ has Lebesgue measure zero and that $f_0^{\pm} = 0$ if $\pm \mathcal{F}^{\pm} > 0$. Now, we fix $(x, p) \in \mathbb{R}^5$ and consider three cases corresponding to the decomposition above. For simplicity, we will suppress the argument (x, p) in the following.

Case 1: $\pm \mathcal{F}^{\pm} < 0, f_0^{\pm} > 0$: Then $\mathcal{E}_0^{\pm} \in [\mathcal{E}_{\min}^{\pm}, \mathcal{E}_{\max}^{\pm}[$ and

$$\mathcal{E}_0^{\pm} = (\vartheta^{\pm})^{-1}(\vartheta^{\pm}(\mathcal{E}_0^{\pm})) = (\vartheta^{\pm})^{-1} \left(\frac{f_0^{\pm}}{\psi^{\pm}(\mathcal{F}^{\pm}, \mathcal{G}^{\pm})} \right) = -\partial_{\tau} \Phi^{\pm}(f_0^{\pm}, \mathcal{F}^{\pm}, \mathcal{G}^{\pm}).$$

By the known regularity of Φ^{\pm} and (5.6), we have

$$\begin{aligned} Q^{\pm} &= \Phi^{\pm}(f^{\pm}, \mathcal{F}^{\pm}, \mathcal{G}^{\pm}) - \Phi^{\pm}(f_0^{\pm}, \mathcal{F}^{\pm}, \mathcal{G}^{\pm}) - \partial_{\tau} \Phi^{\pm}(f_0^{\pm}, \mathcal{F}^{\pm}, \mathcal{G}^{\pm})(f^{\pm} - f_0^{\pm}) \\ &= \lim_{\varepsilon \rightarrow 0^+} (\Phi^{\pm}(f^{\pm} + \varepsilon, \mathcal{F}^{\pm}, \mathcal{G}^{\pm}) - \Phi^{\pm}(f_0^{\pm}, \mathcal{F}^{\pm}, \mathcal{G}^{\pm}) \\ &\quad - \partial_{\tau} \Phi(f_0^{\pm}, \mathcal{F}^{\pm}, \mathcal{G}^{\pm})(f^{\pm} + \varepsilon - f_0^{\pm})) \\ &= \lim_{\varepsilon \rightarrow 0^+} \frac{1}{2} \partial_{\tau}^2 \Phi^{\pm}(\zeta_{\varepsilon}, \mathcal{F}^{\pm}, \mathcal{G}^{\pm})(f^{\pm} + \varepsilon - f_0^{\pm})^2 \geq \frac{c_{\vartheta^{\pm}}}{2\psi^{\pm}(\mathcal{F}^{\pm}, \mathcal{G}^{\pm})} (f^{\pm} - f_0^{\pm})^2, \end{aligned}$$

where ζ_{ε} lies between f_0^{\pm} and $f^{\pm} + \varepsilon$.

Case 2: $\pm \mathcal{F}^{\pm} < 0, f_0^{\pm} = 0$: Then

$$\mathcal{E}_0^{\pm} \geq \mathcal{E}_{\max}^{\pm} = (\vartheta^{\pm})^{-1}(0) = -\partial_{\tau} \Phi^{\pm}(0, \mathcal{F}^{\pm}, \mathcal{G}^{\pm}). \quad (5.9)$$

Thus, similarly as before,

$$Q^{\pm} \geq \Phi^{\pm}(f^{\pm}, \mathcal{F}^{\pm}, \mathcal{G}^{\pm}) - \Phi^{\pm}(0, \mathcal{F}^{\pm}, \mathcal{G}^{\pm}) - \partial_{\tau} \Phi(0, \mathcal{F}^{\pm}, \mathcal{G}^{\pm}) f^{\pm}$$

$$\begin{aligned}
&= \lim_{\varepsilon \rightarrow 0^+} (\Phi^\pm(f^\pm + \varepsilon, \mathcal{F}^\pm, \mathcal{G}^\pm) - \Phi^\pm(\varepsilon, \mathcal{F}^\pm, \mathcal{G}^\pm) - \partial_\tau \Phi^\pm(\varepsilon, \mathcal{F}^\pm, \mathcal{G}^\pm) f^\pm) \\
&= \lim_{\varepsilon \rightarrow 0^+} \frac{1}{2} \partial_\tau^2 \Phi^\pm(\zeta_\varepsilon, \mathcal{F}^\pm, \mathcal{G}^\pm) (f^\pm)^2 \geq \frac{c_{\vartheta^\pm}}{2\psi^\pm(\mathcal{F}^\pm, \mathcal{G}^\pm)} (f^\pm)^2 \\
&= \frac{c_{\vartheta^\pm}}{2\psi^\pm(\mathcal{F}^\pm, \mathcal{G}^\pm)} (f^\pm - f_0^\pm)^2.
\end{aligned}$$

Case 3: $\pm \mathcal{F}^\pm > 0$, $f_0^\pm = 0$: We have

$$Q^\pm = (\pm \mathcal{F}^\pm + \xi(r_0) + \frac{1}{2}|p|^2 \pm U_0) f^\pm.$$

To prove that $Q^\pm \geq 0$ it suffices to show that

$$q^\pm := \pm \mathcal{F}^\pm + \xi(r_0) + \frac{1}{2}|p|^2 \pm U_0 \geq 0.$$

To this end, we consider four sub-cases; notice that we already know that $\pm(p_\varphi \pm A_\varphi^0) > 0$ due to $\pm \mathcal{F}^\pm > 0$, and have in mind (5.7):

Case 3.1: $r \leq r_0$. Clearly,

$$q^\pm \geq \xi(r_0) \pm U_0 \geq 0.$$

Case 3.2: $r > r_0$, $\pm p_\varphi \geq 0$: Here, it holds that

$$q^\pm \geq r A_\varphi^0 \pm U_0 \geq 2|U_0| \pm U_0 \geq 0.$$

Case 3.3: $r > r_0$, $0 < \mp p_\varphi < A_\varphi^0 \leq r$. Since the function $y \mapsto \frac{1}{2}y^2 - ry$ is monotonically decreasing on $[0, A_\varphi^0] \subset [0, r]$, we have

$$\begin{aligned}
q^\pm &\geq r A_\varphi^0 + \frac{1}{2}(\mp p_\varphi)^2 - r(\mp p_\varphi) \pm U_0 \geq r A_\varphi^0 + \frac{1}{2}(A_\varphi^0)^2 - r A_\varphi^0 \pm U_0 = \frac{1}{2}(A_\varphi^0)^2 \pm U_0 \\
&\geq |U_0| \pm U_0 \geq 0.
\end{aligned}$$

Case 3.4: $r > r_0$, $0 < \mp p_\varphi < A_\varphi^0$, $A_\varphi^0 > r$: Since the function $y \mapsto \frac{1}{2}y^2 - ry$ attains its global minimum at $y = r$, we have

$$\begin{aligned}
q^\pm &\geq r A_\varphi^0 + \frac{1}{2}(\mp p_\varphi)^2 - r(\mp p_\varphi) \pm U_0 \geq r A_\varphi^0 + \frac{1}{2}r^2 - r^2 \pm U_0 \geq \frac{1}{2}r^2 \pm U_0 \\
&\geq |U_0| \pm U_0 \geq 0.
\end{aligned}$$

Thus, we always have $Q^\pm \geq 0$ in Case 3.

Combining all three cases, we obtain

$$\mathcal{H}_C(f) - \mathcal{H}_C(f_0) \geq \sum_{\pm} \frac{c_{\vartheta^\pm}}{2} \iint_{\{\pm \mathcal{F}^\pm < 0\}} \frac{(f^\pm - f_0^\pm)^2}{\psi^\pm(\mathcal{F}^\pm, \mathcal{G}^\pm)} dp dx. \quad (5.10)$$

5.1.3 Upper estimates on the expansion

In the following we denote $\mathcal{S}(h) := \max\{\mathcal{R}(h), R\}$ for any axially symmetric $h \in C_c(\mathbb{R}^5)^j$ ($j \in \{1, 2\}$). Now we rewrite (5.8) as

$$\mathcal{H}_C(f) - \mathcal{H}_C(f_0) = \sum_{\pm} \left(\iint_{\{\pm \mathcal{F}^\pm < 0\}} Q^\pm dp dx + \iint_{\{\pm \mathcal{F}^\pm > 0\}} Q^\pm dp dx \right)$$

$$- \int_{\mathbb{R}^2} \int_{\mathbb{R}^2} \ln |x - y| (\rho_f - \rho_0)(y) (\rho_f - \rho_0)(x) \, dy \, dx,$$

and estimate the three terms separately. In the case $\pm \mathcal{F}^\pm < 0$ we have

$$\begin{aligned} Q^\pm &\leq |\Phi^\pm(f^\pm, \mathcal{F}^\pm, \mathcal{G}^\pm) - \Phi^\pm(f_0^\pm, \mathcal{F}^\pm, \mathcal{G}^\pm)| + |\mathcal{E}_0^\pm(f^\pm - f_0^\pm)| \\ &\leq \left(c_\pm + \frac{c_\pm \max\{\|f^\pm\|_\infty, \|f_0^\pm\|_\infty\}}{\psi^\pm(\mathcal{F}^\pm, \mathcal{G}^\pm)} + \frac{1}{2}|p|^2 + \xi(\mathcal{S}(f^\pm 1_{\{\pm \mathcal{F}^\pm < 0\}})) \right) |f^\pm - f_0^\pm| \end{aligned}$$

by (5.5). In the case $\pm \mathcal{F}^\pm > 0$ it holds that

$$\begin{aligned} Q^\pm &\leq |\Phi^\pm(f^\pm, \mathcal{F}^\pm, \mathcal{G}^\pm)| + |\mathcal{E}_0^\pm f^\pm| \\ &\leq \left(\pm \mathcal{F}^\pm + \xi(r_0) + \frac{1}{2}|p|^2 + \xi(\mathcal{R}(f^\pm 1_{\{\pm \mathcal{F}^\pm > 0\}})) \right) f^\pm \\ &= \left(\pm \mathcal{F}^\pm + \xi(r_0) + \frac{1}{2}|p|^2 + \xi(\mathcal{R}(f^\pm 1_{\{\pm \mathcal{F}^\pm > 0\}})) \right) |f^\pm - f_0^\pm|. \end{aligned}$$

Lastly, making use of the assumption that ρ_f and ρ_0 have the same total charge and applying Lemma 2.2(a), we find that

$$\begin{aligned} &- \int_{\mathbb{R}^2} \int_{\mathbb{R}^2} \ln |x - y| (\rho_f - \rho_0)(y) (\rho_f - \rho_0)(x) \, dy \, dx \\ &= - \int_{\mathbb{R}^2} \int_{\mathbb{R}^2} \ln \frac{|x - y|}{2\mathcal{S}(f)} (\rho_f - \rho_0)(y) (\rho_f - \rho_0)(x) \, dy \, dx \\ &\quad - \ln(2\mathcal{S}(f)) \left(\int_{\mathbb{R}^2} (\rho_f - \rho_0) \, dx \right)^2 \\ &= - \int_{\mathcal{B}_{\mathcal{S}(f)}} \int_{\mathcal{B}_{\mathcal{S}(f)}} \ln \frac{|x - y|}{2\mathcal{S}(f)} (\rho_f - \rho_0)(y) (\rho_f - \rho_0)(x) \, dy \, dx \\ &\leq - \int_{\mathcal{B}_{\mathcal{S}(f)}} \int_{\mathcal{B}_{\mathcal{S}(f)}} \ln \frac{|x - y|}{2\mathcal{S}(f)} |\rho_f - \rho_0|(y) |\rho_f - \rho_0|(x) \, dy \, dx \\ &= - \int_{\mathbb{R}^2} \int_{\mathbb{R}^2} \ln |x - y| |\rho_f - \rho_0|(y) |\rho_f - \rho_0|(x) \, dy \, dx + \ln(2\mathcal{S}(f)) \|\rho_f - \rho_0\|_1^2 \\ &\leq \frac{1}{2} \|\rho_f - \rho_0\|_1 \int_{\mathbb{R}^2} |\rho_f - \rho_0| \ln \frac{|\rho_f - \rho_0|}{\|\rho_f - \rho_0\|_1} \, dx + (C + \ln(2\mathcal{S}(f))) \|\rho_f - \rho_0\|_1^2. \end{aligned}$$

In summary, we have

$$\begin{aligned} &\mathcal{H}_{\mathcal{C}}(f) - \mathcal{H}_{\mathcal{C}}(f_0) \\ &\leq \sum_{\pm} \left[\iint_{\{\pm \mathcal{F}^\pm < 0\}} \left(c_\pm + \frac{c_\pm \max\{\|f^\pm\|_\infty, \|f_0^\pm\|_\infty\}}{\psi^\pm(\mathcal{F}^\pm, \mathcal{G}^\pm)} + \frac{1}{2}|p|^2 \right) |f^\pm - f_0^\pm| \, dp \, dx \right. \\ &\quad + \iint_{\{\pm \mathcal{F}^\pm < 0\}} \xi(\mathcal{S}(f^\pm 1_{\{\pm \mathcal{F}^\pm < 0\}})) |f^\pm - f_0^\pm| \, dp \, dx \\ &\quad \left. + \iint_{\{\pm \mathcal{F}^\pm > 0\}} \left(\pm \mathcal{F}^\pm + \xi(r_0) + \frac{1}{2}|p|^2 + \xi(\mathcal{R}(f^\pm 1_{\{\pm \mathcal{F}^\pm > 0\}})) \right) |f^\pm - f_0^\pm| \, dp \, dx \right] \end{aligned}$$

$$+ \frac{1}{2} \|\rho_f - \rho_0\|_1 \int_{\mathbb{R}^2} |\rho_f - \rho_0| \ln \frac{|\rho_f - \rho_0|}{\|\rho_f - \rho_0\|_1} dx + (C + \ln(2\mathcal{S}(f))) \|\rho_f - \rho_0\|_1^2. \quad (5.11)$$

5.1.4 The stability result

For any initial data $\mathring{f} = (\mathring{f}^+, \mathring{f}^-) \in X$, we now write f to denote the unique classical solution of (1.1) to the external vector potential A^0 . Recall that $\mathcal{H}_C(f)$ is conserved in time. We can thus combine (5.10) and (5.11) written for $\mathring{f} \in X$ to obtain a stability estimate on the set $\{\pm \mathcal{F}^\pm < 0\}$ (see (5.13)). Moreover, it is obvious that all L^q -norms of $f^\pm - f_0^\pm$ over $\{\pm \mathcal{F}^\pm \geq 0\}$ are constant time, since $f_0^\pm = 0$ if $\pm \mathcal{F}^\pm > 0$ and $\{\pm \mathcal{F}^\pm \geq 0\}$ is invariant under the characteristic flow of the Vlasov equation with external vector potential A^0 . Altogether, this means that the following theorem is established.

Theorem 5.1. *We consider the θ -pinch configuration and assume that (S1)–(S4) hold. Let f_0 denote the steady state introduced in (5.1).*

For any initial data $\mathring{f} = (\mathring{f}^+, \mathring{f}^-) \in X$ let $f = (f^+, f^-)$ denote the unique classical solution of (1.1).

Then for all $q \in [1, \infty]$,

$$\|f^\pm(t) - f_0^\pm\|_{L^q(\{\pm \mathcal{F}^\pm \geq 0\})} = \|\mathring{f}^\pm - f_0^\pm\|_{L^q(\{\pm \mathcal{F}^\pm \geq 0\})}, \quad (5.12)$$

and there exist constants $C > 0$ independent of all the appearing quantities, and $c_1^\pm, c_2^\pm > 0$ depending only on ϑ^\pm and ψ^\pm such that for all $\mathring{f} \in X$ and $t \geq 0$,

$$\begin{aligned} & \sum_{\pm} c_1^\pm \iint_{\{\pm \mathcal{F}^\pm < 0\}} \frac{(f^\pm(t) - f_0^\pm)^2}{\psi^\pm(\mathcal{F}^\pm, \mathcal{G}^\pm)} dp dx \\ & \leq \sum_{\pm} \left[\iint_{\{\pm \mathcal{F}^\pm < 0\}} \left(c_2^\pm + \frac{c_2^\pm \max\{\|\mathring{f}^\pm\|_\infty, \|f_0^\pm\|_\infty\}}{\psi^\pm(\mathcal{F}^\pm, \mathcal{G}^\pm)} + \frac{1}{2}|p|^2 \right) |\mathring{f}^\pm - f_0^\pm| dp dx \right. \\ & \quad + \iint_{\{\pm \mathcal{F}^\pm < 0\}} \xi(\mathcal{S}(\mathring{f}^\pm 1_{\{\pm \mathcal{F}^\pm < 0\}})) |\mathring{f}^\pm - f_0^\pm| dp dx \\ & \quad \left. + \iint_{\{\pm \mathcal{F}^\pm > 0\}} \left(\pm \mathcal{F}^\pm + \xi(r_0) + \frac{1}{2}|p|^2 + \xi(\mathcal{R}(\mathring{f}^\pm 1_{\{\pm \mathcal{F}^\pm > 0\}})) \right) |\mathring{f}^\pm - f_0^\pm| dp dx \right] \\ & + \frac{1}{2} \|\rho_{\mathring{f}} - \rho_0\|_1 \int_{\mathbb{R}^2} |\rho_{\mathring{f}} - \rho_0| \ln \frac{|\rho_{\mathring{f}} - \rho_0|}{\|\rho_{\mathring{f}} - \rho_0\|_1} dx + (C + \ln(2\mathcal{S}(\mathring{f}))) \|\rho_{\mathring{f}} - \rho_0\|_1^2. \quad (5.13) \end{aligned}$$

Remark 5.2. If ψ^\pm is additionally assumed to be bounded, then from (5.12) and (5.13) the (slightly weaker) estimate

$$\begin{aligned}
& \sum_{\pm} \|f^{\pm}(t) - f_0^{\pm}\|_{L^2(\mathbb{R}^2 \times \mathbb{R}^3)}^2 \\
& \leq c^{\pm} \left\{ \sum_{\pm} \left[\iint_{\{\pm \mathcal{F}^{\pm} < 0\}} \left(1 + \frac{\max\{\|\mathring{f}^{\pm}\|_{\infty}, \|f_0^{\pm}\|_{\infty}\}}{\psi^{\pm}(\mathcal{F}^{\pm}, \mathcal{G}^{\pm})} + \frac{1}{2}|p|^2 \right) |\mathring{f}^{\pm} - f_0^{\pm}| \, dp \, dx \right. \right. \\
& \quad + \iint_{\{\pm \mathcal{F}^{\pm} < 0\}} \xi(\mathcal{S}(\mathring{f}^{\pm} 1_{\{\pm \mathcal{F}^{\pm} < 0\}})) |\mathring{f}^{\pm} - f_0^{\pm}| \, dp \, dx \\
& \quad \left. + \iint_{\{\pm \mathcal{F}^{\pm} > 0\}} \left(\pm \mathcal{F}^{\pm} + \xi(r_0) + \frac{1}{2}|p|^2 + \xi(\mathcal{R}(\mathring{f}^{\pm} 1_{\{\pm \mathcal{F}^{\pm} > 0\}})) \right) |\mathring{f}^{\pm} - f_0^{\pm}| \, dp \, dx \right] \\
& \quad + \frac{1}{2} \|\rho_{\mathring{f}} - \rho_0\|_1 \int_{\mathbb{R}^2} |\rho_{\mathring{f}} - \rho_0| \ln \frac{|\rho_{\mathring{f}} - \rho_0|}{\|\rho_{\mathring{f}} - \rho_0\|_1} \, dx + (C + \ln(2\mathcal{S}(\mathring{f}))) \|\rho_{\mathring{f}} - \rho_0\|_1^2 \Big\} \\
& \quad + \|\mathring{f}^{\pm} - f_0^{\pm}\|_{L^2(\{\pm \mathcal{F}^{\pm} \geq 0\})}^2
\end{aligned}$$

follows, with constants $c^{\pm} > 0$ depending only on ϑ^{\pm} and ψ^{\pm} .

5.2 Stability with respect to perturbations of the magnetic vector potential

The continuous dependence estimate presented in Theorem 3.2 can be used to prove stability of the confined steady state with respect to perturbations of the external magnetic field.

Theorem 5.3. *Let $f_0 = (f_0^+, f_0^-)$ denote an arbitrary steady state (in the sense of Definition 4.1) with external magnetic vector potential A^0 and the associate magnetic field $B^0 = \text{curl}_x A^0$.*

Moreover, let $A \in C^2(\mathbb{R}^2; \mathbb{R}^3)$ be an arbitrary (not necessarily axially symmetric) external magnetic vector potential, let $B = \text{curl}_x A$ denote the associated magnetic field, and let f denote the corresponding classical solution of the Vlasov–Poisson system (1.1) to the the initial condition $f^{\pm}(0) = f_0^{\pm}$ on $\mathbb{R}^2 \times \mathbb{R}^3$.

Then, for any $\gamma > 4$, there exists a real number $q \in (2, \infty)$ depending only on γ as well as a constant $c > 0$ depending only on $\mathcal{H}(f_0)$, $\|f_0^{\pm}\|_1$, $\|f_0^{\pm}\|_{\infty}$ and γ such that for all $t \geq 0$,

$$\sum_{\pm} \|f^{\pm}(t) - f_0^{\pm}\|_2 \leq \alpha_{f_0} \exp(\beta_{f_0} c(1+t)^{\gamma}) \|B - B^0\|_{L^1(0,t; L^2(\mathbb{R}^2))}$$

where

$$\alpha_{f_0} := 2 \max_{\pm} \|p \times \partial_p f_0^{\pm}\|_{L^{\infty}(\mathbb{R}^2; L^2(\mathbb{R}^3))}, \quad \beta_{f_0} := 2 \max_{\pm} \|\partial_p f_0^{\pm}\|_{L^q(\mathbb{R}^2; L^2(\mathbb{R}^3))}.$$

Proof. The assertion follows directly from Theorem 3.2 and the fact that f_0 is constant in time. \square

5.3 Stability with respect to perturbations of both the initial data and the magnetic vector potential

We can now combine our results to obtain the following theorem.

Theorem 5.4. *We consider the θ -pinch configuration and assume that (S1)–(S4) hold and that ψ^\pm is bounded. Let $f_0 = (f_0^+, f_0^-)$ denote the steady state that was introduced in (5.1). Recall that f_0 corresponds to the axially symmetric external magnetic vector potential $A^0 \in C^2(\mathbb{R}^2; \mathbb{R}^3)$ and the associated magnetic field $B^0 = \text{curl}_x A^0$.*

Moreover, let $A \in C^2(\mathbb{R}^2; \mathbb{R}^3)$ be an arbitrary (not necessarily axially symmetric) external magnetic vector potential, and let $B = \text{curl}_x A$ denote the associated magnetic field. For any given initial data $\mathring{f} = (\mathring{f}^+, \mathring{f}^-) \in X$, let $f = (f^+, f^-)$ and $f_ = (f_*^+, f_*^-)$ denote the corresponding classical solutions of (1.1) to the external magnetic potentials A and A^0 , respectively.*

Furthermore, let $\gamma > 4$ be arbitrary. Then there exist

- *a constant $C > 0$ independent of all the appearing quantities,*
- *constants $c^\pm > 0$ depending only on ϑ^\pm and ψ^\pm ,*
- *a real number $q \in (2, \infty)$ depending only on γ ,*
- *a constant $c_* > 0$ depending only on $\mathcal{H}(\mathring{f})$, $\|\mathring{f}^\pm\|_1$, $\|\mathring{f}^\pm\|_\infty$ and γ ,*

such that for all $t \geq 0$,

$$\begin{aligned}
& \sum_{\pm} \|f^\pm(t) - f_0^\pm\|_2 \\
& \leq \left\{ c^\pm \left[\sum_{\pm} \left(\iint_{\{\pm \mathcal{F}^\pm < 0\}} \left(1 + \frac{\max\{\|\mathring{f}^\pm\|_\infty, \|f_0^\pm\|_\infty\}}{\psi^\pm(\mathcal{F}^\pm, \mathcal{G}^\pm)} + \frac{1}{2}|p|^2 \right) |\mathring{f}^\pm - f_0^\pm| \, dp \, dx \right. \right. \\
& \quad + \iint_{\{\pm \mathcal{F}^\pm < 0\}} \xi(\mathcal{S}(\mathring{f}^\pm 1_{\{\pm \mathcal{F}^\pm < 0\}})) |\mathring{f}^\pm - f_0^\pm| \, dp \, dx \\
& \quad + \iint_{\{\pm \mathcal{F}^\pm > 0\}} \left(\pm \mathcal{F}^\pm + \xi(r_0) + \frac{1}{2}|p|^2 + \xi(\mathcal{R}(\mathring{f}^\pm 1_{\{\pm \mathcal{F}^\pm > 0\}})) \right) |\mathring{f}^\pm - f_0^\pm| \, dp \, dx \Big) \\
& \quad + \frac{1}{2} \|\rho_{\mathring{f}} - \rho_0\|_1 \int_{\mathbb{R}^2} |\rho_{\mathring{f}} - \rho_0| \ln \frac{|\rho_{\mathring{f}} - \rho_0|}{\|\rho_{\mathring{f}} - \rho_0\|_1} \, dx + (C + \ln(2\mathcal{S}(\mathring{f}))) \|\rho_{\mathring{f}} - \rho_0\|_1^2 \Big] \\
& \quad + \|\mathring{f}^\pm - f_0^\pm\|_{L^2(\{\pm \mathcal{F}^\pm \geq 0\})}^2 \Big\}^{1/2} \\
& \quad + \alpha_{f_*}(t) \exp(\beta_{f_*}(t) c_*(1+t)^\gamma) \|B - B^0\|_{L^1(0,t; L^2(\mathbb{R}^2))},
\end{aligned}$$

where

$$\mathcal{F}^\pm := r(p_\varphi \pm A_\varphi^0),$$

$$\begin{aligned}
\mathcal{G}^\pm &:= p_3 \pm A_3^0, \\
a_{f_*}(t) &:= 2 \max_{\pm} \max_{s \in [0, t]} \|p \times \partial_p f_*^\pm(s)\|_{L^\infty(\mathbb{R}^2; L^2(\mathbb{R}^3))}, \\
b_{f_*}(t) &:= 2 \max_{\pm} \max_{s \in [0, t]} \|\partial_p f_*^\pm(s)\|_{L^q(\mathbb{R}^2; L^2(\mathbb{R}^3))}.
\end{aligned}$$

Proof. The assertion follows directly from the decomposition $f - f_0 = (f - f_*) + (f_* - f_0)$ and using the estimates of Theorem 3.2 for $f - f_*$ and of Remark 5.2 for $f_* - f_0$. \square

Remark 5.5. Unfortunately, in Theorem 5.4, the constant c_* depends on the perturbed initial data \mathring{f} , and the functions a_{f_*} and b_{f_*} depend on the solution f_* which corresponds to the perturbed initial data \mathring{f} . However, under suitable assumptions, it is possible to show that these quantities depend only on f_0 , A^0 and γ .

(a) Suppose that

$$|\mathcal{H}(\mathring{f}) - \mathcal{H}(f_0)| \leq \delta, \quad \|\mathring{f}^\pm - f_0^\pm\|_1 \leq \delta, \quad \|\mathring{f}^\pm - f_0^\pm\|_\infty \leq \delta \quad (5.14)$$

for a prescribed real number $\delta > 0$. Then, because of

$$|\mathcal{H}(\mathring{f})| \leq |\mathcal{H}(f_0)| + \delta, \quad \|\mathring{f}^\pm\|_1 \leq \|f_0^\pm\|_1 + \delta, \quad \|\mathring{f}^\pm\|_\infty \leq \|f_0^\pm\|_\infty + \delta,$$

the number c_* can be replaced by a positive constant depending only on $\mathcal{H}(f_0)$, $\|f_0^\pm\|_1$, $\|f_0^\pm\|_\infty$, γ and δ .

(b) Proceeding as in [21, Lemma 6], one can show that there exist increasing functions $\alpha, \beta \in C([0, \infty[)$ depending only on A^0 , $\|\mathring{f}^\pm\|_\infty$, and γ such that

$$a_{f_*}(t) \leq \alpha(t) \quad \text{and} \quad b_{f_*}(t) \leq \beta(t)$$

for all $t \geq 0$. However, we remark that due to the Gronwall argument employed in [21, Lemma 6], these functions α and β exhibit exponential growth. Under the assumptions (5.14), α and β could be replaced by similar functions depending only on A^0 , $\|f_0^\pm\|_\infty$, γ and δ .

Acknowledgement This project has received funding from the European Research Council (ERC) under the European Union's Horizon 2020 research and innovation programme (grant agreement no 678698).

References

- [1] J. Batt. Global symmetric solutions of the initial value problem of stellar dynamics. *J. Differential Equations*, 25(3):342–364, 1977.
- [2] J. Batt and K. Fabian. Stationary solutions of the relativistic Vlasov-Maxwell system of plasma physics. *Chinese Ann. Math. Ser. B*, 14(3):253–278, 1993.
- [3] Yu.O. Belyaeva. Stationary solutions of the Vlasov-Poisson system for two-component plasma under an external magnetic field in a half-space. *Math. Model. Nat. Phenom.*, 12(6):37–50, 2017.
- [4] Yu.O. Belyaeva, B. Gebhard, and A.L. Skubachevskii. A general way to confined stationary vlasov-poisson plasma configurations. *Kinet. Relat. Models*, 14(2):257–282, 2021.

- [5] D. Bian, Y. Guo, and I. Tice. Linear instability of Z-pinch in plasma: inviscid case. *Math. Models Methods Appl. Sci.*, 31(2):409–472, 2021.
- [6] P. Braasch, G. Rein, and J. Vukadinović. Nonlinear stability of stationary plasmas—an extension of the energy-Casimir method. *SIAM J. Appl. Math.*, 59(3):831–844, 1999.
- [7] S. Caprino, G. Cavallaro, and C. Marchioro. Time evolution of a Vlasov–Poisson plasma with magnetic confinement. *Kinet. Relat. Models*, 5(4):729–742, 2012.
- [8] S. Caprino, G. Cavallaro, and C. Marchioro. On a magnetically confined plasma with infinite charge. *SIAM J. Math. Anal.*, 46(1):133–164, 2014.
- [9] S. Caprino, G. Cavallaro, and C. Marchioro. On a Vlasov–Poisson plasma confined in a torus by a magnetic mirror. *J. Math. Anal. Appl.*, 427(1):31–46, 2015.
- [10] S. Caprino, G. Cavallaro, and C. Marchioro. A Vlasov–Poisson plasma with unbounded mass and velocities confined in a cylinder by a magnetic mirror. *Kinet. Relat. Models*, 9(4):657–686, 2016.
- [11] E. Carlen and M. Loss. Competing symmetries, the logarithmic HLS inequality and Onofri’s inequality on S^n . *Geom. Funct. Anal.*, 2(1):90–104, 1992.
- [12] J. Dolbeault. Free energy and solutions of the Vlasov–Poisson–Fokker–Planck system: external potential and confinement (large time behavior and steady states). *J. Math. Pures Appl. (9)*, 78(2):121–157, 1999.
- [13] S.S. Dragomir. *Some Gronwall type inequalities and applications*. Nova Science Publishers, Inc., Hauppauge, NY, 2003.
- [14] R. Duan and T. Yang. Stability of the one-species Vlasov–Poisson–Boltzmann system. *SIAM J. Math. Anal.*, 41(6):2353–2387, 2009/10.
- [15] R. Duan, T. Yang, and C. Zhu. Existence of stationary solutions to the Vlasov–Poisson–Boltzmann system. *J. Math. Anal. Appl.*, 327(1):425–434, 2007.
- [16] R.T. Glassey, J. Schaeffer, and Y. Zheng. Steady states of the Vlasov–Poisson–Fokker–Planck system. *J. Math. Anal. Appl.*, 202(3):1058–1075, 1996.
- [17] Y. Guo and C. Grotta Ragazzo. On steady states in a collisionless plasma. *Comm. Pure Appl. Math.*, 49(11):1145–1174, 1996.
- [18] Y. Guo and G. Rein. Stable steady states in stellar dynamics. *Arch. Ration. Mech. Anal.*, 147(3):225–243, 1999.
- [19] D. Han-Kwan. On the confinement of a Tokamak plasma. *SIAM J. Math. Anal.*, 42(6):2337–2367, 2010.
- [20] D.D. Holm, J.E. Marsden, T. Ratiu, and A. Weinstein. Nonlinear stability of fluid and plasma equilibria. *Phys. Rep.*, 123(1-2):116, 1985.
- [21] P. Knopf. Optimal control of a Vlasov–Poisson plasma by an external magnetic field. *Calc. Var. Partial Differential Equations*, 57(5):134, 37, 2018.
- [22] P. Knopf. Confined steady states of a Vlasov–Poisson plasma in an infinitely long cylinder. *Math. Models Methods Appl. Sci.*, 42(18):6369–6384, 2019.
- [23] R. Kurth. Das Anfangswertproblem der Stelldynamik. *Z. Astrophys.*, 30:213–229, 1952.
- [24] N. S. Landkof. *Foundations of modern potential theory*. Springer-Verlag, New York-Heidelberg, 1972. Translated from the Russian by A. P. Doohovskoy, Die Grundlehren der mathematischen Wissenschaften, Band 180.
- [25] M. Lemou, F. Méhats, and P. Raphaël. Orbital stability of spherical galactic models. *Invent. Math.*, 187(1):145–194, 2012.
- [26] E.H. Lieb and M. Loss. *Analysis*, volume 14 of *Graduate Studies in Mathematics*. American Mathematical Society, Providence, RI, second edition, 2001.
- [27] P.-L. Lions and B. Perthame. Propagation of moments and regularity for the 3-dimensional Vlasov–Poisson system. *Invent. Math.*, 105(2):415–430, 1991.
- [28] T.T. Nguyen, T.V. Nguyen, and W. Strauss. Global magnetic confinement for the 1.5D Vlasov–Maxwell system. *Kinet. Relat. Models*, 8(1):153–168, 2015.
- [29] T.T. Nguyen and W. Strauss. Linear stability analysis of a hot plasma in a solid torus. *Arch. Ration. Mech. Anal.*, 211(2):619–672, 2014.

- [30] K. Pfaffelmoser. Global classical solutions of the Vlasov-Poisson system in three dimensions for general initial data. *J. Differential Equations*, 95(2):281–303, 1992.
- [31] G. Rein. Existence of stationary, collisionless plasmas in bounded domains. *Math. Methods Appl. Sci.*, 15(5):365–374, 1992.
- [32] G. Rein. Non-linear stability for the Vlasov-Poisson system—the energy-Casimir method. *Math. Methods Appl. Sci.*, 17(14):1129–1140, 1994.
- [33] G. Rein. Reduction and a concentration-compactness principle for energy-Casimir functionals. *SIAM J. Math. Anal.*, 33(4):896–912, 2001.
- [34] G. Rein. Stability of spherically symmetric steady states in galactic dynamics against general perturbations. *Arch. Ration. Mech. Anal.*, 161(1):27–42, 2002.
- [35] G. Rein. Collisionless kinetic equations from astrophysics—the Vlasov-Poisson system. In *Handbook of differential equations: evolutionary equations. Vol. III*, Handb. Differ. Equ., pages 383–476. Elsevier/North-Holland, Amsterdam, 2007.
- [36] J. Schaeffer. Global existence of smooth solutions to the Vlasov-Poisson system in three dimensions. *Comm. Partial Differential Equations*, 16(8-9):1313–1335, 1991.
- [37] A.L. Skubachevskii. Vlasov-Poisson equations for a two-component plasma in a homogeneous magnetic field. *Uspekhi Mat. Nauk*, 69(2(416)):107–148, 2014.
- [38] W. Stacey. *Fusion plasma physics*. Physics textbook. Wiley-VCH, 2 edition, 2012.
- [39] J. Weber. Confined steady states of the relativistic Vlasov-Maxwell system in an infinitely long cylinder. *Kinet. Relat. Models*, 13(6):1135–1161, 2020.
- [40] K.Z. Zhang. Linear stability analysis of the relativistic Vlasov-Maxwell system in an axisymmetric domain. *SIAM J. Math. Anal.*, 51(6):4683–4723, 2019.
- [41] D. Zhelyazov, D. Han-Kwan, and J.D.M. Rademacher. Global stability and local bifurcations in a two-fluid model for Tokamak plasma. *SIAM J. Appl. Dyn. Syst.*, 14(2):730–763, 2015.