

Nonabelian Extensions and Factor Systems for the Algebras of Loday

Erik Mainellis

Abstract

Factor systems are a tool for working on the extension problem of algebraic structures such as groups, Lie algebras, and associative algebras. Their applications are numerous and well-known in these common settings. We construct \mathcal{P} algebra analogues to a series of results from W. R. Scott's *Group Theory*, which gives an explicit theory of factor systems for the group case. Here \mathcal{P} ranges over Leibniz, Zinbiel, diassociative, and dendriform algebras, which we dub “the algebras of Loday,” as well as over Lie, associative, and commutative algebras. Fixing a pair of \mathcal{P} algebras, we develop a correspondence between factor systems and extensions. This correspondence is strengthened by the fact that equivalence classes of factor systems correspond to those of extensions. Under this correspondence, central extensions give rise to 2-cocycles while split extensions give rise to (nonabelian) 2-coboundaries.


Contents

1	Introduction	1
2	Preliminaries	3
3	Factor Systems of Leibniz Algebras	4
3.1	Belonging	5
3.2	Equivalence	8
3.3	Split Extensions	11
3.4	Abelian A	12
3.5	Central Extensions	13
4	Factor Systems of Diassociative Algebras	14
5	Factor Systems of Dendriform Algebras	20
6	Cohomology	21

1 Introduction

This paper is something of a tribute to Loday, who first generated interest in Leibniz algebras as a generalization of Lie algebras. Loday also defined dual Leibniz algebras [7], which later took the name Zinbiel algebras based on Loday’s pen name G. W. Zinbiel, or “Leibniz” backwards. He wrote under this name in “Encyclopedia of Types of Algebras 2010” [13], a paper that lists various algebras and some of their properties from an “operadic point of view.” Loday also introduced the notions of diassociative algebras (or associative dialgebras) and dendriform algebras in his

paper “Dialgebras” [8]. The algebras of Loday fit nicely into the following commutative diagram of functors between the categories Zinb, Dend, Com, As, Dias, Lie, and Leib of Zinbiel, Dendriform, Commutative, Associative, Diassociative, Lie, and Leibniz algebras respectively. Said diagram depicts the symmetry of their corresponding operads under the Koszul duality, which falls across a vertical line through the category As. We take this diagram from Loday [8], although it has also appeared in a number of other papers concerning operads.



The objective of this paper is to establish an explicit theory of factor systems, also known as nonabelian 2-cocycles, for the algebras of Loday. Factor systems are a classic tool for working on the extension problem of algebraic structures. These concepts originated in the group setting and can be traced back to Schreier’s 1926 paper [11].¹ In particular, consider a pair of \mathcal{P} algebras A and B (throughout, \mathcal{P} will range over the seven algebras above). The extension problem concerns the classification of all algebras L such that $0 \rightarrow A \rightarrow L \rightarrow B \rightarrow 0$ is an extension of A by B . Factor systems yield an explicit multiplication on L and provide concrete tools for dealing with the relations among L , A , and B . These tools are commonly assumed in the well-known settings, although are rarely justified in detail. Nevertheless, the implications of factor systems are considerable, a fact that has been heavily demonstrated in the settings of associative and Lie algebras (and, of course, groups).

The majority of applications for factor systems have involved the special case of central extensions. The corresponding central factor systems, also known as 2-cocycles, are, for example, used in [2] to classify nilpotent associative algebras up to dimension 4. In [1], the authors develop a method for classifying nilpotent Lie algebras that relies on central factor systems. Central factor systems give rise to the second cohomology group $\mathcal{H}^2(B, A)$ of an algebra B with coefficients in a B -module A , yielding a characterization of central extensions by $\mathcal{H}^2(B, A)$. This, in turn, gives rise to nice exact sequences (see [5] for example). The identities for factor systems of Lie algebras appear frequently in Lie theory. One such instance can be found in a section of Jacobson’s *Lie Algebras* [4] which concerns the theorems of Levi and Malcev-Harish-Chandra. On the other hand, a formulation of nonabelian factor systems for associative algebras is given in [3], where the group $\mathcal{H}_{nab}^2(B, A)$ of nonabelian factor systems is shown to be in 1-1 correspondence with Maurer-Cartan elements of a certain differential graded Lie algebra. Such a feat can also be performed in the Leibniz setting (see [6]).

With these applications in mind, the intent of this paper is to serve as a starting point from which to advance extension theory in the settings of less-developed algebras. We continue to find results, some of which have been listed above, that rely on the assumptions of factor systems but have only been established for certain algebras. Furthermore, it is important to construct the more general nonabelian factor systems, which allows for work on noncentral extensions. Indeed, one of our current efforts involves extensions of nilpotent algebras, for which there are known results in the Lie setting that rely on factor systems. We plan to develop said results for all seven algebras in a future paper.

The final motivation for this paper is to appreciate the structure of factor systems themselves. In particular, consider the study of Leibniz algebras. We observe that much of Leibniz theory involves the generalization of Lie-theoretic results, which often carry over to the Leibniz setting

¹See also the introduction of [9] for some discussion of factor systems and the extension problem.

with only slight adjustments. For example, there are entire proofs in Lie theory which hold for Leibniz algebras merely by replacing the bracket Lie algebra $[A, B]$ with the sum Leibniz algebra $AB + BA$. Factor systems, however, are a case in which generalizing gives rise to considerably more complicated structures. One powerful aspect of our functorial diagram is that each northeast movement along its arrows corresponds to a generalization of algebra type. This fact can be reasoned by seeing each algebra as a special case of its northeast-adjacent category. Besides the usual Lie to Leibniz comparison, any Zinbiel algebra can be seen as a dendriform algebra with multiplications satisfying $x < y = y > x$. Next, any commutative algebra is simply an associative algebra in which $xy = yx$. Finally, any associative algebra can be seen as a diassociative algebra in which $x \dashv y = x \vdash y$. Thus, any result that holds for the Leibniz, diassociative, and dendriform cases must necessarily hold for *all seven algebras*. The definitions of all seven factor systems are provided in this paper for the sake of structural comparison.

To develop a theory of factor systems, we take our methodology from a chapter in W. R. Scott's *Group Theory* [12] which develops and investigates the correspondence between factor systems and extensions of groups. This research began as an effort to develop a Leibniz analogue of said chapter in order to work on extensions of nilpotent algebras. We later discovered that the Leibniz case had already appeared in a 2018 paper [6] where a similar list of identities was obtained for nonabelian 2-cocycles. We then turned to the Zinbiel, diassociative, and dendriform cases, which we have not been able to find in written form. We have decided, however, to review the Leibniz case in explicit detail. The purpose is to provide a systematic approach to this theory as well as a self-contained paper with consistent notation. For the more complicated and obscure algebras, the results are thereby easier to follow.

This paper is structured as follows. For preliminaries, we define each type of algebra under consideration and provide a discussion of general extension theory for arbitrary algebras. The main work begins with the Leibniz analogue of the results in [12]. We next derive the diassociative analogue. These results can also be constructed for dendriform algebras via the same process as the diassociative case, replacing \dashv with $<$ and \vdash with $>$, although the identities which appear are uniquely determined by the different algebra structures. Thus do our results hold for all seven algebras via the above discussions. The definitions of the remaining factor systems are provided in the same sections as their generalizations. We conclude this paper with a brief discussion of the usual Leibniz cohomology and show how $\mathcal{H}^2(B, A)$ characterizes extensions. The paper ends with a list of 2-cocycles and their defining identities. Said identities have appeared in [10], but we collect them here for the sake of completeness.

2 Preliminaries

Let \mathbb{F} be a field. All algebras will be \mathbb{F} -vector spaces equipped with bilinear multiplications which satisfy certain identities. We first recall that a *Lie algebra* L has multiplication which is *alternating* and satisfies the *Jacobi identity*. Respectively, this means that $xx = 0$ and $(xy)z + (yz)x + (zx)y = 0$ for all $x, y, z \in L$. A *Leibniz algebra* L is another type of nonassociative algebra whose multiplication satisfies the *Leibniz identity* $x(yz) = (xy)z + y(xz)$ for all $x, y, z \in L$. It is well-known that any multiplication which is alternating is also skew-symmetric (meaning $xy = -yx$). Under skew-symmetry, the Leibniz identity can be rearranged to form the Jacobi identity, and thus Leibniz algebras are famously seen as the non-anticommutative generalization of Lie algebras.

Definition 1. A *Zinbiel algebra* Z is a nonassociative algebra with multiplication satisfying what we will call the *Zinbiel identity* $(xy)z = x(yz) + x(zx)$ for all $x, y, z \in Z$.

Definition 2. A *diassociative algebra* D is a vector space equipped with two associative bilinear products \dashv and \vdash which satisfy the following identities for all $x, y, z \in D$:

- D1. $x \dashv (y \dashv z) = x \dashv (y \vdash z)$,
- D2. $(x \vdash y) \dashv z = x \vdash (y \dashv z)$,
- D3. $(x \dashv y) \vdash z = (x \vdash y) \vdash z$.

Definition 3. A *dendriform algebra* E is a vector space equipped with two bilinear products $<$ and $>$ which satisfy the following identities for all $x, y, z \in E$:

- E1. $(x < y) < z = x < (y < z) + x < (y > z)$,
- E2. $(x > y) < z = x > (y < z)$,
- E3. $(x < y) > z + (x > y) > z = x > (y > z)$.

We now review extensions. Let \mathcal{P} denote a type of algebra (e.g. Lie, Leibniz, diassociative) and let A and B be arbitrary \mathcal{P} algebras. An *extension of A by B* is a short exact sequence of the form $0 \rightarrow A \xrightarrow{\sigma} L \xrightarrow{\pi} B \rightarrow 0$ where L is a \mathcal{P} algebra and σ and π are *homomorphisms*, i.e. linear maps that preserve the \mathcal{P} structure. A *section* of the extension is a linear map $T : B \rightarrow L$ such that $\pi T = I_B$. Two extensions $0 \rightarrow A \xrightarrow{\sigma_1} L_1 \xrightarrow{\pi_1} B \rightarrow 0$ and $0 \rightarrow A \xrightarrow{\sigma_2} L_2 \xrightarrow{\pi_2} B \rightarrow 0$ of A by B are called *equivalent* if there exists an isomorphism $\tau : L_1 \rightarrow L_2$ such that the diagram

$$\begin{array}{ccccccccc} 0 & \longrightarrow & A & \xrightarrow{\sigma_1} & L_1 & \xrightarrow{\pi_1} & B & \longrightarrow & 0 \\ & & \downarrow \text{id} & & \downarrow \tau & & \downarrow \text{id} & & \\ 0 & \longrightarrow & A & \xrightarrow{\sigma_2} & L_2 & \xrightarrow{\pi_2} & B & \longrightarrow & 0 \end{array}$$

commutes, i.e. if $\tau\sigma_1 = \sigma_2$ and $\pi_2\tau = \pi_1$. The isomorphism τ is called an *equivalence*. An extension $0 \rightarrow A \xrightarrow{\sigma} L \xrightarrow{\pi} B \rightarrow 0$ of A by B is said to be *split* if there exists a homomorphism $T : B \rightarrow L$ which is also a section. An extension $0 \rightarrow A \xrightarrow{\sigma} L \xrightarrow{\pi} B \rightarrow 0$ is called *central* if $\sigma(A)$ is contained in the center $Z(L)$ of L , and *abelian* if L is abelian.

It is readily verified that equivalence of extensions is an equivalence relation. Moreover, if an extension splits, then so does every equivalent extension. Indeed, let $0 \rightarrow A \xrightarrow{\sigma_1} L_1 \xrightarrow{\pi_1} B \rightarrow 0$ be a split extension which is equivalent to another extension $0 \rightarrow A \xrightarrow{\sigma_2} L_2 \xrightarrow{\pi_2} B \rightarrow 0$ via an equivalence τ . Then there is a homomorphism $T_1 : B \rightarrow L_1$ such that $\pi_1 T_1 = I_B$, which implies that $T_2 = \tau T_1$ is a homomorphism from B into L_2 satisfying $\pi_2 T_2 = \pi_2 \tau T_1 = \pi_1 T_1 = I_B$. Finally, since an equivalence is an isomorphism, it is straightforward to verify that extensions which are equivalent to abelian extensions are abelian, and extensions which are equivalent to central extensions are central.

3 Factor Systems of Leibniz Algebras

As with the group case, a factor system of algebras is a tuple of maps together with a set of identities that said maps satisfy. In particular, a Leibniz factor system involves three maps and seven defining identities while a Lie factor system involves only two maps and three identities. The Lie identities are, of course, well-known. The definitions of factor systems for both Lie and Leibniz algebras are stated here for the sake of comparison. Recall that ad^l and ad^r denote the left and right multiplication operators respectively; ad^l is simply called ad in the Lie case since $\text{ad}^r = -\text{ad}$.

Definition 4. Let A and B be Leibniz algebras. A *factor system* of A by B is a tuple of maps (φ, φ', f) where

$$\varphi : B \rightarrow \text{Der}(A) \text{ is linear,}$$

$$\varphi' : B \rightarrow \mathcal{L}(A) \text{ is linear,}$$

$$f : B \times B \rightarrow A \text{ is bilinear}$$

such that

1. $m(\varphi(i)n) = (\varphi'(i)m)n + \varphi(i)(mn)$
2. $m(\varphi'(i)n) = \varphi'(i)(mn) + n(\varphi'(i)m)$
3. $\text{ad}_{f(i,j)}^r + \varphi'(ij) = \varphi'(j)\varphi'(i) + \varphi(i)\varphi'(j)$
4. $\varphi(i)(mn) = (\varphi(i)m)n + m(\varphi(i)n)$
5. $\varphi(i)\varphi(j) = \varphi(ij) + \varphi(j)\varphi(i) + \text{ad}_{f(i,j)}^l$
6. $\varphi(i)\varphi'(j) = \varphi'(j)\varphi(i) + \varphi'(ij) + \text{ad}_{f(i,j)}^r$
7. $f(i, jk) + \varphi(i)f(j, k) = f(ij, k) + \varphi'(k)f(i, j) + f(j, ik) + \varphi(j)f(i, k)$

are satisfied for all $m, n \in A$ and $i, j, k \in B$. Note that the fourth identity allows for $\varphi : B \rightarrow \text{Der}(A)$.

Definition 5. Let A and B be Lie algebras with multiplications denoted by bracket and consider the natural Lie algebra structure on $\text{Der}(A)$ under the commutator bracket. A *factor system* of A by B is a pair (φ, f) of functions

$$\varphi : B \rightarrow \text{Der}(A) \text{ linear,}$$

$$f : B \times B \rightarrow A \text{ bilinear}$$

such that

1. $f(i, j) = -f(j, i)$
2. $\varphi[i, j] = [\varphi(i), \varphi(j)] - \text{ad}_{f(i,j)}$
3. $\varphi(k)f(i, j) + \varphi(i)f(j, k) + \varphi(j)f(k, i) = f([i, j], k) + f([j, k], i) + f([k, i], j)$

for all $i, j, k \in B$.

3.1 Belonging

Our first goal is to construct a correspondence between factor systems and extensions. Consider an extension $0 \rightarrow A \xrightarrow{\sigma} L \xrightarrow{\pi} B \rightarrow 0$ of A by B and a section $T : B \rightarrow L$. Consider also the linear maps $\rho : L \rightarrow \text{Der}(\sigma(A))$ and $\rho' : L \rightarrow \mathcal{L}(\sigma(A))$ defined by $\rho(x) = \text{ad}_x^l|_{\sigma(A)}$ and $\rho'(x) = \text{ad}_x^r|_{\sigma(A)}$ respectively for $x \in L$. Put simply, these maps denote the left and right multiplication operators that act on the image of σ in L . We next use ρ and ρ' to define the maps $P : L \rightarrow \text{Der}(A)$ and $P' : L \rightarrow \mathcal{L}(A)$ by $P(x) = \sigma^{-1}\rho(x)\sigma$ and $P'(x) = \sigma^{-1}\rho'(x)\sigma$ respectively, formalizing a way for L to act on A . To work explicitly with these maps, one computes $P(x)m = \sigma^{-1}\rho(x)\sigma(m) = \sigma^{-1}(x\sigma(m))$ and $P'(x)m = \sigma^{-1}\rho'(x)\sigma(m) = \sigma^{-1}(\sigma(m)x)$ for any $m \in A$ and $x \in L$. The maps φ and φ' of a factor system are ways for B to act on A . It is thus natural to compose P and P' with T , as well as to define f in terms of T , which leads to Definition 6. What follows are two converse results that form the framework for our correspondence.

Definition 6. A factor system (φ, φ', f) of A by B belongs to the extension $0 \rightarrow A \xrightarrow{\sigma} L \xrightarrow{\pi} B \rightarrow 0$ and T if $\varphi = PT$, $\varphi' = P'T$, and $\sigma(f(i, j)) = T(i)T(j) - T(ij)$ for all $i, j \in B$.

Theorem 3.1. Given an extension $0 \rightarrow A \xrightarrow{\sigma} L \xrightarrow{\pi} B \rightarrow 0$ of A by B and section $T : B \rightarrow L$, there exists a unique factor system (φ, φ', f) of A by B belonging to the extension and T .

Proof. Let $\varphi = PT$ and $\varphi' = P'T$. To define f , one notes that $T(i)T(j) - T(ij) \in \ker \pi$ for any $i, j \in B$. By exactness, there exists an element $c_{i,j} \in A$ such that $\sigma(c_{i,j}) = T(i)T(j) - T(ij)$. Let f be defined by $f(i, j) = c_{i,j}$. One may verify that $f : B \times B \rightarrow A$ is bilinear by applying σ to perform the computation and then applying σ^{-1} . It remains to verify that (φ, φ', f) is a factor system. We compute:

1.
$$\begin{aligned} m(\varphi(i)n) &= \sigma^{-1}(\sigma(m)(T(i)\sigma(n))) \\ &= \sigma^{-1}((\sigma(m)T(i))\sigma(n) + T(i)(\sigma(m)\sigma(n))) \\ &= (\varphi'(i)m)n + \varphi(i)(mn), \end{aligned}$$
2.
$$\begin{aligned} m(\varphi'(i)n) &= \sigma^{-1}(\sigma(m)(\sigma(n)T(i))) \\ &= \sigma^{-1}((\sigma(m)\sigma(n))T(i) + \sigma(n)(\sigma(m)T(i))) \\ &= \varphi'(i)(mn) + n(\varphi'(i)m), \end{aligned}$$
3.
$$\begin{aligned} mf(i, j) + \varphi'(ij)m &= \sigma^{-1}(\sigma(m)(T(i)T(j)) - \sigma(m)T(ij) + \sigma(m)T(ij)) \\ &= \sigma^{-1}((\sigma(m)T(i))T(j) + T(i)(\sigma(m)T(j))) \\ &= \sigma^{-1}\rho'(T(j))\sigma(\sigma^{-1}\rho'(T(i))\sigma(m)) + \sigma^{-1}\rho(T(i))\sigma(\sigma^{-1}\rho'(T(j))\sigma(m)) \\ &= \varphi'(j)(\varphi'(i)m) + \varphi(i)(\varphi'(j)m), \end{aligned}$$
4.
$$\begin{aligned} \varphi(i)(mn) &= \sigma^{-1}(T(i)(\sigma(m)\sigma(n))) \\ &= \sigma^{-1}((T(i)\sigma(m))\sigma(n) + \sigma(m)(T(i)\sigma(n))) \\ &= (\varphi(i)m)n + m(\varphi(i)n), \end{aligned}$$
5.
$$\begin{aligned} \varphi(i)(\varphi(j)m) &= \sigma^{-1}(T(i)(T(j)\sigma(m))) \\ &= \sigma^{-1}((T(i)T(j))\sigma(m) - T(ij)\sigma(m) + T(j)(T(i)\sigma(m)) + T(ij)\sigma(m)) \\ &= \sigma^{-1}(\sigma(f(i, j))\sigma(m)) + \sigma^{-1}\rho(T(j))\sigma(\sigma^{-1}\rho(T(i))\sigma(m)) + \sigma^{-1}\rho(T(ij))\sigma(m) \\ &= f(i, j)m + \varphi(j)(\varphi(i)m) + \varphi(ij)m, \end{aligned}$$
6.
$$\begin{aligned} \varphi(i)(\varphi'(j)m) &= \sigma^{-1}(T(i)(\sigma(m)T(j))) \\ &= \sigma^{-1}((T(i)\sigma(m))T(j) + \sigma(m)(T(i)T(j)) + \sigma(m)T(ij) - \sigma(m)T(ij)) \\ &= \sigma^{-1}\rho'(T(j))\sigma(\sigma^{-1}\rho(T(i))\sigma(m)) + \sigma^{-1}\rho'(T(ij))\sigma(m) + \sigma^{-1}(\sigma(m)\sigma(f(i, j))) \\ &= \varphi'(j)(\varphi(i)m) + \varphi'(ij)m + mf(i, j), \end{aligned}$$
7.
$$\begin{aligned} f(i, jk) + \varphi(i)f(j, k) &= \sigma^{-1}(T(i)T(jk) - T(ijk) + T(i)(T(j)T(k)) - T(i)T(jk)) \\ &= \sigma^{-1}(T(ij)T(k) - T((ij)k) + (T(i)T(j))T(k) - T(ij)T(k) \\ &\quad + T(j)T(ik) - T(j(ik)) + T(j)(T(i)T(k)) - T(j)T(ik)) \\ &= \sigma^{-1}(\sigma(f(ij, k)) + \sigma(f(i, j))T(k) + \sigma(f(j, ik)) + T(j)\sigma(f(i, k))) \\ &= f(ij, k) + \varphi'(k)f(i, j) + f(j, ik) + \varphi(j)f(i, k). \end{aligned}$$

□

Theorem 3.2. (Converse to Theorem 3.1) Let (φ, φ', f) be a factor system of A by B and let L denote the vector space $A \oplus B$ with multiplication $(m, i)(n, j) = (mn + \varphi(i)n + \varphi'(j)m + f(i, j), ij)$ for $m, n \in A$ and $i, j \in B$. Let $\sigma : A \rightarrow L$ by $\sigma(m) = (m, 0)$, $\pi : L \rightarrow B$ by $\pi(m, i) = i$, and $T : B \rightarrow L$ by $T(i) = (0, i)$. Then

1. L is a Leibniz algebra.
2. $0 \rightarrow A \xrightarrow{\sigma} L \xrightarrow{\pi} B \rightarrow 0$ is an extension.
3. $\pi T = I = I_B$.
4. The factor system (φ, φ', f) belongs to the extension and T .

Proof. For part 1, the multiplication defined on L is clearly linear, and so it suffices to show that the Leibniz identity holds. One computes²

$$\begin{aligned}
(m, i)((n, j)(p, k)) &= \left(\underbrace{m(np)}_{\text{Leib.}} + \underbrace{m(\varphi(j)p)}_1 + \underbrace{m(\varphi'(k)n)}_2 + \underbrace{mf(j, k)}_3 + \underbrace{\varphi(i)(np)}_4 + \underbrace{\varphi(i)(\varphi(j)p)}_5 \right) \\
&\quad + \underbrace{\varphi(i)(\varphi'(k)n)}_6 + \underbrace{\varphi(i)f(j, k)}_7 + \underbrace{\varphi'(jk)m}_3 + \underbrace{f(i, jk)}_7, \underbrace{ijk}_{\text{Leib.}} \Big) \\
&= \left(\underbrace{(mn)p + n(mp)}_{\text{Leib.}} + \underbrace{(\varphi'(j)m)p + \varphi(j)(mp)}_1 + \underbrace{\varphi'(k)(mn) + n(\varphi'(k)m)}_2 \right) \\
&\quad + \underbrace{\varphi'(k)(\varphi'(j)m) + \varphi(j)(\varphi'(k)m)}_3 + \underbrace{(\varphi(i)n)p + n(\varphi(i)p)}_4 \\
&\quad + \underbrace{f(i, j)p + \varphi(ij)p + \varphi(j)(\varphi(i)p)}_5 + \underbrace{\varphi'(k)(\varphi(i)n) + nf(i, k) + \varphi'(ik)n}_6 \\
&\quad + \underbrace{f(ij, k) + \varphi'(k)f(i, j) + f(j, ik) + \varphi(j)f(i, k)}_7, \underbrace{(ij)k + j(ik)}_{\text{Leib.}} \Big)
\end{aligned}$$

where “Leib.” marks the equalities that follow from the Leibniz identities on A and B , and numbers $1, \dots, 7$ mark the equalities that follow from the axioms of the given factor system. Expanding the sum $((m, i)(n, j))(p, k) + (n, j)((m, i)(p, k))$ via multiplication on L yields the same outcome. Hence L is a Leibniz algebra.

For part 2, we first compute $\sigma(mn) = (mn, 0) = (m, 0)(n, 0) = \sigma(m)\sigma(n)$ and

$$\begin{aligned}
\pi((m, i)(n, j)) &= \pi(mn + \varphi(i)n + \varphi'(j)m + f(i, j), ij) \\
&= ij \\
&= \pi(m, i)\pi(n, j)
\end{aligned}$$

which implies that σ and π are homomorphisms. Moreover, the exactness of $0 \rightarrow A \xrightarrow{\sigma} L \xrightarrow{\pi} B \rightarrow 0$ is trivial. Part 3 is also immediate. For part 4, let $m \in A$ and $i, j \in B$. Then

$$\begin{aligned}
PT(i)m &= \sigma^{-1}((0, i)(m, 0)) \\
&= \sigma^{-1}(\varphi(i)m + f(i, 0), 0) \\
&= \varphi(i)m
\end{aligned}$$

implies that $PT = \varphi$. The equality $P'T = \varphi'$ holds by similar computation. Finally, $\sigma(f(i, j)) = (f(i, j), 0) = (0, i)(0, j) - (0, ij) = T(i)T(j) - T(ij)$. Hence our factor system belongs to the extension and T . \square

²It was from this computation that the axioms of factor systems were chosen.

3.2 Equivalence

We now define a relation between factor systems under which a change in T (to an equivalent extension) results in a change in the corresponding factor system to an equivalent factor system. Such a notion establishes equivalence classes of factor systems that correspond to equivalence classes of extensions. Thus Theorem 3.3 strengthens the correspondence of the first two theorems.

Definition 7. Factor systems (φ, φ', f) and (ψ, ψ', g) of A by B are called *equivalent* if there exists a linear transformation $E : B \rightarrow A$ such that

1. $\psi(i) = \varphi(i) + \text{ad}_{E(i)}^l$,
2. $\psi'(i) = \varphi'(i) + \text{ad}_{E(i)}^r$,
3. $g(i, j) = f(i, j) + \varphi'(j)E(i) + \varphi(i)E(j) + E(i)E(j) - E(ij)$

for all $i, j \in B$. The function E is called an *equivalence*.

Theorem 3.3. *If the factor system $(\varphi_1, \varphi'_1, f_1)$ belongs to the extension $0 \rightarrow A \xrightarrow{\sigma_1} L_1 \xrightarrow{\pi_1} B \rightarrow 0$ and T_1 and the factor system $(\varphi_2, \varphi'_2, f_2)$ belongs to the extension $0 \rightarrow A \xrightarrow{\sigma_2} L_2 \xrightarrow{\pi_2} B \rightarrow 0$ and T_2 , then the factor systems are equivalent if and only if the extensions are equivalent.*

Proof. (\implies) Assume the factor systems are equivalent and let E be the corresponding equivalence. Recall that an equivalence of extensions requires an isomorphism $\tau : L_1 \rightarrow L_2$ such that $\tau\sigma_1 = \sigma_2$ and $\pi_2\tau = \pi_1$. We know that any element in L_1 has a unique representation of the form $T_1(i) + \sigma_1(m)$ for $i \in B$ and $m \in A$. Define $\tau(T_1(i) + \sigma_1(m)) = T_2(i) + \sigma_2(-E(i) + m)$. Clearly τ is linear. To show that τ preserves multiplication, consider elements $a, b \in L_1$ with unique representations $a = T_1(i) + \sigma_1(m)$ and $b = T_1(j) + \sigma_1(n)$. We first compute

$$\begin{aligned} \tau(ab) &= \tau(T_1(i)T_1(j) + \sigma_1(m)T_1(j) + T_1(i)\sigma_1(n) + \sigma_1(m)\sigma_1(n)) \\ &= \tau\left(T_1(ij) + \sigma_1(f_1(i, j) + \varphi'_1(j)m + \varphi_1(i)n + mn)\right) && \text{belonging} \\ &= T_2(ij) + \sigma_2(-E(ij) + f_1(i, j) + \varphi'_1(j)m + \varphi_1(i)n + mn). \end{aligned}$$

On the other hand, one computes

$$\begin{aligned} \tau(a)\tau(b) &= T_2(i)T_2(j) + T_2(i)\sigma_2(-E(j)) + T_2(i)\sigma_2(n) + \sigma_2(-E(i))T_2(j) \\ &\quad + \sigma_2(m)T_2(j) + \sigma_2(-E(i))\sigma_2(-E(j)) + \sigma_2(m)\sigma_2(-E(j)) \\ &\quad + \sigma_2(-E(i))\sigma_2(n) + \sigma_2(m)\sigma_2(n) \\ &= T_2(ij) + \sigma_2(f_2(i, j) - \varphi_2(i)E(j) + \varphi_2(i)n - \varphi'_2(j)E(i) \\ &\quad + \varphi'_2(j)m + E(i)E(j) - mE(j) - E(i)n + mn) && \text{belonging} \\ &= T_2(ij) + \sigma_2(p) \end{aligned}$$

where $p \in A$ is the expression in the argument of σ_2 . Since $T_2(ij)$ is the only T_2 term on both sides, it remains to check the σ_2 parts. Compute

$$\begin{aligned} p &= f_1(i, j) + \varphi'_1(j)E(i) + \varphi_1(i)E(j) + E(i)E(j) - E(ij) && \text{equivalence axiom 3} \\ &\quad - \varphi_1(i)E(j) - E(i)E(j) + \varphi_1(i)n + E(i)n && \text{equivalence axiom 1} \\ &\quad - \varphi'_1(j)E(i) - E(i)E(j) + \varphi'_1(j)m + mE(j) && \text{equivalence axiom 2} \\ &\quad + E(i)E(j) - mE(j) - E(i)n + mn \\ &= f_1(i, j) - E(ij) + \varphi_1(i)n + \varphi'_1(j)m + mn. \end{aligned}$$

Thus τ preserves multiplication. The computation

$$\begin{aligned}\pi(T_1(i) + \sigma_1(m)) &= i \\ &= \pi_2(T_2(i) + \sigma_2(-E(i) + m)) \\ &= \pi_2\tau(T_1(i) + \sigma_1(m))\end{aligned}$$

implies that $\pi_1 = \pi_2\tau$. Finally, $\tau\sigma_1(m) = \sigma_2(m)$ for all $m \in A$ by the definition of τ . Hence $\tau\sigma_1 = \sigma_2$ and the extensions are equivalent.

(\Leftarrow) Conversely, assume that the extensions are equivalent. Then there exists an isomorphism $\tau : L_1 \rightarrow L_2$ such that $\tau\sigma_1 = \sigma_2$ and $\pi_2\tau = \pi_1$. The equality $\pi_1\tau^{-1}T_2(i) = \pi_2T_2(i) = \pi_1T_1(i)$ holds for any $i \in B$, yielding an element $\tau^{-1}T_2(i) - T_1(i) \in \ker \pi_1$. By exactness, $\ker \pi_1 = \text{Im } \sigma_1$, and so there exists an element $n_i \in A$ such that $\tau^{-1}T_2(i) = T_1(i) + \sigma_1(n_i)$. Define $E : B \rightarrow A$ by $E(i) = n_i$. It remains to verify that E is an equivalence. One computes:

1. $\begin{aligned}\varphi_2(i)n &= P_2T_2(i)n - n_in + n_in \\ &= \sigma_2^{-1}(T_2(i)\sigma_2(n) - \sigma_2(n_i)\sigma_2(n)) + E(i)n \\ &= \sigma_1^{-1}\tau^{-1}((\tau T_1(i) + \tau\sigma_1(n_i))\tau\sigma_1(n) - \tau\sigma_1(n_in)) + E(i)n \\ &= \sigma_1^{-1}(T_1(i)\sigma_1(n) + \sigma_1(n_in) - \sigma_1(n_in)) + E(i)n \\ &= \varphi_1(i)n + E(i)n,\end{aligned}$
2. $\begin{aligned}\varphi_2'(j)m &= P_2'T_2(j)m - mn_j + mn_j \\ &= \sigma_2^{-1}(\sigma_2(m)T_2(j) - \sigma_2(m)\sigma_2(n_j)) + mE(j) \\ &= \sigma_1^{-1}\tau^{-1}(\tau\sigma_1(m)(\tau T_1(j) + \tau\sigma_1(n_j)) - \tau\sigma_1(mn_j)) + mE(j) \\ &= \sigma_1^{-1}(\sigma_1(m)T_1(j) - \sigma_1(m)\sigma_1(n_j) - \sigma_1(mn_j)) + mE(j) \\ &= \varphi_1'(j)m + mE(j),\end{aligned}$
3. $\begin{aligned}f_2(i, j) &= \sigma_2^{-1}(T_2(i)T_2(j) - T_2(ij)) \\ &= \sigma_1^{-1}\tau^{-1}((\tau T_1(i) + \tau\sigma_1(n_i))(\tau T_1(j) + \tau\sigma_1(n_j)) - \tau T_1(ij) - \tau\sigma_1(n_{ij})) \\ &= \sigma_1^{-1}(T_1(i)T_1(j) + T_1(i)\sigma_1(n_j) + \sigma_1(n_i)T_1(j) + \sigma_1(n_in_j) - T_1(ij) - \sigma_1(n_{ij})) \\ &= \sigma_1^{-1}(\sigma_1(f_1(i, j)) + \rho(T_1(i))\sigma_1(n_j) + \rho'(T_1(j))\sigma_1(n_i)) + n_in_j - n_{ij} \\ &= f_1(i, j) + \varphi_1'(j)E(i) + \varphi_1(i)E(j) + E(i)E(j) - E(ij).\end{aligned}$

Thus the factor systems are equivalent. \square

Two results follow easily from Theorem 3.3. The proofs are stated as one because they are so short.

Corollary 3.4. *Given an extension $0 \rightarrow A \xrightarrow{\sigma} L \xrightarrow{\pi} B \rightarrow 0$, let $T_1 : B \rightarrow L$ and $T_2 : B \rightarrow L$ be linear maps such that $\pi T_1 = I_B = \pi T_2$. Suppose also that (φ, φ', f) is a factor system of A by B which belongs to the extension and T_1 , and (ψ, ψ', g) is a factor system of A by B which belongs to the extension and T_2 . Then (φ, φ', f) is equivalent to (ψ, ψ', g) .*

Corollary 3.5. *Equivalence of factor systems is an equivalence relation.*

Proof. For Corollary 3.4, note first that any extension of A by B is equivalent to itself. By Theorem 3.3, factor systems belonging to this extension (and differing T_i) are equivalent. Corollary 3.5 follows from Theorem 3.3 and the fact that equivalence of extensions is an equivalence relation. \square

We now look to E . Given equivalent factor systems, there may be multiple equivalences between them. On the other hand, *any* linear transformation $E : B \rightarrow A$ defines an equivalence of factor systems, as demonstrated by Theorem 3.6.

Theorem 3.6. *If (φ, φ', f) is a factor system of A by B and E is a linear transformation from B to A , then there exists a factor system (ψ, ψ', g) such that E is an equivalence of (φ, φ', f) with (ψ, ψ', g) . Furthermore, if E is an equivalence, then (ψ, ψ', g) is unique.*

Proof. Let (ψ, ψ', g) be defined by

- i. $\psi(i) = \varphi(i) + \text{ad}_{E(i)}^l$,
- ii. $\psi'(j) = \varphi'(j) + \text{ad}_{E(j)}^r$,
- iii. $g(i, j) = f(i, j) + \varphi'(j)E(i) + \varphi(i)E(j) + E(i)E(j) - E(ij)$

for $i, j \in B$. It is straightforward to check that $\psi, \psi' : B \rightarrow \mathcal{L}(A)$ are linear transformations and that $g : B \times B \rightarrow A$ is a bilinear form. Now to verify that (ψ, ψ', g) is a factor system.

1. $m(\psi(i)n) = m(\varphi(i)n) + m(E(i)n)$
 $= (\varphi'(i)m)n + \varphi(i)(mn) + (mE(i))n + E(i)(mn)$ f.s. axiom 1, Leibniz identity
 $= (\psi'(i)m)n + \psi(i)(mn),$
2. $m(\psi'(i)n) = m(\varphi'(i)n) + m(nE(i))$
 $= \varphi'(i)(mn) + n(\varphi'(i)m) + (mn)E(i) + n(mE(i))$ f.s. axiom 2, Leibniz identity
 $= \psi'(i)(mn) + n(\psi'(i)m),$
3. $mg(i, j) + \psi'(ij)m = \underbrace{mf(i, j)}_3 + \underbrace{m(\varphi'(j)E(i))}_2 + \underbrace{m(\varphi(i)E(j))}_1 + m(E(i)E(j)) - mE(ij)$
 $\quad + \underbrace{\varphi'(ij)m + mE(ij)}_3$
 $= \underbrace{\varphi'(j)(\varphi'(i)m) + \varphi(i)(\varphi'(j)m)}_{\text{f.s. axiom 3}} + \underbrace{\varphi'(j)(mE(i)) + E(i)(\varphi'(j)m)}_{\text{f.s. axiom 2}}$
 $\quad + \underbrace{(\varphi'(i)m)E(j) + \varphi(i)(mE(j))}_{\text{f.s. axiom 1}} + \underbrace{(mE(i))E(j) + E(i)(mE(j))}_{\text{Leibniz}}$
 $= \psi'(j)(\psi'(i)m) + \psi(i)(\psi'(j)m),$
4. $\psi(i)(mn) = \varphi(i)(mn) + E(i)(mn)$
 $= (\varphi(i)m)n + m(\varphi(i)n) + (E(i)m)n + m(E(i)n)$ f.s. axiom 4, Leibniz
 $= (\psi(i)m)n + m(\psi(i)n),$
5. $\psi(i)(\psi(j)m) = \underbrace{\varphi(i)(\varphi(j)m)}_5 + \underbrace{\varphi(i)(E(j)m)}_4 + \underbrace{E(i)(\varphi(j)m)}_1 + \underbrace{E(i)(E(j)m)}_{\text{Leibniz}}$
 $= \underbrace{f(i, j)m + \varphi(ij)m + \varphi(j)(\varphi(i)m)}_{\text{f.s. axiom 5}} + \underbrace{(\varphi(i)E(j))m + E(j)(\varphi(i)m)}_{\text{f.s. axiom 4}}$
 $\quad + \underbrace{(\varphi'(i)E(i))m + \varphi(j)(E(i)m)}_{\text{f.s. axiom 1}} + \underbrace{(E(i)E(j))m + E(j)(E(i)m)}_{\text{Leibniz}}$
 $\quad + E(ij)m - E(ij)m$
 $= \psi(ij)m + \psi(j)(\psi(i)m) + g(i, j)m,$

$$\begin{aligned}
6. \quad \psi(i)(\psi'(j)m) &= \underbrace{\varphi(i)(\varphi'(j)m)}_6 + \underbrace{\varphi(i)(mE(j))}_4 + \underbrace{E(i)(\varphi'(j)m)}_2 + \underbrace{E(i)(mE(j))}_{\text{Leibniz}} \\
&= \underbrace{\varphi'(j)(\varphi(i)m) + mf(i, j) + \varphi'(ij)m}_6 + \underbrace{(\varphi(i)m)E(j) + m(\varphi(i)E(j))}_4 \\
&\quad + \underbrace{\varphi'(j)(E(i)m) + m(\varphi'(j)E(i))}_2 + \underbrace{(E(i)m)E(j) + m(E(i)E(j))}_{\text{Leibniz}} \\
&\quad + mE(ij) - mE(ij) \\
&= \psi'(j)(\psi(i)m) + \psi'(ij)m + mg(i, j),
\end{aligned}$$

$$\begin{aligned}
7. \quad g(i, jk) + \psi(i)g(j, k) &= \underbrace{f(i, jk)}_7 + \underbrace{\varphi'(jk)E(i)}_3 + \varphi(i)E(jk) + \underbrace{E(i)E(jk) - E(ijk)}_{\text{Leibniz}} \\
&\quad + \underbrace{\varphi(i)f(j, k)}_7 + \underbrace{\varphi(i)(\varphi'(k)E(j))}_6 + \underbrace{\varphi(i)(\varphi(j)E(k))}_5 \\
&\quad + \underbrace{\varphi(i)(E(j)E(k))}_4 - \varphi(i)E(jk) + \underbrace{E(i)f(j, k)}_3 \\
&\quad + \underbrace{E(i)(\varphi'(k)E(j))}_2 + \underbrace{E(i)(\varphi(j)E(k))}_1 + \underbrace{E(i)(E(j)E(k))}_{\text{Leibniz}} \\
&\quad - E(i)E(jk) \\
&= \underbrace{f(ij, k)}_7 + \underbrace{\varphi'(k)E(ij)}_5 + \varphi(ij)E(k) + \underbrace{E(ij)E(k) - E((ij)k)}_{\text{Leibniz}} \\
&\quad + \underbrace{\varphi'(k)f(i, j)}_7 + \underbrace{\varphi'(k)(\varphi'(j)E(i))}_3 + \underbrace{\varphi'(k)(\varphi(i)E(j))}_6 \\
&\quad + \underbrace{\varphi'(k)(E(i)E(j))}_2 - \varphi'(k)E(ij) + \underbrace{f(i, j)E(k)}_5 \\
&\quad + \underbrace{(\varphi'(j)E(i))E(k)}_1 + \underbrace{(\varphi(i)E(j))E(k)}_4 + \underbrace{(E(i)E(j))E(k)}_{\text{Leibniz}} \\
&\quad - E(ij)E(k) + \underbrace{f(j, ik)}_7 + \underbrace{\varphi'(ik)E(j)}_6 + \varphi(j)E(ik) \\
&\quad + \underbrace{E(j)E(ik) - E(j(ik))}_{\text{Leibniz}} + \underbrace{\varphi(j)f(i, k)}_7 + \underbrace{\varphi(j)(\varphi'(k)E(i))}_3 \\
&\quad + \underbrace{\varphi(j)(\varphi(i)E(k))}_5 + \underbrace{\varphi(j)(E(i)E(k))}_1 - \varphi(j)E(ik) \\
&\quad + \underbrace{E(j)f(i, k)}_6 + \underbrace{E(j)(\varphi'(k)E(i))}_2 + \underbrace{E(j)(\varphi(i)E(k))}_4 \\
&\quad + \underbrace{E(j)(E(i)E(k))}_{\text{Leibniz}} - E(j)E(ik) \\
&= g(ij, k) + \psi'(k)g(i, j) + g(j, ik) + \psi(j)g(i, k).
\end{aligned}$$

The unmarked terms cancel in part 7. By construction, the two factor systems are equivalent with E as their corresponding equivalence. It is straightforward to verify the uniqueness of (ψ, ψ', g) . \square

3.3 Split Extensions

Before approaching split extensions, we discuss conditions under which $\varphi : B \rightarrow \text{Der}(A)$ is a homomorphism. Let (φ, φ', f) be a factor system of A by B . By axiom 5 of factor systems, we have $\varphi(i)\varphi(j) = \varphi(ij) + \varphi(j)\varphi(i) + \text{ad}_{f(i,j)}^l$ for all $i, j \in B$. Then $\varphi(ij) = [\varphi(i), \varphi(j)]$ holds if and only if $f(i, j) \in Z^l(A)$ for all $i, j \in B$. Hence φ is a homomorphism if and only if $f : B \times B \rightarrow Z^l(A)$. Furthermore, if $Z^l(A) = 0$, then φ is a homomorphism if and only if $f = 0$. Finally, if A is abelian, this ensures that $\text{ad}_m^l = 0$ for all $m \in A$. Hence axiom 5 of factor systems again implies that φ is a homomorphism.

Recall that if an extension splits, then so does every equivalent extension. We thus say that a factor system *splits* if and only if its corresponding extension splits. It follows that if a factor system splits, then so does every equivalent factor system. Now consider a split extension $0 \rightarrow A \xrightarrow{\sigma} L \xrightarrow{\pi} B \rightarrow 0$ of A by B with associated homomorphism $T : B \rightarrow L$ and let (φ, φ', f) be a factor system belonging to this extension. Then $\sigma(f(i, j)) = T(i)T(j) - T(ij) = 0$ for all $i, j \in B$ which implies that $f = 0$ since σ is injective. Axiom 5 of factor systems then implies that φ is a homomorphism.

The following theorem will be quite useful for later proofs.

Theorem 3.7. *Let (φ, φ', f) be a factor system of A by B . The following are equivalent:*

- a. (φ, φ', f) splits;
- b. (φ, φ', f) is equivalent to some factor system (ψ, ψ', g) such that $g = 0$;
- c. there exists a linear transformation $E : B \rightarrow A$ such that $f(i, j) = -\varphi'(j)E(i) - \varphi(i)E(j) - E(i)E(j) + E(ij)$ for all $i, j \in B$.

Proof. (a. \implies b.) We know (φ, φ', f) belongs to a split extension $0 \rightarrow A \xrightarrow{\sigma} L \xrightarrow{\pi} B \rightarrow 0$. By definition, there is an associated homomorphism $T : B \rightarrow L$ such that $\pi T = I_B$. Hence there exists a factor system (ψ, ψ', g) belonging to $0 \rightarrow A \xrightarrow{\sigma} L \xrightarrow{\pi} B \rightarrow 0$ and T which is equivalent to (φ, φ', f) by Corollary 3.4. Since T is a homomorphism, we have $g = 0$.

(b. \implies c.) Let $E : B \rightarrow A$ be an equivalence of (φ, φ', f) with (ψ, ψ', g) where $g = 0$. The third axiom of equivalence gives $0 = g(i, j) = f(i, j) + \varphi'(j)E(i) + \varphi(i)E(j) + E(i)E(j) - E(ij)$ for all $i, j \in B$, which implies the desired equality.

(c. \implies a.) Let E be as in c. By Theorem 3.6, E is an equivalence of (φ, φ', f) with another factor system (ψ, ψ', g) which belongs to an extension $0 \rightarrow A \xrightarrow{\sigma} L \xrightarrow{\pi} B \rightarrow 0$ and $T : B \rightarrow L$. One has $g(i, j) = f(i, j) + \varphi'(j)E(i) + \varphi(i)E(j) + E(i)E(j) - E(ij) = 0$ by assumption. Then, since $\sigma(g(i, j)) = 0$ for all $i, j \in B$, the third axiom of belonging implies that T is a homomorphism. Also, T is injective since $\pi T = I_B$. Hence the extension splits and, therefore, so does the original factor system. \square

It is clear that every semidirect sum yields a split extension. The converse is also true in that every split extension of A by B is equivalent to a semidirect sum. Indeed, let $0 \rightarrow A \rightarrow L \rightarrow B \rightarrow 0$ be a split extension of A by B . By Theorem 3.7, there is an equivalent extension $0 \rightarrow A \rightarrow L_2 \rightarrow B \rightarrow 0$ with associated linear map $T_2 : B \rightarrow L_2$ and a factor system (ψ, ψ', g) belonging to this extension and T_2 such that $g(i, j) = 0$ for all $i, j \in B$. Thus ψ is a homomorphism. The Leibniz algebra construct in Theorem 3.2 is then a semidirect sum of A by B with factor system (ψ, ψ', g) . By Theorem 3.3, the extension built in Theorem 3.2 is equivalent to $0 \rightarrow A \rightarrow L_2 \rightarrow B \rightarrow 0$ and hence to $0 \rightarrow A \rightarrow L \rightarrow B \rightarrow 0$.

3.4 Abelian A

Let A be an abelian Leibniz algebra and (φ, φ', f) be a factor system of A by B . Then φ is a homomorphism. Moreover, suppose a factor system (ψ, ψ', g) of A by B is equivalent to (φ, φ', f) via equivalence E . Then $\varphi(i) = \psi(i) + \text{ad}_{E(i)}^l = \psi(i)$ and $\varphi'(i) = \psi'(i) + \text{ad}_{E(i)}^r = \psi'(i)$ for all $i \in B$ which implies that $\varphi = \psi$ and $\varphi' = \psi'$. We thus fix φ and φ' and define the following constructs.

Let $\text{Fact}(B, A, \varphi, \varphi')$ be the set of bilinear maps $f : B \times B \rightarrow A$ such that (φ, φ', f) is a factor system and let $\mathcal{T}(B, A, \varphi, \varphi')$ be the set of bilinear maps $f : B \times B \rightarrow A$ such that (φ, φ', f) is a split factor system. We denote by $\text{Ext}(B, A, \varphi, \varphi')$ the set of equivalence classes $\text{Fact}(B, A, \varphi, \varphi') / \mathcal{T}(B, A, \varphi, \varphi')$ with fixed φ and φ' .

Theorem 3.8. *If A is abelian, then*

1. $\text{Fact}(B, A, \varphi, \varphi')$ is an abelian Leibniz algebra,
2. $\mathcal{T}(B, A, \varphi, \varphi')$ is an ideal in $\text{Fact}(B, A, \varphi, \varphi')$,
3. factor systems (φ, φ', f) and (φ, φ', g) are equivalent if and only if f and g are in the same coset of $\text{Fact}(B, A, \varphi, \varphi')$ relative to $\mathcal{T}(B, A, \varphi, \varphi')$,
4. the quotient Leibniz algebra $\text{Ext}(B, A, \varphi, \varphi')$ is in one-to-one correspondence with the set of equivalence classes of extensions to which φ and φ' belong.

Proof. Let $f, g \in \text{Fact}(B, A, \varphi, \varphi')$ and c be a scalar. We know $f - cg : B \times B \rightarrow A$ and want to show that $(\varphi, \varphi', f - cg)$ is a factor system. Axioms 1, 2, and 4 are trivial since they do not involve f or g . Axioms 3, 5, and 6 hold since $\text{ad}_{(f-cg)(i,j)}^l = 0$ and $\text{ad}_{(f-cg)(i,j)}^r = 0$ for any $i, j \in B$. Finally, axiom 7 holds by the following computation:

$$\begin{aligned}
(f - cg)(i, jk) + \varphi(i)(f - cg)(j, k) &= f(i, jk) + \varphi(i)f(j, k) - c(g(i, jk) + \varphi(i)g(j, k)) \\
&= f(ij, k) + \varphi'(k)f(i, j) + f(j, ik) + \varphi(j)f(i, k) \\
&\quad - c(g(ij, k) + \varphi'(k)g(i, j) + g(j, ik) + \varphi(j)g(i, k)) \\
&= (f - cg)(ij, k) + \varphi'(k)(f - cg)(i, j) + (f - cg)(j, ik) \\
&\quad + \varphi(j)(f - cg)(i, k).
\end{aligned}$$

Hence $\text{Fact}(B, A, \varphi, \varphi')$ is a vector space. One easily checks that (φ, φ', fg) is a factor system (here juxtaposition denotes $f(i, j)g(i, j)$); indeed, $fg = 0$ since A is abelian. Thus $\text{Fact}(B, A, \varphi, \varphi')$ is a Leibniz algebra with trivial multiplication.

To show that $\mathcal{T}(B, A, \varphi, \varphi')$ is an ideal, it suffices to verify that it is a subspace. Let $f, g \in \mathcal{T}(B, A, \varphi, \varphi')$. We want to show that $(\varphi, \varphi', f - cg)$ is a split factor system for scalar c . By Theorem 3.7, since (φ, φ', f) and (φ, φ', g) split, there exist linear maps $E_1, E_2 : B \rightarrow A$ such that $f(i, j) = -\varphi(j)E_1(i) - \varphi(i)E_1(j) - E_1(i)E_1(j) + E_1(ij)$ and $g(i, j) = -\varphi(j)E_2(i) - \varphi(i)E_2(j) - E_2(i)E_2(j) + E_2(ij)$ for any $i, j \in B$. Define $E : B \rightarrow A$ by $E = E_1 - cE_2$. Then $(f - cg)(i, j) = -\varphi(j)E(i) + \varphi'(i)E(j) - E(i)E(j) + E(ij)$ and so Theorem 3.7 says that $(\varphi, \varphi', f - cg)$ splits.

Suppose factor systems (φ, φ', f) and (φ, φ', g) are equivalent via $E : B \rightarrow A$. Then $g(i, j) = f(i, j) + \varphi'(j)E(i) + \varphi(i)E(j) + E(i)E(j) - E(ij)$ implies that $(f - g)(i, j) = -\varphi'(j)E(i) - \varphi(i)E(j) - E(i)E(j) + E(ij)$. By Theorem 3.7, factor system $(\varphi, \varphi', f - g)$ splits. Hence $f + \mathcal{T}(B, A, \varphi, \varphi') = g + \mathcal{T}(B, A, \varphi, \varphi')$. Conversely, if $(\varphi, \varphi', f - g)$ is a split factor system, then there exists a linear map $E : B \rightarrow A$ such that $(f - g)(i, j) = -\varphi'(j)E(i) - \varphi(i)E(j) - E(i)E(j) + E(ij)$ for all $i, j \in B$ (by Theorem 3.7). Thus E satisfies the third axiom of equivalence between factor systems (φ, φ', f) and (φ, φ', g) . The first two axioms of equivalence hold trivially with $\varphi = \varphi$ and $\varphi' = \varphi'$ since $\text{ad}_m^l = 0$ and $\text{ad}_m^r = 0$ for all $m \in A$.

The final statement follows from Theorem 3.3 and part 3 above. Indeed, part 3 says that two elements of $\text{Ext}(B, A, \varphi, \varphi')$ are equal if and only if their factor systems with fixed φ, φ' are equivalent, and Theorem 3.3 guarantees that the latter statement is true if and only if the two extensions are equivalent. \square

3.5 Central Extensions

Recall that an extension which is equivalent to a central extension is itself central. One may thus refer to equivalence classes of central extensions and to *central factor systems*, i.e. factor systems that belong to central extensions. Once again, let A be an abelian Leibniz algebra and (φ, φ', f) be a factor system of A by B .

Theorem 3.9. (φ, φ', f) is central if and only if $\varphi = 0$ and $\varphi' = 0$.

Proof. By Theorem 3.2, (φ, φ', f) belongs to an extension $0 \rightarrow A \rightarrow L \rightarrow B \rightarrow 0$ where $L = A \oplus B$ with multiplication $(m, i)(n, j) = (mn + \varphi(i)n + \varphi'(j)m + f(i, j), ij)$. The extension is central if and only if $(m, i)(n, 0) = (mn + \varphi(i)n, 0) = (0, 0)$ and $(n, 0)(m, i) = (nm + \varphi'(i)n, 0) = (0, 0)$ for all $m, n \in A$ and $i \in B$. This happens if and only if φ and φ' are zero. \square

Theorem 3.10. The classes of central extensions of A by B form a Leibniz algebra, denoted $\text{Cext}(B, A)$.

Proof. By Theorem 3.8 and Theorem 3.9, said classes form the Leibniz algebra $\text{Ext}(B, A, 0, 0) =: \text{Cext}(B, A)$. \square

Theorem 3.11. Let A and B be abelian Leibniz algebras and let (φ, φ', f) be a central factor system of A by B . Then (φ, φ', f) belongs to an abelian extension if and only if $f = 0$.

Proof. Since (φ, φ', f) is central, we know $\varphi = \varphi' = 0$. In the forward direction, the factor system belongs to an extension $0 \rightarrow A \rightarrow L \rightarrow B \rightarrow 0$ and section T . Since L and B are both abelian, one has $\sigma(f(i, j)) = T(i)T(j) - T(ij) = 0$ for all $i, j \in B$. Conversely, if $f = 0$, then the construction of L in Theorem 3.2 has multiplication $(m, i)(n, j) = (0, 0)$ for all $m, n \in A$ and $i, j \in B$. \square

4 Factor Systems of Diassociative Algebras

This section mimics the structure of the Leibniz section. We begin by stating the definitions of factor systems (for diassociative algebras and their corresponding special cases) and proceed to construct diassociative analogues of the results from [12].

Definition 8. Let A and B be diassociative algebras. A factor system of A by B is a tuple $(\varphi_{\rightarrow}, \varphi_{\leftarrow}, \varphi'_{\rightarrow}, \varphi'_{\leftarrow}, f_{\rightarrow}, f_{\leftarrow})$ of maps such that $\varphi_{\rightarrow}, \varphi_{\leftarrow}, \varphi'_{\rightarrow}, \varphi'_{\leftarrow} : B \rightarrow \mathcal{L}(A)$ are linear, $f_{\rightarrow}, f_{\leftarrow} : B \times B \rightarrow A$ are bilinear, and the following five sets of identities are satisfied for all $m, n, p \in A$ and $i, j, k \in B$:

1. Those resembling D1:

- (a) $m \dashv (\varphi_{\rightarrow}(j)p) = m \dashv (\varphi_{\leftarrow}(j)p)$
- (b) $m \dashv (\varphi'_{\rightarrow}(k)n) = m \dashv (\varphi'_{\leftarrow}(k)n)$
- (c) $m \dashv f_{\rightarrow}(j, k) + \varphi'_{\rightarrow}(j \dashv k)m = m \dashv f_{\leftarrow}(j, k) + \varphi'_{\leftarrow}(j \dashv k)m$
- (d) $\varphi_{\rightarrow}(i)(n \dashv p) = \varphi_{\rightarrow}(i)(n \vdash p)$
- (e) $\varphi_{\rightarrow}(i)(\varphi_{\rightarrow}(j)p) = \varphi_{\rightarrow}(i)(\varphi_{\leftarrow}(j)p)$
- (f) $\varphi_{\rightarrow}(i)(\varphi'_{\rightarrow}(k)n) = \varphi_{\rightarrow}(i)(\varphi'_{\leftarrow}(k)n)$
- (g) $\varphi_{\rightarrow}(i)f_{\rightarrow}(j, k) + f_{\rightarrow}(i, j \dashv k) = \varphi_{\rightarrow}(i)f_{\leftarrow}(j, k) + f_{\rightarrow}(i, j \vdash k)$

2. Those resembling D2:

- (a) $(\varphi_{\leftarrow}(i)n) \dashv p = \varphi_{\leftarrow}(i)(n \dashv p)$
- (b) $(\varphi'_{\leftarrow}(j)m) \dashv p = m \vdash (\varphi_{\rightarrow}(j)p)$
- (c) $f_{\leftarrow}(i, j) \dashv p + \varphi_{\rightarrow}(i \vdash j)p = \varphi_{\leftarrow}(i)(\varphi_{\rightarrow}(j)p)$
- (d) $\varphi'_{\rightarrow}(k)(m \vdash n) = m \vdash (\varphi'_{\rightarrow}(k)n)$
- (e) $\varphi'_{\rightarrow}(k)(\varphi_{\leftarrow}(i)n) = \varphi_{\leftarrow}(i)(\varphi'_{\rightarrow}(k)n)$
- (f) $\varphi'_{\rightarrow}(k)(\varphi'_{\leftarrow}(j)m) = \varphi'_{\leftarrow}(j \dashv k)m$
- (g) $\varphi'_{\rightarrow}(k)f_{\leftarrow}(i, j) + f_{\leftarrow}(i \vdash j, k) = \varphi_{\leftarrow}(i)f_{\rightarrow}(j, k) + f_{\leftarrow}(i, j \vdash k)$

3. Those resembling D3:

- (a) $(\varphi_{\rightarrow}(i)n) \vdash p = (\varphi_{\leftarrow}(i)n) \vdash p$
- (b) $(\varphi'_{\rightarrow}(j)m) \vdash p = (\varphi'_{\leftarrow}(j)m) \vdash p$

- (c) $f_{\dashv}(i, j) \vdash p + \varphi_{\dashv}(i \dashv j)p = f_{\dashv}(i, j) \vdash p + \varphi_{\dashv}(i \vdash j)p$
- (d) $\varphi'_{\dashv}(k)(m \dashv n) = \varphi'_{\dashv}(k)(m \vdash n)$
- (e) $\varphi'_{\dashv}(k)(\varphi_{\dashv}(i)n) = \varphi'_{\dashv}(k)(\varphi_{\dashv}(i)n)$
- (f) $\varphi'_{\dashv}(k)(\varphi'_{\dashv}(j)m) = \varphi'_{\dashv}(k)(\varphi'_{\dashv}(j)m)$
- (g) $\varphi'_{\dashv}(k)f_{\dashv}(i, j) + f_{\dashv}(i \dashv j, k) = \varphi'_{\dashv}(k)f_{\dashv}(i, j) + f_{\dashv}(i \vdash j, k)$

4. Those resembling the associativity of \dashv :

- (a) $m \dashv (\varphi_{\dashv}(j)p) = (\varphi'_{\dashv}(j)m) \dashv p$
- (b) $m \dashv (\varphi'_{\dashv}(k)n) = \varphi'_{\dashv}(k)(m \dashv n)$
- (c) $m \dashv f_{\dashv}(j, k) + \varphi'_{\dashv}(j \dashv k)m = \varphi'_{\dashv}(k)(\varphi'_{\dashv}(j)m)$
- (d) $\varphi_{\dashv}(i)(n \dashv p) = (\varphi_{\dashv}(i)n) \dashv p$
- (e) $\varphi_{\dashv}(i)(\varphi_{\dashv}(j)p) = \varphi_{\dashv}(i \dashv j)p + f_{\dashv}(i, j) \dashv p$
- (f) $\varphi_{\dashv}(i)(\varphi'_{\dashv}(k)n) = \varphi'_{\dashv}(k)(\varphi_{\dashv}(i)n)$
- (g) $\varphi_{\dashv}(i)f_{\dashv}(j, k) + f_{\dashv}(i, j \dashv k) = \varphi'_{\dashv}(k)f_{\dashv}(i, j) + f_{\dashv}(i \dashv j, k)$

5. Those resembling the associativity of \vdash :

- (a) $m \vdash (\varphi_{\vdash}(j)p) = (\varphi'_{\vdash}(j)m) \vdash p$
- (b) $m \vdash (\varphi'_{\vdash}(k)n) = \varphi'_{\vdash}(k)(m \vdash n)$
- (c) $m \vdash f_{\vdash}(j, k) + \varphi'_{\vdash}(j \vdash k)m = \varphi'_{\vdash}(k)(\varphi'_{\vdash}(j)m)$
- (d) $\varphi_{\vdash}(i)(n \vdash p) = (\varphi_{\vdash}(i)n) \vdash p$
- (e) $\varphi_{\vdash}(i)(\varphi_{\vdash}(j)p) = \varphi_{\vdash}(i \vdash j)p + f_{\vdash}(i, j) \vdash p$
- (f) $\varphi_{\vdash}(i)(\varphi'_{\vdash}(k)n) = \varphi'_{\vdash}(k)(\varphi_{\vdash}(i)n)$
- (g) $\varphi_{\vdash}(i)f_{\vdash}(j, k) + f_{\vdash}(i, j \vdash k) = \varphi'_{\vdash}(k)f_{\vdash}(i, j) + f_{\vdash}(i \vdash j, k)$

Definition 9. Let A and B be associative algebras. A *factor system* of A by B is a tuple of maps (φ, φ', f) where $\varphi, \varphi' : B \rightarrow \mathcal{L}(A)$ are linear, $f : B \times B \rightarrow A$ is bilinear, and

1. $\varphi(i)\varphi(j) = \varphi(ij) + \text{ad}_{f(i,j)}^l$
2. $\varphi'(i)\varphi'(j) = \varphi'(ji) + \text{ad}_{f(i,j)}^r$
3. $\varphi(i)\varphi'(j) = \varphi'(j)\varphi(i)$
4. $\varphi'(i)(mn) = m(\varphi'(i)n)$
5. $\varphi(i)(mn) = (\varphi(i)m)n$
6. $(\varphi'(i)m)n = m(\varphi(i)n)$
7. $\varphi(i)f(j, k) + f(i, jk) = \varphi'(k)f(i, j) + f(ij, k)$

are satisfied for all $m, n \in A$ and $i, j, k \in B$.

Definition 10. Let A and B be commutative algebras. A *factor system* of A by B is a tuple of maps (φ, f) where $\varphi : B \rightarrow \mathcal{L}(A)$ is linear, $f : B \times B \rightarrow A$ is bilinear, and

1. $f(i, j) = f(j, i)$
2. $\varphi(i)\varphi(j) = \varphi(ij) + \text{ad}_{f(i,j)}$
3. $\varphi(i)(mn) = m(\varphi(i)n) = (\varphi(i)m)n$
4. $\varphi(i)f(j, k) + f(i, jk) = \varphi(k)f(i, j) + f(ij, k)$

are satisfied for all $m, n \in A$ and $i, j, k \in B$.

Definition 11. Let A and B be diassociative algebras. A factor system $(\varphi_{\dashv}, \varphi_{\vdash}, \varphi'_{\dashv}, \varphi'_{\vdash}, f_{\dashv}, f_{\vdash})$ of A by B belongs to an extension $0 \rightarrow A \xrightarrow{\sigma} L \xrightarrow{\pi} B \rightarrow 0$ and section T if

$$\begin{aligned} \varphi_{\dashv} &= P_{\dashv}T, & \varphi_{\vdash} &= P_{\vdash}T, \\ \varphi'_{\dashv} &= P'_{\dashv}T, & \varphi'_{\vdash} &= P'_{\vdash}T, \\ \sigma(f_{\dashv}(i, j)) &= T(i) \dashv T(j) - T(i \dashv j), & \sigma(f_{\vdash}(i, j)) &= T(i) \vdash T(j) - T(i \vdash j) \end{aligned}$$

for all $i, j \in B$, where

$$\begin{aligned} P_{\dashv}(x)m &= \sigma^{-1}(x \dashv \sigma(m)), & P_{\vdash}(x)m &= \sigma^{-1}(x \vdash \sigma(m)), \\ P'_{\dashv}(x)m &= \sigma^{-1}(\sigma(m) \dashv x), & P'_{\vdash}(x)m &= \sigma^{-1}(\sigma(m) \vdash x) \end{aligned}$$

for any $x \in L, m \in A$.

Theorem 4.1. Let A and B be diassociative algebras. Given an extension $0 \rightarrow A \rightarrow L \xrightarrow{\pi} B \rightarrow 0$ of A by B and section $T : B \rightarrow L$, there exists a unique factor system $(\varphi_{\dashv}, \varphi_{\vdash}, \varphi'_{\dashv}, \varphi'_{\vdash}, f_{\dashv}, f_{\vdash})$ of A by B belonging to the extension and T .

Proof. Set $\varphi_{\dashv} = P_{\dashv}T, \varphi_{\vdash} = P_{\vdash}T, \varphi'_{\dashv} = P'_{\dashv}T$, and $\varphi'_{\vdash} = P'_{\vdash}T$. Next, it is easily checked that $T(i) \dashv T(j) - T(i \dashv j)$ and $T(i) \vdash T(j) - T(i \vdash j)$ are in the kernel of π for all $i, j \in B$. We define f_{\dashv} and f_{\vdash} by $\sigma(f_{\dashv}(i, j)) = T(i) \dashv T(j) - T(i \dashv j)$ and $\sigma(f_{\vdash}(i, j)) = T(i) \vdash T(j) - T(i \vdash j)$ which are clearly bilinear maps $B \times B \rightarrow A$. It is straightforward to verify that $(\varphi_{\dashv}, \varphi_{\vdash}, \varphi'_{\dashv}, \varphi'_{\vdash}, f_{\dashv}, f_{\vdash})$ is a factor system. \square

Theorem 4.2. Let $(\varphi_{\dashv}, \varphi_{\vdash}, \varphi'_{\dashv}, \varphi'_{\vdash}, f_{\dashv}, f_{\vdash})$ be a factor system of A by B and let L denote the vector space $A \oplus B$ with multiplications

$$\begin{aligned} (m, i) \vdash (n, j) &= (m \vdash n + \varphi_{\vdash}(i)n + \varphi'_{\vdash}(j)m + f_{\vdash}(i, j), i \vdash j), \\ (m, i) \dashv (n, j) &= (m \dashv n + \varphi_{\dashv}(i)n + \varphi'_{\dashv}(j)m + f_{\dashv}(i, j), i \dashv j) \end{aligned}$$

for $m, n \in A$ and $i, j \in B$. Let $\sigma : A \rightarrow L$ by $\sigma(m) = (m, 0)$, $\pi : L \rightarrow B$ by $\pi(m, i) = i$, and $T : B \rightarrow L$ by $T(i) = (0, i)$. Then

1. L is a diassociative algebra.
2. $0 \rightarrow A \xrightarrow{\sigma} L \xrightarrow{\pi} B \rightarrow 0$ is an extension.
3. $\pi T = I = I_B$.
4. The factor system $(\varphi_{\dashv}, \varphi_{\vdash}, \varphi'_{\dashv}, \varphi'_{\vdash}, f_{\dashv}, f_{\vdash})$ belongs to the extension and T .

Proof. It takes five direct computations to verify that the vector space $L = A \oplus B$, with multiplications defined in the statement of the theorem, is a diassociative algebra. In particular, one must check D1, D2, D3, and the associativity of both \dashv and \vdash . Said computations follow via the axioms of factor systems and the diassociative structures on A and B . \square

We now define a notion of equivalence for factor systems so that equivalence classes of factor systems will correspond to those of extensions. The subsequent corollaries hold by the same logic as their Leibniz analogues.

Definition 12. Two factor systems $(\varphi_{\dashv}, \varphi_{\vdash}, \varphi'_{\dashv}, \varphi'_{\vdash}, f_{\dashv}, f_{\vdash})$ and $(\psi_{\dashv}, \psi_{\vdash}, \psi'_{\dashv}, \psi'_{\vdash}, g_{\dashv}, g_{\vdash})$ of A by B are *equivalent* if there exists a linear transformation $E : B \rightarrow A$ such that

1. $\psi_{\dashv}(i) = \varphi_{\dashv}(i) + \text{ad}_{\dashv}^l(E(i)),$
2. $\psi'_{\dashv}(i) = \varphi'_{\dashv}(i) + \text{ad}_{\dashv}^r(E(i)),$
3. $\psi_{\vdash}(i) = \varphi_{\vdash}(i) + \text{ad}_{\vdash}^l(E(i)),$
4. $\psi'_{\vdash}(i) = \varphi'_{\vdash}(i) + \text{ad}_{\vdash}^r(E(i)),$
5. $g_{\dashv}(i, j) = f_{\dashv}(i, j) + \varphi'_{\dashv}(j)E(i) + \varphi_{\dashv}(i)E(j) + E(i) \dashv E(j) - E(i \dashv j),$
6. $g_{\vdash}(i, j) = f_{\vdash}(i, j) + \varphi'_{\vdash}(j)E(i) + \varphi_{\vdash}(i)E(j) + E(i) \vdash E(j) - E(i \vdash j)$

for all $i, j \in B$ where $\text{ad}_{\dashv}^l(E(i))m = E(i) \dashv m$, $\text{ad}_{\vdash}^l(E(i))m = E(i) \vdash m$, $\text{ad}_{\dashv}^r(E(i))m = m \dashv E(i)$, and $\text{ad}_{\vdash}^r(E(i))m = m \vdash E(i)$ for all $m \in A$.

Theorem 4.3. *If the factor system $(\varphi_{\dashv}, \varphi_{\vdash}, \varphi'_{\dashv}, \varphi'_{\vdash}, f_{\dashv}, f_{\vdash})$ belongs to the extension $0 \rightarrow A \xrightarrow{\sigma_1} L_1 \xrightarrow{\pi_1} B \rightarrow 0$ and T_1 and the factor system $(\psi_{\dashv}, \psi_{\vdash}, \psi'_{\dashv}, \psi'_{\vdash}, g_{\dashv}, g_{\vdash})$ belongs to the extension $0 \rightarrow A \xrightarrow{\sigma_2} L_2 \xrightarrow{\pi_2} B \rightarrow 0$ and T_2 , then the factor systems are equivalent if and only if the extensions are equivalent.*

Proof. In the forward direction, one defines τ in the same way as the Leibniz case and computes $\tau(a \dashv b) = \tau(a) \dashv \tau(b)$ and $\tau(a \vdash b) = \tau(a) \vdash \tau(b)$ via the axioms of equivalence for diassociative factor systems. In the other direction, one defines $E(i) = n_i$ where $\tau^{-1}T_2(i) = T_1(i) + \sigma_1(n_i)$. There are six axioms to check when verifying that E is an equivalence of factor systems. Otherwise, the theorem follows by similar logic. \square

Corollary 4.4. *Given an extension $0 \rightarrow A \xrightarrow{\sigma} L \xrightarrow{\pi} B \rightarrow 0$, let $T_1 : B \rightarrow L$ and $T_2 : B \rightarrow L$ be linear maps such that $\pi T_1 = I_B = \pi T_2$. Suppose also that $(\varphi_{\dashv}, \varphi_{\vdash}, \varphi'_{\dashv}, \varphi'_{\vdash}, f_{\dashv}, f_{\vdash})$ is a factor system of A by B which belongs to the extension and T_1 , and $(\psi_{\dashv}, \psi_{\vdash}, \psi'_{\dashv}, \psi'_{\vdash}, g_{\dashv}, g_{\vdash})$ is a factor system of A by B which belongs to the extension and T_2 . Then $(\varphi_{\dashv}, \varphi_{\vdash}, \varphi'_{\dashv}, \varphi'_{\vdash}, f_{\dashv}, f_{\vdash})$ is equivalent to $(\psi_{\dashv}, \psi_{\vdash}, \psi'_{\dashv}, \psi'_{\vdash}, g_{\dashv}, g_{\vdash})$.*

Corollary 4.5. *Equivalence of factor systems is an equivalence relation.*

Theorem 4.6. *If $(\varphi_{\dashv}, \varphi_{\vdash}, \varphi'_{\dashv}, \varphi'_{\vdash}, f_{\dashv}, f_{\vdash})$ is a factor system of A by B and E is a linear transformation from B to A , then there exists a factor system $(\psi_{\dashv}, \psi_{\vdash}, \psi'_{\dashv}, \psi'_{\vdash}, g_{\dashv}, g_{\vdash})$ such that E is an equivalence between them. Furthermore, if E is an equivalence, then $(\psi_{\dashv}, \psi_{\vdash}, \psi'_{\dashv}, \psi'_{\vdash}, g_{\dashv}, g_{\vdash})$ is unique.*

Proof. Define

- i. $\psi_{\dashv}(i) = \varphi_{\dashv}(i) + \text{ad}_{\dashv}^l(E(i)),$
- ii. $\psi_{\vdash}(i) = \varphi_{\vdash}(i) + \text{ad}_{\vdash}^l(E(i)),$
- iii. $\psi'_{\dashv}(i) = \varphi'_{\dashv}(i) + \text{ad}_{\dashv}^r(E(i)),$
- iv. $\psi'_{\vdash}(i) = \varphi'_{\vdash}(i) + \text{ad}_{\vdash}^r(E(i)),$
- v. $g_{\dashv}(i, j) = f_{\dashv}(i, j) + \varphi'_{\dashv}(j)E(i) + \varphi_{\dashv}(i)E(j) + E(i) \dashv E(j) - E(i \dashv j),$
- vi. $g_{\vdash}(i, j) = f_{\vdash}(i, j) + \varphi'_{\vdash}(j)E(i) + \varphi_{\vdash}(i)E(j) + E(i) \vdash E(j) - E(i \vdash j)$

for all $i, j \in B$. It is straightforward to verify that $\psi_{-}, \psi_{+}, \psi'_{-}$, and ψ'_{+} are linear transformations and that g_{-} and g_{+} are bilinear forms. One checks that $(\psi_{-}, \psi_{+}, \psi'_{-}, \psi'_{+}, g_{-}, g_{+})$ is a factor system via the identities of $(\varphi_{-}, \varphi_{+}, \varphi'_{-}, \varphi'_{+}, f_{-}, f_{+})$ and the axioms of diassociative algebras. By construction, the two factor systems are equivalent with E as their corresponding equivalence. It is straightforward to verify uniqueness. \square

Theorem 4.7. *Let $(\varphi_{-}, \varphi_{+}, \varphi'_{-}, \varphi'_{+}, f_{-}, f_{+})$ be a factor system of A by B . The following are equivalent:*

- a. $(\varphi_{-}, \varphi_{+}, \varphi'_{-}, \varphi'_{+}, f_{-}, f_{+})$ splits;
- b. $(\varphi_{-}, \varphi_{+}, \varphi'_{-}, \varphi'_{+}, f_{-}, f_{+})$ is equivalent to some factor system $(\psi_{-}, \psi_{+}, \psi'_{-}, \psi'_{+}, g_{-}, g_{+})$ such that $g_{-} = 0$ and $g_{+} = 0$;
- c. there exists a linear transformation $E : B \rightarrow A$ such that

$$\begin{aligned} f_{-}(i, j) &= -\varphi'_{-}(j)E(i) - \varphi_{-}(i)E(j) - E(i) \dashv E(j) + E(i \dashv j), \\ f_{+}(i, j) &= -\varphi'_{+}(j)E(i) - \varphi_{+}(i)E(j) - E(i) \vdash E(j) + E(i \vdash j). \end{aligned}$$

Proof. (a. \implies b.) We know $(\varphi_{-}, \varphi_{+}, \varphi'_{-}, \varphi'_{+}, f_{-}, f_{+})$ belongs to a split extension $0 \rightarrow A \xrightarrow{\sigma} L \xrightarrow{\pi} B \rightarrow 0$. By definition, there is an associated homomorphism $T : B \rightarrow L$ such that $\pi T = I_B$. Hence there exists a factor system $(\psi_{-}, \psi_{+}, \psi'_{-}, \psi'_{+}, g_{-}, g_{+})$ belonging to the extension and T which is equivalent to $(\varphi_{-}, \varphi_{+}, \varphi'_{-}, \varphi'_{+}, f_{-}, f_{+})$ by Corollary 4.4. Since T is a homomorphism, we have $g_{-} = g_{+} = 0$.

(b. \implies c.) Let $E : B \rightarrow A$ be an equivalence of factor systems $(\varphi_{-}, \varphi_{+}, \varphi'_{-}, \varphi'_{+}, f_{-}, f_{+})$ and $(\psi_{-}, \psi_{+}, \psi'_{-}, \psi'_{+}, g_{-}, g_{+})$ where $g_{-} = g_{+} = 0$. Then $0 = g_{-}(i, j) = f_{-}(i, j) + \varphi'_{-}(j)E(i) + \varphi_{-}(i)E(j) + E(i) \dashv E(j) - E(i \dashv j)$ and $0 = g_{+}(i, j) = f_{+}(i, j) + \varphi'_{+}(j)E(i) + \varphi_{+}(i)E(j) + E(i) \vdash E(j) - E(i \vdash j)$ for all $i, j \in B$ by the axioms of equivalence, which implies the desired equalities.

(c. \implies a.) Let E be as in c. By Theorem 4.6, E is an equivalence of $(\varphi_{-}, \varphi_{+}, \varphi'_{-}, \varphi'_{+}, f_{-}, f_{+})$ with another factor system $(\psi_{-}, \psi_{+}, \psi'_{-}, \psi'_{+}, g_{-}, g_{+})$ which belongs to an extension $0 \rightarrow A \xrightarrow{\sigma} L \xrightarrow{\pi} B \rightarrow 0$ and $T : B \rightarrow L$. One has $g_{-}(i, j) = f_{-}(i, j) + \varphi'_{-}(j)E(i) + \varphi_{-}(i)E(j) + E(i) \dashv E(j) - E(i \dashv j) = 0$ and $g_{+}(i, j) = f_{+}(i, j) + \varphi'_{+}(j)E(i) + \varphi_{+}(i)E(j) + E(i) \vdash E(j) - E(i \vdash j) = 0$ by assumption. Then, since $\sigma(g(i, j)) = 0$ for all $i, j \in B$, the axioms of belonging imply that T is a homomorphism. Also, T is injective since $\pi T = I_B$. Hence the extension splits and, therefore, so does the original factor system. \square

Let A be an abelian diassociative algebra and let $(\varphi_{-}, \varphi_{+}, \varphi'_{-}, \varphi'_{+}, f_{-}, f_{+})$ be a factor system of A by B which is equivalent to another factor system $(\psi_{-}, \psi_{+}, \psi'_{-}, \psi'_{+}, g_{-}, g_{+})$. Since A is abelian, all adjoint operators on A are equal to zero. Thus, by the axioms of equivalence for factor systems, $\varphi_{-} = \psi_{-}$, $\varphi_{+} = \psi_{+}$, $\varphi'_{-} = \psi'_{-}$, and $\varphi'_{+} = \psi'_{+}$. We thus fix the first four maps of factor systems and narrow our focus to pairs of bilinear forms. Let $\text{Fact}(B, A, \varphi_{-}, \varphi_{+}, \varphi'_{-}, \varphi'_{+})$ denote the set of all pairs (f_{-}, f_{+}) such that $(\varphi_{-}, \varphi_{+}, \varphi'_{-}, \varphi'_{+}, f_{-}, f_{+})$ is a factor system and let $\mathcal{T}(B, A, \varphi_{-}, \varphi_{+}, \varphi'_{-}, \varphi'_{+})$ denote the set of all pairs (f_{-}, f_{+}) such that $(\varphi_{-}, \varphi_{+}, \varphi'_{-}, \varphi'_{+}, f_{-}, f_{+})$ is a split factor system. For ease of notation, let φ denote the fixed tuple $(\varphi_{-}, \varphi_{+}, \varphi'_{-}, \varphi'_{+})$ and let

$$(\varphi, f_{-}, f_{+})$$

denote the factor system $(\varphi_{-}, \varphi_{+}, \varphi'_{-}, \varphi'_{+}, f_{-}, f_{+})$. We abbreviate the previous sets by Fact_{φ} and \mathcal{T}_{φ} respectively and denote by

$$\text{Ext}(B, A, \varphi_{-}, \varphi_{+}, \varphi'_{-}, \varphi'_{+})$$

the set of equivalence classes $\text{Fact}_{\varphi} / \mathcal{T}_{\varphi}$. For the rest of this section, we assume A is abelian.

Theorem 4.8. *If A is abelian, then*

1. $\text{Fact}(B, A, \varphi_{-}, \varphi_{+}, \varphi'_{-}, \varphi'_{+})$ is an abelian diassociative algebra,
2. $\mathcal{T}(B, A, \varphi_{-}, \varphi_{+}, \varphi'_{-}, \varphi'_{+})$ is an ideal in $\text{Fact}(B, A, \varphi_{-}, \varphi_{+}, \varphi'_{-}, \varphi'_{+})$,
3. factor systems $(\varphi_{-}, \varphi_{+}, \varphi'_{-}, \varphi'_{+}, f_{-}, f_{+})$ and $(\varphi_{-}, \varphi_{+}, \varphi'_{-}, \varphi'_{+}, g_{-}, g_{+})$ are equivalent if and only if (f_{-}, f_{+}) and (g_{-}, g_{+}) are in the same coset of Fact_{φ} relative to \mathcal{T}_{φ} ,
4. the quotient diassociative algebra $\text{Ext}(B, A, \varphi_{-}, \varphi_{+}, \varphi'_{-}, \varphi'_{+})$ is in one-to-one correspondence with the set of equivalence classes of extensions to which $\varphi_{-}, \varphi_{+}, \varphi'_{-}$, and φ'_{+} belong.

Proof. For (f_{-}, f_{+}) and (g_{-}, g_{+}) in Fact_{φ} , one verifies $(f_{-} - cg_{-}, f_{+} - cg_{+}) \in \text{Fact}_{\varphi}$ via the axioms of the factor systems (φ, f_{-}, f_{+}) and (φ, g_{-}, g_{+}) and the fact that multiplication in A is trivial. For the second statement, it suffices to verify that \mathcal{T}_{φ} is a subspace. Consider elements (f_{-}, f_{+}) and (g_{-}, g_{+}) in \mathcal{T}_{φ} , which form split factor systems (φ, f_{-}, f_{+}) and (φ, g_{-}, g_{+}) respectively. By Theorem 4.7, there exist linear transformations $E_f, E_g : B \rightarrow A$ such that

$$\begin{aligned} f_{-}(i, j) &= -\varphi'_{-}(j)E_f(i) - \varphi_{-}(i)E_f(j) - E_f(i) \dashv E_f(j) + E_f(i \dashv j), \\ f_{+}(i, j) &= -\varphi'_{+}(j)E_f(i) - \varphi_{+}(i)E_f(j) - E_f(i) \vdash E_f(j) + E_f(i \vdash j), \end{aligned}$$

and

$$\begin{aligned} g_{-}(i, j) &= -\varphi'_{-}(j)E_g(i) - \varphi_{-}(i)E_g(j) - E_g(i) \dashv E_g(j) + E_g(i \dashv j), \\ g_{+}(i, j) &= -\varphi'_{+}(j)E_g(i) - \varphi_{+}(i)E_g(j) - E_g(i) \vdash E_g(j) + E_g(i \vdash j). \end{aligned}$$

Letting $E = E_f - cE_g$, one has

$$\begin{aligned} (f_{-} - cg_{-})(i, j) &= -\varphi'_{-}(j)E(i) - \varphi_{-}(i)E(j) - E(i) \dashv E(j) + E(i \dashv j), \\ (f_{+} - cg_{+})(i, j) &= -\varphi'_{+}(j)E(i) - \varphi_{+}(i)E(j) - E(i) \vdash E(j) + E(i \vdash j) \end{aligned}$$

which implies that $(\varphi, f_{-} - cg_{-}, f_{+} - cg_{+})$ splits. For the third statement, one observes that the last two axioms of equivalence for factor systems hold if and only if the third condition of Theorem 4.7 holds for the factor system $(\varphi, f_{-} - g_{-}, f_{+} - g_{+})$. Since A is abelian, adjoint operators on A are trivial. The final statement holds as in the Leibniz analogue. \square

Theorem 4.9. $(\varphi_{-}, \varphi_{+}, \varphi'_{-}, \varphi'_{+}, f_{-}, f_{+})$ is central if and only if $\varphi_{-}, \varphi_{+}, \varphi'_{-}, \varphi'_{+} = 0$.

Proof. By Theorem 4.2, the factor system belongs to an extension $0 \rightarrow A \rightarrow L \rightarrow B \rightarrow 0$ which is central if and only if $(m, i) \dashv (n, 0) = (n, 0) \dashv (m, i) = (m, i) \vdash (n, 0) = (n, 0) \vdash (m, i) = (0, 0)$ for all $m, n \in A$ and $i \in B$. But this happens if and only if $\varphi_{-}, \varphi_{+}, \varphi'_{-}, \varphi'_{+} = 0$. \square

Theorem 4.10. *The classes of central extensions of A by B form a diassociative algebra, denoted $\text{Cext}(B, A)$.*

Proof. By Theorem 4.8 and Theorem 4.9, said classes form a diassociative algebra $\text{Cext}(B, A) := \text{Ext}(B, A, 0, 0, 0, 0)$. \square

Theorem 4.11. *Let A and B be abelian diassociative algebras and let (φ, f_{-}, f_{+}) be a central factor system of A by B . Then (φ, f_{-}, f_{+}) belongs to an abelian extension if and only if $f_{-} = 0$ and $f_{+} = 0$.*

Proof. Since (φ, f_{-}, f_{+}) is central, we know all φ maps are zero. In the forward direction, (φ, f_{-}, f_{+}) belongs to an abelian extension $0 \rightarrow A \rightarrow L \rightarrow B \rightarrow 0$ and section T . Since L and B are both abelian, one has $\sigma(f_{-}(i, j)) = T(i) \dashv T(j) - T(i \dashv j) = 0$ and $\sigma(f_{+}(i, j)) = T(i) \vdash T(j) - T(i \vdash j) = 0$ for all $i, j \in B$. Conversely, if f_{-} and f_{+} are trivial, then the construction of L in Theorem 4.2 has trivial multiplications. \square

5 Factor Systems of Dendriform Algebras

The dendriform versions of these results follow by the same logic as the diassociative case with the substitutions of $<$ and $>$ for multiplications \dashv and \vdash respectively. We note again that Zinbiel algebras are a special case of dendriform algebras, and the identities of factor systems condense along these lines.

Definition 13. Let A and B be dendriform algebras. A *factor system* of A by B is a tuple $(\varphi_<, \varphi_>, \varphi'_<, \varphi'_>, f_<, f_>)$ of maps such that $\varphi_<, \varphi_>, \varphi'_<, \varphi'_> : B \rightarrow \mathcal{L}(A)$ are linear, $f_<, f_> : B \times B \rightarrow A$ are bilinear, and the following three sets of identities are satisfied for all $m, n, p \in A$ and $i, j, k \in B$:

1. Those resembling E1:

- (a) $(\varphi_<(i)n) < p = \varphi_<(i)(n < p) + \varphi_<(i)(n > p)$
- (b) $(\varphi'_<(j)m) < p = m < (\varphi_<(j)p) + m < (\varphi_>(j)p)$
- (c) $f_<(i, j) < p + \varphi_<(i < j)p = \varphi_<(i)(\varphi_<(j)p) + \varphi_<(i)(\varphi_>(j)p)$
- (d) $\varphi'_<(k)(m < n) = m < (\varphi'_<(k)n) + m < (\varphi'_>(k)n)$
- (e) $\varphi'_<(k)(\varphi_<(i)n) = \varphi_<(i)(\varphi'_<(k)n) + \varphi_<(i)(\varphi'_>(k)n)$
- (f) $\varphi'_<(k)(\varphi'_<(j)m) = m < f_<(j, k) + \varphi'_<(j < k)m + m < f_>(j, k) + \varphi'_<(j > k)m$
- (g) $\varphi'_<(k)f_<(i, j) + f_<(i < j, k) = \varphi_<(i)f_<(j, k) + f_<(i, j < k) + \varphi_<(i)f_>(j, k) + f_<(i, j > k)$

2. Those resembling E2:

- (a) $(\varphi_>(i)n) < p = \varphi_>(i)(n < p)$
- (b) $(\varphi'_>(j)m) < p = m > (\varphi_<(j)p)$
- (c) $f_>(i, j) < p + \varphi_<(i > j)p = \varphi_>(i)(\varphi_<(j)p)$
- (d) $\varphi'_<(k)(m > n) = m > (\varphi'_<(k)n)$
- (e) $\varphi'_<(k)(\varphi_>(i)n) = \varphi_>(i)(\varphi'_<(k)n)$
- (f) $\varphi'_<(k)(\varphi'_>(j)m) = \varphi'_>(j < k)m + m > f_<(j, k)$
- (g) $\varphi'_<(k)f_>(i, j) + f_<(i > j, k) = \varphi_>(i)f_<(j, k) + f_>(i, j < k)$

3. Those resembling E3:

- (a) $(\varphi_<(i)n) > p + (\varphi_>(i)n) > p = \varphi_>(i)(n > p)$
- (b) $(\varphi'_<(j)m) > p + (\varphi'_>(j)m) > p = m > (\varphi_>(j)p)$
- (c) $f_<(i, j) > p + \varphi_>(i < j)p + f_>(i, j) > p + \varphi_>(i > j)p = \varphi_>(i)(\varphi_>(j)p)$
- (d) $\varphi'_>(k)(m < n) + \varphi'_>(k)(m > n) = m > (\varphi'_>(k)n)$
- (e) $\varphi'_>(k)(\varphi_<(i)n) + \varphi'_>(k)(\varphi_>(i)n) = \varphi_>(i)(\varphi'_>(k)n)$
- (f) $\varphi'_>(k)(\varphi'_<(j)m) + \varphi'_>(k)(\varphi'_>(j)m) = m > f_>(j, k) + \varphi'_>(j > k)m$
- (g) $\varphi'_>(k)f_<(i, j) + f_>(i < j, k) + \varphi'_>(k)f_>(i, j) + f_>(i > j, k) = \varphi_>(i)f_>(j, k) + f_>(i, j > k)$

Definition 14. Let A and B be Zinbiel algebras. A *factor system* of A by B is a tuple of maps (φ, φ', f) where $\varphi, \varphi' : B \rightarrow \mathcal{L}(A)$ are linear, $f : B \times B \rightarrow A$ is bilinear, and

1. $(\varphi(i)n)p = \varphi(i)(np) + \varphi(i)(pn)$
2. $(\varphi'(j)m)p = m(\varphi(j)p) + m(\varphi'(j)p)$
3. $\varphi'(k)(mn) = m(\varphi(k)n) + m(\varphi'(k)n)$
4. $f(i, j)p + \varphi(ij)p = \varphi(i)(\varphi(j)p) + \varphi(i)(\varphi'(j)p)$
5. $\varphi'(k)(\varphi(i)n) = \varphi(i)(\varphi(k)n) + \varphi(i)(\varphi'(k)n)$
6. $\varphi'(k)(\varphi'(j)m) = mf(j, k) + mf(k, j) + \varphi'(jk)m + \varphi'(kj)m$
7. $\varphi'(k)f(i, j) + f(ij, k) = \varphi(i)f(j, k) + \varphi(i)f(k, j) + f(i, jk) + f(i, kj)$

are satisfied for all $m, n, p \in A$ and $i, j, k \in B$.

6 Cohomology

Given a central extension $0 \rightarrow A \rightarrow L \rightarrow B \rightarrow 0$ of A by B , the general construction of cohomology for Leibniz algebras begins with the set $\mathcal{C}^n(B, A)$ of n -linear maps $f : B \times \cdots \times B \rightarrow A$. Elements of $\mathcal{C}^n(B, A) = \text{Mult}(B \times \cdots \times B, A) \cong \text{Hom}_{\mathbb{F}}(B^{\otimes n}, A)$ are called n -cochains. The usual Leibniz coboundary map $d^n : \mathcal{C}^n(B, A) \rightarrow \mathcal{C}^{n+1}(B, A)$ simplifies (in the central case) to:

$$(d^n f)(x_1, \dots, x_{n+1}) = \sum_{1 \leq i < j \leq n+1} (-1)^i f(x_1, \dots, \hat{x}_i, \dots, x_{j-1}, x_i x_j, x_{j+1}, \dots, x_{n+1})$$

for $f \in \mathcal{C}^n(B, A)$. Note specifically that $(d^2 f)(x, y, z) = -f(xy, z) + f(x, yz) - f(y, xz)$. We denote by $\mathcal{Z}^n(B, A)$ the set of all $f \in \mathcal{C}^n(B, A)$ such that $d^n f = 0$ (n -cocycles) and by $\mathcal{B}^n(B, A)$ the set of all $f \in \mathcal{C}^n(B, A)$ such that $d^{n-1} E = f$ for some $E \in \mathcal{C}^{n-1}(B, A)$ (n -coboundaries). It is well known that $d^n \circ d^{n-1} = 0$ and thus $\mathcal{B}^n(B, A) \subseteq \mathcal{Z}^n(B, A)$. Therefore $\mathcal{H}^n(B, A) = \mathcal{Z}^n(B, A)/\mathcal{B}^n(B, A)$ is the n th cohomology group. In particular, $\mathcal{H}^2(B, A)$ arises from our theory of extensions and factor systems. First recall Theorem 3.8 and the construction $\text{Fact}(B, A, 0, 0)$. Given a central extension $0 \rightarrow A \rightarrow L \rightarrow B \rightarrow 0$, the axioms of the corresponding factor systems become trivial except for the seventh one, which reduces to $f(i, jk) = f(ij, k) + f(j, ik)$, the 2-cocycle identity for Leibniz algebras. Thus $\text{Fact}(B, A, 0, 0)$ is the set of all bilinear maps $f : B \times B \rightarrow A$ such that $d^2 f = 0$. By Theorem 3.7, $\mathcal{T}(B, A, 0, 0)$ is the set of all bilinear $f : B \times B \rightarrow A$ such that $f(i, j) = -E(ij)$ for some linear transformation $E : B \rightarrow A$. These sets are thereby the 2-cocycles and 2-coboundaries of our cohomology respectively. In other words, $\mathcal{Z}^2(B, A) = \text{Fact}(B, A, 0, 0)$ and $\mathcal{B}^2(B, A) = \mathcal{T}(B, A, 0, 0)$. Thus $\text{Cext}(B, A)$ is the second cohomology group $\mathcal{H}^2(B, A)$. We refer the reader to Loday's [8] for constructions of (co)homology in the diassociative and dendriform settings.

In conclusion, each set of 2-cocycle identities is the central simplification of the identities for factor systems. The following table lists these identities for each \mathcal{P} algebra, as well as the total numbers $\mu(\mathcal{P})$ of factor system identities. By construction, each cocycle formulation resembles the defining identities of the corresponding \mathcal{P} structure.

\mathcal{P}	$\mu(\mathcal{P})$	2-cocycle form	2-cocycle Identities
Associative	7	f	$f(ij, k) = f(i, jk)$
Leibniz	7	f	$f(i, jk) = f(ij, k) + f(j, ik)$
Zinbiel	7	f	$f(ij, k) = f(i, jk) + f(i, kj)$
Diassociative	35	(f_{\dashv}, f_{\vdash})	$f_{\dashv}(i, j \dashv k) = f_{\dashv}(i, j \vdash k)$ $f_{\dashv}(i \vdash j, k) = f_{\vdash}(i, j \dashv k)$ $f_{\vdash}(i \dashv j, k) = f_{\vdash}(i \vdash j, k)$ $f_{\dashv}(i, j \dashv k) = f_{\dashv}(i \dashv j, k)$ $f_{\vdash}(i, j \vdash k) = f_{\vdash}(i \vdash j, k)$
Dendriform	21	$(f_{<}, f_{>})$	$f_{<}(i < j, k) = f_{<}(i, j < k) + f_{<}(i, j > k)$ $f_{<}(i > j, k) = f_{>}(i, j < k)$ $f_{>}(i < j, k) + f_{>}(i > j, k) = f_{>}(i, j > k)$
Lie	3	f	$f(i, j) = -f(j, i)$ $f([i, j], k) + f([j, k], i) + f([k, i], j) = 0$
Commutative	4	f	$f(i, j) = f(j, i)$ $f(ij, k) = f(i, jk)$

Acknowledgements

The author would like to thank Ernest Stitzinger for the many helpful discussions.

References

- [1] D. V. Millionshchikov, R. Jimenez. “Geometry of central extensions of nilpotent Lie algebras.” May 2019. arXiv:1905.00214.
- [2] De Graaf, Willem A. “Classification of nilpotent associative algebras of small dimension.” *International Journal of Algebra and Computation* Vol. 28, No. 1 (2018), pp. 133–161.
- [3] Gouray, Jean-Baptiste. “A differential graded Lie algebra approach to non abelian extensions of associative algebras.” 2018. arXiv:1802.04641.
- [4] Jacobson, Nathan. *Lie Algebras*. Dover Publications, Inc. New York. 1962.
- [5] Karel Dekimpe, Manfred Hartl, Sarah Wauters. “An Explicit Seven-Term Exact Sequence for the Cohomology of a Lie Algebra Extension.” *Communications in Algebra* Vol. 44, Issue 3 (2016), pp. 1321-1349, DOI: 10.1080/00927872.2015.1027351.
- [6] Jiefeng Liu, Yunhe Sheng, and Qi Wang. “On non-abelian extensions of Leibniz algebras,” *Communications in Algebra*, Vol. 46, No. 2 (2018), pp. 574-587.
- [7] Loday, Jean-Louis. “Cup-product for Leibniz Cohomology and Dual Leibniz Algebras.” *Mathematica Scandinavica*, Vol. 77, No. 2 (1995), pp. 189-196.
- [8] Loday, Jean-Louis. “Dialgebras,” in *Dialgebras and related operads*, pp.7-66, Lecture Notes in Math., 1763, Springer, Berlin 2001. arXiv:math/0102053.
- [9] R. Brown, T. Porter. “On the Schreier theory of nonabelian extensions: generalisations and computations.” *Proceeding Royal Irish Academy* Vol. 96A (1996), pp. 213–227.
- [10] Rakhimov, Isamidin. “On Extensions of Some Classes of Algebras.” *Algebra, Complex Analysis, and Pluripotential Theory*, pp.131-149. Springer Proceedings in Mathematics & Statistics, Vol. 264. 2018.
- [11] Schreier, O. “Über die Erweiterung von Gruppen, I.” *Monatshefte für Mathematik und Physik* Vol. 34 (1926), pp. 165-180.
- [12] Scott, W. R. *Group Theory*. Dover Publications, Inc. New York. 1964.
- [13] Zinbiel, G. W. “Encyclopedia of Types of Algebras 2010.” arXiv:1101.0267.