On α -points of q-analogs of the Fano plane

Michael Kiermaier * Universität Bayreuth

December 31, 2021

Abstract

Arguably, the most important open problem in the theory of q-analogs of designs is the question for the existence of a q-analog D of the Fano plane. It is undecided for every single prime power value $q \geq 2$.

A point P is called an α -point of D if the derived design of D in P is a geometric spread. In 1996, Simon Thomas has shown that there must always exist at least one non- α -point. For the binary case q=2, Olof Heden and Papa Sissokho have improved this result in 2016 by showing that the non- α -points must form a blocking set with respect to the hyperplanes.

In this article, we show that a hyperplane consisting only of α -points implies the existence of a partition of the symplectic generalized quadrangle W(q) into spreads. As a consequence, the statement of Heden and Sissokho is generalized to all primes q and all even values of q.

1 Introduction

Due to the connection to network coding, the theory of subspace designs has gained a lot of interest recently. Subspace designs are the q-analogs of combinatorial designs and arise by replacing the subset lattice of the finite ambient set V by the subspace lattice of a finite ambient vector space V. Arguably the most important open problem in this field is the question for the existence of a q-analog of the Fano plane, which is a subspace design with the parameters 2- $(7,3,1)_q$. This problem has already been stated in 1972 by Ray-Chaudhuri [3, Problem 28]. Despite considerable investigations, its existence remains undecided for every single order q of the base field.

^{*}Universität Bayreuth, Mathematisches Institut, D-95440 Bayreuth, Germany email: michael.kiermaier@uni-bayreuth.de homepage: http://www.mathe2.uni-bayreuth.de/michaelk/

It has been shown that the smallest instance q=2, the binary q-analog of a Fano plane, can have at most a single nontrivial automorphism [6, 18]. A q-analog of the Fano plane would be the largest possible $[7,4;3]_q$ constant dimension subspace code. However, the hitherto best known sizes of such constant dimension subspace codes still leave considerable gaps, namely 333 vs. 381 in the binary case [14] and 6978 vs. 7651 in the ternary case [16].

Another approach has been the investigation of the derived designs of a putative q-analog D of the Fano plane. A derived design exists for each point $P \in PG(6,q)$ and is always a q-design with the parameters 1- $(6,2,1)_q$, which is the same as a line spread of PG(5,q). Following the notation of [13], a point P is called an α -point of D if the derived design in P is the geometric spread, which is the most symmetric and natural one among the 131044 isomorphism types [21] of such spreads. For highest possible regularity, one would expect all points to be α -points. However, this has been shown to be impossible, so there must always be at least one non- α -point of D [28]. For the binary case q = 2, this result has been improved to the statement that each hyperplane contains at least a non- α -point Heden-Sissokho-2016-ArsComb124:161-164, In other words, the non- α -points of of a binary q-analog of the Fano plane form a blocking set with respect to the hyperplanes.

In this article, α -points will be investigated for general values of q, which will lead to the following theorem.

Theorem 1 Let D be a q-analog of the Fano plane and assume that there exists a hyperplane H such that all points of H are α -points of D. Then the following equivalent statements hold:

- (a) The line set of the symplectic generalized quadrangle W(q) is partitionable into spreads.
- (b) The point set of the parabolic quadric Q(4,q) is partitionable into ovoids.

As a consequence, we will get the following generalization of the result of [13].

Theorem 2 Let D be a q-analog of the Fano plane and q be prime or even. Then each hyperplane contains a non- α -point. In other words, the non- α -points form a blocking set with respect to the hyperplanes.

2 Preliminaries

Throughout the article, $q \neq 1$ is a prime power and V is a vector space over \mathbb{F}_q of finite dimension v.

¹The [7, 4; 3]₃ code of size 6977 in [16] can be extended trivially by a further codeword.

2.1 The subspace lattice

For simplicity, a subspace U of V of dimension $\dim_{\mathbb{F}_q}(U) = k$ will be called a k-subspace. The set of all k-subspaces of V is called the $\operatorname{Gra\betamannian}$ and will be denoted by $\begin{bmatrix} V \\ k \end{bmatrix}_q$. Picking the "best of two worlds", we will prefer the algebraic dimension $\dim_{\mathbb{F}_q}(U)$ over the geometric dimension $\dim_{\mathbb{F}_q}(U) - 1$, but we will otherwise make heavy use of geometric notions, such as calling the 1-subspaces of V points, the 2-subspaces lines, the 3-subspaces planes, the 4-subspaces solids and the (v-1)-subspaces hyperplanes. In fact, the subspace lattice $\mathcal{L}(V)$ consisting of all subspaces of V ordered by inclusion is nothing else than the finite projective geometry $\operatorname{PG}(v-1,q) = \operatorname{PG}(V)$. There are good reasons to consider the subset lattice as a subspace lattice over the unary "field" \mathbb{F}_1 [11].

The number of all k-subspaces of V is given by the Gaussian binomial coefficient

$$\# \begin{bmatrix} V \\ k \end{bmatrix}_q = \begin{bmatrix} v \\ k \end{bmatrix}_q = \begin{cases} \frac{(q^v - 1) \cdots (q^{v-k+1} - 1)}{(q^k - 1) \cdots (q-1)} & \text{if } k \in \{0, \dots, v\}; \\ 0 & \text{otherwise.} \end{cases}$$

The Gaussian binomial coefficient $\begin{bmatrix} v \\ 1 \end{bmatrix}_q$ is also known as the q-analog of the number v and will be abbreviated as $[v]_q$.

For $S \subseteq \mathcal{L}(V)$ and $U, W \in \mathcal{L}(V)$, we will use the abbreviations

$$S|_{U} = \{B \in S \mid U \leq B\},\$$

 $S|_{U} = \{B \in S \mid B \leq W\} \text{ and }$
 $S|_{U} = \{B \in S \mid U \leq B \leq W\}.$

For a point P in a plane E, the set of all lines in E passing through P is known as a line pencil.

The subspace lattice $\mathcal{L}(V)$ is isomorphic to its dual, which arises from $\mathcal{L}(V)$ by reversing the order. Fixing a non-degenerate bilinear form β on V, a concrete isomorphism is given by $U \mapsto U^{\perp}$, where $U^{\perp} = \{\mathbf{x} \in V \mid \beta(\mathbf{x}, \mathbf{u}) = 0 \text{ for all } \mathbf{u} \in U\}$. When addressing the dual of some geometric object in $\mathrm{PG}(V)$, we mean its (element-wise) image under this map. Up to isomorphism, the image does not depend on the choice of β .

2.2 Subspace designs

Definition 2.1 Let t, v, k be integers with $0 \le t \le k \le v - t$ and λ another positive integer. A set $D \subseteq {V \brack k}_q$ is called a t- $(v, k, \lambda)_q$ subspace design if each t-subspace of V is contained in exactly λ elements (called blocks) of D. In the important case $\lambda = 1$, D is called a q-Steiner system.

The earliest reference for subspace designs is [10]. However, the idea is older, since it is stated that "Several people have observed that the concept of a t-design can be generalised [...]". They have also been mentioned in a more general context in [12]. The

²In established symbols like PG(v-1,q), the geometric dimension v-1 is not altered.

first nontrivial subspace designs with $t \geq 2$ has been constructed in [27], and the first nontrivial Steiner system with $t \geq 2$ in [4]. An introduction to the theory of subspace designs can be found at [7], see also [25, Day 4].

Subspace designs are interlinked to the theory of network coding in various ways. To this effect we mention the recently found q-analog of the theorem of Assmus and Mattson [9], and that a t- $(v, k, 1)_q$ Steiner systems provides a $(v, 2(k-t+1); k)_q$ constant dimension network code of maximum possible size.

Classical combinatorial designs can be seen as the limit case q=1 of subspace designs. Indeed, quite a few statements about combinatorial designs have a generalization to subspace designs, such that the case q=1 reproduces the original statement [19, 20, 22, 5].

One example of such a statement is the following [26, Lemma 4.1(1)], see also [19, Lemma 3.6]: If D is a t- $(v, k, \lambda)_q$ subspace design, then D is also an s- $(v, k, \lambda_s)_q$ subspace design for all $s \in \{0, \ldots, t\}$, where

$$\lambda_s \coloneqq \lambda \frac{\begin{bmatrix} v-s \\ t-s \end{bmatrix}_q}{\begin{bmatrix} k-s \\ t-s \end{bmatrix}_q}.$$

In particular, the number of blocks in D equals

$$#D = \lambda_0 = \lambda \frac{{v \brack t}_q}{{k \brack t}_q}.$$

So, for a design with parameters t- $(v, k, \lambda)_q$, the numbers λ_s necessarily are integers for all $s \in \{0, \ldots, t\}$ (integrality conditions). In this case, the parameter set t- $(v, k, \lambda)_q$ is called admissible. It is further called realizable if a t- $(v, k, \lambda)_q$ design actually exists. The smallest admissible parameters of a q-analog of a Steiner system are 2- $(7, 3, 1)_q$, which are the parameters of the q-analog of the Fano plane. This explains the significance of the question of its realizability.

The numbers λ_i can be refined as follows. Let i, j be non-negative integers with $i+j \leq t$ and let $I \in {V \brack i}_q$ and $J \in {V \brack v-j}_q$. By [26, Lemma 4.1], see also [7, Lemma 5], the number

$$\lambda_{i,j} \coloneqq \#D|_I^J = \lambda \frac{\left[\begin{smallmatrix} v-i-j \\ k-i \end{smallmatrix} \right]_q}{\left[\begin{smallmatrix} v-t \\ k-t \end{smallmatrix} \right]_q}$$

only depends on i and j, but not on the choice of I and J. The numbers $\lambda_{i,j}$ are important parameters of a subspace design. A further generalization is given by the intersection numbers in [20].

A nice way to arrange the numbers $\lambda_{i,j}$ is the following triangle form.

$$\lambda_{0,0}$$
 $\lambda_{1,0}$
 $\lambda_{0,1}$
 $\lambda_{2,0}$
 $\lambda_{1,1}$
 $\lambda_{0,2}$
 $\lambda_{1,1}$
 $\lambda_{0,2}$
 $\lambda_{t,0}$
 $\lambda_{t-1,1}$
 $\lambda_{t,0}$
 $\lambda_{t,t-1}$
 $\lambda_{t,t-1}$
 $\lambda_{t,t-1}$
 $\lambda_{t,t-1}$

For a q-analog of the Fano plane, we get:

$$\lambda_{0,0}=q^8+q^6+q^5+q^4+q^3+x^2+1 \\ \lambda_{1,0}=q^4+q^2+1 \qquad \qquad \lambda_{0,1}=q^5+q^3+q^2+1 \\ \lambda_{2,0}=1 \qquad \qquad \lambda_{1,1}=q^2+1 \qquad \qquad \lambda_{0,2}=q^2+1$$

As a consequence of the numbers $\lambda_{i,j}$, the dual design $D^{\perp} = \{B^{\perp} \mid B \in D\}$ is a subspace design with the parameters

$$t - \left(v, v - k, \frac{{v - t \brack k}_q}{{v - t \brack k - t}_q}\right).$$

For a point $P \leq V$, the derived design of D in P is

$$\mathrm{Der}_P(D) = \{B/P \mid B \in D|_P\}.$$

By [19], $\operatorname{Der}_P(D)$ is a subspace design with the parameters (t-1)- $(v-1,k-1,\lambda)_q$. In the case of a q-analog of the Fano plane, $\operatorname{Der}_P(D)$ has the parameters 1-(6, 2, 1).

2.3 Spreads

A 1- $(v, k, 1)_q$ Steiner system \mathcal{S} is just a partition of the point set of V into k-subspaces. These objects are better known under the name (k-1)-spread and have been investigated in geometry well before the emergence of subspace designs. A 1-spread is also called *line spread*.

A set S of k-subspaces is called a *partial* (k-1)-spread if each point is covered by at most one element of S. The points not covered by any element are called *holes*. A recent survey on partial spreads is found in [17].

The parameters $1-(v, k, 1)_q$ are admissible if and only v is divisible by k. In this case, spreads do always exist [24, §VI]. An example can be constructed via field reduction: We consider V as a vector space over \mathbb{F}_{q^k} and set $\mathcal{S} = {V \brack 1}_{q^k}$. Switching back to vector spaces over \mathbb{F}_q , the set \mathcal{S} is a (k-1)-spread of V, known as the *Desarguesian* spread.

A (k-1)-spread S is called *geometric* or *normal* if for two distinct blocks $B, B' \in S$, the set $S|^{B+B'}$ is always a (k-1)-spread of B+B'. In other words, S is geometric if every 2k-subspace of V contains either 0, 1 or $\frac{[2k]_q}{[k]_q} = q^k + 1$ blocks of S. It is not hard to see that the Desarguesian spread is geometric. In fact, it follows from [2, Theorem 2] that a (k-1)-spread is geometric if and only if it is isomorphic to a Desarguesian spreads.

The derived designs of a q-analog of the Fano plane D are line spreads in PG(5, 2). These spreads have been classified in [21] into 131044 isomorphism types. The most symmetric one among these spreads is the Desarguesian spread. Following the notation of [13], a point P is called an α -point of the q-analog of the Fano plane D if the derived design in P is the geometric spread.

2.4 Generalized quadrangles

Definition 2.2 A generalized quadrangle is an incidence structure $Q = (\mathcal{P}, \mathcal{L}, I)$ with a non-empty set of points \mathcal{P} , a non-empty set of lines \mathcal{L} , and an incidence relation $I \subseteq \mathcal{P} \times \mathcal{L}$ such hat

- (i) Two distinct points are incident with at most a line.
- (ii) Two distinct lines are incident with at most one point.
- (iii) For each non-incident point-line-pair (P, L) there is a unique incident point-line-pair (P', L') with P I L' and P' I L.

Generalized quadrangles have been introduced in the more general setting of generalized polygons in [29], as a tool in the theory of finite groups.

A generalized quadrangle $Q = (\mathcal{P}, \mathcal{L}, I)$ is called *degenerate* if there is a point P such that each point of Q is incident with a line through P. If each line of Q is incident with t+1 points, and each point is incident with s+1 lines, we say that Q is of order(s,t). The $dual\ Q^{\perp}$ arises from Q by interchanging the role of the points and the lines. It is again a generalized quadrangle. Clearly, $(Q^{\perp})^{\perp} = Q$, and Q is of order (s,t) if and only if Q^{\perp} is of order (t,s).

Furthermore, Q is said to be *projective* if it is embeddable in some Desarguesian projective geometry. This means that there is a Desarguesian projective geometry with point set \mathcal{P}' , line set \mathcal{L}' , and point-line incidence relation I' such that $\mathcal{P} \subseteq \mathcal{P}'$, $\mathcal{L} \subseteq \mathcal{L}'$ and for all $(P, L) \in \mathcal{P} \times \mathcal{L}$ we have P I L if and only if P' I' L'. The non-degenerate finite projective generalized quadrangles have been classified in [8, Th. 1], see also [23, 4.4.8.]. These are exactly the so-called *classical generalized quadrangles* which are associated to a quadratic form or a symplectic or Hermitean polarity on the ambient geometry, see [23, 3.1.1.].

In this article, two of these classical generalized quadrangles will appear.

- (i) The symplectic generalized quadrangle W(q) [23, 3.1.1 (iii)], consisting of the set of points of PG(3,q) together with the totally isotropic lines with respect to a symplectic polarity. Taking the geometry as $PG(\mathbb{F}_q^4)$, the symplectic polarity can be represented by the alternating bilinear form $\beta(\mathbf{x},\mathbf{y}) = x_1y_2 x_2y_1 + x_3y_4 x_4y_3$. The configuration of the lines \mathcal{L} in PG(3,q) is also known as a (general) linear complex of lines, see [23, 3.1.1 (iii)] or [15, Th. 15.2.13]. Under the Klein correspondence, \mathcal{L} is a non-tangent hyperplane section of the Klein quadric.
- (ii) The second one is the parabolic quadric Q(4,q), whose points \mathcal{P} are the zeroes of a parabolic quadratic form in PG(4,q), and whose lines are all the lines contained in \mathcal{P} . Taking the geometry as $PG(\mathbb{F}_q^5)$, the parabolic quadratic form can be represented by $q(\mathbf{x}, \mathbf{y}) = x_1 x_2 + x_3 x_4 + x_5^2$.

Both W(q) and Q(4,q) are of order (q,q). By [23, 3.2.1] they are duals of each other, meaning that $W(q)^{\top} \cong Q(4,q)$.

Let $Q = (\mathcal{P}, \mathcal{L}, I)$ be a generalized quadrangle. As in projective geometries, a set $\mathcal{S} \subseteq \mathcal{L}$ is called a *spread* of Q if each point of Q is incident with a unique line in \mathcal{S} . Dually, a set $\mathcal{O} \subseteq \mathcal{P}$ is called an *ovoid* of Q if each line of Q is incident with a unique point in \mathcal{O} . Clearly, the spreads of Q bijectively correspond to the ovoids of Q^{\perp} . This already shows the equivalence of Part (a) and (b) in Theorem 1.

3 Proof of the theorems

For the remainder of the article, we fix v = 7 and assume that $D \subseteq {V \brack 7}_q$ is a q-analog of the Fano plane. The numbers $\lambda_{i,j}$ are defined as in Section 2.2.

By the design property, the intersection dimension of two distinct blocks $B, B' \in D$ is either 0 or 1. So by the dimension formula, $\dim(B+B') \in \{5,6\}$. Therefore two distinct blocks contained in a common 5-space always intersect in a point. Moreover, a solid S of V contains either a single block of no block at all. We will call S a rich solid in the former case and a poor solid in the latter.

Remark 3.1 By [20, Remark 4.2], the poor solids form a dual 2- $(7,3,q^4)_q$ subspace design. By the above discussion, the $\lambda_{0,2} = q^2 + 1$ blocks in any 5-subspace F form dual partial spread in F. The poor solids contained in F are exactly the holes of that partial spread.

We will call a 5-subspace F such that all the $\lambda_{0,2} = q^2 + 1$ blocks in F pass through a common point P a β -flat with focal point P.

Lemma 3.2 The focal point of a β -flat is uniquely determined.

Proof. Assume that $P \neq Q$ are focal points of a β -flat F. Then all $\lambda_{0,2} = q^2 + 1 > 1$ blocks in F pass through the line P + Q, contradicting the Steiner system property. \square

Lemma 3.3 Let H be a hyperplane and P a point in H. Then P is the focal point of at most one β -flat in H.

Proof. There are $\lambda_{1,1} = q^2 + 1$ blocks in H passing through P. For any β -flat F < H with focal point P, all these blocks are contained in F.

Now assume that there are two such β -flats $F \neq F'$. Then the $q^2 + 1 > 1$ blocks in $D|_P^H$ are contained in $F \cap F'$. This is a contradiction, since $\dim(F \cap F') \leq 4$ and any solid contains at most a single block.

Lemma 3.4 Let $F \in {V \brack 5}_q$ be a β -flat with focal point P.

- (a) Each point $Q \neq P$ is covered by a unique block in F. (In other words: $D|^F/P$ is a line spread of $F/P \cong PG(3,q)$)
- (b) A solid S of F is poor if and only if it does not contain P.
- (c) For all poor solids S of F, the set $\{B \cap S \mid B \in D|^F\}$ is a line spread of S.

Proof. Part (a): As the blocks in $D|^F$ intersect each other only in the point P, the number of points in $\begin{bmatrix} F \\ 1 \end{bmatrix}_q \setminus \{P\}$ covered by these blocks is $(q^2 + 1)(\begin{bmatrix} 3 \\ 1 \end{bmatrix}_q - 1) = q^4 + q^3 + q^2 + q = \begin{bmatrix} 5 \\ 1 \end{bmatrix}_q - 1$. Therefore, all points in F different from P are covered by a single block in D_F .

Part (b): The number of solids in F containing one of the q^2+1 blocks in F is $(q^2+1)\cdot {5-3\brack 4-3}_q=(q^2+1)(q+1)=q^3+q^2+q+1$. These solids are rich. Moreover, the q^4 solids in F not containing P do not contain a block, so they are poor. As $q^4+(q^3+q^2+q+1)={5\brack 4}_q$ is already the total number of solids in F, the poor solids in F are precisely those not containing P.

Part (c): Let S be a poor solid of F. For every block B in F we have $\dim(B \cap S) \leq 2$ as S is poor, and moreover $\dim(B \cap S) \geq \dim(B) + \dim(S) - \dim(F) = 3 + 4 - 5 = 2$ by the dimension formula. So for all blocks B in F we get that B + S = F and $B \cap S$ is a line. By parts (a) and (b), every point of the poor solid S is contained in a unique block in F. Hence $\{B \cap S \mid B \in D \text{ and } B + S = F\}$ is a line spread of S.

Lemma 3.5 Let B, B' be two distinct blocks which intersect in an α -point P. Then P is the focal point of the β -flat B + B'.

Proof. Since $P = B \cap B'$ is a point, F = B + B' is a 5-subspace. Since P is an α -point, we have that $\{B''/P \mid B'' \in D|_P^F\}$ is a line spread of $F/P \cong \mathbb{F}_q^4$. Such a line spread contains $\frac{[4]_q}{[2]_q} = q^2 + 1$ lines, so F contains $q^2 + 1$ blocks passing through P. However, the total number of blocks contained in F is only $\lambda_{0,2} = q^2 + 1$, so F is a β -flat with focal point P.

Lemma 3.6 Let F be a 5-subspace such that all points of F are α -points. Then F is a β -flat.

Proof. The 5-subspace F contains $\lambda_{0,2} = q^2 + 1 > 1$ blocks. Let B, B' be two distinct blocks in F. Then $P = B \cap B'$ is a point and F = B + B'. By the precondition, P is an α -point, so by Lemma 3.5, P is the focal point of the β -flat F.

Remark 3.7 The statement of Lemma 3.6 is still true if F contains a single non- α -point Q. Then either all blocks contained in F pass through Q, or there are two distinct blocks B, B' in F such that $P = B \cap B' \neq Q$. In the latter case, all blocks pass through the α -point P as in the proof of Lemma 3.6.

Lemma 3.8 Let H be a hyperplane and P an α -point contained in H. Then H contains a unique β -flat whose focal point is P.

Proof. There are $\lambda_{1,1} = q^2 + 1 > 1$ blocks in H containing P. Let $B, B' \in D|_P^H$. Then $P = B \cap B'$. By Lemma 3.5, the α -point P is the focal point of the β -flat F = B + B'. By Lemma 3.3, the β -flat F is unique.

³Remember that a solid cannot contain 2 blocks.

Now we fix a hyperplane H of V and assume that all its points are α -points.

By Lemma 3.6, every 5-subspace F is a β -flat. We denote its unique focal point by $\alpha(F)$. Moreover by Lemma 3.8, each point P of H is the focal point of a unique β -flat F in H. We will denote this β -flat by $\beta(P)$. Clearly, the mappings

$$\alpha: \begin{bmatrix} H \\ 5 \end{bmatrix}_q \to \begin{bmatrix} H \\ 1 \end{bmatrix}_q \quad \text{and} \quad \beta: \begin{bmatrix} H \\ 1 \end{bmatrix}_q \to \begin{bmatrix} H \\ 5 \end{bmatrix}_q$$

are inverse to each other. So they provide a bijective correspondence between the points and the 5-subspaces of V.

Lemma 3.9 Let B be a block in H.

- (a) For all points P of B, $B \leq \beta(P)$.
- (b) For all 5-subspaces F in H containing B, $\alpha(F) \leq B$.

Proof. For part (a), let P be a point on B. There are $\lambda_{1,1} = q^2 + 1$ blocks in H passing through P, which equals the number $\lambda_{0,2}$ of blocks in $\beta(P)$ (which all pass through P). Therefore, $B \leq \beta(P)$.

For part (b), let F be a 5-subspace containing B. All blocks in F pass through its focal point $\alpha(F)$.

By Lemma (b), each 5-subspace and a fortiori the hyperplane H contains a poor solid S. This poor solid S is fixed for the remainder of this article.

The set of $\begin{bmatrix} 6-4 \\ 5-4 \end{bmatrix}_q = q+1$ intermediate 5-subspaces F with S < F < H will be denoted by \mathcal{F} . For each $F \in \mathcal{F}$, the set $\mathcal{L}_F := \{B \cap S \mid B \in D \text{ and } B \in D|^F\}$ is a line spread of S by Lemma 3.4(c).

Lemma 3.10 The line spreads \mathcal{L}_F with $F \in \mathcal{F}$ are pairwise disjoint.

Proof. Let $L \in \mathcal{L}_F \cap \mathcal{L}_{F'}$ with $F, F' \in \mathcal{F}$. Then $L = B \cap S = B' \cap S$ with $B \in \mathcal{L}_F$ and $B' \in \mathcal{L}_{F'}$. So B and B' are two blocks passing through the same line L. The Steiner system property gives B = B'. Hence F = B + S = B' + S = F'.

Now let $\mathcal{L} = \bigcup_{F \in \mathcal{F}} \mathcal{L}_F$.

Lemma 3.11 The set \mathcal{L} consists of $q^3 + q^2 + q + 1$ lines of S and is partitionable into q+1 line spreads of S.

Proof. By Lemma 3.10, the sets \mathcal{L}_F are pairwise disjoint, so \mathcal{L} is a set of $\#\mathcal{F} \cdot \#D|^F = (q+1)(q^2+1) = q^3+q^2+q+1$ lines in S which admits a partition into the q+1 line spreads \mathcal{L}_F with $F \in \mathcal{F}$.

Lemma 3.12 For each point P of S, $\mathcal{L}|_P$ is a line pencil in the plane $E_P = \beta(P) \cap S$.

Proof. Let P be a point in S.

By Lemma 3.4(b), the poor solid S is not contained in the 5-subspace $\beta(P)$. Therefore, $\dim(\beta(P)\cap S) \leq 3$. On the other hand, as both S and $\beta(P)$ are contained in H, we have $\dim(\beta(P)+S) \leq \dim(H) = 6$ and therefore by the dimension formula $\dim(\beta(P)\cap S) = \dim(\beta(P)) + \dim(S) - \dim(\beta(P) + S) \geq 3$. Hence $E_P = \beta(P) \cap S$ is a plane.

Let $L \in \mathcal{L}|_P$. Then there is a block $B \in D|^H$ with $B \cap S = L$. By Lemma 3.9(a), $B \leq \beta(P)$. So $L = B \cap S \leq \beta(P) \cap S = E_P$. As the disjoint union of q + 1 line spreads of S, \mathcal{L} contains q + 1 lines passing through P. Therefore, these lines form a line pencil in E_P through P.

Lemma 3.13 The incidence structure $\binom{S}{1}_q, \mathcal{L}, \subseteq$ is a projective generalized quadrangle of order (s,t)=(q,q).

Proof. Clearly, every line in \mathcal{L} contains q+1 points in S. By Lemma 3.11, through every point in S there pass q+1 lines in \mathcal{L} . Now let P be a point in S and $L \in \mathcal{L}$ not containing P.

By Lemma 3.10, there is a unique $F \in \mathcal{F}$ with $L \in \mathcal{L}_F$, and there is a line $L'' \in \mathcal{L}_F$ passing through P. By Lemma 3.12, $L'' < E_P$, so we get $L \not< E_P$ as otherwise L and L'' would be distinct intersecting lines in the spread \mathcal{L}_F . Moreover, L and E_P are both contained in S, so they cannot have trivial intersection. Therefore $L \cap E_P$ is a point.

Now let $P' \in {S \brack 1}_q$ and $L' \in \mathcal{L}$ with $L \cap L' = P'$ and P + P' = L'. Then L' is a line through P, so $L' < E_P$. So necessarily $P' = E_P \cap L$ and L' = P + P', showing that P' and L' are unique.

By Lemma 3.12 indeed $L' \in \mathcal{L}$, as P + P' is a line in E_P containing P. This shows that P' and L' do always exist and therefore, the incidence structure $\binom{S}{1}$, \mathcal{L} is a generalized quadrangle of order (q, q).

Lemma 3.14 $\left(\begin{bmatrix}S\\1\end{bmatrix}_q,\mathcal{L}\right)$ is isomorphic to W(q).

Proof. By Lemma 3.13 we know that $Q = {S \brack 1}_q, \mathcal{L}, \subseteq$ is a finite generalized quadrangle of order (s,t) = (q,q) embedded in PG(S). By the classification in [8, Theorem 1] (see also [23, p. 4.4.8]), we know that Q is a finite classical generalized quadrangle which are listed in [23, p. 3.1.2]. Comparing the orders and the dimension of the ambient geometry, the only possibility for Q is the symplectic generalized quadrangle W(q).

Now we can proof our main result.

Proof of Theorem 1. Part (a) is a direct consequence of Lemma 3.14 and Lemma 3.11. The equivalence of Part (a) and (b) has already been discussed in Section 2.4. \Box

Theorem 2 is now a direct consequence.

Proof of Theorem 2. We show that the statement in Theorem 1(b) is not satisfied.

For q prime, the only ovoids of Q(4,q) are the elliptic quadrics $Q^{-}(3,q)$ [1, Cor. 1]. As any two such quadrics have nontrivial intersection, there is no partition of Q(4,q) into ovoids.

For q even, Q(4,q) does not admit a partition into evoids by [23, 3.4.1 (i)].

Acknowledgement

I would like to thank Jan de Beule for his patient help with the theory of generalized quadrangles during a research stay at the Vrije Universiteit Brussel (VUB) in the framework of WOG Coding Theory and Cryptography.

References

- [1] S. Ball, P. Govaerts, and L. Storme. "On ovoids of parabolic quadrics". In: *Designs, Codes and Cryptography* 38.1 (Jan. 2006), pp. 131–145.
- [2] A. Barlotti and J. Cofman. "Finite Sperner spaces constructed from projective and affine spaces". In: Abhandlungen aus dem Mathematischen Seminar der Universität Hamburg 40 (Mar. 1974), pp. 231–241.
- [3] C. Berge and D. Ray-Chaudhuri. "Unsolved Problems". In: *Hypergraph Seminar*. Hypergraph Seminar (Ohio State University, 1972). Ed. by C. Berge and D. Ray-Chaudhuri. Lecture Notes in Mathematics 411. 1974, pp. 278–287. ISBN: 978-3-540-06846-4.
- [4] M. Braun, T. Etzion, P. R. J. Östergård, A. Vardy, and A. Wassermann. "Existence of q-analogs of Steiner systems". In: Forum of Mathematics. Pi 4 (2016), e7.
- [5] M. Braun, M. Kiermaier, A. Kohnert, and R. Laue. "Large sets of subspace designs". In: *Journal of Combinatorial Theory. Series A* 147 (Apr. 2017), pp. 155–185.
- [6] M. Braun, M. Kiermaier, and A. Nakić. "On the automorphism group of a binary q-analog of the Fano plane". In: European Journal of Combinatorics 51 (2016), pp. 443–457.
- [7] M. Braun, M. Kiermaier, and A. Wassermann. "q-analogs of designs: Subspace designs". In: Network Coding and Subspace Designs. Ed. by M. Greferath, M. O. Pavčević, N. Silberstein, and M. Á. Vázquez-Castro. Signals and Communication Technology. Cham: Springer, 2018, pp. 171–211. ISBN: 978-3-319-70292-6.
- [8] F. Buekenhout and C. Lefèvre. "Generalized quadrangles in projective spaces". In: *Archiv der Mathematik* 25.1 (Dec. 1974), pp. 540–552.
- [9] E. Byrne and A. Ravagnani. "An Assmus-Mattson theorem for rank metric codes". In: SIAM Journal on Discrete Mathematics 33.3 (2019), pp. 1242–1260.
- [10] P. J. Cameron. "Generalisation of Fisher's inequality to fields with more than one element". In: Combinatorics. Proceedings of the British Combinatorial Conference 1973.
 Ed. by T. P. McDonough and V. C. Mavron. London Mathematical Society Lecture Note Series 13. Cambridge: Cambridge University Press, 1974, pp. 9–13. ISBN: 0-521-20454-2.
- [11] H. Cohn. "Projective geometry over \mathbb{F}_1 and the Gaussian binomial coefficients". In: The American Mathematical Monthly 111.6 (June–July 2004), pp. 487–495.
- [12] P. Delsarte. "Association schemes and t-designs in regular semilattices". In: Journal of Combinatorial Theory. Series A 20.2 (1976), pp. 230–243.
- [13] O. Heden and P. A. Sissokho. "On the existence of a (2,3)-spread in V(7,2)". In: Ars Combinatoria 124 (2016), pp. 161–164.

- [14] D. Heinlein, M. Kiermaier, S. Kurz, and A. Wassermann. "A subspace code of size 333 in the setting of a binary q-analog of the Fano plane". In: Advances in Mathematics of Communications 13.3 (Aug. 2019), pp. 457–475.
- [15] J. W. P. Hirschfeld. Finite projective spaces of three dimensions. Oxford Mathematical Monographs. Oxford and New York: The Clarendon Press and Oxford University Press, 1985. ISBN: 0-19-853536-8.
- [16] T. Honold and M. Kiermaier. "On putative q-analogues of the Fano plane and related combinatorial structures". In: *Dynamical Systems, Number Theory and Applications. A Festschrift in Honor of Armin Leutbecher's 80th Birthday.* Ed. by T. Hagen, F. Rupp, and J. Scheurle. World Scientific, Mar. 2016, pp. 141–175. ISBN: 978-981-4699-86-0.
- [17] T. Honold, M. Kiermaier, and S. Kurz. "Partial spreads and vector space partitions". In: *Network Coding and Subspace Designs*. Ed. by M. Greferath, M. O. Pavčević, N. Silberstein, and M. Á. Vázquez-Castro. Signals and Communication Technology. Cham: Springer, 2018, pp. 131–170. ISBN: 978-3-319-70292-6.
- [18] M. Kiermaier, S. Kurz, and A. Wassermann. "The order of the automorphism group of a binary q-analog of the Fano plane is at most two". In: *Designs, Codes and Cryptography* 86.2 (Feb. 2018), pp. 239–250.
- [19] M. Kiermaier and R. Laue. "Derived and residual subspace designs". In: Advances in Mathematics of Communications 9.1 (2015), pp. 105–115.
- [20] M. Kiermaier and M. O. Pavčević. "Intersection numbers for subspace designs". In: *Journal of Combinatorial Designs* 23.11 (2015), pp. 463–480.
- [21] Z. T. Mateva and S. T. Topalova. "Line spreads of PG(5, 2)". In: *Journal of Combinatorial Designs* 17.1 (Jan. 2009), pp. 90–102.
- [22] A. Nakić and M. O. Pavčević. "Tactical decompositions of designs over finite fields". In: *Designs, Codes and Cryptography* 77.1 (Oct. 2015), pp. 49–60.
- [23] S. E. Payne and J. A. Thas. *Finite generalized quadrangles*. 2nd ed. EMS Series of Lectures in Mathematics. European Mathematical Society. ISBN: 978-3-03719-066-1.
- [24] B. Segre. "Teoria di Galois, fibrazioni proiettive e geometrie non desarguesiane". In: Annali di Matematica Pura ed Applicata. Serie Quarta 64.1 (Dec. 1964), pp. 1–76.
- [25] H. Suzuki. "Five Days Introduction to the Theory of Designs". 1989. URL: http://subsite.icu.ac.jp/people/hsuzuki/lecturenote/designtheory.pdf.
- [26] H. Suzuki. "On the inequalities of t-designs over a finite field". In: European Journal of Combinatorics 11.6 (1990), pp. 601–607.
- [27] S. Thomas. "Designs over finite fields". In: Geometriae Dedicata 24.2 (Oct. 1987), pp. 237–242.
- [28] S. Thomas. "Designs and partial geometries over finite fields". In: Geometriae Dedicata 63.3 (Dec. 1996), pp. 247–253.
- [29] J. Tits. "Sur la trialité et certains groupes qui s'en déduisent". In: *Institut des Hautes Études Scientifiques. Publications Mathématiques* 2 (Jan. 1959), pp. 13–60.