Divergence Scaling for Distribution Matching

Gerhard Kramer

Institute for Communications Engineering Technical University of Munich D-80333 Munich, Germany

Abstract—Distribution matchers for finite alphabets are shown to have informational divergences that grow logarithmically with the block length, generalizing a basic result for binary strings.

I. INTRODUCTION

Distribution matchers (DMs) are random number generators with a one-to-one mapping [1]. An important application of DMs is probabilistic amplitude shaping for energy-efficient communication [2]. The objective of this paper is to prove that block (fixed-length) DMs cannot give low informational divergence for finite alphabets, thereby generalizing a basic result of Schulte and Geiger [3] for binary strings. The main tool we use is a large deviation theorem of Bahadur and Ranga Rao [4] that applies more generally than the binomial bounds and identities in [3].

This paper is organized as follows. Sec. II introduces notation and a few lemmas. Sec. III gives bounds on sums and establishes two further results, namely Lemma 5 that bounds the size of optimal DM codebooks and Theorem 2 that bounds a divergence for average empirical distributions. Sec. IV develops the desired scaling results. Appendixes A-D provide proofs of Lemmas and a Theorem.

II. PRELIMINIARIES

A. Basic Notation

Sets are written with calligraphic letters \mathcal{A} and the empty set with \emptyset . The cardinality of \mathcal{A} is $|\mathcal{A}|$ and the n-fold Cartesian product of \mathcal{A} is \mathcal{A}^n . For sequences f(n) and g(n), $n=1,2,\ldots$, the little-o notation f(n)=o(g(n)) means that $\lim_{n\to\infty} f(n)/g(n)=0$, see [5, p. 61].

Random variables (RVs) are written with uppercase letters such as A and their realizations with lowercase letters a. A probability mass function (pmf) of A is denoted by $P_A \in \mathcal{P}$ or $Q_A \in \mathcal{P}$, where \mathcal{P} is the set of pmfs for alphabet \mathcal{A} . We often discard subscripts on pmfs. Pmfs or functions are also written as vectors, e.g., we write pmf Q with the L-letter alphabet $\mathcal{A} = \{1, \ldots, L\}$ as $Q = [Q(1), \ldots, Q(L)]$. The uniform pmf over a set of K elements is denoted by U_K .

A random string $A^n = A_1 A_2 \dots A_n$ has realizations $a^n = a_1 a_2 \dots a_n \in \mathcal{A}^n$. The probability of a set $\mathcal{S} \subseteq \mathcal{A}^n$ of strings with respect to P_{A^n} is written as

$$P_{\mathsf{A}^n}(\mathcal{S}) = \sum_{a^n \in \mathcal{S}} P_{\mathsf{A}^n}(a^n). \tag{1}$$

We write $Q_{A^n} = Q_A^n$ for the pmf of a string of independent and identically distributed (iid) RVs.

The ℓ_1 distance between two pmfs P and Q on \mathcal{A} is

$$d_1(P,Q) = \sum_{a \in A} |P(a) - Q(a)|$$
 (2)

and we have $d_1(P,Q) \leq 2$ with equality if and only if $\operatorname{supp}(P) \cap \operatorname{supp}(Q) = \emptyset$. Of course, one may alternatively use the variational distance [6, Ch. 11.6].

The expectation of a real-valued function f of a RV A with respect to the pmf P is denoted as

$$E_f(P) = \mathbb{E}_P\left[f(\mathsf{A})\right] = \sum_{a \in \text{supp}(P)} P(a)f(a) \tag{3}$$

where $\operatorname{supp}(P)$ is the support of P, i.e., the set of $a \in \mathcal{A}$ with P(a) > 0. The notation $E_f(P)$ is useful because we usually consider P as the variable while f is fixed. Observe that $E_f(P)$ is linear in P.

The variance of f(A) with respect to P is $V_P(f(A))$ and the entropy of P is $H(P) = \mathbb{E}_P[-\log_2 P(A)]$. The cross entropy and divergence of pmfs P and Q with common alphabet \mathcal{A} are the respective

$$X(P||Q) = \mathbb{E}_P\left[-\log_2 Q(\mathsf{A})\right] \tag{4}$$

$$D(P||Q) = \mathbb{E}_P\left[\log_2 \frac{P(\mathsf{A})}{Q(\mathsf{A})}\right] \tag{5}$$

and we have X(P||Q) = H(P) + D(P||Q).

B. Empirical Probability

Let $\mathcal{P}_n \subseteq \mathcal{P}$ be the set of pmfs that have rational values with denominator n, also called the n-types. For a string a^n , let $n_i = n_i(a^n)$ be the number of occurrences of the letter i for $i \in \mathcal{A}$. The empirical pmf, or type, of a^n is

$$\pi(a^n) = \frac{1}{n} [n_1, \dots, n_L]$$
 (6)

and $\pi(a^n) \in \mathcal{P}_n$. The number of a^n with the same empirical pmf $\pi(a^n)$ is given by the multinomial

$$\binom{n}{n_1 \dots n_L} = \frac{n!}{n_1! \dots n_L!}.$$
 (7)

Let $\bar{n}_i = \sum_{a^n \in \mathcal{A}^n} P_{\mathsf{A}^n}(a^n) n_i(a^n)$ be the average number of occurrences of letter i for a specified P_{A^n} . The average empirical pmf is

$$\bar{\pi} = \sum_{a^n \in \mathcal{A}^n} P_{\mathcal{A}^n}(a^n) \, \pi(a^n) = \frac{1}{n} [\bar{n}_1, \dots, \bar{n}_L]$$
 (8)

and for any target pmf Q_{T} we have

$$D(P_{\mathsf{A}^{n}} \| Q_{\mathsf{T}}^{n}) = n X(\bar{\pi} \| Q_{\mathsf{T}}) - H(P_{\mathsf{A}^{n}}). \tag{9}$$

C. Four Lemmas

We state several lemmas; the first three are proved in Appendixes A-C. Let p_{\min} and p_{\max} be the respective minimum and maximum probabilities of P, and define similar notation for the minima and maxima of the pmfs Q and R.

Lemma 1. Let $d_1 = d_1(P, Q)$. If $d_1 < 2p_{\min}$ then

$$D(Q||R) - D(P||R) \le \frac{d_1}{2} \log_2 \left(\frac{p_{\max} + d_1/2}{p_{\min} - d_1/2} \cdot \frac{r_{\max}}{r_{\min}} \right). \tag{10}$$

Next, consider a threshold I and define the pmf and string sets $\mathcal{E}=\{P: E_f(P)\leq I\}$ and $\mathcal{S}=\{a^n: \pi(a^n)\in \mathcal{E}\}$. We further define $P_{\mathsf{A}^n}(a^n)=Q^n(a^n)/Q^n(\mathcal{S})$ for $a^n\in \mathcal{S}$ and consider $\bar{\pi}$ to be a function of I.

Lemma 2. Consider any P^* for which $E_f(P^*) = I$. Then there is a $P \in \mathcal{E} \cap \mathcal{P}_n$ such that

$$d_1(P, P^*) \le \frac{2(L-1)}{n}.$$
 (11)

Lemma 3. $E_f(\bar{\pi})$ is non-decreasing in I.

Next define

$$P^* = \operatorname*{arg\,min}_{P \in \mathcal{E}} D\left(P \| Q\right). \tag{12}$$

We have the following standard result, see e.g. [6, Ch. 11.5].

Lemma 4. Let $f_{\min} = \min_{a \in \text{supp}(Q)} f(a)$ and suppose that $f_{\min} < I < E_f(Q)$. Then the solution of (12) is

$$P^*(a) = \frac{Q(a) \, 2^{-\tau f(a)}}{\sum_{b \in \mathcal{A}} Q(b) \, 2^{-\tau f(b)}}$$
(13)

for $a \in A$, where τ is positive and chosen so that $E_f(P^*) = I$. Observe that $P^*(a) = 0$ if and only if Q(a) = 0.

III. BOUNDS FOR SAMPLE MEANS

Consider the sample mean

$$S_n = \frac{1}{n} \sum_{i=1}^n f(A_i). \tag{14}$$

We are interested in characterizing $\Pr[S_n \leq I]$. Observe that $S_n = E_f(\pi(A^n))$ so that $S_n \leq I$ is the same as requiring $\pi(A^n)$ to lie in the half-space $\mathcal{E} = \{P : E_f(P) \leq I\}$.

A. Deviations of the Sample Mean

We distinguish between two classes of RVs:

- 1) lattice: there are constants c_0 and d > 0 such that $A \subseteq \{c_0 + jd : j \text{ an integer}\};$
- 2) non-lattice.

For example, every real-valued binary RV is lattice because one may choose d as the absolute difference between the two alphabet values. In general, one chooses d as the greatest common divisor of the differences between consecutive possible

values of A, see [4, Sec. 4]. The following large deviation theorem is developed in [4, Thm. 1], see also [7, Ch. 3.7].

Theorem 1 (Bahadur and Ranga Rao). Consider the iid string A^n with pmf Q^n and the real-valued function f with domain A. Let $f_{min} = \min_{a \in \text{supp}(Q)} f(a)$ and consider an I satisfying

$$f_{\min} < I < \mathbb{E}_Q[f(\mathsf{A})] = E_f(Q) \tag{15}$$

and also $Pr[S_n = I] > 0$ if f(A) is a lattice RV. We have

$$\Pr\left[\mathsf{S}_n \le I\right] = \frac{2^{-nD(P^*\|Q)}}{\sqrt{2\pi n}} \, b\left(1 + o(1)\right) \tag{16}$$

where P^* is given by (13). The values

$$b = \begin{cases} \frac{1}{\alpha} \frac{\tau d}{1 - e^{-\tau d}}, & \text{if } f(A) \text{ is a lattice } RV\\ 1/\alpha, & \text{else} \end{cases}$$
 (17)

$$\alpha = \sqrt{V_{P^*} \left(\ln \frac{P^*(\mathsf{A})}{Q(\mathsf{A})} \right)} \tag{18}$$

satisfy $0 < b < \infty$ and $0 < \alpha < \infty$, and τ is as in Lemma 4. Moreover, given I the b, α , and τ are independent of n. However, if f(A) is lattice and $\Pr[S_n = I] = 0$ then one must replace b with $b_n = be^{-\tau d \theta_n}$ where $0 \le \theta_n < 1$. Thus, the right-hand side of (16) is an upper bound on $\Pr[S_n \le I]$.

Corollary 1. Choosing $Q = U_L$ in Theorem 1, (16) becomes

$$\Pr\left[\mathsf{S}_n \le I\right] = \frac{L^{-n} \, 2^{nH(P^*)}}{\sqrt{2\pi n}} \, b\left(1 + o(1)\right) \tag{19}$$

where the value b is as in (17) and

$$\alpha = \sqrt{V_{P^*} \left(\ln P^*(\mathsf{A}) \right)}. \tag{20}$$

Furthermore, $P^*(a) > 0$ for all $a \in A$.

B. Example

We apply Corollary 1 to an example that corresponds to optimal DM [1]. Consider a target pmf Q_T , e.g., a shaping pmf for energy-efficient communication. Suppose we wish to count the number of a^n with at least a specified probability with respect to Q_T^n , i.e., the number of a^n satisfying

$$Q_{\mathsf{T}}^n(a^n) \ge 2^{-nI} \tag{21}$$

and where equality holds for some a^n . The bound (21) is the same as

$$\frac{1}{n} \sum_{i=1}^{n} -\log_2 Q_{\mathsf{T}}(a_i) = X\left(\pi(a^n) \| Q_{\mathsf{T}}\right) \le I \tag{22}$$

which has the same form as (14) with $f(a) = -\log_2 Q_T(a)$. We thus restrict attention to the interval (see (15))

$$-\log_2 t_{\max} < I < E_f(Q) = X(Q||Q_{\mathsf{T}}) \tag{23}$$

where $t_{\max} = \max_{a \in \mathcal{A}} Q_{\mathsf{T}}(a)$.

Now consider the pmf set $\mathcal{E} = \{P : X(P||Q_T) \leq I\}$ and the corresponding string set $\mathcal{S} = \{a^n : \pi(a^n) \in \mathcal{E}\}$. We have

$$|\mathcal{S}| = \sum_{P \in \mathcal{E} \cap \mathcal{P}_n} \binom{n}{nP(1) \dots nP(L)}.$$
 (24)

 $^{^1}$ The greatest common divisor of a set of real numbers is the largest d such that all all numbers in the set are integer multiples of d, e.g., the set $\{1,5/3\}$ has greatest common divisor d=1/3.

Normalizing (24) by L^n , one obtains a probability with respect to the uniform pmf over all strings in \mathcal{A}^n . We thus choose A^n with pmf Q^n where $Q = U_L$ and can write

$$|\mathcal{S}| = L^n \cdot \Pr\left[\mathsf{S}_n \le I\right]. \tag{25}$$

Corollary 1 gives the following result.

Lemma 5. The set S of a^n satisfying (22) has cardinality

$$|S| = \frac{2^{nH(P^*)}}{\sqrt{2\pi n}} b (1 + o(1))$$
 (26)

where P^* is given by (13).

Example 1. Fig. 1 illustrates the probability simplex for $A = \{1, 2, 3\}$ and n = 10. Consider $Q = U_L$ and the lattice RV f(A) with

$$f = -\log_2 Q_{\mathsf{T}} = c_0 + [0, 1, 8/3] \tag{27}$$

where $c_0 \approx 0.7290, \ Q_{\rm T} \approx [0.6033, 0.3017, 0.0950],$ and d=1/3. The bounds (23) specify $c_0 < I < c_0 + 11/9$. Consider $I \approx 1.5290$ and n=10 so that $P_1 = [7,0,3]/10$ and $P_2 = [2,8,0]/10$ are both in \mathcal{P}_n and satisfy the constraint $E_f(P) = X \ (P \| Q_{\rm T}) \le I$ with equality. We compute $P^* \approx [0.4915, 0.3336, 0.1749]$ and $H \ (P^*) \approx 1.4720$. Consider the strings

$$\hat{a}^n = 1, 1, 1, 1, 1, 1, 2, 2, 3, 3 \tag{28}$$

$$\tilde{a}^n = 1, 1, 1, 1, 1, 2, 2, 2, 2, 3$$
 (29)

with empirical pmfs $\pi(\hat{a}^n) = [6,2,2]/10$ and $\pi(\tilde{a}^n) = [5,4,1]/10$. The pmf $\pi(\hat{a}^n)$ maximizes the entropy in $\mathcal{E} \cap \mathcal{P}_n$ and $\pi(\tilde{a}^n)$ has the smallest ℓ_1 distance to P^* in $\mathcal{E} \cap \mathcal{P}_n$. The entropies are $H(\pi(\hat{a}^n)) \approx 1.3710$ and $H(\pi(\tilde{a}^n)) \approx 1.3610$. The ℓ_1 distances are $d_1(P_1,P^*) \approx 0.6672$, $d_1(P_2,P^*) \approx 0.9328$, $d_1(\pi(\hat{a}^n),P^*) \approx 0.2672$, and $d_1(\pi(\tilde{a}^n),P^*) \approx 0.1498$.

C. Convergence of the Average Empirical Distribution

We develop a scaling result for the conditional limit theorem [6, Ch. 11.6]. Let p_{\min}^* and p_{\max}^* be the respective minimum and maximum probabilities of P^* . Suppose that $q_{\min} > 0$ so that $p_{\min}^* > 0$ by Lemma 4. Define the function

$$\Delta(P) = D(P||Q) - D(P^*||Q). \tag{30}$$

Observe that $\Delta(P)$ is bounded and convex in P, and non-negative for $P \in \mathcal{E}$. We have the following scaling result.

Theorem 2. Consider the iid string A^n with pmf Q^n . If I satisfies (15) then we have

$$D(\bar{\pi}||P^*) \le \Delta(\bar{\pi}) < \frac{1}{n} 2^{\tilde{c}+1} (1 + o(1))$$
 (31)

where

$$\tilde{c} = (L-1)\log_2 \frac{p_{\max}^* q_{\max}}{p_{\min}^* q_{\min}}.$$
 (32)

Proof. Observe that $E_f(\bar{\pi}) \leq I$ by the linearity of expectation. The first inequality in (31) thus follows by a "Pythagorean" inequality [6, Ch. 11.6]. The second inequality is proved in Appendix D.




Fig. 1. Probability simplex for $\mathcal{A}=\{1,2,3\}$ with grid lines for n=10. The red shaded area shows the set \mathcal{E} of pmfs P satisfying $X\left(P\|Q_{\mathsf{T}}\right)\leq I$ where $Q_{\mathsf{T}}\approx [0.6033,0.3017,0.0950]$ and $I\approx 1.5290$.

IV. DIVERGENCE FOR DISTRIBUTION MATCHERS

The problem of DM, as described in [1], [3], is to choose a codebook S of strings a^n so that the divergence $D\left(P_{\mathsf{A}^n} \| Q_{\mathsf{T}}^n\right)$ is minimized where $P_{\mathsf{A}^n} = U_{|S|}$. Without loss of essential generality, we assume $\operatorname{supp}(Q_{\mathsf{T}}) = \mathcal{A}$ and $Q_{\mathsf{T}} \neq U_L$. We have $H\left(P_{\mathsf{A}^n}\right) = \log_2 |\mathcal{S}|$, and the rate in bits per symbol is $R_{\inf} = \frac{1}{n} \log_2 |\mathcal{S}|$.

The paper [1, Prop. 3] shows that optimal $\mathcal S$ have all strings a^n satisfying $X\left(\pi(a^n)\|Q_\mathsf{T}\right) \leq I$ for some I, see (22). The paper [3] shows that for L=2 the divergence $D\left(P_{\mathsf{A}^n}\|Q_\mathsf{T}^n\right)$ of the best $\mathcal S$ scales as $\frac12\log_2 n$ with n, i.e., binary DMs cannot provide low divergence. We prove the same result for general discrete alphabets $\mathcal A$ by using Lemmas 1-5 and Theorem 2.

A. Small I

The case $I \leq -\log_2 t_{\max}$ makes sense only if equality holds. S is thus the set of a^n having only letters with maximal probability $Q_T(a) = t_{\max}$. The positive $P_{A^n}(a^n)$ then all have the same value $1/N_{\max}^n$ where N_{\max} is the number of letters a with probability $Q_T(a) = q_{\max}$. Moreover, we compute

$$D(P_{\mathsf{A}^n} || Q_{\mathsf{T}}^n) = n \log_2 \frac{1}{N_{\max} t_{\max}}$$
 (33)

and the divergence grows linearly with n since $N_{\rm max}\,t_{\rm max}<1$ by our assumption that $Q_{\rm T}$ is not uniform.

B. Large I

For $I \geq X(U_L || Q_T)$, the following result is the analog of Lemma 17 in [1, App. B].

Lemma 6. $D(P_{A^n}||Q_{\mathsf{T}}^n)$ scales linearly with n for $I > \log_2 L$.

Proof. We have $\log_2 L < X\left(U_L\|Q_\mathsf{T}\right)$ because $Q_\mathsf{T} \neq U_L$. Moreover, if $\log_2 L < I < X\left(U_L\|Q_\mathsf{T}\right)$ then Theorem 2 gives

 $\bar{\pi} \to P^*$ as $n \to \infty$ and thus $X\left(\bar{\pi}\|Q_{\mathsf{T}}\right) \to X\left(P^*\|Q_{\mathsf{T}}\right) = I$ since $X\left(P^*\|Q_{\mathsf{T}}\right) = I$ by Lemma 4. But $H\left(P_{\mathsf{A}^n}\right) \le n\log_2 L$ and thus the divergence (9) grows linearly with n. Finally, Lemma 3 states that $X\left(\bar{\pi}\|Q_{\mathsf{T}}\right)$ is non-decreasing in I. \square

C. Intermediate I

For
$$-\log_2 t_{\text{max}} < I < X(U_L || Q_T)$$
, Lemma 5 gives

$$H(P_{A^n}) = -\frac{1}{2}\log_2 n + nH(P^*) + \log_2 \frac{b}{\sqrt{2\pi}} + o(1).$$
(34)

Inserting (34) into (9) gives

$$D(P_{\mathsf{A}^n} \| Q_{\mathsf{T}}^n) = \frac{1}{2} \log_2 n - n \left[H(P^*) - H(\bar{\pi}) \right] + nD(\bar{\pi} \| Q_{\mathsf{T}}) - \log_2 \frac{b}{\sqrt{2\pi}} + o(1)$$
(35)

which is the analog of [1, Eq. (107)]. Theorem 2 with ${\cal Q}={\cal U}_L$ now gives

$$0 \le \Delta(\bar{\pi}) = H(P^*) - H(\bar{\pi}) < \frac{1}{n} 2^{\tilde{c}+1} (1 + o(1)).$$
 (36)

Inserting (36) into (35) and taking $n \to \infty$, we have

$$-2^{\tilde{c}+1} - \log_2 \frac{b}{\sqrt{2\pi}}$$

$$\leq \liminf_{n \to \infty} \left(D\left(P_{\mathsf{A}^n} \| Q_{\mathsf{T}}^n \right) - \frac{1}{2} \log_2 n - nD\left(\bar{\pi} \| Q_{\mathsf{T}} \right) \right)$$

$$\leq -\log_2 \frac{b}{\sqrt{2\pi}}.$$
(37)

The divergence $D\left(P_{\mathsf{A}^n} \| Q_{\mathsf{T}}^n\right)$ thus grows at least as $\frac{1}{2} \log_2 n$ with n. The next result shows that one can achieve this growth.

Theorem 3. Finite-alphabet block DMs that minimize the divergence $D(P_{A^n}||Q_T^n)$ have $D(P_{A^n}||Q_T^n)$ that grows as $\frac{1}{2}\log_2 n$ with n and $R_{info} \to H(Q_T)$ as $n \to \infty$.

Proof. We have already proved the converse. For the coding theorem, one may quantize Q_{T} to a $\tilde{Q}_{\mathsf{T}} \in \mathcal{P}_n$ satisfying $d_1\left(\tilde{Q}_{\mathsf{T}},Q_{\mathsf{T}}\right) \leq L/(2n)$ [8, Prop. 2]. Now choose $I=H\left(\tilde{Q}_{\mathsf{T}}\right)$ so that $P^*=\tilde{Q}_{\mathsf{T}}$. Theorem 2 gives $nD\left(\bar{\pi}\|\tilde{Q}_{\mathsf{T}}\right) < 2^{\tilde{c}+1}(1+o(1))$, and using (35) and the left-hand side of (36) we find that the divergence $D\left(P_{\mathsf{A}^n}\|Q_{\mathsf{T}}^n\right)$ grows as $\frac{1}{2}\log_2 n$. Furthermore, $R_{\mathsf{info}} \to H\left(Q_{\mathsf{T}}\right)$ by Lemmas 1 and 5.

ACKNOWLEDGEMENT

The author wishes to thank P. Schulte and D. Lentner for discussions, and the reviewers for comments that improved the presentation. This work was supported by the German Research Foundation (DFG) under Grant KR 3517/9-1.

REFERENCES

- [1] P. Schulte, R. A. Amjad, T. Wiegart, and G. Kramer, "Invertible low-divergence coding," 2020. [Online]. Available: https://arxiv.org/abs/2010.10583
- [2] G. Böcherer, F. Steiner, and P. Schulte, "Bandwidth efficient and rate-matched low-density parity-check coded modulation," *IEEE Trans. Commun.*, vol. 63, no. 12, pp. 4651–4665, Dec. 2015.

- [3] P. Schulte and B. C. Geiger, "Divergence scaling of fixed-length, binaryoutput, one-to-one distribution matching," in *Proc. IEEE Int. Symp. Inf. Theory*, Aachen, Germany, June 2017, pp. 3075–3079.
- [4] R. R. Bahadur and R. Ranga Rao, "On deviations of the sample mean," Ann. Math. Statist., vol. 31, no. 4, pp. 1015–1027, Dec. 1960.
- [5] E. Landau, Handbuch der Lehre von der Verteilung der Primzahlen. Leipzig and Berlin: B. G. Teubner, 1909, vol. 1.
- [6] T. M. Cover and J. A. Thomas, *Elements of Information Theory*, 2nd ed. John Wiley & Sons, Inc., 2006.
- [7] A. Dembo and O. Zeitouni, Large Deviations Techniques and Applications, 2nd ed. Springer, 1998.
- [8] G. Böcherer and B. C. Geiger, "Optimal quantization for distribution synthesis," *IEEE Trans. Inf. Theory*, vol. 62, no. 11, pp. 6162–6172, Nov. 2016

APPENDIX A PROOF OF LEMMA 1

The condition $p_{\min} > d_1/2$ implies $q_{\min} > 0$ and therefore $\operatorname{supp}(P) = \operatorname{supp}(Q) = \mathcal{A}$. Consider the pmfs P and $Q = P + \Delta$ where $\sum_a \Delta(a) = 0$ and $\sum_a |\Delta(a)| = d_1$. Define

$$\Delta_{+} = \sum_{a: \Delta(a) > 0} \Delta(a), \quad \Delta_{-} = \sum_{a: \Delta(a) < 0} \Delta(a)$$
 (38)

and observe that $\Delta_+ = -\Delta_- = d_1/2$. Now expand

$$D(Q||R) - D(P||R) = -D(P||Q) + \sum_{a:\Delta(a)\neq 0} \Delta(a) \log_2 \frac{P(a) + \Delta(a)}{R(a)}$$
(39)

and compute the bounds

$$\sum_{a:\Delta(a)>0} \Delta(a) \underbrace{\log_2 \frac{P(a) + \Delta(a)}{R(a)}}_{\leq \log_2 \frac{p_{\max} + d_1/2}{r_{\min}}} \leq \frac{d_1}{2} \log_2 \frac{p_{\max} + d_1/2}{r_{\min}}$$

$$(40)$$

$$\sum_{a:\Delta(a)<0} \Delta(a) \underbrace{\log_2 \frac{P(a) + \Delta(a)}{R(a)}}_{\geq \log_2 \frac{p_{\min} - d_1/2}{r_{\max}}} \leq -\frac{d_1}{2} \log_2 \frac{p_{\min} - d_1/2}{r_{\max}}.$$
(41)

APPENDIX B PROOF OF LEMMA 2

The proof applies the approach in [8, Sec. III and Sec. VI.C] with small changes. Order the values $P^*(i)$, $i=1,\ldots,L$, such that $f(i) \leq f(j)$ if i < j. Define the pmf P via

$$P(i) = \begin{cases} \lfloor nP^*(i) \rfloor / n, & i = 2, \dots, L \\ 1 - \sum_{i=2}^{L} P(i), & i = 1 \end{cases}$$
 (42)

and note that $P \in \mathcal{P}_n$. Define $e(i) = P^*(i) - P(i)$ for all i. We have $0 \le e(i) < \frac{1}{n}$ for $i = 2, \ldots, L$ and $\sum_{i=1}^L e(i) = 0$ so that $-(L-1)/n < e(1) \le 0$. We thus have (11) and

$$E_f(P) = \underbrace{E_f(P^*)}_{=I} - \sum_{i=1}^{L} e(i) \underbrace{f(i)}_{>f(1)} \le I$$
 (43)

where the last step uses $e(i) \geq 0$ for i = 2, ..., n. We thus have $P \in \mathcal{E} \cap \mathcal{P}_n$.

APPENDIX C PROOF OF LEMMA 3

Suppose we increase I from I_1 to I_2 with the corresponding string sets S_1 and S_2 and average empirical pmfs $\bar{\pi}_1$ and $\bar{\pi}_2$, respectively. Suppose $I_2 - I_1$ is sufficiently large so that $S_2 \setminus S_1 \neq \emptyset$. The definition (8) gives

$$E_{f}(\bar{\pi}_{2}) = \sum_{a^{n} \in \mathcal{S}_{2}} \frac{Q^{n}(a^{n})}{Q^{n}(\mathcal{S}_{2})} E_{f}(\pi(a^{n}))$$

$$= \frac{Q^{n}(\mathcal{S}_{1})}{Q^{n}(\mathcal{S}_{2})} E_{f}(\bar{\pi}_{1}) + \sum_{a^{n} \in \mathcal{S}_{2} \setminus \mathcal{S}_{1}} \frac{Q^{n}(a^{n})}{Q^{n}(\mathcal{S}_{2})} E_{f}(\pi(a^{n}))$$

$$> E_{f}(\bar{\pi}_{1})$$
(44)

where the last step follows by $E_f(\pi(a^n)) > I_1 \ge E_f(\bar{\pi}_1)$ for $a^n \in \mathcal{S}_2 \setminus \mathcal{S}_1$.

APPENDIX D PROOF OF THEOREM 2

There may be no $P \in \mathcal{E} \cap \mathcal{P}_n$ such that $E_f(P) = I$. However, for each n there is an I^* such that $E_f(P) \leq I^* \leq I$ for all $P \in \mathcal{E} \cap \mathcal{P}_n$, and with $E_f(P) = I^*$ for some $P \in \mathcal{E} \cap \mathcal{P}_n$. Define the optimized pmf (note that I^* replaces I)

$$P^* = \underset{P}{\arg\min} \ D\left(P\|Q\right)$$
 subject to $E_f(P) = I^*$. (45)

Next, by convexity of $\Delta(P)$ we have

$$\Delta(\bar{\pi}) \le \sum_{a^n \in \mathcal{S}} P_{\mathsf{A}^n}(a^n) \, \Delta(\pi(a^n)). \tag{46}$$

Now partition S into the sets $\{S_j\}_{j=0}^N$ where S_j is the set of $a^n \in S$ satisfying

$$j\,\delta \le \Delta(\pi(a^n)) < (j+1)\,\delta\tag{47}$$

for small positive δ and sufficiently large N. Let Δ_{\max} be an upper bound on $\Delta(P)$. We select a $N_1 < N$ and use (47) and $\Delta(P) \le \Delta_{\max} < \infty$ to write

$$\Delta(\bar{\pi}) < \left(\sum_{j=0}^{N_1 - 1} P_{\mathsf{A}^n}(\mathcal{S}_j)(j+1)\right) \delta + P_{\mathsf{A}^n}\left(\bigcup_{j=N_1}^{N} \mathcal{S}_j\right) \Delta_{\max}$$

$$= \left(\sum_{j=0}^{N_1 - 1} P_{\mathsf{A}^n}\left(\bigcup_{i=j}^{N_1 - 1} \mathcal{S}_i\right)\right) \delta + P_{\mathsf{A}^n}\left(\bigcup_{j=N_1}^{N} \mathcal{S}_j\right) \Delta_{\max}$$
(48)

where the second step follows because $\{S_j\}_{j=0}^N$ is a partition. Let $\mathcal{T}_j = \bigcup_{i=j}^N S_i$ and note that $\mathcal{T}_j \neq \emptyset$ for small δ and j. Define the following parameters of \mathcal{T}_j :

$$a_j^n = \underset{a^n \in \mathcal{T}_j}{\arg \max} \ E_f(\pi(a^n)), \quad I_j^* = E_f(\pi(a_j^n)),$$
 (49)

$$P_j^* = \underset{P}{\operatorname{arg\,min}} \ D\left(P\|Q\right) \text{ subject to } E_f(P) = I_j^* \qquad (50)$$

and observe that $I_0^* = I^*$ and $P_0^* = P^*$. Observe also that $a^n \in \mathcal{T}_j$ if and only if $E_f(\pi(a^n)) \leq I_j^*$ because $D\left(P_j^* \| Q\right)$

increases as I_j^* decreases. In fact, since τ is the Lagrange multiplier for the optimization problem in (12), we have

$$\frac{dD(P^*||Q)}{dI} = -\tau < 0.$$
 (51)

Let $p_{\min,j}^*$ and $p_{\max,j}^*$ be the respective minimum and maximum probabilities of P_j^* . For $j \, \delta$ not too large we have $p_{\min,j}^* > 0$, see Lemma 4. We further define

$$\hat{a}_j^n = \underset{a^n \in \mathcal{S}_j}{\arg \min} \ D\left(\pi(a^n) \| Q\right) \tag{52}$$

and note that $\hat{a}^n_j \neq a^n_j$ in general, see Fig. 1. Next, if $n>(L-1)/p^*_{\min,j}$ then Lemmas 1 and 2 imply that there is an $\tilde{a}^n_j \in \mathcal{S}_j$ such that

$$d_1\left(\pi(\tilde{a}_j^n), P_j^*\right) \le \frac{2(L-1)}{n} \tag{53}$$

$$D\left(\pi(\tilde{a}_{j}^{n})\|Q\right) - D\left(P_{j}^{*}\|Q\right) \le \frac{c_{j}}{n} \tag{54}$$

where

$$c_{j} = (L-1)\log_{2}\left(\frac{p_{\max,j}^{*} + \frac{L-1}{n}}{p_{\min,j}^{*} - \frac{L-1}{n}} \cdot \frac{q_{\max}}{q_{\min}}\right).$$
 (55)

Note that $\tilde{a}_i^n \neq \hat{a}_i^n$ in general, see Fig. 1.

By definition we have $D\left(\pi(\hat{a}_j^n)\|Q\right) \leq D\left(\pi(\tilde{a}_j^n)\|Q\right)$ so that (47) and (54) give

$$j\delta \le \Delta(\pi(\hat{a}_j^n)) \le \Delta(\pi(\tilde{a}_j^n)) \le \Delta(P_j^*) + \frac{c_j}{n}.$$
 (56)

We can thus use Theorem 1 to write

$$P_{\mathsf{A}^{n}}(\mathcal{T}_{j}) = \frac{\Pr\left(E_{f}(\pi(A^{n})) \leq I_{j}^{*}\right)}{\Pr\left(E_{f}(\pi(A^{n})) \leq I_{0}^{*}\right)}$$

$$\leq 2^{-n\Delta(P_{j}^{*})} \frac{b(I_{j}^{*})}{b(I_{0}^{*})} (1 + o(1)) \tag{57}$$

where the $b(I_j^*)$ are the bs in Theorem 1 corresponding to P_j^* . The inequality in (57) accounts for lattice RVs for which one may have $\Pr[S_n = I_j^*] = 0$ for $j = 1, \ldots, N_1$, see Theorem 1.

We now choose $\delta = 1/n$ and $N_1 = \lceil \sqrt{n} \rceil$ so that the expressions (48), (56), and (57) give

$$\Delta(\bar{\pi}) < \left[\left(\sum_{j=0}^{\infty} 2^{-j} \right) \frac{1}{n} + \Delta_{\max} 2^{-\sqrt{n}} \right] b_r 2^c (1 + o(1))$$
(58)

where

$$b_r = \max_{0 \le j \le N_1} \frac{b(I_j^*)}{b(I_0^*)}, \quad c = \max_{0 \le j \le N_1} c_j.$$
 (59)

Next, since $N_1/n \to 0$ we have $(j+1)\delta \to 0$ in (47) and hence $P_j^* \to P^*$ for all $j=0,1,\ldots,N_1$. We thus have $p_{\min,j}^* \to p_{\min}^*$ and $p_{\max,j}^* \to p_{\max}^*$. Moreover, by continuity of α and τ with respect to the threshold I, we have $b(I_j^*) \to b$ for all $j=0,1,\ldots,N_1$ and therefore $b_r \to 1$. Finally, evaluating the sum in (58) gives (31).