

A note on the Lauricella matrix functions

Ravi Dwivedi* and Vivek Sahai†

September 1, 2021

Abstract

In this paper, we study the solutions of system of bilateral type matrix differential equations and presented these solutions in terms of Lauricella hypergeometric matrix functions of several variables and Srivastava's triple hypergeometric matrix functions of three variables. We also discuss the region of convergence and integral representation for these matrix functions.

Keywords: Matrix differential equations, Matrix functional calculus, Multivariable hypergeometric functions.

AMS Subject Classification: 15A15, 33C65, 33E30

1 Introduction

Jodár and Cortés [13], introduced the concept of a fundamental set of solutions for matrix differential equations of the type

$$X'' = f_1(z)X' + f_2(z)Xf_3(z) + X'f_4(z), \quad (1.1)$$

where f_i are matrix valued functions of complex variable z . A closed form general solution of such bilateral type matrix differential equation is determined in terms of Gauss hypergeometric matrix function. In this paper, we give the systems of partial matrix differential equations of bilateral type in the form

$$U_{x_i x_i} = \sum_{\substack{m, l=1 \\ m < l}}^n f_{iml}(x_1, \dots, x_n) U_{x_m x_l} + \sum_{j=1}^n (U_{x_j} g_{ij}(x_1, \dots, x_n) + h_{ij}(x_1, \dots, x_n) U_{x_j}) \\ + f(x_1, \dots, x_n) U g(x_1, \dots, x_n), \quad 1 \leq i, j \leq n, \quad (1.2)$$

where $U_{x_i x_i} = \frac{\partial^2 U}{\partial x_i^2}$, $U_{x_m x_l} = \frac{\partial^2 U}{\partial x_m \partial x_l}$, $U_{x_j} = \frac{\partial U}{\partial x_j}$ and g_{ij} , h_{ij} , f and g are matrix valued functions of complex variables x_1, \dots, x_n .

We show that the Lauricella matrix functions of n -variables satisfy the systems of bilateral type matrix differential equation of the form (1.2). The region of convergence and integral representation for Lauricella matrix functions of n -variables are also determined. The particular case $n = 3$ leading to fourteen Lauricella matrix functions of three variables has been discussed in detail. The sectionwise treatment is as follows:

In Section 2, we give some basic definitions from special matrix function theory that are needed in the sequel. In Section 3, we find the convergence conditions and system of partial matrix differential equations of bilateral type satisfied by the generalized (n -variable) Lauricella matrix functions. The

*Department of Mathematics, National Institute of Technology, Kurukshetra, India.
E-mail: dwivedir999@gmail.com

†Department of Mathematics and Astronomy Lucknow University Lucknow 226007, India.
E-mail: sahai_vivek@hotmail.com (Corresponding author)

region of convergence and integral representation satisfy by the generalized (n -variable) Lauricella matrix functions are also given here. In Section 4, we give the complete list of fourteen Lauricella matrix functions of three variables and matrix analogue of Srivastava triple hypergeometric functions. Their regions of convergence, integral representations and the system of matrix differential equations of bilateral type satisfied by them are also given. Our notations are standard. For details, see [5, 6, 11, 12, 13].

2 Preliminaries

Let the spectrum of a matrix A in $\mathbb{C}^{r \times r}$, denoted by $\sigma(A)$, be the set of all eigenvalues of A and let $\alpha(A) = \max\{\Re(z) \mid z \in \sigma(A)\}$ and $\beta(A) = \min\{\Re(z) \mid z \in \sigma(A)\}$. Then for a positive stable matrix $A \in \mathbb{C}^{r \times r}$, that is $\beta(A) > 0$, the gamma matrix function is defined as [11]

$$\Gamma(A) = \int_0^\infty e^{-t} t^{A-I} dt$$

and the reciprocal gamma matrix function is defined as [11]

$$\Gamma^{-1}(A) = A(A+I) \dots (A+(n-1)I) \Gamma^{-1}(A+nI), \quad n \geq 1. \quad (2.1)$$

The Pochhammer symbol for $A \in \mathbb{C}^{r \times r}$ is given by [12]

$$(A)_n = \begin{cases} I, & \text{if } n = 0, \\ A(A+I) \dots (A+(n-1)I), & \text{if } n \geq 1. \end{cases} \quad (2.2)$$

This gives

$$(A)_n = \Gamma^{-1}(A) \Gamma(A+nI), \quad n \geq 1. \quad (2.3)$$

If $A \in \mathbb{C}^{r \times r}$ is a positive stable matrix and $n \geq 1$ is an integer, then the gamma matrix function can also be defined in the form of a limit as [11]

$$\Gamma(A) = \lim_{n \rightarrow \infty} (n-1)! (A)_n^{-1} n^A. \quad (2.4)$$

If A and B are positive stable matrices in $\mathbb{C}^{r \times r}$, then the beta matrix function is defined as [11]

$$\mathfrak{B}(A, B) = \int_0^1 t^{A-I} (1-t)^{B-I} dt. \quad (2.5)$$

Furthermore, if $AB = BA$, then the beta matrix function can be written in terms of gamma matrix function as [11]

$$\mathfrak{B}(A, B) = \Gamma(A) \Gamma(B) \Gamma^{-1}(A+B). \quad (2.6)$$

Using the Schur decomposition of A , it follows that [9, 17]

$$\|e^{tA}\| \leq e^{t\alpha(A)} \sum_{k=0}^{r-1} \frac{(\|A\|^{r^{1/2}t})^k}{k!}, \quad t \geq 0. \quad (2.7)$$

We shall use the notation $\Gamma \left(\begin{matrix} A_1, \dots, A_p \\ B_1, \dots, B_q \end{matrix} \right)$ for $\Gamma(A_1) \dots \Gamma(A_p) \Gamma^{-1}(B_1) \dots \Gamma^{-1}(B_q)$.

3 Generalized Lauricella matrix functions

The four Appell matrix functions of two variables [2, 7] can be generalized to the following matrix functions of n -variables.

$$\begin{aligned} & F_{\mathcal{A}}[A, B_1, \dots, B_n; C_1, \dots, C_n; x_1, \dots, x_n] \\ &= \sum_{m_1, \dots, m_n=0}^{\infty} (A)_{m_1+\dots+m_n} (B_1)_{m_1} \cdots (B_n)_{m_n} (C_1)_{m_1}^{-1} \cdots (C_n)_{m_n}^{-1} \frac{x_1^{m_1} \cdots x_n^{m_n}}{m_1! \cdots m_n!}; \end{aligned} \quad (3.1)$$

$$\begin{aligned} & F_{\mathcal{B}}[A_1, \dots, A_n, B_1, \dots, B_n; C; x_1, \dots, x_n] \\ &= \sum_{m_1, \dots, m_n=0}^{\infty} (A_1)_{m_1} \cdots (A_n)_{m_n} (B_1)_{m_1} \cdots (B_n)_{m_n} (C)_{m_1+\dots+m_n}^{-1} \frac{x_1^{m_1} \cdots x_n^{m_n}}{m_1! \cdots m_n!}; \end{aligned} \quad (3.2)$$

$$\begin{aligned} & F_{\mathcal{C}}[A, B; C_1, \dots, C_n; x_1, \dots, x_n] \\ &= \sum_{m_1, \dots, m_n=0}^{\infty} (A)_{m_1+\dots+m_n} (B)_{m_1+\dots+m_n} (C_1)_{m_1}^{-1} \cdots (C_n)_{m_n}^{-1} \frac{x_1^{m_1} \cdots x_n^{m_n}}{m_1! \cdots m_n!}; \end{aligned} \quad (3.3)$$

$$\begin{aligned} & F_{\mathcal{D}}[A, B_1, \dots, B_n; C; x_1, \dots, x_n] \\ &= \sum_{m_1, \dots, m_n=0}^{\infty} (A)_{m_1+\dots+m_n} (B_1)_{m_1} \cdots (B_n)_{m_n} (C)_{m_1+\dots+m_n}^{-1} \frac{x_1^{m_1} \cdots x_n^{m_n}}{m_1! \cdots m_n!}; \end{aligned} \quad (3.4)$$

where $A, A_1, \dots, A_n, B, B_1, \dots, B_n, C, C_1, \dots, C_n$ are matrices in $\mathbb{C}^{r \times r}$ such that $C + kI, C_i + kI, 1 \leq i \leq n$ are invertible for all integers $k \geq 0$ and x_1, \dots, x_n are complex variables.

We give the system of partial matrix differential equations of bilateral type satisfied by generalized Lauricella matrix functions under certain conditions. The following theorem gives the system of partial matrix differential equations of bilateral type satisfied by the matrix function $F_{\mathcal{A}}$ defined in (3.1).

Theorem 3.1. *Let A, B_i, C_i be matrices in $\mathbb{C}^{r \times r}$ such that $B_i B_j = B_j B_i, C_i B_j = B_j C_i$ and $C_i C_j = C_j C_i$, for each $i, j = 1, \dots, n$. Then the matrix function $F_{\mathcal{A}}$ satisfies the following system of partial matrix differential equations of bilateral type*

$$\begin{aligned} & x_i(1-x_i)U_{x_i x_i} - x_i x_1 U_{x_i x_1} - \cdots - x_i x_{i-1} U_{x_i x_{i-1}} - x_i x_{i+1} U_{x_i x_{i+1}} - \cdots - x_i x_n U_{x_i x_n} \\ & - x_i(A+I)U_{x_i} - (x_1 U_{x_1} + \cdots + x_n U_{x_n})B_i + U_{x_i} C_i - A U B_i = 0, \quad i = 1, \dots, n. \end{aligned} \quad (3.5)$$

Proof. Let

$$U = F_{\mathcal{A}}(A, B_1, \dots, B_n; C_1, \dots, C_n; x_1, \dots, x_n) = \sum_{m_1, \dots, m_n=0}^{\infty} \mathcal{A}_{m_1, \dots, m_n} x_1^{m_1} \cdots x_n^{m_n}. \quad (3.6)$$

Then, we have

$$\begin{aligned} & x_i(1-x_i)U_{x_i x_i} - x_i x_1 U_{x_i x_1} - \cdots - x_i x_{i-1} U_{x_i x_{i-1}} - x_i x_{i+1} U_{x_i x_{i+1}} - \cdots - x_i x_n U_{x_i x_n} \\ &= \sum_{m_1, \dots, m_n=0}^{\infty} (A + (m_1 + \cdots + m_n)I) \mathcal{A}_{m_1, \dots, m_n} m_i (B_i + m_i I) (C_i + m_i I)^{-1} \\ & \times x_1^{m_1} \cdots x_n^{m_n} - \sum_{m_1, \dots, m_n=0}^{\infty} m_i (m_i - 1) \mathcal{A}_{m_1, \dots, m_n} x_1^{m_1} \cdots x_n^{m_n} \end{aligned}$$

$$\begin{aligned}
& - \sum_{m_1, \dots, m_n=0}^{\infty} (m_1 m_1 + \dots + m_i m_{i-1} + m_i m_{i+1} + \dots + m_i m_n) \mathcal{A}_{m_1, \dots, m_n} x_1^{m_1} \dots x_n^{m_n} \\
& = A \sum_{m_1, \dots, m_n=0}^{\infty} \mathcal{A}_{m_1, \dots, m_n} x_1^{m_1} \dots x_n^{m_n} B_i - \sum_{m_1, \dots, m_n=0}^{\infty} (A + (m_1 + \dots + m_n)I) \mathcal{A}_{m_1, \dots, m_n} \\
& \quad \times x_1^{m_1} \dots x_n^{m_n} C_i (B_i + m_i I) (C_i + m_i I)^{-1} + (A + I) \sum_{m_1, \dots, m_n=0}^{\infty} \mathcal{A}_{m_1, \dots, m_n} \\
& \quad \times m_i x_1^{m_1} \dots x_n^{m_n} + \sum_{m_1, \dots, m_n=0}^{\infty} (m_i m_1 + \dots + m_i m_n) \mathcal{A}_{m_1, \dots, m_n} x_1^{m_1} \dots x_n^{m_n} B_i \\
& = x_i (A + I) U_{x_i} + (x_1 U_{x_1} + \dots + x_n U_{x_n}) B_i - U_{x_i} C_i + A U B_i. \tag{3.7}
\end{aligned}$$

This completes the proof. \square

In the next three theorems, we give the system of partial matrix differential equations of bilateral type satisfied by remaining three generalized Lauricella matrix functions. The proofs are similar to Theorem 3.1 and are omitted.

Theorem 3.2. *Let A_i, B_i, C be matrices in $\mathbb{C}^{r \times r}$ such that $A_i A_j = A_j A_i$, $B_i B_j = B_j B_i$ and $C B_j = B_j C$, for each $i, j = 1, \dots, n$. Then the matrix function F_B satisfies the following system of partial matrix differential equations of bilateral type*

$$\begin{aligned}
& x_i (1 - x_i) U_{x_i x_i} + x_1 U_{x_i x_1} + \dots + x_{i-1} U_{x_i x_{i-1}} + x_{i+1} U_{x_i x_{i+1}} + \dots + x_n U_{x_i x_n} \\
& - x_i (A_i + I) U_{x_i} - x_i U_{x_i} B_i + U_{x_i} C - A_i U B_i = 0, \quad i = 1, \dots, n. \tag{3.8}
\end{aligned}$$

Theorem 3.3. *Let A, B, C_i be matrices in $\mathbb{C}^{r \times r}$ such that $C_i C_j = C_j C_i$ and $C_j B = B C_j$, for each $i, j = 1, \dots, n$. Then the matrix function F_C satisfies a system of bilateral type partial matrix differential equations*

$$\begin{aligned}
& x_i (1 - x_i) U_{x_i x_i} - x_1^2 U_{x_1 x_1} - \dots - x_{i-1}^2 U_{x_{i-1} x_{i-1}} - x_{i+1}^2 U_{x_{i+1} x_{i+1}} - \dots - x_n^2 U_{x_n x_n} - 2x_1 x_2 \\
& \quad \times U_{x_1 x_2} - \dots - 2x_r x_s U_{x_r x_s} - \dots - 2x_{n-1} x_n U_{x_{n-1} x_n} - (A + I) (x_1 U_{x_1} + \dots + x_n U_{x_n}) \\
& \quad + U_{x_i} C_i - (x_1 U_{x_1} + \dots + x_n U_{x_n}) B - A U B = 0, \quad (r \neq s, r, s, i = 1, \dots, n). \tag{3.9}
\end{aligned}$$

Theorem 3.4. *Let A, B_i, C be matrices in $\mathbb{C}^{r \times r}$ such that $B_i B_j = B_j B_i$ and $C B_j = B_j C$, for each $i, j = 1, \dots, n$. Then the matrix function F_D satisfies a system of bilateral type partial matrix differential equations*

$$\begin{aligned}
& x_i (1 - x_i) U_{x_i x_i} + x_1 (1 - x_1) U_{x_i x_1} + \dots + x_{i-1} (1 - x_{i-1}) U_{x_i x_{i-1}} + x_{i+1} (1 - x_{i+1}) U_{x_i x_{i+1}} \\
& \quad + \dots + x_n (1 - x_n) U_{x_i x_n} - (A + I) x_i U_{x_i} + U_{x_i} C - (x_1 U_{x_1} + \dots + x_n U_{x_n}) B_i \\
& - A U B_i = 0, \quad i = 1, \dots, n. \tag{3.10}
\end{aligned}$$

3.1 Region of convergence

We give here the regions of convergence of the four Lauricella matrix functions. We give below the proof for the Lauricella matrix function F_A and present the remaining results without proof.

Theorem 3.5. *Let $A, B_1, \dots, B_n, C_1, \dots, C_n$ be positive stable matrices in $\mathbb{C}^{r \times r}$ such that $\alpha(A) < 1, \alpha(B_1) < \beta(C_1), \dots, \alpha(B_n) < \beta(C_n)$. Then the series F_A defined in (3.1) converges absolutely for $|x_1| + \dots + |x_n| < 1$.*

Proof. Let $\mathcal{A}_{m_1, \dots, m_n} x_1^{m_1} \dots x_n^{m_n}$ denote the general term of the series F_A . Then using (2.4), we get

$$\| \mathcal{A}_{m_1, \dots, m_n} x_1^{m_1} \dots x_n^{m_n} \| \leq N \| (m_1 + \dots + m_n)^A \| \prod_{i=1}^n (\| (m_i)^{B_i} \| \| (m_i)^{-C_i} \|)$$

$$\times \frac{(m_1 + \cdots + m_n - 1)!}{m_1! \cdots m_n!} |x_1|^{m_1} \cdots |x_n|^{m_n}, \quad (3.11)$$

where $N = \|\Gamma^{-1}(A)\| \|\Gamma^{-1}(B_1)\| \cdots \|\Gamma^{-1}(B_n)\| \|\Gamma(C_1)\| \cdots \|\Gamma(C_n)\|$. The Schur decomposition (2.7) yields

$$\|(m_1 + \cdots + m_n)^A \prod_{i=1}^n (\|(m_i)^{B_i}\| \|(m_i)^{-C_i}\|)\| \leq S (m_1 + \cdots + m_n)^{\alpha(A)} \prod_{i=1}^n (m_i)^{\alpha(B_i) - \beta(C_i)}, \quad (3.12)$$

where

$$S = \sum_{j=0}^{r-1} \frac{1}{j!} \left(\|A\| r^{1/2} \ln(m_1 + \cdots + m_n) \right)^j \prod_{i=1}^n \left(\sum_{j=0}^{r-1} \frac{1}{j!} \left(\max\{\|B_i\|, \|C_i\|\} r^{1/2} \ln m_i \right)^j \right)^2. \quad (3.13)$$

Using Equation (3.12) in (3.11), we get

$$\begin{aligned} & \|\mathcal{A}_{m_1, \dots, m_n} x_1^{m_1} \cdots x_n^{m_n}\| \\ & \leq N S (m_1 + \cdots + m_n)^{\alpha(A) - 1} \prod_{i=1}^n (m_i)^{\alpha(B_i) - \beta(C_i)} (|x_1| + \cdots + |x_n|)^{m_1 + \cdots + m_n}. \end{aligned} \quad (3.14)$$

Hence, for $\alpha(A) < 1, \alpha(B_1) < \beta(C_1), \dots, \alpha(B_n) < \beta(C_n)$ and $|x_1| + \cdots + |x_n| < 1$, $\|\mathcal{A}_{m_1, \dots, m_n} x_1^{m_1} \cdots x_n^{m_n}\| \rightarrow 0$ as $m_1, \dots, m_n \rightarrow \infty$. Therefore the series $F_{\mathcal{A}}$ converges absolutely. \square

We remark that the region of convergence of matrix functions $F_{\mathcal{A}}$ obtained above is same as in [6], where all the matrix considered were commuting. Next three theorems give the regions of convergence for generalized Lauricella matrix functions $F_{\mathcal{B}}, F_{\mathcal{C}}$ and $F_{\mathcal{D}}$. The proofs are similar to Theorem 3.5 and are omitted.

Theorem 3.6. *Let $A_1, \dots, A_n, B_1, \dots, B_n$ and C be positive stable matrices in $\mathbb{C}^{r \times r}$ such that $\alpha(A_1) + \alpha(B_1) < 2, \dots, \alpha(A_n) + \alpha(B_n) < 2, \beta(C) > 1$. Then the series $F_{\mathcal{B}}$ defined in (3.2) converges absolutely for $|x_1|, \dots, |x_n| < 1$.*

Theorem 3.7. *Let A, B, C_1, \dots, C_n be positive stable matrices in $\mathbb{C}^{r \times r}$ such that $\alpha(A) + \alpha(B) < 2, \beta(C_1) > 1, \dots, \beta(C_n) > 1$. Then the series $F_{\mathcal{C}}$ defined in (3.3) converges absolutely for $\sqrt{|x_1|} + \cdots + \sqrt{|x_n|} < 1$.*

Theorem 3.8. *Let $A, B_1, \dots, B_n, C \in \mathbb{C}^{r \times r}$ be positive stable matrices such that $\alpha(A) < \beta(C), \alpha(B_1) < 1, \dots, \alpha(B_n) < 1$. Then the series $F_{\mathcal{D}}$ defined in (3.4) converges absolutely for $|x_1|, \dots, |x_n| < 1$.*

3.2 Integral representations

Theorem 3.9. *Let A, B_i, C_i be matrices in $\mathbb{C}^{r \times r}$ such that $B_i B_j = B_j B_i, C_i B_j = B_j C_i, C_i C_j = C_j C_i$ and $B_i, C_i, C_i - B_i$ are positive stable for each $i, j = 1, \dots, n$. Then for $|x_1| \leq r_1, \dots, |x_n| \leq r_n, r_1 + \cdots + r_n < 1$, the series $F_{\mathcal{A}}$ defined in (3.1) can be put in the integral form as*

$$\begin{aligned} & F_{\mathcal{A}}[A, B_1, \dots, B_n; C_1, \dots, C_n; x_1, \dots, x_n] \\ & = \underbrace{\int_0^1 \cdots \int_0^1}_{n\text{-fold}} (1 - (u_1 x_1 + \cdots + u_n x_n))^{-A} \prod_{i=1}^n u_i^{B_i - I} (1 - u_i)^{C_i - B_i - I} \\ & \quad \times du_1 \cdots du_n \Gamma \left(\begin{matrix} C_1, \dots, C_n \\ B_1, \dots, B_n, C_1 - B_1, \dots, C_n - B_n \end{matrix} \right). \end{aligned} \quad (3.15)$$

Theorem 3.10. Let A_i, B_i, C be matrices in $\mathbb{C}^{r \times r}$ such that $B_i B_j = B_j B_i$, $C B_j = B_j C$ for each $i, j = 1, \dots, n$ and B_1, \dots, B_n, C and $C - (B_1 + \dots + B_n)$ are positive stable. Then for $|x_1|, \dots, |x_n| < 1$, the series F_B defined in (3.2), can be put in the integral form as

$$\begin{aligned} & F_B[A_1, \dots, A_n, B_1, \dots, B_n; C; x_1, \dots, x_n] \\ &= \underbrace{\int \dots \int}_{n\text{-fold}} (1 - u_1 x_1)^{-A_1} \dots (1 - u_n x_n)^{-A_n} u_1^{B_1 - I} \dots u_n^{B_n - I} (1 - (u_1 + \dots + u_n))^{C - (B_1 + \dots + B_n) - I} \\ & \quad \times du_1 \dots du_n \Gamma \left(\begin{matrix} C \\ B_1, \dots, B_n, C - (B_1 + \dots + B_n) \end{matrix} \right), \quad u_1 \geq 0, \dots, u_n \geq 0, \quad u_1 + \dots + u_n \leq 1. \end{aligned} \quad (3.16)$$

Theorem 3.11. Let C be a matrix in $\mathbb{C}^{r \times r}$ such that $C B_i = B_i C$, $A C = C A$, for each i and let A, C and $C - A$ be positive stable. Then for $|x_1|, \dots, |x_n| < 1$, the series F_D defined in (3.4) can be put in the integral form as

$$\begin{aligned} & F_D[A, B_1, \dots, B_n; C; x_1, \dots, x_n] \\ &= \Gamma \left(\begin{matrix} C \\ A, C - A \end{matrix} \right) \int_0^1 u^{A - I} (1 - u)^{C - A - I} (1 - u x_1)^{-B_1} \dots (1 - u x_n)^{-B_n} du. \end{aligned} \quad (3.17)$$

Theorem 3.12. Let A, B_i, C be matrices in $\mathbb{C}^{r \times r}$ such that $B_i B_j = B_j B_i$ and $C B_j = B_j C$, for each $i, j = 1, \dots, n$ and let $B_1, \dots, B_n, C, C - (B_1 + \dots + B_n)$ be positive stable. Then for $|x_1| \leq r_1, \dots, |x_n| \leq r_n, r_1 + \dots + r_n < 1$, the series F_D defined in (3.4) can be put in the integral form as

$$\begin{aligned} & F_D[A, B_1, \dots, B_n; C; x_1, \dots, x_n] \\ &= \underbrace{\int \dots \int}_{n\text{-fold}} (1 - (u_1 x_1 + \dots + u_n x_n))^{-A} u_1^{B_1 - I} \dots u_n^{B_n - I} (1 - (u_1 + \dots + u_n))^{C - (B_1 + \dots + B_n) - I} \\ & \quad \times du_1 \dots du_n \Gamma \left(\begin{matrix} C \\ B_1, \dots, B_n, C - (B_1 + \dots + B_n) \end{matrix} \right), \quad u_1 \geq 0, \dots, u_n \geq 0, \quad u_1 + \dots + u_n \leq 1. \end{aligned} \quad (3.18)$$

The proofs of these results are similar to the corresponding proofs of integral representations of Appell matrix functions F_2, F_3 and F_1 respectively, [5]. As such, the proofs are omitted. We remark that the following lemma is used in the proofs of Theorems 3.10 and 3.12

Lemma 3.1. [6] Let A_1, \dots, A_n, C be commuting matrices in $\mathbb{C}^{r \times r}$ such that $A_1, \dots, A_n, C, A_1 + \dots + A_n + C$ are positive stable and $A_1 + \dots + A_n + C + kI$ is invertible for all integers $k \geq 0$. Then

$$\begin{aligned} & \underbrace{\int \dots \int}_{n\text{-fold}} u_1^{A_1 - I} \dots u_n^{A_n - I} (1 - u_1 - \dots - u_n)^{C - I} du_1 \dots du_n = \Gamma \left(\begin{matrix} A_1, \dots, A_n, C \\ A_1 + \dots + A_n + C \end{matrix} \right) \\ & \quad u_1 \geq 0, \dots, u_n \geq 0, \quad u_1 + \dots + u_n \leq 1. \end{aligned} \quad (3.19)$$

We have not given the integral representation for generalized Lauricella matrix function F_C because the matrix function F_C does not give a simple form for integral in the arguments x_1, \dots, x_n .

4 Lauricella matrix functions

There are fourteen Lauricella matrix functions of three variables denoted by F_1, \dots, F_{14} . Out of these, F_1, F_2, F_5 and F_9 are particular cases of generalized Lauricella matrix functions F_A, F_B, F_C and F_D

respectively for $n = 3$. We give below the definition of remaining ten Lauricella matrix functions, *viz.*, $F_3, F_4, F_6, F_7, F_8, F_{10}, F_{11}, F_{12}, F_{13}, F_{14}$ and discuss their regions of convergence. Let A_i, B_i and C_i , $1 \leq i \leq 3$, be matrices in $\mathbb{C}^{r \times r}$ such that each $C_i + kI$ is invertible for all integers $k \geq 0$. Then the Lauricella matrix functions are defined by

$$\begin{aligned} & F_3(A_1, A_2, A_2, B_1, B_2, B_1; C_1, C_2, C_3; x, y, z) \\ &= \sum_{m,n,p=0}^{\infty} (A_1)_m (A_2)_{n+p} (B_1)_{m+p} (B_2)_n (C_1)_m^{-1} (C_2)_n^{-1} (C_3)_p^{-1} \frac{x^m y^n z^p}{m! n! p!}; \end{aligned} \quad (4.1)$$

$$\begin{aligned} & F_4(A_1, A_1, A_1, B_1, B_2, B_2; C_1, C_2, C_3; x, y, z) \\ &= \sum_{m,n,p=0}^{\infty} (A_1)_{m+n+p} (B_1)_m (B_2)_{n+p} (C_1)_m^{-1} (C_2)_n^{-1} (C_3)_p^{-1} \frac{x^m y^n z^p}{m! n! p!}; \end{aligned} \quad (4.2)$$

$$\begin{aligned} & F_6(A_1, A_2, A_3, B_1, B_2, B_1; C_1, C_2, C_2; x, y, z) \\ &= \sum_{m,n,p=0}^{\infty} (A_1)_m (A_2)_n (A_3)_p (B_1)_{m+p} (B_2)_n (C_1)_m^{-1} (C_2)_{n+p}^{-1} \frac{x^m y^n z^p}{m! n! p!}; \end{aligned} \quad (4.3)$$

$$\begin{aligned} & F_7(A_1, A_2, A_2, B_1, B_2, B_3; C_1, C_1, C_1; x, y, z) \\ &= \sum_{m,n,p=0}^{\infty} (A_1)_m (A_2)_{n+p} (B_1)_m (B_2)_n (B_3)_p (C_1)_{m+n+p}^{-1} \frac{x^m y^n z^p}{m! n! p!}; \end{aligned} \quad (4.4)$$

$$\begin{aligned} & F_8(A_1, A_1, A_1, B_1, B_2, B_3; C_1, C_2, C_2; x, y, z) \\ &= \sum_{m,n,p=0}^{\infty} (A_1)_{m+n+p} (B_1)_m (B_2)_n (B_3)_p (C_1)_m^{-1} (C_2)_{n+p}^{-1} \frac{x^m y^n z^p}{m! n! p!}; \end{aligned} \quad (4.5)$$

$$\begin{aligned} & F_{10}(A_1, A_2, A_1, B_1, B_2, B_1; C_1, C_2, C_2; x, y, z) \\ &= \sum_{m,n,p=0}^{\infty} (A_1)_{m+p} (A_2)_n (B_1)_{m+p} (B_2)_n (C_1)_m^{-1} (C_2)_{n+p}^{-1} \frac{x^m y^n z^p}{m! n! p!}; \end{aligned} \quad (4.6)$$

$$\begin{aligned} & F_{11}(A_1, A_2, A_2, B_1, B_2, B_1; C_1, C_2, C_2; x, y, z) \\ &= \sum_{m,n,p=0}^{\infty} (A_1)_m (A_2)_{n+p} (B_1)_{m+p} (B_2)_n (C_1)_m^{-1} (C_2)_{n+p}^{-1} \frac{x^m y^n z^p}{m! n! p!}; \end{aligned} \quad (4.7)$$

$$\begin{aligned} & F_{12}(A_1, A_2, A_1, B_1, B_1, B_2; C_1, C_2, C_2; x, y, z) \\ &= \sum_{m,n,p=0}^{\infty} (A_1)_{m+p} (A_2)_n (B_1)_{m+n} (B_2)_p (C_1)_m^{-1} (C_2)_{n+p}^{-1} \frac{x^m y^n z^p}{m! n! p!}; \end{aligned} \quad (4.8)$$

$$\begin{aligned} & F_{13}(A_1, A_2, A_2, B_1, B_2, B_1; C_1, C_1, C_1; x, y, z) \\ &= \sum_{m,n,p=0}^{\infty} (A_1)_m (A_2)_{n+p} (B_1)_{m+p} (B_2)_n (C_1)_{m+n+p}^{-1} \frac{x^m y^n z^p}{m! n! p!}; \end{aligned} \quad (4.9)$$

$$\begin{aligned}
& F_{14}(A_1, A_1, A_1, B_1, B_2, B_1; C_1, C_2, C_2; x, y, z) \\
&= \sum_{m,n,p=0}^{\infty} (A_1)_{m+n+p} (B_1)_{m+p} (B_2)_n (C_1)_m^{-1} (C_2)_{n+p}^{-1} \frac{x^m y^n z^p}{m! n! p!}. \tag{4.10}
\end{aligned}$$

The matrix analogues of the three Srivastava's triple hypergeometric functions, [15], are given by

$$H_{\mathcal{A}}(A, B, B'; C, C'; x, y, z) = \sum_{m,n,p=0}^{\infty} (A)_{m+p} (B)_{m+n} (B')_{n+p} (C)_m^{-1} (C')_{n+p}^{-1} \frac{x^m y^n z^p}{m! n! p!}, \tag{4.11}$$

$$\begin{aligned}
& H_{\mathcal{B}}(A, B, B'; C, C', C''; x, y, z) \\
&= \sum_{m,n,p=0}^{\infty} (A)_{m+p} (B)_{m+n} (B')_{n+p} (C)_m^{-1} (C')_n^{-1} (C'')_p^{-1} \frac{x^m y^n z^p}{m! n! p!}, \tag{4.12}
\end{aligned}$$

$$H_{\mathcal{C}}(A, B, B'; C; x, y, z) = \sum_{m,n,p=0}^{\infty} (A)_{m+p} (B)_{m+n} (B')_{n+p} (C)_{m+n+p}^{-1} \frac{x^m y^n z^p}{m! n! p!}, \tag{4.13}$$

where A, B, B', C, C' and C'' are matrices in $\mathbb{C}^{r \times r}$ such that $C + kI, C' + kI$ and $C'' + kI$ are invertible for all integers $k \geq 0$.

We remark that the regions of convergence of Lauricella matrix functions and Srivastava's triple hypergeometric matrix functions defined above remain the same as in [6], with proofs modified as illustrated in Theorem 3.5. As such, these results are omitted. The following theorem gives the region of convergence of Lauricella matrix function F_3 .

Theorem 4.1. *Let $A_1, A_2, B_1, B_2, C_1, C_2$ and C_3 be positive stable matrices in $\mathbb{C}^{r \times r}$ such that $\alpha(A_1) < \beta(C_1)$, $\alpha(A_2) < 1$, $\alpha(B_1) < 1$, $\alpha(B_2) < \beta(C_2)$, $\beta(C_3) > 1$. Then the series F_3 defined in (4.1) converges absolutely for $|x| < r$, $|y| < s$, $|z| < t$, $(1-r)(1-s) = t$.*

Proof. Let $\mathcal{A}_{m,n,p} x^m y^n z^p$ denote the general term of the series F_3 . Then

$$\begin{aligned}
& \|\mathcal{A}_{m,n,p} x^m y^n z^p\| \\
& \leq \|(A_1)_m\| \|(A_2)_{n+p}\| \|(B_1)_{m+p}\| \|(B_2)_n\| \|(C_1)_m^{-1}\| \|(C_2)_n^{-1}\| \|(C_3)_p^{-1}\| \left| \frac{x^m y^n z^p}{m! n! p!} \right| \\
& \leq \left\| \frac{(A_1)_m m^{A_1} m^{-A_1} (m-1)!}{(m-1)!} \right\| \left\| \frac{(A_2)_{n+p} (n+p)^{A_2} (n+p)^{-A_2} (n+p-1)!}{(n+p-1)!} \right\| \\
& \quad \times \left\| \frac{(B_1)_{m+p} (m+p)^{B_1} (m+p)^{-B_1} (m+p-1)!}{(m+p-1)!} \right\| \left\| \frac{(B_2)_n n^{B_2} n^{-B_2} (n-1)!}{(n-1)!} \right\| \\
& \quad \times \left\| \frac{(C_1)_m^{-1} m^{C_1} m^{-C_1} (m-1)!}{(m-1)!} \right\| \left\| \frac{(C_2)_n^{-1} n^{C_2} n^{-C_2} (n-1)!}{(n-1)!} \right\| \left\| \frac{(C_3)_p p^{C_3} p^{-C_3} (p-1)!}{(p-1)!} \right\| \\
& \quad \times \frac{|x|^m |y|^n |z|^p}{m! n! p!}. \tag{4.14}
\end{aligned}$$

Using $\Gamma(A) = \lim_{n \rightarrow \infty} (n-1)! (A)_n^{-1} n^A$, [11], we get

$$\begin{aligned}
\|\mathcal{A}_{m,n,p} x^m y^n z^p\| & \leq N \|m^{A_1}\| \|(n+p)^{A_2}\| \|(m+p)^{B_1}\| \|n^{B_2}\| \|m^{-C_1}\| \|n^{-C_2}\| \|p^{-C_3}\| \\
& \quad \times \frac{(n+p-1)! (m+p-1)!}{m! n! p! (p-1)!} |x|^m |y|^n |z|^p, \tag{4.15}
\end{aligned}$$

where $N = \|\Gamma^{-1}(A_1)\| \|\Gamma^{-1}(A_2)\| \|\Gamma^{-1}(B_1)\| \|\Gamma^{-1}(B_2)\| \|\Gamma(C_1)\| \|\Gamma(C_2)\| \|\Gamma(C_3)\|$. The Schur decomposition, [9, 17], yields

$$\|m^{A_1}\| \|(n+p)^{A_2}\| \|(m+p)^{B_1}\| \|n^{B_2}\| \|m^{-C_1}\| \|n^{-C_2}\| \|p^{-C_3}\|$$

$$\leq S m^{\alpha(A_1)-\beta(C_1)} n^{\alpha(B_2)-\beta(C_2)} p^{-\beta(C_3)} (m+p)^{\alpha(B_1)} (n+p)^{\alpha(A_2)}, \quad (4.16)$$

where

$$\begin{aligned} S &= \left(\sum_{j=0}^{r-1} \frac{1}{j!} \left(\max\{\|A_1\|, \|C_1\|\} r^{1/2} \ln m \right)^j \right)^2 \left(\sum_{j=0}^{r-1} \frac{1}{j!} \left(\max\{\|B_2\|, \|C_2\|\} r^{1/2} \ln n \right)^j \right)^2 \\ &\quad \times \sum_{j=0}^{r-1} \frac{1}{j!} \left(\|C_3\| r^{1/2} \ln p \right)^j \sum_{j=0}^{r-1} \frac{1}{j!} \left(\|A_2\| r^{1/2} \ln(n+p) \right)^j \sum_{j=0}^{r-1} \frac{1}{j!} \left(\|B_2\| r^{1/2} \ln(m+p) \right)^j. \end{aligned} \quad (4.17)$$

Equations (4.15) and (4.16) gives

$$\begin{aligned} \|\mathcal{A}_{m,n,p} x^m y^n z^p\| &\leq N S m^{\alpha(A_1)-\beta(C_1)} n^{\alpha(B_2)-\beta(C_2)} p^{1-\beta(C_3)} (m+p)^{\alpha(B_1)-1} \\ &\quad \times (n+p)^{\alpha(A_2)-1} \frac{(n+p)!(m+p)!}{m!n!p!} |x|^m |y|^n |z|^p. \end{aligned} \quad (4.18)$$

Hence, for $\alpha(A_1) < \beta(C_1)$, $\alpha(A_2) < 1$, $\alpha(B_1) < 1$, $\alpha(B_2) < \beta(C_2)$, $\beta(C_3) > 1$ and $|x| < r$, $|y| < s$, $|z| < t$, $(1-r)(1-s) = t$, $\|\mathcal{A}_{m,n,p} x^m y^n z^p\| \rightarrow 0$ as $m, n, p \rightarrow \infty$. Therefore the series F_3 converges absolutely. \square

We now find the system of partial matrix differential equations of bilateral type obeyed by the Lauricella matrix function F_3 .

Theorem 4.2. *Let $A_1, A_2, B_1, B_2, C_1, C_2, C_3$ be matrices in $\mathbb{C}^{r \times r}$ such that $A_1 A_2 = A_2 A_1$, $B_1 B_2 = B_2 B_1$, $B_i C_j = C_j B_i$ and $C_i C_j = C_j C_i$, for each $i, j = 1, 2, 3$. Then the system of partial matrix differential equations of bilateral type satisfied by the matrix function F_3 defined in (4.1) is*

$$x(1-x)U_{xx} - xzU_{xz} + U_x(C_1 - (B_1 + I)x) - A_1(xU_x + zU_z) - A_1UB_1 = 0, \quad (4.19)$$

$$y(1-y)U_{yy} - yzU_{yz} + U_y(C_2 - (B_2 + I)y) - zU_zB_2 - yA_2U_y - A_2UB_2 = 0, \quad (4.20)$$

$$\begin{aligned} z(1-z)U_{zz} - (xyU_{xy} + yzU_{yz} + xzU_{xz}) + U_z(C_3 - (B_1 + I)z) - yU_yB_1 \\ - A_2(xU_x + zU_z) - A_2UB_1 = 0. \end{aligned} \quad (4.21)$$

Proof. Let $U = F_3(A_1, A_2, A_2, B_1, B_2, B_1; C_1, C_2, C_3; x, y, z) = \sum_{m,n,p=0}^{\infty} U_{m,n,p} x^m y^n z^p$. Since $U(x, y, z)$ converges absolutely for $\alpha(A_1) < \beta(C_1)$, $\alpha(A_2) < 1$, $\alpha(B_1) < 1$, $\alpha(B_2) < \beta(C_2)$, $\beta(C_3) > 1$ and $|x| < r$, $|y| < s$, $|z| < t$, $(1-r)(1-s) = t$, so it is termwise differentiable in this domain and

$$\begin{aligned} U_x &= \sum_{m=1,n,p=0}^{\infty} m U_{m,n,p} x^{m-1} y^n z^p, \quad U_y = \sum_{m=0,n=1,p=0}^{\infty} n U_{m,n,p} x^m y^{n-1} z^p, \\ U_z &= \sum_{m=0,n=0,p=1}^{\infty} p U_{m,n,p} x^m y^n z^{p-1}, \quad U_{xx} = \sum_{m=2,n,p=0}^{\infty} m(m-1) U_{m,n,p} x^{m-2} y^n z^p, \\ U_{xy} &= \sum_{m=1,n=1,p=0}^{\infty} m n U_{m,n,p} x^{m-1} y^{n-1} z^p, \quad U_{xz} = \sum_{m=1,n=0,p=1}^{\infty} m p U_{m,n,p} x^{m-1} y^n z^{p-1}, \\ U_{yy} &= \sum_{m=0,n=2,p=0}^{\infty} n(n-1) U_{m,n,p} x^m y^{n-2} z^p, \quad U_{yz} = \sum_{m=0,n=1,p=1}^{\infty} n p U_{m,n,p} x^m y^{n-1} z^{p-1}, \\ U_{zz} &= \sum_{m=0,n=0,p=2}^{\infty} p(p-1) U_{m,n,p} x^m y^n z^{p-2}. \end{aligned} \quad (4.22)$$

This gives

$$\begin{aligned}
& x(1-x)U_{xx} - xzU_{xz} \\
&= \sum_{m,n,p=0}^{\infty} m(m+1)U_{m+1,n,p} x^m y^n z^p - \sum_{m,n,p=0}^{\infty} m(m-1)U_{m,n,p} x^m y^n z^p \\
&\quad - \sum_{m,n,p=0}^{\infty} mp U_{m,n,p} x^m y^n z^p \\
&= \sum_{m,n,p=0}^{\infty} (A_1 + mI)U_{m,n,p}(B_1 + (m+p)I)x^m y^n z^p - \sum_{m,n,p=0}^{\infty} m(m-1)U_{m,n,p} x^m y^n z^p \\
&\quad - \sum_{m,n,p=0}^{\infty} (A_1 + mI)U_{m,n,p}(B_1 + (m+p)I)(C_1 + mI)^{-1} C_1 x^m y^n z^p \\
&\quad - \sum_{m,n,p=0}^{\infty} mp U_{m,n,p} x^m y^n z^p \\
&= \sum_{m,n=0}^{\infty} (m+p)A_1 U_{m,n,p} x^m y^n z^p + \sum_{m,n,p=0}^{\infty} m U_{m,n,p} x^m y^n z^p (B_1 + I) \\
&\quad - \sum_{m,n,p=0}^{\infty} (A_1 + mI)U_{m,n,p}(B_1 + (m+p)I)(C_1 + mI)^{-1} C_1 x^m y^n z^p \\
&\quad + \sum_{m,n,p=0}^{\infty} A_1 U_{m,n,p} x^m y^n z^p B_1 \\
&= -U_x(C_1 - (B_1 + I)x) + A_1(xU_x + zU_z) + A_1UB_1, \tag{4.23}
\end{aligned}$$

completing the proof of Equation (4.19). Similarly we are able to show that the matrix function F_3 satisfies the bilateral type matrix differential equations (4.20) and (4.21). \square

The bilateral type partial matrix differential equations obeyed by remaining Lauricella matrix functions and Srivastava's triple hypergeometric matrix functions are tabulated in Table 1.

Functions	Systems of Matrix Differential equations	Conditions on Matrices
F_4	$ \begin{aligned} & x(1-x)U_{xx} - (xyU_{xy} + xzU_{xz}) - (yU_y + zU_z)B_1 \\ & \quad + U_x(C_1 - (B_1 + I)x) - xA_1U_x - A_1UB_1 = 0, \\ & y(1-y)U_{yy} - (xyU_{xy} + xzU_{xz} + 2yzU_{yz} + z^2U_{zz}) \\ & \quad - xU_xB_2 + U_y(C_2 - (B_2 + I)y) - A_1(yU_y + zU_z) \\ & \quad \quad - zU_z(B_2 + I) - A_1UB_2 = 0, \\ & z(1-z)U_{zz} - (xyU_{xy} + xzU_{xz} + 2yzU_{yz} + y^2U_{yy}) \\ & \quad - xU_xB_2 + U_z(C_3 - (B_2 + I)z) - A_1(yU_y + zU_z) \\ & \quad \quad - yU_y(B_2 + I) - A_1UB_2 = 0; \end{aligned} $	$ \begin{aligned} & B_1B_2 = B_2B_1, \\ & C_iC_j = C_jC_i, \\ & B_iC_j = C_jB_i \end{aligned} $

F_6	$x(1-x)U_{xx} - xzU_{xz} + U_x(C_1 - (B_1 + I)x)$ $-A_1(xU_x + zU_z) - A_1UB_1 = 0,$ $y(1-y)U_{yy} + zU_{yz} + U_y(C_2 - (B_2 + I)y)$ $-yA_2U_y - A_2UB_2 = 0,$ $z(1-z)U_{zz} - xzU_{xz} + yU_{yz} + U_z(C_2 - (B_1 + I)z)$ $-A_3(xU_x + zU_z) - A_3UB_1 = 0;$	$A_iA_j = A_jA_i$ $B_1B_2 = B_2B_1$ $C_iC_j = C_jC_i,$ $B_iC_j = C_jB_i$
F_7	$x(1-x)U_{xx} + yU_{xy} + zU_{xz} + U_x(C_1 - (B_1 + I)x)$ $-xA_1U_x - A_1UB_1 = 0,$ $y(1-y)U_{yy} + xU_{xy} + zU_{yz} - yzU_{yz} - zU_zB_2$ $+U_y(C_1 - (B_2 + I)y) - yA_2U_y - A_2UB_2 = 0,$ $z(1-z)U_{zz} + xU_{xz} + yU_{yz} - yzU_{yz} - yU_yB_3$ $+U_z(C_1 - (B_3 + I)z) - zA_2U_z - A_2UB_3 = 0;$	$A_1A_2 = A_2A_1,$ $B_iB_j = B_jB_i,$ $B_iC_1 = C_1B_i$
F_8	$x(1-x)U_{xx} - xyU_{xy} - xzU_{xz} - xA_1U_x +$ $U_x(C_1 - (B_1 + I)x) - (yU_y + zU_z)B_1 - A_1UB_1 = 0,$ $y(1-y)U_{yy} - xyU_{xy} - yzU_{yz} + zU_{yz} - yA_1U_y$ $-(xU_x + zU_z)B_2 + U_y(C_2 - (B_2 + I)y) - A_1UB_2 = 0,$ $z(1-z)U_{zz} - xzU_{xz} - yzU_{yz} + yU_{yz} - zA_1U_z$ $-(xU_x + yU_y)B_3 + U_z(C_2 - (B_3 + I)z) - A_1UB_3 = 0;$	$B_iB_j = B_jB_i,$ $C_1C_2 = C_2C_1,$ $B_iC_j = C_jB_i$
F_{10}	$x(1-x)U_{xx} - 2xzU_{xz} - z^2U_{zz} - zU_z(B_1 + I)$ $+U_x(C_1 - (B_1 + I)x) - A_1(xU_x + zU_z) - A_1UB_1 = 0,$ $y(1-y)U_{yy} + zU_{yz} + U_y(C_2 - (B_2 + I)y)$ $-yA_2U_y - A_2UB_2 = 0,$ $z(1-z)U_{zz} - 2xzU_{xz} + yU_{yz} - x^2u_{xx} + U_zC_2$ $-(xU_x + zU_z)(B_1 + I) - A_1(xU_x + zU_z) - A_1UB_1 = 0;$	$A_1A_2 = A_2A_1,$ $B_1B_2 = B_2B_1,$ $C_1C_2 = C_2C_1,$ $B_iC_j = C_jB_i$
F_{11}	$x(1-x)U_{xx} - xzU_{xz} + U_x(C_1 - (B_1 + I)x)$ $-A_1(xU_x + zU_z) - A_1UB_1 = 0,$ $y(1-y)U_{yy} - yzU_{yz} + zU_{yz} - zU_zB_2$ $+U_y(C_2 - (B_2 + I)y) - yA_2U_y - A_2UB_2 = 0,$ $z(1-z)U_{zz} - xyU_{xy} - yzU_{yz} - xzU_{xz} + yU_{yz} - yU_yB_1$ $+U_z(C_2 - (B_1 + I)z) - A_2(xU_x + zU_z) - A_2UB_1 = 0;$	$A_1A_2 = A_2A_1,$ $B_1B_2 = B_2B_1,$ $C_1C_2 = C_2C_1,$ $B_iC_j = C_jB_i$

F_{12}	$ \begin{aligned} & x(1-x)U_{xx} - xyU_{xy} - yzU_{yz} - xzU_{xz} - zU_zB_1 \\ & +U_x(C_1 - (B_1 + I)x) - A_1(xU_x + yU_y) - A_1UB_1 = 0, \\ & y(1-y)U_{yy} - xyU_{xy} + zU_{yz} + U_y(C_2 - (B_1 + I)y) \\ & \quad - A_2(xU_x + yU_y) - A_2UB_1 = 0, \\ & z(1-z)U_{zz} - xzU_{xz} + yU_{yz} + U_z(C_2 - (B_2 + I)z) \\ & \quad - xU_xB_2 - zA_1U_z - A_1UB_2 = 0; \end{aligned} $	$ \begin{aligned} A_1A_2 &= A_2A_1, \\ B_1B_2 &= B_2B_1, \\ C_1C_2 &= C_2C_1, \\ B_iC_j &= C_jB_i \end{aligned} $
F_{13}	$ \begin{aligned} & x(1-x)U_{xx} + yU_{xy} + zU_{xz} - xzU_{xz} \\ & +U_x(C_1 - (B_1 + I)x) - A_1(xU_x + zU_z) - A_1UB_1 = 0, \\ & y(1-y)U_{yy} + xU_{xy} + zU_{yz} - yzU_{yz} - zU_zB_2 \\ & +U_y(C_1 - (B_2 + I)y) - yA_2U_y - A_2UB_2 = 0, \\ & z(1-z)U_{zz} + xU_{xz} + yU_{yz} - xyU_{xy} - yzU_{yz} \\ & \quad - xzU_{xz} + U_z(C_1 - (B_1 + I)z) - yU_yB_1 \\ & \quad - A_2(xU_x + zU_z) - A_2UB_1 = 0; \end{aligned} $	$ \begin{aligned} A_1A_2 &= A_2A_1, \\ B_1B_2 &= B_2B_1, \\ B_iC_1 &= C_1B_i \end{aligned} $
F_{14}	$ \begin{aligned} & x(1-x)U_{xx} - xyU_{xy} - yzU_{yz} - 2xzU_{xz} - z^2U_{zz} \\ & +U_x(C_1 - (B_1 + I)x) - zU_z - (yU_y + zU_z)B_1 \\ & \quad - A_1(xU_x + zU_z) - A_1UB_1 = 0, \\ & y(1-y)U_{yy} - xyU_{xy} + zU_{yz} - yzU_{yz} - zU_zB_2 \\ & -xU_xB_2 + U_y(C_2 - (B_2 + I)y) - yA_1U_y - A_1UB_2 = 0, \\ & z(1-z)U_{zz} - 2xzU_{xz} + yU_{yz} - xyU_{xy} - yzU_{yz} \\ & \quad +U_z(C_2 - (B_1 + I)z) - x^2U_{xx} - xU_x(B_1 + I) \\ & \quad - yU_yB_1 - A_1(xU_x + zU_z) - A_1UB_1 = 0; \end{aligned} $	$ \begin{aligned} B_1B_2 &= B_2B_1, \\ C_1C_2 &= C_2C_1, \\ B_iC_j &= C_jB_i \end{aligned} $
H_A	$ \begin{aligned} & x(1-x)U_{xx} - xyU_{xy} - yzU_{yz} - xzU_{xz} - zU_zB \\ & +U_x(C - (B + I)x) - A(yU_y + xU_x) - AUB = 0, \\ & y(1-y)U_{yy} - xyU_{xy} - xzU_{xz} - yzU_{yz} + zU_{yz} - xU_xB' \\ & +U_y(C' - (B' + I)y) - (yU_y + zU_z)B - UBB' = 0, \\ & z(1-z)U_{zz} - xzU_{xz} - xyU_{xy} - yzU_{yz} + yU_{yz} - xU_xB' \\ & +U_z(C' - (B' + I)z) - A(yU_y + zU_z) - AUB' = 0; \end{aligned} $	$ \begin{aligned} & B, B', C, C' \\ & \text{are commuting} \end{aligned} $
H_B	$ \begin{aligned} & x(1-x)U_{xx} - xyU_{xy} - yzU_{yz} - xzU_{xz} - zU_zB \\ & +U_x(C - (B + I)x) - A(yU_y + xU_x) - AUB = 0, \\ & y(1-y)U_{yy} - xyU_{xy} - xzU_{xz} - yzU_{yz} - xU_xB' \\ & +U_y(C' - (B' + I)y) - (yU_y + zU_z)B - UBB' = 0, \\ & z(1-z)U_{zz} - xzU_{xz} - xyU_{xy} - yzU_{yz} - xU_xB' \\ & +U_z(C'' - (B' + I)z) - A(yU_y + zU_z) - AUB' = 0; \end{aligned} $	$ \begin{aligned} & B, B', C, C', \text{ and} \\ & C'' \text{ are commuting} \end{aligned} $

H_C	$ \begin{aligned} & x(1-x)U_{xx} - xyU_{xy} - yzU_{yz} - xzU_{xz} \\ & + yU_{xy} + zU_{xz} - zU_z B + U_x(C - (B + I)x) \\ & - A(yU_y + xU_x) - AUB = 0, \\ & y(1-y)U_{yy} - xyU_{xy} - xzU_{xz} - yzU_{yz} \\ & + xU_{xy} + zU_{yz} - xU_x B' + U_y(C - (B' + I)y) \\ & - (yU_y + zU_z)B - UBB' = 0, \\ & z(1-z)U_{zz} - xzU_{xz} - xyU_{xy} - yzU_{yz} \\ & + xU_{xz} + yU_{yz} + U_z(C - (B' + I)z) - xU_x B' \\ & - A(yU_y + zU_z) - AUB' = 0. \end{aligned} $	B, B' and C are commuting
-------	--	----------------------------------

Table 1: Systems of partial matrix differential equations of bilateral type satisfied by Lauricella and Srivastava matrix functions

Example 1. We show here that the conditions on the matrices given in the third column of Table 1 are necessary for systems of matrix differential equations of bilateral type to hold.

Consider the differential equation of the bilateral type

$$x(1-x)U_{xx} - xzU_{xz} + U_x(C_1 - (B_1 + I)x) - A_1(xU_x + zU_z) - A_1UB_1 = 0 \quad (4.24)$$

satisfied by the Lauricella matrix function F_3 . Let $U = F_3(A_1, A_2, A_2, B_1, B_2, B_1; C_1, C_2, C_3; x, y, z) = \sum_{m,n,p=0}^{\infty} U_{m,n,p} x^m y^n z^p$. Then, using the partial derivatives (4.22), we have

$$\begin{aligned}
& x(1-x)U_{xx} - xzU_{xz} + U_x C_1 \\
& = \sum_{m,n,p=0}^{\infty} (m+1)U_{m+1,n,p} x^m y^n z^p (C_1 + mI) - \sum_{m,n,p=0}^{\infty} m(m-1)U_{m,n,p} x^m y^n z^p \\
& - \sum_{m,n,p=0}^{\infty} m p U_{m,n,p} x^m y^n z^p
\end{aligned} \quad (4.25)$$

and

$$\begin{aligned}
& xU_x(B_1 + I) + A_1(xU_x + zU_z) + A_1UB_1 \\
& = \sum_{m,n,p=0}^{\infty} mU_{m,n,p} x^m y^n z^p (B_1 + I) + \sum_{m,n,p=0}^{\infty} A_1(m+p)U_{m,n,p} x^m y^n z^p \\
& + \sum_{m,n,p=0}^{\infty} A_1U_{m,n,p} x^m y^n z^p B_1.
\end{aligned} \quad (4.26)$$

In particular for $m = 1$ and $n = p = 0$, we get

$$x(1-x)U_{xx} - xzU_{xz} + U_x C_1 = A_1(A_1 + I)B_1(B_1 + I)C_1^{-1}x \quad (4.27)$$

and

$$xU_x(B_1 + I) + A_1(xU_x + zU_z) + A_1UB_1 = A_1(A_1 + I)B_1C_1^{-1}x(B_1 + I). \quad (4.28)$$

From (4.27) and (4.28), one can conclude that the equation (4.24) will hold if

$$A_1(A_1 + I)B_1(B_1 + I)C_1^{-1}x = A_1(A_1 + I)B_1C_1^{-1}x(B_1 + I). \quad (4.29)$$

This gives $B_1C_1 = C_1B_1$. Similarly, at $m = 0, n = 1, p = 0$, we have

$$x(1-x)U_{xx} - xzU_{xz} + U_xC_1 = A_1A_2B_1B_2C_1^{-1}C_2^{-1}C_1y \quad (4.30)$$

and

$$xU_x(B_1 + I) + A_1(xU_x + zU_z) + A_1UB_1 = A_1A_2B_2C_2^{-1}yB_1, \quad (4.31)$$

which gives $B_1B_2 = B_2B_1$, $C_1C_2 = C_2C_1$ and $B_1C_2 = C_2B_1$. The remaining conditions on the matrices can be obtained by changing the particular values of m, n and p .

In Table 2, we give the integral representations of Lauricella matrix functions of three variables and Srivastava's triple hypergeometric matrix functions.

Functions	Integral Representations	Conditions on Matrices
F_6	$\Gamma\left(\begin{matrix} C_1, C_2 \\ A_1, A_2, A_3, C_1 - A_1, C_2 - A_2 - A_3 \end{matrix}\right) \iiint u^{A_1-I} \\ \times v^{A_2-I} w^{A_3-I} (1-u)^{C_1-A_1-I} (1-v-w)^{C_2-A_2-A_3-I} \\ \times (1-vy)^{-B_2} (1-ux-wz)^{-B_1} du dv dw, \quad 0 \leq u \leq 1, \\ v \geq 0, w \geq 0, v+w \leq 1, r+t < 1, x \leq r, z \leq t, \\ \beta(A_1) > 0, \beta(A_2) > 0, \beta(A_3) > 0, \beta(C_1) > 0, \\ \beta(C_2) > 0, \beta(C_1 - A_1) > 0, \beta(C_2 - A_2 - A_3) > 0.$	$A_i A_j = A_j A_i, \\ B_i C_j = C_j B_i, \\ C_1 C_2 = C_2 C_1, \\ A_i C_j = C_j A_i$
F_7	$\iiint (1-ux)^{-A_1} (1-vy-wz)^{-A_2} u^{B_1-I} v^{B_2-I} w^{B_3-I} \\ \times (1-u-v-w)^{C_1-B_1-B_2-B_3-I} du dv dw \\ \times \Gamma\left(\begin{matrix} C_1 \\ B_1, B_2, B_3, C_1 - B_1 - B_2 - B_3 \end{matrix}\right), u \geq 0, \\ v \geq 0, w \geq 0, u+v+w \leq 1, s+t < 1, y \leq s, \\ z \leq t, \beta(B_1) > 0, \beta(B_2) > 0, \beta(B_3) > 0, \beta(C_1) > 0, \\ \beta(C_1 - B_1 - B_2 - B_3) > 0;$	$B_i B_j = B_j B_i, \\ B_i C_1 = C_1 B_i$
F_8	$\iiint (1-ux-vy-wz)^{-A_1} u^{B_1-I} v^{B_2-I} w^{B_3-I} \\ \times (1-u)^{C_1-B_1-I} (1-v-w)^{C_2-B_2-B_3-I} du dv dw, \\ \times \Gamma\left(\begin{matrix} C_1, C_2 \\ B_1, B_2, B_3, C_1 - B_1, C_2 - B_2 - B_3 \end{matrix}\right), 0 \leq u \leq 1, \\ v \geq 0, w \geq 0, v+w \leq 1, r+s+t < 1, x \leq r, y \leq s, \\ z \leq t, \beta(B_1) > 0, \beta(B_2) > 0, \beta(B_3) > 0, \beta(C_1) > 0, \\ \beta(C_2) > 0, \beta(C_1 - B_1) > 0, \beta(C_2 - B_2 - B_3) > 0.$	$B_i B_j = B_j B_i, \\ C_1 C_2 = C_2 C_1, \\ B_i C_j = C_j B_i$
F_{11}	$\Gamma\left(\begin{matrix} C_1, C_2 \\ A_1, A_2, C_1 - A_1, C_2 - A_2 \end{matrix}\right) \int_0^1 \int_0^1 u^{A_1-I} v^{A_2-I} \\ \times (1-u)^{C_1-A_1-I} (1-v)^{C_2-A_2-I} (1-ux-vz)^{-B_1} \\ \times (1-vy)^{-B_2} du dv, \quad r+t < 1, x \leq r, z \leq t, \\ \beta(A_1) > 0, \beta(A_2) > 0, \beta(C_1) > 0, \beta(C_2) > 0, \\ \beta(C_1 - A_1) > 0, \beta(C_2 - A_2) > 0.$	$A_1 A_2 = A_2 A_1, \\ B_i C_j = C_j B_i, \\ C_1 C_2 = C_2 C_1, \\ A_i C_j = C_j A_i$
F_{12}	$\iiint (1-vy)^{A_1} (1-ux-vy-wz+vwyz)^{-A_1} u^{B_1-I} v^{A_2-I} \\ \times w^{B_2-I} (1-u)^{C_1-B_1-I} (1-v-w)^{C_2-A_2-B_2-I} (1-vy)^{-B_1} \\ \times du dv dw \Gamma\left(\begin{matrix} C_1, C_2 \\ A_2, B_1, B_2, C_1 - B_1, C_2 - A_2 - B_2 \end{matrix}\right), \\ 0 \leq u \leq 1, v \geq 0, w \geq 0, v+w \leq 1, r+s+t < 1+st, \\ x \leq r, y \leq s, z \leq t, \beta(A_2) > 0, \beta(B_1) > 0, \\ \beta(B_2) > 0, \beta(C_1) > 0, \beta(C_2) > 0, \beta(C_1 - B_1) > 0, \\ \beta(C_2 - A_2 - B_2) > 0.$	$C_1, C_2, B_1, B_2 \text{ and } A_2 \\ \text{are commuting}$

F_{13}	$\iint (1-ux)^{-A_1} (1-vy-uz)^{-A_2} u^{B_1-I} v^{B_2-I} \times (1-u-v)^{C_1-B_1-B_2-I} du dv$ $\Gamma\left(\begin{matrix} C_1 \\ B_1, B_2, C_1 - B_1 - B_2 \end{matrix}\right), u \geq 0, v \geq 0,$ $u+v \leq 1, s+t < 1, y \leq s, z \leq t, \beta(B_1) > 0,$ $\beta(B_2) > 0, \beta(C_1) > 0, \beta(C_1 - B_1 - B_2) > 0.$	C_1, B_1 and B_2 are commuting
H_A	$\int_0^1 \int_0^1 (1-ux-vy-vz+v^2yz)^{-A} (1-vy)^A u^{B-I} \times v^{B'-I} (1-u)^{C-B-I} (1-v)^{C'-B'-I} (1-vy)^{-B} du dv$ $\Gamma\left(\begin{matrix} C, C' \\ B, B', C-B, C'-B' \end{matrix}\right), r+s+t < 1+t,$ $ x \leq r, y \leq s, z \leq t, \beta(B) > 0, \beta(B') > 0,$ $\beta(C) > 0, \beta(C') > 0, \beta(C-B) > 0, \beta(C'-B') > 0.$	B, B', C and C' are commuting
H_B	$\Gamma^{-1}(A)\Gamma^{-1}(B)\Gamma^{-1}(B') \int_0^\infty \int_0^\infty \int_0^\infty e^{-(u+v+w)} v^{A-I} \times u^{B-I} w^{B'-I} {}_0F_1(-; C; xuv) {}_0F_1(-; C'; yuw) {}_0F_1(-; C''; zvw) du dv dw,$ $ x \leq r, y \leq s, z \leq t, \max\{r, s, t\} < 1, \beta(A) > 0, \beta(B) > 0, \beta(B') > 0.$	B, B' commutes with A
H_C	$\Gamma\left(\begin{matrix} C \\ A, B, C-A-B \end{matrix}\right) \int_0^1 \int_0^1 u^{A-I} v^{B-I} (1-u)^{C-A-I} (1-v)^{C-A-B-I} (1-ux)^{-B} (1-ux)^{B'} (1-ux-vy-wz+uvy-zxu^2)^{-B'} du dv,$ $r+s+t+rt < 1+s, x \leq r, y \leq s, z \leq t,$ $\beta(A) > 0, \beta(B) > 0, \beta(C) > 0, \beta(C-A-B) > 0.$	A, B, C commutes and $B'C = CB'$

Table 2: Integral Representations of Lauricella and Srivastava Matrix Functions

References

- [1] M. Abdalla, *Special matrix functions: characteristics, achievements and future directions*, Linear Multilinear Algebra (2020), no. 1, 1–28.
- [2] A. Altin, B. Cekim, R. Sahin, *On the matrix versions of Appell hypergeometric functions*. Quaest. Math. 37 (2014), no. 1, 31–38.
- [3] H. Buchholz, *The confluent hypergeometric function*, Springer-Verlag, New York, 1969.
- [4] N. Dunford, J. Schwartz, *Linear Operators*, Part-I, New York: Addison-Wesley, 1957.
- [5] R. Dwivedi, V. Sahai, *On the hypergeometric matrix functions of two variables*. Linear Multilinear Algebra 66 (2018), no. 9, 1819–1837.
- [6] R. Dwivedi, V. Sahai, *On the hypergeometric matrix functions of several variables*, J. Math. Phys. 59 (2018), no. 2, 023505, 15pp.
- [7] R. Dwivedi, V. Sahai, *A note on the Appell matrix functions*. Quaest. Math. 43 (2020), no. 3, 321–334.
- [8] H. Exton, *On certain confluent hypergeometric functions of three variables*, Ganita 21 (1970), no. 2, 79–92.

- [9] G.H. Golub, C.F. van Loan, *Matrix Computations*, Johns Hopkins Univ. Press, Baltimore, MD, 1989.
- [10] G.D. Hu, M. Liu, *The weighted logarithmic matrix norm and bounds of the matrix exponential.*, Linear Algebra Appl. 2004;390:145–154.
- [11] L. Jodar, J.C. Cortes, *Some properties of gamma and beta matrix functions.* Appl. Math. Lett. 11 (1998), no. 1, 89–93.
- [12] L. Jodar, J.C. Cortes, *On the hypergeometric matrix function.* Proceedings of the VIIIth Symposium on Orthogonal Polynomials and Their Applications (Seville, 1997). J. Comput. Appl. Math. 99 (1998), no. 1-2, 205–217.
- [13] L. Jodar, J.C. Cortes, *Closed form general solution of the hypergeometric matrix differential equation.* Mathematical and computer modelling in engineering sciences. Math. Comput. Modelling 32 (2000), no. 9, 1017–1028.
- [14] W. Miller, *Lie Theory and Special Functions*, Academic Press, New York, 1968.
- [15] H. M. Srivastava, H. L. Manocha, *A Treatise on Generating Functions*, Ellis Horwood Series: Mathematics and its Applications. Ellis Horwood Ltd., Chichester; Halsted Press [John Wiley & Sons, Inc.], New York, 1984.
- [16] L.N. Trefethen, M. Embree, *Spectra and pseudospectra: the behaviour of nonnormal matrices and operators.* Princeton, NJ: Princeton University press; 2005.
- [17] C. Van Loan, *The sensitivity of the matrix exponential*, SIAM J. Numer. Anal. 14 (1977), no. 6, 971–981.