

# Bifurcating steady-state flows involving energy dissipation over Hartmann boundary layer

Zhi-Min Chen

*School of Mathematics and Statistics, Shenzhen University, Shenzhen 518060, China*

---

## Abstract

A plane non-parallel vortex flow in a square fluid domain is examined. The energy dissipation of the flow is dominated by viscosity and linear friction effect of a Hartmann layer. This is a traditional Navier-Stokes flow when the linear friction effect is not involved, whereas it is a magnetohydrodynamic flow when the energy dissipation is fundamentally dominated by the friction. It is proved that the critical values of the viscosity and friction with respect to its linearized spectral problem are nonlinear thresholds leading to the onset of secondary steady-state flows, the nonlinear phenomenon observed in laboratory experiments.

**Keywords:** Non-parallel flows, Navier-Stokes equations, bifurcation, vortex flows, linear friction effect of Hartmann layer

---

## 1. Introduction

To study the inverse energy cascade towards large scales [8] of plane flows, Sommeria and Verron [11, 12, 14] presented magnetohydrodynamic experiments by using electronically driven flows in a closed box, containing a thin horizontal layer of liquid metal. The box is bottomed with electromagnets producing a uniform vertical magnetic field. The flow velocity is small so that the upper free surface is negligible. The three-dimensional motion reduces a two-dimensional one as the vertical movement in the thin horizontal layer fluid can be ignored. The energy dissipation of the fluid motion counts for viscosity and the Hartmann layer friction applied on the bottom of liquid metal.

The non-dimensional governing equations of the two-dimensional approximation motion for the velocity  $\mathbf{v} = (v_1, v_2)$  and pressure  $p$  in the domain  $[0, 1] \times [0, 1]$  are [11, 12, 14]

$$\frac{\partial \mathbf{v}}{\partial t} + \mathbf{v} \cdot \nabla \mathbf{v} + \nabla p - \frac{1}{Re} \Delta \mathbf{v} + \frac{\mathbf{v}}{Rh} = \mathbf{f}, \quad \nabla \cdot \mathbf{v} = 0. \quad (1)$$

Here  $Re$  is the Reynolds number,  $Rh$  is the Rayleigh number measuring the Hartmann bottom friction and  $\mathbf{f}$  is the Lorentz driving force defined by electric currents so that

$$\nabla \times \mathbf{f} = \frac{\pi^2}{2} \sin(2\pi x) \sin(2\pi y) \quad \text{and} \quad \int_0^1 \int_0^1 |\nabla \times \mathbf{f}| dx dy = 2.$$

The stream function  $\psi$  and the vorticity of the fluid motion are defined as

$$v_1 = \frac{\partial \psi}{\partial y}, \quad v_2 = -\frac{\partial \psi}{\partial x}, \quad \omega = \nabla \times \mathbf{v} = -\Delta \psi.$$

---

*Email address:* zmchen@szu.edu.cn (Zhi-Min Chen)

The vorticity formulation of (1) expressed as

$$-\frac{\partial \Delta \psi}{\partial t} + J(\psi, \Delta \psi) - \frac{\Delta \psi}{Rh} + \frac{\Delta^2 \psi}{Re} = \frac{\pi^2}{2} \sin(2\pi x) \sin(2\pi y), \quad (2)$$

where the nonlinear convective term is written as the Jacobian

$$J(\psi, \Delta \psi) = \partial_x \psi \partial_y \Delta \psi - \partial_y \psi \partial_x \Delta \psi$$

The basic flow of (2) is dependant on the parameters  $Re$  and  $Rh$ . It is convenient to use the modified system [13] of (2) expressed as

$$-\partial_t \Delta \psi + J(\psi, \Delta \psi) + (-\mu \Delta + \nu \Delta^2)(\psi - \sin x \sin y) = 0, \quad (3)$$

As in [11, 12, 14, 13], it is convenience to accept the free slip boundary condition

$$\psi|_{\partial\Omega} = \Delta \psi|_{\partial\Omega} = 0 \quad (4)$$

for the modified square fluid domain  $\Omega = (0, 2\pi) \times (0, 2\pi)$ , and use the Fourier expansion for the stream function

$$\psi = \sum_{n,m \geq 1} a_{n,m} \sin \frac{nx}{2} \sin \frac{my}{2}. \quad (5)$$

The parameters  $\nu$  and  $\mu$  are defined by the transformations [4]

$$Rh = \frac{2\sqrt{\mu + 2\nu}}{\mu\pi}, \quad Re = \frac{8\pi\sqrt{\mu + 2\nu}}{\nu} \quad (6)$$

With this modification, we have the basic steady-state flow

$$\psi_0 = \sin x \sin y.$$

The basic flow exhibits four vortices in Figure 1. The experiments [11, 12, 14] show the transitions of the basic flow in the scenario of the inverse energy cascade towards to large scales. The principal transition amongst them is the steady-state bifurcation of  $\psi_0$  into secondary flows, the merging of two vortices. The secondary flows observed in the experiments of [12, 14] are sketched in Figure 1.

To the understanding of the transition, Thess [13] demonstrated critical stability parameters  $(\mu_c, \nu_c)$  of the linear spectral problem so that linear stable and unstable domains are defined. The author [4] provided nonlinear stability analysis of vortex flow and studied all possible linear spectral solutions and presented secondary flows as a result of nonlinear saturation of primary linear instability. The instability of vortices with respect to two vortex merging phenomena was also discussed by Meunier *et al.* [9] and Cerretelli and Williamson[2]. The experimental studies [11, 12] of the non-parallel flow  $\sin x \sin y$  is developed from the magnetohydrodynamic experiment of Bondarenko *et al.* [1] on the steady-state bifurcation of the parallel Kolmogorov flow  $\sin x$ . The existence of secondary steady-state flows and temporal periodic flows bifurcating from the Kolmogorov flow has been studied extensively [5, 6, 7].

However, the basic flow  $\psi_0$  is non-parallel and rigorous non-instability analysis for the existence of secondary flows is missing.

It is the purpose of present paper to show the existence of the secondary flows, which is to be contained in the Hilbert space

$$H^4 = \left\{ \psi = \sum_{n,m \geq 1} a_{n,m} \sin \frac{nx}{2} \sin \frac{my}{2} \mid \|\psi\|_{H^4}^2 = \sum_{n,m \geq 1} \left( 1 + \left( \frac{n^2 + m^2}{4} \right)^2 \right) |a_{n,m}|^2 < \infty \right\}.$$




Figure 1: (a) The basic steady-state flow  $\psi_0$ ; (b) the contour lines of the function  $\psi_0 - 0.1 \sin \frac{x}{2} \sin \frac{y}{2}$ , representing a profile of the secondary flow in the magnetohydrodynamic experiment [12].

The secondary flows will be constructed by nonlinear perturbation of the basic flow by using the eigenfunctions of the spectral problem

$$\lambda \Delta \psi = -\mu \Delta \psi + \nu \Delta^2 \psi + J(\psi_0, (2 + \Delta) \psi), \quad \psi = \sum_{n,m \geq 1} a_{n,m} \sin \frac{nx}{2} \sin \frac{my}{2}, \quad (7)$$

which is linearized from (3). The eigenfunctions will be studied in the following three linear orthogonal subspaces of  $H^4$ :

$$E_1 = \left\{ \psi \in H^4 \mid \psi = \sum_{n,m \geq 1; n,m \text{ odd}} a_{n,m} \sin \frac{nx}{2} \sin \frac{my}{2} \right\}, \quad (8)$$

$$E_2 = \left\{ \psi \in H^4 \mid \psi = \sum_{n,m \geq 1; n \text{ odd}; m \text{ even}} a_{n,m} \sin \frac{nx}{2} \sin \frac{my}{2} \right\}, \quad (9)$$

$$E_3 = \left\{ \psi \in H^4 \mid \psi = \sum_{n,m \geq 1; n \text{ even}; m \text{ odd}} a_{n,m} \sin \frac{nx}{2} \sin \frac{my}{2} \right\}. \quad (10)$$

The main result of the present paper reads as:

**Theorem 1.1.** (i). Let  $\nu > 0$ ,  $\mu \geq 0$  and  $\lambda + \mu + \frac{1}{2}\nu > 0$ . Then the spectral problem (7) has at most three linear independent eigenfunctions. These eigenfunctions are contained in the set  $E_1 \cup E_2 \cup E_3$ .

(ii) Assume that the spectral problem (7) admits a critical solution  $(\lambda, \psi, \nu, \mu) = (0, \psi_c, \nu_c, \mu_c)$  for  $\nu_c > 0$ ,  $\mu_c \geq 0$  and  $\psi_c \in E_{i_0}$  for  $1 \leq i_0 \leq 3$ . Then there exist a function  $\psi_{i_0} \in H^4$  and a real  $\delta$  so that the system (3)-(4) has a steady-state solution  $(\psi, \nu, \mu)$  branch off the bifurcation point  $(\psi_0, \nu_c, \mu_c)$  in the directions of  $\psi_c$ :

$$\psi = \psi_0 + \varepsilon \psi_c + \varepsilon^2 \psi_{i_0}, \quad \nu = \nu_c + \varepsilon \delta \nu_c, \quad \mu = \mu_c + \varepsilon \delta \mu_c, \quad (11)$$

provided that  $\varepsilon > 0$  is sufficiently small.

**Remark 1.1.** This theorem shows the secondary flow bifurcating in the direction of  $\psi_c$ . If  $\psi_c$  is replaced by the eigenfunction  $-\psi_c$ , we have another secondary flow bifurcating in the direction of  $-\psi_c$ .

This paper is structured as follows, The spectral analysis for the theoretical base of Theorem 1.1 is established in Section 2, which contains the proof Assertion (i) of Theorem 1.1. The second assertion of this theorem is proven in Section 3 by applying a bifurcation technique of Rabinowitz [10]. To enrich the theoretical result, we display numerical spectral solutions and use a finite difference scheme to locate a secondary flow in accordance with the experimental observation of [12, 14] in Section 4.

## 2. Linear spectral analysis

We begin with the spectral assertion of Theorem 1.1.

### 2.1. Proof of Theorem 1.1 (i)

*Proof.* Let  $(\cdot, \cdot)$  denotes the inner product of real  $L_2$  as

$$(\varphi, \phi) = \frac{1}{\pi^2} \int_0^{2\pi} \int_0^{2\pi} \varphi \phi dx dy.$$

Taking the  $L_2$  inner product of the spectral equation (7) with  $(\Delta + 2)\psi$  and employing the integration by parts, we have

$$\begin{aligned} 0 &= (-\lambda\Delta - \mu\Delta\psi + \nu\Delta^2\psi + J(\psi_0, (\Delta + 2)\psi), (\Delta + 2)\psi) \\ &= (-\lambda\Delta - \mu\Delta\psi + \nu\Delta^2\psi, (\Delta + 2)\psi). \end{aligned} \quad (12)$$

This together with (7) becomes

$$0 = \sum_{n,m \geq 1} \beta_{n,m}(\lambda + \mu + \nu\beta_{n,m})(\beta_{n,m} - 2)|a_{n,m}|^2 \quad \text{for } \beta_{n,m} = \frac{1}{4}(n^2 + m^2), \quad (13)$$

or

$$\begin{aligned} &\sum_{n,m \geq 1; \beta_{n,m} > 2} \beta_{n,m}(\lambda + \mu + \nu\beta_{n,m})(\beta_{n,m} - 2)|a_{n,m}|^2 \\ &= \beta_{1,1}(\lambda + \mu + \nu\beta_{1,1})(2 - \beta_{1,1})|a_{1,1}|^2 + \beta_{1,2}(\lambda + \mu + \nu\beta_{1,2})(2 - \beta_{1,2})|a_{1,2}|^2 \\ &\quad + \beta_{2,1}(\lambda + \mu + \nu\beta_{2,1})(2 - \beta_{2,1})|a_{2,1}|^2. \end{aligned} \quad (14)$$

Since  $\lambda + \mu + \nu\beta_{1,1} = \lambda + \mu + \frac{1}{2}\nu > 0$  and the eigenfunction  $\psi \neq 0$ , the right-side of (14) is positive. We may firstly assume the term involving  $a_{1,1}$  on the right-hand side of (14) being positive or  $a_{1,1} \neq 0$ ,

On the other hand, let  $a_{n,m} = 0$  whenever  $n \leq 0$  or  $m \leq 0$ . The spectral problem (7) is formulated as [4]

$$\begin{aligned} &\sum_{n,m \geq 1} \beta_{n,m}(\lambda + \mu + \nu\beta_{n,m})a_{n,m} \sin \frac{nx}{2} \sin \frac{my}{2} \\ &= - \sum_{n,m \geq -2}^{\infty} \left\{ \frac{n-m}{8} [(\beta_{n-2,m-2} - 2)a_{n-2,m-2} - (\beta_{n+2,m+2} - 2)a_{n+2,m+2}] \right. \\ &\quad \left. + \frac{n+m}{8} [(\beta_{n-2,m+2} - 2)a_{n-2,m+2} - (\beta_{n+2,m-2} - 2)a_{n+2,m-2}] \right\} \sin \frac{nx}{2} \sin \frac{my}{2}. \end{aligned} \quad (15)$$

This implies that the non-zero coefficient  $a_{1,1}$  generating the coefficients  $a_{n,m}$  for odd integers  $n, m \geq 1$ . That is, the eigenfunction  $\psi \in E_1$  and is generated by the mode  $\sin \frac{x}{2} \sin \frac{y}{2}$ . Moreover, the derivation of (14) implies that

$$\begin{aligned} &\beta_{1,1}(\lambda + \mu + \nu\beta_{1,1})(2 - \beta_{1,1})|a_{1,1}|^2 \\ &= \sum_{n,m \geq 1; n,m \text{ odd}; \beta_{n,m} > 2} \beta_{n,m}(\lambda + \mu + \nu\beta_{n,m})(\beta_{n,m} - 2)|a_{n,m}|^2. \end{aligned} \quad (16)$$

Similarly, we may suppose  $\psi \in E_2$  when  $a_{1,2} \neq 0$  and  $\psi \in E_3$  when  $a_{2,1} \neq 0$ . Additionally, the corresponding coefficients are subject to the equations

$$\begin{aligned} & \beta_{1,2}(\lambda + \mu + \nu\beta_{1,2})(2 - \beta_{1,2})|a_{1,2}|^2 \\ &= \sum_{n,m \geq 1; n \text{ odd}; m \text{ even}; \beta_{n,m} > 2} \beta_{n,m}(\lambda + \mu + \nu\beta_{n,m})(\beta_{n,m} - 2)|a_{n,m}|^2 \end{aligned} \quad (17)$$

for  $a_{1,2} \neq 0$ , and

$$\begin{aligned} & \beta_{2,1}(\lambda + \mu + \nu\beta_{2,1})(2 - \beta_{2,1})|a_{2,1}|^2 \\ &= \sum_{n,m \geq 1; n \text{ even}; m \text{ odd}; \beta_{n,m} > 2} \beta_{n,m}(\lambda + \mu + \nu\beta_{n,m})(\beta_{n,m} - 2)|a_{n,m}|^2 \end{aligned} \quad (18)$$

for  $a_{2,1} \neq 0$ .

This completes the proof of Assertion (i).  $\square$

## 2.2. Spectral simplicity property

To construct the secondary flows, we have to use the eigenfunction simplicity property shown in the following result.

**Theorem 2.1.** *Let  $\nu > 0$ ,  $\mu > 0$  and  $\lambda + \mu + \frac{1}{2}\nu > 0$ . Then we have eigenfunction space dimension estimate:*

$$\dim \left\{ \psi \in E_i \mid \lambda \Delta \psi = -\mu \Delta \psi + \nu \Delta^2 \psi + J(\psi_0, (\Delta + 2)\psi) \right\} \leq 1, \quad i = 1, 2, 3. \quad (19)$$

Moreover, if  $(\lambda, \psi, \nu, \mu)$  is a spectral solution of (7) for  $\psi \in E_1 \cup E_2 \cup E_3$ , then we have

$$((-\lambda \Delta - \mu \Delta + \nu \Delta^2)\psi, \psi^*) < 0. \quad (20)$$

Here  $\psi^*$  is the conjugate eigenfunction of  $\psi$  subject to the conjugate equation of (7):

$$\lambda \Delta \psi^* = -\mu \Delta \psi^* + \nu \Delta^2 \psi^* + (-\Delta - 2)J(\psi_0, \psi^*), \quad \psi^* = \sum_{n,m \geq 1}^{\infty} a_{n,m}^* \sin \frac{nx}{2} \sin \frac{my}{2} \quad (21)$$

produced by employing the  $L_2$  pairing  $(\cdot, \cdot)$ .

*Proof.* To show the validity of (19), we suppose that there is a spectral solution  $(\lambda, \psi, \nu, \mu)$  with the eigenfunction  $\psi \in E_1$ . To the contrary, if (19) with  $i = 1$  is not true, there exists an additional spectral solution  $(\lambda, \hat{\psi}, \nu, \mu)$  with the eigenfunction  $\hat{\psi} \in E_1$  linearly independent of  $\psi$  and involving expansion coefficients  $\hat{a}_{n,m}$ .

It follows from (16) that  $\hat{a}_{1,1} \neq 0$ . Therefore, we have the additional spectral solution  $(\lambda, \frac{\hat{a}_{1,1}}{a_{1,1}}\psi - \hat{\psi}, \nu, \mu)$ .

Using the eigenfunction  $\frac{\hat{a}_{1,1}}{a_{1,1}}\psi - \hat{\psi}$  instead of  $\psi$  in (16), we have

$$0 = \sum_{n,m \text{ odd}; \beta_{n,m} > 2} \beta_{n,m}(\lambda + \mu + \nu\beta_{n,m})(\beta_{n,m} - 2)|a_{n,m} - \frac{a_{1,1}}{\hat{a}_{1,1}}\hat{a}_{n,m}|^2, \quad (22)$$

which together with the condition  $\lambda + \mu + \nu\frac{1}{2} > 0$  gives

$$a_{n,m} = \frac{a_{1,1}}{\hat{a}_{1,1}}\hat{a}_{n,m} \quad \text{or} \quad \hat{\psi} = \frac{\hat{a}_{1,1}}{a_{1,1}}\psi_{1,1}.$$

This leads to a contraction and thus (19) holds true for  $i = 1$ .

Arguing in the same way, we obtain (19) for  $i = 2$  and 3.

To verify (20), we first assume the eigenfunction  $\psi \in E_1$ . Consider the conjugate spectral problem (21), which can be formulated in the algebraic equation

$$\sum_{n,m \geq -2; n,m \text{ odd}} \left\{ \beta_{n,m}(\lambda + \mu + \nu \beta_{n,m}) a_{n,m}^* \right. \\ \left. - (\beta_{n,m} - 2) \left\{ \frac{n-m}{8} (a_{n-2,m-2}^* - a_{n+2,m+2}^*) + \frac{n+m}{8} (a_{n-2,m+2}^* - a_{n+2,m-2}^*) \right\} \right\} \sin \frac{nx}{2} \sin \frac{my}{2}.$$

Here the assumption  $a_{n,m}^* = 0$  whenever  $n \leq 0$  or  $m \leq 0$  is used. The previous equation is rewritten as

$$0 = \sum_{n,m \geq -2; n,m \text{ odd}} \left\{ [(\lambda + \mu) \beta_{n,m} + \nu \beta_{n,m}^2] \frac{a_{n,m}^*}{\beta_{n,m} - 2} \right. \\ \left. - \frac{n-m}{8} (a_{n-2,m-2}^* - a_{n+2,m+2}^*) + \frac{n+m}{8} (a_{n-2,m+2}^* - a_{n+2,m-2}^*) \right\} \sin \frac{nx}{2} \sin \frac{my}{2}. \quad (23)$$

Moreover, for  $a'_{m,n} = \frac{a_{n,m}^*}{\beta_{n,m} - 2}$ , equation (23) becomes

$$0 = \sum_{n,m \geq -2; n,m \text{ odd}} \left\{ [(\lambda + \mu) \beta_{n,m} + \nu \beta_{n,m}^2] a'_{n,m} \right. \\ \left. + \frac{n-m}{8} [(\beta_{n-2,m-2} - 2) a'_{n-2,m-2} - (\beta_{n+2,m+2} - 2) a'_{n+2,m+2}] \right. \\ \left. + \frac{n+m}{8} [(\beta_{n-2,m+2} - 2) a'_{n-2,m+2} - (\beta_{n+2,m-2} - 2) a'_{n+2,m-2}] \right\} \sin \frac{mx}{2} \sin \frac{ny}{2}. \quad (24)$$

The algebraic equations for the coefficients of (24) are identical to those of (15), when the supercritical primes are omitted. Thus by (19), we have the relationship between the expansion coefficients of the eigenfunction  $\psi$  and those of its conjugate counterpart  $\psi^*$ :

$$a_{n,m}^* = (\beta_{m,n} - 2) a_{m,n}.$$

Hence, we have

$$(\Delta(-\lambda - \mu + \nu \Delta) \psi, \psi^*) = \sum_{n,m \geq 1; n,m \text{ odd}} \beta_{n,m}(\lambda + \mu + \nu \beta_{n,m})(\beta_{n,m} - 2) a_{n,m} a_{m,n}. \quad (25)$$

Note that  $\psi \neq 0$  implies  $a_{1,1} \neq 0$  due to (16). Moreover, from the algebraic equation defined by the first component of (15) with respect to the mode  $\sin \frac{x}{2} \sin \frac{y}{2}$ , it follows that the coefficient  $a_{1,1}$  is proportional to  $a_{1,3} - a_{3,1}$ . Hence  $a_{3,1} \neq a_{1,3}$  due to  $a_{1,1} \neq 0$ . Therefore the Cauchy inequality

$$2a_{1,3}a_{3,1} < a_{1,3}^2 + a_{3,1}^2 \quad (26)$$

holds true. It follows from (25), (26) and Cauchy inequality that

$$\begin{aligned} & (\Delta(-\lambda - \mu + \nu \Delta) \psi, \psi^*) \\ &= -\beta_{1,1}(\lambda + \mu + \nu \beta_{1,1})(2 - \beta_{1,1})a_{1,1}^2 + \sum_{n,m \geq 1; n,m \text{ odd}; \beta_{n,m} > 2} \beta_{n,m}(\lambda + \mu + \nu \beta_{n,m})(\beta_{n,m} - 2)a_{n,m}a_{m,n} \\ &< -\beta_{1,1}(\lambda + \mu + \nu \beta_{1,1})(2 - \beta_{1,1})a_{1,1}^2 + \sum_{n,m \geq 1; n,m \text{ odd}; \beta_{n,m} > 2} \beta_{n,m}(\lambda + \mu + \nu \beta_{n,m})(\beta_{n,m} - 2)a_{n,m}^2, \end{aligned}$$

which equals zero due to (16). This gives the validity of (20) when  $\psi \in E_1$ .

Moreover, if (7) has a spectral solution  $(\lambda, \psi, \nu, \mu)$  with the eigenfunction

$$\psi = \sum_{n,m \geq 1; n \text{ odd}; m \text{ even}} a_{n,m} \sin \frac{nx}{2} \sin \frac{my}{2} \in E_2,$$

the algebraic equation (15) for the coefficients  $a_{n,m}$  of  $\psi$  reduces to

$$\begin{aligned} 0 = & \sum_{n,m \geq -2; n \text{ odd}; m \text{ even}} \left\{ \beta_{n,m}(\lambda + \mu + \nu \beta_{n,m}) a_{n,m} \right. \\ & + \frac{n-m}{8} [(\beta_{n-2,m-2} - 2) a_{n-2,m-2} - (\beta_{n+2,m+2} - 2) a_{n+2,m+2}] \\ & \left. + \frac{n+m}{8} [(\beta_{n-2,m+2} - 2) a_{n-2,m+2} - (\beta_{n+2,m-2} - 2) a_{n+2,m-2}] \right\} \sin \frac{nx}{2} \sin \frac{my}{2}. \end{aligned} \quad (27)$$

Therefore, the conjugate spectral problem (21) has a solution  $(\lambda, \psi^*, \nu, \mu)$ , subject to the algebraic equation

$$\begin{aligned} 0 = & \sum_{n,m \geq -2; n \text{ odd}; m \text{ even}} \left\{ [(\lambda + \mu) \beta_{n,m} + \nu \beta_{n,m}^2] \frac{a_{n,m}^*}{\beta_{n,m} - 2} \right. \\ & \left. - \frac{n-m}{8} (a_{n-2,m-2}^* - a_{n+2,m+2}^*) + \frac{n+m}{8} (a_{n-2,m+2}^* - a_{n+2,m-2}^*) \right\} \sin \frac{nx}{2} \sin \frac{my}{2}. \end{aligned} \quad (28)$$

Here the assumption  $a_{n,m}^* = 0$  whenever  $n \leq 0$  or  $m \leq 0$  is used. For  $a'_{m,n} = \frac{a_{n,m}^*}{\beta_{n,m} - 2}$ , equation (28) becomes

$$\begin{aligned} 0 = & \sum_{n,m \geq -2; n \text{ even}; m \text{ odd}} \left\{ [(\lambda + \mu) \beta_{n,m} + \nu \beta_{n,m}^2] a'_{n,m} \right. \\ & + \frac{n-m}{8} [(\beta_{n-2,m-2} - 2) a'_{n-2,m-2} - (\beta_{n+2,m+2} - 2) a'_{n+2,m+2}] \\ & \left. + \frac{n+m}{8} [(\beta_{n-2,m+2} - 2) a'_{n-2,m+2} - (\beta_{n+2,m-2} - 2) a'_{n+2,m-2}] \right\} \sin \frac{mx}{2} \sin \frac{ny}{2}. \end{aligned} \quad (29)$$

This together with (19) implies that spectral problem (7) has a spectral solution  $(\lambda, \tilde{\psi}, \nu, \mu)$  with

$$\tilde{\psi} = \sum_{n,m \geq 1; n \text{ even}; m \text{ odd}} a_{n,m} \sin \frac{nx}{2} \sin \frac{my}{2} \in E_3 \quad (30)$$

subject to the equation

$$\begin{aligned} 0 = & \sum_{n,m \geq -2; n \text{ even}; m \text{ odd}} \left\{ [(\lambda + \mu) \beta_{n,m} + \nu \beta_{n,m}^2] a_{n,m} \right. \\ & + \frac{n-m}{8} [(\beta_{n-2,m-2} - 2) a_{n-2,m-2} - (\beta_{n+2,m+2} - 2) a_{n+2,m+2}] \\ & \left. + \frac{n+m}{8} [(\beta_{n-2,m+2} - 2) a_{n-2,m+2} - (\beta_{n+2,m-2} - 2) a_{n+2,m-2}] \right\} \sin \frac{mx}{2} \sin \frac{ny}{2}. \end{aligned} \quad (31)$$

Therefore, we have

$$a_{n,m}^* = (\beta_{n,m} - 2) a_{m,n}. \quad (32)$$

We show that  $a_{n,m} \not\equiv a_{m,n}$ . Otherwise, if  $a_{n,m} \equiv a_{m,n}$ , equation (31) becomes

$$\begin{aligned}
0 = & \sum_{n,m \geq -2; n \text{ odd}; m \text{ even}} \left\{ [(\lambda + \mu)\beta_{n,m} + \nu\beta_{n,m}^2]a_{n,m} \right. \\
& - \frac{n-m}{8} [(\beta_{n-2,m-2} - 2)a_{n-2,m-2} - (\beta_{n+2,m+2} - 2)a_{n+2,m+2}] \\
& \left. - \frac{n+m}{8} [(\beta_{n-2,m+2} - 2)a_{n-2,m+2} - (\beta_{n+2,m-2} - 2)a_{n+2,m-2}] \right\} \sin \frac{mx}{2} \sin \frac{ny}{2}.
\end{aligned} \tag{33}$$

Adding (33) to (27), we have

$$\sum_{n,m \geq 1; n \text{ odd}; m \text{ even}} [(\lambda + \mu)\beta_{n,m} + \nu\beta_{n,m}^2]a_{n,m} = 0$$

or  $a_{n,m} \equiv 0$ . This leads to a contradiction. Hence

$$a_{n,m} \not\equiv a_{m,n}. \tag{34}$$

By (32), we have

$$(\Delta(-\lambda - \mu + \nu\Delta)\psi, \psi^*) = \sum_{n,m \geq 1; n \text{ odd}; m \text{ even}} \beta_{n,m}(\lambda + \mu + \nu\beta_{n,m})(\beta_{n,m} - 2)a_{n,m}a_{m,n} \tag{35}$$

If  $a_{1,2} = a_{2,1}$ , we use (12), (34) and Cauchy inequality to obtain from (35) that

$$\begin{aligned}
& (\Delta(-\lambda - \mu + \nu\Delta)\psi, \psi^*) \\
& < \beta_{1,2}(\lambda + \mu + \nu\beta_{1,2})(\beta_{1,2} - 2)a_{1,2}^2 \\
& + \sum_{n,m \geq 1; n \text{ odd}; m \text{ even}; \beta_{n,m} > 2} \beta_{n,m}(\lambda + \mu + \nu\beta_{n,m})(\beta_{n,m} - 2) \frac{a_{n,m}^2 + a_{m,n}^2}{2} \\
& = \beta_{1,2}(\lambda + \mu + \nu\beta_{1,2})(\beta_{1,2} - 2) \frac{a_{1,2}^2 + a_{2,1}^2}{2} \\
& + \frac{1}{2} \sum_{n,m \geq 1; n \text{ odd}; m \text{ even}; \beta_{n,m} > 2} \beta_{n,m}(\lambda + \mu + \nu\beta_{n,m})(\beta_{n,m} - 2)a_{n,m}^2 \\
& + \frac{1}{2} \sum_{n,m \geq 1; n \text{ even}; m \text{ odd}; \beta_{n,m} > 2} \beta_{n,m}(\lambda + \mu + \nu\beta_{n,m})(\beta_{n,m} - 2)a_{n,m}^2
\end{aligned}$$

which equals zero due to (17) and (18). This gives (20) under the condition  $a_{1,2} = a_{2,1}$ .

Actually, we can assume  $a_{1,2} = a_{2,1} = 1$ , since the spectral problem is linear. This gives (20) for the eigenfunction  $\psi \in E_2$ .

This derivation also implies the validity of (20) when the eigenfunction  $\psi \in E_3$ .

The proof of Theorem 2.1 is completed. □

### 3. Assertion (ii) of Theorem 1.1

*Proof.* Firstly, we introduce function spaces so that Fredholm alternative theory can be applied for the critical eigenfunction  $\psi_c$ .



Theorem 2.1 shows that  $\psi_c \in E_{i_0}$  is a simple eigenfunction. That is,

$$\dim \left\{ \psi \in E_{i_0} \mid 0 = -\mu_c \Delta \psi + \nu_c \Delta^2 \psi + J(\psi_0, (\Delta + 2)\psi) \right\} = 1$$

and (20) holds true.

$E_{i_0}$  is an invariant space of the linear spectral problem. Now we are considering steady-state bifurcation in the nonlinear Navier-Stokes equation system. The principal part of the bifurcating solution is the perturbation of the linear eigenfunction  $\psi_c$ . Therefore, we need to consider the bifurcation in the nonlinear flow invariant space of the Navier-Stokes equation and the space is generated from the linear space  $E_{i_0}$ . To do so, we use the summation notation

$$\begin{aligned} \sum_1 &= \sum_{1 \leq n, m \text{ odd}} + \sum_{2 \leq n, m \text{ even}}, \\ \sum_2 &= \sum_{1 \leq n \text{ odd}; 2 \leq m \text{ even}} + \sum_{2 \leq n, m \text{ even}}, \\ \sum_3 &= \sum_{2 \leq n \text{ even}; 1 \leq m \text{ odd}} + \sum_{2 \leq n, m \text{ even}}. \end{aligned}$$

Now we define Hilbert subspace of  $H^4$  as follows

$$H_i^4 = \left\{ \psi \in H^4 \mid \psi = \sum_i a_{n,m} \sin \frac{nx}{2} \sin \frac{my}{2} \right\} \text{ for } i = 1, 2, 3.$$

We also need the  $L_2$  subspaces

$$H_i = \left\{ \psi = \sum_i a_{n,m} \sin \frac{nx}{2} \sin \frac{my}{2} \mid \|\psi\|_{L_2} = \left( \sum_i a_{n,m}^2 \right)^{\frac{1}{2}} < \infty \right\} \text{ for } i = 1, 2, 3.$$

This definition ensures  $H_i^4 \supset E_i$  for  $i = 1, 2, 3$  and  $H_i^4$  is orthogonal to  $E_j$  if  $i \neq j$ . Hence the assertion of Theorem 2.1 remains valid when  $E_{i_0}$  is replaced by  $H_{i_0}^4$ . That is, the eigenfunction simplicity property holds true in  $H_{i_0}^4$ . The nonlinear flow invariant property of  $H_{i_0}^4$  is valid in the following sense

$$\Delta^{-2} J(\phi, \Delta \phi) \in H_{i_0}^4 \text{ whenever } \phi, \phi \in H_{i_0}^4. \quad (36)$$

Now we rewrite the critical spectral problem as

$$\mathcal{L} \psi_c = 0 \text{ for } \mathcal{L} \psi = -\mu_c \Delta \psi + \nu_c \Delta^2 \psi + J(\psi_0, (\Delta + 2)\psi).$$

We see that  $\mathcal{L}$  maps  $H_{i_0}^4$  into  $H_{i_0}$ . To employ the Fredholm theory, we define the range of  $\mathcal{L}$  as

$$\text{Ran}(\mathcal{L}) = \left\{ \phi \in H_{i_0} \mid \text{there exists } \psi \in H_{i_0}^4 \text{ so that } \mathcal{L} \psi = \phi \right\}.$$

It readily seen that  $\text{Ran}(\mathcal{L})$  is the space orthogonal to  $\psi_c^*$ , the conjugate eigenfunction of  $\psi_c$ , in the following sense:

$$\text{Ran}(\mathcal{L}) = \left\{ \psi \in H_{i_0} \mid (\psi, \psi_c^*) = 0 \right\}.$$

By the Fredholm alternative theory of Laplacian operators,  $\mathcal{L}$  has an inverse operator

$$\mathcal{L}^{-1} : \text{Ran}(\mathcal{L}) \mapsto H_{i_0}^4 \quad (37)$$

so that

$$\|\mathcal{L}^{-1}\psi\|_{H_{i_0}^4} \leq C_1 \|\psi\|_{L_2}, \quad \psi \in \text{Ran}(\mathcal{L}) \quad (38)$$

for a constant  $C_1$ .

Secondly, following Rabinowitz [10] on the Bénard problem, we seek the secondary steady-state solution  $(\psi, v, \mu)$  branching from the bifurcation point  $(\psi_0, v_c, \mu_c)$  in the direction of  $\psi_c$  as

$$\psi = \psi_0 + \varepsilon \psi_c + \varepsilon^2 \psi_{i_0}, \quad v = v_c + \varepsilon \sigma v_c, \quad \mu = \mu_c + \varepsilon \sigma \mu_c \quad (39)$$

for a function  $\psi_{i_0} \in H_{i_0}^4$  and a real  $\sigma$ , provided that  $\varepsilon > 0$  is sufficiently small.

Substitution of the predicted solution (39) into the stationary form of (3), or the equation

$$\begin{aligned} 0 &= (-\mu\Delta + v\Delta^2)(\psi - \psi_0) + J(\psi_0, (2 + \Delta)(\psi - \psi_0)) + J(\psi - \psi_0, \Delta(\psi - \psi_0)) \\ &= (-(\mu - \mu_c)\Delta + (v - v_c)\Delta^2)(\psi - \psi_0) + \mathcal{L}(\psi - \psi_0) + J(\psi - \psi_0, \Delta(\psi - \psi_0)), \end{aligned}$$

produces the equation

$$0 = (-\varepsilon\sigma\mu_c\Delta + \varepsilon\sigma v_c\Delta^2)(\varepsilon\psi_c + \varepsilon^2\psi_{i_0}) + \mathcal{L}(\varepsilon\psi_c + \varepsilon^2\psi_{i_0}) + J(\varepsilon\psi_c + \varepsilon^2\psi_{i_0}, \Delta(\varepsilon\psi_c + \varepsilon^2\psi_{i_0})).$$

Since  $\mathcal{L}\psi_c = 0$ , the previous equation can be rewritten as

$$\sigma(-\mu_c\Delta + v_c\Delta^2)\psi_c + \mathcal{L}\psi_{i_0} = F_\varepsilon(\sigma, \psi_{i_0}) \quad (40)$$

with

$$F_\varepsilon(\sigma, \psi_{i_0}) = -\varepsilon\sigma(-\mu_c\Delta + v_c\Delta^2)\psi_{i_0} - J(\psi_c + \varepsilon\psi_{i_0}, \Delta(\psi_c + \varepsilon\psi_{i_0})).$$

To show the existence of the unknowns  $\psi_{i_0}$  and  $\sigma$ , we take  $L_2$  inner product of (40) with  $\psi_c^*$  to obtain

$$\sigma((- \mu_c\Delta + v_c\Delta^2)\psi_c, \psi_c^*) + (\mathcal{L}\psi_{i_0}, \psi_c^*) = (F_\varepsilon(\sigma, \psi_{i_0}), \psi_c^*). \quad (41)$$

Applying Theorem 2.1 with  $E_{i_0}$  replaced by  $H_{i_0}^4$ , the invariance property (36) and the identity

$$(\mathcal{L}\psi_{i_0}, \psi_c^*) = (\psi_{i_0}, \mathcal{L}^*\psi_c^*) = 0$$

we may rewrite (41) as

$$\sigma = \frac{(F_\varepsilon(\sigma, \psi_{i_0}), \psi_c^*)}{((- \mu_c\Delta + v_c\Delta^2)\psi_c, \psi_c^*)}. \quad (42)$$

The combination of (40) and (42) yields

$$\mathcal{L}\psi_{i_0} = F_\varepsilon(\sigma, \psi_{i_0}) - \frac{(-\mu_c\Delta + v_c\Delta^2)\psi_c(F_\varepsilon(\sigma, \psi_{i_0}), \psi_c^*)}{((- \mu_c\Delta + v_c\Delta^2)\psi_c, \psi_c^*)}. \quad (43)$$

The nonlinear invariance property (36) implies  $F_\varepsilon(\sigma, \psi_{i_0}) \in H_{i_0}$ . It is readily see that the right-hand side of (43) is in  $\text{Ran}(\mathcal{L})$ . Therefore, we may use the inverse of  $\mathcal{L}$  to produce

$$\psi_{i_0} = \mathcal{L}^{-1}\left(F_\varepsilon(\sigma, \psi_{i_0}) - \frac{(-\mu_c\Delta + v_c\Delta^2)\psi_c(F_\varepsilon(\sigma, \psi_{i_0}), \psi_c^*)}{((- \mu_c\Delta + v_c\Delta^2)\psi_c, \psi_c^*)}\right). \quad (44)$$

For simplicity of notation, we rewrite the equations (42) and (44) in the following form

$$(\sigma, \psi_{i_0}) = G_\varepsilon(\sigma, \psi_{i_0}), \quad (45)$$

where the two components of the operator  $G_\varepsilon(\sigma, \psi_{i_0})$  represent respectively the right-hand sides of (42) and (44). Thus, to seek the solution  $(\psi, \nu, \mu)$  in (39) becomes to confirm the existence of the fixed point for the operator  $G_\varepsilon$ .

Finally, it remains to prove that  $G_\varepsilon$  is a contraction operator mapping a complete metric space into itself. The complete metric space is defined as

$$X = \left\{ (\sigma, \psi) \in (-\infty, \infty) \times H_{i_0}^4 \mid \|(\sigma, \psi)\|_X = |\sigma| + \|\psi\|_{H_{i_0}^4} \leq C \right\}$$

Here  $C > 0$  is a constant to be defined afterward.

To show the contraction property, we use the boundedness of  $\mathcal{L}^{-1}$  in (38), the expressions (42) and (44), and the Hölder inequality to produce

$$\|G_\varepsilon(\sigma, \psi_{i_0})\|_X \leq \left( \frac{\|\psi_c^*\|_{L_2}}{|((-\mu_c\Delta + \nu_c\Delta^2)\psi_c, \psi_c^*)|} + C_1 \left( 1 + \frac{\|(-\mu_c\Delta + \nu_c\Delta^2)\psi_c\|_{L_2}\|\psi_c^*\|_{L_2}}{|((-\mu_c\Delta + \nu_c\Delta^2)\psi_c, \psi_c^*)|} \right) \right) \|F_\varepsilon(\sigma, \psi_{i_0})\|_{L_2}.$$

This yields, by renaming the constant bounded by the large brackets in the right-hand side of the previous equation as  $C_2$ ,

$$\|G_\varepsilon(\sigma, \psi_{i_0})\|_X \leq C_2 \|F_\varepsilon(\sigma, \psi_{i_0})\|_{L_2}.$$

Hence, by the Hölder inequality and the Sobolev imbedding inequality, we have

$$\begin{aligned} \|G_\varepsilon(\sigma, \psi_{i_0})\|_X &\leq C_2 \|J(\psi_c + \varepsilon\psi_{i_0}, \Delta(\psi_c + \varepsilon\psi_{i_0})) + \varepsilon\sigma(-\mu_c\Delta + \nu_c\Delta^2)\psi_{i_0}\|_{L_2} \\ &\leq C_2 \left( \|\nabla(\psi_c + \varepsilon\psi_{i_0})\|_{L_4} \|\nabla\Delta(\psi_c + \varepsilon\psi_{i_0})\|_{L_4} + \varepsilon\sigma\|(-\mu_c\Delta + \nu_c\Delta^2)\psi_{i_0}\|_{L_2} \right) \\ &\leq C_3 \left( \|\psi_c\|_{H_{i_0}^4}^2 + 2\varepsilon\|\psi_c\|_{H_{i_0}^4}\|\psi_{i_0}\|_{H_{i_0}^4} + \varepsilon^2\|\psi_{i_0}\|_{H_{i_0}^4}^2 + \varepsilon\sigma\|\psi_{i_0}\|_{H_{i_0}^4} \right) \\ &\leq C_4 \left( 1 + \varepsilon C + \varepsilon^2 C^2 + \varepsilon C^2 \right) \end{aligned}$$

for constants  $C_k$  independent of  $(\sigma, \psi) \in X$  and  $\varepsilon > 0$ . Therefore, we obtain

$$\|G_\varepsilon(\sigma, \psi_{i_0})\|_X \leq C \text{ for } (\sigma, \psi_{i_0}) \in X, \quad (46)$$

provided that

$$\frac{C}{2} = C_4$$

and

$$C_4(\varepsilon C + \varepsilon^2 C^2 + \varepsilon C^2) = C_4(\varepsilon + 2\varepsilon^2 C_4 + 2\varepsilon C_4)C \leq \frac{1}{2}C,$$

by taking  $\varepsilon > 0$  sufficiently small.

The property (46) implies the injection property  $G_\varepsilon : X \mapsto X$ .

Arguing in the same manner, we have the contraction property:

$$\|G_\varepsilon(\sigma, \psi_{i_0}) - G_\varepsilon(\sigma', \psi'_{i_0})\|_X \leq \frac{1}{2} \|(\sigma, \psi_{i_0}) - (\sigma', \psi'_{i_0})\|_X$$

for  $(\sigma, \psi_{i_0}), (\sigma', \psi'_{i_0}) \in X$ , provided that  $\varepsilon > 0$  sufficiently small.

Therefore, by the Banach contraction mapping principle, the operator  $G_\varepsilon$  with small  $\varepsilon > 0$  admits a unique fixed point  $(\sigma, \psi_{i_0}) \in X$ . This confirms the existence of the steady-state solution  $(\psi, \mu, \nu)$  of (3) and (4) in the form of (39).

The proof of Theorem 1.1 is completed. □

#### 4. Numerical understanding of Theorem 1.1

Theorem 1.1 shows that secondary steady-state flows are essentially defined by the linear critical spectral solution  $(\psi_c, v_c, \mu_c)$  of (7) with  $\lambda = 0$ , and at most three linear independent critical eigenfunctions are available. They are belong to respectively the three orthogonal spaces  $E_1, E_2$  and  $E_3$ . We thus begin with the critical spectral solutions.

The critical vector value  $(v_c, \mu_c)$  with respect to  $\psi_c \in E_1$  was obtained numerically in [13, Table I] and form a curve joining the points  $(0.2371, 0)$  and  $(0, 0.2307)$  in Figure 2 (a). This is the neutral line separating the linear stable and linear unstable domains. However, for the eigenfunction  $\psi_c \in E_2$ , the proof of Theorem 2.1 shows the coexistence of two critical orthogonal eigenfunctions. The another one  $\tilde{\psi}_c \in E_3$  is given by (30). The numerical simulation of the two eigenfunctions sharing with the same critical vector values  $(v_c, \mu_c)$  was given in [4]. These critical vector values  $(v_c, \mu_c)$  form the line touching the points  $(0.0415, 0)$  and  $(0, 0.01515)$  inside the linear unstable domain displayed in Figure 2(a).




Figure 2: (a) Critical vector values  $(v_c, \mu_c)$  when  $\psi_c \in E_1$  (the long curve) and when  $\psi_c \in E_2 \cup E_3$  (the short curve); (b) critical eigenfunction  $\psi_c \in E_1$  for  $(v_c, \mu_c) = (0.00054, \mu = 0.2315)$  or  $(Re_c, Rh_c) = (22446, 1.326)$ ; (c) the linear approximate secondary flow  $\psi = \psi_0 - \frac{0.1}{a_{1,1}} \psi_c$  for  $(v_c, \mu_c) = (0.00054, \mu = 0.2315)$ ; (d) the nonlinear secondary flow at  $(v, \mu) = (0.0005, 0.23)$  obtained by a finite difference scheme.

To understand the experimental magnetohydrodynamic flows, we follow [11, 12, 14] to consider almost inviscid flows

so that their energy dissipation is essentially controlled by the Hartmann layer friction  $\mu$  or the Rayleigh number  $Rh$ . When the critical eigenfunction  $\psi_c \in E_2 \cup E_3$ , the secondary flows branching from the corresponding critical vector values in the linear unstable domain is inobservable in laboratory experiments, although they are contributed to the complexity of flow dynamic behaviour towards to turbulence. We only consider the flows related to  $(\psi_c, v_c, \mu_c)$  with  $\psi_c \in E_1$ .

Therefore, we display the critical eigenfunction  $\psi_c \in E_1$  in Figure 2(b) for  $(v_c, \mu_c) = (0.00054, \mu = 0.2315)$  or  $(Re_c, Rh_c) = (22446, 1.326)$ . By Theorem 1.1, we may approximate the secondary flow by employ the linear perturbation

$$\psi \approx \psi_0 - \frac{0.1}{a_{1,1}} \psi_c \text{ for } (v_c, \mu_c) = (0.00054, \mu = 0.2315), \psi_c = \sum_{1 \leq n, m \text{ odd}} a_{n,m} \sin \frac{nx}{2} \sin \frac{my}{2}.$$

This approximation flow is displayed in Figure 2 (c).

The secondary flow from the spectral solution  $(\psi_c, v_c, \mu_c)$  is actually locally stable and can be approximated by finite difference method. We employ a finite difference scheme with a  $80 \times 80$  gridding mesh of the fluid domain  $\Omega$  to approximate numerically the bifurcating flows. Numerical secondary solutions are obtained for  $(v, \mu)$  in a vicinity of the critical condition  $(\mu_c, v_c) = (0.00054, 0.2315)$  or  $(Re_c, Rh_c) = (22446, 1.326)$ . In Figure 2(d), we present nonlinear secondary steady-state flow at  $(v, \mu) = (0.0005, 0.23)$  or  $(Re, Rh) = (24158, 1.33)$ , which represents the secondary flow bifurcating from  $\psi_0$  at  $(\mu_c, v_c) = (0.00054, 0.2315)$ .

The secondary flow in Figure 2(d) shows the topological transition for the merging of two vortices, observed by Sommaria and Verron [12, 14]. Their experimental threshold for the onset of secondary flow is  $(Re_c, Rh_c) = (22700, 1.52)$ , which is close to but higher than the present numeric one  $(Re_c, Rh_c) = (22446, 1.326)$ . This is due to neglect of the energy dissipation inside the lateral boundary layers of the original three-dimensional fluid motion problem (see [13]).

When  $(v, \mu)$  is close to the threshold  $(v_c, \mu_c)$ , the nonlinear secondary flow in Figure 2(d) is comparable with the linearized secondary flow in Figure 2 (c). What is more, it is very close to the truncation form

$$\psi \approx \psi_0 - 0.1 \sin \frac{x}{2} \sin \frac{y}{2}$$

expressed in Figure 1 (b). This is owing to the mode  $\sin \frac{x}{2} \sin \frac{y}{2}$  generating the eigenfunction  $\psi_c$ . By numerical computation and (16), the principal coefficient  $a_{1,1}$  of the principal mode is significantly larger than other coefficients  $a_{n,m}$ .

## References

- [1] N.F. Bondarenko, M.Z. Gak, F.V. Dolzhanskii, Laboratory and theoretical models of a plane periodic flow, *Izv. Akad. Nauk SSSR, Fiz. Atmos. Okeana*, **15** (1979), 1017-1026.
- [2] C. Cerretelli, C.H.K. Williamson, The physical mechanism for vortex merging, *J. Fluid Mech.* **475** (2003), 41-77.
- [3] Z.M. Chen, Bifurcating steady-state solutions of the dissipative quasi-geostrophic equation in Lagrangian formulation, *Nonlinearity* **29** (2016), 3132-3147.
- [4] Z.M. Chen, Instability of two-dimensional square eddy flows, *Phys. Fluids* **31**, 044107 (2019).
- [5] Z.M. Chen, W.G. Price, Supercritical regimes of liquid-metal fluid motions in electromagnetic fields: wall-bounded flows, *Proc. R. Soc. Lond. A* **458** (2002), 2735-2757.
- [6] Z.M. Chen, W.G. Price, Secondary fluid flows driven electromagnetically in a two-dimensional extended duct, *Proc. R. Soc. A* **461** (2005), 1659-1683.
- [7] V.I. Iudovich, Example of the generation of a secondary stationary or periodic flow when there is loss of stability of the laminar flow of a viscous incompressible fluid, *J. Appl. Math. Mech.* **29** (1965), 527-544.
- [8] R.H. Kraichnan, Inertial ranges in two-dimensional turbulence, *Phys. Fluids* **10**, 1417 (1967).

- [9] P. Meunier, S. Le Dizès, T. Leweke, Physics of vortex merging, *Comptes Rendus Physique* **6** (2005), 431-450.
- [10] P.H. Rabinowitz, Existence and nonuniqueness of rectangular solutions of the Bénard problem, *Arch. Rational Mech. Anal.* **29** (1968), 32-57.
- [11] J. Sommeria, Experimental study of the two-dimensional inverse energy cascade in a square box, *J. Fluid Mech.* **170** (1986), 139-168.
- [12] J. Sommeria, J. Verron, An investigation of nonlinear interactions in a two-dimensional recirculating flow, *Phys. Fluids* **27**, 1918 (1984).
- [13] A. Thess, Instabilities in two-dimensional spatially periodic flows. Part II: Square eddy lattice, *Phys. Fluids A* **4**, 1396 (1992).
- [14] J. Verron, J. Sommeria, Numerical simulation of a two-dimensional turbulence experiment in magnetohydrodynamics, *Phys. Fluids* **30**, 732 (1987).