

Existence of exponentially spatially localised breather solutions for lattices of nonlinearly coupled particles: Schauder's fixed point approach

Dirk Hennig and Nikos I. Karachalios

Department of Mathematics, University of Thessaly, Lamia GR 35100, Greece

Abstract: We prove the existence of exponentially localised breather solutions for lattices of coupled particles with nonlinear nearest neighbour interaction. The second derivative of the associated interaction potentials is bounded by a power-law growth. The existence problem is formulated in terms of a fixed point equation considered in weighted sequence spaces, which is solved by means of Schauder's Fixed Point Theorem.

I. INTRODUCTION

Non-uniform structures in nonlinear lattice systems, in the form of spatially localised and time-periodically varying states referred to as breather solutions, have attracted considerable interest from the physical but also from the mathematical viewpoint. Representative rigorous results regarding proofs of existence and stability of discrete breathers are provided in [1]-[18]: several methods and techniques were used, such as continuation of (trivially localised) solutions from the anti-integrable of zero couplings for sufficiently weak couplings, center manifold reduction methods, Lyapunov-Schmidt decomposition and continuum approximation techniques, variational approaches, perturbative arguments for the characteristic exponents of the Floquet monodromy matrices, normal form theorems and Nekroroshev-type proofs.

In this work, we present a comparatively concise and to our knowledge, novel approach compared to the aforementioned methods, for the proof of existence of breathers for a class of lattices with nonlinear interaction between its constituents based on Schauder's fixed point theorem [19]. We consider systems with even potentials whose second derivative is bounded by a power-law growth. A prominent example is the β -Fermi-Pasta-Ulam (FPU) system. Other important examples include the periodic and the bounded soft potential. We use methods from [20]. The approach we present herein, establishes, apart from existence, directly a requested degree of spatial localisation [21], [22], [23]. That is, the breathers are supposed to be exponentially localised as it is the case in physically significant applications (see [24], [25] and references therein).

The presentation of the paper is as follows. In section II we discuss the model systems and main assumptions on the interaction potential. Section III contains the description and properties of the functional setting essentially involving weighted sequence spaces and the proof of the main result. We think that the whole approach can be of general interest: Firstly, it implements a Fourier series representation of the solutions to establish compactness embedding properties of the considered functional spaces; these can be useful generically for the analysis of lattice dynamical systems (existence and dynamics, see [26],[27] and references therein). Secondly, facilitating the Schauder's fixed point approach in this set-up gives rise to physically relevant and consistent restrictions on the characteristics of the breather solutions, namely, their frequency (natural non-resonant conditions) and their magnitude measured in the suitable weighted norm ("energy" upper-bounds); importantly, the latter exhibits dependence on the localisation degree (as dictated by the relevant weighted norm) and the growth of the nonlinearity for the general class of potentials considered herein (see Theorem III.2 and Remark III.1). Finally, section IV summarizes the main findings and give an outlook to extensions for future studies.

II. DESCRIPTION OF THE SYSTEM

The system of interacting particles on infinite lattices is given by the following set of coupled equations

$$\frac{d^2 q_n}{dt^2} = -V'(q_n - q_{n-1}) + V'(q_{n+1} - q_n), \quad n \in \mathbb{Z}, \quad (2.1)$$

where q_n is the displacement of a particle from its equilibrium position. The function $V(x)$ is the interaction potential between nearest neighbours and the prime $'$ stands for the derivative with respect to the argument x . This system has a Hamiltonian structure related to the energy

$$H = \sum_{n \in \mathbb{Z}} \left(\frac{1}{2} p_n^2 + V(q_n - q_{n-1}) \right),$$

and it is time-reversible with respect to the involution $p \mapsto -p$. In addition to the energy $H = E$, the total momentum $P = \sum_n p_n$ is conserved too.

Discrete breathers can be characterised as follows:

$$q_n(t+T) = q_n(t), \quad p_n(t+T) = p_n(t), \quad n \in \mathbb{Z}, \quad (2.2)$$

$$\lim_{n \rightarrow \pm\infty} q_n = c_{\pm}, \quad \lim_{n \rightarrow \pm\infty} p_n = 0, \quad (2.3)$$

and the difference $c_+ - c_-$ is the lattice deformation induced by the breather. It is not a loss of generality to set $c_- = 0$. Furthermore, $q_n(t)$ has zero time average, i.e.

$$\int_0^T q_n(t) dt = 0, \quad n \in \mathbb{Z}. \quad (2.4)$$

The existence theorem will be proved under the following assumption:

- **A:** The interaction potential $V(x)$ has a global minimum at $x = 0$ and is of the form

$$V(x) = \frac{1}{2}x^2 + W(x), \quad (2.5)$$

with $W : \mathbb{R} \rightarrow \mathbb{R}$ being even, $W(x) = W(-x)$, and $W(0) = W'(0) = 0$. Further, we assume that W is non-negative and at least twice continuously differentiable and satisfies for some constant $\bar{K} > 0$ and $\alpha \geq 0$, such that

$$|W''(x)| \leq \bar{K}|x|^\alpha, \quad \forall x \in \mathbb{R}. \quad (2.6)$$

With the parity symmetry of the interaction potential V it can readily be seen that in our case $c_+ = 0$.

Examples of interaction potentials which satisfy condition **A** are given by the β -FPU system with $W(x) = \beta x^4/4$, the periodic potential $W(x) = 1 - \cos x$ [24] and the soft bounded potential $W(x) = 1 - \text{sech}^2 x$, [25],[28],[29].

The solutions of the system obtained when linearising equations (2.1) around the equilibrium $q_n = 0$, are superpositions of plane wave solutions (phonons)

$$q_n(t) = \exp(i(kn - \omega t)),$$

with frequencies

$$\omega^2(k) = 4 \sin^2 \left(\frac{k}{2} \right), \quad k \in [-\pi, \pi].$$

These (extended) states disperse. Therefore, the frequency Ω of a localised time-periodic solution must satisfy the non-resonance condition $\Omega \neq |\omega(k)|/m$ for any integer $m \geq 1$. This requires $\Omega^2 > 4$ as a necessary condition for the existence of localised time-periodic solutions of system (2.1).

For the forthcoming analysis, it is convenient to pass to relative variables defined by

$$x_n = q_n - q_{n-1}, \quad (2.7)$$

$$p_n = y_n - y_{n+1}, \quad (2.8)$$

where the original variables satisfy

$$\sum_{n \in \mathbb{Z}} |p_n| < \infty, \quad \sum_{n \in \mathbb{Z}} |q_n - q_{n-1}| < \infty. \quad (2.9)$$

which imply that

$$\sum_{n \in \mathbb{Z}} |y_n - y_{n+1}| < \infty, \quad \sum_{n \in \mathbb{Z}} |x_n| < \infty. \quad (2.10)$$

From (2.3), it follows that $\lim_{n \rightarrow -\infty} y_n = 0$ so that the change of variables (2.7)-(2.8) is invertible. That is, for given $\{(x_n, y_n)\}_{n \in \mathbb{Z}}$, one gets the relations

$$p_n = y_n - y_{n+1}, \quad q_n = \sum_{l=-\infty}^n x_l. \quad (2.11)$$

The transformation (2.7)-(2.8) is symplectic, because

$$\sum_n dp_n \wedge dq_n = \sum_n (dy_n \wedge dq_{n-1} + dy_n \wedge dx_n - dy_{n+1} \wedge dq_n) = \sum_n dy_n \wedge dx_n, \quad (2.12)$$

and from the Hamiltonian

$$H = \sum_n \left(\frac{1}{2} (y_n - y_{n+1})^2 + V(x_n) \right), \quad (2.13)$$

one derives the canonical equations

$$\dot{x}_n = 2y_n - y_{n+1} - y_{n-1}, \quad \dot{y}_n = -V'(x_n), \quad (2.14)$$

or equivalently

$$\ddot{x}_n = V'(x_{n+1}) + V'(x_{n-1}) - 2V'(x_n). \quad (2.15)$$

In (2.12) the symbol \wedge denotes the standard wedge product. The relations (2.2)-(2.4), satisfied by breather solutions of the original system (2.1), when passing to the new variables, read as

$$\begin{aligned} x_n(t+T) &= x_n(t), \quad y_n(t+T) = y_n(t), \quad n \in \mathbb{Z} \\ \lim_{n \rightarrow \pm\infty} x_n &= 0, \quad \lim_{n \rightarrow \pm\infty} y_n = 0, \\ \int_0^T x_n(t) dt &= 0, \quad n \in \mathbb{Z}. \end{aligned}$$

III. EXISTENCE OF EXPONENTIALLY LOCALISED BREATHERS

In the following we prove the existence of localised periodic solutions of system (2.1) on the infinite one-dimensional lattice. To this end, some appropriate function spaces are introduced on which the original problem is presented as a fixed point problem for a corresponding operator. Utilising Schauder's Fixed Point Theorem, we establish the existence of exponentially localised solutions, extending the approach followed in [20], to a suitable functional spaces-setup.

To establish the required spatial localisation of the solutions, we introduce suitable weighted function spaces. First we consider the exponentially weighted Hilbert space of square-summable sequences, $l_w^2(\mathbb{Z})$, defined as

$$l_w^2 = \left\{ u_n \in \mathbb{R} : \|u\|_{l_w^2}^2 := \sum_n w_n |u_n|^2 \right\}, \quad (3.1)$$

with $w_n = \exp(\lambda|n|)$ and $\lambda \geq 0$. Then, we denote by

$$X_0 = \left\{ u \in L_{per}^2([0, T]; l_w^2) : \int_0^T u_n(t) dt = 0, \quad n \in \mathbb{Z} \right\},$$

the space of T -periodic square integrable functions in time with zero time average, with values in l_w^2 . Evidently, X_0 is a closed convex subspace of $L_{per}^2([0, T]; l_w^2)$. We also consider the Sobolev space

$$X_2 = \left\{ u \in H_{per}^2([0, T]; l_w^2) \right\},$$

containing T -periodic functions of time assuming values in l_w^2 which, together with their weak derivatives up to second order are in X_0 . The above spaces are endowed with the following norms:

$$\begin{aligned} \|u\|_{X_0}^2 &= \frac{1}{T} \int_0^T \|u(t)\|_{l_w^2}^2 dt, \\ \|u\|_{X_2}^2 &= \frac{1}{T} \int_0^T \left(\|u(t)\|_{l_w^2}^2 + \|Du(t)\|_{l_w^2}^2 + \|D^2u(t)\|_{l_w^2}^2 \right) dt. \end{aligned}$$

For an element $u \in X_2$, we may consider the Fourier-series expansion of $u_n(t)$ with respect to time t and space variable n , determined by

$$u_n(t) = \sum_{m \in \mathbb{Z} \setminus \{0\}} \hat{u}_{n,m} \exp(i\Omega m t), \quad t \in [0, T], \quad \hat{u}_{n,m} = \frac{1}{T} \int_0^T u_n(t) \exp(-i\Omega m t) dt, \quad (3.2)$$

$$\hat{u}_{n,m} = \overline{\hat{u}_{n,-m}}, \quad (3.3)$$

and

$$u_n(t) = \frac{1}{2\pi} \int_0^{2\pi} \tilde{u}_k(t) \exp(ikn) dk, \quad \tilde{u}_k(t) = \sum_n u_n(t) \exp(-ikn), \quad (3.4)$$

$$\tilde{u}_k = \overline{\tilde{u}_{-k}}. \quad (3.5)$$

Then, using (3.2)-(3.5), we may have the representation of u , given by

$$u_n(t) = \sum_{m \in \mathbb{Z} \setminus \{0\}} \frac{1}{2\pi} \int_0^{2\pi} \hat{u}_{k,m} \exp(ikn) dk \exp(i\Omega m t). \quad (3.6)$$

The following compactness lemma will be particularly useful for our approach.

Lemma III.1 X_2 is compactly embedded in X_0 (denoted as $X_2 \Subset X_0$).

Proof: The continuous embedding $X_2 \subseteq X_0$ is clear due to the inequality

$$\|u\|_{X_0} \leq \|u\|_{X_2}.$$

Now, in order to show the compactness of the embedding, let $(u_n)_{n \in \mathbb{N}} \in X_2$ be a bounded sequence, i.e. $\|u_n\|_{X_2} \leq M < \infty$ for all $n \in \mathbb{N}$. Then, there exists a subsequence (not relabeled) $u_n \rightharpoonup u \in X_2$ weakly. The continuous embedding $X_2 \subseteq X_0$ implies the weak convergence $u_n \rightharpoonup u$ in X_0 . For the strong convergence $u_n \rightarrow u$ in X_0 , considering the difference $\|u_n - u\|_{X_0}^2$, we have the estimate:

$$\begin{aligned} \|u_n - u\|_{X_0}^2 &= \frac{1}{T} \int_0^T \|u_n(t) - u(t)\|_{l_w^2}^2 dt \\ &= \frac{1}{T} \int_0^T \sum_{k \in \mathbb{Z}} w_k |u_n^k(t) - u^k(t)|^2 dt \\ &= \frac{1}{T} \int_0^T \sum_{|k| \leq K} w_k |u_n^k(t) - u^k(t)|^2 dt + \frac{1}{T} \int_0^T \sum_{|k| > K} w_k |u_n^k(t) - u^k(t)|^2 dt \\ &= \frac{1}{T} \int_0^T \sum_{|k| \leq K} w_k |u_n^k(t) - u^k(t)|^2 dt + \sum_{|k| > K} w_k \sum_{m \neq 0} |\hat{u}_{n,m}^k - \hat{u}_m^k|^2 \\ &\leq \frac{1}{T} \int_0^T \sum_{|k| \leq K} w_k |u_n^k(t) - u^k(t)|^2 dt + \sum_{|k| > K} w_k \sum_{m \neq 0} |\hat{u}_{n,m}^k|^2 + \sum_{|k| > K} w_k \sum_{m \neq 0} |\hat{u}_m^k|^2 \\ &\leq \frac{1}{T} \int_0^T \sum_{|k| \leq K} w_k |u_n^k(t) - u^k(t)|^2 dt + \sum_{|k| > K} w_k |u_n^k|^2 \sum_{m \neq 0} \frac{1}{m^4} + \sum_{|k| > K} w_k |u^k|^2 \sum_{m \neq 0} \frac{1}{m^4} \\ &= \frac{1}{T} \int_0^T \sum_{|k| \leq K} w_k |u_n^k(t) - u^k(t)|^2 dt + 2\zeta(4) \left(\sum_{|k| > K} w_k |u_n^k|^2 + \sum_{|k| > K} w_k |u^k|^2 \right), \end{aligned} \quad (3.7)$$

where $\zeta(s) = \sum_{m=1}^{\infty} m^{-s}$ is the Riemann zeta function, $(u_n^k)_{k \in I} \in l_w^2$, $(u^k)_{k \in I} \in l_w^2$, $I = (-K, \dots, 0, \dots, K)$. We also used that since $u \in H_{per}^2[0, T] \Rightarrow |\hat{u}_m| \leq C/|m|^2$. Then for sufficiently large K we get

$$2\zeta(4) \left(\sum_{|k| > K} w_k |v_n^k|^2 + \sum_{|k| > K} w_k |v^k|^2 \right) \leq \frac{2\epsilon}{3}. \quad (3.8)$$

Exploiting that weak and strong convergence coincide in finite-dimensional spaces, by choosing n sufficiently large, we get for the first term on the right-hand side of (3.7), that

$$\frac{1}{T} \int_0^T \sum_{|k| \leq K} w_k |u_n^k(t) - u^k(t)|^2 dt \leq \frac{\epsilon}{3}. \quad (3.9)$$

Combining (3.8) and (3.9) we get

$$\|u_n - u\|_{X_0}^2 \leq \epsilon, \quad (3.10)$$

and the proof is finished. \square

Let us also recall the two versions of the Schauder's fixed point Theorem [19].

Theorem III.1 1. (First version): Let G be a closed bounded convex subset of a Banach space X . Assume that $f : G \mapsto G$ is compact. Then, f has at least one fixed point in G .

2. (Second version): Let G be a closed convex subset of a Banach space X and f a continuous map of G into a compact subset of G . Then, f has at least one fixed point.

With the preparations above, we may proceed to the statement and proof of the main result.

Theorem III.2 Let condition **A** hold and suppose

$$\Omega^2 > 4. \quad (3.11)$$

Then there exist nonzero sequences $x \equiv \{x_n\}_{n \in \mathbb{Z}} \in H^2([0, T]; l^2)$ with zero time average $\int_0^T x_n(t) dt = 0$ and $\|x\|_{X_0} \leq R$, where

$$R \leq \left(\frac{\Omega^2 - 4}{\overline{K} (1 + \Omega^2) \sqrt{2(1 + \cosh(\lambda))}} \right)^{1/\alpha}. \quad (3.12)$$

These solutions are exponentially localised time-periodic solutions of system (2.15) with period $T = \frac{2\pi}{\Omega}$.

Proof: We shall provide two alternatives of the proof by implementing the two versions of the Schauder's fixed point Theorem III.1. For this purpose, it is convenient to rewrite Eq. (2.15) as:

$$\ddot{x}_n - [x_{n+1} - 2x_n + x_{n-1}] = W'(x_{n+1}) + W'(x_{n-1}) - 2W'(x_n). \quad (3.13)$$

Thus, only the right-hand side of (3.13) features terms nonlinear in x . We shall express (3.13) as a fixed point equation in x .

I. 1st version of the proof: We relate the left-hand side of (3.13) to the linear mapping: $M : X_2 \rightarrow X_0$:

$$M(u_n) = \ddot{u}_n - [u_{n+1} - 2u_n + u_{n-1}].$$

Then, applying the operator M to the Fourier elements $\exp(ikn) \exp(i\Omega mt)$ in the representation (3.6), we get that

$$M \exp(ikn) \exp(i\Omega mt) = \nu_m(k) \exp(ikn) \exp(i\Omega mt),$$

where

$$\nu_m(k) = -\Omega^2 m^2 + 4 \sin^2 \left(\frac{k}{2} \right), \quad m \in \mathbb{Z} \setminus \{0\}, \quad k \in [0, 2\pi].$$

As by assumption (3.11) $\Omega^2 > 4$, it is guaranteed that $\nu_m(k) \neq 0$, for all $m \in \mathbb{Z} \setminus \{0\}$ and for all $k \in [0, 2\pi]$, the mapping M possesses an inverse M^{-1} , obeying $M^{-1} \exp(i(\Omega mt + kn)) = (1/\nu_m(k)) \exp(i(\Omega mt + kn))$. For the norm

of the linear operator $M^{-1} : X_0 \rightarrow X_2$ one derives the upper bound:

$$\begin{aligned}
\|M^{-1}\|_{X_0, X_2}^2 &= \sup_{\|u\|_{X_0}=1} \|M^{-1}u\|_{X_2}^2 \\
&= \sup_{\|u\|_{X_0}=1} \frac{1}{T} \int_0^T \left[\|M^{-1}u(t)\|_{L_w^2}^2 + \|DM^{-1}u(t)\|_{L_w^2}^2 + \|D^2M^{-1}u(t)\|_{L_w^2}^2 \right] dt \\
&= \sup_{\|u\|_{X_0}=1} \frac{1}{T} \int_0^T \sum_{n \in \mathbb{Z}} w_n \left(\left| \sum_{m'} \frac{1}{2\pi} \int_0^{2\pi} \frac{\hat{u}_{k,m}}{\nu_m(k)} \exp(ikn) \exp(i\Omega m t) dk \right|^2 \right. \\
&\quad + \left| \sum_{m'} \frac{1}{2\pi} \int_0^{2\pi} \frac{i\Omega m \hat{u}_{k,m}}{\nu_m(k)} \exp(ikn) \exp(i\Omega m t) dk \right|^2 \\
&\quad + \left. \left| \sum_{m'} \frac{1}{2\pi} \int_0^{2\pi} \frac{(i\Omega m)^2 \hat{u}_{k,m}}{\nu_m(k)} \exp(ikn) \exp(i\Omega m t) dk \right|^2 \right) dt \\
&\leq \sup_{m \in \mathbb{Z} \setminus \{0\}} \sup_{k \in [0, 2\pi]} \frac{1 + (\Omega m)^2 + (\Omega m)^4}{|\nu_m(k)|^2} \\
&\quad \cdot \sup_{\|u\|_{X_0}=1} \frac{1}{T} \int_0^T \sum_{n \in \mathbb{Z}} w_n \left| \sum_{m'} \frac{1}{2\pi} \int_0^{2\pi} \hat{u}_{k,m} \exp(ikn) \exp(i\Omega m t) dk \right|^2 dt \\
&\leq \frac{1 + \Omega^2 + \Omega^4}{(\Omega^2 - 4)^2} \sup_{\|u\|_{X_0}=1} \|u\|_{X_0}^2 = \frac{1 + \Omega^2 + \Omega^4}{(\Omega^2 - 4)^2} \leq \frac{(1 + \Omega^2)^2}{(\Omega^2 - 4)^2} < \infty, \tag{3.14}
\end{aligned}$$

verifying the boundedness of M^{-1} and we used the notation $\sum_{l'} = \sum_{l \in \mathbb{Z} \setminus \{0\}}$.

In order to deal with the nonlinear terms, we attribute to the right-hand side of (3.13) the nonlinear operator $N : X_0 \rightarrow X_0$, as

$$(N(u))_n = W'(u_{n+1}) + W'(u_{n-1}) - 2W'(u_n).$$

To establish that the operator N is continuous on X_0 , we prove that N is Frechet differentiable at any u , with bounded derivative. We observe that

$$N'(u) : h \in X_0 \mapsto N'(u)[h] = W''(u)h \in X_0,$$

where $(W''(u))_n = W''(u_{n+1}) + W''(u_{n-1}) - 2W''(u_n)$. Therefore, we derive the estimate

$$\begin{aligned}
\|N'(u)[h]\|_{X_0}^2 &= \frac{1}{T} \int_0^T \sum_{n \in \mathbb{Z}} w_n |(W''(u_{n+1}(t)) + W''(u_{n-1}(t)) - 2W''(u_n(t))) h_n(t)|^2 dt \\
&\leq \frac{1}{T} \int_0^T \sum_{n \in \mathbb{Z}} w_n \bar{K}^2 (|u_{n+1}(t)|^{2\alpha} + |u_{n-1}(t)|^{2\alpha} + 2|u_n(t)|^{2\alpha}) |h_n(t)|^2 dt \\
&\leq 4\bar{K}^2 \sup_{n \in \mathbb{Z}} \max_{t \in [0, T]} |u_n(t)|^{2\alpha} \|h\|_{X_0}^2 = M^2 \|h\|_{X_0}^2,
\end{aligned}$$

where $M^2 = 4\bar{K}^2 \sup_{n \in \mathbb{Z}} \max_{t \in [0, T]} |u_n(t)|^{2\alpha}$. That is

$$\|N'(u)\|_{\mathcal{L}(X_0, X_0)} \leq M, \tag{3.15}$$

implying the (uniform) boundedness of the differential. Let us now use as the closed convex subset Y_0 of X_0 , its closed ball centered at 0 of radius R ,

$$Y_0 = \{u \in X_0 : \|u\|_{X_0} \leq R\}.$$

For the range of N on Y_0 , we get the bound

$$\begin{aligned}
\|N(u)\|_{X_0}^2 &= \int_0^T \sum_{n \in \mathbb{Z}} w_n |W'(u_{n+1}(t)) + W'(u_{n-1}(t)) - 2W'(u_n(t))|^2 dt \\
&\leq \int_0^T \sum_{n \in \mathbb{Z}} w_n (|W'(u_{n+1}(t))|^2 + |W'(u_{n-1}(t))|^2 + 2|W'(u_n(t))|^2) dt \\
&\leq \overline{K}^2 \int_0^T \sum_{n \in \mathbb{Z}} \left(\frac{w_n}{w_{n+1}} w_{n+1} |u_{n+1}(t)|^{2(\alpha+1)} + \frac{w_n}{w_{n-1}} w_{n-1} |u_{n-1}(t)|^{2(\alpha+1)} + 2w_n |u_n(t)|^{2(\alpha+1)} \right) dt \\
&\leq 2\overline{K}^2 (1 + \cosh(\lambda)) \|u\|_{\tilde{X}_0}^{2(\alpha+1)} \leq 2\overline{K}^2 (1 + \cosh(\lambda)) \|u\|_{X_0}^{2(\alpha+1)} \\
&\leq 2\overline{K}^2 (1 + \cosh(\lambda)) R^{2(\alpha+1)}, \quad \forall u \in Y_0,
\end{aligned} \tag{3.16}$$

where $\tilde{X}_0 = \{u \in L_{per}^2([0, T]; l_w^{2(\alpha+1)}) : \int_0^T u_n(t) dt = 0, n \in \mathbb{Z}\}$ and we used $X_0 \subset \tilde{X}_0$.

In order to apply Theorem III.1, the final step is to express the problem (3.13) as a fixed point equation in terms of a mapping $Y_0 \rightarrow Y_0$:

$$x = M^{-1} \circ N(x) \equiv S(x). \tag{3.17}$$

Clearly, S is continuous on X_0 , as the relations (3.14) and (3.15) establish that its constituents M^{-1} and N are continuous. Next, we show that $S(Y_0) \subseteq Y_0$. Using (3.14) and (3.16), we have

$$\begin{aligned}
\|S(x)\|_{X_0} &= \|M^{-1}(N(x))\|_{X_0} \leq \|M^{-1}\|_{X_0, X_2} \cdot \|N(x)\|_{X_0} \\
&\leq \sqrt{2(1 + \cosh(\lambda))} \overline{K} \frac{1 + \Omega^2}{\Omega^2 - 4} R^{\alpha+1} \leq R, \quad \forall x \in Y^0,
\end{aligned}$$

assuring by assumption (3.12), that indeed

$$S(Y_0) \subseteq Y_0.$$

As M^{-1} maps X_0 to $X_2 \Subset X_0$, one has $S(Y_0) \subseteq Y_0 \cap X_2$. Therefore the map $S = M^{-1} \circ N$, viewed as a map $S : Y_0 \subseteq X_0 \mapsto Y_0 \subseteq X_0$, is compact. The first version of Schauder's fixed point theorem implies then that the fixed point equation $x = S(x)$ has at least one solution.

II. 2nd version of the proof: For the application of the second version of Schauder's fixed point theorem, we verify that the range of S is contained in a compact subset of Y_0 . Representing $N(u) \in Y_0$ in terms of its spatial and temporal Fourier-transforms as

$$(N(u))_n(t) = \sum_{m \in \mathbb{Z} \setminus \{0\}} \frac{1}{2\pi} \int_0^{2\pi} \hat{N}_{k,m} \exp(ikn) dk \exp(i\Omega mt), \tag{3.18}$$

the Fourier coefficients of $M^{-1}(N(u))$ fulfill for all $u \in Y_0$, for all $k \in [0, 2\pi]$, and for all $m \in \mathbb{Z} \setminus \{0\}$

$$\begin{aligned}
\left| \frac{\hat{N}_{k,m}}{-\Omega^2 m^2 + 4 \sin^2\left(\frac{k}{2}\right)} \right|^2 &\leq \frac{\sum_{m \in \mathbb{Z} \setminus \{0\}} \frac{1}{2\pi} \int_0^{2\pi} |\hat{N}_{k,m}|^2 dk}{m^4 (\Omega^2 - 4)^2} \\
&= \frac{\|N(u)\|_{X_{0,0}}^2}{m^4 (\Omega^2 - 4)^2},
\end{aligned} \tag{3.19}$$

where $X_{0,0}$ is determined by

$$X_{0,0} = \left\{ u \in L_{per}^2([0, T]; l^2) \mid \int_0^T u_n(t) dt = 0, n \in \mathbb{Z} \right\}.$$

Since $l_w^2 \subset l^2$ and the inequality $\|u\|_{l^2} \leq \|u\|_{l_w^2}$ holds, we get

$$\left| \frac{\hat{N}_{k,m}}{-\Omega^2 m^2 + 4 \sin^2\left(\frac{k}{2}\right)} \right|^2 \leq \frac{2\overline{K}^2 (1 + \cosh(\lambda)) R^{2(\alpha+1)}}{m^4 (\Omega^2 - 4)^2} \leq \left(\frac{R}{m^2 (1 + \Omega^2)} \right)^2. \tag{3.20}$$

Hence, we conclude that S maps Y_0 into the subset

$$Y_{0,c} = \left\{ u = \{u\}_{n \in \mathbb{Z}} \in Y_0, u_n(t) = \sum_{m \in \mathbb{Z} \setminus \{0\}} \frac{1}{2\pi} \int_0^{2\pi} \hat{u}_{k,m} \exp(ikn) dk \exp(i\Omega mt) \mid |\hat{u}_{k,m}| \leq \frac{R}{m^2(1+\Omega^2)} \right\},$$

which is compact in Y_0 . That is, the operator S maps closed convex subsets $Y_0 \subset X_0 \subset L^2_{per}((0, T); l^2)$ into compact subsets $Y_{0,c}$ of Y_0 . The second version of Schauder's fixed point theorem implies then that the fixed point equation $u = S(u)$ has at least one solution.

We conclude the proof with the following details: According to the (invertible) transformation (2.7)-(2.8), breather solutions to system (2.15) represent breather solutions to the original system (2.1), satisfying the relations (2.2)-(2.3).

Moreover, by the hypothesis (3.11) the values of the frequency of oscillations Ω lie above the upper edge of the continuous (phonon) spectrum determined by 2. Thus the corresponding solutions have to be anharmonic which necessitates nonzero nonlinearity. Only the latter facilitates amplitude-depending tuning of the frequency of oscillations so that the breather frequency Ω lies outside of the phonon spectrum. Thus, it must hold that $\|N(u)\|_{X_0} \neq 0$, which, by condition **A**, is satisfied if and only if $u \neq 0$. That is, the fixed point equation (3.17) possesses a nonzero solution and the proof is finished.

Remark III.1 1. Since the obtained time-periodic H^2 fixed-point-solutions are by Sobolev embeddings C^1 in time and since the operator $x \mapsto W'(x)$ maps C^1 into itself, one concludes from Eq. (3.13) that $\tilde{x} \in C^1$ are classical solutions.

2. The localised solutions on the infinite lattice \mathbb{Z} are represented by (infinite) square-summable sequences, i.e. exponential decay of the solutions for $|n| \rightarrow \infty$ takes place in the sense of the exponentially weighted l^2 norm. Notably, for weight function $w_n \equiv 1$, i.e. $\lambda = 0$, our proof establishes the existence of general localised patterns (e.g. multi-site breathers).
3. From (3.12) we infer that the amplitude of the breathers go to zero as their frequency approaches the upper edge of the phonon band, i.e. $\Omega \rightarrow 2^+$.

IV. CONCLUSIONS

In summary, we have proved the existence of non-trivial, exponentially localised breather solutions for general systems of nonlinearly coupled particles with even interaction potentials, on the infinite lattice \mathbb{Z} . The existence problem has been reformulated as a fixed point equation (involving the relevant linear and nonlinear operators associated to the system), on weighted sequence spaces. This fixed point problem has been solved via the Schauder's Fixed Point Theorem. Future extensions may consider systems with other physically important classes of interaction potentials, Discrete Nonlinear Schrödinger models and different degrees of localisation such as algebraic, which can be relevant for the existence of discrete rational solutions [30],[31]. Such investigations are in progress and will be documented elsewhere.

-
- [1] R.S. MacKay and S. Aubry, *Proof of existence of breathers for time-reversible or Hamiltonian networks of weakly coupled oscillators*, Nonlinearity **7** (1994), 1623-1643.
 - [2] D. Bambusi, *Exponential stability of breathers in Hamiltonian networks of weakly coupled oscillators*, Nonlinearity **9** (1996), 433-457.
 - [3] G. Friesecke and J. A. D. Wattis, *Existence theorem for travelling waves on lattices*, Comm. Math. Phys. **161** (1994), 391-418.
 - [4] M. Willem and D. Smets, *Solitary waves with prescribed speed on lattices*, J. Funct. Anal. **149** (1997), 266-275.
 - [5] D. Bambusi, *Asymptotic stability of breathers in some Hamiltonian networks of weakly coupled oscillators*, Comm. Math. Phys. **324** (2013), 515-547.
 - [6] D. Bambusi, S. Paleari and T. Penati, *Existence and continuous approximation of small amplitude breathers in 1D and 2D Klein-Gordon lattices*, Appl. Anal. **89** (2010), 1313-1334.
 - [7] J. Cuevas, J. C. Eilbeck and N. I. Karachalios, *Thresholds for breather solutions of the discrete nonlinear Schrödinger equation with saturable and power nonlinearity*, Discrete Contin. Dyn. Syst. **21** (2008), 445-475.
 - [8] A.I. Komech, E.A. Kopylova and M. Kunze, *Dispersive estimates for 1D discrete Schrödinger and Klein-Gordon equations*, Appl. Anal. **85** (2006), 1487-1508.

- [9] J. Cuevas, N.I. Karachalios and F. Palmero, *Energy thresholds for the existence of breather solutions and travelling waves on lattices*, Appl. Anal. **89** (2010), 1351-1385.
- [10] S. Paleari and T. Penati, *Long time stability of small amplitude breathers in a mixed FPU-KG model*, ZAMP, **67** (2016), 148, 21 pp.
- [11] D. Pelinovsky and A. Sakovich, *Multi-site breathers in Klein-Gordon lattices: stability, resonances and bifurcations*, Nonlinearity **25** (2012), 3423-3451.
- [12] M. Weinstein, *Excitation thresholds for nonlinear localized modes on lattices*, Nonlinearity **12** (1999), 673-691.
- [13] V. Koukouloyannis and P.G. Kevrekidis, *On the stability of multibreathers in Klein-Gordon chains*, Nonlinearity **22** (2009), 2269-2285.
- [14] R. Livi, M. Spicci and R.S. MacKay, *Breathers on a diatomic FPU chain*, Nonlinearity **10** (1997), 1421-1434.
- [15] G. James, *Existence of breathers on FPU lattices*, C.R. Acad. Sci. Paris, **332** (2001), 581-586.
- [16] G. Iooss and G. James, *Localized waves in nonlinear oscillator chains*, Chaos **15** (2005), 015113, 15 pp.
- [17] N. Karachalios, *A remark on the existence of breather solutions for the discrete nonlinear Schrödinger equation in infinite lattices: The case of site-depenent anharmonic parameters*, Proc. Edinb. Math. Soc. (2) **49** (2006), 115-129.
- [18] R. Mirollo and N. Rosen, *Existence, uniqueness, and non-uniqueness of single-wave-form solutions to Josephson junction systems*, SIAM J. Appl. Math. **60** (2000), 1471-1501.
- [19] J. Schauder, *Der Fixpunktsatz in Funktionalräumen*, Studia Math. **2** (1930), 171-180.
- [20] D. Hennig, *Localised time-periodic solutions of discrete nonlinear Klein-Gordon systems with convex on-site potentials*, J. Fixed Point Theory Appl. **23** (2021), 31-38.
- [21] A. Pankov, *Gap solitons in periodic discrete nonlinear Schrödinger equations*, Nonlinearity **19** (2006), 27-40.
- [22] G. Zhang *Breather solutions of the discrete nonlinear Schrödinger equations with unbounded potentials*, J. Math. Phys. **50** (2009), 013105, 12pp.
- [23] A. Pankov, *Solitary waves on nonlocal Fermi-Pasta-Ulam lattices: Exponential localization*, Nonlinear Analysis: Real World Applications **50** (2019) 603-612.
- [24] O.V. Gendelman and A. V. Savin, *Normal Heat Conductivity of the One-Dimensional Lattice with Periodic Potential of Nearest-Neighbor Interaction*, Phys. Rev. Lett **84** (2000), 2381, 5pp.
- [25] S. Lepri, R. Livi and A. Politi *Thermal conduction in classical low-dimensional lattices*, Phys. Rep. **377**, 1-80.
- [26] N. I. Karachalios and A. N. Yannacopoulos, *Global existence and compact attractors for the discrete nonlinear Schrödinger equation*, J. Differential Equations **217** (2005), 88-123.
- [27] D. Hennig and N. I. Karachalios, *Dynamics of nonlocal and local discrete Ginzburg-Landau equations: global attractors and their congruence* (2021) preprint, <https://arxiv.org/abs/2104.00338>.
- [28] J. A. D. Wattis, A. Pickering and P. R. Gordoa, *Combined breathing-kink modes in the FPU lattice*, Phys. D **240** (2011), 547-553.
- [29] G. Arioli and H. Koch, *Traveling wave solutions for the FPU chain: a constructive approach*, Nonlinearity **33** (2020), 1705-1722.
- [30] A. Ankiewicz, N. Akhmediev, and J. M. Soto-Crespo, *Discrete rogue waves of the Ablowitz-Ladik and Hirota equations*, Phys. Rev. E **82** (2010), 026602.
- [31] N. Akhmediev and A. Ankiewicz, *Modulation instability, Fermi-Pasta-Ulam recurrence, rogue waves, nonlinear phase shift, and exact solutions of the Ablowitz-Ladik equation*, Phys. Rev. E **83** (2011), 046603.